

Crystal Interpretation of Kerov-Kirillov-Reshetikhin Bijection

Atsuo Kuniba, Masato Okado, Reiho Sakamoto,
Taichiro Takagi and Yasuhiko Yamada

ABSTRACT: The Kerov-Kirillov-Reshetikhin (KKR) bijection is the crux as in proving fermionic formulas. It is defined by a combinatorial algorithm on rigged configurations and highest paths. We reformulate the KKR bijection as a vertex operator by purely using combinatorial R in crystal base theory. The result is viewed as a nested Bethe ansatz at $q = 0$ as well as the direct and the inverse scattering (Gel'fand-Levitan) map in the associated soliton cellular automaton.

1. INTRODUCTION

Among many approaches to quantum integrable systems, Bethe ansatz [1] stands as a most efficient tool. In the context of solvable lattice models [2], it produces eigenvectors of transfer matrices from solutions of Bethe equations. Beside exact evaluation of physical quantities, Bethe ansatz has brought a number of applications also in representation theory and combinatorics. A prominent example is the fermionic formula [3, 4], which grew out of the completeness problem, a certain counting of Bethe vectors under string hypotheses. With the advent of the corner transfer matrix [2] and the crystal base theory [5], the fermionic formulas have been reformulated as the so called $X = M$ conjecture for any affine Lie algebra [6, 7]. See [8] for the current status of the conjecture.

The proof of the fermionic formula for $A_n^{(1)}$ [3, 4] may be viewed as a combinatorial version of Bethe ansatz. As a substitute of solutions to Bethe equations, the combinatorial object called *rigged configuration* (RC) is introduced. It is an n -tuple of Young diagrams (configuration) each row of which is assigned with an integer (rigging) obeying a selection rule. The Bethe vectors are replaced by Littlewood-Richardson tableaux, or equivalently, *highest paths*. The latter are A_n highest weight elements in $B_{\mu_1} \otimes \cdots \otimes B_{\mu_m}$, where B_l is the $A_n^{(1)}$ crystal of the l -fold symmetric tensor representation corresponding to a ‘‘local spin’’. Under these setting, Bethe ansatz should produce highest paths from rigged configurations and vice versa. What achieves this and thereby proves the fermionic formula is the celebrated Kerov-Kirillov-Reshetikhin (KKR) bijection. It is defined by a purely combinatorial algorithm on rigged configurations and highest paths, whose meaning however has remained rather mysterious.

The purpose of this paper is to clarify the representation theoretical origin of the KKR bijection in the light of crystal base theory and associated soliton cellular automata [9]. Let p be the highest path that corresponds to a rigged configuration under the KKR bijection. Our main Theorem 2.2 asserts that

$$(1.1) \quad p = \Phi_1 \mathcal{C}_1 \Phi_2 \mathcal{C}_2 \cdots \Phi_n \mathcal{C}_n (p^{(n)}).$$

Here $p^{(n)}$ is a trivial vacuum path and $\Phi_1 \mathcal{C}_1 \Phi_2 \mathcal{C}_2 \cdots \Phi_n \mathcal{C}_n$ is a vertex operator in the sense explained below. Recall that the $A_n^{(1)}$ crystal B_l consists of length l row semistandard tableaux with letters $1, 2, \dots, n+1$. As a set the affine crystal is given by $\text{Aff}(B_l) = \{b[d] \mid b \in B_l, d \in \mathbb{Z}\}$ by assigning the mode d to B_l . The isomorphism $\text{Aff}(B_l) \otimes \text{Aff}(B_m) \simeq \text{Aff}(B_m) \otimes \text{Aff}(B_l)$ is called the *combinatorial R* [10, 11]. We will

We show that under time evolutions the configuration remains unchanged while the rigging flows linearly. In this sense the rigged configurations are action-angle variables of the associated soliton cellular automaton. The first Young diagram $\mu^{(1)}$ in the configuration gives the list of amplitudes of solitons. For example, $t = 4$ state in the above pattern coincides with p in Example 2.9 (apart from redundant 1's) for which $\mu^{(1)} = (4, 3, 1)$ indeed. The meaning of the other Young diagrams is understood similarly along the nested family of cellular automata explained in Section 2.7.

Finally to interpret the formula (1.1), write it as $p = \Phi_1 \mathcal{C}_1(p^{(1)})$. In the above example, one has $p^{(1)} = \boxed{2222} \otimes \boxed{233} \otimes \boxed{4}$, and this is nothing but the list of incoming solitons. The map \mathcal{C}_1 assigns $p^{(1)}$ with the modes in affine crystals that encode the positions of solitons. Thus $\mathcal{C}_1(p^{(1)})$ is the *scattering data* of the soliton cellular automaton. Then Φ_1 plays the role of a vertex operator to create solitons over the “vacuum path” $111\dots$ by injecting the scattering data $\mathcal{C}_1(p^{(1)})$ by using combinatorial R . Again these constructions work inductively along the nested family of crystals for $A_n^{(1)} \supset A_{n-1}^{(1)} \supset \dots \supset A_0^{(1)}$. Thus the composition $\Phi_1 \mathcal{C}_1 \dots \Phi_n \mathcal{C}_n$ is the inverse scattering map that reproduces solitons from the scattering data.

The layout of the paper is as follows. In Section 2, we treat the $A_n^{(1)}$ case. A proof of Theorem 2.2 will be presented in [17]. Section 2.7 includes the solution of the direct scattering problem, namely, a method to determine the rigged configuration from a given highest path by only using combinatorial R and the KKR bijection for \widehat{sl}_2 . It is a crystal theoretical separation of variables. In Section 3, conjectures similar to (1.1) are presented for the bijection in the other non-exceptional affine algebras [18, 19]. They all possess a similar feature to the $A_n^{(1)}$ case, but the items as Φ_a, \mathcal{C}_a need individual descriptions. We do this from Sections 4 to 8. Appendix A is a brief exposition of the crystal base theory. Further applications of the present results to fermionic formulas and soliton cellular automata will be given elsewhere.

2. $A_n^{(1)}$ CASE

2.1. Rigged configurations. Consider the data of the form

$$(2.1) \quad (\mu^{(0)}, (\mu^{(1)}, J^{(1)}), \dots, (\mu^{(n)}, J^{(n)})),$$

where $\mu^{(a)} = (\mu_1^{(a)}, \dots, \mu_{l_a}^{(a)})$ is a partition and $J^{(a)} = (J_1^{(a)}, \dots, J_{l_a}^{(a)}) \in (\mathbb{Z}_{\geq 0})^{l_a}$. Set

$$(2.2) \quad E_j^{(a)} = \sum_{i=1}^{l_a} \min(j, \mu_i^{(a)}).$$

Define the vacancy numbers by

$$(2.3) \quad p_j^{(a)} = E_j^{(a-1)} - 2E_j^{(a)} + E_j^{(a+1)} \quad (1 \leq a \leq n),$$

where $E_j^{(n+1)} = 0$. The data (2.1) is called a rigged configuration if the following conditions are satisfied for all $1 \leq a \leq n$ and $j \in \mathbb{Z}_{>0}$:

$$(2.4) \quad 0 \leq J_i^{(a)} \leq J_{i+1}^{(a)} \leq \dots \leq J_l^{(a)} \leq p_j^{(a)} \quad \text{if } \{i, i+1, \dots, l\} = \{k \mid \mu_k^{(a)} = j\}.$$

The array of partitions $\mu^{(0)}, \dots, \mu^{(n)}$ is called a configuration and the nonnegative integers $J_i^{(a)}$ are called rigging. If $\mu^{(a)}$ is represented as a Young diagram, the vacancy number $p_j^{(a)}$ is assigned to each “cliff” of width j . $J^{(a)}$ may be viewed as assigning

to the cliff a partition whose parts are at most $p_j^{(a)}$. Here is an example of the $A_3^{(1)}$ rigged configuration.

$$(2.5) \quad \begin{array}{cccc} \mu^{(0)} & \mu^{(1)} & \mu^{(2)} & \mu^{(3)} \\ \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \\ \hline \square & & \\ \hline \square & & \\ \hline \square & & \\ \hline \square & & \\ \hline \square & & \\ \hline \square & & \\ \hline \end{array} & \begin{array}{|c|c|c|} \hline \square & \square & 0 \\ \hline \square & \square & 0 \\ \hline \square & & 0 \\ \hline 1 & \square & 1 \\ \hline \end{array} & \begin{array}{|c|c|c|} \hline 1 & \square & 1 \\ \hline \square & \square & \\ \hline \square & & \\ \hline \square & & \\ \hline \square & & \\ \hline \square & & \\ \hline \square & & \\ \hline \square & & \\ \hline \end{array} & \begin{array}{|c|c|} \hline 0 & \square & 0 \\ \hline \square & \square & \\ \hline \square & & \\ \hline \square & & \\ \hline \square & & \\ \hline \square & & \\ \hline \square & & \\ \hline \square & & \\ \hline \end{array} \end{array}$$

The vacancy number is written on the left of the Young diagrams and the rigging is attached to each row. For a partition $\lambda = (\lambda_1, \dots, \lambda_k)$, we let $\text{RC}(\lambda)$ denote the set of rigged configurations (2.1) with $\mu^{(0)} = \lambda$.

2.2. Crystals. We recapitulate basic facts on the $A_n^{(1)}$ crystal B_l . For a general background see Appendix A. The B_l is the crystal base of the l -fold symmetric tensor representation. As the set it is given by

$$(2.6) \quad B_l = \{x = (x_1, \dots, x_{n+1}) \in (\mathbb{Z}_{\geq 0})^{n+1} \mid x_1 + \dots + x_{n+1} = l\}.$$

The Kashiwara operators act as $\tilde{e}_i(x) = x'$, $\tilde{f}_i(x) = x''$ with $x'_j = x_j + \delta_{i,j} - \delta_{i,j+1}$ and $x''_j = x_j - \delta_{i,j} + \delta_{i,j+1}$. Here indices are in \mathbb{Z}_{n+1} and x' and x'' are to be understood as 0 unless they belong to $(\mathbb{Z}_{\geq 0})^{n+1}$. The combinatorial $R : \text{Aff}(B_l) \otimes \text{Aff}(B_m) \rightarrow \text{Aff}(B_m) \otimes \text{Aff}(B_l)$ has the form $R : x[d] \otimes y[e] \mapsto \tilde{y}[e - H(x \otimes y)] \otimes \tilde{x}[d + H(x \otimes y)]$ with

$$(2.7) \quad \tilde{x}_i = x_i + Q_i(x, y) - Q_{i-1}(x, y), \quad \tilde{y}_i = y_i + Q_{i-1}(x, y) - Q_i(x, y),$$

$$(2.8) \quad Q_i(x, y) = \min\left\{\sum_{j=1}^{k-1} x_{i+j} + \sum_{j=k+1}^{n+1} y_{i+j} \mid 1 \leq k \leq n+1\right\},$$

$$(2.9) \quad H(x \otimes y) = \min(l, m) - Q_0(x, y).$$

The energy function H here is normalized so that $0 \leq H \leq \min(l, m)$ and coincides with the ‘‘winding number’’ [11]. The element $x = (x_1, \dots, x_{n+1})$ is also denoted by a row shape semistandard tableau of length l containing the letter i x_i times and $x[d] \in \text{Aff}(B_l)$ by the tableau with index d . For example in $A_3^{(1)}$, the following stand for the same relation under R :

$$(1, 2, 0, 1)[5] \otimes (1, 0, 1, 0)[9] \simeq (0, 1, 0, 1)[8] \otimes (2, 1, 1, 0)[6],$$

$$\boxed{1224}_5 \otimes \boxed{13}_9 \simeq \boxed{24}_8 \otimes \boxed{1123}_6.$$

To save the space we use the notation:

$$(2.10) \quad a^l = \boxed{a \cdots a} \in B_l.$$

The relation $x \otimes y \simeq \tilde{y} \otimes \tilde{x}$ is depicted as

$$(2.11) \quad \begin{array}{c} y \\ | \\ x \text{---} \tilde{x} \\ | \\ \tilde{y} \end{array} \quad \text{or} \quad \begin{array}{c} \tilde{x} \\ | \\ \tilde{y} \text{---} y \\ | \\ x \end{array}$$

Setting

$$(2.12) \quad B_l^{\geq a+1} = \{(x_1, \dots, x_{n+1}) \in B_l \mid x_1 = \dots = x_a = 0\} \quad (0 \leq a \leq n),$$

we have

$$(2.13) \quad B_l = B_l^{\geq 1} \supset B_l^{\geq 2} \supset \cdots \supset B_l^{\geq n+1} = \{(n+1)^l\}$$

as sets. We will need to consider the crystals not only for $A_n^{(1)}$ but also for the nested family $A_0^{(1)}, A_1^{(1)}, \dots, A_{n-1}^{(1)}$. In such a circumstance we realize the crystal B_l for $A_{n-a}^{(1)}$ ($0 \leq a \leq n$) on the set $B_l^{\geq a+1}$ with the Kashiwara operators \tilde{e}_i, \tilde{f}_i ($a \leq i \leq n$). In this convention the highest element with respect to A_{n-a} is $(a+1)^l \in B_l^{\geq a+1}$.

Let

$$(2.14) \quad \mathcal{P}_+(\lambda) = \{p \in B_{\lambda_1} \otimes \cdots \otimes B_{\lambda_k} \mid \tilde{e}_i p = 0, 1 \leq i \leq n\}.$$

be the set of highest elements (paths) with respect to A_n . The bijection [3, 4] between $\text{RC}(\lambda)$ and the Littlewood-Richardson tableaux is translated to the one between $\text{RC}(\lambda)$ and $\mathcal{P}_+(\lambda)$. We call the resulting map the KKR bijection. See [8] for a recent review. It sends the rigged configuration (2.5) to

$$(2.15) \quad \boxed{111} \otimes \boxed{22} \otimes \boxed{3} \otimes \boxed{1} \otimes \boxed{4} \otimes \boxed{2} \otimes \boxed{3}.$$

2.3. Normal ordering. For an element $b_1[d_1] \otimes \cdots \otimes b_m[d_m] \in \text{Aff}(B_{l_1}) \otimes \cdots \otimes \text{Aff}(B_{l_m})$ we call the number d_i the i -th mode. By using the combinatorial R , tensor products can be reordered and the modes are changed accordingly. Given an element $s \in \text{Aff}(B_{l_1}) \otimes \cdots \otimes \text{Aff}(B_{l_m})$, define \mathcal{S}_m to be the set of such reordering as

$$(2.16) \quad \mathcal{S}_m = \{s' \in \bigsqcup_{\sigma \in \mathfrak{S}_m} \text{Aff}(B_{l_{\sigma(1)}}) \otimes \cdots \otimes \text{Aff}(B_{l_{\sigma(m)}}) \mid s' \simeq s\},$$

where \bigsqcup' means the disjoint union over $\sigma \in \mathfrak{S}_m$ satisfying $\sigma(i) < \sigma(j)$ for any i, j such that $i < j$ and $l_i = l_j$. The cardinality of \mathcal{S}_m is $m!$ if l_1, \dots, l_m are distinct. For $i = 2, \dots, m$, let \mathcal{S}_{i-1} be the subset of \mathcal{S}_i having the maximal i -th mode. Then we have

$$(2.17) \quad \emptyset \neq \mathcal{S}_1 \subseteq \mathcal{S}_2 \subseteq \cdots \subseteq \mathcal{S}_m.$$

We call the elements of \mathcal{S}_1 *normal ordered forms* of s . In general the normal ordered form $b_1[d_1] \otimes \cdots \otimes b_m[d_m]$ is not unique but the mode sequence d_1, \dots, d_m is unique by the definition. For type $A_n^{(1)}$, one has $d_1 \leq \cdots \leq d_m$. Any element of \mathcal{S}_1 is denoted by $:s:$.

In the context of soliton cellular automaton explained in Section 1, the restriction on the union \bigsqcup' in (2.16) reflects the fact that solitons of equal velocity(=amplitude) do not collide with each other.

Example 2.1. Take $s = \boxed{1223}_5 \otimes \boxed{34}_8 \otimes \boxed{1}_6$ for $A_3^{(1)}$.

$$\begin{aligned} \mathcal{S}_3 &= \{ \boxed{1223}_5 \otimes \boxed{34}_8 \otimes \boxed{1}_6, \boxed{23}_8 \otimes \boxed{1234}_5 \otimes \boxed{1}_6, \boxed{23}_8 \otimes \boxed{4}_5 \otimes \boxed{1123}_6, \\ &\quad \boxed{3}_5 \otimes \boxed{24}_8 \otimes \boxed{1123}_6, \boxed{3}_5 \otimes \boxed{1224}_5 \otimes \boxed{13}_9, \boxed{1223}_5 \otimes \boxed{4}_5 \otimes \boxed{13}_9 \}, \\ \mathcal{S}_2 = \mathcal{S}_1 &= \{ \boxed{3}_5 \otimes \boxed{1224}_5 \otimes \boxed{13}_9, \boxed{1223}_5 \otimes \boxed{4}_5 \otimes \boxed{13}_9 \}. \end{aligned}$$

2.4. Maps $\mathcal{C}_1, \dots, \mathcal{C}_n$. Pick the color a part $(\mu^{(a)}, J^{(a)})$ of the rigged configuration. Here we simply write it as (μ, J) . Namely $\mu = (\mu_1, \dots, \mu_m)$ is a partition and $J = (J_i)$, where J_i is the rigging attached to the i -th row in μ of length μ_i . For $1 \leq a \leq n$,

let $B_l = B_l^{\geq a+1}$ be the $A_{n-a}^{(1)}$ crystal in the sense explained around (2.13). Define the map \mathcal{C}_a among the $A_{n-a}^{(1)}$ crystals by

$$(2.18) \quad \begin{aligned} \mathcal{C}_a : B_{\mu_1} \otimes \cdots \otimes B_{\mu_m} &\rightarrow : \text{Aff}(B_{\mu_1}) \otimes \cdots \otimes \text{Aff}(B_{\mu_m}) : \quad (1 \leq a \leq n) \\ b_1 \otimes \cdots \otimes b_m &\mapsto : b_1[d_1] \otimes \cdots \otimes b_m[d_m] : \end{aligned}$$

$$(2.19) \quad d_i = J_i + \sum_{0 \leq k < i} H(b_k \otimes b_i^{(k+1)}), \quad b_0 = (a+1)^{\mu_1}.$$

Here $b_i^{(j)} \in B_{\mu_i}$ ($j \leq i$) is defined by bringing b_i to the left by the combinatorial R as

$$(2.20) \quad (b_j \otimes \cdots \otimes b_{i-1}) \otimes b_i \simeq b_i^{(j)} \otimes (\cdots)$$

under the isomorphism $(B_{\mu_j} \otimes \cdots \otimes B_{\mu_{i-1}}) \otimes B_{\mu_i} \simeq B_{\mu_i} \otimes (B_{\mu_j} \otimes \cdots \otimes B_{\mu_{i-1}})$.

The map \mathcal{C}_n involves “ $A_0^{(1)}$ crystal” $B_l^{\geq n+1} = \{(n+1)^l\}$. The following suffices to define \mathcal{C}_n :

$$(2.21) \quad (n+1)^l \otimes (n+1)^m \simeq (n+1)^m \otimes (n+1)^l, \quad H((n+1)^l \otimes (n+1)^m) = \min(l, m).$$

Since the normal ordering in (2.18) is not unique, \mathcal{C}_a is actually multi-valued in general. Here we mean by $\mathcal{C}_a(\cdot)$ to pick any one of the normal ordered forms.

2.5. Maps Φ_1, \dots, Φ_n . Pick the color a and $a-1$ parts of the configuration and denote them simply by $\mu^{(a)} = (\mu_1, \dots, \mu_m)$ and $\mu^{(a-1)} = (\lambda_1, \dots, \lambda_k)$. Set $B_l = B_l^{\geq a+1}$ and $B'_l = B_l^{\geq a}$. We define the map Φ_a from the normal ordered elements in $A_{n-a}^{(1)}$ affine crystals to $A_{n-a+1}^{(1)}$ crystals:


$$(2.22) \quad \begin{aligned} \Phi_a : : \text{Aff}(B_{\mu_1}) \otimes \cdots \otimes \text{Aff}(B_{\mu_m}) &: \rightarrow B'_{\lambda_1} \otimes \cdots \otimes B'_{\lambda_k} \quad (1 \leq a \leq n) \\ b_1[d_1] \otimes \cdots \otimes b_m[d_m] &\mapsto c_1 \otimes \cdots \otimes c_k. \end{aligned}$$

Since $b_1[d_1] \otimes \cdots \otimes b_m[d_m]$ is normal ordered, we know that $d_1 \leq \cdots \leq d_m$. We suppose further that $d_1 \geq 0$, which is the case in the actual use later. Then the image $c_1 \otimes \cdots \otimes c_k$ is determined by the following relation under the isomorphism of $A_{n-a+1}^{(1)}$ crystals: (We write $\mathcal{T}_a^d = \boxed{a}^{\otimes d} \in (B_1^{\geq a})^{\otimes d}$ for short.)

$$(2.23) \quad \begin{aligned} (\mathcal{T}_a^{d_1} \otimes b_1 \otimes \mathcal{T}_a^{d_2-d_1} \otimes b_2 \otimes \cdots \otimes \mathcal{T}_a^{d_m-d_{m-1}} \otimes b_m) \otimes (a^{\lambda_1} \otimes a^{\lambda_2} \otimes \cdots \otimes a^{\lambda_k}) \\ \simeq (c_1 \otimes \cdots \otimes c_k) \otimes \text{tail}, \end{aligned}$$

Here we are regarding $b_i \in B_{\mu_i} = B_{\mu_i}^{\geq a+1}$ as an element of $B'_{\mu_i} = B_{\mu_i}^{\geq a}$ by the natural embedding (2.13) as sets. The tail part has the same structure as $(\mathcal{T}_a^{d_1} \otimes b_1 \otimes \mathcal{T}_a^{d_2-d_1} \otimes \cdots \otimes b_m)$ on the left hand side. In the actual use, it turns out to be $(\mathcal{T}_a^{d_1} \otimes a^{\mu_1} \otimes \mathcal{T}_a^{d_2-d_1} \otimes \cdots \otimes a^{\mu_m})$ containing the letter a only. (This fact will not be used.)

To obtain $c_1 \otimes \cdots \otimes c_k$ using (2.23), one applies the combinatorial R on $B'_j \otimes B'_l$ to carry $(\mathcal{T}_a^{d_1} \otimes b_1 \otimes \cdots \otimes \mathcal{T}_a^{d_m-d_{m-1}} \otimes b_m)$ through $(a^{\lambda_1} \otimes a^{\lambda_2} \otimes \cdots \otimes a^{\lambda_k})$ to the right. The procedure is depicted as



2.6. Main theorem. Define the $A_0^{(1)}$ crystal element

$$(2.24) \quad p^{(n)} = (n+1)^{\mu_1^{(n)}} \otimes \cdots \otimes (n+1)^{\mu_n^{(n)}}.$$

Theorem 2.2. *The image p of the rigged configuration $(\mu^{(0)}, (\mu^{(1)}, J^{(1)}), \dots, (\mu^{(n)}, J^{(n)}))$ under the KKR bijection is given by*

$$(2.25) \quad p = \Phi_1 \mathcal{C}_1 \Phi_2 \mathcal{C}_2 \cdots \Phi_n \mathcal{C}_n(p^{(n)}).$$

The theorem asserts that $\Phi_1 \mathcal{C}_1 \Phi_2 \mathcal{C}_2 \cdots \Phi_n \mathcal{C}_n(p^{(n)})$ is independent of the choices of the possibly non-unique normal ordered forms when applying the maps $\mathcal{C}_1, \dots, \mathcal{C}_n$. A proof is presented in [17].

Set

$$(2.26) \quad p^{(a)} = \Phi_{a+1} \mathcal{C}_{a+1} \cdots \Phi_n \mathcal{C}_n(p^{(n)}) \quad (0 \leq a \leq n-1),$$

which belongs to the $A_{n-a}^{(1)}$ crystal $B_{\mu_1^{(a)}}^{\geq a+1} \otimes \cdots \otimes B_{\mu_n^{(a)}}^{\geq a+1}$. Thus p in (2.25) is $p^{(0)}$.

Corollary 2.3. *For $0 \leq a \leq n-1$, $p^{(a)}$ coincides with the image of the truncated rigged configuration $(\mu^{(a)}, (\mu^{(a+1)}, J^{(a+1)}), \dots, (\mu^{(n)}, J^{(n)}))$ under the KKR bijection.*

Remark 2.4. In a rigged configuration (2.1), one may treat $\mu^{(a)}$ as a composition rather than a partition. In that case, the condition (2.4) should read $0 \leq J_{i_1}^{(a)} \leq \cdots \leq J_{i_m}^{(a)} \leq p_j^{(a)}$ if $\{i_1 < \cdots < i_m\} = \{k \mid \mu_k^{(a)} = j\}$. The other necessary change is only to redefine b_0 in (2.19) as $b_0 = (a+1)^{\max\{\mu_1, \dots, \mu_m\}}$. Actually, $b_0 = (a+1)^M$ with any $M \geq \max\{\mu_1, \dots, \mu_m\}$ leads to the same d_i in (2.19). The same remark applies also for the other algebras treated in this paper.

Example 2.5. Consider the $A_3^{(1)}$ rigged configuration (2.5). According to (2.24) we set

$$p^{(3)} = \boxed{4}, \quad \mathcal{C}_3(p^{(3)}) = \boxed{4}_1.$$

We use (2.23) to find $p^{(2)} = \Phi_3 \mathcal{C}_3(p^{(3)})$. It amounts to calculating $\mathcal{T}_3^1 \otimes \boxed{4} \otimes (\boxed{33} \otimes \boxed{3})$. This is done as

$$\begin{array}{cc} \begin{array}{c} 33 \\ 4 \text{ --- } | \\ 34 \\ 3 \text{ --- } | \\ 33 \end{array} & \begin{array}{c} 3 \\ 3 \text{ --- } | \\ 3 \\ 4 \text{ --- } | \\ 4 \end{array} \end{array}$$

Thus we have

$$p^{(2)} = \boxed{33} \otimes \boxed{4}, \quad \mathcal{C}_2(p^{(2)}) = \boxed{3}_1 \otimes \boxed{34}_3,$$

where we have used $H(\boxed{33} \otimes \boxed{4}) = 0$, $H(\boxed{33} \otimes \boxed{33}) = 2$ and $H(\boxed{33} \otimes \boxed{3}) = 1$. To find $p^{(1)} = \Phi_1 \mathcal{C}_1(p^{(2)})$, we calculate $\mathcal{T}_2^1 \otimes \boxed{3} \otimes \mathcal{T}_2^2 \otimes \boxed{34} \otimes (\boxed{22} \otimes \boxed{22} \otimes \boxed{2} \otimes \boxed{2})$.

$$\begin{array}{cccc} \begin{array}{c} 22 \\ 34 \text{ --- } | \\ 34 \\ 2 \text{ --- } | \\ 24 \end{array} & \begin{array}{c} 22 \\ 22 \text{ --- } | \\ 22 \\ 3 \text{ --- } | \\ 23 \end{array} & \begin{array}{c} 2 \\ 2 \text{ --- } | \\ 2 \\ 2 \text{ --- } | \\ 2 \end{array} & \begin{array}{c} 2 \\ 2 \text{ --- } | \\ 2 \\ 2 \text{ --- } | \\ 2 \end{array} \\ 2 \text{ --- } | & 4 \text{ --- } | & 2 \text{ --- } | & 2 \text{ --- } | \\ 22 & 34 & 2 & 2 \\ 3 \text{ --- } | & 2 \text{ --- } | & 3 \text{ --- } | & 2 \text{ --- } | \\ 23 & 24 & 3 & 2 \\ 2 \text{ --- } | & 3 \text{ --- } | & 4 \text{ --- } | & 3 \text{ --- } | \\ 22 & 23 & 4 & 3 \end{array}$$

Thus we find

$$p^{(1)} = \boxed{22} \otimes \boxed{23} \otimes \boxed{4} \otimes \boxed{3}, \quad \mathcal{C}_1(p^{(1)}) =: \boxed{22}_2 \otimes \boxed{23}_3 \otimes \boxed{4}_1 \otimes \boxed{3}_3:$$

There are three normal ordered forms $\boxed{2}_1 \otimes \boxed{3}_2 \otimes \boxed{22}_3 \otimes \boxed{34}_3$, $\boxed{2}_1 \otimes \boxed{23}_2 \otimes \boxed{2}_3 \otimes \boxed{34}_3$ and $\boxed{2}_1 \otimes \boxed{23}_2 \otimes \boxed{24}_3 \otimes \boxed{3}_3$. Any one of them can be chosen as $\mathcal{C}_1(p^{(1)})$. We illustrate the derivation of $p = \Phi_1 \mathcal{C}_1(p^{(1)})$ along the first one. According to (2.23) we

large L . Set $\mu = \mu^{(1)} = (\mu_1, \dots, \mu_l)$. Consider $(u^{2, \mu_1} \otimes \dots \otimes u^{2, \mu_l}) \otimes p$ and switch the order by the combinatorial R :

$$\begin{aligned} B_\mu^{(2)} \otimes B_\lambda &\longrightarrow B_\lambda \otimes B_\mu^{(2)} \\ (u^{2, \mu_1} \otimes \dots \otimes u^{2, \mu_l}) \otimes p &\longmapsto \tilde{p} \otimes (b_1 \otimes \dots \otimes b_l). \end{aligned}$$

Then we have the following.

- (1) \tilde{p} is an element of $\mathcal{P}_+(\lambda)$ that does not contain letters greater than 2.
- (2) \tilde{p} corresponds to the RC $(\mu^{(1)}, J^{(1)})$.
- (3) The letters in the first row of b_j are all 1 for any j .
- (4) $\bar{b}_1 \otimes \dots \otimes \bar{b}_l$ is an element of $\mathcal{P}_+(\mu)$ that corresponds to the RC $((\mu^{(2)}, J^{(2)}), \dots, (\mu^{(n)}, J^{(n)}))$. Here \bar{b}_j is the second row of b_j .

Proof. Let $RC = ((\mu^{(1)}, J^{(1)}), (\mu^{(2)}, J^{(2)}), \dots, (\mu^{(n)}, J^{(n)}))$ be the RC corresponding to p . From Proposition 2.6 the RC corresponding to $(u^{2, \mu_1} \otimes \dots \otimes u^{2, \mu_l}) \otimes p$ is given by

$$\widehat{RC} = ((\mu^{(1)}, J^{(1)}), (\mu^{(2)}, \hat{J}^{(2)}), \dots, (\mu^{(n)}, J^{(n)}))$$

where $\hat{J}_i^{(2)} = J_i^{(2)} + \sum_{k=1}^l \min(\mu_k, \mu_i^{(2)})$. Since the RC does not change by the application of the combinatorial R (see [21] Lemma 8.5), the RC corresponding to $\tilde{p} \otimes (b_1 \otimes \dots \otimes b_l)$ is also given by \widehat{RC} . Note the following facts.

- (i) A string in \widehat{RC} is singular, if and only if the corresponding string in RC is singular.
- (ii) There is no singular string in $(\mu^{(1)}, J^{(1)})$.

We now apply the procedures ls, lb, lh, i, j, δ successively on RC to obtain $b_1 \otimes \dots \otimes b_l$ and \tilde{p} . By applying ls, i , one can assume $b_l \in B^{2,1}$. By applying lb, j, lh, δ , we then obtain $b_l = \boxed{\begin{smallmatrix} 1 \\ a \end{smallmatrix}}$ ($a \geq 2$). Notice that this process is parallel to applying ls, i, lh, δ to the RC

$$\widetilde{RC} = (\mu, (\mu^{(2)}, J^{(2)}), \dots, (\mu^{(n)}, J^{(n)})).$$

Namely, if $b_l = \boxed{\begin{smallmatrix} 1 \\ a \end{smallmatrix}}$, then the first letter to be obtained from \widetilde{RC} is $a - 1$. By doing these procedures successively until we obtain b_1, \dots, b_l , we can obtain (3), (4). ($\bar{b}_1 \otimes \dots \otimes \bar{b}_l \in \mathcal{P}_+(\mu)$ follows from the fact that it corresponds to an admissible RC.)

These procedures continue until we finish removing letters corresponding to $B_\mu^{(2)}$. We then see \tilde{p} corresponds to $(\mu^{(1)}, J^{(1)})$ and therefore has no letter greater than 2. $\tilde{p} \in \mathcal{P}_+(\lambda)$ follows from the simple fact on crystals that if $b_1 \otimes b_2$ is a highest element, then so is b_1 . \square

Let p be an element of $\mathcal{P}_+(\lambda)$. In view of the above theorem we consider the following procedure for p .

- (1) Supply p with sufficiently many $u^{1,1}$ on the right.
- (2) Compute the image of the map

$$\begin{aligned} (B^{2,1})^{\otimes L} \otimes B_\lambda &\longrightarrow B_\lambda \otimes (B^{2,1})^{\otimes L} \\ (u^{2,1})^{\otimes L} \otimes p &\longmapsto \tilde{p} \otimes p'. \end{aligned}$$

L should be chosen so large that \tilde{p} does not contain letters greater than 2.

- (3) Compute the RC (μ, J) corresponding to \tilde{p} .

- (4) Cut the first rows of p' (for their entries are all 1), decrease each letter by 1 and denote it by p'' . Compute the image of the map

$$\begin{aligned} B_\mu \otimes (B^{1,1})^{\otimes L} &\longrightarrow (B^{1,1})^{\otimes L} \otimes B_\mu \\ (u^{1,\mu_1} \otimes \cdots \otimes u^{1,\mu_\iota}) \otimes p'' &\longmapsto u \otimes p'''. \end{aligned}$$

(In fact, $u = (u^{1,1})^{\otimes L}$.)

- (5) Replace p with p''' and repeat (1)-(4) until p''' has no letter greater than 2.

This procedure is shown to give the RC corresponding to p by Proposition 2.6, Theorem 5.7 of [20], Theorem 2.7 and the following lemma, that can easily be obtained from [22].

Lemma 2.8. *Let $b \in B^{2,l}, b' \in B^{2,m}$. Suppose the letters of the first rows of b, b' are all 1. Let $\tilde{b}' \otimes \tilde{b}$ be the image of $b \otimes b'$ under the map $B^{2,l} \otimes B^{2,m} \rightarrow B^{2,m} \otimes B^{2,l}$. Then the letters of the first rows of \tilde{b}', \tilde{b} are again all 1, and their second rows are given by the image of $b_2 \otimes b'_2$ under $B^{1,l} \otimes B^{1,m} \rightarrow B^{1,m} \otimes B^{1,l}$, where b_2, b'_2 are the second rows of b, b' .*

Example 2.9. $n = 3$. We abbreviate \otimes to \cdot and draw frames of tableaux only for $B^{2,l}$.

$$p = 1 \cdot 1 \cdot 1 \cdot 1 \cdot 2 \cdot 2 \cdot 3 \cdot 2 \cdot 1 \cdot 4 \cdot 3 \cdot 2 \cdot 2 \in B_{(113)}.$$

By $(B^{2,1})^{\otimes 3} \otimes B_{(116)} \rightarrow B_{(116)} \otimes (B^{2,1})^{\otimes 3}$, $(u^{2,1})^{\otimes 3} \otimes (p \otimes (u^{1,1})^{\otimes 3})$ maps to $\tilde{p} \otimes p'$ where

$$\tilde{p} = 1 \cdot 1 \cdot 1 \cdot 1 \cdot 2 \cdot 2 \cdot 1 \cdot 2 \cdot 2 \cdot 1 \cdot 1 \cdot 1 \cdot 2 \cdot 2 \cdot 2 \cdot 2, \quad p' = \begin{array}{|c|} \hline 1 \\ \hline 3 \\ \hline \end{array} \cdot \begin{array}{|c|} \hline 1 \\ \hline 3 \\ \hline \end{array} \cdot \begin{array}{|c|} \hline 1 \\ \hline 4 \\ \hline \end{array}.$$

The RC for \tilde{p} is $(\mu, J) = ((4, 3, 1), (0, 1, 4))$, and $p'' = 2 \cdot 2 \cdot 3$. By $B_\mu \otimes (B^{1,1})^{\otimes 3} \rightarrow (B^{1,1})^{\otimes 3} \otimes B_\mu$, $(u^{1,4} \otimes u^{1,3} \otimes u^{1,1}) \otimes p''$ maps to $1^{\otimes 3} \otimes p'''$ where

$$p''' = 1111 \cdot 122 \cdot 3.$$

Redefine $p = p'''$ and repeat the procedure. We obtain

$$(2.27) \quad \tilde{p} = 1111 \cdot 122 \cdot 1 \cdot 2, \quad p' = \begin{array}{|c|} \hline 1 \\ \hline 3 \\ \hline \end{array},$$

$$(2.28) \quad (\mu, J) = ((2, 1), (0, 0)), \quad p'' = 2, \quad p''' = 11 \cdot 2.$$

The RC for this p''' is $((1), (0))$. Thus we find the RC for the original p as

$$\begin{array}{ccc} \mu^{(1)} & \mu^{(2)} & \mu^{(3)} \\ \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline \end{array} 0 & \begin{array}{|c|c|} \hline & \\ \hline & \\ \hline \end{array} 0 & \begin{array}{|c|} \hline \\ \hline \end{array} 0 \\ \begin{array}{|c|c|c|} \hline & & \\ \hline & & \\ \hline & & \\ \hline \end{array} 1 & \begin{array}{|c|} \hline \\ \hline \end{array} 0 & \\ \begin{array}{|c|} \hline \\ \hline \end{array} 4 & & \end{array}$$

3. CONJECTURES ON OTHER TYPES

There is a bijection between the set of rigged configurations $\text{RC}(\lambda)$ and the highest paths (2.6) for $\mathfrak{g}_n = B_n^{(1)}, C_n^{(1)}, D_n^{(1)}, A_{2n-1}^{(2)}, A_{2n}^{(2)}$ and $D_{n+1}^{(2)}$. It has been established in [18, 19] by a combinatorial algorithm similar to the $A_n^{(1)}$ case. Here we state conjectures analogous to (2.25).

Precise specifications of the data $\text{RC}(\lambda)$, $C_a, \Phi_a, p^{(n)}$ and $\bar{p}^{(n)}$ will be given in subsequent sections. In particular for $\mathfrak{g}_n = A_{2n}^{(2)}$ and $D_{n+1}^{(2)}$, we utilize the embedding ι into $C_n^{(1)}$ specified in (8.5) and (8.8). The relevant combinatorial R for these algebras

has been obtained in [23, 24] by an insertion scheme. A piecewise linear formula is also available for $D_n^{(1)}$ in [25] and the other \mathfrak{g}_n using the embedding described in [26].

Conjecture 3.1. *Given a rigged configuration (μ, J) as in (2.1), let $p \in \mathcal{P}_+(\lambda)$ be the classically highest path (2.14) that corresponds to it under the bijection in [18, 19]. Then the following formulas are valid:*

$$(3.1) \quad p = \Phi_1 \mathcal{C}_1 \cdots \Phi_{n-2} \mathcal{C}_{n-2} (\Phi_{n-1} \mathcal{C}_{n-1} (p^{(n)}) + \Phi_n \mathcal{C}_n (\bar{p}^{(n)})) \quad \text{for } \mathfrak{g}_n = D_n^{(1)},$$

$$(3.2) \quad p = \Phi_1 \mathcal{C}_1 \cdots \Phi_n \mathcal{C}_n (\bar{p}^{(n)}) \quad \text{for } \mathfrak{g}_n = A_{2n-1}^{(2)}, B_n^{(1)}, C_n^{(1)}.$$

For $\mathfrak{g}_n = A_{2n}^{(2)}$ and $D_{n+1}^{(2)}$, let p' be the above $\Phi_1 \mathcal{C}_1 \cdots \Phi_n \mathcal{C}_n (\bar{p}^{(n)})$ for $C_n^{(1)}$ corresponding to the rigged configuration $\iota((\mu, J))$. Then $\iota^{-1}(p')$ exists and is equal to p .

For $\mathfrak{g}_n = A_{2n-1}^{(2)}, B_n^{(1)}$ and $C_n^{(1)}$, define the \mathfrak{g}_{n-a} path $p^{(a)} = \Phi_{a+1} \mathcal{C}_{a+1} \cdots \Phi_n \mathcal{C}_n (\bar{p}^{(n)})$ for $0 \leq a \leq n-1$. Similarly for $\mathfrak{g}_n = D_n^{(1)}$, set $p^{(a)} = \Phi_{a+1} \mathcal{C}_{a+1} \cdots \Phi_{n-2} \mathcal{C}_{n-2} (\Phi_{n-1} \mathcal{C}_{n-1} (p^{(n)}) + \Phi_n \mathcal{C}_n (\bar{p}^{(n)}))$ for $0 \leq a \leq n-2$. Under Conjecture 3.1, $p^{(a)}$ coincides with the image of the truncated rigged configuration $(\mu^{(a)}, (\mu^{(a+1)}, J^{(a+1)}), \dots, (\mu^{(n)}, J^{(n)}))$ under the bijection in [18, 19].

4. $D_n^{(1)}$ CASE

4.1. Rigged configurations. Consider the data of the form (2.1) and define $E_j^{(a)}$ as in (2.2). The vacancy number is specified by (2.3) for $1 \leq a \leq n-3$ and

$$(4.1) \quad \begin{aligned} p_j^{(n-2)} &= E_j^{(n-3)} + E_j^{(n-1)} + E_j^{(n)} - 2E_j^{(n-2)}, \\ p_j^{(a)} &= E_j^{(n-2)} - 2E_j^{(a)} \quad (a = n-1, n). \end{aligned}$$

The data (2.1) is called a $D_n^{(1)}$ rigged configuration if (2.4) is satisfied. It is depicted as in $A_n^{(1)}$ case by the Young diagrams with rigging. The following is an example of $D_4^{(1)}$ rigged configuration.

$$(4.2) \quad \begin{array}{ccccc} \mu^{(0)} & \mu^{(1)} & \mu^{(2)} & \mu^{(3)} & \mu^{(4)} \\ (1^9) & 0 \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array} 0 & 0 \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} 0 & 0 \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} 0 & 0 \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} 0 \\ & 2 & 0 & & \end{array}$$

4.2. Crystals.

$$(4.3) \quad B_l = \{(x_1, \dots, x_n, \bar{x}_n, \dots, \bar{x}_1) \in (\mathbb{Z}_{\geq 0})^{2n} \mid \sum_{i=1}^n (x_i + \bar{x}_i) = l, x_n \bar{x}_n = 0\}.$$

$$(4.4) \quad B_l^{\geq a+1} = \{(x_1, \dots, x_n, \bar{x}_n, \dots, \bar{x}_1) \in B_l \mid x_i = \bar{x}_i = 0 (1 \leq i \leq a)\}$$

for $0 \leq a \leq n-2$. Then

$$B_l = B_l^{\geq 1} \supset B_l^{\geq 2} \supset \cdots \supset B_l^{\geq n-1}$$

as sets. We regard $B_l^{\geq a+1}$ as $D_{n-a}^{(1)}$ crystal and represent the elements by length l semistandard row tableaux over the alphabet $a+1 < \cdots < n < \bar{n} < \cdots < \bar{a+1}$. (Actually the letter n and \bar{n} do not coexist in a tableau.) We use the notation (2.10).

Then the highest element in $B_l^{\geq a+1}$ with respect to D_{n-a} is $(a+1)^l$. The rigged configuration (4.2) corresponds to

$$(4.5) \quad \boxed{1} \otimes \boxed{1} \otimes \boxed{1} \otimes \boxed{1} \otimes \boxed{2} \otimes \boxed{\bar{1}} \otimes \boxed{\bar{1}} \otimes \boxed{3} \otimes \boxed{2} \in \mathcal{P}_+((1^9)).$$

4.3. $\mathcal{C}_{n-1}(p^{(n)})$ and $\mathcal{C}_n(\bar{p}^{(n)})$. Set

$$(4.6) \quad p^{(n)} = n^{\mu_1^{(n-1)}} \otimes \cdots \otimes n^{\mu_{l_{n-1}}^{(n-1)}},$$

$$(4.7) \quad \bar{p}^{(n)} = \bar{n}^{\mu_1^{(n)}} \otimes \cdots \otimes \bar{n}^{\mu_{l_n}^{(n)}},$$

where n^l means the unique element of “ $A_0^{(1)}$ crystal” B_l realized by the tableau with letter n only. (See (2.10).) \bar{n}^l stands for the same object in another copy of $A_0^{(1)}$ crystal B_l . Then $\mathcal{C}_{n-1}(p^{(n)})$ is defined from (2.18) and (2.19) by setting $(\mu, J) = (\mu^{(n-1)}, J^{(n-1)})$, $b_i = n^{\mu_i}$ ($i \geq 1$), $b_0 = n^{\mu_1}$ and regarding B_l there as the $A_0^{(1)}$ crystal with letter n only. Namely, $\mathcal{C}_{n-1}(p^{(n)}) =: n^{\mu_1}[d_1] \otimes \cdots \otimes n^{\mu_m}[d_m]$; where the normal ordering $::$ is done along the $A_0^{(1)}$ combinatorial R (2.21) with $n+1$ replaced by n . Similarly $\mathcal{C}_n(\bar{p}^{(n)})$ is defined by setting $(\mu, J) = (\mu^{(n)}, J^{(n)})$, $b_i = \bar{n}^{\mu_i}$ ($i \geq 1$) and $b_0 = \bar{n}^{\mu_1}$ in (2.18) and (2.19). As the result $\mathcal{C}_{n-1}(p^{(n)})$ and $\mathcal{C}_n(\bar{p}^{(n)})$ are elements of the two independent copies of $A_0^{(1)}$ affine crystals realized with tableaux with letters n only or \bar{n} only. Since the energy function in (2.21) is positive, the i -th mode d_i in the normal ordered form satisfies $0 \leq d_1 \leq \cdots \leq d_m$.

4.4. $\Phi_{n-1}\mathcal{C}_{n-1}(p^{(n)})$, $\Phi_n\mathcal{C}_n(\bar{p}^{(n)})$ and their superposition. Let us write as $(\lambda_1, \dots, \lambda_k) = (\mu_1^{(n-2)}, \dots, \mu_{l_{n-2}}^{(n-2)})$ and $\mathcal{C}_{n-1}(p^{(n)}) = n^{\mu_1}[d_1] \otimes \cdots \otimes n^{\mu_m}[d_m]$. Let $B_l' = \{(x_{n-1}, x_n)' \in (\mathbb{Z}_{\geq 0})^2 \mid x_{n-1} + x_n = l\}$ be the $A_1^{(1)}$ crystal with tableau letters $n-1, n$, where $n^l = \boxed{n \cdots n}$ is identified as the lowest weight element $(0, l)'$. We define $\Phi_{n-1}\mathcal{C}_{n-1}(p^{(n)}) \in B_{\lambda_1}' \otimes \cdots \otimes B_{\lambda_k}'$ to be $c_1 \otimes \cdots \otimes c_k$ determined by (2.23) by setting $b_i = n^{\mu_i}$ and $a = n-1$.

$\Phi_n\mathcal{C}_n(\bar{p}^{(n)})$ is defined similarly by replacing the letter n by \bar{n} . To be precise, we reset the meaning of μ_i and d_i by saying that $\mathcal{C}_n(\bar{p}^{(n)}) = \bar{n}^{\mu_1}[d_1] \otimes \cdots \otimes \bar{n}^{\mu_m}[d_m]$. Let $B_l'' = \{(x_{n-1}, \bar{x}_n)'' \in (\mathbb{Z}_{\geq 0})^2 \mid x_{n-1} + \bar{x}_n = l\}$ be the $A_1^{(1)}$ crystal with tableau letters $n-1, \bar{n}$, where $\bar{n}^l = \boxed{\bar{n} \cdots \bar{n}}$ is identified as the lowest weight element $(0, l)''$. We keep the meaning $\lambda = \mu^{(n-2)}$ as before. Then $\Phi_n\mathcal{C}_n(\bar{p}^{(n)}) \in B_{\lambda_1}'' \otimes \cdots \otimes B_{\lambda_k}''$ is defined to be $c_1 \otimes \cdots \otimes c_k$ determined by (2.23) by setting $b_i = \bar{n}^{\mu_i}$ and $a = n-1$.

We introduce the “superposition” $+$ of the two copies of $A_1^{(1)}$ crystals B_l' and B_l'' :

$$(4.8) \quad \begin{aligned} + : B_l' \times B_l'' &\longrightarrow D_2^{(1)} \text{ crystal } B_l^{\geq n-1} \\ (x_{n-1}, x_n)' \times (y_{n-1}, \bar{y}_n)'' &\longmapsto (0, \dots, 0, z_{n-1}, z_n, \bar{z}_n, \bar{z}_{n-1}, 0, \dots, 0), \\ z_{n-1} &= \min(x_{n-1}, y_{n-1}), z_n = \max(0, y_{n-1} - x_{n-1}), \\ \bar{z}_n &= \max(0, \bar{y}_n - x_n), \bar{z}_{n-1} = \min(x_n, \bar{y}_n). \end{aligned}$$

In terms of the tableaux, this $+$ means to “superpose” the letters $n-1, n$ from B_l' and $n-1, \bar{n}$ from B_l'' as $(n-1, n-1) \mapsto n-1, (n, n-1) \mapsto n, (n-1, \bar{n}) \mapsto \bar{n}$ and $(n, \bar{n}) \mapsto \bar{n}-\bar{1}$. For example in $n=4$ case, one has $\boxed{334} + \boxed{344} = \boxed{343}$. Note that the conditions $z_n \bar{z}_n = 0$ and $z_{n-1} + z_n + \bar{z}_n + \bar{z}_{n-1} = l$ are satisfied. Under these

conditions, the map (4.8) is invertible. Suppose we have

$$\begin{aligned}\Phi_{n-1}\mathcal{C}_{n-1}(p^{(n)}) &= b'_1 \otimes \cdots \otimes b'_k \in B'_{\lambda_1} \otimes \cdots \otimes B'_{\lambda_k}, \\ \Phi_n\mathcal{C}_n(\bar{p}^{(n)}) &= b''_1 \otimes \cdots \otimes b''_k \in B''_{\lambda_1} \otimes \cdots \otimes B''_{\lambda_k},\end{aligned}$$

where $\lambda = \mu^{(n-2)}$ as before. Then we specify their superposition as

$$(4.9) \quad \Phi_{n-1}\mathcal{C}_{n-1}(p^{(n)}) + \Phi_n\mathcal{C}_n(\bar{p}^{(n)}) = (b'_1 + b''_1) \otimes \cdots \otimes (b'_k + b''_k) \in B_{\lambda_1}^{\geq n-1} \otimes \cdots \otimes B_{\lambda_k}^{\geq n-1}.$$

4.5. Maps $\mathcal{C}_1, \dots, \mathcal{C}_{n-2}$. For $1 \leq a \leq n-2$, \mathcal{C}_a are given by (2.18) and (2.19) provided that B_l is regarded as the $D_{n-a}^{(1)}$ crystal $B_l^{\geq a+1}$. The normal ordered form $b[d_1] \otimes \cdots \otimes b_m[d_m]$ is defined in the same way as the $A_n^{(1)}$ case described in Section 2.3. A new feature is that the energy function can be negative, hence $d_1 \leq \cdots \leq d_m$ is no longer valid in general.

4.6. Maps $\Phi_1, \dots, \Phi_{n-2}$. For $1 \leq a \leq n-2$, we define the map Φ_a by (2.22) and (2.23) by regarding $B_l = B_l^{\geq a+1}$ and $B'_l = B_l^{\geq a}$ as the $D_{n-a}^{(1)}$ and $D_{n-a+1}^{(1)}$ crystals, provided $d_1 \leq \cdots \leq d_m$ holds among the modes of the normal ordered element $b_1[d_1] \otimes \cdots \otimes b_m[d_m]$. In contrast to the $A_n^{(1)}$ case, this is no longer valid in general since the energy function can be negative. ($d_1 \geq 0$ can be assured.) To make sense of $\mathcal{T}_a^d = \boxed{a}^{\otimes d}$ for negative d in (2.23), we make the regularization explained in the sequel.

In (2.23), take the largest i such that $d_i > d_{i+1}$. Then its left hand side looks as

$$(4.10) \quad (\mathcal{T}_a^{d_1} \otimes \cdots \otimes b_i \otimes \mathcal{T}_a^{-\delta} \otimes b_{i+1} \otimes \mathcal{T}_a^{d_{i+2}-d_{i+1}} \otimes \cdots \otimes b_m) \otimes (a^{\lambda_1} \otimes a^{\lambda_2} \otimes \cdots \otimes a^{\lambda_k}),$$

which is regarded as an element in the tensor product of $D_{n-a+1}^{(1)}$ crystal $B_l^{\geq a}$'s. We have set $\delta = d_i - d_{i+1} > 0$. There is no problem in carrying the components $(b_{i+1} \otimes \cdots \otimes b_m)$ through $(a^{\lambda_1} \otimes \cdots \otimes a^{\lambda_k})$ to the right to find p_j 's in $(b_{i+1} \otimes \cdots \otimes b_m) \otimes (a^{\lambda_1} \otimes \cdots \otimes a^{\lambda_k}) \simeq (p_1 \otimes \cdots \otimes p_k) \otimes \text{tail}$. Thus we are to define $\mathcal{T}_a^{-\delta} \otimes (p_1 \otimes \cdots \otimes p_k) \otimes \text{tail}$. Actually we declare that $b_i \otimes \mathcal{T}_a^{-\delta} \otimes (p_1 \otimes \cdots \otimes p_k) \otimes \text{tail}$ is to be understood as

$$(4.11) \quad \tilde{b}_i \otimes (\tilde{p}_1 \otimes \cdots \otimes \tilde{p}_k) \otimes \text{tail},$$

where $\tilde{p}_1 \otimes \cdots \otimes \tilde{p}_k$ is determined by sending \mathcal{T}_a^δ from the right to the left by the $D_{n-a+1}^{(1)}$ combinatorial R :

$$(4.12) \quad (p_1 \otimes \cdots \otimes p_k) \otimes \boxed{a}^{\otimes \delta} \simeq \text{head} \otimes (\tilde{p}_1 \otimes \cdots \otimes \tilde{p}_k).$$

In the actual use, $\text{head} = \overline{\boxed{a+1}}^{\otimes \delta}$ seems valid, but this is not necessary for our definition. To specify \tilde{b}_i in (4.11), we use the expression $b_i = (x_1, \dots, x_n, \bar{x}_n, \dots, \bar{x}_1) \in B_{\mu_i}^{\geq a+1}$. Then \tilde{b}_i is given by modifying it as $(x_{a+1}, \bar{x}_a = 0) \rightarrow (x_{a+1} - \delta, \delta)$ leaving the other coordinates unchanged. It is a part of our conjecture that $x_{a+1} \geq \delta$ is always valid as the result of normal ordering hence \tilde{b}_i is well defined.

This regularization of $b_i \otimes \mathcal{T}_a^{d_{i+1}-d_i}$ with $d_i > d_{i+1}$ can be done successively also in the part $(\mathcal{T}_a^{d_1} \otimes b_1 \otimes \cdots \otimes b_{i-1} \otimes \mathcal{T}_a^{d_i-d_{i-1}})$ in (4.10), which leads to the (conjectural) definition of our Φ_a . The change $b_i \rightarrow \tilde{b}_i$ is analogous to the pair annihilation and creation in the $D_n^{(1)}$ automaton described in [26].

4.7. **Example.** Consider the rigged configuration (4.2). We have $p^{(4)} = \boxed{44}$, $\bar{p}^{(4)} = \boxed{\bar{44}}$, $\mathcal{C}_3(p^{(4)}) = \boxed{44}_2$ and $\mathcal{C}_4(\bar{p}^{(4)}) = \boxed{\bar{44}}_2$. From

$$\begin{array}{c} \begin{array}{ccc} 333 & & 33 \\ 44 \text{ --- } | \text{ --- } 33 & \text{ --- } | \text{ --- } 33 \\ 344 & & 33 \\ 3 \text{ --- } | \text{ --- } 4 & \text{ --- } | \text{ --- } 3 \\ 334 & & 34 \\ 3 \text{ --- } | \text{ --- } 4 & \text{ --- } | \text{ --- } 3 \\ 333 & & 44 \end{array} \end{array}$$

we have $\Phi_3\mathcal{C}_3(p^{(4)}) = \boxed{333} \otimes \boxed{44}$. Similarly, $\Phi_4\mathcal{C}_4(\bar{p}^{(4)}) = \boxed{333} \otimes \boxed{\bar{44}}$. Thus the superposition is determined as $p^{(2)} = \Phi_3\mathcal{C}_3(p^{(4)}) + \Phi_4\mathcal{C}_4(\bar{p}^{(4)}) = \boxed{333} \otimes \boxed{\bar{33}}$. Since

$$\boxed{333}_3 \otimes \boxed{\bar{33}}_0 \simeq \boxed{33}_2 \otimes \boxed{3\bar{3}\bar{3}}_1$$

in $D_2^{(1)}$ crystal, we obtain $\mathcal{C}_2(p^{(2)}) = \boxed{33}_2 \otimes \boxed{3\bar{3}\bar{3}}_1$. Next to find $p^{(1)} = \Phi_2\mathcal{C}_2(p^{(2)})$, we compute $\mathcal{T}_2^2 \otimes \boxed{33} \otimes \mathcal{T}_2^{-1} \otimes \boxed{3\bar{3}\bar{3}} \otimes (\boxed{2222} \otimes \boxed{222})$ in $D_3^{(1)}$ crystal as follows:


$$\begin{array}{c} \begin{array}{ccc} 2222 & & 222 \\ 3\bar{3}\bar{3} \text{ --- } | \text{ --- } 222 & \text{ --- } | \text{ --- } 222 \\ 23\bar{3}\bar{3} & & 222 \\ \bar{3} \text{ --- } | \text{ --- } 2 & \text{ --- } | \text{ --- } 2 \\ 223\bar{3} & & 222 \\ 33 \rightsquigarrow 3\bar{2} \text{ --- } | \text{ --- } 23 & \text{ --- } | \text{ --- } 22 \\ 22\bar{2}\bar{2} & & 223 \\ 2 \text{ --- } | \text{ --- } \bar{2} & \text{ --- } | \text{ --- } 2 \\ 222\bar{2} & & 23\bar{2} \\ 2 \text{ --- } | \text{ --- } \bar{2} & \text{ --- } | \text{ --- } 2 \\ 2222 & & 3\bar{2}\bar{2} \end{array} \end{array}$$

Here the symbol $<$ in the horizontal line signifies \mathcal{T}_2^{-1} where the combinatorial R acts from NE to SW. (Our usual convention is from NW to SE as in (2.11).) The change 33 into $3\bar{2}$ due to the pair annihilation and creation is indicated by \rightsquigarrow . We have $p^{(1)} = \boxed{2222} \otimes \boxed{3\bar{2}\bar{2}}$. Under the $D_3^{(1)}$ combinatorial R , one has

$$\boxed{2222}_4 \otimes \boxed{3\bar{2}\bar{2}}_2 \simeq \boxed{222}_4 \otimes \boxed{23\bar{2}\bar{2}}_2.$$

Therefore the both sides are normal ordered forms and we choose the left hand side as $\mathcal{C}_1(p^{(1)})$. Finally to obtain $p = \Phi_1\mathcal{C}_1(p^{(1)})$, we compute $\mathcal{T}_1^4 \otimes \boxed{2222} \otimes \mathcal{T}_1^{-2} \otimes \boxed{3\bar{2}\bar{2}} \otimes (\boxed{1}^{\otimes 9})$ in $D_4^{(1)}$ crystal. It is calculated by applying $\mathcal{T}_1^4 \otimes$ to the output of

the following diagram ($\boxed{1}^{\otimes 9}$ has been replaced with $\boxed{1}^{\otimes 6}$ below):



Thus we get $p = 11112\bar{1}\bar{1}32$ in agreement with (4.5).

5. $A_{2n-1}^{(2)}$ CASE

The $A_{2n-1}^{(2)}$ case can be reduced to $D_{n+1}^{(1)}$ by dropping one term in the superposition in (3.1).

5.1. Rigged configurations. For the data (2.1) define $E_j^{(a)}$ as in (2.2). The vacancy numbers are specified by (2.3) for $1 \leq a \leq n-2$ and

$$\begin{aligned}
 (5.1) \quad p_j^{(n-1)} &= E_j^{(n-2)} - 2E_j^{(n-1)} + 2E_j^{(n)}, \\
 p_j^{(n)} &= E_j^{(n-1)} - 2E_j^{(n)}.
 \end{aligned}$$

The data (2.1) is called an $A_{2n-1}^{(2)}$ rigged configuration if the condition (2.4) is satisfied. The following is an example of $A_5^{(2)}$ rigged configuration.

$$\begin{array}{cccc}
 \mu^{(0)} & \mu^{(1)} & \mu^{(2)} & \mu^{(3)} \\
 (1^7) & 3 \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & 1 \\ \hline \square & 3 \\ \hline \end{array} 3 & 0 \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & 0 \\ \hline \end{array} 0 & 0 \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} 0 \\
 (5.2) & & &
 \end{array}$$

5.2. Crystals.

$$B_l = \{(x_1, \dots, x_n, \bar{x}_n, \dots, \bar{x}_1) \in (\mathbb{Z}_{\geq 0})^{2n} \mid \sum_{i=1}^n (x_i + \bar{x}_i) = l\},$$

$$B_l^{\geq a+1} = \{(x_1, \dots, x_n, \bar{x}_n, \dots, \bar{x}_1) \in B_l \mid x_i = \bar{x}_i = 0 \ (1 \leq i \leq a)\} \quad (0 \leq a \leq n-1).$$

$$B_l = B_l^{\geq 1} \supset B_l^{\geq 2} \supset \dots \supset B_l^{\geq n}$$

as sets. We regard $B_l^{\geq a+1}$ as $A_{2n-2a+1}^{(2)}$ crystal and represent the elements by length l semistandard row tableaux over the alphabet $a+1 < \dots < n < \bar{n} < \dots < \bar{a} + \bar{1}$. We use the notation (2.10). Then the highest element in $B_l^{\geq a+1}$ with respect to the classical part C_{n-a} is $(a+1)^l$. The rigged configuration (5.2) corresponds to the following element in $\mathcal{P}_+((1^7))$:

$$(5.3) \quad \boxed{1} \otimes \boxed{1} \otimes \boxed{2} \otimes \boxed{1} \otimes \boxed{1} \otimes \boxed{\bar{2}} \otimes \boxed{\bar{1}}.$$

5.3. **Maps $\mathcal{C}_1, \dots, \mathcal{C}_n$.** For $1 \leq a \leq n-1$, \mathcal{C}_a is defined by (2.18) and (2.19) provided that B_l is regarded as the $A_{2n-2a-1}^{(2)}$ crystal $B_l^{\geq a+1}$.

We define $\bar{p}^{(n)}$ and $\mathcal{C}_n(\bar{p}^{(n)})$ in the same way as in Section 4.3. The latter is obtained by assigning the modes to the former and normal ordering by using the $A_0^{(1)}$ combinatorial R (2.21) with letter \bar{n} .

5.4. **Maps Φ_1, \dots, Φ_n .** For $1 \leq a \leq n-1$, we define the map Φ_a by (2.22) and (2.23) by regarding $B_l = B_l^{\geq a+1}$ and $B'_l = B_l^{\geq a}$ as the $A_{2n-2a-1}^{(2)}$ and $A_{2n-2a+1}^{(2)}$ crystals. Since the energy function can be negative we employ the same regularization as $D_n^{(1)}$ explained in Section 4.6.


$\Phi_n \mathcal{C}_n(\bar{p}^{(n)})$ is defined similarly to Section 4.4 by changing the letter $n-1$ to n . To be precise, let $\mathcal{C}_n(\bar{p}^{(n)}) = \bar{n}^{\mu_1}[d_1] \otimes \dots \otimes \bar{n}^{\mu_m}[d_m]$. Regard $B_l^{\geq n} = \{(0, \dots, 0, x_n, \bar{x}_n, 0, \dots, 0) \in (\mathbb{Z}_{\geq 0})^2 \mid x_n + \bar{x}_n = l\}$ as the $A_1^{(1)}$ crystal with tableau letters n, \bar{n} , where $\bar{n}^l = \boxed{\bar{n} \cdots \bar{n}}$ is identified as the lowest weight element. Set $\lambda = \mu^{(n-1)}$. Then $\Phi_n \mathcal{C}_n(\bar{p}^{(n)}) \in B_{\lambda_1}^{\geq n} \otimes \dots \otimes B_{\lambda_k}^{\geq n}$ is defined to be $c_1 \otimes \dots \otimes c_k$ determined by (2.23) with $b_i = \bar{n}^{\mu_i}$ and $a = n$.

5.5. **Example.** Consider the rigged configuration (5.2). We have $\bar{p}^{(3)} = \boxed{\bar{3}\bar{3}}$ and $\mathcal{C}_3(\bar{p}^{(3)}) = \boxed{\bar{3}\bar{3}}_2$. To find $\Phi_3 \mathcal{C}_3(\bar{p}^{(3)})$, we calculate $\mathcal{T}_3^2 \otimes \boxed{\bar{3}\bar{3}} \otimes (\boxed{33} \otimes \boxed{33})$ as


$$\begin{array}{ccc}
 & 33 & 33 \\
 \bar{3}\bar{3} & \begin{array}{c} | \\ \hline \\ | \end{array} & 33 \begin{array}{c} | \\ \hline \\ | \end{array} & 33 \\
 & \bar{3}\bar{3} & 33 & \\
 3 & \begin{array}{c} | \\ \hline \\ | \end{array} & \bar{3} \begin{array}{c} | \\ \hline \\ | \end{array} & 3 \\
 & \bar{3}\bar{3} & \bar{3}\bar{3} & \\
 3 & \begin{array}{c} | \\ \hline \\ | \end{array} & \bar{3} \begin{array}{c} | \\ \hline \\ | \end{array} & 3 \\
 & 33 & \bar{3}\bar{3} &
 \end{array}$$

Thus we get $p^{(2)} = \Phi_3 \mathcal{C}_3(\bar{p}^{(3)}) = \boxed{33} \otimes \boxed{\bar{3}\bar{3}}$. In view of $H(\boxed{33} \otimes \boxed{\bar{3}\bar{3}}) = -2$, we have $\mathcal{C}_2(p^{(2)}) = \boxed{33}_2 \otimes \boxed{\bar{3}\bar{3}}_0$. To find $\Phi_2 \mathcal{C}_2(p^{(2)})$, we calculate $\mathcal{T}_2^2 \otimes \boxed{33} \otimes \mathcal{T}_2^{-2} \otimes \boxed{\bar{3}\bar{3}} \otimes$

$(\boxed{22} \otimes \boxed{2} \otimes \boxed{2})$ by using the $A_3^{(2)}$ combinatorial R on letters $2, 3, \bar{3}, \bar{2}$ as



Thus we get $p^{(1)} = \Phi_2 \mathcal{C}_2(p^{(2)}) = \boxed{22} \otimes \boxed{\bar{2}} \otimes \boxed{\bar{2}}$. Assigning the mode and normal ordering, we find $\mathcal{C}_1(p^{(1)}) =: \boxed{22}_5 \otimes \boxed{2}_1 \otimes \boxed{2}_4 := \boxed{2}_2 \otimes \boxed{22}_4 \otimes \boxed{2}_4$. Finally to find $\Phi_1 \mathcal{C}_1(p^{(1)})$ we compute $\mathcal{T}_1^2 \otimes \boxed{2} \otimes \mathcal{T}_1^2 \otimes \boxed{22} \otimes \boxed{\bar{2}} \otimes (\boxed{1}^{\otimes 7})$ by using the $A_5^{(2)}$ combinatorial R on letters $1, 2, 3, \bar{3}, \bar{2}, \bar{1}$. It is by given applying $\mathcal{T}_1^2 \otimes$ to the output of the following diagram ($\boxed{1}^{\otimes 7}$ has been replaced with $\boxed{1}^{\otimes 5}$ below):



Therefore we obtain $\Phi_1 \mathcal{C}_1(p^{(1)}) = 11211\bar{2}\bar{1}$ in agreement with (5.3).

6. $B_n^{(1)}$ CASE

6.1. Rigged configurations. Consider the data of the form (2.1), where $\mu^{(a)}$ with $a < n$ and $2\mu^{(n)}$ are partitions. Thus $\mu_i^{(n)} \in \mathbb{Z}/2$. $J^{(a)} = (J_1^{(a)}, \dots, J_{l_a}^{(a)})$ is taken from $(\mathbb{Z}_{\geq 0})^{l_a}$ for all $1 \leq a \leq n$. Using $E_j^{(a)}$ in (2.2), we define the vacancy numbers

by (2.3) with $1 \leq a \leq n-2$, $j \in \mathbb{Z}_{>0}$ and

$$\begin{aligned} p_j^{(n-1)} &= E_j^{(n-2)} - 2E_j^{(n-1)} + 2E_j^{(n)} \quad (j \in \mathbb{Z}_{>0}), \\ p_j^{(n)} &= 2E_j^{(n-1)} - 4E_j^{(n)} \quad (j \in \mathbb{Z}_{>0}/2). \end{aligned}$$

All the vacancy numbers are integers even though $E_j^{(n)} \in \mathbb{Z}/2$. The data (2.1) is called a $B_n^{(1)}$ rigged configuration if (2.4) is satisfied, where $j \in \mathbb{Z}_{>0}$ for $1 \leq a \leq n-1$ and $j \in \mathbb{Z}_{>0}/2$ for $a = n$.

The following is an example of $B_2^{(1)}$ rigged configuration.

$$(6.1) \quad \begin{array}{ccc} \mu^{(0)} & \mu^{(1)} & \mu^{(2)} \\ (1^7) & 0 \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & 2 \\ \hline 2 & \square \\ \hline \square & 2 \\ \hline \end{array} 0 & 2 \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} 1 \\ & & 0 \begin{array}{|c|} \hline \square \\ \hline \end{array} 0 \end{array}$$

We have depicted $\mu^{(2)}$ by a Young diagram consisting of $1 \times \frac{1}{2}$ elementary blocks.

6.2. Crystals.

$$B_l = \{(x_1, \dots, x_n, x_0, \bar{x}_n, \dots, \bar{x}_1) \in (\mathbb{Z}_{\geq 0})^{2n+1} \mid x_0 + \sum_{i=1}^n (x_i + \bar{x}_i) = l, x_0 = 0, 1\},$$

$$B_l^{\geq a+1} = \{(x_1, \dots, x_n, x_0, \bar{x}_n, \dots, \bar{x}_1) \in B_l \mid x_i = \bar{x}_i = 0 \ (1 \leq i \leq a)\} \quad (0 \leq a \leq n-1).$$

$$B_l = B_l^{\geq 1} \supset B_l^{\geq 2} \supset \dots \supset B_l^{\geq n}$$

as sets. We regard $B_l^{\geq a+1}$ as $B_{n-a}^{(1)}$ crystal and represent the elements by length l semistandard row tableaux over the alphabet $a+1 < \dots < n < 0 < \bar{n} < \dots < \overline{a+1}$. We use the notation (2.10). Then the highest element in $B_l^{\geq a+1}$ with respect to the classical part B_{n-a} is $(a+1)^l$.

The $B_2^{(1)}$ rigged configuration (6.1) corresponds to

$$(6.2) \quad \boxed{1} \otimes \boxed{1} \otimes \boxed{2} \otimes \boxed{2} \otimes \boxed{0} \otimes \boxed{0} \otimes \boxed{0} \in \mathcal{P}_+((1^7)).$$

6.3. Maps $\mathcal{C}_1, \dots, \mathcal{C}_n$. For $1 \leq a \leq n-1$, the map \mathcal{C}_a is specified by (2.18) and (2.19) provided that B_l is regarded as the $B_{n-a}^{(1)}$ crystal $B_l^{\geq a+1}$.

Define $\bar{p}^{(n)}$ and $\mathcal{C}_n(\bar{p}^{(n)})$ in the same way as in Section 4.3 by replacing $\mu_i^{(n)}$ with $2\mu_i^{(n)}$.

6.4. Maps Φ_1, \dots, Φ_n . For $1 \leq a \leq n-1$, we define the map Φ_a by (2.22) and (2.23) by regarding $B_l = B_l^{\geq a+1}$ and $B'_l = B_l^{\geq a}$ as the $B_{n-a}^{(1)}$ and $B_{n-a+1}^{(1)}$ crystals. Since the energy function can be negative we employ the same regularization as $D_n^{(1)}$ explained in Section 4.6.

To define $\Phi_n \mathcal{C}_n(\bar{p}^{(n)})$, we put $b_1[d_1] \otimes \dots \otimes b_m[d_m] = \mathcal{C}_n(\bar{p}^{(n)})$ and $(\lambda_1, \dots, \lambda_k) = 2\mu^{(n-1)}$ in (2.22) regarding B_l as the $A_0^{(1)}$ crystal with letter \bar{n} only and B'_l as the $A_1^{(1)}$ crystal with letters n, \bar{n} . By using the resulting $c_1 \otimes \dots \otimes c_k$ in (2.22), we construct $\Phi_n \mathcal{C}_n(\bar{p}^{(n)})$ as

$$(6.3) \quad \Phi_n \mathcal{C}_n(\bar{p}^{(n)}) = \tilde{c}_1 \otimes \dots \otimes \tilde{c}_k \in B_{\lambda_1}^{\geq n} \otimes \dots \otimes B_{\lambda_k}^{\geq n}.$$

Here for B'_l with even l , \tilde{c}_j is determined from c_j as

$$(6.4) \quad \begin{aligned} \tilde{\cdot} : B'_l &\longrightarrow B_l^{\geq n} \\ (x_n, \bar{x}_n) &\longmapsto (0, \dots, 0, z_n, z_0, \bar{z}_n, 0, \dots, 0) \\ z_n = \left\lfloor \frac{x_n}{2} \right\rfloor, \bar{z}_n = \left\lfloor \frac{\bar{x}_n}{2} \right\rfloor, z_0 = l - z_n - \bar{z}_n, \end{aligned}$$

where $\lfloor x \rfloor$ denotes the largest integer not exceeding x .

6.5. Example. Consider the $B_2^{(1)}$ rigged configuration (6.1). Doubling the width of $\mu^{(2)}$, we have

$$\bar{p}^{(2)} = \boxed{\bar{2}\bar{2}} \otimes \boxed{\bar{2}}, \quad \mathcal{C}_2(\bar{p}^{(2)}) =: \boxed{\bar{2}\bar{2}}_3 \otimes \boxed{\bar{2}}_2 := \boxed{\bar{2}}_1 \otimes \boxed{\bar{2}\bar{2}}_4.$$

To find $\Phi_2 \mathcal{C}_2(\bar{p}^{(2)})$ we first need $\mathcal{T}_2^1 \otimes \boxed{\bar{2}} \otimes \mathcal{T}_2^3 \otimes \boxed{\bar{2}\bar{2}} \otimes (\boxed{2222} \otimes \boxed{22} \otimes \boxed{22} \otimes \boxed{22})$.

$$\begin{array}{cccccccc} & 2222 & & 22 & & 22 & & 22 \\ \bar{2}\bar{2} & \begin{array}{c} | \\ \hline \\ | \end{array} & 22 & \begin{array}{c} | \\ \hline \\ | \end{array} & 22 & \begin{array}{c} | \\ \hline \\ | \end{array} & 22 & \begin{array}{c} | \\ \hline \\ | \end{array} & 22 \\ & 22\bar{2}\bar{2} & & 22 & & 22 & & 22 \\ 2 & \begin{array}{c} | \\ \hline \\ | \end{array} & \bar{2} & \begin{array}{c} | \\ \hline \\ | \end{array} & 2 & \begin{array}{c} | \\ \hline \\ | \end{array} & 2 & \begin{array}{c} | \\ \hline \\ | \end{array} & 2 \\ & 222\bar{2} & & \bar{2}\bar{2} & & 22 & & 22 \\ 2 & \begin{array}{c} | \\ \hline \\ | \end{array} & \bar{2} & \begin{array}{c} | \\ \hline \\ | \end{array} & 2 & \begin{array}{c} | \\ \hline \\ | \end{array} & 2 & \begin{array}{c} | \\ \hline \\ | \end{array} & 2 \\ & 2222 & & \bar{2}\bar{2} & & 22 & & 22 \\ 2 & \begin{array}{c} | \\ \hline \\ | \end{array} & 2 & \begin{array}{c} | \\ \hline \\ | \end{array} & \bar{2} & \begin{array}{c} | \\ \hline \\ | \end{array} & 2 & \begin{array}{c} | \\ \hline \\ | \end{array} & 2 \\ & 2222 & & \bar{2}\bar{2} & & \bar{2}\bar{2} & & 22 \\ \bar{2} & \begin{array}{c} | \\ \hline \\ | \end{array} & 2 & \begin{array}{c} | \\ \hline \\ | \end{array} & \bar{2} & \begin{array}{c} | \\ \hline \\ | \end{array} & 2 & \begin{array}{c} | \\ \hline \\ | \end{array} & 2 \\ & 22\bar{2}\bar{2} & & 22 & & \bar{2}\bar{2} & & 22 \\ 2 & \begin{array}{c} | \\ \hline \\ | \end{array} & \bar{2} & \begin{array}{c} | \\ \hline \\ | \end{array} & 2 & \begin{array}{c} | \\ \hline \\ | \end{array} & \bar{2} & \begin{array}{c} | \\ \hline \\ | \end{array} & 2 \\ & 2222 & & \bar{2}\bar{2} & & \bar{2}\bar{2} & & \bar{2}\bar{2} \end{array}$$

Under $\tilde{\cdot}$, the elements $\boxed{2222}$ and $\boxed{\bar{2}\bar{2}}$ turn into $\boxed{22}$ and $\boxed{0}$. Thus we get

$$p^{(1)} = \Phi_2 \mathcal{C}_2(\bar{p}^{(2)}) = \boxed{22} \otimes \boxed{0} \otimes \boxed{0} \otimes \boxed{0},$$

which is an element of $B_1^{(1)}$ crystal with letters $2, 0, \bar{2}$. In view of $\boxed{22} \otimes \boxed{0} \simeq \boxed{2} \otimes \boxed{20}$ and $H(\boxed{22} \otimes \boxed{0}) = H(\boxed{0} \otimes \boxed{0}) = 0$, we have $\mathcal{C}_1(p^{(1)}) = \boxed{22}_2 \otimes \boxed{0}_3 \otimes \boxed{0}_3 \otimes \boxed{0}_3$. $\Phi_1 \mathcal{C}_1(p^{(1)})$ is derived from $\mathcal{T}_1^2 \otimes \boxed{22} \otimes \mathcal{T}_1^1 \otimes \boxed{0}^{\otimes 3} \otimes (\boxed{1}^{\otimes 7})$. Since the combinatorial R is the identity on $B_1 \otimes B_1$, the part $\mathcal{T}_1^1 \otimes \boxed{0}^{\otimes 3}$ just pushes $\boxed{1}$ to the right. $\Phi_1 \mathcal{C}_1(p^{(1)})$ is obtained by putting $\boxed{1}^{\otimes 2}$ in front of the output of the diagram:

$$\begin{array}{cccccccc} & 1 & & 0 & & 0 & & 0 & & 1 & & 1 & & 1 \\ 22 & \begin{array}{c} | \\ \hline \\ | \end{array} & 12 & \begin{array}{c} | \\ \hline \\ | \end{array} & 10 & \begin{array}{c} | \\ \hline \\ | \end{array} & 10 & \begin{array}{c} | \\ \hline \\ | \end{array} & 10 & \begin{array}{c} | \\ \hline \\ | \end{array} & 11 & \begin{array}{c} | \\ \hline \\ | \end{array} & 11 & \begin{array}{c} | \\ \hline \\ | \end{array} & 11 \\ & 2 & & 2 & & 0 & & 0 & & 0 & & 1 & & 1 \end{array}$$

Thus we get $\Phi_1 \mathcal{C}_1(p^{(1)}) = 1122000$ in agreement with (6.2).

7. $C_n^{(1)}$ CASE

7.1. Rigged configurations. Consider the data of the form (2.1) and define $E_j^{(a)}$ as in (2.2). The vacancy number is specified by (2.3) for $1 \leq a \leq n-1$ and

$$(7.1) \quad p_j^{(n)} = E_j^{(n-1)} - E_j^{(n)}.$$

The data (2.1) is called a $C_n^{(1)}$ rigged configuration if $\mu_i^{(n)} \in 2\mathbb{Z}$ and (2.4) is satisfied. It is depicted as in $A_n^{(1)}$ case by the Young diagrams with rigging. The following is an example of $C_2^{(1)}$ rigged configuration.

$$(7.2) \quad \begin{array}{ccc} \mu^{(0)} & \mu^{(1)} & \mu^{(2)} \\ (1^{10}) & 0 \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \end{array} 0 & 2 \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \end{array} 2 \\ & 1 \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array} 1 & 1 \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} 0 \\ & 6 \begin{array}{|c|} \hline \square \\ \hline \end{array} 2 & & \end{array}$$

7.2. Crystals.

$$(7.3) \quad B_l = \{(x_0, \dots, x_n, \bar{x}_n, \dots, \bar{x}_0) \in (\mathbb{Z}_{\geq 0})^{2n+2} \mid \sum_{i=0}^n (x_i + \bar{x}_i) = l, x_0 = \bar{x}_0\}.$$

$$(7.4) \quad B_l^{\geq a} = \{(x_0, \dots, x_n, \bar{x}_n, \dots, \bar{x}_0) \in B_l \mid x_i = \bar{x}_i = 0 \ (0 \leq i \leq a-1), x_a = \bar{x}_a\}$$

for $0 \leq a \leq n$. Then

$$(7.5) \quad B_l = B_l^{\geq 0} \supset B_l^{\geq 1} \supset \dots \supset B_l^{\geq n} = \begin{cases} \emptyset & l : \text{odd} \\ \{(0, \dots, 0, \frac{l}{2}, \frac{l}{2}, 0, \dots, 0)\} & l : \text{even} \end{cases}$$

as sets. We regard $B_l^{\geq a}$ as $C_{n-a}^{(1)}$ crystal and represent the elements by length l semistandard row tableaux over the alphabet $a < \dots < n < \bar{n} < \dots < \bar{a}$. For example for $n=3, a=2$, the highest elements in $B_5^{\geq 2}$ with respect to C_{n-a} are

$$\boxed{33333}, \boxed{2333\bar{2}}, \boxed{223\bar{2}\bar{2}}.$$

The rigged configuration (7.2) corresponds to

$$(7.6) \quad \boxed{1} \otimes \boxed{1} \otimes \boxed{1} \otimes \boxed{2} \otimes \boxed{2} \otimes \boxed{2} \otimes \boxed{2} \otimes \boxed{1} \otimes \boxed{\bar{1}} \otimes \boxed{\bar{1}} \in \mathcal{P}_+((1^{10})).$$

7.3. Maps $\mathcal{C}_1, \dots, \mathcal{C}_n$. Let $B_l = B_l^{\geq a}$ be the $C_{n-a}^{(1)}$ crystal. Then the map \mathcal{C}_a with $a < n$ is defined in the same way as in Section 2.4. Since the energy function is nonnegative, one has $0 \leq d_1 \leq \dots \leq d_m$ for normal ordered forms $b_1[d_1] \otimes \dots \otimes b_m[d_m]$. Define the $C_0^{(1)}$ crystal element

$$(7.7) \quad \bar{p}^{(n)} = (n\bar{n})^{\mu_1^{(n)}/2} \otimes \dots \otimes (n\bar{n})^{\mu_n^{(n)}/2},$$

where $(n\bar{n})^{l/2}$ denotes the unique element in $B_l^{\geq n}$ with even l in (7.5). (Recall that $\mu_i^{(n)}$ is even for $C_n^{(1)}$ rigged configuration.) Then $\mathcal{C}_n(\bar{p}^{(n)})$ is obtained by assigning the mode in (2.18) as $d_i = J_i + \mu_i/2$ instead of (2.19). For normal ordering we need the $C_0^{(1)}$ combinatorial R . It is formally specified by saying that $H = 0$ on $B_l \otimes B_m$ and $b[d] \otimes b'[d'] \simeq b'[d'] \otimes b[d]$.

7.4. Maps Φ_1, \dots, Φ_n . The map Φ_a is defined in the same way as in Section 2.5 if B_l and B_l' there are replaced by $B_l = B_l^{\geq a}$ and $B_l' = B_l^{\geq a-1}$, respectively.

7.5. **Example.** We start from the $C_2^{(1)}$ rigged configuration (7.2). Then

$$\bar{p}^{(2)} = \boxed{22\bar{2}\bar{2}} \otimes \boxed{2\bar{2}}, \quad C_2(\bar{p}^{(2)}) = \boxed{2\bar{2}}_1 \otimes \boxed{22\bar{2}\bar{2}}_4.$$

In order to find $\Phi_2 C_2(\bar{p}^{(2)})$ we are to compute

$$\mathcal{T}_2^1 \otimes \boxed{2\bar{2}} \otimes \mathcal{T}_2^3 \otimes \boxed{22\bar{2}\bar{2}} \otimes (\boxed{2222} \otimes \boxed{222} \otimes \boxed{2})$$

for $C_1^{(1)}$ crystal with letters $1, 2, \bar{2}, \bar{1}$. From the diagram

$$\begin{array}{cccc} & 2222 & 222 & 2 \\ 22\bar{2}\bar{2} & \text{---} & 2222 & \text{---} & 2222 & \text{---} & 2222 \\ & 22\bar{2}\bar{2} & 222 & 2 \\ 2 & \text{---} & 2 & \text{---} & 2 & \text{---} & 2 \\ & 12\bar{2}\bar{1} & 222 & 2 \\ 2 & \text{---} & 2 & \text{---} & 2 & \text{---} & 2 \\ & 11\bar{1}\bar{1} & 222 & 2 \\ 2 & \text{---} & \bar{2} & \text{---} & 2 & \text{---} & 2 \\ & 122\bar{1} & 22\bar{2} & 2 \\ 2\bar{2} & \text{---} & 1\bar{1} & \text{---} & 22 & \text{---} & 22 \\ & 222\bar{2} & 1\bar{2}\bar{1} & 2 \\ 2 & \text{---} & 2 & \text{---} & \bar{2} & \text{---} & 2 \\ & 122\bar{1} & 1\bar{2}\bar{1} & \bar{2} \end{array}$$

we get $p^{(1)} = \Phi_2 C_2(\bar{p}^{(2)}) = \boxed{122\bar{1}} \otimes \boxed{1\bar{2}\bar{1}} \otimes \boxed{2}$. Assigning the mode to $p^{(1)}$ we have $\mathcal{C}_1(p^{(1)}) = : \boxed{122\bar{1}}_3 \otimes \boxed{1\bar{2}\bar{1}}_4 \otimes \boxed{2}_3 :$, where the normal ordering is to be done along the $C_1^{(1)}$ crystal with letters $1, 2, \bar{2}, \bar{1}$. There are four normal ordered forms:

$$\begin{aligned} & \boxed{2}_3 \otimes \boxed{22\bar{2}}_3 \otimes \boxed{11\bar{1}\bar{1}}_4, \quad \boxed{2}_3 \otimes \boxed{222\bar{2}}_3 \otimes \boxed{1\bar{2}\bar{1}}_4 \\ & \boxed{1\bar{2}\bar{1}}_3 \otimes \boxed{2}_3 \otimes \boxed{11\bar{1}\bar{1}}_4, \quad \boxed{122\bar{1}}_3 \otimes \boxed{2}_3 \otimes \boxed{1\bar{2}\bar{1}}_4. \end{aligned}$$

We pick the first one as $\mathcal{C}_1(p^{(1)})$. Then $\Phi_1 \mathcal{C}_1(p^{(1)})$ is calculated from

$$\mathcal{T}_1^3 \otimes \boxed{2} \otimes \boxed{22\bar{2}} \otimes \mathcal{T}_1^1 \otimes \boxed{11\bar{1}\bar{1}} \otimes (\boxed{1}^{\otimes 10})$$

where for an array $\lambda = (\lambda_i)$, 2λ means $(2\lambda_i)$. In the $D_{n+1}^{(2)}$ case, all the rigging except the n -th one $J^{(n)}$ are doubled.

8.2. Crystals. For $A_{2n}^{(2)}$,

$$(8.6) \quad B_l = \{(x_1, \dots, x_n, x_\emptyset, \bar{x}_n, \dots, \bar{x}_1) \in (\mathbb{Z}_{\geq 0})^{2n+1} \mid x_\emptyset + \sum_{i=1}^n (x_i + \bar{x}_i) = l\}.$$

For $D_{n+1}^{(2)}$,

$$(8.7) \quad B_l = \{(x_1, \dots, x_n, x_0, x_\emptyset, \bar{x}_n, \dots, \bar{x}_1) \in (\mathbb{Z}_{\geq 0})^{2n+2} \mid x_0 + x_\emptyset + \sum_{i=1}^n (x_i + \bar{x}_i) = l, x_0 = 0, 1\}.$$

There is an embedding $(B_l \text{ for } \mathfrak{g}_n) \rightarrow (B_{2l} \text{ for } C_n^{(1)})$ as sets, which will also be denoted by ι :

$$(8.8) \quad \iota(x) = \begin{cases} (x_\emptyset, 2x_1, \dots, 2x_n, 2\bar{x}_n, \dots, 2\bar{x}_1, x_\emptyset) & \text{for } A_{2n}^{(2)}, \\ (x_\emptyset, 2x_1, \dots, 2x_{n-1}, 2x_n + x_0, 2\bar{x}_n + x_0, 2\bar{x}_{n-1}, \dots, 2\bar{x}_1, x_\emptyset) & \text{for } D_{n+1}^{(2)}. \end{cases}$$

Here $x = (x_1, \dots, \bar{x}_1) \in B_l$ is specified as in (8.6) and (8.7). When applying ι^{-1} , the tableau letters are halved (in case the image exists). In particular, a pair of 0 and $\bar{0}$ turns into \emptyset . As for n and \bar{n} , the change is described by (6.4). We extend the map ι to the tensor product of \mathfrak{g}_n crystals by $\iota(p_1 \otimes \dots \otimes p_k) = \iota(p_1) \otimes \dots \otimes \iota(p_k)$.

The $A_4^{(2)}$ rigged configuration (8.3) corresponds to

$$(8.9) \quad \boxed{1} \otimes \boxed{1} \otimes \boxed{\emptyset} \otimes \boxed{\bar{1}} \otimes \boxed{1} \otimes \boxed{\emptyset} \otimes \boxed{2} \otimes \boxed{2} \in \mathcal{P}_+((1^8)).$$

The $D_3^{(2)}$ rigged configuration (8.4) corresponds to

$$(8.10) \quad \boxed{1} \otimes \boxed{\emptyset} \otimes \boxed{2} \otimes \boxed{1} \otimes \boxed{0} \otimes \boxed{1} \otimes \boxed{2} \otimes \boxed{2} \otimes \boxed{\emptyset} \otimes \boxed{0} \in \mathcal{P}_+((1^{10})).$$

8.3. Example. Consider the $A_4^{(2)}$ rigged configuration (8.3). According to (8.5) we double everything horizontally to get

$$(8.11) \quad \begin{array}{ccc} \mu^{(0)} & \mu^{(1)} & \mu^{(2)} \\ \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}^8 & \begin{array}{|c|c|c|c|c|c|c|c|} \hline 0 & \square & \square & \square & \square & \square & \square & 0 \\ \hline 6 & \square & \square & \square & 4 & \square & \square & \square \\ \hline \end{array} & \begin{array}{|c|c|c|c|c|c|} \hline 2 & \square & \square & \square & \square & 0 \\ \hline 0 & \square & \square & \square & 0 & \square \\ \hline \end{array} \end{array}$$

We regard this as a $C_2^{(1)}$ rigged configuration. The vacancy numbers are determined by (7.1) which is the same as (8.1). As the result they are all doubled as well. From (7.7) we set

$$\bar{p}^{(2)} = \boxed{222222} \otimes \boxed{22}, \quad C_2(\bar{p}^{(2)}) = \boxed{22}_1 \otimes \boxed{222222}_3.$$

Then $p^{(1)} := \Phi_2 C_2(\bar{p}^{(2)})$ is determined as

$$p^{(1)} = \boxed{11122\bar{1}\bar{1}} \otimes \boxed{122\bar{1}}, \quad C_1(p^{(1)}) = \boxed{11122\bar{1}\bar{1}}_5 \otimes \boxed{122\bar{1}}_7.$$

$p = \Phi_1 C_1(p^{(1)})$ reads

$$p = \boxed{11} \otimes \boxed{11} \otimes \boxed{00} \otimes \boxed{\bar{1}\bar{1}} \otimes \boxed{11} \otimes \boxed{00} \otimes \boxed{22} \otimes \boxed{22}.$$

Finally applying ι^{-1} in (8.8), we find $\iota^{-1}(p) = 11\emptyset\bar{1}1\emptyset22$, reproducing (8.9).

Next we consider the $D_3^{(2)}$ rigged configuration (8.4). According to (8.5) we double everything horizontally except the rigging attached to $\mu^{(2)}$:

$$(8.12) \quad \begin{array}{ccc} \mu^{(0)} & \mu^{(1)} & \mu^{(2)} \\ \begin{array}{c} \boxed{} \boxed{} \\ 10 \end{array} & \begin{array}{c} \begin{array}{cccc} \boxed{} & \boxed{} & \boxed{} & \boxed{} \\ \boxed{} & \boxed{} & \boxed{} & \boxed{} \\ \boxed{} & \boxed{} & \boxed{} & \boxed{} \\ \boxed{} & \boxed{} & \boxed{} & \boxed{} \end{array} \begin{array}{c} 0 \\ 2 \\ 2 \\ 4 \end{array} \\ 2 \\ 10 \end{array} & \begin{array}{c} \begin{array}{cccc} \boxed{} & \boxed{} & \boxed{} & \boxed{} \\ \boxed{} & \boxed{} & \boxed{} & \boxed{} \\ \boxed{} & \boxed{} & \boxed{} & \boxed{} \\ \boxed{} & \boxed{} & \boxed{} & \boxed{} \end{array} \begin{array}{c} 0 \\ 2 \\ 3 \\ 1 \end{array} \\ 2 \\ 4 \\ 2 \end{array} \end{array}$$

We regard this as a $C_2^{(1)}$ rigged configuration. The vacancy numbers determined by (7.1) instead of (8.2) have been doubled except $p_j^{(2)}$'s which remain unchanged. From (7.7) we set

$$\bar{p}^{(2)} = \boxed{222222} \otimes \boxed{2222} \otimes \boxed{22}, \quad C_2(\bar{p}^{(2)}) = \boxed{22}_2 \otimes \boxed{222222}_3 \otimes \boxed{2222}_5.$$

Then $p^{(1)} = \Phi_2 C_2(\bar{p}^{(2)})$ is determined as

$$p^{(1)} = \boxed{122\bar{1}} \otimes \boxed{2222} \otimes \boxed{122\bar{1}} \otimes \boxed{22}.$$

There are two normal ordered forms for $C_1(p^{(1)})$:

$$\boxed{122\bar{1}}_3 \otimes \boxed{22}_6 \otimes \boxed{2222}_6 \otimes \boxed{122\bar{1}}_7, \quad \boxed{122\bar{1}}_3 \otimes \boxed{2222}_6 \otimes \boxed{22}_6 \otimes \boxed{122\bar{1}}_7.$$

Both of them lead to

$$p = \Phi_1 C_1(p^{(1)}) = \boxed{11} \otimes \boxed{00} \otimes \boxed{22} \otimes \boxed{11} \otimes \boxed{22} \otimes \boxed{11} \otimes \boxed{22} \otimes \boxed{22} \otimes \boxed{00} \otimes \boxed{22}.$$

Thus we obtain $\iota^{-1}(p) = 1 \emptyset 2 1 0 1 \bar{2} 2 \emptyset 0$, in agreement with (8.10).

APPENDIX A. CRYSTALS AND COMBINATORIAL R

The crystals B_l used in the main text are crystal bases of irreducible finite-dimensional representations of a quantum affine algebra $U'_q(\mathfrak{g})$. Let us recall basic facts on them following [5, 10, 27].

Let P be the weight lattice, $\{\alpha_i\}_{0 \leq i \leq n}$ the simple roots, and $\{\Lambda_i\}_{0 \leq i \leq n}$ the fundamental weights of \mathfrak{g} . A crystal B is a finite set with weight decomposition $B = \sqcup_{\lambda \in P} B_\lambda$. The Kashiwara operators \tilde{e}_i, \tilde{f}_i ($i = 0, 1, \dots, n$) act on B as $\tilde{e}_i : B_\lambda \rightarrow B_{\lambda + \alpha_i} \sqcup \{0\}$, $\tilde{f}_i : B_\lambda \rightarrow B_{\lambda - \alpha_i} \sqcup \{0\}$. In particular, these operators are nilpotent. By definition, we have $\tilde{f}_i b = b'$ if and only if $b = \tilde{e}_i b'$. For any $b \in B$, set $\varepsilon_i(b) = \max\{m \geq 0 \mid \tilde{e}_i^m b \neq 0\}$ and $\varphi_i(b) = \max\{m \geq 0 \mid \tilde{f}_i^m b \neq 0\}$. Then we have the weight $\text{wt} b$ of b by $\text{wt} b = \sum_{i=0}^n (\varphi_i(b) - \varepsilon_i(b)) \Lambda_i$.

For two crystals B and B' , one can define the tensor product $B \otimes B' = \{b \otimes b' \mid b \in B, b' \in B'\}$. The operators \tilde{e}_i, \tilde{f}_i act on $B \otimes B'$ by

$$\begin{aligned} \tilde{e}_i(b \otimes b') &= \begin{cases} \tilde{e}_i b \otimes b' & \text{if } \varphi_i(b) \geq \varepsilon_i(b') \\ b \otimes \tilde{e}_i b' & \text{if } \varphi_i(b) < \varepsilon_i(b'), \end{cases} \\ \tilde{f}_i(b \otimes b') &= \begin{cases} \tilde{f}_i b \otimes b' & \text{if } \varphi_i(b) > \varepsilon_i(b') \\ b \otimes \tilde{f}_i b' & \text{if } \varphi_i(b) \leq \varepsilon_i(b'). \end{cases} \end{aligned}$$

Here $0 \otimes b'$ and $b \otimes 0$ should be understood as 0. For crystals we are considering, there exists a unique isomorphism $B \otimes B' \xrightarrow{\sim} B' \otimes B$, *i.e.* a unique map which commutes with the action of Kashiwara operators. In particular, it preserves the weight.

For a crystal B we define its affinization $\text{Aff}(B) = \{b[d] \mid d \in \mathbb{Z}, b \in B\}$ by $\tilde{e}_i(b[d]) = (\tilde{e}_i b)[d - \delta_{i0}]$ and $\tilde{f}_i(b[d]) = (\tilde{f}_i b)[d + \delta_{i0}]$. ($b[d]$ here corresponds to $T^{-d} a f(b)$ in [10].)

The crystal isomorphism $B \otimes B' \xrightarrow{\sim} B' \otimes B$ is lifted up to a map $\text{Aff}(B) \otimes \text{Aff}(B') \xrightarrow{\sim} \text{Aff}(B') \otimes \text{Aff}(B)$ called the combinatorial R . It has the following form:

$$\begin{aligned} R : \text{Aff}(B) \otimes \text{Aff}(B') &\longrightarrow \text{Aff}(B') \otimes \text{Aff}(B) \\ b[d] \otimes b'[d'] &\longmapsto \tilde{b}'[d' - H(b \otimes b')] \otimes \tilde{b}[d + H(b \otimes b')], \end{aligned}$$

where $b \otimes b' \mapsto \tilde{b}' \otimes \tilde{b}$ under the isomorphism $B \otimes B' \xrightarrow{\sim} B' \otimes B$. $H(b \otimes b')$ is called the energy function and determined up to an additive constant by

$$H(\tilde{e}_i(b \otimes b')) = \begin{cases} H(b \otimes b') + 1 & \text{if } i = 0, \varphi_0(b) \geq \varepsilon_0(b'), \varphi_0(\tilde{b}') \geq \varepsilon_0(\tilde{b}), \\ H(b \otimes b') - 1 & \text{if } i = 0, \varphi_0(b) < \varepsilon_0(b'), \varphi_0(\tilde{b}') < \varepsilon_0(\tilde{b}), \\ H(b \otimes b') & \text{otherwise.} \end{cases}$$

Proposition A.1 (Yang-Baxter equation). *The following equation holds on $\text{Aff}(B) \otimes \text{Aff}(B') \otimes \text{Aff}(B'')$:*

$$(R \otimes 1)(1 \otimes R)(R \otimes 1) = (1 \otimes R)(R \otimes 1)(1 \otimes R).$$

We often write the map R simply by \simeq . The combinatorial R is naturally restricted to $B \otimes B'$.

In the main text we are concerned about the crystal B_l corresponding to the l -fold symmetric fusion of the vector representation. We normalize the energy function so that

$$(A.1) \quad \max\{H(b \otimes c) \mid b \otimes c \in B_l \otimes B_m\} = \min(l, m).$$

Under this convention one has

$$(A.2) \quad \min\{H(b \otimes c) \mid b \otimes c \in B_l \otimes B_m\} = \begin{cases} 0 & \mathfrak{g}_n = A_n^{(1)}, C_n^{(1)}, \\ -\min(l, m) & \mathfrak{g}_n \neq A_n^{(1)}, C_n^{(1)}. \end{cases}$$

When $l = m$, the combinatorial R becomes the identity map on $B_l \otimes B_l$ but still acts non-trivially as $R(x[d] \otimes y[e]) = x[e - H(x \otimes y)] \otimes y[d + H(x \otimes y)]$.

Acknowledgments. The authors thank Anne Schilling and Mark Shimozono for kind interest and comments. A.K. and Y.Y. are supported by Grants-in-Aid for Scientific Research JSPS No.15540363 and No.17340047, respectively. R.S. is grateful to Miki Wadati for warm encouragement during the study.

REFERENCES

- [1] H. A. Bethe, Zur Theorie der Metalle, I. Eigenwerte und Eigenfunktionen der linearen Atomkette, *Z. Physik* **71** (1931) 205–231.
- [2] R. J. Baxter, *Exactly solved models in statistical mechanics*, Academic Press, London (1982).
- [3] S. V. Kerov, A. N. Kirillov and N. Yu. Reshetikhin, Combinatorics, the Bethe ansatz and representations of the symmetric group. *J. Soviet Math.* **41** (1988) 916–924.
- [4] A. N. Kirillov and N. Yu. Reshetikhin, The Bethe ansatz and the combinatorics of Young tableaux. *J. Soviet Math.* **41** (1988) 925–955.
- [5] M. Kashiwara, On crystal bases of the q -analogue of universal enveloping algebras. *Duke Math. J.* **63** (1991) 465–516.
- [6] G. Hatayama, A. Kuniba, M. Okado, T. Takagi and Y. Yamada, Remarks on fermionic formula, *Contemp. Math.* **248** (AMS 1999) 243–291.
- [7] G. Hatayama, A. Kuniba, M. Okado, T. Takagi and Z. Tsuboi: Paths, Crystals and Fermionic Formulae, *Prog. in Math. Phys.* **23** (2002) 205–272.
- [8] A. Schilling, X=M Theorem: Fermionic formulas and rigged configurations under review preprint, [math.QA/0512161](https://arxiv.org/abs/math.QA/0512161).
- [9] G. Hatayama, A. Kuniba, M. Okado, T. Takagi and Y. Yamada, Scattering rules in soliton cellular automata associated with crystal bases, *Contemporary Math.* **297** (2002) 151–182.

- [10] S.-J. Kang, M. Kashiwara, K. C. Misra, T. Miwa, T. Nakashima and A. Nakayashiki, Affine crystals and vertex models, *Int. J. Mod. Phys. A* **7** (suppl. 1A), (1992) 449–484.
- [11] A. Nakayashiki and Y. Yamada, Kostka polynomials and energy functions in solvable lattice models, *Selecta Mathematica, New Ser.* **3** (1997) 547–599.
- [12] C. L. Schultz, Eigenvectors of the multicomponent generalization of the six-vertex model, *Physica.* **A122** (1983) 71–88.
- [13] D. Takahashi and J. Satsuma, A soliton cellular automaton, *J. Phys. Soc. Jpn.* **59** (1990) 3514–3519.
- [14] D. Takahashi, On some soliton systems defined by using boxes and balls, *Proceedings of the International Symposium on Nonlinear Theory and Its Applications (NOLTA '93)*, (1993) 555–558.
- [15] G. Hatayama, K. Hikami, R. Inoue, A. Kuniba, T. Takagi and T. Tokihiro, The $A_M^{(1)}$ automata related to crystals of symmetric tensors, *J. Math. Phys.* **42** (2001) 274–308.
- [16] K. Fukuda, M. Okado, Y. Yamada, Energy functions in box ball systems, *Int. J. Mod. Phys. A* **15** (2000) 1379–1392.
- [17] R. Sakamoto, Crystal interpretation of Kerov-Kirillov-Reshetikhin bijection II. Proof for \mathfrak{sl}_n Case, preprint.
- [18] M. Okado, A. Schilling and M. Shimozono, A crystal to rigged configuration bijection for nonexceptional affine algebras, *Algebraic Combinatorics and Quantum Groups*, Ed by N. Jing, World Scientific (2003), 85–124.
- [19] A. Schilling and M. Shimozono, X=M for symmetric powers, *J. Alg.* **295** (2006) 562–610.
- [20] T. Takagi, Separation of colour degree of freedom from dynamics in a soliton cellular automaton, *J. Phys. A* **38** (2005) 1961–1976.
- [21] A. N. Kirillov, A. Schilling and M. Shimozono, A bijection between Littlewood-Richardson tableaux and rigged configurations. *Selecta Math.* **8** (2002) 67–135.
- [22] M. Shimozono, Affine type A crystal structure on tensor products of rectangles, Demazure characters, and nilpotent varieties, *J. Algebraic Combin.* **15** (2002) 151–187.
- [23] G. Hatayama, A. Kuniba, M. Okado and T. Takagi, Combinatorial R matrices for a family of crystals: $C_n^{(1)}$ and $A_{2n-1}^{(2)}$ cases, *Prog. in Math.* **191** (2000) 105–139.
- [24] G. Hatayama, A. Kuniba, M. Okado and T. Takagi, Combinatorial R matrices for a family of crystals: $B_n^{(1)}$, $D_n^{(1)}$, $A_{2n}^{(2)}$, and $D_{n+1}^{(2)}$ cases, *J. Alg.* **247** (2002) 577–615.
- [25] A. Kuniba, M. Okado, T. Takagi and Y. Yamada, Geometric crystal and tropical R for $D_n^{(1)}$, *Int. Math. Res. Notices* **48** (2003) 2565–2620.
- [26] A. Kuniba, T. Takagi and A. Takenouchi, Factorization, reduction and embedding in integrable cellular automata, *J. Phys. A.* **37** (2004) 1691–1709.
- [27] S.-J. Kang, M. Kashiwara and K. C. Misra, Crystal bases of Verma modules for quantum affine Lie algebras, *Compositio Math.* **92** (1994) 299–325.

Atsuo Kuniba:

Institute of Physics, Graduate School of Arts and Sciences, University of Tokyo,
Komaba, Tokyo 153-8902, Japan
atsuo@gokutan.c.u-tokyo.ac.jp

Masato Okado:

Department of Mathematical Science, Graduate School of Engineering Science,
Osaka University, Osaka 560-8531, Japan
okado@sigmath.es.osaka-u.ac.jp

Reiho Sakamoto:

Department of Physics, Graduate School of Science, University of Tokyo, Hongo,
Tokyo 113-0033, Japan
reiho@monet.phys.s.u-tokyo.ac.jp

Taichiro Takagi:

Department of Applied Physics, National Defense Academy, Kanagawa 239-8686,
Japan
`takagi@nda.ac.jp`

Yasuhiko Yamada:
Department of Mathematics, Faculty of Science, Kobe University, Hyogo 657-8501,
Japan
`yamaday@math.kobe-u.ac.jp`