

On the q -analogue of the Maurer-Cartan equation

Mauricio Angel and Rafael Díaz

August 27, 2018

Abstract

We consider deformations of the differential of a q -differential graded algebra. We prove that it is controlled by a generalized Maurer-Cartan equation. We find explicit formulae for the coefficients c_k involved in that equation.

Introduction

Following the work of Kapranov [6] the interest on the theory of N -complexes has risen. A delicate issue in the theory is the appropriate definition of a N -differential algebra. At the moment there exists two options for such a definition, each describing very distinct mathematical phenomena.

In [2] the notion of N -differential graded algebra A has been defined as follows: A must be a graded associative algebra provided with an operator $d : A \rightarrow A$ of degree 1 such that $d(ab) = d(a)b + (-1)^{\bar{a}}ad(b)$ and $d^N = 0$.

The study of the deformations of the differential of a N -dga has also been initiated in [2]. Applications to differential geometry are given in [3]. Applications to the study of A_∞ -algebras of depth N , are contained in [4].

The other possible N -generalization of the notion of a differential algebra (see [7],[8],[9]) goes as follows: One fixed a primitive N -th root of unity q . One considers q -differential graded algebras A defined by: A must be a graded associative algebra provided with an operator $d : A \rightarrow A$ of degree 1 such that $d(ab) = d(a)b + q^{\bar{a}}ad(b)$ and $d^N = 0$.

In this paper we consider deformations of the differential of a q -differential graded algebra. We prove that it is controlled by a generalized Maurer-Cartan equation. We find explicit formulae for the coefficients c_k involved in that equation.

1 q -differential graded algebras and modules

Let \mathbf{k} be a commutative ring with unity. For $q \in \mathbf{k}$ the q -numbers are given by $[]_q : \mathbb{N} \rightarrow \mathbf{k}$, $k \mapsto [k]_q = 1 + q + \cdots + q^{k-1}$ and $[k]_q! = [1]_q [2]_q \cdots [k]_q$. We assume that on \mathbf{k} , for $N \in \mathbb{N}$ fixed, we can choose $q \in \mathbf{k}$ such that $[N]_q = 0$ and $[k]_q$ is invertible for $1 \leq k \leq N-1$, in this case we refer to q as a primitive N -th root of unity.

Definition 1 A q -differential graded algebra or q -dga over \mathbf{k} is a triple (A^\bullet, m, d) where $m : A^k \otimes A^l \rightarrow A^{k+l}$ and $d : A^k \rightarrow A^{k+1}$ are \mathbf{k} -modules morphisms satisfying

- 1) (A^\bullet, m) is a graded associative \mathbf{k} -algebra.
- 2) d satisfies the q -Leibniz rule $d(ab) = d(a)b + q^{\bar{a}}ad(b)$.
- 3) If q is a primitive N -th root of unity, then $d^N = 0$, i.e., (A^\bullet, d) is a N -complex.

Definition 2 Let (A^\bullet, m_A, d_A) be a q -dga and M^\bullet be a \mathbf{k} -module. A q -differential module (q -dgm) over (A^\bullet, m_A, d_A) is a triple (M^\bullet, m_M, d_M) where $m_M : A^k \otimes M^l \rightarrow M^{k+l}$ and $d_M : M^k \rightarrow M^{k+1}$ are \mathbf{k} -module morphisms satisfying the following properties

1. $m_M(a, m_M(b, \phi)) = m_M(m_A(a, b), \phi)$, for all $a, b \in A^\bullet$ and $\phi \in M^\bullet$. We denote $m_M(a, \phi)$ by $a\phi$, for all $a \in A$ and $\phi \in M$.
2. $d_M(a\phi) = d_A(a)\phi + q^{\bar{a}}ad_M(\phi)$ for all $a \in A^\bullet$ and $\phi \in M^\bullet$.
3. If q is a primitive N -th root of unity, then $d_M^N = 0$.

2 N -curvature

Let (M^\bullet, m_M, d_M) be a q -dgm over a q -dga (A^\bullet, m_A, d_A) . For $a \in M^1$ we define a deformation of d_M as follows

$$D\phi = d_M\phi + a\phi, \quad \text{for all } \phi \in M^\bullet.$$

In order to obtain a q -dgm structure on (M^\bullet, m_M, D) we require that $D^N = 0$. In [KN] the consecutive powers of D are calculated and the following formula is obtained

$$D^N\phi = d_M^N\phi + \sum_{k=1}^{N-1} \binom{N}{k}_q (D^{k-1}a)d^{N-k}\phi + (D^{N-1}a)\phi,$$

where $\binom{N}{k}_q$ is the q -binomial coefficients given by

$$\binom{N}{k}_q = \frac{[N]_q!}{[N-k]_q![k]_q!}.$$

For q a primitive N -th root of unity the formula above reduces to

$$D^N \phi = (D^{N-1} a) \phi, \quad (1)$$

because in this case $d_M^N = 0$ and $\binom{N}{k}_q = 0$ for $0 < k < N$, since $[N]_q = 1 + q + \dots + q^{N-1} = \frac{1-q^N}{1-q} = 0$.

Formula (1) is the first step towards the construction of the q -analogue of the Maurer-Cartan equation. A further step is required in order to write down $D^{N-1} a$ in terms of d_M and a only. We shall work with a slightly more general problem: finding an explicit expression for $D^N \phi$ given in terms of d_M and a only. First we review some notations from [AD].

For $s = (s_1, \dots, s_n) \in \mathbb{N}^n$ we set $l(s) = n$ and $|s| = \sum_i s_i$. For $1 \leq i < n$, $s_{>i}$ denotes the vector given by $s_{>i} = (s_{i+1}, \dots, s_n)$, for $1 < i \leq n$, $s_{<i}$ stands for $s_{<i} = (s_1, \dots, s_{i-1})$, we also set $s_{>n} = s_{<1} = \emptyset$. $\mathbb{N}^{(\infty)}$ denotes the set $\bigsqcup_{n=0}^{\infty} \mathbb{N}^n$. By convention $\mathbb{N}^0 = \{\emptyset\}$.

For $e \in \text{End}(M^\bullet)$ and $s \in \mathbb{N}^n$ we define $e^{(s)} = e^{(s_1)} \dots e^{(s_n)}$, where $e^{(l)} = d_{\text{End}}^l(e)$ if $l \geq 1$, $e^{(0)} = e$ and $e^\emptyset = 1$. In the case that $e_a \in \text{End}(M^\bullet)$ is given by

$$e_a(\phi) = a\phi, \quad \text{for } a \in M^1 \text{ fixed and all } \phi \in M^\bullet,$$

then $e_a^{(l)} = d_{\text{End}}^l(e_a)$ reduces to $e_a^{(l)} = e_{d^l(a)}$, thus

$$e_a^{(s)} = e_a^{(s_1)} \dots e_a^{(s_n)} = e_{d^{s_1}(a)} \dots e_{d^{s_n}(a)}.$$

For $N \in \mathbb{N}$ we define $E_N = \{s \in \mathbb{N}^{(\infty)} : |s| + l(s) \leq N\}$ and for $s \in E_N$ we define $N(s) \in \mathbb{Z}$ by $N(s) = N - |s| - l(s)$.

We introduce a discrete quantum mechanical ¹ by

1. $V = \mathbb{N}^{(\infty)}$.

¹a discrete quantum mechanical system is given by the following data (1) A directed graph Γ (finite or infinite). (2) A map $L : E_\Gamma \rightarrow \mathbb{R}$ called the Lagrangian map of the system. An associated Hilbert space $\mathcal{H} = \mathbb{C}^{V_\Gamma}$. Operators $U_n : \mathcal{H} \rightarrow \mathcal{H}$, where $n \in \mathbb{Z}$, given by

$$(U_n f)(y) = \sum_{x \in V_\Gamma} \omega_n(y, x) f(x),$$

where the discretized kernel $\omega_n(y, x)$ admits the following representation

$$\omega_n(y, x) = \sum_{\gamma \in P_n(\Gamma, x, y)} \prod_{e \in \gamma} e^{iL(e)}.$$

$P_n(\Gamma, x, y)$ denotes the set of length n paths in Γ from x to y , i.e., sequences (e_1, \dots, e_n) of edges in Γ such that $s(e_1) = x$, $t(e_i) = s(e_{i+1})$, $i = 1, \dots, n-1$ and $t(e_n) = y$.

2. There is a unique directed edge e from vertex s to t if and only if $t \in \{(0, s), s, (s + e_i)\}$ where $e_i = (0, \dots, \underbrace{1}_{i\text{-th}}, \dots, 0) \in \mathbb{N}^{l(s)}$, in this case we set $source(e) = s$ and $target(e) = t$.
3. Edges are weighted according to the following table

$source(e)$	$target(e)$	$v(e)$
s	$(0, s)$	1
s	s	$q^{ s +l(s)}$
s	$(s + e_i)$	$q^{ s +i-1}$

The set $P_N(\emptyset, s)$ consists of all paths $\gamma = (e_1, \dots, e_N)$, such that $source(e_1) = \emptyset$, $target(e_N) = s$ and $source(e_{l+1}) = target(e_l)$. For $\gamma \in P_N(\emptyset, s)$ we define the weight $v(\gamma)$ of γ as

$$v(\gamma) = \prod_{l=1}^N v(e_l).$$

The following result is proven as Theorem 17 in [AD].

Theorem 3 *Let (M^\bullet, m_M, d_M) be a q -dgm over a q -dga (A^\bullet, m_A, d_A) . For $a \in M^1$ consider the map $D\phi = d_M\phi + a\phi$. Then we have*

$$D^N = \sum_{k=0}^{N-1} c_k d_M^k$$

where

$$c_k = \sum_{\substack{s \in E_N \\ N(s)=k \\ s_i < N}} c_q(s, N) a^{(s)} \quad \text{and} \quad c_q(s, N) = \sum_{\gamma \in P_N(\emptyset, s)} v(\gamma).$$

Suppose that we have a q -dgm (M^\bullet, m_M, d_M) , q a 3-rd primitive root of unity, and we want to deform it to $D = d_M + a$, $a \in M^1$. So we required that $D^3 = 0$. By Theorem 3 we must have $\sum_{k=0}^2 c_k d_M^k = 0$. Let us compute the coefficients c_k . First, notice that

$$E_3 = \{\emptyset, (0), (1), (2), (0, 0), (1, 0), (0, 1), (0, 0, 0)\}$$

We proceed to compute the coefficients c_k for $0 \leq k \leq 2$ using Theorem 3.

$k = 0$

There are four vectors in E_3 such that $N(s) = 0$, these are $(2), (1, 0), (0, 1), (0, 0, 0)$.

For $s = (2)$ the only path from \emptyset to (2) of length 3 is $\emptyset \rightarrow (0) \rightarrow (1) \rightarrow (2)$ with weight 1 and $e_a^{(2)} = d_M^2(e_a)$, thus $c(s, 3) = d_M^2(a)$.

For $s = (1, 0)$ the only path from \emptyset to $(1, 0)$ of length 3 is $\emptyset \rightarrow (0) \rightarrow (0, 0) \rightarrow (1, 0)$ with weight 1 and $e_a^{(1,0)} = d_M(e_a)e_a$, thus $c(s, 3) = d_M(a)e_a$.

For $s = (0, 1)$ the paths from \emptyset to $(0, 1)$ of length 3 are $\emptyset \rightarrow (0) \rightarrow (0, 0) \rightarrow (0, 1)$ with weight q ; and $\emptyset \rightarrow (0) \rightarrow (1) \rightarrow (0, 1)$ with weight 1. Since $e_a^{(0,1)} = e_a d_M(e_a)$, then $c(s, 3) = (1 + q)e_a d_M(e_a)$.

For $s = (0, 0, 0)$ the only path from \emptyset to $(0, 0, 0)$ of length 3 is $\emptyset \rightarrow (0) \rightarrow (0, 0) \rightarrow (0, 0, 0)$ whose weight is 1 and $e_a^{(0,0,0)} = a^3$, thus $c(s, 3) = a^3$.

$k = 1$

There are two vectors in E_3 such that $N(s) = 1$, namely (1) and $(0, 0)$.

For $s = (1)$, the paths from \emptyset to (1) of length 3 are $\emptyset \rightarrow \emptyset \rightarrow (0) \rightarrow (1)$ with weight 1; $\emptyset \rightarrow (0) \rightarrow (0) \rightarrow (1)$ with weight q ; $\emptyset \rightarrow (0) \rightarrow (1) \rightarrow (1)$ with weight q^2 . $e_a^{(1)} = d_M(e_a)$ and thus $c_q(s, 3) = (1 + q + q^2) = 0$.

For $s = (0, 0)$ the paths from \emptyset to $(0, 0)$ of length 3 are $\emptyset \rightarrow (0) \rightarrow (0) \rightarrow (0, 0)$ with weight q . $\emptyset \rightarrow \emptyset \rightarrow (0) \rightarrow (0, 0)$ with weight 1. $\emptyset \rightarrow (0) \rightarrow (0, 0) \rightarrow (0, 0)$ with weight q^2 . $e_a^{(0,0)} = a^2$ and $c_q(s, 3) = 0$.

$k = 2$

(0) is the only vector in E_3 such that $N(s) = 2$.

The paths from \emptyset to (0) of length 3 are $\emptyset \rightarrow \emptyset \rightarrow \emptyset \rightarrow (0)$ with weight 1; $\emptyset \rightarrow \emptyset \rightarrow (0) \rightarrow (0)$ with weight q ; $\emptyset \rightarrow (0) \rightarrow (0) \rightarrow (0)$ with weight q^2 . $e_a^{(0)} = a$ and $c(s, 3) = (1 + q + q^2) = 0$.

So we have proven that the 3-curvature is given by

$$D^3 = d_M^2(a) + d_M(a)a + (1 + q)ad_M(a).$$

Notice that in case of q -commutativity $\alpha\beta = q^{\bar{\alpha}\bar{\beta}}\beta\alpha$, the 3-curvature reduces to $D^3 = d_M^2(a)$.

Suppose now that q is a 4-th primitive root of unity and we required that $D^4 = 0$.

By Theorem 3 we must have $\sum_{k=0}^3 c_k d_M^k = 0$. Notice that

$$E_4 = \{\emptyset, (0), (1), (2), (3), (0, 0), (1, 0), (0, 1), (2, 0), (0, 2), (1, 1), \\ (0, 0, 0), (1, 0, 0), (0, 1, 0), (0, 0, 1), (0, 0, 0, 0)\}$$

We proceed to compute the coefficients c_k for $0 \leq k \leq 3$ using Theorem 3.

$$\underline{k=3}$$

$$\emptyset \xrightarrow{i} \emptyset \rightarrow (0) \xrightarrow{j} (0) \quad \text{the weight is } \sum_{i+j=3} q^j \text{ and } c_3 = (1 + q + q^2 + q^3)a = 0.$$

$$\underline{k=2}$$

$$\emptyset \xrightarrow{i} \emptyset \rightarrow (0) \xrightarrow{j} (0) \rightarrow (0,0) \xrightarrow{k} (0,0) \quad \text{the weight is } \sum_{i+j+k=2} q^j q^{2k} \text{ and}$$

$$c_q((0,0),4)a^{(0,0)} = (1 + q^2)a^2.$$

$$\emptyset \xrightarrow{i} \emptyset \rightarrow (0) \xrightarrow{j} (0) \rightarrow (1) \xrightarrow{k} (1) \quad \text{the weight is } \sum_{i+j+k=2} q^j q^{2k} \text{ and}$$

$$c_q((1),4)a^{(1)} = (1 + q^2)d(a).$$

$$\text{Finally, } c_2 = (1 + q^2)(a^2 + d(a)).$$

$$\underline{k=1}$$

$$\emptyset \xrightarrow{i} \emptyset \rightarrow (0) \xrightarrow{j} (0) \rightarrow (0,0) \xrightarrow{k} (0,0) \rightarrow (0,0,0) \xrightarrow{l} (0,0,0)$$

$$\text{the weight is } \sum_{i+j+k+l=1} q^j q^{2k} q^{3l} \text{ and } c_q((0,0,0),4)a^{(0,0,0)} = (1 + q + q^2 + q^3)a^3 = 0.$$

$$\emptyset \xrightarrow{i} \emptyset \rightarrow (0) \xrightarrow{j} (0) \rightarrow (1) \xrightarrow{k} (1) \rightarrow (0,1) \xrightarrow{l} (0,1)$$

$$\text{the weight is } \sum_{i+j+k+l=1} q^j q^{2k} q^{3l}.$$

$$\emptyset \xrightarrow{i} \emptyset \rightarrow (0) \xrightarrow{j} (0) \rightarrow (0,0) \xrightarrow{k} (0,0) \rightarrow (0,1) \xrightarrow{l} (0,1)$$

$$\text{the weight is } \sum_{i+j+k+l=1} q^j q^{2k} q^{3l} \text{ and } c_q((0,1),4)a^{(0,1)} = ((1 + q + q^2 + q^3) + (1 + q + q^2 + q^3))ad(a) = 0.$$

$$\emptyset \xrightarrow{i} \emptyset \rightarrow (0) \xrightarrow{j} (0) \rightarrow (0,0) \xrightarrow{k} (0,0) \rightarrow (1,0) \xrightarrow{l} (1,0)$$

$$\text{the weight is } \sum_{i+j+k+l=1} q^j q^{2k} q^{3l} \text{ and } c_q((1,0),4)a^{(1,0)} = (1 + q + q^2 + q^3)d(a)a = 0.$$

$$\emptyset \xrightarrow{i} \emptyset \rightarrow (0) \xrightarrow{j} (0) \rightarrow (1) \xrightarrow{k} (1) \rightarrow (2) \xrightarrow{l} (2)$$

$$\text{the weight is } \sum_{i+j+k+l=1} q^j q^{2k} q^{3l} \text{ and } c_q((2),4)a^{(2)} = (1 + q + q^2 + q^3)d^2(a) = 0.$$

$$\text{Finally, } c_1 = c_q((0,0,0),4)a^{(0,0,0)} + c_q((0,1),4)a^{(0,1)} + c_q((1,0),4)a^{(1,0)} + c_q((2),4)a^{(2)} = 0.$$

$$\underline{k=0}$$

$$\emptyset \rightarrow (0) \rightarrow (0,0) \rightarrow (0,0,0) \rightarrow (0,0,0,0)$$

$$\text{the weight is } 1 \text{ and } c_q((0,0,0,0),4)a^{(0,0,0,0)} = a^4.$$

$$\begin{aligned} \emptyset &\longrightarrow (0) \longrightarrow (0, 0) \longrightarrow (0, 0, 0) \longrightarrow (0, 0, 1) \\ \emptyset &\longrightarrow (0) \longrightarrow (1) \longrightarrow (0, 1) \longrightarrow (0, 0, 1) \\ \emptyset &\longrightarrow (0) \longrightarrow (0, 0) \longrightarrow (0, 1) \longrightarrow (0, 0, 1) \end{aligned}$$

the weight is $1 + q + q^2$ and $c_q((0, 0, 1), 4)a^{(0,0,1)} = (1 + q + q^2)a^2d(a)$.

$$\begin{aligned} \emptyset &\longrightarrow (0) \longrightarrow (0, 0) \longrightarrow (1, 0) \longrightarrow (0, 1, 0) \\ \emptyset &\longrightarrow (0) \longrightarrow (0, 0) \longrightarrow (0, 0, 0) \longrightarrow (0, 1, 0) \end{aligned}$$

the weight is $1 + q$ and $c_q((0, 1, 0), 4)a^{(0,1,0)} = (1 + q)ad(a)a$.

$$\emptyset \longrightarrow (0) \longrightarrow (0, 0) \longrightarrow (0, 0, 0) \longrightarrow (1, 0, 0)$$

the weight is 1 and $c_q((1, 0, 0), 4)a^{(1,0,0)} = d(a)a^2$.

$$\emptyset \longrightarrow (0) \longrightarrow (0, 0) \longrightarrow (1, 0) \longrightarrow (2, 0)$$

the weight is 1 and $c_q((2, 0), 4)a^{(2,0)} = d^2(a)a$.

$$\begin{aligned} \emptyset &\longrightarrow (0) \longrightarrow (0, 0) \longrightarrow (0, 1) \longrightarrow (0, 2) \\ \emptyset &\longrightarrow (0) \longrightarrow (1) \longrightarrow (0, 1) \longrightarrow (0, 2) \\ \emptyset &\longrightarrow (0) \longrightarrow (1) \longrightarrow (2) \longrightarrow (0, 2) \end{aligned}$$

the weight is $1 + q + q^2$ and $c_q((0, 2), 4)a^{(0,2)} = (1 + q + q^2)ad^2(a)$.

$$\begin{aligned} \emptyset &\longrightarrow (0) \longrightarrow (0, 0) \longrightarrow (1, 0) \longrightarrow (1, 1) \\ \emptyset &\longrightarrow (0) \longrightarrow (1) \longrightarrow (0, 1) \longrightarrow (1, 1) \end{aligned}$$

the weight is $1 + q^2$ and $c_q((1, 1), 4)a^{(1,1)} = (d(a))^2$.

$$\emptyset \longrightarrow (0) \longrightarrow (1) \longrightarrow (2) \longrightarrow (3)$$

the weight is 1 and $c_q((3), 4)a^{(3)} = d^3(a)$.

Finally, $c_0 = a^4 + (1 + q + q^2)a^2d(a) + (1 + q)ad(a)a + d(a)a^2 + d^2(a)a + (1 + q + q^2)ad^2(a) + (d(a))^2 + d^3(a)$.

So we have that the 4-curvature is given by

$$\begin{aligned} D^4 &= (1 + q^2)(a^2 + d(a))d_M^2 + a^4 + (1 + q + q^2)a^2d(a) + (1 + q)ad(a)a + \\ &\quad d(a)a^2 + d^2(a)a + (1 + q + q^2)ad^2(a) + (d(a))^2 + d^3(a). \end{aligned}$$

In the remainder of the paper we consider infinitesimal deformations of a q -differential.

Theorem 4 Let (M^\bullet, m_M, d_M) be a q -dgm such that $d_M^N = 0$, consider the infinitesimal deformation $D = d_M + te$, where $e \in \text{End}^1(M^\bullet)$ and $t^2 = 0$, then

$$D^N = \sum_{k=0}^{N-1} \left(\sum_{v \in \text{Par}(N, N-k-1)} q^{|v|+w(v)} \right) d_M^{N-k-1},$$

where

$$\text{Par}(N, N-k-1) = \{(v_0, \dots, v_{N-k-1}) / \sum v_i = N\},$$

$$|v| = v_1 + \dots + v_n \text{ and } w(k) = \sum iv_i.$$

Proof: By Theorem 3, $D^N = \sum_{k=0}^{N-1} c_k d_M^k$. Since $t^2 = 0$, then

$$(te)^{(s)} = (te)^{(s_1)} \dots (te)^{(s_l(s))} = t^{l(s)} e^{(s)} = 0 \text{ unless } l(s) \leq 1.$$

Thus E_N is given by

$$E_N = \{(0), (1), \dots, (N-1)\}.$$

For $0 \leq k \leq N-1$, since $N(s) = N - |s| - l(s) = k$ and $l(s) = 1$ we obtain $|s| = N-k-1$, and the only vector s in E_N of length 1 such that $|s| = N-k-1$ is precisely $s = (N-k-1)$. Thus

$$c_k = \sum_{\substack{s \in E_N \\ N(s)=k \\ s_i < N}} c_q(s, N) a^{(s)} = c_q((N-k-1), N) a^{(s)} = c_q((N-k-1), N) d^{N-k-1}(a).$$

Any path from \emptyset to $(N-k-1)$ of length N must be of the form

$$\underbrace{\emptyset \rightarrow \dots \rightarrow \emptyset}_{p_0} \rightarrow \underbrace{(0) \rightarrow \dots \rightarrow (0)}_{p_1} \rightarrow \underbrace{(1) \rightarrow \dots \rightarrow (1)}_{p_2} \rightarrow \dots \rightarrow \underbrace{(N-k-1) \rightarrow \dots \rightarrow (N-k-1)}_{p_{N-k-1}}$$

with $p_0 + p_1 + \dots + p_{N-k-1} = N$. The weight of such path is $q^{p_1} q^{p_2(1+1)} \dots q^{p_{N-k-1}(N-k)} = q^{|p|+w(p)}$, where $p = (p_0, \dots, p_{N-k-1})$. Then

$$c_q(s, N) = \sum_{\gamma \in P_N(\emptyset, s)} v(\gamma) = \sum_{p \in \text{Par}(N, N-k-1)} q^{|p|+w(p)}. \blacklozenge$$

Acknowledgement

We thank Sylvie Paycha and Nicolás Andruskiewitsch.

References

- [1] V. Abramov, R. Kerner, *On certain realizations of the q -deformed exterior differential calculus*, Reports on Math. Phys., 43 (1999) 179-194
- [2] M. Angel, R. Díaz, *N -differential graded algebras*, Preprint math.DG/0504398.
- [3] M. Angel, R. Díaz, *N -flat connections*, Preprint math.DG/0511242.
- [4] M. Angel, R. Díaz, *A_∞^N -algebras*, In preparation.
- [5] M. Dubois-Violette, *Lectures on differentials, generalized differentials and some examples related to theoretical physics*.-Quantum symmetries in theoretical physics and mathematics (Bariloche 2000), pages 59-94, Contemp. Math. 294, A.M.S. 2002.
- [6] M.M. Kapranov, *On the q -analog of homological algebra*.-Preprint q-alg/9611005.
- [7] C. Kassel et M. Wambst, *Algèbre homologique des N -complexes et homologie de Hochschild aux racines de l'unité*.-Publ. Res. Inst. Math. Sci. Kyoto University 34 (1998), n 2, 91-114.
- [8] R. Kerner, B. Niemeier, *Covariant q -differential calculus and its deformations at $q^N = 1$* , Lett. in Math. Phys., 45,161-176, (1998).
- [9] A. Sitarz; *On the tensor product construction for q -differential algebras*.-Lett. Math. Phys. 44 1998.

Mauricio Angel. Universidad Central de Venezuela (UCV). mangel@euler.ciens.ucv.ve
Rafael Díaz. Universidad Central de Venezuela (UCV). rdiaz@euler.ciens.ucv.ve