HIGHER LAME EQUATIONS AND CRITICAL POINTS OF MASTER FUNCTIONS

E. MUKHIN *, V. TARASOV *,*,1, AND A. VARCHENKO **,2

*Department of Mathematical Sciences, Indiana University – Purdue University, Indianapolis, 402 North Blackford St, Indianapolis, IN 46202-3216, USA

> *St. Petersburg Branch of Steklov Mathematical Institute Fontanka 27, St. Petersburg, 191023, Russia

** Department of Mathematics, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599-3250, USA

ABSTRACT. Under certain conditions, we give an estimate from above on the number of differential equations of order r+1 with prescribed regular singular points, prescribed exponents at singular points, and having a quasi-polynomial flag of solutions. The estimate is given in terms of a suitable weight subspace of the tensor power $U(\mathfrak{n}_{-})^{\otimes (n-1)}$, where n is the number of singular points in \mathbb{C} and $U(\mathfrak{n}_{-})$ is the enveloping algebra of the nilpotent subalgebra of \mathfrak{gl}_{r+1} .

Dedicated to Askold Khovanskii on the occasion of his 60th birthday.

1. INTRODUCTION

Consider the differential equation

(1)
$$F(x) u''(x) + G(x) u'(x) + H(x) u(x) = 0,$$

where F(x) is a polynomial of degree n, and G(x), H(x) are polynomials of degree not greater than n-1, n-2, respectively. If F(x) has no multiple roots, then all singular points of the equation are regular singular. Write

(2)
$$F(x) = \prod_{s=1}^{n} (x - z_s), \qquad \frac{G(x)}{F(x)} = -\sum_{s=1}^{n} \frac{m_s}{x - z_s}$$

for suitable complex numbers m_s , z_s . Then 0 and m_s+1 are exponents at z_s of equation (1). If -l is one of the exponents at ∞ , then the other is $l - 1 - \sum_{s=1}^{n} m_s$.

Problem ([Sz], Ch. 6.8) Given polynomials F(x), G(x) as above and a non-negative integer l,

¹ Supported in part by RFFI grant 05-01-00922.

² Supported in part by NSF grant DMS-0244579.

- (a) find a polynomial H(x) of degree at most n-2 such that equation (1) has a polynomial solution of degree l;
- (b) find the number of solutions to Problem (a).

If H(x), u(x) is a solution to Problem (a), then the corresponding equation (1) is called a Lame equation and the polynomial u(x) is called a Lame function.

Example. Let $F(x) = 1 - x^2$, $G(x) = \alpha - \beta + (\alpha + \beta + 2)x$. Then $H = l(l + \alpha + \beta + 1)$ and the corresponding polynomial solution of degree l, normalized by the condition $u(1) = \binom{l+\alpha}{l}$, is called *the Jacobi polynomial* and denoted by $P_l^{(\alpha,\beta)}(x)$.

The following result is classical.

Theorem 1.1 (Cf. [Sz], Ch. 6.8, [St]).

• Let u(x) be a polynomial solution of (1) of degree l with roots t_1^0, \ldots, t_l^0 of multiplicity one. Then $\mathbf{t}^0 = (t_1^0, \ldots, t_l^0)$ is a critical point of the function

$$\Phi_{l,n}(\boldsymbol{t};\boldsymbol{z};\boldsymbol{m}) = \prod_{i=1}^{l} \prod_{s=1}^{n} (t_i - z_s)^{-m_s} \prod_{1 \le i < j \le l} (t_i - t_j)^2 ,$$

where $z = (z_1, ..., z_n)$ and $m = (m_1, ..., m_n)$.

• Let t^0 be a critical point of the function $\Phi_{l,n}(\cdot; \boldsymbol{z}; \boldsymbol{m})$, then the polynomial u(x) of degree l with roots t^0_1, \ldots, t^0_l is a solution of (1) with H(x) = (-F(x)u''(x) - G(x)u'(x))/u(x) being a polynomial of degree at most n-2.

The function $\Phi_{l,n}(\cdot; \boldsymbol{z}; \boldsymbol{m})$ is called the master function. The master function is a symmetric function of the variables t_1, \ldots, t_l . Therefore, the symmetric group S_l naturally acts on the set of critical points of the master function by permuting the coordinates.

By Theorem 1.1, the S_l -orbits of critical points are in one-to-one correspondence with solution H(x), u(x) of Problem (a) such that u(x) has no multiple roots.

The following result is also classical.

Theorem 1.2 (Cf. [Sz], Ch. 6.8, [H], [St]). If z_1, \ldots, z_n are distinct real numbers and m_1, \ldots, m_n are negative numbers, then the number of solutions to Problem (a) is equal to $\binom{l+n-2}{l}$.

Under these conditions on \boldsymbol{z} and \boldsymbol{m} , the master function has exactly $\binom{l+n-2}{l} S_l$ -orbits of critical points, see [Sz] and [V3].

The number $\binom{l+n-2}{l}$ has the following representation theoretical interpretation. The universal enveloping algebra $U(\mathfrak{n}_{-})$ of the nilpotent subalgebra $\mathfrak{n}_{-} \subset \mathfrak{gl}_{2}$ is generated by one element e_{21} and is weighted by powers of the generator. The number $\binom{l+n-2}{l}$ is the dimension of the weight l part of the tensor power $U(\mathfrak{n}_{-})^{\otimes (n-1)}$.

The case of nonnegative integers m_1, \ldots, m_n is interesting for applications to the Bethe ansatz method in the Gaudin model. For a nonnegative integer m, let L_m be the m + 1-dimensional irreducible \mathfrak{sl}_2 -module. If m_1, \ldots, m_n are nonnegative integers and $m_{\infty} = \sum_{s=1}^{n} m_s - 2l$ is a nonnegative integer, then for any distinct z_1, \ldots, z_n the number of S_l -orbits of critical points of the master function function $\Phi_{l,n}(\cdot; \boldsymbol{z}; \boldsymbol{m})$ is not greater than the multiplicity of the \mathfrak{sl}_2 -module $L_{m_{\infty}}$ in the tensor product $\bigotimes_{s=1}^{n} L_{m_s}$. Moreover, for generic z_1, \ldots, z_n , the number of S_l -orbits is equal to that multiplicity, see [ScV].

The goal of this paper is to generalize the formulated results. We consider linear differential equations of order r + 1 with regular singular points only, located at z_1, \ldots, z_n, ∞ , and having prescribed exponents at each of the singular points. We introduce the notion of a quasi-polynomial flag of solutions for such a differential equation. (For a second order differential equation, a quasi-polynomial flag is just a solution of the form $y(x) \prod_{s=1}^{n} (x - z_s)^{\lambda_s}$ where y(x) is a polynomial and $\lambda_1, \ldots, \lambda_n$ are some complex numbers.) We prove two facts:

- Differential equations with a quasi-polynomial flag, prescribed singular points and exponents are in one-to-one correspondence with suitable orbits of critical points of some \mathfrak{gl}_{r+1} -master function.
- Under explicit generic conditions on exponents (they must be *separated*), we show that the number of orbits of critical points of the corresponding master function is not greater than the dimension of a suitable weight subspace of $U(\mathfrak{n}_{-})^{\otimes (n-1)}$, where \mathfrak{n}_{-} is the nilpotent subalgebra of \mathfrak{gl}_{r+1} .

The proofs are based on results from [MV2, BMV].

This paper was motivated by discussions with B. Shapiro of his preprint [BBS] in which another generalization of Problems (a) and (b) is introduced for differential equations of order r + 1. The authors thank B. Shapiro for useful discussions of his preprint.

2. CRITICAL POINTS OF \mathfrak{gl}_{r+1} -MASTER FUNCTIONS

2.1. Lie algebra \mathfrak{gl}_{r+1} . Consider the Lie algebra \mathfrak{gl}_{r+1} with standard generators $e_{a,b}$, $a, b = 1, \ldots, r+1$, and Cartan decomposition $\mathfrak{gl}_{r+1} = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$,

$$\mathfrak{n}_{-} = \bigoplus_{a > b} \mathbb{C} \cdot e_{a,b} , \qquad \mathfrak{h} = \bigoplus_{a=1}^{r+1} \mathbb{C} \cdot e_{a,a} , \qquad \mathfrak{n}_{+} = \bigoplus_{a < b} \mathbb{C} \cdot e_{a,b} .$$

Set $h_a = e_{a,a} - e_{a+1,a+1}$ for $a = 1, \ldots, r$. Let $e_{1,1}^*, \ldots, e_{r+1,r+1}^* \in \mathfrak{h}^*$ be the basis dual to the basis $e_{1,1}, \ldots, e_{r+1,r+1} \in \mathfrak{h}$. Set $\alpha_a = e_{a,a}^* - e_{a+1,a+1}^*$ for $a = 1, \ldots, r$. Fix the scalar product on \mathfrak{h}^* such that $(e_{a,a}^*, e_{b,b}^*) = \delta_{a,b}$.

Let $U(\mathfrak{n}_{-})$ be the universal enveloping algebra of \mathfrak{n}_{-} ,

$$U(\mathbf{n}_{-}) = \bigoplus_{\boldsymbol{l} \in \mathbb{Z}_{>0}^{r}} U(\mathbf{n}_{-})[\boldsymbol{l}] ,$$

where for $\boldsymbol{l} = (l_1, \ldots, l_r)$, the space $U(\boldsymbol{n}_{-})[\boldsymbol{l}]$ consists of elements f such that

$$[f,h] = \langle h, \sum_{i=1}^r l_i \alpha_i \rangle f$$
.

The element $\prod_i e_{a_i,b_i}$ with $a_i > b_i$ belongs to the graded subspace $U(\mathfrak{n}_{-})[l]$, where $l = \sum_i l_i$ with

 $\boldsymbol{l}_{i} = (0, 0, \dots, 0, 1_{b}, 1_{b+1}, \dots, 1_{a-1}, 0, 0, \dots, 0) \; .$

Choose an order on the set of elements $e_{a,b}$ with $r+1 \ge a > b \ge 1$. Then the ordered products $\prod_{a>b} e_{a,b}^{n_{a,b}}$ form a graded basis of $U(\mathfrak{n}_{-})$.

The grading of $U(\mathfrak{n}_{-})$ induces the grading of $U(\mathfrak{n}_{-})^{\otimes k}$ for any positive integer k,

$$U(\mathfrak{n}_{-})^{\otimes k} = \bigoplus_{\boldsymbol{l}} U(\mathfrak{n}_{-})^{\otimes k}[\boldsymbol{l}]$$
.

Denote

$$d(k, \boldsymbol{l}) = \dim U(\boldsymbol{\mathfrak{n}}_{-})^{\otimes k}[\boldsymbol{l}]$$
 .

For a weight $\Lambda \in \mathfrak{h}^*$, denote by L_{Λ} the irreducible \mathfrak{gl}_{r+1} -module with highest weight Λ . Let

$$\mathbf{\Lambda} = (\Lambda_1, \dots, \Lambda_n) , \qquad \Lambda_s \in \mathfrak{h}^* ,$$

be a collection of weights and $\boldsymbol{l} = (l_1, \ldots, l_r)$ a collection of nonnegative integers. Let

$$L_{\mathbf{\Lambda}} = L_{\Lambda_1} \otimes \cdots \otimes L_{\Lambda_n}$$

be the tensor product of irreducible \mathfrak{gl}_{r+1} -modules. Let $L_{\mathbf{\Lambda}} = \bigoplus_{l \in \mathbb{Z}_{\geq 0}^{r}} L_{\mathbf{\Lambda}}[l]$ be its weight decomposition, where $L_{\mathbf{\Lambda}}[(l_{1},\ldots,l_{r})]$ is the subspace of vectors of weight $\sum_{s=1}^{n} \Lambda_{s} - \sum_{i=1}^{r} l_{i}\alpha_{i}$. Let Sing $L_{\mathbf{\Lambda}}[l] \subset L_{\mathbf{\Lambda}}[l]$ be the subspace of singular vectors, i.e. the subspace of vectors annihilated by \mathbf{n}_{+} . Denote

$$\delta(\mathbf{\Lambda}, \mathbf{l}) = \dim \operatorname{Sing} L_{\mathbf{\Lambda}}[\mathbf{l}]$$

It is well-known that for given l and a generic set of weights Λ , we have

$$d(n-1, \mathbf{l}) = \delta(\mathbf{\Lambda}, \mathbf{l})$$
.

For given Λ and l, introduce the weight

$$\Lambda_{\infty} = \sum_{s=1}^{n} \Lambda_{s} - \sum_{i=1}^{r} l_{i} \alpha_{i} \in \mathfrak{h}^{*}$$

and the sequences of numbers $\boldsymbol{m}_1, \ldots, \boldsymbol{m}_{r+1}, \boldsymbol{m}_{\infty}$, where

$$\boldsymbol{m}_s = \{ m_{s,1}, \ldots, m_{s,r+1} \}$$
 and $m_{s,i} = \langle \Lambda_s, e_{i,i} \rangle$.

Having sequences $m_1, \ldots, m_{r+1}, m_{\infty}$, we can recover Λ and l as follows:

(3)
$$\Lambda_s = \sum_{i=1}^{r+1} m_{s,i} e_{i,i}^*$$
 and $\sum_{i=1}^r l_i \alpha_i = \sum_{s=1}^n \Lambda_s - \Lambda_\infty$.

We say that the set of sequences $m_1, \ldots, m_{r+1}, m_{\infty}$ of complex numbers is *admissible*, if all of the numbers $l = (l_1, \ldots, l_r)$, defined by (3), are nonnegative integers.

2.2. Master functions. Let $\boldsymbol{z} = (z_1, \ldots, z_n) \in \mathbb{C}^n$ be a point with distinct coordinates. Let

$$oldsymbol{\Lambda} \;=\; (\Lambda_1,\ldots,\Lambda_n)\;, \qquad \Lambda_s \in \mathfrak{h}^*\;,$$

be a collection of \mathfrak{gl}_{r+1} -weights and $l = (l_1, \ldots, l_r)$ a collection of nonnegative integers. Set $l = l_1 + \cdots + l_r$. Introduce a function of l variables

$$\boldsymbol{t} = (t_1^{(1)}, \dots, t_{l_1}^{(1)}, \dots, t_1^{(r)}, \dots, t_{l_r}^{(r)})$$

by the formula

(4)
$$\Phi(\boldsymbol{t}; \boldsymbol{z}; \boldsymbol{\Lambda}; \boldsymbol{l}) = \prod_{i=1}^{r} \prod_{j=1}^{l_{i}} \prod_{s=1}^{n} (t_{j}^{(i)} - z_{s})^{-(\boldsymbol{\Lambda}_{s}, \alpha_{i})} \prod_{i=1}^{r} \prod_{1 \le j < s \le l_{i}} (t_{j}^{(i)} - t_{s}^{(i)})^{2} \prod_{i=1}^{r-1} \prod_{j=1}^{l_{i}} \prod_{k=1}^{l_{i+1}} (t_{j}^{(i)} - t_{k}^{(j+1)})^{-1} .$$

The function Φ is a (multi-valued) function of t, depending on parameters z, Λ . The function is called *the master function*.

The master functions arise in the hypergeometric solutions of the KZ equations [?, V1] and in the Bethe ansatz method for the Gaudin model [RV, ScV, MV1, MV2, MV3, V2, MTV].

The product of symmetric groups

$$S_l = S_{l_1} \times \cdots \times S_{l_r}$$

acts on variables t by permuting the coordinates with the same upper index. The master function is S_l -invariant.

A point \boldsymbol{t} with complex coordinates is called a *critical point* of $\Phi(\cdot; \boldsymbol{z}; \boldsymbol{\Lambda}; \boldsymbol{l})$ if the following system of l equations is satisfied

(5)
$$\sum_{s=1}^{n} \frac{(\Lambda_s, \alpha_1)}{t_j^{(1)} - z_s} - \sum_{s=1, s \neq j}^{l_1} \frac{2}{t_j^{(1)} - t_s^{(1)}} + \sum_{s=1}^{l_2} \frac{1}{t_j^{(1)} - t_s^{(2)}} = 0 ,$$
$$\sum_{s=1}^{n} \frac{(\Lambda_s, \alpha_i)}{t_j^{(i)} - z_s} - \sum_{s=1, s \neq j}^{l_i} \frac{2}{t_j^{(i)} - t_s^{(i)}} + \sum_{s=1}^{l_{i-1}} \frac{1}{t_j^{(i)} - t_s^{(i-1)}} + \sum_{s=1}^{l_{i+1}} \frac{1}{t_j^{(i)} - t_s^{(i+1)}} = 0 ,$$
$$\sum_{s=1}^{n} \frac{(\Lambda_s, \alpha_r)}{t_j^{(r)} - z_s} - \sum_{s=1, s \neq j}^{l_r} \frac{2}{t_j^{(r)} - t_s^{(r)}} + \sum_{s=1}^{l_{i-1}} \frac{1}{t_j^{(r)} - t_s^{(r-1)}} = 0 ,$$

where $j = 1, ..., l_1$ in the first group of equations, i = 2, ..., r - 1 and $j = 1, ..., l_i$ in the second group of equations, $j = 1, ..., l_r$ in the last group of equations.

In other words, a point t is a critical point if

$$\left(\Phi^{-1}\frac{\partial\Phi}{\partial t_j^{(i)}}\right)(\boldsymbol{t};\boldsymbol{z};\boldsymbol{\Lambda};\boldsymbol{l}) = 0, \qquad i = 1,\ldots,r, \ j = 1,\ldots,l_i.$$

In the Gaudin model, equations (5) are called the Bethe ansatz equations.

The set of critical points is S_l -invariant.

2.3. The case of isolated critical points. We say that the pair Λ , l is separating if

$$(2\Lambda_{\infty} + \sum_{i=1}^{r} c_i \alpha_i, \sum_{i=1}^{r} c_i \alpha_i) + 2\sum_{i=1}^{r} c_i \neq 0$$

for all sets of integers $\{c_1, \ldots, c_r\}$ such that $0 \le c_i \le l_i$, $\sum_i c_i \ne 0$.

For example, if Λ_{∞} is dominant integral, then Λ, l is separating.

Lemma 2.1 (Theorem 16 [ScV], Lemma 2.1 [MV2]). If the pair Λ , \boldsymbol{l} is separating, then the set of critical points of the master function $\Phi(\cdot; \boldsymbol{z}; \Lambda; \boldsymbol{l})$ is finite.

By Lemma 2.1, for given \boldsymbol{l} and generic $\boldsymbol{\Lambda}$, the master function $\Phi(\cdot; \boldsymbol{z}; \boldsymbol{\Lambda}; \boldsymbol{l})$ has finitely many critical points.

Theorem 2.2. Assume that the pair Λ , \boldsymbol{l} is separating, then the number of $S_{\boldsymbol{l}}$ -orbits of critical points counted with multiplicities is not greater than $d(n-1, \boldsymbol{l})$.

The multiplicity of a critical point t is the multiplicity of t as a solution of system (5).

Proof. It is shown in [BMV] that, if $\Lambda_1, \ldots, \Lambda_n, \Lambda_\infty$ is a collection of dominant integral weights, then the number of $S_{\boldsymbol{l}}$ -orbits of critical points counted with multiplicities is not greater than $\delta(\boldsymbol{\Lambda}, \boldsymbol{l})$. Together with equality $d(n-1, \boldsymbol{l}) = d(\boldsymbol{\Lambda}, \boldsymbol{l})$ for generic $\boldsymbol{\Lambda}$, this proves the theorem.

3. DIFFERENTIAL OPERATORS WITH QUASI-POLYNOMIAL FLAGS OF SOLUTIONS

3.1. Fundamental differential operator. For the S_l -orbit of a critical point t of the master function $\Phi(\cdot; \boldsymbol{z}; \boldsymbol{\Lambda}; \boldsymbol{l})$, define the tuple $\boldsymbol{y}^t = (y_1, \ldots, y_r)$ of polynomials in variable x,

$$y_i(x) = \prod_{j=1}^{l_i} (x - t_j^{(i)}), \qquad i = 1, \dots, r.$$

Since t is a critical point, all fractions in (5) are well-defined. Therefore, the tuple y^t has the following properties:

- (6) Every polynomial y_i has no multiple roots.
- (7) Every pair of polynomials y_i and y_{i+1} has no common roots.
- (8) For every i = 1, ..., r and s = 1, ..., n, if $m_{s,i} m_{s,i+1} \neq 0$, then $y_i(z_s) \neq 0$.

A tuple of monic polynomials (y_1, \ldots, y_r) with properties (6)-(8) will be called *off-diagonal*.

Define quasi-polynomials T_1, \ldots, T_{r+1} in x by the formula

(9)
$$T_i(x) = \prod_{s=1}^n (x - z_s)^{-m_{s,i}}$$

Consider the linear differential operator of order r + 1,

$$D_{t} = \left(\frac{d}{dx} - \ln'(\frac{T_{r+1}}{y_{r}})\right) \left(\frac{d}{dx} - \ln'(\frac{y_{r}T_{r}}{y_{r-1}})\right) \dots \left(\frac{d}{dx} - \ln'(\frac{y_{2}T_{2}}{y_{1}})\right) \left(\frac{d}{dx} - \ln'(y_{1}T_{1})\right)$$

where $\ln'(f)$ denotes (df/dx)/f for any f. We say that D_t is the fundamental operator of the critical point t.

Theorem 3.1. Assume that the pair Λ , l is separating. Then

(i) All singular points of D_t are regular and lie in z₁,..., z_n,∞. The exponents of D_t at z_s are

 $-m_{s,1}, -m_{s,2}+1, \ldots, -m_{s,r+1}+r$

for s = 1, ..., n, and the exponents of D_t at ∞ are

$$m_{\infty,1}, m_{\infty,2}-1, \ldots, m_{\infty,r+1}-r$$
.

(ii) The differential equation $D_t u = 0$ has solutions u_1, \ldots, u_{r+1} such that

$$u_1 = y_1 T_1$$

and for $i = 2, \ldots, r+1$ we have

$$\operatorname{Wr}(u_1,\ldots,u_i) = y_i T_i T_{i-1} \ldots T_1$$

where Wr (u_1, \ldots, u_i) denotes the Wronskian of u_1, \ldots, u_i and $y_{r+1} = 1$.

Proof. Part (ii) follows from the presentation of D_t as a product. To prove part (i) consider the operator $\tilde{D}_t = T_1^{-1} \cdot D_t \cdot T_1$, the conjugate of D_t by the operator of multiplication by T_1 . Then

$$\tilde{D}_{t} = \left(\frac{d}{dx} - \ln'(\frac{T_{r+1}}{y_{r}T_{1}})\right) \left(\frac{d}{dx} - \ln'(\frac{y_{r}T_{r}}{y_{r-1}T_{1}})\right) \dots \left(\frac{d}{dx} - \ln'(\frac{y_{2}T_{2}}{y_{1}T_{1}})\right) \left(\frac{d}{dx} - \ln'(y_{1})\right) \,.$$

It is enough to show that

- (iii) All singular points of \tilde{D}_t are regular and lie in $\{z_1, \ldots, z_n, \infty\}$.
- (iv) The exponents of \tilde{D}_t at z_s are

$$0, m_{s,1} - m_{s,2} + 1, \ldots, m_{s,1} - m_{s,r+1} + r$$

for $s = 1, \ldots, n$, and the exponents of \tilde{D}_t at ∞ are

 $-l_1$, $-m_{\infty,1} + m_{\infty,2} - 1 - l_1$, ..., $-m_{\infty,1} + m_{\infty,r+1} - r - l_1$.

If all highest weights in the collection $\Lambda_1, \ldots, \Lambda_n, \Lambda_\infty$ are integral dominant, then statements (iii-iv) are proved in [MV2], Section 5. Hence statements (iii-iv) hold for arbitrary separating $\Lambda_1, \ldots, \Lambda_n$, \boldsymbol{l} .

3.2. Quasi-polynomial flags. Let $\mathbf{z} = (z_1, \ldots, z_n) \in \mathbb{C}^n$ be a point with distinct coordinates. Let $\mathbf{m}_1, \ldots, \mathbf{m}_{r+1}, \mathbf{m}_{\infty}$ be an admissible set of sequences of complex numbers as in Section 2.1. Let

$$D = \frac{d^{r+1}}{dx^{r+1}} + A_1(x)\frac{d^r}{dx^r} + \dots + A_r(x)\frac{d}{dx} + A_{r+1}(x)$$

be a differential operator with rational coefficients.

We say that D is associated with $\boldsymbol{z}, \boldsymbol{m}_1, \ldots, \boldsymbol{m}_{r+1}, \boldsymbol{m}_{\infty}$, if

- All singular points of D are regular and lie in z_1, \ldots, z_n, ∞ .
- The exponents of D at z_s are $-m_{s,1}$, $-m_{s,2}+1$, ..., $-m_{s,r+1}+r$ for $s=1,\ldots,n$.
- The exponents of D at ∞ are $m_{\infty,1}$, $m_{\infty,2} 1$, ..., $m_{\infty,r+1} r$.

Define quasi-polynomials T_1, \ldots, T_{r+1} in x by formula (9).

We say that an operator D, associated with $\boldsymbol{z}, \boldsymbol{m}_1, \ldots, \boldsymbol{m}_{r+1}, \boldsymbol{m}_{\infty}$, has a quasipolynomial flag, if there exists a tuple $\boldsymbol{y} = (y_1, \ldots, y_r)$ of monic polynomials in x, such that

(i) For i = 1, ..., r, deg $y_i = l_i$.

- (ii) The tuple \boldsymbol{y} is off-diagonal in the sense of (6)-(8).
- (iii) The differential equation D u = 0 has solutions u_1, \ldots, u_{r+1} such that

$$u_1 = y_1 T_1$$

and for $i = 2, \ldots, r+1$ we have

$$\operatorname{Wr}(u_1,\ldots,u_i) = y_i T_i T_{i-1} \ldots T_1,$$

where $y_{r+1} = 1$.

Proposition 3.2. An operator D, associated with z, $m_1, \ldots, m_{r+1}, m_{\infty}$, has a quasipolynomial flag if and only there exists a tuple $\boldsymbol{y} = (y_1, \ldots, y_r)$ of monic polynomials in x with properties (*i*-*ii*) and

(iv) The differential operator D can be presented in the form

$$D = \left(\frac{d}{dx} - \ln'(\frac{T_{r+1}}{y_r})\right) \left(\frac{d}{dx} - \ln'(\frac{y_r T_r}{y_{r-1}})\right) \dots \left(\frac{d}{dx} - \ln'(\frac{y_2 T_2}{y_1})\right) \left(\frac{d}{dx} - \ln'(y_1 T_1)\right) \,.$$

Proof. The equivalence of (iii) and (iv) follows from the next lemma.

Lemma 3.3. Let u_1, \ldots, u_{r+1} be any functions such that

$$Wr(u_1, \ldots, u_i) = y_i T_i T_{i-1} \ldots T_1,$$

for i = 1, ..., r + 1. Then $u_1, ..., u_{r+1}$ form a basis of the space of solutions of the equation Du = 0 with

$$D = \left(\frac{d}{dx} - \ln'(\frac{T_{r+1}}{y_r})\right) \left(\frac{d}{dx} - \ln'(\frac{y_r T_r}{y_{r-1}})\right) \dots \left(\frac{d}{dx} - \ln'(\frac{y_2 T_2}{y_1})\right) \left(\frac{d}{dx} - \ln'(y_1 T_1)\right)$$

Proof. It is enough to prove that $Du_i = 0$ for i = 1, ..., r + 1. By induction on i, we obtain that

$$\left(\frac{d}{dx} - \ln'(\frac{y_i T_i}{y_{i-1}})\right) \dots \left(\frac{d}{dx} - \ln'(\frac{y_2 T_2}{y_1})\right) \left(\frac{d}{dx} - \ln'(y_1 T_1)\right) u = \frac{W(u_1, \dots, u_i, u)}{y_i T_i T_{i-1} \dots T_1}$$

from which the statement follows.

Examples of differential operators with quasi-polynomial flags are given by Theorem 3.1. If \boldsymbol{t} is a critical point of the master function $\Phi(\cdot; \boldsymbol{z}; \boldsymbol{\Lambda}; \boldsymbol{l})$, then by Theorem 3.1, the fundamental differential operator $D_{\boldsymbol{t}}$ is associated with $\boldsymbol{z}, \boldsymbol{m}_1, \ldots, \boldsymbol{m}_{r+1}, \boldsymbol{m}_{\infty}$ and has a quasi-polynomial flag.

Having $\boldsymbol{z}, \boldsymbol{m}_1, \ldots, \boldsymbol{m}_{r+1}, \boldsymbol{m}_{\infty}$, define $\boldsymbol{\Lambda}, \boldsymbol{l}$ by (3) and the master function $\Phi(\cdot; \boldsymbol{z}; \boldsymbol{\Lambda}; \boldsymbol{l})$ by (4).

Having a tuple
$$\boldsymbol{y} = (y_1, \dots, y_r), \ y_i(x) = \prod_{j=1}^{l_i} (x - t_j^{(i)}),$$
 denote by

(10)
$$\boldsymbol{t} = (t_1^{(1)}, \dots, t_{l_1}^{(1)}, \dots, t_1^{(r)}, \dots, t_{l_r}^{(r)})$$

a point in \mathbb{C}^l , whose coordinates are roots of the polynomials of \boldsymbol{y} . The tuple \boldsymbol{y} uniquely determines the S_l -orbit of \boldsymbol{t} .

Theorem 3.4. Assume that an operator D is associated with z, $m_1, \ldots, m_{r+1}, m_{\infty}$ and has a quasi-polynomial flag. Let y be the corresponding tuple of polynomials. Then the point t, introduced in (10), is a critical point of the master function $\Phi(\cdot; z; \Lambda; l)$.

Proof. Let u_1, \ldots, u_{r+1} be the set of solutions of the differential equation D u = 0 giving the quasi-polynomial flag. Introduce the functions $\tilde{y}_1, \ldots, \tilde{y}_r$ by the formulas $u_2 = \tilde{y}_1 T_1$ and for $i = 2, \ldots, r$,

 $\tilde{y}_i T_i T_{i-1} \dots T_1 = \operatorname{Wr} (u_1, \dots, u_{i-1}, u_{i+1}) .$

The functions $\tilde{y}_1, \ldots, \tilde{y}_r$ are multi-valued functions.

• Singularities of all branches of all of these functions lie in z_1, \ldots, z_n .

It follows from the Wronskian identities of [MV2], that

• Wr
$$(\tilde{y}_1, y_1) = \frac{T_2}{T_1} y_2$$
, Wr $(\tilde{y}_r, y_r) = \frac{T_{r+1}}{T_r} y_{r-1}$, Wr $(\tilde{y}_i, y_i) = \frac{T_{i+1}}{T_i} y_{i-1} y_{i+1}$

for i = 2, ..., r - 1. These two properties of functions $\tilde{y}_1, ..., \tilde{y}_r$ imply equations (5) for roots of polynomials $y_1, ..., y_r$, see [MV2]. This shows that the point \boldsymbol{t} is a critical point.

The correspondence between critical points and differential operators with quasipolynomial flags is reflexive. If \boldsymbol{t} is a critical point of the master function $\Phi(\cdot; \boldsymbol{z}; \boldsymbol{\Lambda}; \boldsymbol{l})$, which corresponds by Theorem 3.4 to the differential operator D, then D is the fundamental differential operator of the critical point \boldsymbol{t} .

3.3. Conclusion. Let $\boldsymbol{z} = (z_1, \ldots, z_n) \in \mathbb{C}^n$ be a point with distinct coordinates. Let $\boldsymbol{m}_1, \ldots, \boldsymbol{m}_{r+1}, \boldsymbol{m}_{\infty}$ be an admissible set of sequences of complex numbers as in Section 2.1. Define $\Lambda_1, \ldots, \Lambda_n, \Lambda_\infty$, \boldsymbol{l} by (3). Assume that \boldsymbol{l} is fixed and $\Lambda_1, \ldots, \Lambda_n$ are separating (that is generic). Then the number of differential operators D, associated with $\boldsymbol{z}, \boldsymbol{m}_1, \ldots, \boldsymbol{m}_{r+1}, \boldsymbol{m}_{\infty}$ and having a quasi-polynomial flag, is not greater than $d(n-1, \boldsymbol{l})$, see Theorems 2.2 and 3.4.

It is interesting to note that if $\Lambda_1, \ldots, \Lambda_n, \Lambda_\infty$, are dominant integral, then the number of differential operators D, associated with $\boldsymbol{z}, \boldsymbol{m}_1, \ldots, \boldsymbol{m}_{r+1}, \boldsymbol{m}_\infty$ and having a quasipolynomial flag, is not greater than the multiplicity $\delta(\boldsymbol{\Lambda}, \boldsymbol{l})$, which could be a smaller number. See Theorem 3.4 and description in [MV2] of the critical points of the corresponding master functions.

References

- [BMV] P. Belkale, E. Mukhin, and A. Varchenko, Multiplicity of critical points of master functions and Schubert calculus Preprint (2005), 1–23; math.QA/0503132
- [BBS] J. Borcea, P. Branden, and B. Shapiro, On higher Lame equations, Heine-Stieltjes and Van Vleck polynomials, in preparation
- [H] E. Heine, *Handbuch der Kugelfunctionen*, G. Reimer Verlag, Berlin, vol. 1 (1878), 472–479
- [MTV] E. Mukhin, V. Tarasov, and A. Varchenko, The B. and M. Shapiro conjecture in real algebraic geometry and the Bethe ansatz, Preprint (2005), 1–17, math.AG/0512299
- [MV1] E. Mukhin and A. Varchenko, *Remarks on critical points of phase functions and norms of Bethe vectors*, Adv. Stud. Pure Math. **27** (2000), 239–246
- [MV2] E. Mukhin and A. Varchenko, Critical Points of Master Functions and Flag Varieties, Communications in Contemporary Mathematics 6 (2004), no. 1, 111-163
- [MV3] E. Mukhin and A. Varchenko, Norm of a Bethe Vector and the Hessian of the Master Function, Compos. Math. 141 (2005), no. 4, 1012–1028
- [RV] N. Reshetikhin and A. Varchenko, Quasiclassical asymptotics of solutions to the KZ equations, Geometry, Topology and Physics for R. Bott, Intern. Press, 1995, 293–322

- [SV] V V. Schechtman and A. Varchenko, Arrangements of hyperplanes and Lie algebra homology, Invent. Math. 106 (1991), 139-194
- [ScV] I. Scherbak and A. Varchenko, Critical points of functions, \$ and Fuchsian differential equations with only univalued solutions, Dedicated to Vladimir I. Arnold on the occasion of his 65th birthday, Mosc. Math. J. 3 (2003), no. 2, 621–645
- [St] T. Stieltjes, Sur certains polynomes qui verifient une equation differentielle lineaire du second ordre et sur la theorie des fonctions de Lame, Acta Math. 8 (1885), 321–326
- [Sz] G. Szego, Orthogonal Polynomials, AMS, 1939.
- [V1] A. Varchenko, Hypergeometric Functions and Representation Theory of Lie Algebras and Quantum Groups, Advanced Series in Mathematical Physics, Vol. 21, World Scientific, 1995
- [V2] A. Varchenko, Bethe Ansatz for Arrangements of Hyperplanes and the Gaudin Model, Preprint (2004), 1–17; hep-th/0408001
- [V3] A. Varchenko, Critical points of the product of powers of linear functions and families of bases of singular vectors, Compositio Mathematica 97 (1995), 385– 401