HOLOMORPHIC HOROSPHERICAL TRANSFORM ON NON-COMPACTLY CAUSAL SPACES

SIMON GINDIKIN, BERNHARD KRÖTZ AND GESTUR ÓLAFSSON

ABSTRACT. We develop integral geometry for non-compactly causal symmetric spaces. We define a complex horospherical transform and, for some cases, identify it with a Cauchy type integral.

INTRODUCTION

Within the class of semisimple symmetric spaces Y = G/H we focus on those which can be realized as Shilov boundaries of Stein tubes D = D(Y) in the affine complexification $Y_{\mathbb{C}} = G_{\mathbb{C}}/H_{\mathbb{C}}$. These Y come in two different flavors: compactly causal (CC) and noncompactly causal (NCC) symmetric spaces. It is important to mention that one can realize different series of representations of $L^2(Y)$ as holomorphic functions on D: the holomorphic discrete series for CC and a multiplicity one subspace of the most continuous spectrum in the NCC-case.

In [11] we developed integral geometry for D in the CC-case. If we consider the usual (real) horospherical transform on Y, then holomorphic discrete series lie in its kernel. So we considered a complex version of such a transform - horospherical Cauchy transform using a kernel of Cauchy type with singularities on complex horospheres (on $Y_{\mathbb{C}}$) which do not intersect Y. As a result we constructed a dual domain Ξ_+ in the manifold Ξ of complex horospheres on $Y_{\mathbb{C}}$ and our horospherical transform is an intertwining operator from holomorphic functions on D to holomorphic functions on Ξ_+ which admits an explicit inversion.

In this paper we develop holomorphic integral geometry for NCC-spaces. In [11] we constructed for any NCC-space a G-invariant tube type domain D(Y) in $Y_{\mathbb{C}}$ which has Y as Shilov boundary. The crucial point is that D contains the associated Riemannian symmetric space X = G/K as a totally real submanifold. We have $X_{\mathbb{C}} \simeq Y_{\mathbb{C}}$. Let us remark that all spherical functions on X extend holomorphically to D and admit boundary values on Y. Furthermore, in some cases, D coincides with the maximal G-domain of holomorphy of spherical functions on X – the complex crown of X.

Key words and phrases. Radon transform, horospheres, Hardy spaces.

SG was supported in part by NSF-grant DMS-0070816.

BK was supported by the RiP-program in Oberwolfach, and a Heisenberg fellowship of the DFG.

GÓ was supported by the RiP-program in Oberwolfach, NSF grants DMS-0139783 and DMS-0402068.

SIMON GINDIKIN, BERNHARD KRÖTZ AND GESTUR ÓLAFSSON

There are substantial differences in our constructions of holomorphic horospherical transform for CC and NCC-spaces. In the CC-case we used a Cauchy type integral operator on Y with singularities on complex horospheres which do not intersect Y (horospherical Cauchy transform). For NCC-spaces we are able to construct similar operators in some cases, including the important subclass of Cayley type spaces. In general, we give another construction. We use the fact that complex horospheres which do not intersect Y allow real forms in D and we can construct a complex horospherical transformation extending the real one. Let us mention that this construction does not work in CC-case. If we have both type of constructions the second one can be regarded as a residue of first one.

Let us decsribe our results in more detail. For a fixed choice of horospherical coordinates $X \simeq N \times A$ one parameterizes the set $\Xi_{\mathbb{R}}$ of real horospheres on X by G/MN (with $M = Z_K(A)$ as usual). For Schwartz-functions f on X one defines the real Radon transform

$$\mathcal{R}_{\mathbb{R}}: \mathcal{S}(X) \to C^{\infty}(\Xi_{\mathbb{R}}), \quad \mathcal{R}(f)(\xi) = \int_{\xi} f.$$

Our first goal is to give a holomorphic version of $\mathcal{R}_{\mathbb{R}}$. For that we construct a certain Ginvariant submanifold Ξ_+ of the complex horospheres $\Xi = G_{\mathbb{C}}/M_{\mathbb{C}}N_{\mathbb{C}}$ on $X_{\mathbb{C}} = Y_{\mathbb{C}}$. The complex horospheres of Ξ_+ have the property that they do not intersect Y. Let us point out, that in contrast to the CC-case, Ξ_+ is not longer open in Ξ ; it is a CR-submanifold of complex dimension equal to the rank of Y (i.e. dim A). Next we recall the most-continuous Hardy-space $\mathcal{H}^2(D)$ on D from [11]. This is a Hilbert space of holomorphic functions on Dwith an equivariant isometric boundary map $b : \mathcal{H}^2(D) \to L^2_{\mathrm{mc}}(Y)$ into a multiplicity-one subspace of the most-continuous spectrum of Y. We show that there is a natural dense G-subspace $\mathcal{H}^2(D)_0$ of $\mathcal{H}^2(D)$ whose elements restrict to Schwartz-functions on X and for which $\mathcal{R}_{\mathbb{R}}$ allows a holomorphic G-equivariant extension

$$\mathcal{R}: \mathcal{H}^2(D)_0 \to CR^\infty(\Xi_+).$$

We call \mathcal{R} the holomorphic horospherical Radon transform.

The main result of this paper is an interpretation of a (slightly twisted) Radon transform \mathcal{R}_{χ} as a Cauchy-type integral. For the highly symmetric subclass of Cayley-type spaces CT=CC∩NCC we show in Theorem 5.3 that there is Cauchy-type kernel $\mathcal{K}_{\chi}(,) \in CR^{\infty}_{\gamma}(\Xi_{+}) \hat{\otimes} \mathcal{S}(Y)$ such that

(0.1)
$$\mathcal{R}_{\chi}(f)(\cdot) = \frac{1}{(2\pi)^n} \int_Y b(f)(y) \ \mathcal{K}_{\chi}(\cdot, y) \ dy \in CR_{\chi}^{\infty}(\Xi_+) \,.$$

Note that (0.1) is in analogy to our definition of the holomorphic horospherical Radon transform for the CC-case [13]. For the general case of an NCC-space we believe that there is an integral representation of \mathcal{R}_{χ} as in (0.1) but we are not aware of such. We consider the definition of a holomorphic horospherical Radon transform in terms of a Cauchy type integral to be most robust for further generalization to other symmetric spaces or different series of representations.

For the CC-case, following [9], we found an inversion formula for the holomorphic horospherical transform of compelling beauty:

$$f = (\mathcal{LR}f)^{\vee}.$$

Here \mathcal{L} is a differential operator and $\phi \mapsto \phi^{\vee}$ is a dual transform [13] with regard to a double fibration which brings D and Ξ_+ into duality. Let us remark that the situation is different for the NCC-spaces – for the basic example $Y = \text{Sl}(2, \mathbb{R})/\text{SO}(1, 1)$ we show that \mathcal{L} is a pseudo differential but not a differential operator (cf. Section 6).

The paper is concluded with a geometric definition of the most continuous Hardy space $\mathcal{H}^2(D)$ which is of independent interest.

Achknowledgement: We thank the referee for his patient pressure on more precision. It is due to him that the paper is now more readable.

1. Horospheres on symmetric spaces of triangular type

In [11] we associated to every NCC-symmetric space Y = G/H a G-Stein manifold D with the following properties:

- (1) The complex manifold D has a natural G-realization in the complexification $Y_{\mathbb{C}}$ of Y;
- (2) The symmetric space Y = G/H is G-isomorphic to the distinguished (Shilov) boundary of D.

The objective of this section is to study the space $\Xi = G_{\mathbb{C}}/M_{\mathbb{C}}N_{\mathbb{C}}$ of horospheres in $Y_{\mathbb{C}}$ in relation to D. In particular we will introduce a natural G-invariant CR-manifold $\Xi_+ \subset \Xi$ whose elements have the properties that they do non intersect the real space Y, i.e. have no real points.

1.1. Notation. We begin with the construction of NCC-symmetric spaces from the complex geometric point of view.

Let us denote by \mathfrak{g} a simple non-compact real Lie algebra. We fix a Cartan involution $\theta: \mathfrak{g} \to \mathfrak{g}$, write $\mathfrak{g} = \mathfrak{k} + \mathfrak{p}$ for the associated eigenspace decomposition and select a maximal abelian subspace $\mathfrak{a} \subset \mathfrak{p}$. Write $\Pi = \{\alpha_1, \ldots, \alpha_n\}$ for a basis of the restricted root system $\Sigma = \Sigma(\mathfrak{g}, \mathfrak{a})$ and expand the highest root β in terms of Π

$$\beta = k_1 \alpha_1 + \ldots + k_n \alpha_n \qquad (k_i \in \mathbb{Z}_{>0}).$$

One calls α_i miniscule if $k_i = 1$. Let us recall that miniscule elements exist if and only if Σ is not of the type E_8 , F_4 or G_2 . For the remainder we fix a miniscule root α and define $Z \in \mathfrak{a}$ by the requirement $\alpha_i(Z) = \delta_{\alpha_i \alpha}$. Observe that Z gives rise to a triangular decomposition

(1.1)
$$\mathfrak{g} = \mathfrak{g}_{-1} + \mathfrak{g}_0 + \mathfrak{g}_1$$

with $\mathfrak{g}_j = \{X \in \mathfrak{g} \mid [Z, X] = jX\}$. Further the 3-grading (1.1) defines an involution σ of \mathfrak{g} by $\sigma(X) = (-1)^j X$ for $X \in \mathfrak{g}_j$ and eigenspace decomposition

$$\mathfrak{g} = \mathfrak{g}_0 + (\mathfrak{g}_{-1} + \mathfrak{g}_1).$$

By construction σ and θ commute and hence $\tau = \sigma \circ \theta$ defines an involution whose eigenspace decomposition shall be denoted $\mathfrak{g} = \mathfrak{h} + \mathfrak{q}$. The so-obtained symmetric pairs $(\mathfrak{g}, \mathfrak{h})$ we call non-compactly causal (NCC). It is an elementary exercise to classify all NCC-symmetric pairs.

We denote by \mathcal{W} the Weyl group of Σ , set $Z_H = \frac{\pi}{2}Z$ and define $\Omega_H \subseteq \mathfrak{a}$ by

(1.2)
$$\Omega_H = \inf\{\text{convex hull of } \mathcal{W}(Z_H)\}.$$

Observe that $\overline{\Omega}_H$ is a compact convex subset of \mathfrak{a} with extreme points $\mathcal{W}(Z_H)$.

On the global level we fix a linear connected Lie group G with Lie algebra \mathfrak{g} . We denote by $G_{\mathbb{C}}$ the corresponding linear connected complexification of G. Further we request that τ as well is complex linear extension $\tau_{\mathbb{C}}$ exponentiate to involutions on G, resp. $G_{\mathbb{C}}$, and we denote by H, resp. $H_{\mathbb{C}}$ their corresponding fixed point groups. In this way we obtain a totally real embedding

$$Y \hookrightarrow Y_{\mathbb{C}} = G_{\mathbb{C}}/H_{\mathbb{C}}$$

of Y in the Stein symmetric space $Y_{\mathbb{C}}$. We refer to Y as a non-compactly causal symmetric space (NCC).

Attached to Y and Ω_H comes a Stein manifold D which we will now describe. Write K < G for the compact group of θ -fixed elements and X = G/K for the corresponding Riemann symmetric space. As before we obtain a totally real embedding

$$X \hookrightarrow X_{\mathbb{C}} = G_{\mathbb{C}}/K_{\mathbb{C}}$$
.

Recall that $H_{\mathbb{C}}$ and $K_{\mathbb{C}}$ are conjugate, i.e. with $z_H = \exp(iZ_H)$ we have cf. [11]:

(1.3)
$$e^{i\mathrm{ad}(Z_H)}\mathfrak{t}_{\mathbb{C}} = \mathfrak{h}_{\mathbb{C}}$$
 and $\mathrm{Ad}(z_H)K_{\mathbb{C}} = H_{\mathbb{C}}$.

Hence $X_{\mathbb{C}}$ and $Y_{\mathbb{C}}$ are canonically $G_{\mathbb{C}}$ -isomorphic via the map

$$X_{\mathbb{C}} \ni gK_{\mathbb{C}} \mapsto gz_{H}^{-1}H_{\mathbb{C}} \in Y_{\mathbb{C}}.$$

In the sequel we identify $X_{\mathbb{C}}$ with $Y_{\mathbb{C}}$.

We write $x_o = K_{\mathbb{C}} \in X_{\mathbb{C}}$ for the base point in $X_{\mathbb{C}}$ and set

$$D = G \exp(i\Omega_H) \cdot x_o.$$

Note that D was denoted by Ξ_H in our previous article [11]. According to [11], D is an open G-invariant Stein neighborhood of X in $X_{\mathbb{C}} = Y_{\mathbb{C}}$. Moreover, the map $Y = G/H \ni gH \mapsto gz_H \cdot x_o \in X_{\mathbb{C}}$ identifies Y with the distinguished boundary $\partial_d D$ of D (see [11], Section 1, for more details).

In summary, the symmetric Stein manifold $X_{\mathbb{C}} = Y_{\mathbb{C}}$ admits 2 real forms X and Y and a Stein neighborhood D of X with Y as its Shilov boundary.

1.2. Complex horospheres. From now we let Y = G/H be an NCC symmetric space and X = G/K its Riemannian counterpart.

In this section we introduce the G-space of horospheres in the complex manifold $X_{\mathbb{C}} = Y_{\mathbb{C}}$. This was done for CC-spaces in [13].

We begin with some general remarks on convexity which we will use frequently. Let G = NAK be an Iwasawa decomposition of G and $N_{\mathbb{C}}A_{\mathbb{C}}K_{\mathbb{C}} \subsetneq G_{\mathbb{C}}$ its Zariski-open complexification. In particular, $N_{\mathbb{C}}A_{\mathbb{C}} \cdot x_o$ is a Zariski-open subset in the affine variety $X_{\mathbb{C}}$. Define the finite 2-group $F = A_{\mathbb{C}} \cap K_{\mathbb{C}}$ and note that there are well defined holomorphic maps

$$n: N_{\mathbb{C}}A_{\mathbb{C}} \cdot x_o \to N_{\mathbb{C}}, \qquad a: N_{\mathbb{C}}A_{\mathbb{C}} \cdot x_o \to A_{\mathbb{C}}/F$$

such that $z = n(z)a(z) \cdot x_o$ for all $z \in N_{\mathbb{C}}A_{\mathbb{C}} \cdot x_o$. Now, the fact that D is contractible and $D \subset N_{\mathbb{C}}A_{\mathbb{C}} \cdot x_o$, implies that $a|_D$ admits a well defined holomorphic logarithm

 $\log a: D \to \mathfrak{a}_{\mathbb{C}}$.

For $Z \in \Omega_H$, the complex convexity theorem (cf. [10, 18]) then implies that

(1.4)
$$\operatorname{Im} \log a(G \exp(iZ) \cdot x_o) = \operatorname{conv}(\mathcal{W} \cdot Z)$$

where $\operatorname{conv}(\cdot)$ denotes the convex hull of (\cdot) .

Submanifolds of $X_{\mathbb{C}}$ of the type

$$gN_{\mathbb{C}} \cdot x_o \qquad (g \in G_{\mathbb{C}})$$

will be referred as *horospheres*. We denote by $\operatorname{Hor}(X_{\mathbb{C}})$ the set of all horospheres on $X_{\mathbb{C}}$ and note that $\operatorname{Hor}(X_{\mathbb{C}})$ has a natural *G*-structure $(g, hN_{\mathbb{C}} \cdot x_o) \mapsto ghN_{\mathbb{C}} \cdot x_o$.

To understand the space horospheres and the related harmonic analysis it is useful to bring them in the context of a double fibration. Set $M = Z_K(A) \subset M_{\mathbb{C}} = Z_{K_{\mathbb{C}}}(A)$, define

$$\Xi = G_{\mathbb{C}}/M_{\mathbb{C}}N_{\mathbb{C}}$$

and consider:



Then horospheres in $X_{\mathbb{C}}$ are exactly the subsets of $X_{\mathbb{C}}$ of the form

(1.6)
$$E(\xi) = \pi_2(\pi_1^{-1}(\xi)) \quad (\xi \in \Xi).$$

If $\xi_o = M_{\mathbb{C}} N_{\mathbb{C}} \in \Xi$ denotes the base point and $\xi = g \cdot \xi_o \in \Xi$ then, using that $M_{\mathbb{C}} \subset H_{\mathbb{C}}$, we have:

$$E(\xi) = gM_{\mathbb{C}}N_{\mathbb{C}} \cdot x_o = gN_{\mathbb{C}} \cdot x_o \subset X_{\mathbb{C}}$$

Similarly, for $z \in X_{\mathbb{C}}$ we set

(1.7)
$$S(z) = \pi_1(\pi_2^{-1}(z)) .$$

If $z = g \cdot x_o$ for $g \in G_{\mathbb{C}}$, then $S(z) = gK_{\mathbb{C}} \cdot \xi_o$. Moreover, for $z \in X_{\mathbb{C}}$ and $\xi \in \Xi$ one has the incidence relations

(1.8)
$$z \in E(\xi) \iff \pi_1^{-1}(\xi) \cap \pi_2^{-1}(z) \neq \emptyset \iff \xi \in S(z).$$

Proposition 1.1. The map

$$\Xi \to \operatorname{Hor}(X_{\mathbb{C}}), \ \xi \mapsto E(\xi)$$

is a $G_{\mathbb{C}}$ -equivariant bijection.

Proof. $G_{\mathbb{C}}$ -equivariance and surjectivity are clear. The injectivity follows the same way as in the proof of Proposition 2.1 in [13] by replacing $H_{\mathbb{C}}$ by $K_{\mathbb{C}}$.

One of the important features of Ξ is, that there exists a right $A_{\mathbb{C}}$ -action on Ξ that commutes with the left $G_{\mathbb{C}}$ -action. For $\xi = g \cdot \xi_o$ and $a \in A_{\mathbb{C}}$ we set

(1.9)
$$\xi \cdot a = ga \cdot \xi_o$$

Since $A_{\mathbb{C}}$ normalizes $M_{\mathbb{C}}N_{\mathbb{C}}$ it is clear that (1.9) is well defined. From the definition it is also clear that the left $G_{\mathbb{C}}$ -action and the right $A_{\mathbb{C}}$ -action commutes. In this way we obtain an action of $G_{\mathbb{C}} \times A_{\mathbb{C}}$ on Ξ by

$$(G_{\mathbb{C}} \times A_{\mathbb{C}}) \times \Xi \to \Xi, \quad ((g, a), \xi) \mapsto g \cdot \xi \cdot a$$

We conclude this subsection with an alternative characterization of horospheres as level sets of holomorphic functions. For that let $\{\omega_1, \ldots, \omega_n\} \subset \mathfrak{a}^*$ be the set of fundamental K-spherical lowest weights. For each $1 \leq j \leq n$ we write (π_j, V_j) for the corresponding finite dimensional representation of G with lowest weight ω_j . We extend this representation to a holomorphic representation of $G_{\mathbb{C}}$ which we denote by the same symbol. Endow V_j with a complex bilinear pairing \langle , \rangle such that $\langle \pi_j(g)v, w \rangle = \langle v, \pi_j(\theta(g)^{-1})w \rangle$ for all $v, w \in V_j$ and $g \in G_{\mathbb{C}}$. Such a form \langle , \rangle exists as $\pi \circ \theta$ is isomorphic to the representation contragredient to π_j . We write $v_j \in V_j$ for a lowest weight vector and $\eta_j \in V_j$ for a $K_{\mathbb{C}}$ -fixed vector subject to the normalization $\langle \eta_j, v_j \rangle = 1$. Finally, define holomorphic functions on $f_j : G_{\mathbb{C}} \to \mathbb{C}$ by

(1.10)
$$f_j(g) = \langle \pi_j(g)\eta_j, v_j \rangle \qquad (g \in G_{\mathbb{C}}).$$

Note, that we have

(1.11)
$$f_j(nak) = a^{\omega_j}$$

for all $n \in N_{\mathbb{C}}$, $k \in K_{\mathbb{C}}$ and $a \in A_{\mathbb{C}}$. Here, as elsewhere in this article, we use the notation $a^{\mu} = e^{\mu(X)}$ if $a = \exp X \in A_{\mathbb{C}}$. We recall that

(1.12)
$$G_{\mathbb{C}} \setminus N_{\mathbb{C}} A_{\mathbb{C}} K_{\mathbb{C}} = \{ g \in G_{\mathbb{C}} \mid \prod_{j=1}^{n} f_j(g) = 0 \}$$

(see [2], Lemma 3.4).

Lemma 1.2. $N_{\mathbb{C}}K_{\mathbb{C}} = \{g \in G_{\mathbb{C}} \mid f_j(g) = 1 \text{ for all } 1 \leq j \leq n\}.$

Proof. This follows from (1.11) and (1.12).

We will often view f_j , or more generally it left translates, as a function on $X_{\mathbb{C}}$. We will also, without further comments, view the function $g \mapsto f_j(g^{-1}z), z \in X_{\mathbb{C}}$ as a function on Ξ . With that in mind we have:

Lemma 1.3. Let $\xi \in \Xi$ and $x \in X_{\mathbb{C}}$. Then

$$E(\xi) = \{ z \in X_{\mathbb{C}} \mid f_j(\xi^{-1}z) = 1 \quad \text{for all} \quad 1 \le j \le n \}$$

and

$$S(x) = \{ \varrho \in \Xi \mid f_j(\varrho^{-1}x) = 1 \quad for \ all \quad 1 \le j \le n \}$$

Proof. Notice that $E(g \cdot \xi_o) = gE(\xi_o)$ and $S(g \cdot x_o) = gS(x_o)$. We can therefore assume that $\xi = \xi_o$ and $x = x_o$. Now, the claim is a reformulation of Lemma 1.2.

1.3. Some G-submanifolds of Ξ . We define the G-space of real horospheres in X as

$$\Xi_{\mathbb{R}} = G/MN$$

Then $\Xi_{\mathbb{R}} \subset \Xi = G_{\mathbb{C}}/M_{\mathbb{C}}N_{\mathbb{C}}$ is obviously a totally real *G*-submanifold of Ξ and the right *A*-action leaves $\Xi_{\mathbb{R}}$ invariant.

Let $T = \exp(i\mathfrak{a}) \subset G_{\mathbb{C}}$ and note that $A_{\mathbb{C}} = A \times T$; note that $F = K \cap T$. We contrast $\Xi_{\mathbb{R}}$ with the $G \times A_{\mathbb{C}}$ -invariant subset of Ξ

(1.13)
$$\Xi_0 = G \cdot \xi_o \cdot A_{\mathbb{C}} = G A_{\mathbb{C}} \cdot \xi_o.$$

Proposition 1.4. *The following assertion hold:*

(1) The map

(1.14)
$$K/M \times_F A_{\mathbb{C}} \to \Xi_0, \quad [kM, a] \mapsto ka \cdot \xi_o$$

is a real analytic isomorphism.

(2) The map

$$\Xi_{\mathbb{R}} \times_F T \to \Xi_0, \quad [gMN, t] \mapsto gt \cdot \xi_o$$

is a G-equivariant real analytic diffeomorphism.

Proof. (i) follows from the fact that G = KAN and $NA_{\mathbb{C}} \subset A_{\mathbb{C}}N_{\mathbb{C}}$. Finally (ii), is a consequence of (i).

Note that (1.14) describes a natural CR-structure on Ξ_0 of CR-dimension dim A and CR-codimension dim K/M.

Define a tube domain in $A_{\mathbb{C}}$ by

$$T(\Omega_H) = \exp(\mathfrak{a} + i\Omega_H) = A \exp(i\Omega_H) \simeq \mathfrak{a} + i\Omega_H$$

and set

(1.15)
$$\Xi_{+} = G \exp(i\Omega_{H}) \cdot \xi_{o} = KT(\Omega_{H}) \cdot \xi_{o}$$

7

Then Ξ_+ is a real-analytic, *G*-invariant open submanifold of Ξ_0 . In particular Ξ_+ is a *CR*-manifold. The coordinate decomposition of Ξ_0 slightly simplifies for Ξ_+ .

Proposition 1.5. For Ξ_+ the following assertions hold:

(1) The map

$$K/M \times T(\Omega_H) \to \Xi_+, \quad (kM, a) \mapsto ka \cdot \xi_o$$

is a real analytic isomorphism.

(2) The map

$$\Xi_{\mathbb{R}} \times \Omega_H \to \Xi_+, \quad (gMN, Z) \mapsto g \exp(iZ) \cdot \xi_o$$

is a G-equivariant real analytic diffeomorphism.

We conclude this section with a remark on the structure of Ξ_+ .

Remark 1.6. (Shilov boundary of Ξ_+) The map

$$\Xi_{\mathbb{R}} \to \partial \Xi_+, \quad gMN \mapsto gz_H \cdot \xi_o$$

identifies $\Xi_{\mathbb{R}}$ as the Shilov boundary $\partial_S \Xi_+$ of Ξ_+ . In this sense $\Xi_{\mathbb{R}}$ parameterizes the real horospheres on Y (see also Lemma 1.7 below).

1.4. Horospheres without real points. The aim of this subsection is to show that horospheres corresponding to Ξ_+ do not contain real points, i.e., are disjoint form Y.

Recall from Subsection 1.1 that we identify Y = G/H with the (Shilov) boundary orbit $G \cdot y_o \subset X_{\mathbb{C}}$ of $y_o = z_H \cdot x_o$ in $G_{\mathbb{C}}/K_{\mathbb{C}}$. Define the parameter set of horospheres without real points by

(1.16)
$$\Xi_{\rm nr} = \{\xi \in \Xi \mid E(\xi) \cap Y = \emptyset\}.$$

The following statement should be compared to the complex convexity Theorem (1.4); it means that convexity breaks down at the extreme points of Ω_H .

Lemma 1.7. Let $\mathcal{U} = \bigcup_{w \in \mathcal{W}} NAwH$. Then $\mathcal{U} \cdot y_o$ is open and dense in $G \cdot y_o$ and $G \cdot y_o \cap N_{\mathbb{C}}A_{\mathbb{C}} \cdot x_o = \mathcal{U} \cdot y_o = NA\mathcal{W}z_H \cdot x_o$.

Proof. It is a special case of a Theorem by Rossmann-Matsuki (cf. [21]) that \mathcal{U} is dense in G. Hence $\mathcal{U} \cdot y_o$ is dense in $G \cdot y_o$. As $z_H^{-1}H_{\mathbb{C}}z_H = K_{\mathbb{C}}$ (cf. (1.3)), it follows that $\mathcal{U} \cdot y_o = NA\mathcal{W}z_H \cdot x_o$. It remains to show the inclusion " \supseteq " for the first asserted equality. But this follows from (1.12).

We can now prove the main result of this subsection.

Theorem 1.8. $\Xi_+ \subseteq \Xi_{nr}$.

Proof. We argue by contradiction. Note that $E(\xi) \cap Y \neq \emptyset$ for some $\xi \in \Xi_+$ means that there exist $Z \in \Omega_H$ such that

 $Gz_H \cdot x_o \cap \exp(iY) N_{\mathbb{C}} \cdot x_o \neq \emptyset$.

Now $\exp(iY)N_{\mathbb{C}} \cdot x_o = N_{\mathbb{C}}\exp(iY) \cdot x_o \subset N_{\mathbb{C}}\exp(i\Omega_H) \cdot x_o$ and the assertion follows from Lemma 1.7.

1.5. Real forms of $E(\xi)$ and S(z). In this last part of this section we introduce certain G-invariant real forms of the complex manifolds $E(\xi)$ and S(z).

We begin with the horospheres. For $\xi = ga \cdot \xi_o \in \Xi_+$, with $g \in G$ and $a \in \exp(i\Omega_H)$, define

(1.17)
$$E_{\mathbb{R}}(\xi) = gNa \cdot x_o \subset E(\xi) \,.$$

Then $E_{\mathbb{R}}(\xi)$ is well defined, *G*-invariant and a totally real submanifold of $E(\xi)$. Further, the assignment $\Xi_+ \ni \xi \mapsto E_{\mathbb{R}}(\xi)$ is *G*-equivariant.

Next we consider $S(z) \simeq K_{\mathbb{C}}/M_{\mathbb{C}}$. Because of the relation $K_{\mathbb{C}} = z_H^{-1} H_{\mathbb{C}} z_H$ there are two natural real forms. Accordingly we define for $z = ga \cdot x_o \in D$:

(1.18)
$$S_{\mathbb{R}}^{K}(z) = gaK \cdot \xi_{o} \quad \text{and} \quad S_{\mathbb{R}}^{H}(z) = gaz_{H}^{-1}Hz_{H} \cdot \xi_{o}.$$

Obviously $S_{\mathbb{R}}^{K}(z)$ and $S_{\mathbb{R}}^{H}(z)$ are *G*-invariant totally real submanifold of $S(z) = gaK_{\mathbb{C}} \cdot x_{o}$ and the maps $D \ni z \mapsto S_{\mathbb{R}}^{K}(z)$ and $D \ni z \mapsto S_{\mathbb{R}}^{H}(z)$ are *G*-equivariant. Note that $S_{\mathbb{R}}^{H}(z) \simeq H/M$ as manifolds.

1.6. Invariant measure on Y. Lemma 1.7 allows for a natural normalization of the invariant measure on Y. Assume that invariant measures on G, A and N have been fixed and let $\mathcal{W}_H = N_{K \cap H}(\mathfrak{a})/Z_{H \cap K}(\mathfrak{a})$ be the small Weyl group. By Lemma 1.7 the union

$$\mathcal{U} = \bigcup_{w \in \mathcal{W}/\mathcal{W}_H} ANw \cdot y_o$$

is disjoint open and dense in Y. As the complement is an analytic set, it follows that $Y \setminus \mathcal{U}$ has measure zero. We can normalize the invariant measure on Y such that for all $f \in L^1(Y)$:

$$\int_Y f(y) \, dy = \sum_{w \in \mathcal{W}/\mathcal{W}_H} \int_A \int_N f(anw \cdot y_o) \, da \, dn \, .$$

2. The Frechét module $CR^{\infty}(\Xi_+)$

In this section we use the right A-action on $\Xi_{\mathbb{R}}$ and Ξ_+ to define G-submodules of the smooth A-covariant functions on $\Xi_{\mathbb{R}}$ respectively CR-functions on Ξ_+ . Those modules are the standard realization respectively a CR-realization of the space of smooth vectors in the principal series representations given by induction from the right. Note that later we will use the induction from the left.

Recall, that A acts on the space of horospheres from the right. This action induces a right regular representation of A on any function space on $\Xi_{\mathbb{R}}$, Ξ_+ or any other right invariant set of horospheres given by

$$(R(a)f)(\xi) = f(\xi \cdot a).$$

Let $\rho = 1/2 \sum_{\alpha \in \Sigma^+} (\dim \mathfrak{g}^{\alpha}) \cdot \alpha$. Let $\lambda \in \mathfrak{a}_{\mathbb{C}}^*$.

The index λ will denote the subspace of $(\lambda - \rho)$ -covariant functions. In particular

$$C^{\infty}(\Xi_{\mathbb{R}})_{\lambda} = \{ f \in C^{\infty}(\Xi_{\mathbb{R}}) \mid (\forall a \in A) \ R(a)f = a^{\lambda - \rho}f \}.$$

We recall that G acts $C^{\infty}(\Xi_{\mathbb{R}})$ by left translations in the argument

$$(L(g)f)(\xi) = f(g^{-1} \cdot \xi)$$

for $g \in G$, $f \in C^{\infty}(\Xi_{\mathbb{R}})$ and $\xi \in \Xi_{\mathbb{R}}$. The so obtained representation $(L, C^{\infty}(\Xi_{\mathbb{R}})_{\lambda})$ is the smooth model of the spherical principal series with parameter λ .

Write $CR^{\infty}(\Xi_{+})$ for the space of smooth CR-functions on Ξ^{+} and set

$$CR^{\infty}(\Xi_{+})_{\lambda} = \{ f \in CR^{\infty}(\Xi_{+}) \mid (\forall a \in A) \ R(a)f = a^{\lambda-\rho}f \}.$$

As characters on A extend to holomorphic functions on $T(\Omega_H)$, it follows that the restriction map

(2.1)
$$\operatorname{Res}_{\lambda} : CR^{\infty}(\Xi_{+})_{\lambda} \to C^{\infty}(\Xi_{\mathbb{R}})_{\lambda}, \quad f \mapsto f|_{\Xi_{\mathbb{R}}}$$

is a G-equivariant topological isomorphism of G-modules.

2.1. *CR*-realization of the *H*-spherical holomorphic vector. For each $\lambda \in \mathfrak{a}_{\mathbb{C}}^*$ we define a certain *H*-invariant element $f_{\lambda} \in CR^{-\infty}(\Xi_+)_{\lambda}$ which was called the *H*-spherical holomorphic distribution vector in [11]. The generalized function f_{λ} is defined by

$$f_{\lambda}(\xi) = a(\xi^{-1}z_H^{-1})^{\rho-\lambda} \qquad (\xi \in \Xi_+).$$

We notice that on the dense subset

$$\Xi'_{+} = \bigcup_{w \in \mathcal{W}} HwT(\Omega_{H}) \cdot \xi_{o}$$

of Ξ_+ , the function belongs to $CR^{\infty}(\Xi'_+)_{\lambda}$ and is given by

$$f_{\lambda}(hwa \cdot \xi_o) = (w^{-1}z_Hw)^{\lambda-\rho}a^{\lambda-\rho}.$$

For $\Re \lambda \ll 0$, this function is actually continuous on Ξ_+ and the meromorphic continuation in λ as a distribution is achieved with Bernstein's theorem [4]. There are no singularities on the imaginary axis $i\mathfrak{a}^*$ and we arrive at a well defined analytic assignment

$$i\mathfrak{a}^* \to CR^{-\infty}(\Xi_+)^H_\lambda, \ \lambda \mapsto f_\lambda$$

(cf. [11], Th. 2.4.1).

3. The holomorphic horospherical Radon transform

The real Radon transform on X is the G-equivariant injective map

(3.1)
$$\mathcal{R}_{\mathbb{R}}: \mathcal{S}(X) \to C^{\infty}(\Xi_{\mathbb{R}}), \quad \mathcal{R}_{\mathbb{R}}(f)(g \cdot \xi_o) = \int_N f(gn \cdot x_o) \, dn(\quad (g \in G)).$$

The purpose of this section is to show that $\mathcal{R}_{\mathbb{R}}$ has a natural extension to a *G*-equivariant map

$$\mathcal{R}:\mathcal{H}^2(D)_0\to CR^\infty(\Xi_+)$$

which we call the holomorphic horospherical Radon transform. Here $\mathcal{H}^2(D)_0 \hookrightarrow L^2(X)$ is a dense a subspace of the most-continuous Hardy space $\mathcal{H}^2(D) \subset \mathcal{O}(D)$ of $L^2(Y)$ (cf. [11]). On the infinitesimal level this extension is related to the previously established fact (2.1), i.e. $C^{\infty}(\Xi_{\mathbb{R}})_{\lambda}$ is canonically *G*-isomorphic to $CR^{\infty}(\Xi_{+})_{\lambda}$ via restriction.

This section is organized as follows: First we have to recall some facts about the Fourier analysis on X, in particular Arthur's spectral characterization of the Schwartz space C(X). Subsequently we give a brief summary on the most-continuous Hardy space $\mathcal{H}^2(D)$ of [11]. Finally we define the holomorphic horospherical Radon transform \mathcal{R} and discuss some of its properties.

3.1. Fourier analysis on X. We recall the compact realization of the principal series representations. Let $B = M \setminus K$. For $\lambda \in \mathfrak{a}_{\mathbb{C}}^*$ define a representation π_{λ} of G on $L^2(B)$ by

(3.2)
$$\pi_{\lambda}(g)f(Mk) = a(kg)^{\rho-\lambda}f(Mk(kg)).$$

Then $(\pi_{\lambda}, L^2(B))$ is unitary if $\lambda \in i\mathfrak{a}^*$. We write $\mathcal{H}_{\lambda} = L^2(B)$ to indicate the dependence of the *G*-action on λ . Let $\mathfrak{a}^+ = \{H \in \mathfrak{a} \mid (\forall \alpha \in \Delta^+) \alpha(H) > 0\}$ and

$$\mathfrak{a}_{+}^{*} = \left\{ \lambda \in \mathfrak{a} \mid (\forall H \in \mathfrak{a}^{+}) \, \lambda(H) > 0 \right\}.$$

Denote by \hat{G}_{r} the reduced dual of G and by \hat{G}_{rsp} the spherical reduced dual. Then $i\mathfrak{a}_{+}^{*} \ni \lambda \mapsto [\pi_{\lambda}] \in \hat{G}_{rsp}$ is an isomorphism of measure spaces. Here $[\pi_{\lambda}]$ denotes the equivalence class of π_{λ} . We have

(3.3)
$$L^{2}(X) \simeq \int_{i\mathfrak{a}_{+}^{*}}^{\oplus} \mathcal{H}_{\lambda} \frac{d\lambda}{|\mathbf{c}(\lambda)|^{2}}$$

where $\mathbf{c}(\lambda)$ is the Harish-Chandra *c*-function. To explain the above isomorphism, we need some basic fact on the Fourier transform on X. For that, recall that $a: X \to A$ denotes the A-projection with regard to the horospherical coordinates $X = NA \cdot x_o \simeq N \times A$. Set $\mathcal{X} = B \times i\mathfrak{a}^*_+$ and define a Radon measure $d\mu_{\mathcal{X}}$ on \mathcal{X} by

$$d\mu_{\mathcal{X}}(b,\lambda) := db \frac{d\lambda}{|\mathbf{c}(\lambda)|^2}$$

For $f \in L^1(X) \cap L^2(X)$ define its spherical Fourier transform $\hat{f} : \mathcal{X} \to \mathbb{C}$ by

$$\hat{f}(b,\lambda) = \int_X f(x)a(bx)^{\rho-\lambda} dx$$

We will also write $\mathcal{F}_X(f)$ for \hat{f} . We can normalize the left-Haar measure dx on X such that the Fourier transform extends to an unitary isomorphism $\hat{}: L^2(X) \to L^2(\mathcal{X}, d\mu_{\mathcal{X}})$. If fis rapidly decreasing (see exact definition in a moment), then the Fourier inversion formula holds pointwise:

$$f(x) = \int_{\mathcal{X}} \hat{f}(b,\lambda) a(bx)^{\rho+\lambda} d\mu_{\mathcal{X}}(b,\lambda) \qquad (x \in X) \,.$$

For $\lambda \in i\mathfrak{a}^*$ define $\hat{f}_{\lambda} \in L^2(B)$ by $b \mapsto \hat{f}_{\lambda}(b) = \hat{f}(b,\lambda)$. Then the isomorphism in (3.3) is given by

$$L^2(X) \ni f \mapsto (\hat{f}_{\lambda})_{\lambda} \in \int_{i\mathfrak{a}^*_+}^{\oplus} \mathcal{H}_{\lambda} \frac{d\lambda}{|\mathbf{c}(\lambda)|^2}.$$

In the following we will also need the operator valued Fourier transform. If \mathcal{H} is a Hilbert space, then $B_2(\mathcal{H}) \simeq \mathcal{H} \hat{\otimes} \mathcal{H}^*$ denotes the Hilbert space of Hilbert-Schmidt operators on \mathcal{H} . Write

$$L^2(G)_{\mathrm{sph}} = \int_{i\mathfrak{a}^*_+}^{\oplus} B_2(\mathcal{H}_{\lambda}) \; \frac{d\lambda}{|\mathbf{c}(\lambda)|^2}$$

for the K-spherical spectrum in $L^2(G)$. Recall that the isomorphism is given by the operator valued Fourier transform $\mathcal{F}(f)(\lambda) = \int_G f(x)\pi_\lambda(x) \, dx, \ f \in L^1(G) \cap L^2(G)$. The inverse map is

$$f(g) = \int_{i\mathfrak{a}^*_+} \operatorname{Tr}(\pi_{\lambda}(g^{-1})\mathcal{F}(f)(\lambda)) \frac{d\lambda}{|\mathbf{c}(\lambda)|^2}.$$

The constant function $v_{K,\lambda} = \mathbf{1}_B$ defines a normalized K-fixed vector in \mathcal{H}_{λ} . Assume that $f \in L^1(G) \cap L^2(G)_{\text{sph}}$. Then, because $\mathcal{F}(f)(\lambda) = \mathcal{F}(R_k f)(\lambda) = \mathcal{F}(f)(\lambda)\pi_{\lambda}(k)$, it follows, that

(3.4)
$$\mathcal{F}(f)(\lambda)v = \langle v, v_{K,\lambda} \rangle \mathcal{F}(f)(\lambda)v_{K,\lambda} = \langle v, v_{K,\lambda} \rangle \hat{f}_{\lambda} \,.$$

For $x = k_1 \exp(Z)k_2 \in G$, with $Z \in \mathfrak{a}$ and $k_1, k_2 \in K$, let $\sigma(x) = -B(Z, d\theta(\mathbf{1})(\mathbf{Z}))$, where *B* is the Killing form on \mathfrak{g} . Denote by $U(\mathfrak{g})$ the Universal enveloping algebra of \mathfrak{g} and by φ_0 the basic spherical function. For $D, E \in U(\mathfrak{g}), s \in \mathbb{R}$ and $f \in C^{\infty}(G)$, let

$$p_{D,E,s}(f) := \sup_{x \in G} |L_D R_E f(x)| \varphi_0(x)^{-1} (1 + \sigma(x))^s \,.$$

Then $\mathcal{C}(G)$ is the space of smooth functions on G such that $p_{D,E,s}(f) < \infty$ for all such E, D and s, cf. [15], §9. We set

$$\mathcal{C}(G)_{\mathrm{sph}} = \mathcal{C}(G) \cap L^2(G)_{\mathrm{sph}}.$$

For $\tau \in \hat{K}$ we write $|\tau|$ for the norm of the corresponding highest weight. If $\tau \in \hat{K}$, then we write $L^2(B)_{\tau}$ for the subspace of $L^2(B)$ which transform under τ . We denote by $\mathbb{D}(i\mathfrak{a}^*)$ the algebra of all constant coefficient differential operators on \mathfrak{a}^* and set $\mathcal{S}(i\mathfrak{a}^*_+) = \{f|_{i\mathfrak{a}^*_+} | f \in \mathcal{S}(i\mathfrak{a}^*)\}$.

We cite a theorem of Arthur [1], p. 4719, specialized to the spherical case:

Theorem 3.1. The operator valued Fourier transform \mathcal{F} is a topological linear isomorphism from $\mathcal{C}(G)_{\text{sph}}$ onto

$$\{A(\cdot) \in \int_{i\mathfrak{a}^*_+}^{\oplus} B_2(\mathcal{H}_{\lambda}) \frac{d\lambda}{|\mathbf{c}(\lambda)|^2} \mid (\forall v, w \in L^2(B)_{\mathrm{K-fin}}) \langle A(\cdot)v, w \rangle \in \mathcal{S}(i\mathfrak{a}^*_+), \\ \forall D \in \mathbb{D}(i\mathfrak{a}^*), \forall n \in \mathbb{N}_0 \sup_{\substack{\lambda \in i\mathfrak{a}^*_+, \sigma, \tau \in \hat{K} \\ v \in L^2(B)\sigma, w \in L^2(B)\tau}} \frac{|D\langle A(\lambda)v, w \rangle|}{\|v\| \cdot \|w\|} (1+|\lambda|)^n (1+|\tau|)^n (1+|\sigma|)^n < \infty \}.$$

3.2. The most-continuous Hardy space. We recall now the spectral definition of the Hardy space $\mathcal{H}^2(D)$ from [11]. For $v \in \mathcal{H}_{\lambda}$ define an analytic function $f_{v,\lambda}$ on X by

$$f_{v,\lambda}(x) = \langle \pi_{\lambda}(x^{-1})v, v_{K,\lambda} \rangle = \langle v, \pi_{\lambda}(x)v_{K,\lambda} \rangle$$

Let us denote by $z \mapsto \overline{z}$ the complex conjugation in $G_{\mathbb{C}}$ with respect to G and recall that $f_{v,\lambda}$ extends to a holomorphic function $\tilde{f}_{v,\lambda}$ on D via

$$\hat{f}_{v,\lambda}(x) = \langle v, \pi_{\lambda}(\bar{x})v_{K,\lambda} \rangle$$

for $x \in D$, cf. [11] and [12], Proposition 2.2.3. In particular

$$f_{v,\lambda}(ga \cdot x_o) = \langle \pi_{\lambda}(g^{-1})v, \pi_{\lambda}(a^{-1})v_{K,\lambda} \rangle$$

for $g \in G$ and $a \in \exp(i\Omega_H)$

Define a generalized hyperbolic cosine function on $i\mathfrak{a}^*$ by

(3.5)
$$\cosh(\lambda) = \sum_{w \in \mathcal{W}/\mathcal{W}_H} z_H^{2w^{-1}\lambda}$$

for $\lambda \in i\mathfrak{a}^*$. Define a measure μ on $i\mathfrak{a}^*_+$ by

(3.6)
$$d\mu(\lambda) = \frac{d\lambda}{\cosh(\lambda) \cdot |\mathbf{c}(\lambda)|^2}.$$

With this preparation we can define the unitary representation $(L, \mathcal{H}^2(D))$ of G by

$$(L, \mathcal{H}^2(D)) = \int_{i\mathfrak{a}^*_+}^{\oplus} (\pi_\lambda, \mathcal{H}_\lambda) \, d\mu(\lambda) \, .$$

Thus $\mathcal{H}^2(D)$ is the Hilbert space of all measurable functions $s : i\mathfrak{a}^*_+ \to L^2(M \setminus K)$ such that $\|s\|^2 = \int_{i\mathfrak{a}^*_+} \|s(\lambda)\|^2 d\mu(\lambda) < \infty$. In the sequel we often write write s_λ for $s(\lambda)$. Let us denote by $\|\cdot\|_H$ the norm on $\mathcal{H}^2(D)$. Recall from [11] that the map

$$\Phi: \mathcal{H}^2(D) \hookrightarrow \mathcal{O}(D), \quad s = (s_\lambda) \mapsto \left(x \mapsto \int_{i\mathfrak{a}^*_+} \tilde{f}_{s_\lambda,\lambda}(x) \, d\mu(\lambda) \right)$$

is a G-equivariant continuous injection. In the sequel we often view $\mathcal{H}^2(D)$ as a subspace of $\mathcal{O}(D)$; we call $\mathcal{H}^2(D)$ the most-continuous Hardy space of Y. This notion is motivated by the main result of [11] which states that there exists an isometric G-equivariant value mapping

$$b: \mathcal{H}^2(D) \to L^2_{\mathrm{mc}}(Y), \quad f \mapsto b(f)$$

which is isometric onto a multiplicity free subspace of $L^2_{\rm mc}(Y)$.

3.3. The Fourier Transform on X and the Hardy space. The definition of $\mathcal{H}^2(D)$ in the previous subsection does not use the Fourier transform on X. But the following Lemma shows that the space $\mathcal{H}^2(D)$ has a natural description in terms of the Fourier transform.

Lemma 3.2. Let $f \in \mathcal{H}^2(D)$. Then the following assertions hold:

(1) $f|_X \in L^2(X)$ and

$$\begin{split} f(z) &= \int_{i\mathfrak{a}^*_+} \widehat{f|_X}(b,\lambda) \mathbf{cosh}(\lambda) \, a(bz)^{\lambda+\rho} \, d\mu(b,\lambda) \qquad (z \in D) \\ \|f\|_H^2 &= \int_{\mathcal{X}} |\widehat{f|_X}(b,\lambda)|^2 \cdot \mathbf{cosh}(\lambda) \, d\mu_{\mathcal{X}}(b,\lambda) \geq \|f|_X\|_{L^2(X)}^2 \, . \end{split}$$

(2) If $f = \Phi^{-1}(s_{\lambda})$, then

$$(\widehat{f|_X})_{\lambda} = \frac{s_{\lambda}}{\cosh(\lambda)}.$$

(3) For $a = \exp(iY) \in \exp(i\Omega_H)$ let $f_a : G \to \mathbb{C}$ by $f_a(g) = f(ga \cdot x_o)$. Let $Q \subset \exp(i\Omega_H)$ be compact and such that $a \in Q$. Then there exists a constant $C_Q > 0$ such that

$$||f_a||_{L^2(G)} \le C_Q ||f||_H$$

Proof. (1) and (2). Let $f \in \mathcal{H}^2(D)$ and $f = \int_{i\mathfrak{a}^*_+} \tilde{f}_{s_{\lambda},\lambda} d\mu(\lambda)$. Then obviously we have (2), i.e.,

(3.7)
$$f = \int_{i\mathfrak{a}_{+}^{*}} \frac{\tilde{f}_{s_{\lambda},\lambda}}{\cosh(\lambda)} \frac{d\lambda}{|\mathbf{c}(\lambda)|^{2}}$$

and

$$\begin{split} \|f\|_{H}^{2} &= \int_{i\mathfrak{a}_{+}^{*}} \|s_{\lambda}\|_{L^{2}(B)}^{2} d\mu(\lambda) \\ &= \int_{i\mathfrak{a}_{+}^{*}} \left\|\frac{s_{\lambda}}{\cosh(\lambda)}\right\|_{L^{2}(B)}^{2} \cosh(\lambda) \frac{d\lambda}{|\mathbf{c}(\lambda)|^{2}} \\ &\geq \int_{i\mathfrak{a}_{+}^{*}} \left\|\frac{s_{\lambda}}{\cosh(\lambda)}\right\|_{L^{2}(B)}^{2} \frac{d\lambda}{|\mathbf{c}(\lambda)|^{2}} \\ &= \|f|_{X}\|_{L^{2}(X)}^{2} \,. \end{split}$$

Thus $f|_X \in L^2(X)$ and we can write $f|_X = \int_{\mathcal{X}} \widehat{f|_X}(b,\lambda) a(b \cdot)^{\rho+\lambda} db \frac{d\lambda}{|\mathbf{c}(\lambda)|^2}$. Equation (3.7) implies that $\widehat{f|_X}(b,\lambda) = s_{\lambda}(b)/\cosh(\lambda)$ or $s_{\lambda}(b) = \widehat{f|_X}(b,\lambda)\cosh(\lambda)$ for almost all λ . This finishes the proof of (1) and (2).

(3) We recall Faraut's version of the Gutzmer identity [6]

(3.8)
$$\int_{G} |f(ga \cdot x_{o})|^{2} dg = \int_{\mathcal{X}} |\widehat{f|_{X}}(b,\lambda)|^{2} \varphi_{\lambda}(a^{2}) db \frac{d\lambda}{|\mathbf{c}(\lambda)|^{2}}$$

where $\varphi_{\lambda}(a^2)$ is the analytically continued spherical function given by

$$\varphi_{\lambda}(a^2) = \int_K \left| a(ka)^{\rho+\lambda} \right|^2 dk$$

(cf. [19], Sect. 4). Now for a compact subset $Q \subset \exp(i\Omega_H)$ there exists a constant $C_Q > 0$ such that

(3.9)
$$(\forall \lambda \in i\mathfrak{a}^*) \qquad \varphi_{\lambda}(a^2) \le C_Q \mathbf{cosh}(\lambda)$$

for all λ (cf. [17], Lemma 2.1) and the assertion of the lemma follows.

In order to define the Radon transform for functions in the Hardy space we first need a technical fact, interesting on its own.

Let $\left(\int_{i\mathfrak{a}_{+}^{*}}^{\oplus} \mathcal{H}_{\lambda} d\mu(\lambda)\right)_{0}$ denote the space of all sections (s_{λ}) such that for all $v \in L^{2}(B)_{K}$ -finite $i\mathfrak{a}_{+}^{*} \ni \lambda \mapsto \langle s_{\lambda}, v \rangle \in \mathcal{S}(i\mathfrak{a}_{+}^{*})$

(3.10)
$$(\forall D \in \mathbb{D}(i\mathfrak{a}^*), \forall n \in \mathbb{N}_0) \sup_{\substack{\lambda \in i\mathfrak{a}^*_+, \tau \in \hat{K} \\ v \in L^2(B)_\tau}} \frac{|D\langle s_\lambda, v\rangle|}{\|v\|} (1+|\lambda|)^n (1+|\tau|)^n < \infty.$$

Then we set

(3.11)
$$\mathcal{H}^2(D)_0 = \Phi^{-1}\left(\left(\int_{i\mathfrak{a}^*_+}^{\oplus} \mathcal{H}_\lambda \, d\mu(\lambda)\right)_0\right) \,.$$

Theorem 3.3. Let $f \in \mathcal{H}^2(D)_0$. Fix $z \in T(\Omega_H)$. Then the function

 $G \ni g \mapsto f(gz \cdot x_o) \in \mathbb{C}$

belongs to $\mathcal{C}(G)$. Moreover, the following functions are locally bounded on $T(\Omega_H)$:

(1)
$$z \to \int_G |f(gz \cdot x_o)|^2 dg$$

(2) $z \to \int_N |f(nz \cdot x_o)| dn$

Proof. Without loss of generality we may assume that $z = a \in \exp(i\Omega_H)$. In the sequel we often identify f with $f|_X$, a right K-invariant function on G. Let $v, w \in L^2(B)_{K-\text{finite}}$. Then, by (3.4) and Lemma 3.2:

$$\begin{aligned} \langle \mathcal{F}(R_a f)(\lambda) v, w \rangle &= \langle \mathcal{F}(f)(\lambda) \pi_{\lambda}(a) v, w \rangle \\ &= \langle \pi_{\lambda}(a) v, v_{K,\lambda} \rangle \langle \hat{f}_{\lambda}, w \rangle \\ &= \frac{\langle v, \pi_{\lambda}(a) v_{K,\lambda} \rangle}{\cosh(\lambda)} \langle s_{\lambda}, w \rangle \,. \end{aligned}$$

Let us write $F(\lambda, v, w) := \langle \mathcal{F}(R_a f)(\lambda)v, w \rangle$. It is clear that F is smooth in the λ -variable. In order to show that $g \mapsto f(ga \cdot x_0)$ belongs to $\mathcal{C}(G)$ we use Arthur's Theorem 3.3: we have to show for all $n \in \mathbb{N}_0$ and $D \in \mathbb{D}(\mathfrak{ia}^*)$ that

(3.12)
$$\sup_{\substack{\lambda \in ia_{+}^{*}, \sigma, \tau \in \hat{K} \\ v \in L^{2}(B)_{\tau}, w \in L^{2}(B)_{\sigma}}} \frac{|DF(\lambda, v, w)|}{\|v\| \cdot \|w\|} (1 + |\lambda|)^{n} (1 + |\tau|)^{n} (1 + |\sigma|)^{n} < \infty$$

We use the expression for $F(\lambda, v, w)$ derived above to compute its derivatives: For fixed D, we find according to Leibniz a finite set of $D_1, D'_1, \ldots, D_N, D'_N$ in $\mathbb{D}(i\mathfrak{a}^*)$ such that

(3.13)
$$DF(\lambda, v, w) = \sum_{j=1}^{N} \left[D_j \frac{\langle v, \pi_\lambda(a) v_{K,\lambda} \rangle}{\cosh(\lambda)} \right] \cdot \left[D'_j \langle s_\lambda, w \rangle \right]$$

According to (3.10), for all $D \in \mathbb{D}(i\mathfrak{a}^*)$, $n \in \mathbb{N}_0$ the following estimate holds:

(3.14)
$$\sup_{\substack{\lambda \in ia^*_+, \sigma \in \hat{K} \\ w \in L^2(B)_{\sigma}}} \frac{|D\langle s_{\lambda}, w \rangle|}{\|w\|} (1+|\lambda|)^n (1+|\sigma|)^n < \infty$$

Now, after combining (3.12)-(3.14) we are left to show for all $D \in \mathbb{D}(i\mathfrak{a}^*)$, $n \in \mathbb{N}_0$:

(3.15)
$$\sup_{\substack{\lambda \in i\mathfrak{a}^*_+, \tau \in \hat{K} \\ v \in L^2(B)_{\tau}}} \frac{\left| D \frac{\langle v, \pi_\lambda(a) v_{K,\lambda} \rangle}{\cosh(\lambda)} \right|}{\|v\|} (1 + |\tau|)^n < \infty.$$

To establish (3.15), we first note that $\pi_{\lambda}(a)v_{K,\lambda}$ is an analytic vector for the representation. Now if $w \in \mathcal{H}^{\omega}_{\lambda} = C^{\omega}(M \setminus K)$ and $w = \sum_{\tau \in \hat{K}} w_{\tau}$ is its expansion in K-types, then we recall from [20], Th. 2.2 (1), that there exists a $\delta > 0$ such that

$$\|w_{\tau}\| << e^{-\delta|\tau|}$$

To be more precise, given $\delta > 0$ sufficiently small, there exists a ball U arround 1 in $K_{\mathbb{C}}$ such that $\pi_{\lambda}(t)w$ exists for all $t \in U$ and

$$\|w_{\tau}\| \leq \left(\sup_{t \in U} \|\pi_{\lambda}(t)w\|\right) \cdot e^{-\delta/2|\tau|}.$$

Coming back to our original situation where our analytic vector is $w = \pi_{\lambda}(a)v_{K_{\lambda}}$, we obtain that $\pi_{\lambda}(t)\pi_{\lambda}(a)v_{K_{\lambda}}$ exists for all $t \in U$. It follows that $\pi(GUaK_{\mathbb{C}})v_{K_{\lambda}}$ exists. As the crown is a domain of holomorphy for principal series representations (cf. [16]), we obtain, by continuity, a compact neighborhood Q of a in $\exp(i\Omega_H)$ such that $UaK_{\mathbb{C}} \subset GQK_{\mathbb{C}}$. By the very nature of the principal series, it is easy to see that U can be chosen independent from λ . Hence we get a C > 0 such that for all $\lambda \in i\mathfrak{a}^*, \tau \in \hat{K}$ and $v \in L^2(B)_{\tau}$

(3.16)
$$\frac{|\langle v, \pi_{\lambda}(a)v_{K,\lambda}\rangle|}{\|v\|} \le C(1+|\tau|)^{-n} \cdot \sup_{b \in Q} \|\pi_{\lambda}(b)v_{K,\lambda}\|$$

Moreover, $\sup_{b \in Q} \frac{\|\pi_{\lambda}(b)v_{K,\lambda}\|}{\sqrt{\cosh(\lambda)}}$ is uniformly bounded in λ by (3.9). This, in combination with (3.16), proves (3.15) for the case of $D = \mathbf{1}$. Now, if D is of order k, we first observe that

$$(\forall k \in K)(\forall b \in T(\Omega_H)) \quad [D\pi_{\lambda}(b)v_{K,\lambda}](Mk) = p_k((\lambda + \rho)(\log a(kb))(\pi_{\lambda}(b)v_{K,\lambda})(Mk))$$

for a polynomial p_k of degree k and independent from λ . It follows from the complex convexity theorem (1.4) that

$$\sup_{b \in Q} \sup_{k \in K} |p_k(\lambda + \rho)(\log a(kb))| \le C \cdot \sqrt{\cosh(\lambda)}$$

Furthermore the decay of $\frac{1}{\cosh(\lambda)}$ is not affected by differentiation. Therefore we obtain (3.15) for all D.

Moving on to (1) and (2), we observe that statement (1) is Lemma 3.2, part 3; part (2) follows from the already established fact in conjunction with Lemma 22 in [15]. \Box

3.4. The definition of the Radon Transform. Denote by $CR(\Xi_+)$ the vector space of continuous CR-functions on Ξ_+ , i.e. the space of continuous functions on $\Xi_+ \simeq K/M \times T(\Omega_H)$ (cf. (1.14)) which are holomorphic in the second variable.

Lemma 3.4. Let $f \in \mathcal{H}^2(D)_0$. Then the assignment

$$\Xi_+ \ni \xi = ga \cdot \xi_o \mapsto a^{-2\rho} \int_N f(gna \cdot x_o) \ dn \in \mathbb{C} \qquad (g \in G, a \in \exp(i\Omega_H))$$

defines a CR-function on Ξ_+ .

Proof. It follows from Theorem 3.3 that the right hand side is a continuous function. It remains to show that is a CR-function. For that let g = kb for $k \in K$ and $b \in A$. The right hand side becomes

$$a^{-2\rho} \int_{N} f(kbna \cdot x_{o}) \, dn = (ab)^{-2\rho} \int_{N} f(knba \cdot x_{o}) \, dn$$

and the holomorphicity in ab follows with Theorem 3.3.

In view of this lemma, the prescription

$$\mathcal{R}: \mathcal{H}^2(D)_0 \to CR(\Xi_+), \quad f \mapsto \left(\xi = ga \cdot \xi_o \mapsto a^{-2\rho} \int_N f(gna \cdot x_o) \ dn\right)$$

is a well defined and continuous G-equivariant map. We call \mathcal{R} the holomorphic horospherical Radon transform.

Remark 3.5. (a) We recall the real Radon transform on X from (3.1).Now if $f \in \mathcal{H}^2(D)_0$, then $f|_X \in L^2(X)$ by Lemma 3.4. Thus, for $g \in G$, we get

$$\mathcal{R}(f)(g \cdot \xi_o) = \mathcal{R}_{\mathbb{R}}(f|_X)(g \cdot \xi_o) \,.$$

In other words, the holomorphic Radon transform restricted to $\Xi_{\mathbb{R}}$ agrees with the real Radon transform of the restricted function on X.

(b) Notice that $E(\xi) \cap D$ for $\xi \in \Xi_+$ contains the real horosphere $E_{\mathbb{R}}(\xi)$. The holomorphic Radon transform \mathcal{R} then writes as

$$\mathcal{R}(f)(\xi) = \int_{E_{\mathbb{R}}(\xi)} f(\xi') \, d\nu_{\xi}(\xi')$$

with $d\nu_{\xi}$ equals $a^{-2\rho}$ -times the measure on $E_{\mathbb{R}}(\xi)$ obtained by the natural identification of the real horosphere $E_{\mathbb{R}}(\xi)$ with N. It is clear that any other N-orbit in $E(\xi) \cap D$ would yield the same result.

If the function $f \in \mathcal{H}^2(D)_0$ is left K-invariant, then we can define the holomorphic Abeltransform by:

$$\mathcal{A}(f)(z) = z^{-\rho} \int_{N} f(nz \cdot x_{o}) \, dn \qquad (z \in T(\Omega_{H})) \, .$$

Note that \mathcal{A} is just the restriction of the holomorphic Radon transform to K-invariant function (modulo the $z^{-\rho}$ -factor). Further let us remark that \mathcal{A} gives a continuous mapping

$$\mathcal{A}: \mathcal{H}^2(D)_0^K \to \mathcal{O}(T(\Omega_H))^{\mathcal{W}}, \quad \mathcal{A}(f)(z) = z^{-\rho} \int_N f(nz \cdot x_o) \ dn$$

4. The holomorphic Radon transform as Cauchy integral I: The hyperboloid

In this and the next section we will show (for an appropriate class of Y's) that the holomorphic Radon transform on the NCC-space Y can be expressed as a Cauchy type integral. For that purpose it is instructive to explaining the example of the hyperboloid first. For earlier treatments of the hyperboloid with alternative methods we refer to [7], [8]. We start by recalling some standard function spaces on Y.

4.1. Function spaces. Let Y be a NCC-space. For $g \in G$ let $\Theta(g) = \varphi_0(g\tau(g)^{-1})^{1/2}$. Then Θ is left K-invariant and right H-invariant. For $g = k \exp(Z)h$ with $Z \in \mathfrak{a}$ define $\|g \cdot y_0\| := \|Z\|$, where $\|Z\| = \sqrt{\operatorname{Tr} \operatorname{ad}(Z)^2}$. Let $D \in U(\mathfrak{g})$, were $U(\mathfrak{g})$ is the enveloping algebra of $\mathfrak{g}_{\mathbb{C}}$, and view L_D as a differential operator on Y. For $n \in \mathbb{N}$ and $f \in C^{\infty}(Y)$ define

$$p_{n,D}(f) = \sup_{y \in Y} \Theta_G(y)^{-1} (1 + ||y||)^n |L_D f(y)|.$$

Then the Schwartz space $\mathcal{C}(Y)$ is defined as the space of smooth function on Y such that $p_{n,D}(f) < \infty$ for all n and D. It is well known, that $\mathcal{C}(Y) \subset L^2(Y)$, but $\mathcal{C}(Y)$ is not contained in $L^1(Y)$. We will therefore need a smaller space to make sure that the Cauchy integral exists. For that, define for r > 0 the space

$$\mathcal{S}_r(Y) := \{ f \in C^{\infty}(Y) \mid (\forall D \in U(\mathfrak{g})) \sup_{y \in Y} e^{r ||y||} |L_D f(y)| < \infty \}$$

and

$$\mathcal{S}(Y) := \bigcap_{r>0} \mathcal{S}_r(Y) \,.$$

Then $\mathcal{S}(Y) \subset L^1(Y) \cap L^2(Y)$ and $C_c^{\infty}(Y) \subset \mathcal{S}(Y) \subset \mathcal{C}(Y)$. The space $\mathcal{S}(Y)$ is called the (zero) Schwartz space, cf. [5]. It follows from Theorem 3 in [5] and our spectral definition of the space $\mathcal{H}^2(D)$ that

$$\mathcal{H}^2(D)_{00} := \{ f \in \mathcal{H}^2(D)_0 \mid f \text{ K-finite, } b(f) \in \mathcal{S}(Y) \}$$

is dense in $\mathcal{H}^2(D)$. Here used the fact that elements in $\mathcal{H}^2(D)_0$ have boundary values on Y (cf. [11], Sect. 3 for a similar argument). In particular we have that every element $f \in \mathcal{H}^2(D)_{00}$ is integrable on Y, has a holomorphic extension to D, and that the integral $\int_N f(anw \cdot y_0) dn$ is well defined for all $a \in A$ and $w \in \mathcal{W}$. We will use this without comments in the following.

4.2. The Radon transform and Cauchy integral on the hyperboloid. Assume that $n \geq 2$ and let $G = SO_e(1, n)$ be the Lorentz group. Let us fix our choices for the groups A, N and K. For the maximal compact subgroup we take

$$K = \left\{ k_R = \begin{pmatrix} 1 & 0 \\ 0 & R \end{pmatrix} \mid R \in \mathrm{SO}(n) \right\} \simeq \mathrm{SO}(n) \,.$$

Next, for $z \in \mathbb{C}$ we set

$$a_{z} = \begin{pmatrix} \cosh z & 0 & \sinh z \\ 0 & \mathbf{1}_{n-1} & 0 \\ \sinh z & 0 & \cosh z \end{pmatrix}$$

and

$$A = \{a_t \mid t \in \mathbb{R}\} \quad \text{and} \quad A_{\mathbb{C}} = \{a_z \mid z \in \mathbb{C}\}.$$

Note that $\mathfrak{a} = \mathbb{R}Z$ and $a_z = \exp(zZ)$ with $Z = E_{1n+1} + E_{n+11}$. The only positive root is α , determined by $\alpha(Z) = 1$ and hence $Z_H = \frac{\pi}{2}Z$. We also have, that $\rho = \frac{n-1}{2}\alpha$. Further, for $v \in \mathbb{C}^{n-1}$ and $(v, v) = \sum_{j=1} v_j v_j$ we define an unipotent matrix

$$n_v = \begin{pmatrix} 1 + \frac{1}{2}(v, v) & v^T & -\frac{1}{2}(v, v) \\ v & \mathbf{1}_{n-1} & -v \\ \frac{1}{2}(v, v) & v^T & 1 - \frac{1}{2}(v, v) \end{pmatrix}$$

Then N and $N_{\mathbb{C}}$ are given by

$$N = \{n_v \mid v \in \mathbb{R}^{n-1}\} \quad \text{and} \quad N_{\mathbb{C}} = \{n_v \mid v \in \mathbb{C}^{n-1}\}.$$

Define a quadratic form

$$z, w \mapsto z \cdot w = z_0 w_0 - \sum_{j=1}^n z_j w_j$$

on \mathbb{C}^{n+1} and let

$$\Box(\mathbf{z}) = z_0^2 - z_1^2 - \ldots - z_n^2 \qquad (\mathbf{z} = (z_0, \ldots, z_n)^T \in \mathbb{C}^{n+1})$$

be the corresponding quadratic form. We define the real and complex hyperboloids by

$$X = \{ \mathbf{x} \in \mathbb{R}^{n+1} \mid \Box(\mathbf{x}) = 1, x_0 > 0 \}$$

and

$$X_{\mathbb{C}} = \{ \mathbf{z} \in \mathbb{C}^{n+1} \mid \Box(\mathbf{z}) = 1 \}$$

As a common base point for X and $X_{\mathbb{C}}$ we take $\mathbf{x}_o = (1, 0, \dots, 0)^T$ and note that the map

$$G_{\mathbb{C}}/K_{\mathbb{C}} \to X_{\mathbb{C}}, \ gK_{\mathbb{C}} \mapsto g(\mathbf{x}_o)$$

is a $G_{\mathbb{C}}$ -isomorphism which identifies G/K with X.

We have that $a_z \cdot \mathbf{x}_o = (\cosh(z), 0, \dots, 0, \sinh(z))^T$ and hence $\mathbf{y}_0 = z_H \cdot \mathbf{x}_o = (0, \dots, 0, i)^T$. It is clear that the stabilizer of \mathbf{y}_o is

$$H = \left\{ \begin{pmatrix} h & 0\\ 0 & 1 \end{pmatrix} \mid h \in \mathrm{SO}_e(1, n-1) \right\} \simeq \mathrm{SO}_e(1, n-1) \,.$$

We have therefore with this identification:

$$D = \{ \mathbf{z} = \mathbf{x} + i\mathbf{y} \in X_{\mathbb{C}} \mid \Box(\mathbf{x}) > 0, x_0 > 0 \}$$

and

$$Y = G(\mathbf{y}_0) = \{ i \mathbf{y} \in i \mathbb{R}^{n+1} \mid \Box(\mathbf{y}) = -1 \}.$$

 Set

$$\Xi = \{ \zeta \in \mathbb{C}^{n+1} \mid \zeta \neq 0, \Box(\zeta) = 0 \}$$

If $\xi_o = (1, 0, \dots, 0, 1)^T \in \Xi$, then the stabilizer of ξ_o is $M_{\mathbb{C}}N_{\mathbb{C}}$ and the map

$$G_{\mathbb{C}}/M_{\mathbb{C}}N_{\mathbb{C}} \to \Xi, \quad gM_{\mathbb{C}}N_{\mathbb{C}} \mapsto g \cdot \xi_o$$

is a $G_{\mathbb{C}}$ -isomorphism. Now the *CR*-submanifold $\Xi_+ \subset \Xi$ is described as

$$\Xi_+ = G\left\{ (e^{it}, 0, \dots, 0, e^{it})^T \mid |t| < \frac{\pi}{2}, t \in \mathbb{R} \right\}$$

= $\{ \zeta = \xi + i\eta \in \Xi : \Box(\xi) = \Box(\eta) = 0; \ \xi \neq 0 \}_0 .$

We will also use certain G-subdomains of Ξ_+ : for $0 < c \leq \frac{\pi}{2}$ set

$$\Xi_c = G\left\{ (e^{it}, 0, \dots, 0, e^{it})^T \mid |t| < c, t \in \mathbb{R} \right\} \,.$$

In order to define the Cauchy-transform we need to establish a simple, but important, technical fact.

Lemma 4.1. For all $\xi \in \Xi_+$ and $y \in Y$ one has

$$\xi \cdot y
ot\in \mathbb{R}^{ imes}$$
 .

More precisely, for all $0 < c < \frac{\pi}{2}$ there exists a C > 0 such that

$$(\forall y \in Y)(\forall \xi \in \Xi_c)$$
 $|1 - \xi \cdot y| > C.$

Proof. By *G*-equivariance of the form we may assume that $\xi = e^{it}\xi_o$. Let $y = i\mathbf{y}$ for $\mathbf{y} \in \mathbb{R}^{n+1}$. Thus $\xi \cdot y = ie^{it}\xi_o \cdot \mathbf{y}$ with $\xi_o \cdot \mathbf{y} \in \mathbb{R}$. As $|t| < \frac{\pi}{2}$, the assertions follow. We now define the Cauchy-kernel function

$$\mathcal{K}: \Xi_+ \times Y \to \mathbb{C}, \ (\xi, y) \mapsto \frac{1}{1 - \xi \cdot y}.$$

In view of the previous lemma, this function is defined, continuous and bounded on all subsets $\Xi_c \times Y$ for $c < \frac{\pi}{2}$. Moreover, \mathcal{K} is a *CR*-function in the first variable and *G*-invariant, i.e., $\mathcal{K}(g \cdot \xi, g \cdot y) = \mathcal{K}(\xi, y)$. In particular, the function

$$G \ni g \mapsto \mathcal{K}(g) = \mathcal{K}(\xi_o, g \cdot \mathbf{y}_0) \in \mathbb{C}$$

is left N-invariant and right H-invariant, a fact that we will use in a moment. We will therefore identify \mathcal{K} with a function on Y without further comments. A simple calculation shows that

(4.1)
$$\mathcal{K}(g) = \frac{1}{1 - i(g_{0\,n} - g_{n\,n})}$$

We have $\mathcal{W}_H = \{\mathbf{1}\}$, and $\mathcal{W} = \{\mathbf{1}, \epsilon\}$ where $\epsilon = -1$ on \mathfrak{a} . As ϵ corresponds to the matrix $w = \begin{pmatrix} I_{n-1} & 0 \\ 0 & -I_2 \end{pmatrix}$ it follows that

(4.2)
$$\mathcal{K}(a_z) = \frac{1}{1 - ie^{-z}}$$
 and $\mathcal{K}(a_z w) = \frac{1}{1 + ie^{-z}}$.

Write $\mathcal{S}(Y)$ for the Schwartz-space on Y. Henceforth we will make the assumption that n is even and define the *Cauchy* - *transform* by

$$\mathcal{C}: \mathcal{S}(Y) \to CR(\Xi_+), \quad \mathcal{C}(f)(\xi) = \int_Y \frac{f(y)}{1 - \xi \cdot y} \, dy = \int_Y f(y) K(\xi, y) \, dy$$

where dy denotes the *G*-invariant measure on *Y* from Subsection 1.6.

Remark 4.2. In the following we have to raise complex powers of elements $z \in A_{\mathbb{C}}$ which are in the boundary of $\exp(i\Omega_H)$. This is not a problem as the map

$$\exp: 2\Omega_H \to A_{\mathbb{C}}, \quad Y \mapsto \exp(iY)$$

is injective; a fact which holds in full generality.

Theorem 4.3. Let G = SO(n, 1) with n = 2k even. Let $f \in \mathcal{H}^2(D)_{00}$. Then, up to normalization of measures on A, one has

$$\mathcal{C}(f)(\xi) = (-1)^{k-1} 2\pi \cdot \mathcal{R}(f)(\xi) \qquad (\xi \in \Xi_+) \,.$$

Proof. Since both C(f) and $\mathcal{R}(f)$ are CR-functions, it is sufficient to show that both coincide on $G/MN \subset \Xi$. Moreover, by G-equivariance of both maps, it is in fact sufficient to show that

(4.3)
$$\mathcal{C}(f)(\xi_o) = (-1)^{k-1} 2\pi \cdot \mathcal{R}(f)(\xi_o) + \mathcal{R}(f)(\xi_o)$$

Using (4.2) and that $z_{H}^{2\rho} = (-1)^{k-1}i$ and $z_{H}^{-2\rho} = \overline{z_{H}^{2\rho}} = (-1)^{k}i$ we get:

$$\begin{split} \mathcal{C}(f)(\xi_o) &= \int_Y f(y) \cdot \mathcal{K}(y) \, dy \\ &= \sum_{w \in \mathcal{W}} \int_A \int_N f(anw \cdot \mathbf{y}_o) \cdot \mathcal{K}(anw) \, dn \, da \\ &= \sum_{w \in \mathcal{W}} \int_A \int_N f(anw \cdot \mathbf{y}_o) \cdot \mathcal{K}(aw) \, dn \, da \\ &= \sum_{w \in \mathcal{W}} \int_A \int_N f(anw z_H \cdot \mathbf{x}_o) \cdot \mathcal{K}(aw) \, dn \, da \\ &= \sum_{w \in \mathcal{W}} \int_A \int_N f(an z_H^w \cdot \mathbf{x}_o) \cdot \mathcal{K}(aw) \, dn \, da \\ &= \sum_{w \in \mathcal{W}} \int_A \int_N f(an z_H^w \cdot \mathbf{x}_o) \cdot (z_H^w)^{-2\rho} \cdot (z_H^w)^{2\rho} \cdot \mathcal{K}(aw) \, dn \, da \\ &= \sum_{w \in \mathcal{W}} \int_A \mathcal{R}(f)(a z_H^w \cdot \xi_o) \cdot (z_H^w)^{2\rho} \cdot \mathcal{K}(aw) \, dn \, da \\ &= \sum_{w \in \mathcal{W}} \int_A \mathcal{R}(f)(a z_H^w \cdot \xi_o) \cdot (z_H^w)^{2\rho} \cdot \mathcal{K}(aw) \, da \end{split}$$

Consider the strip domain $S = \{z \in \mathbb{C} \mid |\operatorname{Im} z| \leq \frac{\pi}{2}\}$. By our assumption on f, the function

$$S \ni z \mapsto F(z) = \frac{i\mathcal{R}(f)(a_z \cdot \xi_o)}{1 - e^{-z}} \in \mathbb{C}$$

defines a meromorphic function F on int S which extends continuously up to the boundary and which has at most a simple pole at z = 0. Thus the Residue theorem yields that

$$\mathcal{C}(f)(\xi_o) = (-1)^k 2\pi i \cdot \operatorname{Res}(F, 0) = (-1)^{k-1} 2\pi \cdot \mathcal{R}(f)(\xi_o)$$

and this concludes the proof of our theorem.

Remark 4.4. (a) We mention that the geometric pairing $\xi \cdot y$ can be expressed using the previously defined power functions f_j (cf. 1.10):

$$\xi \cdot y = f_1(\xi^{-1}y) \,.$$

(b) The assumption that n is even is not a real restriction as one can slightly modify C so that it works for all parities (see our discussion in the next section).

5. The holomorphic Radon transform as Cauchy integral II: Cayley type spaces

The technique used for the hyperboloid in the previous can be used for NCC-spaces of Cayley type as well. Let us recall that Cayley type spaces are those which are associated

to Euclidean Jordan algebras V: i.e. X is a tube domain associated to V and H is the structure group of the cone of squares in V. In terms of the set of restricted roots Σ , this means that Σ is of type C_n , say

$$\Sigma = \left\{ \frac{1}{2} (\pm \gamma_i \pm \gamma_j) \mid 1 \le i, j \le n \right\} \setminus \{0\}.$$

We assume now that Y = G/H is of Cayley type. Define $T_j \in \mathfrak{a}$ by $\gamma_i(T_j) = \delta_{ij}$, then

(5.1)
$$\Omega_H = \bigoplus_{j=1}^n \left[-\frac{\pi}{2}, \frac{\pi}{2} \right[T_j \, .$$

As a basis of Σ we shall choose

$$\Pi = \left\{ \frac{1}{2} (\gamma_1 - \gamma_2), \dots, \frac{1}{2} (\gamma_{n-1} - \gamma_n), \gamma_n \right\} .$$

Obviously $\omega_1 = -\gamma_1$ is a fundamental spherical lowest weight and accordingly $f_1(g) = \langle \pi_1(g)\eta_j, v_1 \rangle$ defines a holomorphic function on $G_{\mathbb{C}}$ with $f_1(g) = a(g)^{-\gamma_1}$ for $g \in N_{\mathbb{C}}A_{\mathbb{C}}K_{\mathbb{C}}$. The analogue of Lemma 4.1 now reads as follows.

Lemma 5.1. Let the notation be as above. Then the following holds:

- (1) $f_1(\exp(i\Omega_H)Gz_H) \subseteq \mathbb{C} \setminus \mathbb{R}^{\times}$.
- (2) For all $0 \le c < 1$ there exists C > 0 such that

$$(\forall g \in G)(\forall Z \in \Omega_H) \quad |1 - f_1(\exp(icZ)gz_H)| > C.$$

Proof. First it is clear from (5.1) that

(5.2)
$$\exp(i\Omega_H)^{\omega_1} = \{ z \in \mathbb{C} \mid \operatorname{Re} z > 0 \}$$

Next recall that $\bigcup_{w\in\mathcal{W}} NAwH$ is dense in G and that

(5.3)
$$f_1(nawh) = a^{\omega_1} z_H^{w\omega_1} \in \mathbb{R}^+\{-i, i\}$$

Combine (5.2) and (5.3) and the assertions follow.

For 0 < c < 1 define a G-subdomains of Ξ_+ by

(5.4)
$$\Xi_c = G \exp(ic\Omega_H) \cdot \xi_o.$$

Note that $-\gamma_j$, $j \neq 1$ is not a fundamental spherical lowest weight. Therefore, for $1 \leq j \leq n$ we define a meromorphic function on $G_{\mathbb{C}}$ directly by

$$h_j(g) = a(g)^{-\gamma_j}$$
 for $g \in N_{\mathbb{C}} A_{\mathbb{C}} K_{\mathbb{C}}$.

Note that $h_1 = f_1$ and, in the same manner as in Lemma 5.1, one establishes that

(5.5)
$$h_j(\exp(i\Omega_H)Gz_H) \subseteq \mathbb{C} \setminus \mathbb{R}^{\times} \cup \{\infty\}$$

In particular we see that the prescription

$$\mathcal{K}: \Xi_+ \times Y \to \mathbb{C}, \ (\xi, y) \mapsto \prod_{j=1}^n \frac{1}{1 - h_j(\xi^{-1}y)}$$

defines an analytic function, CR in the first variable and bounded on all subsets $\Xi_c \times Y$.

Remark 5.2. Alternatively, the kernel \mathcal{K} can be expressed in Jordan algebra terms. Let V be the Euclidean Jordan algebra associated to X and $W \subset V$ its cone of squares. Form the tube domains $\mathcal{T}^{\pm} = V \pm iW \subset V_{\mathbb{C}}$. Then $X = \mathcal{T}^+$ and $D \simeq \mathcal{T}^+ \times \mathcal{T}^-$ with X realized in D via the map $x \mapsto (x, \overline{x})$. Write Δ_j for the power functions on $V_{\mathbb{C}}$ (i.e. generalized principal minors). Then, on D, one has

$$h_j(z,w) = \frac{\Delta_j(z-w)}{\Delta_{j-1}(z-w)} \quad \text{for } (z,w) \in D$$

with the understanding that $\Delta_0 \equiv 1$ Thus \mathcal{K} , when considered as a function on D, is given by

$$\mathcal{K}(z,w) = \frac{\Delta_1(z-w) \cdot \ldots \cdot \Delta_{n-1}(z-w)}{\prod_{j=1}^n (\Delta_{j-1}(z-w) - \Delta_j(z-w))}$$

However, \mathcal{K} is not our Cauchy-kernel yet and some small modification is needed. For that recall the decomposition $\Xi_+ \simeq G/MN \times \exp(i\Omega_H)$. For a unitary character $\chi \in \widehat{T/F}$, where $F = K \cap T$ is the canonical finite 2-group as usually, define the space of χ -twisted *CR*-functions by

(5.6)
$$CR_{\chi}(\Xi_{+}) = \{f(ga \cdot \xi_{o}) = h(ga \cdot \xi_{o})\chi^{-1}(a) \mid h \in CR(\Xi_{+})\}.$$

Let $\gamma_0 = \gamma_1 + \ldots + \gamma_n$. In the sequel we will fix χ to be

$$\chi = -2\rho + \gamma_0$$

and notice that $\chi = \mathbf{1}$ for $G = \mathrm{Sl}(2, \mathbb{R})$. For $g \in N_{\mathbb{C}}ATK_{\mathbb{C}}$ write $t(g) \in T/F$ for the compact middle part of g and define the Cauchy-kernel \mathcal{K} by

(5.7)
$$\mathcal{K}_{\chi}(\xi, y) = \mathcal{K}(\xi, y)\chi(t(\xi^{-1}y)).$$

and notice that \mathcal{K}_{χ} is defined whenever $\xi^{-1}y \in N_{\mathbb{C}}A_{\mathbb{C}}K_{\mathbb{C}}$ and, when defined, is in CR_{χ} as a function of the first variable. We note that \mathcal{H}_{χ} is *G*-invariant and hence corresponds to a function of one variable $\mathcal{K}_{\chi}(g) = \mathcal{K}_{\chi}(\xi_o, g \cdot y_0)$. As before, \mathcal{K}_{χ} is $N \times H$ -invariant, and will be identified with left *N*-invariant function on *Y*.

We define the *twisted Cauchy transform* by

$$\mathcal{C}_{\chi}: \mathcal{S}(Y) \to CR_{\chi}(\Xi_{+}), \ \ \mathcal{C}_{\chi}(f)(\xi) = \int_{Y} f(y)\mathcal{K}_{\chi}(\xi, y) \ dy$$

and the twisted holomorphic Radon transform by

$$\mathcal{R}_{\chi} : \mathcal{H}^2(D)_{00} \to CR_{\chi}(\Xi_+), \quad f \mapsto \left(\xi = ga \cdot \xi_o \mapsto a^{-\gamma_0} \int_N f(gna \cdot x_o) \ dn\right).$$

We come to the main result of this section.

Theorem 5.3. Suppose that Y = G/H is of Cayley type. Let $f \in \mathcal{H}^2(D)_{00}$. Then

 $\mathcal{C}_{\chi}(f)(\xi) = (-2\pi i)^n \cdot \mathcal{R}_{\chi}(f)(\xi) \qquad (\xi \in \Xi_+) \,.$

Proof. Both $\mathcal{C}_{\chi}(f)$ and $\mathcal{R}_{\chi}(f)$ are *CR*-functions, and so it is sufficient to show that both coincide on $G/MN \subset \Xi$. Next, by *G*-equivariance of both maps, it is enough to show that

(5.8)
$$\mathcal{C}_{\chi}(f)(\xi_o) = (2\pi)^n \cdot \mathcal{R}_{\chi}(f)(\xi_o) \,.$$

We now get, as in the proof of Theorem 4.2, for the left hand side:

$$\begin{aligned} \mathcal{C}_{\chi}(f)(\xi_{o}) &= \int_{Y} f(y) \cdot \mathcal{K}_{\chi}(y) \, dy \\ &= \cdots \\ &= \sum_{w \in \mathcal{W}/\mathcal{W}_{H}} \int_{A} \mathcal{R}_{\chi}(f)(az_{H}^{w} \cdot \xi_{o}) \cdot (z_{H}^{w})^{\gamma_{0}} \cdot \mathcal{K}_{\chi}(az_{H}^{w} \cdot x_{o}) \, da \\ &= \sum_{w \in \mathcal{W}/\mathcal{W}_{H}} \int_{A} \mathcal{R}(az_{H}^{w} \cdot \xi_{o}) \cdot (z_{H}^{w})^{\gamma_{0}} \cdot \mathcal{K}(az_{H}^{w} \cdot x_{o}) \, da \, . \end{aligned}$$

Specifically we have $\mathcal{W}/\mathcal{W}_H \simeq (\mathbb{Z}_2)^n$ and $z_H^{w\gamma_0} = i^n \operatorname{sgn}(w)$

$$\mathcal{C}_{\chi}(f)(\xi_{o}) = i^{n} \sum_{\varepsilon \in (\mathbb{Z}_{2})^{n}} \operatorname{sgn}(\varepsilon) \int_{\mathbb{R}^{n}} \mathcal{R}(f)(\exp\left(\sum_{j=1}^{n} (t_{j} + i\varepsilon_{j}\frac{\pi}{2})T_{j}\right) \cdot \xi_{o}) \cdot \mathcal{K}(\exp\left(\sum_{j=1}^{n} (t_{j} + i\varepsilon_{j}\frac{\pi}{2})T_{j}\right) \cdot x_{o}) dt$$

Next observe that

$$\mathcal{K}(\exp\left(\sum_{j=1}^{n} (t_j + i\varepsilon_j \frac{\pi}{2})T_j\right) \cdot x_o) = \prod_{j=1}^{n} \frac{1}{1 - e^{-(t_j + i\varepsilon_j \frac{\pi}{2})}}$$

Let us consider the multi strip domain $S = \{z \in \mathbb{C}^n \mid |\operatorname{Im} z_j| \leq \frac{\pi}{2}\}$. By our assumption on f, the prescription

$$S \ni z \mapsto \mathcal{R}(f)(\exp\left(\sum_{j=1}^{n} z_j T_j\right) \cdot \xi_o) \cdot \prod_{j=1}^{n} \frac{1}{1 - e^{-z_j}} \in \mathbb{C}$$

defines a meromorphic function F on int S which extends continuously up to the boundary and with at most a simple multi-pole at $z_j = 0$. Thus iteratively applying the Residue theorem yields that

$$\mathcal{C}_{\chi}(f)(\xi_o) = (-2\pi i)^n \cdot \operatorname{Res}(F, 0) = (-2\pi i)^n \cdot \mathcal{R}(f)(\xi_o) = (-2\pi i)^n \mathcal{R}_{\chi}(f)(\xi_o)$$

and it concludes the proof of our theorem.

6. Some remarks on the inversion of the holomorphic Radon transform

The inversion of the real horospherical transform on X can be analytically continued to give the inversion of the holomorphic horospherical transform. The dual transform is given by integration over the real form $S_{\mathbb{R}}^{K}$ of S(z). However, there is a second non-compact real form $S_{\mathbb{R}}^{H}(z)$ of S(z) which gives rise to a different dual transform and inversion. This is the topic of this section. Mainly we will focus on $G = Sl(2, \mathbb{R})$.

We begin with the definition of an appropriate function space. Let us denote by $\mathcal{F}(\Xi_+)$ denote the space of CR-functions on Ξ_+ which extend continuously to $G \exp(i\overline{\Omega}_H) \cdot \xi_o$ such that $H \ni h \mapsto f(ghz_H \cdot \xi_o) \in \mathbb{C}$ is integrable for all $g \in G$. For those functions we define the dual Radon transform by

$$\mathcal{F}(\Xi_+) \to C(Y), \quad \phi \mapsto \phi^{\vee}$$

with

$$\phi^{\vee}(y) = \int_{H} f(ghz_{H} \cdot \xi_{o}) \ dh = \int_{S_{\mathbb{R}}(y)} f \qquad (y = g \cdot y_{o} \in Y) \,.$$

Clearly, this is a *G*-equivariant mapping.

We would like to understand the relation between \mathcal{R} and $\phi \mapsto \phi^{\vee}$. In this context we would like to mention the result in [13] for the holomorphic discrete series: there exists a differential operator \mathcal{L} such that $(\mathcal{LR}(f))^{\vee} = f$. Hence it is natural to ask whether a similar statement would hold true for the most continuous spectrum considered in this paper. It will turn out that the situation different for the most continuous series in the sense that the inverting operator \mathcal{L} is not a differential operator. We give a detailed discussion for the the basic example.

6.1. The example of $G = Sl(2, \mathbb{R})$. For this paragraph we let $G = Sl(2, \mathbb{R})$ with the usual choices

$$A = \left\{ \begin{pmatrix} t & 0\\ 0 & \frac{1}{t} \end{pmatrix} \mid t > 0 \right\}, \quad N = \left\{ \begin{pmatrix} 1 & x\\ 0 & 1 \end{pmatrix} \mid x \in \mathbb{R} \right\},$$

and $K = \mathrm{SO}(2,\mathbb{R})$. Let $f \in \mathcal{H}^2(D)_{00}$ be a K-invariant function. In the sequel we will identify $\mathfrak{a}_{\mathbb{C}}^*$ with \mathbb{C} via the assignment

$$\mathbb{C} \ni \lambda \mapsto \lambda \cdot \rho \in \mathfrak{a}_{\mathbb{C}}^*.$$

In this coordinates one has

$$\mathbf{c}(\lambda) = \pi^{-1/2} \frac{\Gamma(\lambda/2)}{\Gamma((\lambda+1)/2)}$$
 and $|\mathbf{c}(\lambda)|^{-2} = \frac{i\pi\lambda}{2} \tanh\left(\frac{i\pi\lambda}{2}\right)$

We know that $f|_X \in L^2(X)$ and, as f is K-invariant, we can write

$$f(x) = \frac{1}{2} \int_{\mathbb{R}} \hat{f}(i\lambda) \phi_{i\lambda}(x) \frac{d\lambda}{|\mathbf{c}(i\lambda)|^2} \qquad (x \in X) \,.$$

Applying \mathcal{R} yields that

26

$$\mathcal{R}(f)(\xi) = \frac{1}{2} \int_{\mathbb{R}} \hat{f}(i\lambda) a(\xi^{-1})^{\rho(1+i\lambda)} \, d\lambda \,,$$

and thus

$$\mathcal{R}(f)^{\vee}(y_o) = \frac{1}{2} \int_H \int_{\mathbb{R}} \hat{f}(i\lambda) a(z_H^{-1}h)^{\rho(1+i\lambda)} d\lambda dh.$$

Now, for $h = \begin{pmatrix} \cosh t & \sinh t \\ \sinh t & \cosh t \end{pmatrix} \in H = \mathrm{SO}_e(1,1)$ one has $z_H^{-1}h = \begin{pmatrix} e^{-i\frac{\pi}{4}}\cosh t & e^{-i\frac{\pi}{4}}\sinh t \\ e^{i\frac{\pi}{4}}\sinh t & e^{i\frac{\pi}{4}}\cosh t \end{pmatrix}$ and so

$$a(z_H^{-1}h)^{\rho} = \left(\frac{1}{i(\sinh^2 t + \cosh^2 t)}\right)^{\frac{1}{2}} = e^{-i\frac{\pi}{4}} \cdot \frac{1}{(\cosh 2t)^{\frac{1}{2}}}$$

Therefore we obtain that

(6.1)
$$\mathcal{R}(f)^{\vee}(y_o) = \frac{1}{2} \int_{\mathbb{R}} \int_{\mathbb{R}} \hat{f}(i\lambda) \cdot \frac{e^{\frac{\pi}{4}(\lambda-i)}}{(\cosh 2t)^{\frac{1}{2}(1+i\lambda)}} \, d\lambda \, dt \, .$$

Lemma 6.1.

$$\int_{\mathbb{R}} \frac{1}{(\cosh 2t)^{\frac{1}{2}(1+i\lambda)}} dt = \frac{1}{2} B(1/2, (1+i\lambda)/4) = \frac{1}{2} \frac{\Gamma((1+i\lambda)/4)\Gamma(1/2)}{\Gamma((3+i\lambda)/4)}$$

Proof. Let us denote the integral on the left hand side by $I(\lambda)$. With the substitution $u = \cosh 2t$ we obtain

$$I(\lambda) = \int_{1}^{\infty} \frac{1}{u^{\frac{1}{2}(1+i\lambda)}} \frac{1}{(u^{2}-1)^{1/2}} du$$

$$= \frac{1}{2} \int_{1}^{\infty} v^{-\frac{1+i\lambda}{4}} (v-1)^{-1/2} v^{-1/2} dv \qquad (v=u^{2})$$

$$= \frac{1}{2} B(1/2, (1+i\lambda)/4)$$

$$u^{-(p+q)}(u-1)^{p-1} du$$

as $B(p,q) = \int_1^\infty u^{-(p+q)} (u-1)^{p-1} du.$

Using this, we get:

(6.2)
$$\mathcal{R}(f)^{\vee}(y_o) = \frac{1}{4} \int_{\mathbb{R}} \hat{f}(i\lambda) \cdot e^{\frac{\pi}{4}(\lambda-i)} B(1/2, (1+i\lambda)/4) \ d\lambda.$$

Define

$$C_1(\lambda) = e^{\frac{\pi}{4}(\lambda-i)}B(1/2,(1+i\lambda)/4) + e^{-\frac{\pi}{4}(\lambda+i)}B(1/2,(1-i\lambda)/4)$$

and note that $\lambda \mapsto f(i\lambda)$ is an even function. Thus (6.2) yields that

(6.3)
$$\mathcal{R}(f)^{\vee}(y_o) = \frac{1}{2} \int_{\mathbb{R}} \hat{f}(i\lambda) \cdot C_1(\lambda) \ d\lambda \,.$$

By the Fourier inversion formula, we have

(6.4)
$$f(y_o) = \frac{1}{2} \int_{\mathbb{R}} \hat{f}(i\lambda) \phi_{i\lambda}(y_o) \frac{d\lambda}{|\mathbf{c}(\lambda)|^2}.$$

Now, from the special values of the Gauß hypergeometric function we get

$$\phi_{i\lambda}(y_o) = F(1/4 + i\lambda/4, 1/4 - i\lambda/4, 1; 1) = \frac{1}{2B((3 - i\lambda)/4, (3 + i\lambda)/4)}$$

We therefore define

$$C_{2}(\lambda) = \frac{1/2}{B((3-i\lambda)/4, (3+i\lambda)/4) |\mathbf{c}(i\lambda)|^{2}}$$

and note that (6.4) transforms into

(6.5)
$$f(y_o) = \frac{1}{2} \int_{\mathbb{R}} \hat{f}(i\lambda) \cdot C_2(\lambda) \ d\lambda$$

In the next step we want to compare the expressions $C_1(\lambda)$ and $C_2(\lambda)$. If there would exist a differential operator \mathcal{L} , then there should be a polynomial $g(\lambda)$ such that $g \cdot C_1 = C_2$.

We first consider

$$c_1(\lambda) := C_1(\lambda) \cdot \left(\frac{\Gamma(1/2)}{\Gamma(3/4 - i\lambda/4)\Gamma(3/4 + i\lambda/4)}\right)^{-1} = C_1^+(\lambda) + C_1^-(\lambda)$$

with

$$C_1^{\pm}(\lambda) = e^{\pm \frac{\pi\lambda}{4} - i\frac{\pi}{4}} \cdot \frac{\Gamma((1\pm i\lambda)/4)\Gamma(1/2)}{\Gamma(3/4\pm i\lambda/4)} \cdot \left(\frac{\Gamma(1/2)}{\Gamma(3/4 - i\lambda/4)\Gamma(3/4 + i\lambda/4)}\right)^{-1}.$$

We focus on C_1^+ and obtain

$$C_{1}^{+}(\lambda) = e^{\frac{\pi}{4}(\lambda-i)} \cdot \frac{\Gamma((1+i\lambda)/4)\Gamma(1/2)}{\Gamma(3/4+i\lambda/4)} \cdot \left(\frac{\Gamma(1/2)}{\Gamma(3/4-i\lambda/4)\Gamma(3/4+i\lambda/4)}\right)^{-1}$$

$$= e^{\frac{\pi}{4}(\lambda-i)} \cdot \Gamma((1+i\lambda)/4)\Gamma(3/4-i\lambda/4)$$

$$= e^{\frac{\pi}{4}(\lambda-i)} \cdot \Gamma((1+i\lambda)/4)\Gamma(1+(-1/4-i\lambda/4))$$

$$= e^{\frac{\pi}{4}(\lambda-i)} \cdot (-1/4-i\lambda/4)\Gamma((1+i\lambda)/4)\Gamma(-(1+i\lambda)/4)$$

$$= \frac{e^{\frac{\pi}{4}(\lambda-i)}\pi}{\sin\pi(1/4+i\lambda/4)}$$

Likewise we obtain

$$C_1^{-}(\lambda) = C_1^{+}(-\lambda) = \frac{e^{-\frac{\pi}{4}(\lambda+i)}\pi}{\sin\pi(1/4 - i\lambda/4)}$$

and so

$$c_{1}(\lambda) = \frac{e^{\frac{\pi}{4}(\lambda-i)}\pi}{\sin\pi(1/4+i\lambda/4)} + \frac{e^{-\frac{\pi}{4}(\lambda+i)}\pi}{\sin\pi(1/4-i\lambda/4)}$$

$$= \frac{\pi}{i}\frac{e^{\frac{\pi}{4}(\lambda-i)}}{\sinh\pi(\lambda/4-i/4)} + \frac{\pi}{i}\frac{e^{-\frac{\pi}{4}(\lambda+i)}}{\sinh\pi(-\lambda/4-i/4)}$$

$$= \frac{\pi}{i}\frac{\cosh(\pi\lambda 4 - i\frac{\pi}{4}) + \sinh(\frac{\pi\lambda}{4} - i\frac{\pi}{4})}{\sinh\pi(\lambda/4 - i/4)} + \frac{\pi}{i}\frac{\cosh(-\frac{\pi\lambda}{4} - i\frac{\pi}{4}) + \sinh(-\frac{\pi\lambda}{4} - i\frac{\pi}{4})}{\sinh\pi(-\lambda/4 - i/4)}$$

$$= \frac{\pi}{i} \cdot \left(2 + \coth(\frac{\pi}{4}(\lambda-i)) + \coth(-\frac{\pi}{4}(\lambda+i))\right)$$

Now define $g(\lambda)$ by the requirement

$$g(\lambda)c_1(\lambda) = \frac{1}{|\mathbf{c}(i\lambda)|^2} = \frac{\pi\lambda}{2} \tanh\left(\frac{\pi\lambda}{2}\right)$$

Now note that

$$c_1(\lambda + i) = \frac{\pi}{i} \cdot \left(2 + \coth(\frac{\pi\lambda}{4}) + \coth(-\frac{\pi\lambda}{4} - i\frac{\pi}{2})\right)$$
$$= \frac{\pi}{i} \cdot \left(2 + \coth(\frac{\pi\lambda}{4}) - \tanh(\frac{\pi\lambda}{4})\right)$$
$$= \frac{\pi}{i} \cdot \left(2 + \frac{2}{\sinh(\frac{\pi\lambda}{2})}\right)$$

and thus we get

$$g(\lambda+i)\frac{\pi}{i}\cdot\left(2+\frac{2}{\sinh(\frac{\pi\lambda}{2})}\right) = \frac{\pi(\lambda+i)}{2}\coth\left(\frac{\pi\lambda}{2}\right).$$

Further manipulation then yields that

$$g(\lambda) = \frac{i\lambda}{4} \cdot \frac{\sinh(\frac{\pi\lambda}{2})}{1 - \cosh(\frac{\pi\lambda}{2})}$$

and it is obvious that g is not a polynomial function. Since

$$g(\lambda)C_1(\lambda) = C_2(\lambda)$$

it is now clear that there exists *no* differential operator \mathcal{L} which inverts the holomorphic Radon transform. However the function $g(\lambda)$ defines us a spectral multiplier which is a pseudo-differential operator which we now call \mathcal{L} . We summarize our discussion.

Theorem 6.2. Let $G = \operatorname{Sl}(2, \mathbb{R})$ and \mathcal{L} the spectral multiplier defined by the function $g(\lambda) = \frac{i\lambda}{4} \cdot \frac{\sinh(\frac{\pi\lambda}{2})}{1-\cosh(\frac{\pi\lambda}{2})}$. Then for $f \in \mathcal{H}^2(D)_0$ such that $\mathcal{R}(f) \in \mathcal{F}(\Xi_+)$ one has

$$f = (\mathcal{LR}(f))^{\vee}.$$

7. Geometric definition of the Hardy space

This final section deals with the structure of the Hardy space $\mathcal{H}^2(D)$. It allows independent reading and is of independent interest.

Initially, the Hardy space was defined spectrally (see [11]). Below we will show how to define the Hardy space geometrically, i.e. we give geometric definition of the norm on $\|\cdot\|_H$ on $\mathcal{H}^2(D)$ through *G*-orbit integrals on *D*. For that we start by recalling the orbital integral \mathbf{O}_h and the pseudo-differential operator \mathcal{D} introduced also used in [17].

In this section Y = G/H can be an arbitrary NCC-space.

7.1. *G*-orbit integrals on the domain *D*. For a sufficiently decaying functions *h* on *D* we define its *G*-orbit integral on *D* as the following function on $i2\Omega_H$

$$\mathbf{O}_h(iX) = \int_G h(g \exp(i\frac{1}{2}X) \cdot x_o) \, dg \qquad (X \in 2\Omega_H) \, .$$

For $f \in \mathcal{H}^2(D)$ we notice that $|f|^2$ is a sufficiently decaying function on D, i.e. $\mathbf{O}_{|f|^2}(iX)$ is finite for all $X \in 2\Omega_H$. Moreover, in view of (3.8) we see that $\mathbf{O}_{|f|^2}$ has a natural holomorphic extension to a holomorphic function on the abelian tube domain $\mathcal{T}(2\Omega_H) = \mathfrak{a} + i2\Omega_H$, namely

(7.1)
$$\mathbf{O}_{|f|^2}(Z) = \int_{\mathcal{X}} |\hat{f}(b,\lambda)|^2 \varphi_{\lambda}(\exp(Z)) \ d\mu_{\mathcal{X}}(b,\lambda) \qquad (Z \in \mathcal{T}(2\Omega_H))$$

7.2. A certain pseudo-differential operator. Define a space $\mathcal{F}(\mathcal{T}(2\Omega_H))$ of \mathcal{W} -invariant holomorphic functions on the tube domain $\mathcal{T}(2\Omega_H)$ by the following property: $f \in \mathcal{F}(\mathcal{T}(2\Omega_H))$ if f can be written as

$$f(Z) = \int_{i\mathfrak{a}_{+}^{*}} h(\lambda)\varphi_{\lambda}(\exp(Z))\frac{d\lambda}{|\mathbf{c}(\lambda)|^{2}} \qquad (Z \in \mathcal{T}(2\Omega_{H}))$$

where $h \in L^1(i\mathfrak{a}^*_+, \frac{\cosh(\lambda)}{|\mathbf{c}(\lambda)|^2}d\lambda)$. If $Q \subset \mathcal{T}(2\Omega_H)$ is compact, then there exists a constant $C_Q > 0$ such that

$$(\forall \lambda \in i\mathfrak{a}^*)$$
 $\sup_{X \in Q} |\varphi_{\lambda}(\exp(i2X))| \le C_Q \cosh(\lambda).$

As $\frac{1}{|\mathbf{c}(\lambda)|^2}$ is at most of polynomial growth, it follows that f is indeed holomorphic and \mathcal{W} -invariant. Moreover, f is uniquely determined by h. It follows from our discussion that the prescription

$$\mathcal{D}: \mathcal{F}(\mathcal{T}(2\Omega_H)) \to \mathcal{O}(\mathcal{T}(2\Omega_H))^{\mathcal{W}}; \ (\mathcal{D}F)(Z) = \int_{i\mathfrak{a}^*_+} h(\lambda) \sum_{w \in \mathcal{W}} e^{\lambda(wZ)} \frac{d\lambda}{|\mathbf{c}(\lambda)|^2}$$

is a well defined linear mapping.

Remark 7.1. The operator \mathcal{D} is a pseudo-differential operator and a differential operator if all root multiplicities are even. The operator \mathcal{D} is related to the Abel transform as explained in [17], Remark 3.2.

Example 7.2. In this example we discuss the operator \mathcal{D} when the underlying group G is complex. Then \mathcal{D} is a differential operator of a particularly nice form.

If G is complex, then there is an explicit formula for spherical functions, due to Harish-Chandra:

$$\varphi_{\lambda}(\exp(Z)) = \mathbf{c}(\lambda) \frac{\sum_{w \in \mathcal{W}} \varepsilon(w) e^{\lambda(wZ)}}{\prod_{\alpha \in \Sigma^{+}} 2 \sinh \alpha(Z)}$$

for all $Z \in \mathcal{T}(2\Omega_H)$. The **c**-function has the familiar form

$$\mathbf{c}(\lambda) = \frac{\prod_{\alpha \in \Sigma^+} \langle \rho, \alpha \rangle}{\prod_{\alpha \in \Sigma^+} \langle \lambda, \alpha \rangle} \; .$$

For each $\alpha \in \Sigma$ let $A_{\alpha} \in \mathfrak{a}$ be such that $\alpha = \langle \cdot, A_{\alpha} \rangle$. Furthermore let ∂_{α} be the partial derivative on $\mathcal{T}(2\Omega_H)$ in direction A_{α} . Define a partial differential operator on $\mathcal{T}(2\Omega_H)$ by $\partial_{\Sigma^+} = \prod_{\alpha \in \Sigma^+} \partial_{\alpha}$. Finally with $J(Z) = \prod_{\alpha \in \Sigma^+} 2 \sinh \alpha(Z)$ we declare a differential operator on $\mathcal{T}(2\Omega_H)$ by

$$\mathcal{D} = \operatorname{const} \cdot \partial_{\Sigma^+} \circ J \; .$$

with const = $\prod_{\alpha \in \Sigma^+} \langle \rho, \alpha \rangle$. The relation

$$\mathcal{D}(\varphi_{\lambda} \circ \exp)(Z) = \sum_{w \in \mathcal{W}} e^{\lambda(wZ)}$$

is now obvious.

7.3. The geometric norm. For a function $f \in \mathcal{H}^2(D)$ let us write $||f||_H$ for its norm as before. By Lemma 3.2 this norm is given by

(7.2)
$$||f||_{H} = \int_{\mathcal{X}} |\hat{f}(b,\lambda)|^{2} \cosh(\lambda) \ d\mu_{\mathcal{X}}(b,\lambda)$$

The objective of this section is to express $||f||_H$ in terms of the much more geometric orbital integrals $O_{|f|^2}$. Our result is as follows.

Theorem 7.3. Let $f \in \mathcal{H}^2(D)$. Then $\mathbf{O}_{|f|^2} \in \mathcal{F}(\mathcal{T}(2\Omega_H))$ and the Hardy space norm $||f||_H$ of f is given by

(7.3)
$$||f||_{H} = \sup_{X \in 2\Omega_{H}} \frac{\mathcal{D}\mathbf{O}_{|f|^{2}}(iX)}{|\mathcal{W}_{H}|}$$

In particular, the Hardy space $\mathcal{H}^2(D)$ can be defined as

(7.4)
$$\mathcal{H}^2(D) = \{ f \in \mathcal{O}(D) \mid \mathbf{O}_{|f|^2} \in \mathcal{F}(\mathcal{T}(2\Omega_H)) \sup_{X \in 2\Omega_H} \frac{\mathcal{D}\mathbf{O}_{|f|^2}(iX)}{|\mathcal{W}_H|} < \infty \} .$$

Proof. Fix $f \in \mathcal{H}^2(D)$. By equation (7.1) we have

$$\mathbf{O}_{|f|^2}(Z) = \int_{\mathcal{X}} |\hat{f}(b,\lambda)|^2 \varphi_{\lambda}(\exp(Z)) \ d\mu_{\mathcal{X}}(b,\lambda)$$

for all $Z \in \mathcal{T}(2\Omega_H)$. By the spectral definition of $\mathcal{H}^2(D)$ it follows that

$$h(\lambda) := \int_B |\hat{f}(b,\lambda)|^2 \ db$$

defines a function $h \in L^1(i\mathfrak{a}^*_+, \frac{\cosh(\lambda)}{|\mathbf{c}(\lambda)|^2} d\lambda)$. Thus $\mathbf{O}_{|f|^2} \in \mathcal{F}(\mathcal{T}(2\Omega_H))$ and the application of \mathcal{D} to $\mathbf{O}_{|f|^2}$ yields

$$(\mathcal{D}\mathbf{O}_{|f|^2})(Z) = \int_{\mathcal{X}} |\hat{f}(b,\lambda)|^2 \sum_{w \in \mathcal{W}} e^{\lambda(Z)} d\mu_{\mathcal{X}}(b,\lambda).$$

Now notice that

$$\frac{1}{|\mathcal{W}_H|} \sum_{w \in \mathcal{W}} e^{\lambda(iwX)} \le \cosh(\lambda) \qquad (X \in 2\Omega_H)$$

and

$$\sup_{X \in 2\Omega_H} \sum_{w \in \mathcal{W}} e^{\lambda(iwX)} = \lim_{X \to Z_H} \sum_{w \in \mathcal{W}} e^{\lambda(i2wX)} = |\mathcal{W}_H| \cdot \cosh(\lambda) .$$

The claim follows now from the spectral definition of the norm in $\mathcal{H}^2(D)$. Finally, back-tracking the steps of the proof readily yields (7.4).

Remark 7.4. Some comments on the geometric Hardy norm

$$||f||_{H} = \sup_{Z \in \Omega_{H}} \frac{1}{|\mathcal{W}_{H}|} \cdot (\mathcal{D}\mathbf{O}_{|f|^{2}})(Z)$$

seem to be appropriate. Usually, in the theory of Hardy spaces (e.g. Hardy space on the upper half plane) one takes the supremum over a family of L^2 -integrals over totally real submanifolds. In our case one takes a supremum over G-orbits, which for the exception of the orbit through the origin, are never totally real. Secondly, we find the appearance of the pseudo differential operator \mathcal{D} interesting. In the context of Hardy spaces it might be novel.

7.4. The K-invariant case. In this subsection we give another description of the subspace $\mathcal{H}^2(D)^K$ using the Abel transform and the results in Appendix A. We start by noting the following simple connection between the Abel transform and the Fourier transform of a

K-invariant function. For that we note first, that $\hat{f}(b,\lambda)$ is independent of $b \in B$ if f is K-invariant. We write then simply $\hat{f}(\lambda)$ and note that \hat{f} is W-invariant. Furthermore

$$\hat{f}(\lambda) = \int_{X} f(x)a(x)^{\rho-\lambda} dx$$
$$= \int_{A} \int_{N} f(na \cdot x_{o})a^{-\rho-\lambda} dnda$$
$$= \mathcal{F}_{A}(\mathcal{A}(f))(\lambda)$$

where \mathcal{F}_A stands for the Fourier transform on the abelian group A. Recall, that $\mathcal{F} : L^2(A) \to L^2(i\mathfrak{a}^*, |\mathcal{W}|^{-1}d\lambda)$ is a unitary isomorphism. Define a multiplication operator $D_{\mathfrak{a}}$ on $i\mathfrak{a}^*$ by $D_{\mathfrak{a}}(F) = \mathbf{c}(-\lambda)^{-1}F$ and denote the corresponding multiplier operator on A by D_A . Let $\Lambda := D_A \circ \mathcal{A}$. Finally, we define a multiplier m on $\mathcal{W} \times i\mathfrak{a}^*$ by $m(s, \lambda) = \mathbf{c}(-s\lambda)/\mathbf{c}(-\lambda)$. We denote by τ the corresponding representation $\tau(s)f(\lambda) = m(s^{-1},\lambda)f(s^{-1}\lambda)$. Then, cf. [?], Section 1, in particular Lemma 1.4, we have a commutative diagram, where each of the maps is an unitary isomorphism:

(7.5)
$$L^{2}(A, |\mathcal{W}|^{-1}da)^{\mathcal{W}} \xrightarrow{\Lambda} L^{2}(A, |\mathcal{W}|^{-1}da)^{\tau(\mathcal{W})}$$
$$\downarrow^{\mathcal{F}_{X}} \qquad \qquad \qquad \downarrow^{\mathcal{F}_{A}}$$
$$L^{2}(i\mathfrak{a}^{*}, \frac{d\lambda}{|\mathcal{W}||\mathbf{c}(\lambda)|^{2}}) \xrightarrow{}_{D_{\mathfrak{a}}} L^{2}(i\mathfrak{a}^{*}, |\mathcal{W}|^{-1}d\lambda)^{\tau(\mathcal{W})}$$

Recall the Hardy space $\mathcal{H}^2(T(\Omega_H))$ from Appendix A and its spectral description in Theorem 7.9. It follows then from Theorem 3.2, and the obvious renormalization of measures, as we have not included the 2π in the exponential function, that $\Lambda(\mathcal{H}^2(D)^K) \subseteq \mathcal{H}^2(T(\Omega_H))^{\tau(\mathcal{W})}$. As $\Lambda : L^2(A, |\mathcal{W}|^{-1}da)^{\mathcal{W}} \to L^2(A, |\mathcal{W}|^{-1}da)^{\tau(\mathcal{W})}$ is a unitary isomorphism, we get:

Theorem 7.5. The map $\Lambda : \mathcal{H}^2(D)^K \to \mathcal{H}^2(T(\Omega_H))^{\tau(\mathcal{W})}$ is a unitary isomorphism.

Example 7.6. If G has complex structure, then the map Λ is a multiplication operator given by

$$\Lambda(f)(a) = \left(\prod_{\alpha \in \Sigma^+} \sinh \langle \alpha, \log(a) \rangle \right) f(a) \,.$$

We now determine the reproducing kernel for $\mathcal{H}^2(D)^K$. One could easily deduct that from Theorem 7.10 but we will give another proof, that follows similar arguments.

Theorem 7.7. The reproducing kernel $\mathcal{K}(z, w)$ for $\mathcal{H}^2(D)^K$ is given by

$$\mathcal{K}(z,w) = \int_{i\mathfrak{a}_+^*} \varphi_{\lambda}(z) \varphi_{-\lambda}(\overline{w}) \, d\mu(\lambda) \, .$$

Proof. Let $f \in \mathcal{H}^2(D)^K$ and $w \in D$. Recall, that by Lemma 3.2 we have $\Phi(g)(\lambda) = \hat{g}(\lambda) \cosh(\lambda)$ for all $g \in \mathcal{H}^2(D)^K$. Therefore

$$\begin{split} f(w) &= \langle f, K_w \rangle \\ &= \int_{i\mathfrak{a}_+} \hat{f}(\lambda) \mathbf{cosh}(\lambda) \overline{\widehat{K_w}(\lambda) \mathbf{cosh}(\lambda)} \, d\mu(\lambda) \\ &= \int_{i\mathfrak{a}_+^*} \hat{f}(\lambda) \overline{\widehat{K_w}(\lambda) \mathbf{cosh}(\lambda)} \, \frac{d\lambda}{|\mathbf{c}(\lambda)|^2} \\ &= \int_{i\mathfrak{a}_+^*} \hat{f}(\lambda) \varphi_\lambda(w) \, \frac{d\lambda}{|\mathbf{c}(\lambda)|^2} \, . \end{split}$$

Thus

$$\widehat{K_w}(\lambda) = \frac{\varphi_{-\lambda}(\overline{w})}{\cosh(\lambda)}.$$

From this we now get:

$$K(z,w) = \langle K_w, K_z \rangle$$

= $\int_{i\mathfrak{a}^*_+} \widehat{K_w}(\lambda) \operatorname{cosh}(\lambda) \overline{\widehat{K_z}(\lambda) \operatorname{cosh}(\lambda)} d\mu(\lambda)$
= $\int_{i\mathfrak{a}^*_+} \varphi_{-\lambda}(\overline{w}) \varphi_{\lambda}(z) d\mu(\lambda)$

and the claim follows.

APPENDIX: HARDY SPACES ON STRIP DOMAIN

We let V be an Euclidean vector space, e.g. $V = \mathbb{R}^n$ endowed with the standard inner product. Denote by O(V) the orthogonal group of V and let $\mathcal{W} \subset O(V)$ be a finite subgroup which acts irreducibly on V. We fix $y_o \in V$, $y_o \neq 0$ and set

 $\Omega = \operatorname{int} \left(\{ \operatorname{convex} \operatorname{hull} \operatorname{of} \mathcal{W}(y_o) \} \right) .$

Notice that Ω is the interior of a compact polyhedron and that $0 \in \Omega$. Write $V_{\mathbb{C}} := V \otimes_{\mathbb{R}} \mathbb{C} \simeq V + iV$ for the complexification of V and define a tube domain in $V_{\mathbb{C}}$ by

$$T(\Omega) = V + i\Omega \; .$$

Let us denote by dx the measure $(2\pi)^{-\dim V/2}$ times the normalized Lebesgue measure on V. Then the Fourier transform

$$f \mapsto \mathcal{F}f(\lambda) = \int_V f(x)e^{-\langle \lambda, x \rangle} dx$$

is a unitary L^2 -isomorphism. V^* is the dual of V, and $\langle \lambda \cdot x \rangle = \lambda(x)$ denotes the standard duality between V and V^* . Denote by $\mathcal{O}(T(\Omega))$ the space of holomorphic functions on $T(\Omega)$.

The Hardy space $\mathcal{H}^2(T(\Omega))$ is defined by:

$$\mathcal{H}^2(T(\Omega)) := \{ f \in \mathcal{O}(T(\Omega)) \mid ||f||_{\mathcal{H}}^2 = \sup_{y \in \Omega} \int_V |f(x+iy)|^2 \, dx < \infty \}$$

As the Hardy-norm locally dominates the Bergman-norm on $T(\Omega)$, it follows hence $\mathcal{H}^2(T(\Omega))$ is complete, i.e. a Banach space. In fact, $\mathcal{H}^2(T(\Omega))$ is a Hilbert space as we will show in a moment. Then for $f \in \mathcal{H}^2(T(\Omega))$ and $y \in \Omega$ one has

$$\int_{V} |f(x+iy)|^2 dx = \int_{V^*} |\mathcal{F}(f|_V)(\xi)|^2 e^{-2\langle y,\xi\rangle} d\xi$$

which is immediate from [22] Ch. III, §2. It follows that

(7.6)
$$||f||_{\mathcal{H}}^2 = \sup_{y \in \Omega} \int_{V^*} |\mathcal{F}(f|_V)(\xi)|^2 e^{-2\langle y, \xi \rangle} d\xi$$

For $y \in V$ define $\cosh, \cosh_y : V^* \to \mathbb{C}$ by

$$\cosh_{y}(\lambda) = \frac{1}{|W|} \sum_{s \in \mathcal{W}} e^{-2\langle y, s\lambda \rangle} = \frac{1}{|W|} \sum_{s \in \mathcal{W}} e^{-2\langle sy, \lambda \rangle}$$

and

$$\cosh(\lambda) = \cosh_{y_o}(\lambda)$$
 .

As with $\|\cdot\|_{L^2(V)}$ is \mathcal{W} -invariant and \mathcal{F} is \mathcal{W} -equivariant, it follows from (7.6) that

(7.7)
$$||f||^2 = \sup_{y \in \Omega} \int_{V^*} |\mathcal{F}(f|_V)(\xi)|^2 \cosh(y,\xi) \ d\xi \ .$$

Now, for every $\lambda \in V$, the function $y \mapsto \cosh_{\mathbf{y}}(\lambda)$ is strictly convex on $\overline{\Omega}$; hence we have the inequality

$$(\forall y \in \Omega)(\forall \lambda \in V^*)$$
 $\cosh_y(\lambda) \le \cosh(\lambda)$

and so it follows that

(7.8)
$$||f||_{\mathcal{H}}^2 = \int_{V^*} |\mathcal{F}(f|_V)(\lambda)|^2 \mathbf{cosh}(\lambda) \ d\lambda$$

Define a \mathcal{W} -invariant measure μ_0 on V^* by

$$d\mu_0(\lambda) = \cosh(\lambda) d\lambda$$
.

Theorem 7.8. The mapping

$$\mathcal{H}^2(T(\Omega)) \to L^2(V^*, d\mu_0), \quad f \mapsto \mathcal{F}(f|_V)$$

is an isometric isomorphism. In particular, $\mathcal{H}^2(T(\Omega))$ is a Hilbert space.

A (continuous) multiplier on V^* is a continuous map $m: \mathcal{W} \times V^* \to \mathbb{C}$ such that for all $s, w \in \mathcal{W}$ and $\lambda \in V^*$ we have

$$m(sw, \lambda) = m(s, w\lambda)m(w, \lambda)$$
.

Assume from now on that $|m(s,\lambda)| = 1$ for all $s \in \mathcal{W}$ and $\lambda \in V^*$. Then, because of the \mathcal{W} -invariance of $d\mu_0$, we can define a unitary representation of \mathcal{W} on $L^2(V^*, d\mu_0)$ by

(7.9)
$$\tau(s)f(\lambda) = m(s^{-1},\lambda)f(s^{-1}\lambda).$$

As the Fourier transform is a unitary isomorphism, we have a unitary representation, also denoted by τ , of \mathcal{W} on $\mathcal{H}^2(T(\Omega))$ such that the Fourier transform is an intertwining operator. Denote the space of $\tau(\mathcal{W})$ -invariant elements by the superscript $\tau(\mathcal{W})$. Then

Corollary 7.9. The Fourier transform is a unitary isomorphism

 $\mathcal{F}: \mathcal{H}^2(T(\Omega))^{\tau(\mathcal{W})} \to L^2(V^*, d\mu_0)^{\tau(\mathcal{W})}.$

The Hardy space $\mathcal{H}^2(T(\Omega))^{\tau(\mathcal{W})}$ being a Hilbert space of holomorphic functions admits as such a reproducing kernel function K(z, w) often called the Cauchy-Szegö-kernel. Notice that for fixed $w \in T(\Omega)$, the function $K_w(z)$ belongs to $\mathcal{H}^2(T(\Omega))^{\tau(\mathcal{W})}$ and that

$$\langle f, K_w \rangle = f(w)$$
 for all $f \in \mathcal{H}^2(T(\Omega))^{\tau(\mathcal{W})}$

Here of course $\langle \cdot, \cdot \rangle$ denotes the inner product on $\mathcal{H}^2(T(\Omega))^{\tau(\mathcal{W})}$.

We now determine K(z, w) explicitly. For that define for $w \in V$, $\cos_w^m : V^* \to \mathbb{C}$ by

$$\cos_w^m(\lambda) := \frac{1}{|\mathcal{W}|} \sum_{s \in \mathcal{W}} m(s, \lambda)^{-1} e^{i \langle w, s \lambda \rangle}$$

and note that

$$\frac{\cos_w^m}{\cosh} \in \mathcal{H}^2(T(\Omega))^{\tau(\mathcal{W})}$$

Write $(\cdot|\cdot)$ for the inner product on $L^2(V^*, d\mu_0)^{\tau(\mathcal{W})}$. For $f \in \mathcal{H}^2(T(\Omega))^{\tau(\mathcal{W})}$ let $F = \mathcal{F}(f|_V)$. It follows from Corollary 7.9 that $\langle f, K_w \rangle = (F|\mathcal{F}(K_w|_V))$. On the other hand we have

$$\begin{split} f(w) &= \int_{V^*} F(\lambda) e^{-i\langle w,\lambda\rangle} d\lambda \\ &= \int_{V^*} F(\lambda) \frac{e^{-i\langle w,\lambda\rangle}}{\cosh(\lambda)} d\mu_0(\lambda) \\ &= \int_{V^*} m(s^{-1},\lambda) F(s^{-1}\lambda) \frac{e^{-i\langle w,\lambda\rangle}}{\cosh(\lambda)} d\mu_0(\lambda) \\ &= \int_{V^*} F(\lambda) m(s^{-1},s\lambda) \frac{e^{-i\langle w,s\lambda\rangle}}{\cosh(\lambda)} d\mu_0(\lambda) \\ &= \int_{V^*} F(\lambda) \overline{\frac{\cos^m(\lambda)}{\cosh(\lambda)}} d\mu_0(\lambda) \end{split}$$

and thus

$$\mathcal{F}(K_w|_V)(\lambda) = \frac{\cos_w^m(\lambda)}{\cosh(\lambda)}.$$

Theorem 7.10. The reproducing kernel for the Hardy space $\mathcal{H}^2(T(\Omega))^{\tau(\mathcal{W})}$ is given by

(7.10)
$$K(z,w) = \int_{V^*} \frac{\left(\frac{1}{|W|} \sum_{s \in W} \frac{e^{i\langle s(z), \lambda \rangle}}{m(s, \lambda)}\right) \cdot \left(\frac{1}{|W|} \sum_{s \in W} \frac{e^{i\langle s(w), \lambda \rangle}}{m(s, \lambda)}\right)}{\cosh(\lambda)} d\lambda$$
$$= \int_{V^*} \frac{\cos^m_z(\lambda)}{\cosh(\lambda)} \cos^m_{-\overline{w}}(\lambda) d\lambda \,.$$

Example 7.11. The equation (7.10) can be evaluated in the relevant special cases. Let us for example consider the case of $V = \mathbb{R}$, $\Omega =]-1, 1[$, $\mathcal{W} = O(\mathbb{R}) \simeq \{1, -1\}$ and $m(s, \lambda) = 1$. Using the standard measure on \mathbb{R} we get the following from (7.10) and [3], Sect. 1.9, formula (12):

$$K(z,w) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \frac{\cos(z\xi)\cos(\overline{w}\xi)}{\cosh(2\xi)} d\xi$$
$$= \sqrt{\frac{\pi}{2}} \frac{\cosh(\frac{\pi}{4}z) \cdot \cosh(\frac{\pi}{4}\overline{w})}{\cosh(\frac{\pi}{2}z) + \cosh(\frac{\pi}{2}\overline{w})}.$$

References

- J. Arthur, A theorem on the Schwartz space of a reductive Lie group, Proc. Natl. Acad. Sci. USA 72, 4718–4719 (1975)
- [2] E. van den Ban, A convexity theorem for semisimple symmetric spaces, Pacific J. Math. 124 (1)(1986) , 21–55
- [3] A. Erdelyi et al., Tables of Integral Transforms, Volume I, McGraw-Hill, 1954.
- [4] Bernstein, J., Analytic continuation of generalized functions with respect to a parameter, Funkcional. Anal. i Priložen. 6 (1972), no. 4, 26–40.
- [5] P. Delorme and M. Flensted-Jensen, Towards a Paley-Wiener theorem for semisimple symmetric spaces, Acta Math. 167 (1991), 127–151.
- [6] J. Faraut, Formule de Gutzmer pour la complexification d'un espace Riemannien symétrique, Rend. Mat. Acc. Lincei s. 9, v. 13:233-241 (2002).
- [7] Gindikin, S., SO(1; n)-twistors, J. Geom. Phys. 26 (1998), no. 1-2, 26–36.
- [8] —, Integral geometry on SL(2; R), Math. Res. Lett. 7 (2000), no. 4, 417–432.
- [9] —, Horospherical Cauchy-Radon transform on compact symmetric spaces, Moscow Math. J., to appear
- [10] S. Gindikin and B. Krötz, Invariant Stein Domain in Stein Symmetric Spaces and a Nonlinear Complex Convexity Theorem, IMRN 18 (2002), 959–971.
- [11] S. Gindikin, B. Krötz and G. Ólafsson, Holomorphic H-spherical distribution vectors in principal series representations, Invent. Math. 158 (2004), 643–682.
- [12] _____, Erratum: Holomorphic H-spherical distribution vectors in principal series representations, Invent. Math. 158 (2004), 683 - 684.
- [13] _____, Horospherical model for the holomorphic discrete series and the horospherical Cauchy transform, Compositio Math. 142 (2006), 983–1008.

- [14] Harish-Chandra, Spherical Functions on a Semisimple Lie Group I. Amer. J. Math. 80 (1958), 241–310.
- [15] Harish-Chandra, Discrete Series for Semisimple Lie Groups II, Acta Math. 116 (1966), 1–111.
- [16] B. Krötz, Domains of holomorphy for irreducible unitary representations of simple Loe groups, MPIM preprint 2006 (102).
- [17] B. Krötz, G. Ólafsson and R. Stanton, The image of the heat kernel transform on Riemannian symmetric spaces of the noncompact type, IMRN 22 (2005), 1307–1329.
- [18] B. Krötz and M. Otto, A Refinement of the Complex Convexity Theorem via Symplectic Techniques, Proc. Amer. Math. Soc. 134 (2) (2006), 549–558.
- [19] B. Krötz and R. Stanton, Holomorphic extension of representations: (I) automorphic functions, Ann. Math. 159(2004), 641–724.
- [20] M. Kashiwara and M. Vergne, K-types and singular spectrum, Noncommutative harmonic analysis (Proc. Third Colloq., Marseille-Luminy, 1978), 177–200, Lecture Notes in Math., 728, Springer, Berlin, 1979.
- [21] T. Matsuki, The orbits of affine symmetric spaces under the action of minimal parabolic subgroups, J. Math. Soc. Jpn. 31 (1979), 331–357.
- [22] E. Stein and G. Weiss, Introduction to Fourier Analysis on Euclidean Spaces, Princeton University Press, 1971.

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, NEW BRUNSWICK, NJ 08903 E-mail address: gindikin@math.rutgers.edu

MAX-PLANCK-INSTITUT FÜR MATHEMATIK, VIVATSGASSE 7, D-53111 BONN, GERMANY *E-mail address*: kroetz@mpim-bonn.mpg.de

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LA 70803, USA *E-mail address:* olafsson@math.lsu.edu