Coarse geometry and asymptotic dimension

Dissertation zur Erlangung des Doktorgrades der Mathematisch-Naturwissenschaftlichen Fakultäten der Georg-August-Universität zu Göttingen

> vorgelegt von Bernd Grave aus Damme

Göttingen 2005

D7

Referent: Prof. Dr. Thomas SchickKorreferent: Prof. Dr. Ulrich StuhlerTag der mündlichen Prüfung: Montag, 23. Januar 2006

Abstract

We prove that two spaces, whose coarse structures are induced by metrisable compactifications, are coarsely equivalent if and only if their (Higson) coronas are homeomorphic.

We introduce translation C^* -algebras for coarse spaces which admit a countable, uniformly bounded cover using projection-valued measures. This was already done in [Roe03]. Here we give a more complete exposition on the subject including a result about the independence of the translation C^* algebra from the involved projection-valued measure. Moreover, we eliminate some mistakes, e.g. we have to be careful about null sets when defining the support of an operator.

We introduce some characterisations of asymptotic dimension in the general setting of coarse spaces and prove some basic properties such as monotony, a formula for the asymptotic dimension of finite unions and estimates for the asymptotic dimension of the product of two coarse spaces.

We define coarse cell complexes and prove the obvious conjecture about their asymptotic dimension.

We prove $\operatorname{asdim}(X, \mathcal{E}) = \dim(K \setminus X) + 1$ if the coarse structure \mathcal{E} is induced by a metrisable compactification K of X. As an application, we obtain coarse spaces X and Y such that $\operatorname{asdim}(X \times Y) < \operatorname{asdim}(X) + \operatorname{asdim}(Y)$.

For $CAT(\kappa)$ -spaces with $\kappa < 0$ having nicely *n*-covered spheres we prove that the asymptotic dimension is at most *n*. We apply this result to prove that asdim(X) = dim(X) for any complete, simply connected Riemannian manifold X with bounded, strictly negative sectional curvature.

Acknowledgements

First of all I wish to thank my advisor, Thomas Schick, for his support, many enlightening discussions and most especially his patience when I was struggling.

The Graduiertenkolleg Gruppen und Geometrie at Göttingen University provided the financial support and mathematical education which benefited me greatly in this endeavor. And I profited as well from visiting the research group in Noncommutative Geometry and Operator Algebras at Vanderbilt University.

Special thanks to Paul Mitchener for inspiring discussions about coarse geometry, his many helpful suggestions and his very special English lessons, in addition to those taught by Marjory Frauts.

I am also grateful to Elias Kappos, Behnam Norouzizadeh and Moritz Wiethaup for their careful read of this dissertation and their detailed feedback.

Last, but not least, I wish to thank all my friends both in the Göttingen math department and without for their kindness and the part they play in making Göttingen such an attractive place in which to work and live.

Contents

1	Some basics in coarse geometry		3
	1.1	Finitely generated groups from the metric viewpoint	3
	1.2	The coarse category	4
	1.3	Some constructions	9
	1.4	Coarse structures induced by metrisable compactifications	13
2	Translation C^* -algebras via projection-valued measures		
	2.1	Geometric Hilbert spaces and the calculus of supports $\ . \ . \ .$	19
	2.2	Translation C^* -algebras $\ldots \ldots \ldots \ldots \ldots \ldots \ldots \ldots \ldots \ldots \ldots$	23
	2.3	Induced maps	29
3	General theory of asymptotic dimension		33
	3.1	Asymptotic dimension of pseudometric spaces	33
	3.2	Asymptotic dimension of coarse spaces	35
	3.3	Basic properties	40
4	Asy	mptotic dimension of coarse cell complexes	47
5	Asymptotic dimension and the covering dimension of th Higson corona		e 57
6	Asymptotic dimension of $CAT(\kappa)$ -spaces for $\kappa < 0$		63
	6.1	Spaces with nicely covered spheres	63
	6.2	Proof of Theorem 6.5	64
	6.3	Triangulations of bounded distortion	67
	6.4	Complete, simply connected manifolds with bounded, strictly	
		negative sectional curvature	70
	Bibliography		73
	Ind	ex	77
	Cuu	rriculum vitae	79

Chapter 1

Some basics in coarse geometry

1.1 Finitely generated groups from the metric viewpoint

In this section we give an example of a setting where coarse geometry naturally arises. We define the word metric on each finitely generated group. This metric depends on the chosen generating set, but - as we will see soon - the asymptotic behavior does not. So we can associate a coarse structure to each finitely generated group.

Let G be a group with a finite generating set A.

Definition 1.1 (word metric) The distance $d_{(G,A)}(g_1, g_2)$ of $g_1, g_2 \in G$ in the word metric associated to the pair (G, A) is the length of the shortest word in A representing $g_1^{-1}g_2$.

Alternatively we could have used the Cayley graph of (G, A) to define this metric. Compare [BH99] for more details.

Lemma 1.2 If B is another finite generating set of G, there is $\lambda > 0$ such that $d_{(G,A)} \leq \lambda \cdot d_{(G,B)}$.

Proof. For each $b \in B$ choose a word in A of minimal length representing b. Define λ to be the maximum of the length of these words. Given any word in B representing $g \in G$, we replace each letter by the chosen word in A and count letters.

Definition 1.3 Let X, Y be pseudometric¹ spaces and $f: X \to Y$ a (not necessarily continuous) map.

¹Omitting the condition $d(x_1, x_2) > 0$ if $x_1 \neq x_2$ in the definition of a metric space, we get the definition of a pseudometric space. In a pseudometric space being of distance zero

- f is called *coarsely proper* if the inverse image of any bounded set is bounded.
- f is called *coarsely uniform* if for every r > 0 there is s(r) > 0 such that $d(f(x_1), f(x_2)) \le s(r)$ for all $x_1, x_2 \in X$ with $d(x_1, x_2) \le r$.
- f is called a *coarse map* if it is coarsely proper and coarsely uniform.
- Let S be a set. Two maps $f, g: S \to X$ are called *close* if there is D > 0 such that d(f(s), g(s)) < D for all $s \in S$.
- f is called a *coarse equivalence* if it is a coarse map and if there exists another coarse map $g: Y \to X$ such that $g \circ f$ is close to id_X and $f \circ g$ is close to id_Y .
- X and Y are called *coarsely equivalent* if there exists a coarse equivalence from X to Y.

Compare our coarse maps with asymptotically Lipschitz maps in [Dra00].

Consider a finitely generated group. The metric spaces corresponding to different finite generating sets are obviously coarsely equivalent. Thus, every functor from the category of metric spaces and arbitrary maps which is invariant under coarse equivalence gives an invariant of finitely generated groups. The asymptotic dimension will be an example of such an invariant.

1.2 The coarse category

In this section an axiomatic description of the structure needed to do coarse geometry will be given. Compare [Roe96], [Mit01] and [Roe03].

Definition 1.4 (coarse structure) Let X be a set. A collection \mathcal{E} of subsets of $X \times X$ is called a *coarse structure*, and the elements of \mathcal{E} will be called *entourages*, if the following axioms are fulfilled:

- (a) A subset of an entourage is an entourage.
- (b) A finite union of entourages is an entourage.
- (c) The diagonal $\Delta_X := \{(x, x) \mid x \in X\}$ is an entourage.
- (d) The inverse E^{-1} of an entourage E is an entourage.

$$E^{-1} := \{ (y, x) \in X \times X \mid (x, y) \in E \}$$

is an equivalence relation. Equivalent points are contained in exactly the same open sets. The map to the quotient can be used to carry over many properties of metric spaces to pseudometric spaces. From the asymptotic point of view pseudometric spaces should be as good as metric spaces.

(e) The composition E_1E_2 of entourages E_1 and E_2 is an entourage.

$$E_1E_2 := \{(x, z) \in X \times X \mid \exists_{y \in X}(x, y) \in E_1 \text{ and } (y, z) \in E_2\}$$

The pair (X, \mathcal{E}) is called a *coarse space*. Sometimes we will say that $E \subseteq X \times X$ is *controlled* if E is an entourage.

A coarse space is called *connected* if every point of $X \times X$ is contained in an entourage.

Remark 1.5 For $E_1, E_2, F_1, F_2 \subseteq X \times X$ we have

$$(E_1 \cup E_2)(F_1 \cup F_2) = E_1F_1 \cup E_1F_2 \cup E_2F_1 \cup E_2F_2$$
.

Definition 1.6 (bounded sets) Let (X, \mathcal{E}) be a coarse space, $A \subseteq X$ and $E \in \mathcal{E}$. We define

$$E[A] := \{ x \in X \mid (x, a) \in E \text{ for some } a \in A \} .$$

For a point $x \in X$ we will write E(x) instead of $E[\{x\}]$. Sets of the form E(x) with $x \in X$ and $E \in \mathcal{E}$ are called *bounded*.

Proposition 1.7 (properties of bounded sets) Let (X, \mathcal{E}) be a coarse space.

- 1. Subsets of bounded sets are bounded.
- 2. If $B \subseteq X$ is bounded, then $B \times B \in \mathcal{E}$.
- 3. If $B \subseteq X$ is bounded and $E \in \mathcal{E}$, then E[B] is bounded.
- 4. Let $B_1, B_2 \subseteq X$ be bounded sets. The following are equivalent.
 - $B_1 \cup B_2$ is bounded.
 - $B_1 \times B_2 \in \mathcal{E}$
 - There exists an entourage $E \in \mathcal{E}$ such that $E \cap B_1 \times B_2 \neq \emptyset$.

If (X, \mathcal{E}) is a connected coarse space, then any finite union of bounded sets is bounded.

Proof. Observe that for entourages $E_1, E_2 \in \mathcal{E}$ and $A \subseteq X$ we have $E_1E_2[A] = E_1[E_2[A]]$. Therefore E[B] is bounded if $E \in \mathcal{E}$ and $B \subseteq X$ is bounded.

Let $B_1, B_2 \subseteq X$ be bounded sets and let $E \in \mathcal{E}$ such that $E \cap B_1 \times B_2 \neq \emptyset$. Take $(b_1, b_2) \in E \cap B_1 \times B_2$ and observe that $b_1, b_2 \in (E \cup \Delta_X)(b_2)$ and $B_1 \cup B_2 \subseteq (B_1 \times B_1 \cup B_2 \times B_2)(E \cup \Delta_X)(b_2)$. Hence $B_1 \cup B_2$ is bounded. \Box **Definition 1.8** Let X be a set and \mathcal{M} a collection of subsets of $X \times X$. Since any intersection of coarse structures on X is itself a coarse structure, we can make the following definition. By $\mathfrak{S}(\mathcal{M})$ we denote the smallest coarse structure containing \mathcal{M} , i.e. the intersection of all coarse structures containing \mathcal{M} . We call $\mathfrak{S}(\mathcal{M})$ the *coarse structure generated by* \mathcal{M} . In the same way, define the connected coarse structure generated by \mathcal{M} and

denote it by $_{cn}$ CS (\mathcal{M}) .

Remark 1.9 There is an explicit description of the coarse structure $\mathfrak{S}(\mathcal{M})$ generated by \mathcal{M} . Set $\mathcal{M}_0 := \mathcal{M} \cup \{\Delta_X\}$ and for $n \in \mathbb{N}$ define \mathcal{M}_n inductively as follows.

$$\mathcal{M}_{n+1} := \{ M_1 \mid M_2 \in \mathcal{M}_n, M_1 \subseteq M_2 \} \cup \{ M^{-1} \mid M \in \mathcal{M}_n \} \\ \cup \{ M_1 \cup M_2 \mid M_1, M_2 \in \mathcal{M}_n \} \cup \{ M_1 M_2 \mid M_1, M_2 \in \mathcal{M}_n \}$$

Obviously, $\mathcal{M}_{\infty} := \bigcup_{n \in \mathbb{N}} \mathcal{M}_n$ is contained in $\mathfrak{S}(\mathcal{M})$ and since \mathcal{M}_{∞} is indeed a coarse structure, we get $\mathfrak{S}(\mathcal{M}) = \mathcal{M}_{\infty}$.

Remark 1.10 Let (X, \mathcal{E}) be a coarse structure and define $\mathcal{E}_{cn} := {}_{cn} \mathfrak{S}(\mathcal{E})$. For $E \in \mathcal{E}_{cn}$ there are $E' \in \mathcal{E}, k \in \mathbb{N}$ and sets $A_1, \ldots, A_k, B_1, \ldots, B_k \subseteq X$ which are bounded with respect to \mathcal{E} such that

$$E = E' \cup A_1 \times B_1 \cup \cdots \cup A_k \times B_k .$$

Any set which is bounded with respect to \mathcal{E}_{cn} is a finite disjoint union of sets which are bounded with respect to \mathcal{E} .

There are notions of compatibility of coarse structure and topology.

Definition 1.11 Suppose we are given a topological space X. A coarse structure \mathcal{E} on X is said to be *compatible with the topology* if (1) there is a neighborhood of the diagonal Δ_X which is an entourage and (2) the closure of any bounded set is compact.

In [Roe03] and [HR00], the term proper is being used for compatibility of coarse structure and topology. Compatibility of coarse structure and topology in the sense of [Mit01], implies compatibility as defined above.

Example 1.12 (bounded coarse structure)

Let (X, d) be a pseudometric space. Set $\Delta_r := \{(x, y) \in X \times X \mid d(x, y) < r\}$ and define

$$\mathcal{E}_d := \mathbb{C}(\{\Delta_r \mid r > 0\}) = \{E \subseteq X \times X \mid E \subseteq \Delta_r \text{ for some } r > 0\}.$$

It is easy to verify that (X, \mathcal{E}_d) is a connected coarse space compatible with the topology.

Let G be a finitely generated group. Lemma 1.2 tells us that word metrics arising from different finite generating systems induce the same bounded coarse structure \mathcal{E}_G . Thus, in a natural manner, every finitely generated group G is a coarse space.

Example 1.13 (continuously controlled coarse structure)

Let X be a Hausdorff space and \overline{X} a compactification of X, i.e. X is a dense and open subset of the compact set \overline{X} . The collection

$$\mathcal{E}_{\overline{X}} := \{ E \subseteq X \times X \mid \overline{E} \subseteq X \times X \cup \Delta_{\overline{X}} \}$$

of all subsets $E \subseteq X \times X$, whose closure meets the boundary $(\overline{X} \times \overline{X}) \setminus (X \times X)$ only in the diagonal, is a connected coarse structure on X. If \overline{X} is metrisable, the coarse structure $\mathcal{E}_{\overline{X}}$ is compatible with the topology. The proof is not hard. Compare [Roe03].

Definition 1.14 (close maps) Let (X, \mathcal{E}) be a coarse space and S a set. The maps $f: S \to X$ and $g: S \to X$ are called *close* if $\{(f(s), g(s)) | s \in S)\}$ is an entourage.

Compare with Definition 1.3.

Definition 1.15 Let X, Y be coarse spaces and $f: X \to Y$ a map.

- We call *f* coarsely proper if the inverse image of bounded sets is bounded.
- We call f coarsely uniform if the image of each entourage under the map $f \times f : X \times X \to Y \times Y$ is an entourage.
- We call f a *coarse map* if it is coarsely proper and coarsely uniform.
- We call f a coarse embedding if f is coarsely uniform and the inverse image of an entourage under $f \times f$ is an entourage.²

Note that a coarse embedding is a coarse map.

We will denote the category of coarse spaces and coarse maps by C. This is the category used in coarse algebraic topology. Compare [Roe96], [Roe03] and [Mit01].

For some constructions it turns out to be more convenient to work in the category of coarse spaces and coarsely uniform maps. We will denote this category by \mathcal{D} . When referring only to connected coarse structures, we will write \mathcal{C}_{cn} and \mathcal{D}_{cn} respectively. Observe that there are some forgetful functors between these categories.

 $^{^{2}}$ In [Roe03], coarse embeddings are also called rough maps.

A difference between the categories C and D is that in D products and direct limits exist, while in C they do not. We will deal with products, direct limits, etc. in Section 1.3.

Definition 1.16 Let X, Y be coarse spaces and $f: X \to Y$ a map.

- f is called a \mathcal{D} -equivalence if f is coarsly uniform and there exists a coarsely uniform map $g: Y \to X$ such that $g \circ f$ is close to id_X and $f \circ g$ is close to id_Y .
- f is called a *C*-equivalence if f is coarse and there exists a coarse map $g: Y \to X$ such that $g \circ f$ is close to id_X and $f \circ g$ is close to id_Y .

Proposition 1.17 Any C-equivalence is also a D-equivalence. Conversely, any D-equivalence f is a coarse embedding and hence a C-equivalence.

Proof. Obviously, any C-equivalence is a D-equivalence.

Let \mathcal{E} and \mathcal{E}' be the coarse structures for X and Y respectively. We prove that a \mathcal{D} -equivalence $f: X \to Y$ is a coarse embedding. Let $g: Y \to X$ be a coarsely uniform map as in Definition 1.16. Take $M' \in \mathcal{E}'$ and set $M := g \times g(M')$. Define $E := \{(x, g \circ f(x)) \mid x \in X\} \in \mathcal{E}$. It is sufficient to prove $(f \times f)^{-1}(M') \subseteq EME^{-1}$. Let $(x_1, x_2) \in (f \times f)^{-1}(M')$. With $y_i := f(x_i)$ we have $(y_1, y_2) \in M'$. Set $\tilde{x}_i := g(y_i) = g \circ f(x_i)$. Using $(x_i, \tilde{x}_i) \in E$ and $(\tilde{x}_1, \tilde{x}_2) \in M$, we get $(x_1, x_2) \in EME^{-1}$. \Box

Definition 1.18 (coarse equivalence)

We call $f: X \to Y$ a coarse equivalence if f is a *C*-equivalence. A map $g: Y \to X$ as in Definition 1.16 is called a *coarse inverse* of f. We say that X and Y are *coarsely equivalent* if there exists a coarse equivalence from X to Y.

Proposition 1.19 Let $f: X \to Y$ be a map and \mathcal{M} a collection of subsets of $X \times X$. If \mathcal{E} is a coarse structure on Y and $f(M) \in \mathcal{E}$ for all $M \in \mathcal{M}$, then $f: (X, \mathfrak{S}(\mathcal{M})) \to (Y, \mathcal{E})$ is coarsely uniform.

Proof. The proposition follows easily from Remark 1.9 and the fact that $f \times f(E_1 E_2) \subseteq f \times f(E_1) \ f \times f(E_2)$ for all $E_1, E_2 \subseteq X \times X$.

Proposition 1.20 Let (X, \mathcal{E}_X) and (Y, \mathcal{E}_Y) be coarse spaces and suppose that $f_1: X \to Y$ and $f_2: X \to Y$ are close maps. If f_1 is coarsely proper, coarsely uniform or coarse, the same is true for f_2 . If f_1 is a coarse embedding or a coarse equivalence, then so is f_2 .

Proof. Define $M := \{(f_1(x), f_2(x)) \mid x \in X\}$ and note that $M \in \mathcal{E}_Y$. Observe that $f_2 \times f_2(E) \subseteq M^{-1}(f_1 \times f_1(E))M$ for $E \in \mathcal{E}_X$. This implies the statement about coarsely uniform maps. For a bounded set $B \subseteq Y$ we notice that $f_2^{-1}(B)$ is contained in the bounded set $f_1^{-1}(M[B])$. Hence, the claim about coarsely proper maps follows. For $E \in \mathcal{E}_Y$ we have $(f_2 \times f_2)^{-1}(E) \subseteq (f_1 \times f_1)^{-1}(MEM^{-1})$. This implies

For $E \in \mathcal{E}_Y$ we have $(f_2 \times f_2)^{-1}(E) \subseteq (f_1 \times f_1)^{-1}(MEM^{-1})$. This implies the claim about coarse embeddings.

Suppose that f_1 is a coarse equivalence and $g: Y \to X$ a coarse inverse of f_1 . Observe that $f_1 \circ g$ and $f_2 \circ g$ are close maps. Since g is coarsely uniform, we obtain that $g \circ f_1$ and $g \circ f_2$ are close maps. Since "being close" has the same properties as an equivalence relation, it follows that g is also a coarse inverse of f_2 .

1.3 Some constructions in the coarse category

Pull-back of a coarse structure

Let (X, \mathcal{E}) be a coarse space, A an arbitrary set and $f: A \to X$ any map.

Definition 1.21 We call $f^*(\mathcal{E}) := \mathbb{C}\left(\left\{(f \times f)^{-1}(E) \mid E \in \mathcal{E}\right\}\right)$ the *pullback* of \mathcal{E} by f^{3} .

Clearly, $f^*(\mathcal{E})$ is a coarse structure on A such that f is a coarse embedding. If (X, \mathcal{E}) is connected, the same is true for $(A, f^*(\mathcal{E}))$.

Suppose f is a continuous and proper map between topological spaces. If \mathcal{E} is compatible with the topology, the same is true for $f^*(\mathcal{E})$.

Definition 1.22 (restriction of a coarse structure to a subset)

If $A \subseteq X$ and f is the inclusion map, we call $f^*(\mathcal{E})$ the *restriction* of \mathcal{E} to A and write $\mathcal{E}|_A$.

Unions of coarse spaces

Let I be a set, (X_i, \mathcal{E}_i) a coarse space for every $i \in I$ and $X = \bigcup_{i \in I} X_i$. Note that the sets X_i for $i \in I$ are not supposed to be disjoint.

Definition 1.23 We call $\bigvee_{i \in I} \mathcal{E}_i := \mathfrak{S}(\bigcup_{i \in I} \mathcal{E}_i)$ the minimal coarse structure on the union of (X_i, \mathcal{E}_i) .

Remark 1.24 If we are given a coarse structure \mathcal{E} on X such that for all $i \in I$ its restriction to X_i is \mathcal{E}_i , then $\bigvee_{i \in I} \mathcal{E}_i \subseteq \mathcal{E}$.

³Since the collection $\{(f \times f)^{-1}(E) \mid E \in \mathcal{E}\}$ is not necessarily closed under taking subsets, we need to take the coarse structure generated by this collection. Note that this is the only reason for putting \mathfrak{G} here.

Direct limits of coarse spaces

For information about category theory compare [ML98]. Let I be a small category and let $F: I \to \mathcal{D}$ or $F: I \to \mathcal{D}_{cn}$ be a functor. In particular, we are given a coarse space (X_i, \mathcal{E}_i) for each $i \in Obj(I)$ and a coarsely uniform map $F(h): (X_i, \mathcal{E}_i) \to (X_j, \mathcal{E}_j)$ for each $h \in Mor(i, j)$.

Proposition 1.25 (existence of direct limits in \mathcal{D} and \mathcal{D}_{cn}) The categories \mathcal{D} and \mathcal{D}_{cn} are co-complete, i.e. F has a colimit. More precisely,

$$\underset{\longrightarrow}{\lim} (X_i, \mathcal{E}_i) = \left(\underset{\longrightarrow}{\lim} X_i , \mathcal{E}_{\underset{\longrightarrow}{\lim}}\right)$$

where $\varinjlim X_i$ is the corresponding colimit in the category of sets and \mathcal{E}_{\varinjlim} is the intersection of all (connected) coarse structures such that the maps $f_i: X_i \to \varinjlim X_i$ are coarsely uniform for all $i \in Obj(I)$.

Proof. Observe that $\mathcal{E}_{\lim} = \mathbb{C}\left(\bigcup_{i \in Obj(I)} f_i(\mathcal{E}_i)\right)$. With Proposition 1.19 the claim follows easily from the definitions.

The analogue of Proposition 1.25 in C and C_{cn} is not true. See Example 1.26. However, coproducts do exist in the categories C and C_{cn} .

Example 1.26 For each $i \in \mathbb{N}$ take $X_i := \mathbb{R}_+$ with the usual bounded coarse strucure and define $f_{ji}: X_i \to X_j, x \mapsto \max\{0, x-j+i\}$ for $i \leq j$. Then $\lim X_i$ is just a point.

In some cases we can give a more explicit description for the coarse structure of the direct limit.

Proposition 1.27 Let *I* be a directed set⁴. If for all $i \in I$ the map $f_i \colon X_i \to \lim X_i$ is injective, then

$$\mathcal{E}_{\lim} = \{ D \cup f_j \times f_j(L) \mid j \in I, L \in \mathcal{E}_j, D \subseteq \Delta_{\lim X_i} \} .$$

Proof. For $j \in I$, $L \in \mathcal{E}_j$ and $D \subseteq \Delta_{\lim X_i}$ the set $D \cup f_j \times f_j(L)$ is in \mathcal{E}_{\lim} . It remains to prove that $\mathcal{E}_c := \{D \cup f_j \times f_j(L) \mid j \in I, L \in \mathcal{E}_j, D \subseteq \Delta_{\lim X_i}\}$ is a coarse structure. In order to conclude that \mathcal{E}_c is closed under composition of elements, we need the maps f_i to be injective. \Box

Proposition 1.28 Suppose that the identity morphisms are the only morphisms in the category *I*. In this case the colimit is just the coproduct and we denote the coarse structure of the direct limit by \mathcal{E}_{\coprod} . We get $\mathcal{E}_{\coprod} = \{D \cup E_{i_1} \cup \cdots \cup E_{i_k} \mid k \in \mathbb{N}, i_1, \ldots, i_k \in Obj(I), E_{i_j} \in \mathcal{E}_{i_j}, D \subseteq \Delta_{\coprod X_i}\}.$

⁴Compare [Spa95] for the definition of a directed set and the definition of direct limits in this special case.

Infinite disjoint unions of coarse structures

Example 1.29 \mathbb{R} and \mathbb{Z} (both equipped with their usual bounded coarse strucure) are coarsely equivalent, but the corresponding countable coproducts $\prod_{\mathbb{N}} \mathbb{R}$ and $\prod_{\mathbb{N}} \mathbb{Z}$ are not.

Proof. Any map $\coprod_{\mathbb{N}} \mathbb{R} \to \coprod_{\mathbb{N}} \mathbb{Z} \to \coprod_{\mathbb{N}} \mathbb{R}$ will be different from the identity on any copy of \mathbb{R} . Thus, no such map can be close to $id_{\coprod_{\mathbb{N}}}\mathbb{R}$, since any entourage of the coproduct involves only off-diagonal elements of finitely many copies of \mathbb{R} .

Because of the above example one might look for another more geometric coarse structure on the coproduct.

Definition 1.30 (disjoint union of coarse spaces)

Let I be a set and assume for any $i \in I$ we are given a coarse space (X_i, \mathcal{E}_i) . Set $X := \bigsqcup_{i \in I} X_i$. We define

$$\mathcal{E}_{\bigsqcup} := \left\{ A \subseteq \bigsqcup_{i \in I} X_i^2 \mid A \cap X_i^2 \in \mathcal{E}_i \right\}.$$

The underlying sets of coproduct and disjoint union are the same. We just take different coarse structures. For any finite set I, coproduct and disjoint union coincide.

Proposition 1.31 If for every $i \in I$ the coarse spaces (X_i, \mathcal{E}_i) and (Y_i, \mathcal{F}_i) are coarsely equivalent, then $\bigsqcup_{i \in I} (X_i, \mathcal{E}_i)$ and $\bigsqcup_{i \in I} (Y_i, \mathcal{F}_i)$ are also coarsely equivalent.

Proof. If a coarse equivalence $f_i: (X_i, \mathcal{E}_i) \to (Y_i, \mathcal{F}_i)$ is given for all $i \in I$, then $\bigsqcup_{i \in I} f_i: \bigsqcup_{i \in I} (X_i, \mathcal{E}_i) \to \bigsqcup_{i \in I} (Y_i, \mathcal{F}_i)$ is also a coarse equivalence. \Box

Products of coarse spaces

Definition 1.32 For $i \in \{1, \ldots, k\}$ let (X_i, \mathcal{E}_i) be a coarse space. By $p_i: X_1 \times \cdots \times X_k \to X_i$ we denote the projection to the *i*-th factor. The product coarse structure is defined as follows.

$$\mathcal{E}_1 * \cdots * \mathcal{E}_k := \left\{ E \subseteq (X_1 \times \cdots \times X_k)^2 \mid p_i \times p_i(E) \in \mathcal{E}_i \text{ for } i \in \{1, \dots, k\} \right\}$$

If (X, \mathcal{E}) is a coarse space, we will sometimes write \mathcal{E}^{*k} for the product coarse structure on X^k .

It is easy to prove that $\mathcal{E}_1 * \cdots * \mathcal{E}_k$ actually is a coarse structure. The product coarse structure $\mathcal{E}_1 * \cdots * \mathcal{E}_k$ is connected if and only if the coarse structures \mathcal{E}_i are connected. Moreover, we have the following formulas:

$$\Delta_{X_1 \times \dots \times X_k} = \Delta_{X_1} \times \dots \times \Delta_{X_k}$$
$$(E_1 \times \dots \times E_k)(E'_1 \times \dots \times E'_k) = E_1 E'_1 \times \dots \times E_k E'_k$$

One remark on our notation: If $E \subseteq X \times X$, we should not confuse the composition $E^k = E \cdots E$ and the product $E^{\times k} = E \times \cdots \times E$.

Remark 1.33 (compatibility with other products)

• Let (X, d_X) and (Y, d_Y) be metric spaces. Then the formula

 $d((x_1, y_1), (x_2, y_2)) := d_X(x_1, x_2) + d_Y(y_1, y_2)$

defines a metric on $X \times Y$. It is easy to see that $\mathcal{E}_d = \mathcal{E}_{d_X} * \mathcal{E}_{d_Y}$.

• Let X and Y be Hausdorff spaces with compactifications \overline{X} and \overline{Y} respectively. Then $\overline{X} \times \overline{Y}$ is a compactification of $X \times Y$. By $\mathcal{E}_{\overline{X}}$, $\mathcal{E}_{\overline{Y}}$ and $\mathcal{E}_{\overline{X} \times \overline{Y}}$ we denote the coarse structures on X, Y and $X \times Y$ induced by \overline{X} , \overline{Y} and $\overline{X} \times \overline{Y}$ respectively. The definitions easily imply $\mathcal{E}_{\overline{X} \times \overline{Y}} = \mathcal{E}_{\overline{X}} * \mathcal{E}_{\overline{Y}}$.

Remark 1.34 (compatibility with coarse equivalence)

Let (X, \mathcal{E}) , (X', \mathcal{E}') and (Y, \mathcal{F}) be coarse spaces and assume (X, \mathcal{E}) and (X', \mathcal{E}') to be coarsely equivalent. Then the spaces $(X \times Y, \mathcal{E} * \mathcal{F})$ and $(X' \times Y, \mathcal{E}' * \mathcal{F})$ are coarsely equivalent.

Quotients of coarse spaces

Definition 1.35 Let (X, \mathcal{E}) be a coarse space. Let \sim be an equivalence relation on X and $\pi: X \to X/\sim$ the corresponding quotient map. We define a coarse structure on X/\sim by

$$\mathcal{E}_{\sim} := \mathbb{C}\left\{ \left\{ \pi \times \pi(E) \mid E \in \mathcal{E} \right\} \right\} .$$

If \mathcal{E} is connected, \mathcal{E}_{\sim} is connected. The map π is coarsely uniform by definition of the coarse structure on the quotient. If there is an unbounded equivalence class, then π is not coarsely proper.

Attaching coarse spaces

Let (X, \mathcal{E}_X) , (Y, \mathcal{E}_Y) be coarse spaces, $A \subseteq X$ and $f: A \to Y$ coarsely uniform. Attaching X to Y using f, we get $X \cup_f Y := X \sqcup Y / \sim$ and a quotient map $\pi: X \sqcup Y \to X \cup_f Y$. Here \sim is the equivalence relation generated by $a \sim f(a)$ for all $a \in A$. The map π is coarsely uniform, but not necessarily coarsely proper. We denote the coarse structure on the quotient by $\mathcal{E}_{X \cup_f Y}$.

Lemma 1.36 $\mathcal{E}_Y = \mathcal{E}_{X \cup_f Y}|_Y$

Proof. We certainly have $\mathcal{E}_Y \subseteq \mathcal{E}_{X \cup_f Y}|_Y$ and need to prove the converse. Let $L \in \mathcal{E}_{X \cup_f Y}|_Y$. There are $L'_{X,1}, \ldots, L'_{X,k} \in \mathcal{E}_X$ and $L_{Y,1}, \ldots, L_{Y,k} \in \mathcal{E}_Y$, such that

$$L \subseteq \pi \times \pi(L'_{X,1} \cup L_{Y,1}) \cdots \pi \times \pi(L'_{X,k} \cup L_{Y,k}).$$

Define $L_{X,i} \cdots L_{X,i+j} := f \times f(L'_{X,i} \cdots L'_{X,i+j}) \in \mathcal{E}_Y$. This is a partial replacement of the notation of composition which we will apply only in this proof. We conclude that

$$L \subseteq \bigcup_{(\sigma_1,\dots,\sigma_k)\in\{X,Y\}^n} L_{\sigma_1,1}\cdots L_{\sigma_k,k}\in\mathcal{E}_Y .$$

Since $\pi|_{X\setminus A}$ is coarsely uniform, we have $\mathcal{E}_X|_{X\setminus A} \subseteq \mathcal{E}_{X\cup_f Y}|_{X\setminus A}$. But in general these coarse structures do not coincide as the following example shows.

Example 1.37 Let Y be a single point, X' any coarse space such that $X' \times X'$ is not an entourage, $X = X' \times \mathbb{R}_+$ and $A = X' \times \{0\}$. For r > 0 the set $\{((x,t), (\tilde{x}, \tilde{t})) \mid t, \tilde{t} \in (0, r], x, \tilde{x} \in X'\}$ is contained in $\mathcal{E}_{X \cup_f Y}|_{X \setminus A}$ but not in $\mathcal{E}_X|_{X \setminus A}$.

Remark 1.38 If the attaching map f is a coarse embedding, we have $\mathcal{E}_X|_{X\setminus A} = \mathcal{E}_{X\cup fY}|_{X\setminus A}$.

Proof. The map $\pi|_X$ is a coarse equivalence and $\pi|_{X\setminus A}$ is a canonical isomorphism.

1.4 Coarse structures induced by metrisable compactifications

We will briefly recall the definitions of the Higson compactification and the Higson corona. For more detailed explanations see [Roe93], [HR00] or [Roe03].

Let X be a Hausdorff space with a coarse structure \mathcal{E} such that the closure of every bounded set is compact. A bounded continuous function f on X is called a *Higson function* with respect to a coarse structure \mathcal{E} if for all entourages $E \in \mathcal{E}$ the function

$$\mathbf{d}f: X \times X \to \mathbb{C}, \quad (x_1, x_2) \mapsto f(x_1) - f(x_2)$$

restricted to E tends to zero at infinity. The Higson functions form a unital C^{*}-algebra $C_h(X)$. It follows from the Gelfand-Naimark theorem that $C_h(X)$ is the algebra of bounded, continuous functions on a compactification hX of X. We call hX the Higson compactification of (X, \mathcal{E}) . The Higson corona νX is defined to be $hX \setminus X$.

For metrisable compactifications the relation between compactifications and coarse structures is especially simple:

Proposition 1.39 Let X be a Hausdorff space and K a metrisable compactification. The Higson compactification of the continuously controlled coarse structure on X induced by K is homeomorphic to K.

Proof. This is Proposition 2.48 of [Roe03].

Let X be a non-compact Hausdorff space and hX a metrisable compactification of X. By \mathcal{E}_{hX} denote the continuously controlled coarse structure induced by hX.

Choose a metric d on hX and define

$$X_i := \left\{ x \in X \mid d(x, \nu X) \ge \frac{1}{i} \right\}$$

for $i \in \mathbb{N}$ $(X_0 := \emptyset)$. We have defined a sequence of compact subsets of X such that $X_0 \subseteq X_1 \subseteq X_2 \subseteq \cdots \subseteq X$ and $\bigcup_{i \in \mathbb{N}} X_i = X$. Moreover $\{hX \setminus X_i\}_{i \in \mathbb{N}}$ is a basis of neighborhoods for $\nu X := hX \setminus X$.

Lemma 1.40 For $\rho > 0$ define $\Delta_{\rho} := \{(x, y) \in X^2 \mid d(x, y) < \rho\}$. We assert that $E \in \mathcal{E}_{hX}$ if and only if there is a sequence $\{\rho_i\}_{i \in \mathbb{N}}$ of non-negative real numbers converging to zero such that $E \setminus X_i^2 \subseteq \Delta_{\rho_i}$ for all $i \in \mathbb{N}$.

The proof of this lemma is straightforward.

Our intention is to construct a coarse structure \mathcal{E} on $\nu X \times \mathbb{N}$ such that (X, \mathcal{E}_{hX}) is coarsely equivalent to $(\nu X \times \mathbb{N}, \mathcal{E})$. We may assume that $X_1 \neq \emptyset$. Choose a map

$$f: \nu X \times \mathbb{N} \to X \tag{1.1}$$

such that $f(\overline{x}, n) \in X_n$ has minimal distance to \overline{x} among all points in X_n . Since X_n is compact, this definition makes sense. Next, we define a coarse structure on $\nu X \times \mathbb{N}$ by pulling back the coarse structure \mathcal{E}_{hX} . Thus f becomes a coarse embedding.

$$\mathcal{E} := f^*(\mathcal{E}_{hX})$$

By $\pi_{\mathbb{N}} : \nu X \times \mathbb{N} \to \mathbb{N}$ we denote the projection onto \mathbb{N} and by $\mathcal{E}_{\mathbb{N}}$ the continuously controlled coarse structure on \mathbb{N} induced by the one-point compactification. Observe that $\mathcal{E}_{\mathbb{N}}$ is the maximal coarse structure which is compatible with the topology.

Lemma 1.41 We assert that $E \in \mathcal{E}$ if and only if the following two conditions are satisfied.

- (a) $\pi_{\mathbb{N}} \times \pi_{\mathbb{N}}(E) \in \mathcal{E}_{\mathbb{N}}$
- (b) There is a sequence $\{\delta_k\}_{k\in\mathbb{N}}$ of non-negative real numbers converging to zero such that $d(\overline{x},\overline{y}) < \delta_k$ for all $((\overline{x},i),(\overline{y},j)) \in E$ with $\max\{i, j\} > k.$

Proof. Define

$$a_i := \max\{d(X_i, \overline{x}) \mid \overline{x} \in \nu X\}$$

for $i \in \mathbb{N}$. The sequence $\{a_i\}_{i \in \mathbb{N}}$ is monotone decreasing, since $X_i \subseteq X_{i+1}$ for all $i \in \mathbb{N}$. We claim that this sequence converges to zero.

Assume $a_i > a > 0$ for all $i \in \mathbb{N}$. Take the centers of a finite cover of νX with balls of radius $\frac{a}{3}$. For each of these points there is a point in X within distance less than $\frac{a}{3}$. These finitely many points in X form a compact set and hence are contained in some X_k . We get $a_k \leq \frac{2}{3}a$, which contradicts our assumption. Hence $\lim_{i \to \infty} a_i = 0.$

Define

$$n_i := \max\left\{n \in \mathbb{N} \mid a_i < \frac{1}{n}\right\}$$

for all *i* with $a_i < 1$. The equation $\lim_{i \to \infty} a_i = 0$ implies $\lim_{i \to \infty} n_i = \infty$. For $M \subseteq \mathbb{N} \times \mathbb{N}$ we define a completion M^{\square} of M in the following way. Set $M^{\square} := \{ (u,v) \in \mathbb{N}^2 \mid (x,y) \in M \text{ and } (x \le u \le v \le y \text{ or } y \le v \le u \le x) \}.$ Let $E \subseteq (\nu X \times \mathbb{N})^2$ and define

$$b_i := \min\left\{n \in \mathbb{N} \mid (n, i) \in \pi_{\mathbb{N}} \times \pi_{\mathbb{N}} \left(E \cup E^{-1}\right)^{\bowtie}\right\}.$$

Suppose that E satisfies conditions (a) and (b). First we will prove that $f \times f(E) \in \mathcal{E}_{hX}$. Since $\mathcal{E}_{\mathbb{N}}$ is compatible with the topology, condition (a) implies $\lim_{i \to \infty} b_i = \infty$. We define a sequence $\{\rho_i\}_{i \in \mathbb{N}}$.

$$\rho_i := a_i + a_{b_i} + \delta_i$$

Certainly we have $\lim_{i\to\infty} \rho_i = 0.$

Let $((\overline{x}, n), (\overline{y}, m)) \in E$ and define $x := f(\overline{x}, n)$ and $y := f(\overline{y}, m)$. If $x \notin X_i$, then $d(x,\overline{x}) \leq a_i, d(y,\overline{y}) \leq a_{b_i}$ and $d(\overline{x},\overline{y}) \leq \delta_i$. Hence $f \times f(E) \setminus X_i^2 \subseteq \Delta_{\rho_i}$. Now we can apply Lemma 1.40 to conclude $f \times f(E) \in \mathcal{E}_{hX}$. It follows $E \subseteq (f \times f)^{-1}(f \times f(E)) \in \mathcal{E}.$

Suppose $E \in \mathcal{E}$. There is an entourage $E' \in \mathcal{E}_{hX}$ with $E \subseteq (f \times f)^{-1}(E')$. For all $i \in \mathbb{N}$ there is $m_i \in \mathbb{N}$ such that $a_{m_i} < \frac{1}{i}$. Since $\frac{1}{m_i} \leq a_{m_i}$, we get $m_i > i$ for all $i \in \mathbb{N}$. We may assume that the sequence $\{m_i\}_{i \in \mathbb{N}}$ is monotone.

Since X_i is bounded, $E'[X_i]$ is bounded, i.e. $E'[X_i]$ is contained in some X_j . Note that the image of $\nu X \times \{0, \ldots, i\}$ under f is contained in X_i . Observe that $E[\nu X \times \{0, \ldots, i\}] \subseteq f^{-1}(E'[X_i]) \subseteq f^{-1}(X_j)$. It follows that $\pi_{\mathbb{N}}(E[\nu X \times \{0, \ldots, i\}]) \subseteq \pi_{\mathbb{N}}(f^{-1}(X_j)) \subseteq \{0, \ldots, m_j - 1\}$. Together with an analog fact for E^{-1} this implies that E satisfies condition (a). Choose a sequence $\{\rho_i\}_{i\in\mathbb{N}}$ converging to zero and such that $E' \setminus X_i^2 \subseteq \Delta_{\rho_i}$ for all $i \in \mathbb{N}$. Let $((\overline{x}, i), (\overline{y}, j)) \in E$ with $\max\{i, j\} > k$. Assume $i \geq k$. It follows that $j \geq b_k$. We already proved $\pi_{\mathbb{N}} \times \pi_{\mathbb{N}}(E) \in \mathcal{E}_{\mathbb{N}}$. Hence $\lim_{i \to \infty} b_i = \infty$. Define $x := f(\overline{x}, i)$ and $y := f(\overline{y}, j)$. We have $d(x, \overline{x}) \leq a_i \leq a_k < \frac{1}{n_k}$, i.e. $x \notin X_{n_k}$. Therefore $d(x, y) < \rho_{n_k}$. Note that

$$\delta_k := a_k + a_{b_k} + \rho_{n_k} \xrightarrow[k \to \infty]{} 0 .$$

Obviously E satisfies (b) with $\{\delta_k\}_{k\in\mathbb{N}}$ as defined above.

We take a map

$$g\colon X \to \nu X \times \mathbb{N} \tag{1.2}$$

such that for $x \in X_i \setminus X_{i-1}$ the image $g(x) = (\overline{x}, i)$ is chosen in such a way that \overline{x} minimizes the distance to x among all points in νX .

Lemma 1.42 The map g is coarsely uniform.

Proof. Let $E \in \mathcal{E}_{hX}$. Since \mathcal{E}_{hX} is compatible with the topology, $g \times g(E)$ satisfies condition (a) of Lemma 1.41. Let $\{\rho_i\}_{i\in\mathbb{N}}$ be a sequence as in Lemma 1.40 for the entourage E and define $b_i := \min\{n \in \mathbb{N} \mid E[X_i \setminus X_{i-1}] \cap X_n \neq \emptyset\}$. This implies $\lim_{i \to \infty} b_i = \infty$. Defining $\delta_i := \frac{1}{i-1} + \frac{1}{b_i-1} + \rho_{i-1}$ we see that $g \times g(E)$ satisfies condition (b) of Lemma 1.41. \Box

Theorem 1.43 Any map f chosen as in (1.1) is a coarse equivalence and any map g chosen as in (1.2) is a coarse inverse of f.

Proof. We will use some of the notation from the proof of Lemma 1.41. We first consider the map $g \circ f : \nu X \times \mathbb{N} \to \nu X \times \mathbb{N}$. Let $(x,k) \in \nu X \times \mathbb{N}$. Observe that $d(f(x,k),x) \leq d(X_k,x) \leq a_k < \frac{1}{n_k}$. Hence $f(x,k) \in X_k \setminus X_{n_k}$. Defining $(\tilde{x}, \tilde{k}) := g \circ f(x,k)$ we get $n_k + 1 \leq \tilde{k} \leq k$. Moreover $d(x, \tilde{x}) \leq 2a_k$. It follows that conditions (a) and (b) of Lemma 1.41 are satisfied for $\{((x,k),g \circ f(x,k)) \mid x \in \nu X \times \mathbb{N}\}$.

Now we consider the map $f \circ g \colon X \to X$. Let $x \in X_i \setminus X_{i-1}$ and $(\overline{x}, i) := g(x)$. Then $d(x, \overline{x}) < \frac{1}{i-1}$. Thus $d(f \circ g(x), x) \leq \frac{2}{i-1}$, i.e. $f \circ g$ is close to id_X . \Box

Corollary 1.44 Let X and Y be Hausdorff spaces with metrisable compactifications hX and hY respectively. The coarse spaces (X, \mathcal{E}_{hX}) and (Y, \mathcal{E}_{hY}) are coarsely equivalent if and only if the coronas νX and νY are homeomorphic. **Proof.** To see that the Higson coronas of coarsely equivalent spaces are homeomorphic, we refer to Corollary 2.42 of [Roe03]. \Box

To every compact Hausdorff space K we assign a coarse space $\psi(K)$ as follows.

$$\psi(K) := \left(K \times [0,1), \mathcal{E}_{K \times [0,1]}\right)$$

Corollary 1.45 ψ induces a bijection Ψ between homeomorphism classes of metrisable compact spaces⁵ and coarse equivalence classes of spaces whose coarse structure is induced by a metrisable compactification. The inverse of Ψ is induced by the correspondence ν assigning to each coarse space its Higson corona.

We define a "product" of two compact spaces A and B as follows.

$$\begin{split} A \boxtimes B &:= A \times B \times [0,1] \cong A \times \{1\} \times B \times [0,1] \cup A \times [0,1] \times B \times \{1\} \\ &= A \times [0,1] \times B \times [0,1] \setminus A \times [0,1) \times B \times [0,1) \end{split}$$

Here \cong stands for homeomorphism.

Proposition 1.46 If A and B are metrisable compact spaces, then

$$\Psi(A\boxtimes B)=\Psi(A)\times\Psi(B).$$

Here \times denotes the product of coarse spaces as in Definition 1.32.

Proof. Because of Remark 1.34, the right hand side of the equation makes sense. The Higson corona of $\psi(A) \times \psi(B) = A \times [0, 1) \times B \times [0, 1)$ is homeomorphic to $A \boxtimes B$. Thus, Corollary 1.45 implies $\Psi(A \boxtimes B) = \Psi(A) \times \Psi(B)$.

Example 1.47 By \mathcal{E} , we denote the coarse structure on \mathbb{R}_+ induced by the one-point compactification and by \mathcal{E}_{vis} the coarse structure induced on \mathbb{R}^n by the visual corona S^{n-1} . The usual bounded coarse structure on \mathbb{R}^n is contained in \mathcal{E}_{vis} . Proposition 1.46 implies that $(\mathbb{R}_+, \mathcal{E}_.)^n$ and $(\mathbb{R}_+^n, \mathcal{E}_{vis}|_{\mathbb{R}_+^n})$ are coarsely equivalent.

⁵Since every metrisable compact set is homeomorphic to a subset of $[0, 1]^{\mathbb{N}}$, the homeomorphism classes of metrisable compact spaces form a set.

Chapter 2

Translation C^* -algebras via projection-valued measures

To any coarse space (X, \mathcal{E}) which admits a countable, uniformly bounded cover, we will assign translation C^* -algebras $E^*(X, \mathcal{E})$ and $C^*(X, \mathcal{E})$.

This may be considered as an analog of assigning to a locally compact space X the algebra $C_0(X)$ of complex valued, continuous function vanishing at infinity.

In [HR00], a representation of the algebra $C_0(X)$ is used to define translation C^* -algebras for coarse spaces which are at the same time nice topological spaces such that topology and coarse structure are compatible. John Roe proposed in [Roe03] to use projection-valued measures in order to define translation C^* -algebras for more general coarse spaces. In this chapter we give a more complete exposition on the subject and eliminate some mistakes. The idea is to generalize an ad-hoc approach which has been used for defining translation C^* -algebras of discrete coarse spaces.

We prove that the translation C^* -algebras are well defined up to isomorphism of C^* -algebras, we compare the different approaches of defining translation C^* -algebras and we discuss the matter whether a coarse map induces maps between the translation C^* -algebras of the corresponding coarse spaces.

2.1 Geometric Hilbert spaces and the calculus of supports

Let (X, \mathcal{E}) be a coarse space.

Definition 2.1 A family \mathcal{U} of subsets of X is called a *localizing decompo*sition of (X, \mathcal{E}) , if (1) \mathcal{U} is a decomposition, i.e. a cover of X consisting of pairwise disjoint sets, (2) \mathcal{U} is countable and (3) \mathcal{U} is uniformly bounded, i.e. $\Delta_{\mathcal{U}} := \bigcup_{U \in \mathcal{U}} U \times U \in \mathcal{E}$. **Remark 2.2** Let (X, \mathcal{E}) be a coarse space and \mathcal{U} a localizing decomposition. We say that $x, y \in X$ are \mathcal{U} -equivalent if they are contained in the same $U \in \mathcal{U}$. We denote the quotient of X with respect to \mathcal{U} -equivalence by $X/_{\mathcal{U}}$. Using the quotient map $\pi \colon X \to X/_{\mathcal{U}}$, we define the coarse structure $\mathcal{E}/_{\mathcal{U}} := \{\pi \times \pi(E) \mid E \in \mathcal{E}\}$ on $X/_{\mathcal{U}}$.

In this sence, a localizing decomposition \mathcal{U} induces a countable "discretization" $(X/_{\mathcal{U}}, \mathcal{E}/_{\mathcal{U}})$ which is coarsely equivalent to (X, \mathcal{E}) .

Let \mathfrak{A} be a σ -algebra over X.¹

Definition 2.3 A localizing σ -algebra over (X, \mathcal{E}) is a σ -algebra \mathfrak{A} together with a topology on X such that for each entourage $E \in \mathcal{E}$ the closure \overline{E} is still an entourage. Moreover, there has to exist a localizing decomposition $\mathcal{U} \subseteq \mathfrak{A}$ such that $\bigcup_{U \in \mathcal{U}} \mathring{U}$ is dense in X. We will call \mathcal{U} an \mathfrak{A} -localizing decomposition.

For each $x \in X$ we define $\mathfrak{A}(x) \subseteq \mathfrak{A}$ to be all the sets $A \in \mathfrak{A}$ whose interior \mathring{A} contains x. Such an $A \in \mathfrak{A}(x)$ will be called an \mathfrak{A} -neighborhood of $x \in X$.

Example 2.4 Given a coarse space (X, \mathcal{E}) which admits a uniformly bounded cover $\mathcal{V} = \{V_n \mid n \in \mathbb{N}\}$, there are several localizing σ -algebras:

- (1) The σ -algebra $\sigma(\mathcal{V})$ generated by \mathcal{V} together with the discrete topology.
- (2) The σ -algebra $\mathfrak{P}(X)$ consisting of all subsets of X together with the discrete topology.
- (3) If X is a topological space and the coarse structure \mathcal{E} is compatible with the topology, then the Borel algebra $\mathfrak{B}(X)$, i.e. the smallest σ -algebra containing all open sets, is localizing.

The σ -algebra $\mathfrak{A} = \{\emptyset, X\}$ is not localizing if X is unbounded.

Proof. Defining $U_n = V_n \setminus (V_0 \cup \cdots \cup V_{n-1})$ we get a localizing decomposition $\mathcal{U} = \{U_n \mid n \in \mathbb{N}\}$ which is contained in the σ -algebras $\sigma(\mathcal{V})$ and $\mathfrak{P}(X)$.

In order to prove (3), note that from \mathcal{V} we can construct a countable, uniformly bounded cover consisting of open sets as follows: If E is an open entourage containing Δ_X , take the cover $\{E[V_n] \mid n \in \mathbb{N}\}$. From this cover we get a localizing decomposition as before.

Definition 2.5 Let \mathfrak{A} be a σ -algebra over X and H a Hilbert space. A projection-valued measure is a map $\lambda \colon \mathfrak{A} \to \mathcal{B}(H)$ with the following properties:

¹A σ -algebra \mathfrak{A} over a set X is a family of subsets of X such that (1) $X \in \mathfrak{A}$, (2) $X \setminus A \in \mathfrak{A}$ whenever $A \in \mathfrak{A}$ and (3) the union of countably many elements of \mathfrak{A} is in \mathfrak{A} . Compare any book on measure theory.

- (1) For every $A \in \mathfrak{A}$ the operator $\lambda(A)$ is a projection, i.e. $\lambda(A)^* = \lambda(A) = \lambda(A)^2$.
- (2) $\lambda(\emptyset) = 0$ and $\lambda(X) = \mathrm{id}_H$.
- (3) If $A_n \in \mathfrak{A}$ for all $n \in \mathbb{N}$ and $A_i \cap A_j = \emptyset$ whenever $i \neq j$, then

$$\lambda\left(\bigcup_{n\in\mathbb{N}}A_n\right)=\sum_{n\in\mathbb{N}}\lambda(A_n).$$

Remark 2.6 Note that the infinite sum in Definition 2.5 has to be considered as a limit with respect to strong convergence.²

The sequence $\left\{\sum_{n=0}^{k} \lambda(A_n)\right\}_{k \in \mathbb{N}}$ does in general not converge in the operator norm.

Lemma 2.7 Let $\lambda : \mathfrak{A} \to \mathcal{B}(H)$ be a projection-valued measure over X. For $A, B \in \mathfrak{A}$ we have $\lambda(A)\lambda(B) = \lambda(B)\lambda(A) = \lambda(A \cap B)$.

Proof. First suppose $A \cap B = \emptyset$. In order to prove $\lambda(A)\lambda(B) = 0$, observe that the images of $\lambda(A)$ and $\lambda(B)$ are orthogonal, because $\lambda(A) + \lambda(B) = \lambda(A \cup B)$ is a projection.

The general case now follows easily.

$$\lambda(A)\lambda(B) = (\lambda(A \setminus B) + \lambda(A \cap B))(\lambda(B \setminus A) + \lambda(B \cap A))$$

= $\lambda(A \cap B)\lambda(B \cap A) = \lambda(A \cap B)$

The first equality uses additivity of projection-valued measures. The last one requires that a projection P is idempotent, i.e. $P^2 = P$.

Example 2.8 Take the Hilbert space $H = L^2(X, \mu)$ where μ is a measure on some localizing σ -algebra \mathfrak{A} . A projection-valued measure $\lambda \colon \mathfrak{A} \to \mathcal{B}(H)$ is given by

$$\lambda(A)f = \chi_A \cdot f$$
 for $A \in \mathfrak{A}$ and $f \in H$.

Here $\chi_A \colon X \to \mathbb{C}$ is the characteristic function of A, i.e. $\chi_A(x) = 1$ if $x \in A$ and $\chi(x) = 0$ if $x \in X \setminus A$.

Example 2.9 Let (X, \mathcal{E}) be a coarse space and $\mathcal{U} = \{U_i \mid i \in \mathbb{N}\}$ a localizing decomposition of X. Define $\mathfrak{A} := \sigma(\mathcal{U})$ to be the σ -algebra generated by \mathcal{U} . Assume we are given a Hilbert space H_i for each $i \in \mathbb{N}$ such that $H_i = \{0\}$ if $U_i = \emptyset$. In this situation we can define a projection-valued measure $\lambda \colon \mathfrak{A} \to \mathfrak{B}\left(\bigoplus_{i \in \mathbb{N}} H_i\right)$ as follows: For $A \in \mathfrak{A}$ define

$$\lambda(A) = \text{ orthogonal projection onto } \bigoplus_{\substack{i \in \mathbb{N} \\ U_i \subset A}} H_i .$$

²A sequence $\{T_i\}_{i\in\mathbb{N}}$ in $\mathcal{B}(H)$ converges strongly to $T \in \mathcal{B}(H)$, if for any $v \in H$ the sequence $\{T_i(v)\}_{i\in\mathbb{N}}$ converges to T(v) in H.

Definition 2.10 Let (X, \mathcal{E}) be a coarse space. A geometric Hilbert space over (X, \mathcal{E}) is a projection-valued measure $\lambda \colon \mathfrak{A} \to \mathcal{B}(H)$ where \mathfrak{A} is a localizing σ -algebra over (X, \mathcal{E}) and H a separable Hilbert space.

Remark 2.11 The projection-valued measure in Example 2.8 is a geometric Hilbert space if and only if $L^2(X, \mu)$ is separable.

The projection-valued measure in Example 2.9 is a geometric Hilbert space if and only if the Hilbert space H_i is separable for all $i \in \mathbb{N}$.

Definition 2.12 Let $\lambda: \mathfrak{A} \to \mathcal{B}(H)$ be a geometric Hilbert space over (X, \mathcal{E}) . We define the *support* of $u \in H$ and $T \in \mathcal{B}(H)$ as follows:

$$\begin{aligned} \operatorname{Supp}(u) &= \left\{ x \in X \mid \lambda(A)u \neq 0 \text{ for all } A \in \mathfrak{A}(x) \right\} \\ \operatorname{Supp}(T) &= \left\{ (x_1, x_2) \in X \times X \mid \lambda(A_1)T\lambda(A_2) \neq 0 \text{ for all } A_i \in \mathfrak{A}(x_i) \right\} \end{aligned}$$

Remark 2.13 Assume $\lambda(A) = 0$ and $x \in A$. Then $x \notin \text{Supp}(u)$ for all $u \in H$. Similarly $(x, y), (y, x) \notin \text{Supp}(T)$ for $T \in \mathcal{B}(H)$ and $y \in X$.

In particular, consider the case where $\{x\} \in \mathfrak{A}$ and $\lambda(\{x\}) = 0$ for all $x \in X$. If we consider the discrete topology on X or equivalently if we set $\mathfrak{A}(x) := \{A \in \mathfrak{A} \mid x \in A\}$, then $\operatorname{Supp}(u) = \emptyset$ for all $u \in H$ and $\operatorname{Supp}(T) = \emptyset$ for all $T \in \mathcal{B}(H)$. This demonstrates why we introduced \mathfrak{A} -neighborhoods and why we do not use the definition of support given in [Roe03].

In order to get a calculus of supports as in Proposition 2.18, we need λ to fulfill an additional condition.

Remark 2.14 Let $\lambda : \mathfrak{A} \to \mathcal{B}(H)$ be any geometric Hilbert space. For $A, A_1, A_2 \in \mathfrak{A}, u \in H$ and $T \in \mathcal{B}(H)$ the following is always true.

$$\lambda(A)u = 0 \implies \mathring{A} \cap \operatorname{Supp}(u) = \emptyset$$
$$\lambda(A_1) T \lambda(A_2) = 0 \implies \mathring{A_1} \times \mathring{A_2} \cap \operatorname{Supp}(T) = \emptyset$$

Definition 2.15 We say that a geometric Hilbert space leads to *computable* supports if for $A, A_1, A_2 \in \mathfrak{A}, u \in H$ and $T \in \mathcal{B}(H)$ the following is true.

$$A \cap \operatorname{Supp}(u) = \emptyset \implies \lambda(A)u = 0 \tag{2.1}$$

$$A_1 \times A_2 \cap \operatorname{Supp}(T) = \emptyset \implies \lambda(A_1) T \lambda(A_2) = 0 \tag{2.2}$$

Remark 2.16 In the situation of Example 2.9, supports are computable (provided that the Hilbert spaces H_i are separable).

Lemma 2.17 Let $\lambda \colon \mathfrak{A} \to \mathcal{B}(H)$ be a geometric Hilbert space over (X, \mathcal{E}) . Assume that X is σ -compact.³ Then the geometric Hilbert space λ leads to computable supports.

³A topological space is σ -compact if every open cover of X has a countable subcover.

Proof. Assume $A \in \mathfrak{A}$ and $A \cap \operatorname{Supp}(u) = \emptyset$. For every $x \in A$ there is $B_x \in \mathfrak{A}(x)$ such that $\lambda(B_x)u = 0$. Set $C_x = B_x \cap A$. Using Lemma 2.7, we get

$$\lambda(C_x)u = \lambda(C_x)\underbrace{\lambda(B_x)u}_{=0} = 0$$
.

Take a countable subcover $\mathcal{C}' \subseteq \mathcal{C} := \{C_x \mid x \in A\}$ of A. Using σ -additivity of λ , we get $\lambda(A)u = 0$.

The proof of the corresponding claim about the support of T is similar. \Box

The assumption of σ -compactness of X is necessary in Lemma 2.17. Compare Remark 2.13.

Proposition 2.18 Let $\lambda: \mathfrak{A} \to \mathcal{B}(H)$ be a geometric Hilbert space over (X, \mathcal{E}) which leads to computable supports. Let $u, v \in H$ and $T, S \in \mathcal{B}(H)$. Take $\Delta_{\mathcal{U}} = \bigcup_{U \in \mathcal{U}} U \times U \in \mathcal{E}$ for some localizing decomposition \mathcal{U} . We get the following calculus of supports:

$$\operatorname{Supp}(u+v) \subseteq \operatorname{Supp}(u) \cup \operatorname{Supp}(v)$$
 (2.3)

$$\operatorname{Supp}(S+T) \subseteq \operatorname{Supp}(S) \cup \operatorname{Supp}(T)$$
 (2.4)

$$\operatorname{Supp}(Tu) \subseteq \Delta_{\mathcal{U}} \operatorname{Supp}(T) \Delta_{\mathcal{U}}[\operatorname{Supp}(u)]$$
(2.5)

$$\operatorname{Supp}(ST) \subseteq \Delta_{\mathcal{U}} \operatorname{Supp}(S) \Delta_{\mathcal{U}} \operatorname{Supp}(T) \Delta_{\mathcal{U}}$$
(2.6)

$$\operatorname{Supp}(T^*) = \operatorname{Supp}(T)^{-1} \tag{2.7}$$

Proof. The first two inclusions (2.3) and (2.4) follow directly from the definitions. The same is true for (2.7).

In order to prove (2.5), suppose $x \notin \Delta_{\mathcal{U}} \operatorname{Supp}(T) \Delta_{\mathcal{U}} \operatorname{Supp}(u)$ and choose $U_x \in \mathcal{U}$ such that $x \in U_x$. For every $U \in \mathcal{U} \setminus \{U_x\}$ we have

$$U \cap \operatorname{Supp}(u) = \emptyset$$
 or $U_x \times U \cap \operatorname{Supp}(T) = \emptyset$.

Hence $\lambda(U_x)Tu = \sum_{U \in \mathcal{U}} \lambda(U_x)T\lambda(U)u = 0$, i.e. $x \notin \operatorname{Supp}(Tu)$. The prove of (2.6) is similar. Suppose $(x, y) \notin \Delta_{\mathcal{U}} \operatorname{Supp}(S)\Delta_{\mathcal{U}} \operatorname{Supp}(T)\Delta_{\mathcal{U}}$ and choose $U_x, U_y \in \mathcal{U}$ containing x and y respectively. For every $U \in \mathcal{U}$ we have

$$U_x \times U \cap \operatorname{Supp}(S) = \emptyset$$
 or $U \times U_y \cap \operatorname{Supp}(T) = \emptyset$.

Therefore $\lambda(U_x)ST\lambda(U_y) = \sum_{U \in \mathcal{U}} \lambda(U_x)S\lambda(U)T\lambda(U_y) = 0$ and it follows that $(x, y) \notin \operatorname{Supp}(ST)$.

2.2 Translation C*-algebras

In the following let $\lambda \colon \mathfrak{A} \to \mathcal{B}(H)$ be a geometric Hilbert space over (X, \mathcal{E}) with computable supports. From Proposition 2.18 we get the following corollary.

Corollary 2.19 The bounded operators of controlled support on H, i.e. the bounded operators on H whose support is an entourage, form a *-algebra.

Definition 2.20 We denote the closure of the *-algebra of operators with controlled support in the operator norm by $E^*_{\lambda}(X, \mathcal{E})$. We may call it the *big translation* C^* -algebra.

Remark 2.21 The closure of the *-algebra of operators with controlled support with respect to strong convergence is the full algebra of bounded operators $\mathcal{B}(H)$.

Proof. Let $\{U_i\}_{i\in\mathbb{N}}$ be an \mathfrak{A} -localizing decomposition of (X, \mathcal{E}) and set $V_i := U_0 \cup \cdots \cup U_i$. Observe that for $T \in \mathcal{B}(H)$ the sequence $\{\lambda(V_i)T\lambda(V_i)\}_{i\in\mathbb{N}}$ converges strongly to T.

Definition 2.22 An operator $T \in \mathcal{B}(H)$ is *pseudolocal* if $T\lambda(B) - \lambda(B)T$ is a compact operator for every bounded set $B \in \mathfrak{A}$.

Lemma 2.23 The pseudolocal operators with controlled support form a *-algebra.

Proof. The pseudolocal operators are closed under addition, scalar multiplication and adjoints, i.e. if S and T are pseudolocal operators and $z \in \mathbb{C}$, then S+T, $z \cdot T$ and T^* are also pseudolocal. The composition of pseudolocal operators is pseudolocal, as the following computation shows: If S and T are pseudolocal operators and $B \in \mathcal{A}$ is a bounded set, then

$$ST\lambda(B) - \lambda(B)ST = (S\lambda(B) - \lambda(B)S)T + \text{compact operator}$$

is compact.

Definition 2.24 By $D_{\lambda}^{*}(X, \mathcal{E})$ we denote the closure of the *-algebra of pseudolocal operators with controlled support in the operator norm.

Definition 2.25 An operator $T \in \mathcal{B}(H)$ is *locally compact* if $T\lambda(B)$ and $\lambda(B)T$ are compact operators for every bounded set $B \in \mathfrak{A}$.

Lemma 2.26 The locally compact operators with controlled support form a *-algebra.

Proof. The locally compact operators are closed under addition, scalar multiplication and adjoints. The composition of locally compact operators is locally compact, since composing a bounded operator and a compact operator gives a compact operator. $\hfill \Box$

Definition 2.27 The small translation C^* -algebra of (X, \mathcal{E}) with respect to λ is the closure of the *-algebra of locally compact operators with controlled support in the operator norm. We denote it by $C^*_{\lambda}(X, \mathcal{E})$. This algebra is often called *Roe-C^*-algebra*.

Remark 2.28 Observe that every locally compact operator is pseudolocal. Hence $C^*_{\lambda}(X, \mathcal{E}) \subseteq D^*_{\lambda}(X, \mathcal{E}) \subseteq E^*_{\lambda}(X, \mathcal{E})$.

Lemma 2.29 The small translation C^* -algebra $C^*_{\lambda}(X, \mathcal{E})$ is an ideal⁴ in the big translation C^* -algebra $E^*_{\lambda}(X, \mathcal{E})$. The small translation C^* -algebra $C^*_{\lambda}(X, \mathcal{E})$ is also an ideal in $D^*_{\lambda}(X, \mathcal{E})$.

Proof. Let $S \in C^*_{\lambda}(X, \mathcal{E})$ and $T \in E^*_{\lambda}(X, \mathcal{E})$. For a bounded set $B \in \mathfrak{A}$ the operator $\lambda(B)S$ is compact. Therefore $\lambda(B)ST$ is compact.

It remains to prove that $ST\lambda(B_2)$ is compact for any bounded set $B_2 \in \mathfrak{A}$. Let \mathcal{U} be an \mathfrak{A} -localizing decomposition of (X, \mathcal{E}) and observe that $B_1 := \Delta_{\mathcal{U}} \operatorname{Supp}(T)[B_2] \in \mathfrak{A}$ is bounded and $(X \setminus B_1) \times B_2 \cap \operatorname{Supp}(T) = \emptyset$. Using (2.2), we see that

$$ST\lambda(B_2) = \underbrace{S\lambda(B_1)}_{\text{compact}} T\lambda(B_2) + S\underbrace{\lambda(X \setminus B_1) T\lambda(B_2)}_{=0}$$

is compact.

Example 2.30 If (X, \mathcal{E}) is a coarse space such that X is bounded, then $E^*(X, \mathcal{E})$ is the entire algebra of bounded operators on H and $C^*(X, \mathcal{E})$ is the algebra of compact operators. Moreover, $D^*_{\lambda}(X, \mathcal{E}) = \{T \in \mathcal{B}(H) \mid [T, \lambda(A)] \text{ is compact for all } A \in \mathfrak{A}\}.$

Definition 2.31 A geometric Hilbert space $\lambda : \mathfrak{A} \to \mathcal{B}(H)$ over (X, \mathcal{E}) is called *ample* if there is an \mathfrak{A} -localizing decomposition $\mathcal{U} \subseteq \mathfrak{A}$ such that $\lambda(U)$ is not compact for $U \in \mathcal{U}$. Note that for a projection being compact and having finite dimensional image is the same.

Example 2.32 Let (X, \mathcal{E}) be a coarse space and $x_0 \in X$. The geometric Hilbert space $\lambda \colon \mathfrak{A} \to \mathcal{B}(H)$ defined by

$$\lambda(A) = \begin{cases} 0 & \text{if } x_0 \notin A \\ \text{id} & \text{if } x_0 \in A \end{cases}$$

is not ample if X is unbounded.

⁴In the context of C^* -algebras, by an *ideal* we mean a closed, two-sided ideal which is closed unter taking adjoints.

Theorem 2.33 Let $\lambda: \mathfrak{A} \to \mathcal{B}(H_1)$ and $\mu: \mathfrak{B} \to \mathcal{B}(H_2)$ be ample, geometric Hilbert spaces over (X, \mathcal{E}) with computable supports. Then

$$C^*_{\lambda}(X,\mathcal{E}) \cong C^*_{\mu}(X,\mathcal{E})$$
 and $E^*_{\lambda}(X,\mathcal{E}) \cong E^*_{\mu}(X,\mathcal{E}).$

The following two lemmas yield a proof of this theorem.

Lemma 2.34 Let $\lambda : \mathfrak{A} \to \mathcal{B}(H)$ be a geometric Hilbert space over (X, \mathcal{E}) with computable supports and \mathcal{U} an \mathfrak{A} -localizing decomposition. By λ' we denote the restriction of λ to the σ -algebra $\sigma(\mathcal{U})$ generated by \mathcal{U} . Then λ' together with the discrete topology on X is another geometric Hilbert space and we have $C^*_{\lambda}(X, \mathcal{E}) = C^*_{\lambda'}(X, \mathcal{E})$ and $E^*_{\lambda}(X, \mathcal{E}) = E^*_{\lambda'}(X, \mathcal{E})$.

Proof. Whenever in this proof we make reference to a topology we mean the topology coming with \mathfrak{A} . We first claim

$$\operatorname{Supp}_{\lambda}(T) \subseteq \overline{\operatorname{Supp}_{\lambda'}(T)} \quad \text{ for } T \in \mathcal{B}(H).$$

This implies $E_{\lambda'}^*(X, \mathcal{E}) \subseteq E_{\lambda}^*(X, \mathcal{E})$. To prove the claim, set $Y := \bigcup_{U \in \mathcal{U}} \mathring{U}$. If $x \in Y$, then $\sigma(\mathcal{U})(x) = \{U \in \mathcal{U} \mid x \in U\} \subseteq \mathfrak{A}(x)$. Therefore we get $\operatorname{Supp}_{\lambda}(T) \cap Y \times Y \subseteq \operatorname{Supp}_{\lambda'}(T) \cap Y \times Y$.

For $x \in X$ set $\mathcal{U}_x := \{U \in \mathcal{U} \mid x \in \overline{U}\}$ and observe that $\bigcup_{U \in \mathcal{U}_x} U$ is an \mathfrak{A} neighborhood of x. Thus, for $(x, y) \in \operatorname{Supp}_{\lambda}(T)$ there is $U \in \mathcal{U}_x$ and $V \in \mathcal{U}_y$ such that $\lambda(U)T\lambda(V) \neq 0$. Take sequences $\{x_i\}_{i \in \mathbb{N}} \subseteq U$ and $\{y_i\}_{i \in \mathbb{N}} \subseteq V$ converging to x and y respectively. Note that $(x_i, y_i) \in \operatorname{Supp}_{\lambda'}(T)$ for all $i \in \mathbb{N}$ and hence $(x, y) \in \operatorname{Supp}_{\lambda'}(T)$.

Set $\Delta_{\mathcal{U}} = \bigcup_{U \in \mathcal{U}} U \times U$. We claim

$$\operatorname{Supp}_{\lambda'}(T) \subseteq \Delta_{\mathcal{U}} \operatorname{Supp}_{\lambda}(T) \Delta_{\mathcal{U}} \quad \text{ for all } T \in \mathcal{B}(H)$$

This claim implies $E^*_{\lambda}(X, \mathcal{E}) \subseteq E^*_{\lambda'}(X, \mathcal{E})$. To prove the claim, suppose $U, V \in \mathcal{U}$ and $U \times V \cap \operatorname{Supp}_{\lambda}(T) = \emptyset$. Since supports are computable, we get $\lambda(U)T\lambda(V) = 0$. Therefore $U \times V \cap \operatorname{Supp}_{\lambda'}(T) = \emptyset$. So far we proved $E^*_{\lambda}(X, \mathcal{E}) = E^*_{\lambda'}(X, \mathcal{E})$.

In order to prove $\hat{C}^*_{\lambda}(X, \mathcal{E}) \subseteq \hat{C}^*_{\lambda'}(X, \mathcal{E})$, just note that each bounded set in $\sigma(\mathcal{U})$ is also a bounded set in \mathfrak{A} .

Each bounded set $B \in \mathfrak{A}$ is contained in $\widetilde{B} := \Delta_{\mathcal{U}}[B]$ which is a bounded set in $\sigma(\mathcal{U})$. Let T be locally compact with respect to $\sigma(\mathcal{U})$. Using Lemma 2.7, we see that $\lambda(B)T = \lambda(B)\lambda(\widetilde{B})T$ is compact, since $\lambda(\widetilde{B})T$ is compact. \Box

Lemma 2.35 Given localizing decompositions \mathcal{U} and \mathcal{V} of the coarse space (X, \mathcal{E}) let $\lambda : \sigma(\mathcal{U}) \to \mathcal{B}(H_1)$ and $\mu : \sigma(\mathcal{V}) \to \mathcal{B}(H_2)$ be ample, geometric Hilbert spaces over (X, \mathcal{E}) . Then there is an isometric isomorphism $\rho : H_1 \to H_2$ such that $ad_{\rho} : \mathcal{B}(H_1) \to \mathcal{B}(H_2), T \mapsto \rho T \rho^{-1}$ restricts to isomorphisms $E^*_{\lambda}(X, \mathcal{E}) \xrightarrow{\cong} E^*_{\mu}(X, \mathcal{E})$ and $C^*_{\lambda}(X, \mathcal{E}) \xrightarrow{\cong} C^*_{\mu}(X, \mathcal{E})$.

Proof. Defining $\mathcal{W} = \{U \cap V \mid U \in \mathcal{U}, V \in \mathcal{V}\} \setminus \{\emptyset\}$ we get a localizing decomposition refining \mathcal{U} and \mathcal{V} . Set $\mathcal{W}_U = \{W \in \mathcal{W} \mid W \subseteq U\}$ for $U \in \mathcal{U}$ and $\mathcal{W}_V = \{W \in \mathcal{W} \mid W \subseteq V\}$ for $V \in \mathcal{V}$. We will write H_U for the image of $\lambda(U)$ and H_V for the image of $\mu(V)$. For every $U \in \mathcal{U}$ we choose a decomposition $H_U = \bigoplus_{W \in \mathcal{W}_U} H_{1,W}$ such that $H_{1,W}$ is infinite dimensional for all $W \in \mathcal{W}$. Similarly, we choose a decomposition $H_V = \bigoplus_{W \in \mathcal{W}_V} H_{2,W}$ for each $V \in \mathcal{V}$. Now we extend the geometric Hilbert spaces λ and μ to $\tilde{\lambda}: \sigma(\mathcal{W}) \to \mathcal{B}(H_1)$ and $\tilde{\mu}: \sigma(\mathcal{W}) \to \mathcal{B}(H_2)$ respectively. We define $\tilde{\lambda}(W)$ to be the projection onto $H_{1,W}$. Similarly $\tilde{\mu}(W)$ will be the projection onto $H_{2,W}$. Remember that $\lambda, \tilde{\lambda}, \mu$ and $\tilde{\mu}$ have computable supports. We choose an isometric isomorphism $\rho: H_1 \to H_2$ such that the image of $\rho|_{H_{1,W}}$ is $H_{2,W}$ for all $W \in \mathcal{W}$. Then ad_{ρ} is an isomorphism of C^* -algebras.

Using $\rho \widetilde{\lambda} = \widetilde{\mu} \rho$, we see that $\operatorname{Supp}_{\widetilde{\lambda}}(T) = \operatorname{Supp}_{\widetilde{\mu}}(\rho T \rho^{-1})$ for all $T \in \mathcal{B}(H_1)$. Hence ad_{ρ} restricts to an isomorphism $E^*_{\widetilde{\lambda}}(X, \mathcal{E}) \xrightarrow{\cong} E^*_{\widetilde{\mu}}(X, \mathcal{E})$.

Moreover, we note that for $B \in \sigma(\mathcal{W})$ and $T \in \mathcal{B}(H_1)$ the operator $\widetilde{\lambda}(B)T$ is compact if and only if $\widetilde{\mu}(B)ad_{\rho}(T)$ is compact. Therefore ad_{ρ} also restricts to an isomorphism $C^*_{\widetilde{\lambda}}(X, \mathcal{E}) \xrightarrow{\cong} C^*_{\widetilde{\mu}}(X, \mathcal{E})$. Applying Lemma 2.34 now completes the proof.

There are some choices involved in order to get the isometric isomorphism $\rho: H_1 \to H_2$ in the proof of Lemma 2.35. Therefore, the isomorphisms $E^*_{\lambda}(X, \mathcal{E}) \xrightarrow{\cong} E^*_{\mu}(X, \mathcal{E})$ and $C^*_{\lambda}(X, \mathcal{E}) \xrightarrow{\cong} C^*_{\mu}(X, \mathcal{E})$ are not uniquely determined and without further specifications there is no natural choice.

We compare our definition of small translation C^* -algebra with the corresponding definition in Section 6.3 of [HR00].

Proposition 2.36 Let X be a locally compact, separable and metrizable space with a coarse structure \mathcal{E} which is compatible with the topology. Suppose that

$$\rho\colon C_0(X)\to \mathcal{B}(H)$$

is a non-degenerate⁵ and ample⁶ representation of the C^* -algebra $C_0(X)$ on the separable Hilbert space H. There is an ample geometric Hilbert space λ over (X, \mathcal{E}) with computable supports and for any such λ

$$C^*_{\rho}(X,\mathcal{E}) \cong C^*_{\lambda}(X,\mathcal{E})$$

Proof. According to Theorem 2.33 we are free to replace λ by any ample geometric Hilbert space with computable supports. We will construct λ from ρ .

⁵A C^* -representation ρ of a C^* -algebra A on a Hilbert space H is non-degenerate if $\{\rho(a)(v) \mid a \in A, v \in H\}$ is dense in H.

⁶A C^{*}-representation ρ is called ample, if the operator $\rho(f)$ is not compact if $f \neq 0$.

Using Borel functional calculus (compare Remark 1.5.7 of [HR00]), we see that ρ extends to a representation $\tilde{\rho}$ of the C^* -algebra B(X) of bounded Borel functions on X. It is easy to check that

$$\lambda \colon \mathfrak{B}(X) \to \mathcal{B}(H), \quad A \mapsto \widetilde{\rho}(\chi_A)$$

is a projection-valued measure.

Since X is separable and its coarse structure is compatible with the topology, there exists a countable, uniformly bounded, open cover of X. We may even assume that each $U \in \mathcal{U}$ has non-empty interior. (Compare Claim 6.3.14 of [HR00].) Hence λ is a geometric Hilbert space.

Let $U \in \mathcal{U}$ and $f \in C_0(U)$, $f \neq 0$. Since ρ is ample, $\lambda(U)\rho(f) = \rho(f)$ is not compact. Hence $\lambda(U)$ is not compact and λ is ample. Applying Lemma 2.17 we see that λ leads to computable supports.

Now that we have a convenient geometric Hilbert space, we compare the support of a bounded operator T with respect to λ (as defined in Definition 2.12) and the support with respect to ρ (as defined in [HR00]). Observe that

$$\operatorname{Supp}_{\rho}(T) = \left\{ (x_1, x_2) \in X^2 \mid \forall_{U_1 \in \mathfrak{B}(X)(x_1)} \exists_{f_1 \in C_0(\mathring{U}_1)} \rho(f_1) T \rho(f_2) \neq 0 \right\} \ .$$
$$U_2 \in \mathfrak{B}(X)(x_2) \quad f_2 \in C_0(\mathring{U}_2)$$

Let $x_1, x_2 \in X$ and $U_i \in \mathfrak{B}(X)(x_i)$. Suppose there are $f_i \in C_0(\mathring{U}_i)$ such that $\rho(f_1)T\rho(f_2) \neq 0$. Observe the following relations between ρ and λ .

$$\rho(f_1)\lambda(U_1) = \rho(f_1)\widetilde{\rho}(\chi_{U_1}) = \widetilde{\rho}(f_1 \cdot \chi_{U_1}) = \rho(f_1)$$

$$\lambda(U_2)\rho(f_2) = \widetilde{\rho}(\chi_{U_2})\rho(f_2) = \widetilde{\rho}(\chi_{U_2} \cdot f_2) = \rho(f_2)$$

It follows

$$\rho(f_1)\lambda(U_1)T\lambda(U_2)\rho(f_2) = \rho(f_1)T\rho(f_2) \neq 0$$

and hence $\lambda(U_1)T\lambda(U_2) \neq 0$. This proves $\operatorname{Supp}_{\rho}(T) \subseteq \operatorname{Supp}_{\lambda}(T)$. Choose a metric *d* which induces the given topology on *X*. For $U \subseteq X$ and $k \in \mathbb{N}$ define

$$f_{U,k} \colon X \to \mathbb{R}, \quad x \mapsto \max\{0, 1 - k \cdot d(x, U)\}.$$

Let $(x_1, x_2) \in \operatorname{Supp}_{\lambda}(T)$ and $U_i \in \mathfrak{B}(X)(x_i)$. Since (X, d) is a metric space, there is $k_i \in \mathbb{N}$ and $V_i \in \mathfrak{B}(X)(x_i)$ such that $f_i := f_{V_i, k_i} \in C_0(\mathring{U}_i)$. We have the following relations between ρ and λ .

$$\lambda(V_1)\rho(f_1) = \widetilde{\rho}(\chi_{V_1})\rho(f_1) = \widetilde{\rho}(\chi_{V_1} \cdot f_1) = \lambda(V_1)$$

$$\rho(f_2)\lambda(V_2) = \rho(f_2)\widetilde{\rho}(\chi_{V_2}) = \widetilde{\rho}(f_2 \cdot \chi_{V_2}) = \lambda(V_2)$$

These relations imply

$$\lambda(V_1)\rho(f_1)T\rho(f_2)\lambda(V_2) = \lambda(V_1)T\lambda(V_2) \neq 0$$

and hence $\rho(f_1)T\rho(f_2) \neq 0$. It follows $\operatorname{Supp}_{\lambda}(T) \subseteq \operatorname{Supp}_{\rho}(T)$.

We assume that $T \in C^*_{\rho}(X, \mathcal{E})$. Note that $f_A := f_{A,1} \in C_0(X)$ for a bounded set $A \subseteq X$. Since $\rho(f_A)T$ is compact, we conclude that $\lambda(\overline{A})T = \lambda(\overline{A})\rho(f_A)T$ is compact. In the same way we get compactness for $T\lambda(\overline{A})$. This proves $T \in C^*_{\lambda}(X, \mathcal{E})$.

Now let $T \in \mathcal{B}(H)$ be locally compact with respect to λ . If $f \in C_0(X)$, then $M_{f,n} := \{x \in X \mid |f(x)| \ge \frac{1}{n}\}$ is compact for all $n \in \mathbb{N}$ and thus bounded. Continuity of the representation $\tilde{\rho}$ yields

$$\rho(f)\lambda(M_{f,n})T \xrightarrow{\|\cdot\|}{n \to \infty} \rho(f)T$$
.

Since all operators on the left hand side are compact, $\rho(f)T$ is compact. In the same way we get compactness for $T\rho(f)$.

Corollary 2.37 Up to isomorphism, the C^* -algebra $C_{\rho}(X)$ does not depend on the choice of the representation ρ (as long as ρ is an ample and non-degenerate representation on a separable Hilbert space).

2.3 Induced maps

Let (X, \mathcal{E}_X) and (Y, \mathcal{E}_Y) be coarse spaces which admit localizing decompositions and let $f: (X, \mathcal{E}_X) \to (Y, \mathcal{E}_Y)$ be a coarse map. We would like to obtain induced maps $E^*(f)$ and $C^*(f)$ on the translation C^* -algebras such that the following diagramm commutes.

Question 2.38 Let $f: (X, \mathcal{E}_X) \to (Y, \mathcal{E}_Y)$ be a coarse map and \mathfrak{A} a localizing σ -algebra with respect to \mathcal{E}_X . Is there a way of defining an induced localizing σ -algebra $f_*(\mathfrak{A})$?

An attempt to define an induced localizing σ -ring⁷ was made in Remark 4.26 of [Roe03]. Roe defines

$$f_*(\mathfrak{A}) := \{ S \subseteq f(X) \subseteq Y \mid f^{-1}(S) \in \mathfrak{A} \}$$
.

Now $f_*(\mathfrak{A})$ is indeed a σ -ring, but it does not have to be localizing as the following example tells us.

⁷A σ -ring \mathfrak{R} over a set X is a family of subsets of X such that (1) $\emptyset \in \mathfrak{R}$, (2) $A \setminus B \in \mathfrak{R}$ whenever $A, B \in \mathfrak{R}$ and (3) the union of countably many elements of \mathfrak{R} is in \mathfrak{R} .

Example 2.39 Take $(X, \mathcal{E}_X) = (Y, \mathcal{E}_Y) = (\mathbb{R}, \mathcal{E}_{eucl})$ and the σ -algebra \mathfrak{A} generated by the set of half open intervals $\{[n, n+1[| n \in \mathbb{Z}\})\}$. For the coarse map $f: X \to Y$

$$f(x) = \begin{cases} n & \text{if } x \in \left[n - \frac{1}{4}, n + \frac{1}{4}\right] \text{ for some } n \in \mathbb{Z} \\ 2 \cdot x - n - \frac{1}{2} & \text{if } x \in \left[n + \frac{1}{4}, n + \frac{3}{4}\right] \text{ for some } n \in \mathbb{Z} \end{cases}$$

we get $f_*(\mathfrak{A}) = \{\emptyset, \mathbb{R}\}$ which is not localizing (in the sense of Definition 4.20 in [Roe03]), since \mathbb{R} is unbounded.

Roe's definition of $f_*(\mathfrak{A})$ does give a "localizing σ -ring" if the coarse map f "respects an \mathfrak{A} -localizing decomposition", i.e. if f has the following additional property: There is an \mathfrak{A} -localizing decomposition \mathcal{U} of X such that $U_1, U_2 \in \mathcal{U}$ and $U_1 \neq U_2$ implies $f(U_1) \cap f(U_2) = \emptyset$. This is the case for example if f is injective.

We start constructing geometric Hilbert spaces λ_X and λ_Y such that f does induce maps between the corresponding translation C^* -algebras.

Lemma 2.40 There are geometric Hilbert spaces $\lambda_X : \sigma(\mathcal{U}_X) \to \mathcal{B}(H_X)$ and $\lambda_Y : \sigma(\mathcal{U}_Y) \to \mathcal{B}(H_Y)$, a map $F : \mathcal{U}_X \to \mathcal{U}_Y$ and a partial isometry $\varphi : H_X \to H_Y$ such that

- \mathcal{U}_X and \mathcal{U}_Y are localizing decompositions of X and Y respectively,
- λ_X and λ_Y are ample and lead to computable supports,
- the map $F: \mathcal{U}_X \to \mathcal{U}_Y$ satisfies $F(U) \subseteq f(U)$ for all $U \in \mathcal{U}_X$,
- the map $F: \mathcal{U}_X \to \mathcal{U}_Y$ is almost injective, i.e. F(U) = F(U') implies U = U' or $F(U) = \emptyset$,
- the partial isometry satisfies $\varphi(\lambda_X(U)(H_X)) \subseteq \lambda_Y(F(U))(H_Y)$ for all $U \in \mathcal{U}_X$.

Proof. Take any localizing decomposition \mathcal{U}_X of (X, \mathcal{E}_X) . We enumerate the elements of \mathcal{U}_X , denote them by U_0, U_1, U_2, \ldots and set $W_i := f(U_i)$. For $i \in \mathbb{N}$ define $V_i = W_i \setminus \bigcup_{j < i} W_j$ and note that $\mathcal{V} := \{V_0, V_1, V_2, \ldots\}$ is a localizing decomposition of $\operatorname{im}(f)$. The empty set might be an element of \mathcal{V} . For $i \in \mathbb{N}$ define $F(U_i) := V_i$. Let $\widetilde{\mathcal{U}}_Y$ be a localizing decomposition of Yand define

$$\mathcal{U}_Y := \mathcal{V} \cup \left\{ U \setminus \operatorname{im}(f) \mid U \in \widetilde{\mathcal{U}}_Y \right\}.$$

Note that \mathcal{U}_Y is itself a localizing decomposition of Y.

Choose a separable Hilbert space H_U for each $U \in \mathcal{U}_X$ with $U \neq \emptyset$ and similarly choose H_V for each $V \in \mathcal{U}_Y$ with $V \neq \emptyset$. Define $H_{\emptyset} = \{0\}$ if $\emptyset \in \mathcal{U}_X$ or $\emptyset \in \mathcal{U}_Y$. Define $H_X = \bigoplus_{U \in \mathcal{U}_X} H_U$ and $H_Y = \bigoplus_{V \in \mathcal{U}_Y} H_V$. Choose a partial isometry $\varphi \colon H_X \to H_Y$ such that $\varphi(H_U) \subseteq H_{F(U)}$ for all $U \in \mathcal{U}_X$. We consider the geometric Hilbert space $\lambda_X : \sigma(\mathcal{U}_X) \to \mathcal{B}(H_X)$ which is determined by defining $\lambda_X(U)$ to be the projection onto H_U for any $U \in \mathcal{U}_X$. Likewise we define $\lambda_Y : \sigma(\mathcal{U}_Y) \to \mathcal{B}(H_Y)$. Note that λ_X and λ_Y are ample if H_U and H_V are infinite dimensional for all $U \in \mathcal{U}_X$ and $V \in \mathcal{U}_Y$. Moreover λ_X and λ_Y lead to computable supports. (Compare Remark 2.16.)

Lemma 2.41 In the situation of Lemma 2.40, the partial isometry φ induces the map

$$ad_{\varphi} \colon \mathcal{B}(H_X) \to \mathcal{B}(H_Y), \quad T \mapsto \varphi T \varphi^*$$

and ad_{φ} restricts to maps between the translation C^* -algebras.

$$E^*_{\lambda_X}(X, \mathcal{E}_X) \xrightarrow{ad_{\varphi}} E^*_{\lambda_Y}(Y, \mathcal{E}_Y)$$
 (2.8)

$$C^*_{\lambda_X}(X, \mathcal{E}_X) \xrightarrow{ad_{\varphi}} C^*_{\lambda_Y}(Y, \mathcal{E}_Y)$$
 (2.9)

Proof. Let $T \in \mathcal{B}(H_X)$. We will prove

$$\operatorname{Supp}_{\lambda_Y}(ad_{\varphi}(T)) \subseteq f \times f\left(\operatorname{Supp}_{\lambda_X}(T)\right)$$
.

This implies $ad_{\varphi}\left(E^*_{\lambda_X}(X,\mathcal{E}_X)\right)\subseteq E^*_{\lambda_Y}(Y,\mathcal{E}_Y).$

Let $U_1, U_2 \in \mathcal{U}_X$ such that $U'_1 \times U_2 \not\subseteq \operatorname{Supp}_{\lambda_X}(T)$. The definition of λ_X implies $U_1 \times U_2 \cap \operatorname{Supp}_{\lambda_X}(T) = \emptyset$ and hence $\lambda_X(U_1)T\lambda_X(U_2) = 0$, since supports are computable. Observe that

$$\lambda_Y(F(U_1)) \ ad_{\varphi}(T) \ \lambda_Y(F(U_2)) = \lambda_Y(F(U_1)) \ \varphi \ T \ \varphi^* \ \lambda_Y(F(U_2))$$
$$= \ \lambda_Y(F(U_1)) \ \varphi \left(\sum_{U \in \mathcal{U}_X} \lambda_X(U)\right) T \left(\sum_{U \in \mathcal{U}_X} \lambda_X(U)\right) \ \varphi^* \ \lambda_Y(F(U_2))$$
$$= \ \lambda_Y(F(U_1)) \ \varphi \ \underbrace{\lambda_X(U_1) \ T \ \lambda_X(U_2)}_{=0} \ \varphi^* \ \lambda_Y(F(U_2)) = 0 \ .$$

If $(y_1, y_2) \in \operatorname{Supp}_{\lambda_Y}(ad_{\varphi}(T))$, there are $U_1, U_2 \in \mathcal{U}_X$ such that $y_i \in F(U_i)$. Hence $\lambda_Y(F(U_1))ad_{\varphi}(T)\lambda_Y(F(U_2)) \neq 0$. As we have just seen, this implies $U_1 \times U_2 \subseteq \operatorname{Supp}_{\lambda_X}(T)$ and therefore

$$(y_1, y_2) \in F(U_1) \times F(U_2) \subseteq f(U_1) \times f(U_2) \subseteq f \times f(\operatorname{Supp}_{\lambda_X}(T)).$$

It remains to prove $ad_{\varphi}\left(C^*_{\lambda_X}(X,\mathcal{E}_X)\right) \subseteq C^*_{\lambda_Y}(Y,\mathcal{E}_Y)$. For this purpose suppose $T \in C^*_{\lambda_X}(X,\mathcal{E}_X)$. Let $B \in \sigma(\mathcal{U}_Y)$ be a bounded set. Then $\lambda_X\left(\Delta_{\mathcal{U}_X}[f^{-1}(B)]\right)T$ is compact, since $\Delta_{\mathcal{U}_X}[f^{-1}(B)] \in \sigma(\mathcal{U}_X)$ is bounded and T is locally compact. Therefore

$$\lambda_Y(B)ad_{\varphi}(T) = \lambda_Y(B)\varphi T\varphi^* = \lambda_Y(B)\varphi \underbrace{\lambda_X\left(\Delta_{\mathcal{U}_X}[f^{-1}(B)]\right)T}_{\text{compact}}\varphi^*$$

and $ad_{\varphi}(T)\lambda_Y(B)$ are compact operators. This proves that $ad_{\varphi}(T)$ is locally compact.

We denote the maps (2.8) and (2.9) by $E^*_{\lambda_X,\lambda_Y,\varphi}(f)$ and $C^*_{\lambda_X,\lambda_Y,\varphi}(f)$ respectively. Note that these maps do not only depend on λ_X and λ_Y , but also on the partial isometry φ .

Proposition 2.42 Let (X, \mathcal{E}_X) and (Y, \mathcal{E}_Y) be coarse spaces which admit countable, uniformly bounded covers. If (X, \mathcal{E}_X) and (Y, \mathcal{E}_Y) are coarsely equivalent, then $C^*(X, \mathcal{E}_X) \cong C^*(Y, \mathcal{E}_Y)$ and $E^*(X, \mathcal{E}_X) \cong E^*(Y, \mathcal{E}_Y)$.

Proof. Let $f: (X, \mathcal{E}_X) \to (Y, \mathcal{E}_Y)$ be a coarse equivalence with coarse inverse $g: (Y, \mathcal{E}_Y) \to (X, \mathcal{E}_X)$. Take a localizing decomposition \mathcal{W} of $\operatorname{im}(f \circ g) \subseteq Y$ which does not contain the empty set. Define

$$E := \Delta_{\mathcal{W}} \cup \{ (y, f \circ g(y)) \mid y \in Y \setminus \operatorname{im}(f \circ g) \}$$

and observe that $\mathcal{U}_Y := \{E[W] \mid W \in \mathcal{W}\}$ is a localizing decomposition of the coarse space (Y, \mathcal{E}_Y) .

Let $U \in \mathcal{U}_Y$. Since there is $W \in \mathcal{W}$ with $W \subseteq U$, the inverse image of U under f is not empty. Define $\mathcal{U}_X := \{f^{-1}(U) \mid U \in \mathcal{U}_Y\}$. Choose an infinite dimensional, separable Hilbert space H_U for each $U \in \mathcal{U}_Y$ and set $H := \bigoplus_{U \in \mathcal{U}_Y} H_U$. Consider the geometric Hilbert space $\lambda_Y : \sigma(\mathcal{U}_Y) \to \mathcal{B}(H)$ which is determined by setting $\lambda(U)$ to be the projection onto H_U . Furthermore, consider the geometric Hilbert space $\lambda_X : \sigma(\mathcal{U}_X) \to \mathcal{B}(H)$ which is given by defining $\lambda(f^{-1}(U))$ to be the projection onto H_U .

If we define $F: \mathcal{U}_X \to \mathcal{U}_Y$ by $F(f^{-1}(U)) = U$ for all $U \in \mathcal{U}_Y$ and $\varphi = \mathrm{id}_H$, the conclusions of Lemma 2.40 are satisfied. Moreover, the map ad_{φ} is just the identity on $\mathcal{B}(H)$. Hence, Lemma 2.41 yields $E^*_{\lambda_X}(X, \mathcal{E}_X) \subseteq E^*_{\lambda_Y}(Y, \mathcal{E}_Y)$ and $C^*_{\lambda_X}(X, \mathcal{E}_X) \subseteq C^*_{\lambda_Y}(Y, \mathcal{E}_Y)$.

The reverse inclusion for the big translation C^* -algebras follows from

$$\operatorname{Supp}_{\lambda_X}(T) = (f \times f)^{-1} \left(\operatorname{Supp}_{\lambda_Y}(T) \right) \quad \text{for all } T \in \mathcal{B}(H)$$

and the fact that f is a coarse embedding.

Observe that $\lambda_Y(B)T = \lambda_X(f^{-1}(B))T$ for each bounded sets $B \in \sigma(\mathcal{U}_Y)$. We conclude that an operator $T \in \mathcal{B}(H)$ is locally compact with respect to Y if and only if T is locally compact with respect to X.

Chapter 3

General theory of asymptotic dimension

The following definitions are an asymptotic analog of the covering dimension of topological spaces. Asymptotic dimension was first introduced in [Gro93]. For dimension theory of topological spaces see [HW41], [Eng78] and [Fed88].

Finiteness of asymptotic dimension seems to play an important role for some isomorphism conjectures in K-theory. Guoliang Yu proved in [Yu98] that the Baum-Connes assembly map is injective for groups of finite asymptotic dimension admitting a finite classifying space. This implies the Novikov conjecture on the homotopy invariance of higher signatures. Arthur Bartels proved injectivity of the assembly map in algebraic K-theory for the same class of groups. Compare [Bar03].

3.1 Asymptotic dimension of pseudometric spaces

Definition 3.1 (asymptotic dimension of pseudometric spaces) Let (X, d) be a pseudometric space, \mathcal{M} a collection of subsets of X and $n \in \mathbb{N}$.

- The multiplicity of \mathcal{M} is defined to be the maximal number of sets with non-empty intersection and will be denoted by $\mu(\mathcal{M})$.
- The mesh of \mathcal{M} is defined to be the supremum of the diameter of sets from \mathcal{M} .
- The Lebesgue number of a cover \mathcal{U} of X is the largest positive number L such that for every $x \in X$ the open ball of radius L with center x is contained in some $U \in \mathcal{U}$. We will write $L(\mathcal{U})$ for the Lebesgue number of \mathcal{U} .

- $\operatorname{asdim}(X, d) \leq n$ if for all L > 0 there exist D > 0 and a cover \mathcal{U} of X such that
 - (1) $\mu(\mathcal{U}) \le n+1,$
 - (2) $L(\mathcal{U}) \ge L$ and
 - (3) $\operatorname{mesh}(\mathcal{U}) \leq D$, i.e. \mathcal{U} is uniformly bounded.
- $\operatorname{asdim}(X, d) = n$ if $\operatorname{asdim}(X, d) \le n$ and $\operatorname{asdim}(X, d) \le n 1$.
- $\operatorname{asdim}(X, d) = \infty$ if there is no $n \in \mathbb{N}$ with $\operatorname{asdim}(X, d) \le n$.

In Section 3.2 we will generalize this definition to coarse spaces.

As in dimension theory of topological spaces, there are some alternative definitions. For details see [Gro93] and [Dra00]. We will give a second definition of asymptotic dimension, which has been generalized to coarse spaces in [Roe03]. We will prove that both definitions coincide, in the more general setting of coarse spaces.

Definition 3.2 Let (X, d) be a pseudometric space.

- Let L > 0. A family \mathcal{V} of subsets of X is called L-disjoint, if the distance of two sets from \mathcal{V} is always bigger than L.
- $\operatorname{asdim}(X, d) \leq n$ if for all L > 0 there is a cover \mathcal{U} of X such that
 - (1) the cover \mathcal{U} consists of n+1 families $\mathcal{U}_1, \ldots, \mathcal{U}_{n+1}$,
 - (2) each family \mathcal{U}_i is *L*-disjoint and
 - (3) the cover \mathcal{U} is uniformly bounded.

Remark 3.3 (monotony) Let (X, d) be a pseudometric space. If $(A, d|_A)$ is a subspace of (X, d), then $\operatorname{asdim}(A, d|_A) \leq \operatorname{asdim}(X, d)$.

Example 3.4 Let T be a tree and d the natural metric on T. (All edges are supposed to be of length 1.) Then $\operatorname{asdim}(T, d) \leq 1$ and $\operatorname{asdim}(T, d) = 1$ if and only if T is not bounded.

Proof. Choose $x_0 \in vert(T)$. Let L > 0 and define L' to be the smallest natural number bigger than $2 \cdot L$. Consider the map

$$f: T \to \mathbb{N}, \ x \mapsto \left[\frac{d(x_0, x)}{L'}\right].$$

We define $A := \overline{f^{-1}(2 \cdot \mathbb{N})}$ and $B := \overline{f^{-1}(2 \cdot \mathbb{N} + 1)}$, thus $X = A \cup B$. There is an equivalence relation on A: Let $a_1, a_2 \in A$. With γ_i we denote the geodesic path from x_0 to a_i .

$$a_1 \sim a_2 \iff f(a_1) = f(a_2) \text{ and } \gamma_1(\tau) = \gamma_2(\tau)$$

where $\tau := L' \cdot (f(a_1) - \frac{1}{2})$. If $a_1 \not \sim a_2$, then $d(a_1, a_2) \geq L'$. Furthermore, we see diam([a]) $\leq 3 \cdot L'$. Here [a] denotes the equivalence class of $a \in A$. Of course we have the same equivalence relation on B. The open L-neighborhoods of the equivalence clases give an open cover of T with Lebesgue number L and multiplicity two. \Box

Proposition 3.5 Assume $X \subseteq \mathbb{R}^n$ contains arbitrarily big balls, i.e. for each $n \in \mathbb{N}$ there is a point $x_n \in X$ such that the ball $B_{\mathbb{R}^n}(x_n, n)$ with center x_n and radius n is contained in X. In this case $\operatorname{asdim}(X, d_{eucl.}) = n$.

Proof. First we prove $\operatorname{asdim}(\mathbb{R}^n, d_{eucl.}) \leq n$. Then an application of Remark 3.3 leads to $\operatorname{asdim}(X, d_{eucl.}) \leq n$.

By $Q_a(x) := \{y \in \mathbb{R}^n \mid |y_i - x_i| < \frac{a}{2} \text{ for all } i \in \{1, \ldots, n\}\}$ we denote the specified *n*-cube around $x \in \mathbb{R}^n$ with edges of length a > 0. We define $v := (1, \ldots, 1) \in \mathbb{Z}^n$ and consider the following n + 1 families of disjoint, open cubes:

$$Q_i := \left\{ Q_a \left(a \cdot \left(z + \frac{i}{n+1} \cdot v \right) \right) \mid z \in \mathbb{Z}^n \right\} \quad \text{for } i \in \{0, \dots, n\}$$

The cover $\mathcal{U}_a := \mathcal{Q}_0 \cup \cdots \cup \mathcal{Q}_n$ is uniformly bounded and open and has multiplicity $\mu(\mathcal{U}_a) = n+1$. We calculate the Lebesgue number of this cover. The boundaries of the covering sets decompose \mathbb{R}^n into cubes with edges of length $\frac{a}{n+1}$. Any *n*-cube is limited by 2n faces. Now take $x \in \mathbb{R}^n$. Choose the small closed cube x is belonging to. At most n of its limiting faces are at a distance not exceeding $\frac{a}{2\cdot(n+1)}$. There is $i \in \{0, \ldots, n\}$ such that a cube of \mathcal{Q}_i contains all these faces. Therefore $B_{\mathbb{R}^n}\left(x, \frac{a}{2\cdot(n+1)}\right)$ is contained in this cube. If we want the cover \mathcal{U}_a to have Lebesgue number $L(\mathcal{U}_a) \geq L$, we just choose $a \geq 2 \cdot (n+1) \cdot L$. Now suppose $\operatorname{asdim}(X, d_{n-1}) =: k < n$. There is a uniformly bounded open

Now suppose $\operatorname{asdim}(X, d_{eucl.}) =: k < n$. There is a uniformly bounded, open cover \mathcal{U} of X with multiplicity $\mu(X) \leq k + 1 < n + 1$. Let d be the mesh of the cover. Let $\varepsilon > 0$. We get an open cover of $B_{\mathbb{R}^n}(0,1)$ with sets of diameter at most ε and of multiplicity not exceeding k + 1 by translating a ball of diameter $\frac{d}{\varepsilon}$ contained in X to the origin and multiplying with $\frac{\varepsilon}{d}$. Applying Theorem 1.6.12 of [Eng78] yields dim $(B_{\mathbb{R}^n}(0,1)) \leq k < n$, but this is wrong. Thus $\operatorname{asdim}(X, d_{eucl.}) \geq n$.

Corollary 3.6 asdim $(\mathbb{R}^n, d_{eucl.}) = n$

3.2 Asymptotic dimension of coarse spaces

In [Roe03], John Roe generalized Definition 3.2. We are now going to generalize Definition 3.1.

Definition 3.7 (asymptotic dimension of coarse spaces)

Let (X, \mathcal{E}) be a coarse space.

- Let $L \in \mathcal{E}$ be an entourage and \mathcal{U} a cover of X. We say that \mathcal{U} has appetite L if $\[\forall \quad \exists \\ u \in \mathcal{U} \\ U \in \mathcal{U} \\ U \in \mathcal{U} \\ \end{bmatrix} L(x) \subseteq U.$
- We call a cover \mathcal{U} uniformly bounded if $\Delta_{\mathcal{U}} := \bigcup_{U \in \mathcal{U}} U \times U$ is an entourage.
- Let $n \in \mathbb{N}$. We say $\operatorname{asdim}(X, \mathcal{E}) \leq n$ if for every^1 entourage $L \in \mathcal{E}$ there exists a cover \mathcal{U} of X such that
 - (1) the multiplicity $\mu(\mathcal{U})$ is at most n+1,
 - (2) \mathcal{U} has appetite L and
 - (3) \mathcal{U} is uniformly bounded.

Remark 3.8 If (X, d) is a pseudometric space and \mathcal{E}_d the corresponding bounded coarse structure, then $\operatorname{asdim}(X, d)$ as defined in Definition 3.1 and $\operatorname{asdim}(X, \mathcal{E}_d)$ coincide.

Definition 3.9 Let G be a finitely generated group. Lemma 1.2 yields that different word metrics on G all induce the same coarse structure \mathcal{E}_G . We define the asymptotic dimension of the finitely generated group G.

$$\operatorname{asdim}(G) := \operatorname{asdim}(G, \mathcal{E}_G)$$

Proposition 3.10 A cover \mathcal{U} is uniformly bounded if and only if

$$\underset{D \in \mathcal{E}}{\exists} \quad \forall \underset{U \in \mathcal{U}}{\exists} \quad U \subseteq D(x) .$$
 (3.1)

Proof. Let (X, \mathcal{E}) be a coarse space and let \mathcal{U} be a cover of X. Defining $D := \Delta_{\mathcal{U}}$ we see that (3.1) follows from being uniformly bounded.

Suppose conversely that we have an entourage D as in (3.1). Now we conclude that $\bigcup_{U \in \mathcal{U}} U \times U \subseteq DD^{-1} \in \mathcal{E}$, i.e. \mathcal{U} is uniformly bounded. \Box

Example 3.11 Let X be a Hausdorff space. We call the collection

$$\mathcal{T} := \mathbb{C}\left(\left\{K \subseteq X \times X \mid K \text{ compact } \right\}\right)$$

the trivial coarse structure on X. Observe that this coarse structure is not compatible with the topology if X is not compact and points in X are not open. The asymptotic dimension of (X, \mathcal{T}) is zero.

¹An entourage $L \in \mathcal{E}$ is called symmetric if $L = L^{-1}$. We need to consider only symmetric entourages which contain the diagonal, because for any entourage $L \in \mathcal{E}$ we have $L \subseteq L \cup L^{-1} \cup \Delta_X \in \mathcal{E}$.

Proof. The projections $\pi_i \colon X \times X \to X, (x_1, x_2) \mapsto x_i$ are continuous. Let $L \in \mathcal{E}$ be a symmetric entourage and define $K := \pi_1(\overline{L \setminus \Delta_X})$. Then $\mathcal{U} := \{K\} \cup \{\{x\} \mid x \notin K\}$ is a uniformly bounded cover with appetite Land multiplicity zero.

Remark 3.12 Note that the covers in Definition 3.7 do not have to be open. If we demand the covers to be open, we obtain $\operatorname{asdim}_{open}(X, \mathcal{T}) = \infty$ for every connected, non-compact Hausdorff space X, because a cover of X cannot be open and uniformly bounded with respect to \mathcal{T} at the same time.

Comparison with Roe's definition

The following is a redraft of Roe's definition of asymptotic dimension.

Definition 3.13 Let (X, \mathcal{E}) be a coarse space. We say $\operatorname{asdim}_{Roe}(X, \mathcal{E}) \leq n$ if for every entourage $L \in \mathcal{E}$ there is a cover \mathcal{U} of X such that

- (1) $\mathcal{U} = \mathcal{U}_1 \cup \cdots \cup \mathcal{U}_{n+1},$
- (2) each of the families $\mathcal{U}_1, \ldots, \mathcal{U}_{n+1}$ is *L*-disjoint (i.e. whenever $A, B \in \mathcal{U}_i$ and $A \neq B$, then $A \times B \cap L = \emptyset$) and
- (3) \mathcal{U} is uniformly bounded.

There is a small difference between Definition 3.13 and the definition given in [Roe03]. In Roe's original definition the cover \mathcal{U} is supposed to be countable. We will not make any assumptions on the cardinality of \mathcal{U} .

Remark 3.14 Let (X, d) be a pseudometric space and \mathcal{E} the corresponding bounded coarse structure. It is easy to see that $\operatorname{asdim}_{Roe}(X, \mathcal{E})$ and the asymptotic dimension of (X, d) as defined in Definition 3.2 coincide.

A third version of asymptotic dimension will appear in Theorem 3.16.

Definition 3.15 asdim_{fam} $(X, \mathcal{E}) \leq n$ if for every entourage $L \in \mathcal{E}$ there is a cover \mathcal{U} of X such that (1) $\mathcal{U} = \mathcal{U}_1 \cup \cdots \cup \mathcal{U}_{n+1}$ where each of the families \mathcal{U}_i consists of disjoint sets, (2) \mathcal{U} has appetite L and (3) \mathcal{U} is uniformly bounded.

Theorem 3.16 Let (X, \mathcal{E}) be a coarse space. Then

 $\operatorname{asdim}(X, \mathcal{E}) = \operatorname{asdim}_{Roe}(X, \mathcal{E}) = \operatorname{asdim}_{fam}(X, \mathcal{E})$.

Proof. We first prove $\operatorname{asdim}_{Roe} \geq \operatorname{asdim}_{fam}$. Assume $\operatorname{asdim}_{Roe}(X, \mathcal{E}) = n \in \mathbb{N}$. Let L be a symmetric entourage which contains the diagonal. For $L^2 := LL \in \mathcal{E}$, there exists a cover \mathcal{U} as in Definition 3.13. Since $A \times B \cap L^2 = \emptyset$ is equivalent to $L[A] \cap L[B] = \emptyset$, the cover $\mathcal{U}_L := \{L[U] \mid U \in \mathcal{U}\}$ meets

all conditions required in Definition 3.15. Note that $\bigcup_{U \in \mathcal{U}} L[U] \times L[U] \subseteq L(\bigcup_{U \in \mathcal{U}} U \times U) L^{-1} \in \mathcal{E}.$

In a second step we have to prove $\operatorname{asdim}_{fam} \geq \operatorname{asdim}$, but this is obvious, since condition (1) of Definition 3.15 implies condition (1) of Definition 3.7. It remains to prove $\operatorname{asdim} \geq \operatorname{asdim}_{Roe}$. For this purpose we need to construct a uniformly bounded cover \mathcal{V} consisting of *L*-disjoint families from a uniformly bounded cover \mathcal{U} with appetite L^{n+1} . The idea is to take all intersections of n + 1 sets from \mathcal{U} as one family, the intersections of exactly n sets from \mathcal{U} as a second family, etc. However, we have to ensure these families to be *L*-disjoint.

Assume that $\operatorname{asdim}(X, \mathcal{E}) = n \in \mathbb{N}$. Let $L \in \mathcal{E}$ be a symmetric entourage that contains the diagonal. Let \mathcal{U} be a uniformly bounded cover of X with appetite L^{n+1} and multiplicity at most n + 1. For an entourage E and $U \subseteq X$ we define $Int_E(U) := \{x \in X \mid E(x) \subseteq U\}$. Observe that $E_1 \subseteq E_2$ implies $Int_{E_2}(U) \subseteq Int_{E_1}(U)$. Some more definitions are needed to get \mathcal{V} .

$$\begin{aligned} \mathcal{U}_i &:= \{ U_1 \cap \dots \cap U_i \mid U_1, \dots, U_i \in \mathcal{U} \text{ pairwise distinct } \} \\ S_i &:= \bigcup_{U \in \mathcal{U}_i} Int_{L^{n+2-i}}(U) \\ \mathcal{V}_i &:= \{ Int_{L^{n+2-i}}(U) \setminus S_{i+1} \mid U \in \mathcal{U}_i \} \\ \mathcal{V} &:= \mathcal{V}_1 \cup \dots \cup \mathcal{V}_{n+1} \end{aligned}$$

Now \mathcal{V} is a cover of X. Actually, \mathcal{V} is a refinement of the cover \mathcal{U} . Therefore \mathcal{V} is uniformly bounded.

It remains to prove that each of the families $\mathcal{V}_1, \ldots, \mathcal{V}_{n+1}$ is *L*-disjoint. Let $A, B \in \mathcal{V}_i$ such that $A \neq B$. There are $A_1, \ldots, A_i, B_1, \ldots, B_i \in \mathcal{U}$ such that $A = Int_{L^{n+2-i}}(A_1 \cap \cdots \cap A_i) \setminus S_{i+1}$ and $B = Int_{L^{n+2-i}}(B_1 \cap \cdots \cap B_i) \setminus S_{i+1}$. The sets A_1, \ldots, A_i are supposed to be pairwise distinct as are the sets B_1, \ldots, B_i .

Let $(a, b) \in A \times B \cap L$ and observe the following facts.

$$a, b \notin S_{i+1} \tag{3.2}$$

$$a \in A \subseteq Int_{L^{n+1-i}}(A_1 \cap \dots \cap A_i) \tag{3.3}$$

$$b \in B \subseteq Int_{L^{n+1-i}}(B_1 \cap \dots \cap B_i)$$
(3.4)

$$a \in L[B] \subseteq L[Int_{L^{n+2-i}}(B_1 \cap \dots \cap B_i)]$$

$$(3.5)$$

$$b \in L[A] \subseteq L[Int_{L^{n+2-i}}(A_1 \cap \dots \cap A_i)]$$

$$(3.6)$$

Since $L[Int_{L^{j}}(U)] = \{x \mid \exists_{y \in X} L^{j}(y) \subseteq U, x \in L(y)\} \subseteq \{x \mid L^{j-1}(x) \subseteq U\} = Int_{L^{j-1}}(U)$, we get the following conclusions from (3.5) and (3.6):

 $a \in Int_{L^{n+1-i}}(B_1 \cap \dots \cap B_i)$ $b \in Int_{L^{n+1-i}}(A_1 \cap \dots \cap A_i)$

Finally $a, b \in Int_{L^{n+2-(i+1)}}(A_1 \cap \cdots \cap A_i \cap B_1 \cap \cdots \cap B_i)$. Since $A \neq B$, we know that the set $\{A_1, \ldots, A_i, B_1, \ldots, B_i\}$ contains at least i + 1 elements. Thus $a, b \in S_{i+1}$, but this is a contradiction to (3.2).

Remark 3.17 For further reference we record the results of our constructions in the previous proof. Let (X, \mathcal{E}) be a coarse space, $L \in \mathcal{E}$ and $n \in \mathbb{N}$.

- If we are given a uniformly bounded cover \mathcal{U} of X such that $\mathcal{U} = \mathcal{U}_1 \cup \cdots \cup \mathcal{U}_{n+1}$ and each of the families $\mathcal{U}_1, \ldots, \mathcal{U}_{n+1}$ is L^2 -disjoint, we get the uniformly bounded cover $\mathcal{U}_L := \{L[U] \mid U \in \mathcal{U}\}$ consisting of n+1 families of disjoint subsets of X and with appetite L.
- Given a uniformly bounded cover \mathcal{U} of X with multiplicity n + 1 and appetite L^{n+1} , we constructed a uniformly bounded cover \mathcal{V} such that $\mathcal{V} = \mathcal{V}_1 \cup \cdots \cup \mathcal{V}_{n+1}$ and each of the families $\mathcal{V}_1, \ldots, \mathcal{V}_{n+1}$ is L-disjoint.

Suppose X is a metric space and \mathcal{E} its bounded coarse structure. Given a uniformly bounded cover with multiplicity m and Lebesgue number L, the previous construction yields a uniformly bounded cover which consists of m families of disjoint subsets of X and with Lebesgue number $\frac{L}{2n+2}$.

Asymptotic dimension via anti-Čech systems

The asymptotic dimension of proper metric spaces can be characterized using anti-Čech systems. Compare Theoreom 9.9 of [Roe03]. We will see that this characterisation does not work for all coarse spaces. However, it does work for coarse structures induced from metrizable compactifications. We recall some definitions from John Roe's book.

Definition 3.18 Let (X, \mathcal{E}) be a coarse space.

- A uniformly bounded cover \mathcal{U} of X is called *uniform* if each bounded subset of X meets only finitely many elements of \mathcal{U} .
- A collection of uniform covers is called an anti-Cech system if for every entourage $L \in \mathcal{E}$ it contains a cover having appetite L.
- We define a partial order on an anti-Čech system. $\mathcal{U} \leq \mathcal{V}$ if \mathcal{V} has appetite $\Delta_{\mathcal{U}}$. This way, any anti-Čech system becomes a directed set.

Example 3.19 Consider the coproduct $\coprod_{n\in\mathbb{N}}(\mathbb{R}, \mathcal{E}_{eucl.})$. According to Proposition 3.30, this coarse space has asymptotic dimension one. On the other hand, for every uniformly bounded cover there exists a copy of \mathbb{R} where the cover consists of the sets $\{x\}$ for $x \in \mathbb{R}$. Hence, there don't exist uniform covers for this coarse space and the following proposition can not be true in general.²

 $^{^{2}}$ Even if this coarse space does not admit an anti-Čech system, it does have a coarsening sequence in the sense of [Mit03], i.e. this space belongs to the category where coarse homology theory can be defined.

Proposition 3.20 Let X be a Hausdorff space and let K be a metrisable compactification of X.³ Let \mathcal{E} be the coarse structure induced by K. Then $\operatorname{asdim}(X, \mathcal{E}) \leq n$ if and only if there is an anti-Čech system consisting of covers with multiplicity at most n + 1.

Remark 3.21 If we take just uniformly bounded covers in the definition of anti-Čech systems, Proposition 3.20 is true for all coarse spaces - just by definitions.

Proof of Proposition 3.20. Because of Remark 3.21, we only need to prove that we can choose all covers to be uniform.

Let us call a cover \mathcal{V} a nice refinement of a cover \mathcal{U} if we have an injective map $i: \mathcal{V} \to \mathcal{U}$ such that $V \subseteq i(V)$ for all $V \in \mathcal{V}$.

Let L be a symmetric entourage that contains the diagonal and let \mathcal{U} be a uniformly bounded cover of X consisting of n+1 families each of them being L^2 -disjoint. Note that every nice refinement of \mathcal{U} has the same properties.

Since K is metrisable, we can write $X = \bigcup_{i \in \mathbb{N}} K_i$ with K_i compact and $K_1 \subseteq K_2 \subseteq \cdots \subseteq X$. We proceed inductively in order to define a nice refinement of \mathcal{U} which finally will lead to a uniform cover of appetite L. Set $\mathcal{U}_0 := \mathcal{U}$. Observe that $L[K_i]$ can be covered by a finite subcover $\mathcal{U}_{fin,i}$ of \mathcal{U}_{i-1} . Define

$$\mathcal{U}_i := \mathcal{U}_{fin,i} \cup \{ U \setminus L[K_i] \mid U \in \mathcal{U}_{i-1} \setminus \mathcal{U}_{fin,i} \}.$$

The cover \mathcal{U}_i is a nice refinement of \mathcal{U}_{i-1} . Finally, we get a nice refinement

$$\mathcal{U}_{red} := \{ U \mid \exists_{k \in \mathbb{N}} \forall_{i \ge k} U \in \mathcal{U}_{fin,i} \}$$

of the original cover \mathcal{U} . In order to see that \mathcal{U}_{red} is indeed a cover of X, let $x \in K_i \subseteq X$. The union of all $U \in \mathcal{U}_{fin,i}$ containing x is bounded and therefore contained in K_j for some $j \in \mathbb{N}$. It follows that any $U \in \mathcal{U}_{fin,j}$ which contains x is an element of \mathcal{U}_{red} .

Observe that $\mathcal{V}_L := \{L[U] \mid U \in \mathcal{U}_{red}\}$ is a uniformly bounded cover of X with appetite L and multiplicity at most n + 1. Moreover \mathcal{V}_L is uniform. In order to see this, let $B \subseteq X$ be bounded. This implies $B \subseteq K_i$ for some $i \in \mathbb{N}$. Hence $B \cap U \neq \emptyset$ for most N_i different $U \in \mathcal{U}_{red}$ where N_i is the cardinality of $\mathcal{U}_{fin,i}$.

Thus, the covers \mathcal{V}_L for symmetric entourages L which contain the diagonal form the desired anti-Čech system.

3.3 Basic properties of asymptotic dimension

Theorem 3.22 If $f: (X, \mathcal{E}) \to (Y, \mathcal{F})$ is a coarse embedding, then

 $\operatorname{asdim}(X, \mathcal{E}) \leq \operatorname{asdim}(Y, \mathcal{F}).$

³This implies that X is locally compact and paracompact. Note that a compact set K is metrisable if and only if its topology has a countable base.

Proof. Suppose that $n := \operatorname{asdim}(Y, \mathcal{F}) < \infty$. Let $E \in \mathcal{E}$ be an entourage and set $F := f \times f(E)$. Note that $E \subseteq (f \times f)^{-1}(F)$. There is a uniformly bounded cover \mathcal{U} of Y with appetite F and multiplicity at most n + 1. The inverse image of \mathcal{U} is a uniformly bounded cover of X with appetite E and the same multiplicity as \mathcal{U} .

Corollary 3.23 (monotony of asymptotic dimension) Let (X, \mathcal{E}) be a coarse space and $A \subseteq X$. Remember that the inclusion map is a coarse embedding. Hence $\operatorname{asdim}(A, \mathcal{E}|_A) \leq \operatorname{asdim}(X, \mathcal{E})$.

Corollary 3.24 (coarse invariance of asymptotic dimension) Asymptotic dimension is a coarse invariant, i.e. given a coarse equivalence $f: (X, \mathcal{E}_X) \to (Y, \mathcal{E}_Y)$, we have $\operatorname{asdim}(X, \mathcal{E}_X) = \operatorname{asdim}(Y, \mathcal{E}_Y)$.

Definition 3.25 Let (X, \mathcal{E}) be a coarse space and $A \subseteq X$. We call A a substantial part of X if $\operatorname{asdim}(A, \mathcal{E}|_A) = \operatorname{asdim}(X, \mathcal{E})$.

For a coarsely uniform map $(X, \mathcal{E}) \to (Y, \mathcal{F})$ which is also injective, there is no relation between the asymptotic dimensions of (X, \mathcal{E}) and (Y, \mathcal{F}) .

Example 3.26 Let (X, \mathcal{E}) be a coarse space. Observe that the power set $\mathcal{P}(X \times X)$ is a coarse structure on X. The map $id: (X, \mathcal{E}) \to (X, \mathcal{P}(X \times X))$ is coarsely uniform, but $\operatorname{asdim}(X, \mathcal{E}) \ge 0 = \operatorname{asdim}(X, \mathcal{P}(X \times X))$.

Example 3.27 Let $n \in \{2, 3, ...\}$. By \mathcal{E} we denote the coarse structure coming from the one-point compactification of \mathbb{R}^n and by $\mathcal{E}_{eucl.}$ the bounded coarse structure corresponding to the euclidean metric of \mathbb{R}^n . The map $id: (\mathbb{R}^n, \mathcal{E}_{eucl.}) \to (\mathbb{R}^n, \mathcal{E}_{.})$ is coarse, but $\operatorname{asdim}(\mathbb{R}^n, \mathcal{E}_{eucl.}) = n > 1 = \operatorname{asdim}(\mathbb{R}^n, \mathcal{E}_{.}).$

Proof. Since $\mathcal{E}_{eucl.} \subseteq \mathcal{E}_{\cdot}$, the map *id* is coarsely uniform. A set *B* is bounded with respect to $\mathcal{E}_{eucl.}$ if and only if *B* is precompact. The same is true for \mathcal{E}_{\cdot} . Thus *id* is coarsely proper.

It remains to prove $\operatorname{asdim}(\mathbb{R}^n, \mathcal{E}) = 1$. Note that

$$|\cdot| : (\mathbb{R}^n, \mathcal{E}_{\cdot}) \to (\mathbb{R}_+, \mathcal{E}_{\cdot})$$
$$x \mapsto |x|$$

is a coarse equivalence with inverse $i: (\mathbb{R}_+, \mathcal{E}_{\cdot}) \to (\mathbb{R}^n, \mathcal{E}_{\cdot}), r \mapsto (r, 0, \dots, 0)$. In fact $|\cdot| \circ i = id_{\mathbb{R}^n}$ and $i \circ |\cdot|$ is close to $id_{\mathbb{R}^n}$ with respect to \mathcal{E}_{\cdot} . This yields $\operatorname{asdim}(\mathbb{R}^n, \mathcal{E}_{\cdot}) = \operatorname{asdim}(\mathbb{R}_+, \mathcal{E}_{\cdot})$. It remains to calculate the asymptotic dimension of $(\mathbb{R}_+, \mathcal{E}_{\cdot})$. Since $(\mathbb{R}_+, \mathcal{E}_{\cdot})$ is a ray, we can apply Corollary 4.11. As an alternative, we refer to Example 9.7 of [Roe03].

Proposition 3.28 Let (X, \mathcal{E}) be a coarse space and $\mathcal{E}_{cn} = {}_{cn} \mathfrak{C}(\mathcal{E})$ the connected coarse structure generated by \mathcal{E} . Then $\operatorname{asdim}(X, \mathcal{E}) = \operatorname{asdim}(X, \mathcal{E}_{cn})$.

Proof. Suppose $n := \operatorname{asdim}(X, \mathcal{E}_{cn}) < \infty$. Let $E \in \mathcal{E} \subseteq \mathcal{E}_{cn}$. There is a cover \mathcal{U} of X with appetite E and multiplicity at most n + 1 which is uniformly bounded with respect to \mathcal{E}_{cn} . Each $U \in \mathcal{U}$ can be written as the disjoint union of finitely many sets U_1, \ldots, U_k which are bounded with respect to \mathcal{E} and such that the union of any two of the sets U_1, \ldots, U_k is not bounded with respect to \mathcal{E} . Define $\operatorname{comp}(U) := \{U_1, \ldots, U_k\}$ and observe that $\mathcal{U}' := \bigcup_{U \in \mathcal{U}} \operatorname{comp}(U)$ is a cover of X with multiplicity at most n + 1. Furthermore, \mathcal{U}' has appetite E and is uniformly bounded with respect to \mathcal{E} . Hence $\operatorname{asdim}(X, \mathcal{E}) \leq n$.

Set $n := \operatorname{asdim}(X, \mathcal{E})$. Let $E \in \mathcal{E}_{cn}$ be a symmetric entourage. This implies that $E = E' \cup (A_1 \times A_{\sigma(1)}) \cup \cdots \cup (A_k \times A_{\sigma(k)})$ with $E' \in \mathcal{E}, k \in \mathbb{N}, A_1, \ldots, A_k$ bounded subsets of X (not necessarily pairwise distinct) and σ a permutation of the set $\{1, \ldots, k\}$ with the additional property $\sigma \circ \sigma = \operatorname{id}$. Set $M := \Delta_X \cup A_1^2 \cup \cdots \cup A_k^2$ and observe that $E'' := (E' \cup \Delta_X)M \in \mathcal{E}$. Let \mathcal{U}'' be a cover of X which is uniformly bounded with respect to \mathcal{E} and which has appetite E'' and multiplicity at most n + 1. Note that there are sets $U_1, \ldots, U_k \in \mathcal{U}''$ such that $E'[A_i] \cup A_i \subseteq U_i$. We define the cover $\mathcal{U} := \mathcal{U}'' \cup \{U_1 \cup \cdots \cup U_k\} \setminus \{U_1, \ldots, U_k\}$ of X. Observe that \mathcal{U} is uniformly bounded with respect to \mathcal{E}_{cn} and has multiplicity at most n + 1.

Moreover, \mathcal{U} has appetite E. To see this, let $x \in X$. If $x \in A_i$, then $E(x) = E'(x) \cup A_{\sigma(i)} \subseteq U_i \cup U_{\sigma(i)} \subseteq U_1 \cup \cdots \cup U_k$. If $x \notin \{A_1 \cup \cdots \cup A_k\}$, then $E(x) = E'(x) \subseteq E''(x)$.

Proposition 3.29 (asymptotic dimension of finite unions)

Let (X, \mathcal{E}) be a coarse space and $A, B \subseteq X$ with $A \cup B = X$.

 $\operatorname{asdim}(X, \mathcal{E}) = \max\{\operatorname{asdim}(A, \mathcal{E}|_A), \operatorname{asdim}(B, \mathcal{E}|_B)\}$

Proof. The proof of \geq follows from monotony. To see \leq , we generalize an argument of Bell and Dranishnikov (see [BD01]).

Let *n* be the maximum of $\operatorname{asdim}(A, \mathcal{E}|_A)$ and $\operatorname{asdim}(B, \mathcal{E}|_B)$ and take a symmetric entourage $L \in \mathcal{E}$ which contains Δ_X . For $\mathcal{U} \subseteq \mathcal{P}(X)$ and $V \subseteq X$ we define

$$N_L(V, \mathcal{U}) := V \cup \bigcup_{\substack{U \in \mathcal{U} \\ L \cap U \times V \neq \emptyset}} U.$$

There is a uniformly bounded cover $\mathcal{U} = \mathcal{U}_1 \cup \cdots \cup \mathcal{U}_{n+1}$ of A consisting of L-disjoint families \mathcal{U}_i . Moreover, there is a uniformly bounded cover $\mathcal{V} = \mathcal{V}_1 \cup \cdots \cup \mathcal{V}_{n+1}$ of B consisting of $(L\Delta_{\mathcal{U}}L\Delta_{\mathcal{U}}L)$ -disjoint families \mathcal{V}_i . For $i \in \{1, \ldots, n+1\}$ set

$$\mathcal{W}_i := \left\{ N_L(V, \mathcal{U}_i) \mid V \in \mathcal{V}_i \right\} \cup \left\{ U \in \mathcal{U}_i \mid L \cap U \times V = \emptyset \text{ for all } V \in \mathcal{V}_i \right\}.$$

Observe that $N_L(V, \mathcal{U}_i) \subseteq \Delta_{\mathcal{U}} L[V]$. Hence, we get a uniformly bounded cover $\mathcal{W} = \mathcal{W}_1 \cup \cdots \cup \mathcal{W}_{n+1}$ of X where \mathcal{W}_i is L-disjoint for $1 \leq i \leq n+1$. This proves $\operatorname{asdim}(X, \mathcal{E}) \leq n$.

Proposition 3.30 (asymptotic dimension of coproducts)

Let Λ be any set and $(X_{\lambda}, \mathcal{E}_{\lambda})$ a coarse space for every $\lambda \in \Lambda$. Define $X := \coprod_{\lambda \in \Lambda} X_{\lambda}$. If (X, \mathcal{E}) and (X, \mathcal{E}_{cn}) are the coproducts in the categories \mathcal{D} and \mathcal{D}_{cn} respectively, then

asdim
$$(X, \mathcal{E}_{cn})$$
 = asdim (X, \mathcal{E}) = $\sup_{\lambda \in \Lambda} \operatorname{asdim}(X_{\lambda}, \mathcal{E}_{\lambda})$.

Proof. Set $n := \sup_{\lambda \in \Lambda} \operatorname{asdim}(X_{\lambda}, \mathcal{E}_{\lambda})$. Monotony of asymptotic dimension implies $\operatorname{asdim}(X, \mathcal{E}_{cn}) \ge n$ and $\operatorname{asdim}(X, \mathcal{E}) \ge n$.

We will now prove $\operatorname{asdim}(X, \mathcal{E}) \leq n$. Take an entourage $L \in \mathcal{E}$ which contains Δ_X . Then there are $\lambda_1, \ldots, \lambda_k \in \Lambda$ and $L_{\lambda_i} \in \mathcal{E}_{\lambda_i}$ such that $L = L_{\lambda_1} \cup \cdots \cup L_{\lambda_k} \cup \Delta_X$. For $i \in \{1, \ldots, k\}$ choose a uniformly bounded cover \mathcal{U}_{λ_i} of X_{λ_i} with appetite L_{λ_i} and multiplicity at most $\operatorname{asdim}(X_{\lambda_i}, \mathcal{E}_{\lambda_i}) + 1$. For $\lambda \in \Lambda \setminus \{\lambda_1, \ldots, \lambda_k\}$ set $\mathcal{U}_{\lambda} := \{\{x\} \mid x \in X_{\lambda}\}$. The union $\mathcal{U} := \bigcup_{\lambda \in \Lambda} \mathcal{U}_{\lambda}$ is a uniformly bounded cover of X with appetite L whose multiplicity does not exceed n + 1.

For the equality $\operatorname{asdim}(X, \mathcal{E}_{cn}) = \operatorname{asdim}(X, \mathcal{E})$ compare Proposition 3.28. \Box

Proposition 3.31 (asymptotic dimension of disjoint unions)

Let Λ be any set and $(X_{\lambda}, \mathcal{E}_{\lambda})$ a coarse space for every $\lambda \in \Lambda$.

asdim
$$\left(\bigsqcup_{\lambda \in \Lambda} X_{\lambda}, \mathcal{E}_{\sqcup}\right) = \sup_{\lambda \in \Lambda} \operatorname{asdim}(X_{\lambda}, \mathcal{E}_{\lambda})$$

Proof. The proof of \geq follows from monotony of asymptotic dimension. Set $X := \bigsqcup_{\lambda \in \Lambda} X_{\lambda}$ and $n := \sup_{\lambda \in \Lambda} \operatorname{asdim}(X_{\lambda}, \mathcal{E}_{\lambda})$. In order to prove the inequality $\operatorname{asdim}(X, \mathcal{E}_{\sqcup}) \leq n$, let $L \in \mathcal{E}_{\sqcup}$. Set $L_{\lambda} := L \cap X_{\lambda}^2 \in \mathcal{E}_{\lambda}$ and choose uniformly bounded covers \mathcal{U}_{λ} of X_{λ} with appetite L_{λ} and multiplicity at most n + 1. Now $\mathcal{U} := \bigcup_{\lambda \in \Lambda} \mathcal{U}_{\lambda}$ is a uniformly bounded cover of X with appetite L and multiplicity at most n + 1. \Box

Proposition 3.32 (asdim of certain direct limits) Let $j \in I$. If I is a directed set and if for all $k \ge j$ the map $f_k \colon X_k \to \underline{\lim} X_i$ is injective, then

asdim
$$\left(\underbrace{\lim}_{k \in \mathcal{E}_i} (X_i, \mathcal{E}_i) \right) = \sup_{k \geq j} \operatorname{asdim}(X_k, \mathcal{E}_k)$$
.

Proof. With Proposition 1.27 in mind, the proof is not difficult. Observe that injectivity of f_k for all $k \ge j$ implies that f_i is a coarse embedding for $i \ge j$. The proof of \ge now is given by applying Theorem 3.22.

We write (X, \mathcal{E}) for the direct limit. Let $L \in \mathcal{E}$ be an entourage which contains Δ_X . Proposition 1.27 implies that L is the union of Δ_X and an entourage $L_i \in \mathcal{E}_i$ for some $i \in I$. We may assume $i \geq j$. If \mathcal{U}_i is a uniformly bounded cover of X_i with appetite L_i and multiplicity at most $\operatorname{asdim}(X_i, \mathcal{E}_i) + 1$, then $\mathcal{U} := \mathcal{U}_i \cup \{\{x\} \mid x \in X \setminus X_i\}$ is a uniformly bounded cover of X with appetite L. Moreover \mathcal{U} has the same multiplicity as \mathcal{U}_i . \Box

Proposition 3.33 (asymptotic dimension of products)

Let (X, \mathcal{E}_X) and (Y, \mathcal{E}_Y) be coarse spaces.

 $\operatorname{asdim}(X \times Y, \mathcal{E}_X * \mathcal{E}_Y) \leq \operatorname{asdim}(X, \mathcal{E}_X) + \operatorname{asdim}(Y, \mathcal{E}_Y)$ $\operatorname{asdim}(X, \mathcal{E}_X) \leq \operatorname{asdim}(X \times Y, \mathcal{E}_X * \mathcal{E}_Y) \quad \text{if } Y \neq \emptyset$

Proof. Compare [Roe03] for the special case of bounded coarse structures and Remark 5.8 for the case of continuously controlled coarse structures induced by metrisable compactifications.

Set $n := \operatorname{asdim}(X)$ and $m := \operatorname{asdim}(Y)$. Let $E \in \mathcal{E}_X * \mathcal{E}_Y$. There are symmetric entourages $E_X \in \mathcal{E}_X$ and $E_Y \in \mathcal{E}_Y$ containing the diagonals Δ_X and Δ_Y respectively such that $E \subseteq E_X \times E_Y$.

There is a uniformly bounded cover \mathcal{U} of X with appetite E_X^{n+m+1} and multiplicity $\mu(\mathcal{U}) \leq n+1$. There is also a uniformly bounded cover \mathcal{V} of Ywith appetite E_Y^{n+m+1} and multiplicity $\mu(\mathcal{V}) \leq m+1$. We get a uniformly bounded cover $\mathcal{U} \times \mathcal{V} := \{U \times V \mid U \in \mathcal{U}, V \in \mathcal{V}\}$ of $X \times Y$ with appetite E^{n+m+1} and multiplicity $\leq (n+1) \cdot (m+1) = n \cdot m + n + m + 1$. Thus, we need to improve the multiplicity.

We proceed similar as in the proof of Theorem 3.16 and begin with some definitions. Let $k \in \{2, ..., n + m + 2\}$.

$$\mathcal{A}_{k} := \{U_{1} \cap \dots \cap U_{p} \times V_{1} \cap \dots \cap V_{q} \mid p+q = k, U_{i} \in \mathcal{U}, V_{i} \in \mathcal{V} \text{ pairwise distinct } \}$$

$$B_{k} := \bigcup_{A \in \mathcal{A}_{k}} Int_{E^{n+m+3-k}}(A) \quad \text{and} \quad B_{n+m+3} := \emptyset$$

$$\mathcal{W}_{k} := \{Int_{E^{n+m+3-k}}(U) \setminus B_{k+1} \mid U \in \mathcal{A}_{k}\}$$

$$\mathcal{W} := \mathcal{W}_{2} \cup \dots \cup \mathcal{W}_{n+m+2}$$

Notice that \mathcal{W} is a uniformly bounded cover of $X \times Y$ consisting of the n + m + 1 disjoint families $\mathcal{W}_2, \ldots, \mathcal{W}_{n+m+2}$. It remains to prove that \mathcal{W}_k is *E*-disjoint for $k \in \{2, \ldots, n+m+2\}$.

For this purpose let $M, N \in \mathcal{W}_k$ with $M \neq N$ and suppose $M \times N \cap E \neq \emptyset$. Choose $((x_M, y_M), (x_N, y_N)) \in M \times N \cap E$. There are $p_M, q_M \in \mathbb{N}$ with $p_M + q_M = k$ and $M_1, \ldots, M_{p_M} \in \mathcal{U}, M'_1, \ldots, M'_{q_M} \in \mathcal{V}$ such that

$$M = Int_{E^{n+m+3-k}} (\underbrace{M_1 \cap \dots \cap M_{p_M} \times M'_1 \cap \dots \cap M'_{q_M}}_{=:MM'}) \setminus B_{k+1}.$$

Similarly, there are $p_N, q_N \in \mathbb{N}$ with $p_N + q_N = k$ and $N_1, \ldots, N_{p_N} \in \mathcal{U}$, $N'_1, \ldots, N'_{q_N} \in \mathcal{V}$ such that

$$N = Int_{E^{n+m+3-k}} \underbrace{(N_1 \cap \dots \cap N_{p_N} \times N'_1 \cap \dots \cap N'_{q_N})}_{=:NN'} \setminus B_{k+1}$$

Observe that the following relations hold.

$$(x_M, y_M) \notin B_{k+1}$$

$$(x_M, y_M) \in M \subseteq Int_{E^{n+m+2-k}}(MM')$$

$$(x_M, y_M) \in E[N] \subseteq E[Int_{E^{n+m+3-k}}(NN')] \subseteq Int_{E^{n+m+2-k}}(NN')$$

It follows that $(x_M, y_M) \in Int_{E^{n+m+3-(k+1)}}(MM' \cap NN')$. Since $M \neq N$, the set $\{M_1, \ldots, M_{p_M}, M'_1, \ldots, M'_{q_M}, N_1, \ldots, N_{p_N}, N'_1, \ldots, N'_{q_N}\}$ contains at least k + 1 different elements. Hence $(x_M, y_M) \in B_{k+1}$. But this is a contradiction to what we found before. \Box

The equality $\operatorname{asdim}(X \times Y) = \operatorname{asdim}(X) + \operatorname{asdim}(Y)$ is not true in general. Compare Corollary 5.9.

There are no general restrictions on the asymptotic dimensions of quotients as the following three examples show.

Example 3.34 Let $X = \mathbb{R}^2$ be the real plane and \mathcal{E} the bounded coarse structure induced by the euclidean metric on \mathbb{R}^2 . Define an equivalence relation by $(x, y) \sim (x', y')$ if and only if x = x'. The quotient X/\sim is clearly \mathbb{R} and \mathcal{E}_{\sim} is the bounded coarse structure on \mathbb{R} again induced by the euclidean metric on \mathbb{R} . In this case $\operatorname{asdim}(X, \mathcal{E}) > \operatorname{asdim}(X/\sim, \mathcal{E}_{\sim})$.

Example 3.35 Let $\mathbb{F}_2 = \langle a, b | \rangle$ be the free group with two generators. We say that two words are equivalent if the number of letters *a* counted with exponents as signs is equal in both words and the same for the letter *b*. The quotient is the group \mathbb{Z}^2 . In this case $\operatorname{asdim}(\mathbb{F}_2) = 1 < 2 = \operatorname{asdim}(\mathbb{Z}^2)$.

Remark 3.36 Let (X, \mathcal{E}) be a coarse space. Assume there is an equivalence relation \sim on X. If the equivalence relation happens to be an entourage, i.e. $\{(x, y) \mid x \sim y\} \in \mathcal{E}$, then the projection onto the quotient is a coarse equivalence.

Let (X, \mathcal{E}_X) , (Y, \mathcal{E}_Y) be coarse spaces, $A \subseteq X$ and $f: A \to Y$ coarsely uniform. Attaching X to Y using f we get $X \cup_f Y := X \sqcup Y / a \sim f(a)$ and a quotient map $\pi: X \sqcup Y \to X \cup_f Y$. We denote the coarse structure on the quotient by $\mathcal{E}_{X \cup_f Y}$.

Lemma 3.37 Whenever $\operatorname{asdim}(X, \mathcal{E}_X) = \operatorname{asdim}(X \setminus A, \mathcal{E}_{X \cup_f Y}|_{X \setminus A})$, we can conclude that $\operatorname{asdim}(X \cup_f Y) = \max\{\operatorname{asdim}(X), \operatorname{asdim}(Y)\}.$

Proof. Note that $X \cup_f Y = X \setminus A \cup Y$. Thus $\operatorname{asdim}(X \cup_f Y, \mathcal{E}_{X \cup_f Y}) = \max \{\operatorname{asdim}(X \setminus A, \mathcal{E}_{X \cup_f Y}|_{X \setminus A}), \operatorname{asdim}(Y, \mathcal{E}_{X \cup_f Y}|_Y)\}$ by Proposition 3.29. In Lemma 1.36 we proved $\mathcal{E}_{X \cup_f Y}|_Y = \mathcal{E}_Y$. Together with the assumption this implies the claim. \Box

Chapter 4

Asymptotic dimension of coarse cell complexes

Coarse cell complexes

In order to define coarse cell complexes, we need to describe their building blocks. Compare the discussion in [Mit03] for a better understanding of the following definition of a ray.

Definition 4.1 Let \mathcal{E} be a connected coarse structure on \mathbb{R}_+ which is compatible with the standard topology on \mathbb{R}_+ . We call $R = (\mathbb{R}_+, \mathcal{E})$ a ray if the coarse structure \mathcal{E} satisfies the following two conditions.

• If M and N are entourages, the same is true for

$$M + N := \{ (u + x, v + y) \mid (u, v) \in M, (x, y) \in N \} .$$

• If M is an entourage, so is

$$M^{\bowtie} := \{(u,v) \mid (x,y) \in M \text{ and } (x \le u \le v \le y \text{ or } y \le v \le u \le x)\}$$

Proposition 4.2 By $\mathcal{E}_{eucl.}$ we denote the usual bounded coarse structure on \mathbb{R}_+ . The coarse space $(\mathbb{R}_+, \mathcal{E}_{eucl.})$ is a ray and $\mathcal{E}_{eucl.} \subseteq \mathcal{E}$ whenever $(\mathbb{R}_+, \mathcal{E})$ is a ray.

For a proof consult Proposition 2.5 of [Mit03].

Proposition 4.3 By \mathcal{E} we denote the continuously controlled coarse structure on \mathbb{R}_+ induced by the one-point compactification. The coarse space $(\mathbb{R}_+, \mathcal{E})$ is a ray and $\mathcal{E} \subseteq \mathcal{E}$ whenever $(\mathbb{R}_+, \mathcal{E})$ is a ray.

Proof. The inclusion $\mathcal{E} \subseteq \mathcal{E}$. is true for every coarse structure which is compatible with the topology of \mathbb{R}_+ . It remains to prove that $(\mathbb{R}_+, \mathcal{E}_-)$ is a ray. This is straightforward.

Definition 4.4 Let *R* be a ray and $n \in \mathbb{N}$.

$$S^{n-1}R := (R \amalg R)^n \times \{0\}$$
$$D^n R := (R \amalg R)^n \times R$$

We call $S^n R$ a coarse *n*-sphere and $D^n R$ a coarse *n*-cell.

There are two different coarse structures on $R_{\text{II}}R$. There is the coproduct coarse structure \mathcal{E}_{II} which is not connected and there is the connected coarse structure $_{\text{cn}}\mathbb{C}(\mathcal{E}_{\text{II}})$ generated by \mathcal{E}_{II} . The arguments in this chapter work for both coarse structures, but only equipped with the connected coarse structure the spaces $S^n R$ and $D^n R$ deserve the names sphere and cell respectively.

Definition 4.5 (coarse cell complex)

We call (Y, \mathcal{E}) a *coarse cell complex* if it is obtained by "inductively" attaching coarse cells to a disjoint union of coarse cells.

More precisely, suppose that for all $k \in \mathbb{N}$ we are given a coarse space (Y_k, \mathcal{E}_k) and a set I_k . Furthermore, for all $k \in \mathbb{N}$ and $i \in I_k$ there is

- a number $n_{k,i} \in \mathbb{N}$,
- a ray $R_{k,i}$ and
- a coarsely uniform map $f_{k,i} \colon S^{n_{k,i}-1}R_{k,i} \to Y_{k-1}$.

Define $f_k := \coprod_{i \in I_k} f_{k,i} : \coprod_{i \in I_k} S^{n_{k,i}-1} R_{k,i} \to Y_{k-1}$ and observe that f_k is coarsely uniform. If

$$Y_0 = \prod_{i \in I_0} D^{n_{0,i}} R_{0,i}$$
 and $Y_k = \left(\prod_{i \in I_k} D^{n_{k,i}} R_{k,i}\right) \cup_{f_k} Y_{k-1}$,

then $\lim_{k \to \infty} (Y_k, \mathcal{E}_k)$ is a coarse cell complex.

Define $\delta(Y) := \sup\{n_{k,i} \mid k \in \mathbb{N}, i \in I_k\}$ and call $\delta(Y)$ the *cell dimension* of the coarse cell complex Y.

Asymptotic dimension of coarse cell complexes

Our goal is to prove the following theorem.

Theorem 4.6 Let Y be a coarse cell complex. Then asdim $Y = \delta(Y) + 1$ where $\delta(Y)$ is the cell dimension of Y.

In order to get Theorem 4.6, we start with some lemmas.

Let R be a ray, X' any coarse space and $A' \subseteq X'$. Define $X := D^{n-1}R \sqcup X'$ and $A := S^{n-2}R \sqcup A'$ and denote the coarse structure of X by \mathcal{E}_X . Notice that $R^n \subseteq D^{n-1}R$. We will consider the set

$$P_n := \{ (x_1, \dots, x_n) \in \mathbb{R}^n \mid x_n > 0 \text{ and } x_i \le x_n \text{ for } i < n \} \subseteq X \setminus A.$$

Lemma 4.7 For any coarse space (Y, \mathcal{E}_Y) and any coarsely uniform map $f: (A, \mathcal{E}_X|_A) \to (Y, \mathcal{E}_Y)$, we have $\mathcal{E}_X|_{P_n} = \mathcal{E}_{X \cup_f Y}|_{P_n}$.

Proof. It follows directly from the definition of the quotient coarse structure that $\mathcal{E}_X|_{P_n} \subseteq \mathcal{E}_{X \cup_f Y}|_{P_n}$. We need to prove the converse inclusion. Let $E \in \mathcal{E}_{X \cup_f Y}|_{P_n}$. There are symmetric entourages $L_i \in \mathcal{E}_{D^{n-1}R}$ containing the diagonal $\Delta_{D^{n-1}R}$ such that $E \subseteq L_1^A \cdots L_k^A$ where $L_i^A := L_i \cup A \times A$. If we set $L := L_1 \cdots L_k$, we get $E \subseteq L_1^A \cdots L_k^A \subseteq L \cup L(A \times A)L$. Thus, it remains to prove $L(A \times A)L \cap P_n^2 \in \mathcal{E}_X|_{P_n}$.

Since the projection $p_n: P_n \to R, (x_1, \ldots, x_n) \mapsto x_n$ is coarsely uniform, the image $\widetilde{L} := p_n \times p_n(L)$ of L is in \mathcal{E}_R and compatibility with the topology implies $\widetilde{L}[0] \subseteq [0, r_E]$. We conclude $L(A \times A)L \subseteq \{(x_1, \ldots, x_n) \mid x_n \leq r_E\}^2$ and hence $L(A \times A)L \cap P_n^2 \subseteq ([0, r_E]^n)^2 \in \mathcal{E}_X|_{P_n}$. \Box

Lemma 4.8 asdim $(P_n, \mathcal{E}_X|_{P_n}) \ge n$

Proof. Let \mathcal{U} be a uniformly bounded cover of P_n with appetite $\Delta_{P_n,1} := \{(x,y) \in P_n \times P_n \mid d_{eucl.}(x,y) \leq 1\}$. Note that by Proposition 4.2 the set $\Delta_{P_n,1}$ is an entourage. We will prove that the multiplicity of \mathcal{U} is at least n+1.

By $p_i: P_n \to R, (x_1, \ldots, x_n) \mapsto x_i$, we denote the projection of P_n onto the *i*-th factor. Set $E_i := p_i \times p_i(\Delta_{\mathcal{U}}) \in \mathcal{E}_R$ and $\Delta_1 := \{(x, y) \in R^2 \mid |x-y| \leq 1\}$. Choose r > 1 such that $E_n \Delta_1 E_{n-1} E_{n-2} \cdots E_2 E_1[0] \subseteq [0, r)$. Define

$$a_n := (0, \dots, 0, 1) \in P_n$$
 and $a_j := (\underbrace{0, \dots, 0}_{j}, r, \dots, r) \in P_n$.

The points a_0, \ldots, a_n are the vertices of the *n*-simplex

 $S := \{\lambda_0 a_0 + \dots + \lambda_n a_n \mid \lambda_0, \dots, \lambda_n \ge 0 \text{ and } \lambda_0 + \dots + \lambda_n = 1\}.$

By S_j we denote the (n-1)-face of S opposite to a_j . For $x = (x_1, \ldots, x_n)$ the following conclusions hold.

$$\begin{array}{ll} x \in S_0 & \Rightarrow & x_1 = 0 \\ x \in S_j & \Rightarrow & x_j = x_{j+1} \\ x \in S_{n-1} & \Rightarrow & 0 \le x_n - x_{n-1} \le 1 \\ x \in S_n & \Rightarrow & x_n = r \end{array}$$
 $(j = 1, \dots, n-2)$

We claim that for each $U \in \mathcal{U}$ there is a face S_i such that $U \cap S_i = \emptyset$. Suppose there is $U \in \mathcal{U}$ with $U \cap S_i \neq \emptyset$ for all $i \in \{0, \ldots, n\}$. Take an element $x^{(0)} \in U \cap S_0$. Then $x_1^{(0)} = 0$ and hence $p_1(U) \subseteq E_1[0]$. Now take $x^{(1)} \in U \cap S_1$. We have $x_2^{(1)} = x_1^{(1)}$ and therefore $p_2(U) \subseteq E_2E_1[0]$. Inductively, we get $p_j(U) \subseteq E_j \cdots E_1[0]$ for $j \in \{1, \ldots, n-1\}$. Finally, we take $x^{(n-1)} \in U \cap S_{n-1}$. This time we have $x_n^{(n-1)} \in \Delta_1(x_{n-1}^{(n-1)})$ and hence $p_n(U) \subseteq E_n \Delta_1 E_{n-1} \cdots E_1[0] \subseteq [0, r)$. On the other hand, since $U \cap S_n \neq \emptyset$, we get $r \in p_n(U)$. This is a contradiction.

Of course $\mathcal{U}_S := \{U \in \mathcal{U} \mid U \cap S \neq \emptyset\}$ is a cover of S. Take any map $g: \mathcal{U}_S \to \{a_0, \ldots a_n\}$, such that $U \in \mathcal{U}_S$ does not intersect the (n-1)-face opposite to g(U). Defining $U_i := \bigcup_{U \in g^{-1}(a_i)} U$ we obtain the cover $\{U_0, \ldots U_n\}$ of S with Lebesgue number at least 1 and such that U_i contains a_i and $U_i \cap S_i = \emptyset$.

We intend to apply Theorem 4.9. Therefore, take a triangulation K of S such that the diameter of any simplex of K is at most 1. Choose a map $h: \operatorname{vert}(K) \to \{a_0 \ldots, a_n\}$ such that for each $v \in h^{-1}(a_i)$ the star of v, i.e. the union of all simplices containing v, is contained in U_i . Applying Theorem 4.9 we get an n-simplex of K which is contained in $U_0 \cap \cdots \cap U_n$. Hence $U_0 \cap \cdots \cap U_n \neq \emptyset$. This yields $\mu(\mathcal{U}) \ge n+1$.

Theorem 4.9 (Sperner's Lemma) Let K be a triangulation of a closed n-simplex S. Suppose we are given a map $h: vert(K) \to vert(S)$ with the following property: If $v \in vert(K)$ lies in any face of S, then h(v) lies in the same face. Under these assumptions, there is at least one n-simplex T of the triangulation K whose n + 1 vertices correspond under h to the n + 1 different vertices of S.

A proof of Sperner's Lemma can be found in [Fed88]. Sperner's Lemma is a combinatorial version of the fact that the sphere S^{n-1} is not a retract of the closed ball of radius 1 around the origin in \mathbb{R}^n .

Lemma 4.10 We can estimate the asymptotic dimension of coarse spheres and coarse cells as follows. Let R be a ray and $n \in \mathbb{N}$. Then

asdim $D^{n-1}R \le n$ and asdim $S^{n-1}R \le n$.

Proof. Notice that Proposition 3.30 yields

asdim
$$D^{n-1}R$$
 = asdim R^n = asdim $S^{n-1}R$.

For $i \in \{1, \ldots, n\}$ consider the projection $p_i \colon \mathbb{R}^n \to \mathbb{R}, (x_1, \ldots, x_n) \mapsto x_i$ and remember that $\Delta_1 := \{(x, y) \in \mathbb{R} \times \mathbb{R} \mid |x - y| \leq 1\} \in \mathcal{E}_R$. Let $E \in \mathcal{E}_R^{*n}$ be a symmetric entourage. Set $E_i := p_i \times p_i(E) \cup \Delta_1 \in \mathcal{E}_R$ and $\widetilde{E} := (E_1 \cup \cdots \cup E_n)^{\square} \in \mathcal{E}_R$. This implies $E \subseteq \widetilde{E}^{\times n} \in \mathcal{E}_R^{*n}$. We define some bounded sets out of which we will build a cover of \mathbb{R}^n .

$$K_{0} := \{0\}$$

$$K_{i+1} := \widetilde{E}[K_{i}] \quad \text{for } i \in \mathbb{N}$$

$$K_{i} := \emptyset \quad \text{for } i < 0$$

$$\kappa_{i} := \sup K_{i} \quad \text{for } i \in \mathbb{N}$$

Observe that $[0, \kappa_i) \subseteq K_i$. Now we define a uniformly bounded cover \mathcal{U} of \mathbb{R}^n consisting of n + 1 families each of them being *E*-disjoint.

$$U_i := K_i \setminus K_{i-n}$$

$$\mathcal{U}_i := \{U_{i_1} \times \dots \times U_{i_n} \mid i_1, \dots, i_n \equiv i \mod n+1\}$$

$$\mathcal{U} := \mathcal{U}_0 \cup \dots \cup \mathcal{U}_n$$

We first prove that \mathcal{U} is indeed a cover of \mathbb{R}^n . Let $(x_1, \ldots, x_n) \in \mathbb{R}^n$. Choose $i_1, \ldots, i_n \in \mathbb{N}$ such that $(x_1, \ldots, x_n) \in K_{i_1} \setminus K_{i_1-1} \times \cdots \times K_{i_n} \setminus K_{i_n-1}$. There is $i \in \{1, \ldots, n+1\}$ such that for all $k \in \{1, \ldots, n\}$ we have $i_k \not\equiv i \mod n+1$. Define $j_k := \min\{j \in \mathbb{N} \mid j \geq i_k \text{ and } j - n \equiv i \mod n+1\}$. It follows that $j_k - n < i_k \leq j_k$ and hence $K_{i_k} \setminus K_{i_k-1} \subseteq \mathcal{U}_{j_k}$. This implies $(x_1, \ldots, x_n) \in \mathcal{U}_{j_1} \times \cdots \times \mathcal{U}_{j_n} \in \mathcal{U}_{i-1}$.

Since $\Delta_1 \subseteq \widetilde{E}$, the closure of \widetilde{E} (with respect to the usual topology on \mathbb{R}^2) is contained in \widetilde{E}^3 . This implies $(\kappa_i, \kappa_{i+1}) \in \widetilde{E}^3$ for $i \in \mathbb{N}$ and $(x, \kappa_i) \in \widetilde{E}^3$ for $i \in \mathbb{N}$ and $x \in K_i \setminus K_{i-1}$. Hence $\Delta_{\mathcal{U}} \subseteq \left(\widetilde{E}^{\times n}\right)^{3n+6}$, i.e. $\Delta_{\mathcal{U}}$ is uniformly bounded.

Observe that for $i \in \{0, ..., n\}$ the family \mathcal{U}_i is *E*-disjoint. This is a direct consequence of the definitions.

Corollary 4.11 If R is a ray, then asdim $D^n R = \operatorname{asdim} S^n R = n + 1$.

Proof. Apply Lemma 4.8 and Lemma 4.10.

 $\{(x,y) \in \mathbb{R}^2_+ \mid |x-y| \le 1\} \subseteq E_i \text{ for } i \in I_0.$

Lemma 4.12 Let I be a set. For $i \in I$ let R_i be a ray and $n_i \in \mathbb{N}$. Define $X := \coprod_{i \in I} D^{n_i} R_i$ and $A := \coprod_{i \in I} S^{n_i - 1} R_i$. Let (Y, \mathcal{E}_Y) be any coarse space and \mathcal{E}_X the coarse structure of X. Set $n := \sup\{n_i \mid i \in I\}$. If $f : (A, \mathcal{E}_X|_A) \to (Y, \mathcal{E}_Y)$ is coarsely uniform, then

asdim $(X \setminus A, \mathcal{E}_{X \cup_f Y}|_{X \setminus A}) \le \max\{n+1, \operatorname{asdim}(\operatorname{im} f, \mathcal{E}_Y|_{\operatorname{im} f})\}.$

Proof. Suppose $n < \infty$ and set $m := \max\{n+1, \operatorname{asdim}(\operatorname{im} f, \mathcal{E}_Y|_{\operatorname{im} f})\}$. For every entourage $\check{E} \in \mathcal{E}_{X \cup fY}|_X$ there is $E \in \mathcal{E}_X$ and $L \in \mathcal{E}_Y|_{\operatorname{im} f}$ such that $\check{E} \subseteq E \cup E\check{L}E$ where $\check{L} := (f \times f)^{-1}(L)$. Note that far away from A the entourages E and \check{E} coincide, i.e. $E \setminus E[A]^2 = \check{E} \setminus E[A]^2$. We may assume $\Delta_A \subseteq \check{L}$ and $\Delta_X \subseteq E$. Note that $D^{n_i}R_i$ consists of at most 2^n copies of $R_i^{n_i+1}$. Choose one of these copies for each $i \in I$ and denote their union by X'. Furthermore, set $A' := A \cap X'$. With Proposition 3.29 in mind it is enough to prove that $\operatorname{asdim}(X' \setminus A', \mathcal{E}_{X \cup fY}|_{X' \setminus A'}) \leq m$. Similarly as in Lemma 4.10 we can choose $E_i \in \mathcal{E}_{R_i}$ for $i \in I$ such that $E|_{R_i^{n_i+1}} \subseteq E_i^{\times (n_i+1)}$ and $E_i = E_i^{\boxtimes}$. Since X is a coproduct, Proposition 1.28 implies that we can choose E_i in such a way that there is a finite set $I_0 \subseteq I$ such that $E_i \subseteq \Delta_{R_i}$ for all $i \in I \setminus I_0$. Define

$$\begin{aligned}
K_{i,0} &:= \{0\} & \text{for } i \in I_0, \\
K_{i,k} &:= \emptyset & \text{for } k < 0, i \in I_0 \text{ and} \\
K_{i,k+1} &:= E_i^5[K_{i,k}] & \text{for } k \in \mathbb{N}, i \in I_0.
\end{aligned}$$

Furthermore, we define a uniformly bounded cover $\mathcal{U} = \mathcal{U}_0 \cup \cdots \cup \mathcal{U}_m$ of X' as follows: Let $k \in \mathbb{N}, l \in \{1, \ldots, m+1\}$ and $i \in I_0$.

$$U_{i,k} := E_i^2 [K_{i,k} \setminus K_{i,k-m}]$$

$$\mathcal{U}_l := \{U_{i,k_1} \times \dots \times U_{i,k_{n_i+1}} \mid i \in I_0, k_1, \dots, k_{n_i+1} \equiv l \mod m+1\}$$

So far we did not cover the parts of X' corresponding to $I \setminus I_0$. Define

$$\mathcal{U}_0 := \mathcal{U}_{m+1} \cup \{\{x\} \mid x \in R_i^{n_i+1}, i \in I \setminus I_0\}.$$

The proof that \mathcal{U} is a uniformly bounded cover of X' is similar to an argument in the proof of Lemma 4.10. Observe that \mathcal{U} has appetite $(E|_{X'})^2$ and that each of the families $\mathcal{U}_0, \ldots, \mathcal{U}_m$ is *E*-disjoint.

We want to construct a nice cover \mathcal{V} of A'. For $M \subseteq X \times X$ define $f_*M := f \times f(M \cap A \times A)$. Notice that $f_*E, f_*\Delta_{\mathcal{U}} \in \mathcal{E}_Y$. Take a uniformly bounded cover $\widetilde{\mathcal{V}} := \widetilde{\mathcal{V}}_0 \cup \cdots \cup \widetilde{\mathcal{V}}_m$ of $\inf f$ with appetite $f_*E \cup (f_*E)L(f_*E)$ and such that each family $\widetilde{\mathcal{V}}_k$ is $f_*\Delta_{\mathcal{U}}$ -disjoint. By taking the inverse image under $f|_{A'}$ we get a cover $\mathcal{V} = \mathcal{V}_0 \cup \cdots \cup \mathcal{V}_m$ of A' with appetite $(E\check{L}E)|_{A'} = (E \cup E\check{L}E)|_{A'}$ and such that each of the families $\mathcal{V}_0, \ldots, \mathcal{V}_m$ is $\Delta_{\mathcal{U}}$ -disjoint. In general \mathcal{V} will not be uniformly bounded with respect to $\mathcal{E}_X|_{A'}$.

Now we construct a cover of $E[A] \cap X' \setminus A'$ starting from the cover \mathcal{V} of A'. Let $k \in \{0, \ldots, m\}$. For $V \in \mathcal{V}_k$ define

$$V^{\flat} := \bigcup_{i \in I} (V \cap R_i^{n_i}) \times E_i[0] \quad \text{and} \quad V^{\sharp} := \left(V^{\flat} \cup \bigcup_{\substack{U \in \mathcal{U}_k \\ U \cap V^{\flat} \neq \emptyset}} U \right) \setminus A .$$

Observe that $\mathcal{V}_k^{\sharp} := \{ V^{\sharp} \mid V \in \mathcal{V}_k \}$ is uniformly bounded with respect to $\mathcal{E}_{X \cup_f Y}|_{X' \setminus A'}$. From \mathcal{U}_k and \mathcal{V}_k^{\sharp} we build the family

$$\mathcal{W}_k := \{ U \setminus A \mid U \in \mathcal{U}_k, U \cap V^{\flat} = \emptyset \text{ for all } V \in \mathcal{V}_k \} \cup \mathcal{V}_k^{\sharp}.$$

Note that any $U \in \mathcal{U}_k$ does intersect V^{\flat} for at most one $V \in \mathcal{V}_k$. We obtain a cover $\mathcal{W} := \mathcal{W}_0 \cup \cdots \cup \mathcal{W}_m$ of $X' \setminus A'$ which is uniformly bounded with respect to $\mathcal{E}_{X \cup_f Y}|_{X' \setminus A'}$ and has appetite $\check{E}|_{X' \setminus A'}$.

Proof of Theorem 4.6. Let Y_k , I_k , $R_{k,i}$, $n_{k,i}$ and $f_{k,i}$ be as in Definition 4.5. By induction we will prove

asdim
$$Y_k = \sup\{n_{j,i} \mid j \le k, i \in I_j\} + 1.$$
 (4.1)

The case k = 0 follows from Proposition 3.30 and Corollary 4.11.

Set $X := \prod_{i \in I_k} D^{n_{k,i}} R_{k,i}$ and $A := \prod_{i \in I_k} S^{n_{k,i}-1} R_{k,i}$. By Corollary 4.11 the asymptotic dimension of X is $s_k + 1$ where $s_k := \sup\{n_{k,i} \mid i \in I_k\}$. We need information on the asymptotic dimension of $(X \setminus A, \mathcal{E}_{X \cup f_k} Y_{k-1}|_{X \setminus A})$. Observe that

$$X \setminus A = \prod_{i \in I_k} D^{n_{k,i}} R_{k,i} \setminus S^{n_{k,i}-1} R_{k,i} .$$

Using Lemma 4.7, Lemma 4.8 and monotony of asymptotic dimension, we get asdim $(X \setminus A, \mathcal{E}_{X \cup_{f_k} Y_{k-1}} |_{X \setminus A}) \geq s_k + 1$. Applying Lemma 4.12 we get asdim $(X \setminus A, \mathcal{E}_{X \cup_{f_k} Y_{k-1}} |_{X \setminus A}) \leq \max\{s_k + 1, \operatorname{asdim} Y_{k-1}\}$. Lemma 1.36 assures us that there is no ambiguity about the coarse structure of Y_{k-1} . Notice that $Y_k = X \cup_{f_k} Y_{k-1} = X \setminus A \cup Y_{k-1}$. Hence, Proposition 3.29 completes the proof of (4.1).

Since all the maps $Y_k \to Y$ are injective, the claim of Theorem 4.6 now follows from Proposition 3.32.

We conclude this chapter by discussing some possible modifications in the definition of a coarse cell complex and the effect on the asymptotic dimension.

Remark 4.13 We might consider building a coarse cell out of different rays, e.g. $R_{1 \amalg} R_1 \times \cdots \times R_{n \amalg} R_n \times R_{n+1}$.

Using Proposition 4.3, the arguments given in this chapter can be easily adjusted to prove asdim $Y = \delta(Y) + 1$ for a coarse cell complex Y built with modified cells.

Remark 4.14 If we want to use the disjoint union \bigsqcup instead of the coproduct \coprod in Definition 4.5, we have to demand the maps f_k to be coarsely uniform, since this is no longer automatic.

If for all $k \in \mathbb{N}$ the map f_k is coarsely uniform, we can prove the formula of Theorem 4.6 in the same way as before. In the proof of Lemma 4.12 we do not have to distinguish two subsets of I. We may define $I_0 := I$ in this case. We do not need the fact that I_0 is a finite set if we work with disjoint unions instead of coproducts.

Metric coarse cell complexes

If (X, d_X) and (Y, d_Y) are pseudometric spaces, $A \subseteq X$ and $f: A \to Y$ is a (not necessarily continuous) map, then $X \cup_f Y$ is defined as a pseudometric space. We will use $\mathbb{S}^n := \mathbb{R}^{n+1} \times \{0\}$ and $\mathbb{D}^n := \mathbb{R}^n \times \mathbb{R}_+$ with the restrictions of the euclidean metric as coarse *n*-sphere and coarse *n*-cell respectively. **Definition 4.15** Let I be any set and X_i and Y_i pseudometric spaces. A family of maps $\{f_i \colon X_i \to Y_i\}_{i \in I}$ is said to be *coarsely uniform in a uniform way* if there is a (not necessarily continuous) map $s \colon \mathbb{R}_+ \to \mathbb{R}_+$ such that

$$\begin{array}{ccc} \forall & \forall & \forall \\ r > 0 & i \in I & x, \widetilde{x} \in X_i \end{array} \quad d(x, \widetilde{x}) \leq r \ \Rightarrow \ d(f_i(x), f_i(\widetilde{x})) \leq s(r). \end{array}$$

Definition 4.16 Suppose that for all $k \in \mathbb{N}$ there is a pseudometric space (Y_k, d_k) and a set I_k . Moreover, suppose that for all $k \in \mathbb{N}$ and $i \in I_k$ there is a number $n_{k,i} \in \mathbb{N}$ and a map $f_{k,i} \colon \mathbb{S}^{n_{k,i}-1} \to Y_{k-1}$ such that for each $k \in \mathbb{N}$ the family $\{f_{k,i}\}_{i \in I_k}$ is coarsely uniform in a uniform way. Define $f_k := \bigsqcup_{i \in I_k} f_{k,i} : \bigsqcup_{i \in I_k} \mathbb{S}^{n_{k,i}-1} \to Y_{k-1}$ and observe that f_k is coarsely uniform. If

$$Y_0 = \bigsqcup_{i \in I_0} \mathbb{D}^{n_{0,i}}$$
 and $Y_k = \left(\bigsqcup_{i \in I_k} \mathbb{D}^{n_{k,i}}\right) \cup_{f_k} Y_{k-1}$,

then $(Y, d) := \lim_{k \to \infty} (Y_k, d_k)$ is a metric coarse cell complex.

There are two coarse structures on Y. We write \mathcal{E}_d for the bounded coarse structure of Y induced by the pseudometric d. Similarly, we write \mathcal{E}_{d_k} for the bounded coarse structure of Y_k induced by the pseudometric d_k . Denote the coarse structure of $\lim_{k \to \infty} (Y_k, \mathcal{E}_{d_k})$ by \mathcal{E} .

Using Proposition 1.27, we see that $\mathcal{E} \subseteq \mathcal{E}_d$. The following example yields that in general \mathcal{E} and \mathcal{E}_d do not coincide.

Example 4.17 Consider any metric coarse cell complex (Y, d) with $I_k \neq \emptyset$ for all $k \in \mathbb{N}$. Define $\Delta_1 := \{(y_1, y_2) \in Y \times Y \mid |y_1 - y_2| \leq 1\}$ and observe that $\Delta_1 \in \mathcal{E}_d$. Proposition 1.27 implies $\Delta_1 \notin \mathcal{E}$.

Remark 4.18 The arguments given in this chapter can be easily adjusted to prove

asdim $Y_k = \sup\{n_{j,i} \mid j \le k, i \in I_j\} + 1$

for metric coarse cell complexes. Here we use the notations of Definition 4.16.

Proposition 4.19 asdim $(Y, \mathcal{E}) = \delta(Y) + 1$.

Proof. The claim follows from Remark 4.18 and Proposition 3.32. \Box

Proposition 4.20 If the metric coarse cell complex Y can be built in finitely many steps, i.e. if there is $k_0 \in \mathbb{N}$ such that $I_k = \emptyset$ for all $k \geq k_0$, then $\operatorname{asdim}(Y, \mathcal{E}_d) = \delta(Y) + 1$.

Proposition 4.20 follows directly from Remark 4.18.

The following example of a metric coarse cell complex yields that the formula of Theorem 4.6 is not true in general.

Example 4.21 We inductively define a metric coarse cell complex (Y, d)and start with $Y_0 := \mathbb{R}^2 \times \mathbb{R}_+ = \mathbb{D}^2 \subseteq \mathbb{R}^3$. For each $k \in \mathbb{N} \setminus \{0\}$ and $i \in I := \{0\} \times \mathbb{Z} \times \mathbb{N} \subseteq \mathbb{R}^3$ we attach a coarse 1-cell \mathbb{D}^1 to Y_0 using the map

$$\widetilde{f}_{k,i}$$
: $\mathbb{S}^0 \to Y_0, \ (x,0) \mapsto (2^k x, 0, 0) + i$.

Denote the metric space obtained after attaching a cell for all $k \in \{1, \ldots, n\}$ and $i \in I$ as just explained by Y_n . Observe that the inclusion $i_k \colon Y_0 \hookrightarrow Y_k$ is not isometric. On the other hand, all the attaching maps $f_{k,i} := i_{k-1} \circ \tilde{f}_{k,i}$ are 2-Lipschitz.¹ Hence $Y := \lim Y_k$ is a metric coarse cell complex.

The metric coarse cell complex Y is coarsely equivalent to the consecutively defined metric coarse cell complex Y': We start with $Y'_0 := \mathbb{R} \times \mathbb{R}_+ = \mathbb{D}^1$ and for each $k \in \mathbb{N} \setminus \{0\}$ and $y \in \mathbb{Z} \times \mathbb{N} \subset Y'$ we attach a coarse cell \mathbb{D}^1 using the constant attaching map $f'_{k,y} : \mathbb{S}^0 \to Y'_0$ whose image is just $\{y\}$. Finally, we obtain $\operatorname{asdim}(Y, \mathcal{E}_d) = \operatorname{asdim}(Y') = 2 < 2 + 1 = \delta(Y) + 1$.

Note that in the previous example the inclusion $Y_k \hookrightarrow Y$ is not a coarse embedding.

Remark 4.22 Let Y be a metric coarse cell complex such that all attaching maps $f_{k,i}$ are contracting. In this case (Y_k, d_k) is a subspace of (Y, d), i.e. the inclusion map $Y_k \hookrightarrow Y$ is isometric. Hence $\operatorname{asdim}(Y, \mathcal{E}_d) \ge \delta(Y) + 1$.

Proof. Compare Proposition 3.5.

Question 4.23 Does asdim $Y \leq \delta(Y) + 1$ hold for any metric coarse cell complex Y? Does the inequality hold if all attaching maps are contracting?

¹Let X and Y be pseudometric spaces. A map $f: X \to Y$ is called λ -Lipschitz if $d(f(x), f(x')) \leq \lambda \cdot d(x, x')$ for all $x, x' \in X$. The map f is called *contracting* if it is 1-Lipschitz.

Chapter 5

Asymptotic dimension and the covering dimension of the Higson corona

We will need dimension theory of topological spaces in this section, in particular some knowledge about the covering dimension dim. Compare [HW41], [Eng78] and [Fed88].

We refer to the beginning of Section 1.4 for the definitions of the Higson compactification and the Higson corona.

For a proper metric space X, we can recover the bounded coarse structure from its Higson compactification hX, since the continuously controlled coarse structure with respect to hX and the bounded coarse structure coincide. (This is Proposition 2.47 of [Roe03].)

Dranishnikov, Keesling and Uspenskij proved in [DKU98] and [Dra00] the following theorem.

Theorem 5.1 Let (X, d) be a metric space of finite asymptotic dimension and νX its Higson corona. Then $\operatorname{asdim}(X, d) = \dim(\nu X)$.

The following examples show that Theorem 5.1 cannot be generalized to arbitrary coarse spaces.

Example 5.2 Let X be a non-compact, normal Hausdorff space which can be written as a union of countably many compact sets. The Stone-Čech compactification βX induces on X the trivial coarse structure \mathcal{T} , and the Higson compactification of (X, \mathcal{T}) is homeomorphic to βX .

We get the following results about dimensions: On the one hand we know that $\operatorname{asdim}(X, \mathcal{T}) = 0$. On the other hand $\dim(\beta X \setminus X) = 0$ if and only if $\beta X \setminus X$ is totally disconnected. If X has the property that for every compact

subset $K \subseteq X$ the space $X \setminus K$ is connected, then $\beta X \setminus X$ is also connected. Hence, for such X we have $\dim(\beta X \setminus X) \neq 0$.

Proof. Obviously \mathcal{T} is contained in the coarse structure induced by any compactification. We write $\mathcal{E}_{\beta X}$ for the coarse structure on X induced by βX . It remains to prove that $\mathcal{E}_{\beta X} \subseteq \mathcal{T}$. There are compact sets $K_0 \subset K_1 \subset K_2 \subset \cdots \subset X$ such that $X = \bigcup_{i \in \mathbb{N}} K_i$. Let $E \in \mathcal{E}_{\beta X}$. Assume that for all $i \in \mathbb{N}$ there is $(x_i, y_i) \in E \setminus (\Delta_X \cup K_i \times K_i)$. This implies $x_i \neq y_i$. We may assume that $\{x_i \mid i \in \mathbb{N}\} \cap \{y_i \mid i \in \mathbb{N}\} = \emptyset$, since we can choose subsequences with this property. Observe that $\{x_i \mid i \in \mathbb{N}\} \subseteq X$ and $\{y_i \mid i \in \mathbb{N}\} \subseteq X$ are closed sets. Let $(x, y) \in (\beta X \setminus X)^2$ be an element in the closure of the sequence $\{(x_i, y_i)\}_{i \in \mathbb{N}}$. Since the closure of E in $\beta X \times \beta X$ is contained in $X \times X \cup \Delta_{\beta X}$, we conclude x = y. Using Urysohn's Lemma, we get a continuous function $f \colon X \to [0, 1]$ with $f(x_i) = 1$ and $f(y_i) = 0$ for all $i \in \mathbb{N}$. We may as well think of f as a continuous function with domain βX . Since inverse images of closed sets are closed, we get f(x) = 1 and f(y) = 0. This is a contradiction to x = y. Hence $E \subseteq \Delta_X \cup K \times K$ for a compact set $K \subseteq X$, i.e. $E \in \mathcal{T}$.

Every bounded, continuous function on X is a Higson function with respect to \mathcal{T} . Hence, the Higson compactification of (X, \mathcal{T}) is βX .

Compare Example 3.11 and [Fed88], Chapter 1, Section 3.2, Theorem 5 for the statements on dimensions.

Assume that $\beta X \setminus X$ is the disjoint union of two open and non-empty sets A and B. Then the characteristic function χ_A is a continuous projection in $C(\beta X \setminus X) \cong C_b(X)/C_0(X)$.¹ Choose a function $f \in C_b(X)$ whose image under $\pi \colon C_b(X) \to C_b(X)/C_0(X)$ is χ_A . Observe that $g := \frac{1}{2}(f + f^*)$ is a self-adjoint preimage of χ_A with respect to π . In particular g is real-valued. Consider g as an element of $C(\beta X)$. There is a compact set $K \subseteq X$ such that $g(x) \neq \frac{1}{2}$ for all $x \in X \setminus K$. Hence

$$X \setminus K = \left\{ x \in X \setminus K \mid g(x) < \frac{1}{2} \right\} \cup \left\{ x \in X \setminus K \mid g(x) > \frac{1}{2} \right\}$$

is not connected.

Example 5.3 Let X be a proper metric space and \mathcal{E} . the continuously controlled coarse structure induced by the one-point compactification of X. In this case νX is just a single point and $\operatorname{asdim}(X, \mathcal{E}) = 1 = \dim(\nu X) + 1$.

Proof. Compare Example 9.7 of [Roe03] for the calculation of asymptotic dimension. If $X = \mathbb{R}^n$, we can apply Example 3.27.

¹Let M be a locally compact Hausdorff space. We denote the space of continuous functions on M with values in \mathbb{C} by C(M). The space of continuous, bounded functions on M with values in \mathbb{C} is called $C_b(M)$ and by $C_0(M) \subseteq C_b(M)$ we denote the subspace of those functions vanishing at infinity.

Example 5.4 Consider \mathbb{R}^n with the visual corona S^{n-1} and denote by \mathcal{E}_{vis} the coarse structure coming from the corresponding compactification. Then $\operatorname{asdim}(\mathbb{R}^n, \mathcal{E}_{vis}) = n = \dim(S^{n-1}) + 1.$

Proof. Example 1.47, the calculation of the asymptotic dimension of coarse cells and Proposition 3.29 yield the desired formula for the asymptotic dimension. \Box

Let hX be a metrisable compactification of X and denote the corresponding continuously controlled coarse structure by \mathcal{E}_{hX} . The last two examples suggest a relation between the asymptotic dimension of (X, \mathcal{E}_{hX}) and the covering dimension of the corona $\nu X := hX \setminus X$. The goal of this chapter is to prove the following theorem.

Theorem 5.5

$$\operatorname{asdim}(X, \mathcal{E}_{hX}) = \dim(\nu X) + 1$$

The proof of Theorem 5.5 is given by the next two lemmas.

а

Lemma 5.6

$$\dim(\nu X) + 1 \leq \operatorname{asdim}(X, \mathcal{E}_{hX})$$

Proof. Consider the map $f: \nu X \times \mathbb{R}_+ \to \nu X \times \mathbb{N}$, $(x, t) \mapsto (x, [t])$ and denote by \mathcal{E}' the pull-back of the coarse structure \mathcal{E} described in Lemma 1.41. Observe that f is a coarse equivalence. The coarse structure \mathcal{E}' has a description which is very similar to the description of \mathcal{E} in Lemma 1.41. Theorem 1.43 now implies that (X, \mathcal{E}_{hX}) and $(\nu X \times \mathbb{R}_+, \mathcal{E}')$ are coarsely equivalent.

Let $\operatorname{asdim}(X, \mathcal{E}_{hX}) = n$. We have a uniformly bounded cover of $\nu X \times \mathbb{R}_+$ with appetite

$$E := \left\{ \left((x,t), (y,s) \right) \in (\nu X \times \mathbb{R}_+)^2 \mid d(x,y) < \frac{1}{\max\{t,s\}}, \ |t-s| < 1 \right\}$$

and multiplicity $\leq n+1$. Since E((x,t)) is open for all $(x,t) \in \nu X \times \mathbb{R}_+$, this cover has an open refinement which also has appetite E and multiplicity at most n+1. Denote this cover by \mathcal{U} .

Let $\varepsilon > 0$. From \mathcal{U} we will construct an open cover $\mathcal{U}_{\varepsilon}$ of $\nu X \times [0,1]$ with multiplicity $\leq n + 1$ and mesh $(\mathcal{U}_{\varepsilon}) \leq \varepsilon$. This implies $\dim(\nu X \times [0,1]) \leq n$. We start constructing $\mathcal{U}_{\varepsilon}$. Observe that $L := \pi_{\mathbb{R}_+} \times \pi_{\mathbb{R}_+}(\Delta_{\mathcal{U}}) \subseteq \mathbb{R}^2_+$ is symmetric and contains $\Delta_1 := \{(t,s) \in \mathbb{R}^2_+ \mid |t-s| < 1\}$. Choose $t_{\varepsilon} > 0$ such that $d(x,y) \leq \frac{\varepsilon}{2}$ for all $((x,t),(y,s)) \in \Delta_{\mathcal{U}}$ with max $\{s,t\} \geq t_{\varepsilon}$. Consider the continuous map $f_{\varepsilon} : \mathbb{R}_+ \to \mathbb{R}_+$ with $f_{\varepsilon}(0) = t_{\varepsilon}$ and

$$f_{\varepsilon}\left(\frac{n\cdot\varepsilon}{4}\right) = \sup L^n(t_{\varepsilon}) \quad \text{for} \quad n = \{1, 2, \ldots\}$$

which is affine on the interval $\left[\frac{n \cdot \varepsilon}{4}, \frac{(n+1) \cdot \varepsilon}{4}\right]$ for each $n \in \mathbb{N}$. Observe that f_{ε} is strictly monotone increasing. Define $\mathcal{U}_{\varepsilon}$ to be the pull-back of \mathcal{U} under

the continuous map $\mathrm{id}_{\nu X} \times f_{\varepsilon}$. It is clear that $\mathcal{U}_{\varepsilon}$ has the same multiplicity as the cover \mathcal{U} .

Take $V \in \mathcal{U}$ and set $U := (\mathrm{id}_{\nu X} \times f_{\varepsilon})^{-1}(V)$. It remains to check that the diameter of U is at most ε . Let $(x,t), (y,s) \in U$. The definition of t_{ε} implies $d(x,y) \leq \frac{\varepsilon}{2}$. Suppose $s \leq t$. Notice that $f_{\varepsilon}(t) \in L(f_{\varepsilon}(s))$. This implies $|t-s| \leq \frac{\varepsilon}{2}$. Together, we get $d((x,t), (y,s)) \leq \varepsilon$. This completes the construction of the cover $\mathcal{U}_{\varepsilon}$.

Using some results about the covering dimension of products (compare [Fed88], Chapter 2, Section 6.3, Theorem 18 and Chapter 5, Section 5.3, Corollary 5), we get $\dim(\nu X) + 1 = \dim(\nu X \times [0,1]) \leq n$.

Lemma 5.7

$$\operatorname{asdim}(X, \mathcal{E}_{hX}) \le \operatorname{dim}(\nu X) + 1$$

Proof. Assume dim $(\nu X) + 1 = n$. We will prove asdim $(\nu X \times \mathbb{N}, \mathcal{E}) \leq n$ where \mathcal{E} is the coarse structure described in Lemma 1.41.

Let $E \in \mathcal{E}$ be a symmetric entourage containing the diagonal $\Delta_{\nu X \times \mathbb{N}}$ and $\{\delta_k\}_{k \in \mathbb{N}}$ a sequence of non-negative real numbers converging to zero such that $d(x, x') < \delta_k$ for $((x, m), (x', m')) \in E$ with $\max\{m, m'\} \ge k$. We may assume this sequence to be monotone.

For each $k \in \mathbb{N}_+$ there is a finite cover of νX with open balls of radius $\frac{1}{2k}$ and multiplicity at most n. Ostrand's theorem² provides a finite open cover \mathcal{V}_k of νX with mesh $(\mathcal{V}_k) \leq \frac{1}{k}$ consisting of n disjoint families $\mathcal{V}_{k,1}, \ldots, \mathcal{V}_{k,n}$. For $V \in \mathcal{V}_{k,j}$ define $j_V := j$.

Since νX is compact, \mathcal{V}_k has positive Lebesgue number $L(\mathcal{V}_k) > 0$. We define inductively a sequence $\{l_i\}_{i \in \mathbb{N}}$. Set $l_0 := 1$ and for i > 0 take $l_i \in \mathbb{N}$ such that $\frac{1}{l_i} \leq L(\mathcal{V}_{l_{i-1}})$ and $l_i > l_{i-1}$. We choose maps $\phi_i \colon \mathcal{V}_{l_i} \to \mathcal{V}_{l_{i-1}}$ such that $W \subseteq \phi_i(W)$ for all $W \in \mathcal{V}_{l_i}$. For $V \in \mathcal{V}_{l_{i-1}}$ we define $\breve{V} := \bigcup_{W \in \phi_i^{-1}(V)} W$. Observe that $\breve{V} \subset V$.

Consider the projection $\pi_{\mathbb{N}} : \nu X \times \mathbb{N} \to \mathbb{N}$ and define $E_{\mathbb{N}} := \pi_{\mathbb{N}} \times \pi_{\mathbb{N}}(E) \cup \Delta_1$ where $\Delta_1 := \{(i, j) \in \mathbb{N}^2 \mid |i - j| \leq 1\}$. Observe that $E_{\mathbb{N}}$ is an entourage of the continuously controlled coarse structure induced by the one-point compactification of \mathbb{N} . Set $K_0 := \{0\}$ and $K_k := E_{\mathbb{N}}[K_{k-1}]$ and notice that $K_0 \subset K_1 \subset \cdots \subset \mathbb{N}$ and $\bigcup_{k \in \mathbb{N}} K_k = \mathbb{N}$.

We define a sequence $\{k_i\}_{i\in\mathbb{N}}$. Set $k_{-2} := 0$. For all $i \in \mathbb{Z}$ with $i \geq -1$ choose $k_i \in \mathbb{N}$ such that $k_i > k_{i-1} + 2n$ and $\delta_m < \frac{1}{l_{i+1}}$ for all $m \notin K_{k_i-2}$. We need some last preparations. For $i \in \mathbb{N} \setminus \{0\}$ and $j \in \{1, \ldots, n\}$ we set $A_{i,j} := K_{k_i} \setminus K_{k_{i-1}+2j-2}$ and $B_{i,j} := K_{k_i+2j} \setminus K_{k_i}$.

²We use the following version of Ostrand's theorem: A normal space X satisfies dim $X \leq n$ if and only if for every locally finite open cover \mathcal{U} there exists an open cover $\mathcal{V} = \mathcal{V}_1 \cup \cdots \cup \mathcal{V}_{n+1}$ where each of the families \mathcal{V}_i consists of disjoint sets and for each $V \in \mathcal{V}$ there exists $U \in \mathcal{U}$ such that $V \subseteq U$. Compare [Eng78].

Finally, we define a cover \mathcal{U} of $\nu X \times \mathbb{N}$.

$$U_{0} := \bigcup_{V \in \mathcal{V}_{l_{0}}} V \times K_{k_{0}+2j_{V}}$$
$$\mathcal{U}_{i} := \left\{ V \times A_{i,j_{\phi_{i}(V)}} \cup \breve{V} \times B_{i,j_{V}} \mid V \in \mathcal{V}_{l_{i}} \right\}$$
$$\mathcal{U} := \left\{ U_{0} \right\} \cup \bigcup_{i \in \mathbb{N} \setminus \{0\}} \mathcal{U}_{i}$$

To check that \mathcal{U} is uniformly bounded, let $((x, m_x), (y, m_y)) \in \Delta_{\mathcal{U}}$. Suppose that $m_x, m_y \notin K_{k-1}$. There are natural numbers k_x, k_y, i_x and i_y such that the following is true.

$$m_x \in K_{k_x} \setminus K_{k_x-1} \subseteq K_{k_{i_x}} \setminus K_{k_{i_x-1}}$$

$$m_y \in K_{k_y} \setminus K_{k_y-1} \subseteq K_{k_{i_y}} \setminus K_{k_{i_y-1}}$$

We have $|i_x - i_y| \leq 1$. Hence $\pi_{\mathbb{N}} \times \pi_{\mathbb{N}}(\Delta_{\mathcal{U}}) \in \mathcal{E}_{\mathbb{N}}$ where $\mathcal{E}_{\mathbb{N}}$ is the continuously controlled coarse structure on \mathbb{N} induced by the one-point compactification. Without loss of generality, assume that $m_x \leq m_y$. Set $d_m := 1$ if $m \in K_{k_1}$. Set $d_m := \frac{1}{l_{i-2}}$ if $m \in K_{k_i} \setminus K_{k_{i-1}}$ and i > 1. Observe that $\lim_{m \to \infty} d_m = 0$ and $d(x, y) \leq \frac{1}{l_{i_y-2}} = d_{m_y}$.

Now we prove that \mathcal{U} has multiplicity at most n+1. Let $(x,m) \in \nu X \times \mathbb{N}$. If $m \in K_{k_0}$, then $(x,m) \in U_0$ and (x,m) is not contained in any other covering set. Suppose that $m \in K_{k_i} \setminus K_{k_{i-1}}$ with $i \in \mathbb{N} \setminus \{0\}$. If $(x,m) \in U$ for some $U \in \mathcal{U}$, then $U \in \mathcal{U}_i \cup \mathcal{U}_{i-1}$. Observe that x is contained in at most one set of each of the collections $\mathcal{V}_{l_i,1}, \ldots, \mathcal{V}_{l_i,n}, \mathcal{V}_{l_{i-1},1}, \ldots, \mathcal{V}_{l_{i-1},n}$. Moreover, $m \notin A_{l,j}$ if $l \neq i$, and $m \notin B_{l,j}$ if $l \neq i-1$. Hence (x,m) is contained in at most 2n sets of the cover \mathcal{U} . There is $l \in \mathbb{N} \setminus \{0\}$ such that $m \in K_{k_{i-1}+2l} \setminus K_{k_{i-1}+2l-2}$. If $l \geq n$, then $m \notin B_{i-1,j}$ for $j \in \{1, \ldots, n-1\}$. If 1 < l < n, then $m \notin B_{i-1,j}$ for $j \in \{1, \ldots, n-1\}$. If l = 1, then $m \notin A_{i,j}$ for $j \in \{2, \ldots, n\}$. Hence, there are at most n+1 sets of the cover \mathcal{U} containing (x,m).

Finally we prove that \mathcal{U} has appetite E. Let $(x,m) \in \nu X \times \mathbb{N}$. Set $K_k := \emptyset$ for k < 0. There is a unique $k \in \mathbb{N}$ such that $m \in K_k \setminus K_{k-1}$. It follows that $\pi_{\mathbb{N}}(E((x,m))) \subseteq K_{k+1} \setminus K_{k-2}$. There is a unique $i \in \{-1\} \cup \mathbb{N}$ with $k_{i-1} \leq k < k_i$. We conclude that $\pi_{\mathbb{N}}(E((x,m))) \subseteq K_{k_i} \setminus K_{k_{i-1}-2}$. If $i \leq 0$, then $E((x,m)) \subseteq \nu X \times K_{k_0} \subseteq U_0$.

Suppose i > 0. Observe that $\pi_{\nu X}(E((x,m)))$ is contained in the ball $B_{\delta_m}(x)$ of radius δ_m around x. There is $W \in \mathcal{V}_{l_i}$ such that $B_{\delta_m}(x) \subseteq W$. Set $V := \phi_i(W)$ and observe that $W \subseteq V \in \mathcal{V}_{l_{i-1}}$.

It remains to prove that $W \times K_{k+1} \setminus K_{k-2}$ is contained in some $U \in \mathcal{U}$. If $k \ge k_{i-1} + 2j_V$, then

$$W \times K_{k+1} \setminus K_{k-2} \subseteq W \times K_{k_i} \setminus K_{k_{i-1}+2j_V-2}$$

= $W \times A_{i,j_V} \subseteq W \times A_{i,j_V} \cup \breve{W} \times B_{i,j_W} \in \mathcal{U}_i$.

If $k < k_{i-1} + 2j_V$ and i > 1, then

$$W \times K_{k+1} \setminus K_{k-2} \subseteq W \times K_{k_{i-1}+2j_V} \setminus K_{k_{i-1}-2}$$
$$\subseteq V \times A_{i-1,j_{\phi_{i-1}(V)}} \cup \breve{V} \times B_{i-1,j_V} \in \mathcal{U}_{i-1} .$$

If $k < k_{i-1} + 2j_V$ and i = 1, then

$$W imes K_{k+1} \setminus K_{k-2} \subseteq V imes K_{k_0+2j_V} \subseteq U_0 \in \mathcal{U}$$

This completes the proof that the cover \mathcal{U} has appetite E.

Remark 5.8 For any coarse spaces X and Y whose coarse structures are induced by metrisable compactifications, we get another proof of

$$\operatorname{asdim}(X \times Y) \leq \operatorname{asdim}(X) + \operatorname{asdim}(Y).$$

Proof. By Corollary 1.45 the coarse space X is coarsely equivalent to $\psi(\nu X)$ and the same for Y. This and Proposition 1.46 imply that $X \times Y$ is coarsely equivalent to $\psi(\nu X \boxtimes \nu Y)$. Hence, the Higson corona of $X \times Y$ is homeomorphic to $\nu X \boxtimes \nu Y$. Using Theorem 5.5 and the product theorem of dimension theory (i.e. [Fed88], Chapter 2, Section 6.3, Theorem 18), we get $\operatorname{asdim}(X \times Y) = \dim(\nu X \boxtimes \nu Y) + 1 = \dim(\nu X \times \nu Y \times [0, 1]) + 1 \leq \dim(\nu X) + \dim(\nu Y) + 2 = \operatorname{asdim}(X) + \operatorname{asdim}(Y).$

Corollary 5.9 There are coarse spaces X and Y such that

 $\operatorname{asdim}(X \times Y) < \operatorname{asdim}(X) + \operatorname{asdim}(Y).$

Proof. There are metric compacta K and L such that $\dim(K \times L) < \dim(K) + \dim(L)$. Compare the section about Pontryagin's compacta, i.e Capter 5, Section 4.2 of [Fed88]. The same reasoning as in the proof of Remark 5.8 translates this to asymptotic dimensions.

Remark 5.10 In [BL] the authors construct metric spaces X and Y such that $\operatorname{asdim}(X \times Y) < \operatorname{asdim}(X) + \operatorname{asdim}(Y)$.

Chapter 6

Asymptotic dimension of $CAT(\kappa)$ -spaces for $\kappa < 0$

Let $\kappa < 0$. We write M_{κ}^{n} for the *n*-dimensional hyperbolic space with constant sectional curvature κ . We need some basic facts about $CAT(\kappa)$ -spaces and some knowledge in hyperbolic geometry. A good introduction on these subjects can be found in [BH99].

We want to calculate the asymptotic dimension of $CAT(\kappa)$ -spaces. As the following example shows, we should not expect that the asymptotic dimension and the covering dimension of a $CAT(\kappa)$ -space always coincide.

Example 6.1 Consider an *n*-simplex σ in M_{κ}^n . Take a *k*-dimensional face f of σ . There is an isometric embedding $i: M_{\kappa}^k \hookrightarrow M_{\kappa}^n$ such that $f \subseteq \operatorname{im} i$. Now $\sigma \cup \operatorname{im} i$ is a CAT(0)-space with covering dimension n and asymptotic dimension k.

Hence, for every pair of natural numbers (k, n) with $k \leq n$, there is a $CAT(\kappa)$ -space with asymptotic dimension k and covering dimension n.

6.1 Spaces with nicely covered spheres

Let (X, d) be a metric space and $x_0 \in X$ a basepoint. For $r \geq 0$ we define $S_r(x_0) = \{x \in X \mid d(x, x_0) = r\}$ and $D_r(x_0) = \{x \in X \mid d(x, x_0) < r\}$. We consider $S_r(x_0) \subseteq X$ and $D_r(x_0) \subseteq X$ with the metric obtained by restricting the metric d of X.

Definition 6.2 We say that X has nicely n-colored spheres (with respect to the basepoint $x_0 \in X$) if there is $\rho > 0$ and a cover \mathcal{U}_k of $S_{k \cdot \rho}(x_0)$ for all $k \in \mathbb{N}$ such that

- there is a lower bound $\lambda > 0$ on the Lebesgue numbers of \mathcal{U}_k for $k \in \mathbb{N}$,
- there is an upper bound on the mesh of the covers \mathcal{U}_k and

• $\mathcal{U}_k = \mathcal{U}_{k,1} \cup \cdots \cup \mathcal{U}_{k,n}$ with multiplicity $\mu(\mathcal{U}_{k,i}) \leq 1$ for $i \in \{1, \ldots, n\}$.

Definition 6.3 We say that X has nicely n-covered spheres (with respect to the basepoint $x_0 \in X$) if there is $\rho > 0$ and a cover \mathcal{U}_k of $S_{k \cdot \rho}(x_0)$ for all $k \in \mathbb{N}$ such that

- there is a lower bound $\lambda > 0$ on the Lebesgue numbers of \mathcal{U}_k for $k \in \mathbb{N}$,
- there is an upper bound on the mesh of the covers \mathcal{U}_k and
- \mathcal{U}_k has multiplicity at most n.

Lemma 6.4 A metric space X has nicely n-colored spheres if and only if it has nicely n-covered spheres.

Proof. Compare Remark 3.17.

Theorem 6.5 Let $\kappa < 0$. For a CAT(κ)-space X with nicely n-covered spheres, we have $\operatorname{asdim}(X) \leq n$.

Remark 6.6 In some cases the upper bound on the asymptotic dimension of X given by Theorem 6.5 is lower than the covering dimension of X. Compare Example 6.1 and observe that in this case the upper bound provided by Theorem 6.5 is sharp.

6.2 Proof of Theorem 6.5

Because of Lemma 6.4 we may assume that X has nicely n-colored spheres. We will prove Theorem 6.5 by constructing uniformly bounded covers $\mathcal{M}_{\rho,N}$ of X with multiplicity at most n+1. In a second step we prove that for every L > 0 we can adjust the parameters ρ and N so that $\mathcal{M}_{\rho,N}$ has Lebesgue number at least L.

Step 1: Constructing covers

Let x_0 be a basepoint of X and $\rho > 0$ such that for $k \in \mathbb{N}$ the sphere $S_k := S_{k \cdot \rho}(x_0)$ is covered by $\mathcal{U}_k = \mathcal{U}_{k,1} \cup \cdots \cup \mathcal{U}_{k,n}$ with Lebesgue number $L(\mathcal{U}_k) \geq \lambda$, mesh $(\mathcal{U}_k) \leq D$ and $\mu(\mathcal{U}_{k,i}) = 1$ for all $i \in \{1, \ldots, n\}$. Observe that $\mathcal{U}_0 = \{\{x_0\}\}$. For $k \in \mathbb{N}$ we define $D_k := D_{k \cdot \rho}(x_0)$.

Since X is a $CAT(\kappa)$ -space, the space X is uniquely geodesic. We denote the geodesic segment from x to y by [x, y].

Two parameters will appear in the following construction: the distance ρ between spheres and the magnitude of shift $N \in \mathbb{N}$ towards the basepoint. Note that whenever we have a space with nicely *n*-colored spheres of distance ρ , we can choose any integral multiple of ρ as a new distance of spheres, i.e. we may choose ρ as large as we want. For $k \in \mathbb{N}$ we define a map $\theta_k \colon X \setminus D_k \to S_k$ by assigning to $x \in X \setminus D_k$ the intersecting point of the geodesic segment $[x_0, x]$ with S_k .

Lemma 6.7 The map θ_k is contracting for any $k \in \mathbb{N}$.

Proof. For $x \in X$ set $r(x) = d(x_0, x)$. Let $x, y \in X \setminus D_k$. Without loss of generality assume $r(x) \leq r(y)$. Take $z \in [x_0, y]$ with r(z) = r(x). Consider the triangle with vertices x_0 , x and y. Take a comparison triangle in M_{κ}^2 and denote its vertices by $\overline{x_0}, \overline{x}$ and \overline{y} . Then

$$d(\theta_k(x), \theta_k(y)) \le d\left(\overline{\theta_k(x)}, \overline{\theta_k(y)}\right) \le d\left(\overline{x}, \overline{z}\right) \le d\left(\overline{x}, \overline{y}\right) = d(x, y) \ .$$

Here $\overline{\theta_k(x)}$ and $\overline{\theta_k(y)}$ are the points on the chosen comparison triangle corresponding to $\theta_k(x)$ and $\theta_k(y)$ respectively.

For $k \in \mathbb{N}$ we choose a map $\Theta_k : \mathcal{U}_{k+n} \to \mathcal{U}_k$. We will see in Corollary 6.10 that for ρ large enough we may assume that Θ_k has the following property:

$$\theta_k(U) \subseteq \Theta_k(U) \quad \text{for } k \in \mathbb{N} \text{ and } U \in \mathcal{U}_{k+n}$$
(6.1)

Let $N \in \mathbb{N}$, $k \in n \cdot \mathbb{N}$ and $i \in \{1, \ldots, n\}$. For $U \in \mathcal{U}_{k,i}$ we define as follows:

$$A(U) := \begin{cases} D_{N+n} & \text{if } k = 0\\ \theta_k^{-1}(U) \cap D_{k+N+n} \setminus D_{k+N+i-1} & \text{if } k > 0 \end{cases}$$

$$B(U) := \theta_k^{-1}(U) \cap D_{k+N+i} \setminus D_{k+N}$$

$$U^{\#} := A(U) \cup \bigcup_{V \in \Theta_k^{-1}(U)} B(V)$$

We get the following cover of X.

$$\mathcal{M}_{\rho,N} := \left\{ U^{\#} \mid k \in n \cdot \mathbb{N}, \ U \in \mathcal{U}_k \right\}$$

Besides ρ and N this cover does depend on the choice of the maps Θ_k .

Lemma 6.8 The multiplicity of $\mathcal{M}_{\rho,N}$ is at most n + 1. Moreover, if the maps Θ_k have property (6.1), then $\mathcal{M}_{\rho,N}$ is uniformly bounded.

Proof. Let $x \in X$. Choose $k \in n \cdot \mathbb{N}$ and $j \in \{1, \ldots, n\}$ such that $x \in D_{k+N+j} \setminus D_{k+N+j-1}$. Observe that for $i \in \{1, \ldots, n\}$ the point $\theta_k(x)$ belongs to at most one set from $\mathcal{U}_{k,i}$. Hence $x \in A(U)$ for at most j sets $U \in \mathcal{U}_k$. Similarly $x \in B(U)$ for at most n+1-j sets $U \in \mathcal{U}_k$.

Assume that the maps Θ_k have property (6.1). Let $U \in \mathcal{U}_k$ Observe that for $x, y \in U^{\sharp}$ we have

$$d(x,y) \le d(x,U) + \operatorname{diam}(U) + d(U,y) \le (N+2n) \cdot \rho + D + (N+2n) \cdot \rho$$
.

Hence $\mathcal{M}_{\rho,N}$ is uniformly bounded.

Step 2: Adjusting parameters

In order to conclude the proof of Theorem 6.5, it remains to prove that for every L > 0 we can choose the parameters ρ and N such that $\mathcal{M}_{\rho,N}$ has Lebesgue number at least L.

Lemma 6.9 Let $\delta > 0$ and $B \subseteq S_{\rho \cdot k+a}(x_0)$ with diam(B) < a. The map $\theta_k|_B \colon B \to S_k$ is δ -Lipschitz if $a > \frac{2}{\sqrt{-\kappa}} \cdot \max\left\{1, \log \frac{2}{\delta}\right\}$.

Proof. Set $r = \rho \cdot k + \frac{a}{2}$. Let $x, y \in B$. Consider the geodesic triangle with vertices x_0 , x and y. Denote the vertices of a comparison triangle in M_{κ}^2 by $\overline{x_0}, \overline{x}$ and \overline{y} respectively. We write $\overline{\theta_k(x)}$ and $\overline{\theta_k(y)}$ for the points on the chosen comparison triangle corresponding to $\theta_k(x)$ and $\theta_k(y)$ respectively. By \overline{x}_m we denote the point on $[\overline{x_0}, \overline{x}]$ with distance $\frac{a}{2}$ to \overline{x} . Similarly, we define a point \overline{y}_m on the segment $[\overline{x_0}, \overline{y}]$. Let α be the angle at $\overline{x_0}$ between $[\overline{x_0}, \overline{x}]$ and $[\overline{x_0}, \overline{y}]$. By γ_r we denote the circular arc parametrized by arc length from \overline{x}_m to \overline{y}_m in $S_r(\overline{x_0}) \subset M_{\kappa}^2$ and by $\gamma_{\rho \cdot k}$ the corresponding arc from $\overline{\theta_k(x)}$ to $\overline{\theta_k(y)}$ in $S_{\rho \cdot k}(\overline{x_0})$. Observe

$$d(\theta_{k}(x), \theta_{k}(y)) \stackrel{(1)}{\leq} d\left(\overline{\theta_{k}(x)}, \overline{\theta_{k}(y)}\right) \stackrel{(2)}{\leq} \text{ length of } \gamma_{\rho \cdot k}$$

$$\stackrel{(3)}{=} \frac{\alpha}{\sqrt{-\kappa}} \cdot \sinh\left(\sqrt{-\kappa} \cdot \rho \cdot k\right)$$

$$< \frac{\alpha}{2 \cdot \sqrt{-\kappa}} \exp\left(\sqrt{-\kappa} \cdot \rho \cdot k\right)$$

$$= \frac{\alpha}{2 \cdot \sqrt{-\kappa}} \exp\left(\sqrt{-\kappa} \cdot r\right) \cdot \exp\left(-\sqrt{-\kappa} \cdot \frac{a}{2}\right)$$

$$\stackrel{(4)}{\leq} \frac{\alpha}{2 \cdot \sqrt{-\kappa}} \exp\left(\sqrt{-\kappa} \cdot r\right) - \exp\left(-\sqrt{-\kappa} \cdot r\right) \cdot \delta$$

$$= \frac{\alpha}{2 \cdot \sqrt{-\kappa}} \left(\exp\left(\sqrt{-\kappa} \cdot r\right) - \exp\left(-\sqrt{-\kappa} \cdot r\right)\right) \cdot \delta$$

$$= \frac{\alpha}{\sqrt{-\kappa}} \cdot \sinh\left(\sqrt{-\kappa} \cdot r\right) \cdot \delta \stackrel{(3)}{=} \delta \cdot \text{ length of } \gamma_{r}$$

$$\stackrel{(6)}{\leq} \delta \cdot d\left(\overline{x}, \overline{y}\right) = \delta \cdot d(x, y) .$$

Since X is a CAT(κ)-space, we can conclude (1). The fact that $\gamma_{\rho \cdot k}$ is a path from $\overline{\theta(x)}$ to $\overline{\theta(y)}$ implies inequality (2). For (3) compare Proposition 6.17 in Part I of [BH99] about the Riemannian metric of M_{κ}^2 in polar coordinates. The assumption $a > \frac{2}{\sqrt{-\kappa}} \cdot \log \frac{2}{\delta}$ implies $\exp\left(-\sqrt{-\kappa} \cdot \frac{a}{2}\right) < \frac{\delta}{2}$ and hence (4). The assumption $a > \frac{2}{\sqrt{-\kappa}}$ yields $\sqrt{-\kappa} \cdot r > 1$. This implies $\exp(\sqrt{-\kappa} \cdot r) < 2 \cdot \left(\exp\left(\sqrt{-\kappa} \cdot r\right) - \exp\left(-\sqrt{-\kappa} \cdot r\right)\right)$ which proves inequality (5). Since $d(\overline{x}, \overline{y}) \leq a$, the geodesic segment $[\overline{x}, \overline{y}]$ lies in $M_{\kappa}^2 \setminus D_r(\overline{x_0})$. For a

Since $d(\overline{x}, \overline{y}) \leq a$, the geodesic segment $[\overline{x}, \overline{y}]$ lies in $M_{\kappa}^2 \setminus D_r(\overline{x_0})$. For a moment consider $M_{\kappa}^2 \setminus D_r(\overline{x_0})$ and $S_r(\overline{x_0})$ as metric spaces with their length

metric. ¹ Note that the radial projection $M_{\kappa}^2 \setminus D_r(\overline{x_0}) \to S_r(\overline{x_0})$ with center $\overline{x_0}$ is a contracting map which maps the geodesic segment $[\overline{x}, \overline{y}]$ onto γ_r . This implies (6).

Remember that λ is a lower bound on the Lebesgue number and D an upper bound on the mesh of the covers \mathcal{U}_i for $i \in \mathbb{N}$.

Corollary 6.10 If $\rho \cdot n > D$ and $\rho \cdot n > \frac{2}{\sqrt{-\kappa}} \cdot \max\left\{1, \log \frac{2 \cdot D}{\lambda}\right\}$, then we can choose Θ_k such that $\theta_k(U) \subseteq \Theta_k(U)$ for all $k \in \mathbb{N}$ and $U \in \mathcal{U}_{k+n}$.

Proof. Set $\delta = \frac{\lambda}{D}$. Lemma 6.9 implies that θ_k maps any $U \in \mathcal{U}_{k+n}$ into a ball $B \subseteq S_k$ of radius λ . Since \mathcal{U}_k has Lebesgue number λ , we may choose $\Theta_k(U)$ such that $B \subseteq \Theta_k(U)$.

Corollary 6.11 Let L > 0. Choose $\rho > 2 \cdot L$ and such that Corollary 6.10 applies. Choose $N \in \mathbb{N}$ such that $(N-1) \cdot \rho > 2L$ and $(N-1) \cdot \rho > \frac{2}{\sqrt{-\kappa}} \cdot \max\left\{1, \log \frac{2 \cdot L}{\lambda}\right\}$. Under these assumptions $\mathcal{M}_{\rho,N}$ has Lebesgue number at least L.

Proof. Let $x \in X$ and let $j \in \mathbb{N}$ be the smallest natural number such that the ball $D_L(x)$ of radius L around x is contained in D_{j+N} . We claim that $D_L(x) \subseteq X \setminus D_{j+N-2}$. Assume $D_L(x) \cap D_{j+N-2} \neq \emptyset$. This implies $D_L(x) \subseteq \{x \in X \mid d(x, D_{j+N-2}) < 2L\} \subseteq D_{j+N-1}$. Here the last inclusion follows from $\rho > 2L$. But this is a contradiction to the definition of j.

Take $k \in n \cdot \mathbb{N}$ and $l \in \{1, \ldots, n\}$ such that j = k + l. Since θ_{k+N-1} is contracting, the image of $D_L(x)$ is contained in a ball $B \subseteq S_{k+N-1}$ of radius L. Set $\delta = \frac{\lambda}{L}$. Lemma 6.9 implies that $\theta_k(B)$ is contained in a ball $B' \subseteq S_k$ of radius λ . Let $U \in \mathcal{U}_{k,i}$ be a covering set containing B'. Then $D_L(x) \subseteq \theta_k^{-1}(U)$.

Hence $D_L(x) \subseteq A(U)$ if $i < l \leq n$. On the other hand, we get $D_L(x) \subseteq B(U)$ if $1 < l \leq i$. Suppose l = 1. Set $V := \Theta_{k-n}(U)$. Note that Corollary 6.10 yields $\theta_k^{-1}(U) \cap D_{k+N} \setminus D_{k+N-1} \subseteq A(V)$. Hence $D_L(x) \subseteq V^{\#}$. \Box

This completes the proof of Theorem 6.5.

6.3 Triangulations of bounded distortion

Simplicial complexes

A (combinatorial) simplicial complex C consists of a set $\operatorname{vert}(C)$ and a collection $\Sigma(C)$ of finite, non-empty subsets of $\operatorname{vert}(C)$ such that $\Sigma(C)$ is closed under taking subsets, i.e. $\sigma \in \Sigma(C)$ and $\tau \subseteq \sigma$ imply $\tau \in \Sigma(C)$. The collection $\Sigma(C)$ must contain $\{v\}$ for every $v \in \operatorname{vert}(C)$. An element $v \in \operatorname{vert}(C)$

¹Compare [BH99] for details on length metrics.

is called a *vertex* of C and $\sigma \in \Sigma(C)$ is called a *simplex* of C. The dimension of σ is defined to be the cardinality of σ minus one. The dimension of C is defined to be the supremum of the dimensions of its simplices, i.e. $\dim(C) = \sup\{\dim(\sigma) \mid \sigma \in \Sigma(C)\}$. By $C^{(k)}$ we denote the k-skeleton of C, i.e. the simplicial complex obtained from C by removing all simplices of dimension greater than k. A simplicial complex is said to be *locally finite* if each vertex is contained in only finitely many simplices.

By δ_v : vert $(C) \to \mathbb{C}$ we denote the function which maps v to 1 and all the other vertices to 0. We identify vert(C) as follows with a subset of $\ell^2(\text{vert}(C))$:

$$\operatorname{vert}(C) \hookrightarrow \ell^2(\operatorname{vert}(C)) , \quad v \mapsto \frac{1}{\sqrt{2}} \cdot \delta_v$$

Observe that the distance of any two vertices in $\ell^2(\operatorname{vert}(C))$ is one. We write $|\sigma|$ for the convex hull of σ in $\ell^2(\operatorname{vert}(C))$ and call $|\sigma|$ a geometric simplex. The euclidean realization |C| of C is defined to be the union of the geometric simplices.

There are two natural metrics on |C|. The *affine metric* is just the restriction of the standard metric on $\ell^2(\operatorname{vert}(C))$ to |C|. The *geodesic metric* is the length metric associated to the affine metric. We will consider |C| with the geodesic metric. Note that restricted to a single simplex both metrics coincide.

We write $\operatorname{Star}(v)$ for the open *star* of $v \in \operatorname{vert}(C)$, i.e. the interior of the union of all geometric simplices $|\sigma| \subseteq |C|$ with $v \in \sigma$.

The *stability* of a finite dimensional simplicial complex C is defined to be the biggest natural number k such that there exist two different k-simplices $\sigma, \tau \in \Sigma(C)$ such that $\sigma \cap \tau$ is a (k-1)-simplex of C.

Lemma 6.12 Let *C* be a simplicial complex of dimension *n* and stability *k*. The cover $\mathcal{V} = {\text{Star}(v) | v \in \text{vert}(C)}$ of |C| is uniformly bounded and has multiplicity n + 1. Moreover \mathcal{V} has Lebesgue number

$$\lambda_k = \frac{1}{\sqrt{2 \cdot k \cdot (k+1)}} \; .$$

Proof. The mesh of the cover \mathcal{V} is at most 2, i.e. \mathcal{V} is uniformly bounded. The multiplicity of \mathcal{V} is n+1, since a simplex of C has at most n+1 vertices. The Lebesgue number of \mathcal{V} is exactly the distance of the center of a k-simplex to a face of dimension k-1. Hence, an elementary computation yields the claim.

Spaces with triangulations of bounded distortion

Definition 6.13 A triangulation of bounded distortion of a metric space X is a tuple (C, ψ, l_1, l_2) where C is a simplicial complex, $l_1, l_2 > 0$ and

 $\psi \colon |C| \to X$ is a homeomorphism such that for every $\sigma \in \Sigma(C)$ the map $\psi|_{|\sigma|}$ is (l_1, l_2) -bi-Lipschitz, i.e.

$$l_1 \cdot d(x, y) \le d(\psi(x), \psi(y)) \le l_2 \cdot d(x, y)$$

for all $x, y \in |\sigma|$. We say that the triangulation is of dimension dim(C).

Note that according to our definition the simplicial complex of a triangulation of bounded distortion does not have to be locally finite.

Proposition 6.14 Let X be a metric space and (C, ψ, l_1, l_2) a triangulation of bounded distortion such that C is of dimension n and stability k. Under these assumptions $\mathcal{U} = \{\psi(\operatorname{Star}(v)) \mid v \in \operatorname{vert}(C)\}$ is a cover of X with the following properties:

- The cover \mathcal{U} is uniformly bounded. More precisely, the diameter of any $U \in \mathcal{U}$ is at most $2 \cdot l_2$.
- When restricted to a single connected component, the cover \mathcal{U} has Lebesgue number $L(\mathcal{U}) \geq l_1 \cdot \lambda_k$ where λ_k is defined as in Lemma 6.12.
- The cover \mathcal{U} has multiplicity n+1.

For the rest of this section let X be a $CAT(\kappa)$ -space which admits a triangulation of bounded distortion (C, ψ, l_1, l_2) of dimension n. Choose a basepoint $x_0 \in X$. As above we set $S_r := \{x \in X \mid d(x, x_0) = r\}$.

Remark 6.15 By restricting the cover \mathcal{U} defined in Proposition 6.14 to the spheres S_r with r > 0 we see that X has nicely (n + 1)-covered spheres. Hence, Theorem 6.5 yields $\operatorname{asdim}(X) \leq n + 1$.

The original goal is to prove $\operatorname{asdim}(X) \leq n$, but this seems to be much more difficult - at least in full generality. However, if there is a radial structure for X as defined below, we do obtain $\operatorname{asdim}(X) \leq n$.

Definition 6.16 Let $\rho > 0$. For $k \in \mathbb{N}$ define C_k to be the smallest simplicial complex containing as simplices all $\sigma \in \Sigma(C)$ such that $\psi(|\sigma|)$ has non-empty intersection with $S_{k \cdot \rho}$. Let D > 0. Assume that for all $k \in \mathbb{N}$ there is an equivalence relation on $\operatorname{vert}(C_k)$ such that

- in each *n*-simplex of C_k there are two equivalent vertices and
- the number D is an upper bound for the diameter of the image under ψ of any equivalence class.

A family of equivalence relations as described above will be called a *radial* structure for X.

One might try to get a radial structure by defining the endpoints of "radial" edges to be equivalent.

Proposition 6.17 If X has a radial structure, then $\operatorname{asdim}(X) \leq n$.

Proof. Consider the collection $\mathcal{U}_k = \{\psi(\operatorname{Star}(v)) \mid v \in \operatorname{vert}(C_k)\}$. We say $U_1, U_2 \in \mathcal{U}_k$ are equivalent if they contain equivalent vertices. For $U \in \mathcal{U}_k$ define \widetilde{U} to be the union of all $V \in \mathcal{U}_k$ which are equivalent to U. By restricting $\widetilde{\mathcal{U}}_k := \{\widetilde{U} \mid U \in \mathcal{U}_k\}$ to $S_{k \cdot \rho}$ we get a uniformly bounded cover of $S_{k \cdot \rho}$ with multiplicity at most n and Lebesgue number at least $L(\mathcal{U})$. Applying Theorem 6.5 completes this proof.

Question 6.18 Is there a $CAT(\kappa)$ -space with a triangulation of bounded distortion which does not admit a radial structure?

6.4 Complete, simply connected manifolds with bounded, strictly negative sectional curvature

Theorem 6.20 has been conjectured by Mikhael Gromov. Compare Example $1.E'_1$ of [Gro93].

Definition 6.19 Let X be a Riemannian manifold. For $x \in X$ and a twodimensional subspace $p \subseteq T_x X$, we denote the sectional curvature of the plane p by $K(p) \in \mathbb{R}$.

We say that the sectional curvature of X is bounded between κ' and κ or simply that X has bounded sectional curvature if $\kappa' \leq K(p) \leq \kappa$ for all $x \in X$ and all two-dimensional subspaces $p \subseteq T_x X$.

We say that X has strictly negative sectional curvature if there is $\kappa < 0$ such that $K(p) \leq \kappa$ for all $x \in X$ and all two-dimensional subspaces $p \subseteq T_x X$.

Theorem 6.20 For a complete, simply connected Riemannian manifold X with bounded, strictly negative sectional curvature we have

$$\operatorname{asdim}(X) = \dim(X).$$

We will prove this theorem by applying Theorem 6.5. A different approach (using the boundary at infinity and a new invariant called capacity dimension) has been taken in [Buy] and [BL] to prove similar results. These results include $\operatorname{asdim}(X) \leq \dim(X)$ if in addition to our assumptions X is cobounded, i.e. if there is a bounded subset $B \subset X$ such that the orbit of B under the isometry group of X covers X. Compare Theorem 1.1 of [Buy] and Corollary 1.2 and Proposition 6.2 of [BL].

Before proceeding with the proof of Theorem 6.20, we will look at the differential geometry of the spheres in X.

Lemma 6.21 Let X be a complete, simply connected Riemannian manifold of dimension n with bounded, strictly negative sectional curvature. Then there are $\kappa_1, \kappa_2 \in \mathbb{R}$ and $\rho > 0$ such that for all $x_0 \in X$ and $r \geq 1$

- (a) the sectional curvature of the sphere $S_r(x_0) = \{x \in X \mid d(x, x_0) = r\}$ is bounded between κ_1 and κ_2 and
- (b) the injectivity radius of $S_r(x_0)$ is at least ρ .

Proof. Let $x_0 \in X$ and $r \ge 1$.

First we calculate bounds for the sectional curvature of $S_r(x_0)$.

Suppose the sectional curvature of X is bounded between $-\kappa'$ and $-\kappa$. Define $f: X \to \mathbb{R}, x \mapsto d(x_0, x)$. Note that for each $x \in X$ the second fundamental tensor $S_{\text{grad } f}: T_x S_r(x_0) \to T_x S_r(x_0)$ is just the restriction of the Hessian² Hess(f) to the directions tangential to the spheres.

Let $x \in X$ with $f(x) = r \ge 1$. Denote the eigenvalues of Hess(f) at x by $\tau_1(x), \dots, \tau_{n-1}(x)$. Using Riccati comparison arguments, we get

$$\sqrt{\kappa} \le \sqrt{\kappa} \cdot \coth\left(\sqrt{\kappa} \cdot r\right) \stackrel{(\flat)}{\le} \tau_i(x) \stackrel{(\sharp)}{\le} \sqrt{\kappa'} \cdot \coth\left(\sqrt{\kappa'} \cdot r\right) \le \sqrt{\kappa'} \cdot \coth\left(\sqrt{\kappa'}\right).$$

For (b) compare 1.7.3 of [Kar87]. For (\sharp) compare 1.7.1 of [Kar87] or Sections 1.4 and 1.6 of [Mey89]. Hence, the norm of the second fundamental form is bounded in terms of κ' and κ . Using the Gauß equation (see [Car92] or [Jos02]) in order to relate the curvature of X and the spheres, we get universal bounds

$$\kappa_1 := \kappa - \kappa' - \kappa' \cdot \coth^2\left(\sqrt{\kappa'}\right) \quad \text{and} \quad \kappa_2 := -2 \cdot \kappa + \kappa' \cdot \coth^2\left(\sqrt{\kappa'}\right)$$

for the sectional curvature of $S_r(x_0)$.

Applying Corollary 4 of [AL] yields that the injectivity radius of $S_r(x_0)$ is at least $\rho := \frac{\pi}{\sqrt{\kappa' \cdot \coth(\sqrt{\kappa'})}}$.

Proof of Theorem 6.20. First note that X is a $CAT(\kappa)$ -space. Compare Theorem II.1A.6 and the Cartan-Hadamard Theorem (II.4.1) of [BH99]. We need completeness of X to apply the Cartan-Hadamard Theorem.

Lemma 6.21 yields that $M := \bigcup_{n \in \mathbb{N}} S_n(x_0) \subset X$ is a Riemannian manifold of bounded geometry, i.e. M has bounded sectional curvature and positive injectivity radius. Hence, we may apply the following theorem.

Theorem 6.22 If M is an n-dimensional Riemannian manifold of bounded geometry, then M admits a triangulation of bounded distortion and the dimension of this triangulation is n.

²If $f: X \to \mathbb{R}$ is a differentiable function on a Riemannian manifold X, ∇ the Levi-Civita connection and Y a tangent vector field on X, then $\operatorname{Hess}(f)(Y) = \nabla_Y \operatorname{grad} f$.

Theorem 6.22 is an immediate consequence of Theorem 1.14 in [Att94]. Moreover, Theorem 6.22 and Proposition 6.14 imply that X has nicely covered spheres. Applying Theorem 6.5 now yields $\operatorname{asdim}(X) \leq \dim(X)$.

Suppose $\operatorname{asdim}(X) =: k < n := \dim(X)$. This implies that there is a uniformly bounded, open cover \mathcal{U} of X with multiplicity at most k + 1. Set $D := \operatorname{mesh}(\mathcal{U})$. Choose a basepoint $x_0 \in X$. Let $\varepsilon > 0$. We consider the following diffeomorphism $f_{\varepsilon} : X \to X$. Define $f_{\varepsilon}(x)$ to be the point on the geodesic segment from x_0 to x with distance $\frac{\varepsilon}{D} \cdot d(x_0, x)$ to x_0 . Observe that $f_{\varepsilon}(x)$ is $\frac{\varepsilon}{D}$ -Lipschitz, since the corresponding map on \mathbb{R}^n is $\frac{\varepsilon}{D}$ -Lipschitz and X is a CAT(0)-space. Now the image of the cover \mathcal{U} under f_{ε} gives an open cover of X with sets of diameter at most ε and with multiplicity at most k + 1. Applying Theorem 1.6.12 of [Eng78] yields $\dim(X) \leq k < n$. This is a contradiction. Hence $\operatorname{asdim}(X) \geq \dim(X)$.

Remark 6.23 We actually proved $\operatorname{asdim}(X) \ge \dim(X)$ for any complete, simply connected Riemannian manifold with strictly negative sectional curvature.

Question 6.24 Is there a complete, simply connected Riemannian manifold with strictly negative, but unbounded sectional curvature and $\operatorname{asdim}(X) > \operatorname{dim}(X)$?

Bibliography

- [AL] Stephanie B. Alexander and Richard L.Bishop, Gauss equation and injectivity radii for subspaces in spaces of curvature bounded above, available at the arXiv. 6.4
- [Att94] Oliver Attie, Quasi-isometry classification of some manifolds of bounded geometry, Math. Z. 216 (1994), 501–527. 6.4
- [Bar03] Arthur C. Bartels, Squeezing and higher algebraic K-theory, K-Theory 28 (2003), no. 1, 19–37. 3
- [BD01] G. Bell and A. Dranishnikov, On asymptotic dimension of groups, Algebr. Geom. Topol. 1 (2001), 57–71 (electronic). 3.3
- [BD04] _____, On asymptotic dimension of groups acting on trees, Geom. Dedicata **103** (2004), 89–101.
- [BH99] Martin R. Bridson and Andre Haefliger, Metric spaces of non-positive curvature, Grundlehren der mathematischen Wissenschaften ed., Springer, 1999. 1.1, 6, 6.2, 1, 6.4
- [BL] Sergei Buyalo and Nina Lebedeva, Dimensions of locally and asymptotically self-similar spaces, available at the arXiv. 5.10, 6.4
- [Buy] Sergei Buyalo, Asymptotic dimension of a hyperbolic space and capacity dimension of its boundary at infinity, available at the arXiv. 6.4
- [Car92] Manfredo P. do Carmo, *Riemannian geometry*, Birkhäuser, 1992. 6.4
- [DKU98] A.N. Dranishnikov, J. Keesling, and V.V. Uspenskij, On the Higson corona of uniformly contractible spaces., Topology 37 (1998), no. 4, 791–803. 5
- [Dra00] A.N. Dranishnikov, *Asymptotic topology*, Russ. Math. Surv. **55** (2000), no. 6, 1085–1129. 1.3, 3.1, 5

- [Eng78] Ryszard Engelking, Dimension theory, North-Holland Publishing Company, Warsaw, 1978. 3, 3.1, 5, 2, 6.4
- [Fed88] V.V. Fedorchuk, The fundamentals of dimension theory, General topology. I. Basic concepts and constructions. Dimension theory. Encycl. Math. Sci. 17, 91-192 (1990); translation from Itogi Nauki Tekh., Ser. Sovrem. Probl. Mat., Fundam. Napravleniya 17, 111-224, 1988. 3, 4, 5, 5, 5, 5
- [Gro93] Mikhael Gromov, Asymptotic invariants of infinite groups, Lecture Notes Series, no. 182, London Mathematical Society, Cambridge University Press, 1993. 3, 3.1, 6.4
- [HR00] Nigel Higson and John Roe, Analytic K-homology, Oxford Mathematical Monographs, Oxford University Press, Oxford, 2000, Oxford Science Publications. 1.2, 1.4, 2, 2.2, 2.2
- [HW41] Witold Hurewicz and Henry Wallman, Dimension theory, Princeton University Press, 1941. 3, 5
- [Jos02] Jürgen Jost, *Riemannian geometry and geometric analysis*, third ed., Springer, 2002. 6.4
- [Kar87] Hermann Karcher, Riemannian comparison constructions, Studies in Mathematics, no. 27, The Mathematical Association of Amerika, 1987, pp. 170–222. 6.4
- [Mey89] Wolfgang T. Meyer, *Topogonov's theorem and applications*, Lecture Notes, College on Differential Geometry, Triest, 1989. 6.4
- [Mit01] Paul D. Mitchener, *Coarse homology theories*, Algebraic & Geometric Topology **1** (2001), 271–297 (electronic). 1.2, 1.2, 1.2
- [Mit03] _____, Addendum to: "Coarse homology theories" [Algebr. Geom. Topol. 1 (2001)], Algebr. Geom. Topol. 3 (2003), 1089– 1101 (electronic). 2, 4, 4
- [ML98] Saunders Mac Lane, Categories for the working mathematician, second ed., Springer, 1998. 1.3
- [Pet98] Peter Petersen, *Riemannian geometry*, Springer, 1998.
- [Roe93] John Roe, Coarse cohomology and index theory on complete Riemannian manifolds., Mem. Am. Math. Soc. **497** (1993). 1.4
- [Roe96] _____, Index theory, coarse geometry, and topology of manifolds, Regional Conference Series in Mathematics. 90. Providence, RI: American Mathematical Society (AMS), 1996. 1.2, 1.2

- [Roe03] _____, Lectures on coarse geometry, University Lecture Series, vol. 31, American Mathematical Society, Providence, RI, 2003. (document), 1.2, 1.2, 1.13, 1.2, 2, 1.4, 1.4, 1.4, 2, 2.13, 2.3, 2.39, 3.1, 3.2, 3.2, 3.2, 3.3, 3.3, 5, 5
- [Spa95] Edwin H. Spanier, Algebraic topology, Springer-Verlag, 1995. 4
- [Yu98] Guoliang Yu, The Novikov conjecture for groups with finite asymptotic dimension, Annals of Mathematics **147** (1998), 325–355. 3

Lebenslauf

Persönliche Daten

Name	Bernd Grave
Geburtsdatum	6. Oktober 1975
Geburtsort	Damme, Niedersachsen
Staatsangehörigkeit	deutsch

Schulausbildung

1982 - 1988	Grundschule und Orientierungsstufe in Holdorf
1988 - 1995	Gymnasiums in Damme
1995	Abitur

Zivildienst

1995 - 1996	Zivildienst in einem	Altenpflegeheim	in Holdorf
-------------	----------------------	-----------------	------------

universitäre Ausbildung

1996 - 1999	Studium der Mathematik und der Physik an der
	Westfälischen-Wilhelms-Universität Münster
1998	Vordiplom in Mathematik und in Physik
1999 - 2000	Studium der Mathematik an der Universidad
	Complutense de Madrid
2000 - 2001	Studium der Mathematik an der Universität Münster
2002	Diplom in Mathematik
seit 2002	Promotionsstudent im Graduiertenkolleg Gruppen und
	Geometrie an der Georg-August-Universität Göttingen
2004	Forschungsaufenthalt an der Vanderbilt University
	in Nashville, Tennessee

Göttingen, den 19. Dezember 2005

Bernd Grave