

Coassociative 4-folds with Conical Singularities

JASON DEAN LOTAY
 University College
 Oxford

1 Introduction

This paper is dedicated to the study of *deformations of coassociative 4-folds* in a G_2 manifold which have *conical singularities*. Understanding the deformations of such singular coassociative 4-folds should be a useful step towards attempting to prove a 7-dimensional analogue of the SYZ conjecture. The research detailed here is motivated by the work on the deformation theory of special Lagrangian m -folds with conical singularities by Joyce in the series of papers [6], [7], [8], [9] and [10], and the work of the author in [15] on deformations of asymptotically conical coassociative 4-folds.

We begin, in Section 2, by discussing the notions of G_2 structures, G_2 manifolds and coassociative 4-folds. In Section 3 we introduce a distinguished class of singular manifolds known as *CS manifolds*. CS manifolds have conical singularities and their nonsingular part is a *noncompact* Riemannian manifold. We also define what we mean by CS coassociative 4-folds.

In order that we may employ various analytic techniques in the course of our study, we choose to use *weighted Banach spaces* of forms on the nonsingular part of a CS manifold. These spaces are described in §4. We then focus, in Section 5, on a particular linear, elliptic, first-order differential operator acting between weighted Banach spaces in the case of a 4-dimensional CS manifold. The Fredholm and index theory of this operator is discussed using the theory developed in [14].

In Section 6 we stratify the types of deformations allowed into three problems, each with an associated nonlinear first-order differential operator whose kernel gives a local description of the moduli space. The main result for each problem, given in §7, states that the moduli space is locally homeomorphic to the kernel of a smooth map between smooth manifolds. In each case, the map in question can be considered as a projection from the *infinitesimal deforma-*

tion space onto the obstruction space. Thus, when there are no obstructions the moduli space is a smooth manifold. Furthermore, using the material in §5 helps to provide a lower bound on the expected dimension of the moduli space.

The last section shows that, in weakening the condition on the G_2 structure of the ambient 7-manifold, there is a generic smoothness result for the moduli spaces of deformations corresponding to our second and third problems.

Notes

- (a) Manifolds are taken to be nonsingular and submanifolds to be embedded, for convenience, unless stated otherwise.
- (b) We use the convention that the natural numbers $\mathbb{N} = \{0, 1, 2, \dots\}$.

2 Coassociative 4-folds

The key to defining coassociative 4-folds lies with the introduction of a distinguished 3-form on \mathbb{R}^7 .

Definition 2.1 Let (x_1, \dots, x_7) be coordinates on \mathbb{R}^7 and write $d\mathbf{x}_{i_j \dots k}$ for the form $dx_i \wedge dx_j \wedge \dots \wedge dx_k$. Define a 3-form φ_0 by:

$$\varphi_0 = d\mathbf{x}_{123} + d\mathbf{x}_{145} + d\mathbf{x}_{167} + d\mathbf{x}_{246} - d\mathbf{x}_{257} - d\mathbf{x}_{347} - d\mathbf{x}_{356}. \quad (1)$$

The 4-form $*\varphi_0$, where φ_0 and $*\varphi_0$ are related by the Hodge star, is given by:

$$*\varphi_0 = d\mathbf{x}_{4567} + d\mathbf{x}_{2367} + d\mathbf{x}_{2345} + d\mathbf{x}_{1357} - d\mathbf{x}_{1346} - d\mathbf{x}_{1256} - d\mathbf{x}_{1247}. \quad (2)$$

Our choice of expression (1) for φ_0 follows that of [5, Chapter 10]. This form is sometimes known as the G_2 3-form because the Lie group G_2 is the subgroup of $GL(7, \mathbb{R})$ preserving φ_0 .

Definition 2.2 A 4-dimensional submanifold N of \mathbb{R}^7 is coassociative if and only if $\varphi_0|_N \equiv 0$ and $*\varphi_0|_N > 0$.

This definition is not standard but is equivalent to the usual definition in the language of *calibrated geometry* by [3, Proposition IV.4.5 & Theorem IV.4.6].

Remark The condition $\varphi_0|_N \equiv 0$ forces $*\varphi_0$ to be a nonvanishing 4-form on N . Thus, the positivity of $*\varphi_0|_N$ is equivalent to a choice of orientation on N .

So that we may describe coassociative submanifolds of more general 7-manifolds, we make two definitions following [2, p. 7] and [5, p. 243].

Definition 2.3 Let M be an oriented 7-manifold. For each $x \in M$ there exists an orientation preserving isomorphism $\iota_x : T_x M \rightarrow \mathbb{R}^7$. Since $\dim \mathbf{G}_2 = 14$, $\dim \mathrm{GL}_+(T_x M) = 49$ and $\dim \Lambda^3 T_x^* M = 35$, the $\mathrm{GL}_+(T_x M)$ orbit of $\iota_x^*(\varphi_0)$ in $\Lambda^3 T_x^* M$, denoted $\Lambda_+^3 T_x^* M$, is open. A 3-form φ on M is *definite*, or *positive*, if $\varphi|_{T_x M} \in \Lambda_+^3 T_x^* M$ for all $x \in M$. Denote the bundle of definite 3-forms $\Lambda_+^3 T^* M$. It is a bundle with fibre $\mathrm{GL}_+(7, \mathbb{R})/\mathbf{G}_2$ which is *not* a vector subbundle of $\Lambda^3 T^* M$.

Essentially, a definite 3-form is identified with the \mathbf{G}_2 3-form on \mathbb{R}^7 at each point in M . Therefore, to each definite 3-form φ we can uniquely associate a 4-form $*\varphi$ and a metric g on M such that the triple $(\varphi, *\varphi, g)$ corresponds to $(\varphi_0, *\varphi_0, g_0)$ at each point. This leads us to our next definition.

Definition 2.4 Let M be an oriented 7-manifold, let φ be a definite 3-form on M and let g be the metric associated to φ . We call (φ, g) a \mathbf{G}_2 *structure* on M . If φ is closed (or coclosed) then (φ, g) is a *closed* (or *coclosed*) \mathbf{G}_2 structure. A closed and coclosed \mathbf{G}_2 structure is called *torsion-free*.

Our choice of notation here agrees with [2].

Remark There is a 1-1 correspondence between pairs (φ, g) and principal \mathbf{G}_2 subbundles of the frame bundle.

Our definition of torsion-free \mathbf{G}_2 structure is not standard, but agrees with other definitions by the following result [19, Lemma 11.5].

Proposition 2.5 *Let (φ, g) be a \mathbf{G}_2 structure and let ∇ be the Levi-Civita connection of g . The following are equivalent:*

$$d\varphi = d^*\varphi = 0; \quad \nabla\varphi = 0; \quad \text{and} \quad \mathrm{Hol}(g) \subseteq \mathbf{G}_2 \text{ with } \varphi \text{ as the associated 3-form.}$$

Definition 2.6 Let M be an oriented 7-manifold endowed with a \mathbf{G}_2 structure (φ, g) , denoted (M, φ, g) . We say that (M, φ, g) is a φ -*closed*, or φ -*coclosed*, 7-manifold if (φ, g) is a closed, respectively coclosed, \mathbf{G}_2 structure. If (φ, g) is torsion-free, we call (M, φ, g) a \mathbf{G}_2 *manifold*.

We are now able to complete our definitions.

Definition 2.7 A 4-dimensional submanifold N of (M, φ, g) is coassociative if and only if $\varphi|_N \equiv 0$ and $*\varphi|_N > 0$.

We end this section with a result, which follows from [16, Proposition 4.2], that is invaluable in describing the deformation theory of coassociative 4-folds.

Proposition 2.8 *Let N be a coassociative 4-fold in (M, φ, g) . There is an isomorphism between the normal bundle $\nu(N)$ of N in M and $\Lambda_+^2 T^*N$ given by $v \mapsto (v \cdot \varphi)|_{TN}$.*

3 Conical singularities

3.1 CS manifolds

Definition 3.1 Let M be a connected Hausdorff topological space and let $z_1, \dots, z_s \in M$. Suppose that $\hat{M} = M \setminus \{z_1, \dots, z_s\}$ has the structure of a (nonsingular) n -dimensional Riemannian manifold, with Riemannian metric g , compatible with its topology. Then M is a *manifold with conical singularities* (at z_1, \dots, z_s with rate λ) if there exist constants $\epsilon > 0$ and $\lambda > 1$, a compact $(n-1)$ -dimensional Riemannian manifold (Σ_i, h_i) , an open set $U_i \ni z_i$ in M with $U_i \cap U_j = \emptyset$ for $j \neq i$ and a diffeomorphism $\Psi_i : (0, \epsilon) \times \Sigma_i \rightarrow U_i \setminus \{z_i\} \subseteq \hat{M}$, for $i = 1, \dots, s$, such that

$$|\nabla_i^j(\Psi_i^*(g) - g_i)| = O(r_i^{\lambda-1-j}) \quad \text{for } j \in \mathbb{N} \text{ as } r_i \rightarrow 0, \quad (3)$$

where r_i is the coordinate on $(0, \infty)$ on the cone $C_i = (0, \infty) \times \Sigma_i$, $g_i = dr_i^2 + r_i^2 h_i$ is the conical metric on C_i , ∇_i is the Levi-Civita connection derived from g_i and $|\cdot|$ is calculated using g_i . We call C_i the *cone* at the singularity z_i and let the *ends* \hat{M}_∞ of \hat{M} be the disjoint union

$$\hat{M}_\infty = \bigsqcup_{i=1}^s U_i \setminus \{z_i\}.$$

We say that M is *CS* or a *CS manifold* (with rate λ) if it is a manifold with conical singularities which have rate λ and it is compact as a topological space. In these circumstances it may be written as the disjoint union

$$M = K \sqcup \bigsqcup_{i=1}^s U_i,$$

where K is compact as it is closed in M .

The condition $\lambda > 1$ guarantees that the metric on \hat{M} genuinely converges to the conical metric on C_i , as is evident from (3). Since M is supposed to be Hausdorff, the set $U_i \setminus \{z_i\}$ is open in \hat{M} for all i . Moreover, the condition that the U_i are disjoint may be easily satisfied since, if $i \neq j$, z_i and z_j may be separated by two disjoint open sets and, by hypothesis, there are only a finite number of singularities.

Remark If M is a CS manifold, \hat{M} is a *noncompact* manifold.

Definition 3.2 Let M be a CS manifold. Using the notation of Definition 3.1, a *radius function* on \hat{M} is a smooth function $\rho : \hat{M} \rightarrow (0, 1]$, bounded below by a positive constant on $\hat{M} \setminus \hat{M}_\infty$, such that there exist positive constants $c_1 < 1$ and $c_2 > 1$ with

$$c_1 r_i < \Psi_i^*(\rho) < c_2 r_i$$

on $(0, \epsilon) \times \Sigma_i$ for $i = 1, \dots, s$.

If M is CS we may construct a radius function on \hat{M} as follows. Let $\rho(x) = 1$ for all $x \in \hat{M} \setminus \hat{M}_\infty$. Define $\rho_i : \Psi_i((0, \epsilon/2) \times \Sigma_i) \rightarrow (0, 1)$ to be equal to r_i/ϵ for $i = 1, \dots, s$ and then define ρ by interpolating smoothly between its definition on $\hat{M} \setminus \hat{M}_\infty$ and ρ_i on each of the disjoint sets $\Psi_i((\epsilon/2, \epsilon) \times \Sigma_i)$.

3.2 CS coassociative 4-folds

Let $B(0; \eta)$ denote the open ball about 0 in \mathbb{R}^7 with radius $\eta > 0$, i.e. $B(0; \eta) = \{\mathbf{v} \in \mathbb{R}^7 : |\mathbf{v}| < \eta\}$. We define a preferred choice of local coordinates on a G_2 manifold near a finite set of points.

Definition 3.3 Let (M, φ, g) be a G_2 manifold as in Definition 2.6 and let z_1, \dots, z_s be points in M . There exist a constant $\eta > 0$, an open set $V_i \ni z_i$ in M with $V_i \cap V_j = \emptyset$ for $j \neq i$ and a diffeomorphism $\chi_i : B(0; \eta) \subseteq \mathbb{R}^7 \rightarrow V_i$ with $\chi_i(0) = z_i$, for $i = 1, \dots, s$, such that $\zeta_i = d\chi_i|_0 : \mathbb{R}^7 \rightarrow T_{z_i}M$ is an isomorphism identifying the standard G_2 structure (φ_0, g_0) on \mathbb{R}^7 with the pair $(\varphi|_{T_{z_i}M}, g|_{T_{z_i}M})$. We call the set $\{\chi_i : B(0; \eta) \rightarrow V_i : i = 1, \dots, s\}$ a *G_2 coordinate system near z_1, \dots, z_s* .

We say that two G_2 coordinate systems near z_1, \dots, z_s , with maps χ_i and $\tilde{\chi}_i$ for $i = 1, \dots, s$ respectively, are *equivalent* if $d\tilde{\chi}_i|_0 = d\chi_i|_0 = \zeta_i$ for all i .

The definition above is an analogue of the local coordinate system for almost Calabi–Yau manifolds used by Joyce [6, Definition 3.6]. Although the family of G_2 coordinate systems near z_1, \dots, z_s is clearly infinite-dimensional, there are only finitely many equivalence classes, given by the number of possible sets $\{\zeta_1, \dots, \zeta_s\}$. Moreover, the family of choices for each ζ_i is isomorphic to G_2 .

Note Definition 3.3 does not require the G_2 structure (φ, g) to be *torsion-free*.

Definition 3.4 Let (M, φ, g) be a G_2 manifold, let $N \subseteq M$ be compact and connected and let $z_1, \dots, z_s \in N$. We say that N is a 4-fold in M with *conical singularities at z_1, \dots, z_s with rate λ* , denoted a *CS 4-fold*, if $\hat{N} = N \setminus \{z_1, \dots, z_s\}$

is a (nonsingular) 4-dimensional submanifold of M and there exist constants $0 < \epsilon < \eta$ and $\lambda > 1$, a compact 3-dimensional Riemannian submanifold (Σ_i, h_i) of $\mathcal{S}^6 \subseteq \mathbb{R}^7$, where h_i is the restriction of the round metric on \mathcal{S}^6 to Σ_i , an open set $U_i \ni z_i$ in N with $U_i \subseteq V_i$ and a smooth map $\Phi_i : (0, \epsilon) \times \Sigma_i \rightarrow B(0; \eta) \subseteq \mathbb{R}^7$, for $i = 1, \dots, s$, such that $\Psi_i = \chi_i \circ \Phi_i : (0, \epsilon) \times \Sigma_i \rightarrow U_i \setminus \{z_i\}$ is a diffeomorphism and Φ_i satisfies

$$|\nabla_i^j(\Phi_i(r_i, \sigma_i) - \iota_i(r_i, \sigma_i))| = O(r_i^{\lambda-j}) \quad \text{for } j \in \mathbb{N} \text{ as } r_i \rightarrow 0, \quad (4)$$

where $\iota_i(r_i, \sigma_i) = r_i \sigma_i \in B(0; \eta)$, ∇_i is the Levi-Civita connection of the cone metric $g_i = dr_i^2 + r_i^2 h_i$ on $C_i = (0, \infty) \times \Sigma_i$ coupled with partial differentiation on \mathbb{R}^7 , $|\cdot|$ is calculated with respect to g_i and $\{\chi_i : B(0; \eta) \rightarrow V_i : i = 1, \dots, s\}$ is a G_2 coordinate system near z_1, \dots, z_s .

We call C_i the *cone* at the singularity z_i and Σ_i the *link* of the cone C_i . We may write N as the disjoint union

$$N = K \sqcup \bigsqcup_{i=1}^s U_i,$$

where K is compact.

If \hat{N} is coassociative in M , we say that N is a *CS coassociative 4-fold*.

Suppose N is a CS 4-fold at z_1, \dots, z_s with rate λ in (M, φ, g) and use the notation of Definition 3.4. The induced metric on \hat{N} , $g|_{\hat{N}}$, makes \hat{N} into a Riemannian manifold. Moreover, it is clear from (4) that the maps Ψ_i satisfy (3) in Definition 3.1 with the same constant λ . Thus, N may be considered as a CS manifold with rate λ .

It is important to note that, if $\lambda \in (1, 2)$, Definition 3.4 is independent of the choice of G_2 coordinate system near the singularities, up to equivalence. Suppose we have two equivalent coordinate systems defined using maps χ_i and $\tilde{\chi}_i$. These maps must agree up to second order since the zero and first order behaviour of each is prescribed, as stated in Definition 3.3. Therefore, the transformed maps $\tilde{\Phi}_i$ corresponding to $\tilde{\chi}_i$ such that $\tilde{\Psi}_i = \tilde{\chi}_i \circ \tilde{\Phi}_i = \chi_i \circ \Phi_i = \Psi_i$ are defined by:

$$\tilde{\Phi}_i = (\tilde{\chi}_i^{-1} \circ \chi_i) \circ \Phi_i.$$

Hence

$$|\nabla_i^j(\tilde{\Phi}_i(r_i, \sigma_i) - \Phi_i(r_i, \sigma_i))| = O(r_i^{2-j}) \quad \text{for } j \in \mathbb{N} \text{ as } r_i \rightarrow 0,$$

where ∇_i and $|\cdot|$ are calculated as in Definition 3.4. Thus, in order that the terms generated by the transformation of the G_2 coordinate system neither dominate nor be of equal magnitude to the $O(r_i^{\lambda-j})$ terms given in (4), we need $\lambda < 2$.

We now make a definition which also depends only on equivalence classes of G_2 coordinate systems near the singularities.

Definition 3.5 Let N be a CS 4-fold at z_1, \dots, z_s in a G_2 manifold (M, φ, g) . Use the notation of Definitions 3.3 and 3.4. For $i = 1, \dots, s$ define a cone \hat{C}_i in $T_{z_i}M$ by $\hat{C}_i = (\zeta_i \circ \iota_i)(C_i)$. We call \hat{C}_i the *tangent cone* at z_i .

One can show that \hat{C}_i is a tangent cone to N at z_i in the sense of *geometric measure theory* (see, for example, [4, p. 233]). We also have a straightforward result relating to the tangent cones at singular points of CS coassociative 4-folds.

Proposition 3.6 *Let N be a CS coassociative 4-fold at z_1, \dots, z_s in a G_2 manifold (M, φ, g) . The tangent cones at z_1, \dots, z_s are coassociative.*

Proof: Use the notation of Definitions 3.3 and 3.4.

It is enough to show that $\iota_i(C_i)$ is coassociative in \mathbb{R}^7 for all i , since $\zeta_i : \mathbb{R}^7 \rightarrow T_{z_i}M$ is an isomorphism identifying (φ_0, g_0) with $(\varphi|_{T_{z_i}M}, g|_{T_{z_i}M})$. This is equivalent to the condition $\iota_i^*(\varphi_0) \equiv 0$ for $i = 1, \dots, s$.

Note that $\varphi|_{\hat{N}} \equiv 0$ implies that, for all i , $\varphi|_{U_i \setminus \{z_i\}} \equiv 0$. Hence, $\Psi_i^*(\varphi) = \Phi_i^*(\chi_i^*(\varphi))$ vanishes on C_i for all i . Using (4),

$$|\Phi_i^*(\chi_i^*(\varphi)) - \iota_i^*(\chi_i^*(\varphi))| = O(r_i^{\lambda-1}) \quad \text{as } r_i \rightarrow 0$$

for all i . Moreover,

$$|\iota_i^*(\chi_i^*(\varphi)) - \iota_i^*(\varphi_0)| = O(r_i) \quad \text{as } r_i \rightarrow 0$$

since

$$\chi_i^*(\varphi) = \varphi_0 + O(r_i) \quad \text{and} \quad |\nabla \iota_i| = O(1) \quad \text{as } r_i \rightarrow 0.$$

Therefore, because $\lambda > 1$,

$$|\iota_i^*(\varphi_0)| \rightarrow 0 \quad \text{as } r_i \rightarrow 0$$

for all i . As $T_{r_i \sigma_i \iota_i}(C_i) = T_{\sigma_i \iota_i}(C_i)$ for all $(r_i, \sigma_i) \in C_i$, $|\iota_i^*(\varphi_0)|$ is independent of r_i and thus vanishes for all i as required. \square

4 Weighted Banach spaces

For this section let M be an n -dimensional CS manifold and let \hat{M} be its non-singular part as in Definition 3.1. We define *weighted* Banach spaces of forms as in [1, §1], as well as the usual ‘unweighted’ spaces.

Definition 4.1 Let $p \geq 1$ and let $k, m \in \mathbb{N}$ with $m \leq n$. The *Sobolev space* $L_k^p(\Lambda^m T^* \hat{M})$ is the set of m -forms ξ on \hat{M} which are k times weakly differentiable and such that the norm

$$\|\xi\|_{L_k^p} = \left(\sum_{j=0}^k \int_{\hat{M}} |\nabla^j \xi|^p dV_g \right)^{\frac{1}{p}} \quad (5)$$

is finite. The normed vector space $L_k^p(\Lambda^m T^* \hat{M})$ is a Banach space for all $p \geq 1$ and $L_k^2(\Lambda^m T^* \hat{M})$ is a Hilbert space.

We introduce the space of m -forms

$$L_{k, \text{loc}}^p(\Lambda^m T^* \hat{M}) = \{\xi : f\xi \in L_k^p(\Lambda^m T^* \hat{M}) \text{ for all } f \in C_{\text{cs}}^\infty(\hat{M})\}$$

where $C_{\text{cs}}^\infty(\hat{M})$ is the space of smooth functions on \hat{M} with compact support.

Let $\mu \in \mathbb{R}$ and let ρ be a radius function on \hat{M} . The *weighted Sobolev space* $L_{k, \mu}^p(\Lambda^m T^* \hat{M})$ of m -forms ξ on \hat{M} is the subspace of $L_{k, \text{loc}}^p(\Lambda^m T^* \hat{M})$ such that the norm

$$\|\xi\|_{L_{k, \mu}^p} = \left(\sum_{j=0}^k \int_{\hat{M}} |\rho^{j-\mu} \nabla^j \xi|^p \rho^{-n} dV_g \right)^{\frac{1}{p}} \quad (6)$$

is finite. Then $L_{k, \mu}^p(\Lambda^m T^* \hat{M})$ is a Banach space and $L_{k, \mu}^2(\Lambda^m T^* \hat{M})$ is a Hilbert space.

We may note here, trivially, that $L_0^p(\Lambda^m T^* \hat{M})$ is equal to the standard L^p -space of m -forms on \hat{M} . Further, by comparing equations (5) and (6) for the respective norms, $L^p(\Lambda^m T^* \hat{M}) = L_{0, -\frac{n}{p}}^p(\Lambda^m T^* \hat{M})$. In particular,

$$L^2(\Lambda^m T^* \hat{M}) = L_{0, -\frac{n}{2}}^2(\Lambda^m T^* \hat{M}). \quad (7)$$

For the following two definitions we take $C_{\text{loc}}^k(\Lambda^m T^* \hat{M})$ to be the vector space of k times continuously differentiable m -forms.

Definition 4.2 Let ρ be a radius function on \hat{M} , let $\mu \in \mathbb{R}$ and let $k, m \in \mathbb{N}$ with $m \leq n$. The *weighted C^k -space* $C_\mu^k(\Lambda^m T^* \hat{M})$ of m -forms ξ on \hat{M} is the subspace of $C_{\text{loc}}^k(\Lambda^m T^* \hat{M})$ such that the norm

$$\|\xi\|_{C_\mu^k} = \sum_{j=0}^k \sup_{\hat{M}} |\rho^{j-\mu} \nabla^j \xi|$$

is finite. We also define

$$C_\mu^\infty(\Lambda^m T^* \hat{M}) = \bigcap_{k \geq 0} C_\mu^k(\Lambda^m T^* \hat{M}).$$

Then $C_\mu^k(\Lambda^m T^* \hat{M})$ is a Banach space but in general $C_\mu^\infty(\Lambda^m T^* \hat{M})$ is not.

In the next definition we refer to the usual normed vector space $C^k(\Lambda^m T^* \hat{M})$ of k times continuously differentiable m -forms such that the following norm is finite:

$$\|\xi\|_{C^k} = \sum_{j=0}^k \sup_{\hat{M}} |\nabla^j \xi|.$$

Definition 4.3 Let $d(x, y)$ be the geodesic distance between points $x, y \in \hat{M}$ and let ρ be a radius function on \hat{M} . Let $a \in (0, 1)$ and let $k, m \in \mathbb{N}$ with $m \leq n$. Let

$$H = \{(x, y) \in \hat{M} \times \hat{M} : x \neq y, c_1 \rho(x) \leq \rho(y) \leq c_2 \rho(x) \text{ and} \\ \text{there exists a geodesic in } \hat{M} \text{ of length } d(x, y) \text{ from } x \text{ to } y\},$$

where $0 < c_1 < 1 < c_2$ are constant. A section s of a vector bundle V on \hat{M} , endowed with a connection, is *Hölder continuous* (with *exponent* a) if

$$[s]^a = \sup_{(x, y) \in H} \frac{|s(x) - s(y)|_V}{d(x, y)^a} < \infty.$$

We understand the quantity $|s(x) - s(y)|_V$ as follows. Given $(x, y) \in H$, there exists a geodesic γ of length $d(x, y)$ connecting x and y . Parallel translation along γ using the connection on V identifies the fibres over x and y and the metrics on them. Thus, with this identification, $|s(x) - s(y)|_V$ is well-defined.

The *Hölder space* $C^{k, a}(\Lambda^m T^* \hat{M})$ is the set of $\xi \in C^k(\Lambda^m T^* \hat{M})$ such that $\nabla^k \xi$ is Hölder continuous (with exponent a) and the norm

$$\|\xi\|_{C^{k, a}} = \|\xi\|_{C^k} + [\nabla^k \xi]^a$$

is finite. The normed vector space $C^{k, a}(\Lambda^m T^* \hat{M})$ is a Banach space.

We also introduce the notation

$$C_{\text{loc}}^{k, a}(\Lambda^m T^* \hat{M}) \\ = \{\xi \in C_{\text{loc}}^k(\Lambda^m T^* \hat{M}) : f \xi \in C^{k, a}(\Lambda^m T^* \hat{M}) \text{ for all } f \in C_{\text{cs}}^\infty(\hat{M})\}.$$

Let $\mu \in \mathbb{R}$. The *weighted Hölder space* $C_\mu^{k, a}(\Lambda^m T^* \hat{M})$ of m -forms ξ on \hat{M} is the subspace of $C_{\text{loc}}^{k, a}(\Lambda^m T^* \hat{M})$ such that the norm

$$\|\xi\|_{C_\mu^{k, a}} = \|\xi\|_{C_\mu^k} + [\xi]_\mu^{k, a}$$

is finite, where

$$[\xi]_\mu^{k, a} = [\rho^{k+a-\mu} \nabla^k \xi]^a.$$

Then $C_\mu^{k,a}(\Lambda^m T^* \hat{M})$ is a Banach space. It is clear that we have an embedding $C_\mu^{k,a}(\Lambda^m T^* \hat{M}) \hookrightarrow C_\mu^l(\Lambda^m T^* \hat{M})$ whenever $l \leq k$.

We shall need the analogue of the Sobolev Embedding Theorem for weighted spaces, which is adapted from [14, Lemma 7.2] and [1, Theorem 1.2].

Theorem 4.4 (Weighted Sobolev Embedding Theorem) *Let $p, q \geq 1$, $a \in (0, 1)$, $\mu, \nu \in \mathbb{R}$ and $k, l, m \in \mathbb{N}$ with $m \leq n$.*

(a) *If $k \geq l$, $k - \frac{n}{p} \geq l - \frac{n}{q}$ and either*

- (i) *$p \leq q$ and $\mu \geq \nu$ or*
- (ii) *$p > q$ and $\mu > \nu$,*

there is a continuous embedding $L_{k,\mu}^p(\Lambda^m T^ \hat{M}) \hookrightarrow L_{l,\nu}^q(\Lambda^m T^* \hat{M})$.*

(b) *If $k - \frac{n}{p} \geq l + a$, there is a continuous embedding $L_{k,\mu}^p(\Lambda^m T^* \hat{M}) \hookrightarrow C_\mu^{l,a}(\Lambda^m T^* \hat{M})$.*

We shall also require an Implicit Function Theorem for Banach spaces, which follows immediately from [12, Chapter 6, Theorem 2.1].

Theorem 4.5 (Implicit Function Theorem) *Let X and Y be Banach spaces and let $W \subseteq X$ be an open neighbourhood of 0. Let $\mathcal{G} : W \rightarrow Y$ be a C^k map ($k \geq 1$) such that $\mathcal{G}(0) = 0$. Suppose further that $d\mathcal{G}|_0 : X \rightarrow Y$ is surjective with kernel K such that $X = K \oplus A$ for some closed subspace A of X . There exist open sets $V \subseteq K$ and $V' \subseteq A$, both containing 0, with $V \times V' \subseteq W$, and a unique C^k map $\mathcal{V} : V \rightarrow V'$ such that*

$$\text{Ker } \mathcal{G} \cap (V \times V') = \{(x, \mathcal{V}(x)) : x \in V\}$$

in $X = K \oplus A$.

5 The operator $d + d^*$

In this section we let M be a 4-dimensional CS manifold and let \hat{M} be as in Definition 3.1. An essential part of our study is the use of the Fredholm and index theory for the elliptic operator $d + d^*$ acting from $\Lambda_+^2 T^* \hat{M} \oplus \Lambda^4 T^* \hat{M}$ to $\Lambda^3 T^* \hat{M}$. We therefore consider

$$d + d^* : L_{k+1,\mu}^p(\Lambda_+^2 T^* \hat{M} \oplus \Lambda^4 T^* \hat{M}) \rightarrow L_{k,\mu-1}^p(\Lambda^3 T^* \hat{M}), \quad (8)$$

where $p \geq 2$, $k \in \mathbb{N}$ and $\mu \in \mathbb{R}$.

5.1 Fredholm theory

Our first result follows from [14, Theorem 1.1 & Theorem 6.1].

Proposition 5.1 *There exists a countable discrete set $\mathcal{D} \subseteq \mathbb{R}$ such that (8) is Fredholm if and only if $\mu \notin \mathcal{D}$.*

Moreover, we can give an explicit description of \mathcal{D} by a similar argument to [15, p. 13-14], which is for *asymptotically conical* (AC) manifolds, as follows.

Recall the notation of Definition 3.1. Transform the metric on \hat{M} to a conformally equivalent metric which is asymptotically *cylindrical* on the ends \hat{M}_∞ of \hat{M} ; that is, if (t_i, σ_i) are coordinates on $(0, \infty) \times \Sigma_i$, the metric is asymptotic to $dt_i^2 + h_i$. With respect to this new metric, $d + d^*$ corresponds to

$$(d + d^*)_\infty = e^{mt}(d + e^{-2t}d^*)e^{-mt}$$

acting on m -forms on \hat{M} .

Let

$$\Sigma = \bigsqcup_{i=1}^s \Sigma_i.$$

If $\pi : (0, \infty) \times \Sigma \rightarrow \Sigma$ is the natural projection map, the action of $(d + d^*)_\infty$ on $\pi^*(\Lambda^2 T^* \Sigma) \oplus \pi^*(\Lambda^{\text{odd}} T^* \Sigma)$ is:

$$(d + d^*)_\infty = \begin{pmatrix} d + d^* & \frac{\partial}{\partial t} + 3 - m \\ -(\frac{\partial}{\partial t} + m) & -(d + d^*) \end{pmatrix} \quad (9)$$

where m denotes the operator which multiplies m -forms by a factor m . However, we wish only to consider elements of $\Lambda^1 T^* \Sigma \oplus \Lambda^2 T^* \Sigma$ which correspond to self-dual 2-forms on \hat{M} , so we define $V_\Sigma \subseteq \Lambda^2 T^* \Sigma \oplus \Lambda^{\text{odd}} T^* \Sigma$ by

$$V_\Sigma = \bigsqcup_{i=1}^s \{(\alpha, * \alpha + \beta) : \alpha \in \Lambda^2 T^* \Sigma_i, \beta \in \Lambda^3 T^* \Sigma_i\}.$$

Then $\pi^*(V_\Sigma)$ corresponds to $\Lambda_+^2 T^* \hat{M} \oplus \Lambda^4 T^* \hat{M}$.

For $w \in \mathbb{C}$ define a map $(d + d^*)_\infty(w)$ by:

$$(d + d^*)_\infty(w) = \begin{pmatrix} d + d^* & -w + 3 - m \\ w - m & -(d + d^*) \end{pmatrix} \quad (10)$$

acting on $V_\Sigma \otimes \mathbb{C}$. Notice that we have formally substituted w for $-\frac{\partial}{\partial t}$ in (9).

Let

$$W_\Sigma = \bigsqcup_{i=1}^s \{(*\alpha + \beta, \alpha) : \alpha \in \Lambda^2 T^* \Sigma_i, \beta \in \Lambda^3 T^* \Sigma_i\} \subseteq \Lambda^{\text{odd}} T^* \Sigma \oplus \Lambda^2 T^* \Sigma.$$

Define $\mathcal{C} \subseteq \mathbb{C}$ as the set of w for which the map

$$(d + d^*)_\infty(w) : L_{k+1}^p(V_\Sigma \otimes \mathbb{C}) \rightarrow L_k^p(W_\Sigma \otimes \mathbb{C})$$

is *not* an isomorphism. By the proof of [14, Theorem 1.1], $\mathcal{D} = \{\operatorname{Re} w : w \in \mathcal{C}\}$. By [17, Lemma 6.1.13], the corresponding sets $\mathcal{C}(\Delta^m)$, where Δ^m is the Laplacian on m -forms, are all real for an asymptotically conical manifold. Since the same will be true for the CS case, we deduce that $\mathcal{C} \subseteq \mathbb{R}$. Hence $\mathcal{C} = \mathcal{D}$.

The symbol, hence the index ind_w , of $(d + d^*)_\infty(w)$ is independent of w . Furthermore, $(d + d^*)_\infty(w)$ is an isomorphism for generic values of w since \mathcal{D} is countable and discrete. Therefore $\operatorname{ind}_w = 0$ for all $w \in \mathbb{C}$; that is,

$$\dim \operatorname{Ker}(d + d^*)_\infty(w) = \dim \operatorname{Coker}(d + d^*)_\infty(w),$$

so that (10) is not an isomorphism precisely when it is not injective.

The condition $(d + d^*)_\infty(w) = 0$, using (10), corresponds to the existence of $\alpha \in C^\infty(\Lambda^2 T^* \Sigma_i)$ and $\beta \in C^\infty(\Lambda^3 T^* \Sigma_i)$, for some i , satisfying

$$d\alpha = w\beta \quad \text{and} \quad d^* \alpha + d^* \beta = (w - 2)\alpha. \quad (11)$$

Notes

- (a) The equations above imply that

$$dd^* \beta = \Delta \beta = w(w - 2)\beta.$$

Since eigenvalues of the Laplacian on Σ_i must necessarily be positive, $\beta = 0$ if $w \in (0, 2)$.

- (b) If $w = 0$ and we take $\alpha = 0$, (11) forces β to be coclosed. As there are nontrivial coclosed 3-forms on Σ_i , $(d + d^*)_\infty(0)$ is not injective, so $0 \in \mathcal{D}$.
- (c) Suppose that $w = 2$ lies in \mathcal{D} . Then (11) gives $[\beta] = 0$ in $H_{\text{dR}}^3(\Sigma_i)$. We know that β is harmonic so, by Hodge theory, $\beta = 0$. Therefore $2 \in \mathcal{D}$ if and only if there exists a nonzero closed and coclosed 2-form on Σ_i for some i .

We state a proposition which follows from the work above.

Proposition 5.2 *Let M be a 4-dimensional CS manifold. Use the notation of Definition 3.1. For $i = 1, \dots, s$ let $D(\mu, i) = \{(\alpha, \beta) \in C^\infty(\Lambda^2 T^* \Sigma_i \oplus \Lambda^3 T^* \Sigma_i) : d\alpha = \mu\beta, d^* \alpha + d^* \beta = (\mu - 2)\alpha\}$. The set \mathcal{D} of real numbers μ such that (8) is not Fredholm is given by:*

$$\mathcal{D} = \bigcup_{i=1}^s \{\mu \in \mathbb{R} : D(\mu, i) \neq 0\}.$$

Remark A perhaps more illuminating way to characterise $D(\mu, i)$ is by:

$$(\alpha, \beta) \in D(\mu, i) \iff \xi = (r^{\mu-2}\alpha + r^{\mu-1}dr \wedge * \alpha, r^{\mu-3}dr \wedge \beta)$$

is an $O(r^\mu)$ solution of $(d + d^*)\xi = 0$ on C_i ,

using the notation of Definition 3.1.

Lockhart and McOwen [14, §10] study the Laplacian on m -forms on a manifold with a conical singularity. From this work, which can easily be extended to manifolds with more than one singularity, we can make an important observation about the set \mathcal{D} .

Proposition 5.3 *In the notation of Proposition 5.2, $\mathcal{D} \cap (-2, -1] = \emptyset$.*

Proof: Let

$$\Delta^m : L_{k+1, \mu}^p(\Lambda^m T^* \hat{M}) \longrightarrow L_{k-1, \mu-2}^p(\Lambda^m T^* \hat{M})$$

be the Laplacian on m -forms and denote the set of μ such that it is not Fredholm by $\mathcal{D}(\Delta^m)$. Since $\mu > -2$ and $p \geq 2$ we see that $L_{k+1, \mu}^p \hookrightarrow L_{0, -2}^2 = L^2$ by Theorem 4.4 and (7).

We then apply [14, Theorem 10.2] for the Laplacian on 2-forms and 4-forms on a 4-dimensional CS manifold to see that

$$\mathcal{D}(\Delta^2) \cap (-2, -1] = \mathcal{D}(\Delta^4) \cap (-2, -1] = \emptyset.$$

Note that our rate μ is related to the weighting factor in [14, §10], which we may denote as ν , by $\mu = -\nu - 2$. As it is clear that $\mathcal{D} \subseteq (\mathcal{D}(\Delta^2) \cup \mathcal{D}(\Delta^4))$, the result follows. \square

5.2 Index theory

We begin with some definitions following [14].

Definition 5.4 Use the notation of §5.1. Let $\mu \in \mathcal{D}$. Define $d(\mu)$ to be the dimension of the vector space of solutions of $(d + d^*)_\infty \xi = 0$ of the form

$$\xi(t, \sigma) = e^{-\mu t} p(t, \sigma)$$

where $p(t, \sigma)$ is a polynomial in $t \in (0, \infty)$ with coefficients in $C^\infty(V_\Sigma \otimes \mathbb{C})$.

The next result is immediate from [14, Theorem 1.2].

Theorem 5.5 *Let $\lambda, \lambda' \notin \mathcal{D}$ with $\lambda' \leq \lambda$, where \mathcal{D} is given in Proposition 5.2. For any $\mu \notin \mathcal{D}$ let $\text{ind}_\mu(d + d^*)$ denote the Fredholm index of (8). Then*

$$\text{ind}_{\lambda'}(d + d^*) - \text{ind}_\lambda(d + d^*) = \sum_{\mu \in \mathcal{D} \cap (\lambda', \lambda)} d(\mu).$$

We make a key observation, which shall be used on a number of occasions in later sections.

Proposition 5.6 *Let $\lambda, \lambda' \in \mathbb{R}$ such that $\lambda' \leq \lambda$ and $[\lambda', \lambda] \cap \mathcal{D} = \emptyset$. The kernels, and cokernels, of (8) when $\mu = \lambda$ and $\mu = \lambda'$ are equal.*

Proof: Denote the dimensions of the kernel and cokernel of (8), for $\mu \notin \mathcal{D}$, by $k(\mu)$ and $c(\mu)$ respectively. Since $[\lambda', \lambda] \cap \mathcal{D} = \emptyset$, $k(\lambda) - c(\lambda) = k(\lambda') - c(\lambda')$ and hence

$$k(\lambda) - k(\lambda') = c(\lambda) - c(\lambda'). \quad (12)$$

We know that $k(\lambda) \leq k(\lambda')$ because $L_{k+1, \lambda}^p \hookrightarrow L_{k+1, \lambda'}^p$ by Theorem 4.4 as $\lambda \geq \lambda'$. Similarly, since $c(\mu)$ is equal to the dimension of the kernel of the formal adjoint operator acting on a Sobolev space with weight $-3 - \mu$, $c(\lambda) \geq c(\lambda')$. Noting that the right-hand side of (12) is non-negative and the left-hand side is less than or equal to zero, we conclude that both must be zero. The result follows from the fact that the kernel of $d + d^*$ in $L_{k+1, \lambda}^p$ is contained in the kernel of $d + d^*$ in $L_{k+1, \lambda'}^p$, and vice versa for the cokernels. \square

We can now go further and give a more explicit description of the quantity $d(\mu)$ in Definition 5.4.

Proposition 5.7 *Using the notation of Proposition 5.2 and Definition 5.4, $d(\mu) = \sum_{i=1}^s \dim D(\mu, i)$ for $\mu \in \mathcal{D}$.*

This is an analogue of [15, Proposition 5.4], which is for the AC scenario, and can be proved in exactly the same manner.

6 The deformation problems

We have a common notation for the next three sections. Let N be a CS coassociative 4-fold at z_1, \dots, z_s with rate λ in a G_2 manifold (M, φ, g) . Suppose $\lambda \in (1, 2) \setminus \mathcal{D}$, where \mathcal{D} is defined in Proposition 5.2, and the cone at z_i is C_i with link Σ_i . We shall then use the notation of Definitions 3.4 and 3.5. In particular, we let $\{\chi_i : B(0; \eta) \rightarrow V_i : i = 1, \dots, s\}$, with $d\chi_i|_0 = \zeta_i$ for all i ,

be the G_2 coordinate system near z_1, \dots, z_s used to define N and let \hat{C}_i be the tangent cone at z_i . Recalling that N is a CS manifold, in the sense of Definition 3.1, we have a radius function ρ on \hat{N} as in Definition 3.2.

We consider deformations of N which are CS coassociative 4-folds at s points with rate λ in (M, φ, g) with the same cones at the singularities as N , but the singularities need not be at the same points, nor have identical tangent cone. We also, eventually, consider deforming the G_2 structure on the ambient 7-manifold M .

6.1 Problem 1: fixed singularities and G_2 structure

The first deformation problem we consider is where the deformations of N have identical singular points to N with the same rate, cones and tangent cones, and the G_2 structure of M is fixed.

Definition 6.1 The *moduli space of deformations* $\mathcal{M}_1(N, \lambda)$ for Problem 1 is the set of N' in (M, φ, g) which are CS coassociative 4-folds at z_1, \dots, z_s with rate λ , having cone C_i and tangent cone \hat{C}_i at z_i for all i , such that there exists a homeomorphism $h : N \rightarrow N'$, isotopic to the identity, with $h(z_i) = z_i$ for $i = 1, \dots, s$ and such that $h|_{\hat{N}} : \hat{N} \rightarrow N' \setminus \{z_1, \dots, z_s\}$ is a diffeomorphism.

We begin our formulation of a local description of $\mathcal{M}_1(N, \lambda)$ with a result which is immediate from the proof of [11, Chapter IV, Theorem 9] since M is a Riemannian manifold.

Theorem 6.2 *Let P be a closed embedded submanifold of M . There exist an open subset V of the normal bundle $\nu(P)$ of P in M , containing the zero section, and an open set S in M containing P , such that the exponential map $\exp|_V : V \rightarrow S$ is a diffeomorphism.*

Note The proof of this result relies entirely on the observation that $\exp|_{\nu(P)}$ is a local isomorphism upon the zero section.

This information provides us with a useful corollary.

Corollary 6.3 *For $i = 1, \dots, s$ choose $\Phi_i : (0, \epsilon) \times \Sigma_i \rightarrow B(0; \eta) \subseteq \mathbb{R}^7$ uniquely by imposing the condition that*

$$\Phi_i(r_i, \sigma_i) - \iota_i(r_i, \sigma_i) \in (T_{r_i \sigma_i} \iota_i(C_i))^\perp$$

for all $(r_i, \sigma_i) \in (0, \epsilon) \times \Sigma_i$, which can be achieved by making ϵ smaller and K larger if necessary. Let $P_i = \iota_i((0, \epsilon) \times \Sigma_i)$, $Q_i = \Phi_i((0, \epsilon) \times \Sigma_i)$ and define

$n_i : \nu(P_i) \rightarrow \mathbb{R}^7$ by $n_i(r_i\sigma_i, v) = v + \Phi_i(r_i, \sigma_i)$. For all i , there exist an open subset \hat{V}_i of $\nu(P_i)$ in \mathbb{R}^7 , containing the zero section, and an open set \hat{S}_i in $B(0; \eta) \subseteq \mathbb{R}^7$ containing Q_i such that $n_i|_{\hat{V}_i} : \hat{V}_i \rightarrow \hat{S}_i$ is a diffeomorphism. Moreover, \hat{V}_i and \hat{S}_i can be chosen to grow like r_i on $(0, \epsilon) \times \Sigma_i$, for all i , and such that $P_i \subseteq \hat{S}_i$.

Proof: Note that n_i takes the zero section of $\nu(P_i)$ to Q_i . By the definition of Φ_i , we see that n_i is a local isomorphism upon the zero section. Thus, the proof of Theorem 6.2 gives open sets \hat{V}_i and \hat{S}_i such that $n_i|_{\hat{V}_i} : \hat{V}_i \rightarrow \hat{S}_i$ is a diffeomorphism. We can ensure that \hat{S}_i lies in $B(0; \eta)$ by making \hat{V}_i smaller if necessary.

Furthermore, since $\Phi_i - \iota_i$ is orthogonal to $(0, \epsilon) \times \Sigma_i$, it can be identified with a small section of the normal bundle and hence P_i lies in \hat{S}_i as long as \hat{S}_i grows at $O(r_i)$ as $r_i \rightarrow 0$. As we can form \hat{S}_i and \hat{V}_i in a translation equivariant way because we are working on a portion of the cone C_i , we can construct our sets with this decay rate as $r_i \rightarrow 0$ and such that they do not collapse as $r_i \rightarrow \epsilon$. \square

Corollary 6.3 helps us in establishing the next proposition.

Proposition 6.4 *There exist an open set $\hat{U} \subseteq \Lambda_+^2 T^* \hat{N}$ containing the zero section, an open set $\hat{T} \subseteq M$ containing \hat{N} and a diffeomorphism $\delta : \hat{U} \rightarrow \hat{T}$ which takes the zero section to \hat{N} . Moreover, \hat{U} and \hat{T} can be chosen to grow with order $O(\rho)$ as $\rho \rightarrow 0$ and δ is compatible with the identifications $U_i \setminus \{z_i\} \cong (0, \epsilon) \times \Sigma_i$ for all i and the isomorphism $j : \nu(\hat{N}) \rightarrow \Lambda_+^2 T^* \hat{N}$ given in Proposition 2.8.*

Proof: Use the notation of Corollary 6.3 and define $\hat{T}_i = \chi_i(\hat{S}_i)$. Then \hat{T}_i is an open set in M such that $U_i \setminus \{z_i\} \subseteq \hat{T}_i \subseteq V_i$, since $\chi_i(Q_i) = U_i \setminus \{z_i\}$, and which grows with order $O(\rho)$ as $\rho \rightarrow 0$.

Consider the bundle $(\Lambda_+^2)_{\chi_i^*(g)} T^*((0, \epsilon) \times \Sigma_i)$, where the notation $(\Lambda_+^2)_h$ indicates that the Hodge star is calculated using the metric h and we consider $(0, \epsilon) \times \Sigma_i \cong P_i \subseteq \mathbb{R}^7$. Then

$$\begin{aligned} J_i : \nu(P_i) &\longrightarrow (\Lambda_+^2)_{\chi_i^*(g)} T^* P_i \\ v|_{r_i\sigma_i} &\longmapsto (v|_{r_i\sigma_i} \cdot \chi_i^*(\varphi)|_{\Phi_i(r_i, \sigma_i)})|_{T_{r_i\sigma_i} P_i} \end{aligned}$$

is an isomorphism because $U_i \setminus \{z_i\}$ is coassociative and thus P_i is, with respect to the metric $\chi_i^*(g)$ and 3-form $\chi_i^*(\varphi)$, and hence we may apply Proposition 2.8.

Note also that

$$\Psi_i^* : (\Lambda_+^2)_g T^*(U_i \setminus \{z_i\}) \longrightarrow (\Lambda_+^2)_{\chi_i^*(g)} T^*((0, \epsilon) \times \Sigma_i)$$

is clearly a diffeomorphism. Therefore, let $\hat{U}_i \subseteq (\Lambda_+^2)_g T^*(U_i \setminus \{z_i\})$ be such that $\Psi_i^*(\hat{U}_i) = j_i(\hat{V}_i)$. Note, by construction, that \hat{U}_i grows with order $O(\rho)$ as $\rho \rightarrow 0$.

Define a diffeomorphism $\delta_i : \hat{U}_i \rightarrow \hat{T}_i$ such that the following diagram commutes:

$$\begin{array}{ccc} \hat{U}_i & \xrightarrow{\Psi_i^*} & j_i(\hat{V}_i) \\ \delta_i \downarrow & & \downarrow j_i^{-1} \\ & & \hat{V}_i \\ & & \downarrow n_i \\ \hat{T}_i & \xleftarrow{\chi_i} & \hat{S}_i. \end{array} \quad (13)$$

Interpolating smoothly over K , we extend $\bigcup_{i=1}^s \hat{U}_i$ and $\bigcup_{i=1}^s \hat{T}_i$ to \hat{U} and \hat{T} as required and extend the diffeomorphisms δ_i smoothly to a diffeomorphism $\delta : \hat{U} \rightarrow \hat{T}$ such that δ acts as the identity on \hat{N} , which is identified with the zero section in $\Lambda_+^2 T^* \hat{N}$.

Note that we have a splitting $T\hat{U}|_{(x,0)} = T_x \hat{N} \oplus \Lambda_+^2 T_x^* \hat{N}$ for all $x \in \hat{N}$. Thus we can consider $d\delta$ at \hat{N} as a map from $T\hat{N} \oplus \Lambda_+^2 T^* \hat{N}$ to $T\hat{N} \oplus \nu(\hat{N}) \cong TM|_{\hat{N}}$. Hence, we require in our extension of δ from δ_i to ensure that, in matrix notation,

$$d\delta|_{\hat{N}} = \begin{pmatrix} I & A \\ 0 & j^{-1} \end{pmatrix}, \quad (14)$$

where I is the identity and A is arbitrary. This can be achieved because of the definition of δ_i .

The compatibility of δ with j and Ψ_i for all i , mentioned in the statement of the proposition, is given by (13) and the behaviour of $d\delta|_{\hat{N}}$ stipulated in (14). \square

We now define our deformation map for Problem 1. Let $C_{\text{loc}}^k(\hat{U}) = \{\alpha \in C_{\text{loc}}^k(\Lambda_+^2 T^* \hat{N}) : \alpha \in \hat{U}\}$, where \hat{U} is given in Proposition 6.4, and adopt similar notation to define subsets of the spaces of forms described in §4.

Definition 6.5 Use the notation of Proposition 6.4. Let Γ_α be the graph of $\alpha \in C_{\text{loc}}^1(\hat{U})$ and let $\pi_\alpha : \hat{N} \rightarrow \Gamma_\alpha$ be given by $\pi_\alpha(x) = (x, \alpha(x))$. Let $f_\alpha = \delta \circ \pi_\alpha$ and let $\hat{N}_\alpha = f_\alpha(\hat{N}) \subseteq \hat{T}$. Define a map F_1 from $C_{\text{loc}}^1(\hat{U})$ to $C_{\text{loc}}^0(\Lambda^3 T^* \hat{N})$ by:

$$F_1(\alpha) = f_\alpha^* \left(\varphi|_{\hat{N}_\alpha} \right).$$

By [16, p. 731], which we are allowed to use by our choice of δ , the linearisation of F_1 at 0 is

$$dF_1|_0(\alpha) = L_1(\alpha) = d\alpha$$

for all $\alpha \in C_{\text{loc}}^1(\Lambda_+^2 T^* \hat{N})$.

Remark The operator L_1 is *not* elliptic.

By Proposition 2.7, $\text{Ker } F_1$ is the set of $\alpha \in C_{\text{loc}}^1(\hat{U})$ such that \hat{N}_α is coassociative.

However, we want CS coassociative deformations with singularities at the same points with the same tangent cones. Suppose $\alpha \in C_{\text{loc}}^1(\hat{U})$ and $N_\alpha = \hat{N}_\alpha \cup \{z_1, \dots, z_s\}$ is such a deformation. Then there exist smooth maps $(\Phi_\alpha)_i : (0, \epsilon) \times \Sigma_i \rightarrow B(0; \eta)$ satisfying (4) such that $(\Psi_\alpha)_i = \chi_i \circ (\Phi_\alpha)_i$ is a diffeomorphism onto an open subset of \hat{N}_α for all i as in Definition 3.4. Note that we are free to use χ_i because the tangent cones at the singularities of N_α must be the same as for N , so any G_2 coordinate system near the singularities used to define N_α must be equivalent to the one given by χ_i for $i = 1, \dots, s$. Choose $(\Phi_\alpha)_i$ uniquely such that

$$(\Phi_\alpha)_i(r_i, \sigma_i) - \iota_i(r_i, \sigma_i) \in (T_{r_i \sigma_i} \iota_i(C_i))^\perp$$

for all $(r_i, \sigma_i) \in (0, \epsilon) \times \Sigma_i$.

Use the notation of Corollary 6.3 and the proof of Proposition 6.4. Since

$$\Phi_i(r_i, \sigma_i) - \iota_i(r_i, \sigma_i) \in (T_{r_i \sigma_i} P_i)^\perp \cong \nu_{r_i \sigma_i}(P_i),$$

$\Phi_i - \iota_i$ can be identified using j_i with the graph of $\beta_i \in (\Lambda_+^2)_{\chi_i^*(g)} T^*((0, \epsilon) \times \Sigma_i)$. Thus,

$$|\nabla_i^j \beta_i| = O(r_i^{\lambda-j}) \quad \text{for } j \in \mathbb{N} \text{ as } r_i \rightarrow 0 \quad (15)$$

by (4) and therefore $\beta_i \in C_\lambda^\infty((\Lambda_+^2)_{\chi_i^*(g)} T^*((0, \epsilon) \times \Sigma_i))$.

We may similarly deduce, by the definition of δ , Φ_i and $(\Phi_\alpha)_i$, that $(\Phi_\alpha)_i - \iota_i = ((\Phi_\alpha)_i - \Phi_i) + (\Phi_i - \iota_i)$ corresponds to the graph of $\Psi_i^*(\alpha) + \beta_i$ on $(0, \epsilon) \times \Sigma_i$, recalling that

$$\Psi_i^* : \Lambda_+^2 T^*(U_i \setminus \{z_i\}) \rightarrow (\Lambda_+^2)_{\chi_i^*(g)} T^*((0, \epsilon) \times \Sigma_i)$$

is a diffeomorphism for all i . Since N_α has the same types of singularities as N , both β_i and $\Psi_i^*(\alpha) + \beta_i$ lie in $C_\lambda^\infty((\Lambda_+^2)_{\chi_i^*(g)} T^*((0, \epsilon) \times \Sigma_i))$ for each i . Thus α must lie in $C_\lambda^\infty(\Lambda_+^2 T^* \hat{N})$.

We conclude that \hat{N}_α is a sufficiently nearby deformation of \hat{N} with the same conical singularities if and only if $\alpha \in C_\lambda^\infty(\hat{U}) \subseteq C_\lambda^\infty(\Lambda_+^2 T^* \hat{N})$. We state this as a proposition.

Proposition 6.6 *The moduli space of deformations for Problem 1 is locally homeomorphic to $\text{Ker } F_1 = \{\alpha \in C_\lambda^\infty(\hat{U}) : F_1(\alpha) = 0\}$.*

We define an associated map G_1 to F_1 .

Definition 6.7 Define $G_1 : C_{\text{loc}}^1(\hat{U}) \times C_{\text{loc}}^1(\Lambda^4 T^* \hat{N}) \rightarrow C_{\text{loc}}^0(\Lambda^3 T^* \hat{N})$ by:

$$G_1(\alpha, \beta) = F_1(\alpha) + d^* \beta.$$

Then G_1 is a first order *elliptic* operator at $(0, 0)$ since

$$dG_1|_{(0,0)} = d + d^* : C_{\text{loc}}^1(\Lambda_+^2 T^* \hat{N} \oplus \Lambda^4 T^* \hat{N}) \longrightarrow C_{\text{loc}}^0(\Lambda^3 T^* \hat{N}).$$

Note If $G_1(\alpha, \beta) = 0$ and $\beta \in C_\lambda^\infty(\Lambda^4 T^* \hat{N})$, $*\beta$ is a harmonic function which decays with order $O(\rho^\lambda)$ as $\rho \rightarrow 0$. Since $\lambda > 1$, $*\beta \rightarrow 0$ as $\rho \rightarrow 0$ and hence, by the Maximum Principle for harmonic functions, it must be 0.

We therefore deduce the following.

Proposition 6.8 $\text{Ker } F_1 \cong \{(\alpha, \beta) \in C_\lambda^\infty(\hat{U}) \times C_\lambda^\infty(\Lambda^4 T^* \hat{N}) : G_1(\alpha, \beta) = 0\}$.

We conclude this subsection by stating and proving two results on *regularity* which are analogous to [15, Proposition 4.3] and the argument in [15, p. 22-24] respectively.

Proposition 6.9 *The map F_1 given in Definition 6.5 can be written as*

$$F_1(\alpha)(x) = d\alpha(x) + P_{F_1}(x, \alpha(x), \nabla\alpha(x)) \quad (16)$$

for $x \in \hat{N}$, where $P_{F_1} : \{(x, y, z) : (x, y) \in \hat{U}, z \in T_x^* \hat{N} \otimes \Lambda_+^2 T_x^* \hat{N}\} \rightarrow \Lambda^3 T^* \hat{N}$ is a smooth map such that $P_{F_1}(x, y, z) \in \Lambda^3 T_x^* \hat{N}$. For $\alpha \in C_\lambda^\infty(\hat{U})$ with $\|\alpha\|_{C_1^1}$ sufficiently small, denoting $P_{F_1}(x, \alpha(x), \nabla\alpha(x))$ by $P_{F_1}(\alpha)(x)$, $P_{F_1}(\alpha) \in C_{2\lambda-2}^\infty(\Lambda^3 T^* \hat{N}) \subseteq C_{\lambda-1}^\infty(\Lambda^3 T^* \hat{N})$, as $\lambda > 1$. Moreover, for each $k \in \mathbb{N}$, if $\alpha \in C_\lambda^{k+1}(\hat{U})$ and $\|\alpha\|_{C_1^1}$ is sufficiently small, $P_{F_1}(\alpha) \in C_{2\lambda-2}^k(\Lambda^3 T^* \hat{N})$ and there exists a constant $c_k > 0$ such that

$$\|P_{F_1}(\alpha)\|_{C_{2\lambda-2}^k} \leq c_k \|\alpha\|_{C_\lambda^{k+1}}^2.$$

Proof: Firstly, by the definition of F_1 , $F_1(\alpha)(x)$ relates to the tangent space to Γ_α at $\pi_\alpha(x)$. Note that $T_{\pi_\alpha(x)}\Gamma_\alpha$ depends on both $\alpha(x)$ and $\nabla\alpha(x)$ and hence so must $F_1(\alpha)(x)$. We may then define P_{F_1} by (16) such that it is a smooth function of its arguments as claimed.

We argued above that we may identify $\Phi_i - \iota_i$ on $(0, \epsilon) \times \Sigma_i$ with

$$\beta_i \in C_\lambda^\infty((\Lambda_+^2)_{\mathcal{X}_i^*(g)} T^*((0, \epsilon) \times \Sigma_i))$$

for $i = 1, \dots, s$. Recall that

$$\Psi_i^* : \Lambda_+^2 T^*(U_i \setminus \{z_i\}) \rightarrow (\Lambda_+^2)_{\mathcal{X}_i^*(g)} T^*((0, \epsilon) \times \Sigma_i)$$

is a diffeomorphism. Let $k \in \mathbb{N}$, $\alpha \in C_\lambda^{k+1}(\hat{U})$, $\alpha_i = \alpha|_{U_i \setminus \{z_i\}}$ and $\gamma_i = \Psi_i^*(\alpha_i)$.

For each i , define a function $F_{C_i}(\gamma_i + \beta_i)$ on $(0, \epsilon) \times \Sigma_i$ by

$$F_{C_i}(\gamma_i + \beta_i)(r_i, \sigma_i) = F_1(\alpha_i)(\Psi_i(r_i, \sigma_i)). \quad (17)$$

Define a smooth function P_{C_i} by an equation analogous to (16):

$$\begin{aligned} F_{C_i}(\gamma_i + \beta_i)(r_i, \sigma_i) &= d(\gamma_i + \beta_i)(r_i, \sigma_i) \\ &\quad + P_{C_i}((r_i, \sigma_i), (\gamma_i + \beta_i)(r_i, \sigma_i), \nabla(\gamma_i + \beta_i)(r_i, \sigma_i)). \end{aligned} \quad (18)$$

We notice that F_{C_i} and P_{C_i} are only dependent on the cone C_i and, rather trivially, on ϵ . Therefore, because of this fact and our choice of δ in Proposition 6.4, these functions have scale equivariance properties. We may therefore derive equations and inequalities on $\{\epsilon\} \times \Sigma_i$ and deduce the result on all of $(0, \epsilon) \times \Sigma_i$ by introducing an appropriate scaling factor of r .

Now, since $\alpha = 0$ corresponds to our coassociative 4-fold \hat{N} , $F_1(0) = 0$. So, by (17),

$$F_{C_i}(\beta_i) = d\beta_i + P_{C_i}(\beta_i) = 0, \quad (19)$$

adopting similar notation for $P_{C_i}(\beta_i)$ as for $P_{F_1}(\alpha_i)$. Using (16)-(19), we deduce that

$$\begin{aligned} P_{F_1}(\alpha_i) &= d\beta_i + P_{C_i}(\gamma_i + \beta_i) = d\beta_i + P_{C_i}(\gamma_i + \beta_i) - (d\beta_i + P_{C_i}(\beta_i)) \\ &= P_{C_i}(\gamma_i + \beta_i) - P_{C_i}(\beta_i). \end{aligned} \quad (20)$$

We then calculate

$$\begin{aligned} P_{C_i}(\gamma_i + \beta_i) - P_{C_i}(\beta_i) &= \int_0^1 \frac{d}{dt} P_{C_i}(t\gamma_i + \beta_i) dt \\ &= \int_0^1 \gamma_i \cdot \frac{\partial P_{C_i}}{\partial y}(t\gamma_i + \beta_i) + \nabla \gamma_i \cdot \frac{\partial P_{C_i}}{\partial z}(t\gamma_i + \beta_i) dt, \end{aligned} \quad (21)$$

recalling that P_{C_i} is a function of three variables x , y and z . Using Taylor's Theorem,

$$P_{C_i}(\gamma_i + \beta_i) = P_{C_i}(\beta_i) + \gamma_i \cdot \frac{\partial P_{C_i}}{\partial y}(\beta_i) + \nabla \gamma_i \cdot \frac{\partial P_{C_i}}{\partial z}(\beta_i) + O(r^{-2}|\gamma_i|^2 + |\nabla \gamma_i|^2) \quad (22)$$

when $|\gamma_i|$ and $|\nabla\gamma_i|$ are small. Since $dF_1|_0(\alpha_i) = d\alpha_i$, $dF_{C_i}|_{\beta_i}(\gamma_i + \beta_i) = d\gamma_i$ and hence $dP_{C_i}|_{\beta_i} = 0$. Thus, the first derivatives of P_{C_i} with respect to y and z must vanish at β_i by (22). Therefore, given small $\nu > 0$ there exists a constant $A_0 > 0$ such that

$$\begin{aligned} \left| \frac{\partial P_{C_i}}{\partial y}(t\gamma_i + \beta_i) \right| &\leq A_0(r^{-2}|\gamma_i| + r^{-1}|\nabla\gamma_i|); \text{ and} \\ \left| \frac{\partial P_{C_i}}{\partial z}(t\gamma_i + \beta_i) \right| &\leq A_0(r^{-1}|\gamma_i| + |\nabla\gamma_i|) \end{aligned} \quad (23)$$

for $t \in [0, 1]$ whenever

$$r^{-1}|\gamma_i|, r^{-1}|\beta_i|, |\nabla\gamma_i| \text{ and } |\nabla\beta_i| \leq \nu. \quad (24)$$

By (15), $r^{-1}|\beta_i|$ and $|\nabla\beta_i|$ tend to zero as $r \rightarrow 0$. We can thus ensure that (24) is satisfied by the β_i components by making ϵ smaller. Hence, (24) holds if $\|\gamma_i\|_{C_1^1} \leq \nu$. Therefore, putting estimates (23) in (21) and using (20),

$$|P_{F_1}(\alpha_i)| = |P_{C_i}(\gamma_i + \beta_i) - P_{C_i}(\beta_i)| \leq A_0(r^{-1}|\gamma_i| + |\nabla\gamma_i|)^2 \quad (25)$$

whenever $\|\gamma_i\|_{C_1^1} \leq \nu$. As $r \rightarrow 0$ the terms in the bracket on the right-hand side of (25) are of order $O(r^{\lambda-1})$ by (15). Thus, $|P_{F_1}(\alpha_i)|$ is of order $O(r^{2\lambda-2})$, hence $O(r^{\lambda-1})$ since $\lambda > 1$, as $r \rightarrow 0$ for $i = 1, \dots, s$. We deduce that $|P_{F_1}(\alpha)|$ is of order $O(\rho^{2\lambda-2})$ as $\rho \rightarrow 0$ for all $\alpha \in C_\lambda^1(\hat{U})$ with $\|\alpha\|_{C_1^1}$ sufficiently small.

Similar calculations give analogous results to (25) for derivatives of P_{F_1} , but we shall explain the method by considering the first derivative. From (21) we calculate

$$\begin{aligned} &\nabla(P_{C_i}(\gamma_i + \beta_i) - P_{C_i}(\beta_i)) \\ &= \int_0^1 \nabla \left(\gamma_i \cdot \frac{\partial P_{C_i}}{\partial y}(t\gamma_i + \beta_i) + \nabla\gamma_i \cdot \frac{\partial P_{C_i}}{\partial z}(t\gamma_i + \beta_i) \right) dt \\ &= \int_0^1 \nabla\gamma_i \cdot \frac{\partial P_{C_i}}{\partial y} + \gamma_i \cdot \left(\nabla(t\gamma_i + \beta_i) \cdot \frac{\partial^2 P_{C_i}}{\partial y^2} + \nabla^2(t\gamma_i + \beta_i) \cdot \frac{\partial^2 P_{C_i}}{\partial y \partial z} \right) \\ &\quad + \nabla^2\gamma_i \cdot \frac{\partial P_{C_i}}{\partial z} + \nabla\gamma_i \cdot \left(\nabla(t\gamma_i + \beta_i) \cdot \frac{\partial^2 P_{C_i}}{\partial z \partial y} + \nabla^2(t\gamma_i + \beta_i) \cdot \frac{\partial^2 P_{C_i}}{\partial z^2} \right) dt. \end{aligned}$$

Whenever $\|\gamma_i\|_{C_1^1} \leq \nu$ there exists a constant $A_1 > 0$ such that (23) holds with A_0 replaced by A_1 and, for $t \in [0, 1]$,

$$\left| \frac{\partial^2 P_{C_i}}{\partial y^2}(t\gamma_i + \beta_i) \right|, \left| \frac{\partial^2 P_{C_i}}{\partial y \partial z}(t\gamma_i + \beta_i) \right| \text{ and } \left| \frac{\partial^2 P_{C_i}}{\partial z^2}(t\gamma_i + \beta_i) \right| \leq A_1,$$

since the second derivatives of P_{C_i} are continuous functions defined on the closed bounded set given by $\|\gamma_i\|_{C_1^1} \leq \nu$. We deduce that

$$|\nabla(P_{F_1}(\alpha_i))| = |\nabla(P_{C_i}(\gamma_i + \beta_i) - P_{C_i}(\beta_i))| \leq A_1 \left(\sum_{j=0}^2 r^{j-2} |\nabla^j \gamma_i| \right)^2$$

whenever $\|\gamma_i\|_{C_1^1} \leq \nu$. Therefore $|\nabla(P_{F_1}(\alpha_i))|$ is of order $O(r^{2\lambda-3})$, hence $O(r^{\lambda-2})$, as $r \rightarrow 0$.

In general we have the estimate

$$|\nabla^l(P_{F_1}(\alpha_i))| \leq A_l \left(\sum_{j=0}^{l+1} r^{j-(l+1)} |\nabla^j \gamma_i| \right)^2$$

for some $A_l > 0$ whenever $\|\gamma_i\|_{C_1^1} \leq \nu$. The result follows. \square

We now consider the regularity of solutions to the *nonlinear* elliptic equation $G_1(\alpha, \beta) = 0$ near $(0, 0)$.

Proposition 6.10 *Let $(\alpha, \beta) \in L_{k+1, \lambda}^p(\hat{U}) \times L_{k+1, \lambda}^p(\Lambda^4 T^* \hat{N})$ for some $p > 4$ and $k \geq 2$. If $G_1(\alpha, \beta) = 0$ and $\|\alpha\|_{C_1^1}$ is sufficiently small, $(\alpha, \beta) \in C_\lambda^\infty(\hat{U}) \times C_\lambda^\infty(\Lambda^4 T^* \hat{N})$.*

Proof: Suppose that $(\alpha, \beta) \in L_{k+1, \lambda}^p(\hat{U}) \times L_{k+1, \lambda}^p(\Lambda^4 T^* \hat{N})$ for some $p > 4$ and $k \geq 2$. Then α and β lie in C_{loc}^1 by Theorem 4.4, since $\frac{k}{4} > \frac{1}{p}$.

Suppose further that $G_1(\alpha, \beta) = 0$ and that $\|\alpha\|_{C_1^1}$ is sufficiently small. Since F_1 smoothly depends on α and $\nabla\alpha$, G_1 is a smooth function of $\alpha, \beta, \nabla\alpha$ and $\nabla\beta$. We apply [18, Theorem 6.8.1], which is a general regularity result for nonlinear elliptic equations, to conclude that α and β are smooth. However, we want more than this: the derivatives of α and β must decay at the required rates.

Recall the note after Definition 6.7 that $G_1(\alpha, \beta) = 0$ implies that $\beta = 0$. Thus $\beta \in C_\lambda^\infty(\Lambda^4 T^* \hat{N})$ trivially.

For the following argument we find it useful to work with weighted Hölder spaces. By Theorem 4.4, $\alpha \in C_\lambda^{k, a}(\hat{U})$ with $a = 1 - 4/p \in (0, 1)$ since $p > 4$. We also know that $d^*(G_1(\alpha, \beta)) = d^*(F_1(\alpha)) = 0$, which is a nonlinear elliptic equation on α . Using the notation and results of Proposition 6.9, $d^*d\alpha + d^*(P_{F_1}(\alpha)) = 0$ and $d^*(P_{F_1}(\alpha)) \in C_{2\lambda-3}^{k-2, a}(\Lambda^2 T^* \hat{N})$. We see that

$$d^*(F_1(\alpha))(x) = R(x, \alpha(x), \nabla\alpha(x))\nabla^2\alpha(x) + E(x, \alpha(x), \nabla\alpha(x)),$$

where $R(x, \alpha(x), \nabla\alpha(x))$ and $E(x, \alpha(x), \nabla\alpha(x))$ are smooth functions of their arguments. Define

$$S_\alpha(\gamma)(x) = R(x, \alpha(x), \nabla\alpha(x))\nabla^2\gamma(x)$$

for $\gamma \in C_{\text{loc}}^2(\Lambda_+^2 T^* \hat{N})$. Then S_α is a smooth, *linear, elliptic*, second-order operator, if $\|\alpha\|_{C_1^1}$ is sufficiently small, whose coefficients depend on x , $\alpha(x)$ and $\nabla\alpha(x)$. These coefficients therefore lie in $C_{\text{loc}}^{k-1, a}$. We also notice that

$$S_\alpha(\alpha)(x) = -E(x, \alpha(x), \nabla\alpha(x)) \in C_{2\lambda-3}^{k-2, a}(\Lambda^2 T^* \hat{N}) \subseteq C_{\lambda-2}^{k-2, a}(\Lambda^2 T^* \hat{N}),$$

since $\lambda > 1$. However, $E(x, \alpha(x), \nabla\alpha(x))$ only depends on α and $\nabla\alpha$, and is at worst quadratic in these quantities by Proposition 6.9, so it must in fact lie in $C_{\lambda-2}^{k-1, a}(\Lambda^2 T^* \hat{N})$ since we are given control on the decay of the first k derivatives of α as $\rho \rightarrow 0$.

The work in [17, §6.1.1] on asymptotically conical manifolds gives regularity results for smooth linear elliptic operators acting between weighted Hölder spaces. These results can easily be adapted to the CS scenario. In particular, if $\gamma \in C_\lambda^2(\Lambda_+^2 T^* N)$ and $S_\alpha(\gamma) \in C_{\lambda-2}^{k-1, a}(\Lambda^2 T^* N)$, we have that $\gamma \in C_\lambda^{k+1, a}(\Lambda_+^2 T^* N)$. Since $k \geq 2$, α and $S_\alpha(\alpha)$ satisfy these conditions by the discussion above. We deduce that $\alpha \in C_\lambda^{k+1, a}(\Lambda_+^2 T^* N)$ only knowing a priori that $\alpha \in C_\lambda^{k, a}(\Lambda_+^2 T^* N)$. We proceed by induction to show that $\alpha \in C_\lambda^{l, a}(\Lambda_+^2 T^* N)$ for all $l \geq 2$. \square

6.2 Problem 2: moving singularities and fixed G_2 structure

For this problem we again consider deformations of N in (M, φ, g) which are CS coassociative 4-folds at s points with the same rate and cones at the singularities, but now we allow the singular points and tangent cones at those points to differ from those of N . However, we still assume that the G_2 structure on M is fixed.

Definition 6.11 The *moduli space of deformations* $\mathcal{M}_2(N, \lambda)$ for Problem 2 is the set of N' in (M, φ, g) which are CS coassociative 4-folds at z'_1, \dots, z'_s with rate λ , having cone C_i and tangent cone \hat{C}'_i at z'_i for all i , such that there exists a homeomorphism $h : N \rightarrow N'$, isotopic to the identity, with $h(z_i) = z'_i$ for $i = 1, \dots, s$ and such that $h|_{\hat{N}} : \hat{N} \rightarrow N' \setminus \{z'_1, \dots, z'_s\}$ is a diffeomorphism.

Here it is more difficult to create a local description of the moduli space which is compatible with the analytic framework in which our study is made. What one would consider more ‘intuitive’ approaches do not, as far as the author is aware, bear fruit. We therefore follow what is, at first sight, a slightly indirect route.

For each $i = 1, \dots, s$ let B_i be an open set in M containing z_i such that $B_i \cap B_j = \emptyset$ for $i \neq j$. Let $B = \prod_{i=1}^s B_i$. For each $\mathbf{z}' = (z'_1, \dots, z'_s) \in B$,

we have a family $I(\mathbf{z}')$ of choices of s -tuples $\zeta' = (\zeta'_1, \dots, \zeta'_s)$ of isomorphisms $\zeta'_i : \mathbb{R}^7 \rightarrow T_{z'_i}M$ identifying (φ_0, g_0) with $(\varphi|_{T_{z'_i}M}, g|_{T_{z'_i}M})$. Clearly, for each $\mathbf{z}' \in B$, $I(\mathbf{z}') \cong G_2$. We thus make the following definition.

Definition 6.12 The *translation space* is

$$\mathcal{T} = \{(\mathbf{z}', \zeta') : \mathbf{z}' \in B, \zeta' \in I(\mathbf{z}')\}.$$

It is a principal G_2^s bundle over B and hence is a smooth manifold.

Let H_i denote the Lie subgroup of G_2 preserving $\iota_i(C_i)$ in \mathbb{R}^7 for $i = 1, \dots, s$ and let $H = \prod_{i=1}^s H_i \subseteq G_2^s$. Then H acts freely on \mathcal{T} by

$$(\mathbf{z}', \zeta') \mapsto (\mathbf{z}', (\zeta'_1 \circ A_1^{-1}, \dots, \zeta'_s \circ A_s^{-1})),$$

where $(A_1, \dots, A_s) \in H$. Thus there exists an H -orbit through (\mathbf{z}, ζ) in \mathcal{T} , where

$$\mathbf{z} = (z_1, \dots, z_s) \quad \text{and} \quad \zeta = (\zeta_1, \dots, \zeta_s).$$

Define $\hat{\mathcal{T}}$ to be a small open ball in \mathbb{R}^n containing 0, where $n = \dim \mathcal{T} - \dim H$, and let $h_{\hat{\mathcal{T}}} : \hat{\mathcal{T}} \rightarrow \mathcal{T}$ be an embedding with $h_{\hat{\mathcal{T}}}(0) = (\mathbf{z}, \zeta)$ such that $h_{\hat{\mathcal{T}}}(\hat{\mathcal{T}})$ is transverse to the H -orbit through (\mathbf{z}, ζ) . Write $h_{\hat{\mathcal{T}}}(t) = (\mathbf{z}(t), \zeta(t))$ for $t \in \hat{\mathcal{T}}$, with $\mathbf{z}(0) = \mathbf{z}$ and $\zeta(0) = \zeta$.

Notes

- (a) If $t, t' \in \hat{\mathcal{T}}$, with $t \neq t'$, are such that $\mathbf{z}(t) = \mathbf{z}(t')$, the s -tuples of tangent cones, $\{\hat{C}_1(t), \dots, \hat{C}_s(t)\}$ and $\{\hat{C}_1(t'), \dots, \hat{C}_s(t')\}$, are distinct.
- (b) $\hat{\mathcal{T}}$ is an open ball in $\mathbb{R}^n \cong T_0\hat{\mathcal{T}}$ and hence can be considered as an open subset of $T_0\hat{\mathcal{T}}$.

We use $\hat{\mathcal{T}}$ to extend N to a family of nearby CS 4-folds and provide an analogue to Proposition 6.4 for Problem 2. In defining N we chose a G_2 coordinate system $\{\chi_i : B(0; \eta) \rightarrow V_i : i = 1, \dots, s\}$ with $d\chi_i|_0 = \zeta_i$ for $i = 1, \dots, s$. Extend this to a smooth family of G_2 coordinate systems

$$\left\{ \{\chi_i(t) : B(0; \eta) \rightarrow V_i(t) : i = 1, \dots, s\} : t \in \hat{\mathcal{T}} \right\},$$

where $V_i(t)$ is an open set in M containing $z_i(t)$, $\chi_i(t)(0) = z_i(t)$, $d\chi_i(t)|_0 = \zeta_i(t)$, $\chi_i(0) = \chi_i$ and $V_i(0) = V_i$ for $i = 1, \dots, s$.

Proposition 6.13 *Use the notation of Proposition 6.4 and Definition 6.12.*

(a) There exists a family $\mathcal{N} = \{N(t) : t \in \hat{\mathcal{T}}\}$ of CS 4-folds in M , with $N(0) = N$, such that $N(t)$ has singularities at $z_1(t), \dots, z_s(t)$ with rate λ , cones C_1, \dots, C_s and tangent cones $\hat{C}_1(t), \dots, \hat{C}_s(t)$ defined by $\hat{C}_i(t) = (\zeta_i(t) \circ \iota_i)(C_i)$.

(b) Let $\hat{N}(t) = N(t) \setminus \{z_1(t), \dots, z_s(t)\}$ and write

$$N(t) = K(t) \sqcup \bigsqcup_{i=1}^s U_i(t)$$

where $K(t)$ is compact and $U_i(t) \setminus \{z_i(t)\} \cong (0, \epsilon) \times \Sigma_i$ for all i , in the obvious way, ensuring that $K(0) = K$ and $U_i(0) = U_i$. For $t \in \hat{\mathcal{T}}$, there exist open sets $\hat{T}(t) \subseteq M$ containing $\hat{N}(t)$ and diffeomorphisms $\delta(t) : \hat{U} \rightarrow \hat{T}(t)$ taking the zero section to $\hat{N}(t)$, varying smoothly in t , with $\hat{T}(0) = \hat{T}$ and $\delta(0) = \delta$. Moreover, $\hat{T}(t)$ can be chosen to grow with order $O(\rho)$ as $\rho \rightarrow 0$ and $\delta(t)$ is compatible with the identifications $U_i(t) \setminus \{z_i(t)\} \cong (0, \epsilon) \times \Sigma_i$ for all i .

Remark The family \mathcal{N} does *not* necessarily consist of CS *coassociative* 4-folds and $\delta(t)$ is not required to be compatible with the isomorphism $\nu(\hat{N}) \cong \Lambda_+^2 T^* \hat{N}$ for $t \neq 0$.

Proof. Use the notation from the proof of Proposition 6.4. For $t \in \hat{\mathcal{T}}$, define $\hat{T}_i(t) = \chi_i(t)(\hat{S}_i)$ and

$$U_i(t) = \left(\chi_i(t) \circ \Phi_i((0, \epsilon) \times \Sigma_i) \right) \cup \{z_i(t)\}$$

for $i = 1, \dots, s$. Then $\hat{T}_i(t)$ contains $U_i(t) \setminus \{z_i(t)\}$. Define a diffeomorphism $\delta_i(t)$ such that the following diagram commutes:

$$\begin{array}{ccc} \hat{U}_i & \xrightarrow{\Psi_i^*} & j_i(\hat{V}_i) \\ \delta_i(t) \downarrow & & \downarrow j_i^{-1} \\ & & \hat{V}_i \\ & & \downarrow n_i \\ \hat{T}_i(t) & \xleftarrow{\chi_i(t)} & \hat{S}_i. \end{array} \quad (26)$$

We then interpolate smoothly over K to extend $\bigcup_{i=1}^s \hat{T}_i(t)$ to $\hat{T}(t)$ and $\delta_i(t)$ to $\delta(t)$ as required. Note by construction that $\hat{T}(t)$ grows with order $O(\rho)$ as $\rho \rightarrow 0$.

Let $e(t) = \delta(t)|_{\hat{N}}$ and define $\hat{N}(t) = e(t)(\hat{N})$. Then $e(t) : \hat{N} \rightarrow \hat{N}(t)$ is a diffeomorphism for all $t \in \hat{\mathcal{T}}$ and $e(0)$ is the identity. Let $N(t) = \hat{N}(t) \cup \{z_1(t), \dots, z_s(t)\}$. We then have a family $\mathcal{N} = \{N(t) : t \in \hat{\mathcal{T}}\}$ as claimed. Note that $K(t) = e(t)(K)$.

By the construction of $\delta(t)$ and the family \mathcal{N} , it is clear that the proposition is proved, where the compatibility conditions on $\delta(t)$ are given by (26). \square

The next definition is analogous to Definition 6.5.

Definition 6.14 Use the notation of Proposition 6.13. Let Γ_α be the graph of $\alpha \in C_{\text{loc}}^1(\hat{U})$ and let $\pi_\alpha : \hat{N} \rightarrow \Gamma_\alpha$ be given by $\pi_\alpha(x) = (x, \alpha(x))$. For $t \in \hat{\mathcal{T}}$, let $f_\alpha(t) = \delta(t) \circ \pi_\alpha$ and let $\hat{N}_\alpha(t) = f_\alpha(t)(\hat{N})$. Define F_2 from $C_{\text{loc}}^1(\hat{U}) \times \hat{\mathcal{T}}$ to $C_{\text{loc}}^0(\Lambda^3 T^* \hat{N})$ by:

$$F_2(\alpha, t) = f_\alpha(t)^* \left(\varphi|_{\hat{N}_\alpha(t)} \right).$$

The linearisation of F_2 at $(0, 0)$ acts as

$$dF_2|_{(0,0)} : (\alpha, t) \longmapsto d\alpha + L_2(t),$$

where $\alpha \in C_{\text{loc}}^1(\Lambda_+^2 T^* \hat{N})$, $t \in T_0 \hat{\mathcal{T}}$ and L_2 is a linear map into the space of smooth exact 3-forms on \hat{N} since φ is exact near \hat{N} .

Remark By construction $F_2(\alpha, 0) = F_1(\alpha)$ as given in Definition 6.5.

Clearly, $\text{Ker } F_2$ is the set of $\alpha \in C_{\text{loc}}^1(\hat{U})$ and $t \in \hat{\mathcal{T}}$ such that $\hat{N}_\alpha(t)$ is coassociative. However, we have not yet encoded the information that $N_\alpha(t)$ is CS with *rate* λ . This is the subject of the next proposition.

Proposition 6.15 *The moduli space of deformations for Problem 2 is locally homeomorphic to $\text{Ker } F_2 = \{(\alpha, t) \in C_\lambda^\infty(\hat{U}) \times \hat{\mathcal{T}} : F_2(\alpha, t) = 0\}$.*

Proof: For each $t \in \hat{\mathcal{T}}$, we are in the situation of Problem 1 in the sense that we want coassociative deformations $\hat{N}_\alpha(t)$ of $\hat{N}(t)$, defined by a self-dual 2-form α , which have the same singular points, cones and tangent cones as $\hat{N}(t)$. It is thus clear that $\alpha \in C_\lambda^\infty(\hat{U})$ by Proposition 6.6. \square

We now introduce an associated map G_2 to F_2 .

Definition 6.16 Define $G_2 : C_{\text{loc}}^1(\hat{U}) \times C_{\text{loc}}^1(\Lambda^4 T^* \hat{N}) \times \hat{\mathcal{T}} \rightarrow C_{\text{loc}}^0(\Lambda^3 T^* \hat{N})$ by:

$$G_2(\alpha, \beta, t) = F_2(\alpha, t) + d^* \beta.$$

Then $dG_2|_{(0,0,0)} : (\alpha, \beta, t) \longmapsto d\alpha + d^* \beta + L_2(t)$, in the notation of Definition 6.14.

We then have an analogous result to Proposition 6.8, which follows in exactly the same fashion because $F_2(\alpha, t)$ is exact.

Proposition 6.17

$$\text{Ker } F_2 \cong \{(\alpha, \beta, t) \in C_\lambda^\infty(\hat{U}) \times C_\lambda^\infty(\Lambda^4 T^* \hat{N}) \times \hat{\mathcal{T}} : G_2(\alpha, \beta, t) = 0\}.$$

The next result studies the regularity of the kernel of G_2 near $(0, 0, 0)$ and is the analogue of Proposition 6.10.

Proposition 6.18 *Let $(\alpha, \beta, t) \in L_{k+1, \lambda}^p(\hat{U}) \times L_{k+1, \lambda}^p(\Lambda^4 T^* \hat{N}) \times \hat{\mathcal{T}}$, where $p > 4$ and $k \geq 2$. If $G_2(\alpha, \beta, t) = 0$ and $\|\alpha\|_{C_1^1}$ and t are sufficiently small, $(\alpha, \beta) \in C_\lambda^\infty(\hat{U}) \times C_\lambda^\infty(\Lambda^4 T^* \hat{N})$.*

Proof: Note that $dG_2(\alpha, \beta, t) = \Delta\beta = 0$ implies that $\beta = 0$ by the Maximum Principle for harmonic functions and $d^*G_2(\alpha, \beta, t) = d^*F_2(\alpha, t) = 0$ is an elliptic equation at 0 on α . Using similar notation to the proof of Proposition 6.10,

$$d^*F_2(\alpha, t)(x) = R_t(x, \alpha(x), \nabla\alpha(x))\nabla^2\alpha(x) + E_t(x, \alpha(x), \nabla\alpha(x)),$$

where R_t and E_t are smooth functions of their arguments. If we define

$$S_{(\alpha, t)}(\gamma)(x) = R_t(x, \alpha(x), \nabla\alpha(x))\nabla^2\gamma(x),$$

then $S_{(\alpha, t)}$ is a smooth linear differential operator on $\gamma \in C_{\text{loc}}^2(\Lambda_+^2 T^* \hat{N})$. The ellipticity of $S_\alpha = S_{(\alpha, 0)}$ results from the coassociativity of \hat{N} . Ellipticity is an *open* condition so, although $\hat{N}(t)$ is not necessarily coassociative, the fact that it is ‘close’ to being coassociative means that $S_{(\alpha, t)}$ is elliptic, as long as we shrink $\hat{\mathcal{T}}$ as necessary to make t sufficiently small.

The regularity results for $S_{(\alpha, t)}$ follow in the same way as in the proof of Proposition 6.10 since $F_2(\alpha, t)$ depends smoothly on t and $\hat{N}(t)$ is asymptotically coassociative near the singular points, which validates the use of the theory from [17, §6.1.1]. Recall that $L_{k+1, \lambda}^p \hookrightarrow C_\lambda^{k, a}$ where $a = 1 - 4/p$. Thus, if $S_{(\alpha, t)}(\gamma) \in C_{\lambda-2}^{k-1, a}$ and $\gamma \in C_\lambda^2(\hat{U})$, then $\gamma \in C_\lambda^{k+1, a}(\hat{U})$.

Since $E_0 = E$ maps into $C_{\lambda-2}^{k-1, a}$, as argued in the proof of Proposition 6.10, and F_2 depends smoothly on t , E_t maps into $C_{\lambda-2}^{k-1, a}$ for t sufficiently small. Hence,

$$S_{(\alpha, t)}(\alpha)(x) = -E_t(x, \alpha(x), \nabla\alpha(x)) \in C_{\lambda-2}^{k-1, a}.$$

We deduce that $\alpha \in C_\lambda^{k+1, a}$, given only that $\alpha \in C_\lambda^{k, a}$. Induction gives the result. \square

6.3 Problem 3: moving singularities and varying G_2 structure

For our final problem we consider CS deformations N' of N with the same rate and cones at s singularities, but with possibly different singular points and tangent cones there, such that N' is coassociative under a deformation of the G_2 structure on M .

We begin with the following.

Proposition 6.19 *Use the notation of Proposition 6.4. Let*

$$T = \hat{T} \cup \bigcup_{i=1}^s V_i \supseteq N.$$

By making \hat{T} and V_i , for $i = 1, \dots, s$, smaller if necessary, T retracts onto N . There exists an isomorphism $\Xi : H_{\text{dR}}^3(T) \rightarrow H_{\text{cs}}^3(\hat{N})$.

Proof: Let $[\xi] \in H_{\text{dR}}^3(T)$. Since the sets V_i retract onto $\{z_i\}$ for $i = 1, \dots, s$, ξ can be chosen such that $\xi|_{V_i} = 0$. Therefore, $\xi|_{U_i \setminus \{z_i\}} = 0$ which implies that the support of $\xi|_{\hat{N}}$ is contained in K , which is compact. Hence $[\xi|_{\hat{N}}]$ is a well-defined element of $H_{\text{cs}}^3(\hat{N})$. Define Ξ by $[\xi] \mapsto [\xi|_{\hat{N}}]$. We show that Ξ is well-defined. Suppose that $\xi' = \xi + dv$, for $v \in C^\infty(\Lambda^2 T^*T)$, such that $\xi'|_{V_i} = 0$ for all i . Then $dv|_{V_i} = 0$ for all i . Since V_i retracts onto $\{z_i\}$ we can choose v such that $v|_{V_i} = 0$ without affecting dv by smoothly interpolating over \hat{T} . Thus $v|_{\hat{N}}$ is compactly supported on \hat{N} and $\xi|_{\hat{N}} + d(v|_{\hat{N}}) = \xi'|_{\hat{N}}$. Hence Ξ is well-defined and injective.

Any closed form on \hat{N} with support in K can be extended smoothly to a closed form on T which vanishes on V_i for all i . Thus, any cohomology class in $H_{\text{cs}}^3(\hat{N})$ has a representative γ that can be lifted to a form ξ on T such that $\Xi([\xi]) = [\gamma]$, which implies that Ξ is surjective. \square

Notes The reason for this result is two-fold.

- (a) The condition that $\Xi([\varphi|_T]) = 0$ in $H_{\text{cs}}^3(\hat{N})$ is implied by the coassociativity of \hat{N} and it forces $[\varphi|_{\hat{N}}] = 0$ in $H_{\text{cs}}^3(\hat{N})$. This is stronger than the seemingly more natural condition of $[\varphi|_{\hat{N}}] = 0$ in $H_{\text{dR}}^3(\hat{N})$, which would be the correct requirement if \hat{N} were compact by the work of McLean [16].
- (b) If a G_2 structure (φ', g') on M is such that $\Xi([\varphi'|_T]) \neq 0$ then $\varphi'|_{\hat{N}'} \neq 0$ for any nearby deformation \hat{N}' of \hat{N} , so there are *no* coassociative deformations.

Proposition 6.19 allows us to define a distinguished family of ‘nearby’ G_2 structures to (φ, g) .

Definition 6.20 Let $\hat{\mathcal{F}}$ be a small open ball about 0 in \mathbb{R}^m for some m . Let

$$\mathcal{F} = \{(\varphi^f, g^f) : f \in \hat{\mathcal{F}}\}$$

be a family of torsion-free G_2 structures, with $(\varphi^0, g^0) = (\varphi, g)$, such that $\Xi([\varphi^f|_T]) = 0$ in $H_{cs}^3(\hat{N})$ and the map $h_{\hat{\mathcal{F}}} : \hat{\mathcal{F}} \rightarrow \mathcal{F}$ given by $h_{\hat{\mathcal{F}}}(f) = (\varphi^f, g^f)$ is an embedding.

Note $\hat{\mathcal{F}}$ can be considered as an open subset of $T_0\hat{\mathcal{F}}$.

We now describe the moduli space for Problem 3.

Definition 6.21 The *moduli space of deformations* $\mathcal{M}_3(N, \lambda)$ for Problem 3 is the set of pairs (N', f) of $f \in \hat{\mathcal{F}}$ and N' in (M, φ^f, g^f) which are CS coassociative 4-folds at z'_1, \dots, z'_s with rate λ , having cone C_i and tangent cone \hat{C}'_i at z'_i for all i , such that there exists a homeomorphism $h : N \rightarrow N'$, isotopic to the identity, with $h(z_i) = z'_i$ for $i = 1, \dots, s$ and such that $h|_{\hat{N}} : \hat{N} \rightarrow N' \setminus \{z'_1, \dots, z'_s\}$ is a diffeomorphism.

We have a projection map $\pi_{\hat{\mathcal{F}}} : \mathcal{M}_3(N, \lambda) \rightarrow \hat{\mathcal{F}}$, with $\pi_{\hat{\mathcal{F}}}(N', f) = f$, whose fibres $\pi_{\hat{\mathcal{F}}}^{-1}(f)$ are equal to the moduli space for Problem 2 defined using the G_2 structure (φ^f, g^f) .

We must adapt our translation space from Problem 2 to incorporate the varying G_2 structure.

Definition 6.22 Use the notation of Definitions 6.12 and 6.20. For $f \in \hat{\mathcal{F}}$ and $\mathbf{z}' \in B$ let $I^f(\mathbf{z}')$ denote the family of choices of s -tuples $\zeta' = (\zeta'_1, \dots, \zeta'_s)$ of isomorphisms $\zeta'_i : \mathbb{R}^7 \rightarrow T_{z'_i}M$ identifying (φ_0, g_0) with $(\varphi^f|_{T_{z'_i}M}, g^f|_{T_{z'_i}M})$.

The *translation space* corresponding to $\hat{\mathcal{F}}$ is

$$\mathcal{T}^{\hat{\mathcal{F}}} = \{(\mathbf{z}', \zeta', f) : \mathbf{z}' \in B, f \in \hat{\mathcal{F}}, \zeta' \in I^f(\mathbf{z}')\}.$$

It is a principal G_2^s bundle over $B \times \hat{\mathcal{F}}$.

There is a natural free action of H on $\mathcal{T}^{\hat{\mathcal{F}}}$ and hence an H -orbit through $(\mathbf{z}, \zeta, 0)$. Therefore, we may embed $\hat{\mathcal{T}} \times \hat{\mathcal{F}}$ into $\mathcal{T}^{\hat{\mathcal{F}}}$ by $h_{\hat{\mathcal{T}} \times \hat{\mathcal{F}}} : (t, f) \mapsto (\mathbf{z}(t, f), \zeta(t, f), f)$ such that $h_{\hat{\mathcal{T}} \times \hat{\mathcal{F}}}(\hat{\mathcal{T}} \times \hat{\mathcal{F}})$ is transverse to this H -orbit, $h_{\hat{\mathcal{T}} \times \hat{\mathcal{F}}}(t, 0) = h_{\hat{\mathcal{T}}}(t)$ for all t and $\mathbf{z}(0, f) = \mathbf{z}$ for all f .

Use the notation introduced before Proposition 6.13. Extend the G_2 coordinate system near z_1, \dots, z_s used to define N to a smooth family of G_2 coordinate

systems

$$\left\{ \{ \chi_i(t, f) : B(0; \eta) \rightarrow V_i(t, f) : i = 1, \dots, s \} : (t, f) \in \hat{\mathcal{T}} \times \hat{\mathcal{F}} \right\}$$

such that $V_i(t, f)$ is an open set in M containing $z_i(t, f)$, $\chi_i(t, f)(0) = z_i(t, f)$, $d\chi_i(t, f)|_0 = \zeta_i(t, f)$, $\chi_i(t, 0) = \chi_i(t)$, $V_i(0, f) = V_i$ and $V_i(t, 0) = V_i(t)$ for $i = 1, \dots, s$. We state the analogue of Proposition 6.13.

Proposition 6.23 *Use the notation of Propositions 6.4 and 6.13 and Definition 6.22.*

- (a) *There exists a family $\mathcal{N}^{\hat{\mathcal{F}}} = \{N(t, f) : (t, f) \in \hat{\mathcal{T}} \times \hat{\mathcal{F}}\}$ of CS 4-folds in M , with $N(0, f) = N$ and $N(t, 0) = N(t)$, such that $N(t, f)$ has singularities at $z_1(t, f), \dots, z_s(t, f)$ with rate λ , cones C_1, \dots, C_s and tangent cones $\hat{C}_1(t, f), \dots, \hat{C}_s(t, f)$ defined by $\hat{C}_i(t, f) = (\zeta_i(t, f) \circ \iota_i)(C_i)$.*
- (b) *Let $\hat{N}(t, f) = N(t, f) \setminus \{z_1(t, f), \dots, z_s(t, f)\}$ and write*

$$N(t, f) = K(t, f) \sqcup \bigsqcup_{i=1}^s U_i(t, f)$$

where $K(t, f)$ is compact and $U_i(t, f) \setminus \{z_i(t, f)\} \cong (0, \epsilon) \times \Sigma_i$ for all i , in the obvious way, ensuring that $K(0, f) = K$, $K(t, 0) = K(t)$, $U_i(0, f) = U_i$ and $U_i(t, 0) = U_i(t)$. For $(t, f) \in \hat{\mathcal{T}} \times \hat{\mathcal{F}}$, there exist open sets $\hat{T}(t, f) \subseteq M$ containing $\hat{N}(t, f)$ and diffeomorphisms $\delta(t, f) : \hat{U} \rightarrow \hat{T}(t, f)$ taking the zero section to $\hat{N}(t, f)$, varying smoothly in t and f , with $\hat{T}(0, f) = \hat{T}$, $\hat{T}(t, 0) = \hat{T}(t)$ and $\delta(t, 0) = \delta(t)$. Moreover, $\hat{T}(t, f)$ can be chosen to grow with order $O(\rho)$ as $\rho \rightarrow 0$ and $\delta(t, f)$ is compatible with the identifications $U_i(t, f) \setminus \{z_i(t, f)\} \cong (0, \epsilon) \times \Sigma_i$ for $i = 1, \dots, s$.

The proof is almost identical to that of Proposition 6.13 and so we omit it. The compatibility conditions on $\delta(t, f)$ are given by similar commutative diagrams to (26).

Remark $\delta(t, f)$ is not required to be compatible with the isomorphism $\nu(\hat{N}) \cong \Lambda_+^2 T^* \hat{N}$ for $(t, f) \neq (0, 0)$.

We proceed by defining our final deformation map.

Definition 6.24 Use the notation of Proposition 6.23. Let Γ_α be the graph of $\alpha \in C_{\text{loc}}^1(\hat{U})$ and let $\pi_\alpha : \hat{N} \rightarrow \Gamma_\alpha$ be given by $\pi_\alpha(x) = (x, \alpha(x))$. For

$(t, f) \in \hat{\mathcal{T}} \times \hat{\mathcal{F}}$, let $f_\alpha(t, f) = \delta(t, f) \circ \pi_\alpha$ and let $\hat{N}_\alpha(t, f) = f_\alpha(t, f)(\hat{N})$. Define F_3 from $C_{\text{loc}}^1(\hat{U}) \times \hat{\mathcal{T}} \times \hat{\mathcal{F}}$ to $C_{\text{loc}}^0(\Lambda^3 T^* \hat{N})$ by:

$$F_3(\alpha, t, f) = f_\alpha(t, f)^* \left(\varphi^f|_{\hat{N}_\alpha(t, f)} \right).$$

The linearisation of F_3 at $(0, 0, 0)$ acts as

$$dF_3|_{(0,0,0)} : (\alpha, t, f) \longmapsto d\alpha + L_2(t) + L_3(f),$$

where $\alpha \in C_{\text{loc}}^1(\Lambda_+^2 T^* \hat{N})$, $(t, f) \in T_0 \hat{\mathcal{T}} \oplus T_0 \hat{\mathcal{F}}$, L_2 is given in Definition 6.14 and L_3 is a linear map into the space of smooth exact 3-forms on \hat{N} by the condition imposed on φ^f in Definition 6.20.

Note $F_3(\alpha, t, 0) = F_2(\alpha, t)$ as given in Definition 6.14.

Now, $\text{Ker } F_3$ corresponds to choices of $\hat{N}_\alpha(t, f)$ which are coassociative with respect to (φ^f, g^f) . The next result is then clear from considering the proof of Proposition 6.15.

Proposition 6.25 *The moduli space of deformations for Problem 3 is locally homeomorphic to $\text{Ker } F_3 = \{(\alpha, t, f) \in C_\lambda^\infty(\hat{U}) \times \hat{\mathcal{T}} \times \hat{\mathcal{F}} : F_3(\alpha, t, f) = 0\}$.*

We again have an associated map to our deformation map.

Definition 6.26 Define $G_3 : C_{\text{loc}}^1(\hat{U}) \times C_{\text{loc}}^1(\Lambda^4 T^* \hat{N}) \times \hat{\mathcal{T}} \times \hat{\mathcal{F}} \rightarrow C_{\text{loc}}^0(\Lambda^3 T^* \hat{N})$ by:

$$G_3(\alpha, \beta, t, f) = F_3(\alpha, t, f) + d^* \beta.$$

Then $dG_3|_{(0,0,0,0)} : (\alpha, \beta, t, f) \longmapsto d\alpha + d^* \beta + L_2(t) + L_3(f)$, in the notation of Definition 6.24.

The next result is analogous to Propositions 6.8 and 6.17 and may be immediately deduced from the exactness of $F_3(\alpha, t, f)$, which follows from the condition imposed on φ^f in Definition 6.20.

Proposition 6.27

$$\text{Ker } F_3 \cong \{(\alpha, \beta, t, f) \in C_\lambda^\infty(\hat{U}) \times C_\lambda^\infty(\Lambda^4 T^* \hat{N}) \times \hat{\mathcal{T}} \times \hat{\mathcal{F}} : G_3(\alpha, \beta, t, f) = 0\}.$$

The argument used to prove the regularity result Proposition 6.18 is easily generalised to the map G_3 , so we end the section with the following.

Proposition 6.28 *Let $(\alpha, \beta, t, f) \in L_{k+1, \lambda}^p(\hat{U}) \times L_{k+1, \lambda}^p(\Lambda^4 T^* \hat{N}) \times \hat{\mathcal{T}} \times \hat{\mathcal{F}}$, where $p > 4$ and $k \geq 2$. If $G_3(\alpha, \beta, t, f) = 0$ and $\|\alpha\|_{C_1^1}$, t and f are sufficiently small, $(\alpha, \beta) \in C_\lambda^\infty(\hat{U}) \times C_\lambda^\infty(\Lambda^4 T^* \hat{N})$.*

7 The deformation and obstruction spaces

In this section we describe the infinitesimal deformation and obstruction spaces for each of our problems and show in each scenario that, if the obstruction space is zero, we get a *smooth* moduli space of deformations. We recollect the common notation introduced at the start of §6. In addition, fix some $p > 4$ and integer $k \geq 2$.

7.1 Problem 1

Recall the maps F_1 and G_1 given in Definitions 6.5 and 6.7 respectively. Their kernels give a local description for the moduli space $\mathcal{M}_1(N, \lambda)$ by Propositions 6.6 and 6.8. Therefore the kernels of $dF_1|_0$ and $dG_1|_{(0,0)}$ describe the *infinitesimal deformations*.

Definition 7.1 The *infinitesimal deformation space* for Problem 1 is

$$\begin{aligned} \mathcal{I}_1(N, \lambda) &= \{\alpha \in C_\lambda^\infty(\Lambda_+^2 T^* \hat{N}) : d\alpha = 0\} \\ &\cong \{(\alpha, \beta) \in C_\lambda^\infty(\Lambda_+^2 T^* \hat{N} \oplus \Lambda^4 T^* \hat{N}) : d\alpha + d^* \beta = 0\}. \end{aligned}$$

The equivalence of the spaces follows by Proposition 6.8 or, more simply, by the Maximum Principle for harmonic functions.

Using Proposition 6.10,

$$\mathcal{I}_1(N, \lambda) \cong \{(\alpha, \beta) \in L_{k+1, \lambda}^p(\Lambda_+^2 T^* \hat{N} \oplus \Lambda^4 T^* \hat{N}) : d\alpha + d^* \beta = 0\}.$$

Therefore, $\mathcal{I}_1(N, \lambda)$ is finite-dimensional.

We turn to possible *obstructions* to the deformation theory and start with the following.

Proposition 7.2 The map F_1 takes $L_{k+1, \lambda}^p(\hat{U})$ into $d(L_{k+1, \lambda}^p(\Lambda^2 T^* \hat{N}))$.

Proof: Let $\alpha \in L_{k+1, \lambda}^p(\hat{U})$ and let T be as in Proposition 6.19. As noted after that proposition, $[\varphi|_T] = 0$ in $H_{\text{dR}}^3(T)$ and hence $\varphi|_T$ is exact. Thus, $\varphi|_T = d\psi$ for some $\psi \in C^\infty(\Lambda^2 T^* T)$. However, we want to select ψ in a particular way near the singularities. On $B(0; \eta) \subseteq \mathbb{R}^7$, for each $i = 1, \dots, s$,

$$\chi_i^*(\varphi) = \varphi_0 + O(r_i).$$

If v is the dilation vector field on \mathbb{R}^7 , given in coordinates (x_1, \dots, x_7) by

$$v = x_1 \frac{\partial}{\partial x_1} + \dots + x_7 \frac{\partial}{\partial x_7},$$

we can choose ψ to satisfy

$$\chi_i^*(\psi) = \frac{1}{3}(v \cdot \varphi_0) + O(r_i^2)$$

on V_i , since $d(v \cdot \varphi_0) = 3\varphi_0$, then extend ψ smoothly to a form on T such that $d\psi = \varphi|_T$. Note that

$$(v \cdot \varphi_0)|_{\iota_i(C_i)} = v \cdot (\varphi_0|_{\iota_i(C_i)}) = 0$$

as $v \in T(\iota_i(C_i))$. Hence $\chi_i^*(\psi) = O(r_i^2)$ on $\iota_i(C_i)$, for all i , and similar results hold for the derivatives of ψ . Define

$$H_1(\alpha) = f_\alpha^* \left(\psi|_{\hat{N}_\alpha} \right)$$

so that $F_1(\alpha) = d(H_1(\alpha))$. Note that $\chi_i^*(\psi)|_{\iota_i(C_i)} = O(r_i^2)$ is dominated by $O(r_i^\lambda)$ terms as $r_i \rightarrow 0$ since $\lambda < 2$. Further, $f_\alpha^*(\psi|_{\hat{N}_\alpha})$ has the same growth as $\chi_i^*(\psi)|_{(\Phi_\alpha)_i((0,\epsilon) \times \Sigma_i)}$ as $r_i \rightarrow 0$, using the notation preceding Proposition 6.6. However,

$$\chi_i^*(\psi)|_{(\Phi_\alpha)_i((0,\epsilon) \times \Sigma_i)} = \chi_i^*(\psi)|_{((\Phi_\alpha)_{i-\iota_i})((0,\epsilon) \times \Sigma_i)} + \chi_i^*(\psi)|_{\iota_i((0,\epsilon) \times \Sigma_i)}.$$

The first term on the right-hand side depends on $|(\Phi_\alpha)_i - \iota_i|$ and hence is $O(r_i^\lambda)$ as $r_i \rightarrow 0$. This dominates the second term by our observation above. Hence, $H_1(\alpha) \in L_{k,\lambda}^p$ because H_1 depends on α and $\nabla\alpha$. Note that $H_1(\alpha)$ has one degree of differentiability less than expected.

Recalling that $\lambda \notin \mathcal{D}$, we deduce that $F_1(\alpha)$ lies in $d(L_{k,\lambda}^p(\Lambda^2 T^* \hat{N}))$ and hence is L^2 -orthogonal to elements of the kernel of

$$d + d^* : L_{l+1, -3-\lambda}^q(\Lambda^3 T^* \hat{N}) \rightarrow L_{l, -4-\lambda}^q(\Lambda^2 T^* \hat{N} \oplus \Lambda^4 T^* \hat{N}),$$

where $q > 1$ such that $1/p + 1/q = 1$. We show that

$$d(L_{k,\lambda}^p(\Lambda^2 T^* \hat{N})) \oplus d^*(L_{k,\lambda}^p(\Lambda^4 T^* \hat{N})) \subseteq L_{k-1, \lambda-1}^p(\Lambda^3 T^* \hat{N})$$

is *characterised* as the subspace which is L^2 -orthogonal to this kernel.

Consider

$$d + d^* : L_{k,\lambda}^p(\Lambda^{\text{even}} T^* \hat{N}) \rightarrow L_{k-1, \lambda-1}^p(\Lambda^{\text{odd}} T^* \hat{N}).$$

This elliptic map has image which comprises precisely of those elements of $L_{k-1, \lambda-1}^p(\Lambda^{\text{odd}} T^* \hat{N})$ which are L^2 -orthogonal to the kernel \mathcal{K} of

$$d + d^* : L_{l+1, -3-\lambda}^q(\Lambda^{\text{odd}} T^* \hat{N}) \rightarrow L_{l, -4-\lambda}^q(\Lambda^{\text{even}} T^* \hat{N}).$$

The space \mathcal{K} can be written as the direct sum $\mathcal{K} = \mathcal{K}^1 \oplus \mathcal{K}^3 \oplus \mathcal{K}^m$, where

$$\mathcal{K}^j = \mathcal{K} \cap L_{l+1, -3-\lambda}^q(\Lambda^j T^* \hat{N})$$

for $j = 1$ and 3 and \mathcal{K}^m is some transverse subspace. Then

$$d(L_{k, \lambda}^p(\Lambda^2 T^* \hat{N})) \oplus d^*(L_{k, \lambda}^p(\Lambda^4 T^* \hat{N})) = \{\alpha_3 : \exists \alpha_1 \text{ such that } (\alpha_1, \alpha_3) \in \mathcal{K}^\perp\},$$

where we take the orthogonal complement in $L_{k-1, \lambda-1}^p$. Note that the projection $\pi_1(\mathcal{K}^m)$ of \mathcal{K}^m onto the space of 1-forms must meet \mathcal{K}^1 in the zero form since, if $(\alpha_1, \alpha_3) \in \mathcal{K}^m$ and $\alpha_1 \in \mathcal{K}^1$ then $\alpha_3 \in \mathcal{K}^3$, which contradicts the direct sum decomposition of \mathcal{K} . Therefore, $\pi_1(\mathcal{K}^m)$ and \mathcal{K}^1 are transverse finite-dimensional subspaces of $L_{l+1, -3-\lambda}^q(\Lambda^1 T^* \hat{N})$. Hence, there exists a space \mathcal{A} of smooth compactly supported 1-forms on \hat{N} which is L^2 -orthogonal to \mathcal{K}^1 and such that $\mathcal{A} \times \mathcal{K}^m \rightarrow \mathbb{R}$ given by $(\gamma, \xi) \mapsto (\gamma, 0) \cdot \xi$ is a dual pairing. If $\alpha_3 \in L_{k-1, \lambda-1}^p(\Lambda^3 T^* \hat{N})$ such that $\alpha_3 \in (\mathcal{K}^3)^\perp$, there exists a unique $\alpha_1 \in \mathcal{A}$ such that $(\alpha_1, 0) \cdot \xi = -(0, \alpha_3) \cdot \xi$ for all $\xi \in \mathcal{K}^m$, which implies that $(\alpha_1, \alpha_3) \in (\mathcal{K}^m)^\perp$. We conclude that

$$\begin{aligned} (\mathcal{K}^3)^\perp &= \{\alpha_3 \in (\mathcal{K}^3)^\perp : \exists \alpha_1 \in (\mathcal{K}^1)^\perp \text{ such that } (\alpha_1, \alpha_3) \in (\mathcal{K}^m)^\perp\} \\ &= \{\alpha_3 : \exists \alpha_1 \text{ such that } (\alpha_1, \alpha_3) \in \mathcal{K}^\perp\} \\ &= d(L_{k, \lambda}^p(\Lambda^2 T^* \hat{N})) \oplus d^*(L_{k, \lambda}^p(\Lambda^4 T^* \hat{N})) \subseteq L_{k-1, \lambda-1}^p(\Lambda^3 T^* \hat{N}). \end{aligned}$$

However, \mathcal{K}^3 is independent of k , and hence $F_1(\alpha)$ must lie in the image of $d + d^*$ from $L_{k+1, \lambda}^p$, since $F_1(\alpha)$ lies in $L_{k, \lambda-1}^p$. We may thus write $F_1(\alpha) = d\gamma + d^*\beta$ for some $\gamma \in L_{k+1, \lambda}^p(\Lambda^2 T^* \hat{N})$ and $\beta \in L_{k+1, \lambda}^p(\Lambda^4 T^* \hat{N})$. Moreover, $dd^*\beta = 0$ and so β is harmonic and $O(\rho^\lambda)$ as $\rho \rightarrow 0$. By the Maximum Principle, (noting that $*\beta$ is a harmonic function on \hat{N}), $\beta = 0$. The proposition is thus proved. \square

We deduce from Propositions 6.6, 6.8, 6.10 and 7.2 that $\mathcal{M}_1(N, \lambda)$ is locally homeomorphic to the kernel of

$$G_1 : L_{k+1, \lambda}^p(\hat{U}) \times L_{k+1, \lambda}^p(\Lambda^4 T^* \hat{N}) \rightarrow d(L_{k+1, \lambda}^p(\Lambda^2 T^* \hat{N})) \oplus d^*(L_{k+1, \lambda}^p(\Lambda^4 T^* \hat{N})).$$

Therefore, our deformation theory will be obstructed if and only if the map

$$d : L_{k+1, \lambda}^p(\Lambda_+^2 T^* \hat{N}) \rightarrow d(L_{k+1, \lambda}^p(\Lambda^2 T^* \hat{N}))$$

is *not* surjective. This leads us to the next result and definition.

Proposition 7.3 *There exists a finite-dimensional subspace $\mathcal{O}_1(N, \lambda)$ of $L_{k, \lambda-1}^p(\Lambda^3 T^* \hat{N})$ such that*

$$d(L_{k+1, \lambda}^p(\Lambda^2 T^* \hat{N})) = d(L_{k+1, \lambda}^p(\Lambda_+^2 T^* \hat{N})) \oplus \mathcal{O}_1(N, \lambda).$$

Proof: The Fredholmness of $d + d^*$ implies that the images of $L_{k+1, \lambda}^p(\Lambda_+^2 T^* \hat{N}) \oplus L_{k+1, \lambda}^p(\Lambda^4 T^* \hat{N})$ and $L_{k+1, \lambda}^p(\Lambda^2 T^* \hat{N}) \oplus L_{k+1, \lambda}^p(\Lambda^4 T^* \hat{N})$ under $d + d^*$ are both closed and have finite codimension in $L_{k, \lambda-1}^p(\Lambda^3 T^* \hat{N})$. Since

$$\begin{aligned} \{0\} &= d(L_{k+1, \lambda}^p(\Lambda^2 T^* \hat{N})) \cap d^*(L_{k+1, \lambda}^p(\Lambda^4 T^* \hat{N})) \\ &= d(L_{k+1, \lambda}^p(\Lambda_+^2 T^* \hat{N})) \cap d^*(L_{k+1, \lambda}^p(\Lambda^4 T^* \hat{N})) \end{aligned}$$

by the Maximum Principle, we deduce that

$$d(L_{k+1, \lambda}^p(\Lambda_+^2 T^* \hat{N})) \quad \text{and} \quad d(L_{k+1, \lambda}^p(\Lambda^2 T^* \hat{N}))$$

are both closed and that the former has finite codimension in the latter. Thus, $\mathcal{O}_1(N, \lambda)$ can be chosen as stated. \square

Definition 7.4 The *obstruction space* for Problem 1 is

$$\mathcal{O}_1(N, \lambda) \cong \frac{d(L_{k+1, \lambda}^p(\Lambda^2 T^* \hat{N}))}{d(L_{k+1, \lambda}^p(\Lambda_+^2 T^* \hat{N}))}.$$

We proceed as follows. Define

$$\begin{aligned} U_1 &= L_{k+1, \lambda}^p(\hat{U}) \times L_{k+1, \lambda}^p(\Lambda^4 T^* \hat{N}), \\ X_1 &= L_{k+1, \lambda}^p(\Lambda_+^2 T^* \hat{N} \oplus \Lambda^4 T^* \hat{N}), \\ Y_1 &= \mathcal{O}_1(N, \lambda) \subseteq L_{k, \lambda-1}^p(\Lambda^3 T^* \hat{N}) \quad \text{and} \\ Z_1 &= d(L_{k+1, \lambda}^p(\Lambda^2 T^* \hat{N})) \oplus d^*(L_{k+1, \lambda}^p(\Lambda^4 T^* \hat{N})). \end{aligned}$$

Then X_1 , Y_1 and Z_1 are Banach spaces and U_1 is an open neighbourhood of $(0, 0)$ in X_1 because $L_{k+1, \lambda}^p \hookrightarrow C_\lambda^0$ by Theorem 4.4 and \hat{U} grows with order $O(\rho)$ as $\rho \rightarrow 0$ by Proposition 6.4. Thus, $W_1 = U_1 \times Y_1$ is an open neighbourhood of $(0, 0, 0)$ in $X_1 \times Y_1$. Define $\mathcal{G}_1 : W_1 \rightarrow Z_1$ by:

$$\mathcal{G}_1(\alpha, \beta, \gamma) = G_1(\alpha, \beta) + \gamma.$$

Then \mathcal{G}_1 is well-defined by Propositions 7.2 and 7.3 and its derivative at $(0, 0, 0)$ acts from $X_1 \times Y_1$ to Z_1 as

$$d\mathcal{G}_1|_{(0,0,0)} : (\alpha, \beta, \gamma) \longmapsto d\alpha + d^*\beta + \gamma.$$

Clearly, $d\mathcal{G}_1|_{(0,0,0)}$ is surjective by construction and its kernel, using the fact that $(d + d^*)(X_1) \cap Y_1 = \{0\}$, is given by:

$$\begin{aligned} \text{Ker } d\mathcal{G}_1|_{(0,0,0)} &= \{(\alpha, \beta, \gamma) \in X_1 \times Y_1 : d\alpha + d^*\beta + \gamma = 0\} \\ &\cong \{(\alpha, \beta) \in X_1 : d\alpha + d^*\beta = 0\} \cong \mathcal{I}_1(N, \lambda). \end{aligned}$$

The conclusion, by implementing the Implicit Function Theorem for Banach spaces (Theorem 4.5), is that $\text{Ker } \mathcal{G}_1$ is a smooth manifold near zero which may be identified with an open neighbourhood $\hat{\mathcal{M}}_1(N, \lambda)$ of 0 in $\mathcal{I}_1(N, \lambda)$. Formally, if we write $X_1 = \mathcal{I}_1(N, \lambda) \oplus A$ for some closed subspace A of X_1 , there exist open sets $\hat{\mathcal{M}}_1(N, \lambda) \subseteq \mathcal{I}_1(N, \lambda)$, $V_A \subseteq A$, $V_Y \subseteq Y_1$, all containing 0, with $\hat{\mathcal{M}}_1(N, \lambda) \times V_A \subseteq U_1$, and smooth maps $\mathcal{V}_A : \hat{\mathcal{M}}_1(N, \lambda) \rightarrow V_A$ and $\mathcal{V}_Y : \hat{\mathcal{M}}_1(N, \lambda) \rightarrow V_Y$ such that

$$\text{Ker } \mathcal{G}_1 \cap (\hat{\mathcal{M}}_1(N, \lambda) \times V_A \times V_Y) = \{(x, \mathcal{V}_A(x), \mathcal{V}_Y(x)) : x \in \hat{\mathcal{M}}_1(N, \lambda)\}.$$

If we define a smooth map $\pi_1 : \hat{\mathcal{M}}_1(N, \lambda) \rightarrow \mathcal{O}_1(N, \lambda)$ by $\pi_1(x) = \mathcal{V}_Y(x)$, the moduli space $\mathcal{M}_1(N, \lambda)$ near N is locally homeomorphic to the kernel of π_1 near 0. We can think of π_1 as a map on an open neighbourhood of $(0, 0, 0)$ in $\text{Ker } \mathcal{G}_1$ which projects onto the obstruction space. We write these results as a theorem.

Theorem 7.5 *Use the notation of Definitions 6.1, 7.1 and 7.4. There exists a smooth manifold $\hat{\mathcal{M}}_1(N, \lambda)$, which is an open neighbourhood of 0 in $\mathcal{I}_1(N, \lambda)$, and a smooth map $\pi_1 : \hat{\mathcal{M}}_1(N, \lambda) \rightarrow \mathcal{O}_1(N, \lambda)$, with $\pi_1(0) = 0$, such that an open neighbourhood of 0 in $\text{Ker } \pi_1$ is homeomorphic to an open neighbourhood of N in $\mathcal{M}_1(N, \lambda)$.*

We deduce from this theorem that, if the obstruction space is zero, the moduli space is a smooth manifold near N of dimension equal to that of the infinitesimal deformation space. We expect the obstruction space to be zero for generic choices of N and the G_2 structure on M .

7.2 Problem 2

Recall the notation introduced in Definitions 6.12, 6.14 and 6.16. We begin by defining the infinitesimal deformation space for this problem.

Definition 7.6 *The infinitesimal deformation space for Problem 2 is*

$$\begin{aligned} \mathcal{I}_2(N, \lambda) &= \{(\alpha, t) \in C_\lambda^\infty(\Lambda_+^2 T^* \hat{N}) \oplus T_0 \hat{\mathcal{T}} : d\alpha + L_2(t) = 0\} \\ &\cong \{(\alpha, \beta, t) \in C_\lambda^\infty(\Lambda_+^2 T^* \hat{N} \oplus \Lambda^4 T^* \hat{N}) \oplus T_0 \hat{\mathcal{T}} : d\alpha + d^* \beta + L_2(t) = 0\}. \end{aligned}$$

The equivalence in the definition follows from Proposition 6.17 or from the observation that $d\alpha + L_2(t)$ is exact and so $\beta = 0$ by the Maximum Principle.

By Proposition 6.18,

$$\mathcal{I}_2(N, \lambda) \cong \{(\alpha, \beta, t) \in L_{k+1, \lambda}^p(\Lambda_+^2 T^* \hat{N} \oplus \Lambda^4 T^* \hat{N}) \oplus T_0 \hat{\mathcal{T}} : d\alpha + d^* \beta + L_2(t) = 0\}.$$

Therefore, $\mathcal{I}_2(N, \lambda)$ is finite-dimensional.

Note There is a subspace of $\mathcal{I}_2(N, \lambda)$ which is isomorphic to $\mathcal{I}_1(N, \lambda)$.

To start our consideration of obstructions, we have the generalisation of Proposition 7.2.

Proposition 7.7 *The map F_2 takes $L_{k+1, \lambda}^p(\hat{U}) \times \hat{\mathcal{T}}$ into $d(L_{k+1, \lambda}^p(\Lambda^2 T^* \hat{N}))$.*

Proof: Use the notation from Proposition 6.13 and its proof and from the proof of Proposition 7.2. Recall that we have an open set $T \supseteq \hat{T}$ in M containing N , which retracts onto N , and $\psi \in C^\infty(\Lambda^2 T^* T)$ such that $d\psi = \varphi|_T$. We may similarly construct open sets $T(t) \supseteq \hat{T}(t)$ in M , with $T(0) = T$, which contain $N(t)$ and retract onto it, varying smoothly with t . We also have $\psi(t) \in C^\infty(\Lambda^2 T^* T(t))$, with $\psi(0) = \psi$, such that $d\psi(t) = \varphi|_{T(t)}$, using the fact that φ is exact on $N(t)$. Again, the $\psi(t)$ vary smoothly with t . Formally, let

$$T(t) = \hat{T}(t) \cup \bigcup_{i=1}^s V_i(t).$$

By making $\hat{T}(t)$ and $V_i(t)$ smaller if necessary, $T(t)$ will be an open set as stated. We may choose $\psi(t)$ such that

$$\chi_i(t)^*(\psi(t)) = \frac{1}{3}(v \cdot \varphi_0) + O(r_i^2)$$

on $V_i(t)$ and then extend smoothly to a form $\psi(t)$ on $T(t)$ as required. Define

$$H_2(\alpha, t) = f_\alpha(t)^* \left(\psi(t)|_{\hat{N}_\alpha(t)} \right).$$

Then $d(H_2(\alpha, t)) = F_2(\alpha, t)$. Moreover, by the same reasoning that $H_1(\alpha) \in L_{k, \lambda}^p$ in the proof of Proposition 7.2, $H_2(\alpha, t)$ lies in $L_{k, \lambda}^p$. Therefore, $F_2(\alpha, t)$ lies in $d(L_{k, \lambda}^p(\Lambda^2 T^* \hat{N}))$. However, because $F_2(\alpha, t) \in L_{k, \lambda-1}^p(\Lambda^3 T^* \hat{N})$, the argument at the end of the proof of Proposition 7.2 implies that $F_2(\alpha, t) \in d(L_{k+1, \lambda}^p(\Lambda^2 T^* \hat{N}))$ as required. \square

We now define the obstruction space.

Definition 7.8 From Propositions 7.3 and 7.7, since L_2 is a linear map on a finite-dimensional vector space, there exists a finite-dimensional subspace $\mathcal{O}_2(N, \lambda)$ of $L_{k, \lambda-1}^p(\Lambda^3 T^* \hat{N})$ such that

$$d(L_{k+1, \lambda}^p(\Lambda^2 T^* \hat{N})) = (d(L_{k+1, \lambda}^p(\Lambda_+^2 T^* \hat{N})) + L_2(T_0 \hat{\mathcal{T}})) \oplus \mathcal{O}_2(N, \lambda).$$

We define $\mathcal{O}_2(N, \lambda)$ to be the *obstruction space* for Problem 2.

Note $\mathcal{O}_2(N, \lambda)$ may be chosen to be contained in $\mathcal{O}_1(N, \lambda)$.

Following the scheme for Problem 1, we let

$$\begin{aligned} U_2 &= L_{k+1, \lambda}^p(\hat{U}) \times L_{k+1, \lambda}^p(\Lambda^4 T^* \hat{N}) \times \hat{\mathcal{T}}, \\ X_2 &= L_{k+1, \lambda}^p(\Lambda_+^2 T^* \hat{N} \oplus \Lambda^4 T^* \hat{N}) \oplus T_0 \hat{\mathcal{T}}, \\ Y_2 &= \mathcal{O}_2(N, \lambda) \subseteq L_{k, \lambda-1}^p(\Lambda^3 T^* \hat{N}) \text{ and} \\ Z_2 &= d(L_{k+1, \lambda}^p(\Lambda^2 T^* \hat{N})) \oplus d^*(L_{k+1, \lambda}^p(\Lambda^4 T^* \hat{N})). \end{aligned}$$

Recall that $\hat{\mathcal{T}} \subseteq \mathbb{R}^n \cong T_0 \hat{\mathcal{T}}$ is open. Then X_2, Y_2 and Z_2 are Banach spaces, U_2 is an open neighbourhood of $(0, 0, 0)$ in X_2 and hence $W_2 = U_2 \times Y_2$ is an open neighbourhood of $(0, 0, 0, 0)$ in $X_2 \times Y_2$. Define $\mathcal{G}_2 : W_2 \rightarrow Z_2$ by:

$$\mathcal{G}_2(\alpha, \beta, t, \gamma) = G_2(\alpha, \beta, t) + \gamma.$$

Then $d\mathcal{G}_2|_{(0,0,0,0)} : X_2 \times Y_2 \rightarrow Z_2$ acts as

$$(\alpha, \beta, t, \gamma) \mapsto d\alpha + d^*\beta + L_2(t) + \gamma.$$

By construction, $d\mathcal{G}_2|_{(0,0,0,0)}$ is surjective and, using the fact that the image of $d\mathcal{G}_2|_{(0,0,0)}$ meets Y_2 at 0 only,

$$\begin{aligned} \text{Ker } d\mathcal{G}_2|_{(0,0,0,0)} &= \{(\alpha, \beta, t, \gamma) \in X_2 \times Y_2 : d\alpha + d^*\beta + L_2(t) + \gamma = 0\} \\ &\cong \{(\alpha, \beta, t) \in X_2 : d\alpha + d^*\beta + L_2(t) = 0\} \cong \mathcal{I}_2(N, \lambda). \end{aligned}$$

As for Problem 1, Theorem 4.5 gives us that $\text{Ker } \mathcal{G}_2$ is a smooth manifold near zero which may be identified with an open neighbourhood $\hat{\mathcal{M}}_2(N, \lambda)$ of $(0, 0)$ in $\mathcal{I}_2(N, \lambda)$. We can again define a smooth map $\pi_2 : \hat{\mathcal{M}}_2(N, \lambda) \rightarrow \mathcal{O}_2(N, \lambda)$ such that $\text{Ker } \pi_2$ is locally homeomorphic near $(0, 0)$ to an open neighbourhood of N in $\mathcal{M}_2(N, \lambda)$. We thus have the following theorem.

Theorem 7.9 *Use the notation of Definitions 6.11, 7.6 and 7.8. There exists a smooth manifold $\hat{\mathcal{M}}_2(N, \lambda)$, which is an open neighbourhood of $(0, 0)$ in $\mathcal{I}_2(N, \lambda)$, and a smooth map $\pi_2 : \hat{\mathcal{M}}_2(N, \lambda) \rightarrow \mathcal{O}_2(N, \lambda)$, with $\pi_2(0, 0) = 0$, such that an open neighbourhood of zero in $\text{Ker } \pi_2$ is homeomorphic to an open neighbourhood of N in $\mathcal{M}_2(N, \lambda)$.*

We deduce that, if $\mathcal{O}_2(N, \lambda) = \{0\}$, the moduli space for Problem 2 is a smooth manifold near N of dimension $\dim \mathcal{I}_2(N, \lambda) = \dim \mathcal{I}_1(N, \lambda) + \dim \hat{\mathcal{T}}$, which we expect to occur for generic choices of N and the torsion-free \mathbb{G}_2 structure on M . We shall see, in §9, that if we choose a suitable generic *closed* \mathbb{G}_2 structure on M we may drop the assumption that N is generic and still obtain a smooth moduli space.

7.3 Problem 3

We presume in this subsection that the reader is sufficiently familiar with the schemata we have used in the previous two subsections to be able to generalise them to Problem 3. This allows us to present a tidier treatment of the problem.

Recall the notation of Definitions 6.20, 6.24 and 6.26.

Definition 7.10 The *infinitesimal deformation space* $\mathcal{I}_3(N, \lambda)$ for Problem 3 is

$$\begin{aligned} \mathcal{I}_3(N, \lambda) &= \{(\alpha, t, f) \in C_\lambda^\infty(\Lambda_+^2 T^* \hat{N}) \oplus T_0 \hat{\mathcal{T}} \oplus T_0 \hat{\mathcal{F}} : d\alpha + L_2(t) + L_3(f) = 0\} \\ &\cong \{(\alpha, \beta, t, f) \in C_\lambda^\infty(\Lambda_+^2 T^* \hat{N} \oplus \Lambda^4 T^* \hat{N}) \oplus T_0 \hat{\mathcal{T}} \oplus T_0 \hat{\mathcal{F}} \\ &\quad : d\alpha + d^* \beta + L_2(t) + L_3(f) = 0\}. \end{aligned}$$

By Proposition 6.28,

$$\begin{aligned} \mathcal{I}_3(N, \lambda) &\cong \{(\alpha, \beta, t, f) \in L_{k+1, \lambda}^p(\Lambda_+^2 T^* \hat{N} \oplus \Lambda^4 T^* \hat{N}) \oplus T_0 \hat{\mathcal{T}} \oplus T_0 \hat{\mathcal{F}} \\ &\quad : d\alpha + d^* \beta + L_2(t) + L_3(f) = 0\}. \end{aligned}$$

In considering obstructions, we first have the generalisation of Propositions 7.2 and 7.7.

Proposition 7.11 $F_3(L_{k+1, \lambda}^p(\hat{U}) \times \hat{\mathcal{T}} \times \hat{\mathcal{F}}) \subseteq d(L_{k+1, \lambda}^p(\Lambda^2 T^* \hat{N}))$.

The proposition is proved in a similar way to Proposition 7.7 and so we omit the details. The result leads us to define our final obstruction space.

Definition 7.12 From Propositions 7.3 and 7.11, since L_2 and L_3 are linear maps on finite-dimensional vector spaces, there exists a finite-dimensional subspace $\mathcal{O}_3(N, \lambda)$ of $L_{k, \lambda-1}^p(\Lambda^3 T^* \hat{N})$ such that

$$d(L_{k+1, \lambda}^p(\Lambda^2 T^* \hat{N})) = (d(L_{k+1, \lambda}^p(\Lambda_+^2 T^* \hat{N})) + L_2(T_0 \hat{\mathcal{T}}) + L_3(T_0 \hat{\mathcal{F}})) \oplus \mathcal{O}_3(N, \lambda).$$

We define $\mathcal{O}_3(N, \lambda)$ to be the *obstruction space* for Problem 3.

Note We may choose our obstruction spaces such that $\mathcal{O}_3(N, \lambda) \subseteq \mathcal{O}_2(N, \lambda) \subseteq \mathcal{O}_1(N, \lambda)$.

The use of the Implicit Function Theorem (Theorem 4.5) in the derivation of Theorems 7.5 and 7.9 can be easily generalised to give the following.

Theorem 7.13 *Use the notation of Definitions 6.21, 7.10 and 7.12. There exists a smooth manifold $\hat{\mathcal{M}}_3(N, \lambda)$, which is an open neighbourhood of $(0, 0, 0)$ in $\mathcal{I}_3(N, \lambda)$, and a smooth map $\pi_3 : \hat{\mathcal{M}}_3(N, \lambda) \rightarrow \mathcal{O}_3(N, \lambda)$, with $\pi_3(0, 0, 0) = 0$, such that an open neighbourhood of zero in $\text{Ker } \pi_3$ is homeomorphic to an open neighbourhood of $(N, 0)$ in $\mathcal{M}_3(N, \lambda)$.*

We deduce that, if $\mathcal{O}_3(N, \lambda) = \{0\}$, $\mathcal{M}_3(N, \lambda)$ is a smooth manifold near $(N, 0)$ of dimension $\dim \mathcal{I}_3(N, \lambda) = \dim \mathcal{I}_2(N, \lambda) + \dim \hat{\mathcal{F}}$. Moreover, the projection map $\pi_{\hat{\mathcal{F}}} : \mathcal{M}_3(N, \lambda) \rightarrow \hat{\mathcal{F}}$ is smooth near $(N, 0)$. We expect this to occur for generic choices of N and the torsion-free G_2 structure on M . If we allow ourselves to work with *closed* G_2 structures on M , we shall show in §9 that we may drop our genericity assumptions for N and (φ, g) and still get a smooth moduli space.

8 Dimension calculations

We shall relate the expected dimension of the moduli space for Problem 1 to the index of $d + d^*$ as discussed in §5.2. Recall that $p > 4$, $k \geq 2$ and $\lambda \in (1, 2) \setminus \mathcal{D}$.

Definition 8.1 Define

$$\mathcal{H}^m = \{\xi \in L^2(\Lambda^m T^* \hat{N}) : d\xi = d^* \xi = 0\}.$$

The Hodge star maps \mathcal{H}^2 into itself, so there is a splitting $\mathcal{H}^2 = \mathcal{H}_+^2 \oplus \mathcal{H}_-^2$ where

$$\mathcal{H}_\pm^2 = \mathcal{H}^2 \cap C^\infty(\Lambda_\pm^2 T^* \hat{N}).$$

Let $\mathcal{J} = j\left(H_{\text{cs}}^2(\hat{N})\right)$, where $H_{\text{cs}}^m(\hat{N})$ is the m th *compactly supported* cohomology group on \hat{N} and $j : H_{\text{cs}}^2(\hat{N}) \rightarrow H_{\text{dR}}^2(\hat{N})$ is the inclusion map. If $\alpha, \beta \in \mathcal{J}$, there exist compactly supported closed 2-forms ξ and η such that $\alpha = [\xi]$ and $\beta = [\eta]$. We define a product on $\mathcal{J} \times \mathcal{J}$ by

$$\alpha \cup \beta = \int_{\hat{N}} \xi \wedge \eta. \quad (27)$$

Suppose that ξ' and η' are also compactly supported with $\alpha = [\xi']$ and $\beta = [\eta']$. Then there exist 1-forms χ and ζ such that $\xi - \xi' = d\chi$ and $\eta - \eta' = d\zeta$. Therefore,

$$\begin{aligned} \int_{\hat{N}} \xi' \wedge \eta' &= \int_{\hat{N}} (\xi - d\chi) \wedge (\eta - d\zeta) = \int_{\hat{N}} \xi \wedge \eta - d\chi \wedge \eta - \xi' \wedge d\zeta \\ &= \int_{\hat{N}} \xi \wedge \eta - d(\chi \wedge \eta) - d(\xi' \wedge \zeta) = \int_{\hat{N}} \xi \wedge \eta, \end{aligned}$$

as both $\chi \wedge \eta$ and $\xi' \wedge \zeta$ have compact support. The product (27) on $\mathcal{J} \times \mathcal{J}$ is thus well-defined and is a symmetric topological product with a signature (a, b) . By [13, Example (0.16)], $\mathcal{H}^2 \cong \mathcal{J}$ and the isomorphism is given by $\xi \mapsto [\xi]$. Thus, $\dim \mathcal{H}_+^2 = a$ and hence is a topological number.

Definition 8.2 Let

$$(d_+ + d^*)_\lambda = d + d^* : L_{k+1, \lambda}^p(\Lambda_+^2 T^* \hat{N} \oplus \Lambda^4 T^* \hat{N}) \rightarrow L_{k, \lambda-1}^p(\Lambda^3 T^* \hat{N}).$$

By Definition 7.1, $\mathcal{I}_1(N, \lambda)$ is isomorphic to the kernel of this map. Define the adjoint map by

$$(d_+^* + d)_{-3-\lambda} = d_+^* + d : L_{l+1, -3-\lambda}^q(\Lambda^3 T^* \hat{N}) \longrightarrow L_{l, -4-\lambda}^q(\Lambda_+^2 T^* \hat{N} \oplus \Lambda^4 T^* \hat{N}),$$

where $q > 1$ such that $1/p + 1/q = 1$ and $l \geq 4$. The cokernel of $(d_+ + d^*)_\lambda$ is then isomorphic to the kernel of $(d_+^* + d)_{-3-\lambda}$.

Note The choice of $l \geq 4$ in Definition 8.2 ensures that $L_{l+1, -3-\lambda}^q \hookrightarrow C_{-3-\lambda}^{1, a}$ for $0 < a \leq 4 - \frac{4}{q} = \frac{4}{p} < 1$ by Theorem 4.4.

We now study the dimension of the kernel and cokernel of $(d_+ + d^*)_\mu$.

Proposition 8.3 *The kernel of $(d_+ + d^*)_{-2}$ is isomorphic to \mathcal{H}_+^2 . Furthermore, if $\mu > -2$ is such that $(-2, \mu] \cap \mathcal{D} = \emptyset$, $\dim \text{Ker} (d_+ + d^*)_\mu = \dim \mathcal{H}_+^2$.*

Proof: Using (7) and the Maximum Principle,

$$\begin{aligned} \mathcal{H}_+^2 &= \{\alpha \in L^2(\Lambda^2 T^* \hat{N}) \cap C^\infty(\Lambda_+^2 T^* \hat{N}) : d\alpha = d^* \alpha = 0\} \\ &= \{\alpha \in L_{0, -2}^2(\Lambda_+^2 T^* \hat{N}) \cap C^\infty(\Lambda_+^2 T^* \hat{N}) : d\alpha = 0\} \\ &\cong \{(\alpha, \beta) \in L_{0, -2}^2(\Lambda_+^2 T^* \hat{N} \oplus \Lambda^4 T^* \hat{N}) : \alpha \in C^\infty(\Lambda_+^2 T^* \hat{N}), d\alpha + d^* \beta = 0\}. \end{aligned}$$

This gives the first part of the proposition.

If $-2 \notin \mathcal{D}$, $[-2, \mu] \cap \mathcal{D} = \emptyset$ and thus, by Proposition 5.6, $\dim \text{Ker} (d_+ + d^*)_\mu = \dim \text{Ker} (d_+ + d^*)_{-2}$.

Suppose now that $-2 \in \mathcal{D}$ and that (α, β) corresponds to a self-dual 2-form and 4-form on \hat{N} which are subtracted from the kernel of $(d_+ + d^*)_\nu$ as ν crosses -2 from below. By the work in [14, §3 & §4] this occurs if and only if (α, β) is asymptotic to an $O(r^{-2})$ form ξ on C_i , for some i , satisfying $(d + d^*)\xi = 0$. (The form ξ is determined by an element of $D(-2, i)$, using the notation of Proposition 5.2.) Therefore, (α, β) is of order $O(\rho^{-2})$ as $\rho \rightarrow 0$ and thus lies in L^2 . We deduce that $(\alpha, \beta) \in \text{Ker} (d_+ + d^*)_{-2}$, implying that the function $k(\nu) = \dim \text{Ker} (d_+ + d^*)_\nu$ is upper semi-continuous at -2 by Proposition 5.6.

The second part of the proposition is thus proved. \square

Proposition 8.4 *If $\mu < -1$ is such that $[\mu, -1) \cap \mathcal{D} = \emptyset$, the cokernel of $(d_+ + d^*)_\mu$ is isomorphic to $H_{\text{dR}}^1(\hat{N})$.*

Proof: By Theorem 5.1, there exists a countable discrete subset \mathcal{D}' of rates ν such that

$$d + d^* : L_{k+1, \nu}^p(\Lambda^{\text{even}}T^*\hat{N}) \rightarrow L_{k, \nu-1}^p(\Lambda^{\text{odd}}T^*\hat{N}) \quad (28)$$

is not Fredholm. Clearly, $\mathcal{D}' \supseteq \mathcal{D}$. For $\nu \notin \mathcal{D}'$ with $\nu < -1$, so that $-3 - \nu > \nu - 1$,

$$L_{k, \nu-1}^p(\Lambda^{\text{odd}}T^*\hat{N}) = (d + d^*)(L_{k+1, \nu}^p(\Lambda^{\text{even}}T^*\hat{N})) \oplus \mathcal{K},$$

where \mathcal{K} is the kernel of the adjoint map

$$d + d^* : L_{l+1, -\nu-3}^q(\Lambda^{\text{odd}}T^*\hat{N}) \rightarrow L_{l, -\nu-4}^q(\Lambda^{\text{even}}T^*\hat{N}),$$

for $1/p + 1/q = 1$ and $l \geq 4$, which is graded and closed under the Hodge star.

If $\gamma \in L_{k, \nu-1}^p(\Lambda^3T^*\hat{N})$ then $(*\gamma, \gamma) \in L_{k, \nu-1}^p(\Lambda^{\text{odd}}T^*\hat{N})$ and hence there exist some $\gamma_m \in L_{k+1, \nu}^p(\Lambda^mT^*\hat{N})$, for $m = 0, 2, 4$, and $\eta \in \mathcal{K}$ such that

$$(*\gamma, \gamma) = (d + d^*)(\gamma_0, \gamma_2, \gamma_4) + \eta.$$

By applying the Hodge star,

$$(*\gamma, \gamma) = (d + d^*)(*\gamma_4, *\gamma_2, *\gamma_0) + *\eta.$$

Adding the above formulae and averaging gives:

$$\gamma = d\left(\frac{\gamma_2 + *\gamma_2}{2}\right) + d^*\left(\frac{*\gamma_0 + \gamma_4}{2}\right) + \tilde{\eta}$$

where $\tilde{\eta} \in \mathcal{K} \cap L_{k, \nu-1}^p(\Lambda^3T^*\hat{N})$. We deduce that

$$L_{k, \nu-1}^p(\Lambda^3T^*\hat{N}) = \left(d(L_{k+1, \nu}^p(\Lambda_+^2T^*\hat{N})) + d^*(L_{k+1, \nu}^p(\Lambda_+^4T^*\hat{N}))\right) \oplus \mathcal{K}^3,$$

where $\mathcal{K}^3 = \mathcal{K} \cap L_{k, \nu-1}^p(\Lambda^3T^*\hat{N})$. Moreover, for $\nu \notin \mathcal{D}'$, $\nu < -1$,

$$d(L_{k+1, \lambda}^p(\Lambda^2T^*\hat{N})) = d(L_{k+1, \lambda}^p(\Lambda_+^2T^*\hat{N})).$$

We must surely have that the images are equal for $\nu \notin \mathcal{D}$, $\nu < -1$, as well.

Thus, the cokernel of $(d_+ + d^*)_\mu$ is isomorphic to the kernel of

$$(d^* + d)_{-3-\mu} = d^* + d : L_{l+1, -3-\mu}^q(\Lambda^3T^*\hat{N}) \longrightarrow L_{l, -4-\mu}^q(\Lambda^2T^*\hat{N} \oplus \Lambda^4T^*\hat{N}). \quad (29)$$

Using (7) as in the proof of Proposition 8.3, the kernel of $(d^* + d)_{-3-(-1)} = (d^* + d)_{-2}$ is isomorphic to \mathcal{H}^3 . By [13, Example (0.16)], $\mathcal{H}^3 \cong H_{\text{dR}}^1(\hat{N})$ and the isomorphism is given by $\gamma \mapsto [* \gamma]$. Since $[\mu, -1] \cap \mathcal{D} = \emptyset$, there are no changes in the cokernel in $[\mu, -1]$ by Proposition 5.6. Moreover, the dimension of the

cokernel is lower semi-continuous in μ at -1 ; this fact can be demonstrated using similar methods to those employed in the proof of Proposition 8.3. The result follows. \square

By Proposition 5.3, $(-2, -1] \cap \mathcal{D} = \emptyset$. Therefore, for any $\mu \in (-2, -1]$,

$$\dim \text{Ker}(d_+ + d^*)_\mu = \dim \mathcal{H}_+^2 \quad \text{and} \quad \dim \text{Coker}(d_+ + d^*)_\mu = b^1(\hat{N}),$$

using Propositions 8.3 and 8.4. Knowing the index of $(d_+ + d^*)_\mu$ for $\mu \in (-2, -1]$, we can calculate it for all growth rates using Theorem 5.5.

Proposition 8.5 *Use the notation of Propositions 5.2 and 5.7. If $\lambda \in (1, 2)$, $\lambda \notin \mathcal{D}$, the index of $(d_+ + d^*)_\lambda$ is given by:*

$$\text{ind}(d_+ + d^*)_\lambda = \dim \mathcal{H}_+^2 - b^1(\hat{N}) - \sum_{\mu \in (-1, \lambda) \cap \mathcal{D}} d(\mu).$$

However, the obstruction space $\mathcal{O}_1(N, \lambda)$ given in Definition 7.4 is a subspace of the cokernel of $(d_+ + d^*)_\lambda$, so we must relate their dimensions.

Proposition 8.6 *The following inequality holds:*

$$\dim \mathcal{O}_1(N, \lambda) \leq \dim \text{Coker}(d_+ + d^*)_\lambda - b^1(\hat{N}).$$

Proof: From the proof of Proposition 7.2, the image of

$$(d + d^*)_\lambda = d + d^* : L_{k+1, \lambda}^p(\Lambda^2 T^* \hat{N} \oplus \Lambda^4 T^* \hat{N}) \rightarrow L_{k, \lambda-1}^p(\Lambda^3 T^* \hat{N})$$

is characterised as the subspace of $L_{k, \lambda-1}^p(\Lambda^3 T^* \hat{N})$ which is L^2 -orthogonal to the kernel \mathcal{K} of $(d^* + d)_{-3-\lambda}$ defined by (29). Furthermore, as noticed in the proof of Proposition 7.3, $\text{Image}(d + d^*)_\lambda$ has finite codimension in $L_{k, \lambda-1}^p(\Lambda^3 T^* \hat{N})$. Therefore, we may choose a finite-dimensional space \mathcal{C} of smooth compactly supported 3-forms on \hat{N} such that

$$L_{k, \lambda-1}^p(\Lambda^3 T^* \hat{N}) = \text{Image}(d + d^*)_\lambda \oplus \mathcal{C}$$

and so that the product $\mathcal{C} \times \mathcal{K} \rightarrow \mathbb{R}$ given by $(\gamma, \eta) \mapsto \langle \gamma, \eta \rangle_{L^2}$ is nondegenerate.

We may similarly deduce that the image of $(d_+ + d^*)_\lambda$ is the subspace of $L_{k, \lambda-1}^p(\Lambda^3 T^* \hat{N})$ which is L^2 -orthogonal to the kernel \mathcal{K}' of $(d_+^* + d)_{-3-\lambda}$. Then $\mathcal{K}' \supseteq \mathcal{K}$ and \mathcal{K} consists of closed and coclosed 3-forms, whereas \mathcal{K}' consists of 3-forms η such that $d\eta = d_+^* \eta = 0$. Hence, we may choose a subspace \mathcal{K}'' of \mathcal{K}' , transverse to \mathcal{K} , comprising 3-forms which are *not* coclosed and such that $\mathcal{K}' = \mathcal{K} \oplus \mathcal{K}''$.

The next stage is to extend \mathcal{C} to a space $\mathcal{C}' = \mathcal{C} \oplus \mathcal{C}''$, where \mathcal{C}'' consists of smooth *exact* compactly supported 3-forms on \hat{N} , such that

$$L_{k, \lambda-1}^p(\Lambda^3 T^* \hat{N}) = \text{Image}(d_+ + d^*)_\lambda \oplus \mathcal{C}'$$

and such that the product $\mathcal{C}'' \times \mathcal{K}'' \rightarrow \mathbb{R}$ given by $(\gamma, \eta) \mapsto \langle \gamma, \eta \rangle_{L^2}$ is nondegenerate, which is possible as \mathcal{K}'' comprises forms which are not coclosed. By construction, \mathcal{C}'' is a valid choice for $\mathcal{O}_1(N, \lambda)$ by Proposition 7.3. Therefore,

$$\dim \mathcal{O}_1(N, \lambda) = \dim \mathcal{C}' - \dim \mathcal{C} = \dim \text{Coker}(d_+ + d^*)_\lambda - \dim \mathcal{K}.$$

If γ lies in the kernel of (29) for rate $\mu = -1$ then $\gamma \in \mathcal{K}$ for $\lambda \in (1, 2)$ by Theorem 4.4. Thus, the map from \mathcal{K} to $H_{\text{dR}}^1(\hat{N})$ given by $\gamma \mapsto [* \gamma]$ is surjective. This gives the result. \square

We may now calculate a lower bound for the expected dimension of $\mathcal{M}_1(N, \lambda)$ using Propositions 8.5 and 8.6.

Proposition 8.7 *Using the notation of Propositions 5.2 and 5.7,*

$$\dim \mathcal{I}_1(N, \lambda) - \dim \mathcal{O}_1(N, \lambda) \geq \dim \mathcal{H}_+^2 - \sum_{\mu \in (-1, \lambda) \cap \mathcal{D}} d(\mu).$$

Recalling that the dimension of \mathcal{T} given in Definition 6.12 is $21s$, we derive analogous results for our other problems.

Proposition 8.8 *Using the notation of Definitions 6.12 and 6.20 and Propositions 5.2 and 5.7,*

$$\dim \mathcal{I}_2(N, \lambda) - \dim \mathcal{O}_2(N, \lambda) \geq \dim \mathcal{H}_+^2 + 21s - \dim \text{H} - \sum_{\mu \in (-1, \lambda) \cap \mathcal{D}} d(\mu).$$

and

$$\dim \mathcal{I}_3(N, \lambda) - \dim \mathcal{O}_3(N, \lambda) \geq \dim \mathcal{H}_+^2 + 21s - \dim \text{H} + \dim \hat{\mathcal{F}} - \sum_{\mu \in (-1, \lambda) \cap \mathcal{D}} d(\mu).$$

We note that Propositions 5.3, 8.4, 8.6 and 8.7 imply the following bound on $\dim \mathcal{O}_1(N, \lambda)$.

Proposition 8.9 *In the notation of Propositions 5.2 and 5.7,*

$$\dim \mathcal{O}_1(N, \lambda) \leq \sum_{\mu \in (-1, \lambda) \cap \mathcal{D}} d(\mu)$$

We also know that, in Problem 2, we remove the obstructions which correspond to translations of the singularities and G_2 transformations of the tangent cones. These obstructions occur, respectively, at rates 0 and 1. Hence, $d(0) \geq 7s$, $d(1) \geq 14s - \dim H$ and we have the following stronger bound on the dimension of $\mathcal{O}_2(N, \lambda)$.

Proposition 8.10 *In the notation of Definition 6.12 and Propositions 5.2 and 5.7,*

$$\dim \mathcal{O}_2(N, \lambda) \leq -21s + \dim H + \sum_{\mu \in (-1, \lambda) \cap \mathcal{D}} d(\mu).$$

9 φ -Closed 7-manifolds

For our deformation problems we have assumed the ambient manifold (M, φ, g) is a G_2 manifold; that is, M is endowed with a G_2 structure such that $d\varphi = d^*\varphi = 0$. However, the results of McLean [16] we have used, which are based upon the linearisation of the map we denoted F_1 in Definition 6.5, still hold if this condition on φ is relaxed to just $d\varphi = 0$. Thus, our deformation theory results hold if (M, φ, g) is a φ -closed 7-manifold in the sense of Definition 2.6.

Remark The effect of $*\varphi$ not being closed on M means that coassociative 4-folds in M are no longer necessarily volume minimizing in their homology class. This does not, however, affect our discussion.

The use of φ -closed 7-manifolds (M, φ, g) is that closed G_2 structures occur in infinite-dimensional families, since the set of closed definite 3-forms on M , in the sense of Definition 2.3, is open. We show that we can choose a family \mathcal{F} , in a similar fashion to Definition 6.20 of Problem 3, of closed G_2 structures on M such that $\dim \mathcal{F} = \dim \mathcal{O}_1(N, \lambda)$ and, further, such that $\mathcal{O}_3(N, \lambda) = \{0\}$. In other words, we have enough freedom in our choice of \mathcal{F} to ensure that $dF_3|_{(0,0,0)}$, as given in Definition 6.24, maps onto $d(L_{k+1}^p, \lambda(\Lambda^2 T^* \hat{N}))$. Then $\mathcal{M}_3(N, \lambda)$ is a smooth manifold near $(N, 0)$ by Theorem 7.13 and $\pi_{\hat{\mathcal{F}}} : \mathcal{M}_3(N, \lambda) \rightarrow \hat{\mathcal{F}}$ is a smooth map near $(N, 0)$.

Sard's Theorem [11, p. 173] states that, if $f : X \rightarrow Y$ is a smooth map between finite-dimensional manifolds, the set of $y \in Y$ with some $x \in f^{-1}(y)$ such that $df|_x : T_x X \rightarrow T_y Y$ is *not* surjective is of measure zero in Y . Therefore, $f^{-1}(y)$ is a submanifold of X for almost all $y \in Y$.

By Sard's Theorem, $\pi_{\hat{\mathcal{F}}}^{-1}(f)$ is a smooth manifold near (N, f) for almost all $f \in \hat{\mathcal{F}}$. As observed in Definition 6.21, $\pi_{\hat{\mathcal{F}}}^{-1}(f)$ corresponds to the moduli space of deformations for Problem 2 defined using the G_2 structure (φ^f, g^f) . Thus,

for any given N , a generic perturbation of the closed G_2 structure within \mathcal{F} ensures that $\mathcal{M}_2(N, \lambda)$ is smooth near N .

We thus prove the following, which is similar to the result [7, Theorem 9.1].

Theorem 9.1 *Let (M, φ, g) be a φ -closed 7-manifold in the sense of Definition 2.6 and let N in (M, φ, g) be a CS coassociative 4-fold at z_1, \dots, z_s with rate $\lambda \in (1, 2) \setminus \mathcal{D}$, where \mathcal{D} is defined in Proposition 5.2. Use the notation of Definitions 6.12, 7.4 and 7.12 and Proposition 5.7. Let $m = \dim \mathcal{O}_1(N, \lambda)$ and let $\hat{\mathcal{F}}$ be an open ball about 0 in \mathbb{R}^m . There exists a smooth family $\mathcal{F} = \{(\varphi^f, g^f) : f \in \hat{\mathcal{F}}\}$ of closed G_2 structures on M such that $\mathcal{O}_3(N, \lambda) = \{0\}$. Hence, the moduli space of deformations for Problem 3 is a smooth manifold near $(N, 0)$ of dimension greater than or equal to*

$$\dim \mathcal{H}_+^2 + 21s - \dim \mathbb{H} + \dim \mathcal{O}_1(N, \lambda) - \sum_{\mu \in (-1, \lambda) \cap \mathcal{D}} d(\mu).$$

Moreover, for generic $f \in \hat{\mathcal{F}}$, the moduli space of deformations in (M, φ^f, g^f) for Problem 2 is a smooth manifold near N of dimension greater than or equal to

$$\dim \mathcal{H}_+^2 + 21s - \dim \mathbb{H} - \sum_{\mu \in (-1, \lambda) \cap \mathcal{D}} d(\mu).$$

Proof: Use the notation in the proof of Proposition 8.6. Recall that we have a subspace \mathcal{K}'' of $L_{l+1, -3-\lambda}^q(\Lambda^3 T^* \hat{N})$ consisting of forms η such that $d\eta = d_+^* \eta = 0$ but $d^* \eta \neq 0$. Moreover, $\mathcal{O}_1(N, \lambda)$ can be chosen to be a space of smooth compactly supported exact 3-forms γ such that $\langle \gamma, \eta \rangle_{L^2} = 0$ for all $\eta \in \mathcal{K}'' \setminus \{0\}$ implies that $\gamma = 0$. Therefore $\mathcal{K}'' \cong (\mathcal{O}_1(N, \lambda))^*$ and hence has dimension m .

Let $\{\eta_1, \dots, \eta_m\}$ be a basis for \mathcal{K}'' and choose a basis $\{dv_1, \dots, dv_m\}$ for $\mathcal{O}_1(N, \lambda)$, where v_j is a smooth compactly supported 2-form for all j , such that $\langle dv_i, \eta_j \rangle_{L^2} = \delta_{ij}$. This is possible because the L^2 product on $\mathcal{O}_1(N, \lambda) \times \mathcal{K}''$ is nondegenerate. For $f = (f_1, \dots, f_m) \in \mathbb{R}^m$ define

$$v_f = \sum_{j=1}^m f_j v_j.$$

Using the notation of Proposition 6.19 define (φ^f, g^f) , for f in a sufficiently small open ball $\hat{\mathcal{F}}$ about 0 in \mathbb{R}^m , to be a closed G_2 structure on M such that $\Xi([\varphi^f|_T]) = 0$ in $H_{cs}^3(\hat{N})$ and $\varphi^f|_{\hat{N}} = dv_f$. Recall from Definitions 6.24 and 7.12 that we have a linear map $L_3 : T_0 \hat{\mathcal{F}} \cong \mathbb{R}^m \rightarrow d(L_{k+1, \lambda}^p(\Lambda^2 T^* \hat{N}))$ arising from $dF_3|_{(0,0,0)}$. By construction, $L_3(f) = dv_f$ for $f \in \mathbb{R}^m$ and hence L_3 maps

onto $\mathcal{O}_1(N, \lambda)$. Proposition 7.3 and Definition 7.12 imply that $\mathcal{O}_3(N, \lambda) = \{0\}$ as required.

The latter parts of the theorem follow from the discussion preceding it and Proposition 8.8. \square

Acknowledgements Many thanks go to Dominic Joyce for his enormous help with this project. I would also like to thank Alexei Kovalev for useful comments and suggestions and EPSRC for providing the funding for this research.

References

- [1] R. Bartnik, *The Mass of an Asymptotically Flat Manifold*, Comm. Pure Appl. Math. **39**, 661-693, 1986.
- [2] R. L. Bryant, *Some Remarks on G_2 -Structures*, preprint, <http://www.arxiv.org/math.DG/0305124>, 2003
- [3] R. Harvey and H. B. Lawson, *Calibrated Geometries*, Acta Math. **148**, 47-152, 1982.
- [4] H. Federer, *Geometric Measure Theory*, Grundlehren Math. Wiss. **153**, Springer-Verlag, Berlin, 1969.
- [5] D. D. Joyce, *Compact Manifolds with Special Holonomy*, OUP, Oxford, 2000.
- [6] D. D. Joyce, *Special Lagrangian Submanifolds with Isolated Conical Singularities. I. Regularity*, Ann. Global Ann. Geom. **25**, 201-251, 2004.
- [7] D. D. Joyce, *Special Lagrangian Submanifolds with Isolated Conical Singularities. II. Moduli Spaces*, Ann. Global Ann. Geom. **25**, 301-352, 2004.
- [8] D. D. Joyce, *Special Lagrangian Submanifolds with Isolated Conical Singularities. III. Desingularization, The Unobstructed Case*, Ann. Global Ann. Geom. **26**, 1-58, 2004.
- [9] D. D. Joyce, *Special Lagrangian Submanifolds with Isolated Conical Singularities. IV. Desingularization, Obstructions and Families*, Ann. Global Ann. Geom. **26**, 117-174, 2004.
- [10] D. D. Joyce, *Special Lagrangian Submanifolds with Isolated Conical Singularities. V. Survey and Applications*, J. Differential Geom. **63**, 299-347, 2003.

- [11] S. Lang, *Differentiable Manifolds*, Addison–Wesley, Reading, Massachusetts, 1972.
- [12] S. Lang, *Real Analysis*, Second Edition, Addison–Wesley, Reading, Massachusetts, 1983.
- [13] R. B. Lockhart, *Fredholm, Hodge and Liouville Theorems on Noncompact Manifolds*, Trans. Amer. Math. Soc. **301**, 1-35, 1987.
- [14] R. B. Lockhart and R. C. McOwen, *Elliptic Differential Operators on Noncompact Manifolds*, Ann. Sc. Norm. Super. Pisa Cl. Sci. **12**, 409-447, 1985.
- [15] J. Lotay, *Deformation Theory of Asymptotically Conical Coassociative 4-folds*, preprint, <http://www.arxiv.org/math.DG/0411116>, 2004.
- [16] R. C. McLean, *Deformations of Calibrated Submanifolds*, Comm. Anal. Geom. **6**, 705-747, 1998.
- [17] S. P. Marshall, *Deformations of Special Lagrangian Submanifolds*, DPhil thesis, Oxford University, 2002.
- [18] C. B. Morrey, *Multiple Integrals in the Calculus of Variations*, Grundlehren Series Volume 130, Springer–Verlag, Berlin, 1966.
- [19] S. Salamon, *Riemannian Geometry and Holonomy Groups*, Pitman Research Notes in Mathematics **201**, Longman, Harlow, 1989.