# A certain class of standard subalgebras of affine Kac-Moody algebras

Bouaazza ES SAADI

Laboratoire de Mathématiques et Applications, Université de Haute Alsace, Mulhouse, France

> Email : Bouaazza.Es-saadi@uha.fr

#### Abstract

The aim of this paper is to extend the theory of standard subalgebras of finite dimensional simple Lie algebras to infinite dimensional Lie algebras. We construct and characterize a class of standard subalgebras of affine Kac-Moody algebra.

AMS Classification 2000 : 17B20, 17B30, 17B65, 17B67. Keywords : Lie algebras - Standard subalgebra - Kac-Moody algebra

# Introduction

Let  $\mathfrak{g}$  be a finite dimensional simple Lie algebra over  $\mathbb{C}$ , of rank p and let  $\mathfrak{h}$  be a Cartan subalgebra of  $\mathfrak{g}$ . We denote by  $\Delta$  the roots system of the couple  $(\mathfrak{g}, \mathfrak{h})$  and by  $\Pi = \{\alpha_1, \dots, \alpha_p\}$  the simple roots of  $\Delta$ .

Let A be a generalized Cartan matrix of affine type of order p + 1 associated to finite dimensional simple Lie algebra  $\mathfrak{g}$ .

Let S(A) (resp.  $S({}^{t}A)$ ) be the Dynkin diagram associated to A (resp.  ${}^{t}A$ ) and  $\mathfrak{B} = \{\alpha_0, \dots, \alpha_p\}$  (resp.  $\mathfrak{B}^{\vee} = \{\alpha_0^{\vee}, \dots, \alpha_p^{\vee}\}$ ) be the set of the vertex of diagram S(A) (resp.  $S({}^{t}A)$ ) called roots basis (resp. coroots basis). Let  $a_0, a_1, \dots, a_p$  be the numerical labels of S(A) and  $a_0^{\vee}, \dots, a_p^{\vee}$  be the numerical labels of  $S({}^{t}A)$ .

We denote by  $\mathcal{L} := \mathbb{C}[t, t^{-1}]$  the algebra of Laurent polynomials in t. Recall that the residue of a Laurent polynomial  $P = \sum_{i \in \mathbb{Z}} c_i t^i$  (where all but a finite number of  $c_i$ 

are 0) is defined by  $ResP = c_{-1}$ .

Consider the infinite dimensional Lie algebra, called the Loop algebra

$$\mathcal{L}(\mathfrak{g}) := \mathcal{L} \otimes \mathfrak{g}$$

We denote by [.,.] the bracket on  $\mathfrak{g}$ . The bracket on  $\mathcal{L}(\mathfrak{g})$  is defined as follows, for all  $(x, y) \in \mathfrak{g} \times \mathfrak{g} : [t^n \otimes x, t^m \otimes y]_{\mathcal{L}} = t^{n+m} \otimes [x, y].$ 

Fix a nondegenerate invariant symmetric bilinear  $\mathbb{C}$ -valued form (./.) on  $\mathfrak{g}$ . We extend this form by linearity to an  $\mathcal{L}$ -valued bilinear form  $(./.)_L$  on  $\mathcal{L}(\mathfrak{g})$  which is defined by

$$(P \otimes x/Q \otimes y)_L = PQ(x/y)$$

Now, we define a  $\mathbb{C}$ -valued 2-cocycle on the Lie algebra  $\mathcal{L}(\mathfrak{g})$  by

$$\psi(a,b) = Res(\frac{da}{dt},b)_t$$
 for all  $(a,b) \in \mathcal{L}(\mathfrak{g}) \times \mathcal{L}(\mathfrak{g})$ 

We denote by  $\tilde{\mathcal{L}}(\mathfrak{g})$  the extension of the Lie algebra  $\mathcal{L}(\mathfrak{g})$  by 1-dimensional center, associated to the cocycle  $\psi$ . Namely,  $\tilde{\mathcal{L}}(\mathfrak{g}) = \mathcal{L}(\mathfrak{g}) \oplus \mathbb{C}K$  where  $K = \sum_{i=0}^{i=p} a_i^{\vee} \alpha_i^{\vee}$  is a generator of the 1-dimensional center.

Finally, the affine Kac-Moody algebra associated to matrix A is

$$\mathfrak{g}(A) := \mathcal{L}(\mathfrak{g}) \oplus \mathbb{C} \mathrm{K} \oplus \mathbb{C} \mathrm{d}$$

where **d** is the derivation defined by :  $\mathbf{d} = t \frac{d}{dt}$  on  $\mathcal{L}(\mathbf{g}), \mathbf{d}(\mathbf{K}) = 0$ . For more details of this construction see [8, chap. 7].

The aim of this paper is to construct a certain class of standard subalgebras of affine Kac-Moody algebras  $\mathfrak{g}(A)$ . This uses standard subalgebras of the complex simple Lie algebra  $\mathfrak{g}$ .

Let  $\tau$  be a subalgebra of  $\mathfrak{g}$ . If the normalizer of  $\tau$  is a parabolic subalgebra of  $\mathfrak{g}$ then  $\tau$  is called standard subalgebra of g. Several papers were devoted to the study of standard subalgebras for finite dimensional case, see [6], [9], [13], [10], [11].

These subalgebras were characterized in [9] and [13] using a noncomparable roots with respect to the partial order relation "  $\leq$  " defined on the dual vector space  $\mathfrak{h}^*$ by  $\omega_1 \leq \omega_2$  if  $\omega_2 - \omega_1$  is linear combination of simple roots of  $\Pi$  with non-negative coefficients.

Let  $\mathfrak{R}$  be a set of positive roots two by two noncomparable. Consider the set  $\mathfrak{R}_1 = \{\beta \in \Delta_+ : \omega \leq \beta \text{ for certain } \omega \in \mathfrak{R}\}.$  Then the subalgebra  $\mathfrak{m} = \sum_{\beta \in \mathfrak{R}_1} \mathfrak{g}_\beta$  is

nilpotent standard subalgebra of  $\mathfrak{g}$ .

Setting  $\rho(\mathfrak{m})$  be the normalizer of  $\mathfrak{m}$ . Let  $\mathfrak{r}$  be an ideal of the Levi reductive subalgebra  $\mathfrak{h} + \sum_{\alpha \in \Omega_1} \mathfrak{g}_{\alpha}$ , lying in the Levi reductive subalgebra associated to parabolic subalgebra  $\rho(\mathfrak{m})$  where  $\Omega_1$  is the set of roots which are linear combination of  $\Pi \setminus (\Pi \cap$  $\mathfrak{R}$ ).

Any standard subalgebra  $\tau$  of  $\mathfrak{g}$  has the form

$$\tau = \mathfrak{m} + \mathfrak{r}$$

and the normalizer  $\rho(\tau) = \rho(\mathfrak{m})$ .

In this paper, theorem 1.1 states that there exists standard subalgebra  $\overline{\tau}$  of  $\mathfrak{g}(A)$ associated to a given standard subalgebra  $\tau$  of finite dimensional simple Lie algebra  $\mathfrak{g}$  and a certain vector subspace  $V_{\tau} = [\tau, \mathfrak{g}]$  of  $\mathfrak{g}$  associated to  $\tau$ . The standard subalgebra  $\overline{\tau}$  has the form

$$\overline{\tau} = t^n \otimes \tau + t^{n+1} \otimes V_{\tau} + t^{n+2} \mathbb{C}[t] \otimes \mathfrak{g} + \mathbb{C}K \quad \text{for } n \in \mathbb{N}^*$$

The subspace  $V_{\tau}$  is called "appui" subspace of  $\mathfrak{g}$ .

The standard subalgebra  $\overline{\tau}$  is determined by the finite dimensional standard subalgebra  $\tau$  and the appui subspace  $V_{\tau}$ . Explicit formulas of  $V_{\tau}$  are established for different cases of  $\tau$ .

In the case where  $\tau$  is nilpotent standard subalgebra. Then,  $\tau$  is associated to subset  $\mathfrak{R} \subset \Delta_+$  of positive roots two by two noncomparable. The theorem 2.1 states that the appui subspace  $V_{\tau}$  has the following form :

$$V_{\tau} = \sum_{\alpha \in \Delta_+ \backslash < \mathfrak{R}_3 >^+} \mathfrak{g}_{\alpha} + \sum_{\alpha \in \mathfrak{R}_1 \cup \mathfrak{R}_2} \mathbb{C}[e_{\alpha}, e_{-\alpha}] + \sum_{\alpha \in \mathfrak{R}_2} \mathfrak{g}_{-\alpha}$$

where  $\mathfrak{R}_1, \mathfrak{R}_2$  and  $\mathfrak{R}_3$  are defined by the relation 1, 3 and 4.

In the second case where  $\tau$  is standard subalgebra of nilpotent radical. Then,  $\tau = \mathfrak{m} + \mathfrak{r}_0$  where  $\mathfrak{m}$  is nilpotent part of  $\tau$  and  $\mathfrak{r}_0$  is semisimple part of  $\tau$ . Therefor,  $\mathfrak{r}_0$  is constructed from a certain common connected subsystem  $\psi$  of  $\Pi \setminus (\Pi \cap \mathfrak{R})$  and  $\Pi \setminus (\bigcup_{\omega \in \mathfrak{R}} S^{\omega})$  where  $S^{\omega} = \{\gamma \in \Pi : \omega = \gamma \text{ or } \omega - \gamma \in \Delta\}$ . One may write  $\mathfrak{r}_o = \sum_{\alpha \in \langle \Psi \rangle} \mathfrak{g}_{\alpha} + \sum_{\alpha \in \langle \Psi \rangle} \mathbb{C}[e_{\alpha}, e_{-\alpha}]$  where  $\langle \psi \rangle$  is the set of roots which are linear combination of elements of  $\psi$ . We prove in theorem 2.2 that the appui subspace  $V_{\tau}$  has the following form :

$$V_{\tau} = V_{\mathfrak{m}} \quad \text{if} \quad P_{\Psi}^{-} = \varnothing$$
$$V_{\tau} = \mathfrak{g} \quad \text{if} \quad P_{\Psi}^{-} \neq \varnothing$$

where  $P_{\Psi}^{-} = \sum_{\alpha \in \langle \Psi \rangle} \sum_{\beta \in \Delta_{+} \setminus \Re_{2}} [\mathfrak{g}_{\alpha}, \mathfrak{g}_{-\beta}]$  and  $V_{\mathfrak{m}}$  is the appui subspace associated to  $\mathfrak{m}$ .

# 1 Standard subalgebras of affine Kac-Moody algebra

Let  $\mathfrak{g}$  be a simple Lie algebra of rank p. Let A be its extended Cartan matrix of affine type and  $\mathfrak{g}(A)$  be the affine Kac-Moody algebra associated to A. We consider the element  $\delta$  defined by

$$\delta = \sum_{i=0}^{i=p} a_i \alpha_i$$

Let  $\Delta^{aff} = \Delta^{re} \cup \Delta^{im}$  be a roots system of the couple  $(\mathfrak{g}(A), \mathfrak{h} + \mathbb{C}K + \mathbb{C}d)$  where  $\Delta^{re} = \{\alpha + j\delta : \alpha \in \Delta \text{ and } j \in \mathbb{Z}\}$  is the set of real roots and  $\Delta^{im} = \{j\delta : j \in \mathbb{Z}\}$  is the set of imaginary roots.

We denote by [.,.] the bracket on  $\mathfrak{g}$ . The bracket  $[.,.]_{aff}$  on  $\mathfrak{g}(A)$  is defined as follows : for all  $(x, y) \in \mathfrak{g} \times \mathfrak{g}$ ,  $(\lambda, \mu, \lambda_1, \mu_1) \in \mathbb{C}^4$ ,  $(n, m) \in \mathbb{Z}^2$ :

$$[t^n \otimes x + \lambda \mathbf{K} + \mu \mathbf{d}, t^m \otimes y + \lambda_1 \mathbf{K} + \mu_1 \mathbf{d}]_{aff} = t^{n+m} \otimes [x, y] + \mu m t^m \otimes y - \mu_1 n t^n \otimes x + n \delta_{n, -m}(x|y) \mathbf{K} + \mu_1 \mathbf{d} \mathbf{d}_{n, -m}(x|y) \mathbf$$

In the particular case where m and n are two natural numbers, we have :

$$[t^n \otimes x + \lambda \mathbf{K} + \mu \mathbf{d}, t^m \otimes y + \lambda_1 \mathbf{K} + \mu_1 \mathbf{d}]_{aff} = t^{n+m} \otimes [x, y] + \mu m t^m \otimes y - \mu_1 n t^n \otimes x$$

**Definition 1** We introduce the following definitions :

1. A subalgebra of  $\mathfrak{g}(A)$  is called parabolic if it contains a Borel subalgebra of  $\mathfrak{g}(A)$ .

2. A subalgebra of  $\mathfrak{g}(A)$  is called standard if its normalizer is a parabolic subalgebra of  $\mathfrak{g}(A)$ .

Let  $\tau$  be a standard subalgebra of  $\mathfrak{g}$  of normalizer  $\rho(\tau)$  and let  $V_{\tau} = [\tau, \mathfrak{g}]$  be the associated subspace.

**Proposition 1.1** The appui subspace  $V_{\tau}$  satisfies the following relation :  $[V_{\tau}, \rho(\tau)] \subset V_{\tau}$ .

**Proof**: Using the Jacobi identity, we have  $[V_{\tau}, \rho(\tau)] = [[\tau, \mathfrak{g}], \rho(\tau)] = [\tau, [\mathfrak{g}, \rho(\tau)]] + [[\tau, \rho(\tau)], \mathfrak{g}] \subset [\tau, \mathfrak{g}] + [\tau, \mathfrak{g}] \subset [\tau, \mathfrak{g}] = V_{\tau}.$ 

The following main theorem gives a sufficient condition for the existence of a class of standard subalgebras of  $\mathfrak{g}(A)$ .

**Theorem 1.1** Let  $\tau$  be a standard subalgebra of  $\mathfrak{g}$  of normalizer  $\rho(\tau)$  and  $V_{\tau} = [\tau, \mathfrak{g}]$  be the appui subspace associated to  $\tau$ .

Then, the subalgebra

$$\overline{\tau} = t^n \otimes \tau + t^{n+1} \otimes V_\tau + t^{n+2} \mathbb{C}[t] \otimes \mathfrak{g} + \mathbb{C}K \ (n \in \mathbb{N}^*)$$

is a standard subalgebra of  $\mathfrak{g}(A)$  with normalizer  $\rho(\overline{\tau}) = \rho(\tau) + t\mathbb{C}[t] \otimes \mathfrak{g} + \mathbb{C}K + \mathbb{C}d$ .

**Proof**: We show that the parabolic subalgebra  $\rho(\overline{\tau})$  is the normalizer of  $\overline{\tau}$ .

In  $\mathfrak{g}(A)$ , the subalgebra  $\overline{\tau}$  is an ideal of parabolic subalgebra  $\rho(\tau) + t\mathbb{C}[t] \otimes \mathfrak{g} + \mathbb{C}K + \mathbb{C}d$ .

Let  $\rho_1$  be a parabolic subalgebra of  $\mathfrak{g}$  such that  $\rho_1 + t\mathbb{C}[t] \otimes \mathfrak{g} + \mathbb{C}K + \mathbb{C}d$  is a normalizer of  $\overline{\tau}$ . We have  $[\overline{\tau}, \rho_1 + t\mathbb{C}[t] \otimes \mathfrak{g} + \mathbb{C}K + \mathbb{C}d]_{aff} \subset \overline{\tau}$  then  $[\tau, \rho_1] \subset \tau$ and  $\rho_1 \subset \rho(\tau)$ . Since  $\rho_1 + t\mathbb{C}[t] \otimes \mathfrak{g} + \mathbb{C}K + \mathbb{C}d$  is a normalizer of  $\overline{\tau}$  then, we have  $\rho_1 + t\mathbb{C}[t] \otimes \mathfrak{g} + \mathbb{C}K + \mathbb{C}d$  included in  $\rho(\tau) + t\mathbb{C}[t] \otimes \mathfrak{g} + \mathbb{C}K + \mathbb{C}d$  and  $\rho(\tau)$  included in  $\rho_1$ . Furthermore,  $\rho_1 = \rho(\tau)$  and  $\rho(\tau) + t\mathbb{C}[t] \otimes \mathfrak{g} + \mathbb{C}K + \mathbb{C}d$  is a normalizer of  $\overline{\tau}$ .

Corollary 1.1 The subalgebra

 $\tau + t \otimes \mathfrak{g} + t^2 \mathbb{C}[t] \otimes \mathfrak{g} + \mathbb{C}K + \mathbb{C}d$ 

is a standard subalgebra of  $\mathfrak{g}(A)$  of normalizer  $\rho(\tau) + t\mathbb{C}[t] \otimes \mathfrak{g} + \mathbb{C}K + \mathbb{C}d$ .

**Corollary 1.2** Let V be a subspace of  $\mathfrak{g}$  such that  $V_{\tau} \subset V$  and  $[V, \rho(\tau)] \subset V$  then the subalgebra

$$t^n \otimes \tau + t^{n+1} \otimes V + t^{n+2} \mathbb{C}[t] \otimes \mathfrak{g} + \mathbb{C}K \ (n \in \mathbb{N}^*)$$

is standard subalgebra of  $\mathfrak{g}(A)$ .

**Remark 1.1** We deduce from the theorem 1.1 that the subalgebra  $\overline{V}_{\tau} = t^{n+1} \otimes$  $V_{\tau} + t^{n+2}\mathbb{C}[t] \otimes \mathfrak{g} + \mathbb{C}K$  is also standard subalgebra of  $\mathfrak{g}(A)$ .

**Eexample 1.1** Let  $\mathfrak{g}$  be a simple Lie algebra of type  $B_4$  and  $\Pi = \{\alpha_1, \alpha_2, \alpha_3, \alpha_4\}$  be a basis of roots system  $\Delta$ .

We set  $\mathfrak{R}_1 = \{\alpha_2 + 2\alpha_3 + 2\alpha_4, \alpha_1 + \alpha_2 + 2\alpha_3 + 2\alpha_4, \alpha_1 + 2\alpha_2 + 2\alpha_3 + 2\alpha_4\}$ . We consider  $\tau = \sum_{\alpha \in \mathfrak{R}_1} \mathfrak{g}_{\alpha}$  and  $V_{\tau} = \sum_{\alpha \in \Delta_+ \setminus \{\alpha_4\}} \mathfrak{g}_{\alpha} + \mathfrak{h} + \mathfrak{g}_{-\alpha_1} + \mathfrak{g}_{-\alpha_2} + \mathfrak{g}_{-\alpha_1-\alpha_2}$  respectively the standard subalgebra of  $\mathfrak{g}$  and the appui subspace associated to  $\tau$ .

Then  $\overline{\tau} = t \otimes \tau + t^2 \otimes V + t^3 \mathbb{C}[t] \otimes \mathfrak{g} + \mathbb{C}K$  is a standard subalgebra of  $\mathfrak{g}(A)$  with normalizer  $\rho(\overline{\tau}) = (\sum_{\alpha \in \Delta_+} \mathfrak{g}_{\alpha} + \mathfrak{h} + \mathfrak{g}_{-\alpha_1} + \mathfrak{g}_{-\alpha_2} + \mathfrak{g}_{-\alpha_1-\alpha_2} + \mathfrak{g}_{-\alpha_4}) + t\mathbb{C}[t] \otimes \mathfrak{g} + \mathbb{C}\mathbf{K} + \mathbb{C}\mathbf{d}.$ 

The standard subalgebra  $\overline{\tau}$  may be written using the real and imaginary roots as follows :

$$\overline{\tau} = \sum_{\alpha \in \mathfrak{R}_1} \mathfrak{g}(A)_{\alpha+\delta} + \sum_{\alpha \in \mathfrak{R}_\tau} \mathfrak{g}(A)_{\alpha+2\delta} + \mathfrak{g}(A)_{2\delta} + \sum_{n \geqslant 3} \{\sum_{\alpha \in \Delta} \mathfrak{g}(A)_{\alpha+n\delta} + \mathfrak{g}(A)_{n\delta}\} + \mathbb{C}\mathrm{K}$$

with  $\mathfrak{R}_{\tau} = \Delta_{+} \setminus \{\alpha_{4}\} \cup \{-\alpha_{1}, -\alpha_{2}, -\alpha_{1} - \alpha_{2}\}, \mathfrak{g}(A)_{\alpha + n\delta} = t^{n} \otimes \mathfrak{g}_{\alpha} \text{ and } \mathfrak{g}(A)_{n\delta} = t^{n} \otimes \mathfrak{h}$ 

Theorem 1.1 proves that the existence of this class depends on the description of the standard subalgebras  $\tau$  and the appui subspaces  $V_{\tau}$  of the Lie algebra g.

In the next section, we characterize the appui subspaces  $V_{\tau}$  when  $\tau$  is the nilpotent standard subalgebra then when  $\tau$  is the standard subalgebra.

#### 2 Appui subspaces

Let  $\mathfrak{g}$  be a complex simple Lie algebra,  $\mathfrak{h}$  be its Cartan subalgebra and  $\Delta$  be the roots system of the couple  $(\mathfrak{g}, \mathfrak{h})$ . Let  $\tau$  be the standard subalgebra of  $\mathfrak{g}$ . The aim of this section is to give an explicit formulas for the appui subspaces  $V_{\tau} = [\tau, \mathfrak{g}]$ .

#### 2.1Case of nilpotent standard subalgebra

#### Main notations.

1. For any positive root  $\beta$ , we denote by  $S^{\beta}$  the set of simple roots  $\gamma$  of  $\Pi$  such that  $\beta = \gamma$  or  $\beta - \gamma \in \Delta$ . The set  $S^{\beta}$  is called set of extremal roots of  $\beta$ .

2. For any root  $\alpha = \sum_{\gamma \in \Pi} \alpha_{\gamma} \gamma$ , we denote by  $C_{\alpha}$  the set of the simple roots  $\gamma$  in  $\Pi$ such that  $\alpha_{\gamma} \neq 0$ .

3. Let B be the subset of the simple roots  $\Pi$ . We denote by  $\langle B \rangle$  the set of roots which are linear combination of elements of B and by  $\langle B \rangle^+$  (resp.  $\langle B \rangle^-$ ) the positive roots (resp. the negative roots) of  $\langle B \rangle$ .

Let  $\tau$  be a nilpotent standard subalgebra of  $\mathfrak{g}$  associated to the subsystem  $\mathfrak{R}$  of positive roots two by two non comparable. We set

$$\mathfrak{R}_{1} = \{ \beta \in \Delta_{+} : \quad \omega \leqslant \beta \quad \text{for a certain } \omega \in \mathfrak{R} \}$$
(1)

The subalgebra  $\tau$  has the form  $\tau = \sum_{\beta \in \mathfrak{R}_1} \mathfrak{g}_{\beta}$ .

We set  $S_2 = \bigcup_{\omega \in \mathfrak{R}} S^{\omega}$ ,  $\Delta_1 = \langle \Pi \backslash S_2 \rangle$  and  $\Delta_2 = \Delta \backslash \Delta_1$ . The normalizer of  $\tau$  is defined by  $S_2$  and has the following form  $\rho(\tau) = \sum_{\alpha \in \Delta_+} \mathfrak{g}_{\alpha} + \mathfrak{h} + \sum_{\alpha \in \Delta_+^+} \mathfrak{g}_{-\alpha}$ .

We can write  $\mathfrak{g} = \rho(\tau) + \sum_{\nu \in \Delta_2^+} \mathfrak{g}_{-\nu}$ . Then, we have

$$V_{\tau} = [\tau, \mathfrak{g}] \subset \tau + \sum_{\beta \in \mathfrak{R}_1} \sum_{\nu \in \Delta_2^+} [\mathfrak{g}_{\beta}, \mathfrak{g}_{-\nu}].$$
<sup>(2)</sup>

**Lemma 2.1** Let  $\alpha$  be a positive root. Then  $\mathfrak{g}_{\alpha}$  is included in  $V_{\tau}$  if and only if there exists  $\beta \in \mathfrak{R}_1$  such that  $C_{\alpha} \cap S^{\beta} \neq \emptyset$ .

**Proof**:  $(\Longrightarrow)$  Let  $\alpha \in \Delta_+$  such that  $\mathfrak{g}_{\alpha} \subset V_{\tau}$ . There are two cases : First case :  $\alpha \in \mathfrak{R}_1$  then there exists  $\beta \in \mathfrak{R}$  such that  $\beta \leq \alpha$ . This implies that  $C_{\alpha} \cap S^{\beta} \neq \emptyset$ .

Second case :  $\alpha \notin \Re_1$ , using the formula 2, there exists a root  $\beta \in \Re_1$  such that  $\beta - \alpha$  is a root of  $\Delta_2^+$ . One we can write  $\alpha$  in the form  $\alpha = \alpha_1 + \cdots + \alpha_k$  with  $\alpha_1, \cdots, \alpha_k$  simple roots and each partial sum  $\alpha_1 + \cdots + \alpha_l$  is a root, for  $1 \leq l \leq k$ . We want to prove by induction on k that there exists  $i_0 \in [\![1, k]\!]$  such that  $\alpha_{i_0}$  is in  $S^{\beta}$ .

If k = 1. We have  $\beta - \alpha_1 \in \Delta$ . Then  $\alpha_1 \in S^{\beta}$ . Suppose that k > 1, then we have  $-\alpha'_{k-1} - \alpha_k + \beta$  is a root where  $\alpha'_{k-1} = \alpha_1 + \cdots + \alpha_{k-1}$ . By applying Jacobi identity to the roots  $-\alpha'_{k-1}, -\alpha_k$  and  $\beta$ , we have  $\beta - \alpha'_{k-1}$  or  $\beta - \alpha_k$  is a root.

If  $\beta - \alpha'_{k-1} \in \Delta$ , by the assumption of recurrence, there exists  $j_0 \in [\![1, k-1]\!]$  such that  $\alpha_{j_0} \in S^{\beta}$  and  $C_{\alpha} \cap S^{\beta} \neq \emptyset$ .

If  $\beta - \alpha_k \in \Delta$ , we have  $\alpha_k \in S^{\beta}$  and  $C_{\alpha} \cap S^{\beta} \neq \emptyset$ .

( $\Leftarrow$ ) Let  $\alpha$  be a root of  $\Delta_+$  such that there exists  $\beta$  in  $\Re_1$  with  $C_{\alpha} \cap S^{\beta} \neq \emptyset$ . We set  $\alpha = \alpha_1 + \cdots + \alpha_n$  such that for all  $i \in [\![1, n]\!]$ ,  $\alpha_i \in \Pi$  and for  $r \in [\![1, n]\!]$  each partial sum  $\alpha_1 + \cdots + \alpha_r$  is a root.

Let J be a subset of  $\llbracket 1, n \rrbracket$  such that  $\alpha_j \in C_\alpha \cap S^\beta$  for any  $j \in J$ . It is clear that the subset J is not empty. Let q be an element of J. We have  $\beta - \alpha_q$  is a root and  $\mathfrak{g}_{\alpha_q} = [\mathfrak{g}_\beta, \mathfrak{g}_{-\beta+\alpha_q}] \subset [\tau, \mathfrak{g}] = V_\tau$ .

Using the proposition 1.1, we may deduce that

$$\mathfrak{g}_{\alpha} = \mathfrak{g}_{\alpha_1 + \dots + \alpha_n} = [[\mathfrak{g}_{\alpha_q}, \mathfrak{g}_{\alpha_1 + \dots + \alpha_{q-1}}], \mathfrak{g}_{\alpha_{q+1}}, \cdots, \mathfrak{g}_{\alpha_n}] \subset [V_{\tau}, \rho(\tau)] \subset V_{\tau}$$

**Lemma 2.2** Let  $\alpha$  be a positive root. Then  $\mathfrak{g}_{-\alpha}$  is included in  $V_{\tau}$  if and only if there exists  $\beta \in \mathfrak{R}_1$  such that  $\alpha + \beta \in \Delta_+$ .

**Proof**:  $(\Longrightarrow)$  Let  $\alpha \in \Delta_+$  such that  $\mathfrak{g}_{-\alpha} \subset V_{\tau}$ . Then, by relation 2, there exists a root  $\beta \in \mathfrak{R}_1$  such that  $\alpha + \beta$  is a root of  $\Delta_2^+$ .

 $(\Leftarrow)$  Let  $\beta$  be a root in  $\mathfrak{R}_1$  such that  $\alpha + \beta$  is a root. Then we have  $\mathfrak{g}_{-\alpha} = [\mathfrak{g}_{\beta}, \mathfrak{g}_{-\beta-\alpha}] \subset [\tau, \mathfrak{g}] = V_{\tau}$ .

We consider the set  $\mathfrak{R}_2$  of  $\Delta$  satisfying the following property :

If 
$$\alpha \in \Delta_+, \beta \in \mathfrak{R}_1$$
, and  $\alpha + \beta \in \Delta$  then  $\alpha \in \mathfrak{R}_2$  (3)

And the set  $\mathfrak{R}_3$  of  $\Pi$  satisfying the following property :

If 
$$\alpha \in \Pi, \beta \in \mathfrak{R}_1$$
, and  $\beta - \alpha \notin \Delta$  then  $\alpha \in \mathfrak{R}_3$  (4)

In other words  $\mathfrak{R}_3 = \Pi \setminus S_2$ 

The next theorem characterizes the appui subspace  $V_{\tau}$  when  $\tau$  is a nilpotent standard subalgebra of  $\mathfrak{g}$ .

**Theorem 2.1** Let  $\mathfrak{g}$  be a complex simple Lie algebra and  $\tau$  be a nilpotent standard subalgebra associated to a subsystem  $\mathfrak{R}$  of positive roots two by two non comparable. Then, the appui subspace associated to  $\tau$  has the form

$$V_{\tau} = \sum_{\alpha \in \Delta_+ \backslash < \mathfrak{R}_3 >^+} \mathfrak{g}_{\alpha} + \sum_{\alpha \in \mathfrak{R}_1 \cup \mathfrak{R}_2} \mathbb{C}[e_{\alpha}, e_{-\alpha}] + \sum_{\alpha \in \mathfrak{R}_2} \mathfrak{g}_{-\alpha}$$

**Proof**: By applying the lemma 2.1 and 2.2, it is enough to prove that  $\sum_{\alpha \in \Re_1 \cup \Re_2} \mathbb{C}[e_{\alpha}, e_{-\alpha}]$  is included in  $V_{\tau}$ .

If  $\alpha \in \mathfrak{R}_1$  then  $\mathbb{C}[e_{\alpha}, e_{-\alpha}] \subset [\tau, \mathfrak{g}] = V_{\tau}$ . If  $\alpha \in \mathfrak{R}_2$  then, we have  $\mathfrak{g}_{-\alpha}$  is included in  $V_{\tau}$  and by applying the proposition 1.1, we have  $\mathbb{C}[e_{-\alpha}, e_{\alpha}] \subset [V_{\tau}, \rho(\tau)] \subset V_{\tau}$ .

**Remark 2.1** We denote by  $\theta$  the highest root of  $\Delta$ . If the  $\Re$  is included in basis  $\Pi$ , the nilpotent standard subalgebra  $\tau$  is called complete standard subalgebra of  $\mathfrak{g}$  and it is easy to prove that the appui subspace  $V_{\tau}$  has the form

$$V_{\tau} = \sum_{\alpha \in \Delta_+} \mathfrak{g}_{\alpha} + \mathfrak{h} + \sum_{\alpha \in \Delta_+ \backslash \mathfrak{R}_C} \mathfrak{g}_{-\alpha}$$

with  $\mathfrak{R}_C = \{ \alpha \in \Delta_+ : \text{ for all root } \beta \in \mathfrak{R} , \alpha_\beta = \theta_\beta \}.$ 

**Remark 2.2** We consider  $m = \sum_{\alpha \in \Delta} \mathfrak{g}_{\alpha}$  the nilpotent standard subalgebra associated to the basis  $\Pi$ . It is the maximal ad-nilpotent ideal of Borel subalgebra of  $\mathfrak{g}$ . Then, the appui subspace is

$$V_{\tau} = \sum_{\alpha \in \Delta_{+}} \mathfrak{g}_{\alpha} + \mathfrak{h} + \sum_{\alpha \in \Delta_{+} \setminus \{\theta\}} \mathfrak{g}_{-\alpha}$$

### 2.2 General case

In this section, we establish an explicit formula for the appui subspaces  $V_{\tau}$  when  $\tau$  is a standard subalgebra of  $\mathfrak{g}$  not necessary nilpotent.

Let  $\tau = \mathfrak{m} + \mathfrak{r}_0$  be a standard subalgebra of  $\mathfrak{g}$  of nilpotent radical where  $\mathfrak{m}$  is its nilpotent part and  $\mathfrak{r}_0$  is its semisimple part. The nilpotent part  $\mathfrak{m}$  of  $\tau$  is a nilpotent standard subalgebra of  $\mathfrak{g}$ , then  $\mathfrak{m}$  is associated to a certain subset  $\mathfrak{R}$  of positive roots two by two noncomparable.

We recall that  $S_1 = \Re \cap \Pi$ ,  $S_2 = \bigcup_{\omega \in \Re} S^{\omega}$  and  $\Delta_1 = \langle \Pi \setminus S_2 \rangle$ . We set  $\Omega_1 = \langle \Pi \setminus S_1 \rangle$ . We have  $\mathfrak{r}_0$  an ideal of the Levi reductive subalgebra  $\mathfrak{r}_1 = \mathfrak{h} + \sum_{\alpha \in \Omega_1} \mathfrak{g}_{\alpha}$  lying in the Levi reductive subalgebra  $\mathfrak{r}_2 = \mathfrak{h} + \sum_{\alpha \in \Delta_1} \mathfrak{g}_{\alpha}$ . Since  $\mathfrak{r}_0$  is semisimple algebra, then there exists  $\Psi$  a common connected subsystem of  $\Pi \setminus S_1$  and  $\Pi \setminus S_2$  such that  $\mathfrak{r}_{\mathfrak{o}} = \sum_{\alpha \in \langle \Psi \rangle} \mathfrak{g}_{\alpha} + \sum_{\alpha \in \langle \Psi \rangle} \mathbb{C}[e_{\alpha}, e_{-\alpha}].$ 

Setting the subspace  $P_{\Psi}^{-} = \sum_{\alpha \in \langle \Psi \rangle} [\mathfrak{g}_{\alpha}, \mathfrak{n}_{2}^{-}]$  and the subspace  $P_{\Psi}^{+} = \sum_{\alpha \in \langle \Psi \rangle} [\mathfrak{g}_{\alpha}, \mathfrak{n}_{2}^{+}]$ where  $\mathfrak{n}_{2}^{+} = \sum_{\beta \in \Delta_{+} \setminus \mathfrak{R}_{2}} \mathfrak{g}_{\beta}$  and  $\mathfrak{n}_{2}^{-} = \sum_{\beta \in \Delta_{+} \setminus \mathfrak{R}_{2}} \mathfrak{g}_{-\beta}$ .

Using the properties of  $\mathfrak{m}$  and  $\mathfrak{r}_0$ , one proves that

$$V_{\tau} = V_{\mathfrak{m}} + P_{\Psi}^{-} + [h_{\Psi}, \mathfrak{n}_{2}^{-}]$$

with  $h_{\Psi} = \sum_{\alpha \in \langle \Psi \rangle} \mathbb{C}[e_{\alpha}, e_{-\alpha}]$ .

**Lemma 2.3** If  $\Psi$  is a common connected component of  $\Pi \setminus S_1$  and  $\Pi \setminus S_2$  then  $P_{\Psi}^+$  is included in  $\mathfrak{n}_2^+$ .

**Proof**: Let  $\alpha$  be a root such that  $\mathfrak{g}_{\alpha} \subset P_{\psi}^+$ . Then, there exists  $\gamma \in \langle \psi \rangle^+$  and  $\beta \in \Delta_+ \backslash \mathfrak{R}_2$  such that  $\alpha = \pm \gamma + \beta \in \Delta_+$ . We want to prove that  $\mathfrak{g}_{\alpha} \subset \mathfrak{n}_2^+$ .

We assume that  $\alpha$  is a root of  $\Re_2$ . Then, there exists  $\omega \in \Re_1$  such that  $\pm \gamma + \beta + \omega \in \Delta$ . Applying the Jacobi identity to  $\pm \gamma$ ,  $\beta$  and  $\omega$ , we have two cases : First case : if  $\pm \gamma + \omega \in \Delta$ , since  $\pm \gamma \in \langle \psi \rangle \subset \Delta_1$  then  $\pm \gamma + \omega \in \Re_1$ . We have  $\pm \gamma + \omega + \beta$  then  $\beta \in \Re_2$ . Therefor, we have a contradiction.

Second case : if  $\beta + \omega \in \Delta$  then  $\beta \in \mathfrak{R}_2$ . Therefor, we have a contradiction. Finally,  $\alpha$  is a root of  $\Delta_+ \setminus \mathfrak{R}_2$  and  $\mathfrak{g}_\alpha \subset \mathfrak{n}_2^+$ .

**Proposition 2.1** Let  $\alpha$  and  $\beta$  be two roots of  $\Delta_+ \setminus \Re_2$  such that  $\alpha + \beta \in \Delta_+$ . 1. If  $\mathfrak{g}_{\alpha+\beta} \subset P_{\Psi}^+$  then  $\mathfrak{g}_{\beta} \subset P_{\Psi}^+$  and  $\mathfrak{g}_{\beta} \subset P_{\Psi}^+$ . 2. If  $\mathfrak{g}_{\alpha} \subset P_{\psi}^+$  or  $\mathfrak{g}_{\beta} \subset P_{\psi}^+$  then  $\mathfrak{g}_{\alpha+\beta} \subset P_{\psi}^+$ .

**Proof**: We have  $\alpha + \beta \in \Delta_+ \setminus \mathfrak{R}_2$  then, it is necessary that  $\alpha \in \langle \Pi \setminus S_1 \rangle$  and  $\beta \in \langle \Pi \setminus S_1 \rangle$  because if no, we have  $\alpha$  and  $\beta$  are two roots of  $\mathfrak{R}_1$ . So, by definition of the set  $\mathfrak{R}_2$ , we have  $\alpha$  and  $\beta$  in  $\mathfrak{R}_2$ , contradiction.

1. We have  $\mathfrak{g}_{\alpha+\beta} \subset P_{\psi}^+$  then, there exists  $\gamma_0 \in \langle \Psi \rangle^+$  such that  $\alpha + \beta \pm \gamma_0 \in \Delta_+ \backslash \mathfrak{R}_2$ . Applying the Jacobi identity to roots  $\alpha$ ,  $\beta$  and  $\pm \gamma_0$ , we obtain two cases :

First case : if  $\alpha \pm \gamma_0 \in \Delta$  then  $\alpha \pm \gamma \in \langle \psi \rangle$  because  $\psi$  is the connected subsystem of  $\Pi \setminus S_1$ . Therefore, we have  $[\mathfrak{g}_{-\alpha \mp \gamma_0}, \mathfrak{g}_{\beta+\alpha\pm\gamma_0}] = \mathfrak{g}_{\beta} \subset P_{\psi}^+$  and  $[\mathfrak{g}_{\mp\gamma_0}, \mathfrak{g}_{\alpha\pm\gamma_0}] = \mathfrak{g}_{\alpha} \subset P_{\psi}^+$ . Second case : if  $\beta \pm \gamma_0 \in \Delta$  then  $\beta \pm \gamma_0 \in \langle \psi \rangle$  because  $\psi$  is the connected subsystem of  $\Pi \setminus S_1$ . Therefor, we have  $[\mathfrak{g}_{-\beta\mp\gamma_0}, \mathfrak{g}_{\alpha+\beta\pm\gamma_0}] = \mathfrak{g}_{\alpha} \subset P_{\psi}^+$  and  $[\mathfrak{g}_{\mp\gamma_0}, \mathfrak{g}_{\beta\pm\gamma_0}] = \mathfrak{g}_{\beta} \subset P_{\psi}^+$ .

2. Since  $\mathfrak{g}_{\alpha} \subset P_{\psi}^{+}$  then, there exists  $\alpha_{0} \in \Delta \backslash \mathfrak{R}_{2}$  and  $\omega_{0} \in \langle \psi \rangle^{+}$  such that  $\alpha = \alpha_{0} \pm \omega_{0}$ . Applying the Jacobi identity to  $\alpha_{0}, \pm \omega_{0}$  and  $\beta$ , we have two cases : First case : if  $\alpha_{0} + \beta \in \Delta_{+} \backslash \mathfrak{R}_{2}$  then  $\mathfrak{g}_{\alpha+\beta} \subset P_{\psi}^{+}$ . Second case : if  $\beta \pm \omega_{0} \in \Delta$  then  $\beta \pm \omega_{0} \in \langle \psi \rangle$  and  $\mathfrak{g}_{\alpha+\beta} \subset P_{\psi}^{+}$ . If  $\mathfrak{g}_{\beta} \subset P_{\psi}^{+}$ , we remplace  $\alpha$  by  $\beta$  and with the same reason we prove that  $\mathfrak{g}_{\alpha+\beta} \subset P_{\psi}^{+}$ .

The next theorem gives the explicit formula of appui subspace  $V_{\tau}$  associated to standard subalgebra  $\tau$  of  $\mathfrak{g}$ .

**Theorem 2.2** Let  $\mathfrak{g}$  be a complex simple Lie algebra and  $\tau = \mathfrak{m} + \mathfrak{r}_{\mathfrak{o}}$  be a standard subalgebra of  $\mathfrak{g}$  of nilpotent radical, where  $\mathfrak{m}$  is a nilpotent standard subalgebra associated to a subsystem  $\mathfrak{R}$  of positive roots two by two non comparable and  $\mathfrak{r}_0$  is an ideal of  $\mathfrak{r}_1$  contained in  $\mathfrak{r}_2$ .

Then, we have :

1.  $V_{\tau} = V_{\mathfrak{m}}$  if  $P_{\Psi}^{-} = \varnothing$ . 2.  $V_{\tau} = \mathfrak{g}$  if  $P_{\Psi}^{-} \neq \varnothing$ .

### **Proof** :

1. If  $P_{\Psi}^{-} = \emptyset$  then,  $[h_{\Psi}, \mathfrak{n}_{2}^{-}] = \emptyset$ . Therefore,  $V_{\tau} = [\tau, \mathfrak{g}] = V_{\mathfrak{m}}$  and we have the result.

2. If  $P_{\Psi}^{-} \neq \emptyset$ . We have  $V_{\tau} = V_{\mathfrak{m}} + P_{\Psi}^{-} + [h_{\Psi}, \mathfrak{n}_{2}^{-}]$  and  $\mathfrak{g} = V_{\mathfrak{m}} + \mathfrak{n}_{2}^{-}$ . So, it is enough to show that  $P_{\Psi}^{-} = \mathfrak{n}_{2}^{-}$ . It is the same to show that  $P_{\Psi}^{+} = \mathfrak{n}_{2}^{+}$ . By using the lemma 2.3, we have  $P_{\Psi}^{+} \subset \mathfrak{n}_{2}^{+}$ , then it left to prove that  $\mathfrak{n}_{2}^{+} \subset P_{\Psi}^{+}$ .

We consider  $\theta$  the highest root of  $\Delta$ . In the first, we prove that  $\mathfrak{g}_{\theta} \subset P_{\psi}^+$ . We have  $P_{\Psi}^+ \neq \emptyset$  then, there exists  $\alpha$  a root of  $\Delta_+ \backslash \mathfrak{R}_2$  such that  $\mathfrak{g}_{\alpha} \subset P_{\Psi}^+$ . Since  $\alpha \leq \theta$ , hence there exists  $\theta_1, \cdots, \theta_k$  a positive roots such that  $\theta = \alpha + \theta_1 + \cdots + \theta_k$ and  $\alpha + \theta_1 + \cdots + \theta_j \in \Delta$  for each  $j \in [\![1, k]\!]$ . By induction on k, we prove that  $\mathfrak{g}_{\theta} \subset P_{\Psi}^+$ .

If k = 1, then  $\theta = \alpha + \theta_1$  and by applying the proposition 2.1, we have  $\mathfrak{g}_{\theta} \subset P_{\psi}^+$ . If  $k \ge 1$ . One may write  $\theta = \underbrace{\alpha + \theta_1 + \cdots + \theta_{k-1}}_{\theta'} + \theta_k$ . By induction  $\mathfrak{g}_{\theta'} \subset P_{\psi}^+$  and

by applying the proposition 2.1, we have  $\mathfrak{g}_{\theta} \subset P_{\psi}^+$ .

Now, let  $\beta$  be a root such that  $\mathfrak{g}_{\beta} \subset \mathfrak{n}_{2}^{+}$ . We have  $\beta \leq \theta$  then there exists  $\beta_{1}, \dots, \beta_{k}$  a positive roots such that  $\theta = \beta + \beta_{1} + \dots + \beta_{s}$  and  $\beta + \beta_{1} + \dots + \beta_{j} \in \Delta$  for each  $j \in \llbracket 1, s \rrbracket$ . We have  $\theta = \underbrace{\beta + \beta_{1} + \dots + \beta_{s-1}}_{\beta'} + \beta_{s}$ , then by applying the

proposition 2.1, we have  $\mathfrak{g}_{\beta'} \subset P_{\psi}^+$ . Then, we have  $\mathfrak{g}_{\beta} \subset P_{\psi}^+$ . Therefore, we prove that  $\mathfrak{n}_2^+ \subset P_{\psi}^+$  and this implies that  $\mathfrak{n}_2^+ = P_{\psi}^+$  and  $\mathfrak{n}_2^- = P_{\psi}^-$ .

Finally,  $[\tau, \mathfrak{g}] = V_{\mathfrak{m}} + \mathfrak{n}_2^- = \mathfrak{g}$ . So the second result of the theorem is proved.

**Eexample 2.1** Let  $\mathfrak{g}$  be a simple Lie algebra of type  $F_4$  and  $\Pi = \{\alpha_1, \alpha_2, \alpha_3, \alpha_4\}$  be a basis of the roots system of  $F_4$ . We consider the nilpotent standard subalgebra  $\mathfrak{m}$ associated to the subsystem  $\mathfrak{R} = \{\alpha_3\}$ . We have  $\Pi \setminus S_2 = \Pi \setminus S_1 = \{\alpha_1, \alpha_2, \alpha_4\}$  and  $\Delta_+ \setminus \mathfrak{R}_2 = \{\alpha_1 + 2\alpha_2 + 4\alpha_3 + 2\alpha_4, \alpha_1 + 3\alpha_2 + 4\alpha_3 + 2\alpha_4, 2\alpha_1 + 3\alpha_2 + 4\alpha_3 + 2\alpha_4\}.$ 

- 1. If we consider a standard subalgebra defined by  $\mathfrak{m}$  and an ideal  $\mathfrak{r}_{\mathfrak{o}}$  associated to common connected component  $\Psi = \{\alpha_4\}$ . We have  $P_{\Psi}^- = \emptyset$ , in this case  $V_{\tau} = V_{\mathfrak{m}}$ .
- 2. If we consider a standard subalgebra defined by  $\mathfrak{m}$  and an ideal  $\mathfrak{r}_{\mathfrak{o}}$  associated to common connected component  $\Psi = \{\alpha_1, \alpha_2\}$ . We have  $P_{\Psi}^- \neq \emptyset$ , more precisely,  $P_{\Psi}^- = \mathfrak{n}_2^-$ . In this case  $V_{\tau} = \mathfrak{g}$ .

# 3 A class of standard subalgebra

In this section, we prove that any standard subalgebra of  $\mathfrak{g}(A)$ , under certain supplementary condition, is of the form given by theorem 1.1.

Let  $\mathfrak{m}$  be a nilpotent standard subalgebra of  $\mathfrak{g}$  associated to subsystem  $\mathfrak{R} \subset \Delta_+$  of roots two by two non comparable.

**Proposition 3.1** Let  $V_{\mathfrak{m}}$  be the appui subspace associated to  $\mathfrak{m}$  then  $[V_{\mathfrak{m}}, \mathfrak{g}] = \mathfrak{g}$ .

**Proof**: We consider  $\mathfrak{n}$  the nilradical part of normalizer of  $\mathfrak{m}$ . The subalgebra  $\mathfrak{n}$  is complete standard subalgebra associated to subsystem  $S_2 = \bigcup_{\omega \in \mathfrak{R}} S^{\omega}$ . We have  $V_{\mathfrak{n}} = [\mathfrak{n}, \mathfrak{g}] \subset [V_{\mathfrak{m}}, \mathfrak{g}]$ . Since  $V_{\mathfrak{n}} = \sum_{\alpha \in \Delta_+} \mathfrak{g}_{\alpha} + \mathfrak{h} + \sum_{\alpha \in \Delta_+ \setminus \mathfrak{R}_2^{\mathfrak{n}}} \mathfrak{g}_{-\alpha}$  with  $\mathfrak{R}_2^{\mathfrak{n}} = \{\alpha \in \Delta_+ : \alpha_{\gamma} = \theta_{\gamma} \text{ for all } \gamma \in S_2\}$  (remark 2.1) then, it is enough to prove that  $\mathfrak{g}_{-\alpha}$  is included in  $[V_{\mathfrak{m}}, \mathfrak{g}]$ , for any root  $\alpha \in \mathfrak{R}_2^{\mathfrak{n}}$ . Now, let  $\beta$  be the root of  $\mathfrak{R}_2^{\mathfrak{n}}$  then for all  $\gamma \in S^{\beta} : ht_{\gamma}(\alpha) \ge ht_{\gamma}(\beta)$  where  $ht_{\gamma}(\alpha)$  and  $ht_{\gamma}(\beta)$  is the multiplicity of  $\gamma$  in the decomposition of simple roots of  $\alpha$  and of  $\beta$ . Therefore, we have  $\beta \le \alpha$  and this implies that  $\alpha \in \mathfrak{R}_1$ . We deduce that  $\mathfrak{g}_{-\alpha} = [[e_{\alpha}, e_{-\alpha}], \mathfrak{g}_{-\alpha}] \subset [V_{\mathfrak{m}}, \mathfrak{g}]$ . This proves that  $V = \mathfrak{g}$ .

**Remark 3.1** In proposition 3.1, if we take  $\tau$  a standard subalgebra not necessarily nilpotent, we have also  $[V_{\tau}, \mathfrak{g}] = \mathfrak{g}$ .

Let *n* be a natural number. Let  $\{I_{n+j}\}_{j\geq 0}$  be a family of vectors subspaces of  $\mathfrak{g}$ such that the subspace  $\overline{\tau} = \sum_{j\geq 0} t^{n+j} \otimes I_{n+j} + \mathbb{C}K$  is a standard subalgebra of  $\mathfrak{g}(A)$  of normalizer  $\overline{\rho} = \rho + t\mathbb{C}[t] \otimes \mathfrak{g} + \mathbb{C}K + \mathbb{C}d$  with  $\rho$  is the parabolic subalgebra of  $\mathfrak{g}$ .

We have  $[\overline{\tau},\overline{\rho}] = t^n \otimes [I_n,\rho] + t^{n+1} \otimes ([I_n,\mathfrak{g}] + [I_{n+1},\rho]) + t^{n+2} \otimes ([I_n,\mathfrak{g}] + [I_{n+1},\mathfrak{g}] + [I_{n+2},\rho]) + t^{n+3} \otimes (\cdots + [I_{n+2},\mathfrak{g}] + \cdots) + \cdots \subset \overline{\tau}.$ 

**Lemma 3.1** We have the following relations :

- 1.  $[I_n, \rho] \subset I_n$ .
- 2.  $[I_n, \mathfrak{g}] \subset I_{n+1}$  and  $[I_{n+1}, \rho] \subset I_{n+1}$ .
- 3.  $[I_{n+1}, \mathfrak{g}] \subset I_{n+2}$  and  $[I_{n+2}, \rho] \subset I_{n+2}$ .
- 4. For all  $j \ge 3$ :  $[I_{n+2}, \mathfrak{g}] \subset I_{n+j}$ .

**Proposition 3.2** If  $I_n$  is a Lie subalgebra of  $\mathfrak{g}$  then  $I_n$  is included in  $\rho$ .

**Proof**: We can write  $\rho = \sum_{\alpha \in \Delta_+} \mathfrak{g}_{\alpha} + \mathfrak{h} + \sum_{\alpha \in \langle \Pi \setminus T \rangle^+} \mathfrak{g}_{-\alpha}$ , for a certain subset T of

Π.

We suppose that  $I_n$  is not in  $\rho$  and we set  $T_1 = \{\beta \in T, \ \mathfrak{g}_{-\beta} \subset I_n\}$ . We consider  $\rho_1 = \sum_{\alpha \in \Delta_+} \mathfrak{g}_{\alpha} + \mathfrak{h} + \sum_{\alpha \in \langle \Pi \setminus (T \setminus T_1) \rangle^+} \mathfrak{g}_{-\alpha}$  the parabolic subalgebra of  $\mathfrak{g}$  associated to  $(T \setminus T_1)$ . By definition  $\rho$  is included in  $\rho_1$ .

We want to prove that  $\overline{\tau}$  is an ideal of  $\rho_1 + t\mathbb{C}[t] \otimes \mathfrak{g} + \mathbb{C}K + \mathbb{C}d$ . First, we prove that  $I_n$  is an ideal of  $\rho_1$ . Let  $\alpha$  and  $\beta$  be two roots such that  $\mathfrak{g}_{\alpha} \subset I_n$ ,  $\mathfrak{g}_{\beta} \subset \rho_1$  and  $\alpha + \beta$  is the root.

If  $\beta$  is a root of  $\Delta_+ \cup \langle \Pi \setminus T \rangle^-$  then  $\mathfrak{g}_\beta$  is included in  $\rho$  and  $[\mathfrak{g}_\alpha, \mathfrak{g}_\beta] \subset [I_n, \rho] \subset I_n$ . If  $\beta$  is a root of  $\langle \Pi \setminus (T \setminus T_1) \rangle^-$  with  $C_\beta \cap T_1 \neq \emptyset$ . There exists  $\beta_1, \cdots, \beta_r$  a simple roots such that  $\beta = -\beta_1 - \cdots - \beta_r$  and for all  $j \in [\![1, r]\!]$ ,  $\beta_1 + \cdots + \beta_j \in \Delta$ . Let i be the smallest index such that  $\beta_i \in T_1$ . We have  $\beta'_{i-1} = \beta_1 + \cdots + \beta_i \in \langle \Pi \setminus T \rangle$  and  $\mathfrak{g}_{-\beta'_i} = [\mathfrak{g}_{-\beta_i}, \mathfrak{g}_{-\beta'_{i-1}}] \subset [I_n, \rho] \subset I_n$ . Now taking the simple root  $\beta_{i+1}$ , we have two cases :

If  $\beta_{i+1} \in T_1$ , we have  $\mathfrak{g}_{-\beta'_{i+1}} = [\mathfrak{g}_{-\beta'_i}, \mathfrak{g}_{-\beta_{i+1}}] \subset [I_n, I_n] \subset I_n$  (because  $I_n$  is subalgebra).

If  $\beta_{i+1} \in \Pi \setminus T$ , we have  $\mathfrak{g}_{-\beta'_{i+1}} = [\mathfrak{g}_{-\beta_{i+1}}, \mathfrak{g}_{-\beta'_i}] \subset [\rho, I_n] \subset I_n$ . With similar argument, we show that  $\mathfrak{g}_{\beta} \subset I_n$ .

Therefore,  $\mathfrak{g}_{\alpha+\beta} = [\mathfrak{g}_{\alpha}, \mathfrak{g}_{\beta}] \subset [I_n, I_n] \subset I_n$ . Then,  $I_n$  is the standard subalgebra of  $\mathfrak{g}$  of normalizer  $\rho_1$ .

Now, we have  $V_{I_n} = [I_n, \mathfrak{g}] \subset I_{n+1}$  and  $[I_{n+1}, \rho_1] \subset [I_{n+1}, \rho] + [I_{n+1}, I_n] \subset I_{n+1}$ .

Since  $I_n$  is a standard subalgebra then, by applying the proposition 3.1 an the relation 3 of the lamma 3.1, we have  $\mathfrak{g} = [V_{I_n}, \mathfrak{g}] \subset [I_{n+1}, \mathfrak{g}] \subset I_{n+2}$ . Then  $I_{n+2} = \mathfrak{g}$ . Since  $\mathfrak{g}$  is the simple Lie algebra, for  $j \ge 3$ , we have  $[I_{n+2}, \mathfrak{g}] = [\mathfrak{g}, \mathfrak{g}] = \mathfrak{g} \subset I_{n+j}$ . This implies that  $I_{n+j} = \mathfrak{g}$ .

Finally, we have proved that  $\overline{\tau}$  is an ideal of the parabolic subalgebra  $\rho_1 + t\mathbb{C}[t] \otimes \mathfrak{g} + \mathbb{C}K + \mathbb{C}d$  and this subalgebra contains the normalizer  $\rho + t\mathbb{C}[t] \otimes \mathfrak{g} + \mathbb{C}K + \mathbb{C}d$  of  $\overline{\tau}$ , contradiction. Hence  $I_n$  is included in  $\rho$ .

**Remark 3.2** The proposition 3.2 proves that  $I_n$  is the standard subalgebra of normalizer  $\rho$ .

The following theorem determines a class of standard subalgebra of  $\mathfrak{g}(A)$  under certain condition.

**Theorem 3.1** Let  $\overline{\tau} = \sum_{j \ge 0} t^{n+j} \otimes I_{n+j} + \mathbb{C}K$  be a standard subalgebra of  $\mathfrak{g}(A)$  of normalizer  $\overline{\rho} = \rho + t\mathbb{C}[t] \otimes \mathfrak{g} + \mathbb{C}K + \mathbb{C}\mathfrak{d}$  and  $I_n \neq 0$ .

If  $I_n$  is a Lie subalgebra of  $\mathfrak{g}$  then  $\overline{\tau} = t^n \otimes \tau + t^{n+1} \otimes V + t^{n+2}\mathbb{C}[t] \otimes \mathfrak{g} + \mathbb{C}K$ where  $\tau = I_n$  is a standard subalgebra of  $\mathfrak{g}$  of normalizer  $\rho$ , V is a subspace of  $\mathfrak{g}$ contain the subspace  $V_{\tau} = [\tau, \mathfrak{g}]$  and  $I_{n+j} = \mathfrak{g}$ , for all  $j \ge 2$ .

**Proof**: Let *n* be a natural number. We set  $I_n = \tau$ . We write  $\overline{\tau} = t^n \otimes \tau + \sum_{j \ge 1} t^{n+j} \otimes I_{n+j} + \mathbb{C}K$ .

By applying the proposition 3.2, we have  $\tau$  is included in  $\rho$ . Then,  $\tau$  is the standard subalgebra of  $\mathfrak{g}$  of normalizer  $\rho$ .

We set  $V = I_{n+1}$ . The relation 2. in the lemma 3.1 proves that  $V_{\tau}$  is included in V where  $V_{\tau}$  is the appui subspace associated to  $\tau$ .

We prove by induction on  $j \ge 2$  that  $I_{n+j} = \mathfrak{g}$ . We use the relation 3 in the lemma 3.1, then we have  $[V_{\tau}, \mathfrak{g}] \subset I_{n+2}$ . By applying the proposition 3.1 and remark 3.1, we have  $I_{n+2} = \mathfrak{g}$ .

Since  $\mathfrak{g}$  is the simple Lie algebra, then we have  $[\mathfrak{g},\mathfrak{g}] = \mathfrak{g}$ . Using the relation 4 in the lemma 3.1, we have  $I_{n+j} = \mathfrak{g}$ . Therefore  $\overline{\tau} = t^n \otimes \tau + t^{n+1} \otimes V + t^{n+2} \mathbb{C}[t] \otimes \mathfrak{g}$ .

**Corollary 3.1** Let  $\overline{\tau} = I_0 + \sum_{k \ge 1} t^k \otimes I_k + \mathbb{C}K + \mathbb{C}d$  be the standard subalgebra of  $\mathfrak{g}(A)$ of normalizer  $\rho(\overline{\tau}) = \rho + t\mathbb{C}[t] \otimes \mathfrak{g} + \mathbb{C}K + \mathbb{C}d$ .

Then  $\tau = I_0$  is a standard subalgebra of  $\mathfrak{g}$  of normalizer  $\rho$  and  $\overline{\tau} = \tau + t\mathbb{C}[t] \otimes \mathfrak{g} + \mathbb{C}K + \mathbb{C}\mathfrak{d}$ .

**Proof**: It is an immediate consequence of the previous theorem.

# References

- [1] BOURBAKI, N. : Groupe et Algèbre de Lie, ch 1. Hermann Paris 1960.
- [2] BOURBAKI, N. : Groupe et Algèbre de Lie, ch 4, 5, 6. Hermann Paris, 1968.
- [3] CELLINI, P., PAPI, P. : Ad-nilpotent ideals of a Borel subalgebras. Journal of algebras. 225 (2000), 130-141.
- [4] FAVRE, G., L. SANTHAROUBANE, L. : Nilpotent Lie algebras of classical type. Journal of algebras. 202, 1998.
- [5] GOZE, M., KHAKIMDJANOV, Y. : Nilpotent Lie algebras. Kluwer Academic Publishers 1996.
- [6] GUREVICH, G. B. : Standard Lie algebras. Math. Sbornik. 1954.
- [7] HUMPHREYS, J. E. : Introduction to Lie lgebras and representation theory. Springer-Verlag New York Inc 1972.
- [8] Victor, G. KAC. : Infinite dimensional Lie algebras. Cambridge university press 1990.
- [9] KHAKIMDJANOV, Y. : Standard subalgebras of reductive Lie algebras. Moscow University Mathematics Bulletin. 29 (1974).
- [10] KRATTENTHALER, C., ORSINA, L., PAPI, P. : Enumeration of ad-nilpotent b-ideal for simple Lie algebras. Adv. in App. Math. 28 (2002), 478-522.
- [11] ORSINA, L., PAPI, P. : Enumeration of ad-nilpotent ideals of a Borel subalgebras in type A by class of nilpotence. C.R.Acad.Sci.Paris Seri I Math. 330 (2000), 651-655.
- [12] ES SAADI, B. : Sous algèbres de Lie standard d'algèbre de Kac-Moody affine. PhD thesis, Université de Haute Alsace Juin 2004.
- [13] ES SAADI, B., KHAKIMDJANOV, YU., MAKHLOUF, A. : Standard subalgebras of semisimple Lie algebras and computer-aided for enumeration. Annales Mathematiques Blaise Pascal. 10 (2003), 315-326.
- [14] SERRE, J-P. : Algbres de Lie smi-simples complexes. Benjamin New york 1966.