# BETHE ANSATZ AND INVERSE SCATTERING TRANSFORM IN A PERIODIC BOX-BALL SYSTEM

A. KUNIBA, T. TAKAGI, AND A. TAKENOUCHI

ABSTRACT. We formulate the inverse scattering method for a periodic boxball system and solve the initial value problem. It is done by a synthesis of the combinatorial Bethe ansätze at q = 1 and q = 0, which provides the ultradiscrete analogue of quasi-periodic solutions in soliton equations, e.g., action-angle variables, Jacobi varieties, period matrices and so forth. As an application we establish explicit formulas counting the states characterized by conserved quantities and the generic and fundamental period under the commuting family of time evolutions.

### 1. INTRODUCTION

In [1, 2], a class of periodic soliton cellular automata on one dimensional lattice is introduced. They are associated with crystal basis of non-exceptional quantum affine algebras  $U_q(\mathfrak{g}_n)$  [3]. In this paper we focus on the simplest case  $\mathfrak{g}_n = A_1^{(1)}$ known as the periodic box-ball system [4], and solve the initial value problem by the inverse scattering method. As a result, all the properties and formulas conjectured in [1, 2] are established in this case. Our approach is a synthesis of the two versions of the combinatorial Bethe ansatz at q = 1 [5, 6] and q = 0 [7]. We provide a selfcontained proof for all the important statements together with several examples. Here is a typical time evolution pattern on the length 13 lattice:

 $\begin{array}{l}t=0:&2&2&2&1&1&1&1&2&2&1&1&2&1\\t=1:&1&1&1&2&2&2&1&1&2&2&2&2&2\\t=2:&2&2&1&1&1&2&2&2&2&1&1&2&1\\t=3:&1&1&2&2&2&1&1&1&1&2&2&1&2\\t=4:&2&2&1&1&1&2&2&2&1&2&1&1&2&2\\t=5:&1&1&2&2&1&1&1&1&2&2&2&1&2\\t=6:&2&2&1&1&2&2&2&1&1&1&2&2&1&1\\t=7:&1&1&2&2&1&1&1&2&2&2&1&1&2\\t=8:&2&2&1&1&2&2&1&1&1&2&2&1\\t=9:&1&1&2&2&1&1&2&2&2&1&1&2\\\end{array}$ 

Regarding the letter 1 as the background, one observes the three solitons traveling periodically to the right with the velocity equal to the amplitudes 3, 2 and 1 as long as they stay away from each other. Under the collisions they do not smash into pieces nor glue together. This is due to the underlying quantum group symmetry. The time evolution from t = 3 to t = 4 states for example have been determined from the following diagram:

$$1 \\ 122 \\ + \\ 112 \\ + \\ 112 \\ + \\ 111 \\ + \\ 112 \\ + \\ 112 \\ + \\ 112 \\ + \\ 112 \\ + \\ 122 \\ + \\ 122 \\ + \\ 122 \\ + \\ 122 \\ + \\ 122 \\ + \\ 112 \\ + \\ 111 \\ + \\ 111 \\ + \\ 111 \\ + \\ 112 \\ + \\ 122 \\ + \\ 122 \\ + \\ 122 \\ + \\ 122 \\ + \\ 122 \\ + \\ 122 \\ + \\ 122 \\ + \\ 122 \\ + \\ 122 \\ + \\ 122 \\ + \\ 122 \\ + \\ 122 \\ + \\ 112 \\ + \\ 111 \\ + \\ 111 \\ + \\ 112 \\ + \\ 122 \\ + \\$$

Here the local vertices stand for the  $U_q(A_1^{(1)})$  quantum R matrix sending the spin  $\frac{3}{2} \otimes \frac{1}{2}$  states on the NW edges to the spin  $\frac{1}{2} \otimes \frac{3}{2}$  states on the SE edges. This is a standard diagram for the row transfer matrix in the  $3 \times 1$  fusion vertex model. The peculiar feature here is the specialization to q = 0, which makes the R a deterministic map called the combinatorial R [8, 9]. The above ones correspond to the l = 3 case in Figure 2.1, and the associated time evolution is named  $T_3$ . On the horizontal edge, the leftmost and the rightmost ones are the same element 122 reflecting the periodic boundary condition. Denote the array on the top and the bottom lines by p and p', respectively. It turns out that p' is uniquely determined from p by the combinatorial R and the boundary condition said above. The resulting map defines a time evolution  $p' = T_3(p)$ . The periodic box-ball system is a dynamical system endowed with such time evolutions  $T_1, T_2, \ldots$ , which are essentially the commuting family of fusion transfer matrices at q = 0. In other words, it is a solvable vertex model [10] at q = 0 on the periodic lattice. It is the periodic extension of the original box-ball system on the infinite lattice [11].

In this paper we solve the initial value problem of the periodic box-ball system. It is done by a synthesis of the Bethe ansätze [12] at q = 1 [5, 6] and q = 0 [7], which yields a periodic ultradiscretization of the inverse scattering method in soliton theory [13, 14]. As an application, we are able to answer the interesting question; what is the fundamental period of p under the time evolutions  $T_1, T_2, \ldots$ ? Namely, the minimum positive integer  $N_l$  satisfying  $T_l^{N_l}(p) = p$ . For instance for the above p, one has  $N_1 = 13, N_2 = 91, N_3 = 273^1$ . An explicit formula for the fundamental period under  $T_{\infty}$  has been discovered in [15] by a combinatorial argument. Here we take the inverse scattering approach and establish the results (4.22) and (4.26) for general  $T_l$ . Under our scheme, the period  $\mathcal{N}$  acquires the intrinsic characterization  $\mathcal{N}\vec{h} \in \Gamma$  (4.31) in terms of straight motions on the set

$$(\mathcal{I}_{m_{j_1}} \times \cdots \times \mathcal{I}_{m_{j_s}})/\Gamma.$$

This is an 'ultradiscrete Jacobi variety'. See (4.30). The conjectural formula for the generic period in the most general  $A_n^{(1)}$  case [2] also admits a similar interpretation.

Let us explain the content of the paper in some detail. In Section 2, we begin with the rudiments of the crystal basis theory, formulate the periodic box-ball system and explain its basic properties. The fundamental role is played by the combinatorial R. It governs the local dynamics of the system and ultimately its whole aspect together with the global constraint imposed by the periodic boundary condition. The commuting family of time evolutions  $T_l$  and the conserved quantity  $E_l$  called energy are constructed. It is done by a simple extension of the arguments in [16, 17, 18] to periodic versions and supplements the original description in [4]. The both  $T_l$  and the  $E_l$  enjoy the symmetry under the extended affine Weyl group

<sup>&</sup>lt;sup>1</sup>It is a general feature of the system that  $T_l(p) = T_{\infty}(p)$  hence  $N_l = N_{\infty}$  for  $l \ge l_0$ , where  $l_0$  is the amplitude of the largest soliton involved in p. See Proposition 2.5 and the subsequent remark.

 $\widetilde{W}(A_1^{(1)})$  (Proposition 2.3). The time evolution  $T_l$  with sufficiently large l admits a simple description as a translation in  $\widetilde{W}(A_1^{(1)})$  (Proposition 2.5). Although quite straightforward, the proof of these properties announced in [1, 2] are presented here for the first time.

Section 3 is a main part of the paper where we present our inverse scattering formalism. We invoke the Kerov-Kirillov-Reshetikhin (KKR) bijection between the rigged configurations and highest paths. We call it the KKR theory or combinatorial Bethe ansatz at q = 1 in this paper. The necessary facts are available in Appendix A as well as in the original papers [5, 6, 28] and the latest review [19].

Roughly speaking, our action-angle variables are the rigged configurations. However, the KKR theory works only for highest paths whereas the periodic boundary condition brings all the paths into the game. To reconcile them in a remarkable harmony via the special prescription (3.12) is the heart of our inverse scattering formalism. Proposition 3.7 is the crux to guarantee the well-definedness of the resulting direct and inverse scattering map  $\Phi$  (3.15). Our solution of the initial value problem is presented in Theorem 3.12. It is also applicable, with a drastic simplification, to the original box-ball system [11] by taking the system size infinite.

In Section 4 we explain the origin of our inverse scattering formalism in the other framework of the combinatorial Bethe ansatz at q = 0 [7]. The central role is played by the linear congruence equation called the string center equation (4.3). It is the Bethe equation on the string centers at q = 0 whose off-diagonal solutions yield the weight multiplicities of the  $sl_2$  module  $(\mathbb{C}^2)^{\otimes L}$ . By the map  $\Psi$  (4.18), we link the logarithmic branch in the string center equation with the angle variable in Section 3. Then our key construction (3.12) is nothing but the equivalence relation among the off-diagonal solutions to the string center equation (Theorem 4.3). Through our direct and inverse scattering transforms, the nonlinear dynamics in the periodic box-ball system becomes a straight motion of the Bethe roots or angle variables. With these features (Corollary 4.4) in hand, it is straightforward to derive explicit formulas for the generic (4.22) and the fundamental period (4.26)under any time evolution  $T_l$ . The number of states characterized by conserved quantities (4.21) are obtained and the Bethe eigenvalue is shown to be a root of unity related to the generic period (Proposition 4.11). The number of disjoint orbits under the commuting family of time evolutions is also determined in (4.35). Our angle variables live in the set (4.30), which serves as an ultradiscrete analog of the Jacobi variety in the classical theory of quasi-periodic solutions to soliton equations [20, 21].

The results in Sections 3 and 4 uncover a significant interplay between the KKR theory and the string center equation. They substantially achieve a synthesis of the combinatorial Bethe ansätze at q = 1 and q = 0, which was already foreseen in the earlier works [1, 2] on generalized periodic box-ball systems. Section 5 is a summary.

Appendix A summarizes necessary facts on the rigged configurations and their bijective correspondence with the highest paths. A piecewise linear formula for the KKR bijection  $\phi^{-1}$  is available in (A.5).

Appendix B contains the proof of Proposition 3.4. For any path not necessarily highest, it essentially identifies the two descriptions of the conserved quantity in terms of the energy and the configuration. The former is related to the soliton content and the latter to the string content. Appendix C is devoted to a proof of Proposition 3.7, which assures the welldefinedness of the direct/inverse scattering map  $\Phi$  (3.15). The key is to investigate the relation between the two rigged configurations corresponding to the paths  $q \otimes r$ and  $r \otimes q$  with q and r both being highest.

Appendix D is the proof of Theorem 3.12, which asserts the linearization of the time evolution in terms of the angle variables. Our main idea is to realize non highest periodic paths as a segment of a large highest path having the structure  $p \otimes p \otimes \cdots$  and apply the KKR theory to the latter.

### 2. Periodic box-ball system

2.1. Crystals and combinatorial R. We recapitulate the basic facts in the crystal basis theory [22, 8, 3, 9, 18]. Let  $B_l$  the the crystal of the l-fold symmetric tensor representation of  $U_q(A_1^{(1)})$ . As a set it is given by  $B_l = \{x = (x_1, x_2) \in (\mathbb{Z}_{\geq 0})^2 \mid x_1 + x_2 = l\}$ . The element  $(x_1, x_2)$  will also be expressed as the length l row shape semistandard tableau containing the letter  $i \ x_i$  times. For example,  $B_1 = \{\underbrace{1}, \underbrace{2}\}, B_2 = \{\underbrace{11}, \underbrace{12}, \underbrace{22}\}$ . (We omit the frames of the tableaux hereafter.) The action of Kashiwara operators  $\tilde{f}_i, \tilde{e}_i : B \to B \sqcup \{0\} (i = 0, 1)$  reads  $(\tilde{f}_i x)_j = x_j - \delta_{j,i} + \delta_{j,i+1}$  and  $(\tilde{e}_i x)_j = x_j + \delta_{j,i} - \delta_{j,i+1}$ , where all the indices are in  $\mathbb{Z}_2$ , and if the result does not belong to  $(\mathbb{Z}_{\geq 0})^2$ , it should be understood as 0. The classical part of the weight of  $x = (x_1, x_2) \in B_l$  is wt $(x) = l\Lambda_1 - x_2\alpha_1 = (x_1 - x_2)\Lambda_1$ , where  $\Lambda_1$  and  $\alpha_1 = 2\Lambda_1$  are the fundamental weight and the simple root of  $A_1$ .

For any  $b \in B$ , set

$$\varepsilon_i(b) = \max\{m \ge 0 \mid \tilde{e}_i^m b \neq 0\}, \quad \varphi_i(b) = \max\{m \ge 0 \mid f_i^m b \neq 0\}$$

By the definition one has  $\varepsilon_i(x) = x_{i+1}$  and  $\varphi_i(x) = x_i$  for  $x = (x_1, x_2) \in B_l$ .

For two crystals B and B', one can define the tensor product  $B \otimes B' = \{b \otimes b' \mid b \in B, b' \in B'\}$ . The operators  $\tilde{e}_i, \tilde{f}_i$  act on  $B \otimes B'$  by

(2.1) 
$$\tilde{e}_i(b \otimes b') = \begin{cases} \tilde{e}_i b \otimes b' & \text{if } \varphi_i(b) \ge \varepsilon_i(b') \\ b \otimes \tilde{e}_i b' & \text{if } \varphi_i(b) < \varepsilon_i(b'), \end{cases}$$

(2.2) 
$$\tilde{f}_i(b \otimes b') = \begin{cases} \tilde{f}_i b \otimes b' & \text{if } \varphi_i(b) > \varepsilon_i(b') \\ b \otimes \tilde{f}_i b' & \text{if } \varphi_i(b) \le \varepsilon_i(b'). \end{cases}$$

Here  $0 \otimes b'$  and  $b \otimes 0$  should be understood as 0. The tensor product  $B_{l_1} \otimes \cdots \otimes B_{l_k}$  is obtained by repeating the above rule. The classical part of the weight of  $b \in B$  for any  $B = B_{l_1} \otimes \cdots \otimes B_{l_k}$  is given by wt $(b) = (\varphi_1(b) - \varepsilon_1(b))\Lambda_1 = (\varepsilon_0(b) - \varphi_0(b))\Lambda_1$ .

The crystal  $B_l$  admits the affinization  $\operatorname{Aff}(B_l)$ . It is the infinite set  $\operatorname{Aff}(B_l) = \{\zeta^{db} \mid b \in B_l, d \in \mathbb{Z}\}$  endowed with the crystal structure  $\tilde{e}_i(\zeta^{db}) = \zeta^{d-\delta_{i,0}}(\tilde{e}_ib), \tilde{f}_i(\zeta^{db}) = \zeta^{d+\delta_{i,0}}(\tilde{f}_ib)$ . The parameter  $\zeta$  is called the spectral parameter. The isomorphism of the affine crystal  $\operatorname{Aff}(B_l) \otimes \operatorname{Aff}(B_k) \xrightarrow{\sim} \operatorname{Aff}(B_k) \otimes \operatorname{Aff}(B_l)$  is the unique bijection that commutes with Kashiwara operators (up to a constant shift of H below). It is the q = 0 analogue of the quantum R and called the combinatorial R. Explicitly it is given by  $R : \zeta^d x \otimes \zeta^e y \mapsto \zeta^{e+H(x \otimes y)} \tilde{y} \otimes \zeta^{d-H(x \otimes y)} \tilde{x}$  with

$$\tilde{x}_{i} = x_{i} + Q_{i}(x, y) - Q_{i-1}(x, y), \quad \tilde{y}_{i} = y_{i} + Q_{i-1}(x, y) - Q_{i}(x, y), 
Q_{i}(x, y) = \min(x_{i+1}, y_{i}), 
H(x \otimes y) = -Q_{0}(x, y).$$
(2.3)

Here  $x \otimes y \simeq \tilde{y} \otimes \tilde{x}$  under the isomorphism  $B_l \otimes B_k \simeq B_k \otimes B_l$ . The relation is depicted as

For example  $B_l \otimes B_1 \simeq B_1 \otimes B_l$  is listed as follows:



FIGURE 2.1. Combinatorial  $R: B_l \otimes B_1 \simeq B_1 \otimes B_l$ 

Let  $\omega : (x_1, x_2) \mapsto (x_2, x_1)$  be the involutive Dynkin digram automorphism of  $B_l$ . The combinatorial R enjoys the symmetry:

(2.4) 
$$(\omega \otimes \omega)R = R(\omega \otimes \omega) \text{ on } B_l \otimes B_k.$$

We write the highest element  $(l, 0) \in B_l$  as  $u_l$ . The energy function H (2.3) is normalized so as to attain the maximum at  $H(u_l \otimes u_k) = 0$  and ranges over  $-\min(l,k) \leq H \leq 0$  on  $B_l \otimes B_k$ . The combinatorial R satisfies the Yang-Baxter relation:

$$(2.5) (1 \otimes R)(R \otimes 1)(1 \otimes R) = (R \otimes 1)(1 \otimes R)(R \otimes 1)$$

on  $\operatorname{Aff}(B_j) \otimes \operatorname{Aff}(B_l) \otimes \operatorname{Aff}(B_k)$ . Let  $\varrho(b_1 \otimes \cdots \otimes b_k) = b_k \otimes \cdots \otimes b_1$  be the reverse ordering of the tensor product for any k. The combinatorial R has the property:

(2.6) 
$$R \varrho = \varrho R \quad \text{on } B_l \otimes B_k.$$

2.2. Time evolution and conserved quantities. We fix the integer  $L \in \mathbb{Z}_{\geq 1}$  corresponding to the system size throughout. Set

(2.7) 
$$\mathcal{P} = B_1^{\otimes L}, \quad \mathcal{P}_+ = \{ p \in \mathcal{P} \mid \tilde{e}_1 p = 0 \}.$$

We will also write  $\operatorname{Aff}(\mathcal{P}) = \operatorname{Aff}(B_1)^{\otimes L}$ . An element of  $\mathcal{P}(\mathcal{P}_+)$  is called a path (highest path). The condition  $\tilde{e}_1 p = 0$  on the path  $p = b_1 \otimes \cdots \otimes b_L$  is equivalent to the simple postulate

The weight of the path  $p = b_1 \otimes \cdots \otimes b_L$  is given by  $\operatorname{wt}(p) = \operatorname{wt}(b_1) + \cdots + \operatorname{wt}(b_L)$ . We write  $\operatorname{wt}(p) > 0$  (wt(p) < 0) when it belongs to  $\mathbb{Z}_{>0}\Lambda_1$  ( $\mathbb{Z}_{<0}\Lambda_1$ ).

The periodic box-ball system is a dynamical system on  $\mathcal{P}$  equipped with the commuting family of time evolutions  $T_1, T_2, \ldots$ , which we shall now introduce.

**Proposition 2.1.** For any path  $p = b_1 \otimes \cdots \otimes b_L \in \mathcal{P}$  and  $l \in \mathbb{Z}_{\geq 1}$ , there exists an element  $v_l \in B_l$  such that  $(e = -d_1 - \cdots - d_L)$ 

(2.9) 
$$\zeta^0 v_l \otimes (\zeta^0 b_1 \otimes \cdots \otimes \zeta^0 b_L) \simeq (\zeta^{d_1} b'_1 \otimes \cdots \otimes \zeta^{d_L} b'_L) \otimes \zeta^e v_l$$

for some  $\zeta^{d_1}b'_1 \otimes \cdots \otimes \zeta^{d_L}b'_L \in \operatorname{Aff}(\mathcal{P})$  under the isomorphism  $\operatorname{Aff}(B_l) \otimes \operatorname{Aff}(\mathcal{P}) \simeq \operatorname{Aff}(\mathcal{P}) \otimes \operatorname{Aff}(B_l)$ . Such  $v_l$  is unique except  $\operatorname{wt}(p) = 0$  case, where  $\zeta^{d_1}b'_1 \otimes \cdots \otimes \zeta^{d_L}b'_L$  is independent of the possibly non unique choice of  $v_l$ .

*Proof.* Suppose the relation  $(x_1, x_2) \otimes p \simeq p' \otimes (y_1, y_2)$  holds under the isomorphism  $B_l \otimes \mathcal{P} \simeq \mathcal{P} \otimes B_l$ . Setting  $p' = b'_1 \otimes \cdots \otimes b'_L$ , we depict it as Figure 2.2.



### FIGURE 2.2

Fixing p and l, we regard it as the functional relation  $y_2 = y_2(x_2)$ . Then from Figure 2.1 we see that  $y_2(x_2 + 1) = y_2(x_2)$  or  $y_2(x_2 + 1) = y_2(x_2) + 1$ . Moreover, the latter holds only if all the intermediate vertices in Figure 2.2 are the bottom two types in Figure 2.1, namely wt(p) = -wt(p').

Let M be the number of  $2 \in B_1$  contained in p. We first consider the case L-M > M, i.e.,  $\operatorname{wt}(p) > 0$ . Set  $c = y_2(0)$ . Suppose that  $y_2(a) = c$  and  $y_2(a+1) = c + 1$  hold. From the above observation, the number of 2 contained in p' (for  $(x_1, x_2) = (l - a, a)$  in Figure 2.2) must be L - M. Then the weight conservation demands that a + M = c + L - M. By the assumption M < L - M, this can not happen for  $0 \le a \le c$ . This implies that  $y_2(a) = c$  for  $0 \le a \le \min(c+1, l)$ , which confirms the sought assertion. The case L - M < M can be shown similarly by interchanging the role of the letters 1 and 2. Next we consider the case L - M = M, i.e.,  $\operatorname{wt}(p) = 0$ . By the same argument as above, we find that  $y_2(a) = c$  for  $0 \le a \le c$ , and  $y_2(a) = a$  can happen for some interval  $c < a \le c'(\le l)$ . When  $c \le a < c'$ , all the intermediate vertices in Figure 2.2 are the bottom two types in Figure 2.1, whose energy H and the vertical  $B_1$  part are independent of a. This verifies the existence of  $v_l$  and uniqueness of  $\zeta^{d_1} b'_1 \otimes \cdots \otimes \zeta^{d_L} b'_L$ .

We define the time evolution  $T_l$  and the energy  $E_l$  of a path p by  $T_l(p) = b'_1 \otimes \cdots \otimes b'_L$  ( $\in \mathcal{P}$ ) and  $E_l(p) = e (\in \mathbb{Z}_{\geq 0})$  using the notation in (2.9). Proposition 2.1 assures that they are solely determined from p and l. The definition is summarized by the relation

(2.10) 
$$\zeta^0 v_l \otimes p \simeq T_l(p) \otimes \zeta^{E_l(p)} v_l$$

omitting the spectral parameters attached to p and  $T_l(p)$ . Here  $v_l \in B_l$  can be constructed from  $p \in \mathcal{P}$  by

(2.11) 
$$\begin{aligned} u_l \otimes p \simeq p^* \otimes v_l & \text{if } \operatorname{wt}(p) \ge 0, \\ \omega(u_l) \otimes p \simeq p^* \otimes v_l & \text{if } \operatorname{wt}(p) < 0 \end{aligned}$$

for some  $p^* \in \mathcal{P}$  under the isomorphism  $B_l \otimes \mathcal{P} \simeq \mathcal{P} \otimes B_l$ , where  $u_l = (l, 0) \in B_l$  as defined after (2.4). One may either use the latter relation to define  $v_l$  when wt(p) =

0. Clearly the time evolutions are weight preserving, i.e.,  $wt(T_l(p)) = wt(p)$ . They are all invertible. By using (2.6), the inverse can be found by

(2.12) 
$$T_l^{-1}(p) = \varrho T_l(\varrho(p)).$$

In the rest of the paper we will use, often without explicitly mentioning, the fact that (2.10) determines  $T_l(p)$  and  $E_l(p)$  unambiguously even though  $v_l$  is not unique in general. If p is a highest path, the affinization of (2.11) is available in Lemma B.1.

Note from Figure 2.1 that the combinatorial R on  $B_1 \otimes B_1$  is the identity map. It follows that  $T_1$  acts as the cyclic shift or 'exp  $(\sqrt{-1} \text{ (momentum)})$ ':

(2.13) 
$$T_1(b_1 \otimes b_2 \otimes \cdots \otimes b_L) = b_L \otimes b_1 \otimes \cdots \otimes b_{L-1}$$

 $T_1$  will play a special role in our inverse scattering formalism in Section 3.

**Theorem 2.2.** The commutativity  $T_lT_k(p) = T_kT_l(p)$  and the conservation  $E_l(T_k(p)) = E_l(p)$  hold.

The commutativity with  $T_1$  (2.13) is the origin of the adjective "periodic".

Proof. Take  $v_k$  for p and  $v_l$  for  $T_k(p)$  as in (2.11). Set  $R(\zeta^0 v_l \otimes \zeta^0 v_k) = \zeta^{\delta} \overline{v}_k \otimes \zeta^{-\delta} \overline{v}_l$ and regard p as an element of  $\operatorname{Aff}(\mathcal{P})$ . By using the combinatorial R, one can reorder  $\zeta^0 v_l \otimes \zeta^0 v_k \otimes p$  in two ways along the isomorphism  $\operatorname{Aff}(B_l) \otimes \operatorname{Aff}(B_k) \otimes \operatorname{Aff}(\mathcal{P}) \simeq \operatorname{Aff}(\mathcal{P}) \otimes \operatorname{Aff}(B_k) \otimes \operatorname{Aff}(B_l)$  as follows:



where the equality of the result is due to the Yang-Baxter equation (2.5). Here we have identified the outputs with  $T_k T_l(p)$ ,  $\zeta^{E_k(T_l(p))+\delta} v_k$ , etc. In particular Proposition 2.1 guarantees that  $\overline{v}_k \otimes T_l(p) \simeq T_k T_l(p) \otimes \overline{v}_k$  and  $\overline{v}_l \otimes p \simeq T_l(p) \otimes \overline{v}_l$  up to the spectral parameter. The sought relations  $T_l T_k(p) = T_k T_l(p)$  and  $E_l(T_k(p)) = E_l(p)$ are obtained by comparing the two sides.

2.3. Extended affine Weyl group invariance. Let  $s_i$  (i = 0, 1) be the Weyl group operator [22] acting on any crystal B as

$$s_i(b) = \begin{cases} \tilde{f}_i^{\varphi_i(b) - \varepsilon_i(b)}(b) & \varphi_i(b) \ge \varepsilon_i(b), \\ \tilde{e}_i^{\varepsilon_i(b) - \varphi_i(b)}(b) & \varphi_i(b) \le \varepsilon_i(b) \end{cases}$$

for  $b \in B$ . We extend  $\omega$  introduced around (2.4) to any  $B = B_{l_1} \otimes \cdots \otimes B_{l_k}$  by  $\omega(B) = \omega(B_{l_1}) \otimes \cdots \otimes \omega(B_{l_k})$ . Then  $\widetilde{W}(A_1^{(1)}) = \langle \omega, s_0, s_1 \rangle$  acts on  $\mathcal{P}$  as the extended affine Weyl group of type  $A_1^{(1)}$ .

The action of  $\tilde{f}_i, \tilde{e}_i$  and  $s_i$  is determined in principle by (2.1) and (2.2). Here we explain the *signature rule* to find the action on any  $B_{l_1} \otimes \cdots \otimes B_{l_k}$  which is of great practical use. It will be the basic ingredient in proving Proposition 2.3. The  $\varepsilon_i(b) \qquad \varphi_i(b)$ 

*i*-signature of an element  $b \in B_l$  is the symbol  $\overline{-\cdots - + \cdots +}$ . The *i*-signature of the tensor product  $b_1 \otimes \cdots \otimes b_k \in B_{l_1} \otimes \cdots \otimes B_{l_k}$  is the array of the *i*-signature of each  $b_j$ . Here is an example from  $B_5 \otimes B_2 \otimes B_1 \otimes B_4$ :

$$\begin{array}{cccc} & 11112 \otimes 12 \otimes 2 \otimes 1122 \\ \text{0-signature} & -+++ & -+ & -++ \\ \text{1-signature} & -+++ & -+ & --++ \end{array}$$

where 1122 for example represents  $1122 \in B_4$  and not  $1 \otimes 1 \otimes 2 \otimes 2 \in B_1^{\otimes 4}$ , etc. In the *i*-signature, one eliminates the neighboring pair +- (not -+) successively to finally reach the pattern  $\overbrace{-\cdots + \cdots +}^{\alpha}$  called reduced *i*-signature. The result is

to finally reach the pattern  $-\cdots -+\cdots +$  called reduced *i*-signature. The result is independent of the order of the eliminations when it can be done simultaneously in more than one places. The reduced *i*-signature tells that  $\varepsilon_i(b_i \otimes \cdots \otimes b_k) = \alpha$  and  $\varphi_i(b_i \otimes \cdots \otimes b_k) = \beta$ . In the above example, we get

$$\begin{array}{c} 11112 \otimes 12 \otimes 2 \otimes 1122 \\ ----- \\ -+ \\ ++ \\ ++ \end{array}$$

Thus  $\varepsilon_0 = 4$ ,  $\varphi_0 = 2$ ,  $\varepsilon_1 = 1$  and  $\varphi_1 = 3$ . Finally  $\tilde{f}_i$  hits the component that is responsible for the leftmost + in the reduced *i*-signature making it -. Similarly,  $\tilde{e}_i$  hits the component corresponding to the rightmost – in the reduced *i*-signature making it +. If there is no such + or – to hit, the result of the action is 0.  $s_i$  acts so as to change the reduced *i*-signature  $\overbrace{-\cdots + \cdots +}^{\alpha}$  into  $\overbrace{-\cdots - + \cdots +}^{\beta}$ . In the above example, we have

$$p = 11112 \otimes 12 \otimes 2 \otimes 1122$$
$$\tilde{f}_0(p) = 11112 \otimes 12 \otimes 2 \otimes 1112$$
$$\tilde{f}_1(p) = 11122 \otimes 12 \otimes 2 \otimes 1122$$
$$\tilde{e}_0(p) = 11122 \otimes 12 \otimes 2 \otimes 1122$$
$$\tilde{e}_1(p) = 11111 \otimes 12 \otimes 2 \otimes 1122$$
$$s_0(p) = 11222 \otimes 12 \otimes 2 \otimes 1122$$
$$s_1(p) = 11122 \otimes 12 \otimes 2 \otimes 1222.$$

For both i = 0 and 1, note that  $wt(s_i(p)) = -wt(p)$  for any p, and  $s_i(p) = p$  if wt(p) = 0. In order that  $\tilde{e}_1 p = 0$  to hold for a path  $p \in \mathcal{P}$ , it is necessary and sufficient that the reduced 1-signature to become  $+\cdots +$ , which is equivalent to the condition (2.8). We note that elimination of the +- pairs is described by successive applications of the rule

$$\overbrace{+\cdots+}^{\alpha} \xrightarrow{\beta} \qquad \qquad \overbrace{+\cdots+}^{(\alpha-\beta)_{+}} (\overbrace{\beta-\alpha)_{+}},$$

where  $(c)_{+} = \max(c, 0)$ . The time evolutions  $T_l$  and the energy  $E_l$  enjoy the extended affine Weyl group symmetry.

**Proposition 2.3.** For any  $w \in \widetilde{W}(A_1^{(1)})$ ,  $p \in \mathcal{P}$  and  $l \in \mathbb{Z}_{\geq 1}$ , the commutativity  $wT_l(p) = T_l(w(p))$  and the invariance  $E_l(w(p)) = E_l(p)$  are valid.

This property persists in the most general  $A_n^{(1)}$  case as announced in [2].

*Proof.* Let the reduced *i*-signature of p be  $\overbrace{-\cdots-+\cdots+}^{\alpha}$  and the one for  $T_l(p)$  be

 $\alpha'$   $\beta'$  $\beta'$ . First we show the assertion for  $w = s_i$ . We assume  $\alpha < \beta$ , hence  $s_i(p) = \tilde{f}_i^{\beta-\alpha}(p)$  and  $s_i(T_l(p)) = \tilde{f}_i^{\beta-\alpha}(T_l(p))$ . The proof for the case  $\alpha \ge \beta$  is parallel. Let  $v_l \in B_l$  be the element specified by (2.10) from p and set  $a = \varepsilon_i(v_l)$  and  $b = \varphi_i(v_l)$ . The *i*-signature of (2.10) reads

(2.14) 
$$\overbrace{-\cdots-+\cdots+}^{a} |\overbrace{-\cdots-+\cdots+}^{\alpha} \xrightarrow{\beta} = \overbrace{-\cdots-++\cdots+}^{\alpha'} |\overbrace{-\cdots-+\cdots+}^{\beta'} |\overbrace{-\cdots-+\cdots+}^{a} ,$$

where | signifies the position of  $\otimes$  that separates  $v_l$  with p or  $T_l(p)$ .

(i) Case  $\alpha \ge b$ . Comparing the reduced + signature, we see  $(\alpha <)\beta = b + (\beta' - a)_+ \le \alpha + (\beta' - a)_+$ , compelling  $\beta' > a$ . Thus we find  $\beta' - a \ge \beta - \alpha$ . By taking these facts into account, (2.14) is reduced to

$$\overbrace{-\cdots-}^{a}|\overbrace{-\cdots-}^{\alpha-b}\overbrace{+\cdots+}^{\beta} = \overbrace{-\cdots-}^{\alpha'}\overbrace{+\cdots+}^{\beta'-a}|\overbrace{+\cdots+}^{b}|$$

In view of this and  $\beta' - a \geq \beta - \alpha$ , application of  $\tilde{f}_i^{\beta - \alpha}$  to (2.10) yields  $v_l \otimes \tilde{f}_i^{\beta - \alpha}(p) \simeq \tilde{f}_i^{\beta - \alpha}(T_l(p)) \otimes \zeta^{E_l(p)}v_l$ . Therefore  $v_l \otimes s_i(p) \simeq s_i(T_l(p)) \otimes \zeta^{E_l(p)}v_l$ , saying  $T_l(s_i(p)) = s_i(T_l(p))$  and  $E_l(s_i(p)) = E_l(p)$ .

(ii) Case  $\alpha < b$ . Comparing the reduced + signature in (2.14), we see  $(b < b + \beta - \alpha = b + (\beta' - a)_+$ , compelling  $\beta' - a > 0$ . Thus we find  $\beta' - a = \beta - \alpha$ . By taking these facts into account, (2.14) is reduced to

$$\overbrace{-\cdots-+\cdots+}^{a}|\overbrace{+\cdots+}^{\beta}|=\overbrace{-\cdots-+\cdots+}^{\alpha'}|\overbrace{+\cdots+}^{\beta-\alpha}|$$

In view of this, the application of  $\tilde{f}_i^{b+\beta-2\alpha}$  to (2.10) yields  $\zeta^{\delta}v'_l \otimes \tilde{f}_i^{\beta-\alpha}(p) \simeq \tilde{f}_i^{\beta-\alpha}(T_l(p)) \otimes \zeta^{\delta+E_l(p)}v'_l$  with  $v'_l = \tilde{f}_i^{b-\alpha}v_l$  and  $\delta = (b-\alpha)\delta_{i,0}$ . This is equivalent to  $v'_l \otimes s_i(p) \simeq s_i(T_l(p)) \otimes \zeta^{E_l(p)}v'_l$  saying again that  $T_l(s_i(p)) = s_i(T_l(p))$  and  $E_l(s_i(p)) = E_l(p)$ .

Next we show the assertion for  $w = \omega$ . Due to the symmetry (2.4), the relation (2.10) implies  $\omega(v_l) \otimes \omega(p) \simeq \omega(T_l(p)) \otimes \omega(v_l)$  at least under  $B_l \otimes \mathcal{P} \simeq \mathcal{P} \otimes B_l$ forgetting the spectral parameter. Thus we obtain  $T_l(\omega(p)) = \omega(T_l(p))$ . In (2.10), imagine the process of sending  $v_l$  to the right through p along Figure 2.2 using the local rule listed in Figure 2.1. By the definition,  $-E_l(p)$  is the sum of H attached to the intermediate vertices. Upon application of  $\omega$ , the four types of vertices in Figure 2.1 are interchanged horizontally. But this does not alter the sum of H since the number of bottom two types are equal because of wt $(p) = \operatorname{wt}(T_l(p))$ . **Example 2.4.** The following is a commutative diagram among the paths in  $\mathcal{P} = B_1^{\otimes 6}$ .

For all the paths here, we have  $E_l(p) = 2$  for any  $l \ge 1$ .

The time evolution  $T_l$  defined by (2.10) has a simple description for l sufficiently large.

**Proposition 2.5.** For any path  $p \in \mathcal{P}$ , there exists  $k \in \mathbb{Z}_{\geq 1}$  such that  $T_l(p)$  is independent of l for  $l \geq k$ . Denoting it by  $T_{\infty}(p)$ , one has

(2.15) 
$$T_{\infty}(p) = \begin{cases} \omega(s_0(p)) & \text{if } \operatorname{wt}(p) \ge 0, \\ \omega(s_1(p)) & \text{if } \operatorname{wt}(p) \le 0. \end{cases}$$

In particular,  $T_{\infty}(p) = \omega(p)$  is valid if wt(p) = 0.

The combination  $\omega s_i$  represents a translation in  $\widetilde{W}(A_1^{(1)})$ . On account of (3.14) and Theorem 3.12, the minimum of such k is given by  $k = j_s$  with  $j_s$  specified in (3.8). This is the amplitude of the largest soliton involved in p. The key to the connection of the combinatorial R and the signature rule is

**Lemma 2.6.** Let  $(l - a, a) \otimes b \simeq b' \otimes (l - a', a')$  under the combinatorial  $R : B_l \otimes B_1 \simeq B_1 \otimes B_l$ . If  $0 \le a < l$ , then a' and b' are expressed as

(2.16) 
$$a' = (a - \varepsilon_0(b))_+ + \varphi_0(b), \quad b' = \omega(\tilde{e}_0^{(\varepsilon_0(b) - a)_+}(b))$$

*Proof.* In Figure 2.1, the top right pattern does not occur. Then (2.16) is directly checked case by case.

Proof of Proposition 2.5. In view of  $\omega(s_1(p)) = s_0(\omega(p))$  for any p, the two formulas in (2.15) are equivalent. Henceforth we assume wt $(p) \ge 0$ , i.e.,  $\varepsilon_0(p) \ge \varphi_0(p)$ . Notice that a' in (2.16) is the number of + in the reduced 0-signature of  $(l-a, a) \otimes b \in B_l \otimes B_1$ . Consider Figure 2.2 with  $p = b_1 \otimes \cdots \otimes b_L$  and  $(x_1, x_2) = (l - x_2, x_2) \in B_l$ with any fixed p and  $x_2$ . It determines  $b'_1, \ldots, b'_L$  and  $(y_1, y_2)$  successively from the left by the local rule specified in Lemma 2.6 by taking l sufficiently large. From the above fact and the signature rule, we find that the choice  $(x_1, x_2) = (l, 0)$  leads to  $(y_1, y_2) = (l - \varphi_0(p), \varphi_0(p))$  which is called  $v_l$  in (2.11). Moreover the latter formula in (2.16) implies that the choice  $(x_1, x_2) = (l - a, a)$  yield  $b'_1 \otimes \cdots \otimes b'_L = \omega(\tilde{e}_0^{\varepsilon_0(p)-a}(p))$  provided  $a \leq \varepsilon_0(p)$ . By the assumption we are allowed to take  $a = \varphi_0(p)$ , leading to  $b'_1 \otimes \cdots \otimes b'_L = \omega(s_0(p))$ . But this must be  $T_l(p)$  due to (2.10).

**Example 2.7.** Take the path  $p = 121221221111 \in B_1^{\otimes 12}$  satisfying wt(p) > 0. To compute  $T_{\infty}(p) = \omega(s_0(p))$ , we display the 0-signature:

$$p = \underbrace{1}_{-} \underbrace{2}_{(+-)} \underbrace{1}_{(+-)} \underbrace{2}_{(+-)} \underbrace{1}_{(+-)} \underbrace{2}_{(+-)} \underbrace{1}_{(+-)} \underbrace{1$$

where the (+-) pairs to be eliminated successively to get the reduced 0-signature are indicated by parentheses. Those paired remain unchanged under  $s_0$ , and we thus find

Interchanging 1 and 2 here we obtain  $T_{\infty}(p) = 112112112221$ . In this example the composition  $\omega s_0$  has the effect of doing nothing for the unpaired 1 and interchanging the paired 1 and 2. When there remain unpaired + in the reduced 0-signature like p = 11122, the effect of  $\omega s_0$  is described in the same manner if those + are paired with - cyclically. Thus the formula (2.15) reproduces the description of  $T_{\infty}$  in terms of the "arc rule" in [15].

#### 3. Inverse scattering method

Here we use the rigged configurations and their bijective correspondence  $\phi$  with the highest paths  $\mathcal{P}_+$  [5, 6] summarized in Appendix A.

3.1. Action variable. The set of states  $\mathcal{P} = B_1^{\otimes L}$  of the periodic box-ball system is decomposed into the disjoint union according to the value of the conserved quantities  $\{E_l \mid l \in \mathbb{Z}_{\geq 1}\}$  which we called energy. Our aim here is to determine their spectrum by making a connection with rigged configurations. It will be attained in Proposition 3.4 and (3.5).

**Lemma 3.1.** For any path  $p \in \mathcal{P}$  with  $wt(p) \ge 0$ , there is an integer  $d \in \mathbb{Z}$  and a highest path  $p_+ \in \mathcal{P}_+$  such that  $p = T_1^d(p_+)$ .

The pair  $(d, p_+)$  is not unique in general even if d is restricted to  $0 \le d < L$ . The proof is elementary, and is illustrated along

**Example 3.2.** Take p = 2211221112122111221 with length L = 19. Regard the letters 1 and 2 in the doubled path  $p \otimes p$  as the arrows  $\nearrow$  and  $\searrow$  respectively, and construct a length 2L trail by following them. In the example, one has the following:



Now we introduce

(3.1) 
$$\mathcal{M} = \{ m = (m_j)_{j \ge 1} \mid m_j \in \mathbb{Z}_{\ge 0}, \sum_{j \ge 1} jm_j \le L/2 \}.$$

By the definition  $m_j = 0$  for  $j \gg 1$ , and we identify  $(m_1, \ldots, m_l, 0, 0, \ldots)$  with  $(m_1, \ldots, m_l)$ .  $\mathcal{M}$  is a finite set. We call its elements *action variables*.  $m = (m_j)$  is identified with the Young diagram in which  $m_j$  is the number of length j rows. We call the  $m_j \times j$  rectangle consisting of the length j rows a *block*.

Based on the KKR bijection we define the map

$$(3.2) \qquad \begin{array}{c} \mu: \ \mathcal{P} \longrightarrow \mathcal{M} \\ p \longmapsto m \end{array}$$

as follows. Find  $(d, p_+) \in \mathbb{Z} \times \mathcal{P}_+$  such that  $p = T_1^d(p_+)$  if  $\operatorname{wt}(p) \ge 0$  and  $\omega(p) = T_1^d(p_+)$  if  $\operatorname{wt}(p) < 0$  according to Lemma 3.1. Then m is obtained as the configuration part of the rigged configuration  $(m, J) = \phi(p_+)$ . As noted in Lemma 3.1, the choice of  $(d, p_+)$  is not unique. Therefore to make sense of the above definition, we have to guarantee that if  $T_1^d(p_+) = T_1^{d'}(p'_+)$  and  $\phi(p_+) = (m, J), \phi(p'_+) = (m', J')$ , then m = m' holds. To see this, note from the highest condition (2.8) that the situation  $T_1^{d-d'}(p_+) = p'_+$  can happen only when  $p_+ = r \otimes q$  and  $p'_+ = q \otimes r$  for some shorter highest paths  $q \in B_1^{\otimes d-d'}$  and  $r \in B_1^{\otimes L-d+d'}$ . (Without loss of generality,  $0 \le d-d' < L$  may be assumed.) Then the assertion m = m' is included in Lemma C.3.

**Example 3.3.** Consider p in Example 3.2, which can be expressed by the highest paths  $p_1, p_2$  and  $p_3$ . Computing  $\phi(p_i)$ , one finds that they all lead to  $m = (m_1, m_2, m_3) = (2, 2, 1)$ . The result is depicted as follows:



Let us relate the action variables m, namely the configurations in the KKR theory, to the energy  $E_l(p)$  of a path p determined by (2.10).

**Proposition 3.4.** For any path  $p \in \mathcal{P}$ , its energy is expressed as

(3.3) 
$$E_l(p) = \sum_{k \ge 1} \min(l, k) m_k$$

in terms of the action variable  $m = (m_j) = \mu(p) \in \mathcal{M}$ .

The proof reduces to the highest case  $p \in \mathcal{P}_+$ , and is given in Appendix B. According to Remark B.2,  $m = (m_j)$  has the meaning of *soliton content*, i.e., there are  $m_j$  solitons with length j. See also Example 4.5. The right hand side of (3.3) is the number of boxes in the first l columns of the Young diagram corresponding to m. Proposition 3.4 determines the range of the energy, i.e., the spectrum of the periodic box-ball system, in terms of the action variable  $m \in \mathcal{M}$  (3.1). In particular it implies the property:

$$0 \le E_1(p) < E_2(p) < \dots < E_s(p) = E_{s+1}(p) = \dots = M,$$

where  $M = \sum_{k\geq 1} km_k$  and s is the greatest integer such that  $m_s > 0$ , or equivalently, the length of the longest (top) row of the Young diagram for m. The relation (3.3) also determines m from  $\{E_l\}$ . Combining Proposition 3.4 with Theorem 2.2 and Proposition 2.3, we obtain

**Corollary 3.5.** The action variable is invariant under the time evolutions and the extended affine Weyl group. Namely,  $\mu(T_l(p)) = \mu(w(p)) = \mu(p)$  for any  $l \in \mathbb{Z}_{\geq 1}$  and  $w \in \widetilde{W}(A_1^{(1)})$ .

For each  $m = (m_j) \in \mathcal{M}$ , we introduce the corresponding "level set", namely, the set of paths p characterized by  $\mu(p) = m$ . Rephrasing the condition in terms of the energy by Proposition 3.4, we put (3.4)

$$\widehat{\mathcal{P}}(m) = \{ p \in \mathcal{P} \mid E_l(p) = \sum_{k \ge 1} \min(l, k) m_k \text{ for any } l \}, \quad \mathcal{P}_+(m) = \widehat{\mathcal{P}}(m) \cap \mathcal{P}_+.$$

One has the disjoint union decomposition:

(3.5) 
$$\mathcal{P} = \sqcup_{m \in \mathcal{M}} \widehat{\mathcal{P}}(m), \quad \mathcal{P}_+ = \sqcup_{m \in \mathcal{M}} \mathcal{P}_+(m).$$

From Corollary 3.5, we see that each  $\widehat{\mathcal{P}}(m)$  is invariant under time evolutions  $T_l$  as well as the extended affine Weyl group  $\widetilde{W}(A_1^{(1)})$ . (On the other hand highest paths can not remain highest under them in general.) In the KKR algorithm,  $M = \sum_{k \ge 1} km_k$  is the number of  $2 \in B_1$  contained in a highest (hence wt $(p) \ge 0$ ) path. Thus we find

$$\operatorname{wt}(\widehat{\mathcal{P}}(m)) = \{p_{\infty}\Lambda_1, -p_{\infty}\Lambda_1\}, \quad \operatorname{wt}(\mathcal{P}_+(m)) = \{p_{\infty}\Lambda_1\},\$$

where  $p_{\infty} = L - 2M \ge 0$  is the limiting (minimum) value of the vacancy number (3.7). Thus we have a further decomposition with respect to the weights:

$$\widehat{\mathcal{P}}(m) = \begin{cases} \mathcal{P}(m) \sqcup \omega(\mathcal{P}(m)) & \text{if } L > 2M, \\ \mathcal{P}(m) & \text{if } L = 2M, \end{cases}$$

where  $M = \sum_{k>1} km_k$  as above and

$$\mathcal{P}(m) = \{ p \in \widehat{\mathcal{P}}(m) \mid \mathrm{wt}(p) \ge 0 \} = \{ p \in \widehat{\mathcal{P}}(m) \mid \mathrm{wt}(p) = p_{\infty} \Lambda_1 \}$$

is a fixed weight subset of  $\widehat{\mathcal{P}}(m)$ . One has  $\mathcal{P}(m) \supset \mathcal{P}_+(m)$  and

(3.6) 
$$\{p \in \mathcal{P} \mid \operatorname{wt}(p) \ge 0\} = \sqcup_{m \in \mathcal{M}} \mathcal{P}(m),$$

$$|\mathcal{P}(m)| = \frac{|\widehat{\mathcal{P}}(m)|}{|\mathrm{wt}(\widehat{\mathcal{P}}(m))|}$$

for any  $m \in \mathcal{M}$ . This cardinality will be evaluated explicitly in (4.21) and (4.8). The set  $\mathcal{P}(m)$  is still invariant under any time evolution  $T_l$ . Now Lemma 3.1 is refined into

**Lemma 3.6.** For any path  $p \in \mathcal{P}(m)$ , there is an integer  $d \in \mathbb{Z}$  and a highest path  $p_+ \in \mathcal{P}_+(m)$  such that  $p = T_1^d(p_+)$ . Conversely, for any  $p_+ \in \mathcal{P}_+(m)$  and  $d \in \mathbb{Z}$ , one has  $T_1^d(p_+) \in \mathcal{P}(m)$ .

3.2. Angle variable. Here we construct the set of angle variables  $\mathcal{J}(m)$  for each prescribed value of the action variable  $m \in \mathcal{M}$ . We introduce the vacancy numbers

(3.7) 
$$p_j = L - 2\sum_{k>1} \min(j,k)m_k.$$

Given an element  $m \in \mathcal{M}$ , put

(3.8) 
$$H := \{ j \in \mathbb{Z}_{\geq 1} \mid m_j > 0 \} = \{ j_1 < j_2 < \dots < j_s \}$$

The vacancy numbers satisfy  $L = p_0 > p_1 > p_2 > \cdots > p_{j_s} = \cdots = p_{\infty} \ge 0$ . We introduce

(3.9) 
$$\mathcal{J} = \mathcal{J}(m) = \mathcal{J}_{j_1} \times \mathcal{J}_{j_2} \times \cdots \times \mathcal{J}_{j_s},$$

(3.10) 
$$\mathcal{J}_{j} = \{ (J_{i})_{i \in \mathbb{Z}} \mid J_{i} \in \mathbb{Z}, J_{i} \leq J_{i+1}, J_{i+m_{j}} = J_{i} + p_{j} \text{ for all } i \}.$$

Note that only  $m_j$  of  $(J_i)$ , say  $J_1, J_2, \ldots, J_{m_j}$ , are independent, and the Young diagram for the partition  $(J_{i+m_j} - J_i, J_{i+m_j-1} - J_i, \ldots, J_{i+1} - J_i)$  is contained in the  $m_j \times p_j$  rectangle for any *i*.

For  $k \in \mathbb{Z}_{\geq 1}$  we introduce a map

(3.11) 
$$\begin{aligned} \sigma_k : & \mathcal{J}_j \longrightarrow \mathcal{J}_j \\ & (J_i)_{i \in \mathbb{Z}} \longmapsto (J'_i)_{i \in \mathbb{Z}}, \\ & J'_i = J_{i+\delta_{j,k}} + 2\min(j,k). \end{aligned}$$

We extend  $\sigma_k$  to the map  $\overline{\mathcal{J}} \to \overline{\mathcal{J}}$  via  $\sigma_k(\overline{\mathcal{J}}) = \sigma_k(\mathcal{J}_{j_1}) \times \sigma_k(\mathcal{J}_{j_2}) \times \cdots \times \sigma_k(\mathcal{J}_{j_s})$ . Obviously the inverse  $\sigma_k^{-1}$  exists and  $\sigma_k \sigma_l = \sigma_l \sigma_k$  holds for any  $k, l \in \mathbb{Z}_{\geq 1}$ . Thus the set of elements  $\{\sigma_{j_1}^{n_1} \sigma_{j_2}^{n_2} \cdots \sigma_{j_s}^{n_s} \mid n_1, n_2, \dots, n_s \in \mathbb{Z}\}$  forms an infinite Abelian group  $\mathcal{A}$  isomorphic to  $\mathbb{Z}^s$ . We call an element of  $\mathcal{A}$  a *slide*, having in mind the index shift  $i \to i + \delta_{j,k}$  in (3.11). The origin of this curious map will be clarified in Lemma 4.2 in connection with the Bethe ansatz. Given two elements  $J, K \in \overline{\mathcal{J}}$ , say that J and K are *equivalent* and denote by  $J \simeq K$  if  $J = \sigma(K)$  for some  $\sigma \in \mathcal{A}$ . Now we define our main object  $\mathcal{J} = \mathcal{J}(m)$  by

$$(3.12) \mathcal{J} = \overline{\mathcal{J}} / \simeq .$$

An element of  $\mathcal{J}$  is called an *angle variable*. For  $J \in \overline{\mathcal{J}}$ , we will mostly use the same symbol  $J \in \mathcal{J}$  to denote its image in  $\mathcal{J}$  and [J] when emphasis is favorable. They are the data of the form  $J = (J_i^{(j)})$  with  $i \in \mathbb{Z}$  and  $j \in H$ . For  $J \in \mathcal{J}$  (resp.  $J \in \overline{\mathcal{J}}$ ), it is easily seen that  $J + d := (J_i^{(j)} + d)_{i \in \mathbb{Z}}$  with  $d \in \mathbb{Z}$  remains in  $\mathcal{J}$  (resp.  $\overline{\mathcal{J}}$ ). For  $J \in \overline{\mathcal{J}}(m)$  it is a good exercise to show

(3.13) 
$$\sigma_{j_1}^{m_{j_1}} \cdots \sigma_{j_s}^{m_{j_s}}(J) = J + L.$$

Therefore [J] = [J + L] represent the same angle variable.  $\mathcal{J}$  is a finite set. Its cardinality, namely the 'volume of the iso-level subset in the phase space' will be determined in (4.21) and (4.6). We introduce the time evolution of angle variables as follows:

(3.14) 
$$\begin{array}{cccc} T_l: & \mathcal{J}(m) & \longrightarrow & \mathcal{J}(m) \\ & & (J_i^{(j)})_{i \in \mathbb{Z}, j \in H} & \longmapsto & (J_i^{(j)} + \min(j, l))_{i \in \mathbb{Z}, j \in H}. \end{array}$$

This is a linear flow on the angle variables. It enjoys the commutativity  $T_l T_k = T_k T_l$ . The  $T_l$  here will be identified with the time evolution (2.10) on the paths  $\mathcal{P}$  in Theorem 3.12. Note that  $T_1^d(J) = J + d$  hence  $T_1^L(J) = J \in \mathcal{J}(m)$ . We also define  $T_l$  on  $\overline{\mathcal{J}}(m)$  by the same formula as (3.14).

It is convenient to depict the element  $(J_i^{(j)})_{i \in \mathbb{Z}, j \in H} \in \overline{\mathcal{J}}(m)$  as the Young diagram m whose rows are assigned with these numbers as follows:



This finite data is enough to recover the whole sequence  $(J_i^{(j)})_{i\in\mathbb{Z}}$  on account of the quasi-periodicity  $J_{i+m_j}^{(j)} = J_i^{(j)} + p_j$  (3.10). Then the slide  $\sigma_{j_k}^l$  (3.11) translates the infinite sequence  $(J_i^{(j_k)})_{i\in \mathbb{Z}}$  by *l* beside the block-wise shift  $2\min(j_k,\alpha)$  on  $\mathcal{J}_{\alpha}$ . It is helpful to depict the action as



Here  $\Delta_j = 2(j_k - j)l$ . Observe that  $\sigma_{j_k}^l$  only causes the uniform shift  $2j_k l$  on the upper blocks  $\mathcal{J}_{j_{k+1}}, \ldots, \mathcal{J}_{j_s}$ , whereas it relatively induces the back flow  $\Delta_{\alpha}$  on the lower blocks  $\mathcal{J}_{\alpha} = \mathcal{J}_{j_1}, \ldots, \mathcal{J}_{j_{k-1}}$ . A rigged configuration (m, J) with J = 0 $(J_i^{(j)})_{1 \leq i \leq m_i, j \in H} \in \operatorname{Rig}(m)$  (A.3) also has the data structure that can be depicted as in the above picture.

3.3. Direct and inverse scattering transforms. The direct and inverse scattering transforms are the maps between the path  $\mathcal{P}$  and the action-angle variables. In Section 3.1 we have constructed the map  $\mu$  (3.2) to find the action variable  $m \in \mathcal{M}$ for a given path  $p \in \mathcal{P}$ . It can be found either from the energy  $E_l(p)$  by (3.3) or from the rigged configuration of a highest path described under (3.2). It remains invariant under any time evolution as noted in Corollary 3.5.

In the remainder of Section 3, we shall only consider angle variables supposing that the action variable m has been determined. We assume that wt(p) > 0 and formulate the inverse scattering method for each  $\mathcal{P}(m)$  appearing in the decomposition (3.6). The other case wt(p) < 0 is reduced to it by

$$T_l(p) = \omega T_l(\omega(p)),$$

owing to Proposition 2.3.

First we formulate the direct scattering transform  $\Phi$  as follows:

(3.15) 
$$\begin{aligned} \Phi: \ \mathcal{P}(m) \longrightarrow \mathbb{Z} \times \mathcal{P}_{+}(m) \longrightarrow \overline{\mathcal{J}}(m) \longrightarrow \mathcal{J}(m) \\ p \longmapsto (d, p_{+}) \longmapsto \iota(J) + d \longmapsto [\iota(J) + d] \end{aligned}$$

Here the pair  $(d, p_+)$  is the one satisfying  $p = T_1^d(p_+)$  whose existence is assured in Lemma 3.6. Then the rigging  $J \in \operatorname{Rig}(m)$  is specified by the KKR bijection  $\phi(p_+) = (m, J)$ . The appearance of m here is guaranteed by Lemma 3.6. The map  $\iota$  is defined by

(3.16) 
$$\iota: \qquad \operatorname{Rig}(m) \longrightarrow \overline{\mathcal{J}}(m) \\ \left( (J_i^{(j_1)})_{1 \le i \le m_{j_1}}, \dots, (J_i^{(j_s)})_{1 \le i \le m_{j_s}} \right) \mapsto \left( (J_i^{(j_1)})_{i \in \mathbb{Z}}, \dots, (J_i^{(j_s)})_{i \in \mathbb{Z}} \right),$$

where the infinite sequence  $(J_i^{(j)})_{i\in\mathbb{Z}}$  is the one that extends  $(J_i^{(j)})_{1\leq i\leq m_j}$  quasiperiodically as  $J_{i+m_j}^{(j)} = J_i^{(j)} + p_j$  for all  $i \in \mathbb{Z}$ . The resulting sequence  $(J_i^{(j)})_{i\in\mathbb{Z}}$ automatically satisfies (3.10). The map  $\iota$  is just the embedding of the quasi-periodic extension, hence an injection.

Note that the  $(d, p_+)$  satisfying  $p = T_1^d(p_+)$  is not unique for a given p. Therefore to assure the well-definedness of  $\Phi$ , one has to show the  $\Rightarrow$  part of

**Proposition 3.7.** Let  $p_+, p'_+ \in \mathcal{P}_+(m)$  be the highest paths and  $J, J' \in \operatorname{Rig}(m)$  be the corresponding rigging, namely,  $\phi(p_+) = (m, J)$  and  $\phi(p'_+) = (m, J')$ . Then the following relation is valid:

(3.17) 
$$T_1^d(p_+) = T_1^{d'}(p'_+) \Longleftrightarrow \iota(J) + d \simeq \iota(J') + d' \in \overline{\mathcal{J}}(m),$$

The proof is available in Appendix C.

**Example 3.8.** Consider the length L = 25 path p = 2122112211221111222111122. According to Lemma 3.1, it can be expressed as  $p = T_1^{12}(p_+) = T_1^{19}(p'_+)$  in terms of the highest paths:

 $p_{+} = 1111222111122212211221122$   $p'_{+} = 111122212211221122111222.$ 

One applies the KKR bijection to get  $\phi(p_+) = (m, J)$  and  $\phi(p'_+) = (m, J')$ . The elements  $\iota(J) + 12$  and  $\iota(J') + 19$  of  $\overline{\mathcal{J}}(m)$  are depicted as



where 1, 3, 7, 15 on the left are the vacancy numbers (3.7) exhibited for convenience. The equivalence  $\iota(J) + 12 \simeq \iota(J') + 19$  is realized by  $\sigma_3(\iota(J) + 12) = \iota(J') + 19$ .

Next we show that the direct scattering map  $\Phi$  (3.15) is invertible, which yield the inverse scattering map  $\Phi^{-1}$ . For this two properties are to be established. First, any element in  $\overline{\mathcal{J}}(m)$  must be equivalent to the form  $\iota(J) + d$  for some  $J \in \operatorname{Rig}(m)$  and  $d \in \mathbb{Z}$ , which ensures the existence of an inverse image for the middle arrow in (3.15).  $(\phi(p_+) = (m, J).)$  Second, any two equivalent forms  $\iota(J) + d \simeq \iota(J') + d' \in \overline{\mathcal{J}}(m)$  must be pulled back to the same path in  $\mathcal{P}(m)$  in (3.15). The second property is nothing but the  $\Leftarrow$  part of Proposition 3.7. Thus it remains to verify the first property. This is done in

**Lemma 3.9.** For any  $\overline{J} \in \overline{\mathcal{J}}(m)$ , there exist  $d \in \mathbb{Z}$  and  $J \in \operatorname{Rig}(m)$  such that  $\overline{J} \simeq \iota(J) + d$ .

Proof. Let  $H = \{j_1 < \cdots < j_s\}$  be the list of lengths of rows in m as in (3.8). We give a concrete algorithm to repair  $\overline{J}$  by a slide so that  $\sigma_{j_1}^{n_1} \cdots \sigma_{j_s}^{n_s} \overline{J} = \iota(J) + d$  holds for some  $J \in \operatorname{Rig}(m)$  and  $d \in \mathbb{Z}$ . Such a slide is not unique in general but this relation can always be achieved gradually from the longer rows in m as explained below. First concentrate only on the two blocks  $\mathcal{J}_{j_{s-1}}$  and  $\mathcal{J}_{j_s}$ . In the slide  $\sigma_{j_{s-1}}$  (3.11), the quantity  $2\min(j, j_{s-1})$  is a common constant  $2j_{s-1}$  on them. Therefore by applying  $\sigma_{j_{s-1}}^n$  for some n, one can make the resulting  $(J_i^{(j_s-1)})_i \in \mathcal{J}_{j_{s-1}}$  and  $(J_i^{(j_s)})_i \in \mathcal{J}_{j_s}$  to satisfy  $d \leq J_1^{(j_{s-1})} \leq \cdots \leq J_{m_{j_{s-1}}}^{(j_{s-1})} \leq d + p_{j_{s-1}}$  and  $d \leq J_1^{(j_s)} \leq \cdots \leq J_{m_{j_s}}^{(j_s)} \leq d + p_{j_s}$  for some d. The effect of the common change

 $2 \min(j, j_{s-1})$  can be absorbed into d. Next one uses  $\sigma_{j_{s-2}}$  similarly to adjust  $\mathcal{J}_{j_{s-2}}$  to the amended  $\mathcal{J}_{j_{s-1}}, \mathcal{J}_{j_s}$  up to a redefinition of d without violating the foregoing adjustment among the latter two. This process, although the adjustment is not unique in general, works through until  $d \leq J_1^{(j)} \leq \cdots \leq J_{m_j}^{(j)} \leq d + p_j$  is achieved for all  $j \in H$  for some d.

## Example 3.10.



Here the vacancy numbers 1, 3, 9 have been shown only in the leftmost diagram.

From Lemma 3.9 and Proposition 3.7, it follows that the  $\Phi$  in (3.15) is a welldefined and invertible map. Taking the disjoint union over the action variable  $m \in \mathcal{M}$ , we obtain

**Theorem 3.11.** The map  $\Phi$  in (3.15) gives the bijection among the set of paths  $\sqcup_{m \in \mathcal{M}} \mathcal{P}(m)$  and the set of action-angle variables  $\sqcup_{m \in \mathcal{M}} \{(m, J) \mid J \in \mathcal{J}(m)\}$ .

3.4. Solution of initial value problem. Now we present our main theorem in this paper.

**Theorem 3.12.** The following commutative diagram is valid:

(3.18) 
$$\begin{array}{ccc} \mathcal{P}(m) & \stackrel{\Phi}{\longrightarrow} & \mathcal{J}(m) \\ T_l & & & \downarrow T_l \\ \mathcal{P}(m) & \stackrel{\Phi}{\longrightarrow} & \mathcal{J}(m) \end{array}$$

Here  $T_l$  on the left and the right are given by (2.10) and (3.14), respectively.

The proof is included in Appendix D. Since the map  $\Phi$  is invertible, Theorem 3.12 completes the solution of the initial value problem of the periodic box-ball system by the inverse scattering method. The time evolution on the paths  $\mathcal{P}(m)$  has been linearized in terms of the angle variables  $\mathcal{J}(m)$ . The number of computational steps required for executing  $\Phi^{-1} \circ T_l^t \circ \Phi$  is independent of t.

It also contains the solution of the initial value problem in the box-ball system on the semi-infinite lattice  $B_1 \otimes B_1 \otimes \cdots$  as the case  $L \to \infty$ . This limit is well-defined and drastically simplifies our construction so far. For any path  $b_1 \otimes b_2 \otimes \cdots$  obeying the boundary condition  $b_i = 1$  for  $i \gg 1$ , one can make  $1^{\otimes n} \otimes b_1 \otimes b_2 \otimes \cdots$  highest for some n. Thus the degree  $d \in \mathbb{Z}$  in (3.15) is always frozen to d = 0. In view of  $p_j \to \infty$  as  $L \to \infty$ , we do not make a non-trivial identification (3.12) under any slide (3.11). Consequently (3.15) just becomes  $\Phi : p \mapsto J$ , which is nothing but  $\phi$ . Namely, the direct and the inverse scattering transforms are the KKR bijection itself (without an upper bound on the rigging), reproducing the results in [23, 24] essentially.

**Example 3.13.** Let us take the length L = 19 path p in Example 3.2 and derive the time evolution

(3.19) 
$$T_2^{1000}(p) = 1211221112122211221, \quad T_3^{1000}(p) = 2112221211221112112$$

based on the inverse scattering formalism in Theorem 3.12. As noted in Example 3.2, we have the expression  $p = T_1^2(p_1)$  in terms of the highest path  $p_1$ . By computing the image of the KKR bijection  $\phi$  of  $p_1$ , one finds

$$p \xrightarrow{\Phi} 2 + \begin{array}{c} 1 \\ 3 \\ 9 \\ 8 \\ 4 \end{array} \begin{array}{c} 1 \\ 1 \\ 0 \\ 9 \\ 4 \end{array}$$

The vacancy numbers displayed on the left of the diagram for convenience will be omitted in the sequel. By using the linearized time evolution (3.14) one has

$$T_2^{1000}\Phi(p) = 2 + \begin{array}{|c|c|c|c|c|} & 2001 & \sigma_1^{222} & & & 1\\ & 2001 & \simeq & 2446 + & & 1\\ & 1008 & & & & 7\\ & 1004 & & & & 3 \end{array}$$

The last rigged configuration corresponds to the highest path p' = 1122112112211211221222. Therefore the map  $\Phi^{-1}$  sends the above scattering data to  $T_1^{2446}(p') = T_1^{14}(p')$  yielding the first result in (3.19). Similarly the calculation of  $T_3^{1000}(p)$  goes as follows:

The last rigged configuration corresponds to the highest path p'' = 1211221112112211222. Therefore the map  $\Phi^{-1}$  sends the above scattering data to  $T_1^{7150}(p'') = T_1^6(p'')$  yielding the second result in (3.19).

**Example 3.14.** We take the length L = 26 path p = 121122111221112211112211111 and compute  $T_3^{130}(p)$ . p is already highest.

$$p \xrightarrow{\Phi} 10 \xrightarrow{5} 0$$

$$10 \xrightarrow{5} 0$$

$$16 \xrightarrow{8} 0$$

$$16 \xrightarrow{8} 0$$

$$16 \xrightarrow{5} 0$$

$$16 \xrightarrow{8} 0$$

$$T_3^{130} \Phi(p) = 390 \xrightarrow{265} 260 \xrightarrow{719} 572 + 30$$

$$8 \xrightarrow{138} 0$$

$$138 \xrightarrow{130} 0$$

Since  $572 = 22 \times L$ , we find  $T_3^{130}(p) = p$ .

# 4. Relation with Bethe ansatz at q = 0

Here we show that our inverse scattering formalism originates in the Bethe ansatz at q = 0 [7]. The angle variables stem from the logarithmic branch of the string center equation and the time evolution is the straight motion of its solution. 4.1. Bethe ansatz at q = 0. Let us quickly recall the relevant results from the Bethe ansatz at q = 0. For the precise definitions and statements, we refer to [7]. The Bethe equation for the spin 1/2 one dimensional XXZ chain on length L periodic lattice reads [10]

(4.1) 
$$\left(\frac{\sin \pi (u_i + \sqrt{-1}\hbar)}{\sin \pi (u_i - \sqrt{-1}\hbar)}\right)^L = -\prod_{j=1}^M \frac{\sin \pi (u_i - u_j + 2\sqrt{-1}\hbar)}{\sin \pi (u_i - u_j - 2\sqrt{-1}\hbar)}$$

for  $1 \leq i \leq M$ , where  $0 \leq M \leq L/2$  is the number of down spins preserved by the Hamiltonian. The system is associated with the quantum affine algebra  $U_q(A_1^{(1)})$  with  $q = e^{-2\pi\hbar}$ . Fix  $m = (m_j) \in \mathcal{M}$ . String solutions are the ones in which  $\{u_1, \ldots, u_M\}$  are arranged as

(4.2) 
$$\bigcup_{j\geq 1} \bigcup_{1\leq \alpha \leq m_j} \bigcup_{u_{\alpha}^{(j)} \in \mathbb{R}} \{u_{\alpha}^{(j)} + \sqrt{-1}(j+1-2k)\hbar + \epsilon_{\alpha k}^{(j)} \mid 1 \leq k \leq j\},$$

where  $M = \sum_{j} jm_{j}$  and  $\epsilon_{\alpha k}^{(j)}$  stands for a small deviation.  $u_{\alpha}^{(j)}$  is the string center of the  $\alpha$  th string of length j. In this context, the data  $m \in \mathcal{M}$  is referred as the *string content*. Let H be as in (3.8). For the generic string solution, the Bethe equation is linearized at q = 0 into a logarithmic form called the *string center equation*:

(4.3) 
$$\sum_{k \in H} \sum_{\beta=1}^{m_k} A_{j\alpha,k\beta} u_{\beta}^{(k)} \equiv \frac{1}{2} (p_j + m_j + 1) \mod \mathbb{Z}$$

for  $j \in H$  and  $1 \leq \alpha \leq m_j$ . Here  $p_j$  is the vacancy number (3.7) and

(4.4) 
$$A_{j\alpha,k\beta} = \delta_{j,k}\delta_{\alpha,\beta}(p_j + m_j) + 2\min(j,k) - \delta_{j,k}$$

From (4.11), the matrix  $A = (A_{j\alpha,k\beta})$  is invertible under the condition  $m \in \mathcal{M}$ .

There are a number of conditions which the solutions of the string center equation (4.3) are to satisfy or to be identified thereunder. First, the Bethe vector depends on  $u_i$  only via  $e^{2\pi\sqrt{-1}u_i}$ . Therefore the string center should be understood as  $u_{\alpha}^{(j)} \in \mathbb{R}/\mathbb{Z}$  rather than  $\mathbb{R}$ . Second, the original Bethe equation (4.1) is symmetric with respect to  $u_1, \ldots, u_M$ , but their permutation does not lead to a new Bethe vector. Consequently, we should regard

$$(u_1^{(j)}, u_2^{(j)}, \dots, u_{m_j}^{(j)}) \in (\mathbb{R}/\mathbb{Z})^{m_j} / \mathfrak{S}_{m_j}$$

for each j. Last, we prohibit  $u_{\alpha}^{(j)} = u_{\beta}^{(j)}$  for  $1 \leq \alpha \neq \beta \leq m_j$  for any j. This is a remnant of the well-known constraint on the Bethe roots so that the associated Bethe vector does not vanish. To summarize, we consider *off-diagonal solutions*  $(u_{\alpha}^{(j)})$  to the string center equation (4.3) that live in

(4.5) 
$$(u_1^{(j)}, u_2^{(j)}, \dots, u_{m_j}^{(j)}) \in \left( (\mathbb{R}/\mathbb{Z})^{m_j} - \delta_{m_j} \right) / \mathfrak{S}_{m_j} \quad \text{for each } j,$$

where  $\delta_n = \{(v_1, \ldots, v_n) \in (\mathbb{R}/\mathbb{Z})^n \mid v_\alpha = v_\beta \text{ for some } 1 \leq \alpha \neq \beta \leq n\}$ . For simplicity we will often say *Bethe roots* to mean the off-diagonal solutions to the string center equation. Let  $\mathcal{U}(m)$  be the set of the Bethe roots having the string content m. For  $m \in \mathcal{M}$ , we introduce

(4.6) 
$$\Omega(m) = (\det F) \prod_{j \in H} \frac{1}{m_j} \binom{p_j + m_j - 1}{m_j - 1}.$$

(4.7) 
$$F = (F_{j,k})_{j,k \in H}, \quad F_{j,k} = \delta_{j,k} p_j + 2\min(j,k)m_k.$$

In case  $H = \emptyset$  (i.e., m = (0, 0, ...)), we put  $\Omega(m) = 1$ . By expanding the determinant, it is easy to see  $\Omega(m) \in \mathbb{Z}$ . Moreover, (4.13) below tells that  $\Omega(m) \in \mathbb{Z}_{\geq 1}$  and (4.6) is also expressed as

(4.8) 
$$\Omega(m) = \frac{L}{p_{j_s}} \prod_{j \in H} \binom{p_j + m_j - 1}{m_j},$$

where  $p_{j_s} = L - 2M$  with  $M = \sum_j jm_j$ . In case  $p_{j_s} = 0$ , the combination  $\frac{L}{p_{j_s}} {p_{j_s} + m_{j_s} - 1 \choose m_{j_s}}$  is to be understood as  $\frac{L}{m_{j_s}}$ . Note that  $\Omega(m) = \det F$  given in (4.13) if  $m_j = 1$  for all  $j \in H$ .

**Theorem 4.1** ([7] Theorems 3.5, 4.9).

(4.9) 
$$\Omega(m) = |\mathcal{U}(m)| \quad m \in \mathcal{M},$$

(4.10) 
$$\sum_{m \vdash M} \Omega(m) = \begin{pmatrix} L \\ M \end{pmatrix} \quad 0 \le M \le L/2,$$

where the sum extends over  $m_1, m_2, \ldots \in \mathbb{Z}_{\geq 0}$  such that  $\sum_j jm_j = M$ .

The derivation of (4.9) is due to the Möbius inversion trick. Actually (4.10) is known to hold for any  $M \ge 0$  if the symbol  $\binom{\alpha}{\beta}$  is interpreted as the generalized binomial coefficient  $\alpha(\alpha-1)\cdots(\alpha-\beta+1)/\beta!$ . The combined identity  $\sum_m |\mathcal{U}(m)| = \binom{L}{M}$  is called the combinatorial completeness of the string hypothesis at q = 0.

We include the formulas needed here and in Sections 4.2 and 4.3. Fix  $l \in \mathbb{Z}_{\geq 1}$ .

(4.11) 
$$\det A = (\det F) \prod_{j \in H} (p_j + m_j)^{m_j - 1}.$$

(4.12) 
$$\det A[k\beta] = (\det F[k]) \prod_{j \in H} (p_j + m_j)^{m_j - 1} \ (k \in H, \ 1 \le \beta \le m_k),$$

(4.13) 
$$\det F = L p_{j_1} p_{j_2} \cdots p_{j_{s-1}},$$

(4.14)

$$\det F[j_{n+1}] - \det F[j_n] = \frac{p_{i_s}(i_{n+1} - i_n)}{p_{i_{n+1}}p_{i_n}} \det F \quad (0 \le n \le s - 1).$$

Here the matrix  $A[k\beta]$  is obtained from  $A = (A_{j\alpha,k\beta})$  (4.4) by replacing the  $k\beta$  th column by  $\vec{h}$ . In (4.14), we are using the notation  $i_0 = 0$  and  $i_n = \min(l, j_n)$  for  $1 \leq n \leq s$ .  $p_{i_0} = p_0 = L$  as noted after (3.8). The matrix F[k] is obtained from  $F = (F_{j,k})$  (4.7) by replacing the column for  $k \in H$  by the column vector  $\vec{h}'$ . (We understand  $F[j_0] = 0$ .) Here the vectors  $\vec{h}$  and  $\vec{h}'$  are given by

(4.15) 
$$\vec{h} = (\min(j,l))_{j \in H, 1 \le \alpha \le m_j} \in \mathbb{Z}^{m_{j_1} + \dots + m_{j_s}}, \quad \vec{h}' = (\min(j,l))_{j \in H} \in \mathbb{Z}^s,$$

which are dependent on l. Note that (4.14) is vanishing unless  $0 \le n \le t$ , where t is the maximum integer such that  $i_{t+1} > i_t$ . By the definition  $0 \le t \le s-1$ . The

relations (4.11) and (4.13) are derived by noting

(4.16) 
$$\sum_{k \in H, 1 \le \beta \le m_k} A_{j\alpha,k\beta} = \sum_{k \in H} F_{j,k} = L$$

See also eq. (3.8) and (3.10) in [7]. Combining (4.11) and (4.12), we find

(4.17) 
$$\frac{\det A[j\alpha]}{\det A} = \frac{\det F[j]}{\det F} \quad (j \in H, \ 1 \le \alpha \le m_j).$$

The property det  $A \ge 0$  is obvious from (4.11) and (4.13) under the condition  $m \in \mathcal{M}$ .

4.2. Solitons as strings. Let us uncover the origin of our inverse scattering formalism in the Bethe ansatz. Note that the action variable  $m \in \mathcal{M}$  has emerged in two independent contexts in  $\mathcal{J}(m)$  and  $\mathcal{U}(m)$ . In the former it represents the soliton content of a path whereas in the latter it is the string content of the Bethe equation. The key to their link is the map:

(4.18) 
$$\begin{array}{cccc} \Psi : \ \overline{\mathcal{J}}(m) & \longrightarrow & \mathcal{U}(m) \\ & J & \longmapsto & \vec{u}, \end{array}$$

where  $\vec{u} = (u_{\alpha}^{(j)})_{j \in H, 1 \leq \alpha \leq m_j}$  is determined from  $J = (J_i^{(j)})_{i \in \mathbb{Z}, j \in H} \in \overline{\mathcal{J}}(m)$  as the solution of the linear equation:

$$(4.19) \quad \sum_{k \in H} \sum_{\beta=1}^{m_k} A_{j\alpha,k\beta} u_{\beta}^{(k)} = \frac{1}{2} (p_j + m_j + 1) + J_{\alpha}^{(j)} + \alpha - 1 \quad (j \in H, \ 1 \le \alpha \le m_j).$$

We write it as  $A\vec{u} = \vec{c} + \vec{J} + \vec{\rho}$ , where  $\vec{c} = (c_{\alpha}^{(j)})$  corresponding to  $(p_j + m_j + 1)/2$ is a constant vector having the components that are independent of the index  $\alpha$ .  $\vec{\rho} = (\rho_{\alpha}^{(j)} = \alpha - 1)$  is also a constant vector<sup>2</sup>. The equation (4.19) is the string center equation (4.3) that corresponds to a prescribed logarithmic branch of the Bethe equation. To make sense of (4.18), we are to check the off-diagonal condition (4.5) on  $\vec{u}$ . To do this it is useful to grasp the structure of the matrix A. For example in case  $H = \{1, 2, 3\}$ , it looks as  $(\mathcal{P}_j = p_j + m_j \text{ for short})$ 

(	$\mathcal{P}_1 + 1$		1	2		2	2		2  )	١
	:	·	:	:		:	:		:	
	1		$\mathcal{P}_1 + 1$	2		2	2		2	
	2		2	$\mathcal{P}_2 + 3$		3	4	• • •	4	
	:		:	:	·	:	:		:	
	2		2	3		$\mathcal{P}_{2} + 3$	4		4	
	2		2	4		4	$\mathcal{P}_3 + 5$	• • •	5	
	:		:	:	·	:	:		:	
ĺ	2		2	4		4	5		$\mathcal{P}_3 + 5$ /	/

which consists of 9 sub-matrices of size  $m_i \times m_j$   $(1 \le i, j \le 3)$ . Thus (4.19) leads to  $(p_j + m_j)(u_{\alpha}^{(j)} - u_{\beta}^{(j)}) = J_{\alpha}^{(j)} - J_{\beta}^{(j)} + \alpha - \beta$ . From this and the condition (3.10), we confirm that  $\Psi$  produces a specific array of real numbers  $(u_{\alpha}^{(j)})$  that obey

<sup>&</sup>lt;sup>2</sup>The vector  $\vec{c}$  plays a role only in Proposition 4.11, and  $\alpha - 1$  in (4.19) can be replaced with  $\alpha + a$  for any integer a.

 $a \leq u_1^{(j)} < u_2^{(j)} < \cdots < u_{m_j}^{(j)} < a+1$  for some  $a \in \mathbb{R}$ , which may indeed be viewed as a representative element of  $\mathcal{U}(m)$ .

Varying  $J \in \overline{\mathcal{J}}(m)$  to J' changes the image  $\vec{u} = \Psi(J)$  to  $\vec{u}' = \Psi(J')$  as an array of real numbers. But  $\vec{u} = (u_{\alpha}^{(j)})$  and  $\vec{u}' = (u_{\alpha}^{\prime(j)})$  can be regarded as the same element in  $\mathcal{U}(m)$  under the identification by  $\mathfrak{S}_{m_j}$  in (4.5)<sup>3</sup>. From the above property of  $\Psi$ , such an event takes place if and only if  $\vec{u}' = \tilde{\sigma}_{j_1}^{n_1} \cdots \tilde{\sigma}_{j_s}^{n_s}(\vec{u})$  for some  $n_1, \ldots, n_s \in \mathbb{Z}$ , where  $\tilde{\sigma}_k$  is defined by

$$\tilde{\sigma}_k: (u_1^{(j)}, u_2^{(j)}, \dots, u_{m_j}^{(j)}) \longmapsto \begin{cases} (u_2^{(j)}, \dots, u_{m_j}^{(j)}, u_1^{(j)} + 1) & \text{if } j = k, \\ (u_1^{(j)}, u_2^{(j)}, \dots, u_{m_j}^{(j)}) & \text{if } j \neq k. \end{cases}$$

Since the matrix A is invertible, the effect of  $\tilde{\sigma}_k$  is translated to that on  $J \in \overline{\mathcal{J}}(m)$ .

**Lemma 4.2.** If  $\vec{u} = \Psi(J)$ , then  $\tilde{\sigma}_k(\vec{u}) = \Psi(\sigma_k(J))$ , where  $\sigma_k$  is the slide defined in (3.11).

*Proof.* Set  $\vec{u}' = \tilde{\sigma}_k(\vec{u}) = \Psi(J')$ . By the definition we have  $A\vec{u} = \vec{c} + \vec{J} + \vec{\rho}$  and  $A\vec{u}' = \vec{c} + \vec{J}' + \vec{\rho}$ . In view of the structure of A, we find for  $j \neq k$  that  $J_{\alpha}^{(j)} = J_{\alpha}^{(j)} + A_{j\alpha,km_k} = J_{\alpha}^{(j)} + 2\min(j,k)$ . On the other hand, the k th block of  $\vec{c} + \vec{J}' + \vec{\rho} = A\vec{u}'$  is evaluated as  $(c = c_1^{(k)})$ 

$$\begin{pmatrix} c+J_{1}^{\prime(k)} \\ c+J_{2}^{\prime(k)}+1 \\ \vdots \\ c+J_{m_{k}}^{\prime(k)}+m_{k}-1 \end{pmatrix}$$

$$= \begin{pmatrix} p_{k}+m_{k}+2k-1 & \cdots & 2k-1 \\ \vdots & \ddots & \vdots \\ 2k-1 & \cdots & p_{k}+m_{k}+2k-1 \end{pmatrix} \begin{pmatrix} u_{2}^{(k)} \\ \vdots \\ u_{m_{k}}^{(k)} \\ u_{1}^{(k)}+1 \end{pmatrix} + \sum_{j(\neq k), 1 \leq \beta \leq m_{j}} \begin{pmatrix} A_{k1,j\beta} \\ A_{k2,j\beta} \\ \vdots \\ A_{km_{k},j\beta} \end{pmatrix} u_{\beta}^{(j)}$$

$$= \begin{pmatrix} c+J_{2}^{(k)}+1+2k-1 \\ \vdots \\ c+J_{m_{k}}^{(k)}+m_{k}-1+2k-1 \\ \vdots \\ c+J_{1}^{(k)}+p_{k}+m_{k}+2k-1 \end{pmatrix},$$

where the last equality is due to the k th block of the relation  $A\vec{u} = \vec{c} + \vec{J} + \vec{\rho}$ . Noting the quasi-periodicity  $J_1^{(k)} + p_k = J_{m_k+1}^{(k)}$ , this result is expressed as  $J_i^{\prime(k)} = J_{i+1}^{(k)} + 2k$  for  $1 \leq i \leq m_k$ . Unifying the formulas we find that  $J' = (J_i^{\prime(j)})$  is obtained from  $J = (J_i^{(j)})$  by  $J_i^{\prime(j)} = J_{i+\delta_{j,k}}^{(j)} + 2\min(j,k)$  in agreement with (3.11).

We have established that  $\Psi(J)$  and  $\Psi(J')$  represent the same element in  $\mathcal{U}(m)$  if and only if  $J \simeq J' \in \overline{\mathcal{J}}(m)$ . From the definition (3.12), we obtain

<sup>&</sup>lt;sup>3</sup>A proper treatment of this is to introduce the set of array of real numbers  $\overline{\mathcal{U}}(m)$  that projects onto  $\mathcal{U}(m)$  under the identification scheme in an analogous way from  $\overline{\mathcal{J}}(m)$  to  $\mathcal{J}(m)$ . However we skip it here supposing no confusion might arise.

**Theorem 4.3.** The map  $\Psi$  (4.18) induces the bijection between the set of angle variables  $\mathcal{J}(m)$  and the set of off-diagonal solutions  $\mathcal{U}(m)$  to the string center equation (4.3).

The induced bijection will also be denoted by  $\Psi$ . Now we are able to introduce the time evolution  $T_l$  of the Bethe root  $\vec{u} = \Psi(J) \in \mathcal{U}(m)$  by  $T_l(\vec{u}) = \Psi(T_l(J))$ using (3.14).

**Corollary 4.4.** Theorem 3.12 is extended to the following commutative diagram:

(4.20) 
$$\begin{array}{ccc} \mathcal{P}(m) & \stackrel{\Phi}{\longrightarrow} & \mathcal{J}(m) & \stackrel{\Psi}{\longrightarrow} & \mathcal{U}(m) \\ T_l & & T_l & & T_l \\ \mathcal{P}(m) & \stackrel{\Phi}{\longrightarrow} & \mathcal{J}(m) & \stackrel{\Psi}{\longrightarrow} & \mathcal{U}(m) \end{array}$$

Their cardinality is given by

(4.21) 
$$|\mathcal{P}(m)| = |\mathcal{J}(m)| = |\mathcal{U}(m)| = \Omega(m).$$

The result  $|\mathcal{P}(m)| = \Omega(m)$  endows the character formula in [7] with the quasiparticle interpretation. The quantity  $\Omega(m)$  originally associated with the string content *m* turns out to be the number of states in the periodic box-ball system having the prescribed soliton content *m*. Then the identity (4.10) implies that weight spaces of the  $sl_2$  module  $(\mathbb{C}^2)^{\otimes L}$  are decomposed into subspaces spanned by isoenergy states in the periodic box-ball system. These observations and Proposition 3.4 are quantitative supports to identify strings with solitons. The identification also agrees with the physical picture [12] that the strings represent bound states of magnons over the ferromagnetic vacuums  $111 \dots 111$  or  $222 \dots 222$ . The fact  $|\mathcal{P}(m)| = \Omega(m)$  has also been shown in [15] by a different approach.

**Example 4.5.** Consider the length L = 8 paths p with wt(p) = 0. There are  $\binom{8}{4} = 70$  such paths, which are grouped into  $\mathcal{P}(m)$  according to the 5 elements  $m = (m_1, m_2, \ldots) \in \mathcal{M}$  (3.1). Their cardinality  $|\mathcal{P}(m)| = \Omega(m)$  are given as follows:

The paths in  $\mathcal{P}(m)$  are listed as follows:

 $(m_1, m_2, m_3, m_4) \mid \mathcal{P}(m)$ 

-

=, =, =,	
(4,0,0,0)	12121212, 21212121
(2,1,0,0)	$T_1^n(12121122), T_1^n(12112122), T_1^n(11212122)$
(0,2,0,0)	11221122, 21122112, 22112211, 12211221
(1,0,1,0)	$T_1^n(12111222), T_1^n(11211222), T_1^n(11121222), T_1^n(11122122)$
(0,0,0,1)	$T_1^n(11112222)$

where  $T_1^n(p)$  stands for the 8 paths  $(n \in \mathbb{Z}_8)$  obtained by cyclic shifts of p. The path 11122122 for m = (1, 0, 1, 0) is the intermediate stage of the collision of the length 3 and 1 solitons.

4.3. **Periodicity.** The time evolution  $T_l$  is invertible as seen in (2.12) and the set of states  $\mathcal{P}$  is finite. Therefore every path  $p \in \mathcal{P}$  possesses the property  $T_l^N(p) = p$ for some integer  $N \geq 1$ . We say any such integer a *period* of p. The minimum period is called the *fundamental period* of p and denoted by  $\mathcal{N}^* = \mathcal{N}_l^*(p)$ . Every period is a multiple of the fundamental period  $\mathcal{N}^*$ . Here we establish a formula for the fundamental period under any  $T_l$  taking advantage of the linearization scheme (4.20). We assume  $p \in \sqcup_{m \in \mathcal{M}} \mathcal{P}(m)$  with no loss of generality thanks to (3.6) and Proposition 2.3.

For  $p \in \mathcal{P}(m)$ , let  $J = \Phi(p) \in \mathcal{J}(m)$ . The fundamental period  $\mathcal{N}^*$  of p under  $T_l$ is a function of l, the action variable m and the angle variable J. To be expository, we approach the fundamental period in two steps. First we show in Theorem 4.6 that there exists a value  $\mathcal{N} = \mathcal{N}_l(m)$  that is independent of J nonetheless making  $T_l^{\mathcal{N}}(p) = p$  hold for all  $p \in \mathcal{P}(m)$ . We call it the *generic period* inherent to the whole  $\mathcal{P}(m)$ . In general  $\mathcal{N}$  is yet a multiple of the fundamental period  $\mathcal{N}^*$ . Second we analyze the accidental symmetry gained by special J that makes  $\mathcal{N}^*$  a divisor of  $\mathcal{N}$ . It improves the idea of Theorem 4.6 and leads to the final result in Theorem 4.9. We begin by presenting the formula for the generic period  $\mathcal{N}$ .

(4.22) 
$$\mathcal{N} = \mathrm{LCM}\left(1, \bigcup_{j\alpha}' \frac{\mathrm{det}A}{\mathrm{det}A[j\alpha]}\right) = \mathrm{LCM}\left(1, \bigcup_{j}' \frac{\mathrm{det}F}{\mathrm{det}F[j]}\right)$$

where the matrices  $A, A[j\alpha]$  and F, F[j] are defined in (4.4), (4.7) and after (4.14). In particular,  $A[j\alpha]$  and F[j] are dependent on l. The union  $\bigcup'_{j\alpha}$  extends over those  $j \in H, 1 \leq \alpha \leq m_j$  such that det  $A[j\alpha] \neq 0$ . The union  $\bigcup'_j$  does over those  $j \in H$  such that det  $F[j] \neq 0$ . The least common multiple  $\operatorname{LCM}(r_1, \ldots, r_n)$  of nonzero rational numbers  $r_1, \ldots, r_n$  is the minimum positive rational number in  $\mathbb{Z}r_1 \cap \cdots \cap \mathbb{Z}r_n$ . Thus  $\operatorname{LCM}(1, r_1, \ldots, r_n)$  denotes the minimum positive integer in  $\mathbb{Z}r_1 \cap \cdots \cap \mathbb{Z}r_n$ . The equality of the two expressions is due to (4.17).

**Theorem 4.6.** For any path  $p \in \mathcal{P}(m)$ ,  $T_l^{\mathcal{N}}(p) = p$  is valid.

To be instructive we present two proofs utilizing  $\mathcal{U}(m)$  and  $\mathcal{J}(m)$  in the linearization scheme (4.20) although they are essentially the same. In both proofs we first consider a sufficient condition for  $T_l^{\mathcal{N}}(p) = p$  by regarding  $\mathcal{N}$  as an unknown, and confirm afterwards that the choice (4.22) fulfills it. The identities and definitions (4.11)–(4.15) work effectively.

Proof based on  $\mathcal{U}(m)$ . Suppose  $p \stackrel{\Phi}{\mapsto} J \stackrel{\Psi}{\mapsto} \vec{u}$  and  $T_l^{\mathcal{N}}(p) \stackrel{\Phi}{\mapsto} J' \stackrel{\Psi}{\mapsto} \vec{u}'$ . We have  $A\vec{u} = \vec{c} + \vec{J} + \vec{\rho}$  and  $A\vec{u}' = \vec{c} + \vec{J}' + \vec{\rho}$ . From (3.14) and (4.15), this leads to  $A(\vec{u}' - \vec{u}) = \vec{J}' - \vec{J} = \mathcal{N}\vec{h}$ . A sufficient condition for  $T_l^{\mathcal{N}}(p) = p$  is that all the components of  $\vec{v} := \vec{u}' - \vec{u}$  are integers. Since  $\vec{v} = \mathcal{N}A^{-1}\vec{h}$ , its  $j\alpha$  th component is  $v_{\alpha}^{(j)} = \mathcal{N} \det A[j\alpha]/\det A$ . Therefore the  $\mathcal{N}$  in (4.22) satisfies the condition.  $\Box$ 

Proof based on  $\mathcal{J}(m)$ . Take  $J \in \overline{\mathcal{J}}(m)$  such that  $[J] = \Phi(p) \in \mathcal{J}(m)$ . We are to consider a sufficient condition for  $[T_l^{\mathcal{N}}(J)] = [J]$ . This indeed happens if there exist integers  $n_1, \ldots, n_s$  such that  $\sigma_{j_1}^{n_1} \cdots \sigma_{j_s}^{n_s}(J) = T_l^{\mathcal{N}}(J) \in \overline{\mathcal{J}}(m)$ . We look for them by further presuming that  $r_a = n_a/m_{j_a}$  is an integer. Then from (3.10) and (3.11), the above relation on the block  $\mathcal{J}_{j_k}$  is expressed as

(4.23) 
$$p_{j_k}r_k + 2\sum_{a=1}^{s}\min(j_k, j_a)m_{j_a}r_a = \mathcal{N}\min(j_k, l).$$

In terms of the column vectors  $\vec{r} = {}^t(r_1, \ldots, r_s)$ ,  $\vec{h}'$  (4.15) and the matrix F (4.7), this is written as  $F\vec{r} = \mathcal{N}\vec{h}'$ . Its solution is given as  $r_a = \mathcal{N} \det F[j_a]/\det F$  by using the matrix F[k] appearing in (4.12). Thus the  $\mathcal{N}$  in (4.22) assures the existence of  $\forall r_a \in \mathbb{Z}$ .

The expressions (4.22) can be simplified. We employ the notation used in (4.14), e.g.,  $H = \{j_1, \ldots, j_s\}$  as in (3.8),  $i_n = \min(j_n, l)$   $(1 \le n \le s), i_0 = 0$  and  $0 \le t \le s - 1$  is the maximum integer such that  $i_{t+1} > i_t$ . Since  $F[j_{t+1}] = F[j_{t+2}] = \cdots = F[j_s]$  from (4.14), we have

$$(4.24)$$

$$\mathcal{N} = \operatorname{LCM}\left(1, \frac{\det F}{\det F[j_1]}, \frac{\det F}{\det F[j_2]}, \dots, \frac{\det F}{\det F[j_{t+1}]}\right)$$

$$= \operatorname{LCM}\left(1, \frac{\det F}{\det F[j_1]}, \frac{\det F}{\det F[j_2] - \det F[j_1]}, \dots, \frac{\det F}{\det F[j_{t+1}] - \det F[j_t]}\right)$$

$$= \operatorname{LCM}\left(1, \bigcup_{n=0}^{t} \frac{p_{i_{n+1}}p_{i_n}}{(i_{n+1} - i_n)p_{i_s}}\right),$$

where the second equality is an elementary property of LCM. The last line is obtained by (4.14). The union  $\cup_n'$  extends over those n such that  $\frac{p_{i_{n+1}}p_{i_n}}{(i_{n+1}-i_n)p_{i_s}}$  is finite. This caution is needed only if  $\operatorname{wt}(p) = 0$  and  $l \geq j_s$ , where one encounters  $p_{i_s} = p_{j_s} = 0$  in the denominator. In this case t = s - 1 and only n = s - 1 is allowed in the union so as to cancel the zero by  $p_{i_{n+1}} = 0$  in the numerator. This leads to  $\mathcal{N} = \operatorname{LCM}(1, \frac{p_{j_{s-1}}}{j_s - j_{s-1}})$ . Noting that  $p_{j_{s-1}} = p_{j_{s-1}} - p_{j_s} = 2(j_s - j_{s-1})m_{j_s}$ , we get  $\mathcal{N} = 2m_{j_s}$ . Thus  $T_l^{\mathcal{N}}(p) = p$  is certainly valid, for Proposition 2.5 and the remark following it imply that  $T_l^2(p) = p$ . Another simplification of (4.22) occurs at l = 1, where  $i_n = 1(n > 0)$  hence t = 0. In this case we have  $\mathcal{N} = L$ , which is again consistent with  $T_1^L(p) = p$  for any path p.

Being able to simplify (4.22) into (4.24) is gratifying. However as we will see in (4.31) in Section 4.5, it is instead the expression (4.22) that makes the generic period  $\mathcal{N}$  conceptual and elucidates the essence of the game. We note that the formula (4.22) is the simplest case of the most general one for  $A_n^{(1)}$  conjectured in eq.(8) in [2]. The expression (4.24) with  $l = \infty$  was first obtained in [15].

**Example 4.7.** Take the path p = 21121111221122111111222 of length L = 23. It belongs to  $\mathcal{P}(m)$  with  $m = \mu(p) = (m_1, m_2, m_3, m_4) = (1, 2, 0, 1)$  hence  $H = \{1, 2, 4\}$  and s = 3 in (3.8). The vacancy numbers (3.7) are  $(p_1, p_2, p_3, p_4) = (15, 9, 7, 5)$ . For  $1 \le l \le 4$ , the data  $i_1, i_2, i_3$  and t are listed as follows:

l	$i_1$	$i_2$	$i_3$	t
1	1	1	1	0
2	1	2	2	1
3	1	2	3	2
4	1	2	4	2
		25		

By convention we also have  $i_0 = 0$  and  $p_0 = L = 23$ . According to the last expression in (4.24),  $\mathcal{N}$  is calculated as

$$l = 1 : LCM(1, p_0) = LCM(1, 23) = 23,$$
  

$$l = 2 : LCM\left(1, \frac{p_1 p_0}{p_2}, p_1\right) = LCM\left(1, \frac{115}{3}, 15\right) = 345,$$
  

$$l = 3 : LCM\left(1, \frac{p_1 p_0}{p_3}, \frac{p_2 p_1}{p_3}, p_2\right) = LCM\left(1, \frac{345}{7}, \frac{135}{7}, 9\right) = 3105,$$
  

$$l = 4 : LCM\left(1, \frac{p_1 p_0}{p_4}, \frac{p_2 p_1}{p_4}, \frac{p_2}{2}\right) = LCM\left(1, 69, 27, \frac{9}{2}\right) = 621.$$

Actually these generic period  $\mathcal{N}$  coincide with the fundamental period  $\mathcal{N}^*$  of p under the respective time evolutions  $T_l$ . The reason will be explained in (4.27).

**Example 4.8.** Take the path p = 2122112211122111122 of length L = 25. This is the same path as that considered in Example 3.8. It belongs to  $\mathcal{P}(m)$  with  $m = \mu(p) = (m_1, m_2, m_3, m_4) = (1, 2, 1, 1)$ . The vacancy numbers read  $(p_1, p_2, p_3, p_4) = (15, 7, 3, 1)$ . This time we calculate  $\mathcal{N}$  along (4.22). The matrix F (4.7) reads

$$F = \begin{pmatrix} p_1 + 2m_1 & 2m_2 & 2m_3 & 2m_4 \\ 2m_1 & p_2 + 4m_2 & 4m_3 & 4m_4 \\ 2m_1 & 4m_2 & p_3 + 6m_3 & 6m_4 \\ 2m_1 & 4m_2 & 6m_3 & p_4 + 8m_4 \end{pmatrix} = \begin{pmatrix} 17 & 4 & 2 & 2 \\ 2 & 15 & 4 & 4 \\ 2 & 8 & 9 & 6 \\ 2 & 8 & 6 & 9 \end{pmatrix}.$$

If l = 3 for instance, the matrix F[j] is obtained from F by replacing its j th column with  $\vec{h}' = {}^t(1, 2, 3, 3)$  (4.15). Thus F[1], F[2], F[3] and F[4] look as

$$\begin{pmatrix} 1 & 4 & 2 & 2 \\ 2 & 15 & 4 & 4 \\ 3 & 8 & 9 & 6 \\ 3 & 8 & 6 & 9 \end{pmatrix}, \begin{pmatrix} 17 & 1 & 2 & 2 \\ 2 & 2 & 4 & 4 \\ 2 & 3 & 9 & 6 \\ 2 & 3 & 6 & 9 \end{pmatrix}, \begin{pmatrix} 17 & 4 & 1 & 2 \\ 2 & 15 & 2 & 4 \\ 2 & 8 & 3 & 6 \\ 2 & 8 & 3 & 9 \end{pmatrix}, \begin{pmatrix} 17 & 4 & 2 & 1 \\ 2 & 15 & 4 & 2 \\ 2 & 8 & 9 & 3 \\ 2 & 8 & 6 & 3 \end{pmatrix},$$

which lead to

$$\left(1, \frac{\det F}{\det F[1]}, \frac{\det F}{\det F[2]}, \frac{\det F}{\det F[3]}, \frac{\det F}{\det F[4]}\right) = \left(1, 125, \frac{875}{32}, \frac{875}{157}, \frac{875}{157}\right)$$

Taking the LCM of them, we get  $\mathcal{N} = 875$ . We list the result of such calculations for  $1 \leq l \leq 4$  in the following table.

l			LCM			$\mathcal{N}$
1	1,	25,	25,	25,	25	25
2	1,	$\frac{375}{7}$ ,	$\frac{375}{32}$ ,	$\frac{375}{32}$ ,	$\frac{375}{32}$	375
3	1,	125,	$\frac{875}{32}$ ,	$\frac{875}{157}$ ,	$\frac{875}{157}$	875
4	1,	375,	$\frac{2625}{32}$ ,	$\frac{2625}{157}$ ,	$\frac{875}{344}$	2625

For each l,  $T_l^{\mathcal{N}}(p) = p$  holds. Actually the  $\mathcal{N}$  listed here coincides with the fundamental period of p. See (4.27) for the reason.

As in Examples 4.7 and 4.8, the generic period  $\mathcal{N} = \mathcal{N}_l(m)$  (4.22) turns out to be the fundamental period  $\mathcal{N}_l^*(p)$  for a majority of paths  $p \in \mathcal{P}(m)$ . The event  $\mathcal{N}_l(m)/\mathcal{N}_l^*(p) \in \mathbb{Z}_{\geq 2}$  indicates some extra symmetry in the path p. The structure of our angle variable provides a lucid picture on the nature of the extra symmetry, and improves (4.22) into a formula for the fundamental period.

Consider the block  $J = (J_i)_{i \in \mathbb{Z}} \in \overline{\mathcal{J}}_j(m)$  (3.10) for  $j \in H = \{j_1, \ldots, j_s\}$ . Seek the maximum positive integer  $g = g_j$  such that

(4.25) 
$$\frac{p_j}{g}, \ \frac{m_j}{g} \in \mathbb{Z}_{\geq 1}, \quad J_{i+\frac{m_j}{g}} - J_i = \frac{p_j}{g} \quad \text{for any } i \in \mathbb{Z},$$

which is consistent with the condition  $J_{i+m_j} = J_i + p_j$  in (3.10). We say that such J has the order g symmetry, which means the finer quasi-periodicity than the original one by factor g. The case g = 1 corresponds to the previous treatment.

**Theorem 4.9.** For any path  $p \in \mathcal{P}(m)$ , let  $(J^{(j)})_{j \in H} = \Phi(p)$  be the angle variable. Suppose  $J^{(j)}$  has the order  $g_j$  symmetry. Then the fundamental period of p is given by

(4.26) 
$$\mathcal{N}^* = \mathrm{LCM}\Big(1, \bigcup_{j \in H}' \frac{\det F}{g_j \det F[j]}\Big),$$

where  $\cup_{j}'$  is the union over those j such that det $F[j] \neq 0$ .

Obviously  $\mathcal{N} \geq \mathcal{N}^*$  and the equality holds if (but not only if)  $g_j = 1$  for all  $j \in H$ .

*Proof.* A slight modification of the argument around (4.23) suffices. From the definition (4.25) of the symmetry of J, to validate  $T_l^N(p) = p$  or equivalently  $\sigma_{j_1}^{n_1} \cdots \sigma_{j_s}^{n_s}(J) = T_l^N(J)$ , it is necessary for  $r_a = n_a/(m_{j_a}/g_{j_a})$  to be an integer and to satisfy

$$\frac{p_{j_k}}{g_{j_k}}r_k + 2\sum_{a=1}^s \min(j_k, j_a)\frac{m_{j_a}}{g_{j_a}}r_a = N\min(j_k, l).$$

This is precisely (4.23) with  $r_a$  replaced by  $r_a/g_{j_a}$  and  $\mathcal{N}$  by N. Thus N must be chosen so that  $r_a = Ng_{j_a} \det F[j_a]/\det F$  be an integer for all  $1 \leq a \leq s$ . The  $\mathcal{N}^*$  in (4.26) is the minimum of such N.

Beside the generic case  $\forall g_j = 1$ , the next simplest situation is  $\forall g_j = g \geq 2$ ). It only happens when  $L/g \in \mathbb{Z}$ , and corresponds to the path of the form  $p = q^{\otimes g}$  for some  $q \in B_1^{\otimes L/g}$  without any such symmetry. (This statement is justified from Lemma C.1.) In this case,  $\mathcal{N}^*$  (4.26) coincides with the generic period  $\mathcal{N}$  (4.22) for q, which is consistent with  $T_l(p) = T_l(q)^{\otimes g}$ .

**Example 4.10.** We consider the path p treated in Example 3.14. We know that  $T_3^{130}(p) = p$ , and actually 130 is the fundamental period under  $T_3$ . Looking at the angle variable  $J = (J^{(1)}, J^{(2)}, J^{(3)}) = \Phi(p)$  given there, we find that  $J^{(1)}$  and  $J^{(2)}$  possess the order 2 symmetry. Thus  $(g_1, g_2, g_3) = (2, 2, 1)$  in the notation in (4.26). We have  $H = \{1, 2, 3\}, (m_1, m_2, m_3) = (2, 2, 1)$  and  $(p_1, p_2, p_3) = (16, 10, 8)$ . Thus the matrix F (4.7) reads

$$F = \begin{pmatrix} p_1 + 2m_1 & 2m_2 & 2m_3 \\ 2m_1 & p_2 + 4m_2 & 4m_3 \\ 2m_1 & 4m_2 & p_3 + 6m_3 \end{pmatrix} = \begin{pmatrix} 20 & 4 & 2 \\ 4 & 18 & 4 \\ 4 & 8 & 14 \end{pmatrix}.$$

We consider the time evolution  $T_3$ , for which the vector  $\vec{h}'(4.15)$  reads  $\vec{h}' = {}^t(1,2,3)$ . Replacing the columns of F with this, we get

$$F[1] = \begin{pmatrix} 1 & 4 & 2 \\ 2 & 18 & 4 \\ 3 & 8 & 14 \end{pmatrix}, \quad F[2] = \begin{pmatrix} 20 & 1 & 2 \\ 4 & 2 & 4 \\ 4 & 3 & 14 \end{pmatrix}, \quad F[3] = \begin{pmatrix} 20 & 4 & 1 \\ 4 & 18 & 2 \\ 4 & 8 & 3 \end{pmatrix}.$$

Substituting the determinants (det F, det F[1], det F[2], det F[3]) = (4160, 80, 288, 704) into the formulas for the generic period  $\mathcal{N}$  (4.22) and the fundamental period  $\mathcal{N}^*$ (4.26), we find (l = 3)

$$\mathcal{N} = \mathrm{LCM}\left(1, \frac{4160}{80}, \frac{4160}{288}, \frac{4160}{704}\right) = \mathrm{LCM}\left(1, 52, \frac{130}{9}, \frac{65}{11}\right) = 260,$$
$$\mathcal{N}^* = \mathrm{LCM}\left(1, \frac{4160}{2 \times 80}, \frac{4160}{2 \times 288}, \frac{4160}{704}\right) = \mathrm{LCM}\left(1, 26, \frac{65}{9}, \frac{65}{11}\right) = 130.$$

From (4.22), (4.25) and (4.26), we find

(4.27) 
$$\mathcal{N} = \mathcal{N}^*$$
 if  $\operatorname{GCD}(p_j, m_j) = 1$  for all  $j \in H$ ,

where we employ the convention  $\text{GCD}(0, m_{j_s}) = 1$  for the greatest common divisor when  $p_{j_s} = 0$ . Note that (4.27) is a sufficient but not a necessary condition for  $\mathcal{N} = \mathcal{N}^*$ . It explains the reason for  $\mathcal{N} = \mathcal{N}^*$  in Examples 4.7 and 4.8. For  $l = \infty$ , the fundamental period has also been studied in [15].

4.4. Bethe eigenvalue. The time evolution  $T_l$  in the periodic box-ball system is the q = 0 limit of the row transfer matrix  $T_l(\zeta)$ . Its eigenvalues are given by the analytic Bethe ansatz [25, 26]. Let  $Q(\theta) = \prod_{k=1}^{M} \sinh \pi(\theta - \sqrt{-1}u_k)$  be Baxter's Qfunction, where  $\{u_k\}$  satisfy the Bethe equation (4.1). We set  $q = e^{-2\pi\hbar}$  as before,  $\zeta = e^{2\pi\theta}$  and assume  $M \leq L/2$ . For the string solution (4.2), the  $q \to 0$  limit of the Bethe eigenvalue ([26] eq.(1.8)) coincides with that of the top term ([26] eq.(2.12)):

(4.28) 
$$\lim_{q \to 0} \frac{Q(\theta - l\hbar)}{Q(\theta + l\hbar)} = \zeta^{-E_l} \Lambda_l, \quad \Lambda_l = \exp\left(2\pi\sqrt{-1}\sum_{j\alpha} \min(j,l)(u_\alpha^{(j)} + \frac{1}{2})\right)$$

under an appropriate normalization. Here  $E_l = \sum_k \min(l,k)m_k$  is the right hand side of (3.3), and the sum  $\sum_{j\alpha}$  extends over  $j \in H$  and  $1 \le \alpha \le m_j$ .

**Proposition 4.11.** For any solution  $\vec{u} = (u_{\alpha}^{(j)})$  to the string center equation (4.3), the equality  $\Lambda_l^{\mathcal{N}} = 1$  is valid, where  $\mathcal{N}$  is the generic period (4.22).

Proof. Put  $\vec{v} = (v_{\alpha}^{(j)}) = \mathcal{N}A^{-1}\vec{h}$ , where  $\vec{h}$  is specified in (4.15). Its component is expressed as  $v_{\alpha}^{(j)} = \mathcal{N} \det A[j\alpha]/\det A = \mathcal{N} \det F[j]/\det F$ , where we have used (4.17). This is independent of the index  $\alpha$ . Moreover from (4.22), we may set  $v_{\alpha}^{(j)} = x_j$  for some  $x_j \in \mathbb{Z}$ . Further define the vectors  $\vec{c} = (c^{(j)})_{j\alpha}$  with  $c^{(j)} = (p_j + m_j + 1)/2$ , and  $\vec{e} = (\frac{1}{2})_{j\alpha}$ . Then the string center equation (4.3) is written as  $A\vec{u} = \vec{I} + \vec{c}$  for some integer vector  $\vec{I}$ . The  $\Lambda_l$  (4.28) is expressed as  $\Lambda_l = \exp(2\pi\sqrt{-1}t\vec{h}(\vec{u} + \vec{e})) = \exp(2\pi\sqrt{-1}t\vec{h}A^{-1}(\vec{I} + \vec{c} + A\vec{e}))$ . Since A is a symmetric matrix, we have  $\mathcal{N}(t\vec{h}A^{-1})_{j\alpha} = v_{\alpha}^{(j)} = x_j$ . Thus  $\Lambda_l^{\mathcal{N}} = \exp(2\pi\sqrt{-1}\sum_{j\alpha}x_j(\vec{I} + \vec{c} + A\vec{e}))$  is calculated using (4.4) as

$$\sum_{j\alpha} x_j (\vec{c} + A\vec{e})_{j\alpha} = \frac{1}{2} \sum_{j\alpha} x_j (p_j + m_j + 1 + \sum_{k\beta} A_{j\alpha,k\beta})$$
  
=  $\frac{1}{2} \sum_{j\alpha} x_j (p_j + m_j + 1 + p_j + m_j + 2 \sum_k \min(j,k)m_k - m_j)$   
=  $\sum_j x_j m_j (p_j + \frac{m_j + 1}{2} + \sum_k \min(j,k)m_k) \equiv 0 \mod \mathbb{Z}.$ 

Proposition 4.11 does not serve as a proof of Theorem 4.6. Nevertheless it has opened a route to create a conjectural formula for the generic period in a large class of generalized periodic box-ball systems [1, 2]. For  $l = \infty$ , Proposition 4.11 has been shown also in [27] independently.

4.5. **Discussion.** We write the string center equation (4.19) in the matrix form:

(4.29) 
$$A\vec{u} = \vec{c} + \vec{I} \in \mathbb{Z}^{\gamma},$$

where  $\gamma = m_{j_1} + \cdots + m_{j_s}$  for  $H = \{j_1, \ldots, j_s\}$  as in (3.8). The shifted array  $\vec{I} = \vec{J} + \vec{\rho}$  will also be called the angle variable here. See around (4.19) for  $\vec{\rho}$ . From (4.5) and (4.19), the  $\vec{I}$  belongs to the set

(4.30) 
$$(\mathcal{I}_{m_{j_1}} \times \cdots \times \mathcal{I}_{m_{j_s}})/\Gamma,$$

where  $\mathcal{I}_n = (\mathbb{Z}^n - \Delta_n)/\mathfrak{S}_n$  is the *n* dimensional lattice without the diagonal points  $\Delta_n = \{(z_1, \ldots, z_n) \in \mathbb{Z}^n \mid z_\alpha = z_\beta \text{ for some } 1 \leq \alpha \neq \beta \leq n\}$  identified under the permutations  $\mathfrak{S}_n$ .  $\Gamma = \bigoplus_{k \in H, 1 \leq \beta \leq m_k} \mathbb{Z} \vec{A}_{k\beta}$  is the  $\gamma$  dimensional lattice generated by the column vectors  $\vec{A}_{k\beta}$  of the matrix  $A = (A_{j\alpha,k\beta})$  (4.4) that characterizes the string center equation (4.29)<sup>4</sup>. The division by  $\Gamma$  originates in the identification of  $\vec{u}$  under  $u_\beta^{(k)} \to u_\beta^{(k)} + 1$  in (4.29). The time evolution  $T_l$  (3.14) on the angle variable  $\vec{I}$  is expressed as the linear flow  $T_l(\vec{I}) = \vec{I} + \vec{h}$  in terms of the *l*-dependent vector  $\vec{h} \in \mathbb{Z}^{\gamma}$  defined in (4.15). According to Theorem 4.3, it induces the time evolution of the Bethe roots  $T_l(\vec{u}) = \vec{u} + A^{-1}\vec{h}$ . They are summarized in the following table:

	Be the roots $\vec{u}$	Angle variables $\vec{I}$
mod	$\mathbb{Z}^\gamma$	$\Gamma = A \mathbb{Z}^{\gamma}$
$T_l$	$A^{-1}\vec{h}$	$ec{h}$

Here mod is the lattice under which the respective variables are to be identified save the permutations  $\mathfrak{S}_{m_{j_1}} \times \cdots \times \mathfrak{S}_{m_{j_s}}$ . Now it is transparent under what condition the time evolution  $T_l^{\mathcal{N}}$  becomes trivial. According to the above table, it is presented in the three equivalent forms:

(4.31) 
$$\mathcal{N}A^{-1}\vec{h} \in \mathbb{Z}^{\gamma}, \quad \mathcal{N}\vec{h} \in \Gamma, \quad \mathcal{N}F^{-1}\vec{h}' \in \mathbb{Z}^{s},$$

<sup>&</sup>lt;sup>4</sup>The column vectors  $\vec{A}_{k\beta}$  are independent because of detA > 0 under the condition  $m \in \mathcal{M}$  as noted in the end of Section 4.1.

where the last one is a contracted version of the first due to (4.17). See (4.15) for  $\vec{h'}$ . Similarly, the formula (4.26) for the fundamental period is rephrased as

(4.32) 
$$\mathcal{N}^* F^{-1} \vec{h}' \in G^{-1} \mathbb{Z}^s, \quad G = \operatorname{diag}(g_{j_1}, \dots, g_{j_s}).$$

In this way we arrive at the intrinsic meaning of the generic period (4.22) (rather than in the form (4.24)) and the fundamental period (4.26). Namely, they are the smallest positive integers  $\mathcal{N}$  and  $\mathcal{N}^*$  that make (4.31) and (4.32) valid. Depending on the choice of l in (4.15), the vectors  $\vec{h}$  and  $\vec{h}'$  (like Hamiltonian) encode various direction and speed of the straight motions in the set (4.30) corresponding to the time evolution  $T_l$ . The simplest among them is  $\vec{h} = {}^t(1, 1, \ldots, 1) \in \mathbb{Z}^{\gamma}$  for l = 1, for which  $L\vec{h} \in \Gamma$  holds because of (4.16). This fact corresponds to the simple property  $T_1^L(p) = p$  for any p. See (2.13). More generally, the generic and fundamental period under the combined time evolution  $T = \prod_l T_l^{\beta_l}$  ( $\beta_l \in \mathbb{Z}$ ) can be obtained by replacing  $\vec{h} = \vec{h}_l$  and  $\vec{h}' = \vec{h}'_l$  (4.15) with  $\sum_l \beta_l \vec{h}_l$  and  $\sum_l \beta_l \vec{h}'_l$  in (4.22) and (4.26), or equivalently, in (4.31) and (4.32). The set (4.30) is an analogue of the Jacobi variety on which the nonlinear dynamics on  $\mathcal{P}$  looks as a straight motion as in the classical theory of quasi-periodic solutions to soliton equations [20, 21]. The coefficient A in the string center equation plays the role of the period matrix. Its size  $\gamma$  is the total number of solitons, which is equal to the first energy  $E_1$  (3.3).

Under any set of selected time evolutions  $T_{l_1}, \ldots, T_{l_k}$ , one can describe the decomposition of  $\mathcal{P}(m)$  into the disjoint union of orbits:

$$\mathcal{P}(m) = \bigsqcup \{ \text{orbit under } T_{l_1}, \dots, T_{l_k} \}$$

From (4.30), each orbit here is in one to one correspondence with an element of

(4.33) 
$$(\mathcal{I}_{m_{j_1}} \times \cdots \times \mathcal{I}_{m_{j_s}})/(\Gamma + \mathbb{Z}\tilde{h}_{l_1} + \cdots + \mathbb{Z}\tilde{h}_{l_k}).$$

Equivalently, in terms of the original angle variable, the orbits are labeled by  $\mathcal{J}(m)/(T_{l_1}^{\mathbb{Z}}\cdots T_{l_k}^{\mathbb{Z}})$ , where the division means the identification of the elements connected by the time evolution (3.14).

Under the single time evolution  $T_l$ , the number of orbits contained in  $\mathcal{P}(m)$ is  $\Omega(m)/\mathcal{N}$  if the action variable  $m \in \mathcal{M}$  satisfies the generic condition (4.27). See (4.21) and (4.6) for  $\Omega(m)$ . Another simple situation is to include the whole family  $T_1, T_2, \ldots$ , which maximizes the orbits and minimizes their number. From (3.9)–(3.12) and (3.14) it is easy to see that  $\mathcal{J}(m)/(T_1^{\mathbb{Z}}T_2^{\mathbb{Z}}\cdots)$  is factorized as

(4.34) 
$$\widehat{\mathcal{J}}_{j_1} \times \widehat{\mathcal{J}}_{j_2} \times \cdots \times \widehat{\mathcal{J}}_{j_s}.$$

Here  $\widehat{\mathcal{J}}_j = \mathcal{J}_j/\sim$  is obtained from  $\mathcal{J}_j = \{(J_i)_{i\in\mathbb{Z}} \mid J_i \in \mathbb{Z}, J_i \leq J_{i+1}, J_{i+m_j} = J_i + p_j$  for all  $i\}$  in (3.10) by the identification  $(J_i) \sim (J'_i)$  defined by  $J'_i = J_{i+1}$  for all i or  $J'_i = J_i + 1$  for all i. The difference  $d_i = J_{i+1} - J_i + 1$  satisfies  $d_i \in \mathbb{Z}_{\geq 1}$  and  $d_{i+1} + d_{i+2} + \cdots + d_{i+m_j} = p_j + m_j$ , especially  $d_i = d_{i+m_j}$  for all  $i \in \mathbb{Z}$ . Thus the set  $\widehat{\mathcal{J}}_j$  is in one to one correspondence with the arrangements of  $p_j + m_j$  letters 1 and  $m_j$  letters 2 as

$$\overbrace{1\ldots1}^{d_1} 2 \overbrace{1\ldots1}^{d_2} 2 \overbrace{1\ldots1}^{d_3} \cdots \overbrace{1\ldots1}^{d_{m_j}} 2$$

that are inequivalent under the periodic boundary condition. No two letters 2 are allowed to be adjacent because of  $d_i \ge 1$ . Such arrangements are the states of

the periodic box-ball system containing amplitude 1 solitons only. In this way we obtain

(4.35) 
$$|\mathcal{J}(m)/(T_1^{\mathbb{Z}}T_2^{\mathbb{Z}}\cdots)| = C(p_{j_1}, m_{j_1})C(p_{j_2}, m_{j_2})\cdots C(p_{j_s}, m_{j_s}),$$

where C(p,m)  $(p \ge 0, m \ge 1)$  is the number of orbits in the size p + 2m periodic box-ball system containing m solitons of amplitude 1 only. Namely, the count of orbits under the entire family  $T_1, T_2, \ldots$  splits into individual blocks wherein all the solitons behave effectively as amplitude 1. The quantity C(p,m) is characterized as the number of orbits of the monomials  $x_1^{\mu_1} \cdots x_m^{\mu_m}$  under the cyclic shift  $x_i \to x_{i+1} (x_{i+m} = x_i)$  in the complete symmetric function  $h_p(x_1, \ldots, x_m)$ .

Finally we comment on the motion of the Bethe roots  $\vec{u} \mapsto T_l(\vec{u}) = \vec{u} + A^{-1}\vec{h}$ . The fusion transfer matrix corresponding to  $T_l$  does not change the Bethe vectors up to an overall scalar nor the associated Bethe roots. This does not contradict the motion of  $\vec{u}$  considered here since each path that we associate to  $\vec{u}$  is a monomial in  $(\mathbb{C}^2)^{\otimes L}$ , which is *not* a Bethe vector at q = 0 in general.

# 5. Summary

In this paper we solved the initial value problem in the periodic box-ball system by a unification of the combinatorial Bethe ansätze at q = 1 and q = 0. Section 2 gives the formulation of the periodic box-ball system in terms of crystal basis theory. The commutativity, energy conservation (Theorem 2.2) and the invariance under the extended affine Weyl group (Propositions 2.3 and 2.5) in the periodic setting are firstly shown explicitly in this paper. In Section 3, we introduced the action and angle variables in (3.2) and (3.12) and the linear time evolution (3.14) on the latter. The direct/inverse scattering map is defined in (3.15), which linearizes the dynamics as in Theorem 3.12. In Section 4, our inverse scattering formalism are linked with the Bethe ansatz by the key relation (4.18). The action-angle variables are in one to one correspondence with the off-diagonal solutions (4.5) to the string center equation (4.3) as summarized in Corollary 4.4. It has led to the explicit formula (4.21), (4.6) counting the states characterized either by soliton content (energy) or string content (configuration), which we identified in Proposition 3.4 and Section 4.2. As further applications, the generic (4.22) and fundamental (4.26)period and the number of disjoint orbits (4.35) under the commuting family of time evolutions are obtained (Theorems 4.6, 4.9). The Bethe eigenvalue is shown to be a root of unity related to the generic period (Proposition 4.11). These results are derived and understood most naturally from the intrinsic picture on the dynamics as a straight motion in the 'ultradiscrete Jacobi variety' (4.30). We expect that the essential features explored in this paper persist in the generalized periodic box-ball systems [1, 2].

**Acknowledgments** This work is partially supported by Grand-in-Aid for Scientific Research JSPS No.15540363.

### Appendix A. KKR bijection

Put  $B = B_{l_1} \otimes \cdots \otimes B_{l_L}$ . The set  $\mathcal{P}$  of our periodic box-ball system (2.7) corresponds to the B with the choice  $l_1 = \cdots = l_L = 1$ . Let

(A.1) 
$$\mathcal{P}_{+} = \{ p \in B \mid \tilde{e}_{1}p = 0 \}$$

be the set of highest elements. For the array of nonnegative integers  $m = (m_j) = (m_1, m_2, \ldots)$ , we put  $H = \{j \in \mathbb{Z}_{\geq 1} \mid m_j > 0\}$ . The data *m* is called a *configuration* if  $p_j \geq 0$  for all  $j \in H$ , where

(A.2) 
$$p_j = \sum_{i=1}^{L} \min(j, l_i) - 2 \sum_{k \ge 1} \min(j, k) m_k$$

is called the vacancy number. The set H is necessarily finite and we parameterize it as  $H = \{j_1 < \cdots < j_s\}$ . The data m is identified with the Young diagram containing  $m_j$  rows of length j, i.e., the  $m_j \times j$  rectangular block for each  $j \in H$ . We let  $\mathcal{M}$  denote the set of all the configurations m. These definitions agree with the earlier ones for  $\mathcal{P}_+$  (2.7),  $\mathcal{M}$  (3.1),  $p_j$  (3.7) and H (3.8) when  $l_1 = \cdots = l_L = 1$ . Define

(A.3)

$$\operatorname{Rig}(m) = \{ (J_i^{(j)})_{1 \le i \le m_j, j \in H} \in \mathbb{Z}^{m_{j_1}} \times \dots \times \mathbb{Z}^{m_{j_s}} \mid 0 \le J_1^{(j)} \le \dots \le J_{m_j}^{(j)} \le p_j \}.$$

When  $l_1 = \cdots = l_L = 1$  and the dependence on L is important, we write  $\operatorname{Rig}_L(m)$  in Appendices C and D.

The purpose of this appendix is to describe the Kerov-Kirillov-Reshetikhin (KKR) bijection  $\phi = \phi_B$  [5, 6]:

(A.4) 
$$\phi: \mathcal{P}_+ \longrightarrow \sqcup_{m \in \mathcal{M}} \{ (m, J) \mid J \in \operatorname{Rig}(m) \}.$$

An element of  $\operatorname{Rig}(m)$  is called *rigging*. Each value  $J_i^{(j)}$  is also said rigging. Similarly  $p_j - J_i^{(j)}$  is called co-rigging. The combined data (m, J) is called the rigged configuration. It is customary to depict it as the Young diagram m with each row attached with the rigging. We often exhibit the vacancy number  $p_j$  on the left of the block of width j. The riggings are arranged so as to decrease weakly downward within a block. Note that these definitions depend on the choice of B although we do not exhibit it explicitly. The choice other than  $l_1 = \cdots = l_L = 1$  is needed only in Proposition A.4.

**Example A.1.** For the highest paths  $p_1, p_2$  and  $p_3$  in Example 3.2, one has  $B = B_1^{\otimes 19}$  and



The original KKR bijection [5, 6] is the one between the rigged configurations and the Littlewood-Richardson tableaux. The bijection (A.4) is obtained through a simple transformation of the Littlewood-Richardson tableaux and the highest paths [9]. Here we illustrate  $\phi$  and  $\phi^{-1}$  casually along two examples rather than the systematic description which is already available in [5, 6, 28, 19]. Our convention here is opposite from [5] in the role of the rigging and co-rigging, and opposite from [19] in the order of tensor product.

Regard a rigged configuration as a multi set of the pairs (row length, attached rigging). For example, the leftmost one in Example A.1 is regarded as  $\{(3, 1), (2, 1), (2, 0), (1, 8), (1, 4)\}$ . Each element  $(j, \alpha)$  of the rigged configuration is called a *string* with length j and rigging  $\alpha$ . A string is *singular* if the co-rigging is zero, namely  $\alpha = p_j$ , which is the maximum allowed value in (A.3). We first illustrate the map  $\phi^{-1}$ .

**Example A.2.** We begin with the basic case  $B = B_1^{\otimes L}$ , where the vacancy number (A.2) reduces to (3.7). Consider the configuration m = (2, 1) for example. For L = 8, there are 6 rigged configurations depicted in the leftmost column of the following:



The procedure to obtain the highest paths by applying  $\phi^{-1}$  has been shown. Reading the numbers on the arrows backward, we find the image of those rigged configurations under  $\phi^{-1}$  as follows:

 $\begin{array}{l} 1\otimes 2\otimes 1\otimes 2\otimes 1\otimes 1\otimes 1\otimes 2\otimes 2,\\ 1\otimes 2\otimes 1\otimes 1\otimes 2\otimes 1\otimes 2\otimes 2,\\ 1\otimes 2\otimes 1\otimes 1\otimes 2\otimes 2\otimes 1\otimes 2\otimes 2,\\ 1\otimes 2\otimes 1\otimes 1\otimes 2\otimes 2\otimes 1\otimes 2\otimes 2\otimes 1\otimes 2,\\ 1\otimes 1\otimes 2\otimes 1\otimes 2\otimes 1\otimes 2\otimes 2\otimes 1\otimes 2\otimes 2,\\ 1\otimes 1\otimes 2\otimes 2\otimes 1\otimes 2\otimes 2\otimes 1\otimes 2,\\ 1\otimes 1\otimes 2\otimes 2\otimes 2\otimes 1\otimes 2\otimes 1\otimes 2.\end{array}$ 

All these paths satisfy the highest condition (2.8). The KKR algorithm for obtaining  $\phi^{-1}$  proceeds recursively as  $\phi_{B_1^{\circ L}}^{-1}(\mathrm{rc}) = \phi_{B_1^{\circ L-1}}^{-1}(\mathrm{rc}') \otimes a$  with  $a \in B_1 = \{1, 2\}$ . This relation is depicted as  $\mathrm{rc} \xrightarrow{a} \mathrm{rc}'$  in the above. We have  $a = 1 \in B_1$  and  $\mathrm{rc}' = \mathrm{rc}$  if the rigged configuration rc is free from singular strings. If there exist singular strings in rc, we set  $a = 2 \in B_1$ . In that case the new rigged configuration  $\mathrm{rc}'$  is obtained by replacing any one of the shortest singular string  $(j, \alpha = p_j)$  by  $(j-1, p'_{j-1})$ , where  $p'_{j-1}$  is the vacancy number (3.7) with L replaced by L-1 and  $m_i$  replaced  $m_i - \delta_{i,j} + \delta_{i,j-1}$ . Namely one removes a box from the shortest singular string and assings a new rigging to the shortened one so that it again becomes singular in the new environment. (If j = 1, just eliminate it.) Even when rc = rc', the vacancy number (3.7) is lowered by one by the change  $B_1^{\otimes L} \to B_1^{\otimes L-1}$ . So one must revise  $p_j$  in each step and keep track of L, which we did on the top line.

To apply our inverse scattering method, the description in Example A.2 suffices for  $\phi^{-1}$ . In order to cover the content of Proposition A.4 which is used in Proposition 3.4 and Lemma D.1, one needs to go beyond  $B = B_1^{\otimes L}$ . We explain it along

**Example A.3.** Take  $B = B_2 \otimes B_1 \otimes B_2 \otimes B_3 \otimes B_1$ . We show the procedure for obtaining a highest path in B.



This time we have drawn a pair of diagrams in each step. The right ones are the rigged configurations. The left ones keep track of the 'shape' of the paths. For example in the top left diagram, the list of its row lengths 2, 1, 2, 3, 1 encodes the indices in  $B = B_2 \otimes B_1 \otimes B_2 \otimes B_3 \otimes B_1$  where we start from. They are removed one by one from the bottom right as marked with •. The process corresponds to the reduction of the path shape as

$$B_{2} \otimes B_{1} \otimes B_{2} \otimes B_{3} \otimes B_{1} \to B_{2} \otimes B_{1} \otimes B_{2} \otimes B_{3} \to B_{2} \otimes B_{1} \otimes B_{2} \otimes B_{2} \to B_{2} \otimes B_{1} \otimes B_{2} \otimes B_{1} \to B_{2} \otimes B_{1} \to B_{2} \otimes B_{1} \otimes B_{2} \to B_{2} \otimes B_{1} \otimes B_{2} \to B_{2} \otimes B_{1} \to B_{2} \otimes B_{1} \to B_{2} \to B_{1} \to \emptyset.$$

In each step the vacancy numbers are revised by adopting the intermediate path shape as  $\{l_i\}$  in (A.2). When a box (marked with •) is removed from the k th column, in accordance with  $(\cdots) \otimes B_k \to (\cdots) \otimes B_{k-1}$ , the rigged configuration is left unchanged if there is no singular string with length  $\geq k$ . In this case we proceed to the next step by  $\xrightarrow{1}$ . If there exist singular strings with length  $\geq k$ , we have  $\xrightarrow{2}$ and remove a box (as marked with  $\star$ ) from any one of the shortest such string. The shortened new string shall be attached with the rigging so that it becomes singular in the new environment. The algorithm ends up with the pair of  $\emptyset$ . Reading the letters on the arrows backward, we get the sequence 11|2|11|221|2, where the symbol | separates those letters coming from different rows in the shape diagrams. Reversing the letters within  $|\cdot|$ , we find the image under  $\phi_B^{-1}$  as  $11 \otimes 2 \otimes 11 \otimes 122 \otimes 2 \in B$ , which is highest.

The algorithm illustrated in Example A.3 reduces to the simpler one in Example A.2 when  $l_1 = \cdots = l_L = 1$ .

We have seen that the KKR algorithm for  $\phi_B^{-1}$  is a removal process of a rigged configuration creating a highest path from its rightmost component. Naturally, the map  $\phi_B$  is an addition process building a rigged configuration by using the information of a highest path from its leftmost component. The rule of addition is easily inferred by looking at Example A.3 backward. One regards the given path  $11 \otimes 2 \otimes 11 \otimes 122 \otimes 2 \in B$  as the word 11|2|11|221|2 and adds a box (•) carrying these letters one by one from the left to form the prescribed shape diagram having the row lengths 2, 1, 2, 3, 2 from the top. If the letter is 1, one does nothing on the rigged configuration. If the letter is 2 and the change of the path shape is  $(\cdots) \otimes B_k \to (\cdots) \otimes B_{k+1}$ , one adds a box ( $\star$ ) to any one of the *longest* singular string  $(j, \alpha = p_j)$  among those  $j \geq k$ . The new string of length j+1 should be assigned with the maximal rigging so as to become singular in the new environment. If there is no such strings, one creates a singular string of length 1.

For the highest path  $p \in \mathcal{P} = B_1^{\otimes L}$  such that  $\phi_{\mathcal{P}}(p) = (m, J)$ , it is easy to see

$$\phi_{\mathcal{P}\otimes B_1^{\otimes n}}(p\otimes \overbrace{1\otimes \cdots \otimes 1}^n) = (m,J).$$

**Proposition A.4.** Let  $p = b_1 \otimes \cdots \otimes b_L \in \mathcal{P} = B_1^{\otimes L}$  be a highest path such that  $\phi(p) = (m, J)$ . For n sufficiently large, define  $\xi \in \mathcal{P} \otimes B_1^{\otimes n}$  by  $u_l \otimes (p \otimes 1^{\otimes n}) \simeq \xi \otimes u_l$ . Then  $\phi_{\mathcal{P} \otimes B_1^{\otimes n}}(\xi) = (m, I)$ , where the rigging  $I = (I_i^{(j)})$  is given by  $I_i^{(j)} = J_i^{(j)} + \min(l, j)$ .

In  $\xi$ , sufficiently many components on the right are also  $1 \in B_1$ .

*Proof.* We first show that  $\phi_{B_l \otimes \mathcal{P} \otimes B_1^{\otimes n}}(u_l \otimes p \otimes 1^{\otimes n}) = (m, I)$ . In fact, the change from  $\phi_{\mathcal{P} \otimes B_1^{\otimes n}}$  to  $\phi_{B_l \otimes \mathcal{P} \otimes B_1^{\otimes n}}$  increases the vacancy number  $p_j$  (A.2) to  $p_j$ +min(l, j). Exactly the same increment has occurred from the rigging J to I. Therefore the co-rigging of (m, J) for  $\mathcal{P} \otimes B_1^{\otimes n}$  and the co-rigging of (m, I) for  $B_l \otimes \mathcal{P} \otimes B_1^{\otimes n}$  coincide, leading to  $\phi_{B_l \otimes \mathcal{P} \otimes B_1^{\otimes n}}(u_l \otimes p \otimes 1^{\otimes n}) = (m, I)$ . On the other hand we have  $\phi_{B_l \otimes \mathcal{P} \otimes B_1^{\otimes n}}(u_l \otimes p \otimes 1^{\otimes n}) = \phi_{\mathcal{P} \otimes B_1^{\otimes n} \otimes B_l}(\xi \otimes u_l)$  by Lemma 8.5 in [28]. In view of the KKR algorithm, the last one is equal to  $\phi_{\mathcal{P} \otimes B_2^{\otimes n}}(\xi)$ . □

The image

$$\phi_B^{-1}((m,J)) = (l_1 - x_1, x_1) \otimes \cdots \otimes (l_L - x_L, x_L) \in B = B_{l_1} \otimes \cdots \otimes B_{l_L}$$

can also be described by the piecewise linear formula:

(A.5)  
$$x_n = \tau_1(n) - \tau_1(n-1) - \tau_0(n) + \tau_0(n-1),$$
$$\tau_i(n) = \max_{\nu} \{ \sum_k \left( \sum_{g=1}^n \min(l_g, \nu_k) + (i-2)\nu_k - I_k \right) - 2 \sum_{j \le k} \min(\nu_j, \nu_k) \},$$

where the maximum is taken over the subset  $\nu = \{(\nu_1, I_1), (\nu_2, I_2), \ldots\}$  of the rigged configuration (m, J) regarded as the multi set of (row length, attached rigging).

#### Appendix B. Proof of Proposition 3.4

Since  $E_l(\omega(p)) = E_l(p)$  by Proposition 2.3, we may assume that  $wt(p) \ge 0$ . Such a path can be expressed as  $p = T_1^d(p_+)$  for some  $d \in \mathbb{Z}$  and  $p_+ \in \mathcal{P}$  due to Lemma 3.1. Then  $E_l(p) = E_l(T_1^d(p_+)) = E_l(p_+)$ , where the last equality is due to Theorem 2.2. Therefore it suffices to show (3.3) for  $p \in \mathcal{P}_+$ , which we shall assume in the sequel.

For any element of the form  $b \in B = B_{l_1} \otimes \cdots \otimes B_{l_k}$  we define the quantity  $D_l(b) \in \mathbb{Z}_{\geq 0}$  by

$$\zeta^0 u_l \otimes b \simeq b^* \otimes \zeta^{D_l(b)} w \quad (b^* \in B, \ w \in B_l)$$

under the isomorphism  $\operatorname{Aff}(B_l) \otimes \operatorname{Aff}(B) \simeq \operatorname{Aff}(B) \otimes \operatorname{Aff}(B_l)$ . (We have omitted the spectral parameters for  $b, b^*$ .) When p is highest, its energy  $E_l(p)$  emerges not only in (2.10) but already in the relation (2.11) that produces  $v_l$ . Namely,

**Lemma B.1.** For any highest path  $p \in \mathcal{P}_+$ , one has  $D_l(p) = E_l(p)$ .

The following proof is direct but not intrinsic.

*Proof.* Fixing a highest path  $p = b_1 \otimes \cdots \otimes b_L$ , we regard  $D_l(p)$  and  $y_2$  in Figure 2.2 as functions of  $x_2$  there. The quantity  $D_l(p)$  is the number of vertices of the bottom right type in Figure 2.1. Call them scoring vertices. We keep track of the scoring vertices appearing in Figure 2.2 as  $x_2$  is increased from 0 to  $c = y_2(l) \le l$ .  $D_l(p) = E_l(p)$  is shown if the scoring vertices for  $x_2 = 0$  remain scoring and no new scoring vertices are created during the increment of  $x_2$ . To see this, recall the property  $y_2(x_2 + 1) = y_2(x_2)$  or  $y_2(x_2 + 1) = y_2(x_2) + 1$  as noted in the proof of the Proposition 2.1. Since  $b_1, \ldots, b_L$  are fixed, a little inspection of Figure 2.1 tells that the increment of  $x_2$  never transforms the other type of vertices into scoring ones. On the other hand a scoring vertex may change into the top right type in Figure 2.1 if a = l - 1. Suppose this firstly happened as  $x_2 = r$  is increased to  $x_2 = r + 1$  for some  $r < c \le l$ . Consider the leftmost such vertex that has ceased to be scoring at  $x_2 = r + 1$ . Such a situation is realized only if all the vertices (at  $x_2 = r + 1$ ) on its left are bottom two types in Figure 2.1. Let  $\alpha$  and  $\beta$  be the number of the bottom left types and the bottom right (scoring) types in them. Then one has  $r + 1 - \alpha + \beta = l$ . On the other hand from the highest path condition (2.8), one also has  $\alpha \geq \beta + 1$ , where the last +1 is the contribution of the very vertex that has ceased to be scoring. Thus we obtain  $r = l + \alpha - \beta - 1 \ge l$ , which is a contradiction.  $\square$ 

Proof of Proposition 3.4. Let  $p \in \mathcal{P}_+$  be a highest path such that  $\phi(p) = (m, J)$ hence  $\mu(p) = m$ . Suppose that  $u_a^{\otimes A} \otimes p \otimes \overbrace{1 \otimes \cdots \otimes 1}^n \simeq p^* \otimes 1 \otimes \cdots \otimes 1 \otimes u_a^{\otimes A}$  is valid. Here we take  $1 \ll A \ll n$ . In general  $p^*$  is a highest path longer than L(but much shorter than n). By Lemma B.1 and  $H(u_l \otimes u_k) = 0$ , we have  $E_l(p) = D_l(p) = D_l(u_a^{\otimes A} \otimes p \otimes 1 \otimes \cdots \otimes 1) = D_l(p^* \otimes 1 \otimes \cdots \otimes 1 \otimes u_a^{\otimes A}) = D_l(p^* \otimes 1 \otimes \cdots \otimes 1)$ . By Proposition A.4, we know that  $\phi(p^* \otimes 1 \otimes \cdots \otimes 1) = (m, I)$ , where  $I = (I_i^{(j)})$ reads  $I_i^{(j)} = J_i^{(j)} + A\min(a, j)$ . Therefore by taking A and a sufficiently large, one can achieve the situation  $1 \ll \cdots \ll I_1^{(j)} \le \cdots \le I_{m_j}^{(j)} \ll I_1^{(j+1)} \le \cdots \le I_{m_{j+1}}^{(j+1)} \ll$  $\cdots$ . For such a rigged configuration (m, I), the KKR algorithm produces the path  $p^* \otimes 1 \otimes \cdots \otimes 1 = \phi^{-1}((m, I))$  of the form:

(B.1) 
$$\dots 2^{\otimes j} \dots 2^{\otimes j} \dots 2^{\otimes j} \dots 2^{\otimes j} \dots 2^{\otimes j+1} \dots 2^{\otimes j+1} \dots 2^{\otimes j+1} \dots 2^{\otimes j+1} \dots$$

Here ... means an array of  $1 \in B_1$  and  $2^{\otimes j} \in B_1^{\otimes j}$ . The number of appearance of  $2^{\otimes j}$  is  $m_j$ . One can satisfy  $L_j \gg l$  for all  $L_j$  by taking A large. Moreover from the KKR algorithm, there are at least  $1^{\otimes j}$  between any two  $2^{\otimes j}$ . For such a path  $p^* \otimes 1 \otimes \cdots \otimes 1$ , it is straightforward to check  $D_l(p^* \otimes 1 \otimes \cdots \otimes 1) = \sum_{j \ge 1} \min(l, j)m_j$ .

Remark B.2. In the pattern like (B.1),  $2^{\otimes j}$  is a soliton with length (amplitude) j. In this context the data  $m = (m_j)$  tells that there are  $m_j$  solitons with length j.

Appendix C. Proof of Proposition 3.7

**Lemma C.1.** Let  $q \in B_1^{\otimes d}$  be a highest path of length d and  $r \in B_1^{\otimes L-d}$  be a highest path of length L-d. Suppose that their rigged configurations are  $\phi(q) = (l, I)$  and  $\phi(r) = (n, K)$ . Then the rigged configuration of the highest path  $q \otimes r \in B_1^{\otimes L}$  is given by  $\phi(q \otimes r) = (l \cup n, I \cup K')$ , where  $K' = (K_i'^{(j)})$  is given by

(C.1) 
$$K_i^{\prime(j)} = K_i^{(j)} - 2\sum_k \min(j,k)l_k + d,$$

where  $(l \cup n, I \cup K')$  means the union regarding (l, I) and (n, K') as multi-sets of rows assigned with rigging.

*Proof.* The new vacancy number  $p'_j$  for the width j block in  $(l \cup n, I \cup K')$  reads  $p'_j = L - 2\sum_k \min(j,k)l_k - 2\sum_k \min(j,k)n_k$ . When applying the KKR map  $\phi^{-1}$  to  $(l \cup n, I \cup K')$ , the co-rigging of the *i*-th row of the width j block in (n, K') is

(C.2) 
$$p'_j - K'^{(j)}_i = (L - d - 2\sum_k \min(j,k)n_k) - K^{(j)}_i.$$

The right hand side is equal to the co-rigging of the same row in the rigged configuration (n, K). Meanwhile the co-rigging of the (l, I) part is not less than that in the original q plus the vacancy number of r. Therefore the algorithm of the map  $\phi^{-1}$  proceeds as  $\phi^{-1}((l \cup n, I \cup K')) = \phi^{-1}((l, I)) \otimes r = q \otimes r$ .

**Example C.2.** Take q = 1112122, r = 111221221122, hence d = 7, L = 19. The highest path  $p_2$  in Examples 3.2 and A.1 is expressed as  $p_2 = q \otimes r$ .

$$\phi(q) = \boxed{\begin{array}{c} 1 \\ 2 \end{array}} 1 \qquad \phi(r) = \boxed{\begin{array}{c} 1 \\ 3 \end{array}} 0$$

The rigged configuration for  $\phi(p_2)$  in Example A.1 contains this  $\phi(q)$  indeed. The rest of it corresponding to (n, K') is related to  $\phi(r) = (n, K)$  as

in agreement with (C.1).

(C.3)

**Lemma C.3.** Let  $q \in B_1^{\otimes d}$  be a highest path of length d and  $r \in B_1^{\otimes L-d}$  be a highest path of length L-d. Set  $\phi(r \otimes q) = (m, J)$  and  $\phi(q \otimes r) = (m', J')$ . Then m = m' and  $\iota(J') \simeq \iota(J) + d \in \overline{\mathcal{J}}(m)$  are valid.

See (3.12) and (3.16) for the definitions of  $\simeq$  and  $\iota$ .

*Proof.* We use the same notation as in Lemma C.1. Thus  $m' = l \cup n$  and  $J' = I \cup K'$ . Applying Lemma C.1 with (l, I) and (n, K) interchanged, we find  $m = l \cup n = m'$ and  $J = I' \cup K$  with  $I'_{i}(j) = I^{(j)}_{i} - 2\sum_{k} \min(j, k)n_{k} + L - d$ . In terms of the vacancy number  $p'_{j}$  for m in the proof of Lemma C.1, the relation between the rigging Jand J' is summarized as

(C.4) 
$$I_i^{\prime(j)} = I_i^{(j)} + 2\sum_k \min(j,k)l_k + p_j' - d,$$

(C.5) 
$$K_i^{(j)} = K_i^{\prime(j)} + 2\sum_k \min(j,k)l_k - d_k$$

See the picture after (C.8). Now switch to the extended sequences  $\iota(J)$  and  $\iota(J')$  by (3.16). Apart from -d, this is exactly the effect of the slide  $\prod_k \sigma_k^{l_k}$  (3.11) on the width j block of  $\iota(J')$ . Therefore we conclude  $\iota(J) = (\prod_k \sigma_k^{l_k})\iota(J') - d$ .  $\Box$ 

Proof of  $\Rightarrow$  in Proposition 3.7. Without loss of generality we may assume d' = 0and  $0 \leq d < L$ . Since  $p_+$  and  $p'_+$  are highest paths, they must have the form  $p_+ = r \otimes q$  and  $p'_+ = q \otimes r$ , where the both paths  $q \in B_1^{\otimes d}$  and  $r \in B_1^{\otimes L-d}$  are highest ones. Then the assertion follows from Lemma C.3.

To show  $\Leftarrow$  in Proposition 3.7, we explore the full implication of the right hand side of (3.17). Let  $H = \{j_1 < \cdots < j_s\}$  be the set of lengths of rows of m as in (3.8).

**Lemma C.4.** Suppose  $\iota(J) + e \simeq \iota(J') + e' \in \overline{\mathcal{J}}(m)$  for some  $m = (m_j) \in \mathcal{M}$ ,  $J, J' \in \operatorname{Rig}(m)$  and  $e, e' \in \mathbb{Z}$ . Then there exist  $0 \leq l_j \leq m_j$   $(j \in H)$  and  $0 \leq d < L$  such that the relation  $\sigma_{j_1}^{l_{j_1}} \cdots \sigma_{j_s}^{l_{j_s}} \iota(J') = \iota(J) + d$  holds.

Proof. From the assumption, there exist  $l_j$ 's and d such that  $\sigma_{j_1}^{l_{j_1}} \cdots \sigma_{j_s}^{l_{j_s}} \iota(J') = \iota(J) + d$ . Our task is to show that they can always be chosen within the range  $0 \leq l_j \leq m_j \ (j \in H)$  and  $0 \leq d < L$ . From the last comment in Section 3.2, to achieve the relation  $\sigma_{j_1}^{l_{j_1}} \cdots \sigma_{j_s}^{l_{j_s}} \iota(J') = \iota(J) + d$  by the successive transformations  $\iota(J') \to \sigma_{j_s}^{l_{j_s}} \iota(J') \to \sigma_{j_s-1}^{l_{j_s}} \sigma_{j_s}^{l_{j_s}} \iota(J') \to \cdots \to \sigma_{j_1}^{l_{j_1}} \cdots \sigma_{j_s}^{l_{j_s}} \iota(J')$ , one must apply  $\sigma_{j_k}^{l_{j_k}}$  so that the blocks  $\mathcal{J}_{j_k}, \mathcal{J}_{j_{k+1}}, \ldots, \mathcal{J}_{j_s}$  in  $\sigma_{j_k}^{l_{j_k}} \cdots \sigma_{j_s}^{l_{j_s}} \iota(J')$  already coincides with that in  $\iota(J)$  up to an overall additive constant. Having this in mind we proceed to showing  $0 \leq l_j \leq m_j$ .

Certainly,  $l_{j_s}$  can be taken as  $0 \leq l_{j_s} < m_{j_s}$  by using (3.13). Since the slides are invertible, one may also assume that the first non-zero number in the sequence  $l_{j_s}, l_{j_{s-1}}, \ldots$  is positive, namely,  $l_{j_s} = l_{j_{s-1}} = \cdots = l_{j_{t+1}} = 0$  and  $l_{j_t} > 0$ . If there is no such t, it follows that  $\iota(J') = \iota(J) + d$  and we are done because  $d \geq 0$  may be assumed without loss of generality and then d < L is obvious from  $L > p_j$  (3.7) and (A.3). Henceforth we assume that  $l_{j_t} > 0$  exists for some  $1 \leq t \leq s$ . We first claim that  $l_{j_t} \leq m_{j_t}$ . In fact, the case t = s is within our assumption. If  $l_{j_t} > m_{j_t}$ for t < s, the upper blocks  $\mathcal{J}_{\beta}(\beta > j_t)$  acquire the uniform shift  $2j_t l_{j_t}$  under  $\sigma_{j_t}^{l_{j_t}}$ . On the other hand the rigging  $J_{m_{j_t}}^{\prime(j_t)}$  gets shifted at least by  $2p_{j_t} + 2j_t l_{j_t}$ . Therefore to adjust  $\sigma_{j_t}^{l_{j_t}} \iota(J')$  to some  $\iota(J)$ , one must extract at least  $p_{j_t} + 2j_t l_{j_t}$  from the new rigging on the upper blocks  $\mathcal{J}_{\beta}$  as an overall constant. But this fails since the original rigging there is not greater than  $p_{\beta}(< p_{j_t})$  hence the result of the above extraction is not greater than  $(p_{\beta} + 2j_t l_{j_t}) - (p_{j_t} + 2j_t l_{j_t}) < 0$ . Thus we have verified  $l_{j_t} \leq m_{j_t}$ .

Next we prove  $0 \leq l_{j_a} \leq m_{j_a}$   $(1 \leq a \leq t)$  by induction on a assuming that  $0 \leq l_{j_b} \leq m_{j_b}$  for all  $a + 1 \leq b \leq t$ . Setting  $K = (K_i^{(j)}) = \sigma_{j_a}^{l_{j_a}} \cdots \sigma_{j_t}^{l_{j_t}} \iota(J')$ , we have  $K_i^{(j_b)} = J_{i+l_{j_b}}^{\prime(j_b)} + 2\sum_{c=a}^t \min(j_b, j_c) l_{j_c}$ . Note that

(C.6) 
$$K_{m_{j_t}}^{(j_t)} \ge p_{j_t} + 2\sum_{b=a}^t \min(j_t, j_b) l_{j_b},$$

(C.7) 
$$K_1^{(j_s)} \le p_{j_s} + 2\sum_{b=a}^t \min(j_s, j_b) l_{j_b},$$

where the former follows from  $l_{j_t} > 0$  and the latter does from  $l_{j_s} < m_{j_s}$  and  $J_{m_{j_s}}^{\prime(j_s)} \leq p_{j_s}$ . As explained above, the blocks  $\mathcal{J}_{\beta}$  ( $\beta \geq j_a$ ) in  $K - \tilde{d}$  should already coincide with those in  $\iota(J)$  for some  $\tilde{d}$ . To show  $l_{j_a} \geq 0$ , suppose  $l_{j_a} < 0$  on the contrary. Then we have  $K_1^{(j_a)} \leq J_{m_{j_a}}^{\prime(j_a)} - p_{j_a} + 2\sum_{b=a}^t \min(j_a, j_b) l_{j_b} \leq 2\sum_{b=a}^t \min(j_a, j_b) l_{j_b}$ . From this and (C.6),  $K - \tilde{d}$  can coincide with some  $\iota(J)$  only if

$$2\sum_{b=a}^{t} \min(j_t, j_b) l_{j_b} \le \tilde{d} \le 2\sum_{b=a}^{t} \min(j_a, j_b) l_{j_b}.$$

But this is impossible because of  $l_{j_t} > 0$  and the induction assumption, verifying  $l_{j_a} \ge 0$ . To show  $l_{j_a} \le m_{j_a}$ , suppose  $l_{j_a} > m_{j_a}$  on the contrary. Then we have  $K_{m_{j_a}}^{(j_a)} \ge 2p_{j_a} + 2\sum_{b=a}^t \min(j_a, j_b)l_{j_b}$ . From this and (C.7),  $K - \tilde{d}$  can coincide with some  $\iota(J)$  only if

$$p_{j_a} + 2\sum_{b=a}^t \min(j_a, j_b) l_{j_b} \le \tilde{d} \le p_{j_s} + 2\sum_{b=a}^t \min(j_s, j_b) l_{j_b}.$$

Again it is easy to check that this is impossible by the reason similar to the previous inequality, proving  $l_{j_a} \leq m_{j_a}$ .

Finally we consider d in the relation  $\iota(J)+d = \sigma_{j_1}^{l_{j_1}} \cdots \sigma_{j_t}^{l_{j_t}} \iota(J') = \sigma_{j_1}^{l_{j_1}} \cdots \sigma_{j_s}^{l_{j_s}} \iota(J')$ . From (C.6) and (C.7) with a = 1, we get  $J_{m_{j_t}}^{(j_t)} + d \ge p_{j_t} + 2\sum_{b=1}^t \min(j_t, j_b)l_{j_b}$  and  $J_1^{(j_s)} + d \le p_{j_s} + 2\sum_{b=1}^t \min(j_s, j_b)l_{j_b}$ . Therefore in order that  $(J_i^{(j)}) \in \operatorname{Rig}(m)$  to hold, d must satisfy  $0 \le 2\sum_{b=1}^t \min(j_t, j_b)l_{j_b} \le d \le p_{j_s} + 2\sum_{b=1}^t \min(j_s, j_b)l_{j_b} < p_{j_s} + 2\sum_{b=1}^s \min(j_s, j_b)m_{j_b} = L$ . In particular, d = 0 can happen only if  $\forall l_{j_b} = 0$ .

Proof of  $\Leftarrow$  in Proposition 3.7. By Lemma C.4, J and J' are connected by the relation  $(\prod_k \sigma_k^{l_k})\iota(J') = \iota(J) + d$  for some  $0 \leq l_j \leq m_j$  and  $0 \leq d < L$ . If  $\forall l_k = 0$ , the assertion follows easily from the definition of the KKR bijection  $\phi$ . Henceforth we assume that  $l_j > 0$  for some  $j \in H$ . Let  $n = (n_j)$  and  $l = (l_j)$  be the Young diagrams giving the decomposition  $m = l \cup n$ , where  $0 \leq n_j \leq m_j$  is

defined by  $n_j = m_j - l_j$ . By the last remark in the proof of Lemma C.4 we have 0 < d < L.

Define 
$$I^{(j)} = (I_i^{(j)})_{1 \le i \le l_j}$$
 and  $K^{(j)} = (K_i^{(j)})_{1 \le i \le n_j}$  by  
(C.8)  $(I_1^{(j)}, \dots, I_{l_j}^{(j)}) = (J_1^{\prime(j)}, \dots, J_{l_j}^{\prime(j)}), \quad (K_1^{(j)}, \dots, K_{n_j}^{(j)}) = (J_1^{(j)}, \dots, J_{n_j}^{(j)}).$ 

Then define further  $(I'^{(j)}_i)_{1 \le i \le l_j}$  and  $(K'^{(j)}_i)_{1 \le i \le n_j}$  by (C.4) and (C.5), respectively. By construction we have  $J = I' \cup K$  and  $J' = I \cup K'$ , where  $(J_{n_j+1}^{(j)}, \dots, J_{m_j}^{(j)}) =$  $(I_1^{\prime(j)}, \ldots, I_{l_j}^{\prime(j)})$  and  $(J_{l_j+1}^{\prime(j)}, \ldots, J_{m_j}^{\prime(j)}) = (K_1^{\prime(j)}, \ldots, K_{n_j}^{\prime(j)})$ . The following figure is helpful to grasp these relations.



We claim that  $I \in \operatorname{Rig}_d(l)$  and  $K \in \operatorname{Rig}_{L-d}(n)$ . In fact,  $0 \leq I_1^{(j)} \leq \cdots \leq I_{l_j}^{(j)}$  is obvious and  $I_{l_j}^{(j)} = I_{l_j}^{(j)} - 2\sum_k \min(j,k)l_k - p'_j + d \le d - 2\sum_k \min(j,k)l_k$  which is the vacancy number for  $B_1^{\otimes d}$ . (Here  $p'_j = L - 2\sum_k \min(j,k)m_k$  is the same as that in the proof of Lemmas C.1 and C.3.) Similarly,  $0 \le K_1^{(j)} \le \cdots \le K_{n_j}^{(j)}$  is obvious and  $K_{n_j}^{(j)} = K_{n_j}^{\prime(j)} + 2\sum_k \min(j,k)l_k - d \le p'_j + 2\sum_k \min(j,k)l_k - d = L - d - 2\sum_k \min(j,k)n_k$  which is the vacancy number for  $B_1^{\otimes L-d}$ . Set  $q = \phi^{-1}((l,I)) \in B_1^{\otimes d}$  and  $r = \phi^{-1}((n,K)) \in B_1^{\otimes L-d}$ . Then Lemma C.1 tells that the highest paths  $p_+$  and  $p'_+$  in Proposition 3.7 are given by  $p_+ = \phi^{-1}((m,J)) = r \otimes q$  and  $p'_+ = \phi^{-1}((m,J')) = q \otimes r$ . Therefore we obtain  $p'_{-1} = T^d(n_j)$ 

 $p'_{+} = T_1^d(p_{+}).$ 

The proof of Proposition 3.7 is finished.

### Appendix D. Proof of Theorem 3.12

For distinction we write the time evolution of the angle variable (3.14) as  $\tau_l$ within this appendix. Obviously  $\tau_l \tau_k = \tau_k \tau_l$  is valid.

It is straightforward to check the commutativity of the diagram (3.18) for l = 1. In fact, if  $p = T_1^d(p_+)$  for  $p_+ \in \mathcal{P}_+(m)$  with  $\phi(p_+) = (m, J)$ , we have  $\Phi(T_1(p)) =$  $\Phi(T_1^{d+1}(p_+)) = [\iota(J) + d + 1].$  On the other hand due to  $\tau_1(\iota(J)) = \iota(J) + 1$ , we obtain  $\tau_1(\Phi(p)) = \tau_1([\iota(J) + d]) = [\iota(J) + d + 1].$ 

The commutativity  $\tau_1 \Phi = \Phi T_1$  and Lemma 3.6 reduce the proof of Theorem 3.12 to the highest paths  $p_+ \in \mathcal{P}_+(m)$ . In fact, the equality  $\Phi T_l(p) = \tau_l \Phi(p)$  for general path  $p = T_1^d(p_+) \in \mathcal{P}(m)$  is deduced from  $\Phi T_l(p_+) = \tau_l \Phi(p_+)$  by multiplying  $\tau_1^d$  on the both sides and using the commutativity  $T_1T_l = T_1T_l$  and  $\tau_1\tau_l = \tau_l\tau_1$ .

In the remainder of this appendix we fix the Young diagram  $m = (m_j) \in \mathcal{M}$  and assume that  $p \in \mathcal{P}_+(m) \subset B_1^{\otimes L}$ .  $(m_j \text{ is the number of the length } j \text{ rows in } m \text{ as in } m$  the main text.) Then we determine J, K, I, d and q successively as follows:

(D.1) 
$$\phi(p) = (m, J), \quad J \in \operatorname{Rig}_L(m),$$

(D.2) 
$$\tau_{l}\iota(J) = K \in \overline{\mathcal{J}}(m),$$

(D.3) 
$$[K] = [\iota(I) + d] \in \mathcal{J}(m), \quad I \in \operatorname{Rig}_L(m), \quad d \ge 0,$$

(D.4)  $\phi^{-1}((m, I)) = q \in \mathcal{P}_+(m),$ 

where the definition of  $\iota$  is available in (3.16). The role of these objects will be seen clearly in (D.8). In (D.3), the choice of I and d is not unique. However what matters in the following proof is the combination  $T_1^d(q)$  whose uniqueness has been assured by Proposition 3.7.

For  $N \in \mathbb{Z}_{\geq 1}$  denote by  $m^N = (Nm_j)$  the N-fold repetition of m and define the map

$$\pi_N: \qquad \overline{\mathcal{J}}(m) \longrightarrow \mathbb{Z}^{Nm_{j_1}} \times \cdots \times \mathbb{Z}^{Nm_{j_s}} \\ \left( (J_i^{(j_1)})_{i \in \mathbb{Z}}, \dots, (J_i^{(j_s)})_{i \in \mathbb{Z}} \right) \mapsto \left( (J_i^{(j_1)})_{1 \le i \le Nm_{j_1}}, \dots, (J_i^{(j_s)})_{1 \le i \le Nm_{j_s}} \right).$$

In what follows we set  $B = B_1^{\otimes \mathcal{L}}$  and assume that  $\mathcal{L} > (N+1)L$ .

**Lemma D.1.** For  $\pi_N(\iota(J)) \in \operatorname{Rig}_{NL}(m^N)$ , one has

(D.6) 
$$\phi_B^{-1}((m^N, \pi_N(\iota(J)))) = p \otimes p \otimes p \otimes p \otimes p \otimes p \otimes 1 \otimes 1 \otimes \dots \otimes 1,$$

(D.7) 
$$\phi_B^{-1}((m^N, \pi_N(K))) = p^* \otimes \overline{T_l(p) \otimes T_l(p)} \otimes \cdots \otimes \overline{T_l(p)} \otimes p^{**} \otimes 1 \otimes \cdots \otimes 1,$$

where  $p^* \in B_1^{\otimes L}$  is defined by (2.11) and  $v_l \otimes 1^{\otimes L} \simeq p^{**} \otimes u_l$ . *Proof.* The relation (D.6) is derived by repeated use of Lemma C.1. The

*Proof.* The relation (D.6) is derived by repeated use of Lemma C.1. To see (D.7), note from (2.11) that

$$u_l \otimes \overbrace{p \otimes \cdots \otimes p}^N \otimes 1 \otimes \cdots \otimes 1 \simeq p^* \otimes \overbrace{T_l(p) \otimes \cdots \otimes T_l(p)}^{N-1} \otimes p^{**} \otimes 1 \otimes \cdots \otimes 1 \otimes u_l$$

From this and (D.6), the relation (D.7) is obtained by applying Proposition A.4 with p and (m, J) replaced with  $p^{\otimes N}$  and  $(m^N, \pi_N(\iota(J)))$ .

Take the slide  $\sigma = \sigma_{j_1}^{l_{j_1}} \cdots \sigma_{j_s}^{l_{j_s}} \in \mathcal{A}$  that achieves (D.3) via  $\sigma(K) = \iota(I) + d$ . By the argument similar to the proof of Lemma C.4 (especially after (C.7)), one can show that  $l_j$ 's can be chosen as  $0 \leq l_j \leq m_j$ . We introduce the Young diagram  $\tilde{m}^N = (\tilde{m}_j^N)$  by  $\tilde{m}_j^N = Nm_j - l_j$  and let  $\tilde{\pi}_N$  be the map (D.5) with  $Nm_j$  replaced by  $\tilde{m}_j^N$ .

**Lemma D.2.** As  $\mathcal{L}$  and N grow large satisfying  $\mathcal{L} > NL \gg 1$ , the path  $\phi_B^{-1}((\tilde{m}^N, \tilde{\pi}_N(\sigma(K))))$ takes the form  $\xi \otimes \eta$ , where the left part  $\xi$  is independent of  $\mathcal{L}, N$  and the right part  $\eta$  growing with  $\mathcal{L}, N$  is the same as the corresponding part in  $\phi_B^{-1}((m^N, \pi_N(K)))$ (D.7).

*Proof.* From the definition (3.11) and  $\sigma = \sigma_{j_1}^{l_{j_1}} \cdots \sigma_{j_s}^{l_{j_s}}$ , the rigged configuration  $(\tilde{m}^N, \tilde{\pi}_N(\sigma(K)))$  is obtained from  $(m^N, \pi_N(K))$  by removing the bottom  $l_j$  rows and adding  $2\sum_{b=1}^s \min(j, j_b)l_{j_b}$  to the rigging for each block of length j rows. This

operation does not change the co-rigging of the remaining part. Therefore the assertion follows from the KKR algorithm.  $\hfill \Box$ 

*Proof of Theorem 3.12.* We are to check  $\stackrel{?}{=}$  in the following diagram:

(D.8)  

$$p \xrightarrow{\Phi} [\iota(J)]$$
  
 $T_l \downarrow \qquad \qquad \downarrow \tau_l$   
 $T_l(p) \stackrel{?}{=} T_1^d(q) \xrightarrow{\Phi^{-1}} [K] = [\iota(I) + d]$ 

Consider the rigged configuration  $(\tilde{m}^N, \tilde{\pi}_N(\sigma(K))) = (\tilde{m}^N, \tilde{\pi}_N(\iota(I) + d))$ . The latter expression tells that this is also obtained from  $(m^N, \pi_N(\iota(I) + d))$  by removing the top  $l_j$  rows in each block of length j rows. Thus from the KKR algorithm, the paths  $\phi_B^{-1}((\tilde{m}^N, \tilde{\pi}_N(\iota(I) + d)))$  and  $\phi_B^{-1}((m^N, \pi_N(\iota(I) + d)))$  are the same except only some components that are located on the right and remain unchanged with growing  $\mathcal{L}, N$ . On the other hand from the definition (D.4) and Lemma D.1 (D.6) with (p, J) replaced by (q, I), we find

$$\phi_B^{-1}((m^N, \pi_N(\iota(I) + d))) = \underbrace{1 \otimes \cdots \otimes 1}^d \otimes \underbrace{q \otimes q \otimes \cdots \otimes q}^N \otimes 1 \otimes \cdots \otimes 1.$$

Combining the foregoing argument and Lemma D.2, we can equate the middle part of this with that in (D.7), leading to  $T_l(p) = T_1^d(q)$ .

### References

- [1] A. Kuniba and A. Takenouchi, Bethe ansatz at q = 0 and periodic box-ball systems, J. Phys. A: Math. Gen. in press. (nlin.SI/0509001)
- [2] A. Kuniba and A. Takenouchi, Periodic cellular automata and Bethe ansatz, (math-ph/0511013) to appear in Nankai Tracks in Math. Proceedings of XXIII International Conference of DGMTP.
- [3] S.-J. Kang, M. Kashiwara and K. C. Misra, Crystal bases of Verma modules for quantum affine Lie algebras, Compositio Math. 92 (1994) 299–325.
- [4] F. Yura and T. Tokihiro, On a periodic soliton cellular automaton, J. Phys. A: Math. Gen. 35 (2002) 3787–3801.
- [5] S.V.Kerov, A.N.Kirillov and N.Yu.Reshetikhin, Combinatorics, Bethe ansatz, and representations of the symmetric group, Zap. Nauch. Semin. LOMI. 155 (1986) 50–64.
- [6] A. N. Kirillov and N. Yu. Reshetikhin, The Bethe ansatz and the combinatorics of Young tableaux. J. Soviet Math. 41 (1988) 925–955.
- [7] A. Kuniba and T. Nakanishi, The Bethe equation at q = 0, the Möbius inversion formula, and weight multiplicities: I. The sl(2) case, Prog. in Math. **191** (2000) 185–216.
- [8] S.-J. Kang, M. Kashiwara, K. C. Misra, T. Miwa, T. Nakashima and A. Nakayashiki, Affine crystals and vertex models, Int. J. Mod. Phys. A 7 (suppl. 1A), (1992) 449–484.
- [9] A. Nakayashiki and Y. Yamada, Kostka polynomials and energy functions in solvable lattice models, Selecta Mathematica, New Ser. 3 (1997) 547–599.
- [10] R. J. Baxter, Exactly solved models in statistical mechanics, Academic Press, London (1982).
- [11] D. Takahashi and J. Satsuma, A soliton cellular automaton, J. Phys. Soc. Jpn. 59 (1990) 3514–3519.
- [12] H. A. Bethe, Zur Theorie der Metalle, I. Eigenwerte und Eigenfunktionen der linearen Atomkette, Z. Physik 71 (1931) 205–231.
- [13] C. S. Gardner, J. M. Greene, M. D. Kruskal and R. M. Miura, Method for solving the Korteweg-de Vries equation, Phys. Rev. Lett. 19 (1967) 1095–1097.
- [14] M. J. Ablowitz and H. Segur, Solitons and the inverse scattering transform, SIAM Studies in Appl. Math. 4. Philadelphia Pa. (1981).

- [15] D. Yoshihara, F. Yura and T. Tokihiro, Fundamental cycle of a periodic box-ball system, J. Phys. A: Math. Gen. 36 (2003) 99–121.
- [16] G. Hatayama, K. Hikami, R. Inoue, A. Kuniba, T. Takagi and T. Tokihiro, The A<sub>M</sub><sup>(1)</sup> Automata related to crystals of symmetric tensors, J. Math. Phys. 42 (2001) 274–308.
- [17] K. Fukuda, M. Okado, Y. Yamada, Energy functions in box ball systems, Int. J. Mod. Phys. A 15 (2000) 1379–1392.
- [18] G. Hatayama, A. Kuniba, M. Okado, T. Takagi and Y. Yamada, Scattering rules in soliton cellular automata associated with crystal bases, Contemporary Math. 297 (2002) 151–182.
- [19] A. Schilling, X=M Theorem: Fermionic formulas and rigged configurations under review, (math.QA/0512161).
- [20] E. Date and S. Tanaka, Periodic multi-soliton solutions of Korteweg-de Vries equation and Toda lattice, Prog. Theoret. Phys. Suppl. 59 (1976) 107–125.
- [21] B. A. Dubrovin, V. B. Matveev and S. P. Novikov, Nonlinear equations of Korteweg-de Vries type, finite-band linear operators and Abelian varieties Russian Math. Surveys **31** (1976) 59–146.
- [22] M. Kashiwara, Crystal bases of modified quantized universal enveloping algebra, Duke Math. 73 (1994) 383–413.
- [23] A. Kuniba, M. Okado, T. Takagi and Y. Yamada, Vertex operators and partition functions in the box-ball systems (Japanese), RIMS Kôkyûroku 1302 (2003) 91–107.
- [24] T. Takagi, Inverse scattering method for a soliton cellular automaton, Nucl. Phys. B707 (2005) 577–601.
- [25] N. Yu. Reshetikhin, The functional equation method in the theory of exactly soluble quantum systems, Sov. Phys. JETP 57 (1983) 691–696.
- [26] A. Kuniba and J. Suzuki, Analytic Bethe ansatz for fundamental representations of Yangians, Commun. Math. Phys. 173 (1995) 225–264.
- [27] J. Mada, M. Idzumi and T. Tokihiro, Fundamental cycle of a periodic box-ball system and solvable lattice models, preprint (Nov. 2005).
- [28] A. N. Kirillov, A. Schilling and M. Shimozono, A bijection between Littlewood-Richardson tableaux and rigged configurations, Selecta Math. 8 (2002) 67–135.

INSTITUTE OF PHYSICS, UNIVERSITY OF TOKYO, TOKYO 153-8902, JAPAN *E-mail address*: atsuo@gokutan.c.u-tokyo.ac.jp

DEPARTMENT OF APPLIED PHYSICS, NATIONAL DEFENSE ACADEMY, KANAGAWA 239-8686, JAPAN

E-mail address: takagi@nda.ac.jp

INSTITUTE OF PHYSICS, UNIVERSITY OF TOKYO, TOKYO 153-8902, JAPAN *E-mail address*: takenouchi@gokutan.c.u-tokyo.ac.jp