## Polynomials, roots, and interlacing Version 2

Steve Fisk Bowdoin College Brunswick, Me 04011

November 26, 2024

## Contents

Contents	i
List of Figures	ix
List of Tables	xi

### I Polynomials with all real roots

1	Polyn	omials in One Variable 2
	1.1	Notation and basic definitions
	1.2	Sign interlacing 5
	1.3	A quantitative version of sign interlacing
	1.4	Elementary interlacings
	1.5	Linear combinations of polynomials 20
	1.6	Basic topology of polynomials
	1.7	Linear transformations that preserve interlacing 24
	1.8	Linear combinations and determinants
	1.9	Polynomials from determinants
	1.10	A relation weaker than interlacing
	1.11	Polytopes and polynomials
	1.12	The Faà di Bruno problem 39
	1.13	Root Trajectories
	1.14	Recursive sequences of polynomials 43
	1.15	The positive and negative parts
	1.16	Polynomials without all real roots
2	Polyn	omials with all positive coefficients 51
	2.1	Basic Properties 51
	2.2	Interlacing in $\mathbf{P}^{\text{pos}}$
	2.3	Linear combinations in $\mathbf{P}^{\text{pos}}$
	2.4	Various linear combinations 56
	2.5	The positivity hierarchy

3	Matri	ices that preserve interlacing	59
	3.1	Totally positive matrices	59
	3.2	Constant matrices preserving mutual interlacing	60
	3.3	Determinants and mutual interlacing	67
	3.4	Interlacing polynomials and the Hurwitz matrix	70
	3.5	The division algorithm	71
	3.6	Integrating families of polynomials	75
	3.7	Families, root zones, and common interlacings	80
	3.8	Two by two matrices preserving interlacing	82
	3.9	Matrices of polynomials that preserve interlacing	87
	3.10	Matrices preserving interlacing in $\mathbf{P}^{pos}$	89
	3.11	Linear 2 by 2 matrices preserving interlacing	93
	3.12	Polynomial sequences from matrices	96
	3.13	Interlacing via the complexes	98
	3.14	Mutually interlacing polynomials	100
4	Hom	ogeneous Polynomials	103
	4.1	Introduction	103
	4.2	Polar derivatives	104
	4.3	Newton's inequalities	106
	4.4	More inequalities and bounds	108
	4.5	The matrix of a transformation	111
	4.6	More coefficient inequalities	115
	4.7	Log concavity in $\mathbf{P}^{pos}$	118
	4.8	A pair of recursions	118
5	Analy	ytic Functions	121
	5.1	The uniform closure of $\mathbf{P}^{pos}$	121
	5.2	The uniform closure of $\mathbf{P}$	123
	5.3	The Hadamard Factorization Theorem	125
	5.4	Polynomials with rapidly decreasing coefficients	128
	5.5	Interlacing of analytic functions	131
	5.6	Characterizing $\hat{\mathbf{P}}$	135
	5.7	<b>P</b> -positive semi-definite matrices	135
6	Linea	ar Transformations of Polynomials	138
	6.1	Characterizing transformations	138
	6.2	The domain and range of a linear transformation	148
	6.3	Transformations determined by products	151
	6.4	Composition	153
	6.5	Recurrence relations for orthogonal polynomials	154
	6.6	Linear transformations satisfying a recursion	156
	6.7	Möbius transformations	156
	6.8	Transformations and commutativity properties	160
	6.9	Singular points of transformations	163
	6 10	Appell sequences	164

	6.11	The diamond product		165
	6.12	Other root preserving transformations		168
7	Linea	ar transformations that preserve roots		170
	7.1	Multiplier transformations		170
	7.2	Homogeneous transformations		173
	7.3	The Hadamard product		174
	7.4	Differential operators		177
	7.5	Transformations based on differentiation		181
	7.6	The falling factorial		182
	7.7	The rising factorial		183
	7.8	Hermite polynomials		186
	7.9	Charlier polynomials		189
	7.10	Laguerre polynomials		190
	7.11	The integration and exponential transformations		191
	7.12	Binomial transformations		194
	7.13	Eulerian polynomials		196
	7.14	Euler-Frobenius polynomials		198
	7.15	Even and odd parts		198
	7.16	Chebyshev and Jacobi Polynomials		201
	7.17	Commuting diagrams of spaces		203
	7.18	The Pincherlet derivative		204
	7.19	Hypergeometric Polynomials		205
	7.20	Eigenpolynomials of linear transformations		207
	7.21	Classical multiple orthogonal polynomials		209
	7.22	Laurent Orthogonal polynomials		210
	7.23	Biorthogonal polynomials	• •	212
8	Affin	e Transformations of Polynomials		213
	8.1	Introduction		213
	8.2	The affine derivative		215
	8.3	Sign interlacing for $\mathbf{P}^{\mathbb{A}}$		220
	8.4	The affine exponential		221
	8.5	A-interlacing and the symmetric derivative		223
	8.6	Linear transformations		225
	8.7	The case $\mathbb{A}x = x + 1$		226
	8.8	A multiplication in $\mathbf{P}^{sep}$		231
	8.9	The analytic closure of $\mathbf{P}^{sep}$		233
	8.10	Integrating families of polynomials in $\mathbf{P}^{sep}$ and $\mathbf{P}^{\mathbb{A}}$		236
	8.11	Determinants and $\mathbf{P}^{\mathbb{A}}$		239
	8.12	The case $Ax = qx$ with $q > 1$		241
	8.13	The case $Ax = qx$ with $0 < q < 1$		242
	8.14	The q-exponential		243
	8.15	q-transformations		244
	8.16	Newton's inequalities for $Ax = qx$ and $Ax = x + 1$		246
	8.17	The case $Ax = -x$		249

8.18	The case $Ax = 0$																									25	1
0.10	1110 0000 11/0 0	• •	•	•	•	•	•	•	•	•	•	•	•	• •	•	•	•	•	•	•	•	•	•	•	•		-

### II Polynomials in several variables

9	Polyn	omials in two variables	254
	9.1	The substitution property and determinants	254
	9.2	Interlacing in $\mathbf{Sub}_2$	259
	9.3	Polynomials in two variables	261
	9.4	Inequalities for coefficients of a quadratic in $P_2$	263
	9.5	Solution curves	266
	9.6	Interlacing in $\mathbf{P}_2$	271
	9.7	Linear transformations on $P_2$	273
	9.8	Applications to linear transformations on <b>P</b>	276
	9.9	Properties of $P_2$	279
	9.10	The analog of $\mathbf{P}^{pos}$	280
	9.11	The representation of $P_2$ by determinants	282
	9.12	When some coefficients are zero	284
	9.13	The Hadamard product	285
	9.14	Some differential operators	288
	9.15	Realizing interlacing	291
	9.16	Karlin's conjecture	294
	9.17	Determinants of coefficients	296
	9.18	Generic polynomials and $\mathbf{P}_2^{\text{sep}}$	301
	9.19	Integrating generic polynomials	305
	9.20	Recursive sequences of polynomials	306
	9.21	Bezout matrices	307
10	Dolum	amiala in coveral veriables	011
10		The substitution property	<b>311</b> 211
	10.1	The substitution property	212
	10.2	Departies of D	216
	10.5	The apples of $\mathbf{P}_d^{pos}$	210
	10.4	Coefficient inequalities hives and Hern	319
	10.5	Log conceptity in $\mathbf{P}^{\text{pos}}$	326
	10.0 10.7	Even ding two polynomials $\mathbf{F}_{2}$	320
	10.7	Symmetric Polynomials	320
	10.0	I in ear transformations on P,	334
	10.7	The graph of polynomials in $P_1$	336
	10.10	Differential operators	337
	10.11	Higher order Newton inequalities	340
	10.12	Negative subdefinite matrices	342
	10.15	Ouadratic Forms	346
	10.14	The analog of xD	348
	10.16	Generalized Hadamard products	349
	10.10	THE BMV conjecture	351
	10.17		551

10.18 10.19 10.20	<ul> <li>B The composition conjecture</li></ul>	352 357 358
10.21	Polynomials from matrices	359
10.22	Extending three interlacing polynomials	361
10.23	Subsets of $P_d$	363
10.24	Higher dimensional recurrences	365
11 The	polynomial closure of $P_d$	369
11.1	The polynomial closure of $\mathbb{P}_2$	369
11.2	Reversal and cancellation in $P_2$	376
11.3	Polynomials linear in y	378
11.4	Constrained polynomials with zero coefficients	383
11.5	The polynomial closure of $P_d$	384
11.6	Multiaffine polynomials	389
11.7	Multiaffine polynomials with complex coefficients	392
11.8	Nearly multiaffine polynomials	394
11.9	The polar derivative and the Schur-Szegö map	395
11.10	Determining the direction of interlacing	397
11.11	Reversal in $P_d$	400
11.12	Induced transformations	401
11.13	Linear Transformations	404
11.14	The diamond product	406
11.15	Substituting into quadratic forms	409
11.16	Simultaneous negative pairs of matrices	410
11.17	' The interior of $\mathbf{P}_{d}$	413
11.18	<sup>3</sup> The cone of interlacing	416
11.19	Products of linear factors	419
11.20	The interlacing of products	421
11.21	Characterizing transformations	423
11.22	? Orthogonal polynomials in $\mathbf{P}_d$	424
11.23	Arrays of interlacing polynomials	428
12 Poly	nomials satisfying partial substitution	431
12 1019	Polynomials satisfying partial substitution in two variables	431
12.1	Fxamples	433
12.2	Induced transformations	435
12.0	Applications to transformations	437
12.1	Möbius transformations	440
12.0	Partial substitution in several variables	442
12.0	Quadratic forms	444
12.7	z	

13 T	The a	nalytic closure of P <sub>d</sub>	447
1	3.1	The analytic closure	447
1	3.2	Differential operators	453
1	3.3	Limits and transformations	454
1	3.4	Totally positive functions	456
1	3.5	Hermite polynomials	459
1	3.6	Hermite polynomials in two variables	460
14 E	Exten	ding P <sub>d</sub>	464
1	4.1	Taking the image of $\mathbf{P}_2$	464
1	4.2	Polynomials in several variables with positive coefficients	468
1	4.3	Epsilon substitution	476
15 C	Gene	rating functions	481
1	5.1	The idea of generating function	481
1	5.2	Multiplier transformations	487
1	5.3	Linear transformations $\mathbf{P} \longrightarrow \mathbf{P}$	488
1	5.4	Linear transformations $\mathbf{P} \longrightarrow \mathbf{P}$ - general properties $\dots \dots$	491
1	5.5	Applications: Linear transformations $\mathbf{P} \longrightarrow \mathbf{P} \dots \dots \dots$	495
1	15.6	Applications: the diamond product	499
1	15.7	Applications: generalized Hadamard products	500
1	15.8	Generating functions on $\mathbf{P}(n)$	502
1	15.9	Higher dimensional generating functions	504
1	5.10	Higher dimensional multiplier transformations	505
1	5.11		509
1	5.12	Generating functions for maps $\mathbf{P}_2 \longleftrightarrow \mathbf{P}$	511
1	5.13	Linear transformations $\mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}^{\text{cos}} \dots \dots \dots \dots \dots \dots$	512
1	15.14	More generating functions	515
16 F	Recur	rences with polynomial coefficients	518
1	6.1	The general construction	519
1	6.2	Recursions from $f(y)$ and $g(x, y)$	520
1	6.3	Recursions from $f(y, z)$ and $g(x, y, z)$	521
1	6.4	Recursions from $f(x, -D_y)$ and $g(x, y)$	524
1	6.5	Recurrences from $f(-D_x, y)$ and $g(x, y)$	525
1	6.6	Recurrences from mutually interlacing polynomials	526
1	6.7	Alternative approaches	527
17 N	Matri	ces	529
1	7.1	Basics	529
1	7.2	Principle submatrices	530
1	7.3	The Schur complement	534
1	7.4	Families of matrices	535
1	7.5	Permanents	537
1	7.6	Matrices from polynomials	538

18	Matri	x Polynomials	540
	18.1	Introduction	540
	18.2	Interlacing	543
	18.3	Common interlacings	547
	18.4	Orthogonal-type recurrences	552
	18.5	Problems with products	553
	18.6	Sign Interlacing	554
	18.7	Root zones	556
	18.8	Quadratics and Newton inequalities	559
	18.9	Newton inequalities in $Hyp_2$	560
	18.10	Singular polynomials	561
	18.11	Matrix polynomials with determinants in $\overline{\mathbf{P}}_d$	562

### III Polynomials with complex roots

19	Polyn	omials non-vanishing in a region	565
	19.1	Fundamental properties	566
	19.2	Differentiation	569
	19.3	Interlacing	571
	19.4	Some determinant properties	573
	19.5	Determinants	575
	19.6	Linear transformations	577
	19.7	Analytic functions	578
	19.8	Homogeneous polynomials	579
	19.9	Möbius transformations	581
	19.10	Properties of polynomials with zeros in a strip	584
	19.11	Polynomials non-vanishing on the open unit ball	585
	19.12	Polynomials non-vanishing in a sector	587
20	Uppe	r half plane polynomials	589
20	<b>Uppe</b> 20.1	r half plane polynomials Properties of upper half plane polynomials	<b>589</b> 589
20	<b>Uppe</b> 20.1 20.2	r half plane polynomialsProperties of upper half plane polynomialsConstructions from determinants	<b>589</b> 589 591
20	<b>Uppe</b> 20.1 20.2 20.3	r half plane polynomialsProperties of upper half plane polynomialsConstructions from determinants $SL_2(\mathbb{R}[x])$	<b>589</b> 589 591 593
20	<b>Uppe</b> 20.1 20.2 20.3 20.4	r half plane polynomialsProperties of upper half plane polynomialsConstructions from determinants $SL_2(\mathbb{R}[x])$ Matrices of nearly quadratic polynomials	<b>589</b> 589 591 593 596
20	Uppe 20.1 20.2 20.3 20.4 20.5	r half plane polynomialsProperties of upper half plane polynomialsConstructions from determinants $SL_2(\mathbb{R}[x])$ Matrices of nearly quadratic polynomials $\mathcal{J}$ , a subset of $U_1(\mathbb{C})$	<b>589</b> 589 591 593 596 598
20	Uppe 20.1 20.2 20.3 20.4 20.5 20.6	r half plane polynomialsProperties of upper half plane polynomialsConstructions from determinants $SL_2(\mathbb{R}[x])$ Matrices of nearly quadratic polynomials $\mathfrak{I}$ , a subset of $U_1(\mathbb{C})$ Some simple consequences	<b>589</b> 591 593 596 598 603
20	Uppe 20.1 20.2 20.3 20.4 20.5 20.6 20.7	r half plane polynomialsProperties of upper half plane polynomialsConstructions from determinants $SL_2(\mathbb{R}[x])$ Matrices of nearly quadratic polynomials $\mathfrak{I}$ , a subset of $U_1(\mathbb{C})$ Some simple consequencesInterlacing in $\mathfrak{I}$	<b>589</b> 591 593 596 598 603 604
20	Uppe 20.1 20.2 20.3 20.4 20.5 20.6 20.7 20.8	r half plane polynomialsProperties of upper half plane polynomialsConstructions from determinants $SL_2(\mathbb{R}[x])$ Matrices of nearly quadratic polynomials $\Im$ , a subset of $U_1(\mathbb{C})$ Some simple consequencesInterlacing in $\Im$ Particular interlacings	<b>589</b> 591 593 596 598 603 604 608
20	Uppe 20.1 20.2 20.3 20.4 20.5 20.6 20.7 20.8 20.9	r half plane polynomialsProperties of upper half plane polynomialsConstructions from determinants $SL_2(\mathbb{R}[x])$ Matrices of nearly quadratic polynomials $\Im$ , a subset of $U_1(\mathbb{C})$ Some simple consequencesInterlacing in $\Im$ Particular interlacingsThe geometry of interlacing in $\Im$	<b>589</b> 591 593 596 598 603 604 608 610
20	Uppe 20.1 20.2 20.3 20.4 20.5 20.6 20.7 20.8 20.9 20.10	r half plane polynomialsProperties of upper half plane polynomialsConstructions from determinants $SL_2(\mathbb{R}[x])$ Matrices of nearly quadratic polynomials $\mathfrak{I}$ , a subset of $U_1(\mathbb{C})$ Some simple consequencesInterlacing in $\mathfrak{I}$ Particular interlacingsThe geometry of interlacing in $\mathfrak{I}$ Orthogonal type recurrences	<b>589</b> 591 593 596 598 603 604 608 610 612
20	Uppe 20.1 20.2 20.3 20.4 20.5 20.6 20.7 20.8 20.9 20.10 20.11	r half plane polynomialsProperties of upper half plane polynomialsConstructions from determinants $SL_2(\mathbb{R}[x])$ Matrices of nearly quadratic polynomials $\mathfrak{I}$ , a subset of $U_1(\mathbb{C})$ Some simple consequencesInterlacing in $\mathfrak{I}$ Particular interlacingsThe geometry of interlacing in $\mathfrak{I}$ Orthogonal type recurrencesThe transformation $x \mapsto exp(-\iota x)$	<b>589</b> 591 593 596 598 603 604 608 610 612 615

21	Stabl	e polynomials	619
	21.1	Stable polynomials in d variables	620
	21.2	Real stable polynomials	621
	21.3	Polynomials (not) in $\mathcal{H}_d$	623
	21.4	Stable polynomials with complex coefficients	625
	21.5	The analytic closure	630
	21.6	Real stable polynomials in one variable	631
	21.7	Coefficient inequalities in $\mathcal{H}_1 \dots \dots \dots \dots \dots \dots \dots$	636
	21.8	Positive interlacing	638
	21.9	Linear transformations	645
	21.10	The totally stable matrix conjecture	648
	21.11	Determinants of Hankel matrices	650
	21.12	Constructing stable polynomials	653
	21.13	Stable matrix polynomials	655
22	Trans	formations in the complex plane	659
	22.1	The derivative and interlacing	660
	22.2	Properties of algebraic Interlacing	661
	22.3	The map $\phi$	663
	22.4	Lines, rays, and sectors	665
	22.5	The set of roots	667
	22.6	Complex Transformations involving $f(x + i)$	667
	22.7	The Hermite-Biehler problem	670
	22.8	The Bôcher-Walsh problem	674
	22.9	Newton's inequalities in the complex plane	675
23	Fauix	variant Polynomials	679
20	23.1	Basic definitions and properties	679
	23.1	Interlacing	681
	23.3	Creating equivariant polynomials and the Hurwitz theorem	682
	23.4	Hermite polynomials	684
~ 4	<b>D</b> 1		<b>60</b>
24	Polyn	iomials modulo an ideal	685
	24.1	Initial segments of P	685
	24.2	Initial segments of $P_2$	687
	24.3		688
	24.4	Inere are many extensions	689
	24.5	Interlacing	690
	24.6	Linear transformations	690
	24.7	Stable polynomials	693
	24.ð	Coefficients of extensions	094 607
	∠4.7		07/

### Appendix

Glossary of polynomials, spaces, and interlacing	702
Tables of transformations	707
Empirical tables of transformations	710
Tables of determinants and integrals	713
Tables of Polynomials	716
Questions	721
Notes	745
Bibliography	750
Index	759

# List of Figures

1.1	A quintic with 4 critical points, and at most 3 roots
1.2	Two interlacing polynomials
1.3	Sign interlacing doesn't imply interlacing
1.4	Interlacing when the degrees are equal
1.5	Intervals containing roots of $f + \alpha g$ , $\alpha > 0$
1.6	Roots of $f + g$ where the degrees of f, g differ by 2
1.7	Graphs of interval polynomials
1.8	The possibilities for a quadratic
1.9	An impossible configuration
2.1	The positivity hierarchy
3.1	Roots of mutually interlacing polynomials
3.2	Impossible solution curves
3.3	The graph of $(a - b)^2 - 2(a + b) + 1 = 0 \dots 94$
3.4	Mutually interlacing polynomials evaluated at $\sigma$ 100
4.1	The Newton graph of a polynomial in $\mathbf{P}^{pos}$

4.2	The Newton graphs of two interlacing polynomials $\ldots \ldots$		. 119
5.1	A polynomial with rapidly decreasing coefficients		. 129
6.1 6.2	A Möbius transformation acting on a linear transformation The graph of the Chebyshev polynomial $T_5(x)$	 	. 158 . 169
7.1	The even-odd tree of $(x+1)^8$ $\hdots$		. 201
8.1 8.2	Understanding $\mathbb{A}$	 	. 220 . 228
9.1 9.2 9.3 9.4 9.5 9.6 9.7 9.8 9.9	The graph of a polynomial satisfying y but not x-substitution. Meets all lines through the origin, fails substitution The graph of a determinant	<ul> <li>.</li> <li>.&lt;</li></ul>	255 256 257 267 269 270 271 271 272 309
10.1	A polynomial in $\mathtt{P}_2(4)$ $\hfill \ldots$ . 		. 342
11.1 11.2 11.3	The graph of of a polynomial in $\overline{P}_2$	  	. 371 . 407 . 422
12.1 12.2 12.3 12.4 12.5 12.6 12.7 12.8 12.9 12.10	$\begin{array}{l} L_5(x;y) \text{ and its derivative (dashed) with respect to } y \ \ldots \$	<ul> <li>.</li> <li>.&lt;</li></ul>	. 432 . 434 . 435 . 437 . 438 . 438 . 438 . 439 . 440 . 441 . 443
13.1	The graph of $H_{2,3}$		. 461
14.1 14.2 14.3 14.4	The graph of $T_*f$ The graph of $\partial_y^6(x^2 + y^2 - 1)^6 = 0$ The graph of $p_{6,6,12}$ The graph of $\partial_y^6(x^2 + y^2 - 1)^6 = 0$ and its y derivative	· · · · · ·	. 467 . 478 . 478 . 479

18.2	The graph of a matrix product not in $Hyp_1$	553
18.3	The graph of a hyperbolic matrix polynomial	554
18.4	Overlapping root zones	557
18.5	The graphs of $v^t$ fv as v varies	558
18.6	Positive semi-definite leading coefficient	561
19.1	Location of roots of $ -xI + A + Bi $ if $B > A$ .	576
19.2	The image of the circle under a Möbius transformation	587
• • •		< 0 <b>0</b>
20.1	Trajectories for $f + t_1g$	602
20.2	Interlacing example	611
20.3	An argument plot for interlacing polynomials	611
20.4	Non-interlacing example	611
20.5	An argument plot for non-interlacing polynomials	612
20.6	Roots of non-interlacing polynomials	612
20.7	Coefficient ratios of $f \in \mathbf{P}_2^{\text{pos}}$ and $\sigma \in \mathbf{Q}_1$	617
	2	
21.1	The graph of a stable polynomial in two variables	625
22.1	Where the roots are	663
22.2	Image of the unit circle under Laguerre transformation	668
1		<b>F</b> 4 1
1	The imaginary part of a polynomial in $J_2$	/41

## List of Tables

6.1 6.2	Realizing possible domains and ranges $\dots \dots \dots$	48 51
8.1	Polynomials of the form $Q_n^{\alpha,\beta}$	26
9.1	Properties of $ Ax + By + C $	58
15.1	Generating function data	82
15.1	Concerning functions for the software time $\mathbf{D}$ $\mathbf{D}$	00
15.2	Generating runctions for transformations $\mathbf{r} \longrightarrow \mathbf{r}$ 40	89
15.3	Generating Functions in higher dimensions	06
15.4	Generating Functions for $\mathbf{P}_2 \longrightarrow \mathbf{P}$	12
15.5	Generating functions for $\mathbf{P}^{alt} \longrightarrow \mathbf{P}^{alt} \dots \dots$	14

#### List of Tables

15.6	Generating functions for $\mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}$	514
15.7	Miscellaneous generating functions	516
19.1	Properties of cones	567
20.1	Intervals where $(x + \iota)^n < (x + s \iota)^{n-1}$	610
20.2	Recurrences for $f_n + \iota f_{n-1} \dots \dots \dots \dots \dots \dots \dots \dots \dots$	614
1	Purported range of $\mathbf{P}^{pos}$ under the commutator map $\ldots \ldots \ldots$	727

Part I

# Polynomials with all real roots

CHAPTER



## **Polynomials in One Variable**

#### 1.1 Notation and basic definitions

We begin with the definitions of the various kinds of interlacing, the classes of polynomials of interest, and the properties of linear transformations that we shall be using.

If f is a polynomial of degree n with all real roots, then define

$$roots(f) = (a_1, \ldots, a_n)$$

where  $a_1 \leq a_2 \leq \cdots \leq a_n$  are the roots of f(x) = 0. If we write roots(f) then we are assuming that f actually has all real roots. Given two polynomials f, g with  $roots(f) = (a_1, \ldots, a_n)$  and  $roots(g) = (b_1, \ldots, b_m)$  then we say that f and g *interlace* if these roots alternate. If all the roots are distinct then there are only four ways that this can happen:

$$\begin{split} g \ll f & a_1 < b_1 < a_2 < b_2 < \cdots < a_n < b_n \\ g \leqslant f & b_1 < a_1 < b_2 < a_2 < \cdots < b_n \\ f \ll g & b_1 < a_1 < b_2 < a_2 < \cdots < b_n < a_n \\ f \leqslant q & a_1 < b_1 < a_2 < b_2 < \cdots < a_n. \end{split}$$
(1.1.1)

We define  $g \gg f$  to mean that  $f \ll g$ , and g > f to mean f < g. Notice that when we write  $f \ll g$  we assume that both f and g have all real roots, which are necessarily distinct. If two polynomials interlace then their degrees differ by at most one.

If the inequalities in (1.1.1) are relaxed, so that all inequalities "<" are replaced by " $\leq$ ", then we write  $g \leq f$ ,  $g \leq f$ ,  $f \leq g$  and  $f \leq g$  respectively. We call interlacings with  $\leq$  or  $\ll$  *strict interlacings*. It's easy to see that non-strict interlacings are limits of strict interlacings - see Remark 9.104 for a non ad hoc argument.

If f and g are two polynomials such that there is a polynomial h with  $f \leq h$ ,  $g \leq h$ , (or  $h \leq g$ ,  $h \leq f$ ) then we say that f, g have a *common interlacing*. Unless stated otherwise, we assume that f and g have the same degree.

The two interlacings  $f \leq g$  and  $f \leq g$  have very similar properties. To eliminate many special cases, we introduce the notation

 $f \leftarrow g$  which means  $f \underline{\ll} g$  or  $f \underline{\lessdot} g$ 

One way to remember this notation is to notice that  $f \leftarrow g$  means that

- f and g interlace
- The largest root belongs to f.
- The arrow points to the polynomial with the largest root.

There are several important subclasses of polynomials with all real roots.

 $\mathbf{P} = \{ all polynomials with all real roots \}$ 

 $\mathbf{P}^{pos} = \{ all polynomials with all real roots, all negative, \}$ 

and all coefficients are the same sign}

 $\mathbf{P}^{alt} = \{ all polynomials with all real roots, all positive, \}$ 

and the coefficients alternate in sign}

$$\mathbf{P}^{\pm} = \mathbf{P}^{alt} \cup \mathbf{P}^{pos}$$

 $\mathbf{P}^{I} = \{ all polynomials with all real roots, all roots lying in a set I \}$ 

$$\overline{\mathbf{P}^{\pm}} = \mathbf{P}^{(-\infty,0]} \cup \mathbf{P}^{[0,\infty)}$$

It is helpful to remember that that  $\mathbf{P}^{\text{pos}}$  has a superscript *pos* because whenever the leading coefficient is positive all the coefficients are positive. If we want to only consider polynomials of degree n then we write  $\mathbf{P}(n)$ ,  $\mathbf{P}^{\text{pos}}(n)$ , and so on.

We are particularly interested in properties of linear transformations of polynomials. Suppose that T is a linear transformation that maps polynomials to polynomials.

- T *preserves roots* if whenever f has all real roots, so does Tf. Using the notation above, we can write this as T: P → P.
- T preserves interlacing if whenever f and g interlace, so do Tf and Tg.
- T preserves degree if the degrees of f and Tf are the same.
- T preserves the sign of the leading coefficient if f and Tf have leading coefficients of the same sign.

The quintessential interlacing theorem is a well known consequence of Rolle's Theorem. The proof given below analyzes the graph of f and f' - see Remark 9.68 and Remark 15.28 for proofs that do not involve any analysis of the graph.

**Theorem 1.1.** *If* f *is a polynomial with all real roots, and all these roots are distinct, then*  $f \leq f'$ .

*Proof.* By Rolle's Theorem, there is a root of f' between consecutive roots of f. If the degree of f is n, then there are n - 1 intervals between roots of f, so f' has at least n - 1 roots. Since f' has degree n - 1, it has exactly n - 1 roots.  $\Box$ 

If f has all real roots, but all the roots of f are not distinct, then f and f' have roots in common. In this case we have that f < f'.

The derivative operator Df = f' is of fundamental importance. It is a linear transformation  $P \longrightarrow P$ ,  $P^{alt} \longrightarrow P^{alt}$ , and  $P^{pos} \longrightarrow P^{pos}$ . We just observed that D preserves roots. We will see in Theorem 1.47 that it also preserves interlacing.

We will occasionally make use of the stronger version of Rolle's theorem that states that there are an *odd* number of roots (counting multiplicity) of f' between any two consecutive distinct roots of f.

**Example 1.2.** If f is a polynomial with all real roots, then it is not necessarily true that some anti-derivative of f has all real zeros. Equivalently, the derivative does not map P(n) *onto* P(n - 1). A quintic p(x) such as in Figure 1.1 has four critical points, so p' has four zeros. However, no horizontal line intersects p in more than three points, and so no integral of p' will have more than three roots. For instance, (x + 1)(x + 2)(x + 4)(x + 5) is not the derivative of a quintic with all real roots.



Figure 1.1: A quintic with 4 critical points, and at most 3 roots

We introduce a relation that is weaker than interlacing. See Section 1.10 for more properties.

Definition 1.3. Suppose that

$$\begin{aligned} f(x) &= (x-r_1)\cdots(x-r_n) & r_1 \leqslant r_2 \cdots \leqslant r_n \\ g(x) &= (x-s_1)\cdots(x-s_m) & s_1 \leqslant s_2 \cdots \leqslant s_m \end{aligned}$$

We say that  $g \blacktriangleleft f$  if  $m \le n$ , and  $s_i \le r_i$  for i = 1, ..., m.

In other words, each root of g is at most the corresponding root of f. Note that the degrees of f and g do not need to be equal.

Common interlacing and  $\triangleleft$  imply interlacing. The proof of the Lemma is immediate.

Lemma 1.4. If

- 1. f **∢** g
- 2. f and g have the same degree
- 3. f and g have a common interlacing

then f  $\leq g$ .

Common interlacings can be split into two interlacings.

**Lemma 1.5.** If f, g have the same degree and have a common interlacing then there are  $f_1, f_2, g_1, g_2 \in \mathbf{P}$  such that

$f_1 \longleftarrow g_1$	$f=f_1f_2$
$g_2 \longleftarrow f_2$	$g = g_1 g_2$

*Proof.* Choose  $h \leq g$  and  $h \leq f$ . There is one root  $r_i$  of f and one root  $s_i$  of g between the ith and (i+1)st roots of h. We may assume that the roots of f and g are distinct. The desired polynomials are defined by

$roots(f_1) = \{r_i \mid r_i < s_i\}$	$roots(\mathfrak{g}_1) = \{ \mathfrak{s}_i \mid r_i < \mathfrak{s}_i \}$
$roots(f_2) = \{r_i \   \ r_i > s_i\}$	$\text{roots}(g_2) = \{s_i \   \ r_i > s_i\}$

*Remark* 1.6. Here are a few important properties about polynomials. See [141]. First of all, the roots of a polynomial are continuous functions of the coefficients. Consequently, if a sequence of polynomials  $\{f_n\}$  converges to the polynomial f and all  $f_n$  have all real roots, then so does f. In addition if there is a polynomial g such that  $g \leftarrow f_n$  for all n then  $g \leftarrow f$ . We should also mention that roots of polynomials with real coefficients come in conjugate pairs. It follows that if a polynomial of degree n has n - 1 real roots, then it has all real roots. Lastly, if g has all real roots, then so do f and g.

#### 1.2 Sign interlacing

The easiest way to show that two polynomials with all real roots interlace is to show that one *sign interlaces* the other. The *sign* of a number c is defined as

$$sgn c = \begin{cases} 1 & if c > 0 \\ -1 & if c < 0 \\ 0 & if c = 0 \end{cases}$$

We say that f *sign interlaces* g if the sign of f alternates on the roots of g. More precisely, there is an  $\epsilon = \pm 1$  so that if r is the i-th largest root of g then the sign of f(r) is  $\epsilon(-1)^i$ .

We usually assume that polynomials have positive leading coefficient. In this case, if  $f \ll g$  then there is exactly one root of f that is larger than the largest root r of g. Since f has positive leading coefficient, it must be the case that f(r) is negative. Consequently, if g has degree n then the sign of f at the i-th largest root of g is  $(-1)^{i+n+1}$ .

It should be clear that if f and g interlace then f sign interlaces g, and g sign interlaces f. The converse is not always true, and is discussed in the first three lemmas. Figure 1.2 shows two interlacing polynomials  $f \ll g$ , where f is the dashed line, and g the solid line. The solid circles indicate the value of f at the zeros of g - notice that these circles alternate above and below the x-axis.



Figure 1.2: Two interlacing polynomials

Lemma 1.7. Suppose f, g are polynomials satisfying

- f has all real roots, all distinct.
- $\deg(f) = \deg(g) + 1$
- g sign interlaces f

#### *then* $f \triangleleft g$ .

*Proof.* Assume that f has n roots. Since the sign of g alternates between consecutive roots of f, we find n - 1 roots of g, one between each pair of consecutive roots of f.

If we knew that f sign interlaced g in Lemma 1.7, then it does not follow that  $f \leq g$ . For instance, take  $f = x^3$  and g = (x - 1)(x + 1), so that the roots of f and g are  $\{0, 0, 0\}$  and  $\{-1, 1\}$ . The reason that g and f do not interlace is that although we have roots of f between roots of g, there aren't any roots of f outside of the roots of (g). In order to find these roots, we must control the behaviors of f and g by specifying appropriate signs at the largest and smallest roots of g.

Figure 1.3 shows graphically that if f (solid) sign interlaces g (dashed) then it is possible that f is not interlaced by g.



Figure 1.3: Sign interlacing doesn't imply interlacing.

Lemma 1.8. Suppose f, g are polynomials satisfying

- f has all real roots, all distinct.
- $\deg(f) = \deg(g) 1$
- g has positive leading coefficient.
- g sign interlaces f.
- Let  $\alpha$  be the smallest root of f, and  $\beta$  the largest root of f. If either
  - 1. sgn  $f(\beta) = -1$  or
  - 2. sgn  $f(\alpha) = 1$

*then*  $g \leq f$ .

*Proof.* Since g sign interlaces f there is a root of g between adjacent roots of f. This determines deg(f) – 2 roots, so it remains to find where the last two roots are. If  $sgn(g(\beta)) = -1$  then since  $\lim_{x\to\infty} sgn(g(x)) = 1$  there are an odd number of roots (hence exactly one) to the right of  $\beta$ . Similarly there is a root to the left of  $\alpha$ . See Figure 1.3.

If f and g both have the same degree and they sign interlace, then either  $f \ll g$  or  $g \ll f$ . Again, we need extra information about the behavior on the extremes to determine which case holds. Figure 1.4 shows the two possibilities.

Lemma 1.9. Suppose f, g are polynomials satisfying



Figure 1.4: Interlacing when the degrees are equal.

- f has all real roots, all distinct.
- g has positive leading coefficient.
- deg(f) = deg(g) = n
- g sign interlaces f.
- Let  $\alpha$  be the smallest root of f and  $\beta$  the largest root. Then
  - 1. If sgn  $g(\beta) = 1$  then  $f \ll g$ .
  - 2. If sgn  $g(\beta) = -1$  then  $g \ll f$ .
  - 3. If sgn  $g(\alpha) = -(-1)^n$  then  $g \ll f$ .
  - 4. If sgn  $g(\alpha) = (-1)^n$  then  $f \ll g$ .

*Proof.* Assume that f has n roots. Since the sign of g alternates between consecutive roots of f, we find n - 1 roots of g, one between each pair of consecutive roots of f. Now g has an odd number of roots in each of these intervals, and there cannot be three roots of g in any of them, for then g would have at least n + 2 roots, so there is one more root of g, to either the left of the smallest root of f, or to the right of the largest root. Thus, we have either  $f \ll g$  or  $g \ll f$ .

All four parts are similar, so we will only establish the first one. Assume that sgn  $g(\beta) = 1$ . Counting the number of alternations of sign of g gives

$$\begin{split} \lim_{x \to -\infty} \operatorname{sgn} \mathfrak{g}(x) &= (-1)^n \\ & \operatorname{sgn} \mathfrak{g}(\alpha) &= (-1)^{n-1} \operatorname{sgn} \beta = (-1)^{n-1} \end{split}$$

and hence there is a root of g in  $(-\infty, \alpha)$ . This shows that  $f \ll g$ .

As a first consequence of these elementary but fundamental results, we choose

**Lemma 1.10.** If f, g have positive leading coefficients and there is an h so that  $h \leftarrow f$ , g then for all positive  $\alpha$  and  $\beta$  the linear combination  $\alpha f + \beta g$  has all real roots and  $h \leftarrow \alpha f + \beta g$ .

*Proof.* f, g both sign interlace h, and since they have positive leading coefficients, they are the same sign on all the roots of h. Consequently,  $\alpha f + \beta g$  has the same sign on a root of h as does f or g. The result follows from Lemmas 1.9 and 1.8.

Although the sum of two arbitrary polynomials with all real roots does not necessarily have all real roots, they might if one of them is sufficiently small.

**Lemma 1.11.** Suppose  $f \in P$  has degree n and all distinct roots. If  $g \in P$  has degree n + 1 then  $f + \epsilon g \in P$  for sufficiently small  $\epsilon$ .

*Proof.* The roots of  $f + \epsilon g$  converge to the roots of f as  $\epsilon$  goes to zero. Since the n roots of f are all distinct,  $f + \epsilon g$  has n real roots for  $\epsilon$  sufficiently small. Since there is only one root unaccounted for, it must be real.

The next well-known lemma is easy to see geometrically.

#### Lemma 1.12.

- 1. If there is an  $\alpha$  such that  $f(x) + c \in P$  for all  $|c| > \alpha$  then f is linear.
- 2. If there is an  $\alpha$  such that  $f(x) + c \in P$  for all  $c > \alpha$  then f is quadratic with negative leading coefficient.
- 3. If there is an  $\alpha$  such that  $f(x) + c \in P$  for all  $c < \alpha$  then f is quadratic with positive leading coefficient.

Sign interlacing determines the intervals where sums of polynomials have roots. In general the roots lie in alternating intervals, but which set of intervals depends on the behavior at infinity. This is in turn determined by the sign of the leading coefficient of the sum. For instance Figure 1.5 shows two polynomials f (dashed) and g (solid). The intervals on the x-axis where  $f + \alpha g$  has roots for positive  $\alpha$  are solid; the intervals where they do not have a root (since both functions have the same sign) are dotted.



Figure 1.5: Intervals containing roots of  $f + \alpha g$ ,  $\alpha > 0$ .

**Lemma 1.13.** Assume that polynomials f, g, h have positive leading coefficients. If f and g interlace, and  $\begin{vmatrix} f & g \\ g & h \end{vmatrix} < 0$  for all x then g and h interlace. In more detail we can say that

- If  $f \leq g$  then  $g \leq h$ .
- If  $f \ll g$  then  $g \leftarrow h$ .
- If f > g then  $g \longrightarrow h$ .

#### • If $f \gg g$ then $g \gg h$ .

*Proof.* Assume that the degree of g is n. If r is a root of g then upon evaluating the determinant at r we find that f(r)h(r) < 0. Since f sign interlaces g, so also does h sign interlace g. This implies that the degree of h is at least n - 1. It is not possible that the degree of fh is greater than 2n, for then  $fh - g^2$  will be positive for large values of x, since the leading coefficient of fh is positive. We now consider the four cases in turn.

If  $f \leq g$  then the degree of f is n + 1. Since the degree of h is at least n - 1, and the degree of fh is at most 2n, the degree of h is exactly n - 1. Since h sign interlaces g, we can apply Lemma 1.7 to conclude that  $g \leq h$ .

If  $f \ge g$  then the degree of f is n - 1, and the degree of h is either n + 1, n - 1 or n. The sign of f at the largest root r of g is positive, so the sign of h at r is negative. Since the leading coefficient of h is positive, there is a root of h in  $(r, \infty)$ , and so the degree of h can not be n - 1. Lemma 1.9 now implies that  $h \le g$  if the degree of h is n + 1. If the degree of h is n then  $h \ll g$ .

When  $f \gg g$  then the degree of h is n or n - 1. Again the sign of f on the largest root r of g is positive, so there is a root of h in  $(r, \infty)$ . Thus h has exactly n roots and  $g \gg h$ .

If  $f \ll g$  then a similar argument applies.

**Corollary 1.14.** If f < g have positive leading coefficients and h = af + (bx + c)g then h < g if and only if the sign of the leading coefficient of h is sign(a).

*Proof.* Since f sign interlaces g it follows that h sign interlaces g. If  $\beta$  is the largest root of g then  $f(\beta) < 0$  so the sign of  $h(\beta)$  is -sign(a). Use Lemma 1.8.

As a consequence we get a special case that will prove useful later.

**Corollary 1.15.** If  $g \in P(n)$  has positive leading coefficient and  $a \notin (0, n)$  then  $xg' - ag \in P$ .

If  $a \in (0, n)$  and g has p positive roots and m negative roots then xg' - ag has at least p - 1 positive roots and at least m - 1 negative roots. It is possible that  $xg' - ag \notin P$ .

*Proof.* We may assume that the leading coefficient of g is 1. The leading coefficient of xg' - ag is (n - a). If a > n then we can apply Corollary 1.14. If a < 0 then note that  $xg \leq g$  and  $xg \leq xg'$  and therefore  $xg \leq xg' + (-a)g$ .

If  $a \in (0, n)$  then xg' - ag sign interlaces the positive roots of g and the negative roots of g. This implies that it has at least p - 1 positive roots, and m - 1 negative roots, but it does not imply that the remaining two roots are real. For example, if  $g = x^2 - 1$  and a = 1 then  $xg' - g = x^2 + 1$  which has no real roots.

The locations of the roots of an arbitrary linear combination can be nearly any value. For instance,

**Lemma 1.16.** *If* f, g *interlace, are not multiples of one another, and* (a, b) *is any non-empty interval then there is an*  $\alpha$  *such that* 

- $f + \alpha g \in P$ .
- $f + \alpha g$  has a root in (a, b).

*Proof.* Choose an  $r \in (a, b)$  that is not a root of g. If we define  $\alpha = -\frac{f(r)}{g(r)}$  then r is a root of  $f + \alpha g$ . Now  $f + \alpha g$  is not the zero polynomial since f and g are not multiples of one another, so  $f + \alpha g$  is in **P**.

However, in the case of the derivative, we can give precise information about the location of the roots of a linear combination.

**Lemma 1.17.** If  $f \in \mathbf{P}^{(a,b)}(n)$  and  $\alpha > 0$  then  $f + \alpha f' \in \mathbf{P}^{(a-\alpha n,b)}$ .

*Proof.* The only question is the location of the smallest root. Assume f has positive leading coefficient. Since sgn  $f(a - \alpha n) = (-1)^n$  it suffices to show that

$$\frac{1}{\alpha} > \frac{-f'}{f}(\alpha - \alpha n) \tag{1.2.1}$$

since this implies sgn  $(f + \alpha f')(a - \alpha n) = (-1)^n$ . Suppose that the roots of f are  $r_1, \ldots, r_n$ . Then

$$\frac{-f'}{f}(a-\alpha n) = \sum \frac{1}{\alpha n - a + r_i}$$

Now since  $r_i > a$  it follows that  $\alpha n - a + r_i > \alpha n$  and so

$$\frac{1}{\alpha} = n \frac{1}{\alpha n} > \sum \frac{1}{\alpha n - a + r_i}$$

		L
		L
_		-

The following lemma combines addition and subtraction, since  $f^2 - g^2 = (f + g)(f - g)$ . It is not a coincidence that the terms come from the binomial expansion  $(f + g)^2$  - the result will be generalized in Lemma 20.36.

**Lemma 1.18.** Suppose that  $f \ll g$ , the leading coefficient of f is  $c_f$ , and the leading coefficient of g is  $c_g$ .

If	$ c_{f}  >  c_{g} $	then	$f^2 - g^2$	$\ll$	2fg
İf	$ c_{f}  <  c_{g} $	then	$f^{2} - g^{2}$	$\gg$	2fg
İf	$ c_{f}  =  c_{q} $	then	$f^2 - q^2$	≫	2 fg

*Proof.* The proof is a standard sign interlacing argument. For example, if the degree is 3, then the diagram below shows where the roots of f + g (marked "+") and f - g (marked "-") are:

$$g \stackrel{+}{-\!\!-\!\!-} f \stackrel{-}{-\!\!-} g \stackrel{+}{-\!\!-\!\!-} f \stackrel{-}{-\!\!-} g \stackrel{+}{-\!\!-} f$$

In the third case this is all of the roots. In the first case there is a root on the far right, and in the second case the remaining root is on the far left.  $\Box$ 

The following convolution lemma is a non-obvious consequence of sign interlacing. It will be generalized to a wider class of polynomials in Lemma 3.16.

**Lemma 1.19.** Choose  $a_1 \leq \cdots \leq a_n$ ,  $b_1 \leq \cdots \leq b_n$  and define  $f = \prod (x - a_i)$ ,  $g = \prod (x - b_i)$ . The following polynomial has all real roots

$$h = \sum_{i=1}^{n} \left( \frac{f}{x - a_i} \right) \cdot \left( \frac{g}{x - b_{n+1-i}} \right)$$

*Proof.* Without loss of generality we may assume that the roots of fg are  $r_1 < \cdots < r_{2n}$ . We will show that the sign of  $h(r_m)$  only depends on m. More precisely, we show that  $sgn(h(r_m)) = (-1)^{m+1}$  if  $m \leq n$ , and  $(-1)^m$  if m > n. This shows that there are 2n - 2 intervals between roots where the sign at the endpoints is positive and negative, so h has 2n - 2 roots, and hence is in **P** since its degree is 2n - 2.

We may assume that  $r_m = a_i$ . Almost all terms of h are zero when evaluated at  $r_m$ :

$$h(\mathbf{r}_{\mathfrak{m}}) = \left(\frac{f}{x - a_{\mathfrak{i}}} \cdot g \cdot \frac{1}{x - b_{\mathfrak{n}+1-\mathfrak{i}}}\right)(a_{\mathfrak{i}})$$

We investigate each of the three factors in turn. The first factor is simply  $f'(a_i)$ , and the sign of this is  $(-1)^{i+n}$ .

Next, since  $g(a_i) = \prod_j (a_i - b_j)$ , the sign of  $g(a_i)$  depends on the number of  $b_j$ 's that are greater than  $a_i$ . Since  $a_i$  is the m-th largest root, there are exactly m-i roots of g that are smaller than  $a_i$ , and hence there are n-(m-i)roots of g that are greater than  $a_i$ . Therefore,  $sgn(g(a_i) = (-1)^{n-m+i}$ .

Finally, consider the sign  $\epsilon$  of  $a_i - b_{n+1-i}$ . The largest j such that  $b_j$  is less than  $a_i$  is j = m - i. Consequently, the sign  $\epsilon$  is negative whenever (n + 1 - i) > m - i, and so  $\epsilon = 1$  if  $m \le n$  and -1 if m > n. Combining these calculations shows that the sign of  $h(r_m)$  is  $(-1)^{i+n}(-1)^{n-m+i}\epsilon$  which is as claimed.

#### **1.3** A quantitative version of sign interlacing

If f, g are two polynomials of the same degree n that sign interlace, then we know the signs of g evaluated at the n roots of f. This is similar to interpolation, where we are given the *values* of g, not just the *signs*, at these roots and use this information to determine g. We can combine these two ideas to give a quantitative version of sign interlacing.

Let f be a polynomial with distinct roots  $roots(f) = (a_1, ..., a_n)$ . The n polynomials

$$\frac{f(x)}{x-a_1}, \cdots, \frac{f(x)}{x-a_n}$$

are of degree n - 1 and form a basis for the polynomials of degree n - 1. If we add any polynomial of degree n (e.g. f itself) then we get a basis for the polynomials of degree n.

**Lemma 1.20.** Assume that f is a polynomial of degree n, with positive leading coefficient, and with roots  $\{a_1, \ldots, a_n\}$ . Suppose that g is a polynomial with positive leading coefficient.

1. If g has degree n - 1 and we write

$$g(x) = c_1 \frac{f(x)}{x - a_1} + \dots + c_n \frac{f(x)}{x - a_n}$$
(1.3.1)

then  $f \leq g$  if and only if all  $c_i$  are positive.

2. If g has degree n and we write

$$g(x) = cf(x) + c_1 \frac{f(x)}{x - a_1} + \dots + c_n \frac{f(x)}{x - a_n}$$
(1.3.2)

then c is positive, and

- $f \ll g$  if and only if all  $c_i$  are positive.
- $f \gg g$  if and only if all  $c_i$  are negative.
- 3. If g has degree n + 1 and we write

$$g(x) = (dx + c)f(x) + c_1 \frac{f(x)}{x - a_1} + \dots + c_n \frac{f(x)}{x - a_n}$$
(1.3.3)

then d is positive, and  $g \leq f$  if and only if all  $c_i$  are negative.

*Proof.* Evaluate g (in all three cases) at the roots of f:

$$g(a_i) = c_i(a_i - a_1) \cdots (a_i - a_{i-1})(a_i - a_{i+1}) \cdots (a_i - a_n)$$

so  $\operatorname{sgn} g(a_i) = \operatorname{sgn}(c_i)(-1)^{i+n}$ . In the first case we know that  $\operatorname{deg}(g) = \operatorname{deg}(f) - 1$ . If all the  $c_i$  have the same sign then g sign interlaces f, and the result follows from Lemma 1.7.

In the second case, if all  $c_i$  are positive, and deg(g) = deg(f), then g sign interlaces f. Since the sign of g at the largest root  $a_n$  of f is the sign of the leading coefficient of f, and the leading coefficient of g is positive, the result follows from Lemma 1.9.

In the third case, sgn  $g(a_n) = \text{sgn } c_n$  and since the leading coefficient of g is positive, we find  $g \leq f$  by Lemma 1.8.

Conversely, if  $f \leq g$ , then g sign interlaces f, and so the signs of  $g(a_i)$  alternate. Thus, all  $c_i$ 's have the same sign, which must be positive since the leading coefficient of g is positive. The converses of the second and third cases are similar, and omitted.

*Remark* 1.21. We can extend Lemma 1.20 by noticing that we do not require the coefficients  $c_i$  to be constant. If  $c_i(x)$  is positive (or negative) on an interval containing the roots of f, then g sign interlaces f.

One special case of (1.3.1) is when all the  $c_1$  are 1. In this case, g is exactly the derivative of f.

If we relax the positivity and negativity conditions of the parameters  $c_i$ 's in Lemma 1.20, then we still have interlacing. For example, when the degree of g is n - 1, then  $f \leq g$  if and only if all  $c_i$  are non-negative. This case where some of the coefficients  $c_i$  are zero follows by continuity.

#### **Corollary 1.22.** *If* $f \leq g$ *have positive leading coefficients then*

 $xf - bg \leq f$  for any positive b. In addition,  $xg + bf \leq g$ . If b is negative then it is possible that xf - bg and xg + bf are not in **P**.

Proof. The first is immediate from Lemma 1.20 if we write

 $g = \sum c_i \frac{f}{x - a_i}$ . In the second case we can write  $f = \alpha xg + \beta g - r$  where  $g \leq r$ . Since  $(xg + bf) + \gamma g = (\alpha + b)g + (b\beta + \gamma)g - br$  the conclusion follows.  $\Box$ 

We can also get interlacings such as  $xg + bf \leq g$  if we restrict g to  $\mathbf{P}^{pos}$ . See Lemma 2.15.

The equation

$$(-1)^{i+n}\operatorname{sgn} g(a_i) = \operatorname{sgn} c_i \tag{1.3.4}$$

shows that the signs of the  $c_i$ 's encode the sign of g evaluated at the roots of f. We can easily determine the  $c_i$ 's in terms of g and f. For instance, if  $f \leq g$  then  $c_i = g(a_i)/f'(a_i)$ .

There is another condition on the signs of the coefficients in (1.3.1) that implies that g has all real roots.

**Lemma 1.23.** Suppose that the roots of the monic polynomial f are  $a_1 < a_2 < \cdots < a_n$ . If  $c_1, \ldots c_n$  in (1.3.1) has exactly one sign change then g(x) in (1.3.1) has all real roots. If  $c_1, \ldots, c_n$  has exactly two sign changes,  $c_1 > 0$ , and  $\sum c_i < 0$  then g(x) has all real roots.<sup>1</sup>

*Proof.* If the  $c_i$ 's have exactly one sign change, then assume that the terms of the sequence  $c_1, \ldots, c_r$  are positive, and  $c_{r+1}, \ldots, c_n$  are negative. (1.3.1) shows that g sign interlaces f between  $a_1, \ldots, a_r$  giving r - 1 roots, and also between  $a_{r+1}, \ldots, a_n$  giving n - r - 1 roots for a total of n - 2 roots. Since g has degree n - 1 it follows that g has all real roots.

If there are exactly two sign changes then the above argument shows that there are at least n - 3 real roots. The  $c_i$ 's are initially positive, then negative, and then positive again. The leading coefficient of g is  $\sum c_i$  which is negative, and since sgn  $g(a_n) = \text{sgn } c_n > 0$  there is a root of g in the interval  $(a_n, \infty)$ , which again implies that g has all real roots.

<sup>&</sup>lt;sup>1</sup>The condition  $\sum_{i} c_i < 0$  is necessary. If f(x) = (x+1)(x+2)(x+3) then a quick computation shows that  $\frac{f}{x+1} - \frac{f}{x+2} \neq P$ , and  $\sum_{i} c_i = 1$ .

Common interlacing puts no restrictions on the signs of the coefficients.

**Lemma 1.24.** *Any sign pattern can be realized by a pair of polynomials with a common interlacing.* 

*Proof.* This means that if we are given any sequence  $\varepsilon_i$  of pluses and minuses then we can find two polynomials  $g_1, g_2$  with a common interlacing such that if  $g_2 = \prod (x - r_i)$  and

$$g_1 = bg_2 + \sum a_i \frac{g_2(x)}{x - r_i}$$

then  $sign(a_i) = \epsilon_i$ .

We prove this constructively; the initial cases are trivial. Assume we are given  $g_1, g_2$  as above, and choose  $\alpha > \beta$  to both be smaller than the smallest root of  $g_1g_2$ . We will show that the sign sequence of

$$f_1 = (x - \alpha)g_1, \qquad f_2 = (x - \beta)g_2$$

is

$$+, \epsilon_1, \ldots, \epsilon_n$$

By construction  $f_1$  and  $f_2$  have a common interlacing. Now  $a_i = g_1(r_i)/g'_2(r_i)$ . Thus the sequence of signs determined by  $f_1, f_2$  is

$$\frac{f_1(\beta)}{f'_2(\beta)}, \frac{f_1(r_1)}{f'_2(r_1)}, \dots, \frac{f_1(r_n)}{f'_2(r_n)}$$

Now

$$\begin{aligned} \frac{f_1(\beta)}{f'_2(\beta)} &= (\beta - \alpha) \frac{g_1(\beta)}{g'_2(\beta)} \\ \frac{f_1(r_i)}{f'_2(r_i)} &= \frac{(r_i - \alpha)g_1(r_i)}{(r_i - \beta)g_2(r_i) + g'_2(r_i)} = (r_i - \alpha) \frac{f_1(r_i)}{f'_2(r_i)} \end{aligned}$$

Since  $\beta$  is smaller than all the roots of  $g_1$  and  $g_2$ , the signs of  $g_1(\beta)$  and  $g'_2(\beta)$  are opposite. Also  $r_i - \alpha$  is positive, so the signs are as claimed.

If we choose  $\alpha > \beta$  instead, we get a minus sign for the sign, so all possibilities can happen.

The next result is a simple consequence of quantitative sign interlacing, and is useful for establishing many interlacing results.

**Lemma 1.25.** Assume  $f, g, j, k \in P$  have positive leading coefficients, and f < g,  $k \ll f \ll j$ . Define

$$\mathbf{h} = \ell_1 \mathbf{f} + \ell_2 \mathbf{j} + \ell_3 \mathbf{k} + \mathbf{q} \mathbf{g}$$

where  $\ell_1, \ell_2, \ell_3$  are linear, and q is at most quadratic. The interlacing  $h \leq f$  holds if these three conditions are true:

- 1. The leading coefficients of  $l_1$ ,  $l_2$ ,  $l_3$  are positive.
- 2. q and  $\ell_2$  are negative on the roots of f.
- 3.  $l_3$  is positive on the roots of f.

*Proof.* Assume that  $roots(f) = (a_1, ..., a_n)$  and write

$$g = \sum \alpha_{i} \frac{f}{x - a_{i}}$$
$$j = \beta f + \sum \beta_{i} \frac{f}{x - a_{i}}$$
$$k = \gamma f + \sum \gamma_{i} \frac{f}{x - a_{i}}$$

where  $\alpha_i$ ,  $\beta_i$ ,  $\beta$ ,  $\gamma$  are positive, and  $\gamma_i$  is negative. From the definition

$$h = (\ell_1 + \beta \ell_2 + \gamma \ell_3)f + \sum (\beta_i \ell_2 + \gamma_i \ell_3 + \alpha_i q) \frac{f}{x - \alpha_i}$$

The hypotheses guarantee that the leading coefficient of  $\ell_1 + \beta \ell_2 + \gamma \ell_3$  is positive, and that  $\beta_i \ell_2 + \gamma_i \ell_3 + \alpha_i q$  is negative on the roots of f. The result now follows from Lemma 1.20.

We will see consequences of the theorem in § 2.2.4. As a quick example,

$$\mathbf{x}\mathbf{f} - \mathbf{f} - \mathbf{f}' \in \mathbf{P}$$
 if  $\mathbf{f} \in \mathbf{P}$  (1.3.5)

because  $\ell_1 = x - 1$ ,  $\ell_2 = \ell_2 = 0$ , q = -1 satisfy conditions 1, 2, 3. The hypotheses on  $\ell_1$ ,  $\ell_2$ ,  $\ell_3$  can be weakened by only requiring that  $\ell_1 + \beta \ell_2 + \gamma \ell_3$  has positive leading coefficient.

If we assume that all roots of f are positive (or negative) we can get more specific interlacing results. These interlacing facts will be useful in showing that various linear transformations preserve interlacing. See § 2.2.4.

If polynomials f, g are in the relation f  $\leftarrow$  g, then g behaves like a derivative of f.

**Lemma 1.26** (Leibnitz Rule). Suppose that  $f, f_1, g, g_1$  are polynomials with positive leading coefficients, and with all real roots. Assume that f and g have no common roots.

• If  $f \leftarrow f_1$  and  $g \leftarrow g_1$  then  $f_1g$ ,  $fg_1$ ,  $fg \leftarrow fg_1 + f_1g \leftarrow f_1g_1$ 

In particular,  $fg_1 + f_1g$  has all real roots. Conversely, if  $f, g, f_1, g_1$  are polynomials that satisfy  $fg \leftarrow f_1g + fg_1$  then

- 1. f, g, f<sub>1</sub>, g<sub>1</sub>  $\in$  *P*
- $2. \ f \longleftarrow f_1$
- 3.  $g \leftarrow g_1$

*Proof.* We only consider the case where the left hand side is fg, the other cases are similar. Let  $f = a \prod_{1}^{n} (x - a_i)$ ,  $g = b \prod_{1}^{n} (x - b_i)$  If we write

$$f_1(x) = cf(x) + c_1 \frac{f(x)}{x - a_1} + \dots + c_n \frac{f(x)}{x - a_n}$$
(1.3.6)

$$g_1(x) = dg(x) + d_1 \frac{g(x)}{x - b_1} + \dots + d_m \frac{g(x)}{x - b_m}$$
(1.3.7)

where all the  $c_i$ 's and  $d_i$ 's are positive and c, d are non-negative then

$$fg_{1} + f_{1}g = (c + d)f(x)g(x) + c_{1}\frac{f(x)g(x)}{x - a_{1}} + \dots + c_{n}\frac{f(x)g(x)}{x - a_{n}} + d_{1}\frac{f(x)g(x)}{x - b_{1}} + \dots + d_{m}\frac{f(x)g(x)}{x - b_{m}}$$
(1.3.8)

and the result follows from Lemma 1.20.

Conversely, if  $fg \leftarrow fg_1 + f_1g$  then we first observe that  $f, g \in \mathbf{P}$ . We can write  $f_1$  in terms of f as in (1.3.6), and  $g_1$  in terms of g as in (1.3.7), but we do not know the signs of the  $c_i$ 's and the  $d_i$ 's. However, from (1.3.8) we know all the coefficients are negative. Thus the  $c_i$ 's and  $d_i$ 's are negative. The coefficients c, d are non-negative since all the polynomials have positive leading coefficients. Thus  $f \leftarrow f_1$  and  $g \leftarrow g_1$ .

**Corollary 1.27.** Suppose that  $f \leq g$  have positive leading coefficients, and a, c are positive.

If  $ad \leq bc$  then  $(ax + b)f + (cx + d)g \in P$ .

*Proof.* Note that  $ad \leq bc$  is equivalent to  $cx + d \leq ax + b$ . Apply Lemma 1.26.

*Remark* 1.28. The Lamé differential equation is

$$f(x)y'' + g(x)y' + h(x)y = 0$$
(1.3.9)

Heine and Stieltjes proved that if  $g \in \mathbf{P}(p)$ , and  $f \leq g$  then there are exactly  $\binom{n+p-1}{n}$  polynomials h(x) such that the equation has a solution y which is a polynomial of degree n. Moreover, both h and y are in  $\mathbf{P}$ . The polynomials h are called Van Vleck polynomials, and the corresponding solution is a Heine-Stieltjes polynomial.

Although h and f do not interlace, we do have [151] that  $fy' \leq hy$ . To see this, note that  $y' \leq y''$  and  $f \leq g$ , so by Leibnitz we have

$$fy' \leq fy'' + gy' = -hy$$

This also shows that if  $y \in \mathbf{P}$  then  $h \in \mathbf{P}$ .

Recall that a *cone* is a set in a vector space that is closed under non-negative linear combinations. Lemma 1.20 describes all polynomials that interlace a given polynomial. If we restrict ourselves to polynomials with positive leading coefficients then this set of polynomials is a cone. This is also true for polynomials in several variables - see Chapter 11.11.18.

#### 1.4 Elementary interlacings

In this section we use an algebraic definition of  $f \leq g$  to derive many fundamental interlacings. The following is a simple consequence of Lemma 1.20.

**Corollary 1.29.** If f, g have positive leading coefficients then  $f \leftarrow g$  if and only there is a polynomial h with positive leading coefficient and non-negative  $\alpha$  such that  $f \leq h$  and  $g = \alpha f + h$ . In addition, if  $g \leftarrow f$  then there is a positive  $\alpha$  and an h with positive leading coefficient satisfying  $f \leq h$  such that  $g = \alpha f - h$ . If the interlacing  $f \leftarrow g$  is strict then f < h.

If  $f \leftarrow g$  then for any  $\alpha$ ,  $\beta$  the linear combination  $\alpha f + \beta g$  interlaces f and g. The exact direction depends on  $\alpha$  and  $\beta$ . The converse is also true and of fundamental importance – see Proposition 1.35. This result is useful, for it gives us a simple way to show that the sum of certain polynomials with all real roots has all real roots.

**Corollary 1.30.** Suppose  $f \leftarrow g$ , both f and g are monic, and  $\alpha$ ,  $\beta$  are non-zero.

 $\mathit{lf}\, f \,\underline{\ll}\, g$ 

 $\alpha f + \beta g \longrightarrow f$  if  $\beta$  and  $\alpha + \beta$  have the same sign  $\alpha f + \beta g \longleftarrow f$  if  $\beta$  and  $\alpha + \beta$  have opposite signs or  $\alpha + \beta = 0$   $\alpha f + \beta g \longleftarrow g$  if  $\alpha$  and  $\alpha + \beta$  have the same sign  $\alpha f + \beta g \longrightarrow g$  if  $\alpha$  and  $\alpha + \beta$  have opposite signs or  $\alpha + \beta = 0$ 

If  $f \leq g$  and  $\alpha$ ,  $\beta$  have the same sign

$$f \longleftarrow \alpha f + \beta g \longleftarrow g$$
$$f \longrightarrow \alpha f - \beta g \longleftarrow g$$

*Proof.* Since f, g are monic we can write g = f + h as in Corollary 1.29. We prove the first two assertions; the remainder are similar. Substitution yields  $\alpha f + \beta g = (\alpha + \beta)f + \beta h$ . Since  $(\alpha + \beta)$  and  $\beta$  are the same sign, the desired interlacing follows since  $f \rightarrow f$  and  $h \rightarrow f$ .

For the second one, write in terms of g:  $\alpha f + \beta g = (\alpha + \beta)g - \alpha h$ . If  $\alpha + \beta$  and  $\alpha$  have opposite signs then  $g \to f$  and  $h \to f$ .

We have seen that we can add interlacings. This is a fundamental way of constructing new polynomials with all real roots. Here are some variations.

**Lemma 1.31.** Suppose f, g, h have positive leading coefficients, and  $\alpha$ ,  $\beta$  are positive.

- 1. If  $f \leftarrow g$  and  $h \leftarrow g$  then  $\alpha f + \beta h \leftarrow g$ .
- 2. *If*  $f \leftarrow g$  *and*  $f \leftarrow h$  *then*  $f \leftarrow \alpha g + \beta h$ .
- 3. If  $f \leftarrow g \leq h$  then  $\alpha f \beta h \leftarrow g$ .

*Proof.* Immediate from the representations of Lemma 1.20 and Corollary 1.29.  $\Box$ 

An *interlacing square* is a collection of four polynomials f, g, h, k that interlace as follows:  $f \swarrow k$ . If we have such a square then there are interlacings between certain products. It is interesting to note that although we do not have the interlacing fg — gk nor fg — fh it is true that the sum interlaces: fg — gk + fh.





*Proof.* The interlacings involving fk + gh follow from Corollary 1.30, as do the right arrows from fh + gk. We now show that fg and gk + fh interlace. Let f and g have roots  $\{a_1, \ldots, a_n\}$  and  $\{b_1 \leq \cdots \leq b_n\}$ . Interlacing follows from these evaluations at the roots of f and g:

$$sgn(gk + fh)(a_i) = sgn g(a_i)k(a_i) = 1$$
  

$$sgn(gk + fh)(b_i) = sgn f(b_i)h(b_i) = -1$$

Since  $f \leftarrow g$  the sign of gk + fh at the largest root  $b_n$  of fg is negative, and so  $fg \leftarrow gk + fh$ . The remaining case is similar.

*Remark* 1.33. We can use the representation of Corollary 1.29 to derive a simple relationship between the coefficients of interlacing polynomials that will be generalized later. Suppose  $f \ll g$  have positive leading coefficients and  $f = a_n x^n + a_{n-1} x^{n-1} + \cdots$ ,  $g = b_n x^n + b_{n-1} x^{n-1} + \cdots$ . We can write  $g = \alpha f + r$  where  $\alpha = b_n/a_n$ , and the leading coefficient of r is  $b_{n-1} - \frac{b_n}{a_n}a_{n-1}$ . As long as g is not a multiple of f we know that the leading coefficient of r is positive. Using the fact that  $a_n$  is positive, we conclude that  $\left| \begin{array}{c} a_n a_{n-1} \\ b_n b_{n-1} \end{array} \right| > 0$ . In Corollary 4.31 we will show that all such determinants of consecutive coefficients are positive.

Squares don't interlace.

**Lemma 1.34.** *If*  $f, g \in P$ , and  $f^2$  and  $g^2$  interlace then f is a constant multiple of g.

*Proof.* If g has a root that is not a root of f then  $g^2$  has two consecutive roots that are not separated by a root of f. Thus g divides f, and f divides g, so they are constant multiples.

#### **1.5** Linear combinations of polynomials

When do all linear combinations of two polynomials with all real roots have all real roots? The answer is quite simple: they interlace. This important result is central to the generalizations of interlacing to polynomials in several variables that is developed in Chapter 9, and of polynomials with complex roots. The following proposition has been (re)discovered many times, see [141, page 210].

**Proposition 1.35.** Suppose relatively prime polynomials f, g have all real roots, all distinct, with deg(f)  $\ge$  deg(g). Suppose that for all positive  $\alpha$ ,  $\beta$  the linear combination  $\alpha f + \beta g$  has all real roots. Then either

- 1. deg(f) = deg(g) and there is a polynomial k such that  $k \leq f, k \leq g$ .
- 2. deg(f) = deg(g) + 1 and then there is a polynomial k such that  $k \leq g$ ,  $k \gg f$  or  $k \ll f$ .
- 3. deg(f) = deg(g) + 2 and there is a k so that  $f \le k \le g$ .

If  $\alpha f + \beta g$  has all real roots for all positive and negative  $\alpha$ ,  $\beta$ , then either

- 4.  $\deg(f) = \deg(g)$  and either  $f \ll g$  or  $g \ll f$ .
- 5.  $\deg(f) = \deg(g) + 1$  and  $f \leq g$ .

In addition, the converses of all the cases are valid.

*Proof.* Assume that the roots of fa nd g are  $a_1 \leq \cdots \leq a_n$  and  $b_1 \leq \cdots \leq b_m$ . These n + m points determine n + m + 1 open intervals, and f and g have constant sign on the interior of each of these intervals. Enumerate these intervals from the right, so that  $I_1$  is either  $(a_n, \infty)$  or  $(b_m, \infty)$ . On adjacent intervals, one of f, g has the same sign on both intervals, and one has different sign on the two intervals.

If f, g have leading coefficients with the same sign then they have the same sign on I<sub>1</sub>, and hence on all odd numbered intervals. Thus, the roots of  $\alpha f + \beta g$  lie in the even numbered intervals I<sub>2</sub>, I<sub>4</sub>,.... By the continuity of the roots as a function of  $\alpha$ ,  $\beta$ , there are the same number of roots of  $\alpha f + \beta g$  in each interval. By choosing  $\alpha = 1$  and  $\beta$  small, we see that each interval has at most one root. Thus, since  $\alpha f + \beta g$  has n roots, the number of even numbered intervals, |(n + m + 1)/2|, is at least n. This implies that n = m or n = m + 1.

If we assume that f, g have leading coefficients with opposite signs, then the roots lie in the odd intervals of which there are  $\lfloor (n + m + 2)/2 \rfloor$ . Consequently, n = m, n = m + 1 or n = m + 2.

If both endpoints of an interval are roots of f (say) then g is the same sign on the closed interval. If we choose  $\beta$  sufficiently small, then we can force f +  $\beta$ g to have the same sign on the interval, and hence there are no roots there. If we can choose  $\beta$  to be any sign, then choosing  $\beta$  large gives at least two roots of  $\alpha$ f +  $\beta$ g on the interval. Consequently, when  $\alpha$ ,  $\beta$  are unrestricted in sign, intervals have one endpoint a root of f and the other a root of g. This implies (4) and (5).

If deg(f) = deg(g), and their leading coefficients have the same sign then the n roots of  $\alpha f + \beta g$  lie in the n even intervals. By the above paragraph each interval has one endpoint a root of f and one a root of g. Choose k to have one root in each of the odd intervals.

If the leading coefficients differ in sign, then the the roots lie in the n + 1 odd intervals. Since  $I_1$  and  $I_{2n+1}$  each have only one root of f, g, and every other interval has one root of f and one root of g, choose k to have one root in each even interval.

If  $\deg(f) = \deg(g) + 2$  then there are n odd intervals and n - 1 even intervals, so all the roots are in the odd intervals. It follows that from the right we have a root of f, then a root of f and of g in some order, a root of f and of g in some order and so on until the leftmost root which is a root of f. We choose k to have roots in all the even intervals.

The case deg(f) = deg(g) + 1 is similar and omitted.

The only unfamiliar case is when the degree differs by two. The thick lines in Figure 1.6 shows where the roots of  $f + \beta g$  are located in case f has degree 4 and g degree 2, and  $\beta$  is positive. A similar discussion shows that if  $f - \alpha g \in \mathbf{P}$  for all positive  $\alpha$ , then f and g have a common interlacing.



Figure 1.6: Roots of f + g where the degrees of f, g differ by 2

**Corollary 1.36.** Assume that the monic polynomials  $f_1, \ldots f_n$  all have degree n, and have all real roots. The following are equivalent:

1. For all positive  $\alpha_1, \ldots, \alpha_n, \alpha_1 f_1 + \cdots + \alpha_n f_n$  has all real roots.

#### 2. $f_1, \ldots, f_n$ have a common interlacing.

*Proof.* By choosing all but two of the  $\alpha$ 's sufficiently small, we find from Proposition 1.35 that for all  $i \neq j$ ,  $f_i$ ,  $f_j$  has a common interlacing. It is easy to see that this implies that they all have a common interlacing. (See p. 80.)

If we have three polynomials whose linear combinations always have roots, then the third is a linear combination of the other two.

**Corollary 1.37.** Suppose that f, g, h are three distinct polynomials such that for all  $\alpha$ ,  $\beta$ ,  $\gamma$  the polynomial  $\alpha f + \beta g + \gamma h$  has all real roots. Then two of them interlace, and the third is a linear combination of the other two.

Otherwise stated, the largest vector space contained in *P* has degree 2.

*Proof.* If we set one of  $\alpha$ ,  $\beta$ ,  $\gamma$  to zero then by Proposition 1.35 we know that any two of f, g, h interlace. After renaming, we may assume that  $f \leq g \leq h$ . Fix an  $\alpha$  and consider the two polynomials  $k_{\alpha} = f + \alpha h$  and g. Since every linear combination of  $k_{\alpha}$  and g has all real roots, we know that  $k_{\alpha}$  and g interlace. Moreover,  $k_0 \leq g$  and  $k_{\alpha} \geq g$  when  $\alpha$  is large. There is an  $\alpha$  for which the interlacing switches from  $\leq$  to  $\geq$ ; at this value  $k_{\alpha}$  and g have the same roots. Thus, for this choice of  $\alpha$  the linear combination  $f + \alpha h$  is a multiple of g.

Next we consider the case of linear combinations where each coefficient is a polynomial. Notice that the order of interlacing is reversed.

**Lemma 1.38.** Suppose h, k are polynomials of the same degree with positive leading coefficients. If for every pair of polynomials  $f \leq g$  with positive leading coefficients the polynomial fh+gk has all real roots, then  $k \leq h$ . If k and h have different degrees then  $k \leftarrow h$ .

*Proof.* We may assume that k and h have no factors in common. We will only need to assume that the conclusion only holds for *linear* f, g. By continuity, fh + gk holds when f and g are multiples of one another. Choosing f = ax, g = bx shows that x(ah + bk) has all real roots for all positive a, b. From Proposition 1.35 we conclude that h, k have a common interlacing.

Next, choose f = a(x - r) and g = b(x - s) where r is larger than all roots of hk, and s is smaller than all the roots of hk. Since  $f \ll g$  we know from the hypotheses that all positive linear combinations of fh and gk lie in **P**, and hence fh and gk have a common interlacing. Combined with the fact that h and k have a common interlacing, it follows that  $k \ll h$ .

The proof of the second statement is similar and omitted.

#### **1.6 Basic topology of polynomials**

In this section we note a few facts about the topology of polynomials and of interlacing. The set V of all polynomials of degree n is a vector space of dimension n + 1, and is homeomorphic to  $\mathbb{R}^{n+1}$ . The set **P**(n) of all polynomials

of degree n in **P** is a subset of V, and so has a topological structure. We will determine the boundary and interior of P(n).

Since roots are continuous functions of the coefficients, if a polynomial f has all real roots and they are all distinct, then there is an open neighborhood of f in which all the polynomials have all real roots, all distinct. This is half of the next theorem:

#### Theorem 1.39.

- *The interior of* **P**(n) *consists of polynomials with all distinct roots.*
- *The boundary of* **P**(n) *consists of polynomials with repeated roots.*

*Proof.* Suppose that f has a repeated root, so that we can write  $f = (x - a)^2 g$  for some polynomial g in **P**. For any positive  $\epsilon$  the polynomial  $((x - a)^2 + \epsilon)g$  has two complex roots. Since  $\epsilon$  can be arbitrarily small, every neighborhood of f has a polynomial that is not in **P**. Thus, f is in the boundary of **P**(n). Since polynomials with distinct roots are in the interior of **P**(n), this establishes the theorem.

**P** satisfies a closure property.

**Lemma 1.40.** If f is a polynomial, and there are polynomials  $f_i \in P$  such that  $\lim f_i = f$  then  $f \in P$ .

*Proof.* Since the roots of a polynomial are continuous functions of the coefficients, if f had a complex root then some  $f_i$  would have a complex root as well. Thus, all roots of f are real, and  $f \in \mathbf{P}$ .

If we allow sequences of polynomials whose degree can increase without bound then we can get analytic functions such as  $e^x$  as the limits of polynomials in **P**. The proof of the lemma shows that such limits will never have complex roots. Such functions are considered in Chapter 5.

We can use Proposition 1.35 to show

#### **Corollary 1.41.** Limits preserve interlacing.

Suppose that  $f_1, f_2, ...$  and  $g_1, g_2, ...$  are sequences of polynomials with all real roots that converge to polynomials f and g respectively. If  $f_n$  and  $g_n$  interlace for n = 1, 2, ... then f and g interlace.

*Proof.* Since  $f_n$  and  $g_n$  interlace,  $f_n + \alpha g_n \in \mathbf{P}$  for all  $\alpha$ . Taking the limit, we see that  $f + \alpha g$  has all real roots for all  $\alpha$ . We apply Proposition 1.35 to conclude that f and g interlace.

The nature of interlacing can change under limits. For instance, as  $n \to \infty$  the interlacing

$$(1-\frac{x}{n})x(1+\frac{x}{n}) < (x-1)(x+1)$$
converges to

$$\mathbf{x} \ge (\mathbf{x} - 1)(\mathbf{x} + 1)$$

However, if the degrees don't change then the direction is preserved.

**Lemma 1.42.** Suppose that  $\lim f_i = f$ ,  $\lim g_i = g$  are all polynomials of degree n. If  $f_i \leq g_i$  for all i then  $f \leq g$ .

*Proof.* Write  $g_i = \alpha_i f_i + h_i$  where  $f_i \leq h_i$ ,  $h_i$  has positive leading coefficient and  $\alpha_i > 0$ . Clearly the limits lim  $h_i = h$  and lim  $\alpha_i = \alpha$  exist. Since  $g = \alpha f + h$ where  $f \leq h$  and h has non-negative leading coefficient we have  $f \leq g$ . 

An alternative argument is to consider the location of the roots.

#### 1.7 Linear transformations that preserve interlacing

In this section we establish some properties of a linear transformation that preserves interlacing. We begin by observing that if a linear transformation maps **P** to itself then it preserves interlacing. This is an important result, for it frees us from having to try to prove that a linear transformation preserves interlacing - all we need to do is to show that the linear transformation maps **P** to itself.

**Theorem 1.43.** If a linear transformation T maps **P** to itself then T preserves interlacing.

If we also assume that T preserves degree, then  $f \leq g$  implies  $Tf \leq Tg$ . In addition, *let*  $h \ll k$ .

- 1. If the leading coefficients of T(x),  $T(x^2)$ ,  $T(x^3)$ ,... all have the same sign, then  $Th \ll Tk$ .
- 2. If the leading coefficients of T(x),  $T(x^2)$ ,  $T(x^3)$ , ... have alternating signs, then  $Th \gg Tk$ .

If T maps P bijectively to P then all interlacings  $\underline{\ll}$  above can be replaced with  $\ll$  . *More generally, if* T *is a* 1 - 1 *open map then all*  $\underline{\ll}$  *and*  $\underline{\triangleleft}$  *can be replaced with*  $\ll$ and  $\triangleleft$ .

*Proof.* If f and g are in **P** then for all  $\alpha$  the polynomial f +  $\alpha$ g lies in **P**. Since T maps **P** to itself,  $T(f + \alpha g) = T(f) + \alpha T(g)$  is in **P**. We now apply Proposition 1.35 to conclude that Tf and Tg interlace.

Now assume T preserves degree. We may assume that h and k have positive leading coefficients. We write k = bh + m where b is non-negative,  $h \leq m$ , and m has positive leading coefficient. Since Tk = bTh + Tm and Th < Tm, the results (1) and (2) follow from Corollary 1.29.

It remains to show that if T maps P onto itself and preserves degree then T preserves strict interlacing. T is an onto map from the set of polynomials of degree n to itself. Choose disjoint opens sets  $f \in \mathcal{O}_1, g \in \mathcal{O}_2$  of polynomials so that  $h \in \mathcal{O}_1$  and  $k \in \mathcal{O}_2$  implies  $h \leq k$ . Then  $T(\mathcal{O}_1)$  and  $T(\mathcal{O}_2)$  are disjoint open sets whose elements interlace, so  $Tf \ll Tg$ . Note that this only used 1 - 1 and open, so the last assertion follows.

*Remark* 1.44. The two conditions for the signs of the leading coefficients of the sequence  $T(x), T(x^2), T(x^3), \ldots$  are the only possibilities. See Corollary 1.50 below.

*Remark* 1.45. If a transformation merely maps  $\mathbf{P}^{\text{pos}}$  to  $\mathbf{P}$ , then T does not necessarily preserve interlacing. We will see (Lemma 8.84) that the linear transformation  $T(x^n) = (-1)^{n(n+1)/2} x^n$  maps  $\mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}$ . However,

$$T(x + 1) = 1 - x$$
 has root (1)  

$$T(x + 1)^{2} = 1 - 2x - x^{2}$$
 has roots (-2.4, .4)  

$$T(x + 1)^{2} \leq T(x + 1)$$

**Corollary 1.46.** If T is a linear transformation such that  $T(xf) \leq T(f)$  for all  $f \in \mathbf{P}^{I}$  (where I is an interval containing 0) and maps polynomials with positive leading coefficients to polynomials with positive leading coefficients then T defines a linear transformation  $\mathbf{P}^{I} \longrightarrow \mathbf{P}$  that preserves interlacing.

*Proof.* We first show that if  $h_r(x - \alpha)h_r(x - \beta)h \in \mathbf{P}^I$  where  $\beta < \alpha$  then  $T(x - \alpha)h \leq T(x - \beta)h$ . By hypothesis we see that  $T(x - \alpha)h \leq Th$ , and so the conclusion follows from

$$T(x - \beta)h = T(x - \alpha)h + (\alpha - \beta)T(h)$$

Next, assume  $f \leq g$  are in  $\mathbf{P}^{I}$ , let r and s be the largest roots of f and g, and define h = f/(x - r). Since we have the interlacings  $(x - r)h \leq g$  and  $g \leq (x - s)h$  it follows from the remark above that we can find a chain of interlacing polynomials

$$(x-r)h \underline{\ll} \cdots \underline{\ll} g \underline{\ll} \cdots \underline{\ll} (x-s)h$$

where consecutive polynomials satisfy the conditions of the first paragraph. Thus

$$T(x-r)h\underline{\ll}\cdots\underline{\ll}T(g)\underline{\ll}\cdots\underline{\ll}T(x-s)h$$

Since  $(x - r)h \ll (x - s)h$  it follows that  $T(x - r)h \ll T(x - s)h$  and hence the sequence of polynomials is mutually interlacing. In particular,  $T(f) = T(x - \alpha)h \ll T(g)$ .

The next result is fundamental.

**Theorem 1.47.** *Differentiation preserves interlacing.* 

*Proof.* The linear transformation T(f) = f' maps **P** to itself, and therefore preserves interlacing. The derivative does not map **P**(n) onto **P**(n - 1). (see Chapter 7.7.11). However, it's easy to see that the derivative is an open map, so strict interlacing is preserved.

The converse to Theorem 1.47 does not  $hold^2$ .

The simplest linear transformations mapping **P** to itself are of the form  $x^i \mapsto a_i x^i$ . Such linear transformations are well understood, see § 5.15.2. Here we determine their image on **P**<sup>pos</sup>.

**Lemma 1.48.** Suppose the linear transformation  $T(x^i) = a_i x^i$  maps P to itself. Then

- (1) All non-zero terms of the sequence  $a_0, a_1, \ldots$  are consecutive.
- (2) Either all non-zero  $a_i$  have the same sign, or the signs alternate.
- (3) T maps  $P^{\text{pos}}$  to either  $P^{\text{pos}}$  or to  $P^{\text{alt}}$ .

*Proof.* If the non-zero  $a_i$ 's aren't consecutive and there are at least two consecutive zeros then there are r, s such that s-r > 2,  $a_r \neq 0$ ,  $a_{r+1} = \cdots = a_{s-1} = 0$ ,  $a_s \neq 0$ . In this case  $T(x^r(1+x)^{s-r}) = x^r(a_r + a_s x^{s-r})$ , and the latter polynomial is not in **P**. If there is an isolated zero, then s - r = 2, and so  $a_r + a_{r+2}x^2$  has all real roots. This implies that  $a_r$  and  $a_{r+2}$  have opposite signs.

Next,  $T(x^r(x^2 - 1)) = (a_{r+2} - a_r x^2)x^r$ , so if  $a_r$  and  $a_{r+2}$  are non-zero then they are the same sign. Consequently, there are no isolated zeros. Thus, the signs of the first two non-zero terms determine the signs of the rest of the terms, so we can conclude (1) and (2). If all the signs are the same then  $T(\mathbf{P}^{\text{pos}}) \subseteq \mathbf{P}^{\text{pos}}$ , and if they alternate then  $T(\mathbf{P}^{\text{pos}}) \subseteq \mathbf{P}^{\text{alt}}$ 

If a linear transformation maps  $\mathbf{P}$  to itself, then the leading coefficients of the images of  $x^i$  also determine a linear transformation that maps  $\mathbf{P}$  to itself.

**Lemma 1.49.** If  $T: P \longrightarrow P$  is a linear transformation that preserves degree, and the leading coefficient of  $T(x^i)$  is  $c_i$ , then the linear transformation  $S(x^i) = c_i x^i$  also maps P to itself.

*Proof.* Let  $T(x^i) = g_i(x)$ , and choose  $f = \sum_0^n a_i x^i \in \mathbf{P}$ . If we substitute  $\alpha x$  for x, and apply T, we find that  $\sum_0^n a_i \alpha^i g_i(x) \in \mathbf{P}$ . Next, substitute  $x/\alpha$  for x to find that  $\sum_0^n a_i \alpha^i g_i(x/\alpha) \in \mathbf{P}$ . Since  $g_i(x)$  is a polynomial of degree i we see that  $\lim_{\alpha \to 0} \alpha^i g_i(x/\alpha) = c_i x^i$ , and so

$$\lim_{\alpha\to 0}\sum_{i=0}^n a_i\alpha^i g_i(x/\alpha) = \sum a_i c_i x^i = S(f) \in \mathbf{P}.$$



<sup>&</sup>lt;sup>2</sup> If f = x(x-1)(x-2) and  $g = (x-.5)^2$  then  $f' \leq g'$  but it is not true that  $f \leq g$ .

**Corollary 1.50.** If  $T: P \longrightarrow P$  is a linear transformation that preserves degree then the leading coefficients of  $T(x^i)$  either all have the same sign, or their signs alternate.

*Proof.* Use Lemma 1.48 and Lemma 1.49.

#### **1.8** Linear combinations and determinants

Suppose that we have an interlacing square  $f \begin{pmatrix} g \\ k \end{pmatrix}$  k. There are close con-

nections between the determinant  $\begin{vmatrix} f & g \\ h & k \end{vmatrix}$  and the interlacings  $\alpha f + \beta g \lessdot \alpha h + \beta k$  for all  $\alpha, \beta \in \mathbb{R}$ .

**Lemma 1.51.** Assume that f, g, h, k are polynomials with positive leading coefficients whose degrees satisfy deg(f) = deg(g) + 1 = deg(h) + 1 = deg(k) + 2. If for all  $\alpha$ ,  $\beta$ , not both zero, we have that  $\alpha f + \beta g < \alpha h + \beta k$  then

2.  $\left| \begin{smallmatrix} f & g \\ h & k \end{smallmatrix} \right| < 0$  for all x.

Similar results hold if we replace " $\ll$ " by " $\ll$ ".

*Proof.* We know that  $h \leq k$  and  $f \leq g$  by Proposition 1.35, and the assumptions on degree. Setting  $\alpha = 0$  or  $\beta = 0$  shows that  $f \leq h$  and  $g \leq k$ . If the determinant is zero then there is an r such that f(r)k(r) = g(r)h(r).

There are two cases. First, assume that f(r) = 0. Since h and f have no roots in common, we know  $h(r) \neq 0$ . If we set  $\alpha = k(r)$ ,  $\beta = -h(r)$  then for this choice we get the contradiction that

$$(\alpha f + \beta g)(r) = 0 = (\alpha h + \beta k)(r).$$

The second case is  $f(r) \neq 0$ . The choices  $\alpha = -g(r)$ ,  $\beta = f(r)$  yield another contradiction.

If we evaluate the determinant at a root r of k we find  $h(r)g(r) \neq 0$  Thus k and h have no roots in common, and so  $h \leq k$ . Similarly,  $f \leq g$ .

Since the determinant is never zero, it always has the same sign. We determine the sign of the determinant by evaluating it at the largest root r of f. The result is -g(r)h(r) which is negative since f < g and f < h.

There is a converse.

**Lemma 1.52.** Suppose f, g, h, k are polynomials in P with positive leading coefficients. If they satisfy

1. f ≪ g

Π

- 2. h∢k
- 3. f < h
- 4.  $\begin{vmatrix} f & g \\ h & k \end{vmatrix}$  is never zero.

then  $f + \alpha g \lt h + \alpha k$  for all  $\alpha$ . A similar result holds if  $\lt$  is replaced by  $\ll$ .

*Proof.* Assumptions (1) and (2) imply that  $f + \alpha g$  and  $h + \alpha k$  have all real roots for all  $\alpha$  and that the degree of  $f + \alpha g$  is always one more than the degree of  $h + \alpha k$ . Assumption (3) implies that the conclusion is true for  $\alpha = 0$ . If  $(f + \alpha g)(r) = (h + \alpha k)(r)$  then eliminating  $\alpha$  shows that (fk - gh)(r) = 0. By assumption (4) the determinant is never zero, so as  $\alpha$  varies  $f + \alpha g$  and  $h + \alpha k$  never have a root in common. Consequently, at all times they must satisfy  $f + \alpha g < h + \alpha k$ . The case with  $\ll$  is similar.

**Example 1.53.** It is easy to construct examples of interlacing squares whose determinants are always negative. We will see more of these determinants in Chapter 8. If we apply Theorem 9.113 to the product

$$f(x, y, z) = (1 + x + y + z)(2 + x + 3y + z)(5 + x + 2y + 2z)$$

then we get the interlacing square

$$\begin{array}{rcl} (1+x)(2+x)(5+x) & \leqslant & 19+19\,x+4\,x^2 \\ 29+31\,x+6\,x^2 & \leqslant & 36+16x \end{array}$$

and the determinant is always negative. If we apply the theorem to the determinant

1 + x + y + z	0	0
0	2 + y + 2z	1
0	1	1 + 3y + z

we get an interlacing square where all interlacings are  $\ll$ :

and the determinant arising from this square is always negative.

Remark 1.54. If we differentiate the entries of the first square

$$M = \begin{pmatrix} (1+x)(2+x)(5+x) & 19+19x+4x^2\\ 29+31x+6x^2 & 36+16x \end{pmatrix}$$
$$M' = \begin{pmatrix} 3x^2+16x+17 & 8x+19\\ 12x+31 & 16 \end{pmatrix}$$

then we have that

- 1. |M| > 0 for all x.
- 2. |M'| > 0 for all x.
- 3.  $|M + \alpha M'| > 0$  for all x and  $\alpha \ge 0$ .

The last condition suggests that |M| and |M'| "positively interlace" in some sense. Indeed, M and M' arise from polynomials f and  $\frac{\partial f}{\partial x}$  that do interlace, as polynomials of three variables. The reason why we require  $\alpha$  to be non-negative is that  $M - \alpha M'$  can have negative coefficients, and the result we are using requires that all the coefficients are positive.

The following lemma arises naturally when considering the linear transformation  $f \mapsto f + \alpha f'$  for polynomials with complex coefficients (Corollary 20.38).

**Lemma 1.55.** If  $f \leq g$  (or  $f \ll g$ ) have positive leading coefficients then  $\begin{vmatrix} f & g \\ f' & g' \end{vmatrix} < 0$ . If  $f \leq g$  then the determinant is also negative except at common roots of f and g where it is zero. Thus, if  $f \leq g$  then  $f + \alpha f' \leq g + \alpha g'$  for all  $\alpha \in \mathbb{R}$ .

*Proof.* We give two proofs. Since  $f \leq g$  it follows from Corollary 1.29 that  $f + \alpha g$  has all distinct roots for all  $\alpha$ . Since differentiation preserves  $\leq$  we can apply Lemma 1.51 to the interlacing  $f + \alpha g \leq f' + \alpha g'$ .

If we don't use Lemma 1.51 then we can use partial fractions. If f < g we write  $g = \sum b_i \frac{f}{x - a_i}$  where  $f(x) = a \prod (x - a_i)$ , and all  $b_i$  are positive. Now

$$\frac{\mathrm{f}\mathrm{g}'-\mathrm{g}\mathrm{f}'}{\mathrm{f}^2} = \left(\frac{\mathrm{g}}{\mathrm{f}}\right)' = \left(\sum \frac{\mathrm{b}_{\mathrm{i}}}{\mathrm{x}-\mathrm{a}_{\mathrm{i}}}\right)' = -\sum \frac{\mathrm{b}_{\mathrm{i}}}{(\mathrm{x}-\mathrm{a}_{\mathrm{i}})^2} < 0$$

If x is not a root of f then  $\left| \begin{array}{c} f & g \\ f' & g' \end{array} \right| < 0$ . If x is a root of f then the determinant is equal to -f'(x)g(x) which is negative since f' and g interlace f.

If the interlacing is not strict, then write  $f = hf_0$  and  $g = hg_0$  where  $f_0 < g_0$ . Since  $\begin{vmatrix} f & g \\ f' & g' \end{vmatrix} = h^2 \begin{vmatrix} f_0 & g_0 \\ f'_0 & g'_0 \end{vmatrix}$  the result follows from the first part.

If  $f \ll g$  then there is a positive  $\alpha$  and polynomial h with positive leading coefficient such that  $g = \alpha f + h$  and f < h. The result now follows from the first case since

$$\begin{vmatrix} f & g \\ f' & g' \end{vmatrix} = \begin{vmatrix} f & \alpha f + h \\ f' & \alpha f' + h' \end{vmatrix} = \begin{vmatrix} f & h \\ f' & h' \end{vmatrix}$$

An immediate corollary (using  $f \leq f'$ ) is a well known property of derivatives called the Laguerre inequality.

**Corollary 1.56.** If f has all real roots, all distinct, then  $\left| \begin{array}{c} f & f' \\ f' & f'' \end{array} \right| < 0$ .

There is a converse to Lemma 1.55.

**Lemma 1.57.** If  $f \in P$ , the degree of g is at least one less than the degree the of f, and  $\begin{vmatrix} f & g \\ f' & g' \end{vmatrix} \neq 0$  for all x then f and g interlace.

*Proof.* It suffices to show that for all  $\alpha$  the polynomial  $f + \alpha g$  is in **P**. When  $\alpha = 0$  we know that  $f \in \mathbf{P}$ . If there is an  $\alpha$  such that  $f + \alpha g \notin \mathbf{P}$  then there is an  $\alpha$  for which  $f + \alpha g$  has a double root, r. In this case we have that

$$0 = f(r) + \alpha g(r)$$
$$0 = f'(r) + \alpha g'(r)$$

so the determinant  $\begin{vmatrix} f & g \\ f' & g' \end{vmatrix}$  is zero for this value of r.

If all linear combinations of f and g have all real roots, then the corollary tells us what direction they interlace.

**Corollary 1.58.** *If the following three conditions hold then*  $f \leq g$ :

- 1. f, g have the same degree.
- 2.  $f + \alpha g \in \mathbf{P}$  for all  $\alpha \in \mathbb{R}$ .
- 3.  $\begin{vmatrix} f & g \\ f' & g' \end{vmatrix} \leq 0$  for all x.

The next lemma allows us to conclude that two polynomials interlace from interlacing information, and knowledge of a determinant.

**Lemma 1.59.** Suppose that  $f \leftarrow g$ ,  $h \leftarrow k$  are polynomials with positive leading coefficients whose degrees satisfy  $deg(f) \ge deg(h)$  and  $deg(g) \ge deg(k)$  where all " $\leftarrow$ " are " $\ll$ " or all are " $\leq$ ". The following are equivalent



2. The determinant  $\begin{vmatrix} f & g \\ h & k \end{vmatrix}$  is negative at roots of fghk.

*Proof.* If r is a root of g then the determinant is f(r)k(r) which is negative by the interlacing assumptions. The argument is similar for the roots of g, h, k.

Conversely, if the determinant is negative at the i-th root r of g then we have that f(r)k(r) < 0. Since f sign interlaces g, so does k sign interlace g. Similarly we see that h sign interlaces f. The degree assumptions imply that these sign interlacings are actually interlacings.

A determinant condition gives information about the signs of the leading coefficients.

**Lemma 1.60.** Assume that  $f \leftarrow h \leftarrow k$ , and that there is a polynomial g such that  $\left| \begin{smallmatrix} f & g \\ h & k \end{smallmatrix} \right| < 0$ . Then the signs of the leading coefficients of f and k are the same.

*Proof.* Without loss of generality we may assume that the leading coefficient of h is positive. Suppose that r is the largest root of h, the leading coefficient of f is  $c_f$ , and the leading coefficient of k is  $c_k$ . Since the determinant is negative we know that f(r)k(r) is negative. Since  $f \leftarrow h$  the sign of f(r) is  $-sgn(c_f)$ , and similarly the sign of k(r) is  $sgn(c_k)$ . This implies that  $c_f$  and  $c_k$  have the same sign.

#### 1.9 Polynomials from determinants

We can construct polynomials from determinants. Although the constructions yield nothing new in one variable, they are an essential ingredient for the construction of polynomials in two variables. We emphasize the properties of principle submatrices that correspond to properties of  $f(x)/(x - r_i)$  where  $f(x) = \prod (x - r_k)$ .

We consider determinants of matrices of the form xI + C where C is a symmetric matrix and I is the identity matrix. More generally we consider xD + C where D is a positive definite matrix, but these are really no different. Since D has a positive definite square root,

$$|\mathbf{x}\mathbf{D} + \mathbf{C}| = |\mathbf{D}^{-1}| |\mathbf{x}\mathbf{I} + \mathbf{D}^{-1/2}\mathbf{C}\mathbf{D}^{-1/2}|$$

and  $D^{-1/2}CD^{-1/2}$  is symmetric.

The determinants |xI + C| generalize the product representation of a polynomial. If  $f(x) = (x-r_1)\cdots(x-r_n)$  then f(x) = |xI+C| where C is the diagonal matrix whose diagonal is  $\{-r_1, \ldots, -r_n\}$ .

One important well known fact is that |xI + C| has all real roots. Indeed, since C is symmetric, its eigenvalues are all real, and the roots of |xI + C| are the negatives of these eigenvalues.

If r is a root of f then the analog of f(x)/(x - r) is a principle submatrix. If M is a d by d matrix, and  $\alpha \subset \{1, 2, ..., d\}$  then  $M\{\alpha\}$  is the (sub)matrix of M formed by the entries in the rows and columns indexed by  $\alpha$ . We let M[i] denote the submatrix resulting from the deletion of the ith row and column of M. Such a submatrix is called a *principle submatrix*.

The maximal factor  $f(x)/(x - r_i)$  interlaces f(x). The analogous statement is also true for principle submatrices. The following is well known, but the proof below is elementary.

**Theorem 1.61.** If A is an n by n Hermitian matrix, and B is any principle submatrix of order n - 1 then their characteristic polynomials interlace.

*Proof.* Choose  $\alpha \in \mathbb{R}$ , partition  $A = \begin{pmatrix} B & c \\ \hline c^* & d \end{pmatrix}$  and consider the following equation that follows from linearity of the determinant:

$$\left| \begin{array}{c|c} B-xI & c \\ \hline c^* & d-x+\alpha \end{array} \right| = \left| \begin{array}{c|c} B-xI & c \\ \hline c^* & d-x \end{array} \right| + \left| \begin{array}{c|c} B-xI & c \\ \hline 0 & \alpha \end{array} \right|$$

Since the matrix on the left hand side is the characteristic polynomial of a Hermitian matrix,  $|A - xI| + \alpha |B - xI|$  has all real roots for any  $\alpha$ , and hence the eigenvalues interlace.

The derivative of  $f(x) = \prod_1^n (x - r_n)$  can be written as

$$\frac{\mathrm{d}}{\mathrm{d}x} f(x) = \frac{f(x)}{x - r_1} + \dots + \frac{f(x)}{x - r_n}$$

There is a similar formula for determinants. Suppose that C is a n by n matrix.

$$\frac{d}{dx}|xI + C| = |M[1]| + \dots + |M[n]|$$
(1.9.1)

The factors of a polynomial with all distinct roots have all distinct roots. The same holds for principle submatrices. This is immediate since the roots of the submatrix interlace the roots of the original matrix.

Finally, we have some ways in which principle submatrices don't generalize properties of maximal factors.

**Example 1.62.** If the roots of a polynomial  $f(x) = \prod_{i=1}^{n} (x - r_i)$  are distinct, then the polynomials  $f(x)/(x - r_i)$  are linearly independent, and span the space of all polynomials of degree n-1. The matrix  $\begin{pmatrix} x & 1 \\ 1 & x \end{pmatrix}$  has two distinct roots  $\{1, -1\}$ , yet the two principle submatrices are equal.

**Example 1.63.** If the roots of f(x) are ordered  $r_1 \leq \cdots \leq r_n$  then the factors are mutually interlacing:

$$\frac{f(x)}{x-r_1} \leq \frac{f(x)}{x-r_2} \leq \cdots \leq \frac{f(x)}{x-r_n}$$

Although the n principle submatrices have characteristic polynomials that interlace the characteristic polynomial of M, it is not the case that these submatrices even have interlacing characteristic polynomials, much less mutually interlacing ones. They only have a common interlacing.

We can interpret (1.3.3) in terms of determinants. This representation of g(x) will prove useful when we consider polynomials in two variables. Since the  $c_i$  in (1.3.3) are negative the vector

$$\boldsymbol{v} = (\sqrt{-c_1}, \dots, \sqrt{-c_n})$$

is real. We assume that f(x) is monic so that  $f(x) = \prod (x - a_i)$ , and define the diagonal matrix A whose diagonal entries are  $a_1, \ldots, a_n$ . Let I be the n by n identity matrix, and |W| the determinant of a matrix W. Expanding by the first row shows that

$$\begin{vmatrix} x+c & v \\ v^{t} & xI-A \end{vmatrix} = \begin{vmatrix} x+c & \sqrt{-c_{1}} & \sqrt{-c_{2}} & \dots & \sqrt{-c_{n}} \\ \sqrt{-c_{1}} & x-a_{1} & 0 & \dots & 0 \\ \sqrt{-c_{2}} & 0 & x-a_{2} & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \sqrt{-c_{n}} & 0 & 0 & \dots & x-a_{n} \end{vmatrix}$$
(1.9.2)
$$= (x+c)f(x) + c_{1}\frac{f(x)}{x-a_{1}} + \dots + c_{n}\frac{f(x)}{x-a_{n}}$$
$$= g(x)$$

Equation (1.9.2) shows that  $g \in \mathbf{P}$  since g is the characteristic polynomial of the symmetric matrix  $\begin{pmatrix} c & \nu \\ \nu^t & -A \end{pmatrix}$ .

### 1.10 A relation weaker than interlacing

The relation  $\blacktriangleleft$  is weaker than interlacing, yet still has some of the properties of interlacing. In particular, it is sometimes preserved by differentiation, and there are inequalities for 2 by 2 determinants of coefficients.

Consider some simple examples and properties.

- 1.  $(x+1)^n \triangleleft x^n$
- 2. If  $h \triangleleft g$  and  $g \triangleleft f$  then  $h \triangleleft f$ .
- 3. If  $g \blacktriangleleft f$  and  $s \leqslant r$  then  $(x s) g \blacktriangleleft (x r) f$ .
- 4. If  $g \blacktriangleleft f$  and  $h \in \mathbf{P}$  then  $hg \blacktriangleleft hf$ .
- 5. If  $g \blacktriangleleft f$  and  $f \in \mathbf{P}^{pos}$  then  $g \blacktriangleleft x^k f$  for any positive integer k.
- 6. If  $f \leq g$  then g < f.
- 7. If  $f \leq g$  then we generally do not have  $g \triangleleft f$  since the smallest root belongs to f.
- $g \triangleleft f$  determines a chain of interlacings.

**Lemma 1.64.** If  $g \triangleleft f$  have the same degree then there are  $h_1, \ldots, h_k$  such that

If f and g have different degrees then we can write f = hk where  $g \blacktriangleleft h$ , and h has the same degree as g.

*Proof.* Let  $g = \prod_{i=1}^{m} (x - s_i)$  and  $f = \prod_{i=1}^{n} (x - r_i)$ . The following interlacings give the desired sequence of polynomials.

$$\begin{split} g &\geq \left(\prod_{2}^{m}(x-s_{\mathfrak{i}})\right) \left(\prod_{1}^{1}(x-r_{\mathfrak{i}})\right) \geq \left(\prod_{3}^{m}(x-s_{\mathfrak{i}})\right) \left(\prod_{1}^{2}(x-r_{\mathfrak{i}})\right) \geq \cdots \\ &\geq \left(\prod_{m}^{m}(x-s_{\mathfrak{i}})\right) \left(\prod_{1}^{m-1}(x-r_{\mathfrak{i}})\right) \geq \prod_{1}^{m}(x-r_{\mathfrak{i}}) = f \end{split}$$

For the second part, we take k to be the polynomial whose roots are the n - m largest roots of f. 

Differentiation does not preserve ◀ if the degrees are different. For example, if f = (x + 1)(x + 2) and g = x(x + 1)(x + 2), then  $f \blacktriangleleft g$  but  $f' \blacktriangle g'$  fails. We do have

#### **Lemma 1.65.** If $g \triangleleft f$ have the same degree then $g' \triangleleft f'$ .

.

*Proof.* The result follows by differentiating the chain of interlacings in the previous lemma. 

**Lemma 1.66.** Suppose that  $g \blacktriangleleft f$ , f and g have all positive coefficients, and f = $\sum_{1}^{n} a_{i} x^{i}$ ,  $g = \sum_{1}^{m} b_{i} x^{i}$ . Then

$$\begin{vmatrix} b_i & b_j \\ a_i & a_j \end{vmatrix} \geqslant 0 \quad \textit{for } 1 \leqslant i < j \leqslant n$$

Proof. We first assume that f and g have the same degree. In order to show that  $b_i a_j \ge b_j a_i$ , we note that this is equivalent to  $\frac{b_i}{a_i} \ge \frac{b_j}{a_j}$ . Thus it suffices to take j = i + 1 since then

$$\frac{b_i}{a_i} \geqslant \frac{b_{i+1}}{a_{i+1}} \geqslant \cdots \geqslant \frac{b_j}{a_j}.$$

Since ◀ is preserved by differentiation, we can differentiate i times, and so assume that i = 0 and j = 1. Write  $f = \prod_{1}^{m} (x + r_i)$ ,  $g = \prod_{1}^{m} (x + s_i)$  where  $0 \leqslant r_{\mathfrak{i}} \leqslant s_{\mathfrak{i}}$  for  $1 \leqslant \mathfrak{i} \leqslant \mathfrak{m}.$  Then

$$\begin{vmatrix} b_0 & b_1 \\ a_0 & a_1 \end{vmatrix} = \begin{vmatrix} \prod_1^m s_i & \prod_1^m \left( \sum_1^m \frac{1}{s_i} \right) \\ \prod_1^m r_i & \prod_1^m \left( \sum_1^m \frac{1}{r_i} \right) \end{vmatrix} = \left( \prod s_i \right) \left( \prod r_i \right) \sum_1^m \left( \frac{1}{r_i} - \frac{1}{s_i} \right)$$

Now assume that the degrees are different. Note that if  $p_1 = \sum \alpha_i x^i$ ,  $p_2 = \sum \beta_i x^i$  and  $p_3 = \sum \gamma_i x^i$  then

$$\begin{vmatrix} \beta_{i} & \beta_{j} \\ \alpha_{i} & \alpha_{j} \end{vmatrix} > 0 \quad \text{and} \quad \begin{vmatrix} \gamma_{i} & \gamma_{j} \\ \beta_{i} & \beta_{j} \end{vmatrix} > 0 \quad \Longrightarrow \quad \begin{vmatrix} \alpha_{i} & \alpha_{j} \\ \beta_{i} & \beta_{j} \end{vmatrix} > 0$$

By Lemma 1.64 it suffices to prove the lemma for g and  $(x + \alpha)g$  where  $\alpha > 0$ . If  $g = \sum b_i x^i$  then the determinant is

$$\begin{vmatrix} b_i & b_j \\ \alpha b_i + b_{i-1} & \alpha b_j + b_{j-1} \end{vmatrix} = \begin{vmatrix} b_i & b_j \\ b_{i-1} & b_{j-1} \end{vmatrix}$$

As above, it suffices to show that  $\begin{vmatrix} b_i & b_{i+1} \\ b_{i-1} & b_i \end{vmatrix} \ge 0$ , but this is Newton's inequality (Theorem 4.8).

A matrix is TP<sub>2</sub> is all 2 by 2 submatrices have non-negative determinant. **Lemma 1.67.** If  $f_1 \triangleleft f_2 \triangleleft \cdots \triangleleft f_k$  are in  $P^{pos}$  then the matrix of coefficients of

$$\begin{pmatrix} f_1 \\ f_2 \\ \vdots \\ f_k \end{pmatrix}$$

is  $TP_2$ .

*Proof.* Since transitivity of  $\blacktriangleleft$  implies that  $f_i \blacktriangleleft f_j$  for i < j the result follows from Lemma 1.66.

Here's a typical corollary:

**Corollary 1.68.** If  $g \triangleleft f$  are in  $P^{pos}$  then

g						$b_0$	$\mathfrak{b}_1$	$\mathfrak{b}_2$	•••				
	g					0	$b_0$	$\mathfrak{b}_1$	$b_2$	• • •			
		g				0	0	$\mathfrak{b}_0$	$\mathfrak{b}_1$	$b_2$	•••		
		f			=	0	0	$a_0$	$\mathfrak{a}_1$	a <sub>2</sub>	•••		
			f			0	0	0	$a_0$	$a_1$	$a_2$	• • •	
				f		0	0	0	0	$a_0$	$\mathfrak{a}_1$	a <sub>2</sub>	

is  $TP_2$ .

*Proof.* The rows are the coefficients of the sequence of polynomials that satisfy the relations

$$g \triangleleft xg \triangleleft x^2g \triangleleft x^2f \triangleleft x^3f \triangleleft x^4f$$

The result now follows from Lemma 1.67.

**Example 1.69.** Here is a example that uses  $\blacktriangleleft$  to get an inequality for the location of the roots of the derivative. Suppose that  $f = \prod_{1}^{n} (x - r_k)$ . Since  $f \blacktriangleleft (x - r_k)^k (x - r_n)^{n-k}$  we differentiate to find that

$$\mathsf{f}' \blacktriangleleft (\mathsf{x} - \mathsf{r}_k)^{k-1} \, (\mathsf{x} - \mathsf{r}_n)^{n-k-1} \, (\mathsf{n} \mathsf{x} - \mathsf{k} \mathsf{r}_n - (\mathsf{n} - \mathsf{k}) \mathsf{r}_k)$$

It follows that the kth root of f' lies in the interval

$$[\mathbf{r}_k,\mathbf{r}_k+\frac{\mathbf{k}}{\mathbf{n}}(\mathbf{r}_n-\mathbf{r}_k)].$$

#### **1.11** Polytopes and polynomials

A polynomial  $a_0 + a_1x + \cdots + a_nx^n$  corresponds to the point  $(a_0, \ldots, a_n)$  in  $\mathbb{R}^{n+1}$ . Thus, we can relate geometric properties to algebraic properties. In particular, we study two questions:

- 1. When does a polytope 𝒫 in ℝ<sup>n+1</sup> have the property that all points in 𝒫 determine polynomials with all real roots?
- 2. When do polytopes P and P' have the property that every polynomial determined by points in P interlaces every polynomial determined by points in P'?

There are simple answers when the polytopes are products of intervals, and the proofs are based on sign interlacing. If we have vectors  $\mathbf{a} = (a_i)$  and  $\mathbf{b} = (b_i)$  then we define

$$poly(\mathbf{a}) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + a_4 x^4 + \cdots$$
$$poly(\mathbf{a}, \mathbf{b}) = a_0 + b_1 x + a_2 x^2 + b_3 x^3 + a_4 x^4 + \cdots$$

**Lemma 1.70.** Suppose that  $0 \le a \le b$ , and that there exists an f satisfying the interlacing  $f \le poly(a, b)$  and  $f \le poly(b, a)$ . Then, for all c satisfying  $a \le c \le b$  we have that  $f \le poly(c)$ .

*Proof.* If the roots of f are  $r_1 \leq \cdots \leq r_n$ , then  $\mathsf{poly}(\mathbf{a}, \mathbf{b})$  and  $\mathsf{poly}(\mathbf{b}, \mathbf{a})$  sign interlace f, and so the sign of either of them at  $r_i$  is  $(-1)^{n+i}$ . The roots are negative since all the coefficients are positive, so the conclusion will follow from

$$\operatorname{poly}(\mathbf{a},\mathbf{b})(-\alpha) \leqslant \operatorname{poly}(\mathbf{c})(-\alpha) \leqslant \operatorname{poly}(\mathbf{b},\mathbf{a})(-\alpha) \quad \text{for } \alpha \ge 0$$

Expanded, this is

$$\begin{aligned} a_0 - b_1 \,\alpha + a_2 \,\alpha^2 - b_3 \,\alpha^3 + a_4 \,\alpha^4 \cdots \leqslant c_0 - c_1 \,\alpha + c_2 \,\alpha^2 - c_3 \,\alpha^3 + c_4 \,\alpha^4 \cdots \\ \leqslant b_0 - a_1 \,\alpha + b_2 \,\alpha^2 - a_3 \,\alpha^3 + b_4 \,\alpha^4 \cdots \end{aligned}$$

which follows from

$$\begin{split} a_{2i} \, \alpha^{2i} \leqslant c_{2i} \, \alpha^{2i} \leqslant b_{2i} \, \alpha^{2i} \\ -b_{2i+1} \, \alpha^{2i+1} \leqslant -c_{2i+1} \, \alpha^{2i+1} \leqslant -a_{2i+1} \, \alpha^{2i+1}. \end{split}$$

Figure 1.7 shows the configuration of poly(a, b), poly(b, a) and poly(c).

Next, we show that we don't need to assume common interlacing.

**Lemma 1.71.** Suppose that  $0 \le a \le b$ , and that both poly(a, b) and poly(b, a) have all real roots. Then poly(a, b) and poly(b, a) have a common interlacing.



Figure 1.7: Graphs of interval polynomials

*Proof.* We prove this by induction on the degree n. We assume that  $\mathbf{a} \neq \mathbf{b}$ . From the proof of Lemma 1.70 we know that the graph of  $\mathsf{poly}(\mathbf{b}, \mathbf{a})$  lies strictly above the graph of  $\mathsf{poly}(\mathbf{a}, \mathbf{b})$  for negative x. When n = 2 there are two possibilities for the graphs of  $\mathsf{poly}(\mathbf{a}, \mathbf{b})$  and  $\mathsf{poly}(\mathbf{b}, \mathbf{a})$  (Figure 1.8)



Figure 1.8: The possibilities for a quadratic

Both of the roots of one quadratic can not be to either the left or the right of the roots of the other quadratic, for then there is an intersection. Thus, they must be arranged as in the right hand picture, and there is clearly a common interlacing.

We now consider the general case. The key point is that we can differentiate. If we define

$$\mathbf{a}' = (a_1, 2a_2, 3a_3, \cdots, na_{n-1})$$
  $\mathbf{b}' = (b_1, 2b_2, 3a_3, \cdots, na_{n-1})$ 

then we have the relations

$$\operatorname{poly}(\mathbf{a},\mathbf{b})' = \operatorname{poly}(\mathbf{b}',\mathbf{a}') \qquad \operatorname{poly}(\mathbf{b},\mathbf{a})' = \operatorname{poly}(\mathbf{a}',\mathbf{b}').$$

The inductive hypothesis implies that poly(a, b)' and poly(b, a)' have a common interlacing.

Suppose that poly(a, b) has roots  $r_1 \cdots < r_n$ . Both poly(a, b) and poly(b, a) are positive on the intervals

$$(r_{n-2}, r_{n-1}), (r_{n-4}, r_{n-3}), \ldots$$

and so the only place that poly(b, a) can have roots is in the intervals

$$(r_{n-1}, r_n), (r_{n-3}, r_{n-2}) \dots$$

We now show that none of these intervals can have more than two roots of poly(**b**, **a**). Figure 1.9 is an example of what can not happen. We observe the



Figure 1.9: An impossible configuration

simple property of polynomials g, h with a common interlacing:

For any  $x \in \mathbb{R}$  the number of roots of g greater than x and the number of roots of h greater than x differ by at most one.

We now see that Figure 1.9 is impossible, since the two largest roots of poly(a, b)' are greater than all the roots of poly(b, a)'. Moreover, if there were two roots of poly(b, a) in the rightmost solid interval labeled Q, then there are 5 roots of poly(b, a)' to the right of the point p, but only 3 roots of poly(a, b)' to the right.

Continuing to the left, suppose up to some point that we find all intervals with poly(a, b) negative have two roots of poly(b, a). The next interval to the left can't have 0 roots, for the same reason as above. Also, it can't have more than two roots, for then there would be too many roots of poly(b, a)'.

Combining the last two lemmas we get

**Proposition 1.72.** Suppose that  $0 \leq a \leq b$ , and that both poly(a, b) and poly(b, a) have all real roots. Then, for all c satisfying  $a \leq c \leq b$  we have that  $poly(c) \in P$ .

*Remark* 1.73. If we drop the hypotheses that that the coefficients are positive, then there might be no common interlacing, and the conclusion fails. If we take  $\mathbf{a} = (1, -2, 1)$ ,  $\mathbf{b} = (1, 2, 1)$  and  $\mathbf{c} = (1, 0, 1)$  then  $\mathbf{a} \leq \mathbf{c} \leq \mathbf{b}$ . However, there is no common interlacing, and

$poly(a, b) = 1 + 2x + x^2$	$\in \mathbf{P}$
$poly(\mathbf{b},\mathbf{a}) = 1 - 2\mathbf{x} + \mathbf{x}^2$	$\in \mathbf{P}$
$poly(\mathbf{c}) = 1 + x^2$	$ ot\in \mathbf{P}$

**Corollary 1.74.** If  $0 \le a \le b$ , poly(a, b) and poly(b, a) have all real roots then all points in the product of intervals

$$\mathbf{a} \times \mathbf{b} = (\mathfrak{a}_0, \mathfrak{b}_0) \times (\mathfrak{a}_1, \mathfrak{b}_1) \times \cdots \times (\mathfrak{a}_n, \mathfrak{b}_n)$$

determine polynomials in **P**.

**Lemma 1.75.** Suppose that  $0 \leq \mathbf{a} \leq \mathbf{c} \leq \mathbf{b}$  and  $0 \leq \mathbf{a}' \leq \mathbf{c}' \leq \mathbf{b}'$ . If

$$\left. \begin{array}{c} \text{poly}(a,b) \\ \text{poly}(b,a) \end{array} \right\} \hspace{0.2cm} \leq \hspace{0.2cm} \left\{ \begin{array}{c} \text{poly}(a',b') \\ \text{poly}(b',a') \end{array} \right.$$

then  $poly(\mathbf{c}) \leq poly(\mathbf{c}')$ .

Proof. Two applications of Lemma 1.70 show that

$$poly(a, b) \leq poly(c')$$
  $poly(b, a) \leq poly(c')$ 

The proof of the lemma holds if we replace  $\leq$  by  $\geq$ , so we conclude that  $poly(c) \leq poly(c')$ .

The Lemma shows that all points in the box  $\mathbf{a} \times \mathbf{b}$  determine polynomials that interlace all the polynomials corresponding to points in  $\mathbf{a}' \times \mathbf{b}'$ .

#### 1.12 The Faà di Bruno problem

The Faà di Bruno formula is an expression for the m'th derivative of a composition. We conjecture that certain polynomials associated with this formula have all real roots. We are able to establish this in the first few cases, for  $e^x$ , and for  $x^d$ . If f and g are polynomials then the Faà di Bruno formula[97] is

$$\frac{d^{\mathfrak{m}}}{dx^{\mathfrak{m}}}(f(g(x)) = \sum \frac{m!}{b_{1}!\cdots b_{\mathfrak{m}}!}f^{(k)}(g(x))\left(\frac{g'(x)}{1!}\right)^{b_{1}}\left(\frac{g^{(2)}(x)}{2!}\right)^{b_{2}}\cdots\left(\frac{g^{(\mathfrak{m})}(x)}{\mathfrak{m}!}\right)^{b_{\mathfrak{m}}}$$

where the sum is over all  $b_1 + 2b_2 + \cdots + mb_m = n$  and  $b_1 + \cdots + b_m = k$ .

For example, the first few are

m	
0	f(g(x))
1	f'(g(x)) g'(x)
2	$f''(g(x)) g'(x)^2 + f'(g(x)) g''(x)$
3	$f^{(3)}(g(x)) g'(x)^3 + 3f''(g(x)) g''(x)g'(x) + f'(g(x)) g^{(3)}(x)$
4	$f^{(4)}(g(x)) g'(x)^4 + 6g''(x) f^{(3)}(g(x)) g'(x)^2 +$
	$4f''(g(x)) g^{(3)}(x) g'(x) + 3f''(g(x)) g''(x)^2 + f'(g(x)) g^{(4)}(x)$

We can simplify these expressions by writing

$$\frac{d^{\mathfrak{m}}}{dx^{\mathfrak{m}}}(f(\mathfrak{g}(x)) = \sum f^{(k)}(\mathfrak{g}(x))A_{\mathfrak{m},k}(x)$$

and defining

$$F_{\mathfrak{m}}(\mathbf{x},\mathbf{y}) = \sum \mathbf{y}^{k} A_{\mathfrak{m},k}(\mathbf{x})$$

Here are the first few values of  $F_m(x, y)$ 

m	$F_m(x,y)$
0	1
1	g'(x) y
2	$g'(x)^2 y^2 + g''(x) y$
3	$g'(x)^3 y^3 + 3g''(x)g'(x)y^2 + g^{(3)}(x)y$
4	$g'(x)^4 y^4 + 6g''(x)g'(x)^2 y^3 + [4g^{(3)}(x)g'(x) + 3g''(x)^2]y^2 + g^{(4)}(x)y$

Note that  $F_m$  only depends on g, and not on f. If  $\alpha$  and m are fixed the transformation  $g \mapsto F_m(\alpha, y)$  is *not* a linear transformation. It follows from [95] that  $F_m$  satisfies the recurrence

$$F_{m+1}(x,y) = \frac{d}{dx}F_m(x,y) + g'(x)yF_m(x,y)$$
(1.12.1)

If we iterate this we get the simple formula

$$F_{\mathfrak{m}} = \left(\mathsf{D}_{x} + \mathfrak{g}'(x)\mathfrak{y}\right)^{\mathfrak{m}}(1)$$

We can recover  $\frac{d^m}{dx^m}(f(g(x)) \text{ from } F_m \text{ as follows:}$ 

$$\frac{d^{\mathfrak{m}}}{dx^{\mathfrak{m}}}(f(\mathfrak{g}(x)) = \left(\sum \mathsf{D}_{y}^{k} A_{\mathfrak{m},k}(x)\right) \left(\sum f^{(k)}(\mathfrak{g}(x)) \frac{y^{k}}{k!}\right)$$

Note that the term on the left is  $F_m(x, D_y)$  and the term on the right is the Taylor series of f(g(x) + y).

**Conjecture 1.76.** *Suppose that*  $g \in \widehat{P}$ *. For any*  $\alpha \in \mathbb{R}$  *and*  $\mathfrak{m} = 0, 1, ...,$ 

- 1.  $F_m(\alpha, y)$  has all real roots.
- 2.  $F_{m+1}(\alpha, y) \leq F_m(\alpha, y)$ .

We can verify this in the first few cases.

**m=1** We need to show that  $g'(\alpha)y \in \mathbf{P}$ , which is trivial.

**m=2** This is again trivial, since

$$F_2(\alpha, y) = y(g''(\alpha) + g(\alpha)^2 y$$

m=3 Factoring out y we have a quadratic whose discriminant is

$$(3 g''(\alpha) g'(\alpha))^2 - 4(g'(\alpha))^3 g'''(\alpha) = g'(\alpha)^2 [9g''(\alpha)^2 - 4g'(\alpha)g'''(\alpha)]$$

and this is non-negative by Newton's inequalities.

It takes a bit more work to verify interlacing  $F_3 \leq F_2$ . Factoring out y we need to show that for all  $\alpha$ 

$$(g')^{3}y^{2} + 3g''g'y + g''' \leq (g')^{2}y + g''$$

If we substitute the root of the right hand side into the left hand side and multiply by  $(g')^2$  we get

$$g'(g'')^2 - 3(g'')^2g' + g'''(g')^2 = g'(g'g''' - 2(g'')^2)$$

By Newton's inequalities this has sign opposite from the leading coefficient of  $F_3$ , so  $F_3 \leq F_2$ .

The conjecture holds for  $g(x) = e^x$ .

**Lemma 1.77.** If  $g = e^x$  then Conjecture 1.76 holds.

*Proof.* The key fact, which follows from [97], is that  $F_m$  has a simple form if  $g = e^x$ . That is,

$$F_{\mathfrak{m}}(\mathbf{x},\mathbf{y}) = B_{\mathfrak{m}}(e^{\mathbf{x}}\mathbf{y})$$

where  $B_m$  is the Bell polynomial. We know that  $B_m(x) \leq B_{m-1}(x)$ , and therefore for all  $\alpha$  we have that  $B_m(\alpha x) \leq B_{m-1}(\alpha x)$ .

The conjecture also holds for x<sup>d</sup>.

**Lemma 1.78.** If  $g = x^d$  then Conjecture 1.76 holds.

*Proof.* To prove the lemma it suffices to show that there are polynomials  $\mathsf{H}_n \in \mathbf{P}$  satisfying

$$x^{n}F_{n}(x,y) = H_{n}(x^{d}y)$$

and  $H_n \leq H_{n-1}$ . For example, if d = 3 we compute using (1.12.1) that

n	$x^{n}F_{n}(x,y)$	$H_n(z)$
0	1	1
1	$x(3x^2y)$	3z
2	$x^2(9y^2x^4 + 6yx)$	$9z^2 + 6z$
3	$x^{3}(27y^{3}x^{6}+54y^{2}x^{3}+6y)$	$27z^3 + 54z^2 + 6z$
4	$x^4(81y^4x^8 + 324y^3x^5 + 180y^2x^2)\\$	$81z^4 + 324z^3 + 180z^2$

We define  $H_m$  by the recursion

$$H_0 = 1 H_{m+1} = -mH_m + dz (H_m + H'_m)$$
(1.12.2)

We will prove that  $F_m = H_m(x^d y)$  by induction, and it holds for m = 0.

$$\begin{split} x^{m+1} F_m &= x^{m+1} (F'_m + dx^{d-1} y F_m) \\ &= x^{m+1} F'_m + dx^{d+m} y F_m \\ H_{m+1} (x^d y) &= -m H_m (x^d y) + dx^d y (H_m (x^d y) + H'(z) [x^d y]) \\ &= -m H_m (x^d y) + dx^d y H_m (x^d y) + x H'_m (x^d y) \\ &= -m x^m F_m + dx^d y x^m F_m + x (x^m F_m)' \\ &= dx^d y x^m F_m + x^{m+1} F'_m \end{split}$$

Since  $F_m$  has all positive coefficients  $H_m$  has all positive coefficients, and by induction  $H_m \in \mathbf{P}^{pos}$ . The recursion (1.12.2) shows that  $H_m \leq H_{m-1}$ .

#### 1.13 Root Trajectories

If we are given polynomials  $p(x, \alpha)$  where  $\alpha$  is a parameter, then we can investigate the movement of the roots of  $p(x, \alpha)$  as  $\alpha$  varies. There is a large body of literature in the case that  $p(x, \alpha)$  is an orthogonal polynomial [50, 128]. We limit ourselves to two simple cases. We first look at the effect of small perturbations of the roots of a polynomial on the roots of the derivative, and then consider the root trajectories of  $f + \alpha g$  where  $f \leftarrow g$ . See Figure 20.1 for trajectories of  $f + \iota t g$  in  $\mathbb{C}$ .

**Lemma 1.79.** *If*  $f \in P$ , and we increase the roots of f then the roots of f' also increase. [4]

*Proof.* Without loss of generality we may assume that the roots of f are distinct. If g is the polynomial resulting from increasing the roots of f by a sufficiently small amount, then  $g \ll f$ . Consequently  $g' \ll f'$ , so each root of g' is larger than the corresponding root of f'.

**Lemma 1.80.** If  $f \leftarrow g$  have positive leading coefficients then the roots of  $f + \alpha g$  decrease as  $\alpha$  increases.

*Proof.* We'll give three proofs! The simplest proof requires results from Chapter 3 (Corollary 3.8): if  $\alpha > \beta$  then the matrix  $\begin{pmatrix} 1 & \beta \\ \alpha & 1 \end{pmatrix}$  preserves interlacing, and hence  $f + \beta g \leq f + \alpha g$ .

Next, a proof that uses the product representation of f. We assume that f, g have no roots in common. Since  $f \leftarrow g$  we can write

$$g = \beta f + \sum_{i} b_{i} \frac{f}{x - a_{i}}$$

where  $\beta \ge 0$ , the  $b_i$  are non-negative, and the  $a_i$  are the roots of f. A root r of  $f + \alpha g = 0$  satisfies

$$0 = f(r) \left( 1 + \alpha \beta + \alpha \sum_{i} b_{i} \frac{1}{r - a_{i}} \right)$$

Since  $f(r) \neq 0$ , we remove the factor of f(r), and differentiate (remember r is a function of  $\alpha$ )

$$0 = \beta + \sum_{i} b_{i} \frac{1}{r - a_{i}} - \alpha \sum_{i} \frac{b_{i}}{(r - a_{i})^{2}} \frac{dr}{d\alpha}$$

and solving for r' shows that r' is negative

$$r'(\alpha) = -\left(\alpha^2 \sum \frac{b_i}{(r-a_i)^2}\right)^{-1}$$

Finally, a conceptual proof. We assume that  $\alpha$  is non-negative; the case where it is non-positive is similar. When  $\alpha = 0$  the roots of  $f + \alpha g$  are the roots of g. As  $\alpha \to \infty$  the roots go the roots of g. The roots are continuous functions of  $\alpha$ . Since every real number r is the root of at most one equation  $f + \alpha g = 0$  (choose  $\alpha = -f(r)/g(r)$ ) it follows that the roots start at roots of f for  $\alpha = 0$ , decrease to the roots of g, and never back up.

#### **1.14** Recursive sequences of polynomials

We give two different ways of constructing sequences of interlacing polynomials. The first is to use a recurrence; the second is the use of log concavity. The interlacing constructions of previous sections can be iterated to create infinite sequences of interlacing polynomials. **Lemma 1.81.** Suppose  $\{a_i\}, \{b_i\}, \{c_i\}$  are sequences where all  $a_i$  and  $c_i$  are positive, and the  $b_i$  are unrestricted. Define a sequence of polynomials recursively by

$$p_{-1} = 0$$
  

$$p_0 = 1$$
  

$$p_n = (a_n x + b_n)p_{n-1} - c_n p_{n-2} \quad for n \ge 1$$

*These polynomials form an infinite interlacing sequence:* 

$$p_1 \gg p_2 \gg p_3 \gg \cdots$$

*Proof.* We first note that  $p_2$  evaluated at the root of  $p_1$  is  $-c_2$  which is negative. This shows that  $p_2 < p_1$ ; the general case now follows by induction from Lemma 1.25.

It is important to note that the sequences of Lemma 1.81 are exactly the sequences of orthogonal polynomials; see [168, page 106]. These polynomials have the property that the degree of  $p_n$  is n; we can also construct sequences of interlacing polynomials that all have the same degree.

We can sometimes replace the constant  $c_n$  by a quadratic:

**Lemma 1.82.** Suppose that  $f_n \in P^{pos}$  is a sequence of polynomials satisfying the recurrence

$$f_{n+1}(x) = \ell_n(x) f_n(x) - q_n(x) f_{n-1}(x)$$

where

- 1.  $l_n$  is linear with positive leading coefficient.
- 2.  $q_n$  is quadratic with positive leading coefficient.
- 3.  $q_n$  is positive at the roots of  $f_n$ .

*then*  $f_{n+1} \leq f_n$ .

*Proof.* This follows from Lemma 1.25.

Here's an example where it is easy to verify the third condition.

**Corollary 1.83.** Suppose that  $f_n \in P^{p \circ s}$  is a sequence of polynomials satisfying the recurrence

$$f_{n+1}(x) = (a_n x + b_n) f_n(x) + x(c_n x + d_n) f_{n-1}(x)$$

where  $c_n \leq 0$  and  $d_n \geq 0$ . Then  $f_{n+1} \leq f_n$ .

*Proof.* Since  $f_n \in \mathbf{P}^{pos}$  its roots are negative, and when r < 0 we have that  $r(rc_n + d_n) < 0$ . Now apply the lemma.

**Example 1.84.** The Narayana polynomials  $N_n$  are defined by the formula

$$N_n = \sum_{i=0}^n \frac{1}{n} \binom{n}{k} \binom{n}{k-1} x^k.$$

and satisfy the recurrence

$$N_{n} = \frac{1}{n+1} \bigg( (2n-1)(x+1) N_{n-1} - (n-2)(x-1)^{2} N_{n-2} \bigg).$$

It follows from Lemma 1.82 that  $N_n \lt N_{n-1}$ .

**Example 1.85.** We show that the sequence of commutators determined by the Narayana polynomials is an interlacing sequence. Let  $T(x^n) = N_n$ , and define

$$f_n = [\mathsf{D},\mathsf{T}](x^n) = \mathsf{N}'_n - n\mathsf{N}_{n-1}$$

The  $f'_n s$  satisfy the recurrence relation

$$f_{n} = \frac{2n-1}{(n+1)(n-1)^{2}} \left(n^{2}-n-1+\left(n^{2}-n+1\right)x\right) f_{n-1}$$
$$-\frac{(n-2)n^{2}}{(n-1)^{2}(n+1)} (x-1)^{2} f_{n-2}$$

Now  $f_3 = 1 + 3x$ , so using induction and the recurrence we conclude that  $f_n(1) \neq 0$ . It follows from Lemma 1.82 that

$$\cdots \lessdot f_n \lessdot f_{n-1} \lessdot f_{n-2} \cdots$$

Example 1.86. In [116] they considered the recurrence relation

$$p_{n+1} = 2x p_n - (x^2 + (2n-1)^2) p_{n-1}.$$

This satisfies all the hypotheses of Lemma 1.82, so we conclude that all  $p_n$  have all real roots, and consecutive ones interlace.

A small modification to the recurrence in Lemma 1.81 determines a series of polynomials that are known as orthogonal Laurent polynomials .[46]

**Lemma 1.87.** If all coefficients  $a_n$ ,  $b_n$ ,  $c_n$  are positive,  $a_n > c_n$ , and

$$\begin{aligned} p_{-1} &= 0\\ p_0 &= 1\\ p_n &= (a_n x + b_n) p_{n-1} - c_n x p_{n-2} \qquad \textit{for } n \ge 1 \end{aligned}$$

*then*  $p_n \in \boldsymbol{P}^{alt}$ *, and*  $p_1 \gg p_2 \gg p_3 \gg \cdots$ 

*Proof.* We proceed by induction. The case  $p_2 < p_1$  is easy, so assume that  $p_{n-1} < p_{n-2}$ . The hypotheses on the coefficients implies that the coefficients of all  $p_k$  alternate in sign, so  $p_{n-1}$  and  $p_{n-2}$  have all positive roots. Consequently

$$a_n x p_{n-1} \underline{\lessdot} p_{n-1} \ll c_n x p_{n-2}$$

and so

$$p_n = a_n x p_{n-1} - c_n x p_{n-2} \lessdot p_{n-1}$$

Note that from  $f \leq g \ll h$  we can not always conclude that  $f - h \leq g$ . We can do so here because the leading coefficient of  $a_n x p_{n-1}$  is greater than the leading coefficient of  $c_n x p_{n-2}$  by the hypothesis that  $a_n > c_n$ .

*Remark* 1.88. The reverse of a polynomial f(x) of degree n is  $x^n f(1/x)$ . The reverse of a (Laurent) orthogonal polynomials also satisfies a simple recurrence. For orthogonal polynomials as in Lemma 1.81 the recurrence is

$$p_n^{rev} = (a_n + b_n x) p_{n-1}^{rev} - c_n x^2 p_{n-2}^{rev}$$

and for Laurent polynomials as in Lemma 1.87

$$p_n^{rev} = (a_n + b_n x) p_{n-1}^{rev} - c_n x p_{n-2}^{rev}$$

If we modify the signs of coefficients in Lemma 1.81 then we still know the location of the roots. See page 50 for more results.

**Lemma 1.89.** Suppose  $\{a_i\}$ ,  $\{c_i\}$  are sequences of positive terms. Define a sequence of polynomials recursively by

$$p_{-1} = 0$$
  

$$p_{0} = 1$$
  

$$p_{n} = a_{n}x p_{n-1} + c_{n}p_{n-2} \quad for n \ge 1$$

All the roots of  $p_n$  are purely imaginary.

*Proof.* It is easy to see that  $p_n$  only has terms of degree with the same parity as n. Consequently, if we define  $q_n(x) = \iota^n p_n(\iota x)$  then  $q_n$  has all real coefficients, and satisfies the recurrence

$$q_n = a_n x q_{n-1} - c_n q_{n-2}$$

Thus  $q_n$  has all real roots, and so  $p_n$  has purely imaginary roots.

Example 1.90. The d-dimensional cross polytope is

$$\mathcal{O}_{\mathbf{d}} = \{ (\mathbf{x}_1, \dots, \mathbf{x}_d) \in \mathbb{R}^d \mid |\mathbf{x}_1| + \dots + |\mathbf{x}_d| \leq n \}$$

The number of integer points  $p_d$  in  $\mathcal{O}_d$  having exactly k non-zero entries is  $2^k \binom{d}{k} \binom{n}{k}$  since there are

- 2<sup>k</sup> choices of sign for the non-zero entries.
- $\binom{d}{k}$  choices for the non-zero coordinates
- $\binom{n}{k}$  positive integer solutions to  $x_1 + \cdots + x_k \leq n$

and therefore  $p_d = \sum_{k=0}^{n} 2^k {d \choose k}$ ,  $p_d$  is known as the Ehrhart polynomial of  $\mathcal{O}_d$ . If we define  $q_d = \sum_{k=0}^{n} 2^k {d \choose k} {x-1/2 \choose k}$  then we can verify that  $q_d$  satisfies the recurrence relation

$$q_d = \frac{2}{d} x q_{d-1} + \frac{d-1}{d} q_{d-2}$$

and thus  $q_d$  has all imaginary roots. It follows from Lemma 1.89 that the real part of all the roots of  $p_d$  is -1/2. (See [26].)

**Example 1.91.** Assume that  $f_0$  has all positive roots and define a sequence by the recurrence  $f_{n+1} = f_n + x f'_n$ . All these polynomials have the same degree, all real roots, and (1.14.1) is satisfied. (See Lemma 2.11.)

We now briefly discuss log-concavity.

A function f is *strictly concave* if for every choice of distinct a, b in the domain of f we have

$$\frac{f(a)+f(b)}{2} < f\left(\frac{a+b}{2}\right).$$

We say that f is *strictly log concave* if its logarithm is strictly concave. This is equivalent to

$$f(a)f(b) < f\left(\frac{a+b}{2}\right)^2$$
.

If we restrict f to integer values (i.e., a sequence), then at the points i - 1, i, i + 1 a strictly log concave function g satisfies

$$g(i-1) g(i+1) < g(i)^2.$$

A sequence of functions  $f_1, f_2, \ldots$  is strictly log concave if the sequence determined by evaluating at any point is strictly log concave. This is equivalent to

For all x, 
$$f_{n-1}(x)f_{n+1}(x) < f_n(x)^2$$
.

We can extend Lemma 1.13 to chains of interlacings.

**Corollary 1.92.** Assume that  $f_1, f_2, ...$  is a sequence of polynomials with positive leading coefficients satisfying

- 1. The sequence is strictly log concave.
- 2. The leading coefficients have the same sign, or alternating signs.
- $3. \ f_1 \longleftarrow f_2$

then

$$f_1 \longleftarrow f_2 \longleftarrow f_3 \longleftarrow f_4 \longleftarrow \cdots$$
 (1.14.1)

The property of log-concavity is not much stronger than interlacing.

**Lemma 1.93.** Suppose f, g, h are polynomials of the same degree with positive leading coefficients and  $f \ll g$ . Then,  $g \ll h$  if and only if there is a positive constant c such that f, g, ch is strictly log-concave.

*Proof.* If f, g, ch is strictly log-concave then  $g \ll h$  by Corollary 1.92. Conversely, if  $g \ll h$  then there is a c such that f, g, ch is log-concave if and only if  $fh/g^2$  is bounded above. Since f, g, h have the same degree, the limit of  $fh/g^2$  is finite as x goes to  $\pm \infty$ . Consequently, it suffices to show that  $fh/g^2$  is bounded above at each root of g. Now  $g^2$  is positive near a root a of g, so in order for  $fh/g^2$  to be bounded above near a it must be the case that f(a)h(a) < 0. If the roots of g are  $\{a_1 < \cdots < a_n\}$ , then  $\text{sgn} f(a_i) = (-1)^{n+i}$ , and  $\text{sgn} h(a_i) = (-1)^{n+i+1}$ , so  $\text{sgn} f(a_i)h(a_i) = -1$ . We conclude that  $fh/g^2$  is bounded above.

#### 1.15 The positive and negative parts

We consider properties of the positive and negative parts of a polynomial, and apply the properties to a recurrence.

If  $f \in \mathbf{P}$ , then  $f^{neg}$  is the monic polynomial whose roots are the negative roots of f, and  $f^{pos}$  the monic polynomial whose roots are the positive roots of f. For example, if we know that  $f \ll g$  then exactly one of these possibilities holds:

$$\begin{aligned} f^{neg} \ll g^{neg} & \& f^{pos} \ll g^{pos} \\ f^{neg} > g^{neg} & \& f^{pos} < g^{pos} \end{aligned}$$

Similarly, if we know that  $f \leq g$  then there are exactly two possibilities:

$$f^{neg} \gg g^{neg}$$
 &  $f^{pos} \ll g^{pos}$   
 $f^{neg} \ll g^{neg}$  &  $f^{pos} \ll g^{pos}$ 

**Lemma 1.94.** The notation  $\gg \ll$  describes the interlacing of the positive and negative parts of a pair of polynomials. If the polynomials are f, g, then this example means that  $f^{neg} \gg g^{neg}$  and  $f^{pos} \ll g^{pos}$ . An arrow between boxes means the interlacing of f, g is described in the first box, and the interlacing of f + xg, f is described in the target box. Then, we have





*Proof.* There are six assertions to be verified. We describe in detail how to prove one of them, namely

$$\langle \rangle \gg \longleftarrow \langle \langle \langle \langle \rangle \rangle$$

The diagram to be proved is the following statement:

$$f^{neg} \ll g^{neg} \& f^{pos} \ll g^{pos} \implies h^{neg} \lessdot f^{neg} \& h^{pos} \gg f^{pos}$$

where h = f + xg. We use a sign interlacing argument, which we illustrate with an example where f has two positive and three negative roots. These assumptions imply the arrangement of the roots in the diagram below. We can determine the sign of g and f + xg at the roots of f; they are given in the second and third lines of the diagram. There is a root in the leftmost segment since f and f + xg have asymptotes of opposite sign as  $x \to -\infty$ . It follows that there is a root of f + xg in every segment of the first line of the diagram. This tells us where all the roots of f + xg are, and implies the conclusion.

We can apply this result to a recurrence.

**Corollary 1.95.** Consider a sequence satisfying the recursion  $p_k = p_{k-1} + xp_{k-2}$ . If  $\{p_1, p_2\}$  satisfy any of the six interlacing conditions of Lemma 1.94 then all  $p_k$  are in **P**.

The recurrence  $q_n = xq_{n-1} + q_{n-2}$  looks like the recurrence for orthogonal polynomials, but the sign is wrong. We show that if  $q_0 = 1$ ,  $q_1 = x$  then the roots are all on the imaginary axis, and the roots of consecutive terms interlace. (See Lemma 1.89.)

By induction we see that  $q_{2n}$  only has even powers of x, and  $q_{2n+1}$  only has odd powers. Write

$$q_{2n}(x) = p_{2n}(x^2)$$
  $q_{2n+1}(x) = x p_{2n+1}(x^2).$ 

Here are the first few values:

$$\begin{array}{ccccc} k & q_k & p_k \\ 0 & 1 & 1 \\ 1 & x & 1 \\ 2 & 1+x^2 & 1+x \\ 3 & 2x+x^3 & 2+x \\ 4 & 1+3x^2+x^4 & 1+3x+x^2 \end{array}$$

The recurrence for q translates into two recurrences for p:

$$p_{2k+1}(x) = p_{2k}(x) + p_{2k-1}(x)$$
  
$$p_{2k}(x) = xp_{2k-1}(x) + p_{2k-2}(x)$$

and as matrix equations these are

$$\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} p_{2k} \\ p_{2k-1} \end{pmatrix} = \begin{pmatrix} p_{2k+1} \\ p_{2k} \end{pmatrix}$$
$$\begin{pmatrix} x & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} p_{2k+1} \\ p_{2k} \end{pmatrix} = \begin{pmatrix} p_{2k+2} \\ p_{2k+1} \end{pmatrix}$$

We claim that we have the interlacings

$$p_{2k} \ll p_{2k+1} \gg p_{2k+2} \tag{1.15.1}$$

The first is trivial since  $p_{2k} < p_{2k-1}$ . The second follows from Lemma 1.20. In particular, all  $p_k$  are in  $\mathbf{P}^{pos}$ .

Now  $q_{2n}(x) = p_{2n}(x^2)$ , so all roots of  $q_{2n}$  lie on the imaginary axis. The roots of  $q_{2n+1}$  also lie on the imaginary axis, and include 0. The interlacings (1.15.1) imply that the roots of the  $q_k$  interlace.

If we had that  $q_n = xq_{n-1} + \alpha_n q_{n-2}$  where  $\alpha_n$  is positive then the same argument applies, and the roots interlace on the imaginary axis.

#### **1.16** Polynomials without all real roots

We have seen many ways of showing that polynomials have all real roots; here we note that certain polynomials do *not* have all real roots.

- 1.  $p_n = 1 + x + \frac{x^2}{2} + \dots + \frac{x^n}{n!}$  does not have all real roots if n > 1. Since  $p'_n = p_{n-1}$ , if  $p_n$  had all real roots, so would  $p_{n-1}$ . The quadratic formula shows that  $p_2$  does not have all real roots, so  $p_n$  does not either.
- 2.  $q_{2n} = 1 + \frac{x^2}{2} + \dots + \frac{x^{2n}}{(2n)!}$  and  $q_{2n+1} = x + \frac{x^3}{3!} + \dots + \frac{x^{2n+1}}{(2n+1)!}$  do not have all real roots for n > 0. Again, note that  $q_2$  does not have all real roots, and  $q'_n = q_{n-1}$ .
- 3.  $p_n = 1!x + 2!x^2 + \cdots + n!x^n$  does not have all real roots. If it did then its (n-2)th derivative would have two real roots, but

$$\left(\frac{d}{dx}\right)^{n-2} p_n = (n-1)!(n-2)!\left(\frac{1}{n-1} + x + \frac{n^2}{2}x^2\right)$$

and this polynomial has no real roots. A more conceptual proof is that the exponential map (see Chapter 7.7.11) takes  $p_n$  to  $1+x+x^2+\cdots+x^n$ . This polynomial isn't in **P**, yet the exponential map sends **P** to itself.

4. We can use Lemma 1.55 to show that there are no real roots. For example,

If f and g interlace and have no roots in common then the polynomial f g' - f' g has no real roots.

If a monic polynomial does not have *any* real roots then it is positive for all real values. Such polynomials are often stable - see Chapter 21.

#### CHAPTER



# Polynomials with all positive coefficients

In this chapter we consider  $\mathbf{P}^{\text{pos}}$ , the collection of all polynomials that have all negative roots. If the leading coefficient is positive then all coefficients are positive. We introduce an interlacing relation  $\stackrel{+}{\sim}$  for such polynomials that is weaker than interlacing for **P**, but is the natural definition for  $\mathbf{P}^{\text{pos}}$ . We call this relation *positive interlacing*.

#### 2.1 **Basic Properties**

We establish some simple properties of  $\mathbf{P}^{pos}$ .

**Lemma 2.1.** Suppose that  $f \in P^{pos}$ .

- 1. If g in **P** has all positive coefficients then  $g \in P^{pos}$ .
- 2. If  $g \in \mathbf{P}$  and  $f \leq g$  then  $g \in \mathbf{P}^{pos}$ .
- 3. If  $g(x) \in P$  then there is a positive  $\alpha$  such that  $g(x + \alpha) \in P^{pos}$ .
- 4.  $f(x) \in \mathbf{P}^{\text{pos}}$  iff  $f(-x) \in \mathbf{P}^{\text{alt}}$ .
- 5.  $f(-x^2)$  has all real roots if and only if  $f(x) \in \mathbf{P}^{pos}$ .
- 6.  $f(x^2)$  has all real roots if and only if  $f \in \mathbf{P}^{alt}$ .

*Proof.* If  $f \in \mathbf{P}$  has all positive coefficients, and r is positive, then f(r) is a sum of positive terms, and hence r is not a root of f. In addition, 0 is not a root since f(0) is the constant term which is positive by assumption.

If  $f \leq g$ , the roots of f are negative, and are interlaced by the roots of g, so the roots of g are also all negative. Thus,  $g \in \mathbf{P}^{\text{pos}}$ . For the next part, take  $\alpha$  to be larger than the absolute value of the smallest root of f.

The roots of  $f(-x^2)$  are the square roots of the negatives of the roots of f(x), so the roots of f(x) are all negative. The case that  $f \in \mathbf{P}^{alt}$  is similar.

The next result constructs a polynomial in **P** from two interlacing polynomials in  $\mathbf{P}^{\text{pos}}$ . See Theorem 7.64 for a converse.

**Lemma 2.2.** If  $f \leq g$ ,  $f, g \in P^{pos}$ , then

- 1.  $f(-x^2) \leq xg(-x^2)$ .
- 2.  $f(-x^2) + xg(-x^2) \in \mathbf{P}$ .

*Proof.* If the roots of f(x) are  $(-r_1^2, ..., -r_n^2)$  then the roots of the new polynomial  $f(-x^2)$  are  $(\pm r_1, ..., \pm r_n)$ . Consequently,  $f(-x^2) \leq xg(-x^2)$ , which implies the lemma.

Here's a different way to tell if a polynomial is in  $\mathbf{P}^{\pm}$ .

**Lemma 2.3.** If f and g have no factors in common,  $g(0) \neq 0$ , and f interlaces both g and xg then g is in  $\mathbf{P}^{alt}$  or  $\mathbf{P}^{pos}$ .

*Proof.* If we remove a zero from the roots of xg then there will be two consecutive roots of f with no root of g in between unless 0 is the largest or smallest of the roots of xg.

Recall (Corollary 1.50) that if  $T: \mathbf{P} \longrightarrow \mathbf{P}$  then the leading coefficients are all the same sign, or alternate in sign. If  $T: \mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}$  then there are more complicated sign patterns. See Question 9.

**Lemma 2.4.** If either  $T: \mathbb{P}^{pos} \longrightarrow \mathbb{P}^{pos}$  or  $T: \mathbb{P}^{alt} \longrightarrow \mathbb{P}^{alt}$ , T preserves degree, and T(1) is positive then the leading coefficient of  $T(x^n)$  is positive.

If  $T: \mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}^{\text{alt}}$  or  $T: \mathbf{P}^{\text{alt}} \longrightarrow \mathbf{P}^{\text{pos}}$ , and T preserves degree then the signs of the leading coefficients of  $T(x^n)$  alternate.

*Proof.* If  $T(x^n) = \sum c_{i,n} x^i$  then for any  $\alpha$ 

$$T((x + \alpha)^{n}) = T(x^{n}) + n\alpha T(x^{n-1}) + \cdots$$
$$= c_{n,n}x^{n} + (c_{n-1,n} + n\alpha c_{n-1,n-1})x^{n-1} + \cdots$$

If  $T: \mathbf{P}^{pos} \longrightarrow \mathbf{P}^{pos}$  then for any  $\alpha > 0$  the polynomial  $(x + \alpha)^n$  is in  $\mathbf{P}^{pos}$ , and so  $T((x + \alpha)^n)$  is also in  $\mathbf{P}^{pos}$ . It follows that for any positive  $\alpha$  both  $c_{n,n}$  and  $c_{n-1,n} + n\alpha c_{n-1,n-1}$  have the same sign. Choosing  $\alpha$  sufficiently large shows that the leading coefficients  $c_{n,n}$  and  $c_{n-1,n-1}$  have the same sign.

If  $T: \mathbf{P}^{alt} \longrightarrow \mathbf{P}^{alt}$  then we choose  $\alpha < 0$  and the argument is similar. For the last part replace x by -x.

If all the roots are positive then the coefficients alternate. Here's another alternation result.

**Lemma 2.5.** If a polynomial has all real roots and all the coefficients of odd degree are zero then the signs of the non-zero coefficients alternate.

*Proof.* If f(x) is such a polynomial then the non-zero terms all have even degree, so f(x) = f(-x). Thus, the roots of f come in pairs  $r_i$ ,  $-r_i$ , which implies

$$f(x) = \prod_{i} (x^2 - r_i^2)$$

and therefore the non-zero coefficients alternate in sign.

There is an interesting condition that guarantees that the signs of the coefficients alternate in pairs.

**Lemma 2.6.** If f(x) and f(-x) interlace then the sign pattern of the coefficients of f is  $\cdots + + - - + + - - \cdots$ .

*Proof.* We note that if

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots$$

then

$$f(x) + f(-x) = 2(a_0 + a_2x^2 + \cdots) = 2g(x^2)$$

Since the left hand side is in **P** it follows that g(x) has all positive roots. Similarly

$$f(x) - f(-x) = 2x(a_1 + a_3x^2 + \cdots) = 2xh(x^2)$$

implies h has all positive roots. Since g and h have all positive roots, their coefficients alternate, and so the coefficients of f have pattern  $\dots + + - - + + \dots$  since it is the intertwining of two alternating sequences.

### 2.2 Interlacing in P<sup>pos</sup>

Our definition of interlacing for **P** requires that all linear combinations of a pair of interlacing polynomials lie in **P**. Such a definition can not work if we replace **P** by  $\mathbf{P}^{\text{pos}}$ , because if we choose a negative coefficient then the resulting polynomial could have both positive and negative coefficients. This motivates our definition:

**Definition 2.7.** If  $f, g \in \mathbf{P}$  and  $\deg(f) > \deg(g)$  then we say that  $f \stackrel{+}{\sim} g$  if and only if  $f + \alpha g$  is in  $\mathbf{P}$  for all *positive*  $\alpha$ . We say that  $f \stackrel{+}{\sim} g$  if there are open neighborhoods  $\mathcal{O}_f, \mathcal{O}_g$  of f and g such that for  $f_0 \in \mathcal{O}_f$  and  $g_o \in \mathcal{O}_g$  we have  $f_0 \stackrel{+}{\sim} g_0$ .

We write  $f \underline{\ll}^+ g$  if there is a positive  $\beta$  and  $h \in \mathbf{P}$  with positive leading coefficient such that  $f \stackrel{+}{\sim} h$  and  $g = \beta f + h$ .

It is clear from the definition that if  $f, g \in \mathbf{P}^{p \circ s}$  satisfy  $f \leq g$  then  $f \stackrel{+}{\sim} g$ . We can express  $\stackrel{+}{\sim}$  in terms of  $\ll$ .

**Lemma 2.8.** Suppose that  $f, g \in P$  have positive leading coefficients. The following are equivalent

- 1. f ∼ g
- 2. There is  $k \in P$  such that  $k \leq g$ . That is, f and g have a common interlacing.

*Proof.* If there is such a polynomial k then Lemma 1.31 implies that all positive linear combinations are in  $\mathbf{P}^{\text{pos}}$ . The converse follows from Proposition 1.35.

Linear transformations that map  $\mathbf{P}^{\text{pos}}$  to itself preserve  $\stackrel{+}{\sim}$ . The proof is the same as in Theorem 1.43.

**Lemma 2.9.** Suppose that T is a linear transformation that maps  $\mathbf{P}^{\text{pos}}$  to itself, and maps polynomials with all positive coefficients to polynomials with all positive coefficients. Then  $f \stackrel{+}{\sim} g$  implies that  $Tf \stackrel{+}{\sim} Tg$ .

Unlike  $\leq$ , it is not true that f  $\stackrel{+}{\sim}$  g, h implies that g + h  $\in$  **P**. For example, if

$$f = (x+2)(x+6)(x+7)$$
  $g = (x+1)(x+3)$   $h = (x+4)(x+5)$ 

then  $f \stackrel{+}{\sim} g$ , h but g + h has imaginary roots<sup>1</sup>. In addition, even though  $f \stackrel{+}{\sim} h$ , f and f + h do not interlace.

Next, we have a simple property of the derivative that will be generalized to homogeneous polynomials in Corollary 4.3, and to polynomials in two variables in Corollary 9.93.

**Corollary 2.10.** If  $f \leq g$  both have positive leading coefficients then  $f \stackrel{+}{\sim} -g'$ . Equivalently,  $f - ag' \in P$  for a > 0.

*Proof.* Apply Lemma 1.31 to the interlacings  $f \leq g \leq g'$ .

**Lemma 2.11.** If  $f \in P$  then  $f \underline{\ll}^+ xf'$ . If  $f \in P^{alt}$  then  $f \underline{\ll} xf'$ , and if  $f \in P^{pos}$  then  $xf' \underline{\ll} f$ .

*Proof.* Since the leading coefficient of xf' is positive it suffices to show that  $f + \alpha xf' \in \mathbf{P}$  for all positive  $\alpha$ . So write  $f = \prod (x - r_i)$  and let g = xf. Then

$$f + \alpha x f' = \frac{g}{x - 0} + \sum \alpha \frac{g}{x - r_i}$$

<sup>&</sup>lt;sup>1</sup>But see p. 639

where we recall that

$$f' = \sum \frac{f}{x - r_i}$$

This shows that  $f + \alpha x f'$  interlaces xf for positive  $\alpha$ , and hence is in **P**.

The second part follows since  $f \leq f'$ , and 0 is either greater or less than all the roots of f.

### 2.3 Linear combinations in P<sup>pos</sup>

In Proposition 1.35 we found that if  $\alpha f + \beta g$  have all real roots for all nonnegative  $\alpha$ ,  $\beta$  then f and g have a common interlacing. The assumptions were not strong enough to conclude that f and g interlace. The next result gives assumptions about non-constant combinations that lead to interlacing.

**Lemma 2.12.** Suppose that polynomials f, g are in  $P^{\text{pos}}$  and have positive leading coefficients. In addition, assume that for all non-negative  $\alpha$ ,  $\beta$ ,  $\gamma$  the polynomial ( $\alpha x + \beta$ )f +  $\gamma$ g has all real roots. Then g  $\leftarrow$  f.

*Proof.* If we apply Proposition 1.35 we see that for all positive  $\beta$  the polynomials  $(x + \beta)f$  and g have a common interlacing. If we choose  $-\beta$  smaller than the smallest root of f and g then since  $(x + \beta)f$  and g have a common interlacing, it follows that the smallest root of g is less than the smallest root of f. If the combined roots of f and g have two consecutive roots of f (or g) then upon choosing  $-\beta$  in between these roots we find that there is no common interlacing. It is possible to make these choices with positive  $\beta$ 's since all roots of f and g are negative.

**Lemma 2.13.** If  $f + \alpha g$  and  $xf + \alpha g$  are in  $P^{pos}$  for all positive  $\alpha$  then  $g \leftarrow f$ . Equivalently, if  $f \stackrel{+}{\sim} g$  and  $xf \stackrel{+}{\sim} g$  where  $f, g \in P^{pos}$  then  $g \leftarrow f$ .

*Proof.* The only was that common interlacing of f and g, and of xf and g, can occur is if f and g interlace.  $\Box$ 

It is not easy for linear combinations of polynomials to remain in  $\overline{\mathbf{P}^{\alpha lt}}$  or  $\overline{\mathbf{P}^{pos}}$ .

**Lemma 2.14.** Suppose that f, g have the same degree, and that  $f + \alpha g \in \overline{P^{\alpha lt}} \cup \overline{P^{\rho \circ s}}$  for all  $\alpha \in \mathbb{R}$ . Then at least one of the following is true

- 1. g is a constant multiple of f.
- 2. There are constants a, b, c, d such that  $f = (ax + b)x^r$ ,  $g = (cx + d)x^r$ .

*Proof.* We may assume that the degree is greater than 1. If it is not the case that g is a constant multiple of f then by Lemma 1.16 we can find  $\alpha$ ,  $\beta$  such that f +  $\alpha$ g has a positive root, and f +  $\beta$ g has a negative root. Since by hypothesis

we have that  $f + \alpha g \in \overline{\mathbf{P}^{\alpha lt}}$ , and  $f + \beta g \in \overline{\mathbf{P}^{p \circ s}}$ , it follows that for some  $\alpha \leq \gamma \leq \beta$  we have that  $f + \gamma g \in \overline{\mathbf{P}^{\alpha lt}} \cap \overline{\mathbf{P}^{p \circ s}}$ . Now the only polynomials in  $\overline{\mathbf{P}^{\alpha lt}} \cap \overline{\mathbf{P}^{p \circ s}}$  are multiples of powers of x, so we find that  $f + \gamma g = ex^r$  for some e, r. Substituting shows that  $ex^r + (\alpha - \gamma)g \in \mathbf{P}$  for all  $\alpha$ . Consequently, g and  $x^r$  interlace. This implies that  $g = (cx + d)x^r$  or  $g = cx^{r-1}$ . The latter is not possible since f has degree at least r, so  $g = (cx + d)x^r$ . The result now follows by substituting for g.

If  $f \leq g$  then the largest root belongs to f, and the smallest to g. Thus, if  $f, g \in \mathbf{P}^{\text{pos}}$  we have  $xg \leq f$ , and if  $f, g \in \mathbf{P}^{\text{alt}}$  we know  $xf \leq g$ .

**Lemma 2.15.** Suppose  $f \ll g$  have positive leading coefficients. If  $f, g \in \mathbf{P}^{pos}$  then xg + f < f. If  $f, g \in \mathbf{P}^{alt}$  then xf - g < g.

*Proof.* Since  $g \in \mathbf{P}^{a l t}$  we know that  $xf \leq g$ , and therefore  $xf - g \leq g$  by Corollary 1.30. The second case follows by replacing x by -x.

**Corollary 2.16.** *If*  $a \notin (0, n)$  *then* 

If 
$$f \in P^{pos}(n)$$
 then  $-af + xf' \begin{cases} \leq f & a > n \\ \geq f & a = n \\ \geq f & a < 0 \end{cases}$   
If  $f \in P^{alt}(n)$  then  $-af + xf' \begin{cases} \geq f & a > n \\ \geq f & a = n \\ \leq f & a < 0 \end{cases}$ 

*Proof.* If  $f \in \mathbf{P}$  and  $a \notin (0, n)$  then  $-af + xf' \in \mathbf{P}$  by Corollary 1.15. The remaining parts follow the usual arguments.

#### 2.4 Various linear combinations

In this section we first look at linear combinations of the form

$$\mathbf{h} = (\mathbf{a}\mathbf{x} + \mathbf{b})\mathbf{f} + (\mathbf{c}\mathbf{x}^2 + \mathbf{d}\mathbf{x} + \mathbf{e})\mathbf{g}$$

where  $f \leq g$ . We are interested in what restrictions we may place on a, b, c, d, *e* to conclude that h interlaces f, or that h is in  $P^{pos}$  or  $P^{alt}$ . The results are all easy consequences of Lemma 1.25, or or Lemma 1.20.

**Theorem 2.17.** Let  $Tf = (ax + b)f + (cx^2 + dx + e)g$  where  $c \neq 0$ ,  $f \in P^{pos}(n)$  has positive leading coefficient, and  $f \leq g$ . Assume that  $cx^2 + dx + e$  has constant sign  $\epsilon$  on all roots of f and  $(a + nc)\epsilon < 0$  for all positive integers n.

- 1. Tf < f.
- 2. If  $f \in \mathbf{P}^{(-\infty,\alpha)}$  and  $(a + nc) \cdot (Tf)(\alpha) > 0$  then  $Tf \in \mathbf{P}^{(-\infty,\alpha)}$ .

- 3. If  $f \in \mathbf{P}^{(\alpha,\infty)}$  and  $(-1)^n(a + nc) \cdot (Tf)(\alpha) > 0$  then  $Tf \in \mathbf{P}^{(\alpha,\infty)}$ .
- 4. If  $f \in P^{pos}$  and (a + nc), b, e have the same sign then  $Tf \in P^{pos}$ .
- 5. If  $f \in \mathbf{P}^{alt}$  and (a + nc), b, e have the sign then  $Tf \in \mathbf{P}^{alt}$ .

*Proof.* In order to show that  $Tf \le f$  we need to show that Tf sign interlaces f, and that Tf evaluated at the largest root of f has sign opposite to the leading coefficient of Tf (Lemma 1.8). If z is a root of f, then  $(Tf)(z) = g(z) \cdot (cz^2 + dz + e)$ . Since g sign interlaces f, we must have that  $cz^2 + dz + e$  has constant sign on any interval containing roots(f).

If z denotes the largest root of f, then the sign of (Tf)(z) equals  $sgn g(z) sgn(az^2+bz+c)$ , and since f has positive leading coefficient, sgn g(z) is positive. The leading coefficient of Tf is a + nc, so a sufficient condition for Tf < f is that  $sgn(a + nc) sgn(az^2 + bz + c)$  is negative.

Now assume that  $f \in \mathbf{P}^{(-\infty,\alpha)}$ . Since  $Tf \leq f$ , the only root of Tf that could possibly be greater than  $\alpha$  is the largest root. If  $(Tf)(\alpha)$  has the same sign as the leading coefficient a + nc of Tf then there are no roots greater than  $\alpha$ .

In case  $f \in \mathbf{P}^{(\alpha,\infty)}$ , we follow the preceding argument; the condition is that  $(Tf)(\alpha)$  has the same behavior as (Tf)(x) as  $x \longrightarrow -\infty$ .

For last two parts we use the fact that (Tf)(0) = bf(0) + ef'(0). If  $f \in \mathbf{P}^{pos}$  then f(0) and f'(0) are positive. If  $f \in \mathbf{P}^{alt}$  then f(0) and f'(0) have sign  $(-1)^n$ .

The corollary below follows immediately from Lemma 1.20.

**Corollary 2.18.** *Suppose that* f, g, h, k *have positive leading coefficients and satisfy*  $f \ll g \ll h, g < k$ . Then for all a and positive  $\alpha, \beta, \gamma$  we have

- 1. If  $g \in \mathbf{P}^{pos}$  then  $xf + ag + \alpha h \beta f \gamma k \leq g$
- 2. If  $g \in \mathbf{P}^{alt}$  then  $xh + ag \alpha h + \beta f + \gamma k \lt g$
- 3.  $xg + \beta h \ll g$

#### 2.5 The positivity hierarchy

If we are given a polynomial with all positive coefficients then there are many properties that it might have, and we should always check to see which properties it does have. These properties are given in Figure 2.1. We assume that  $f = a_0 + \cdots + a_n x^n$  where all  $a_i$  are positive. The conditions get weaker as we move downward.



Figure 2.1: The positivity hierarchy

#### CHAPTER



## Matrices that preserve interlacing

This chapter is concerned with the question:

Suppose that M is a matrix, either a constant matrix, or with polynomial entries. Suppose that v is a vector of polynomials lying in **P**, or perhaps in **P**<sup>pos</sup>. When is Mv also a vector of polynomials in **P**?

The answer when M is constant is roughly that M satisfy a property called total positivity. Such matrices are discussed in the first section, and the characterization is proved in the following section. The next section discusses two by two matrices that preserve interlacing, and Section 4 covers some results about general M. The final section has some examples of matrices that preserve interlacing for polynomials in  $P^{pos}$ .

#### 3.1 Totally positive matrices

In this section we recall properties of totally positive matrices. We will see that these matrices occur whenever we consider linear combinations of interlacing polynomials. See [5] and [61] for the proofs of the theorems in this section.

A matrix is *strictly totally positive* if all its minors are positive. If all the minors are non-negative we say it is *totally positive*. A simple consequence of the Cauchy-Binet formula for the determinant of the product of two matrices is that products of totally positive matrices are totally positive, and products of strictly totally matrices are strictly totally positive.

A weakening of this definition is often sufficient. We say that a matrix is is *strictly totally positive*<sub>2</sub> if all elements and all two by two determinants are positive. It is *totally positive*<sub>2</sub> if all elements and all two by two determinants are non-negative.

The first theorem is due to Whitney:

**Theorem 3.1** (Whitney). Every totally positive matrix is the limit of a sequence of strictly totally positive matrices.
The next theorem describes a useful decomposition of totally non-negative matrices:

**Theorem 3.2** (Loewner-Whitney). *Any invertible totally positive matrix is a product of elementary Jacobi matrices with non-negative matrix entries.* 

There are three kinds of elementary Jacobi matrices. Let  $E_{i,j}$  be the matrix whose i, j entry is 1 while all other entries are 0. If I is the identity matrix then the elementary Jacobi matrices are  $I + tE_{i,i+1}$ ,  $I + tE_{i+1,i}$ , and  $I + (t-1)E_{i,i}$  where t is positive. For example, in the case i = 2 for 4 by 4 matrices they are

/1	0	0	0)	/1	0	0	0)	/1	0	0	0\		
0	1	t	0	0	1	0	0	0	t	0	0	(0.1	1)
0	0	1	0	0	t	1	0	0	0	1	0	(3.1.	1)
0	0	0	1/	0	0	0	1/	0	0	0	1/		

If  $\begin{pmatrix} a & c \\ b & d \end{pmatrix}$  is strictly totally positive, then a, b, c, d > 0, and ad -bc > 0. An explicit factorization into Jacobi matrices is

$$\begin{pmatrix} a & c \\ b & d \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & b \end{pmatrix} \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & \frac{ad-bc}{ab} \end{pmatrix} \begin{pmatrix} 1 & 0 \\ \frac{ab}{ad-bc} & 1 \end{pmatrix} \begin{pmatrix} 1 & \frac{c}{a} \\ 0 & 1 \end{pmatrix}$$
(3.1.2)

## 3.2 Constant matrices preserving mutual interlacing

Suppose that we have a vector v of polynomials in **P**. If A is a constant matrix when is Av also a vector of polynomials in **P**?

Even if we are only considering three polynomials, a condition such as  $f_1 \ll f_2 \ll f_3$  is not strong enough to draw any conclusions. The reason is that when we form linear combinations, we will need to know the relationship between  $f_1$  and  $f_3$  as well. Thus, we assume that  $f_1$ ,  $f_2$ ,  $f_3$  is mutually interlacing:

**Definition 3.3.** A sequence of polynomials  $f_1, \ldots, f_n$  is *mutually interlacing* if and only if for all  $1 \le i < j \le n$  we have  $f_i \le f_j$ , and each  $f_i$  has positive leading coefficient.

The roots of a mutually interlacing have a simple ordering. If we denote the ordered roots of  $f_i$  by  $r_i^j$  then the roots of  $f_1, \ldots, f_n$  are ordered

$$r_1^1 \leqslant r_2^1 \leqslant \cdots \leqslant r_n^1 \ \leqslant \ r_1^2 \leqslant r_2^2 \leqslant \cdots \leqslant r_n^2 \ \leqslant \ r_1^3 \leqslant r_2^3 \leqslant \cdots \leqslant r_n^3 \ \leqslant \ \cdots$$

Here are a few examples of mutually interlacing sequences of polynomials:

#### Lemma 3.4.

1. If g(x) is a polynomial with roots  $a_1 \leq \cdots \leq a_n$ , then the sequence below is mutually interlacing:

$$\frac{g}{x-a_1}$$
,  $\frac{g}{x-a_2}$ ,  $\cdots$ ,  $\frac{g}{x-a_{n-1}}$ ,  $\frac{g}{x-a_n}$ 

- 2. Any subsequence of a mutually interlacing sequence is mutually interlacing.
- 3. If f<sub>1</sub>, f<sub>2</sub>,..., f<sub>n</sub> is a mutually interlacing sequence then the sequences below are mutually interlacing

 $f_1, f_1 + f_2, f_2, f_2 + f_3, \dots, f_{n-1}, f_{n-1} + f_n, f_n$   $f_1, f_1 + f_2, f_1 + f_2 + f_3, \dots, f_1 + f_2 + \dots + f_n$  $f_1, f_2, f_2, f_2, \dots, f_2$ 

Proof. The roots of the first sequence of polynomials are

 $(a_2, a_3, \ldots, a_n), (a_1, a_3, \ldots, a_n), \ldots, (a_1, \ldots, a_{n-2}, a_n), (a_1, \ldots, a_{n-2}, a_{n-1})$ 

which are interlacing. The remaining assertions follow easily from Corollary 1.30.  $\hfill \Box$ 

We first determine when a linear combination of mutually interlacing polynomials has all real roots. The answer is simple: the coefficients have at most one change of sign.

**Lemma 3.5.** Suppose that  $a_1, \ldots, a_n$  are non-zero constants, and  $f_1 \leq \cdots \leq f_n$  is a mutually interlacing sequence of polynomials. The following are equivalent:

- 1. For all possible  $f_i$  as above  $\sum a_i f_i$  has all real roots.
- 2. There is an integer k so that all the  $a_i$  where  $i \leq k$  have the same sign, and all  $a_i$  where i > k have the same sign.

*Proof.* We can write  $f_i = f_k + \varepsilon_i r_i$  where  $f_i \leq r_i$ ,  $r_i$  has positive leading coefficient, and  $\varepsilon_i = -1$  if i < k and  $\varepsilon_i = 1$  if i > k. Now

$$f = \sum a_i f_i = (\sum_i a_i) f_k + \sum_i (a_i \varepsilon_i r_i)$$

and by hypothesis the coefficients  $a_i \varepsilon_i$  all have the same sign. It follows from Lemma 1.10 that  $f \in P$ .

Conversely, consider the three mutually interlacing polynomials

$$g_1 = (x - 8)(x - 24)$$
  

$$g_2 = (x - 5)(x - 21)$$
  

$$g_3 = (x - 2)(x - 13)$$

where  $g_1 \ll g_2 \ll g_3$ . Observe that

$$g_1 - g_2 + g_3 = x^2 - 21x + 13 \notin \mathbf{P}$$

Now embed  $g_1, g_2, g_3$  in a mutually interlacing sequence so that  $f_1 = (1/|a_1|)g_1$ ,  $f_i = (1/|a_i|)g_2$ , and  $f_n = (1/|a_n|)g_3$ . Since

$$a_1f_1 + a_if_i + a_nf_n = sgn(a_1)g_1 + sgn(a_i)g_2 + sgn(a_n)g_3$$

we know that if the left hand side has all real roots then we can not have that  $sgn(a_1) = sgn(a_n) \neq sgn(a_i)$ . This implies that there can be at most one sign change.

The following obvious corollary will be generalized in the next section where sums are replaced by integrals.

**Corollary 3.6.** Suppose that  $a_1, ..., a_n$  are positive constants, and  $f_1 \leq \cdots \leq f_n$  is a mutually interlacing sequence of polynomials. Then

$$f_1 \underline{\ll} a_1 f_1 + \dots + a_n f_n \underline{\ll} f_n \tag{3.2.1}$$

Our first theorem is a simple consequence of the general theorems of the previous section.

**Theorem 3.7.** Suppose  $f_1, \ldots, f_n$  are mutually interlacing, and  $A = (a_{i,j})$  is an n by n matrix of constants such that

$$\mathsf{A} \cdot (\mathsf{f}_1, \ldots, \mathsf{f}_n) = (\mathsf{g}_1, \ldots, \mathsf{g}_n)$$

If A is totally positive, then  $g_1, \ldots, g_n$  is mutually interlacing.

*Proof.* Since the limit of a mutually interlacing sequence is mutually interlacing, and any non-invertible totally positive matrix is the limit of strictly totally positive matrices, we may assume that A is invertible. By the Loewner-Whitney theorem we can assume that A is an elementary Jacobi matrix. The three classes of matrices applied to  $f_1 \leq \cdots \leq f_n$  give three easily verified mutually interlacing sequences:

$f_1$	• • •	$f_{i-1}$	$f_i + tf_{i+1}$	$f_{i+1}$	• • •	fn
$f_1$		$f_{\mathfrak{i}-1}$	fi	$tf_{\mathfrak{i}} + f_{\mathfrak{i}+1}$		fn
$f_1$		$f_{\mathfrak{i}-1}$	tfi	$f_{i+1}$		fn

The two by two case for all positive entries follows from the theorem. However, we can determine the action of an arbitrary two by two matrix.

**Corollary 3.8.** Suppose that  $f_1 \ll f_2$  (or  $f_1 \ll f_2$ ) have positive leading coefficients, and  $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} f_1 \\ f_2 \end{pmatrix} = \begin{pmatrix} g_1 \\ g_2 \end{pmatrix}$ , where without loss of generality we may assume that a is positive. If the determinant ad - bc is positive then  $g_1 \ll g_2$ , and if it is negative  $g_2 \ll g_1$ .

*Proof.* Assume that the determinant ad - bc is non-zero. If r is a common root of  $g_1$  and  $g_2$  then since the matrix is invertible we conclude that  $f_1(r) = f_2(r) = 0$  which contradicts the assumption that  $f_1$  and  $f_2$  have no common factors. Now any linear combination of  $g_1$  and  $g_2$  is a linear combination of  $f_1$  and  $f_2$ , and thus has all real roots. Consequently,  $g_1$  and  $g_2$  interlace. The direction of interlacing will not change if we continuously change the matrix  $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$  as long as the determinant never becomes zero. Consequently, if the determinant is positive we can deform the matrix to the identity, in which case  $g_1 \ll g_2$ . If the determinant is negative we can deform the matrix to  $\begin{pmatrix} 1 & 0 \\ 1 & -1 \end{pmatrix}$ , and since  $f - g \ll f$  we are done.

**Example 3.9.** Consider  $\begin{pmatrix} 1 & b \\ 1 & d \end{pmatrix} \begin{pmatrix} f \\ g \end{pmatrix} = \begin{pmatrix} f+bg \\ f+dg \end{pmatrix}$ . If  $f \leq g$  and 0 < b < d then  $f + bg \leq f + dg$ .

The examples of mutually interlacing sequences in Lemma 3.4(3) are given by the following totally positive matrices, where we take n = 3 for clarity, and the "." are zeros.

/1		• )		/1	\
1	1		$(1 \cdot .)$	$\begin{pmatrix} 1 & \cdot \\ & 1 \end{pmatrix}$	• )
	1				·
1.	1	1	$(1 \ 1 \ 1)$		•
(.		1/		$\langle \cdot   1 \rangle$	•/

An immediate corollary of Corollary 3.8 is that the inverse of a totally positive matrix sometimes preserves interlacing.

#### Corollary 3.10. If

- 1. M is a 2 by 2 totally positive invertible matrix with non-negative entries
- 2. f, g, u,  $v \in P$  have positive leading coefficients
- 3.  $M\left(\begin{smallmatrix}f\\ g\end{smallmatrix}\right) = \left(\begin{smallmatrix}u\\ v\end{smallmatrix}\right)$

then  $f \leq g$  if and only if  $u \leq v$ .

*Remark* 3.11. The inverses of the 3 by 3 Jacobi matrices (except the diagonal ones) do not satisfy the conclusions of the lemma. However, we do have that the matrices of the form

$$\begin{pmatrix} 1 & 0 & -a \\ 0 & 1 & 0 \\ -b & 0 & 1 \end{pmatrix}$$

satisfy the conclusions if  $0 \le ab < 1$ . That is, if  $0 \le a, b$  and  $0 \le ab < 1$ ,

$$\begin{pmatrix} 1 & 0 & -a \\ 0 & 1 & 0 \\ -b & 0 & 1 \end{pmatrix} \begin{pmatrix} f \\ g \\ h \end{pmatrix} = \begin{pmatrix} u \\ v \\ w \end{pmatrix}$$

f, g, h is mutually interlacing, and f, g, h, u, v, w have positive leading coefficients then u, v, w is mutually interlacing. This is equivalent to showing that

$$f-ah \underline{\ll} g \qquad g \underline{\ll} - bf + h \qquad f-ah \underline{\ll} h - bf$$

The first follows from  $f \leq g \leq h$ . For the second, write  $f = \alpha g - r$ ,  $h = \beta g + s$  where  $g \leq r$ , s. Then  $g \leq (\beta - b\alpha)g + s + br$  since the leading coefficient of the right hand side is positive by hypothesis. Finally, the last one is a two by two matrix  $\begin{pmatrix} 1 & -\alpha \\ -b & 1 \end{pmatrix} \begin{pmatrix} f \\ h \end{pmatrix} = \begin{pmatrix} f-\alpha h \\ h-bf \end{pmatrix}$ . The lemma shows that the interlacing is satisfied if the determinant is positive, which is true since  $0 \leq ab < 1$ .

Surprisingly, we do not need the full force of total positivity to conclude that a matrix preserves interlacing, but just the non-negativity of all two by two submatrices.

**Theorem 3.12.** Suppose that  $A = (a_1, ..., a_n)$  and  $B = (b_1, ..., b_n)$  are vectors of positive real numbers, and  $f = (f_1, ..., f_n)$  is a vector of mutually interlacing polynomials. Then

Af  $\leq$  Bf for all f if and only if  $\begin{pmatrix} A \\ B \end{pmatrix}$  is totally positive<sub>2</sub>.

Proof. Consider the mutually interlacing set of polynomials

$$c_1 f_1, c_2 f_2, \ldots, c_n f_n$$

where the  $c_i$  are positive. By continuity of roots, if we set  $c_i = c_j = 1$ , and all other c's to zero, then

$$a_i f_i + a_j f_j \underline{\ll} b_i f_i + b_j f_j.$$

We can rewrite this as  $\begin{pmatrix} a_i & a_j \\ b_i & b_j \end{pmatrix} \begin{pmatrix} f_i \\ f_j \end{pmatrix} = \begin{pmatrix} h \\ k \end{pmatrix}$  where  $h \leq k$ . By Corollary 3.8 we must have  $\begin{vmatrix} a_i & a_j \\ b_i & b_j \end{vmatrix} \ge 0$ .

Conversely, given A and B construct a totally positive n by n matrix C whose first row is A, second row is B, and whose remaining rows are all 0. If we apply C to the mutually interlacing sequence f, g, g,  $\cdots$ , g the conclusion follows from Theorem 3.7.

A small modification of this argument shows

**Corollary 3.13.** Suppose  $f_1, ..., f_n$  are mutually interlacing, and  $A = (a_{i,j})$  is an n by m matrix of constants such that

$$A \cdot (f_1, \ldots, f_n) = (g_1, \ldots, g_m)$$

If A is totally positive<sub>2</sub>, then  $g_1, \ldots, g_m$  is mutually interlacing.

If we fix the constants rather than the polynomials, then we need a determinant condition on the polynomials. Lemma 3.14. Suppose that we have the following interlacing diagram

where

- 1. The f<sub>i</sub> are mutually interlacing.
- 2. The g<sub>i</sub> are mutually interlacing.

3. 
$$\left| \begin{array}{c} f_{\mathfrak{i}} & f_{\mathfrak{i}+1} \\ g_{\mathfrak{i}} & g_{\mathfrak{i}+1} \end{array} \right| < 0$$
 for  $1 \leq \mathfrak{i} < \mathfrak{n}$ 

then for any non-negative  $\alpha_1, \ldots, \alpha_n$  we have  $\sum \alpha_i f_i \ll \sum \alpha_i g_i$ 

*Proof.* We may assume that all  $\alpha_i$  are positive. We proceed by induction. The case n = 2 is Lemma 1.51. let  $h_1 = \alpha f_1 + \alpha_2 f_2$  and  $k_1 = \alpha g_1 + \alpha_2 g_2$  then by the lemma we know that  $h_1 \ll k_1$  and  $\{h_1, f_2, \ldots, f_n\}$  and  $\{k_1, g_2, \ldots, g_n\}$  are mutually interlacing. The determinant condition is satisfied since

$$\begin{vmatrix} h_1 & f_2 \\ k_1 & g_2 \end{vmatrix} = \begin{vmatrix} \alpha_1 f_1 + \alpha_2 f_2 & f_2 \\ \alpha_1 g_1 + \alpha_2 g_2 & g_2 \end{vmatrix} = \alpha_1 \begin{vmatrix} f_1 & f_2 \\ g_1 & g_2 \end{vmatrix} < 0$$

We continue combining terms until done.

Mutually interlacing sequences have a nice interpretation when expressed in the interpolation basis. Suppose  $f_1, \ldots, f_n$  is a sequence of mutually interlacing polynomials of degree d with positive leading coefficients. All the smallest roots of  $f_1, \ldots, f_n$  are smaller than all the second smallest roots of  $f_1, \ldots, f_n$ , and so on. Thus, we can always find a g such that  $g \leq f_i$  for all i. It follows that there are non-negative constants  $a_{i,j}$  such that

$$f_i = \sum a_{i,j} \frac{g(x)}{x - b_j}.$$

If we set A =  $(a_{i,j})$ , F =  $(f_1, \dots, f_n)$  and G =  $(\frac{g}{x - b_j})$  then AG = F.

The next lemma characterizes those A's that always yield mutually interlacing sequences.

**Corollary 3.15.** *Suppose* G *and* A *are as above. For all choices of* G *the sequence* AG *is mutually interlacing if and only if* A *is totally positive*<sub>2</sub>*.* 

*Proof.* We simplify what needs to be proved. First of all, it suffices to assume that A has only two rows since the conclusion only involves two by two matrices. Thus we will show that if for all g we have

$$\sum a_i \frac{g(x)}{x - b_i} \leq \sum c_i \frac{g(x)}{x - b_i}$$

then  $\begin{vmatrix} a_i & a_{i+1} \\ c_i & c_{i+1} \end{vmatrix} \ge 0$ . Next, notice that we can multiply any column of A by any positive number, and the determinants will still be non-negative. By continuity we can set all other columns to zero, and so we assume that

$$a_{i}\frac{g(x)}{x-b_{i}} + a_{i+1}\frac{g(x)}{x-b_{i+1}} \leq c_{i}\frac{g(x)}{x-b_{i}} + c_{i+1}\frac{g(x)}{x-b_{i+1}}$$

Next, we choose  $g = x^{n-2}(x - \alpha)(x - \beta)$  where  $b_i = \alpha$  and  $b_{i+1} = \beta$ , and  $\beta > \alpha$ . We can cancel the powers of x, and so we have

$$a_{i}(x-\beta) + a_{i+1}(x-\alpha) \underline{\ll} c_{i}(x-\beta) + c_{i+1}(x-\alpha)$$

The root of the left hand side is not less than the right hand side, so

$$\frac{c_{i+1}\alpha + c_{i}\beta}{c_{i} + c_{i+1}} \leqslant \frac{a_{i+1}\alpha + a_{i}\beta}{a_{i} + a_{i+1}}$$

and from this we find that

$$(\alpha - \beta)(a_{i+1}c_i - a_ic_{i+1}) \ge 0$$

which is the desired conclusion.

The converse follows from Corollary 3.13 since the set  $\left\{\frac{g(x)}{x-b_i}\right\}$  is mutually interlacing.

The following property of mutually interlacing polynomials generalizes Lemma 1.19, and will be extended to integrals in the next section (Lemma 8.60).

**Lemma 3.16.** If  $f_1, \ldots, f_n$  and  $g_1, \ldots, g_n$  are two sequences of mutually interlacing polynomials with positive leading coefficients, then

$$\mathsf{f}_1\,\mathsf{g}_n+\mathsf{f}_2\,\mathsf{g}_{n-1}+\cdots+\mathsf{f}_n\,\mathsf{g}_1\in \boldsymbol{P}.$$

*Equivalently,*  $\sum f_i(x)g_i(-x) \in \mathbf{P}$ .

*Proof.* Assume the f's have degree r and the g's have degree s. We may assume that the roots are distinct. It is helpful to visualize the location of these roots in two dimensions, so we plot the roots of  $f_i$  and  $g_{n+1-i}$  with y coordinate i. For instance, if r = 3, s = 2, n = 4 then a possible diagram of the roots is

The roots of the f's lie in three increasing groups, because each  $f_i$  has three roots, and the  $f_i$  are mutually interlacing. Since the g's occur in the opposite order, their groups are decreasing. In addition, since the f's are mutually interlacing, each group of four roots on a line does not vertically overlap any other group of four roots.

In this diagram the dotted lines represent values of x for which each of the terms  $f_1 g_n, \dots, f_n g_1$  have the same sign slightly to the left of x. To see this, consider a term  $f_i g_{n+1-i}$ . At any value x and sufficiently small  $\epsilon$  the sign of  $f_i(x - \epsilon) g_{n+1-i}(x - \epsilon)$  is (-1) to the number of roots to the right of x. The



Figure 3.1: Roots of mutually interlacing polynomials

doted lines arise at the intersection of upward and downward lines, or at gaps where there are no up or down lines.

The general case is no different. There are n levels, r upward lines, and s downward lines. It is easy, but notationally complicated, to see that there are r + s + 1 vertical lines where all signs are the same. Between each pair of signs there is a root of the sum, accounting for all the roots.

The final statement follows from the first since the interlacing direction of  $\{g_i(-x)\}$  is the reverse of the interlacing direction of  $\{g_i(x)\}$ .

# 3.3 Determinants and mutual interlacing

If  $f_1, \ldots, f_d$  is a sequence of mutually interlacing polynomials of degree n, and  $f_i = \sum a_{i,j} x^j$  then the matrix  $(a_{d-i,j})$  is called the *matrix of coefficients* of  $f_1, \ldots, f_d$ . If the matrix of coefficients of a sequence of mutually interlacing polynomials is TP<sub>k</sub> (or totally positive), we say that the sequence is TP<sub>k</sub> (or totally positive). We show that any mutually interlacing sequence is TP<sub>2</sub>, but not necessarily TP<sub>3</sub>. In addition, we show that certain mutually interlacing set of polynomials are totally positive.

**Example 3.17.** Here are three mutually interlacing polynomials

$$\begin{split} f_1 &= (2+x) \left(9+x\right) (14+x) \\ f_2 &= (5+x) \left(10+x\right) (15+x) \\ f_3 &= (6+x) \left(12+x\right) (18+x) \end{split}$$

The matrix of coefficients is

$$\begin{pmatrix} 1296 & 396 & 36 & 1 \\ 750 & 275 & 30 & 1 \\ 252 & 172 & 25 & 1 \end{pmatrix}$$

It's easy to see that it is  $TP_2$ , but the four three by three determinants are all negative, so it is not  $TP_3$ .

We first have two useful facts.

Lemma 3.18.

- 1. Suppose that C is the matrix of coefficients of  $V = (f_1, f_2, ..., f_n)$ , and M is a compatible matrix, then the matrix of coefficients of MV is MC.
- 2. If  $g \in P^{pos}$ , and the mutually interlacing sequence  $V = (f_1, \dots, f_d)$  is totally positive, then the sequence  $(g f_1, g f_2, \dots, g f_n)$  is also totally positive.

*Proof.* The first is immediate from the definition. For the second, since we can factor g into positive factors, it suffices to assume that  $g = x + \alpha$ , where  $\alpha > 0$ . If  $f_1 = \sum a_i x^i$  and  $f_2 = \sum b_i x^i$  then the matrices of coefficients satisfy

$$\begin{pmatrix} a_0 & a_1 & a_2 & \cdots \\ b_0 & b_1 & b_2 & \cdots \\ \vdots & & & \end{pmatrix} \begin{pmatrix} \alpha & 1 & & & \\ & \alpha & 1 & & \\ & & \alpha & 1 & \\ & & & & \ddots \end{pmatrix} = \\ \begin{pmatrix} \alpha & a_0 & \alpha & a_1 + a_0 & \alpha & a_2 + a_1 & \cdots \\ \alpha & b_0 & \alpha & b_1 + b_0 & \alpha & b_2 + b_1 & \cdots \\ \vdots & & & & & \end{pmatrix}$$

The first matrix is the matrix of coefficients of V, the second is totally positive, and the third is the matrix of coefficients of  $(x + \alpha)V$ . The conclusion follows.

If  $f = \prod (x + r_i)$  where  $r_1 < r_2 < \cdots < r_n$  then we say that the sequence

$$\frac{f(x)}{x+r_n}, \frac{f(x)}{x+r_{n-1}}, \cdots, \frac{f(x)}{x+r_1}$$

of mutually interlacing polynomials is determined by f.

**Proposition 3.19.** If  $V = (f_1, ..., f_n)$  is a sequence of mutually interlacing polynomials determined by  $f \in \mathbf{P}^{pos}$ , and M is a totally positive matrix, then the matrix of coefficients of MV is totally positive.

*Proof.* Since the matrix of coefficients of MV is M times the matrix of coefficients of V, it suffices to show that V is totally positive. We thus may assume that  $f_i = f/(x+r_{n+1-i})$ , where  $f(x) = (x+r_1)\cdots(x+r_n)$  and  $0 < r_1 < \cdots < r_n$ . The matrix of coefficients is an n by n matrix. Consider the submatrix determined by a set R of d rows and a set of d columns. Define

$$g(x) = \prod_{k \notin R} (x + r_k) \qquad h(x) = \prod_{k \in R} (x + r_k)$$

Note  $f_i = g(x) \cdot h(x)/(x + r_i)$ . We want to prove that all the determinants of size d by d are positive for the following set of polynomials

$$\{g(\mathbf{x}) \cdot \mathbf{h}(\mathbf{x})/(\mathbf{x}+\mathbf{r}_k) \mid k \in \mathbf{R}\}$$

By the lemma, it suffices to show that the determinant of the matrix of coefficients of

$${h(x)/(x+r_k) \mid k \in R}$$

is positive. For instance, if d = 4 and  $R = \{1, 2, 3, 4\}$  then the matrix of coefficients is

 $\begin{pmatrix} r_2 r_3 r_4 & r_2 r_3 + r_2 r_4 + r_3 r_4 & r_2 + r_3 + r_4 & 1 \\ r_1 r_3 r_4 & r_1 r_3 + r_1 r_4 + r_3 r_4 & r_1 + r_3 + r_4 & 1 \\ r_1 r_2 r_4 & r_1 r_2 + r_1 r_4 + r_2 r_4 & r_1 + r_2 + r_4 & 1 \\ r_1 r_2 r_3 & r_1 r_2 + r_1 r_3 + r_2 r_3 & r_1 + r_2 + r_3 & 1 \end{pmatrix}$ 

First of all, the determinant is a polynomial of degree  $\binom{k}{2}$  in the  $r_i$ . If  $r_i = r_j$  then two rows are equal, and the determinant is zero. Thus,  $r_i - r_j$  divides the determinant. Consequently,

$$\Delta = \prod_{i < j} (r_j - r_i)$$

divides the determinant. But this is a polynomial of degree  $\binom{k}{2}$ , so the determinant is a constant multiple of  $\Delta$ . We can check that that the constant of proportionality is 1. Since  $r_i < r_j$ , all terms of the product in  $\Delta$  are positive, so the determinant is positive.

If the matrix of coefficients of a sequence V of mutually interlacing polynomials is not totally positive, then V is not obtained by multiplying the sequence determined by the factors of some polynomial by a totally positive matrix. For instance, the polynomials in Example 3.17 form such a sequence.

**Lemma 3.20.** If  $f_1, \ldots, f_d$  is a sequence of mutually interlacing polynomials in  $P^{pos}$  then the matrix of coefficients is  $TP_2$ .

*Proof.* It suffices to prove that all two by two determinants are non-negative for  $f \ll g$ . Write  $g = af + \sum \alpha_i f/(x + r_i)$  where  $f = \prod (x + r_i)$ , and all  $a_i$  are non-negative. If we write this equation in terms of matrices

$\begin{pmatrix} 1\\ \mathfrak{a} \end{pmatrix}$	$\begin{array}{c} 0 \\ \alpha_1 \end{array}$	$0 \\ \alpha_2$	· · · )	$\begin{pmatrix} f \\ f/(x+r_n) \\ \vdots \\ f/(x+r_n) \end{pmatrix}$	$= \begin{pmatrix} f \\ g \end{pmatrix}$
				$f/(x + r_1)$	

we see the matrix on the left is totally positive, and the middle matrix is totally positive by the Proposition and taking limits, so the conclusion follows.

We will prove this same result in a different way in Corollary 4.31. If we apply the Lemma to xf and f where  $f = \sum a_i x^i \in \mathbf{P}^{pos}$ , then the matrix of coefficients is

$$\begin{pmatrix} a_0 & a_1 & a_2 & \cdots \\ 0 & a_0 & a_1 & \cdots \end{pmatrix}$$

The two by two determinants show that  $a_k^2 \ge a_{k-1}a_{k+1}$ . This is a special case of Newton's inequality (p. 106).

# 3.4 Interlacing polynomials and the Hurwitz matrix

A pair of interlacing polynomials determines a totally positive matrix. We prove it inductively; the key step is given in the following lemma.

**Lemma 3.21.** *Define*  $T_c(f, g) = (xg + cf, f)$ *, and* 

 $\mathcal{D} = \{(f,g) \mid f \leq g \text{ or } f \ll g \text{ in } \mathbf{P}^{pos} \text{ with positive leading coefficients } \}.$ 

**1-1:**  $T_c : \mathfrak{D} \longrightarrow \mathfrak{D}$  is one to one.

**onto:** Given  $\alpha \in D$  there is a positive c and  $\beta \in D$  such that  $T_{c}(\beta) = \alpha$ .

*Proof.* The first part is clear. For the second choose  $(h, k) \in D$ , and set c = h(0)/k(0) and g = (h - ck)/x. Since h and ck have the same constant term, g is a polynomial. If  $h \le k$  then  $gx = h - ck \le k$  so  $T_c(k, g) = (h, k)$  and  $k \ll g$ .

If  $h \ll k$  then write h = ak - l where a is positive and h < l and follow the previous argument to conclude that k < g.

If we iterate the lemma, we see that we can find positive  $c_{\mathfrak{i}}$  and positive  ${\mathfrak{b}}$  such that

$$(f,g) = T_{c_1} \cdots T_{c_n} H(b,0).$$
 (3.4.1)

This factorization can be turned into a factorization of matrices. If  $f = \sum a_i x^i$ ,  $g = \sum b_i x^i$  and we define

$$J(\mathbf{c}) = \begin{pmatrix} \mathbf{c} & 1 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 1 & 0 & 0 & 0 & \dots \\ 0 & 0 & \mathbf{c} & 1 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 1 & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix} \quad H(\mathbf{f}, \mathbf{g}) = \begin{pmatrix} a_0 & a_1 & a_2 & \dots & \\ 0 & b_0 & b_1 & b_2 & \dots \\ 0 & a_0 & a_1 & a_2 & \dots \\ 0 & 0 & b_0 & b_1 & b_2 & \dots \\ 0 & 0 & a_0 & a_1 & a_2 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$
(3.4.2)

then J(c)H(f,g) = H(xg + cf, f). The matrix H(f,g) is sometimes called the *Hurwitz matrix*.

**Proposition 3.22.** If  $f \leq g$  in  $P^{pos}$  have positive leading coefficients then

- 1. H(f, g) is totally positive.
- 2. There are positive  $c_i$  and b such that

$$H(f,g) = J(c_1) \cdots J(c_n)H(b,0).$$

*Proof.* Since all J(c) and H(b, 0) are totally positive, the first statement is a consequence of the second. The second is a consequence of the factorization (3.4.1) and the lemma. 

*Remark* 3.23. If we write out the matrix determined by  $f \leq f'$ 

$$\begin{pmatrix} a_1 & 2a_2 & 3a_3 & \dots & \\ a_0 & a_1 & a_2 & a_3 & \dots & \\ 0 & a_1 & 2a_2 & 3a_3 & \dots & \\ 0 & a_0 & a_1 & a_2 & a_3 & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots & \end{pmatrix}$$

、

then we see that  $\begin{vmatrix} k \alpha_k & (k+1)\alpha_{k+1} \\ \alpha_{k-1} & \alpha_k \end{vmatrix} \ge 0$ . This is equivalent to

$$\mathfrak{a}_{k}^{2} \geqslant \frac{k+1}{k} \mathfrak{a}_{k-1} \mathfrak{a}_{k+1}$$

and this is a weak form of Newton's inequality (See p. 106.).

We will prove the following corollary in a very different way later - see Theorem 4.23.

**Corollary 3.24.** If  $f = \sum a_i x^i$  is in  $P^{pos}$  then the matrix below is totally positive.

$(a_0)$	$a_1$	$a_2$		• )
.	$a_0$	$a_1$	$a_2$	
.	•	$a_0$	$a_1$	
(:				·)

*Proof.* Apply the proposition to  $f \leq f'$  and select the submatrix whose rows come from f. 

#### 3.5 The division algorithm

We study the division algorithm applied to interlacing polynomials. The basic result is the following lemma:

**Lemma 3.25.** Suppose that f, g have positive leading coefficients, and  $f \leq g$  in  $P^{\text{pos}}$ . *If we divide* f by g

f = (ax + b)g - r where deg(r) < deg(g)

then

1. g < r.

- 2.  $r \in P^{pos}$
- 3. r has positive leading coefficient.

4.  $\deg(r) = n - 2$ 

*Proof.* Since f sign interlaces g, it follows that r sign interlaces f. The degree of r is less than the degree of g so we have that  $g \leq r$ . The interlacing is strict, since any common factor of g and r is a common factor of f and g. Since f strictly interlaces g, they have no common factors. Since  $g \in \mathbf{P}^{\text{pos}}$ , so is r.

Now we determine the leading coefficient of r. We may assume that f, g are monic, so

$$f = x^{n} + a_{1}x^{n-1} + a_{2}x^{n-2} + \cdots$$
  

$$g = x^{n-1} + b_{1}x^{n-2} + b_{2}x^{n-3} + \cdots$$
  

$$r = (x + a_{1} - b_{1})g - f$$
  

$$= (a_{1}b_{1} + b_{2} - b_{1}^{2} - a_{2})x^{n-2} + \cdots$$

Using Proposition 3.22 and reversing f, g we see that

$$0 \leqslant \begin{vmatrix} 1 & b_1 & b_2 \\ 1 & a_1 & a_2 \\ 0 & 1 & b_1 \end{vmatrix} = a_1 b_1 + b_2 - b_1^2 - a_2$$

Because r interlaces g the degree of r is n - 2, and so the leading coefficient is positive.

*Remark* 3.26. If we apply the division algorithm to  $f \leq f'$  where  $f \in \mathbf{P}^{pos}$  then all coefficients of r must be positive. The resulting inequalities are either Newton's inequalities, or simple consequences of them.

It's curious that the lemma does not depend on the positivity of f and g.

**Lemma 3.27.** Suppose that f, g have positive leading coefficients, and  $f \leq g$  in **P**. If we divide f by g

f = (ax + b)g - r where deg(r) < deg(g)

then

1. g ⊲ r.

2. r has positive leading coefficient.

3.  $\deg(r) = n - 2$ 

Proof. If we write

$$f(x) = \prod_1^n (x - r_i) \qquad g = \sum_1^n \alpha_i \, \frac{f}{x - r_i}$$

where  $\alpha_i \ge 0$  then a computation shows that the coefficient of r is

$$-\frac{1}{(\alpha_1+\dots+\alpha_n)^2}\sum_{i< j}\alpha_i\alpha_j(r_i-r_j)^2$$

**Example 3.28.** We can apply this in the usual way to solve Ff - Gg = 1. For example,

$$f = (x + 1)(x + 3)(x + 6)$$
  

$$g = (x + 2)(x + 4)$$
  

$$f = (x + 4)[g] - [5x + 14]$$
  

$$g = \frac{5x + 16}{25}[(5x + 4)] - \frac{24}{25}$$

and recursively solving we get

$$1 = \underbrace{\frac{39 + 36x + 5x^2}{24}}_{G} [(x+2)(x+4)] - \underbrace{\frac{5x+16}{24}}_{F} [(x+1)(x+3)(x+6)]$$

Notice that in this example F and G are in  $\mathbf{P}^{pos}$ . In general, if f has degree n then the algorithm yields F, G with deg(F) = deg(G) - 1 = deg(f) - 2.

**Lemma 3.29.** If  $f \leq g$  in  $P^{pos}$ , deg(F) = deg(G)-1 = deg(f)-2, and fF-gG = 1 then

- 1. F,  $G \in P^{pos}$ .
- 2.  $f \leq g, G and g, G \leq F$

*Proof.* If  $\alpha$  is a root of f then  $g(\alpha)(G(\alpha) = 1$  so G and g have the same sign at roots of f. Since deg(G) = deg(f) - 1 it follows that  $f \leq G$  - in particular,  $G \in \mathbf{P}$ . The same arguments show  $G \leq F$  and  $g \leq F$ . All interlacings are strict since any common divisor divides 1.

The identity fF - gG = 1 is known as the *Bezout identity*.

**Corollary 3.30.** If  $f \leq g$  in  $P^{pos}$  then there exist  $F, G \in P^{pos}$  such that

- 1. fF gG = 1
- 2.  $f \leq g, G \text{ and } g, G \leq F$

Restated in terms of the matrix  $\begin{pmatrix} f & g \\ F & G \end{pmatrix}$ 

**Corollary 3.31.** A strict interlacing pair (f, g) in  $P^{pos}$  can be extended to an interlacing matrix of determinant 1.

**Example 3.32.** If we assume that the anti-diagonal elements are (x - a)(x - b) and 1 then there are exactly two solutions to the Bezout identity:

$$\begin{pmatrix} x - \frac{a+b\pm\sqrt{(a-b)^2-4}}{2} & (x-a)(x-b) \\ 1 & x - \frac{a+b\pm\sqrt{(a-b)^2-4}}{2} \end{pmatrix}$$

Here is a family of matrices satisfying the Bezout identity

$$\left(\begin{array}{cc} x+2 & kx^2+2kx+2x+3\\ 1 & kx+2 \end{array}\right)$$

*Remark* 3.33. If M is a matrix as in the lemma, then M and  $M^{-1}$  both map interlacing polynomials to interlacing polynomials. (See p. 83.) This is different from linear transformations on *one* polynomial, where the only linear transformations T for which T and T<sup>-1</sup> map **P** to itself are affine transformations.

Next we show that all elements of  $SL_2(\mathbb{R})$  can be realized by evaluating polynomials.

**Lemma 3.34.** If  $\begin{vmatrix} a & b \\ c & d \end{vmatrix} = 1$  then there are f, g, h,  $k \in \mathbf{P}(n)$  such that

$$\begin{vmatrix} f & g \\ h & k \end{vmatrix} = 1 \qquad \begin{pmatrix} f(0) & g(0) \\ h(0) & k(0) \end{pmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

Proof. Define

S is the set of all  $\begin{pmatrix} f & g \\ h & k \end{pmatrix}$  such that  $\begin{vmatrix} f & g \\ h & k \end{vmatrix} = 1$  and all polynomials in the same row or column interlace.

Note that if  $\begin{vmatrix} f & g \\ h & k \end{vmatrix} = 1$  and polynomials in one row or column interlace then  $\begin{pmatrix} f & g \\ h & k \end{pmatrix} \in S$ . We now claim that multiplication by  $\begin{pmatrix} \alpha & 0 \\ \beta & 1/\alpha \end{pmatrix}$  and  $\begin{pmatrix} \alpha & \beta \\ 0 & 1/\alpha \end{pmatrix}$  map S to itself. For instance,

$$\begin{pmatrix} \alpha & 0 \\ \beta & 1/\alpha \end{pmatrix} \begin{pmatrix} f & g \\ h & k \end{pmatrix} = \begin{pmatrix} \alpha f & \alpha g \\ \beta f + h/\alpha & \beta g + k/\alpha \end{pmatrix}$$

The resulting matrix has determinant one, and one interlacing row, so it is in S.

From Corollary 3.30 we know that S is non-zero, so we choose  $\begin{pmatrix} f & g \\ h & k \end{pmatrix}$  in S. The element  $\begin{pmatrix} f(0) & g(0) \\ h(0) & k(0) \end{pmatrix}$  is in  $SL_2(\mathbb{R})$ . Since  $SL_2(\mathbb{R})$  is generated by upper and lower triangular matrices as above, all elements of  $SL_2(\mathbb{R})$  have the desired representation.

Following the usual matrix representation of the Euclidean algorithm we get a unique representation for interlacing polynomials.

**Lemma 3.35.** If  $f \leq g$  have positive leading coefficients then there are unique positive  $a_1, \ldots, a_n$  and positive  $\alpha$  such that

$$\begin{pmatrix} f\\g \end{pmatrix} = \alpha \begin{pmatrix} a_n x + b_n & -1\\1 & 0 \end{pmatrix} \cdots \begin{pmatrix} a_1 x + b_1 & -1\\1 & 0 \end{pmatrix} \begin{pmatrix} 1\\0 \end{pmatrix}$$

*Proof.* Simply note that if f = (ax + b)g - r then

$$\begin{pmatrix} f \\ g \end{pmatrix} = \begin{pmatrix} ax+b & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} g \\ r \end{pmatrix}$$

## **3.6** Integrating families of polynomials

We can extend the concept of a mutually interlacing sequence  $f_1, \ldots, f_n$  of polynomials to an interlacing family  $\{f_t\}$ , where t belongs to some interval. Most results of section 3.3 have straightforward generalizations, where summation is replaced by integration, and totally positive matrices are replaced by totally positive functions.

We first introduce a quantitative measure of root separation, and use it to integrate families of polynomials. If  $f \in \mathbf{P}$ , then  $\delta(f)$  is the minimum distance between roots of f. If f has repeated roots then  $\delta(f) = 0$ , and so f has all distinct roots if and only if  $\delta(f) > 0$ . It is easy to see that an equivalent definition is

 $\delta(f) = \text{ largest t such that } f(x) \leq f(x+t)$ 

It easily follows from the definition of  $\delta(f)$  that

$$0 \leqslant s \leqslant t \leqslant \delta(f) \implies f(x+s) \underline{\ll} f(x+t)$$
(3.6.1)

This last property is the motivation for the following definitions.

**Definition 3.36.** Suppose that I is an interval, and for each  $t \in I$  we have a polynomial  $f_t(x)$  with all real roots such that the assignment  $t \mapsto f_t$  is continuous. We say  $\{f_t\}$  is a *continuous family of polynomials*. If  $f_t \leq f_s$  for every  $t \leq s$  in I, then  $\{f_t\}$  is an *interlacing family*. The *interval of interlacing at a point* v, written  $\rho(f_t, v)$ , is the largest  $\epsilon$  such that  $f_v \leq f_s$  for all  $s \in (v, v + \epsilon)$ . If there is no such  $\epsilon$  we set  $\rho(f_t, v) = 0$ . A continuous family of polynomials  $\{f_t\}$  is *locally interlacing* if  $\rho(f_t, v)$  is positive for all v in the interior of I. The interval of interlacing of a locally interlacing family,  $\rho(f_t)$ , is the largest  $\epsilon$  such that if  $s, v \in I$  and  $v < s < v + \epsilon$ , then  $f_v \leq f_s$ . If there is no such  $\epsilon$  then the interval of interlacing is 0.

The following is a generalization of Corollary 3.6.

**Proposition 3.37.** Suppose  $\{f_t\}$  is a continuous family of interlacing polynomials on [a, b], and that w(t) is a positive function on [a, b]. The polynomial

$$p(\mathbf{x}) = \int_{a}^{b} f_{t}(\mathbf{x}) w(t) dt \qquad (3.6.2)$$

*has all real roots, and*  $f_a \leq p \leq f_b$ .

*Proof.* Choose  $a = t_0 < t_1 < \cdots < t_n = b$ , and consider the function

$$s(\mathbf{x}, \mathbf{t}) = f_{t_0}(\mathbf{x})w(\mathbf{t})\chi_{[t_0, t_1]} + \dots + f_{t_{n-1}}(\mathbf{x})w(\mathbf{t})\chi_{[t_{n-1}, t_n]}$$

where  $\chi_{[c,d]}$  is the characteristic function of the interval [c, d]. It suffices to show that the polynomial  $\int_a^b s(x, t) dt$  has all real roots, since (3.6.2) is the limit of such functions. Note that

$$\int_{a}^{b} s(x,t) dt = f_{t_{0}} \cdot \int_{t_{0}}^{t_{1}} w(t) dt + \dots + f_{t_{n-1}} \cdot \int_{t_{n-1}}^{t_{n}} w(t) dt$$

Each of the coefficients of the polynomials  $f_{t_i}$  is positive, and the set of polynomials  $f_{t_0}, \ldots, f_{t_n}$  is mutually interlacing since  $\{f_t\}$  is an interlacing family, so  $\int_a^b s(x, t) dt$  has all real roots by Lemma 3.4.

**Example 3.38.** Since  $\delta(\underline{x})_n = 1$ , if we let w(t) = 1 then  $\int_0^1 (\underline{x} + \underline{t})_{(n)} dt$  is in **P**. More generally, for any w(t) that is positive on (0, 1) we have

$$(\underline{\mathbf{x}})_{n} \leq \int_{0}^{1} (\underline{\mathbf{x}} + \underline{\mathbf{t}})_{(n)} w(\underline{\mathbf{t}}) d\underline{\mathbf{t}} \leq (\underline{\mathbf{x}} + \underline{\mathbf{1}})_{(n)}.$$

**Example 3.39.** Given any mutually interlacing sequence of polynomials we can find an interlacing family that contains it. Suppose that  $f_1 \leq \cdots \leq f_n$  is a mutually interlacing family. Define an interlacing family on (1, n) such that  $F_i = f_i$  by

$$F_t = (i+1-t)f_i + (t-i)f_{i+1} \text{ for } i \leqslant t \leqslant i+1.$$

We can easily compute the integral

$$\begin{split} \int_{1}^{n} F_{t} dt &= \sum_{i=1}^{n-1} \int_{i}^{i+1} (i+1-t) f_{i} + (t-i) f_{i+1} dt \\ &= \sum_{i=1}^{n-1} (\frac{1}{2} f_{i} + \frac{1}{2} f_{i+1}) \\ &= \frac{1}{2} f_{1} + f_{2} + \dots + f_{n-1} + \frac{1}{2} f_{n} \end{split}$$

Here is a similar example, where the sequence of mutually interlacing polynomials is determined by the roots of a fixed polynomial. Suppose that  $f(x) = (x - r_1) \cdots (x - r_n)$  where  $r_1 \leq \cdots \leq r_n$ , and let  $f_i = f/(x - r_i)$ . We know that  $f_n \leq \cdots \leq f_1$  is a mutually interlacing sequence. Define an interlacing family on  $(r_1, r_n)$  by

$$F_t = \frac{f(x)}{(x - r_i)(x - r_{i+1})} \left( x - r_i - r_{i+1} + t \right) \text{ on } r_i \leqslant t \leqslant r_{i+1}$$

This family satisfies  $F_{r_i} = f_i$ . The integral is equal to

$$\begin{split} \int_{r_1}^{r_n} F_t \, dt &= \sum_{i=1}^{n-1} \int_{r_i}^{r_{i+1}} \frac{f(x)}{(x-r_i)(x-r_{i+1})} \left( x - r_i - r_{i+1} + t \right) \\ &= \sum_{i=1}^{n-1} (r_{i+1} - r_i) \, \frac{f(x)}{(x-r_i)(x-r_{i+1})} (x - (r_i + r_{i+1})/2) \end{split}$$

The integral interlaces f, since the integral is a positive linear combination of polynomials interlacing f.

In neither of these examples is  $F_t$  differentiable; the constructions involve piecewise linear functions. In Lemma 11.18 we will find a differentiable family for the sequence in the second example.

The next result is the continuous analog of Lemma 3.14. The proof is omitted.

**Lemma 3.40.** Suppose that  $f_t$  and  $g_t$  are interlacing families on [a, b]. If

$$\begin{array}{l} 1. \ f_t(x) \ll g_t(x) \text{ for all } a \leqslant t \leqslant b \\ 2. \ \left| \begin{array}{c} f(s) & f(t) \\ g(s) & g(t) \end{array} \right| < 0 \text{ for } a \leqslant s < t \leqslant b \end{array}$$

then

$$\int_a^b f_t(x) \, dt \underline{\ll} \int_a^b g_t(x) \, dt$$

The conclusion of Proposition 3.37 was only that p weakly interlaced  $f_a$  and  $f_b$ , since the limit of interlacing polynomials is only weakly interlacing polynomials. The next lemma is used to deduce strict interlacing for integrals.

**Lemma 3.41.** Suppose  $\{f_t\}$  is a continuous family of interlacing polynomials on [a, b], and that w(t) is a positive function on [a, b]. Choose  $c_1, c_2$  such that  $a < c_1 < c_2 < b$  and consider the the polynomials

$$p_1 = \int_a^{c_1} f_t(x) w(t) dt$$
$$p_2 = \int_{c_2}^b f_t(x) w(t) dt$$

*Then*,  $f_a \underline{\ll} p_1 \underline{\ll} p_2 \underline{\ll} f_b$ .

*Proof.* From the previous theorem and the hypothesis we know that

 $f_{\mathfrak{a}} \underline{\ll} \mathfrak{p}_1 \underline{\ll} f_{c_1} \underline{\ll} f_{c_2} \underline{\ll} \mathfrak{p}_2 \underline{\ll} f_b$ 

Since  $\{f_t\}$  is interlacing on [a, b], we see that  $p_1 \ll p_2$ .

We can integrate locally interlacing families to get new ones. We first hold the interval constant.

**Proposition 3.42.** *If*  $\{f_t\}$  *is a locally interlacing family on* [a, b]*,* w(x) *positive on* [a, b]*, and*  $r \leq \rho(f)/2$ *, then* 

$$g_{s}(x) = \int_{s}^{s+r} f_{t}(x)w(t) dt$$

*is a locally interlacing family on* [a, b - r] *with*  $\rho(g_s) \ge r$ .

*Proof.* Certainly all members  $g_s$  have roots, so it remains to show that there is local interlacing. Consider  $g_s$  and  $g_{s+\epsilon}$ , where  $0 < \epsilon \leq r$ . By the previous lemma, the three integrals

$$p_{1} = \int_{s}^{s+\epsilon} f_{t}(x)w(t) dt$$
$$p_{2} = \int_{s+\epsilon}^{s+r} f_{t}(x)w(t) dt$$
$$p_{3} = \int_{s+r}^{s+r+\epsilon} f_{t}(x)w(t) dt$$

satisfy  $p_1 \underline{\ll} p_2 \underline{\ll} p_3$  and  $p_1 \underline{\ll} p_3$ . Since  $g_s = p_1 + p_2$ , and  $g_{s+\epsilon} = p_2 + p_3$ , it follows that  $g_s \underline{\ll} g_{s+\epsilon}$ .

Next, we vary the domain of integration.

**Lemma 3.43.** If  $\{f_t(x)\}$  is a locally interlacing family on  $\mathbb{R}$ , w(x) a non-negative function, and

$$g_s(x) = \begin{cases} \int_0^s f_t(x)w(t) dt & s > 0\\ f_0(x) & s = 0 \end{cases}$$

*then*  $\{g_s\}$  *is a locally interlacing family with*  $\rho(g) \ge \rho(f)$ *.* 

*Proof.* We need to show that  $g_u \leq g_v$  for  $0 \leq u \leq v \leq \rho(f)$ . If we consider an approximating sum for  $g_v$  then all the terms are mutually interlacing, which easily implies the result.

If the weight is a function of two variables, then integrating a locally interlacing family against the weight gives a new family whose members all have roots. However, we need an extra condition to guarantee that it is locally interlacing, and total positivity is what we need.

**Lemma 3.44.** Suppose that  $\{f_t\}$  is a family of interlacing polynomials on [a, b], and u(x), w(x) are positive on [a, b]. Set

$$p_{u} = \int_{a}^{b} f_{t}(x) u(t) dt$$
$$p_{w} = \int_{a}^{b} f_{t}(x) w(t) dt.$$

 $\textit{If} \left|\begin{smallmatrix} \mathfrak{u}(s) & \mathfrak{u}(t) \\ \mathfrak{w}(s) & \mathfrak{w}(t) \end{smallmatrix}\right| \textit{ is positive for all } a \leqslant s < t \leqslant b, \textit{ then } p_u \underline{\ll} p_w$ 

*Proof.* For any  $a < t_1 < \cdots < t_n < b$ , the approximations to  $p_u$  and  $p_w$  are

$$\sum_{i=1}^{n-1} \left( \int_{t_i}^{t_{i+1}} u(t) \, dt \right) f_{t_i}$$
$$\sum_{i=1}^{n-1} \left( \int_{t_i}^{t_{i+1}} w(t) \, dt \right) f_{t_i}$$

Now since

$$\left( \int_{t_{i}}^{t_{i+1}} w(t) \, dt \right) \left( \int_{t_{j}}^{t_{j+1}} u(t) \, dt \right) - \left( \int_{t_{j}}^{t_{j+1}} w(t) \, dt \right) \left( \int_{t_{i}}^{t_{i+1}} u(t) \, dt \right) = \int_{t_{i}}^{t_{i+1}} \int_{t_{j}}^{t_{j+1}} (w(t)u(s) - u(t)w(s)) \, dt \, ds > 0$$

and the sequence of polynomials  $f_{t_1}, \ldots, f_{t_n}$  is mutually interlacing, the result follows from Theorem 3.12. Note that we can allow *w* and *u* to have infinitely many zeros on [a, b] since all the integrals will still be positive.

**Definition 3.45.** A function w(s, t) is *strongly totally positive of order* 2 (or STP<sub>2</sub>) on an interval [a, b] if for all t < t', s < s' in [a, b] the determinant

and all its entries are positive. For example,  $e^{xy}$  is STP<sub>2</sub>.

**Corollary 3.46.** Suppose that  $\{f_t\}$  is an interlacing family on [a, b], and that w(s, t) is STP<sub>2</sub>. The family of polynomials

$$g_s = \int_a^b f_t(x) w(t,s) \, dt$$

is a interlacing family.

*Proof.* The lemma shows that it is locally interlacing, and it follows that it is an interlacing family since  $f_a \leq g_s \leq f_b$ .

As an example, consider the family

$$g_t(x) = \int_0^1 e^{ty} (\underline{x+y})_n \, dy$$

Since  $\delta(\underline{x} + \underline{y})_n = 1$  and  $e^{ty}$  is STP<sub>2</sub> we know that  $g_t$  is an interlacing family. Since  $g_t$  is defined for all t, we have an interlacing family on  $\mathbb{R}$ .

#### 3.7 Families, root zones, and common interlacings

In this section we consider families of polynomials that are indexed by a set S. We say that a family  $\{f_t\}_{t\in S}$  has a common interlacing if there is a  $g \in \mathbf{P}$  such that  $f_t \leq g$  for all  $t \in S$ . The basic question is

What properties do we need in order to have a common interlacing?

Here's an example. Consider the family  $\{F_t\}_{t \ge 0}$  where  $F_t(x) = f(x) + tg(x)$ and deg(f) = deg(g) + 1. If  $F_t \in \mathbf{P}$  for all positive t then there is a common interlacing. This is Proposition 1.35.

It is useful to consider all the zeros of a family. Suppose that  $\{f_t\}_{t \in S}$  is a family of polynomials of degree n. The k'th *root zone* is

 $\Delta_{k} = \{ w \in \mathbb{R} \mid \exists t \in S \& w \text{ is the k'th largest root of } f_{t} \}$ 

Since we assume that families are continuous, it follows that if S is connected then all the root zones are intervals. The following is elementary, but points out how root zones can be useful.

**Lemma 3.47.** If a family F consists of polynomials in P of degree n, and all root zones  $\Delta_1, \ldots, \Delta_n$  are disjoint then F has a common interlacing. The family has a common interlacing iff the intersection of any two root zones has at most one element.

*Proof.* Pick the roots of a common interlacing to lie between consecutive root zones.  $\Box$ 

**Corollary 3.48.** *If any two polynomials of a family have a common interlacing, then the family has a common interlacing.* 

*Proof.* The hypotheses imply that all elements in  $\Delta_k$  are less than or equal to all elements of  $\Delta_{k+1}$ .

**Example 3.49.** The root zones of the family  $\{f + tg\}_{t \ge 0}$  mentioned above have a simple description. Suppose that f and g have degree n, and let  $roots(f) = (r_i)$  and  $roots(g) = (s_i)$ . Since f, g have a common interlacing, it follows that the root zones are

$$\Delta_{k} = \left| \min(\mathbf{r}_{k}, \mathbf{s}_{k}), \max(\mathbf{r}_{k}, \mathbf{s}_{k}) \right|.$$

If all members of the family  $\{f + \alpha g\}_{\alpha \ge 0}$  have all real roots then the family has a common interlacing. We can rephrase this in a way that leads to an important generalization:

If  $\beta^2 f + g \in \mathbf{P}$  for all real  $\beta$  then  $\{\beta^2 f + g\}_{\beta \in \mathbb{R}}$  has a common interlacing.

The next lemma adds a linear term.

**Lemma 3.50.** Suppose that f, g, h are polynomials with positive leading coefficients such that deg(f) = deg(g) = deg(h) + 1

If  $\beta^2 f + \beta h + g \in P$  for all real  $\beta$  then  $\{\beta^2 f + \beta h + g\}_{\beta \in \mathbb{R}}$  has a common *interlacing*.

*Proof.* Define  $F(x, \beta) = \beta^2 f + \beta h + g$ . If deg(f) = n then for each  $\beta$  the polynomial  $F(x, \beta)$  has exactly n roots, since  $F(x, \beta)$  has degree n for all  $\beta$ . If  $r_i(\beta)$  is the i'th largest root of  $F(x, \beta)$  then the continuous function  $\beta \mapsto r_i(\beta)$  is called the i'th solution curve.

The i'th solution curve meets the x-axis in the i'th largest root of g since F(x, 0) = g. In addition,  $\lim_{|\beta| \to \infty} r_i(\beta)$  is the i'th largest root of f. Thus, the i'th solution curve is asymptotic to the vertical line through the i'th largest root of f, and meets the x-axis in the i'th largest root of g.

Consideration of the geometry will finish the proof. For any  $x_0$  there are at most two points on the graph with x-coordinate  $x_0$ . There can be no vertical line that meets two solution curves, for then there would be at least four intersections on some vertical line. See Figure 3.2, where the dashed lines are the vertical asymptotes. Thus, the root zones are separated, and we can apply Lemma 3.47.



Figure 3.2: Impossible solution curves

Corollary 3.51. Suppose that f, h are two polynomials satisfying

- 1.  $f \in P$  has positive leading coefficient.
- 2. h has degree less than n, and does not necessarily have all real roots.
- 3.  $f + \alpha h \in \mathbf{P}$  for  $0 \leq \alpha \leq 1$

*then*  $\{f + \alpha h\}_{0 \le \alpha \le 1}$  *has a common interlacing.* 

Proof. Consider the following

$$\begin{aligned} f + \alpha h \in \mathbf{P} & \text{if } |\alpha| \leq 1 \\ f + \frac{2\beta}{\beta^2 + 1} h \in \mathbf{P} & \text{if } \beta \in \mathbb{R} \text{ since } |\frac{2\beta}{\beta^2 + 1}| \leq 1 \\ 3^2 f + 2\beta h + f \in \mathbf{P} & \text{if } \beta \in \mathbb{R} \end{aligned}$$

We can now apply the lemma.

# 3.8 Two by two matrices preserving interlacing

When does a two by two matrix with polynomial entries preserve interlacing? First, a precise definition.

**Definition 3.52.** Suppose that M is a two by two matrix with polynomial entries, and M  $\begin{pmatrix} f \\ a \end{pmatrix} = \begin{pmatrix} r \\ s \end{pmatrix}$ . Then

- M preserves interlacing if  $f \leq g$  or  $f \ll g \implies r \leq s$  or  $r \ll s$ .
- M preserves interlacing for  $\lt$  if  $f \lt g \implies r \lt s$  or  $r \ll s$ .
- M preserves interlacing for  $\ll$  if  $f \ll g \implies r \lt s$  or  $r \ll s$ .
- M weakly preserves interlacing if all interlacings are  $\leq$  or  $\leq$ .

A number of earlier results can be cast as matrix statements. Assume that  $\alpha$ ,  $\beta$  are positive.

$f \longleftarrow g \implies \alpha f \longleftarrow \beta g$	$\left( \begin{smallmatrix} \alpha & 0 \\ 0 & \beta \end{smallmatrix} \right)$ preserves interlacing.
$f \longleftarrow g \implies xf \longleftarrow xg$	$\begin{pmatrix} x & 0 \\ 0 & x \end{pmatrix}$ weakly preserves interlacing.
$f \lessdot g \implies xf - \alpha g \lessdot \beta f$	$\begin{pmatrix} x & -\alpha \\ \beta & 0 \end{pmatrix}$ preserves interlacing for $\ll$ .
$f \lessdot g \implies xg + \alpha f \lessdot \beta g$	$\begin{pmatrix} \alpha & x \\ 0 & \beta \end{pmatrix}$ preserves interlacing for $\ll$ .

We can combine these matrices using the following lemma:

**Lemma 3.53.** Suppose that u, v, w are vectors of length two. If  $\begin{pmatrix} u \\ v \end{pmatrix}$  and  $\begin{pmatrix} u \\ w \end{pmatrix}$  preserve interlacing, then for any positive  $\alpha, \beta$  the matrix  $\begin{pmatrix} u \\ \alpha v+\beta w \end{pmatrix}$  preserves interlacing. If one of them weakly preserves interlacing, then the conclusion still holds.

*Proof.* Let  $f \leftarrow g$ ,  $\phi = \begin{pmatrix} f \\ g \end{pmatrix}$ ,  $\begin{pmatrix} u \\ v \end{pmatrix} \phi = \begin{pmatrix} r \\ s \end{pmatrix}$  and  $\begin{pmatrix} u \\ w \end{pmatrix} \phi = \begin{pmatrix} r \\ t \end{pmatrix}$ . Then  $\begin{pmatrix} u \\ \alpha v + \beta w \end{pmatrix} \phi = \begin{pmatrix} r \\ \alpha s + \beta t \end{pmatrix}$ . Since  $r \leftarrow s$  and  $r \leftarrow t$  and at least one of these is strict, the result follows by adding interlacings.

**Corollary 3.54.** *Suppose that*  $\alpha$ *,*  $\beta$  *are positive.* 

- 1.  $\begin{pmatrix} \alpha & \alpha \\ 0 & \beta \end{pmatrix}$  preserves interlacing.
- 2.  $\begin{pmatrix} x & 0 \\ \alpha & x \end{pmatrix}$  preserves interlacing.
- 3.  $\begin{pmatrix} x & -\alpha \\ 0 & x \end{pmatrix}$  preserves interlacing.
- 4.  $\begin{pmatrix} x & -\alpha \\ \beta & x \end{pmatrix}$  preserves interlacing.

*Proof.* The two matrices on the left satisfy the conditions of the lemma, so the matrix on the right preserves interlacing:

$$\begin{pmatrix} \alpha & 0 \\ 0 & \beta \end{pmatrix} & \begin{pmatrix} 0 & x \\ 0 & \beta \end{pmatrix} \implies \begin{pmatrix} \alpha & x \\ 0 & \beta \end{pmatrix} \\ \begin{pmatrix} x & 0 \\ 0 & x \end{pmatrix} & \& \begin{pmatrix} x & 0 \\ \alpha & 0 \end{pmatrix} \implies \begin{pmatrix} x & 0 \\ \alpha & x \end{pmatrix}$$

In the next one, the second matrix represents the interlacing  $\exp - \alpha g \leq g$ , which follows from  $\exp \leq xg \leq g$ . We then take the limit as  $\epsilon \to 0^+$ .

$$\begin{pmatrix} x & 0 \\ 0 & x \end{pmatrix} \& \begin{pmatrix} 0 & \epsilon x - \alpha \\ 0 & x \end{pmatrix} \Longrightarrow \begin{pmatrix} x & -\alpha \\ 0 & x \end{pmatrix}$$

The next one follows from previous ones:

$$\begin{pmatrix} x & -\alpha \\ \beta & 0 \end{pmatrix} \& \begin{pmatrix} x & -\alpha \\ 0 & x \end{pmatrix} \implies \begin{pmatrix} x & -\alpha \\ \beta & x \end{pmatrix}$$

The condition that a matrix preserves interlacing is not much more than invertibility. Note that the signs of the leading coefficients are not relevant.

**Lemma 3.55.** If the two by two matrix M is invertible for all x, and there are polynomials r < s and u < v such that  $M({s \atop s}) = ({u \atop v})$  then M preserves interlacing for <.

*Proof.* Assume that there is an  $\alpha$  such that  $u(\alpha) = v(\alpha) = 0$ . Since M evaluated at  $\alpha$  is invertible, it follows that  $r(\alpha) = s(\alpha) = 0$ . Since r and s have no roots in common, it follows that u and v also have no roots in common. Thus, whatever the order the zeros of u and v, it is the same order for all u and v. Since there is one example where they interlace, they always interlace.

**Proposition 3.56.** Suppose that  $M = \begin{pmatrix} f & g \\ h & k \end{pmatrix}$  is a matrix of polynomials with positive leading coefficients. M preserves interlacing if and only if

- 1.  $f \longrightarrow g$ ,  $h \longrightarrow k$ , and both interlacings are strict.
- 2. The determinant  $\begin{vmatrix} f & g \\ h & k \end{vmatrix}$  is never zero.

*Proof.* Assume that M preserves interlacing, and let  $M(\begin{smallmatrix} r \\ s \end{smallmatrix}) = (\begin{smallmatrix} u \\ v \end{smallmatrix})$ . If  $r \ll s$  then  $fr + gs \in \mathbf{P}$ . It follows from Lemma 1.38 that  $g \leq f$ . If g and f had a common factor, then u and v would have a common factor. Thus  $g \ll f$ . Similarly,  $k \ll h$ . Suppose that  $\rho$  is a root of |M|. It follows that  $\begin{pmatrix} u(\rho) \\ v(\rho) \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$  which contradicts the hypothesis that u and v have no roots in common.

Conversely, by Leibnitz (Lemma 1.26) both u = fr + gs and v = hr + ks have all real roots. By the preceding lemma, it suffices to show that there is one example where the roots of u and v interlace. For this purpose, choose r = g and s = f and apply Lemma 1.32.

The cases with  $\triangleleft$  are similar and omitted.

**Corollary 3.57.** If  $f \leq g$  and the absolute value of the leading coefficient of f is greater than the absolute value of the leading coefficient of g then  $\begin{pmatrix} f & -g \\ g & f \end{pmatrix}$  preserves interlacing for  $\leq$ .

*Proof.* Since f < g the determinant is always positive. The following interlaces by Lemma 1.18, and so proves the corollary:

$$\begin{pmatrix} f & -g \\ g & f \end{pmatrix} \begin{pmatrix} f \\ g \end{pmatrix} = \begin{pmatrix} f^2 - g^2 \\ 2 f g \end{pmatrix}$$

**Example 3.58.** If we pick  $f \ll g$  then  $\begin{vmatrix} g & f \\ g' & f' \end{vmatrix} \neq 0$ . Consequently,  $\begin{pmatrix} g & f \\ g' & f' \end{pmatrix}$  preserves interlacing. For instance,  $\begin{pmatrix} 2x & x^2 \\ 1 & 2x \end{pmatrix}$  weakly preserves interlacing.

**Example 3.59.** If we expand  $(x+\iota)^n$  we can write it as  $f_n(x) + g_n(x)\iota$  where  $f_n$  and  $g_n$  have all real coefficients. Multiplication by  $x+\iota$  leads to the recurrence

$$\begin{pmatrix} x & -1 \\ 1 & x \end{pmatrix} \begin{pmatrix} f_n \\ g_n \end{pmatrix} = \begin{pmatrix} f_{n+1} \\ g_{n+1} \end{pmatrix}$$

By Corollary 3.54 it follows that  $f_n \leq g_n$ . This is an elementary fact about polynomials whose roots all have negative imaginary part. See Chapter 20.

**Example 3.60.** Here is another example where matrix multiplication naturally arises (see Theorem 7.64). Suppose that f(x) is a polynomial. Write  $f(x) = f_e(x^2) + xf_o(x^2)$ . The even part of f is  $f_e(x)$ , and the odd part is  $f_o(x)$ . For example, consider powers of x + 1:

	$(x + 1)^3$	$(x + 1)^4$	$(x + 1)^5$
even part	3x + 1	$x^2 + 6x + 1$	$5x^2 + 10x + 1$
odd part	x + 3	4x + 4	$x^2 + 10x + 5$

It's clear from the definition that if  $g(x) = (x + \alpha) f(x)$  then

$$\begin{split} g(x) &= \alpha \, f_e(x^2) + x^2 \, f_o(x^2) + x \, f_e(x^2) + \alpha \, x \, f_o(x^2) \\ g_e(x) &= \alpha \, f_e(x) + x f_o(x) \\ g_o(x) &= f_e(x) + \alpha \, f_o(x) \end{split}$$

which we can express as the matrix product

$$\begin{pmatrix} \alpha & x \\ 1 & \alpha \end{pmatrix} \begin{pmatrix} f_e \\ f_o \end{pmatrix} = \begin{pmatrix} g_e \\ g_o \end{pmatrix}$$

We will see in Example 3.74 that if  $h \leftarrow k$  have positive leading coefficients, and h, k lie in  $\mathbf{P}^{pos}$  then

$$\begin{pmatrix} 1 & x \\ 1 & 1 \end{pmatrix} \begin{pmatrix} h \\ k \end{pmatrix} = \begin{pmatrix} r \\ s \end{pmatrix}$$

implies that  $r \leftarrow s$ . However, the matrix does not preserve arbitrary pairs of interlacing polynomials:

$$\begin{pmatrix} 1 & x \\ 1 & 1 \end{pmatrix} \begin{pmatrix} (x+2)(x-2) \\ (x-1)(x+3) \end{pmatrix} = \begin{pmatrix} -4-3x+3x^2+x^3 \\ -7+2x+2x^2 \end{pmatrix}$$

The latter two polynomials are in **P**, but they don't interlace. However, it is worth noticing that

$$\begin{pmatrix} 1 & x \\ 1 & 1 \end{pmatrix}^2 \begin{pmatrix} (x+2)(x-2) \\ (x-1)(x+3) \end{pmatrix} = \begin{pmatrix} -4 - 10 & x + 5 & x^2 + 3 & x^3 \\ -11 - x + 5 & x^2 + x^3 \end{pmatrix}$$

and the latter two polynomials do interlace. This is a general fact:

**Lemma 3.61.** If  $\alpha > 0$  then the matrix  $\left(\begin{smallmatrix} \alpha & x \\ 1 & \alpha \end{smallmatrix}\right)^2$  weakly preserves interlacing.

*Proof.* First of all,  $\begin{pmatrix} \alpha & x \\ 1 & \alpha \end{pmatrix}^2 = \begin{pmatrix} x + \alpha^2 & 2\alpha x \\ 2\alpha & x + \alpha^2 \end{pmatrix}$ . We notice that

$$\begin{pmatrix} x + \alpha^2 & 2\alpha x \\ 2\alpha & x + \alpha^2 \end{pmatrix} \begin{pmatrix} r \\ s \end{pmatrix} = \begin{pmatrix} (x + \alpha^2)r + 2\alpha xs \\ 2\alpha r + (x + \alpha^2)s \end{pmatrix} = \begin{pmatrix} u \\ v \end{pmatrix}$$

and both u and v are in **P**.

If we follow the proof of Proposition 3.56 then  $\begin{vmatrix} x+\alpha^2 & 2\alpha x \\ 2\alpha & x+\alpha^2 \end{vmatrix} = \begin{vmatrix} \alpha & x \\ 1 & \alpha \end{vmatrix}^2 = (x - \alpha)^2$  has a zero at  $x = \alpha$ , and is otherwise positive. As long as u and v do not have  $\alpha$  for a zero then the argument of the proposition applies. This is the case if and only if  $r(\alpha) + s(\alpha) = 0$ . If we choose  $\epsilon$  so that  $s_{\epsilon}(x) = s(x + \epsilon)$  satisfies  $r \ll s_{\epsilon}$ ,  $r(\alpha) + s_{\epsilon}(\alpha) \neq 0$ , then we can apply the argument of the proposition. Taking limits finishes the proof.

## Polynomials of equal degree

For the rest of this section we will always assume that

$$deg(f) = deg(g) = deg(h) = deg(k)$$
$$\begin{pmatrix} f & g \\ h & k \end{pmatrix} \begin{pmatrix} r \\ s \end{pmatrix} = \begin{pmatrix} u \\ v \end{pmatrix}$$

Lemma 3.66 in the next section shows that in order to get non-trivial results we must assume that f, g, h, k, r, s all have positive leading coefficients. We consider two different ways that interlacing can be preserved.

( $\underline{\ll}$ ) r $\underline{\ll}$ s implies u $\underline{\ll}v$ 

These conditions put constraints on f, g, h, k.

**Lemma 3.62.** Suppose  $\begin{pmatrix} f & g \\ h & k \end{pmatrix} \begin{pmatrix} r \\ s \end{pmatrix} = \begin{pmatrix} u \\ v \end{pmatrix}$ , and consider the two interlacings

$$f + \alpha g \le h + \alpha k \text{ for } \alpha > 0 \tag{3.8.1}$$

$$xf + \alpha g \le xh + \alpha k \text{ for } \alpha > 0 \tag{3.8.2}$$

$$\begin{array}{ccc} r \underline{\ll} \ s \ implies \ \mathfrak{u} \underline{\ll} \ \nu & \Longrightarrow & (3.8.1) \\ r \underline{\lessdot} \ s \ implies \ \mathfrak{u} \underline{\lessdot} \ \nu & \Longrightarrow & (3.8.2) \end{array}$$

*Proof.* These cases are all similar. When  $(\underline{\ll})$  holds then we take r = x and  $s = \alpha x$ . When  $(\underline{\ll})$  holds then we set  $r = x^2$  and  $s = \alpha x$ . Since *s* must have a positive leading coefficient we require that  $\alpha$  is positive. The results follow by computing u, v and dividing the interlacing  $u \leftarrow v$  by x.

If (3.8.1) holds then we see from considering the left side that f and g have a common interlacing. If both (3.8.1) and (3.8.2) then Lemma 2.13 shows that we have true interlacing.

*Remark* 3.63. If we only require that  $f \leq s$  implies that  $u \leq v$  and do not require that the degrees of f, g, h, k are equal then we do not get (3.8.2), and so have more possibilities. In particular note that if  $r \leq s$  and  $\begin{pmatrix} x & -b \\ 1 & 0 \end{pmatrix} \begin{pmatrix} r \\ s \end{pmatrix} = \begin{pmatrix} u \\ v \end{pmatrix}$  then  $u \leq v$ . However, then entries of  $\begin{pmatrix} x & -b \\ 1 & 0 \end{pmatrix}$  do not satisfy (3.8.2).

When (3.8.1) and (3.8.2) hold we can combine the restrictions into one polynomial.

**Lemma 3.64.** *If* (3.8.1) *and* (3.8.2) *hold then for any*  $y, z \in \mathbb{R}$  *we have that* 

$$g + zf + yk + yzh \in P$$

*Proof.* For any positive  $\alpha$  and any  $y \in \mathbb{R}$  we know that

$$g + \alpha f + y(k + \alpha h) \in \mathbf{P}$$
$$g + \alpha x f + y(k + \alpha x h) \in \mathbf{P}$$

Rewriting these yields

$$(g + yk) + \alpha(f + yh) \in \mathbf{P}$$
  
 $(g + yk) + \alpha x(f + yh) \in \mathbf{P}$ 

Since  $g + yk \in P$  and  $f + yh \in P$  we apply Lemma 2.13 to conclude that for any  $z \in \mathbb{R}$ 

$$(g + yk) + z(f + yh) \in \mathbf{P}$$

We will study such polynomials in Chapter 14.

### 3.9 Matrices of polynomials that preserve interlacing

We investigate the set \$ of all matrices M with polynomial entries such that if v is a vector of mutually interlacing polynomials then so is Mv. \$ is a semigroup since it's closed under multiplication, and \$ contains all totally positive matrices. In this section we give a few general constructions.

We do not have a complete analog to Lemma 3.5, but just a necessary condition. Notice that the order of interlacing is reversed.

**Corollary 3.65.** Suppose  $f_1, \ldots, f_n$  is a sequence of polynomials of the same degree and all positive leading coefficients. If for all mutually interlacing polynomials  $g_1 \leq \cdots \leq g_n$  with positive leading coefficients the polynomial  $f_1g_1 + \cdots + f_ng_n$  has all real roots, then the  $f_i$  are mutually interlacing, and

$$f_n \underline{\ll} \cdots \underline{\ll} f_2 \underline{\ll} f_1$$

*Proof.* By taking limits we may assume that all but  $g_i$  and  $g_j$ , i < j, are zero. Now apply Lemma 1.38 to find that  $g_j \leq g_i$ .

We have seen an example of this reverse order in Lemma 1.19. Note that we assumed that all the polynomials had positive leading coefficients. This is not a convenience:

**Lemma 3.66.** Suppose that  $f \ll g$  have positive leading coefficients. If for every  $r \ll s$  we have that  $fs + gr \in \mathbf{P}$ , then r and s have leading coefficients of the same sign.

*Proof.* Choose r = f' and s = -g'. From Lemma 1.55 we know that  $fs + gr = \left| \begin{array}{c} f & g \\ f' & g' \end{array} \right|$  is never zero, and so has no real roots.

We do not have a satisfactory answer to the problem of when mutual interlacing is preserved. We will reconsider this question in § 24.24.6 where we find relationships with polynomials in three variables. Here are some necessary conditions; they are easily seen not to be sufficient.

**Lemma 3.67.** Suppose that  $A = (p_{i,j})$  is a m by n matrix of polynomials of the same degree with the property that for all vectors  $f = (f_1, ..., f_n)$  of mutually interlacing polynomials, the polynomials  $Af = (g_1, ..., g_m)$  are mutually interlacing. Then

• The polynomials in any column of A are mutually interlacing.

$$p_{1,j} \underline{\ll} p_{2,j} \underline{\ll} \dots \underline{\ll} p_{n,j}$$

• The polynomials in any row of A are mutually interlacing in the reverse order:

$$p_{j,1} \underline{\gg} p_{j,2} \underline{\gg} \dots \underline{\gg} p_{j,n}$$

*Proof.* By continuity, we can apply A to the vector (f, 0, ..., 0). This implies that all columns are mutually interlacing. The interlacing in the rows follows from Corollary 3.65.

The next lemma determines some necessary conditions that depend on the coefficients.

**Lemma 3.68.** Suppose that M is a matrix of polynomials of degree n and positive leading coefficients that preserves mutually interlacing polynomials. The matrices of constant terms and of leading coefficients of M are both totally positive<sub>2</sub>.

*Proof.* It suffices to prove this for two by two matrices. So, assume that  $M = \begin{pmatrix} f_1 & f_2 \\ f_3 & f_4 \end{pmatrix}$  where  $f_i$  has constant term  $c_i$  and leading coefficient  $d_i$ . Let  $M_r$  be the result of substituting rx for x in M. If r is positive then  $M_r$  preserves interlacing. Consequently,  $\lim_{r\to 0} M_r = \begin{pmatrix} c_1 & c_2 \\ c_3 & c_4 \end{pmatrix}$  preserves mutually interlacing polynomials. Similarly,  $\lim_{r\to\infty} r^{-n}M_r = \begin{pmatrix} d_1 & d_2 \\ d_3 & d_4 \end{pmatrix}$  preserves mutually interlacing polynomials. The conclusion now follows from Theorem 3.12.

We can use the Leibnitz lemma (Lemma 1.26) to get a some classes of elementary matrices that are analogous to the Jacobi matrices (3.1.1). Choose  $g \leq f$  with positive leading coefficients. Assume that  $r \leq s$  and compute

$$\begin{pmatrix} f & g \\ 0 & f \end{pmatrix} \begin{pmatrix} r \\ s \end{pmatrix} = \begin{pmatrix} fr+gs \\ fs \end{pmatrix}.$$

Lemma 1.26 implies that  $fr + gs \leq fs$ , so  $\begin{pmatrix} f & g \\ 0 & f \end{pmatrix}$  preserves interlacing. Similarly, the matrix  $\begin{pmatrix} g & 0 \\ f & g \end{pmatrix}$  also preserves interlacing. We can generalize this construction.

The first elementary matrix  $E_i$  has f on its diagonal, g in position i, i + 1 for some i, and the remaining entries are 0. The second elementary matrix  $E_i^0$  is not the transpose, but has g on its diagonal, f in position i + 1, i, and the remaining entries are 0. The last elementary matrix is the diagonal matrix fI. The previous paragraph shows that these matrices preserve vectors of mutually interlacing polynomials. For example with n = 4 and i = 2 these elementary matrices are

$$\mathsf{E}_{2} = \begin{pmatrix} \mathsf{f} & 0 & 0 & 0 \\ 0 & \mathsf{f} & \mathsf{g} & 0 \\ 0 & 0 & \mathsf{f} & 0 \\ 0 & 0 & 0 & \mathsf{f} \end{pmatrix} \quad \mathsf{E}_{2}^{0} = \begin{pmatrix} \mathsf{g} & 0 & 0 & 0 \\ 0 & \mathsf{g} & 0 & 0 \\ 0 & \mathsf{f} & \mathsf{g} & 0 \\ 0 & 0 & 0 & \mathsf{g} \end{pmatrix} \quad \mathsf{f} \, \mathsf{I} = \begin{pmatrix} \mathsf{f} & 0 & 0 & 0 \\ 0 & \mathsf{f} & 0 & 0 \\ 0 & 0 & \mathsf{f} & 0 \\ 0 & 0 & 0 & \mathsf{f} \end{pmatrix}$$

With these matrices we can construct examples of transformations that preserve mutually interlacing sequences. For instance, if n is 4 then

$$(E_1 E_2 E_3) (E_1 E_2) E_1 = \begin{pmatrix} f^6 & 3 f^5 g & 3 f^4 g^2 & f^3 g^3 \\ 0 & f^6 & 2 f^5 g & f^4 g^2 \\ 0 & 0 & f^6 & f^5 g \\ 0 & 0 & 0 & f^6 \end{pmatrix}$$

and consequently if  $h_1 \ll h_2 \ll h_3 \ll h_4$  is mutually interlacing then (after factoring out  $f^3$ ) the following are mutually interlacing.

$$\begin{aligned} f^{3}h_{1} + 3 f^{2} gh_{2} + 3 f g^{2}h_{3} + g^{3}h_{4} &\leq f^{3}h_{2} + 2 f^{2} gh_{3} + f g^{2}h_{4} \\ &\leq f^{3}h_{3} + f^{2} gh_{4} \\ &\leq f^{3}h_{4} \end{aligned}$$

In general,

$$(\mathsf{E}_1\cdots\mathsf{E}_r)(\mathsf{E}_1\cdots\mathsf{E}_{r-1})\cdots(\mathsf{E}_2\mathsf{E}_1)\mathsf{E}_1=\left(\mathfrak{g}^{j-\mathfrak{i}}\mathfrak{f}^{d+\mathfrak{i}-j}\binom{d+1-\mathfrak{i}}{\mathfrak{j}-\mathfrak{i}}\right)$$

and consequently the following sequence of polynomials is mutually interlacing

$$\sum_{j=i}^{d} g^{j-i} f^{d+i-j} \binom{d+1-i}{j-i} h_i \qquad i=1,\ldots,d$$

**Example 3.69.** By taking coefficients of polynomials in  $\overline{\mathbf{P}}_3$  we can construct matrices of polynomials such that all rows and columns are interlacing, and all two by two determinants are positive. Here's an example of a  $3 \times 3$  matrix of linear terms

$$\begin{pmatrix} 86x+1 & 625x+76 & 1210x+260 \\ 876x+61 & 3953x+666 & 4707x+1351 \\ 1830x+204 & 5208x+1128 & 3216x+1278 \end{pmatrix}$$

and one with quadratic terms

```
\begin{pmatrix} 644x^2 + 114x + 1 & 9344x^2 + 3019x + 81 & 30483x^2 + 13579x + 466 \\ 7816x^2 + 2102x + 78 & 52036x^2 + 24710x + 1206 & 96288x^2 + 73799x + 4818 \\ 19820x^2 + 7918x + 417 & 72132x^2 + 58752x + 4114 & 59610x^2 + 110860x + 11777 \end{pmatrix}
```

# **3.10** Matrices preserving interlacing in P<sup>pos</sup>

The class of matrices preserving interlacing in  $\mathbf{P}^{\text{pos}}$  contains more than the totally positive matrices. For example, if  $f \leftarrow g$  in  $\mathbf{P}^{\text{pos}}$ , then  $xg \leftarrow f$ . The corresponding matrix  $\begin{pmatrix} 0 & x \\ 1 & 0 \end{pmatrix}$  preserves interlacing in  $\mathbf{P}^{\text{pos}}$ , but not in  $\mathbf{P}$ . In this section we will restrict ourselves to the following matrices:

An NX matrix is a matrix whose entries are either non-negative constants, or positive multiples of x.

We begin by determining which NX vectors map to **P**<sup>pos</sup>.

**Lemma 3.70.** v is an NX 1 by n matrix that maps all mutually interlacing sequences of polynomials in  $\mathbf{P}^{pos}$  to  $\mathbf{P}^{pos}$  if and only if all multiples of x in v occur to the right of all the positive constants.

*Proof.* If a multiple of x is to the left of a positive constant, then we may ignore all other coordinates, and just take v = (x, 1). In this case

$$f = x$$
  $g = x + 1$   $f < g$   $xf + g = x^2 + x + 1 \notin \mathbf{P}$ 

Conversely, assume that  $v = (a_1, ..., a_r, x a_{r+1}, ..., x a_n)$ , and let  $F = (f_1, ..., f_n)$  be mutually interlacing. Define

$$g_1 = \sum_{k=1}^r a_k f_k \qquad g_2 = \sum_{k=r+1}^n a_k f_k$$

Note that  $g_1 \leq g_2$  since the  $f_i$  are mutually interlacing. Consequently,  $xg_2 \leq g_1$  since all  $f_i$  are in  $\mathbf{P}^{pos}$ , and so  $vF = g_1 + xg_2 \in \mathbf{P}^{pos}$ .

Next we look at two by two matrices in NX.

**Lemma 3.71.** The two by two matrices listed below preserve mutually interlacing polynomials in  $P^{\text{pos}}$  if they satisfy

$1:({ightarrow}{bightarrow}{ightarrow}{bightarrow}{ightarrow$	$2: \begin{pmatrix} X & X \\ X & X \end{pmatrix}$	determinant is $\ge 0$
$3: \begin{pmatrix} x & x \\ \bullet & \bullet \end{pmatrix}$	$4: \left(\begin{smallmatrix} \bullet & X \\ \bullet & X \end{smallmatrix}\right)$	determinant is $\leqslant 0$
5:(•X)	$6: \begin{pmatrix} X \\ \bullet \\ X \end{pmatrix}$	no restriction

*Proof.* The • stands for a non-negative constant, and X stands for a positive multiple of x. Since two by two matrices with positive determinant preserve interlacing, the first two matrices preserve interlacing. The third and fourth cases follow from the factorizations

$$\begin{pmatrix} ax & bx \\ c & d \end{pmatrix} = \begin{pmatrix} 0 & x \\ 1 & 0 \end{pmatrix} \begin{pmatrix} c & d \\ a & b \end{pmatrix} \qquad \begin{pmatrix} a & cx \\ b & dx \end{pmatrix} = \begin{pmatrix} c & a \\ d & b \end{pmatrix} \begin{pmatrix} 0 & x \\ 1 & 0 \end{pmatrix}$$

The last two cases are easy and omitted.

**Proposition 3.72.** An NX matrix preserves mutually interlacing sequences of polynomials in  $P^{\text{pos}}$  if and only if

- 1. All two by two submatrices satisfy the conditions of Lemma 3.71.
- 2. All elements that lie above or to the right of a multiple of x are also multiples of x.

*Proof.* It suffices to take two rows, and show that they determine interlacing polynomials. Choose mutually interlacing polynomials  $(f_1, \ldots, f_n)$ . Let  $a_r$  be the rightmost nonnegative constant in the first row, and  $a_s$  the rightmost nonnegative constant in the second row. By the second hypothesis we know that  $r \leq s$ . We denote the two rows by

$$(a_1, a_2, \dots, a_r, x a_{r+1}, \dots, x a_n)$$
  $(b_1, b_2, \dots, b_s, x b_{s+1}, \dots, x b_n)$ 

Define

$$g_{1} = \sum_{i=1}^{r} a_{i} f_{i} \qquad g_{2} = \sum_{i=r+1}^{s} a_{i} f_{i} \qquad g_{3} = \sum_{i=s+1}^{n} a_{i} f_{i}$$
$$h_{1} = \sum_{i=1}^{r} b_{i} f_{i} \qquad h_{2} = \sum_{i=r+1}^{s} b_{i} f_{i} \qquad h_{3} = \sum_{i=s+1}^{n} b_{i} f_{i}$$

We need to show that

$$g_1 + x g_2 + x g_3 \leq h_1 + h_2 + x h_3$$

The interlacings  $h_1 \underline{\ll} g_2$ ,  $g_1 \underline{\ll} h_2$ ,  $g_2 \underline{\ll} g_3$ ,  $g_2 \underline{\ll} h_3$  follow from the mutual interlacing of the  $f_i$ 's. The interlacing  $h_2 \underline{\ll} g_2$  follows from the hypothesis that  $\begin{pmatrix} X & X \\ \bullet & \bullet \end{pmatrix}$  has negative determinant. We now show that  $g_1 + x g_3 \underline{\ll} h_1 + x h_3$ . Note that

$$\mathbf{F} = (\mathbf{x}\,\mathbf{f}_{s+1},\ldots,\mathbf{x}\,\mathbf{f}_n,\mathbf{f}_1,\ldots,\mathbf{f}_r)$$

is a mutually interlacing sequence. The assumptions on the two by two matrices imply that

$$M = \begin{pmatrix} a_{s+1} & \dots & a_n & a_1 & \dots & a_r \\ b_{s+1} & \dots & b_n & b_1 & \dots & b_r \end{pmatrix}$$

is TP<sub>2</sub>. Since  $MF = (g_1 + x g_3, h_1 + x h_3)$ , we find

$$g_1 + x g_3 \leq h_1 + x h_3$$

$$x g_2 \leq h_1 + x h_3$$

$$g_1 + x g_2 + x g_3 \leq h_1 + x h_3$$

$$g_1 + x g_2 + x g_3 \leq h_2$$

Adding these last two yields the conclusion.

Conversely, we know from Lemma 3.70 that all the x's occur to the right of the constants. If (1) is satisfied, and (2) is not, then r < s. It follows from the first part that the interlacing is in the wrong direction.

Suppose that a two by two matrix listed in Lemma 3.71 does not satisfy the condition. In the first three cases we know the interlacing is reversed. In the fourth case we take the matrix to be  $\begin{pmatrix} 1 & x \\ 1 & dx \end{pmatrix}$  and assume that 0 < c < d. Applying this to  $x \leq x$  yields  $x(x + 1) \gg x(c + dx)$  since 0 < c < d.

**Example 3.73.** The matrix below preserves mutually interlacing sequences in  $P^{pos}$  since all the determinants are zero, and the x's satisfy the second condi-

tion.

$$\begin{pmatrix} 1 & x & x & \dots & x & x \\ 1 & 1 & x & \dots & x & x \\ 1 & 1 & 1 & \dots & x & x \\ \vdots & & \ddots & & \vdots \\ 1 & 1 & 1 & \dots & 1 & x \\ 1 & 1 & 1 & \dots & 1 & 1 \end{pmatrix}$$

**Example 3.74.** Suppose that M is a totally positive matrix, and we form a new matrix N by replacing the upper right corner element by x. All the hypothesis are met, so N preserves mutual interlacing in **P**<sup>pos</sup>.

**Example 3.75.** We can construct matrices with more complicated polynomials by multiplying these examples together. For instance, if

	/1	χ	x)		/1	0	x)
$\mathfrak{m}_1 =$	1	1	χ	$\mathfrak{m}_2 =$	1	1	0
	$\backslash 1$	1	1/		$\backslash 1$	2	1/

then the product  $m_1 m_2 m_1$  below preserves mutually interlacing polynomials in  $\mathbf{P}^{pos}$ :

$$\begin{pmatrix} 1+7x & 6x+2x^2 & 3x+5x^2 \\ 3+5x & 1+6x+x^2 & 5x+3x^2 \\ 7+x & 4+4x & 1+7x \end{pmatrix}$$

**Example 3.76.** Here's an example where the matrix is not immediately obvious [70, Theorem 2.8]. Suppose r is a positive integer and that  $f_0(x), \ldots, f_{r-1}(x)$  are mutually interlacing. If

$$\begin{aligned} (1+x+\dots+x^{r-1})\left(f_0(x^r)+xf_1(x^r)+\dots+x^{r-1}f_{r-1}(x^r)\right) \\ &=g_0(x^r)+xg_1(x^r)+\dots+x^{r-1}g_{r-1}(x^r) \end{aligned}$$

then  $g_0(x),\ldots,g_{r-1}(x)$  are also mutually interlacing. This follows from the observation that

/1	χ	χ		. x)	(	$f_0(x)$		$\int g_0(\mathbf{x}) $
1	1	χ		. x		$f_1(x)$		$g_1(\mathbf{x})$
1	1	1		. x		$f_2(x)$		$g_2(x)$
:			·	:		:	=	:
1	1	1	1	l x		$f_{r-2}(x)$		$g_{r-2}(x)$
$\backslash 1$	1	1	1	l 1/	/ \	$f_{r-1}(x)$		$\left\langle g_{r-1}(\mathbf{x})\right\rangle$

**Example 3.77.** Here's a similar example. Choose a positive integer r, and suppose that  $g(x) = (x + \alpha)f(x)$ . Write

$$f(x) = \sum_{k=0}^{r-1} x^{i} f_{i}(x^{r}) \qquad \qquad g(x) = \sum_{k=0}^{r-1} x^{i} g_{i}(x^{r})$$

If  $\{f_0, \dots, f_{r-1}\}$  are mutually interlacing then so are  $\{g_0, \dots, g_{r-1}\}$ . The proof is the observation that

1	α	0	0		x \	$\int f_0(\mathbf{x})$		$\int g_0(\mathbf{x}) $
I	1	α	0		0	$f_1(x)$		$g_1(x)$
I	0	1	α		0		_	:
I	:			•.	:	f(x)		
l	•	0	0	•	• ]	$\int_{r=2}^{r} (x)$		$g_{r-2(x)}$
1		0	0	• • •	$\alpha$	$\langle t_{r-1}(x) \rangle$		$\langle g_{r-1}(x) \rangle$

This example can be used to give another proof of Theorem 7.65.

# 3.11 Linear 2 by 2 matrices preserving interlacing

We are interested in determining when the matrix

$$\begin{pmatrix} a_1 x + b_1 & a_2 x + b_2 \\ a_3 x + b_3 & a_4 x + b_4 \end{pmatrix}$$
(3.11.1)

preserves interlacing in  $\mathbf{P}^{\text{pos}}$ . We give a nice construction due to Liu and Wang [117]. There are constructions for matrices of higher degree that preserve interlacing (See p. 693.).

We begin by specializing to the matrix  $M = \begin{pmatrix} x+\alpha & x \\ 1 & x+b \end{pmatrix}$ . If we apply M to the vector  $(x, \alpha x)$  where  $\alpha > 0$  then the result is a pair of interlacing polynomials. After factoring out x

$$\begin{split} x+a+x\alpha &\leq 1+(b+x)\alpha\\ -a/(1+\alpha) \geqslant (-1-b\,\alpha)/\alpha\\ P=\alpha^2 b+\alpha(b+1-a)+1 \geqslant 0 \quad \text{for all } \alpha \geqslant 0 \end{split}$$

Next, apply M to  $(x^2, \alpha x)$ . Factoring out x yields

$$\mathbf{x}(\mathbf{x} + \mathbf{a} + \alpha) \leq \mathbf{x}(1 + \alpha) + \mathbf{b}\alpha$$

Substituting the root of the second term into the first gives

$$\frac{-b\alpha}{1+\alpha}(\frac{-b\alpha}{1+\alpha}+a+\alpha) \leqslant 0$$
$$R = \alpha^2 + (a+1-b)\alpha + a \ge 0 \quad \text{for all } \alpha \ge 0$$

Note that R and P have the same discriminant  $Q = (a-b)^2 - 2(a+b) + 1$ . The graph of Q = 0 is a parabola, and is given in Figure 3.3.

**Lemma 3.78.**  $\binom{x+a}{1} \xrightarrow{x}{x+b}$  preserves interlacing in  $\mathbf{P}^{\text{pos}}$  if and only (a, b) is either in the parabola, or in the finite region bounded by the axes and parabola. See Figure 3.3.

*Proof.* Q intersects the axes at (1,0) and (0,1) and is negative in the region inside the parabola, so P and R are non-negative for those (a, b). If (a, b) is in the unbounded region below the parabola, then Q is positive, and b + 1 - a is negative. Thus, P has two positive roots, and so is not positive for positive  $\alpha$ . R has two positive roots in the region above the parabola. Thus, if M preserves interlacing then (a, b) can't be in either unbounded region outside the parabola.

Lemma 3.79 shows that all points in the gray region of Figure 3.3 determine matrices preserving interlacing. We can use Example 11.61 to show that all (a, b) in the parabola determine interlacing preserving matrices.



Figure 3.3: The graph of  $(a - b)^2 - 2(a + b) + 1 = 0$ 

**Lemma 3.79.** If a and b are positive, and  $|a - b| \leq 1$  then  $\binom{x+a}{1} \binom{x}{x+b}$  preserves interlacing in  $P^{\text{pos}}$ .

*Proof.* We first claim that  $\begin{pmatrix} x+t & x \\ 1 & x+t \end{pmatrix}$  preserves interlacing for positive t. If we write

$$\begin{pmatrix} x+t & x \\ 1 & x+t \end{pmatrix} \begin{pmatrix} f \\ g \end{pmatrix} = \begin{pmatrix} (x+t)f + xg \\ f + (x+t)g \end{pmatrix}$$

then each of the two terms on the bottom interlaces each of the two terms of the top. Similarly,  $\alpha(f+xg) \leq f+(x+t)g$ , so  $\begin{pmatrix} \alpha & \alpha x \\ 1 & x+t \end{pmatrix}$  preserves interlacing for any positive  $\alpha$ .

Since  $\begin{pmatrix} \alpha & \alpha x \\ 1 & x+t \end{pmatrix}$  and  $\begin{pmatrix} x+t & x \\ 1 & x+t \end{pmatrix}$  have the same second row, the matrix  $\begin{pmatrix} x+t+\alpha & (\alpha+1)x \\ 1 & x+t \end{pmatrix}$  preserves interlacing. Multiplying the first row and the second column by  $1/(\alpha + 1)$ , and replacing x by  $(\alpha + 1)x$  shows that

$$\begin{pmatrix} x + \frac{t+\alpha}{\alpha+1} & x\\ 1 & x + \frac{t}{\alpha+1} \end{pmatrix}$$
(3.11.2)

preserves interlacing. Suppose that  $b + 1 \ge a \ge b$ . If we choose t = b/(b + 1 - a) and  $\alpha = (a - b)/(b + 1 - a)$  then t and  $\alpha$  are both non-negative, and (3.11.2) reduces to  $\begin{pmatrix} x+a & x \\ 1 & x+b \end{pmatrix}$ . The case  $a + 1 \ge b \ge a$  is similar.

We have seen that the criterion for a  $2 \times 2$  matrix of polynomials to preserve interlacing is that the determinant is always positive. If we restrict ourselves to  $\mathbf{P}^{\text{pos}}$  then we only need that the determinant is positive for negative x. The following result is due to Liu and Wang[117].

**Lemma 3.80.** Let  $M = \begin{pmatrix} \ell_1 & q \\ c & \ell_2 \end{pmatrix}$  where c > 0,  $\ell_1, \ell_2 \in P^{pos}(1)$  and  $q \in P^{pos}(2)$ , where  $\ell_1, \ell_2, q$  have positive leading coefficients. If  $|M| \ge 0$  for all  $x \le 0$  then M preserves interlacing in  $P^{pos}$ .

*Proof.* Let  $M(f,g)^t = (F,G)^t$  where f < g in  $\mathbf{P}^{pos}$ . The root  $\alpha$  of  $\ell_1$  is negative, and since the determinant is non-negative at  $\alpha$ , it follows that  $q(\alpha) \leq 0$  and so  $q \leq \ell_1$ . From Leibnitz (p. 16) we know that F and G are in  $\mathbf{P}^{pos}$ .

Since G has positive coefficients and negative roots, |M| is non-negative at the roots of G. In addition, we have  $G = cf + \ell_2 g \leq g$ . The identity

$$cF = \ell_1 G - |M| g$$

shows that F sign interlaces G. Since the degree of F is greater than the degree of G we conclude that  $F \leq G$ .

If q is linear then we get

**Corollary 3.81.** Let  $M = \begin{pmatrix} \ell_1 & \ell_3 \\ c & \ell_2 \end{pmatrix}$  where c > 0 and  $\ell_1, \ell_2, \ell_3 \in P^{pos}(1)$  have positive leading coefficients. If  $|M| \ge 0$  for all  $x \le 0$  then M preserves interlacing in  $P^{pos}$ .

Note that the corollary provides another proof of Lemma 3.78, since  $\begin{vmatrix} x+a & x \\ 1 & x+b \end{vmatrix}$  is (x + a)(x + b) - x, and this determines the same region that is described in the proposition.

Here is a simple consequence.

**Corollary 3.82.**  $\begin{pmatrix} x & x \\ 1 & x+\gamma \end{pmatrix}$  and  $\begin{pmatrix} x+\gamma & x \\ 1 & x \end{pmatrix}$  preserve interlacing in  $\mathbf{P}^{\text{pos}}$  if and only if  $0 \leq \gamma \leq 1$ .

*Proof.* Existence follows from the lemma. If  $\gamma > 1$  then  $(0, \gamma)$  and  $(\gamma, 0)$  lie outside the parabola.

**Lemma 3.83.** Suppose that a, b, c, d are positive, and  $\begin{vmatrix} a & b \\ c & d \end{vmatrix} \ge 0$ .

- 1.  $\begin{pmatrix} a & x+b \\ c & d \end{pmatrix}$  preserves interlacing in  $P^{\text{pos}}$ .
- 2.  $\begin{pmatrix} a & x+b \\ c & x+d \end{pmatrix}$  preserves interlacing  $P^{pos}$  if and only if  $c \ge a$ ,  $d \ge b$ .
- 3.  $\binom{x+a}{c} \binom{x+b}{d}$  preserves interlacing  $P^{pos}$  if and only if  $a \ge b, c \ge d$ .
- $\begin{array}{l} \text{4. } \left(\begin{smallmatrix} x+a & x+b \\ x+c & x+d \end{smallmatrix}\right) \text{ preserves interlacing } \textbf{P}^{\text{pos}} \text{ if and only if} \\ c \geqslant a \geqslant b \text{, and } (c-a)(a-b) \geqslant ad-bc \geqslant 0. \end{array}$

*Proof.* The first one follows from the decomposition  $\begin{pmatrix} 0 & x \\ c & d \end{pmatrix} + \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ . If  $c \ge a$  and  $d \ge b$  then

$$\begin{pmatrix} a & x+b \\ c & x+d \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} a & x+b \\ c-a & d-b \end{pmatrix}$$
shows that  $\begin{pmatrix} a & x+b \\ c & x+d \end{pmatrix}$  preserves interlacing. Conversely, if interlacing is preserved, then applying  $\begin{pmatrix} a & x+b \\ c & x+d \end{pmatrix}$  to  $(1, \alpha)$  yields

$$a + \alpha(x+b) \leq c + \alpha(x+d)$$

Taking roots shows that  $\alpha b + a \leq \alpha d + c$  for all positive  $\alpha$  which implies  $c \geq a$  and  $d \geq b$ .

The third one is similar: sufficiency follows from

$$\begin{pmatrix} x+a & x+b \\ c & d \end{pmatrix} = \begin{pmatrix} a-b & x+b \\ c-d & d \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

and necessity is a consequence of applying  $\begin{pmatrix} x+\alpha & x+b \\ c & d \end{pmatrix}$  to  $(x, \alpha)$ . For the last one, we first compute

$$\begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} x+a & x \\ c-a & \frac{ad-bc}{a-b} \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & a-b \end{pmatrix} \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} x+a & x+b \\ x+c & x+d \end{pmatrix}$$

By Corollary 3.10 we see that  $\begin{pmatrix} x+a & x+b \\ x+c & x+d \end{pmatrix}$  preserves interlacing if and only if  $\begin{pmatrix} x+a & x \\ c-a & \frac{ad-bc}{a-b} \end{pmatrix}$  does. From (3) this is the case if and only if  $c-a \ge \frac{ad-bc}{a-b}$ .

#### 3.12 Polynomial sequences from matrices

A totally positive or NX  $n \times n$  matrix determines sequences of polynomials obeying an n'th order recurrence relation. We first recall how to get sequences satisfying a recurrence relation from a matrix. Suppose that M is a matrix, v is a vector, and the characteristic polynomial of M is

$$a_0 + a_1y + a_2y^2 + \cdots + a_ny^n$$
.

Since M satisfies its characteristic polynomial, we have

$$a_0(v) + a_1(Mv) + a_2(M^2v) + \dots + a_n(M^nv) = 0.$$

If we fix an integer k and define  $w_i$  to be the kth coordinate of  $M^i v$  then the  $w_i$  also satisfy the same recurrence relation

$$a_0 w_0 + a_1 w_1 + a_2 w_2 + \dots + a_n w_n = 0.$$
 (3.12.1)

We now let M be a totally positive matrix, and let v be a matrix of mutually interlacing polynomials. Since  $M^i v = v_i$  is a vector of mutually interlacing polynomials, all the coordinates of  $v_i$  are are in **P**. These polynomials satisfy the recurrence relation (3.12.1). All these polynomials have the same degree but consecutive  $v_i$  do not necessarily interlace.

**Example 3.84.** Consider an example where  $M = \begin{pmatrix} 1 & 2 & 1 \\ 1 & 3 & 3 \\ 1 & 4 & 6 \end{pmatrix}$  and we choose  $v_0$  to be three mutually interlacing polynomials of degree 2.

$$\begin{array}{lll} \nu_0 &= \left( \begin{array}{ccc} (1+x)(4+x) & (2+x)(5+x) & (3+x)(6+x) \end{array} \right) \\ \nu_1 &= \left( \begin{array}{ccc} 2(21+14x+2x^2) & 88+53x+7x^2 & 152+87x+11x^2 \end{array} \right) \\ \nu_2 &= \left( \begin{array}{ccc} 370+221x+29x^2 & 2(381+224x+29x^2) & 2(653+381x+49x^2) \end{array} \right) \end{array}$$

If we just consider the second column, and observe that the characteristic polynomial is  $1 - 12y + 10y^2 - y^3$ , then we conclude that all polynomials defined by the sequence below are in  $\mathbf{P}^{\text{pos}}$ .

$$p_{n+3} = 10p_{n+2} - 12p_{n+1} + p_n$$
$$p_0 = (2+x)(5+x)$$
$$p_1 = 88 + 53x + 7x^2$$
$$p_2 = 2(381 + 224x + 29x^2)$$

We now consider NX matrices. These are more interesting, since the degrees of the polynomials can increase.

**Example 3.85.** We take the matrix  $M = \begin{pmatrix} 1 & x \\ 1 & 1 \end{pmatrix}$ . If we begin with  $v_0 = (x + 1, x + 2)$  then the second column yields that  $p_n \in \mathbf{P}^{pos}$ , where

$$p_{n+2} = 2p_{n+1} + (x-1)p_n$$
  
 $p_0 = x + 2$   
 $p_1 = 2x + 3$ 

**Example 3.86.** We now take the matrix  $M = \begin{pmatrix} 1 & x & x \\ 1 & 1 & x \\ 1 & 1 & 1 \end{pmatrix}$ . If we begin with  $v_0 = (x + 1, x + 2, x + 3)$  then the third column yields that all  $p_n \in \mathbf{P}^{pos}$ , where

$$p_{n+3} = 3p_{n+2} + 3(x-1)p_{n+1} + (x-1)^2p_n$$
  

$$p_0 = x + 3$$
  

$$p_1 = 3x + 6$$
  

$$p_2 = 3x^2 + 14x + 10$$

**Example 3.87.** In this example we let M be the cube of the previous M.

$$M = \begin{pmatrix} 1+7x+x^2 & 6x+3x^2 & 3x+6x^2 \\ 3+6x & 1+7x+x^2 & 6x+3x^2 \\ 6+3x & 3+6x & 1+7x+x^2 \end{pmatrix}$$

Since every row has a quadratic term, the degree of the  $p_n$ 's goes up by two at each step, so consecutive  $p_n$ 's don't interlace. If we again start with  $v_0 = (x + 1, x + 2, x + 3)$  then all  $p_n$  are in  $\mathbf{P}^{pos}$ :

$$p_{n+3} = 3(1 + 7x + x^2)p_{n+2} - 3(x - 1)^4 p_{n+1} + (x - 1)^6 p_n$$
  

$$p_0 = x + 3$$
  

$$p_1 = 5 + 42x + 30x^2 + 4x^3$$
  

$$p_2 = 8 + 260x + 947x^2 + 784x^3 + 181x^4 + 7x^5$$

#### 3.13 Interlacing via the complexes

In this section we show how we can use properties of complex numbers to establish interlacing. Let UHP be the open upper half plane  $\{z \mid \Im(z) > 0\}$ . Note that

$$f \in \mathbf{P} \Leftrightarrow f(\alpha) \neq 0$$
 for all  $\alpha \in \mathsf{UHP}$ .

Our first result connects interlacing and quotients.

**Lemma 3.88.** *Suppose that* f, g *are relatively prime and have positive leading coefficients. The following are equivalent* 

1.  $f \longleftarrow g$ 2.  $\frac{f}{g}: UHP \longrightarrow UHP$ 

Proof. Suppose 1) holds. If we write

$$g = a_0 f + \sum a_i \frac{f}{x - r_i}$$

where  $roots(f) = (r_i)$  then

$$\frac{g}{f}(\alpha) = a_0 + \sum a_i \frac{1}{\alpha - r_i}$$

If  $\alpha \in UHP$  then  $\alpha - r_i \in UHP$ , so  $1/(\alpha - r_i) \in -UHP$ , and therefore  $(g/f)(\alpha) \in -UHP$ . Taking inverses shows that  $(f/g)(\alpha) \in UHP$ .

Conversely, if 2) holds and  $f + tg \notin P$  for some  $t \in \mathbb{R}$  then there is  $\alpha \in UHP$  such that  $f(\alpha) + tg(\alpha) = 0$ . Thus,  $(f/g)(\alpha) \notin UHP$ . Thus f and g interlace, and the previous paragraph shows that the direction must be  $f \leftarrow g$ .

The upper half plane is a cone - it is closed under positive sums. This additivity explains two fundamental results about interlacing.

Lemma 3.89. Suppose that f, g, h, k in P have positive leading coefficients.

- 1. If  $f \leq g$ , h then  $f \leq g + h$ .
- 2. If  $f \leq g$ ,  $h \leq k$  then  $fh \leq fk + gh \leq gk$ .

*Proof.* If  $f \leq g$ , h and  $\sigma$  is in UHP, then  $\frac{g+h}{f}(\sigma) \in -UHP$  since  $\frac{g}{f}(\sigma)$  and  $\frac{h}{f}(\sigma)$  are, and -UHP is a cone. It follows that  $f/(g+h)(\sigma) \in UHP$ .

The second one is similar; it's just the addition of fractions:

$$\frac{g}{f}(\sigma) + \frac{k}{h}(\sigma) = \frac{gh + fk}{fh}(\sigma).$$

Note that the region  $\mathbb{C} \setminus (-\infty, 0)$  is *not* a cone, so we should not expect  $f \stackrel{+}{\sim} g$ , h to imply  $g + h \in \mathbf{P}$ . Indeed, it's false.

All points in the upper half plane can be realized by ratios of interlacing polynomials.

**Lemma 3.90.** If  $\alpha$  and  $\sigma$  are in UHP then there are  $f \leq g$  in **P** with positive leading coefficients such that  $\frac{f(\sigma)}{g(\sigma)} = \alpha$ .

*Proof.* We will show that the set

$$S = \left\{ \begin{pmatrix} f(\sigma) \\ g(\sigma) \end{pmatrix} \mid f \lessdot g \right\}$$

equals UHP, considered as a subset of complex projective space. We know all elements of S lie in UHP. Choose any f < g and let  $f(\sigma)/g(\sigma) = \beta$ . Since  $SL_2(\mathbb{R})$  is transitive on UHP we can find  $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  in  $SL_2(\mathbb{R})$  such that  $M(\beta) = \alpha$ . Then

$$\begin{pmatrix} af+bg\\ cf+dg \end{pmatrix}(\sigma) = M \, \begin{pmatrix} f\\ g \end{pmatrix}(\sigma) = M(\beta) = \alpha$$

Since M has determinant one we know that  $af + bg \lt cf + dg$ .

Matrices preserving interlacing are the same as matrices mapping UHP to UHP. The next lemma provides an alternative proof of Corollary 3.8, and clarifies why there is a simple determinant condition. Recall that if M has all real coefficients then M maps UHP to itself if and only if |M| > 0.

**Lemma 3.91.** *Suppose that* M *is a* 2 *by* 2 *matrix of polynomials with positive leading coefficients. The following are equivalent* 

- 1. M preserves interlacing.
- 2.  $M(\sigma)$  maps UHP to UHP for all  $\sigma \in UHP$ .

*Proof.* Assume that M preserves interlacing, and choose any  $\sigma$  and  $\alpha$  in UHP. By Lemma 3.90 we can find  $f \leq g$  such that  $f(\sigma)/g(\sigma) = \alpha$ . Then

$$\left[\mathsf{M}(\sigma)\right](\alpha) = \left[\mathsf{M}(\sigma)\right] \left(\begin{smallmatrix} \mathsf{f}(\sigma) \\ \mathfrak{g}(\sigma) \end{smallmatrix}\right) = \left[\mathsf{M}\left(\begin{smallmatrix} \mathsf{f} \\ \mathfrak{g} \end{smallmatrix}\right)\right](\sigma) \in \mathsf{UHP}$$

since  $M\begin{pmatrix} f\\g \end{pmatrix}$  is a pair of interlacing polynomials.

Conversely, if  $f \leq g$  and  $M\begin{pmatrix} f \\ g \end{pmatrix} = \begin{pmatrix} F \\ G \end{pmatrix}$  then for all  $\sigma \in UHP$  we know  $\begin{pmatrix} f(\sigma) \\ g(\sigma) \end{pmatrix} \in UHP$ . Thus  $\begin{pmatrix} F(\sigma) \\ G(\sigma) \end{pmatrix} \in UHP$ , which implies that  $F \leq G$ .  $\Box$ 



Figure 3.4: Mutually interlacing polynomials evaluated at  $\sigma$ 

# 3.14 Mutually interlacing polynomials

We will generalize the following three equivalent statements that hold for a real 2 by 2 matrix M to mutually interlacing polynomials and strictly totally positive matrices<sub>2</sub>.

- 1. M has positive determinant.
- 2. M maps the upper half plane to itself.
- 3. M preserves interlacing polynomials.

First of all, mutually interlacing polynomials have a pretty geometric interpretation. Suppose that  $f_1, \ldots, f_n$  are mutually interlacing. For any  $\sigma \in UHP$  we know that  $f_i(\sigma)/f_j(\sigma) \in UHP$  for i < j. In terms of angles this means that the angle from  $f_j(\sigma)$  to  $f_i(\sigma)$  is less than  $\pi$ . It follows that

All the points  $\{f_i(\sigma)\}\$  lie in a half plane, and are , in counterclockwise order,  $f_1(\sigma), \ldots, f_n(\sigma)$ . (Figure 3.4)

A 2 by 2 matrix with real entries preserves the upper half plane iff it preserves interlacing. We know that an n by n matrix preserves interlacing if and only if it is totally positive<sub>2</sub>. We now introduce the structure preserved by strictly totally positive<sub>2</sub> matrices.

$$\mathsf{H}_d^+ = \{(\mathfrak{u}_1, \dots, \mathfrak{u}_d) \ | \ \frac{\mathfrak{u}_i}{\mathfrak{u}_j} \in \mathsf{UHP} \text{ for } 1 \leqslant i < j \leqslant d\}.$$

For instance,

$$\mathsf{H}_3^+ = \{(\mathfrak{u}, \mathfrak{v}, \mathfrak{w}) \mid \frac{\mathfrak{u}}{\mathfrak{v}} \in \mathsf{UHP}, \frac{\mathfrak{u}}{\mathfrak{w}} \in \mathsf{UHP}, \frac{\mathfrak{v}}{\mathfrak{w}} \in \mathsf{UHP}\}$$

The space  $H_2^+$  is a subset of  $P^2(\mathbb{C})$ , and is usually identified with UHP. A particular point in  $H_d^+$  is  $u = (\zeta^{d-1}, \ldots, \zeta, 1)$  where  $\zeta = e^{\pi \iota/d}$ . The characterization of mutual interlacing is the same as for interlacing, and follows easily from the case d = 2.

**Lemma 3.92.** Assume  $(f_1, \ldots, f_d)$  is a sequence of polynomials. The following are equivalent:

- 1.  $(f_1, \ldots, f_d)$  is strictly mutually interlacing.
- 2.  $(f_1(\sigma), \ldots, f_d(\sigma)) \in H_d^+$  for all  $\sigma \in UHP$ .

Let  $\mathcal{T}_d$  be the semigroup of all invertible matrices with non-negative entries for which the determinants of all two by two submatrices are positive. For example,  $\mathcal{T}_2$  consists of all two by two matrices with non-negative entries and positive determinant.

**Lemma 3.93.**  $T_d$  acts transitively on  $H_d^+$ .

*Proof.* We know that  $\mathcal{T}_2$  acts transitively on UHP. Choose two elements  $(\mathfrak{u}_1, \ldots, \mathfrak{u}_d)$  and  $(\nu_1, \ldots, \nu_d)$  of  $H_d^+$ . The block matrix

$$\begin{pmatrix} I & 0 & 0 \\ 0 & M & 0 \\ 0 & 0 & I \end{pmatrix}$$

where M is a 2 by 2 matrix in  $\mathcal{T}_2$  preserves  $H_d^+$ . Since  $\mathcal{T}_2$  is transitive we first carry  $u_1, u_2$  to  $v_1, v_2$ , then leaving the first coordinate fixed we carry  $v_2, u_3$  to  $v_2, v_3$  and so on.

**Lemma 3.94.** For any  $u \in H_d^+$  and  $\sigma \in UHP$  there is a vector f of strictly mutually interlacing polynomials such that  $f(\sigma) = u$ .

*Proof.* Choose any vector g of mutually interlacing polynomials. Since  $g(\sigma) \in H_d^+$  there is an  $M \in T_d$  such that  $M(g(\sigma)) = u$ . Thus  $u = (Mg)(\sigma)$ , so Mg is the desired vector.

**Proposition 3.95.** *Suppose that* M *is a* d *by* d *matrix of positive real numbers. The following are equivalent* 

- 1. M preserves mutually interlacing polynomials.
- 2. M is strictly totally positive<sub>2</sub>.
- 3. M maps  $H_d^+$  to itself.

*Proof.* We have already seen the first two equivalences. Assume M maps  $H_d^+$  to itself. For any  $\sigma \in UHP$  and mutually interlacing  $f_1 \ll f_2 \cdots \ll f_d$  we know that

$$|\mathsf{M} \cdot (\mathsf{f}_1, \dots, \mathsf{f}_d)|(\sigma) = \mathsf{M} \cdot (\mathsf{f}_1(\sigma), \dots, \mathsf{f}_d(\sigma)) \in \mathsf{H}_d^+$$

since  $(f_1(\sigma), \dots, f_d(\sigma)) \in H_d^+$ . Thus, M preserves interlacing, and so is totally positive<sub>2</sub>.

Conversely, suppose that M is totally positive<sub>2</sub>, and  $u = (u_1, \ldots, u_d)$  is in  $H_d^+$ . Choose mutually interlacing  $f = (f_1, \ldots, f_d)$  such that  $f(\sigma) = u$  for some  $\sigma \in UHP$ . Then

$$\mathsf{Mu} = \mathsf{M}(\mathsf{f}(\sigma)) = (\mathsf{Mf})(\sigma) \in \mathsf{H}^+_d$$

since Mf consists of strictly totally interlacing polynomials.

*Remark* 3.96. We give a direct proof that (2) implies (3). It suffices to take M to be the 2 by d matrix  $\begin{pmatrix} a \\ b \end{pmatrix}$ . Assume  $a = (a_i)$ ,  $b = (b_i)$  and  $v = (v_i)$ . We show that  $Mv \in UHP$  by computing the imaginary part of the quotient of Mv.

$$\Im\left(\frac{\sum a_{i}\nu_{i}}{\sum b_{i}\nu_{i}}\right) = \frac{\Im\left(\sum a_{i}\nu_{i}\right)\left(\sum b_{i}\overline{\nu_{i}}\right)}{\left|\sum b_{i}\nu_{i}\right|^{2}}$$

Ignoring the positive denominator, the imaginary part is

$$\Im\left[\sum a_i b_i |v_i|^2 + \sum_{i < j} a_i b_j v_i \overline{v_j} + a_j b_i v_j \overline{v_i}\right] = \sum_{i < j} \begin{vmatrix} a_i & a_j \\ b_j & b_j \end{vmatrix} \Im(v_i \overline{v_j})$$

All the two by two determinants are positive by hypothesis, and  $\Im(\nu_i \overline{\nu_j})$  is positive since  $\nu_i / \nu_j \in UHP$ .

CHAPTER



# **Homogeneous Polynomials**

We begin the study of polynomials in two variables with a very special case: homogeneous polynomials. These results are generally well known, and will be generalized in later chapters.

#### 4.1 Introduction

A homogeneous polynomial of degree n is a polynomial in x, y such that the degree of every term is n. A polynomial f of degree n in one variable determines a homogeneous polynomial of degree n:

$$F(x,y) = y^n f(x/y).$$

The coefficients of F are those of f. If

$$f(x) = a_0 + a_1 x + \dots + a_n x^n$$
(4.1.1)

(4.1.2)

then the homogenized polynomial is

$$F(x, y) = a_0 y^n + a_1 x y^{n-1} + \dots + a_n x^n$$
(4.1.3)

We can recover f from F since f(x) = F(x, 1).

The key relation for us between homogeneous polynomials and polynomials in one variable is that factorization for homogeneous polynomials corresponds to the property of having all real roots.

**Lemma 4.1.** Let F(x, y) be homogeneous, and set f = F(x, 1). We can factor F into linear factors if and only if  $f \in P$ .

*Proof.* If 
$$f = \prod (a_i x + b_i)$$
 then  $F = \prod (a_i x + b_i y)$  and conversely.

The reversal of f, written f<sup>REV</sup>, is

$$a_n + a_{n-1}x + \dots + a_0x^n.$$
 (4.1.4)

We can express  $f^{REV}$  in terms of F since  $f^{REV}(x) = F(1, x)$ . This shows that if f is in **P** then  $f^{REV}$  is also in **P**.

# 4.2 Polar derivatives

Since a polynomial in x and y has derivatives with respect to both x and y we can construct a new derivative operator for polynomials in one variable. From (4.1.3) we find

$$\frac{\partial}{\partial y}F = na_0y^{n-1} + (n-1)a_1y^{n-2}x + \dots + a_nx^{n-1}.$$

If we substitute y = 1, then the right hand side becomes

$$na_0 + (n-1)x + \dots + a_{n-1}x^{n-1}$$
 (4.2.1)

which can be written as

$$n(a_0 + a_1x + \dots + a_nx^n) - x(a_1 + 2a_2x + \dots + nx_{n-1})$$
  
= nf - xf'

This is called the *polar derivative* of f and is written  $\frac{\partial}{\partial y}f$ . The degree of  $\frac{\partial}{\partial y}f$  is n - 1, and the formula for  $\frac{\partial}{\partial y}f$  has an n in it. The polar derivative is *not* not defined on **P**, but only on **P**(n). It is useful to see that the polar derivative can be defined as follows:

Since these operations preserve **P**, the polar derivative is in **P**.

If we assume  $f = \prod (x - a_i)$  then

$$\frac{\partial}{\partial y}f = \sum (-a_i) \frac{f(x)}{x - a_i}$$
(4.2.2)

which is similar to the representation for the derivative

$$\frac{\partial}{\partial x}f = \sum \frac{f(x)}{x - a_i}$$

If  $f \in \mathbf{P}^{\pm}$  then the polar derivative has all real roots<sup>1</sup> by Lemma 2.11. Since polar derivatives act like derivatives, it is not surprising that we have this result (due to Laguerre). In fact, in Chapter 9 we will see that they are exactly derivatives.

<sup>&</sup>lt;sup>1</sup>It fails to have all real roots if there are positive and negative roots - e.g.  $(x + 1)^2(x - 2)^2$ 

**Lemma 4.2.** If  $f \in \mathbf{P}^{\pm}$  and f has all distinct roots, then  $f \leq \frac{\partial}{\partial y} f$ . The polar derivative maps  $\mathbf{P}^{\text{pos}}$  to itself, and  $\mathbf{P}^{\text{alt}}$  to itself.

*Proof.* The result follows from Lemma 1.20 and (4.2.2). The mapping properties follow from (4.2.1).  $\Box$ 

**Corollary 4.3.** If 
$$f \in \mathbf{P}^{\pm}$$
 then  $f - \mathfrak{a} \frac{\partial}{\partial x} \frac{\partial}{\partial u} f \in \mathbf{P}$  for  $\mathfrak{a} > 0$ .

Proof. Use Lemma 4.2 and Corollary 2.10.

**Lemma 4.4.** If  $f = a_0 + \cdots + a_n x^n$  is in P then  $\frac{\partial}{\partial x} f$  and  $\frac{\partial}{\partial y} f$  interlace. If f has positive leading coefficient then

- If  $a_{n-1}$  is negative then  $\frac{\partial}{\partial x} f \leq \frac{\partial}{\partial u} f$ .
- If  $a_{n-1}$  is positive then  $\frac{\partial}{\partial x} f \ge \frac{\partial}{\partial u} f$ .

*Proof.* After dividing by  $a_n$  we may assume that f is monic. Since  $\frac{\partial}{\partial y}f = -xf' + nf$  we see that  $\frac{\partial}{\partial y}f$  sign interlaces f'. The sign of  $\frac{\partial}{\partial y}f$  at the largest root of f' is negative. Since the leading coefficient of  $\frac{\partial}{\partial y}f$  is  $a_{n-1}$ , the result follows.

Polar derivatives preserve interlacing, but the direction depends on certain of the coefficients.

**Lemma 4.5.** Suppose that  $f = \sum_{0}^{n} a_{i}x^{i}$  and  $g = \sum_{0}^{n} b_{i}x^{i}$  satisfy  $f \ll g$ .

- If  $a_{n-2}/a_{n-1} > b_{n-2}/b_{n-1}$  then  $\frac{\partial}{\partial y}f \ll \frac{\partial}{\partial y}g$
- If  $a_{n-2}/a_{n-1} < b_{n-2}/b_{n-1}$  then  $\frac{\partial}{\partial y}g \ll \frac{\partial}{\partial y}f$
- If f,  $g \in \mathbf{P}^{\text{alt}}$  or f,  $g \in \mathbf{P}^{\text{pos}}$  then  $\frac{\partial}{\partial y} f \ll \frac{\partial}{\partial y} g$

*Proof.* Since  $\frac{\partial}{\partial y}$  maps **P** to **P**, it follows from Theorem 1.43 that  $\frac{\partial}{\partial y}$  f and  $\frac{\partial}{\partial y}$  g interlace. If we compute  $\frac{\partial}{\partial y}$  f by reversing f, differentiating, and then reversing, we see that if f has roots all the same sign then  $\frac{\partial}{\partial y}$  preserves interlacing. It remains to find the direction of interlacing.

In general, if  $\frac{\partial}{\partial y} f \ll \frac{\partial}{\partial y} g$ , then we can differentiate n - 2 times to find that

$$n!a_{n-1}x + (n-1)!a_{n-2} \ll n!b_{n-1}x + (n-1)!b_{n-2}$$

If  $\frac{\partial}{\partial y}g \ll \frac{\partial}{\partial y}f$  then this interlacing would be reversed. This implies the result.

We can consider the poset of all partial derivatives of a polynomial f. In case f is in  $\mathbf{P}^{\pm}$  then we have predictable interlacings. For instance,

**Corollary 4.6.** If  $f \in P^{pos}$  is a polynomial of degree n, then whenever  $i + j \leq n$  we have

$$\frac{\partial^{i}}{\partial x^{i}}\frac{\partial^{j}}{\partial y^{j}}f \ll \frac{\partial^{i-1}}{\partial x^{i-1}}\frac{\partial^{j+1}}{\partial y^{j+1}}f$$

The poset of partial derivatives is log concave.

**Lemma 4.7.** If  $f \in \mathbf{P}^{\text{pos}}$  has all distinct roots then  $\left| \begin{array}{c} f & \frac{\partial}{\partial x} f \\ \frac{\partial}{\partial y} f & \frac{\partial}{\partial x} \frac{\partial}{\partial y} f \end{array} \right| < 0$ 

Proof. From the definition we have

$$\begin{vmatrix} f & \frac{\partial}{\partial x} f \\ \frac{\partial}{\partial y} f & \frac{\partial}{\partial x} \frac{\partial}{\partial y} f \end{vmatrix} = \begin{vmatrix} f & f' \\ nf - xf & nf' - (xf')' \end{vmatrix} = - \begin{vmatrix} f & f' \\ xf' & (xf')' \end{vmatrix}$$

Since  $f \in \mathbf{P}^{pos}$  we know  $f \gg xf'$ , and the result follows from Lemma 1.55.  $\Box$ 

#### 4.3 Newton's inequalities

We are going to derive relations among the coefficients of one or two polynomials whose roots have appropriate properties. The idea is to differentiate the polynomials until we get a polynomial of degree 1 or 2, and then use properties of linear and quadratic polynomials.

It is immediate from the quadratic formula that if that  $f(x) = ax^2 + bx + c$  has two distinct real roots then  $b^2 > 4ac$ . We can use the quadratic formula to derive inequalities for the coefficients of a polynomial that has all real roots. Suppose that we are given a fourth degree polynomial  $f = a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4$  with all distinct roots. Then,  $\frac{\partial}{\partial x}f$  has all distinct roots. Differentiating once more, the second derivative  $2a_2+6a_3x+12a_4x^2$  has two distinct roots, so we conclude that  $36a_3^2 > 96a_2a_4$ , or  $a_3^2 > (4/3)a_2a_4$ . Newton generalized this argument.

**Theorem 4.8** (Newton's Inequalities). Suppose that the polynomial  $a_n x^n + \cdots + a_0$  has all real roots.

1. If the roots are all distinct then

$$a_k^2 > (1 + \frac{1}{k})(1 + \frac{1}{n-k})a_{k-1}a_{k+1}$$
, for  $k = 1, ..., n-1$ . (4.3.1)

2. If the roots are not necessarily distinct, then

$$a_k^2 > a_{k-1}a_{k+1}$$
 for  $k = 1, \dots, n-1.$  (4.3.2)

*unless*  $a_k = a_{k-1} = 0$  *or*  $a_k = a_{k+1} = 0$ 

*Proof.* Let F be the homogeneous polynomial corresponding to f. Since partial derivatives preserve distinct real roots, if f has all distinct roots then the polynomial

$$\frac{\partial^{k-1}}{\partial x^{k-1}}\frac{\partial^{n-k-1}}{\partial y^{n-k-1}}\mathsf{F}=n!\left(\frac{a_{k-1}}{\binom{n}{k-1}}x^2+2\frac{a_k}{\binom{n}{k}}xy+\frac{a_{k+1}}{\binom{n}{k+1}}y^2\right).$$

has all real roots, and they are all distinct. Consequently,

$$\left(\frac{a_{k}}{\binom{n}{k}}\right)^{2} > \frac{a_{k-1}}{\binom{n}{k-1}} \cdot \frac{a_{k+1}}{\binom{n}{k+1}}$$
(4.3.3)

which reduces to (4.3.1).

If the roots of f aren't distinct, then the inequality in (4.3.1) is replaced by  $\geq$ . If both  $a_{k-1}$  and  $a_{k+1}$  are non-zero, then (4.3.2) follows. If one of them is zero then the only way for (4.3.2) to fail is if  $a_k = 0$ .

There is a kind of converse to Newton's inequalities, see §5.5.4. If all the coefficients are non-zero then we can restate the conclusion in terms of the *Newton quotients*:

$$\frac{a_k^2}{a_{k-1}a_{k+1}} \ge \frac{k+1}{k} \frac{n-k+1}{n-k}$$
(4.3.4)

**Example 4.9.** We can use the Newton quotient to get inequalities for more widely separated coefficients. Suppose  $f(x) = \sum a_i x^i \in \mathbf{P}^{pos}(n)$ . From (4.3.4) we get

$$\begin{split} a_{k}^{4} &\geqslant \left(\frac{k+1}{k}\frac{n-k+1}{n-k}\right)^{2} a_{k-1}^{2} a_{k+1}^{2} \\ &\geqslant \left(\frac{k+1}{k}\frac{n-k+1}{n-k}\right)^{2} \frac{k}{k-1}\frac{n-k+2}{n-k+1} a_{k} a_{k-2} \frac{k+2}{k+1}\frac{n-k}{n-k-1} a_{k+2} a_{k} \\ &\qquad \frac{a_{k}^{2}}{a_{k-2}a_{k+2}} \geqslant \sqrt{\frac{(k+2)(k+1)}{k(k-1)}\frac{(n-k+2)(n-k+1)}{(n-k)(n-k-1)}} \end{split}$$

If we iterate this inequality, then for any r satisfying  $k \ge 2^r$ ,  $n - k \ge 2^r$ 

$$\frac{a_{k}^{2}}{a_{k-2^{r}}a_{k+2^{r}}} \geqslant \sqrt[2^{r}]{\frac{\langle \underline{k}+1 \rangle_{2^{r}}}{(\underline{k})_{2^{r}}}} \frac{\langle \underline{n}-\underline{k}+1 \rangle_{2^{r}}}{(\underline{n}-\underline{k})_{2^{r}}}$$
(4.3.5)

A similar argument to the above puts an interesting restriction on consecutive coefficients of a polynomial with all real roots.

**Lemma 4.10.** If a polynomial f has all real roots, and two consecutive coefficients of f are zero, then we can write  $f = x^2 g$  where g is a polynomial with all real roots.

*Proof.* Write  $f = \sum a_i x^i$ , and let F be the corresponding homogeneous polynomial. Since partial derivatives preserve roots, the polynomial

$$G(x,y) = \frac{\partial^{k-3}}{\partial x^{k-3}} \frac{\partial^{n-k}}{\partial y^{n-k}} F$$
  
=  $c_1 a_k x^3 + c_2 a_{k-1} x^2 y + c_3 a_{k-2} x y^2 + c_4 a_{k-3} y^3$ 

has all real roots, where  $c_1$ ,  $c_2$ ,  $c_3$ ,  $c_4$  are non-zero multiples of the original coefficients only depending on n and k. If consecutive coefficients  $a_{k-1} = a_{k-2} = 0$ , then  $G(x, 1) = c_1 a_k x^3 + c_3 a_{k-3}$ . This polynomial does not have all real roots unless  $a_{k-3} = 0$  or  $a_k = 0$ . A similar argument shows that if we have r consecutive coefficients that are 0, then if they do not include the constant term we can find a block of r + 1 consecutive zero coefficients. Continuing, we see that at the very least we must have that the constant term and the coefficient of x are 0.

**Corollary 4.11.** If  $f(x) = a_0 + \cdots + a_n x^n$  has all real roots and not both  $a_0$  and  $a_1$  are zero then  $a_i^2 - a_{i+1}a_{i-1} > 0$  for i = 1, ..., n - 1.

We can read Lemma 4.10 to say that if two consecutive coefficients are zero for  $f \in \mathbf{P}$ , then all earlier ones are also zero. We can slightly extend this to two variables.

**Lemma 4.12.** Suppose 
$$f(x, y) = \cdots + g(x)y^r + by^{r+1} + cy^{r+2} + \cdots$$
 satisfies

 $f(\alpha, y) \in \boldsymbol{P}$  for all  $\alpha \in \mathbb{R}$ 

then g has even degree, and  $cg(x) \leq 0$  for all x.

*Proof.* As is usual in these arguments, we differentiate r times with respect to y, reverse, and differentiate and reverse again until we have a quadratic. It has the form  $\alpha g(x) + \beta by + \gamma cy^2$  for positive  $\alpha$ ,  $\beta$ ,  $\gamma$ . Since this has roots for all values of x we must have that  $g(x)c \leq 0$ , which implies g(x) must have even degree.

This phenomenon is possible - for instance, take  $f = x^4 - y^2$ . The corollary below is a two variable analog of Lemma 4.10.

**Corollary 4.13.** If  $f(x, y) = \sum f_i(x)y^i$  satisfies

- 1.  $f(\alpha, y) \in \mathbf{P}$  for all  $\alpha \in \mathbb{R}$ .
- 2. the degree of any  $f_i$  is at most 1.
- 3. Some two consecutive  $f_i$  are constant.

then all earlier  $f_i$  are constant.

*Proof.* Using the lemma inductively shows that all earlier coefficients are constant.

### 4.4 More inequalities and bounds

There are many inequalities beyond the Newton inequalities. It's known that the set of polynomials of fixed degree with real coefficients is a semi-algebraic set, and so is determined by a finite set of inequalities. Such a set is given by Theorem 4.23. In this section we show how to get inequalities using majorization and the Arithmetic-Geometric inequality. We apply these to determining a bound for polynomials on an interval.

The following well-known inequality uses the Arithmetic-Geometric inequality. I don't know if it follows from Newton's inequalities. Lemma 4.15 uses a weaker inequality that holds for all the coefficients.

**Lemma 4.14.** If  $a_0 + \cdots + a_n x^n \in P^{pos}$  then

1. 
$$a_{n-1}^n \ge n^n a_0 a_n^{n-1}$$
  
2.  $a_1 \left(\frac{a_1}{n a_0}\right)^{n-1} \ge a_n$ 

*The inequality is strict unless*  $f = a(x + b)^n$  *for constants* a, b.

*Proof.* If the roots are  $-r_1, \ldots, -r_n$  then

$$\frac{a_0}{a_n} = r_1 \cdots r_n$$
$$\frac{a_{n-1}}{a_n} = r_1 + \cdots + r_n$$

Since all r<sub>i</sub> are positive the Arithmetic-Geometric inequality implies that

$$\frac{1}{n}\frac{a_{n-1}}{a_n} = \frac{r_1 + \dots + r_n}{n} \ge (r_1 \cdots r_n)^{1/n} = \left(\frac{a_0}{a_n}\right)^{1/n}$$

The second inequality follows from the first by reversing f. Equality holds in the AGI if and only if all roots are equal.  $\Box$ 

We can use bounds on the coefficients to bound the size of a polynomial with all real roots. We begin with a simple consequence of MacLaurin's inequality. The important point is that the upper bound does not depend on the degree of the polynomial.

**Lemma 4.15.** If 
$$f(x) = \sum_{0}^{n} a_i x^i \in P^{pos}$$
 then  $\sup_{|x| \leq r} |f(x)| \leq |a_0| e^{ra_1/a_0}$ 

*Proof.* Since  $f \in \mathbf{P}^{pos}$  we can apply Newton's inequalities, and so

$$\frac{a_1}{a_0} > 2\frac{a_2}{a_1} > 2\left(\frac{3}{2}\frac{a_3}{a_2}\right) > 3\left(\frac{4}{3}\frac{a_4}{a_3}\right) > \cdots k\frac{a_k}{a_{k-1}}$$

This gives a version of MacLaurin's inequality:

$$a_k < a_1 \frac{1}{k!} \left(\frac{a_1}{a_0}\right)^{k-1}$$

and therefore for r > 1

$$\sup_{|x|\leqslant r} |f(x)| \leqslant r^n \sum_{k=0}^n a_k < a_0 \sum_0^n \left(\frac{a_1}{a_0}\right)^k r^n / k! < a_0 \exp(ra_1/a_0)$$

To find bounds for arbitrary polynomials in **P** we need a result of [167].

**Lemma 4.16.** If  $f(x) = a_m x^m + \cdots + a_n x^n \in P$  and  $a_m a_n \neq 0$  then

$$\sup_{|x|\leqslant r} |f(x)|\leqslant |a_m|r^m \exp \left( r \left| \frac{a_{m+1}}{a_m} \right| + r^2 \left| \frac{a_{m+1}^2}{a_m^2} \right| + 3 \, r^2 \left| \frac{a_{m+2}}{a_m} \right| \right)$$

**Definition 4.17.** Given two sequences  $\mathbf{a} = (a_1 \leq a_2 \leq \cdots \leq a_n)$  and  $\mathbf{b} = (b_1 \leq b_2 \leq \cdots \leq b_n)$  then we say  $\mathbf{a}$  majorizes  $\mathbf{b}$ , written  $\mathbf{b} \prec_m \mathbf{a}$ , if

There is an important inequality between monomial symmetric functions due to Muirhead [78, 164].

#### **Theorem 4.18** (Muirhead). *Given two sequences* $\mathbf{a} \prec_{\mathrm{m}} \mathbf{b}$ *where*

 $\mathbf{a} = (a_1 \leqslant a_2 \leqslant \cdots \leqslant a_n)$  and  $\mathbf{b} = (b_1 \leqslant b_2 \leqslant \cdots \leqslant b_n)$  then for all positive  $x_1, \ldots, x_n$ 

$$\sum_{\sigma \in Sym_{\mathfrak{n}}} x_{\sigma(1)}^{a_1} x_{\sigma(2)}^{a_2} \cdots x_{\sigma(n)}^{a_n} \leqslant \sum_{\sigma \in Sym_{\mathfrak{n}}} x_{\sigma(1)}^{b_1} x_{\sigma(2)}^{b_2} \cdots x_{\sigma(n)}^{b_n}$$
(4.4.1)

**Example 4.19.** Consider a cubic  $f = (x + r_1)(x + r_2)(x + r_3)$  where  $r_1, r_2, r_3$  are positive. We also write  $f = c_0 + c_1x + c_2x^2 + c_3x^3$ . We list the four index sequences that sum to 4 and the expansion (after homogenizing) of the sum in Muirhead's theorem in terms of the coefficients:

Name		expansion
$M_{004}$	0,0,4	$2(c_2^4 - 4c_1c_2^2 c_3 + 2 c_1^2 c_3^2 + 4 c_0 c_2 c_3^2)$
$M_{013}$	0,1,3	$-c_3\left(2c_3c_1^2-c_2^2c_1+c_0c_2c_3\right)$
$M_{022}$	0,2,2	$2(c_1^2 - 2c_0c_2)c_3^2$
M <sub>112</sub>	1,1,2	$2c_0c_2c_3^2$

For example,

$$M_{112} = r_1 r_2 r_3^2 + r_1 r_2^2 r_3 + r_1^2 r_2 r_3$$
  
=  $r_1 r_2 r_3 (r_1 + r_2 + r_3)$   
=  $c_0 c_2$ 

To homogenize we form  $c_3^4 \frac{c_0}{c_3} \frac{c_2}{c_3} = c_0 c_2 c_3^2$ . There are four majorizations that yield (after simplifying) inequalities

$M_{013} \prec_{\mathfrak{m}} M_{004}$	$2c_2^4 - 9c_1c_3c_2^2 + 9c_0c_3^2c_2 + 6c_1^2c_3^2$	$\geqslant 0$
$M_{022} \prec_m M_{004}$	$c_2^4-4c_1c_3c_2^2+6c_0c_3^2c_2+c_1^2c_3^2\\$	$\geqslant 0$
$M_{112} \prec_{\mathfrak{m}} M_{013}$	$-3c_0c_2 + c_1^2$	$\geqslant 0$
$M_{112} \prec_m M_{022}$	$-2c_3c_1^2+c_2^2c_1-3c_0c_2c_3$	$\geqslant 0$

Although the third is Newton's inequality, calculations show that the first one is not a consequence of the two Newton inequalities.

## 4.5 The matrix of a transformation

If T is a transformation on polynomials, then we define  $\varphi(T)$  to be the matrix of T in the basis {1, x, x<sup>2</sup>, x<sup>3</sup>,...}. We investigate the relation between the properties of T and the properties of  $\varphi(T)$ .

It is often the case that  $\varphi(T)$  is totally positive. However,  $T: \mathbf{P} \longrightarrow \mathbf{P}$  does not imply  $\varphi(T)$  is totally positive, nor is the converse true. Consider the two examples:

**Example 4.20.** Suppose that  $T(x^i) = a_i x^i$ , and D is the diagonal matrix whose i, i entry is  $a_i$ . Clearly D is totally positive, yet most transformations of this form do not preserve roots - see Theorem 15.22.

**Example 4.21.** If  $T(f) = f(D)x^2$  then we know that  $T: P \longrightarrow P$ . However,

$$\phi(\mathsf{T}) = \begin{pmatrix} 0 & 0 & 1 & \dots \\ 0 & 2 & \dots & \\ 2 & 0 & \dots & \\ \vdots & & & \end{pmatrix}$$

where all the remaining entries are zero. This matrix is not totally positive.

Although a transformation that preserves roots does not necessarily determine a totally positive matrix, we do have a result for two by two submatrices if the direction of interlacing is preserved.

Lemma 4.22. If the linear transformation T satisfies

1. T:  $P^{\text{pos}} \longrightarrow P^{\text{pos}}$ 

- 2. T preserves the sign of the leading coefficient.
- 3.  $f \leq g$  implies  $T(f) \leq T(g)$  for  $f \in \mathbf{P}^{pos}$ .

then all two by two submatrices of  $\phi(T)$  are non-negative. (Thus  $\phi(T)$  is TP<sub>2</sub>.)

Proof. It suffices to show that all two by two submatrices composed of adjacent entries have non-negative determinants. [61] Since  $T(x^n) \leq T(x^{n-1})$ , it follows from Corollary 4.31 that the determinants are non-negative. 

We now consider several examples where  $\varphi(T)$  is totally positive. If  $M_f(g) = f \cdot g$  is the multiplication transformation, then we write  $\varphi(f)$  for  $\varphi(M_f)$ . One important case is when f = a + bx.

$$\phi(a+bx) = \begin{pmatrix} a & b & 0 & 0 & \dots \\ 0 & a & b & 0 & \dots \\ 0 & 0 & a & b & \dots \\ 0 & 0 & 0 & a & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$
(4.5.1)

For another example, take  $f(x) = a_0 + a_1x + a_2x^2 + a_3x^3$ .

$$\varphi(f) = \begin{pmatrix} a_0 & a_1 & a_2 & a_3 & 0 & 0 & 0 & \dots \\ 0 & a_0 & a_1 & a_2 & a_3 & 0 & 0 & \dots \\ 0 & 0 & a_0 & a_1 & a_2 & a_3 & 0 & \dots \\ 0 & 0 & 0 & a_0 & a_1 & a_2 & a_3 & \dots \\ \vdots & \vdots & & & \ddots & \ddots & \ddots \end{pmatrix}$$
(4.5.2)

The matrix of a derivative transformation is similar:

$$\phi(\mathbf{f} \mapsto \alpha \mathbf{f} + \mathbf{f}') = \begin{pmatrix} \alpha & 0 & 0 & 0 & \dots \\ 1 & \alpha & 0 & 0 & \dots \\ 0 & 2 & \alpha & 0 & \dots \\ 0 & 0 & 3 & \alpha & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$
(4.5.3)

If  $f(x) = a_0 + a_1x + a_2x^2 + a_3x^3$  then

$$\phi(g \mapsto f(\mathsf{D})g) = \begin{pmatrix} a_0 & 0 & \dots & & \\ a_1 & a_0 & 0 & \dots & \\ 2a_2 & 2a_1 & a_0 & 0 & \dots & \\ 6a_3 & 6a_2 & 3a_1 & a_0 & 0 & \dots & \\ 24a_4 & 24a_3 & 12a_2 & 4a_1 & a_0 & 0 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

,

Finally, the map of an affine transformation leads to binomial coefficients:

$$\varphi(f(x)\mapsto f(x+1)) = \begin{pmatrix} 1 & 0 & 0 & 0 & \dots \\ 1 & 1 & 0 & 0 & \dots \\ 1 & 2 & 1 & 0 & \dots \\ 1 & 3 & 3 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

The first result is well known [25].

**Theorem 4.23.** If a polynomial f is in  $P^{\text{pos}}$  then  $\varphi(f)$  is totally positive. [25]

*Proof.* We show by induction on the degree of f that  $\varphi(f)$  is totally positive. If  $M_f$  is multiplication by f then the composition of  $M_f$  and  $M_g$  is  $M_{fg}$ , and consequently  $\varphi(fg) = \varphi(f)\varphi(g)$ . From (4.5.1) it is immediate that  $\varphi(a + bx)$  is a totally positive matrix if a and b are non-negative. Since every polynomial in  $\mathbf{P}^{\text{pos}}$  can be written as a product of factors of the form a + bx where a, b are positive, the conclusion follows from the fact that the product of two totally positive matrices is totally positive.

**Theorem 4.24.** If  $f \in P^{pos}$  and T(g) = f(D)g then  $\varphi(T)$  is totally positive.

*Proof.* Same as above. Just note that f(D) is a composition of operators of the form  $f \mapsto \alpha f + \beta f'$ . From (4.5.3) it is clear that  $\varphi(f \mapsto \alpha f + \beta f')$  is totally positive.

*Remark* 4.25. These last two results are equivalent since the two matrices  $\varphi(f)$  and  $\varphi(g \mapsto f(D)g)$  are similar. This isn't surprising since differentiation and multiplication are adjoint. If F is the infinite diagonal matrix with ith diagonal entry i! then it is easy to see that

$$\varphi(\mathfrak{g} \mapsto \mathfrak{f}(\mathsf{D})\mathfrak{g}) = \mathsf{F}\,\varphi(\mathfrak{f})^{\mathsf{t}}\,\mathsf{F}^{-1}$$

*Remark* 4.26. It is actually the case that the converse of Theorem 4.23 is also true: if  $\varphi(f)$  is totally positive, then  $f \in \mathbf{P}^{pos}$ . This is known as the Aissen-Schoenberg-Whitney Theorem [2]. Also, note that the two by two determinants give Newton's inequalities.

The general form of Hurwitz's theorem (see Theorem 7.64) follows immediately from this remark.

**Corollary 4.27.** If  $\sum_{i} a_i x_i$  is in  $P^{pos}$  then for any positive d the polynomial  $\sum_{i} a_{di} x^i$  is also in  $P^{pos}$ .

*Proof.* The totally positive matrix corresponding to  $\sum a_{di}x^i$  is formed by taking the submatrix of (4.5.2) whose rows and columns are divisible by d.

We can apply Theorem 4.23 to determinants involving evaluations at integers. **Corollary 4.28.** Suppose  $f \in P^{pos}$ , d is a positive integer, and  $\alpha > 0$ . Then

$$\begin{vmatrix} \frac{1}{d!}f(\alpha+d) & \frac{1}{(d-1)!}f(\alpha+d-1) & \dots & f(\alpha) \\ \vdots & & \vdots \\ \frac{1}{(2d)!}f(\alpha+2d) & \frac{1}{(2d-1)!}f(\alpha+2d-1) & \dots & \frac{1}{d!}f(\alpha+d) \end{vmatrix} \ge 0$$

*Proof.* Note that  $f(x + \alpha) \in \mathbf{P}^{pos}$ . From Lemma 7.33 we know that

$$\sum_{k=0}^{\infty} \frac{f(i+\alpha)}{i!} x^i \in \widehat{I\!\!P^{\text{pos}}}$$

We can now apply Theorem 4.23.

If  $f = \sum \alpha_i x^i$  then we can define an infinite matrix which is just  $\varphi(f)$  (see (4.5.2)) bordered by powers of x. For instance, if f has degree three then

$$N(f) = \begin{pmatrix} 1 & \alpha_0 & \alpha_1 & \alpha_2 & \alpha_3 & 0 & 0 & 0 & \dots \\ x & 0 & \alpha_0 & \alpha_1 & \alpha_2 & \alpha_3 & 0 & 0 & \dots \\ x^2 & 0 & 0 & \alpha_0 & \alpha_1 & \alpha_2 & \alpha_3 & 0 & \dots \\ x^3 & 0 & 0 & 0 & \alpha_0 & \alpha_1 & \alpha_2 & \alpha_3 & \dots \\ x^4 & 0 & 0 & 0 & 0 & \alpha_0 & \alpha_1 & \alpha_2 & \dots \\ \vdots & \vdots & & & \ddots & \ddots & \ddots & \ddots \end{pmatrix}$$
(4.5.4)

**Proposition 4.29.** Suppose  $f(x) = \sum \alpha_i x^i \in \mathbf{P}^{pos}$ . If d is odd then all d by d submatrices of N(f) are non-negative for positive x.

*Proof.* We proceed by induction on the degree n of f. If n is 0 then

$$N(f) = \begin{pmatrix} 1 & \alpha_0 & 0 & 0 & 0 & \dots \\ x & 0 & \alpha_0 & 0 & 0 & \dots \\ x^2 & 0 & 0 & \alpha_0 & 0 & \dots \\ x^3 & 0 & 0 & 0 & \alpha_0 & \dots \\ \vdots & \vdots & & & \ddots \end{pmatrix}$$

We need to check that every d by d submatrix has non-negative determinant. If the first column isn't part of the submatrix then this is trivial. If the first column is part of the submatrix then it's easy to see that the only way the determinant is not zero is if the submatrix looks like (for d = 5):

$$\begin{pmatrix} x^m & \alpha_0 & 0 & 0 & 0 \\ x^{m+1} & 0 & \alpha_0 & 0 & 0 \\ x^{m+2} & 0 & 0 & \alpha_0 & 0 \\ x^{m+3} & 0 & 0 & 0 & \alpha_0 \\ x^{m+4} & 0 & 0 & 0 & 0 \end{pmatrix}$$

The determinant is  $(-1)^{d-1} x^{m+4} \alpha_0^{d-1}$ . Since d is odd, this is positive for positive x.

If we multiply N(f) on the left by  $\varphi(a + bx)$  (see (4.5.2)) then the result is

 $\begin{pmatrix} a + bx & a\alpha_0 & a\alpha_1 + b\alpha_0 & a\alpha_2 + b\alpha_1 & \dots \\ (a + bx)x & 0 & a\alpha_0 & a\alpha_1 + b\alpha_0 & \dots \\ (a + bx)x^2 & 0 & 0 & a\alpha_0 & \dots \\ \vdots & \vdots & & & \end{pmatrix}$ (4.5.5)

This is nearly N((a + bx)f), except that the first column of (4.5.5) is the first column of N((a + bx)f) multiplied by a + bx.

Assume that the proposition holds true for N(f). Any submatrix of N((a + bx)f) not containing the first column has non-negative determinant. If a submatrix contains the first column, then its determinant is (a + bx) times the corresponding determinant in  $\varphi(a+bx)N(f)$ , which is positive by Cauchy-Binet and the induction hypothesis.

*Remark* 4.30. The case of d = 3 yields the cubic Newton inequalities (Corollary 4.39) for positive x.

# 4.6 More coefficient inequalities

An important consequence of Lemma 1.55 concerns the coefficients of interlacing polynomials.

**Corollary 4.31.** If  $f = a_0 + \cdots + a_n x^n$ ,  $g = b_0 + \cdots + b_n x^n$ ,  $f \leftarrow g$ , there is no index k such that  $a_k = b_k = 0$ , and f, g have positive leading coefficients then for  $i = 0, 1, \ldots, n - 1$  we have

$$\frac{b_i \ b_{i+1}}{a_i \ a_{i+1}} > 0$$
 (4.6.1)

Proof. From Lemma 1.55 we have that

 $0 > \left|\begin{smallmatrix} f(0) & g(0) \\ f'(0) & g'(0) \end{smallmatrix}\right| = \left|\begin{smallmatrix} a_0 & a_1 \\ b_0 & b_1 \end{smallmatrix}\right|.$ 

since zero is not a common root. In the general case we differentiate  $f \ll g$  exactly i - 1 times. The result is the interlacing

$$i!a_i + (\underline{i+1})_i a_{i+1} + \dots + (\underline{n})_i x^{n-i+1} \ll i!b_i + (\underline{i+1})_i b_{i+1} + \dots + (\underline{n})_i x^{n-i+1}$$

We can now apply the case i = 0 to complete the argument.

*Remark* 4.32. There is a useful picture to help remember this result that will reappear in higher dimensions. If we write the coefficients of f and g in the diagram



then in each pair of adjacent triangles of a square the product of the coefficients at the diagonally adjacent vertices is greater than the product of the coefficients at the remaining two vertices.

*Remark* 4.33. We can derive these inequalities from Newton's inequalities. In fact, if  $f = \prod_i (x - a_i)$ , and  $f_j = f/(x - a_j)$ , then Newton's inequalities imply that the determinant (4.6.1) is positive for the functions f and  $f_j$ . Now the determinant is linear in the coefficients of g, so the coefficients of any convex combination of f,  $f_1$ ,... also satisfy (4.6.1). These convex combinations describe all the possible g's by Lemma 1.20.

If we restrict f, g to lie in  $\mathbf{P}^{pos}$  then

**Lemma 4.34.** If  $f = a_0 + \cdots + a_n x^n$ ,  $g = b_0 + \cdots + b_n x^n$ ,  $f \ll g$ ,  $f, g \in P^{\text{pos}}$  and f, g have positive leading coefficients then for all  $0 \leq r < s \leq n$ :

$$\begin{vmatrix} b_r & b_s \\ a_r & a_s \end{vmatrix} > 0$$

*Proof.* If  $f, g \in \mathbf{P}^{pos}$  then all coefficients are positive, so from Corollary 4.31 we have the inequalities

$$\frac{a_{r+1}}{a_r} > \frac{b_{r+1}}{b_r} \qquad \frac{a_{r+2}}{a_{r+1}} > \frac{b_{r+2}}{b_{r+1}} \qquad \cdots \qquad \frac{a_s}{a_{s-1}} > \frac{b_s}{b_{s-1}}$$

Multiplying these inequalities yields the conclusion.

In Lemma 4.34 we extended the Corollary 4.31 to determinants of the form  $\begin{vmatrix} a_{i+d} & a_i \\ b_{i+d} & b_i \end{vmatrix}$ . Next, we observe that the coefficients of interlacing polynomials can not agree very often:

**Corollary 4.35.** *If*  $f \ll g$  *are both in*  $P^{pos}$ *, then* f *and* g *can not agree at any two coefficients.* 

*Proof.* If f, g have coefficients as above, and  $a_k = b_k$ ,  $a_{k+d} = b_{k+d}$ , then Lemma 4.34 is contradicted, for the determinant is 0.

Here's another variation.

**Lemma 4.36.** If  $f = \sum a_i x^i$  and  $g = \sum b_i x^i$  interlace, are linearly independent, and  $\begin{vmatrix} a_0 & b_0 \\ a_1 & b_1 \end{vmatrix} = 0$  then  $a_0 = b_0 = 0$ .

*Proof.* Since the determinant is zero there is a  $\lambda$  such that  $\lambda(a_0, a_1) = (b_0, b_1)$ . If  $\lambda \neq 0$  then by linear independence  $\lambda f - g \neq 0$ , and f and  $\lambda f - g$  interlace. Since  $\lambda f - g$  has a factor of  $x^2$ , f has a factor of x. Considering g and  $\lambda f - g$  shows that g has a factor of x.

If  $\lambda = 0$  then g has a factor of  $x^2$ , so f has a factor of x.

*Remark* 4.37. If we form the matrix of coefficients of three mutually interlacing quadratics  $f \ll g \ll h$ , then we know that all two by two determinants are positive (Corollary 4.31), but the three by three determinant can be either positive or negative. In particular, the matrix of coefficients is not totally positive.

h = (4 + x)(12 + x)	$\mathbf{h} = (8 + \mathbf{x})(12 + \mathbf{x})$
g = (3+x)(9+x)	g = (7 + x)(10 + x)
f = (2+x)(8+x)	f = (6+x)(9+x)
48 16 1	96 20 1
$-2 = \begin{vmatrix} 27 & 12 & 1 \end{vmatrix}$	$4 = egin{bmatrix} 70 & 17 & 1 \end{bmatrix}$
16 10 1	54 15 1

Next we have a kind of reverse Newton's inequalities for two by two determinants of interlacing polynomials.

**Lemma 4.38.** Suppose that  $f = \sum a_i x^i$  and  $g = \sum b_i x^i$  are in P(n). If  $f \leq g$  then

$$\begin{vmatrix} b_k & b_{k+2} \\ a_k & a_{k+2} \end{vmatrix}^2 \leqslant 4 \, \cdot \, \frac{(k+2)(n-k)}{(k+1)(n-k-1)} \, \cdot \, \begin{vmatrix} b_k & b_{k+1} \\ a_k & a_{k+1} \end{vmatrix} \begin{vmatrix} b_{k+1} & b_{k+2} \\ a_{k+1} & a_{k+2} \end{vmatrix}$$

*Proof.* Since  $f \leq g$  we know that  $f + \alpha g \in \mathbf{P}$  for all  $\alpha$ . If we let  $\beta = (k+2)(n-k)/((k+1)(n-k-1))$  then Newton's inequalities state that for all  $\alpha$  we have

$$(\mathfrak{a}_{k+1} + \alpha \mathfrak{b}_{k+1})^2 \geq \beta (\mathfrak{a}_k + \alpha \mathfrak{b}_k)(\mathfrak{a}_{k+2} + \alpha \mathfrak{b}_{k+2})$$

Since the difference is non-negative for all  $\alpha$  the discriminant is non-positive. A computation of the discriminant shows that

$$\beta^{2} \begin{vmatrix} b_{k} & b_{k+2} \\ a_{k} & a_{k+2} \end{vmatrix}^{2} - 4\beta \begin{vmatrix} b_{k} & b_{k+1} \\ a_{k} & a_{k+1} \end{vmatrix} \begin{vmatrix} b_{k+1} & b_{k+2} \\ a_{k+1} & a_{k+2} \end{vmatrix} \leqslant 0$$

and the conclusion follows.

An equivalent formulation is that the determinant below has all real roots.

$$\begin{vmatrix} 1 & \sqrt{\beta} x & x^2 \\ b_k & b_{k+1} & b_{k+2} \\ a_k & a_{k+1} & a_{k+2} \end{vmatrix}$$

Rosset [147] called the following corollary the cubic Newton inequality **Corollary 4.39.** If  $f(x) = \sum a_i x^i \in P^{pos}(n)$  and  $0 \le k \le n-3$  then

$$4 \left( a_{k+1}^{2} - a_{k} a_{k+2} \right) \left( a_{k+2}^{2} - a_{k+1} a_{k+3} \right) \ge \left( a_{k+1} a_{k+2} - a_{k} a_{k+3} \right)^{2}$$

*Proof.* Apply the lemma to xf and f.

# 4.7 Log concavity in P<sup>pos</sup>

The logarithms of the coefficients of a polynomial in  $\mathbf{P}^{\text{pos}}$  have useful and well-known concavity properties. We will revisit these ideas when we consider polynomials in two variables.

To begin, suppose that  $f = a_0 + \cdots + a_n x^n$  has all positive coefficients and is in  $\mathbf{P}^{pos}$ . We define the Newton graph of f to be the collection of segments joining the points

 $(0, \log a_0), (1, \log a_1), \cdots, (n, \log a_n).$ 

For instance, Figure 4.7 is the Newton graph for n = 4.



Figure 4.1: The Newton graph of a polynomial in **P**<sup>pos</sup>

The Newton graph is concave. Indeed, the slope of the segment between  $(k, \log a_k)$  and  $(k + 1, \log a_{k+1})$  is  $\log(a_{k+1}/a_k)$ . Taking logs of Newton's inequality yields

$$\log \frac{a_{k+1}}{a_k} \ge \log \frac{a_{k+2}}{a_{k+1}},$$

which means that the slopes are decreasing. This implies that the Newton graph is concave.

Next, we consider the Newton graphs of two interlacing polynomials. Assume that  $f \ll g$  in  $\mathbf{P}^{pos}$ . In order to compare coefficients we will assume that both are monic, so that their Newton graphs agree at the right hand endpoint. The Newton graph of f lies below the Newton graph of g, and the gap decreases as the index increases. For example, Figure 4.7 shows f and  $g = \sum b_i x^i$ . The dotted lines decrease in length as we move to the right.

We use Lemma 4.34 which says that  $\begin{vmatrix} b_k & b_{k+1} \\ a_k & a_{k+1} \end{vmatrix} \ge 0$ . Taking logs shows that

$$\log \frac{b_{k+1}}{b_k} \leq \log \frac{a_{k+1}}{a_k} \qquad \qquad \log \frac{b_{k+1}}{a_{k+1}} \leq \log \frac{b_k}{a_k}$$

and these inequalities imply the assertions of the last paragraph.

#### 4.8 A pair of recursions

In [188] there is an interesting pair of recursions



Figure 4.2: The Newton graphs of two interlacing polynomials

$$\begin{split} p_{n+1}(z) &= z \, p_n(z) - \alpha_n \, z^n \, q_n(1/z) \\ q_{n+1}(z) &= z \, q_n(z) - \beta_n \, z^n \, p_n(1/z). \end{split}$$

If we homogenize these equations then we can find recurrences for  $p_n$  and  $q_n$ . Using these these recurrences we can easily show that  $p_n$  and  $q_n$  are in **P** for various choices of  $\alpha_n$  and  $\beta_n$ . So consider P(x, y) and Q(x, y) satisfying the recurrences

$$\begin{aligned} P_{n+1}(x,y) &= x P_n(x,y) - \alpha_n y Q_n(y,x) \\ Q_{n+1}(x,y) &= x Q_n(x,y) - \beta_n y P_n(y,x) \end{aligned}$$

with the initial conditions  $P_0(x,y) = Q_0(x,y) = 1$ . We compute  $Q_{n+1}(y,x)$  using each equation

$$\frac{1}{x \, \alpha_{n+1}} \left( y \, \mathsf{P}_{n+1}(y, x) - \mathsf{P}_{n+2}(y, x) \right) = \frac{1}{x \, \alpha_n} \left( y \, \mathsf{P}_n(y, x) - \mathsf{P}_{n+1}(y, x) \right) - \beta_n \, y \, \mathsf{P}_n(y, x)$$

Interchange x and y and simplify

$$P_{n+2}(x,y) = \left(x + \frac{\alpha_{n+1}}{\alpha_n}y\right)P_{n+1}(x,y) + \left(\alpha_{n+1}\beta_n - \frac{\alpha_{n+1}}{\alpha_n}\right)xyP_n(x,y)$$

Since  $P_n$  is homogeneous  $P_n(x, 1) = p_n(x)$  and

$$p_{n+2}(x) = \left(x + \frac{\alpha_{n+1}}{\alpha_n}\right)p_{n+1}(x) + \left(\alpha_{n+1}\beta_n - \frac{\alpha_{n+1}}{\alpha_n}\right)xp_n(x)$$

If the coefficient of  $p_n$  is negative then this defines a sequence of orthogonal Laurent polynomials (p. 45). In the special case that  $\alpha_n = \beta_n = \alpha$  for all n then  $p_n = q_n = f_n$  and we get the simple recurrence

$$f_{n+2} = (x+1)f_{n+1} + x(\alpha^2 - 1)f_n$$
(4.8.1)

**Lemma 4.40.** If  $|\alpha| \leq 1$ ,  $f_0 = 1$ ,  $f_1 = x - \alpha$  and f satisfies (4.8.1) then  $f_n \in P^{\alpha lt}$  and  $f_{n+1} \leq f_n$ .

Another interesting case is when  $\alpha_n = \beta_n = -q^{(n+1)/2}$ . The resulting polynomials are known as the Rogers-Szego polynomials. The recurrence is

$$f_{n+2} = (x+1)f_{n+1} + x(q^{n+1} - 1)f_n$$
(4.8.2)

It follows that

**Lemma 4.41.** If  $q \ge 1$ ,  $f_0 = 1$ ,  $f_1 = x + q$  and f satisfies (4.8.2) then  $f_n \in P^{alt}$  and  $f_{n+1} \le f_n$ .

CHAPTER



# **Analytic Functions**

We introduce some classes of analytic functions that are the uniform closure of certain sets of polynomials with all real roots. We are interested in the relationship between these classes, and the set of generating functions of linear transformations of the form  $x^i \mapsto a_i x^i$  that preserve roots.

In later chapters we will generalize these results to more general linear transformations, and to polynomials in several variables.

# 5.1 The uniform closure of P<sup>pos</sup>

We begin with the simplest class of analytic functions, the closure of polynomials with all positive coefficients and all real roots.

**Definition 5.1.**  $\widehat{P^{pos}}$  is the set of all functions that are in the closure of  $P^{pos}$  under uniform convergence on compact domains.

From basic complex analysis we know

- 1. All functions in  $\widehat{\mathbf{P}^{pos}}$  are analytic.
- 2. All functions in  $\mathbf{P}^{\text{pos}}$  are *entire* this means that they are defined on the entire complex plane.
- 3. All derivatives of functions in  $\widehat{\mathbf{P}^{\text{pos}}}$  are in  $\widehat{\mathbf{P}^{\text{pos}}}$ .
- 4. The product of two functions in  $\widehat{\mathbf{P}^{\text{pos}}}$  is in  $\widehat{\mathbf{P}^{\text{pos}}}$ .
- 5. Every function in  $\widehat{\mathbf{P}^{pos}}$  has all real roots.

It is obvious that  $\mathbf{P}^{\text{pos}} \subset \widehat{\mathbf{P}^{\text{pos}}}$ . The limit of  $\mathbf{P}^{\text{alt}}$  is  $\widehat{\mathbf{P}^{\text{alt}}}$ , and satisfies the above properties, except that  $\mathbf{P}^{\text{alt}} \subset \widehat{\mathbf{P}^{\text{alt}}}$ . We now introduce a few important non-polynomial functions in  $\widehat{\mathbf{P}^{\text{pos}}}$ .

**Example 5.2.** The exponential function.  $e^x$  is the most important non-polynomial function in  $\widehat{\mathbf{P}^{\text{pos}}}$ . To see that it is in  $\widehat{\mathbf{P}^{\text{pos}}}$  note that

$$\lim_{n\to\infty}\left(1+\frac{x}{n}\right)^n = e^x$$

so  $e^x$  is the limit of polynomials that have n-fold zeros at -n. These zeros go off to infinity, so  $e^x$  has no zeros. More generally,  $e^{ax}$  is in  $\widehat{\mathbf{P}^{\text{pos}}}$  for any positive a. If a is negative then  $e^{ax} \in \widehat{\mathbf{P}^{\text{alt}}}$ .

Also, since

$$\lim_{n\to\infty}\left(1+x+\frac{x^2}{2!}+\cdots+\frac{x^n}{n!}\right) = e^x$$

we see that  $e^x$  is also a limit of polynomials that are *not* in **P**, so  $e^x$  is on the boundary of  $\widehat{\mathbf{P}^{\text{pos}}}$ .

**Example 5.3.** *The* q*-exponential function.* There is a q-analog of the exponential function. We first need to introduce the Pochhammer symbols

$$\begin{aligned} (\mathbf{a};\mathbf{q})_{n} &= (1-\mathbf{a})(1-\mathbf{a}\mathbf{q})\cdots(1-\mathbf{a}\mathbf{q}^{n-1})\\ (\mathbf{a};\mathbf{q})_{\infty} &= (1-\mathbf{a})(1-\mathbf{a}\mathbf{q})\cdots(1-\mathbf{a}\mathbf{q}^{n-1})\cdots\end{aligned}$$

The following basic identity [67, (1.3.16)] is valid for |q| < 1:

$$\sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}}}{(q;q)_n} z^n = (-z;q)_{\infty}$$
(5.1.1)

We denote this function by  $\text{Exp}_q(z)$ . If we write out the right hand side of (5.1.1)

$$(-z;q)_{\infty} = (1+z)(1+qz)(1+q^2z)\cdots$$

then we see that for 0 < q < 1 all the roots of  $\text{Exp}_q(z)$  are negative, and so  $\text{Exp}_q(z)$  is in  $\widehat{\mathbf{P}^{\text{pos}}}$ . We will later use this variation (see [6]) on the q-exponential function, where  $[n] = (q^n - 1)(/(q - 1))$ :

$$E_{q}(z) = \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}}}{[n]!} z^{n} = ((1-q)z;q)_{\infty} \qquad |q| < 1$$
(5.1.2)

*Remark* 5.4.  $\widehat{\mathbf{P}^{\text{pos}}}$  is closed under products, so we can construct new elements in  $\widehat{\mathbf{P}^{\text{pos}}}$  by multiplying appropriate factors. First of all,  $x^{\text{r}}$  and  $e^{\alpha x}$  are always in  $\widehat{\mathbf{P}^{\text{pos}}}$  for r a non-negative integer, and a positive. If the infinite product  $\prod_{1}^{\infty}(1+c_{i}x)$  converges, where all  $c_{i}$  are positive, then the following is in  $\widehat{\mathbf{P}^{\text{pos}}}$ :

$$cx^{r}e^{\alpha x}\prod_{i=1}^{\infty}(1+c_{i}x)$$
(5.1.3)

A famous result due to Pólya-Schur [14, 138] states that this representation describes all the elements of  $\widehat{\mathbf{P}^{\text{pos}}}$ .

# 5.2 The uniform closure of P

 $\widehat{\mathbf{P}}$  is the uniform closure of  $\mathbf{P}$  on compact subsets. It satisfies the same general analytic properties that  $\widehat{\mathbf{P}^{\text{pos}}}$  does. Here are some examples of analytic functions in  $\widehat{\mathbf{P}}$ .

**Example 5.5.** *The Gaussian density.* Since

$$e^{-x^{2}} = \lim_{n \to \infty} \left(1 - \frac{x^{2}}{n}\right)^{n} = \lim_{n \to \infty} \left(1 - \frac{x}{\sqrt{n}}\right)^{n} \left(1 + \frac{x}{\sqrt{n}}\right)^{n}$$

we see that

$$e^{-x^2} = 1 - \frac{x^2}{1!} + \frac{x^4}{2!} - \frac{x^6}{3!} + \dots$$

is in  $\widehat{P} \setminus \widehat{P^{pos}}$  since it is the limit of polynomials with both positive and negative roots.

**Example 5.6.** *Sine and Cosine.* The sine and cosine are in  $\hat{\mathbf{P}}$  since they both have infinite products with positive and negative roots:

$$\sin(x) = x \prod_{k=1}^{\infty} \left( 1 - \frac{x^2}{k^2 \pi^2} \right)$$
$$= \frac{x}{1} - \frac{x^3}{3!} + \frac{x^5}{5!} - \cdots$$
$$\cos(x) = \prod_{k=0}^{\infty} \left( 1 - \frac{4x^2}{(2k+1)^2 \pi^2} \right)$$
$$= 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \cdots$$

**Example 5.7.** *Bessel Functions.* The Bessel function of the first kind,  $J_i(z)$ , for i a positive integer, is defined as

$$J_{i}(z) = \frac{z^{i}}{2^{i}} \sum_{k=0}^{\infty} (-1)^{k} \frac{z^{2k}}{2^{2k}k!(k+i)!}$$

There is an infinite product representation [72, page 980], where we denote the zeros of  $J_i(z)$  by  $z_{i,m}$ :

$$J_{i}(z) = \left(\frac{z}{2}\right)^{i} \frac{1}{i!} \prod_{m=1}^{\infty} \left(1 - \frac{z^{2}}{z_{i,m}^{2}}\right)$$

The product representation shows that  $J_i(z)$  is in  $\widehat{\mathbf{P}}$ .

Inequalities for elements in  $\widehat{\mathbf{P}}$  follow by taking limits of inequalities for **P**. For instance, Newton's inequalities (4.3.2) become  $a_k^2 \ge a_{k-1}a_{k+1}$ . Using

this observation, we derive Turan's inequality: If  $P_n(x)$  is the n-th Legendre polynomial, then

$$P_n(x)^2 - P_{n+1}(x)P_{n-1}(x) \ge 0 \quad \text{ for } |x| < 1 \tag{5.2.1}$$

To see this, we need the identity

$$\sum_{n=0}^{\infty} \mathsf{P}_n(x) \frac{y^n}{n!} = e^{xy} J_0(y\sqrt{1-x^2})$$

Thus, if  $|\alpha| < 1$  the series  $\sum P_n(\alpha) \frac{y^n}{n!}$  is in  $\widehat{\mathbf{P}}$ . An application of Newton's inequalities (4.3.3) yields (5.2.1). See [169].

**Example 5.8.** *The Gamma function* Although the Gamma function is meromorphic, its reciprocal is an entire function with the product [138, page 242]:

$$\frac{1}{\Gamma(z+1)} = e^{\gamma z} \left(1+\frac{z}{1}\right) e^{-\frac{z}{1}} \left(1+\frac{z}{2}\right) e^{-\frac{z}{2}} \cdots \left(1+\frac{z}{n}\right) e^{-\frac{z}{n}} \cdots$$

where  $\gamma$  is Euler's constant. The roots are at the negative integers  $-1, -2, \ldots$ . Moreover, if we choose a positive integer k then the quotient

$$G(z) = \frac{\Gamma(z+1)}{\Gamma(kz+1)}$$

is an entire function since the poles of the numerator are contained among the poles of the denominator. In addition, G(z) is in  $\hat{\mathbf{P}}$  since it has a product formula.

*Remark* 5.9. There is an explicit description [14, 138] of the elements of  $\widehat{\mathbf{P}}$  that is slightly more complicated than the description of the elements of  $\widehat{\mathbf{P}^{\text{pos}}}$ . The elements of  $\widehat{\mathbf{P}}$  have the product representation

$$cx^{r}e^{ax-bx^{2}}\prod_{i=1}^{\infty}(1+c_{i}x)e^{-c_{i}x}$$
 (5.2.2)

where  $a, c_1, \ldots$  are real, r is a non-negative integer, and b is non-negative. Notice that  $e^{x^2}$  is not in  $\widehat{\mathbf{P}}$ .

We can use trigonometric identities to find some non-obvious members of  $\widehat{\mathbf{P}}.$  Recall that

$$\frac{1}{2}\sin\frac{x+y}{2}\,\cos\frac{x-y}{2} = \sin x + \sin y$$

If we choose a so that  $|a| \leq 1$ , then we can find y so that sin y = a, and hence

$$\sin(x) + a = \frac{1}{2}\sin\frac{x+a}{2}\,\cos\frac{x-a}{2}$$

Since the right hand side is a product of members of  $\widehat{\mathbf{P}}$ , it follows that  $a + \sin x \in \widehat{\mathbf{P}}$ . Multiplying several of these together yields half of

**Lemma 5.10.** Suppose that f(x) is a polynomial. Then,  $f(x) \in \mathbf{P}^{[-1,1]}$  if and only if  $f(\sin x) \in \widehat{\mathbf{P}}$ .

*Proof.* Conversely, let  $r_1, \ldots, r_n$  be the roots of f(x). Then the roots of  $f(\sin x)$  are solutions to  $\sin x = r_i$ . If any  $r_i$  has absolute value greater than 1 or is not real, then  $\arcsin r_i$  is also not real, so  $f(\sin x)$  has non-real roots.

### 5.3 The Hadamard Factorization Theorem

How can we show that a function f is in  $\hat{\mathbf{P}}$  or  $\hat{\mathbf{P}^{\text{pos}}}$ ? It is not enough to know that all the roots are real - for example  $e^{x^2}$  has no complex roots and yet is not in  $\hat{\mathbf{P}}$ . If all roots of f are real we can determine if f is in  $\hat{\mathbf{P}}$  using the coefficients of the Taylor series.

The Hadamard factorization theorem represents an arbitrary entire function as an infinite product times an exponential factor. We first summarize the necessary background (see [149] or [15]), and then use the machinery to exhibit a collection of functions whose exponential part satisfies the necessary conditions to belong to  $\widehat{\mathbf{P}}$ .

We begin with definitions. If p is a non-negative integer, we define the *Weierstrass primary factor of order* p by

$$E(z, 0) = (1 - z)$$
$$E(z, p) = (1 - z) \exp(z + \frac{z^2}{2} + \dots + \frac{z^p}{p})$$

For a set Z of complex numbers, *the genus of* Z, p = p(Z) is defined by

$$p(Z) = \inf \left\{ q : q \text{ is an integer, } \sum_{z_i \in Z} \frac{1}{|z_i|^{q+1}} < \infty \right\}$$

The canonical product over Z is defined by

$$\mathsf{P}_{\mathsf{Z}}(\mathsf{z}) = \prod_{z_{\mathsf{i}} \in \mathsf{Z}} \mathsf{E}\left(\frac{\mathsf{z}}{\mathsf{z}_{\mathsf{i}}}, \mathsf{p}\right)$$

where p = p(Z). The maximum modulus of f(z) on the circle |z| = r is denoted M(r). The *order*  $\rho = \rho(f)$  *of* f(z) is

$$\rho = \limsup_{r \to \infty} \frac{\log \log M(r)}{\log r}$$

We can now state

**Theorem 5.11** (Hadamard Factorization Theorem). *If* f(z) *is an entire function of order*  $\rho$  *with an* m*-fold zero at the origin, we have* 

$$f(z) = z^{m} e^{Q(z)} P_{Z}(z)$$

where Q(z) is a polynomial of degree at most  $\rho$ , and Z is the set of zeros of f.

The particular case we are interested in is  $\rho < 1$  and p = 0. In this case

$$f(z) = z^{m} e^{\alpha z} \prod_{i} \left( 1 - \frac{z}{z_{i}} \right)$$
(5.3.1)

If all the roots  $z_i$  are negative then  $f \in \widehat{\mathbf{P}^{\text{pos}}}$ , and  $f \in \widehat{\mathbf{P}}$  if there are positive and negative roots. The main difficulty is showing that all the roots are real.

How can we determine the two parameters  $\rho$  and p? First of all, it is known that  $p \leq \rho$ . Second, it is possible to find  $\rho$  from the Taylor series of f. Suppose that

$$f(z) = \sum_{i=0}^{\infty} a_i z^i$$

If the limit

$$\limsup_{n \to \infty} \frac{n \log n}{\log(1/|a_n|)}$$
(5.3.2)

is finite then its value is  $\rho$ .

We are in a position to determine the exponential part. We begin with the Bessel function  $J_0(-i\sqrt{x})$ .

#### Example 5.12. The Bessel function.

$$J_0(-i\sqrt{z}) = \sum_{i=0}^{\infty} \frac{z^i}{i!\,i!} \in \widehat{\mathbf{P}^{\text{pos}}}$$

We first recall Stirling's formulas:

$$n! \sim \sqrt{2\pi n} n^n e^{-n}$$
$$\log n! \sim \frac{1}{2} \log(2\pi n) + n \log n - n$$

The limit (5.3.2) is

$$\limsup_{n \to \infty} \frac{n \log n}{\log(n! n!)} = \limsup_{n \to \infty} \frac{n \log n}{2(\frac{1}{2}\log(2\pi n) + n \log n - n)} = \frac{1}{2}$$

Thus  $\rho = \frac{1}{2}$ , and hence p = 0. This shows that we have the desired representation (5.3.1), and since all terms are positive there are no positive zeros. Since the Bessel function is known to have all real zeros the series is in  $\widehat{\mathbf{P}^{\text{pos}}}$ .

The general Bessel function has a series expansion

$$J_{\nu}(z) = \sum_{k=0}^{\infty} \frac{(-1)^k}{\Gamma(k+\nu+1)\,k!} \,\left(\frac{z}{2}\right)^{\nu+2k}$$
(5.3.3)

 $J_{\nu}(z)$  has infinitely many real roots for all  $\nu$ , and all of its roots are real for  $\nu \ge -1$ . Since a similar computation shows that p = 0 for  $J_{\nu}(z)$ , it follows that  $J_{\nu} \in \widehat{\mathbf{P}}$  for  $\nu \ge -1$ .

We now give some examples of computations of  $\rho$ ; since we don't know if the functions have all real roots, we can't conclude that they are in  $\mathbf{P}$ .

**Example 5.13.** Suppose that  $a_1, \ldots, a_r$  and  $b_1, \ldots, b_s$  are positive. Then

$$\sum_{n=0}^{\infty} \frac{(a_1n)!\cdots(a_rn)!}{(b_1n)!\cdots(b_sn)!} z^n \text{ has } p < 1$$

if  $(\sum_{1}^{s} \mathfrak{b}_{k}) - (\sum_{1}^{r} \mathfrak{a}_{k}) > 1$ . Compute the limit using Stirling's formula:

$$\limsup \frac{n \log n}{\log \prod(b_k n)! - \log \prod(a_k n)!} = \limsup \frac{n \log n}{\sum b_k n \log b_k n - \sum a_k n \log a_k n}$$
$$= \frac{1}{\sum b_k - \sum a_k}$$

If the hypothesis are satisfied, then the last expression is less than 1, and p < 1.

**Example 5.14.** Here is a similar example. Suppose that  $a_1, \ldots, a_r$  and  $b_1, \ldots, b_s$  are positive. We define the hypergeometric series

$${}_{r}F_{s}(a_{1},\ldots,a_{r};b_{1},\ldots,b_{s};z) = \sum \frac{\langle \underline{a_{1}} \rangle_{n} \cdots \langle \underline{a_{r}} \rangle_{n}}{\langle \underline{b_{1}} \rangle_{n} \cdots \langle \underline{b_{s}} \rangle_{n}} \frac{z^{n}}{n!}$$

If we compute the limit as above, we find that it is  $\frac{1}{1+s-r}$ . If s-r > 0 then p < 1. Hurwitz showed that the series

$$_{0}\mathsf{F}_{\mathsf{s}}(;\mathsf{b}_{1},\ldots,\mathsf{b}_{\mathsf{s}};z) = \sum \frac{1}{\left\langle \underline{\mathsf{b}}_{1}\right\rangle_{\mathsf{n}}\cdots\left\langle \underline{\mathsf{b}}_{\mathsf{s}}\right\rangle_{\mathsf{n}}} \frac{z^{\mathsf{n}}}{\mathsf{n}!} \tag{5.3.4}$$

has all real roots if  $b_1 \ge -1$  and all the other  $b_i$  are positive.

Example 5.15. A q-series. Consider

$$\sum_{\mathfrak{i}=0}^{\infty} \frac{z^{\mathfrak{i}}}{\mathfrak{q}^{\binom{\mathfrak{i}}{2}} \mathfrak{i}!}$$

The series converges if |q| > 1, or q = -1. The limit (5.3.2) in either case is 0:

$$\limsup_{n \to \infty} \frac{n \log n}{\log(n! q^{\binom{n}{2}})} = \limsup_{n \to \infty} \frac{n \log n}{n \log n + \frac{n(n-1)}{2} \log q} = 0$$

and so  $p = \rho = 0$  and again the representation (5.3.1) holds. If q > 1 all terms are positive, and if q = -1 there are positive and negative signs.

**Example 5.16.** A q-factorial. Consider the two series for q > 1.

$$\sum_{i=0}^{\infty} \frac{z^{i}}{(q;q)_{i} i!} \quad \text{and} \quad \sum_{i=0}^{\infty} \frac{z^{i}}{[i]! i!}$$

The limit for the first series is

 $\limsup_{n \to \infty} \frac{n \log n}{n \log n + \sum_{i=0}^n \log(1-\mathfrak{q}^i)} = \limsup_{n \to \infty} \frac{n \log n}{n \log n + \frac{n(n-1)}{2} \log \mathfrak{q}} = 0.$ 

The terms alternate in sign the first series. If q < -1 then all terms of the second series are positive.

**Example 5.17.** Suppose r > 0. If we compute (5.3.2) for

$$(1-x)^{-r} = \sum_{n=0}^{\infty} \binom{r+n-1}{r} x^n$$

we find that  $\rho$  is infinite, which isn't surprising since this is not an entire function. However,

$$EXP(1-x)^{-r} = \sum_{n=0}^{\infty} {\binom{r+n-1}{r}} \frac{x^n}{n!}$$

has  $\rho = 1$ . A result of Pólya-Schur (Lemma 15.25) says that if  $f \in \widehat{\mathbf{P}}$ , and all coefficients are non-negative then  $f \in \widehat{\mathbf{P}^{\text{pos}}}$ . If we knew that  $\text{EXP}(1-x)^{-r} \in \widehat{\mathbf{P}}$  then we could conclude that  $\text{EXP}(1-x)^{-r} \in \widehat{\mathbf{P}^{\text{pos}}}$ .

*Remark* 5.18. Consider  $f = \sum (\underline{\alpha})_i x^i$ . Since  $EXPf = (1+x)^{\alpha}$  we see that  $EXPf \notin \widehat{\mathbf{P}}$ . However, when  $\alpha = 1/2$  we have the identity that  $EXP^2f = e^{x/2}I_0(x/2)$  where  $I_0$  is the modified Bessel function of the first kind. Since

$$I_0(x) = \sum_{i=0}^{\infty} \frac{x^i}{i! \, i!} = \exp e^x$$

we have that  $EXP^2 f \in \widehat{\mathbf{P}}$ .

#### 5.4 Polynomials with rapidly decreasing coefficients

If the coefficients of a polynomial are decreasing sufficiently rapidly then the polynomial has all real roots. To motivate this, suppose that  $f \in \mathbf{P}(n)$ , and define  $g(x) = f + \alpha x^{n+1}$ . If  $\alpha$  is very small then the graph of g looks like Figure 5.1. There are n roots of g that are close to the roots of f, so g has all real roots.

The following theorem of Kurtz [110] gives a sufficient condition for the existence of all real roots.

**Theorem 5.19.** Let  $f = \sum a_i x^i$  be a polynomial of degree n > 1 with all positive coefficients. If

$$a_i^2 - 4a_{i-1}a_{i+1} > 0$$
  $i = 1, 2, ..., n-1$  (5.4.1)

then all roots of f are real and distinct.



Figure 5.1: A polynomial with rapidly decreasing coefficients

If we have equality in (5.4.1) and  $a_0 = 1$  then the polynomial is

$$f(x) = \sum_{k=0}^{n} x^k 4^{-\binom{n}{2}}$$

and the coefficients are rapidly decreasing.

**Corollary 5.20.** If  $\alpha \ge 4$  then the series below is in  $\hat{P}$  and all of its partial sums are in P.

$$\sum_{k=0}^{\infty} x^k \alpha^{-\binom{n}{2}}$$

**Corollary 5.21.** If  $\alpha \ge 2$  then then the series below is in  $\widehat{P}$  and all of its partial sums are in P.

$$\sum_{k=0}^{\infty} x^k \alpha^{-k^2}$$

*Proof.* Use the fact that  $\alpha \ge 2$  to check that

$$(a^{-k^2})^2 \ge 4a^{-(k-1)^2}a^{-(k+1)^2}$$

**Corollary 5.22.** If  $q \ge 4$  then then the series below is in  $\hat{P}$  and all of its partial sums are in P.

$$\sum_{k=0}^{\infty} \frac{x^k}{[k]!}$$

Proof. The Newton quotient is

$$\frac{[n+2]!\,[n]!}{([n+1]!)^2} = \frac{[n+2]}{[n+1]} = \frac{q^{n+2}-1}{q^{n+1}-1} > q \ge 4.$$

Corollary 5.20 is in [138, V 176]. In [139, III 200] they show that there is exactly one root in each annulus  $\alpha^{2n-2} < |z| < \alpha^{2n}$  for n = 1, 2, ... There is a similar result due to Kuperberg [109].

**Lemma 5.23.** If  $q \ge 4$  then the series below is in  $\widehat{P}$ , and all partial sums are in P.

$$\sum_{k=0}^{\infty} \frac{x^{k}}{(q;q)} = \sum_{k=0}^{\infty} \frac{x^{k}}{(1-q)(1-q^{2})\cdots(1-q^{k})}$$

*Proof.* Kuperberg proves this without reference to Kurtz's result. He shows that if  $F_{q,n}$  is the sum of the first n terms of the series and  $q \ge 4$  then  $F_{q,n}(q^{k+\frac{1}{2}})$  alternates in sign as k ranges from 0 to n.

If all the partial sums of an infinite series have all real roots, then the coefficients satisfy a quadratic exponential bound, but the constant is  $\sqrt{2}$ , not 4 as we would like.

**Lemma 5.24.** If  $\sum a_i x^i$  has all positive terms, and all partial sums have all real roots then

$$a_{n} \leqslant a_{0} \left(\frac{a_{1}}{a_{0}}\right)^{n} \left(2^{\binom{n}{2}}n!\right)^{-1}$$
(5.4.2)

*Proof.* Since  $a_0 + a_1x + \cdots + a_nx^n$  has all real roots we can apply Newton's inequalities (4.3.2)

$$\begin{aligned} \frac{a_{n-1}^2}{a_n a_{n-2}} &\ge 2\frac{n}{n-1} \\ \frac{a_{n-1}}{a_n} &\ge 2\frac{n}{n-1} \frac{a_{n-2}}{a_{n-1}} \ge 2\frac{n}{n-1} 2\frac{n-1}{n-2} \frac{a_{n-3}}{a_{n-2}} \ge \cdots \\ \frac{a_{n-1}}{a_n} &\ge 2^{n-1} n \frac{a_0}{a_1} \\ a_n &\leqslant \left(\frac{a_1}{a_0}\right) 2^{-n-1} \frac{1}{n} a_{n-1} \leqslant \left(\frac{a_1}{a_0}\right)^2 2^{-n-1-(n-2)} \frac{1}{n(n-1)} a_{n-2} \leqslant \cdots \end{aligned}$$

from which (5.4.2) follows.

We end the section with a result where the terms are increasing.

**Lemma 5.25.** Suppose  $0 \leq a_0 \leq a_1 \leq a_2 \cdots \leq \cdots$ . Then the polynomials

$$f_{n} = \sum_{i=0}^{n} a_{i} \left(-1\right)^{i} \left(\underline{x}\right)_{i} \quad \textit{and} \quad g_{n} = \sum_{i=0}^{n} a_{i} \left< \underline{x} \right>_{i}$$

*have all real roots. In addition,*  $f_{n+1} \leq f_n$  *and*  $g_{n+1} \leq g_n$ .

*Proof.* The second case follows from the first upon substituting -x for x in the first case. Note that if k is an integer satisfying  $1 \le k \le n$  then  $f_n(k) = \sum_{0}^{k-1} a_i(-1)^i(\underline{k})_i$ . As a function of i the sequence  $\{a_i\}$  is non-decreasing, and  $\{(\underline{k})_i\}$  is increasing, so the sign of  $f_n(k)$  is the sign of the largest term, and hence is  $(-1)^{k-1}$ . Consequently,  $f_n$  sign interlaces  $(\underline{x})_n$ , and so in in **P**. Since  $f_{n+1} = f_n + a_{n+1}(-1)^{n+1}(\underline{x})_{n+1}$  it follows that  $f_{n+1} \le f_n$ .

### 5.5 Interlacing of analytic functions

We use Proposition 1.35 to motivate our definition of interlacing for analytic functions in  $\hat{\mathbf{P}}$ .

**Definition 5.26.** If f,  $g \in \widehat{\mathbf{P}}$  then we say that f and g *interlace* if  $f + \alpha g$  is in  $\widehat{\mathbf{P}}$  for all real  $\alpha$ .

Later in this section we will relate interlacing of analytic functions to the interlacing properties of their zeros. Although it might be hard to verify directly that a linear combination is in  $\hat{\mathbf{P}}$  we can establish interlacing using limits.

**Lemma 5.27.** Suppose that  $f, g \in \widehat{P}$  and we have sequences of polynomials  $(f_n), (g_n)$  where the degree of  $f_n$  and  $g_n$  is n such that  $f_n \longrightarrow f$  and  $g_n \longrightarrow g$ . If  $f_n$  and  $g_n$  interlace for every n then f and g interlace.

*Proof.* Since  $f_n$  and  $g_n$  interlace we know that  $f_n + \alpha g_n$  is in **P** for all  $\alpha$ . Taking limits, we see that  $f + \alpha g$  is in  $\widehat{\mathbf{P}}$  and hence f, g interlace.

Just as for polynomials, the derivative is interlacing.

**Lemma 5.28.** If  $f \in \widehat{P}$  then  $f' \in \widehat{P}$ , and f and f' interlace.

*Proof.* If  $f_n$  is a sequence of polynomials that converge to f, then the derivatives  $f'_n$  converge to f'. Since  $f_n + \alpha f'_n$  converges to  $f + \alpha f'$  it follows that f and f' interlace.

*Remark* 5.29. Exponential functions  $e^{\alpha x}$  and  $e^{\beta x}$  only interlace if  $\alpha = \beta$ . Indeed, if they did interlace, then  $e^{\alpha x} + e^{\beta x} \in \widehat{\mathbf{P}}$ . However,

$$e^{\alpha x} + e^{\beta x} = e^{\alpha x} (1 + e^{(\beta - \alpha)x})$$

has complex roots if  $\alpha \neq \beta.$  The next three results generalize this simple observation.

We first see that no linear combination of polynomials and exponentials is in  $P_2$ .

**Lemma 5.30.** If f(x), g(x) are polynomials and  $g(x)e^{x} + f(x) \in \widehat{P}$  then f = 0 and  $g \in P$ .
*Proof.* We first show that if  $c \in \mathbb{R}$  then  $e^x + c \notin \mathbf{P}$ . If it were then  $(e^{\mathsf{D}} + c)(x^n) = (x+1)^n + cx^n \in \mathbf{P}$ , but the roots are complex for  $n \ge 3$ .

Next, if  $e^x + f(x) \in \widehat{\mathbf{P}}$  where f has degree d then the d'th derivative is in  $\widehat{\mathbf{P}}$  and is of the form  $e^x + c$ . This contradicts the first paragraph.

If  $g(x)e^x + c \in \widehat{\mathbf{P}}$  then replacing x by -x and multiplying by  $e^x$  shows that  $g(-x) + c e^x \in \widehat{\mathbf{P}}$ , which we know is not possible.

Finally, if  $g(x)e^{x}+f(x) \in \widehat{\mathbf{P}}$  then the d'th derivative is of the form  $G(x)e^{x}+c$ , so f must be zero. Multiplying by  $e^{-x}$  shows that  $g \in \mathbf{P}$ .

Next, it is easy to describe the polynomials interlacing  $e^x$ .

**Lemma 5.31.** If  $f \sim e^x$  then  $f = (\alpha x + \beta)e^x$ .

*Proof.* The hypothesis means that  $f(x) + \alpha e^x \in \widehat{\mathbf{P}}$  for all real  $\alpha$ . Thus  $(f + \alpha e^x)(\mathbf{D})x^n \in \mathbf{P}$  for all positive integers n. It follows that

$$f(\mathsf{D})x^{\mathsf{n}} \sim e^{\mathsf{D}}x^{\mathsf{n}} = (x+1)^{\mathsf{n}}$$

We can determine f since we know all polynomials interlacing  $(x + 1)^n$ . Since  $f(D)x^n$  has degree at most n it follows that there are  $a_n$ ,  $b_n$  such that

$$f(D)x^{n} = a_{n}(x+1)^{n} + b_{n}(x+1)^{n-1}.$$

If we write  $f(x) = \sum d_i x^i$  then equating terms yields

$$d_{i}\mathsf{D}^{i}x^{n} = d_{i}x^{n-i}\frac{n!}{(n-i)!} = x^{n-i}\left[a_{n}\binom{n}{n-i} + b_{n}\binom{n-1}{n-i}\right]$$

and so

$$\mathbf{d}_{\mathbf{i}} = \frac{\mathbf{a}_{\mathbf{n}}}{\mathbf{i}!} + \frac{\mathbf{b}_{\mathbf{n}}}{\mathbf{n}} \frac{1}{(\mathbf{i}-1)!}.$$

Setting i = 0 shows that  $a_n = d_0$ , and i = 1 yields  $b_n/n = d_1 - d_0$ . Thus

$$d_{\mathfrak{i}} = d_0 \frac{1}{\mathfrak{i}!} + (d_1 - d_0) \frac{1}{(\mathfrak{i} - 1)!}$$

and consequently

$$f(x) = d_0 e^x + (d_1 - d_0) x e^x.$$

More generally we have

**Lemma 5.32.** If  $f(x) \sim g(x)e^x$  where g(x) is a polynomial then there is a polynomial  $h \sim g$  such that  $f(x) = h(x)e^x$ .

*Proof.* We first show that it suffices to show that  $f(x) = H(x)e^x$  for some polynomial. If so, then since  $f + \alpha ge^x \in \widehat{\mathbf{P}}$  it follows that  $(H + \alpha g)e^x \in \widehat{\mathbf{P}}$ , and therefore  $g + \alpha H \in \mathbf{P}$ . Thus, g and H interlace.

Assume that g has degree r, and write  $f = \sum d_i x^i$ . If n > r + 1 then  $g(D)e^Dx^n = g(D)(x+1)^n$  has a factor of  $(x+1)^2$ . Since  $f(D)x^n \sim g(D)(x+1)^n$  it follows that -1 is a root of  $f(D)x^n$ . This gives us the equations, valid for n > r + 1:

$$\left(\sum d_{i}\mathsf{D}^{i}x^{n}\right)(-1) = n! \sum d_{i}\frac{(-1)^{n-i}}{(n-i)!} = 0$$

that we view as an infinite matrix equation. For example, if r = 1 we have

$$\begin{pmatrix} -1/6 & 1/2 & -1 & 1 & 0 & 0 & \dots \\ 1/24 & -1/6 & 1/2 & -1 & 1 & 0 & \dots \\ -1/120 & 1/24 & -1/6 & 1/2 & -1 & 1 & \dots \\ \vdots & & & & \ddots \end{pmatrix} \begin{pmatrix} d_0 \\ d_1 \\ d_2 \\ \vdots \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ \vdots \end{pmatrix}$$
(5.5.1)

In this example it is clear that we can solve for  $d_3, d_4, \ldots$  in terms of  $d_0, d_1, d_2$ . If M is the lower triangular matrix whose ij entry is  $(-1)^{i-j}/(i-j)!$  then the matrix above if formed by removing the first three rows of M. Now the inverse of M has entries 1/(i-j)!, and if we multiply (5.5.1) by  $M^{-1}$  we recove the identity matrix after r + 1 columns. In the example, the result of multiplying by  $M^{-1}$  is

$$\begin{pmatrix} -1/6 & 1/2 & -1 & 1 & 0 & 0 & 0 & \dots \\ -1/8 & 1/3 & -1/2 & 0 & 1 & 0 & 0 & \dots \\ -1/20 & 1/8 & -1/6 & 0 & 0 & 1 & 0 & \dots \\ -1/72 & 1/30 & -1/24 & 0 & 0 & 0 & 1 & \dots \\ \vdots & \vdots & \vdots & \ddots & & \end{pmatrix} \begin{pmatrix} d_0 \\ d_1 \\ d_2 \\ d_3 \\ \vdots \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ \vdots \end{pmatrix}$$

Consequently we find

Adding up the series determines f:

$$\begin{split} d_0 + d_1 x + d_2 x^2 + d_2 \sum_{n=3}^{\infty} \frac{x^n}{(n-2)!} - d_1 \sum_{n=3}^{\infty} x^n \frac{n-2}{(n-1)!} + d_0 \sum_{n=3}^{\infty} x^n \frac{\binom{n-1}{2}}{n!} \\ &= e^x \left( d_2 x^2 + d_1 (x - x^2) + d_0 (1 - x + \frac{1}{2} x^2) \right) \end{split}$$

which shows that f is  $e^x$  times a quadratic.

In the general case we follow the same approach, and find that f equals

$$e^{x} \sum_{i=0}^{r+1} d_{i} x^{i} \left( \sum_{k=0}^{r+1-i} (-1)^{k} \frac{x^{k}}{k!} \right)$$

which is a polynomial in x of degree r + 1 times an exponential, as desired.

We now define interlacing of zeros:

**Definition 5.33.** If f and g are in  $\hat{\mathbf{P}}$  then we say that *the zeros of* f *and* g *interlace* if there is a zero of f between every pair of zeros of g, and a zero of g between every two zeros of f. If the set of zeros of f and g have least zeros  $z_f$  and  $z_g$  then we say that  $f \leq g$  if f and g interlace, and  $z_f \leq z_g$ .

If the zeros of functions in  $\widehat{\mathbf{P}^{\text{pos}}}$  interlace and the exponential factors are the same, then the functions interlace. We would like to prove the converse.

#### Lemma 5.34. Suppose that

$$\mathsf{f} = z^{\mathrm{r}} e^{\gamma x} \prod_{i=1}^{\infty} (1 + c_i x) \qquad \mathsf{g} = x^{\mathrm{r}} e^{\gamma x} \prod_{i=1}^{\infty} (1 + d_i x)$$

are in  $\widehat{P^{pos}}$  where  $c_1 > c_2 > \cdots > 0$  and  $d_1 > d_2 > \cdots > 0$ . Then f and g interlace if

 $c_1 > d_1 > c_2 > d_2 \dots \quad \textit{or} \quad d_1 > c_1 > d_2 > c_2 \dots$ 

*Proof.* Since the  $c_i$ 's and the  $d_i$ 's interlace then the partial products

$$f_n = \prod_{i=1}^n (1 + c_i x) \qquad g_n = \prod_{i=1}^n (1 + d_i x)$$

interlace, and converge to interlacing functions. If we then multiply these interlacing functions by  $x^r e^{\gamma x}$  we find that f and g interlace.

We can generalize part of Lemma 1.20.

**Lemma 5.35.** Suppose that  $f(x) = \prod (1-x/a_i)$  is in  $\widehat{P^{pos}}$ . Choose positive  $\alpha_i$  such that  $\sum \alpha_i < \infty$ . Then

1. 
$$g = \sum \alpha_i \frac{f}{1 - x/a_i} \in \widehat{P^{pos}}$$
  
2.  $f \text{ and } \sum \alpha_i \frac{f}{1 - x/a_i} \text{ interlace.}$ 

Proof. Consider

$$f_n = \prod_{i=1}^n (1 - x/a_i)$$
 and  $g_n = \sum_{i=1}^n \alpha_i \frac{f}{1 - x/a_i}$ 

We know that  $f_n \leq g_n$ . The assumption on  $\alpha_i$  implies that  $g_n$  converges uniformly to g. The result now follows by taking limits.

# 5.6 Characterizing $\widehat{P}$

The following theorem shows that the functions in  $\mathbf{P}$  are exactly the right functions to generalize polynomials.

**Theorem 5.36.** An analytic function f satisfies  $f(D)P \subset P$  if and only if f is in  $\widehat{P}$ . It satisfies  $f(D)P^{\text{pos}} \subset P^{\text{pos}}$  if and only if it is in  $\widehat{P^{\text{pos}}}$ .

*Proof.* If f is in  $\hat{\mathbf{P}}$ , then it is the uniform limit of polynomials. The containment is true for each of the polynomials by Corollary 7.15, so it is true for f. The converse is more interesting, and follows an argument due to Pólya and Schur[137]. First write  $f(x) = \sum a_i x^i$ , and set  $q_n = f(D)x^n$ . Expanding the series

$$q_n = \sum_{i=0}^n a_i(\underline{n})_i x^{n-i}$$

and reversing  $q_n$  gives a polynomial in **P** 

$$p_n = \sum_{i=0}^n a_i(\underline{n})_i x^i$$

If we now replace x by x/n we find

$$p_n(x/n) = \sum_{i=0}^n a_i x^{n-i} \prod_{k=1}^i (1 - \frac{k}{n})$$

The polynomials  $p_n(x/n)$  converge uniformly to f, and so f is in  $\widehat{\mathbf{P}}$ . The second case is similar.

#### 5.7 P-positive semi-definite matrices

We derive some simple inequalities for quadratic forms involving the coefficients of a polynomial with all real roots. It is interesting that these inequalities are valid for a broader class of functions than just  $\hat{\mathbf{P}}$ .

Define an n by n matrix Q to be *P*-positive semi-definite if for every  $f = \sum_{0}^{n} a_{i}x^{i} \in \mathbf{P}$  and  $A = (a_{0}, ..., a_{n-1})$ , we have  $AQA^{t} \ge 0$ . Of course, any positive semi-definite matrix is also **P**-positive semi-definite, but the converse is not true. In particular, **P**-positive semi-definite matrices can have negative entries. Indeed, Newton's inequalities imply that the matrix  $\begin{pmatrix} 0 & 0 & -2 \\ 0 & 1 & 0 \\ -2 & 0 & 0 \end{pmatrix}$  is **P**-positive semi-definite.

We first note that it is easy to construct new **P**-positive semi-definite matrices from old ones. It's obvious that **P**-positive semi-definite matrices form a cone. In addition,

**Lemma 5.37.** If Q is a **P**-positive semi-definite matrix, and B is the diagonal matrix of coefficients of a polynomial in  $P^{\text{pos}}$ , then BQB is also **P**-positive semi-definite.

*Proof.* If A is the vector of coefficients of  $f \in \mathbf{P}$ , then AB is the vector of coefficients of the Hadamard product of f and g. By Theorem 9.84 the Hadamard product is in  $\mathbf{P}$ , it follows that  $A(BQB)A^t = (AB)Q(AB)^t \ge 0$ .

Here is a construction of a **P**-positive semi-definite matrix. Note that it applies to polynomials in **P**.

**Proposition 5.38.** Suppose that

$$f(y) = \sum_{i=0}^{\infty} a_i y^i = e^{-g(y) + h(y)} \prod_j \left( 1 - \frac{y}{r_j} \right)$$

where h(y) is a polynomial with only terms of odd degree, g(y) has all non-negative coefficients, and all exponents of g are  $\equiv 2 \pmod{4}$ . If m is a positive integer then

$$\sum_{j=-m}^m (-1)^{m+j} a_{m-j} a_{m+j} \geqslant 0$$

Since all real rooted polynomials satisfy the hypothesis, the 2m + 1 by 2m + 1 matrix whose anti-diagonal is alternating  $\pm 1$  beginning and ending with 1 is **P**-positive semi-definite.

*Proof.* The coefficient of  $y^{2m}$  in

$$f(y)f(-y) = \left(\sum a_{s}y^{s}\right)\left(\sum a_{r}(-1)^{r}y^{r}\right)$$

is

$$\sum_{j=-\mathfrak{m}}^{\mathfrak{m}}(-1)^{j}\mathfrak{a}_{\mathfrak{m}-j}\mathfrak{a}_{\mathfrak{m}+j}$$

On the other hand, since h(y) + h(-y) = 0 and g(y) = g(-y) by hypothesis, we have

$$f(\mathbf{y})f(-\mathbf{y}) = \left(e^{-g(\mathbf{y}) + h(\mathbf{y})} \prod \left(1 - \frac{\mathbf{y}}{r_{i}}\right)\right) \left(e^{-g(-\mathbf{y}) + h(-\mathbf{y})} \prod \left(1 + \frac{\mathbf{y}}{r_{i}}\right)\right)$$
$$= e^{-2g(\mathbf{y})} \prod \left(1 - \frac{\mathbf{y}^{2}}{r_{i}^{2}}\right)$$
(5.7.1)

Since  $r_i^2$  is positive, the sign of the coefficient of  $y^{2m}$  in the infinite product is  $(-1)^m$ . The coefficient of  $y^{2m}$  in  $e^{-2g(y)}$  is either  $(-1)^m$  or 0. Consequently, the sign of the coefficient of  $y^{2m}$  in (5.7.1) is also  $(-1)^m$  or 0, which finishes the proof.

The case m = 1 in the following corollary is the Laguerre inequality.

**Corollary 5.39.** *Suppose that*  $f \in P$ *, and* m *is a positive integer. Then* 

$$\sum_{j=-m}^{m} (-1)^{m+j} \frac{f^{(m-j)}(x)}{(m-j)!} \frac{f^{(m+j)}(x)}{(m+j)!} \ge 0$$

More generally, if  $Q = (q_{ij})$  is a **P**-positive semi-definite matrix, then

$$\sum_{i,j} q_{ij} \frac{f^{(i)}}{i!} \frac{f^{(j)}}{j!} \ge 0$$

*Proof.* Use the Taylor series

$$f(x+y) = \sum_{s=0}^{\infty} f^{(s)}(x) \frac{y^s}{s!}$$

**Corollary 5.40.** If  $P_n(x)$  is the Legendre polynomial, m is a positive integer, and |x| < 1 then

$$\sum_{j=-m}^{m} (-1)^{m+j} \frac{P_{m-j}(x)}{(m-j)!} \frac{P_{m+j}(x)}{(m+j)!} \ge 0$$

*Proof.* The following identity is valid for  $|\mathbf{x}| < 1$ .

$$\sum_{n=0}^{\infty} \frac{P_n(x)}{n!} y^n = e^{xy} J_0(y\sqrt{1-x^2})$$

Now apply Proposition 5.38.

*Remark* 5.41. Krasikov[107] essentially shows that the matrix below (derived from his  $V_4(f)$ ) is **P**-positive semi-definite.

$$\begin{pmatrix} 0 & 0 & 0 & 0 & -2 \\ 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 2 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 \\ -2 & 0 & 0 & 0 & 0 \end{pmatrix}$$

To see this directly, notice we can write it as a sum of **P**-positive semi-definite matrices. (0, 0, 0, 0, 1) = (0, 0, 0, 0, 2)

10	0	0	0	1)		10	0	0	0	-31
0	0	0	-1	0		0	0	0	0	0
0	0	1	0	0	+	0	0	1	0	0
0	-1	0	0	0		0	0	0	0	0
$\backslash 1$	0	0	0	0/		\_3	0	0	0	0 /

The first one is **P**-positive semi-definite by the proposition. From equation (4.3.5) we have  $a_2^2 \ge 6a_0a_4$  which shows the second matrix is **P**-positive semi-definite. More generally, if 0 < k < n then Newton's inequalities imply that the matrix with 1 in the (k, k) entry, and  $-\frac{1}{2} \frac{k+1}{k} \frac{n-k+1}{n-k}$  in the (k+1, k-1) and (k-1, k+1) entries is a **P**-positive semi-definite matrix.

CHAPTER



# Linear Transformations of Polynomials

The aim of this chapter is to establish properties of linear transformations that preserve roots. In Section 6.1 we show that if we put assumptions on T such as  $Tf \leq f$  for all  $f \in P$  then Tf is the derivative. If T is bijection on P, then T is affine. We see how to determine the possible domain and range of a linear transformation. In the next section we study transformations of the form  $f \mapsto f(T)(1)$  where T is a linear transformation. The following two sections consider properties of linear transformations that are defined by recurrence relations. We then consider the effect of Möbius transformations on the the roots of polynomials and on transformations. In the final section we begin the study of transformations of the form  $x^n \mapsto \prod_{i=1}^{n} (x + a_i)$ .

#### 6.1 Characterizing transformations

Many linear transformations can be characterized by their mapping and interlacing properties. The derivative can be characterized by its fundamental interlacing property  $f \leq f'$ . This result relies on the fact that the only monic polynomial f such that  $(x + a)^n \leq f$  is  $f = (x + a)^{n-1}$ .

**Theorem 6.1.** If T is a linear transformation on polynomials such that  $f \leq T(f)$  for all polynomials f in  $\mathbf{P}^{\text{alt}}$  (or for all polynomials in  $\mathbf{P}^{\text{pos}}$ ), then T is a multiple of the derivative.

*Proof.* Choosing  $f = x^m$ , we find  $x^m \leq T(x^m)$ , and hence there is a  $c_m$  so that  $T(x^m) = c_m x^{m-1}$ . From  $(x + 1)^m \leq T(x + 1)^m$  we see that there an  $\alpha_m$  such that

$$\sum_{i=1}^{m} \binom{m}{i} c_{i} x^{i-1} = T(x+1)^{m} = \alpha_{m} (x+1)^{m-1} = \alpha_{m} \sum_{i=0}^{m-1} \binom{m-1}{i} x^{i}$$

Equating coefficients shows that  $c_i = (\alpha_m/m)i$ , and so T(f) is a multiple of f'.

**Theorem 6.2.** If T is a linear transformation with the property that Tf and f have the same degree, and interlace for all f in  $P^{alt}$  (or for all f in  $P^{pos}$ ) then there are constants a, b, c such that

$$Tf = af + bxf' + cf'$$

If Tf and f interlace for all  $f \in P$  then Tf = af + cf'.

*Proof.* By continuity  $x^n$  and  $T(x^n)$  interlace, and so there are constants  $a_n$ ,  $b_n$  such that  $T(x^n) = a_n x^n + b_n x^{n-1}$ . Apply T to the polynomial  $f = x^r (x - \alpha)^2$ . We compute

$$Tf = T(x^{r+2} - 2\alpha x^{r+1} + \alpha^2 x^r)$$

$$= a_{r+2}x^{r+2} + b_{r+2}x^{r+1} - 2\alpha a_{r+1}x^{r+1} - 2\alpha b_{r+1}x^r + \alpha^2 a_r x^r + \alpha^2 b_r x^{r-1}$$
(6.1.1)

Now f has a double root at  $\alpha$ , and Tf interlaces f, so Tf must have  $\alpha$  as a root. Substituting  $x = \alpha$  in (6.1.1) gives the recurrence relation

$$0 = \alpha^{r+2}(a_{r+2} - 2a_{r+1} + a_r) + \alpha^{r+1}(b_{r+2} - 2b_{r+1} + b_r)$$

This equation holds for infinitely many  $\alpha$ , so we get the two recurrences

$$0 = a_{r+2} - 2a_{r+1} + a_r$$
  
$$0 = b_{r+2} - 2b_{r+1} + b_r$$

Since these recurrences hold for  $1 \le r \le n-2$ , we solve these equations and find that constants a, b, c, d so that  $a_r = a + br$ ,  $b_r = d + cr$ . Substituting these values gives

$$\mathsf{T}(\mathsf{x}^{\mathsf{r}}) = (\mathsf{a} + \mathsf{b}\mathsf{r})\mathsf{x}^{\mathsf{r}} + (\mathsf{d} + \mathsf{c}\mathsf{r})\mathsf{x}^{\mathsf{r}-1}$$

Since  $rx^r = xDx^r$ , it remains to show that d = 0. We know that for any negative  $\alpha$  the polynomials  $(x + \alpha)^3$  and  $T(x + \alpha)^3$  interlace, and hence  $(x + \alpha)^2$  divides  $T(x + \alpha)^3$ . A computation shows that the remainder of  $T(x + \alpha)^3$  upon division by  $(x + \alpha)^2$  is  $d\alpha(x + 2\alpha)$ . As this remainder is 0 we conclude that d = 0. The case where  $f \in \mathbf{P}^{pos}$  is similar and omitted.

Finally, assume that Tf and f interlace for all  $f \in \mathbf{P}$ . If b is not zero then we may divide Tf by b, and thus assume that Tf = af + xf' + cf'. Choose  $\alpha > |c|$  and consider the polynomial

$$f = (x - \alpha)(x + \alpha)(x - \alpha - 1)(x + \alpha + 1)$$

A computation shows that

$$\begin{aligned} (\mathsf{Tf})(-\alpha) &= 2(\mathsf{c}-\alpha)\alpha(1+2\alpha)\\ (\mathsf{Tf})(\alpha) &= -2(\mathsf{c}+\alpha)\alpha(1+2\alpha) \end{aligned}$$

The signs of the left hand side must alternate since f and Tf sign interlace, and so  $(c - \alpha)(c + \alpha) > 0$  which implies  $|c| > \alpha$ . Since we chose  $\alpha > |c|$  this is a contradiction, and hence b = 0.

The remaining case is  $Tf \leq f$ . Using an argument entirely similar to that of Theorem 6.2 we find

**Theorem 6.3.** *If* T *is a linear transformation such that*  $Tf \leq f$  *for all*  $f \in P$  *then there are constants* a, b, c *where* a *and* c *have the same sign such that* 

$$Tf = axf + bf + cf'$$

*Remark* 6.4. If we restrict the domain then we can find transformations that are not the derivative that satisfy  $Tf \leq f$ . Here are three examples:

$$T(f) = (x^{2} - 1)f' on P^{(-1,1)}$$
  

$$T(f) = D((x^{2} - 1)f) on P^{(-1,1)}$$
  

$$T(x^{n}) = \frac{x^{n+1}}{n+1} on EXP(P)$$

We can generalize Theorem 6.1 by restricting the domain of the transformation to polynomials of degree n. Under these restrictions the polar derivative makes an appearance.

**Theorem 6.5.** *Fix a positive integer* n *and suppose that* T *is a linear transformation that maps homogeneous polynomials of degree* n *to homogeneous polynomials of degree* n - 1 *with the property that*  $F \leq T(F)$  *for all*  $f \in P(n)$  *for which* F(x, 1) = f(x). *There are* b, c *where* bc  $\geq 0$  *such that* 

$$T(F) = b\frac{\partial F}{\partial x} + c\frac{\partial F}{\partial y}$$

*Proof.* Since T is continuous we observe that if F is the limit of polynomials of degree n then F and TF interlace. Considering that

$$\lim_{\epsilon \to 0} x^{r} (1 + \epsilon x)^{n-r} = x^{r}$$

we conclude that  $x^r$  and  $\mathsf{T}(x^r)$  interlace for all  $r\leqslant n.$  Thus there are  $\mathfrak{a}_r, \mathfrak{b}_r, c_r$  such that

$$T(x^{r}) = x^{r-1}(a_{r} + b_{r}x + c_{r}x^{2})$$

Since  $T(x^r(x - \alpha)^2)$  is interlaced by  $x^r(x - \alpha)^2$ , it has  $\alpha$  for a root, and we find

$$0 = \alpha^{r+2}(a_{r+2} + b_{r+2}\alpha + c_{r+2}\alpha^2) - 2\alpha^{r+1}(a_{r+1} + b_{r+1}\alpha + c_{r+1}\alpha^2) + \alpha^{r}(a_r + b_r\alpha + c_r\alpha^2)$$

Since this is true for all  $\alpha$  we find that

$$0 = a_{r+2} - 2a_{r+1} + a_r$$
  

$$0 = b_{r+2} - 2b_{r+1} + b_r$$
  

$$0 = c_{r+2} - 2c_{r+1} + c_r$$

Solving these recurrences yields  $a_r = A_0 + A_1 r$ ,  $b_r = B_0 + B_1 r$ ,  $c_r = C_0 + C_1 r$  which implies

$$\mathsf{T}(x^r) = (\mathsf{A}_0 + \mathsf{A}_1 r) x^{r-1} + (\mathsf{B}_0 + \mathsf{B}_1 r) x^r + (\mathsf{C}_0 + \mathsf{C}_1 r) x^{r+1}$$

Since  $T(x^n)$  is a polynomial of degree n-1 we find  $B_0+B_1n=0$  and  $C_0+C_1n=0$  so that

$$T(x^{r}) = (A_{0} + A_{1}r)x^{r-1} + B_{1}(r-n)x^{r} + C_{1}(r-n)x^{r+1}$$

Since

$$\begin{split} \mathsf{T}(x^{n-1}(x+1)) &= (\mathsf{A}_0 + \mathsf{A}_1 \mathfrak{n})(x^{n-1}) + (\mathsf{A}_0 + \mathsf{A}_1(\mathfrak{n}-1))x^{n-2} - \\ & \mathsf{B}_1 x^{n-1} - \mathsf{C}_1 x^n \end{split}$$

has degree n - 1 we see that  $C_1 = 0$ . It remains to see that  $A_0 = 0$ . If not, we may assume  $A_0 = 1$  and consider

$$T((x - \alpha)^3) = (x^2 - 3x\alpha + 3) + 3A_1(x - \alpha)^2 - B_1(-3x^2\alpha + 6x\alpha^2 - 3\alpha^3)$$

whose discriminant is  $3\alpha^2(-1 - 4A_1 - 4B_1\alpha)$ . Since  $\alpha$  is arbitrary this can be negative unless  $B_1 = 0$ . Thus  $T(x^r) = x^{r-1} + A_1rx^{r-1}$ . We see that TF has degree r - 1 so F < TF. The proof of Theorem 6.1 shows that this can not happen, so  $A_0$  is 0, and  $T(x^r) = A_1(rx^{r-1}) + B_1(r-n)x^r$  which implies the conclusion by linearity.

If f is a polynomial then  $f(D)(x + a)^n$  is a sum of constants times powers of x + a. The next lemma is a converse.

**Lemma 6.6.** Suppose  $T: P \longrightarrow P$  satisfies

$$\mathsf{T}(x+a)^k = \sum_{i=0}^k \alpha_{k,i} \, (x+a)^i$$

for all  $a \in \mathbb{R}$ , where  $\alpha_{k,i}$  are constants not depending on a. Then there is an  $f \in \widehat{P}$  such that T = f(D).

*Proof.* Continuity of T implies that D and T commute:

$$TD(x + a)^{k} = T \lim_{\varepsilon \to 0} \frac{(x + a + \varepsilon)^{k} - (x + a)^{k}}{\varepsilon}$$
$$= \lim_{\varepsilon \to 0} \frac{\sum_{i} \alpha_{k,i} (x + a + \varepsilon)^{i} - \sum_{i} \alpha_{k,i} (x + a)^{i}}{\varepsilon}$$
$$= \sum_{i} \alpha_{k,i} (x + a)^{k-1} = DT(x + a)^{k}$$

Now, any linear operator commuting with differentiation is a power series f(x) in D. Since f(D) maps P to itself, it follows from Theorem 5.36 that  $f \in \widehat{P}$ .

The proof of the following lemma (due to Carncier-Pinkus-Peña[30]) clearly shows the significance of the monic hypothesis. The proof reduces to the fact that we know all the polynomials interlacing  $(x + 1)^n$ .

**Lemma 6.7.** *If*  $T : P(n) \longrightarrow P(n)$  *satisfies* 

1.  $T(x^k) = x^k + terms$  of lower degree for  $0 \le k \le n$ .

2. 
$$T(\mathbf{x}^k) \in \mathbf{P}^{pos}$$
 for  $0 \leq k \leq n$ .

then there is a polynomial f such that

- 1. T = f(D)
- 2.  $EXP(f) \in \mathbf{P}$

*Proof.* Let  $T(x^k) = \sum_{0}^{k} a_{k,i} x^o$  where  $a_{k,k} = 1$ . We first describe the argument for n = 3. Consider  $T_*(x + y)^3$ :

$$y^{3}$$

$$3a_{02}y^{2} \quad 3xy^{2}$$

$$4a_{01}y \quad 6a_{11}xy \quad 3x^{2}y$$

$$a_{00} \quad a_{10}x \quad a_{20}x^{2} \quad x^{3}$$

The monic hypothesis implies that the rightmost diagonal is  $(x+y)^3$ . Since  $T_*(x+y)^3 \in \mathbf{P}_2^{pos}$  the diagonals interlace, so the adjacent diagonal is a multiple of  $(x+y)^2$ , and so on. Thus  $T_*(x+y)^3 =$ 

$$y^{3}$$
  
 $a_{20}y^{2}$   $3xy^{2}$   
 $a_{10}y$   $2a_{20}xy$   $3x^{2}y$   
 $a_{00}$   $a_{10}x$   $a_{20}x^{2}$   $x^{3}$ 

If 
$$f = \sum b_i x^i$$
 then  $f(D)(x + y)^3 =$ 

$$y^{3} b_{0}$$
  
 $3y^{2} b_{1} \quad xy^{2} b_{0}$   
 $6y b_{2} \quad 6xy b_{1} \quad x^{2}y b_{0}$ 

 $\begin{array}{cccc} 6\,b_3 & 6x\,b_2 & 3x^2\,b_1 & x^3\,b_0\\ \\ \text{Equating coefficients shows that if }g=a_{00}+a_{10}x+a_{20}x^2+a_{30}x^3 \text{ then} \end{array}$ 

$$\mathsf{T} = \frac{1}{6} \big( \exp^{-1} \, \mathfrak{g}^{\mathsf{rev}} \big) (\mathsf{D})$$

In general, the rightmost diagonal of  $T_*(x+y)^n$  is  $(x+y)^n$ , so all the parallel diagonals are powers of x + y. Thus

$$T_*(x+y)^n = \sum_{i=0}^n \alpha_{i0}(x+y)^i$$

If  $g = \sum_{i=0}^{n} a_{i0} x^{i}$  then

$$\left(\frac{1}{n!} \exp^{-1} g^{rev}\right) (\mathsf{D})(x+y)^n = \left(\sum_{i=0}^n \frac{(n-i)!}{n!} a_{n-i,0} \mathsf{D}^i\right) (x+y)^n$$
$$= \sum_{i=0}^n a_{i0} (x+y)^i = \mathsf{T}_* (x+y)^n$$

	I
	I
	I

Affine transformations are the only transformations whose inverses also preserve roots.

**Theorem 6.8.** If a degree preserving linear transformation and its inverse both preserve roots then the linear transformation is an affine transformation. More precisely, if T is a linear transformation such that

- $\mathsf{T}(\mathbf{P}) \subset \mathbf{P}$
- $\mathsf{T}^{-1}(\mathbf{P}) \subset \mathbf{P}$
- $deg(T(x^n)) = n$

then there are constants a, b, c such that Tf(x) = cf(ax + b).

*Proof.* Using the three constants, we may assume that T(1) = 1 and T(x) = x. With these normalizations, it suffices to show that Tf = f. Given a polynomial  $f \in \mathbf{P}$ , we can use Theorem 1.43 to apply  $T^{-1}$  to the interlacing  $Tf \leq (Tf)'$  to find that  $f \leq T^{-1}(Tf)'$ . Since  $T^{-1}(Tf)'$  is a linear transformation defined on all of **P**, there is a constant *w* such that  $T^{-1}(Tf)' = wf'$ , or equivalently wT(f') = (Tf)'.

We now can show that  $T(x^2)=x^2.$  Assume that  $T(x^2)=\alpha x^2+\beta x+\gamma.$  Since

$$T(x^2)' = 2\alpha x + \beta$$
$$wT(2x) = 2wx$$

we find that  $T(x^2) = wx^2 + \gamma$ . Solving for  $T^{-1}$  gives  $T^{-1}(x^2) = (1/w)(x^2 - \gamma)$ . If  $\gamma$  is non-zero, then it is not possible that both  $T(x^2)$  and  $T^{-1}(x^2)$  have real roots, so  $T(x^2) = wx^2$ . Since  $T((x - 1)^2) = wx^2 - 2x + 1$  has all real roots, the discriminant is non-negative which implies that  $w \ge 1$ . Similarly, since  $T^{-1}(x^2)$  has all real roots then  $(1/w) \ge 1$ , and so w = 1.

If we assume by induction that  $T(x^{n-1}) = x^{n-1}$  then applying T(f') = (Tf)' shows that  $T(x^n) = x^n + \alpha$ . This polynomial does not have all real roots unless  $\alpha = 0$ , so  $T(x^n) = x^n$  and thus Tf = f.

*Remark* 6.9. There is an alternate argument for the end of the proof. If T commutes with differentiation D then T = f(D) for some  $f \in \hat{\mathbf{P}}$ . Similarly  $T^{-1} = f^{-1}(D)$ , so f is invertible and has no zeros. The only functions of this form in  $\hat{\mathbf{P}}$  are  $f(x) = \alpha e^{bx}$ , and  $\alpha e^{bD}g(x) = \alpha g(x + b)$ .

**Corollary 6.10.** *If* T *is an invertible linear transformation such that* TP = P *then* Tf = cf(ax + b).

*Remark* 6.11. Although the only bijections on **P** are trivial, there are non-trivial bijections on  $\mathbf{P}^{\text{alt}}$ . If  $\tilde{L}_n^{(\alpha)}(x)$  is the monic Laguerre polynomial, (§7.7.10) then [146] the linear transformation  $x^n \mapsto \tilde{L}_n^{\alpha}(x)$  satisfies  $T = T^{-1}$ . Since T maps  $\mathbf{P}^{\text{alt}}$  to itself (Corollary 7.47) it follows that T is a bijection on  $\mathbf{P}^{\text{alt}}$ .

**Lemma 6.12.** Suppose that T is a degree preserving linear transformations from polynomials to polynomials. If f and Tf have a common root for all f then Tf is a multiple of f.

*Proof.* We can assume that T(1) = 1, and let  $T(x^n) = \sum a_{n,i}x^i$ . Since  $T(x^n)$  and  $x^n$  have zero for a common root the constant terms  $a_{n,0}$  are zero for n > 0. Next, x + t and  $T(x+t) = a_{1,1}x + t$  have -t for a common root, so  $a_{1,1} = 1$  and T(x) = x.

Now assume that  $T(x^k) = x^k$  for k < n.

$$T(x+t)^{n} = T(x^{n}) + \sum_{k=1}^{n-1} x^{k} \binom{n}{k} t^{k} = T(x^{n}) + (x+t)^{n} - x^{n}$$

Since -t is the only root of  $(x + t)^n$  we see that  $T(x^n)(-t) = (-t)^n$  for all t and therefore  $T(x^n) = x^n$ .

If f and Tf have a common interlacing then Tf has a simple form.

**Lemma 6.13.** *If* T *is a degree preserving linear transformation such that* Tf *and* f *have a common interlacing for all*  $f \in \mathbf{P}$  *then there are*  $\alpha$ ,  $\beta$ ,  $\gamma$  *so that* 

$$T(f) = (\alpha x + \beta)f' + \gamma f$$

*Proof.* Since  $x^{n-1}$  is the only polynomial that interlaces  $x^n$ , we see that if  $n \ge 2$  then  $T(x^n)$  has a n-1 fold root at zero. Write

$$T(1) = b_0$$
  
$$T(x^n) = a_n x^n + b_n x^{n-1} \qquad n \ge 1$$

Now we use the fact that  $T(x-t)^n$  must have a root at t if  $t \geqslant 0.$  For n=2 this means that

$$0 = \mathsf{T}(x-t)^2[t] = \mathsf{T}(x^2)[t] - 2t\mathsf{T}(x)[t] + t^2\mathsf{T}[1]$$

which implies

$$\mathsf{T}(\mathsf{x}^2) = 2\mathsf{x}\mathsf{T}(\mathsf{x}) - \mathsf{x}^2\mathsf{T}(1)$$

Thus  $a_2 = 2a_1 - b_0$  and  $b_2 = 2b_1$ . We now follow the same argument, and show by induction that

$$a_n = na_1 - (n-1)b_0 \qquad b_n = nb_{n-1}$$

which implies that

$$T(x^{n}) = x^{n}(na_{1} - (n-1)b_{0}) + nx^{n-1}b_{1}$$
  
=  $a_{1}x(x^{n})' - b_{0}(x(x^{n})' - x^{n}) + b_{1}(x^{n})'$ 

and by linearity

$$T(f) = a_1 x f' - b_0 x f' + b_0 f + b_1 f'$$
  
= ((a\_1 - b\_0)x + b\_1) f' + b\_0 f.

Compositions mapping **P** to itself are very simple.

**Lemma 6.14.** Suppose g(x) is a polynomial.

- (1) If  $f(g(x)) \in P$  for all  $f \in P$  then g is linear.
- (2) If  $f(g(x)) \in P$  for all  $f \in P^{pos}$  then g is either linear, or is quadratic with negative leading coefficient.
- (3) If  $g(x) + c \in P$  for all  $c \in \mathbb{R}$  then g is linear.
- (4) If  $g(x) + c \in P$  for all c > 0 then g is quadratic with negative leading coefficient.

*Proof.* If we take f to be linear, then (1) and (2) follow from (3) and (4). If the degree of g is odd then the horizontal line y = -c will intersect the graph of g(x) in only one point if c is sufficiently large. Thus, in this case the degree of g is 1.

If the degree of g is even then the horizontal line y = -c with c large and positive will intersect the graph of g in at most two points, and hence the degree of g is two. Since these horizontal lines are below the x-axis, the parabola must open downward, and so the leading coefficient is negative.

Here is another characterization of affine transformations.

**Lemma 6.15.** Suppose that  $T: P \longrightarrow P$  is a linear transformation. T distributes over multiplication ( $T(f \cdot g) = Tf \cdot Tg$ ) if and only if T is an affine transformation.

*Proof.* If p = T(x) then by induction  $T(x^n) = p^n$ . Since  $T(x + c) = p + c \in \mathbf{P}$  for all  $c \in \mathbb{R}$ , we see that p is linear, so write p = qx + b. If  $f(x) = \sum a_i x^i$  then the conclusion follows from

$$T(f) = \sum a_i T(x^i) = \sum a_i p^i = f(p) = f(qx + b).$$

The next lemma describes those transformations whose translations also preserve roots.

**Lemma 6.16.** If T is a linear transformation on polynomials such that

- 1.  $(T + a)P \subset P$  for all  $a \in \mathbb{R}$ .
- 2.  $deg(T(x^n)) = n$

then there are constants c, d such that Tf = cf + df'.

*Proof.* For any polynomial  $g \in \mathbf{P}$ ,  $(T + a)g = Tg + ag \in \mathbf{P}$ . By Proposition 1.35 Tg and g interlace. The conclusion now follows from Theorem 6.2.

A similar argument shows

**Lemma 6.17.** If T is a linear transformation on polynomials such that

- 1.  $(T + a)P^{alt} \subset P$  for all  $a \in \mathbb{R}$
- 2.  $deg(T(x^n)) = n$

then Tf(x) = cf + (d + ex)f' for certain  $c, d, e \in \mathbb{R}$ .

It is difficult to tell when two linear transformations T, S have the property that Tf and Sf interlace for all  $f \in \mathbf{P}$ , but it is easy if T, S are multiplier transformations. Such transformations map  $x^n$  to a constant multiple of  $x^n$ .

**Lemma 6.18.** Suppose  $T(x^n) = t_n x^n$  and  $S(x^n) = s_n x^n$  both map P to itself. Assume that  $t_n$  is always positive and  $\begin{vmatrix} t_n & t_{n+1} \\ t_{n+1} & t_{n+2} \end{vmatrix} \le 0$ . If Tf and Sf interlace for all  $f \in P$  then there are  $a, b \in \mathbb{R}$  so that Sf = aTf + bx(Tf)'. *Proof.* We use the same approach as we have been following, and choose a test polynomial f such that Tf has a multiple root - this root must also be a root of Sf, and this will give us a relation between T and S. We let  $f = x^n(x-a)(x-1)$  so that

$$Tf(x) = (a t_n - (1+a) x t_{1+n} + x^2 t_{2+n}) x^n$$

Ignoring the factor of  $x^n$ , the discriminant of f is

$$\Delta = (-1 - a)^2 t_{1+n}^2 - 4 a t_n t_{2+n}$$

so if we choose a to make  $\Delta$  zero

$$a = \frac{-t_{1+n}^2 + 2 t_n t_{2+n} - 2 \sqrt{-(t_n t_{1+n}^2 t_{2+n}) + t_n^2 t_{2+n}^2}}{t_{1+n}^2}$$

then we can solve Tf=0 with this value of a. The hypotheses on  $t_n$  ensure that  $a\in\mathbb{R}.$  The root is

$$r = \frac{t_n t_{2+n} - \sqrt{-(t_n t_{1+n}^2 t_{2+n}) + t_n^2 t_{2+n}^2}}{t_{1+n t_{2+n}}}$$

and if we compute (Sf)(r) we get

$$\begin{split} 0 &= -\left(\frac{s_{2+n}\,t_n\,t_{1+n} + (-2\,s_{1+n}\,t_n + s_n\,t_{1+n})\,\,t_{2+n}}{t_{1+n}^3\,t_{2+n}}\right)\,\times \\ &\left(t_{1+n}^2 - 2\,t_n\,t_{2+n} + 2\,\sqrt{t_n\,t_{2+n}\,\left(-t_{1+n}^2 + t_n\,t_{2+n}\right)}\right) \end{split}$$

If the second factor is zero then simple algebra shows that  $t_n = 0$ , contradicting our hypothesis. Consequently

$$0 = s_{2+n} t_n t_{1+n} + (-2 s_{1+n} t_n + s_n t_{1+n}) t_{2+n}$$

If we define  $u_n = s_n/t_n$  then this relation simplifies to

$$0 = u_{n+2} - 2u_{n+1} + u_n$$

This implies that  $u_n = a + bn$  and hence  $s_n = (a + bn)t_n$ . It follows easily that Sf = aTf + b(Tf)'.

The transformation  $x^n \mapsto \frac{\langle x \rangle_n}{n!}$  was shown in [142] to map polynomials whose imaginary part is 1/2 to polynomials whose imaginary part is 1/2. Such a transformation can not map **P** to itself, as the next lemma shows.

**Lemma 6.19.** Suppose that T maps polynomials with complex coefficients to polynomials with complex coefficients. Suppose that  $L_1$  and  $L_2$  are intersecting lines in the complex plane with the property that T:  $P^{L_1} \longrightarrow P^{L_1}$  and T:  $P^{L_2} \longrightarrow P^{L_2}$ . Then, T is a shifted multiplier transformation. That is, if  $\alpha = L_1 \cap L_2$  then there are constants  $a_n$  such that

$$T(x - \alpha)^n = a_n (x - \alpha)^n$$
 for all integers n

*Proof.* All polynomials in  $\mathbf{P}^{L_1} \cap \mathbf{P}^{L_2}$  are constant multiples of powers of  $(\mathbf{x} - \alpha)$ . Since T maps  $\mathbf{P}^{L_1} \cap \mathbf{P}^{L_2}$  to itself, the conclusion follows.

If the transformation considered in [142] also mapped **P** to itself, then  $T(x-1/2)^n$  would be a multiple of  $(x-1/2)^n$ , for all n. This is false for n = 2.

# 6.2 The domain and range of a linear transformation

If T is a linear transformation that maps  $\mathbf{P}^{\mathbf{I}}$  to  $\mathbf{P}^{\mathbf{J}}$  where  $\mathbf{I}, \mathbf{J}$  are intervals, then there are restrictions on the kinds of intervals that  $\mathbf{I}, \mathbf{J}$  may be. Up to affine transformations there are three types of intervals: all of  $\mathbb{R}$ , half infinite intervals (e.g  $(-\infty, 1)$ ), and finite intervals (e.g. (0, 1)). The only restriction in the next lemma is that T is non-trivial - that is, the image of T is not all constant multiples of a fixed polynomial.

**Lemma 6.20.** If **J** is an interval, and  $T: \mathbf{P}^{\mathbb{R}} \longrightarrow \mathbf{P}^{J}$  is non-trivial then  $J = \mathbb{R}$ .

*Proof.* Suppose that  $T: \mathbf{P} \longrightarrow \mathbf{P}^{J}$ . Choose  $f \leq g$  such that Tf and Tg are not constant multiples of one another. Since f and g interlace, so do Tf and Tg, and hence  $Tf + \alpha Tg \in \mathbf{P}^{J}$  for all  $\alpha \in \mathbb{R}$ .

For any r that is not a root of Tg the polynomial  $h = Tf + \frac{-(Tf)(r)}{(Tg)(r)} Tg$  has r for a root. h is not the zero polynomial since Tf and Tg are not constant multiples of one another. Consequently there are polynomials in **P**<sup>J</sup> with arbitrarily large and small roots, so J must be  $\mathbb{R}$ .

Table 6.2 shows that all cases not eliminated by the lemma are possible. For instance, the reversal map  $f \mapsto f^{rev}$  satisfies  $\mathbf{P}^{(1,\infty)}(n) \longrightarrow \mathbf{P}^{(0,1)}$ .

	all of $\mathbb R$	half infinite	finite
all of $\mathbb R$	$x^{i}\mapsto H_{i}$	_	
half infinite	$\mathbf{x}^{\mathbf{i}} \mapsto (-1)^{\binom{\mathbf{i}}{2}} \mathbf{x}^{\mathbf{i}}$	$x^{i} \mapsto (\underline{x})_{i}$	$f\mapsto f^{re\nu}$
finite	$x^i \mapsto {}_q H_i$	$x^i \mapsto A_i$	$x^i \mapsto T_i$

Table 6.1: Realizing possible domains and ranges

If  $T: \mathbf{P}^{[a,b]}(n) \longrightarrow \mathbf{P}$  then we can easily find upper and lower bounds for the range of T.

**Lemma 6.21.** If  $T: P^{[\alpha,b]}(n) \longrightarrow P(n)$  and preserves interlacing then  $T: P^{[\alpha,b]}(n) \longrightarrow P^{[r,s]}(n)$  where s is the largest root of  $T(x - b)^n$  and r is the smallest root of  $T(x - a)^n$ . In addition, for any  $f \in P^{(\alpha,b)}(n)$  we have

$$T(x-a)^n \triangleleft T(f) \triangleleft T(x-b)^n$$

*Proof.* Choose  $f \in \mathbf{P}^{(a,b)}(n)$ . If we join  $(x - b)^n$  and  $(x - a)^n$  by a sequence of interlacing polynomials containing f (Lemma 1.64) then applying T yields

$$T((x-b)^n) \underline{\ll} \cdots \underline{\ll} Tf \underline{\ll} \cdots \underline{\ll} T(x-a)^n$$

This shows that the largest root of  $T(x-b)^n$  is at least as large as the largest root of T(f), and the smallest root of  $T(x-a)^n$  is at most the smallest root of Tf.

If  $f \leftarrow g$  then the k'th root of f is bounded above by the k'th root of g, for all k up to the degree of g. Here's a simple consequence of this idea.

**Lemma 6.22.** Suppose that  $T: P \longrightarrow P$  preserves degree and the sign of the leading coefficient. If  $f \in P(n)$  and  $a_k$  is the k'th largest root of  $f(1 \le k \le n)$  then

k'th largest root of 
$$T(f) \leq k'$$
th largest root of  $T((x - a_k)^{\kappa})$ 

*Proof.* Suppose  $roots(f) = (a_i)$  and define the polynomials

$$g_{r} = (x - a_{1})(x - a_{2}) \cdots (x - a_{r})$$
  

$$h_{r} = (x - a_{r})(x - a_{r+1}) \cdots (x - a_{k})^{r} \qquad (1 \leq r \leq k)$$

We have interlacings

$$f = g_n \underline{\lessdot} g_{n-1} \underline{\lessdot} \cdots \underline{\lessdot} g_k = h_1 \underline{\lessdot} h_2 \cdots \underline{\lessdot} h_k = (x - a_k)^k$$

Since T preserves the direction of interlacing, and the degree,

$$T(f) \leq \cdots \leq T(g_k) = T(h_1) \leq \cdots \leq T((x - a_k)^k)$$

The conclusion now follows from the observation above.

If we apply this kind of argument to the derivative, then we can use the extra information that  $f \leq f'$ . When we replace  $h_r$  with

$$h_r = (x - a_r)(x - a_{r+1}) \cdots (x - a_k)^k (x - a_{k+1})$$

then a similar argument shows

**Lemma 6.23** (Peyson[136]). If  $f \in P(n)$  and  $1 \le k < n$  then the k'th root of f' lies in the interval

$$\left[a_{k} + \frac{a_{k+1} - a_{k}}{n - k + 1}, a_{k+1} - \frac{a_{k+1} - a_{k}}{k + 1}\right]$$

We can integrate the image of a linear transformation over an interval if the images of the endpoints interlace.

**Lemma 6.24.** If  $T: P^{(a,b)} \longrightarrow P$  and

1) T preserves interlacing  
2) 
$$T(x-b)^n \leq T(x-a)^n$$
  $n = 1, 2, \cdots$   
then  $\int_a^b T(x-t)^n dt \in P$ 

*Proof.* From Lemma 1.64 we can join  $(x - b)^n$  and  $(x - a)^n$  by a sequence of interlacing polynomials that contains f and g. Since T preserves interlacing we know that

$$T(x-b)^{n} \underline{\ll} \cdots \underline{\ll} T(g) \underline{\ll} \cdots \underline{\ll} T(f) \underline{\ll} \cdots \underline{\ll} T(x-a)^{n}$$

Now the endpoints of this interlacing sequence interlace, so it follows that the sequence is a mutually interlacing sequence. In particular, we know that  $T(x - t)^n$  is a family of mutually interlacing polynomials on (a, b), so the conclusion follows from Proposition 3.37.

It is not easy to have a finite image. Some multiplier transformations have finite images. For instance, the identity is a multiplier transformation. Less trivially, the map  $f \mapsto xf'$  maps  $\mathbf{P}^{[0,1]}$  to itself, and is the multiplier transformation  $x^i \mapsto ix^i$ . There's a simple restriction on such transformations.

**Lemma 6.25.** Suppose  $T: x^i \mapsto a_i x^i$  maps  $P^{[0,1]}$  to itself. Then

$$\mathfrak{a}_0 \leqslant \mathfrak{a}_1 \leqslant \mathfrak{a}_2 \leqslant \cdots$$

*Proof.* We may assume that all  $a_i$  are positive. Consequently, the roots of  $T(x^n - x^{n-1})$  lie in [0, 1], and so

$$0 \leq \mathsf{T}(x^{n} - x^{n-1})(1) = (a_{n}x^{n} - a_{n-1}x^{n-1})(1) = a_{n} - a_{n-1}.$$

If the constants decrease sufficiently rapidly, then no finite interval can be preserved.

**Lemma 6.26.** Suppose that the multiplier transformation  $T: x^i \mapsto a_i x^i$  maps  $P^{pos}$  to itself, and  $\limsup |a_n|^{1/n} = 0$ . Then there are no finite intervals I, J such that  $T(P^I) \subset P^J$ .

*Proof.* It suffices to show that the absolute value of the largest root of  $T(x+1)^n$  goes to infinity as  $n \to \infty$ . If  $T(x+1)^n = a_n x^n + \cdots + a_0$ , then the product of the roots is  $a_0/a_n$ . Since there is a root of absolute value at least  $|a_0/a_n|^{1/n}$ , the conclusion follows since  $|a_n|^{1/n}$  goes to 0.

For instance, since  $\limsup (1/n!)^{1/n} = 0$ , it follows that the exponential transformation  $x^n \mapsto x^n/n!$  does not preserve any finite interval.

*Remark* 6.27. By Corollary 1.50, the leading coefficients of a linear transformation that preserves degree are either all the same sign, or they alternate. If T alternates the signs of the leading coefficients, and we define S(f)(x) = T(f)(-x) then S preserves the sign of the leading coefficient. Thus, we usually only consider linear transformations T:  $P \longrightarrow P$  that preserve the sign of the leading coefficient.

Table 6.2 shows that there are many possibilities for the degrees of  $T(1), T(x), T(x^2), T(x^3), \ldots$ . Since  $x^n + \alpha x^{n-1} \in \mathbf{P}$  we know  $T(x^n)$  and  $T(x^{n-1})$  interlace, and so the degrees of  $T(x^n)$  and  $T(x^{n-1})$  differ by at most one, provided neither is zero.

Trans	formation	degree of $T(1)$ , $T(x)$ , $T(x^2)$ ,
$g\mapsto$	fg	n, n + 1, n + 2, n + 3,
$g\mapsto$	f * g	$0, 1, \ldots, n-1, n, 0, 0, 0, \ldots$
$g\mapsto$	$D^{k}f(D)g$	0,,0,1,2,3,
$g\mapsto$	g(D)f	$n, n-1, \ldots, 3, 2, 1, 0, 0, \ldots$
$x^k\mapsto$	$c_kf + d_kf'$	0, n or n − 1

Table 6.2: Degrees of a linear transformation where  $f \in \mathbf{P}(n)$ .

It is important to note that the transformation  $T: g \mapsto g(D)f$  maps  $P \longrightarrow P$ , yet it does not preserve the sign of the leading coefficient, nor do the signs of the leading coefficient alternate. Instead, the sign of the leading coefficient of T(g) depends on g, for it is the sign of g(0). This doesn't contradict Lemma 1.50 since T doesn't preserve degree.

The last transformation maps  $x^k$  to a linear combination of f and f', which might be constant, so the degree is 0, n or n - 1.

#### 6.3 Transformations determined by products

When does a linear transformation of the form

$$T(x^{n}) = \prod_{i=1}^{n} (x + a_{i})$$
(6.3.1)

preserve real roots? We can not answer this question but we can show that such a non-trivial T can not map **P** to itself. Next we will observe that if T in (6.3.1) maps  $\mathbf{P}^{\text{pos}}$  to **P** then there are constraints on the parameters  $a_i$ . There are examples of such root preserving transformations, for in Corollary 7.35 we will see that the choice  $a_i = i - 1$  maps  $\mathbf{P}^{\text{pos}}$  to itself.

**Lemma 6.28.** If T is given in (6.3.1) and T maps P to itself then all  $a_i$  are equal.

*Proof.* Choose  $b \in \mathbb{R}$ , assume T preserves roots, and consider

$$T((x-b)^2) = b^2 - 2b(x+a_1) + (x+a_1)(x+a_2)$$
(6.3.2)

The discriminant is

$$-(a_2-a_1)(4b+a_1-a_2)$$

If  $a_1 \neq a_2$  then we can choose b to make the discriminant negative, so such a linear transformation T does not preserve roots. More generally

$$\mathsf{T}(x^{r}(x-b)^{2}) = \prod_{i=1}^{r} (x+a_{i}) \cdot (b^{2} - 2b(x+a_{r+1}) + (x+a_{r+1})(x+a_{r+2}))$$
(6.3.3)

We can apply the same argument as above to conclude that  $a_{r+1} = a_{r+2}$ , and hence if T preserves roots then all  $a_i$  are equal.

Note that the conclusion of the lemma implies that there is an a such that T(f) = f(x + a). We restate the conclusion:

**Corollary 6.29.** If T:  $P \longrightarrow P$  maps monic polynomials to monic polynomials, and  $T(x^i)$  divides  $T(x^{i+1})$  for all  $i \ge 0$  then  $T(x^i) = (x + a)^i$  for some a.

**Lemma 6.30.** If  $\top$  is given in (6.3.1) and  $\top: P^{\text{pos}} \longrightarrow P$  then

$$a_1 \leqslant a_2 \leqslant a_2 \leqslant \cdots$$

*Moreover, if*  $a_{n+1} = a_n$  *for some* n *them*  $a_m = a_n$  *for all*  $m \ge n$ .

*Proof.* Choose negative b so that  $T((x-b)^2)$  is in **P**. If we choose |b| sufficiently large in (6.3.2) then the sign of the discriminant is the sign of  $a_2 - a_1$  and hence  $a_1 \leq a_2$ . The same argument applied to (6.3.3) shows that  $a_{r+1} \leq a_{r+2}$ .

Next, suppose that  $a_{n+1} = a_n$ . If  $f = (x + a)^3 x^{n-1}$  then

$$T(f) = \prod_{1}^{n+2} (x + a_i) + 3a \prod_{1}^{n+1} (x + a_i) + 3a^2 \prod_{1}^{n} (x + a_i) + a^3 \prod_{1}^{n-1} (x + a_i)$$
  
= 
$$\prod_{1}^{n-1} (x + a_i) \times (x + a_{n+2})(x + a_{n+1})(x + a_n) + 3a(x + a_{n+1})(x + a_n) + 3a^2(x + a_n) + a^3$$

and without loss of generality we may set  $a_n = a_{n+1} = 0$ . The resulting cubic polynomial

 $(x+a_{n+2})x^2+3ax^2+3a^2x+a^3\\$ 

is in **P** for all positive a. If we substitute  $a = a_{n+2}$  then the roots are  $(-.42 \pm .35i)a_{n+2}$  and  $-3.1a_{n+2}$ . Consequently,  $a_{n+2} = 0$ . We can now continue by induction.

For other examples of such transformations: see Lemma 12.26.

# 6.4 Composition

If T is a linear transformation, and f, g are polynomials then we can construct a new polynomial f(T)g. In this section observe that such composition is ubiquitous, and give conditions for f(T)g to define a map  $\mathbf{P}^{\mathbf{I}} \times \mathbf{P} \longrightarrow \mathbf{P}$ .

**Lemma 6.31.** If T is a linear transformation that preserves degree then there is a linear transformation W and a constant c so that T(f) = cf(W)(1) for every polynomial f.

*Proof.* Since T preserves degree the polynomials  $T(x^0), T(x^1), T(x^2), ...$  form a basis for the space of all polynomials. Define the linear transformation W by  $W(T(x^i)) = T(x^{i+1})$ , and set c = T(1). We show by induction that for this choice of W and c we have Tf = cf(W)(1). It suffices to establish this for the basis above. We note that  $T(x^0) = c = cW^0(1)$ . The inductive step is

$$T(x^{n+1}) = W(T(x^n)) = W(cW^n(1)) = cW^{n+1}(1).$$

We can also define W by W(Tf) = T(xf). Since T is invertible we also have that  $W(g) = T(x(T^{-1}g))$  This representation shows that if T is the identity then W is multiplication by x. The inverse of a transformation expressed as a composition has an inverse that is easily expressed as a composition.

**Lemma 6.32.** *If* W *is a linear transformation that increases degree by one,* T(1) = 1*, and* T(f) = f(W)(1) *then* 

$$T^{-1}(f) = f(T^{-1}WT)(1)$$

*Proof.* It is enough to verify the conclusion on a basis, and a good choice for basis is  $\{W^n(1)\}$ . We note

$$T^{-1}W^{n}(1) = (T^{-1}WT)^{n}T^{-1}(1) = (T^{-1}WT)^{n}(1)$$

We now look at some general properties of compositions.

**Theorem 6.33.** Suppose that the linear transformation S has the property that  $(S - \alpha)h \in P$  for all  $h \in P$  and all  $\alpha$  in an interval I. The bilinear map  $f \times g \mapsto f(S)g$  defines a bilinear transformation  $P^{I} \times P \longrightarrow P$ .

*Proof.* We may write  $f = (x - a_1) \cdots (x - a_n)$  where all  $a_i$  are in I. Since

$$T(g) = f(S)g = (S - a_1) \cdots (S - a_n)g$$

we see that the hypothesis on S guarantees that  $Tg \in \mathbf{P}$ .

A similar argument yields

**Corollary 6.34.** Suppose that  $T: \mathbf{P}^{pos} \longrightarrow \mathbf{P}^{pos}$  and for any  $f \in \mathbf{P}^{pos}$  we have that  $f \longleftarrow Tf$ . The map  $f \times g \mapsto f(T)g$  maps  $\mathbf{P}^{pos} \times \mathbf{P}^{pos}$  to  $\mathbf{P}^{pos}$ . If  $g \in \mathbf{P}^{pos}$  is fixed the map  $f \mapsto f(T)g$  preserves interlacing.

# 6.5 Recurrence relations for orthogonal polynomials

All orthogonal polynomials satisfy a *three term recurrence* (6.5.1). We use this recurrence to establish recurrence relations for linear transformations based on orthogonal polynomials. We also get recurrence relations for the inverses of these linear transformations. Finally, we specialize to polynomials of Meixner type.

Assume that  $p_1, p_2, \ldots$  is a sequence of orthogonal polynomials satisfying the three term recurrence

$$p_{n+1} = (a_n x + b_n)p_n + c_n p_{n-1}$$
(6.5.1)

and define the linear transformations

$$T : x^{n} \mapsto p_{n}$$

$$T^{-1} : p_{n} \mapsto x^{n}$$

$$A : x^{n} \mapsto a_{n}x^{n}$$

$$B : x^{n} \mapsto b_{n}x^{n}$$

$$C : x^{n} \mapsto c_{n}x^{n-1}$$

From the recurrence we find that

$$T(x x^{n}) = a_{n} x p_{n} + b_{n} p_{n} + c_{n} p_{n-1}$$
  
= x T(A(x<sup>n</sup>)) + T(B(x<sup>n</sup>)) + T(C(x<sup>n</sup>))

Since this holds for all n, and T is linear, we have for all polynomials f

$$T(xf) = xTA(f) + TB(f) + TC(f)$$
(6.5.2)

Similarly we have a recurrence for the inverse of T

$$\begin{aligned} \mathsf{T}^{-1}(\mathsf{x}\mathfrak{p}_n) &= \mathsf{T}^{-1}((1/a_n)\mathfrak{p}_{n+1} - (\mathfrak{b}_n/a_n)\mathfrak{p}_n - (\mathfrak{c}_n/a_n)\mathfrak{p}_{n-1}) \\ &= (1/a_n)\mathfrak{x}^{n+1} - (\mathfrak{b}_n/a_n)\mathfrak{x}^n - (\mathfrak{c}_n/a_n)\mathfrak{x}^{n-1} \\ &= \mathfrak{x} \mathfrak{A}^{-1}\mathsf{T}^{-1}(\mathfrak{p}_n) - \mathsf{B} \mathfrak{A}^{-1}\mathsf{T}^{-1}(\mathfrak{p}_n) - \mathsf{C} \mathfrak{A}^{-1}\mathsf{T}^{-1}(\mathfrak{p}_n) \end{aligned}$$

and consequently we have for all polynomials f

$$T^{-1}(xf) = xA^{-1}T^{-1}(f) - BA^{-1}T^{-1}(f) - CA^{-1}T^{-1}(f)$$
(6.5.3)  
= (xA^{-1} - BA^{-1} - CA^{-1})T^{-1}f

If T(1) = 1 then we have an explicit composition representation of  $T^{-1}$ 

$$T^{-1}(f) = f(xA^{-1} - BA^{-1} - CA^{-1})(1)$$
(6.5.4)

Notice that this last representation is not well defined. However, we take (6.5.4) to mean that we do not multiply f out, and then substitute, but rather we use the product representation that follows from (6.5.3).

These recurrences are too general to be useful. If our goal is to establish that a certain T preserves interlacing, then we must have that  $T(xf) \leq T(f)$ . This necessitates that there is some sort of interlacing relationship between A, B, C. We have seen that if A is the identity then it's probable that B, C are functions of derivatives. There is a well known class of polynomials for which this is true.

The Meixner class of orthogonal polynomials [68, page 165] consists of those orthogonal polynomials  $p_n$  that satisfy a recurrence

$$p_{n+1} = (x - a - \alpha n)p_n - (bn + \beta n(n-1))p_{n-1}$$
(6.5.5)

Upon specializing (6.5.3) and (6.5.4), where A is the identity,  $B = -(a + \alpha xD)$ ,  $C = -(bD + \beta xD^2)$ , we find the recurrences

$$T(xf) = xTf - T((a + \alpha xD + bD + \beta xD^{2})f)$$
(6.5.6)

$$T^{-1}(f) = f(x + a + \alpha x D + b D + \beta x D^{2})(1)$$
(6.5.7)

We have seen several examples of orthogonal polynomial families. Although *families* of orthogonal polynomials have many special properties, there is nothing special about any particular orthogonal polynomial, as the next theorem shows.

**Theorem 6.35.** Every  $f \in P$  is an orthogonal polynomial. In other words, there is a measure that determines an orthogonal family of polynomials, and f is a member of that family.

*Proof.* Assume  $f \in \mathbf{P}(n)$ . We will show that there is a sequence of polynomials  $p_0, p_1, \ldots, p_n = f$  where the degree of  $p_i$  is i, and there are non-negative constants  $a_i$  such that  $p_{i+1} = (a_ix + b_i)p_i - p_{i-1}$  for  $1 \le i < n$ . It follows from Favard's Theorem [168] that the sequence  $p_0, \ldots, p_n$  is a set of orthogonal polynomials determined by a measure.

We construct the  $p_i$  inductively. Assume that f has positive leading coefficient. Let  $p_n = f$ , and  $p_{n-1} = f'$ . Since  $p_n \leq p_{n-1}$  we can find a  $p_{n-2}$  such that  $p_n = (a_n x + b_n)p_{n-1} - p_{n-2}$ . Since  $p_{n-1} \leq p_{n-2}$  we continue inductively.  $\Box$ 

#### 6.6 Linear transformations satisfying a recursion

If a linear transformation satisfies a recursion then we can sometimes conclude that the transformation maps  $P^{\alpha lt}$  or  $P^{p \circ s}$  to itself.

**Theorem 6.36.** Suppose that linear transformations A, B, C, D, E, F, H map polynomials with positive leading coefficients to polynomials with positive leading coefficients. Suppose that for all  $f \in \mathbf{P}^{pos}$  we have

$$Af \leq f \leq Bf$$
  $Cf \leq f \leftarrow Df$   $Ef \leq f \leftarrow Hf \leq f$  (6.6.1)

then the linear transformation defined by

$$T(xf) = (HT + xTB + T(C - D) + (E - F)T) f$$

maps  $P^{\text{pos}}$  to itself and preserves interlacing. If the interlacing assumptions (6.6.1) hold for all  $f \in P^{\text{alt}}$  then

$$S(xf) = (HS + xSA + S(C - D) + (E - F)S)f$$

maps  $P^{\text{alt}}$  to itself and preserves interlacing.

*Proof.* We prove the theorem by induction on the degree of f, so assume that T maps  $\mathbf{P}^{\text{pos}}(n)$  to itself and preserves interlacing. By Corollary 1.46 it suffices to show that  $T(xf) \leq Tf$  for all  $f \in \mathbf{P}^{\text{pos}}(n)$ . If  $f \in \mathbf{P}^{\text{pos}}(n)$  then  $f \leq Bf$  and so  $Tf \leq TBf$ . Now all roots of TBf are negative, so  $xTBf \leq Tf$ . Next, we apply our assumptions (6.6.1) to the polynomial  $Tf \in \mathbf{P}^{\text{pos}}(n)$  and find

$$\begin{array}{ll} \mathsf{HTf} \leq \mathsf{Tf} \\ \mathsf{TCf} \leq \mathsf{Tf} \\ \mathsf{ETf} \ll \mathsf{Tf} \\ \mathsf{Tf} \ll \mathsf{FTf} \\ \end{array}$$

Finally, use Lemma 1.31 to conclude

$$T(xf) = (HT + xTB + T(C - D) + (E - F)T)f \leq Tf$$

and hence  $T(xf) \leq Tf$  which implies that T maps  $\mathbf{P}^{\text{pos}}(n+1)$  to itself and preserves interlacing. The case for  $f \in \mathbf{P}^{\text{alt}}$  is similar.

#### 6.7 Möbius transformations

A Möbius transformation determines a linear transformation on polynomials of fixed degree, and also an action of the space of all linear transformations. Recall that a Möbius transformation (or linear fractional transformation) is a transformation of the form

$$M: z \mapsto \frac{az+b}{cz+d}.$$
 (6.7.1)

To construct a map between polynomials we restrict ourselves to polynomials of a fixed degree n, and homogenize:

$$\widetilde{M}: x^{i} \mapsto (ax+b)^{i}(cx+d)^{n-i}.$$
(6.7.2)

We can express  $\widetilde{M}$  in terms of M:

$$\widetilde{\mathsf{M}}(\mathsf{f}) = (\mathsf{c}\mathsf{x} + \mathsf{d})^{\mathsf{n}}\mathsf{f}(\mathsf{M}\mathsf{x}).$$

We have already seen an example of a Möbius transformation in (4.1.4). If we take M(z) = 1/z then  $\widetilde{M}(f) = z^n f(1/z)$  which is the reverse of f. If I is an interval, then we define M(I) to be the image of I. With this notation, we have the elementary lemma

**Lemma 6.37.** If M is a Möbius transformation (6.7.2) then  $\widetilde{M}$  is a linear transformation that maps  $P^{I}(n)$  to  $P^{M^{-1}(I)}(n)$  bijectively.

*Proof.* If  $f \in \mathbf{P}^{I}$  then  $\widetilde{M}(f) = (cx+d)^{n} f(Mx)$ . A root r of  $\widetilde{M}(f)$  satisfies f(Mr) = 0 so  $r = M^{-1}s$  where s is a root of f, and thus  $s \in I$ . The inverse of  $\widetilde{M}$  is  $\widetilde{M^{-1}}$  and so  $\widetilde{M}$  is a bijection.

**Lemma 6.38.** If *I*, *J* are intervals, and T is a linear transformation T:  $P^I \longrightarrow P^J$  and  $\alpha \in \mathbb{R}$  then the linear transformation  $S(x^i) = \alpha^i T(x^i)$  maps  $P^{I/\alpha} \longrightarrow P^J$ .

Proof. The result follows from the commuting diagram



The next lemma shows that only the type of interval matters.

**Lemma 6.39.** Assume I and J are intervals, and that  $\mathbb{A}_1$  and  $\mathbb{A}_2$  are affine transformations. There is a 1-1 correspondence between linear transformations mapping  $\mathbf{P}^I$ to  $\mathbf{P}^J$  and linear transformations mapping  $\mathbf{P}^{\mathbb{A}_1I}$  to  $\mathbf{P}^{\mathbb{A}_2J}$ 

*Proof.* The correspondence is given by the diagram



As another application, the next lemma shows two linear transformations that are related by a Möbius transformation.



Figure 6.1: A Möbius transformation acting on a linear transformation

**Lemma 6.40.** Choose a positive integer n, and define linear transformations  $T(x^i) = \binom{x+n-i}{n}$ ,  $S(x^i) = \binom{x}{i}$ , and the Möbius transformation  $M(z) = \frac{z}{z+1}$ . Then  $T = S\widetilde{M}$  on P(n).

*Proof.* It suffices to verify that  $T = S\widetilde{M}$  for a basis of P(n), where  $\widetilde{M}(x^i) = x^i(x+1)^{n-i}$ , so consider

$$\begin{split} S\widetilde{M}(x^{i}) &= S\left(x^{i}(1+x)^{n-i}\right) \\ &= S\left(\sum_{j=0}^{n-i} \binom{n-i}{j} x^{i+j}\right) \\ &= \sum_{j=0}^{n-i} \binom{n-i}{j} \binom{x}{i+j} \\ &= \sum_{j=0}^{n-i} \binom{n-i}{n-(i+j)} \binom{x}{i+j} \end{split}$$

and the last expression equals  $T(x^i)$  using the Vandermonde identity

$$\binom{a+b}{c} = \sum_{r+s=c} \binom{a}{r} \binom{b}{s}$$

We will later show that  $S: \mathbf{P}^{alt} \longrightarrow \mathbf{P}$ . M maps (0,1) bijectively to  $(0,\infty)$ , so we have a commuting diagram (Figure 6.7) of maps that preserve roots.

The second action determined by a Möbius transformation is in the space of linear transformations. Let T be a linear transformation that preserves degree. If M is given in (6.7.1) then we define a new linear transformation by

$$T_{M}(x^{r}) = (cx + d)^{r} (T(x^{r}))(Mx)$$
(6.7.3)

If  $T(x^r) = p_r$  then we can write this as

$$\mathsf{T}_{\mathsf{M}}(\mathsf{x}^{\mathsf{r}}) = (\mathsf{c}\mathsf{x} + \mathsf{d})^{\mathsf{r}} \, \mathsf{p}_{\mathsf{r}}(\frac{\mathsf{a}\mathsf{x} + \mathsf{b}}{\mathsf{c}\mathsf{x} + \mathsf{d}})$$

Note that  $T_M \neq \widetilde{M}T$ , since on  $\mathbf{P}(n)$ 

$$\widetilde{\mathsf{MT}}(\mathbf{x}^{\mathsf{r}}) = (\mathbf{c}\mathbf{x} + \mathbf{d})^{\mathsf{n}} \, \mathsf{p}_{\mathsf{r}}(\frac{\mathbf{a}\mathbf{x} + \mathbf{b}}{\mathbf{c}\mathbf{x} + \mathbf{d}})$$

If  $p_r(0) \neq 0$  then both  $T(x^r)$  and  $T_M(x^r)$  have degree r. We first note that this action respects composition.

**Lemma 6.41.** If M, N are Möbius transformations then  $(T_M)_N = T_{NM}$ .

Proof. The proof follows from the commuting diagram



Consider some examples.

1. If T is the identity then  $T_M(x^n) = (ax + b)^n$ . Equivalently,  $T_M(f) = f(ax + b)$ .

- 2. If T is any degree preserving linear transformation, Mz = az + b, and we let  $T(x^n) = p_n$  then  $T_M(x^n) = p_n(ax + b)$ . Consequently  $T_M(f) = (Tf)(ax + b)$ . Moreover, if **I**, **J** are intervals and T:  $\mathbf{P}^{\mathbf{I}} \longrightarrow \mathbf{P}^{\mathbf{J}}$  then  $T_M : \mathbf{P}^{M^{-1}\mathbf{I}} \longrightarrow \mathbf{P}^{\mathbf{J}}$ .
- 3. The most interesting (and difficult) case is when the denominator is not constant. Because of the composition property (Lemma 6.41), and the examples above, we only consider the case M(z) = 1/z. Since  $x^n f(1/x)$  is the reverse of f when f has degree n we see that

$$\mathsf{T}_{1/z}(\mathsf{x}^{\mathfrak{n}}) = \mathsf{T}(\mathsf{x}^{\mathfrak{n}})^{\mathrm{REV}}.$$

For example, suppose that

$$\mathsf{T}(\mathsf{x}^{\mathsf{n}}) = (\mathsf{x} - 1)(\mathsf{x} - 2) \cdots (\mathsf{x} - \mathsf{n}).$$

Applying M(z) = 1/z yields

$$T_{1/z}(x^n) = (1-x)(1-2x)\cdots(1-nx)$$

**Example 6.42.** We can combine Möbius transformations with other root preserving linear transformations. For instance, we will see (Corollary 7.35) that  $\langle \underline{x} \rangle_n \mapsto x^n$  maps  $\mathbf{P}^{alt}$  to itself and (Lemma 7.2)  $x^k \mapsto (\underline{\alpha + n})_k x^k$  maps  $\mathbf{P}^{alt}$  to itself. If M(z) = z/(1-z) the composition

$$\mathbf{P}^{\operatorname{alt}}(\mathfrak{n}) \xrightarrow{\langle \underline{\mathbf{x}} \rangle_{k} \mapsto \mathbf{x}^{k}} \mathbf{P}^{\operatorname{alt}}(\mathfrak{n}) \xrightarrow{\mathbf{x}^{k} \mapsto (\underline{\alpha} + \underline{\mathfrak{n}})_{k} \mathbf{x}^{k}} \mathbf{P}^{\operatorname{alt}}(\mathfrak{n}) \xrightarrow{\widetilde{\mathsf{M}}} \mathbf{P}^{(0,1)}$$

is the linear transformation  $\langle \underline{x} \rangle_k \mapsto (\underline{\alpha + n})_k x^k (1 - x)^{n-k}$  and maps  $\mathbf{P}^{\mathfrak{alt}}(\mathfrak{n})$  to  $\mathbf{P}^{(0,1)}$ . This is Example 1 in [90].

# 6.8 Transformations and commutativity properties

We first characterize transformations that commute with  $x \mapsto \alpha x$ , and those that commute with  $x \mapsto x + 1$ . We also characterize some transformations defined by commuting diagrams. Finally, we list some diagrams that are satisfied by various linear transformations.

**Lemma 6.43.** Suppose that the linear transformation T preserves degree. Then T commutes with  $x \mapsto \alpha x$  where  $\alpha \neq 1$  if and only if there is a  $g(x) = \sum \alpha_i x^i$  such that Tf = f \* g. Equivalently, T commutes with  $x \mapsto \alpha x$  if and only if T is a multiplier transformation.

*Proof.* Simply consider the action of T on  $x^n$ . Assume that  $T(x^n) = \sum_{k=0}^n b_i x^i$ . Commutativity implies that

$$\alpha^n T(x^n) = \alpha^n \sum_{k=0}^n b_i x^i = T(x^n)(\alpha x) = \sum_{k=0}^n b_i \alpha^i x^i$$

This implies that all  $b_i$  are zero except for  $b_n$ , so T is a multiplier transformation.

**Lemma 6.44.** Suppose that the linear transformation T preserves degree. Then T commutes with  $\mathbb{A} : x \mapsto x + 1$  if and only if there is a  $g(x) = \sum a_i x^i$  such that

$$T(x^{n}) = \sum_{i=0}^{n} a_{n-i} {n \choose i} x^{i}.$$
 Equivalently,  
$$T(f) = (q * Af)^{rev}.$$

*Proof.* That  $T(x^n)$  has that form can be proved by induction. Conversely, note that the conclusion is linear in g, so we may take  $g(x) = x^s$ . It is easy to see that the diagram commutes.

$$\begin{array}{c|c} x^{n} & \xrightarrow{T} & \sum_{i=0}^{n} {n \choose i} a_{n-i} x^{i} \\ x \mapsto x+1 & & & & \\ \sum_{j=0}^{n} {n \choose j} x^{j} & \xrightarrow{T} & \sum_{i,j=0}^{n} {n \choose i} a_{n-i} {i \choose k} x^{k} \end{array}$$

*Remark* 6.45. If we ask that T commutes with the seemingly more general transformation  $x \mapsto x + y$ , we find that T is the same as is described in Lemma 6.43. For example, the Bernoulli polynomials satisfy this commutativity property. However, this is not useful for us as most Bernoulli polynomials do not have all real roots.

If we choose  $g(x) = e^{-x}$  then

$$T(x^{n}) = \sum {\binom{n}{k}} \frac{-1)^{k}}{k!} x^{k} = L_{n}^{rev}(x)$$

so we have a commuting square in two variables

$$x \mapsto x + y \not\downarrow \xrightarrow{x^{n} \mapsto L_{n}^{rev}(x)} \downarrow \xrightarrow{x \mapsto x + y} \downarrow (6.8.1)$$

**Example 6.46.** If  $T(x^k) = \frac{\langle x \rangle_k}{k!}$  and  $\mathbb{A} f(x) = f(1-x)$  then  $\mathbb{A}T = T\mathbb{A}$ .

**Example 6.47.** If  $T(x^k) = \frac{(\underline{x})_k}{k!} = {\binom{x}{k}}$  and  $\mathbb{A} f(x) = f(-x-1)$  then  $\mathbb{A}T = T\mathbb{A}$ . **Example 6.48.** T:  $x^n \mapsto H_n$  satisfies

$$T(f)(x + y) = T_*f(x + 2y)$$
(6.8.2)

which leads to the diagram



**Example 6.49.** T:  $x^k \mapsto H_k(x)x^{n-k}$  satisfies

$$T(x-2+\alpha)^n = H_n(\frac{\alpha x}{2}) = (-1)^n T(x-2-\alpha)^n$$
 (6.8.3)

If we define S(f) = T(f(x-2)) then



**Example 6.50.** T:  $x^k \mapsto \langle \underline{x} \rangle_k (\underline{x - \alpha})_{n-k}$  satisfies the identity on polynomials of degree n



**Example 6.51.** T:  $x^i \mapsto (\underline{x+n-i})_n$  satisfies

$$Tf = (Tf^{rev})(-1-x)$$
 (6.8.5)

since  $(\underline{-x-1+i})_n = (-1)^n (\underline{x+n-i})_n$ . Equivalently, reverse  $\sqrt[T]{T}$ 

**Example 6.52.** The Laguerre transformation  $T: x^n \mapsto L_n(x)$  satisfies several identities. This square is the Laguerre identity (7.10.4).



T also satisfies



We can combine the Laguerre and Hermite polynomials. Define the linear transformation T acting on P(n), and two induced transformations:

$$\begin{split} \mathsf{T}(z^k) &= \mathsf{H}_k(z) \mathsf{L}_{n-k}(z) \\ \mathsf{T}_x(x^r \mathfrak{y}^s) &= \mathsf{H}_r(x) \mathsf{L}_{n-r}(x) \mathfrak{y}^s \\ \mathsf{T}_y(x^r \mathfrak{y}^s) &= x^r \, \mathsf{H}_s(\mathfrak{y}) \mathsf{L}_{n-s}(\mathfrak{y}) \end{split}$$

We have the following commutative diagram

$$(x + y)^n$$

$$T_x (x + y)^n \leftarrow T_y$$

$$T_y (x + y)^n \leftarrow T_y (x + y)^n$$

which if written out is

$$\sum_{k=0}^{n} \binom{n}{k} H_{k}(x) L_{n-k}(x) y^{n-k} = \sum_{k=0}^{n} \binom{n}{k} H_{n-k}(y/2) L_{k}(y/2) (2x)^{k}$$

**Example 6.53.** The falling factorials  $(\underline{x})_k$  and  $L_n/n!$  are polynomials of binomial type, and satisfy

(6.8.6)

$$\begin{aligned} (\underline{\mathbf{x}} + \underline{\mathbf{y}})_n &= \sum_{k=0}^n \binom{n}{k} (\underline{\mathbf{x}})_k (\underline{\mathbf{y}})_{n-k} \\ \frac{\mathbf{L}_n (\mathbf{x} + \mathbf{y})}{n!} &= \sum_{k=0}^n \binom{n}{k} \frac{\mathbf{L}_k (\mathbf{x})}{k!} \frac{\mathbf{L}_{n-k} (\mathbf{y})}{(n-k)!} \end{aligned}$$

A polynomial family  $\{p_n\}$  of binomial type satisfies

$$p_{n}(x+y) = \sum_{k=0}^{n} \binom{n}{k} p_{k}(x)p_{n-k}(y)$$

For a general family of polynomials of binomial type, the transformation  $T: x^n \mapsto p_n(x)$  satisfies



where the transformations  $T_x$ ,  $T_y$  are the induced transformations. Considering the inverse transformations yields



#### 6.9 Singular points of transformations

If T is a transformation, and  $f \in \mathbf{P}$  is a polynomial for which the degree of Tf is different from the degree of Tg where g is arbitrarily close to f, then we say that f is a singular value for T. We are interested in transformations for which  $(x + a)^n$  is a singular value for infinitely many n. It is often the case that the behavior of such a transformation changes at a, and that there is some sort of symmetry around x = a. These symmetries are captured in the commuting diagrams of the previous section. If T is a linear transformation that preserves degree, then there can be no singularities, but there can be symmetries. We know of the following examples:

1. T: 
$$x^k \mapsto \langle \underline{x} \rangle_k (\underline{x - \alpha})_{n-k}$$
  $T(x-1)^n = \langle \underline{\alpha} \rangle_n$ 

2. T: 
$$x^{i} \mapsto (\underline{x+n-i})_{n}$$
 T(x+1)<sup>n</sup> = (-1)<sup>n</sup>

3. $x^k \mapsto H_k(x)H_{n-k}(x)$	$T(x-1)^n = 2^{n/2}H_n(0)$
4. $x^k \mapsto H_k(x)H_{n-k}(x)$	$T(x-\iota)^n=2^n(1+\iota)^n$
5. $x^k \mapsto L_k(x)L_{n-k}(x)$	$T(x-1)^n = 0$ if n is odd
6. T: $x^k \mapsto H_k(x)x^{n-k}$	$T\left(x-2\right)^n=H_n(0)$
7. T: $x^k \mapsto (x)_k(x)_{n-k}$	$T(x-1)^n = 0$ if n is odd

# 6.10 Appell sequences

Some of the properties of special functions and commuting diagrams are easy for the general class of Appell sequences. A sequence of polynomials  $\{p_n\}$  is an *Appell sequence* [148] if the degree of  $p_n$  is n, and

$$p'_{n}(x) = np_{n-1}.$$
 (6.10.1)

Such sequences are in 1 - 1 correspondence with formal power series g(x) [146] and satisfy

$$g(D)x^n = p_n(x)$$
 for  $n = 0, 1, ...$  (6.10.2)

**Proposition 6.54.** Suppose  $\{p_n\}$  is an Appell sequence determined by g(x). Then,  $g(x) \in \widehat{P}$  iff the linear transformation  $x^n \mapsto p_n$  maps P to itself.

*Proof.* The linear transformation is simply  $f \mapsto g(D)f$ .

If a sequence of polynomials satisfies  $p_n = \alpha_n n p_{n-1}$ , then the polynomials  $q_n = p_n/(\alpha_0 \alpha_1 \cdots \alpha_n)$  form an Appell sequence since

$$\mathfrak{q}'_{\mathfrak{n}} = \mathfrak{p}'_{\mathfrak{n}}/(\alpha_{0}\cdots\alpha_{\mathfrak{n}}) = \alpha_{\mathfrak{n}}\mathfrak{n}\mathfrak{p}_{\mathfrak{n}-1}/(\alpha_{0}\cdots\alpha_{\mathfrak{n}}) = \mathfrak{n}\mathfrak{p}_{\mathfrak{n}-1}/(\alpha_{0}\cdots\alpha_{\mathfrak{n}-1}) = \mathfrak{n}\mathfrak{q}_{\mathfrak{n}-1}/(\alpha_{0}\cdots\alpha_{\mathfrak{n}-1}) = \mathfrak{n}\mathfrak{q}_{\mathfrak{n}-1}/(\alpha_{0}\cdots\alpha_\mathfrak{n}-1) = \mathfrak{n}\mathfrak{q}_{\mathfrak{n}-1}/(\alpha_{0}\cdots\alpha_\mathfrak{n}-1}) = \mathfrak{n}\mathfrak$$

The Hermite polynomials are nearly Appell sequences. Since they satisfy  $H'_n = 2nH_{n-1}$ ,  $H_n(x/2)$  is an Appell sequence. For a more interesting example, consider the transformation T:  $f \mapsto f(x+\iota) + f(x-\iota)$ . Now  $Tf = 2\cos(D)f$ , and so the sequences

$$p_n = (x + \iota)^n + (x - \iota)^r$$

are Appell sequences with  $g(x) = \cos x$ .

The Euler polynomials form an Appell sequence with  $g(x) = \frac{1}{2}(e^x + 1)$ . This function is not in  $\hat{\mathbf{P}}$ , so we can not conclude that the linear transformation determined by the Euler polynomials maps  $\mathbf{P}$  to itself. Indeed, it doesn't. Similarly, the Bernoulli polynomials have  $g(x) = (e^x - 1)/x$ , and the corresponding linear transformation doesn't map  $\mathbf{P}$  to itself.

Any Appell sequence satisfies the identity

$$p_{n}(x+y) = \sum_{k=0}^{n} \binom{n}{k} p_{k}(x) y^{n-k}$$
(6.10.3)

which we have already seen for the Hermite polynomials in (6.8.2). Equation (6.10.3) implies that we have the commutative diagram



If  $T: x^k \mapsto p_k(x)x^{n-k}$  then the lemma below shows that  $T_*(x - y)^r$  has degree n for all y other than y = 1, so 1 is a singular point.

**Lemma 6.55.** If  $\{p_n\}$  is Appell and  $T: x^k \mapsto p_k(x)x^{n-k}$ , then

$$\mathsf{T}_*(\mathsf{x}-1+\mathsf{y})^{\mathsf{r}}=\mathsf{x}^{\mathsf{n}-\mathsf{r}}\mathsf{p}_{\mathsf{r}}(\mathsf{x}\mathsf{y})$$

*Proof.* Since the conclusion is a polynomial identity, it suffices to prove it for y an integer. It's trivially true for y = 1, and the case y = 0 follows from

$$T(x-1)^{r} = \sum_{k=0}^{r} \binom{r}{k} (-1)^{r-k} p_{k}(x) x^{r-k} = x^{n-r} p_{n}(x-x) = x^{n-r} p_{n}(0).$$

Assume that it is true for y, and consider y + 1

$$T_*(x-1+y+1)^r = \sum_{k=0}^r \binom{r}{k} T_*(x-1+y)^k = \sum_{k=0}^r \binom{r}{k} x^{n-k} p_k(xy) = x^{n-r} \sum_{k=0}^r \binom{r}{k} x^{r-k} p_k(xy) = x^{n-r} p_r(xy+x) = x^{n-r} p_r(x(y+1)).$$

#### 6.11 The diamond product

If we are given an invertible linear transformation T on polynomials then we can form a new bilinear transformation.



We call  $f \diamond g = T^{-1} (Tf \cdot Tg)$  a diamond product. The product can be defined for any number of terms in a simple way. For instance, observe that

$$f \Diamond (g \Diamond h) = f \Diamond [T^{-1}(Tg \cdot Th)] = T^{-1}(Tf \cdot T[T^{-1}(Tg \cdot Th)]) = T^{-1}(Tf \cdot Tg \cdot Th)$$

Consequently, we see that the diamond product is well defined for any number of factors, is associative and commutative, and equals

$$f_1 \Diamond f_2 \Diamond \cdots \Diamond f_n = T^{-1} (Tf_1 \cdot Tf_2 \cdots Tf_n)$$

We can now prove a general result:

**Proposition 6.56.** Suppose that T is a linear transformation such that

1. T:  $P^I \longrightarrow P^J$ 2. T<sup>-1</sup>:  $P^J(1) \longrightarrow P^K(1)$ 

*Then every*  $f \in P^{I}$  *can be written the form* 

$$c(x-a_1) \Diamond (x-a_2) \Diamond \cdots \Diamond (x-a_n)$$

where c is a constant and each  $x - a_i$  is in  $P^K$ .

*Proof.* From (1) we see that  $T(f) \in \mathbf{P}^{J}$ , and consequently can be factored  $T(f) = c_0(x - b_1) \cdots (x - b_n)$ . Write  $T^{-1}(x - b_i) = c_i(x - a_i)$  where each  $x - a_i$  is in  $\mathbf{P}^{K}$  by (2). If  $c = c_0 \cdots c_n$  then

$$\begin{split} c(x-a_1)\otimes(x-a_2)\otimes\cdots\otimes(x-a_n) &= c_0T^{-1}(c_1T(x-a_1)\cdots c_nT(x-a_n))\\ &= T^{-1}(c_0(x-b_1)\cdots(x-b_n))\\ &= T^{-1}(Tf) = f \end{split}$$

Here are some examples of diamond products.

**binomial** Let  $T^{-1}(x^n) = \binom{x}{n}$ . Then

$$\begin{pmatrix} x \\ r \end{pmatrix} \Diamond \begin{pmatrix} x \\ s \end{pmatrix} = \mathsf{T}^{-1} \left( \mathsf{T} \begin{pmatrix} x \\ r \end{pmatrix} \right) \mathsf{T} \left( \begin{pmatrix} x \\ s \end{pmatrix} \right) \right)$$
$$= \mathsf{T}^{-1} (x^{r+s})$$
$$= \begin{pmatrix} x \\ r+s \end{pmatrix}$$

This was considered by Wagner [178] who introduced the name diamond product. See Proposition 11.100.

falling factorial Let  $T^{-1}(\underline{x})_n = x^n$ . Then

$$(\underline{\mathbf{x}})_{\mathbf{n}} \diamondsuit (\underline{\mathbf{x}})_{\mathbf{m}} = (\underline{\mathbf{x}})_{\mathbf{n}+\mathbf{m}}$$

This product is discussed in §8.8.8.

**exponential** If  $T(x^n) = x^n/n!$  then

$$x^{n} \Diamond x^{m} = \binom{n+m}{n} x^{n+m}$$

See Question 43.

We can generalize the diamond product construction by replacing multiplication by some other bilinear map. Given  $\mathfrak{m} : \mathbf{P} \times \mathbf{P} \longrightarrow \mathbf{P}$  then we form a new bilinear transformation

$$\mathbf{f} \diamondsuit \mathbf{g} = \mathsf{T}^{-1}\left(\mathsf{m}(\mathsf{T}(\mathsf{f}),\mathsf{T}(\mathsf{g}))\right)$$

See Proposition 11.101 for an application.

Although affine transformations distribute over multiplication, they do not generally distribute over diamond multiplication. We need a connection between the affine transformation and the defining transformation.

**Lemma 6.57.** Suppose  $f \Diamond g = T^{-1}(Tf \cdot Tg)$  and  $\mathbb{A}$  is an affine transformation. If  $\mathbb{A}$  commutes with T then  $\mathbb{A}$  distributes over  $\Diamond$ . Conversely, if  $\mathbb{A}$  distributes over  $\Diamond$  and  $\mathbb{A}$  commutes with T for polynomials of degree 1, then  $\mathbb{A}$  commutes with T.

*Proof.* The first part is just definition:

$$\begin{split} \mathbb{A}(\mathsf{f} \Diamond \mathsf{g}) &= \mathbb{A}\mathsf{T}^{-1}(\mathsf{T}\mathsf{f} \cdot \mathsf{T}\mathsf{g}) = \mathsf{T}^{-1}\mathbb{A}(\mathsf{T}\mathsf{f} \cdot \mathsf{T}\mathsf{g}) = \mathsf{T}^{-1}(\mathbb{A}\mathsf{T}\mathsf{f} \cdot \mathbb{A}\mathsf{T}\mathsf{g}) \\ &= \mathsf{T}^{-1}(\mathsf{T}\mathbb{A}\mathsf{f} \cdot \mathsf{T}\mathbb{A}\mathsf{g}) = \mathbb{A}\mathsf{f} \Diamond \mathbb{A}\mathsf{g}. \end{split}$$

Conversely, using induction

$$\begin{split} \mathbb{A} T(fg) &= \mathbb{A} (Tf \Diamond Tg) = \mathbb{A} Tf \Diamond \mathbb{A} Tg = T\mathbb{A} f \Diamond T\mathbb{A} g \quad [induction] \\ &= T(\mathbb{A} f \cdot \mathbb{A} g) = T\mathbb{A} (fg) \end{split}$$
# 6.12 Other root preserving transformations

There are linear transformations that increase the degree of a polynomial by more than 1 and still preserve roots. Such transformations *W* will not preserve interlacing if the degrees of *W*f and *W*g differ by more than 1.

A trivial example is  $Tf = p \cdot f$  where  $p \in P$ . This transformation does preserve interlacing. A more interesting example that does not preserve interlacing is

$$\mathsf{T}: \mathfrak{x}^{\mathfrak{n}} \mapsto (\mathfrak{x}^2 - 1)^{\mathfrak{n}}$$

so that deg(Tf) = 2deg(f). This transformation can also be expressed as  $Tf = f(x^2 - 1)$ . If  $f \in \mathbf{P}^{(-1,\infty)}$  and  $roots(f) = (a_1, \ldots, a_n)$  then  $roots(Tf) = (\ldots, \pm (a_i + 1)^{1/2}, \ldots)$ . Thus, T preserves roots for polynomials in  $\mathbf{P}^{(-1,\infty)}$ . We can generalize this idea.

**Lemma 6.58.** Suppose that  $g \in P(r)$  has the property that no relative maximum or minimum of g has value lying in (-1, 1). If  $f \in P^{(-1,1)}(n)$  then  $f(g(x)) \in P^{(-1,1)}(nr)$ .

*Proof.* Since g has no extrema with value in (-1, 1) it follows that every line y = s where  $-1 \le s \le 1$  meets the graph of g in r distinct points.

Without loss of generality we may assume that f has all distinct roots. If f(t) = 0 then  $-1 \le t \le 1$  and we get r roots of f(g(x)) from the r solutions to the equation g(x) = t. This accounts for all nr of the roots of f(g(x)).

**Corollary 6.59.** If  $f \in \mathbf{P}^{(-1,1)}(n)$  and  $T_r$  is the Chebyshev polynomial, then  $f(T_r(x))$  is in  $\mathbf{P}^{(-1,1)}(nr)$ .

*Proof.* The graph of the Chebyshev polynomial  $T_r$  (see Figure 6.2) on the interval (-1, 1) oscillates between its r - 1 relative extrema of 1 and -1. Now apply the lemma.

For example, since  $T_2 = 2x^2 - 1$ , the transformation  $f(x) \mapsto f(2x^2 - 1)$  maps  $\mathbf{P}^{(-1,1)}$  to itself.



Figure 6.2: The graph of the Chebyshev polynomial  $T_5(x)$ 

### CHAPTER



# Linear transformations that preserve roots

We apply the results of the previous chapters to show that particular linear transformations preserve roots.

# 7.1 Multiplier transformations

The simplest linear transformations are *multiplier transformations*, which are of the form  $x^i \mapsto a_i x^i$ . We will see in Theorem 15.22 that if the series  $\sum a_i \frac{x^i}{i!}$  is in  $\widehat{\mathbf{P}^{\text{pos}}}$  then the multiplier transformation maps  $\mathbf{P}$  to itself. In § 5.5.3 we showed that various series were in  $\widehat{\mathbf{P}^{alt}}$  or  $\widehat{\mathbf{P}^{\text{pos}}}$ . Consequently we have

**Theorem 7.1.** The following linear transformations map P to P and  $P^{\text{pos}}$  to  $P^{\text{pos}}$ .

1. 
$$x^{i} \mapsto \frac{x^{i}}{i!}$$
  
2.  $x^{i} \mapsto \frac{x^{i}}{\Gamma(i + \alpha)}$  for  $\alpha \ge 0$   
3.  $x^{i} \mapsto \frac{x^{i}}{\langle \alpha \rangle_{i}}$  for positive  $\alpha$ .  
4.  $x^{i} \mapsto \frac{x^{i}}{\langle \alpha i \rangle!}$  for positive integer  $\alpha$ .  
5.  $x^{i} \mapsto q^{\binom{i}{2}} x^{i}$  for  $|q| \le 1/4$ .  
6.  $x^{i} \mapsto \frac{x^{i}}{[i]!}$  for  $|q| > 4$ .

7.  $x^i \mapsto (\underline{d})_i x^i$  for d a positive integer.

8.  $x^{i} \mapsto {d+i-1 \choose i} x^{i}$  for d a positive integer.

9. 
$$x^{i} \mapsto \frac{i!q^{\binom{i}{2}}}{(q;q)_{i}}x^{i}$$
 for  $|q| < 1$ 

10. 
$$x^{i} \mapsto \frac{q^{\binom{i}{2}}}{(q;q)_{i}} x^{i}$$
 for  $|q| < 1$ .

Proof. The first two follow from (5.3.3), the third one from (5.3.4), and the fourth from [138, 62,#162]. The generating function of  $(\underline{d})_i$  is in  $\mathbf{P}^{\text{pos}}$ :

$$\sum_{i=0}^{\infty} (\underline{d})_i \frac{x^i}{i!} = \sum_{i=0}^{\infty} {\binom{d}{i}} x^i = (1+x)^d.$$

The generating function for  $x^n/[n]!$  is in  $\widehat{P}$  - see 5.4. The generating function for  $\binom{d+i-1}{i}$  is

$$\sum_{i=0}^{\infty} \binom{d+i-1}{i} \frac{x^i}{i!} = e^x L_{d-1}(-x)$$

where  $L_{d-1}$  is the Laguerre polynomial (see Section 7.10). Since  $e^x \in \widehat{\mathbf{P}^{pos}}$  and  $L_{d-1}(-x) \in \mathbf{P}^{pos}$  their product is in  $\widehat{\mathbf{P}^{pos}}$ . 

The last two use (5.1.1) and part (1).

We can extend (7) of the last result to non-integer values of n, but surprisingly there are restrictions.

**Lemma 7.2.** If n is a positive integer and  $\alpha > n - 2 \ge 0$  then

- 1. the map  $x^i \mapsto (\underline{\alpha})_i x^i$  maps  $P^{pos}(n)$  to itself, and  $P^{alt}(n)$  to itself.
- 2. the map  $x^i \mapsto {\binom{\alpha}{i}} x^i$  maps  $P^{pos}(n)$  to itself, and  $P^{alt}(n)$  to itself.

*Proof.* We find a recurrence and then proceed by induction. First, let  $T(x^n) =$  $(\underline{\alpha})_n x^n$ . Then

$$\begin{split} \mathsf{T}(\mathbf{x} \cdot \mathbf{x}^{i}) &= \left(\underline{\alpha}\right)_{i} (\alpha - i) \mathbf{x}^{i+1} \\ &= \alpha \mathbf{x} \left(\underline{\alpha}\right)_{i} \mathbf{x}^{i} - \mathbf{x} i \left(\underline{\alpha}\right)_{i} \mathbf{x}^{i} \\ &= \alpha \mathbf{x} \mathsf{T}(\mathbf{x}^{i}) - \mathbf{x} \mathsf{T}(\mathbf{x}(\mathbf{x}^{i})') \\ \mathsf{T}(\mathbf{x}f) &= \alpha \mathbf{x} \mathsf{T}(f) - \mathbf{x} \mathsf{T}(\mathbf{x}f'). \end{split}$$

We now show by induction on m that T maps  $\mathbf{P}^{pos}(m)$  to  $\mathbf{P}^{pos}(m)$  for m < n and to  $\mathbf{P}(m)$  for m = n. For m = 1 we take f = x + r where r > 0 and then  $Tf = r + \alpha x$  is also in  $\mathbf{P}^{pos}(1)$ .

Assume that T maps  $\mathbf{P}^{pos}(m)$  to itself and m < n. If we choose g = (x+r)f where f has positive leading coefficient and r > 0 then

$$\mathsf{T}(\mathsf{g}) = \alpha \mathsf{x} \mathsf{T}(\mathsf{f}) - \mathsf{x} \mathsf{T}(\mathsf{x}\mathsf{f}') + \mathsf{r} \mathsf{T}(\mathsf{f}).$$

Since  $f \in \mathbf{P}^{pos}$  we know that  $xf' \underline{\ll} f$  and hence by induction  $T(xf') \underline{\ll} T(f)$ . It follows that  $T(g) = (\alpha x + r)T(f) - xT(xf') \underline{\ll} T(f)$ . If m = n - 1 this shows that T maps  $\mathbf{P}^{pos}(n)$  to  $\mathbf{P}(n)$ .

Now assume that m < n - 1. In order to show that T(g) is in  $\mathbf{P}^{\text{pos}}$  we need to show that (Tg)(0) and the leading coefficient of T(g) have the same sign. First, (Tg)(0) = r(Tf)(0) is positive since r > 0 and T(f) has positive leading coefficient and is in  $\mathbf{P}^{\text{pos}}$ . Next, the leading coefficient of T(g) is  $s(\underline{\alpha})_{m+1}$  where s > 0 is the leading coefficient of f. Consequently, the leading coefficient of T(g) is positive since  $\alpha > n - 2 \ge m$  and the conclusion now follows.

The second part follows from the first by applying the exponential transformation (Theorem 7.1(1)). The results for  $\mathbf{P}^{alt}$  follow from applying the above to the composition  $f(x) \mapsto T(f(-x))(-x)$ .

If the generating function lies in  $\widehat{\mathbf{P}}$  but not in  $\widehat{\mathbf{P}^{alt}}$  or  $\widehat{\mathbf{P}^{pos}}$  then the corresponding linear transformation maps  $\mathbf{P}^{\pm}$  to  $\mathbf{P}$  by Theorem 15.22.

**Lemma 7.3.** The following linear transformations map  $P^{\pm}$  to P.

1. 
$$x^{i} \mapsto \begin{cases} 0 & i \text{ odd} \\ (-1)^{n} \frac{(2n)!}{n!} x^{2n} & i = 2n \end{cases}$$
  
2.  $x^{i} \mapsto \begin{cases} 0 & i \text{ even} \\ (-1)^{n} x^{i} & i = 2n+1 \end{cases}$   
3.  $x^{i} \mapsto \begin{cases} 0 & i \text{ odd} \\ (-1)^{n} x^{i} & i = 2n \end{cases}$ 

*Proof.* The correspond the the functions  $e^{-x^2}$ , sin(x) and cos(x).

The q-exponential function was defined in Example 5.3. We define the

*~*···

q-exponential transformation (0 < q < 1)

$$\operatorname{Exp}_{q}(x^{i}): x^{i} \mapsto \frac{q^{\binom{i}{2}}}{(q;q)_{i}} x^{i}$$
(7.1.1)

Theorem 7.1 shows that  $Exp_q$  maps **P** to itself.

#### 7.2 Homogeneous transformations

Some linear transformations are only defined on polynomials of bounded degree. For instance, if

$$\mathsf{T}(\mathsf{x}^{\mathsf{i}}) = \mathsf{x}^{\mathsf{i}}(1-\mathsf{x})^{\mathsf{n}-\mathsf{i}}$$

then T is a linear transformation on all polynomials of degree at most n.

**Lemma 7.4.** If a and c are not both zero then the linear transformation T given by

$$\mathsf{T}(\mathsf{x}^{\mathsf{i}}) = (\mathfrak{a}\mathsf{x} + \mathfrak{b})^{\mathsf{i}}(\mathfrak{c}\mathsf{x} + \mathfrak{d})^{\mathsf{n}-\mathsf{i}}$$

*maps* P(n) *to itself.* 

*Proof.* Let F be the homogeneous polynomial corresponding to f. If  $f(x) = (x - a_1) \cdots (x - a_n)$  then  $F(x, y) = (x - a_1 y) \dots (x - a_n y)$ . The image of f under the transformation T is simply F(ax + b, cx + d). In order to see this, we only need to check it for a basis, namely  $x^i$ , in which case the result is clear.

We therefore have the factorization

$$T(f) = F(ax + b, cx + d) = (ax + b - a_1(cx + d)) \cdots (ax + b - a_n(cx + d))$$

which shows that Tf has all real roots.

This lemma also follows from the properties of Möbius transformations in § 6.6.7 - the argument here shows the factorization of the transformation.

**Corollary 7.5.** The linear transformation  $T: x^i \mapsto x^i(1-x)^{n-i}$  maps  $P^{alt}(n)$  to  $P^{(0,1)}(n)$ .

*Proof.* This is a special case of Lemma 6.37 where M = z/(1-z) since M maps (0, 1) to  $(0, \infty)$ . Alternatively, if  $\alpha$  is a root of  $f \in \mathbf{P}^{alt}(n)$  then the corresponding root of Tf is  $\alpha/(\alpha + 1)$ .

The reversal operator is the special case of this construction where we set a = 0, b = 1, c = 1, d = 0. See (4.1.4).

**Lemma 7.6.** If the polynomial f(x) has all real roots then  $f^{\text{REV}}$  has all real roots. If  $f \in P^{\text{pos}}$  (resp.  $P^{\text{alt}}$ ) then  $f^{\text{REV}} \in P^{\text{pos}}$  (resp.  $P^{\text{alt}}$ ).

If  $f \ll g$  and 0 is a not root of fg then  $g^{REV} \ll f^{REV}$ . If  $f \ll g$  and  $f \in \mathbf{P}^{\pm}$  then  $f^{REV} \ll g^{REV}$ .

*Proof.* Without loss of generality we assume that 0 is not a root. The reverse of f is  $x^n f(\frac{1}{x})$ , so its roots are the reciprocals of the non-zero roots of f. The rest of the properties are immediate from this observation.

If f has 0 as a root, then the degree of  $f^{REV}$  is less than n. If 0 is not a root of f, then  $(f^{REV})^{REV} = f$ . However, if 0 is a d-fold root, then  $x^d(f^{REV})^{REV} = f$ 

If we have a linear transformation T on P(n) then we can conjugate T by the reversal map to create a new linear transformation  $T^{rev}$  on P(n).



Here are some examples:

- 1. If D is the derivative, then  $D^{\text{REV}}$  is the polar derivative.
- 2. If T is the transformation  $x^i \mapsto c_i x^i$  then  $T^{\text{REV}}(x_i) = c_{n-i} x^i$ .
- 3. If  $T(x^i) = p_i$  where  $p_i$  is symmetric ( $p_i = (p_i)^{\text{REV}}$ ) then  $T^{\text{REV}}(x^i) = p_{n-i}$ . This is the case for the Hermite polynomials (see § 7.7.8).

# 7.3 The Hadamard product

The generalized Hadamard product of two polynomials is the coordinate wise product with an extra coefficient per coordinate:

$$\begin{aligned} (a_0 + a_1 x + a_2 x^2 + \dots + a_n x^n) \circledast (b_0 + b_1 x + b_2 x^2 + \dots + b_n x^n) \\ &= a_0 b_0 c_0 + a_1 b_1 c_1 x + a_2 b_2 c_2 x^2 + \dots + a_n b_n c_n x^n \end{aligned}$$

Equivalently, we can define it in terms of monomials:

$$x^{i} \circledast x^{j} \mapsto egin{cases} c_{i} x^{i} & i = j \ 0 & i \neq j \end{bmatrix}$$
 and  $c_{i} > 0$ 

With the help of the method of generating functions (p. 500), we will prove that

The generalized Hadamard product  $\circledast$  determines a map  $\mathbf{P}^{pos}$   $\times$ 

$$\mathbf{P} \mapsto \mathbf{P} \text{ if and only if } \sum_{1}^{\infty} c_i \frac{x^i}{i!i!} \in \widehat{\mathbf{P}^{\text{pos}}}.$$
(7.3.1)

If all  $c_i$  are 1 we call it the Hadamard product and write f \* g in place of  $f \circledast g$ . If  $c_i = i!$  then we write f \*' g. It follows from the above result that \* and  $*' map P^{p \circ s} \times P$  to P. There is a an alternative proof for the Hadamard product (see § 9.9.13) that uses an identification of it as a coefficient of a polynomial in two variables.

The generalized Hadamard preserves interlacing.

**Lemma 7.7.** If  $\circledast$  satisfies (7.3.1),  $f \leq g \in P^{pos}$ ,  $h \in P$  then  $f \circledast h \leq g \circledast h$ .

*Proof.* If  $g = \alpha f + r$  where  $f \leq r$  then  $f \circledast h \leq \alpha f \circledast h$  and  $f \circledast h \leq r \circledast h$  since all coefficients are positive. Adding these interlacings gives the result.

Our goal is to prove a multiplicative property of  $\circledast.$  We begin with a special case.

**Lemma 7.8.** If  $f \in P^{pos}$ ,  $h \in P$ ,  $\alpha, \beta > 0$  and  $\circledast$  satisfies (7.3.1) then

$$(x + \alpha) f \circledast (x + \beta) h \leq f \circledast h$$

Proof. We have the interlacings

xf⊛xh <u>∢</u> f⊛h	since $xf \circledast xh = x(f \circledast h)$
$xf \circledast \beta h \leq f \circledast h$	Lemma 7.7
$\alpha f \circledast xh \leq f \circledast h$	Lemma 7.7
αf ⊛ βh≪f ⊛ h	

and the conclusion follows upon adding the interlacings.

It's surprising that the generalized Hadamard product allows us to multiply interlacings. In the case of \* there is a natural interpretation using polynomials with complex coefficients (p. 606).

**Lemma 7.9.** If  $f \leq g$  in P,  $h \leq k$  in  $P^{pos}$  and  $\circledast$  satisfies (7.3.1) then

$$f \circledast h \lessdot g \circledast k$$

Proof. Using Lemma 1.20 we write

$$g = \sum_{i} \frac{a_i f}{x - r_i}$$
 and  $k = \sum_{j} \frac{b_j h}{x - s_j}$ 

where the r's and s's are negative, and the a's and b's are positive. From the above lemma we know that for all relevant i, j

$$f \circledast h \leq \frac{a_i f}{x - r_i} \circledast \frac{b_j h}{x - s_j}$$

Adding these interlacings gives the conclusion.

**Lemma 7.10.** If m is a positive integer then

- 1. the map  $x^i \mapsto \binom{m}{i} x^i$  maps **P** to itself.
- 2. the map  $x^i \mapsto (\underline{m})_i x^i$  maps **P** to itself.

*Proof.* The first statement is the map  $f \mapsto (1 + x)^m * f$ . The second statement follows from the first statement and Theorem 9.87. Another proof uses generating functions -see Theorem 7.1.

It is obvious that the Hadamard product extends to a bilinear map  $\widehat{P} \times P^{\pm} \longrightarrow \widehat{P}$ . Since

$$e^{x^2} * (x+1)^2 = x^2 + 1$$

it follows that  $e^{x^2} \notin \widehat{\mathbf{P}}$ . Of course, we know that  $e^{-x^2} \in \widehat{\mathbf{P}}$ .

The effect of scaling on the Hadamard product is easy to describe.

**Lemma 7.11.** Suppose that  $\circledast$  is a generalized Hadamard product, and choose constants  $\alpha$ ,  $\beta$ . If S, T, U are regions satisfying

$$\circledast: \mathbf{P}^{\mathsf{S}} \times \mathbf{P}^{\mathsf{T}} \longrightarrow \mathbf{P}^{\mathsf{U}}$$

then

$$\circledast: \mathbf{P}^{\alpha S} \times \mathbf{P}^{\beta T} \longrightarrow \mathbf{P}^{\alpha \beta U}$$

Proof. It follows from the definition of generalized Hadamard product that

$$f(\alpha x) \circledast g(\beta x) = (f \circledast g)(\alpha \beta x)$$

Consideration of the diagram below yields the conclusion.

$$\begin{array}{c} \mathbf{P}^{\alpha S} \times \mathbf{P}^{\beta T} \xrightarrow{(\circledast)} \mathbf{P}^{\alpha \beta U} \\ x \times y \mapsto \alpha x \times \beta y \\ \mathbf{P}^{S} \times \mathbf{P}^{T} \xrightarrow{(\circledast)} \mathbf{P}^{U} \end{array}$$

*Remark* 7.12. Differentiating a Hadamard product has no nice properties in general. However, we do have the following properties that are easily verified, where m is non-negative.

$$\frac{d}{dx} [(1+x)^m \circledast x f] = m(1+x)^{m-1} \circledast f$$
$$\frac{d}{dx} [e^x \circledast x f] = e^x \circledast f$$
$$\left(\frac{d}{dx}\right)^m [e^x \circledast x^m f] = e^x \circledast f$$

Thus, if  $f \in \mathbf{P}$  then we can find polynomials  $p_0, p_1, \ldots$  so that

- 1.  $p_0 = e^x \circledast f$ .
- 2. all  $p_i$  are in **P**.
- 3.  $(d/dx)p_i = p_{i-1}$  for i = 1, 2, ...

Such polynomials have been called *very hyperbolic*. Here is a different construction of very hyperbolic polynomials due to Chebeterov.

**Lemma 7.13.** If  $f = \sum_{0}^{\infty} a_i x^i \in \widehat{P}$  and  $f_n = \sum_{0}^{n} a_i \frac{x^{n-i}}{(n-i)!}$  then all  $f_n$  have all real roots and  $f'_n = f_{n-1}$ .

*Proof.* Clearly  $f'_n = f_{n-1}$ ; we need to see that  $f_n \in \mathbf{P}$ . Now

$$(1+x)^{n} *' f = \sum_{0}^{n} {n \choose i} i! a_{i} x^{i} = n! \sum_{0}^{n} a_{i} \frac{x^{i}}{(n-i)!}$$

is in **P**, and taking the reverse shows  $f_n \in \mathbf{P}$ .

For more properties of the Hadamard product, see § 9.9.13.

# 7.4 Differential operators

The most general differential operators on polynomials that we consider have the form

$$g \mapsto \sum_{i=0}^{n} f_i(x) g^{(i)}(x) \tag{7.4.1}$$

where the  $f_i(x)$  are polynomials. We first determine some restrictions on the coefficient functions for operators that map **P** to **P**. We then look at some general composition properties of polynomials, and apply these to some particular operators. In Proposition 15.66 we will characterize, modulo some positivity and degree conditions, those differential operators (7.4.1) that map **P** to **P**.

It is useful to view the differential operator as being determined by the two variable polynomial  $f(x, y) = \sum f_i(x)y^i$ . These polynomials satisfy a substitution condition:

**Lemma 7.14.** If  $f(x, y) = \sum f_i(x)y^i$  determines a differential operator (7.4.1) that maps P to itself then  $f(x, \alpha) \in P$  for all  $\alpha \in \mathbb{R}$ .

*Proof.* If the corresponding differential operator T maps **P** to **P** then T maps  $\widehat{\mathbf{P}}$  to  $\widehat{\mathbf{P}}$ . If we apply T to  $e^{\alpha x} \in \widehat{\mathbf{P}}$  we find

$$\left(\sum_{i=0}^{n} f_{i}(x) \mathsf{D}^{i}\right) e^{\alpha x} = \left(\sum_{i=0}^{n} f_{i}(x) \alpha^{i}\right) e^{\alpha x}$$

Since the right hand side is in  $\hat{\mathbf{P}}$  we can multiply by  $e^{-\alpha x}$  and remain in  $\hat{\mathbf{P}}$ . The result follows.

We next consider two particular differential operators. Choose a polynomial  $h(x) = \sum a_i x^i$  and define  $h(D)g = \sum a_i g^{(i)}$ . In this case the function  $f_i(x)$  of (7.4.1) is just the constant  $a_i$ . We also define

$$h(xD)g = \sum_{i=0}^{n} a_i \left(x \frac{d}{dx}\right)^i g(x)$$

If we expand  $(xD)^ig$  we see we get a differential operator of the form (7.4.1).

**Proposition 7.15.** *The map*  $f \times g \mapsto f(D)g$  *defines a bilinear map*  $P \times P \longrightarrow P$ .

*Proof.* It follows from Corollary 7.23 that  $(D - \alpha)f \in P$  for all f. Apply Corollary 6.34.

**Example 7.16.** If we choose  $f = x^2 - 1$  then for any  $g \in \mathbf{P}$  it follows that  $g - g'' \in \mathbf{P}$ . Moreover, if  $g \leq h$  then  $g - g'' \leq h - h''$ .

If n is a positive integer then the linear transformation  $f \mapsto \sum_{i=0}^{n} {n \choose r} f^{(r)}$ maps **P** to itself - simply choose  $f = (1 + x)^{n}$ .

Suppose f = x+1 and g = x+2. Although  $f \ll g$ , substituting the derivative reverses the direction, e.g.  $x + 1 = f(D)x \gg g(D)x = 2x + 1$ . This is not true in general, but depends on the sign of the constant terms.

**Proposition 7.17.** *If*  $f \leq g$  *and*  $h \in P$  *and if* 

$$\begin{cases} f(0)g(0) > 0 \\ f(0)g(0) < 0 \end{cases} \quad \text{then} \begin{cases} f(D)h \ge g(D)h \\ f(D)h \le g(D)h \end{cases}$$

*Proof.* Linearity implies that f(D)h and g(D)h interlace, but we don't know which direction. If we write  $f(D)h = \beta g(D)h + k$  then we need to determine the sign of the leading coefficient of k, and the signs of the leading coefficients of f(D)h and g(D)h. Compute

$$\begin{split} h(x) &= x^{n} + a_{n-1}x^{n-1} + \cdots \\ f(x) &= x^{m} + \cdots + b_{1}x + b_{0} \\ g(x) &= x^{m} + \cdots + c_{1}x + c_{0} \\ f(D)h &= b_{0}x^{n} + (b_{0}a_{n-1} + nb_{1})x^{n-1} + \cdots \\ g(D)h &= c_{0}x^{n} + (c_{0}a_{n-1} + nc_{1})x^{n-1} + \cdots \\ f(D)h &= \frac{b_{0}}{c_{0}}g(D)h + n\frac{b_{1}c_{0} - b_{0}c_{1}}{c_{0}}x^{n-1} + \cdots \end{split}$$

Without loss of generality we assume that  $f \ll g$ . Consequently, we know the numerator  $b_1c_0 - b_0c_1$  is positive.

If  $b_0c_0 > 0$  then assume that  $b_0 > 0$  and  $c_0 > 0$ . The leading coefficients of f(D)h and g(D)h are positive, and the leading coefficient of k is also positive. This implies  $g(D)h \leq f(D)h$ .

If  $b_0c_0 < 0$  then assume that  $b_0 > 0$  and  $c_0 < 0$ . The leading coefficient of f(D)h is positive, the leading coefficient of g(D)h is negative, and the leading coefficient of k is negative. This implies  $g(D)h \ge f(D)h$ .

The remaining cases are similar.

**Example 7.18.** We know that  $f \leq T(g)$  for all  $g \in \mathbf{P}$  implies that T is the derivative. However, we can find many different linear transformations S, T so that  $S(g) \leq T(g)$  for all  $g \in \mathbf{P}$ . Choose  $f_1 < f_2$  where  $f_2(0) = 0$  and define

$$S(g) = f_1(D)g$$
  $T(g) = f_2(D)g$ .

The proposition above implies that S(g) and T(g) interlace, and since  $f_2(0) = 0$  the degree of T(g) is one less than the degree of S(g). Thus,  $S(g) \leq T(g)$ .

**Corollary 7.19.** The transformation<sup>1</sup>  $f \times g \mapsto f(xD)g$  defines a bilinear map

- 1.  $P^{\text{pos}} \times P^{\pm} \longrightarrow P^{\pm}$ .
- 2.  $\mathbf{P}^{\mathrm{I}}(\mathfrak{n}) \times \mathbf{P} \longrightarrow \mathbf{P}$  if  $\mathrm{I} = (-\infty, 0) \cup (\mathfrak{n}, \infty)$ .
- 3.  $P^{\text{pos}} \times P^{(0,1)} \longrightarrow P^{(0,1)}$ .
- 4.  $\mathbf{P}^{\text{pos}} \times \mathbf{P}^{(-1,0)} \longrightarrow \mathbf{P}^{(-1,0)}$ .

*Proof.* To prove the first part for  $f \in \mathbf{P}^{p \circ s}$  it suffices to show that  $f(xD)g \in \mathbf{P}$  when f = x + a and a is positive. Since f(xD)g = ag + xg' this follows from Corollary 2.16. The second part also follows from the corollary. The third part holds since  $ag + xg' \in \mathbf{P}^{(0,1)}$  when a is positive and  $g \in \mathbf{P}^{(0,1)}$ . The fourth part is similar.

We can extend this to analytic functions:

**Corollary 7.20.** If  $f \in \hat{P}$  has no roots in [0, n] then T(g) = f(xD)g defines a linear transformation  $T: P(n) \longrightarrow P(n)$ .

*Proof.* The reason this is not immediate is that f is not necessarily presented as a limit of polynomials with no roots in [0, n]. However, if a sequence  $f_i$  of polynomials converges uniformly to f, then the zeros of the  $f_i$  converge uniformly to the zeros of f. If f has no roots in [0, n], then the roots of  $f_i$  must go to to roots that are not in [0, n] as i goes to infinity. If  $g \in \mathbf{P}(n)$  then for i sufficiently large no roots of  $f_i$  are in [0, n], and we can apply Corollary 7.27 as above.

It's easy to express  $f(x\mathsf{D})g$  in terms of coefficients. If  $g(x)=b_0+\dots+b_nx^n$  then

$$f(xD)g = f(0)b_0 + f(1)b_1x + \dots + f(n)b_nx^n.$$
(7.4.2)

To see this, note that the conclusion is linear in both f and g so it suffices to establish it for the case  $f = x^r$  and  $g = x^n$ . In this case we see that

$$f(\mathbf{x}\mathsf{D})g = (\mathbf{x}\mathsf{D})^r \mathbf{x}^n = n^r \mathbf{x}^n = f(n)g.$$

We can interpret (7.4.2) as a Hadamard product, but the factor corresponding to f is not in  $\hat{\mathbf{P}}$ . Given a polynomial f, we form the infinite series

$$F(x) = f(0) + f(1)x + f(2)x^2 + \cdots$$

We see from (7.4.2) that f(xD)g = F \* g. Moreover, we will see in §7.7.13 that this series is a rational function of the form  $h(x)/(1-x)^r$ , and so F is not in  $\widehat{\mathbf{P}}$ . It

<sup>&</sup>lt;sup>1</sup> It is not true that the transformation maps  $\mathbf{P}^{alt} \times \mathbf{P}^{\pm}$  to  $\mathbf{P}$ . For example, if f = (x - .7)(x - .1) and g = (x + .7)(x + .1) then  $f(xD)g = .0049 + .216x + 2.47x^2$  has no real roots.

is interesting that there are series that map **P** to **P** under the Hadamard product, but these series are not uniform limits of polynomials. This phenomenon is explained in Theorem 9.87.

Using Corollary 7.19 and (7.4.2) we get these corollaries.

**Corollary 7.21.** If  $g = b_0 + b_1 x + \cdots + b_n x^n$  is in **P** and r is a positive integer then the following polynomials are in **P**.

$$b_0 + 1^r b_1 x + \dots + n^r b_n x^n$$
$$(\underline{r})_r b_0 + (\underline{r+1})_r b_1 x + \dots + (\underline{n+r})_r b_n x^n$$

*Proof.* Choose  $f = x^r$  in the first case, and  $f = (x + 1) \cdots (x + r)$  in the second.

We end this section with a discussion of the Bernstein polynomials. Given a function, the n-th Bernstein polynomial is defined by

$$B_{n}(f) = \sum_{k=0}^{n} f\left(\frac{k}{n}\right) \binom{n}{k} x^{k} (1-x)^{n-k}$$

and in particular the polynomial corresponding to x<sup>r</sup> is

$$\mathsf{B}_{\mathsf{n}}(\mathsf{x}^{\mathsf{r}}) = \sum_{k=0}^{\mathsf{n}} \frac{k^{\mathsf{r}}}{\mathsf{n}^{\mathsf{r}}} \binom{\mathsf{n}}{k} \mathsf{x}^{\mathsf{k}} (1-\mathsf{x})^{\mathsf{n}-\mathsf{k}}$$

Note that

$$\sum_{k=0}^{n} \frac{k^{r}}{n^{r}} \binom{n}{k} x^{k} = \left(\frac{x\mathsf{D}}{n}\right)^{r} (1+x)^{n}$$

We therefore have the diagram

$$f(x) \xrightarrow[f \mapsto B_n(f)]{} f(x) \xrightarrow[f \mapsto B_n(f)]{} f(x) \xrightarrow[f \mapsto B_n(f)]{} f(x) \xrightarrow[f \mapsto B_n(f)]{} f(x)$$

Now from Corollary 7.19 we know that T:  $f \mapsto f(\frac{xD}{n})(1+x)^n$  determines a map  $\mathbf{P}^{pos} \longrightarrow \mathbf{P}^{(-1,0)}$ . In addition, since  $f \in \mathbf{P}^{(1,\infty)}$  implies that the roots of f(x/n) are greater than n, we find that T also maps  $\mathbf{P}^{(1,\infty)}$  to  $\mathbf{P}^{(-\infty,-1)}$ . Moreover, it follows from the Corollary that we actually have  $\mathbf{P}^{\mathbb{R}\setminus(0,1)} \longrightarrow \mathbf{P}$ . Combined with the Möbius transformation  $z \mapsto z/(1-z)$ , we get the commuting diagram of spaces



In summary,

**Lemma 7.22.** The transformation  $f \mapsto B_n(f)$  satisfies

- 1.  $\boldsymbol{P}^{\mathbb{R}\setminus(0,1)}\longrightarrow \boldsymbol{P}$
- 2.  $P^{(-\infty,0)} \longrightarrow P^{(-\infty,0)}$
- 3.  $P^{(1,-\infty)} \longrightarrow P^{(1,\infty)}$

# 7.5 Transformations based on differentiation

We now look at specific transformations that are based on differentiation. The proofs are all easy consequences of the results in Chapter 1 and Chapter 6.

**Corollary 7.23.** If  $Tf = (\alpha + D)f$  then  $T: P \longrightarrow P$ . If  $\alpha > 0$  then  $T: P^{pos} \longrightarrow P^{pos}$ . If  $\alpha < 0$  then  $T: P^{\alpha lt} \longrightarrow P^{\alpha lt}$ .

**Corollary 7.24.** If  $\alpha, \gamma > 0$ ,  $Tf = (\alpha x - \beta - \gamma x D)f$ , then  $T: P \longrightarrow P$ . If  $f \in P^{alt}$  then  $h \leq f$ . If we know in addition that  $\beta > 0$  then  $T: P^{alt} \longrightarrow P^{alt}$ .

**Corollary 7.25.** If a and e are positive then the transformation  $f \times g \mapsto f(ax - eD)g$  defines a bilinear map  $P \times P \longrightarrow P$ .

Proof. Apply Corollary 7.24.

**Corollary 7.26.** The map  $f \times g \mapsto f(x+xD)g$  defines a bilinear map  $P^{pos} \times P^{alt} \longrightarrow P^{alt}$ .

Proof. Apply Corollary 7.24.

See Corollary 2.16 for the details about the next transformation.

**Corollary 7.27.** *If*  $a \notin (0, n)$  *and*  $\mathsf{Tf} = (-a + xD)\mathsf{f}$  *then*  $\mathsf{T}: P^{\pm}(n) \longrightarrow P^{\pm}(n)$  *and*  $\mathsf{T}: P(n) \longrightarrow P(n)$ .

**Corollary 7.28.** If  $\mathsf{Tf} = (\beta + (1 - x^2)D)\mathsf{f}$  where  $\beta > 0$  and  $\mathsf{f} \in P^{(-1,1)}$  then  $\mathsf{Tf} \leq \mathsf{f}$  and  $\mathsf{T} \colon P^{(-1,1)} \longrightarrow P^{(-1,\infty)}$ .

*Proof.* Note that  $1 - x^2$  is positive on the roots of f. Let  $h = (\beta + (1 - x^2)D)f$ . Since the leading coefficient of h is negative it follows from Lemma 1.20 that  $h \leq f$ . At x = 1 we see that  $h(1) = \beta f(1)$ , so h has at exactly one root greater than 1. There are no roots less that -1, so  $h \in \mathbf{P}^{(-1,\infty)}$ .

**Corollary 7.29.** If Tf = axf + x(1-x)f' then T maps  $P^{(-\infty,0]}$  to itself.

*Proof.* Apply Lemma 1.20 after factoring out x.

A similar argument shows that

**Corollary 7.30.** *If*  $Tf = ((2x + \alpha) - (1 - x^2)D)f$  *where*  $|\alpha| < 1$  *then* T *maps*  $P^{(-1,1)}$  *to itself.* 

#### 7.6 The falling factorial

Recall  $(\underline{x})_n = x(x-1)\cdots(x-n+1)$ . Since  $(\underline{x})_{n+1} = (x-n)(\underline{x})_n$  the falling factorials are of Meixner type with parameters  $(a, \alpha, b, \beta) = (0, 1, 0, 0)$ . If T:  $x^n \mapsto (\underline{x})_n$  then it follows from (6.5.6) and (6.5.7) that

$$T(xf) = xT(f) - T(xf')$$
(7.6.1)  
$$T^{-1}(xf) = xT^{-1}(f) + xT^{-1}(f)'$$

The polynomials  $T^{-1}(x^n)$  are also known as the *exponential polynomials* [148, page 69] or the *Bell polynomials* 

**Proposition 7.31.** If  $T(x^n) = (\underline{x})_n = x(x-1)...(x-n+1)$ , then T maps  $P^{alt}$  to itself.  $T^{-1}$  maps  $P^{pos}$  to itself.

*Proof.* From (7.6.1), we can write T(xf) = xT(f) - TB(f) where B(f) = xf'. Since  $Bf \leq f$  for any f with all positive roots, Theorem 6.36 shows that T preserves interlacing for polynomials with all positive roots.

We can also write  $T^{-1}(xf) = xT^{-1}(f) + BT^{-1}f$ , so that the second statement now follows from Theorem 6.36.

*Remark* 7.32. We can use the falling factorial to explicate a subtlety of Theorem 6.3. If T is a transformation that preserves interlacing, then we can define S by T(xf) = S(Tf), and  $Sg \le g$  for g = Tf. From Theorem 6.3 we see that if  $Sg \le g$  then S has a certain form, namely  $Sg = (ax + b)g + (cx^2 + dx + e)g'$ . This would seem to contradict the fact that T is arbitrary.

In the case of the falling factorial, where  $T(x^n) = (\underline{x})_n$ , we can show that  $S(x^n) = x(x-1)^n$ . In particular,  $S(x^n)$  is *not* interlaced by  $x^n$ . Moreover, S does not have the simple form given above. The reason is that we have ignored the domains of definition of these results. The explicit form of Sg was determined by substituting  $x^n$  for g. However, we only know that  $Sg \leq g$  when g = Tf, and  $f \in \mathbf{P}^{nlt}$ , yet  $T^{-1}x^n$  is a polynomial in  $\mathbf{P}^{pos}$ .

To establish that  $S(x^n) = x(x-1)^n$ , define L(f(x)) = f(x-1). The map S is defined by  $S((\underline{x})_n) = (\underline{x})_{n+1}$ , and this can be rewritten as  $S((\underline{x})_n) = xL((\underline{x})_n)$ . By linearity, Sf = xLf, and substituting  $x^n = f$  shows that  $S(x^n) = x(x-1)^n$ .

The falling factorial transformation is connected to a series identity.

**Lemma 7.33.** If  $T(x^n) = (\underline{x})_{n'}$  then for any polynomial f we have

$$\sum_{\mathfrak{i}=0}^{\infty} \frac{f(\mathfrak{i})}{\mathfrak{i}!} x^{\mathfrak{i}} = e^{x} \operatorname{T}^{-1}(\mathfrak{f}).$$

*Proof.* By linearity we may assume that  $f(x) = x^n$ . If we let  $h_n(x) = \sum \frac{i^n}{i!} x^i$ , then it is easy to verify that  $xh'_n(x) = h_{n+1}$ . Since induction and integration by parts show that there is a polynomial  $g_n$  such that  $h_n = g_n e^x$ , we find that  $g_n$  satisfies  $g_{n+1} = x(g_n + g'_n)$ .

We can now show by induction that  $g_n = T^{-1}(x^n)$ . It is clearly true for n = 0, and the recursion (7.6.1) shows that  $T^{-1}(x^{n+1}) = x(g_n + g'_n) = g_{n+1}$ .

This is an ad hoc argument. See [148, page 70] for a proof based on the general principles of the operator calculus.  $\hfill \Box$ 

From (6.5.7) we can express  $T^{-1}$  as a composition:

$$\Gamma^{-1}(f) = f(x + xD)(1).$$
(7.6.2)

We could have applied Corollary 7.26 to deduce that  $T^{-1}$  preserves interlacing.

**Corollary 7.34.** *If a polynomial* f *has all negative roots, then for all positive* x

$$f(x)^2 \ge f(x-1)f(x+1).$$

*Proof.* First, assume that x is a positive integer. Using Lemma 7.33 we see if  $T(x^n) = (\underline{x})_n$  and  $f \in \mathbf{P}^{pos}$  then  $T^{-1}f \in \mathbf{P}^{pos}$ , and so  $e^x T^{-1}(f)$  is in  $\widehat{\mathbf{P}^{pos}}$ . The inequality is Newton's inequality (Theorem 4.8) for  $\widehat{\mathbf{P}}$ .

In general, let  $\alpha = r + \varepsilon$  where r is an integer and  $0 \le \varepsilon < 1$ . If  $g(x) = f(x + \varepsilon)$  then the desired inequality follows by applying the previous paragraph to g(x) since  $g(r) = f(\alpha)$ .

# 7.7 The rising factorial

Recall  $\langle \underline{x} \rangle_n = x(x+1) \cdots (x+n-1)$ . If  $T: x^n \mapsto \langle \underline{x} \rangle_n$  then

$$T(xf) = xT(f) + T(xf')$$
(7.7.1)  
$$T^{-1}(xf) = xT^{-1}(f) - xT^{-1}(f)'$$

**Corollary 7.35.** If  $T(x^n) = x(x+1)...(x+n-1)$ , then  $T^{-1}$  maps  $P^{alt}$  to itself. T maps  $P^{pos}$  to itself.

*Proof.* The proofs of the first two assertions are like the proofs for the falling factorial, and uses the recurrences of (7.7.1).

The rising factorial satisfies a different recurrence than the falling factorial, and it leads to a composition formula. Let<sup>2</sup> Af(x) = f(x + 1). Then

$$\begin{split} \langle \underline{\mathbf{x}} \rangle_{n+1} &= \mathbf{x}(\mathbf{x}+1) \cdots (\mathbf{x}+n) \\ &= \mathbf{x} \cdot \mathbb{A}(\mathbf{x} \cdots (\mathbf{x}+n-1)) \\ &= \mathbf{x} \mathbb{A} \langle \underline{\mathbf{x}} \rangle_n \end{split}$$

Consequently, if we define  $T(x^n) = \langle \underline{x} \rangle_n$  then

$$\mathsf{T}(\mathsf{x}^{\mathsf{n}}) = \langle \underline{\mathsf{x}} \rangle_{\mathsf{n}} = (\mathsf{x}\mathbb{A}) \langle \underline{\mathsf{x}} \rangle_{\mathsf{n}-1} = \dots = (\mathsf{x}\mathbb{A})^{\mathsf{n}}(1)$$

<sup>&</sup>lt;sup>2</sup>see § 8.8.7 for the motivation.

and so we have the useful composition formula

$$T(f) = f(xA)(1)$$
 (7.7.2)

For more information about the rising factorial, see §8.8.7.

Sometimes the rising factorial transformation is hidden. The following lemma is based on the definition of L found in [3].

**Lemma 7.36.** If  $g = \sum a_i x^i$  then define  $L(f,g) = \sum x^i a_i \Delta^i f(0)$  where  $\Delta g(x) = g(x+1) - g(x)$ . If  $f \in \mathbf{P}^{p \circ s}$  and  $g \in \mathbf{P}$  then  $L(f,g) \in \mathbf{P}$ .

*Proof.* Write  $f = \sum b_i \langle \underline{x} \rangle_i$ . Since  $\Delta \langle \underline{x} \rangle_i = i \langle \underline{x} \rangle_{i-1}$  and  $\Delta^i \langle \underline{x} \rangle_i = i!$  we see that  $b_i = \frac{(\Delta^i f)(0)}{i!}$  and hence  $L(f, g) = \sum x^i a_i b_i i!$ . Define  $T: \langle \underline{x} \rangle_i \mapsto x^i$  and set  $h = T(f) = \sum b_i x^i$ . If  $h \in \mathbf{P}^{\text{pos}}$  and  $g \in \mathbf{P}$  then by Theorem 9.87  $\sum x^i a_i b_i i! \in \mathbf{P}$ . From Corollary 7.35 we know  $T: \mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}^{\text{pos}}$ , so  $f \in \mathbf{P}^{\text{pos}}$  implies  $h \in \mathbf{P}^{\text{pos}}$ . Thus we are done.

The maps  $x^n \mapsto \langle \underline{x} \rangle_n$  and  $x^n \mapsto (\underline{x})_n$  are conjugates by the map  $x \mapsto -x$ , as expressed in the diagram



The diagram commutes since

$$\mathbf{x}^{\mathbf{n}} \mapsto (-1)^{\mathbf{n}} \mathbf{x}^{\mathbf{n}} \mapsto (-1)^{\mathbf{n}} (\underline{\mathbf{x}})_{\mathbf{n}} \mapsto (\underline{-1})_{\mathbf{n}} (-\mathbf{x})^{\mathbf{n}} = \langle \underline{\mathbf{x}} \rangle_{\mathbf{n}}.$$

If we compose the falling and rising factorial transformations we get

#### Lemma 7.37.

- 1. The transformation  $(\underline{x})_n \mapsto \langle \underline{x} \rangle_n$  maps  $P^{pos} \longrightarrow P^{pos}$ .
- 2. The transformation  $\langle \underline{x} \rangle_n \mapsto (\underline{x})_n$  maps  $P^{\text{alt}} \longrightarrow P^{\text{alt}}$ .
- 3. The transformation  $(\underline{x})_k \mapsto \langle \underline{x} \rangle_{n-k}$  maps  $P^{pos}(n) \longrightarrow P^{pos}(n)$ .
- 4. The transformation  $x^k \mapsto \langle \underline{x} \rangle_{n-k}$  maps  $P^{alt}(n) \longrightarrow P^{alt}(n)$ .
- 5. The transformation  $(\underline{x})_k \mapsto (\underline{\alpha})_k x^k$  maps  $P^{pos}(n) \longrightarrow P^{pos}(n)$  for  $\alpha > n-2$ .

*Proof.* The first one follows from the diagram



and the second is similar. The next follows from the diagram



The next two use the diagrams



Given any linear transformation T, the map  $T(x^i) \mapsto T(x^{i+1})$  usually does not preserve roots. The only non-trivial cases known are when  $T(x^n) = (\underline{x})_n$ ,  $\langle \underline{x} \rangle_n$ , or  $H_n$  (Corollary 7.44).

**Lemma 7.38.** The linear transformations  $(\underline{x})_i \mapsto (\underline{x})_{i+1}$  and  $\langle \underline{x} \rangle_i \mapsto \langle \underline{x} \rangle_{i+1}$  map  $P \longrightarrow P$ .

*Proof.* If T(f) = xf and S(f) = f(x + 1) then the fact that the diagram



commutes shows that the desired transformation is  $TS^{-1}$ . Now both  $S^{-1}$  and T map **P** to itself, and so the same is true for  $TS^{-1}$ . Incidentally, this shows  $T(x^n) = x(x-1)^n$ . The proof for  $\langle \underline{x} \rangle_i$  is similar.

**Lemma 7.39.** Let  $T(x^n) = \langle \underline{x} \rangle_n / n!$ .

$$T: \mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}^{\text{pos}}$$
$$T: \mathbf{P}^{(1,\infty)} \longrightarrow \mathbf{P}^{(1,\infty)}$$

*Proof.* We have the identity

$$(T(f))(1-x) = T(f(1-x))$$

which is proved by taking  $f(x) = x^k$ , and using

$$\frac{\langle \underline{1-x} \rangle_{k}}{k!} = \sum_{i=0}^{k} \frac{\langle \underline{x} \rangle_{i}}{i!} \binom{k}{i}$$

Now we know that  $x^n \mapsto \langle \underline{x} \rangle_n$  maps  $\mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}^{\text{pos}}$ , and upon application of the exponential map, so does T. The identity above implies that we have the following commuting diagram



*Remark* 7.40. T above does not  $\mathbf{P}^{(0,1)}$  to  $\mathbf{P}$ . Indeed,  $T(x - 1/2)^2$  has complex roots. It is known [26] that T maps  $\mathbf{P}^{\iota+\mathbb{R}}$  to itself. See also Question 59.

If we need to prove a result about polynomials without all real roots, then sign interlacing is useful. Consider

**Lemma 7.41.** *If* g *has* p *positive roots, and*  $\alpha$  *is positive, then*  $h = (x - \alpha)g - xg'$  *has at least* p + 1 *positive roots.* 

*Proof.* Suppose that the positive roots of g are  $r_1 < \cdots < r_p$ . The sign of  $h(r_i)$  is the sign of  $-g'(r_i)$  and so is  $(-1)^{p+i+1}$ . This shows that h has at least p roots, one between each positive root of g, and one to the right of the largest root of g. The sign of h(0) is the sign of  $-\alpha h(r_1)$ . Since  $\alpha$  is positive, there is a root between 0 and  $r_1$ , for a total of at p + 1 roots.

**Lemma 7.42.** If  $T^{-1} : \langle \underline{x} \rangle_n \mapsto x^n$ , and  $f \in P$  has p positive roots, then  $T^{-1}(f)$  has at least p positive roots.

*Proof.* We prove this by induction on the number p of positive roots of f. There is nothing to prove if p = 0. Let  $g(x) = T^{-1}(f)$ . By induction we assume that g has p positive roots. If  $h(x) = T^{-1}(x - \alpha)f$  where  $\alpha$  is positive, then the recurrence (7.7.1) shows that  $h = (x - \alpha)g - xg'$ . Lemma 7.41 shows that h has at least p + 1 positive roots.

#### 7.8 Hermite polynomials

The Hermite polynomials are the orthogonal polynomials for the weight function  $e^{-x^2}$  on  $(-\infty, \infty)$ . There is an explicit formula (the Rodrigues' formula) for the Hermite polynomials [168, page 25]:

$$H_{n} = (-1)^{n} e^{x^{2}} \left(\frac{d}{dx}\right)^{n} e^{-x^{2}}$$
(7.8.1)

an expansion in a series

$$H_{n} = \sum_{k=0}^{n/2} \frac{n!}{(n-2k)!} (-1)^{k} (2x)^{n-2k}$$
(7.8.2)

and a recurrence relation

$$H_n = 2xH_{n-1} - 2nH_{n-2} \tag{7.8.3}$$

An addition formula for Hermite polynomials is

$$H_{n}(x+y) = \sum_{k=0}^{n} {\binom{n}{k}} H_{k}(x)(2y)^{n-k}$$
(7.8.4)

The Hermite polynomials are not of Meixner type since they aren't monic, but if we rescale them to make them monic then we get a recursion of the form  $p_{n+1} = xp_n - np_{n-1}$  which is Meixner with parameters  $(a, \alpha, b, \beta) = (0, 0, 1, 0)$ . From the recurrence (7.8.3) it is easy to verify that  $H'_n = 2nH_{n-1}$ . There is also a differential recurrence

$$H_{n+1} = 2xH_n - (H_n)' = (2x - D)H_n$$
(7.8.5)

and consequently

$$H_n = (2x - D)^n (1)$$
(7.8.6)

If we set  $T(x^n) = H_n$  then we have the composition

$$T(f) = f(2x - D)(1)$$
(7.8.7)

It is easy to verify that

$$T^{-1}(f) = f(x/2 + D)(1)$$
 (7.8.8)

We need to be careful here because multiplication by x does not commute with D. We define  $(2x - D)^n(1)$  inductively:

$$(2x - D)^{n}(1) = (2x - D) ((2x - D)^{n-1}(1))$$

and for  $f = \sum \alpha_i x^i$  we use linearity:

$$f(2x - D)(1) = \sum a_i (2x - D)^i(1)$$

In this light the following lemma establishes an interesting relation between these two definitions. The proof is by induction and omitted.

**Lemma 7.43.** If  $f(x) = (ax + bD)^n(1)$  then

$$f(cx + dD)(1) = (acx + (ad + b/c)D)^{n}(1)$$

The linear recurrences satisfied by the transformation  $T(x^n) = H_n$  follow from (7.8.3), and are

$$T(xf) = 2xT(f) - T(f')$$
 (7.8.9)

$$\mathsf{T}^{-1}(\mathsf{x}\mathsf{f}) = (1/2)\mathsf{x}\mathsf{T}^{-1}(\mathsf{f}) + (\mathsf{T}^{-1}\mathsf{f})' \tag{7.8.10}$$

**Corollary 7.44.** Let  $H_n$  be the Hermite polynomial.

- 1. If  $T(x^n) = H_n$  then  $T: P \longrightarrow P$ .
- 2. The linear transformation  $x^i \mapsto H_{n-i}$  maps P(n) to itself.
- 3. The linear transformation  $H_i \mapsto H_{i+1}$  maps P(n) to itself.

*Proof.* For the first part apply Corollary 7.25 to (7.8.7). The next map is the conjugate of T by the reversal map – see § 7.7.2. The map  $H_i \mapsto H_{i+1}$  is equivalent to  $f \mapsto (2x - D)f$  by (7.8.5), and this maps **P** to itself by Lemma 1.20. As an aside, the map  $H_n \to H_{n+1}$  evaluated at  $x^n$  is  $x^{n-1}(2x^2 - n)$ .

There are also generalized Hermite polynomials [146, page 87] that are defined for any positive  $\nu$ . They satisfy  $H_0^{\nu} = 1$  and

$$\mathsf{H}_{n+1}^{\mathsf{v}} = \mathsf{x} \mathsf{H}_{n}^{\mathsf{v}} - \mathsf{v} \mathfrak{n} \mathsf{H}_{n-1}^{\mathsf{v}}$$

The corresponding linear transformation preserves roots just as the ordinary Hermite transformation does.

Different representations of orthogonal polynomials can lead to different proofs of Corollary 7.44. For instance, using (7.8.2) the Hermite polynomials can be defined in terms of the derivative operator. [57]

$$H_n = 2^n \exp(-D^2/4)(x^n)$$
(7.8.11)

The transformation  $T: x^n \longrightarrow H_n$  can be factored as  $T_1T_2$  where  $T_1(x^n) = (2x)^n$  and  $T_2(x_n) = \exp(-D^2/4)(x^n)$ . Since  $\exp(-x^2/4)$  is in  $\widehat{\mathbf{P}}$ , we can apply Theorem 5.36, and conclude that  $T_2(f) \in \mathbf{P}$ . Thus, T maps  $\mathbf{P}$  to itself.

The action of T:  $x^n \longrightarrow H_n(x)$  on sin and cos is surprisingly simple. It's easiest to compute T on  $e^{ix}$ .

$$T(e^{\iota x}) = \sum_{k=0}^{\infty} H_k(x) \frac{\iota^k}{k!}$$
  
=  $e^{-2\iota x+1}$  (from Table 15.2)  
=  $e(\cos(2x) - \iota \sin(2x))$   
=  $T(\cos(x) + \iota \sin(x))$ 

and so we conclude that

$$T(\cos(x)) = e\cos(2x)$$
$$T(\sin(x)) = -e\sin(2x)$$

# 7.9 Charlier polynomials

The monic Charlier polynomials with parameter  $\alpha$  are orthogonal polynomials that satisfy the recurrence formula

$$C_{n+1}^{\alpha}(x) = (x - n - \alpha)C_n^{\alpha}(x) - \alpha nC_{n-1}^{\alpha}(x)$$
(7.9.1)

The first five Charlier polynomials are in Appendix 24.9. Their recurrence shows them to be of Meixner type, with parameters  $(a, \alpha, b, \beta) = (\alpha, 1, \alpha, 0)$ . They satisfy the difference equation (analogous to the differential equation for the Hermite polynomials)

$$\Delta C_n^{\alpha}(\mathbf{x}) = n C_{n-1}^{\alpha}(\mathbf{x}) \tag{7.9.2}$$

where  $\Delta(f) = f(x + 1) - f(x)$ . Consequently, the Charlier polynomials give an infinite sequence of polynomials with all real roots that is closed under differences, just as the infinite sequence of Hermite polynomials is closed under differentiation.

The corresponding linear transformation is  $\mathsf{T}(x^n)=\mathsf{C}_n^\alpha(x).$  T satisfies the recurrences

$$T(xf) = (x - \alpha)Tf - T(xf') - \alpha T(f')$$

$$T^{-1}(xf) = (x + \alpha)T^{-1}(f) + x(T^{-1}f)' + \alpha(T^{-1}f)'$$

$$T^{-1}(f) = f((x + \alpha)(D + 1))(1)$$
(7.9.3)

**Corollary 7.45.** If  $T(x^n) = C_n^{\alpha}(x)$ , the n-th Charlier polynomial, then T maps  $P^{\text{alt}}$  to itself.  $T^{-1}$  maps  $P^{\text{pos}}$  to itself.

*Proof.* If we set A(f) = xf' and D(f) = f', then the Charlier polynomials satisfy

$$T(\mathbf{x}\mathbf{f}) = (\mathbf{x} - \alpha)T(\mathbf{f}) - TA(\mathbf{f}) - \alpha TD(\mathbf{f})$$
$$T^{-1}(\mathbf{x}\mathbf{f}) = (\mathbf{x} + \alpha)T^{-1}(\mathbf{f}) + AT^{-1}(\mathbf{f}) + \alpha DT^{-1}(\mathbf{f})$$

Since  $A(f) \ll f$  and  $f \ll Df$  the results follow from Theorem 6.36.

Since  $T^{-1}$  maps  $\mathbf{P}^{pos}$  to itself, we know that  $B_{n+1} \stackrel{+}{\sim} B_n$ . Using the identity

$$B_{n+1} = x(B_n + B'_n)$$

we see that actually  $B_{n+1} \leq B_n$ . (See p. 724.)

*Remark* 7.46. The Charlier polynomials satisfy a formula that is analogous to the composition formula for Hermite polynomials. This will be used in § 8.8.6. If we define  $\mathbb{A}^{-1} f(x) = f(x - 1)$ , then the transformation T above satisfies

$$T(x^{n}) = C_{n}^{\alpha}(x)$$
  
=  $(x\mathbb{A}^{-1} - \alpha)C_{n-1}^{\alpha}(x)$  by (7.9.1) and (7.9.2)  
=  $(x\mathbb{A}^{-1} - \alpha)^{n}(1)$  by induction

and so

$$T(f) = f(xA^{-1} - \alpha)(1)$$
(7.9.4)

# 7.10 Laguerre polynomials

As with so many special functions there are variations in the definition of the Laguerre polynomials. The usual notation is  $L_n^{\alpha}(x)$ ; we will see that the  $\alpha$  is quite important, and so we also write  $L_n(x;y) = L_n^y(-x)$ . The definition of that we use is in [72], but we set the leading coefficient of  $L_n(x;y)$  to 1/n!. The Rodrigures formula is

$$L_{n}^{a}(x) = \frac{1}{x^{a}e^{-x}} \frac{1}{n!} \mathsf{D}^{n}(e^{-x}x^{n+a})$$
(7.10.1)  
$$L_{n}(x;y) = \frac{1}{x^{y}e^{x}} \frac{1}{n!} \mathsf{D}^{n}(e^{x}x^{n+y})$$

and a series expansion is

$$L_{n}(x;y) = \sum_{k=0}^{n} {\binom{n+y}{n-k} \frac{x^{k}}{k!}}$$
(7.10.2)

The Hermite polynomials (7.8.11) and the Laguerre polynomials have nice representations in terms of the exponential of a quadratic differential.

$$e^{\alpha \partial_{x} \partial_{y}} (x^{n} y^{n}) = n! \alpha^{n} L_{n}(\frac{-xy}{\alpha})$$
(7.10.3)

The homogenized Laguerre polynomials have an expansion

$$(y+1)^{n}L_{n}(\frac{x}{y+1}) = \sum_{k=0}^{n}L_{k}(x)\binom{n}{k}y^{n-k}$$
(7.10.4)

The expansion (7.10.2) shows that  $L_n(x; y)$  is a polynomial in x and y with positive coefficients.

An alternative definition is found in [146] with leading coefficient  $(-1)^n$ . We denote these alternative polynomials by  $\tilde{L}_n^{\alpha}(x)$ , and note that we have the simple relationship  $\tilde{L}_n^{\alpha}(x) = n! L_n^{\alpha}(x)$ . The three term recurrence is not useful for us, but we will use the composition formula [146, page 110]

$$\begin{split} \tilde{\mathsf{L}}_{\mathfrak{n}}^{(\alpha)}(\mathbf{x}) &= (\mathbf{x}\mathsf{D} - \mathbf{x} + \alpha + 1)\tilde{\mathsf{L}}_{\mathfrak{n}}^{\alpha+1} \\ &= (\mathbf{x}\mathsf{D} - \mathbf{x} + \alpha + 1)(\mathbf{x}\mathsf{D} - \mathbf{x} + \alpha + 2)\cdots(\mathbf{x}\mathsf{D} - \mathbf{x} + \alpha + n)(1) \\ &= \left\langle \underline{\mathbf{x}}\mathsf{D} - \mathbf{x} + \alpha + 1\right\rangle_{\mathfrak{n}}(1). \end{split}$$
(7.10.5)

**Corollary 7.47.** If T is the Laguerre transformation  $T(x^n) = \tilde{L}_n^{(\alpha)}(x)$  where  $\alpha \ge -1$  then  $T: \mathbf{P}^{pos} \longrightarrow \mathbf{P}^{alt}$ .

*Proof.* If we define  $S(f) = (xD-x+\alpha+1)(f)$  then from (7.10.5) the composition of maps

$$x^{n} \mapsto \langle \underline{x} \rangle_{n} \mapsto \langle \underline{S} \rangle_{n}(1)$$

is the Laguerre transformation. If  $U(x^n) = \langle \underline{x} \rangle_n$  and V(f) = f(S)(1), then T = VU. From Corollary 7.35 U maps  $\mathbf{P}^{pos}$  to  $\mathbf{P}^{pos}$ . By Corollary 7.24 V maps  $\mathbf{P}^{pos}$  to  $\mathbf{P}$ . Consequently, VU maps  $\mathbf{P}^{pos}$  to  $\mathbf{P}$ . Since the coefficients of T(f) are alternating in sign, we see that T maps to  $\mathbf{P}^{alt}$ .

If we define  $T(x^n) = \tilde{L}_n^{\alpha}(-x)$ , and  $S(x^n) = L_n(x; \alpha)$  then  $S = T \circ EXP$ . It follows that S maps  $\mathbf{P}^{pos}$  to itself. We will see in Lemma 15.41 that S actually maps  $\mathbf{P}$  to itself.

The Laguerre polynomials are also related to the linear transformation

$$R: f \mapsto D^n(x^n f)$$

where n is a fixed positive integer, whose action on monomials is  $R(x^i) = (n+i)_i x^i$ . The exponential generating function corresponding to R is

$$\sum_{i=0}^{\infty} \frac{R(x^{i})}{i!} = \sum_{i=0}^{\infty} \frac{(n+i)!}{i! \, i!} x^{i} = e^{x} \tilde{L}_{n}(-x).$$
(7.10.6)

Since both  $e^x$  and  $\tilde{L}_n(-x)$  are in  $\widehat{\mathbf{P}^{\text{pos}}}$ , so is their product. The fact that the generating function of R is in  $\widehat{\mathbf{P}^{\text{pos}}}$  also follows from Theorem 15.22 since R is a multiplier transformation that maps  $\mathbf{P}^{\text{pos}}$  to itself.

# 7.11 The integration and exponential transformations

The exponential transformation  $x^n \mapsto x^n/n!$  is a fundamental linear transformation that maps **P** to **P**. Although not all polynomials in **P** have integrals that are in **P** (see Example 1.2) we can find some polynomials for which *all* integrals have a root. These polynomials are exactly the image of the exponential transformation. We first define the integral ( $\int$ ) and exponential (EXP) transformations. Define

$$exp: x^{n} \mapsto x^{n}/n!$$

$$\int : x^{n} \mapsto x^{n+1}/n + 1$$

$$\int^{k} : x^{n} \mapsto x^{n+k}/\langle \underline{n+1} \rangle_{k}$$

$$\int^{k} exp: x^{n} \mapsto x^{n+k}/(n+k)!$$

Since  $e^x \in \widehat{\mathbf{P}}$ , we can apply Theorem 15.22 to conclude that EXP maps **P** to itself. The integral doesn't preserve all real roots as Example 1.2 shows, but we do have

**Lemma 7.48.** If f is a polynomial with all real roots, and for every  $k \ge 0$  we define  $T(f) = \int^k EXP \ f$  then

- 1. T maps **P** to **P**.
- 2. T maps  $P^{\text{pos}}$  to  $P^{\text{pos}}$ .
- 3. T maps  $P^{\text{alt}}$  to  $P^{\text{alt}}$ .
- 4. T preserves interlacing.

*Proof.* It follows from the definition of  $\int_{k}^{k} EXP$  that

$$\int^{k} EXP f = e^{x} * (x^{k}f)$$

where "\*" is the Hadamard product. The results now follow from the fact that  $e^x \in \widehat{\mathbf{P}^{\text{pos}}}$  and Hadamard products preserve interlacing.

When we apply various exponential transformations, we get some interesting transformations that we express in terms of coefficients:

**Corollary 7.49.** If  $f = \sum_{i=0}^{n} a_i x^i$  is in **P** then the following polynomials also have all real roots:

1. 
$$\sum_{i=0}^{n} \frac{a_{i}}{i!} x^{i}$$
  
2. 
$$\sum_{i=0}^{n} \frac{a_{i}}{(n-i)!} x^{i}$$
  
3. 
$$\sum_{i=0}^{n} \frac{a_{i}}{i!(n-i)!} x^{i}$$
  
4. 
$$\sum_{i=0}^{n} {n \choose i} a_{i} x^{i}$$
  
5. 
$$\sum_{i=0}^{n} (\underline{n})_{i} a_{i} x^{i}$$

*Proof.* The first is EXP applied to f, the second is EXP applied to the reverse, and the third is EXP applied to the second. Multiplying the third by n! gives the fourth, and the fifth is n! times the second.

We can prove a more general exponential result using the gamma function:

**Lemma 7.50.** For any positive integer k the transformations  $x^i \mapsto \frac{i!}{(ki)!}x^i$  and  $x^i \mapsto \frac{x^i}{(ki)!}$  map P to itself.

*Proof.* In Example 5.8 we saw that  $\Gamma(z + 1)/\Gamma(kz + 1)$  is in  $\mathbf{P}$ , and has all negative roots. The result now follows from Corollary 7.20 and (7.4.2) since  $\Gamma(k+1)/\Gamma(ki+1) = i!/(ki)!$ .

It is not an accident that EXP occurs along with  $\int$ .

**Theorem 7.51.** If f is a polynomial such that  $\int^k f$  has all real roots for infinitely many positive integers k, then there is a polynomial g with all real roots such that f = EXP(g).

Proof. If we write 
$$f(x) = \sum_{i=0}^{n} a_i \frac{x^i}{i!}$$
 then  

$$\int^k f(x) = \sum_{i=0}^{n} a_i \frac{x^{i+k}}{(i+k)!}$$
(7.11.1)

The polynomial on the right hand side of (7.11.1) has all real roots. Replace x by kx:

$$\sum_{i=0}^n \alpha_i \frac{(kx)^{i+k}}{(i+k)!} = \frac{k^k x^k}{k!} \sum_{i=0}^n \alpha_i \frac{k}{k+1} \cdots \frac{k}{k+i} x^k$$

so the n-th degree polynomial

$$\sum_{i=0}^n a_i \frac{k}{k+1} \cdots \frac{k}{k+i} x^k$$

has all real roots. Taking the limit as  $k \to \infty$  we see that  $\sum_{i=0}^{n} a_i x^i$  has all real roots. This last polynomial is the desired g.

The Laplace transform, when restricted to polynomials, is closely related to the exponential transformation. If f(t) is any function then the Laplace transform of f(t) is

$$\mathcal{L}(f)(x) = \int_0^\infty e^{-xt} f(t) \, dt.$$

If x is positive then  $\mathcal{L}(t^n)(x) = \frac{n!}{x^{n+1}}$ . We say that the Laplace transform  $\mathcal{L}(f)$  of a polynomial  $f = \sum a_i x^i$  is  $\sum a_i \frac{i!}{x^{i+1}}$ . Consequently, we can express  $\mathcal{L}(f)$  in terms of EXP:

$$\mathcal{L}(f)(\mathbf{x}) = \frac{1}{\mathbf{x}} E X P^{-1}(f)(\frac{1}{\mathbf{x}})$$

since  $EXP^{-1}(x^n)(\frac{1}{x}) = \frac{n!}{x^n}$ . If f is a polynomial of degree n and  $\mathcal{L}(f)(x)$  has n real roots, then 0 is not a root, so  $EXP^{-1}(f)(\frac{1}{x})$  has all real roots. Taking the reverse shows that  $EXP^{-1}(f)(x)$  has all real roots, and hence applying EXP yields that  $f \in \mathbf{P}$ . Summarizing,

**Lemma 7.52.** If f is a polynomial of degree n and  $\mathcal{L}(f)$  has n real roots then  $f \in \mathbf{P}(n)$ .

### 7.12 **Binomial transformations**

In this section we study the transformation T:  $x^n \mapsto \binom{x}{n}$  and its inverse. The many identities satisfied by binomial coefficients yield recurrences for these transformations.

**Corollary 7.53.** The linear transformation  $x^i \mapsto \binom{x}{i}$  maps

$$\mathbf{P}^{(0,\infty)} \longrightarrow \mathbf{P}^{(0,\infty)}$$
  
 $\mathbf{P}^{(-\infty,-1)} \longrightarrow \mathbf{P}^{(-\infty,-1)}$ 

*Proof.* The map T is the composition  $x^n \longrightarrow x^n/n! \longrightarrow {\binom{x}{n}}$  where the second map is the falling factorial (see Proposition 7.31) and the first is the exponential map. Since the falling factorial maps  $P^{\alpha lt}$  to itself we have proved the first part. For the second one we use the identity  $\mathbb{A}T = T\mathbb{A}$  where  $\mathbb{A}f(x) = f(-x-1)$ . This yields the communicative diagram



since we know the top row, and  $\mathbb{A}^{-1} = \mathbb{A}$ 

**Corollary 7.54.** The map  $T(x^i) = \binom{x+n-i}{n}$  maps  $P^{(0,1)}$  to  $P^{alt}$ .

*Proof.* This follows from Lemma 6.40 and the diagram following. See Lemma 8.30 for a more general result.

It is not the case that T maps  $\mathbf{P}^{-1,0}$  to itself - even  $T(x+1/2)^2$  has imaginary roots. We are now going to show that  $T^{-1}$  maps  $\mathbf{P}^{-1,0}$  to itself.

We use a simple binomial identity to get a recurrence for  $T^{-1}$ .

$$\begin{aligned} x \begin{pmatrix} x \\ n \end{pmatrix} &= (n+1) \begin{pmatrix} x \\ n+1 \end{pmatrix} + n \begin{pmatrix} x \\ n \end{pmatrix} \\ T^{-1}(x \begin{pmatrix} x \\ n \end{pmatrix}) &= (n+1)x^{n+1} + nx^n \\ &= x \cdot x^n + x(x+1) \cdot nx^{n-1} \\ &= x \cdot T^{-1} \begin{pmatrix} x \\ n \end{pmatrix} + x(x+1) \left( T^{-1} \begin{pmatrix} x \\ n \end{pmatrix} \right)' \end{aligned}$$

By linearity it follows that

$$T^{-1}(xf) = xT^{-1}f + x(x+1)(T^{-1}f)'$$

**Lemma 7.55.** The map  $x^n \mapsto \binom{x}{n}$  satisfies  $T^{-1} : \mathbf{P}^{(-1,0)} \longrightarrow \mathbf{P}^{(-1,0)}$  and preserves interlacing.

*Proof.* If  $T^{-1}f \in \mathbf{P}^{(-1,0)}$  then  $T^{-1}f \leq (T^{-1}f)'$  and so it follows that  $x(x + 1)T^{-1}f \leq T^{-1}f$  since all roots lie in (-1,0). Using the recurrence yields  $T^{-1}(xf) \leq Tf$  and we can follow the argument of Corollary 1.46 to finish the proof.

The behavior of  $T^{-1}$  at the endpoints of (-1, 0) is unusual:

**Lemma 7.56.**  $T^{-1}(x^n) \leq T^{-1}(x+1)^n$ . In fact,

$$(x+1)T^{-1}(x^n) = xT^{-1}(x+1)^n$$

*Proof.* If we define the affine transformation  $\mathbb{B}f(x) = f(x + 1)$  then we will prove the more general result  $(x+1)T^{-1}f = xT^{-1}\mathbb{B}f$ . It suffice to prove this for a basis, so choose  $f = \binom{x}{n}$ . The computation below establishes the lemma.

$$xT^{-1}\mathbb{B}f = xT^{-1}\mathbb{B}\binom{x}{n} = xT^{-1}\binom{x+1}{n} = xT^{-1}\binom{x}{n} + \binom{x}{n-1}$$
$$= x(x^{n} + x^{n-1}) = (x+1)x^{n} = (x+1)T^{-1}f$$

**Corollary 7.57.** The map  $T: x^n \mapsto \binom{x}{n}$  satisfies

$$\int_0^1 \mathsf{T}^{-1}(\mathsf{x}+\mathsf{t})^n \, d\mathsf{t} \in \boldsymbol{P} \quad \textit{for } \mathsf{n} = 1, 2, \cdots.$$

*Proof.* We can apply Proposition 3.37 once we show that  $T^{-1}$  satisfies  $T^{-1}(x + a)^n \leq T^{-1}(x + b)^n$  if  $0 \leq b \leq a \leq 1$ . We show more generally that if the roots of g are all at least as large as the corresponding roots of f then  $T^{-1}g \leq T^{-1}f$ . Since  $T^{-1}$  preserves interlacing we know that there is a sequence of interlacing polynomials from  $T^{-1}x^n$  to  $T^{-1}(x + 1)^n$  that includes f and g. Since the endpoints of this sequence interlace by the preceding lemma, we have a mutually interlacing sequence, and therefore  $T^{-1}g \leq T^{-1}f$ .

We now consider the transformation T:  $x^n \mapsto \langle \underline{x} \rangle_n / n!$ . Since  $\langle \underline{-x} \rangle_n = (-1)^n {x \choose n}$  we can apply the results of this section to conclude

**Lemma 7.58.** The map  $T: x^n \mapsto \langle \underline{x} \rangle_n / n!$  satisfies

- 1. T:  $\mathbf{P}^{(-\infty,0)} \cup \mathbf{P}^{(1,\infty)} \longrightarrow \mathbf{P}$ .
- 2.  $\mathsf{T}^{-1}: \mathbf{P}^{(0,1)} \mapsto \mathbf{P}$ .

If  $\mathbb{A}f(x) = f(1-x)$  then  $(x - 1/2)^k$  is invariant up to sign under T and its roots are invariant under A. The following shows that there are complex roots, and they all have real part 1/2.

**Lemma 7.59.** If  $T: x^n \mapsto \langle \underline{x} \rangle_n / n!$  then all roots of  $T(x - 1/2)^n$  have real part 1/2. If n > 1 then there is a complex root.

*Proof.* If we define

$$S(f(x)) = (T(x - 1/2))(x + 1/2)$$

then S satisfies the recurrence

$$S(x^{n+1}) = \frac{1}{n+1}S(x^n) + \frac{n}{4(n+1)}S(x^{n-1})$$

Since S(1) = 1 and S(x) = x, it follows that  $S(x^n)$  has purely imaginary roots. The first part now follows from  $S(x^n)(x - 1/2) = T(x - 1/2)^n$ .

If  $T(x - 1/2)^n$  had all real roots then they would all be 1/2, but it is easy to check that  $T(x - 1/2)^n$  is not a multiple of  $(x - 1/2)^n$ .

# 7.13 Eulerian polynomials

The Eulerian polynomials can be defined as the numerator of the sum of an infinite series. If r is a non-negative integer, then the Eulerian polynomial  $A_r(x)$  is the unique polynomial of degree r such that

$$\sum_{i=0}^{\infty} i^{r} x^{i} = \frac{A_{r}(x)}{(1-x)^{r+1}}.$$
(7.13.1)

If we differentiate (7.13.1) and multiply by x, we get

$$\sum_{i=0}^{\infty} i^{r+1} x^{i} = x \frac{(r+1)A_{r} + (1-x)A'_{r}}{(1-x)^{r+2}}.$$

and so

$$A_{0} = 1 \qquad A_{1} = x$$
  

$$A_{r} = x \left( rA_{r-1} + (1-x)A'_{r-1} \right)$$
(7.13.2)

 $A_n$  has all non-negative coefficients and Corollary 7.29 shows that the roots of  $A_r$  are all non-positive, and  $A_r \leq A_{r-1}$ . Using (7.13.2), we can write the recurrence for the transformation T:  $x^n \mapsto A_n$ :

$$T(xf) = xTf + xT(xf') + x(1-x)(Tf)'$$
(7.13.3)

There is a Möbius transformation that simplifies this recurrence.

**Lemma 7.60.** If  $Mz = \frac{z+1}{z}$  then  $T_M : \mathbf{P}^{(0,1)} \mapsto \mathbf{P}^{(-1,0)}$  where  $T_M$  is defined in (6.7.3).

*Proof.* Let  $B_n = x^n A_n(\frac{x+1}{x})$ . A bit of calculation using (7.13.2) shows that  $B_0 = 1$  and

$$B_{n+1} = (x+1)(B_n + xB'_n)$$

This leads to the recurrence for  $T_M$  :

$$\mathsf{T}_{\mathsf{M}}(\mathsf{x}\mathsf{f}) = (\mathsf{x}+1)(\mathsf{T}_{\mathsf{M}}(\mathsf{f}) + \mathsf{x}\mathsf{T}_{\mathsf{M}}(\mathsf{f})')$$

The conclusion now follows easily by induction using this recurrence.

**Corollary 7.61.** If  $T: x^n \mapsto A_n$  then T maps  $P^{(-1,0)}$  to  $P^{pos}$ .

*Proof.* If V is a linear transformation and M is a Möbius transformation then write  $(M) \circ V = V_M$ . Setting  $S = T_{(z+1)/z}$  then with this notation we have that

$$\mathsf{T} = (z-1) \circ (\frac{1}{z}) \circ \mathsf{T}_{(z+1)/z}$$

since the composition  $(z-1) \circ (1/z) \circ ((z+1)/z)$  is the identity. We now apply Lemma 7.60 and the general results of §6.6.7 to deduce the commutative diagram

$$\mathbf{P}^{(0,1)} - - - \stackrel{\mathrm{T}}{-} - \mathbf{P}^{\operatorname{pos}} \\ \bigvee_{\gamma}^{\mathsf{S}} \qquad (z-1) \\ \mathbf{P}^{(-1,0)} \xrightarrow{(1/z)} \mathbf{P}^{(-\infty,-1)}$$

It appears that the roots of  $A_n$  go to minus infinity, and hence the roots of polynomials in the image of  $\mathbf{P}^{(-1,0)}$  are not contained in any finite interval.

If f is a polynomial of degree n, then we define the transformation W by

$$\sum_{i=0}^{\infty} f(i)x^{i} = \frac{(Wf)(x)}{(1-x)^{n+1}}$$
(7.13.4)

If  $f = \sum_{1}^{n} b_{i} x^{i}$  then

$$Wf = \sum_{1}^{n} b_i (1-x)^{n-i} A_i$$

This shows that Wf is a polynomial of degree n. W can be realized as a composition with T and a homogeneous transformation

$$W: x^k \mapsto A_k \mapsto (1-x)^{n-k} A_k$$

**Corollary 7.62.** The map W defines a linear transformation from  $P^{(-1,0)}$  to P.

*Proof.* If  $f \in \mathbf{P}^{(-1,0)}$  then  $\mathsf{T} f \in \mathbf{P}^{\mathfrak{alt}}$  where T is given in Corollary 7.61. Since  $W(x^n) = A_n \in \mathbf{P}^{pos}$  has leading coefficient 1 we can apply Lemma 12.27 to conclude that  $Wf \in \mathbf{P}$ .

### 7.14 Euler-Frobenius polynomials

The Euler-Frobenius polynomials [51] arise in interpolation problems. These polynomials satisfy the recurrence

$$P_{n+1} = 2xP_n - (1 - x^2)P'_n$$

$$= (2x + (x^2 - 1)D)P_n$$

$$= D((x^2 - 1)P_n)$$
(7.14.1)

If we define the linear transformations  $T(x^n)=P_n$  and  $S(f)=\mathsf{D}\left((x^2-1)f\right)$  then

$$\mathsf{T}(\mathsf{x}^{\mathsf{n}}) = \mathsf{S}^{\mathsf{n}}(1)$$

and for any polynomial f we have

$$\mathsf{T}(\mathsf{f}) = \mathsf{f}(\mathsf{S})(1)$$

By Corollary 7.30 the transformation  $2x + (x^2 - 1)D + \alpha$  maps  $P^{(-1,1)}$  to itself for  $|\alpha| < 1$  so we conclude

**Lemma 7.63.** If  $T(x^n) = P_n$  then T maps  $P^{(-1,1)}$  to itself.

There is a modification of the Euler-Frobenius polynomials that are also called Euler-Frobenius polynomials [183]. They are defined by applying a Möbius transformation to  $P_n$ :

$$\mathsf{E}_{\mathsf{n}}(\mathsf{x}) = (\mathsf{x}-1)^{\mathsf{n}} \mathsf{P}_{\mathsf{n}}\left(\frac{\mathsf{x}+1}{\mathsf{x}-1}\right)$$

These polynomials satisfy the recurrence

$$E_n = (1 + nx)E_{n-1} + x(1 - x)E'_{n-1}$$
(7.14.2)

It is not obvious that all the coefficients of  $E_n$  are positive, but this can be shown [183] by explicitly determining the coefficients. The recurrence (7.14.2) then shows that all the roots of  $E_n$  are negative, and that  $E_n \leq E_{n-1}$ . We can use the fact that  $P_n = S^n(1)$  to derive a "Rodrigues' formula" for  $E_n$ :

$$E_n = \frac{(1-x)^{n+2}}{x} (x\mathsf{D})^n \frac{x}{(1-x)^2}$$

#### 7.15 Even and odd parts

When does a polynomial with all real coefficients have roots whose real parts are all negative? The answer is given by Hurwitz: write a polynomial f in

terms of its even and odd parts. The polynomial f has all roots with negative real part iff the polynomials corresponding to the even and odd parts interlace. See [138, V171.4] or Proposition 22.31.

In this section we generalize the transformations that assign either the even or odd part to a polynomial. Under appropriate assumptions, these transformations preserve roots and form mutually interlacing sequences.

Write  $f(x) = f_e(x^2) + xf_o(x^2)$ . The even part of f is  $f_e(x)$ , and the odd part is  $f_o(x)$  so we define two linear transformations by  $T_e(f) = f_e(x)$  and  $T_o(f) = f_o(x)$  (see Page 84). It is clear that  $T_e$  and  $T_o$  are linear, and satisfy the recurrences

$$T_e(xf) = xT_o(f)$$
(7.15.1)  
$$T_o(xf) = T_e(f).$$

**Theorem 7.64** (Hurwitz's theorem). If  $f \in P^{pos}$  then  $T_e(f)$  and  $T_o(f)$  are in  $P^{pos}$ . In addition,  $xT_o(f)$  is interlaced by  $T_e(f)$ . Thus,

$$if deg(f) is \begin{cases} even & then T_{e}(f) \leq T_{o}(f) \\ odd & then T_{e}(f) \leq T_{o}(f). \end{cases}$$

We will derive this theorem from the following more general result. If f(x) is a polynomial and d is a positive integer, then we can uniquely write

$$f(\mathbf{x}) = f_0(\mathbf{x}^d) + \mathbf{x}f_1(\mathbf{x}^d) + \dots + \mathbf{x}^{d-1}f_{d-1}(\mathbf{x}^d)$$
(7.15.2)

**Theorem 7.65.** If  $f \in P^{\text{pos}}$ , d is a positive integer, and  $f_0, \ldots, f_{d-1}$  are given in (7.15.2) then  $f_0, f_1, \ldots, f_{d-1}$  is a mutually interlacing sequence. In particular, all  $f_i$  are in  $P^{\text{pos}}$ .

*Proof.* The idea is to use a certain matrix H such that f(H) contains all the  $f_i$ 's as entries. We then show that f(H) preserves mutually interlacing sequences. This yields  $f_i \in \mathbf{P}^{\text{pos}}$ , and the mutual interlacing of the sequence.

Assume that f is monic, and write

$$f(x) = \sum_{i=0}^n a_i x^i = \prod_{i=1}^n (x + r_i)$$

where all  $r_i$  and  $a_i$  are positive. Define the d by d matrix H as follows. The upper right corner is x, the lower diagonal is all 1's, and the remaining entries are zero. For instance, if d is three then H, H<sup>2</sup>,..., H<sup>6</sup> are

$$\begin{pmatrix} 0 & 0 & x \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & x & 0 \\ 0 & 0 & x \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} x & 0 & 0 \\ 0 & x & 0 \\ 0 & 0 & x \end{pmatrix}, \begin{pmatrix} x & 0 & 0 \\ 0 & 0 & x \end{pmatrix}, \begin{pmatrix} 0 & x^2 & 0 \\ 0 & 0 & x^2 \\ x & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & x^2 & 0 \\ 0 & 0 & x^2 \\ x & 0 & 0 \end{pmatrix}, \begin{pmatrix} x^2 & 0 & 0 \\ 0 & x^2 & 0 \\ 0 & 0 & x^2 \end{pmatrix},$$

If I is the identity matrix then

$$\prod_{i=1}^n (H+r_iI) = \sum_{i=0}^n \mathfrak{a}_i H^i$$

An important property of H is that  $H^d = xId$ , as we can see from the above matrices when d = 3. Upon consideration of the patterns in the powers of H we see that

$$\sum_{i=0}^{n} a_{i} H^{i} = \begin{pmatrix} f_{0} & xf_{d-1} & xf_{d-2} & \dots & xf_{1} \\ f_{1} & f_{0} & xf_{d-1} & \dots & xf_{2} \\ f_{2} & f_{1} & \ddots & & \vdots \\ \vdots & \vdots & & \ddots & xf_{d-1} \\ f_{d-1} & f_{d-2} & \dots & f_{1} & f_{0} \end{pmatrix}$$

where all terms are either of the form  $f_i(x)$  or  $xf_i(x)$ .

Since  $(\sum a_i H^i)(1, 0, ..., 0)^t = (f_0, f_1, ..., f_{d-1})$  it suffices to show that H+rI preserves mutually interlacing sequences. This follows from Example 3.74.

*Remark* 7.66. The even part of a polynomial in  $\mathbf{P}^{\text{pos}}$  has all real roots. Is every polynomial in  $\mathbf{P}^{\text{pos}}$  the even part of a polynomial in  $\mathbf{P}^{\text{pos}}$ ? We use Newton's inequalities to show that the answer is no.

Suppose that  $f = f_e(x^2) + x f_o(x^2) \in \mathbf{P}^{pos}(2n)$ , and let  $f_e = \sum a_i x^i$ . From (4.3.5) with k = 2 and r = 1 we see that

$$\frac{a_1^2}{a_0 a_2} \ge \frac{4 \cdot 3}{2 \cdot 1} \frac{(2n)(2n-1)}{(2n-2)(2n-3)} > 3\left(2 \cdot \frac{n}{n-1}\right)$$
(7.15.3)

Now Newton's inequality for  $f_e$  is

$$\frac{a_1^2}{a_0a_2} \ge 2\frac{n}{n-1}$$

Thus, if  $f_e$  is the even part of a polynomial in  $\mathbf{P}^{pos}$ , then it must satisfy (7.15.3) which is stronger than the usual Newton inequality. In particular,  $(x + 1)^n$  is not the even part of a polynomial in  $\mathbf{P}^{pos}$  for  $n \ge 2$ .

**Example 7.67.** If we begin with a polynomial in  $\mathbf{P}^{\text{pos}}$  (or a stable polynomial, see Chapter 21) then we can form a tree by recursively splitting into even and odd parts. For instance, if we begin with  $(x + 1)^8$  we get Figure 7.1. The  $2^k$  polynomials in the kth row are mutually interlacing (but not left to right).

There is an elementary reason why the even and odd parts never have a common factor.

**Lemma 7.68.** If  $f \in P^{pos}$  then  $f_e$  and  $f_o$  have no common factor.



Figure 7.1: The even-odd tree of  $(x + 1)^8$ 

*Proof.* Assume that they do. If  $f_e = (x + a)g(x)$  and  $f_o = (x + a)h(x)$  then

$$f(x) = f_e(x^2) + xf_o(x^2) = (x^2 + a)(g(x^2) + xh(x^2))$$

This is not possible, since  $f \in \mathbf{P}^{pos}$  implies that a is positive, and  $x^2 + a \notin \mathbf{P}$ .

# 7.16 Chebyshev and Jacobi Polynomials

The Chebyshev polynomials  $T_n$  are orthogonal polynomials given by the recurrence formula  $T_0 = 1$ ,  $T_1 = x$ , and  $T_{n+1} = 2xT_n - T_{n-1}$ . We study some Möbius transformations associated to the Chebyshev and Jacobi polynomials.

The transformations associated to the reversal of the Chebyshev and Legendre polynomials satisfy some simply stated identities. We can use these to establish mapping properties for the reversal of Chebyshev polynomials.

For our purposes we need an explicit formula:

$$T_{n}(x) = 2^{-n} \sum_{i=0}^{n} {2n \choose 2i} (x+1)^{i} (x-1)^{n-i}.$$
 (7.16.1)

**Lemma 7.69.** If T is the linear transformation  $x^n \mapsto T_n$  and  $Mz = \frac{z+1}{z-1}$  then  $T_M : \mathbf{P}^{(0,1)} \to \mathbf{P}^{pos}$ .

*Proof.* Using (7.16.1) we find a simple expression for  $T_M(x^n)$ :

$$\begin{split} \mathsf{T}_{\mathsf{M}} \left( x^{n} \right) &= (x-1)^{n} \mathsf{T}_{\mathsf{n}} \left( \frac{x+1}{x-1} \right) \\ &= (x-1)^{n} 2^{-n} \sum_{i=0}^{n} \binom{2n}{2i} \left( \frac{x+1}{x-1} + 1 \right)^{i} \left( \frac{x+1}{x-1} - 1 \right)^{n-i} \\ &= \sum_{i=0}^{n} \binom{2n}{2i} x^{i} \\ &= \text{ even part of } (1+x)^{2n} \end{split}$$

where the even part is defined in §7.7.15. Consequently

 $T_M (f) = \text{ even part of } f((1+x)^2).$ 

If  $f \in \mathbf{P}^{(0,1)}$  then  $f((1+x)^2)$  is in  $\mathbf{P}^{(-2,0)}$  and by Theorem 7.64 the even part is in  $\mathbf{P}^{p\,o\,s}$  as well. Thus  $T_M$  maps  $\mathbf{P}^{(0,1)}$  to  $\mathbf{P}^{p\,o\,s}$ .

Formulas analogous to (7.16.1) hold for all Jacobi polynomials. From [168, (4.3.2)]

$$P_{n}^{\alpha,\beta}(x) = \sum_{k=0}^{n} \binom{n+\alpha}{n-k} \binom{n+\beta}{k} \left(\frac{x-1}{2}\right)^{k} \left(\frac{x+1}{2}\right)^{n-k}$$

If we apply the Möbius transformation M of Lemma 7.69 to the linear transformation  $J:x^n\mapsto P_n^{\alpha,\beta}$  we get

$$J_{M}(x^{n}) = \sum_{k=0}^{n} \binom{n+\alpha}{n-k} \binom{n+\beta}{k} x^{k}$$
(7.16.2)

In particular, the Legendre polynomials are  $P_n^{0,0}$  so defining  $L(x^n) = P_n^{0,0}$  we have the elegant formula

$$L_{M}(x^{n}) = \sum_{k=0}^{n} {\binom{n}{k}}^{2} x^{k}.$$
(7.16.3)

It follows from Corollary 13.17 that

**Lemma 7.70.** The map  $x^n \mapsto L_M(x^n)$  given in (7.16.3) maps  $P^{\text{alt}}$  to itself.

We now state two identities for reverse polynomials, where  $T_n$  and  $P_n$  are the Chebyshev and Legendre polynomials respectively.

$$T(x^{n}) = (T_{n})^{re\nu} \quad T(1-x)^{n} = \begin{cases} 0 & n \text{ odd} \\ (1-x^{2})^{n/2} & n \text{ even} \end{cases}$$
(7.16.4)

$$S(x^{n}) = (P_{n})^{re\nu} \quad S(1-x)^{n} = \begin{cases} 0 & n \text{ odd} \\ \binom{n-1}{n/2} 2^{1-n} (1-x^{2})^{n/2} & n \text{ even} \end{cases}$$
(7.16.5)

**Lemma 7.71.** If  $\mathsf{T}$  is given above, then  $\mathsf{T}: P^{\mathsf{alt}} \cup P^{\mathsf{pos}} \longrightarrow P$ .

*Proof.* The proof relies on the commutative diagram that holds for  $(x - 1)^n$ which is a consequence of the identity (7.16.4), and by linearity it holds for all polynomials. The map even returns the even part of f.



Since we know that the even part determines a map from **P**<sup>pos</sup> to itself, the claim in the top line of the following diagram follows by commutativity.



#### 7.17 Commuting diagrams of spaces

The earlier commuting diagrams of transformations in §6.6.8 give rise to transformations between spaces of polynomials. These are useful to us because we generally understand three of the four transformations, and so can deduce information about the fourth.

**Example 7.72.** T:  $x^n \mapsto H_n$  satisfies (Corollary 7.44)



**Example 7.73.** T:  $x^k \mapsto H_k(x)H_{n-k}(x)$  satisfies (Corollary 9.53)


**Example 7.74.** T:  $x^k \mapsto L_k(x)L_{n-k}(x)$  satisfies (Corollary 9.53) a similar diagram as the Hermite transformation above, but the image is smaller.



**Example 7.75.** For the transformation  $T: x^k \mapsto \langle \underline{x} \rangle_k (\underline{x} - \underline{\alpha})_{n-k}$  the following diagram commutes at the function level, but we are not able to prove either the top or the bottom. If one were true, then the other follows from the commutativity.



**Example 7.76.** The transformation T:  $x^i \mapsto (\underline{x+d-i})_d$  actually maps  $\mathbf{P}^{alt} \longrightarrow \mathbf{P}^{sep}$  (Lemma 8.30).



**Example 7.78.** T:  $x^k \mapsto H_k(x)x^{n-k}$  acts (Lemma 9.60) on polynomials of degree n:



## 7.18 The Pincherlet derivative

If we are given a linear transformation T on polynomials we can form the *Pincherlet* derivative

$$T'(f) = T(xf) - xT(f) = [M, T](f)$$

where M(f) = xf, and [M, T] is the commutator TM - MT. In some circumstances T' will map **P** to itself.

**Lemma 7.79.** If  $f \leq g$ , f has leading coefficient a, g has leading coefficient b, then  $f - xg \in P$  if  $a \geq b$ .

*Proof.* Immediate from Lemma 1.20.

**Corollary 7.80.** Suppose  $T: P \longrightarrow P$  maps polynomials with positive leading coefficients to polynomials with positive leading coefficients. If the sequence of leading coefficients of  $T(x^i)$  is strictly increasing then  $T': P \longrightarrow P$ .

*Proof.* Choose monic f. Then since  $xf \leq f$  and since T preserves interlacing, T(xf) < T(f). By hypothesis T(xf) and T(f) meet the conditions of the lemma, so  $T(xf) - xT(f) = T'(f) \in \mathbf{P}$ .

**Corollary 7.81.** The linear transformation  $x^n \mapsto H_{n+1} - xH_n$  maps  $P \longrightarrow P$ .

*Proof.* The leading coefficient of  $H_n$  is  $2^n$ .

If T maps **P**<sup>pos</sup> to itself, then we have weaker assumptions.

**Lemma 7.82.** Suppose  $T: \mathbf{P}^{pos} \longrightarrow \mathbf{P}^{pos}$  maps polynomials with positive leading coefficients to polynomials with positive leading coefficients, and preserves degree. Then,  $T': \mathbf{P}^{pos} \longrightarrow \mathbf{P}^{pos}$ .

*Proof.* Choose  $f \in \mathbf{P}^{pos}$ . Then  $T(xf) \leq Tf$  and since  $T(xf) \in \overline{\mathbf{P}^{pos}}$  it follows that  $xT(f) \leq T(xf)$  and hence  $T(xf) - xT(f) \in \mathbf{P}^{pos}$ .

**Corollary 7.83.** The linear transformation  $x^n \mapsto n \langle \underline{x} \rangle_n$  maps  $P^{\text{pos}}$  to itself.

*Proof.* If  $T(x^n) = \langle \underline{x} \rangle_n$  then  $T'(x^n) = \langle \underline{x} \rangle_{n+1} - x \langle \underline{x} \rangle_n = n \langle \underline{x} \rangle_n$ . Alternatively, this is the composition

$$x^{n} \xrightarrow{\text{differentiation}} nx^{n-1} \xrightarrow{\text{mult. by } x} nx^{n} \xrightarrow{x^{n} \mapsto \langle \underline{x} \rangle_{n}} n\langle \underline{x} \rangle_{n}$$

## 7.19 Hypergeometric Polynomials

Hypergeometric series are a natural class of functions that are sometimes polynomials. We show that for certain values of the parameters these polynomials are in **P**. We begin with the confluent hypergeometric function  $_1F_1$  which is defined by

$${}_{1}F_{1}(a,b;z) = 1 + \frac{a}{b}\frac{z}{1!} + \frac{a(a+1)}{b(b+1)}\frac{z^{2}}{2!} + \cdots = \sum_{i=0}^{\infty}\frac{\langle \underline{a} \rangle_{i}}{\langle \underline{b} \rangle_{i}}\frac{z^{i}}{i!}$$

We are interested in the case that a is a negative integer, and b is a positive integer. Since  $\langle \underline{a} \rangle_i$  is zero if a is a negative integer and i > |a|, it follows

that  $_1F_1(a,b;z)$  is a polynomial in these cases, of degree |a|. For example,  $_1F_1(-4,b;z)$  is equal to

$$1 - \frac{4z}{b} + \frac{6z^2}{b(1+b)} - \frac{4z^3}{b(1+b)(2+b)} + \frac{z^4}{b(1+b)(2+b)(3+b)}$$

and we can manipulate this into a more familiar form

$$\frac{1}{(\underline{b+4})_4} \left( (\underline{b+4})_4 - 4z(\underline{b+4})_3 + 6z^2(\underline{b+4})_2 - 4z^3(\underline{b+4})_1 + z^4(\underline{b+4})_0 \right)$$

In general when d a positive integer we can write

$$_{1}F_{1}(-d,b;-z) = \frac{1}{(b+d)_{d}} \sum_{i=0}^{a} \left(\frac{b+d}{i}\right)_{i} {d \choose i} z^{i}$$

Applying Lemma 7.2 twice to this representation shows that

Lemma 7.84. If a is a positive integer, and b is positive then

$$_{1}F_{1}(-a,b;-z) \in P^{pos}$$

There are many relationships involving hypergeometric functions - the one below shows that the above result also follows from the fact that the generalized Laguerre polynomial  $L_n^{\lambda}$  has all real roots for  $\lambda > -1$ .

$$L_{n}^{\lambda}(z) = \frac{\langle \underline{1+\lambda} \rangle_{n}}{\Gamma(1+\nu)} \, _{1}F_{1}(-n,1+\lambda,z)$$

<sub>1</sub>F<sub>1</sub> satisfies recurrences in each parameter:

$${}_{1}F_{1}(a, b, z) = \frac{2+2 a - b + z}{1+a-b} {}_{1}F_{1}(1+a, b, z) - \frac{a+1}{1+a-b} {}_{1}F_{1}(2+a, b, z)$$
$${}_{1}F_{1}(a, b; z) = \frac{b+z}{b} {}_{1}F_{1}(a, b+1; z) + \frac{(a-b-1)z}{b(b+1)} {}_{1}F_{1}(a, b+2; z)$$

These recurrences can be used to establish this interlacing square for positive integral a, and positive b:

$$\begin{array}{ccc} {}_{1}F_{1}(-(a+1),b;z) & \stackrel{<}{\scriptstyle{\scriptstyle{\frown}}} & {}_{1}F_{1}(-a,b;y) \\ & & & & & \\ & & & & \\ \downarrow \ll & & & & \\ {}_{1}F_{1}(-(a+1),b+1;z) & \stackrel{<}{\scriptstyle{\leftarrow}} & {}_{1}F_{1}(-a,b+1;y) \end{array}$$

If the hypergeometric series has only one factor in the numerator then we can apply the same argument. For instance

$$_{1}F_{2}(a;b,c;z) = \sum_{i=0}^{\infty} \frac{\langle \underline{a} \rangle_{i}}{\langle \underline{b} \rangle_{i} \langle \underline{c} \rangle_{i}} \frac{z^{i}}{i!}$$

and if we take d to be a positive integer then

$${}_{1}F_{2}(-d; b, c; -z) = \frac{1}{\langle \underline{b} + d \rangle_{d} \langle \underline{c} + d \rangle_{d}} \sum_{i=0}^{d} {\binom{d}{i}} (\underline{b} + d)_{i} (\underline{c} + d)_{i} z^{i}$$

and we can again apply Lemma 7.2 to yield that  ${}_{1}F_{2}(-d; b, c; z) \in \mathbf{P}^{alt}$ .

#### 7.20 Eigenpolynomials of linear transformations

If  $T: \mathbf{P} \longrightarrow \mathbf{P}$  is a linear transformation then we can look for eigenpolynomials of T. These are polynomials *e* for which there is a  $\lambda \in \mathbb{R}$  so that  $Te = \lambda e$ . We consider several questions: does T have eigenpolynomials? If so, are they in **P**? Do successive eigenpolynomials interlace? What are explicit examples?

**Lemma 7.85.** Suppose  $T: P \longrightarrow P$  maps polynomials of degree n to polynomials of degree n. Let  $c_n$  be the leading coefficient of  $T(x^n)$ .

- 1. If all  $c_n$  are distinct then T has a unique eigenpolynomial of degree n with eigenvalue  $c_n$ .
- 2. If  $|c_n| > |c_r|$  for  $0 \le r < n$  then T has an eigenpolynomial of degree n that is in **P**.

*Proof.* If we let V be the vector space of all polynomials of degree at most n then the assumption on T implies that T maps V to itself. In addition, the matrix representation M of T is upper triangular. The r-th diagonal element of M is the coefficient  $c_r$  of  $x^r$  in  $T(x^r)$ . If they are all distinct then T has an eigenvector corresponding to each eigenvalue.

We now use the power method to find the eigenvalue. Choose any initial value  $v_0$  and define

$$v_{k+1} = \frac{1}{|v_k|} M v_k$$

where  $|v_k|$  is any vector norm. Since M has a dominant eigenvalue, the  $v_k$  converge to an eigenvector v of M, provided that  $v_0$  is not orthogonal to v.

Recasting this in terms of polynomials, we choose an initial polynomial  $p_0$  in  $\mathbf{P}(n)$ . Since T maps  $\mathbf{P}$  to itself, all the  $p_k$  are in  $\mathbf{P}$ . If this happens to converge to zero, then  $p_0$  was orthogonal to p. Simply perturb  $p_0$ , and we get an eigenpolynomial in  $\mathbf{P}$ .

The sequence of possible eigenvalues is quite restricted.

**Corollary 7.86.** Suppose that  $T: P \longrightarrow P$  preserves degree and has positive eigenvalues  $\lambda_n$  for n = 0, 1, ... Then

$$\sum_{n=0}^{\infty}\frac{\lambda_n}{n!}\mathbf{x}^n\in\widehat{\boldsymbol{P}}.$$

*Proof.* The  $\lambda_n$  are the leading coefficients of  $T(x^n)$ . Apply Lemma 15.33.

**Lemma 7.87.** Suppose that T is a linear transformation  $P \longrightarrow P$ ,  $p_n$  and  $p_{n+1}$  are eigenpolynomials of degree n and n + 1 with eigenvalues  $\lambda_n$  and  $\lambda_{n+1}$ . If  $|\lambda_{n+1}| > |\lambda_n|$  and the roots of  $p_n$  are distinct then  $p_{n+1} \leq p_n$ .

*Proof.* We show that  $g(x) = p_n + \alpha p_{n+1} \in \mathbf{P}$  for any  $\alpha$ . If we define

$$g_{k}(\mathbf{x}) = \lambda_{n}^{-k} + \alpha \lambda_{n+1}^{-k} p_{n+1}$$

then  $T^kg_k = g$  so it suffices to show that  $g_k \in \mathbf{P}$ . Now

$$g_{k}(x) = \lambda_{n}^{-k} \left( p_{n} + (\lambda_{n}/\lambda_{n+1})^{k} p_{n+1} \right)$$

Since  $|\lambda_n/\lambda_{n+1}| < 1$  we can apply Lemma 1.11 for k sufficiently large since the roots of  $p_n$  are distinct. It follows that  $g_k \in \mathbf{P}$ , and hence  $g \in \mathbf{P}$ .

*Remark* 7.88. T(f) = f - f'' is a simple example of a linear transformation mapping  $\mathbf{P} \longrightarrow \mathbf{P}$  that has only two eigenpolynomials. If  $T(f) = \lambda f$  then clearly  $\lambda = 1$  and f'' = 0, so f has degree 0 or 1. The matrix representing T has all 1's on the diagonal, so the lemma doesn't apply.

We next show that multiplier transformations generally have no interesting eigenpolynomials, and then we determine the eigenpolynomials for the Hermite transformation.

**Lemma 7.89.** If  $T(x^n) = t_n x^n$  where all the  $t_i$  are distinct and non-zero, then the only eigenpolynomials are multiples of  $x^i$ .

*Proof.* If  $Te = \lambda e$  where  $e = \sum a_i x^i$  then  $a_i = \lambda t_i a_i$ . Every non-zero  $a_i$  uniquely determines  $\lambda = 1/t_i$ , so there is exactly one non-zero  $a_i$ .

Some simple linear transformations that map **P** to itself have no eigenpolynomials. For instance, consider  $Tf = f + \alpha f'$ . If  $f + \alpha f' = \lambda f$  then since the degree of f' is less than the degree of f we see  $\lambda = 1$ , and hence  $\alpha = 0$ .

The eigenpolynomials of the Hermite transformation  $T(x^n) = H_n$  are given by a composition, and are also orthogonal polynomials.

**Lemma 7.90.** If  $T(x^n) = H_n$  then the eigenpolynomials of T are given by the composition

$$g_n = (3x - 2D)^n (1)$$

and the corresponding eigenvalue is  $2^{n}$ .

*Proof.* Since T(f) = f(2x - D)(1) we can apply Lemma 7.43 to conclude that

$$\Gamma(g_n) = g_n (2x - D)(1)$$
$$= (6x - 4D)^n (1)$$
$$= 2^n g_n$$

It isn't known how to determine if a linear transformation T has all its eigenpolynomials in **P**. If T eventually maps into **P** then all the eigenpolynomials are in **P**.

**Lemma 7.91.** If T is a linear transformation such that for any polynomial f there is an integer n such that  $T^n f \in P$ , then all eigenpolynomials corresponding to non-zero eigenvalues are in P.

*Proof.* If  $Tf = \lambda f$  then choose n as in the hypothesis and note that  $T^n f = \lambda^n f$ . If  $\lambda$  is non-zero it follows that  $f \in \mathbf{P}$ .

## 7.21 Classical multiple orthogonal polynomials

It is possible to construct polynomials that are orthogonal to several weight functions [170]. The resulting polynomials are called *multiple orthogonal polynomials* and all have all real roots. In this section we look at the corresponding differential operators, and also observe some interlacing properties.

#### Lemma 7.92.

- 1. Suppose  $\alpha > -1$ . The operator  $T_{n,\alpha} : f \mapsto x^{-\alpha} D^n x^{n+\alpha} f$  maps  $P \longrightarrow P$  and  $P^{pos}$  to  $P^{pos}$ .
- 2. If  $f \in \mathbf{P}^{\pm}$  then  $T_{n+1,\alpha}f \leq T_{n,\alpha}f$ .
- 3. Suppose  $b \ge 0$ . The operator  $S_{n,b,c} : f \mapsto e^{bx^2 cx} D^n e^{-bx^2 + cx} f$  maps  $P \longrightarrow P$ .

*Proof.* The first one follows from a more general result (Proposition 11.140) in several variables. To check interlacing we note that

$$\begin{split} \beta T_{n,\alpha} f + T_{n+1,\alpha} f &= x^{-\alpha} \mathsf{D}^n \left( \beta x^{\alpha+n} f + \mathsf{D} x^{n+1+\alpha} f \right) \\ &= x^{-\alpha} \mathsf{D}^n x^{n+\alpha} \left( (\beta+n+\alpha+1) f + x f' \right) \end{split}$$

Since  $f\in I\!\!P^\pm$  we know that  $h(x)=(\beta+n+\alpha+1)f+xf'\in I\!\!P$  which implies that

$$\beta T_{n,\alpha}f + T_{n+1,\alpha}f = x^{-\alpha}D^n x^{n+\alpha}h$$

where  $h \in P$ . By the first part this is in P for all  $\beta$  and so the interlacing follows.

For the Hermite case it suffices to show that  $e^{bx^2-cx}D(e^{-bx^2+cx}f)$  is in **P** if f is in **P**. Note that we know that  $D(e^{-bx^2+cx}f)$  is in  $\widehat{\mathbf{P}}$  and has the form  $e^{-bx^2+cx}g$ , but we can't multiply by  $e^{bx^2-cx}$  since  $e^{bx^2}$  is not in  $\widehat{\mathbf{P}}$ . We just directly compute

$$e^{bx^2-cx}D(e^{-bx^2+cx}f) = (f'-(2bx-c)f)$$

and f' - (2bx - c)f is in **P** by Lemma 1.20.

If f = 1 in (1) then we have Laguerre polynomials, and if f = 1, c = 0, b = 1 in (2) we have Hermite polynomials.

If we iterate these operators we get, up to a constant factor, the classical multiple orthogonal polynomials of [170]. Let  $n = (n_1, ..., n_r)$  be positive integers, and  $\alpha = (\alpha_0, ..., \alpha_r)$  where  $\alpha_1, ..., \alpha_r$  are all at least -1. The following polynomials are all in **P**. The interlacing property above shows that the Jacob-Piñeiro polynomials  $P_n^{\alpha}$  and  $P_m^{\alpha}$  interlace if n and m are equal in all coordinates, and differ by one in the remaining coordinate.

Jacobi-Piñeiro 
$$(1-x)^{-\alpha} \prod_{i=1}^{r} \left[ x^{-\alpha_i} \frac{d^{n_i}}{dx^{n_i}} x^{n_i+\alpha_i} \right] (1-x)^{\alpha_0+n_1+\dots+n_r}$$
  
Laguerre-I  $e^x \prod_{i=1}^{r} \left[ x^{-\alpha_i} \frac{d^{n_i}}{dx^{n_i}} x^{n_i+\alpha_i} \right] e^{-x}$   
Laguerre-II  $x^{-\alpha_0} \prod_{i=1}^{r} \left[ e^{\alpha_i x} \frac{d^{n_i}}{dx^{n_i}} e^{-\alpha_i x} \right] x^{\alpha_0+n_1+\dots+n_r}$   
Hermite  $\prod_{i=1}^{r} \left( e^{bx^2-cx} \mathsf{D}_i^n e^{-bx^2+cx} \right) 1$ 

#### 7.22 Laurent Orthogonal polynomials

Laurent orthogonal polynomials satisfy the recurrence

$$\begin{split} p_{-1} &= 0 \\ p_0 &= 1 \\ p_n &= (a_n x + b_n) p_{n-1} - c_n x \, p_{n-2} \qquad \text{for } n \geqslant 1 \end{split}$$

where all  $a_n$ ,  $b_n$ ,  $c_n$  are positive. There are three simple recurrences that appear to have many mapping properties (see Appendix 24.9). We are able to prove some mapping properties for two of them. Notice that the recurrence implies that the coefficients are alternating, and hence if the roots are all real they are all positive.

**Lemma 7.93.** If  $p_n$  is the sequence of Laurent orthogonal polynomials arising from the recurrence  $p_{n+1} = x p_n - n x p_{n-1}$  then the transformation  $T: x^n \mapsto p_n$  satisfies  $T: P^{alt} \longrightarrow P^{alt}$  and preserves interlacing.

The transformation  $T: x^n \mapsto p_n^{rev} maps P^{alt}$  to  $P^{alt}$ .

Proof. Using linearity

$$\begin{split} T(x \cdot x^n) &= xp_n - nxp_{n-1} \\ &= x \, T(x^n) - x T(x^{n \prime}) \\ T(xf) &= x \, T(f - f'). \end{split}$$

We now prove the result by induction - the base case is easy and omitted. Since  $f - f' \leq f$  it follows by induction that  $T(f - f') \leq T(f)$ . Since the roots are positive by induction, we know that

$$\mathsf{T}(\mathsf{x}\mathsf{f}) = \mathsf{x}\mathsf{T}(\mathsf{f} - \mathsf{f}') \leq \mathsf{T}(\mathsf{f})$$

and the conclusion follows from Corollary 1.46.

For the second part, we have the recurrence T(xf) = T(f) - xT(f'). We can't apply Corollary 1.46 since T(f) does not have positive leading coefficients. However, if we let S(f) = T(f)(-x) then we can apply the lemma to S. Since S satisfies the recurrence

$$S(xf) = S(f) + x S(f')$$

we see that by induction

$$S(xf) = S(f) + x S(f) \leq S(f)$$

and so  $S: \mathbf{P}^{alt} \longrightarrow \mathbf{P}^{pos}$ .

**Lemma 7.94.** If  $p_n$  is the sequence of Laurent orthogonal polynomials satisfying the recurrence  $p_{n+1} = (x + n) p_n - n x p_{n-1}$  then the transformation  $T: x^n \mapsto p_n$  satisfies  $T: P^{alt} \longrightarrow P^{alt}$  and preserves interlacing.

*Proof.* The proof is similar to the previous one. We check that

$$T(xf) = xT(f + xf' - f')$$

Now we know that

$$f \leq f + xf' \leq f'$$
  
so  $f + xf' - f' \leq f$ 

By induction

$$T(xf) = xT(f + xf' - f') \leq T(f)$$

and we apply Corollary 1.46 again to complete the proof.

#### 7.23 Biorthogonal polynomials

In this section I would like to sketch a completely different approach to the problem of showing that a linear transformation preserves real roots. The main results are by Iserles and Norsett, and can be found in [90]. The idea is to begin with a parametrized measure, and to use it to associate to a polynomial a *biorthogonal* polynomial that is determined by the roots of the polynomial. There are a number of non-degeneracy conditions that must be met. In general the resulting transformation is highly non-linear, but there are quite a number of measures for which we get linearity.

Choose a function  $\omega(x, \alpha)$  and a measure dµ, and define a linear functional

$$\mathcal{L}_{\alpha}(f) = \int f(x) \omega(x, \alpha) \, d\mu$$

We start with a monic polynomial g(x) with all real roots  $r_1, \ldots, r_n$ . Each root determines a linear functional  $\mathcal{L}_{r_i}$ . We next try to construct a monic polynomial p(x) of degree n that satisfies

$$\mathcal{L}_{r_1}(p) = 0, \quad \mathcal{L}_{r_2}(p) = 0, \quad \cdots \quad \mathcal{L}_{r_n}(p) = 0$$

This is a set of linear equations. There are n unknown coefficients, and n conditions, so as long as the determinant of the system is non-zero we can uniquely find p, which is called a *biorthogonal* polynomial.

Next, we want to show that p has all real roots. The proof of this is similar to the proof that a single orthogonal polynomial has all real roots, and requires that the  $w(x, \alpha)$  satisfy an interpolation condition.

Finally, we define the transformation T(g) = p. It is quite surprising that there are *any* choices that make T linear. The coefficients of p are symmetric functions of the roots, and so can be expressed in terms of the coefficients of g, but in general there is no reason to suppose that there is linearity.

CHAPTER



# Affine Transformations of Polynomials

In this chapter we investigate properties of polynomials associated to affine transformations. Recall that an affine transformation  $\mathbb{A}$  satisfies  $\mathbb{A}(x) = qx+b$ , and the action on polynomials is  $\mathbb{A}(f(x)) = f(qx + b)$ . In order not to burden the reader (and the writer) with lots of similar but different cases, we will generally confine ourselves to the affine transformations of the form  $\mathbb{A}(x) = qx+b$  where  $q \ge 1$ . We are interested in those polynomials that interlace their affine transformations. In particular we will study the set of polynomials

$$\mathbf{P}^{\mathbb{A}} = \{ \mathbf{f} \in \mathbf{P} \mid \mathbf{f} \underline{\ll} \mathbb{A}\mathbf{f} \}.$$

#### 8.1 Introduction

We begin with general properties of these affine transformations for q > 1. A acts on the real line, and preserves order. Since q > 1 we see that A has a unique fixed point  $\frac{b}{1-q}$ . Consequently A is a bijection restricted to the two intervals

$$\mathbb{A}^- = (-\infty, \frac{b}{1-q}) \qquad \qquad \mathbb{A}^+ = (\frac{b}{1-q}, \infty)$$

Since  $\mathbb{A}$  and  $\mathbb{A}^{-1}$  preserve order, it is easy to see that  $\mathbb{A}$  is increasing on  $\mathbb{A}^+$  and it is decreasing on  $\mathbb{A}^-$ . More precisely,

$$\alpha < \mathbb{A}\alpha$$
 iff  $\alpha \in (\frac{b}{1-q}, \infty)$   
 $\mathbb{A}\alpha < \alpha$  iff  $\alpha \in (-\infty, \frac{b}{1-q})$ 

The action of  $\mathbb{A}$  on the roots of a polynomial is easy to determine. If  $f(\alpha) = 0$  then  $(\mathbb{A}f)(\mathbb{A}^{-1}\alpha) = f(\alpha) = 0$  so that the roots of  $\mathbb{A}f$  are  $\mathbb{A}^{-1}$  applied to the roots of f. Since  $\mathbb{A}^{-1}$  preserves order it follows that  $f \longleftarrow g$  implies  $\mathbb{A}f \longleftarrow \mathbb{A}g$ .

If  $f \leq A f$  and  $\alpha$  is a root of f then we must have that  $\mathbb{A}^{-1}\alpha \leq \alpha$ , and hence  $\alpha \leq \mathbb{A}\alpha$ . We conclude that if  $f \in \mathbf{P}^{\mathbb{A}}$  then the roots of f must lie in  $(\frac{b}{1-q}, \infty)$ . For the three examples considered above we have

$\mathbb{A}(\mathbf{x}) = 2\mathbf{x}$	$f\in\mathbf{P}^{\mathbb{A}}$	$\implies$	roots of $f \in (0, \infty)$
$\mathbb{A}(x) = 2x + 1$	$f\in \mathbf{P}^{\mathbb{A}}$	$\implies$	roots of $f \in (-1, \infty)$
$\mathbb{A}(\mathbf{x}) = \mathbf{x} + 1$	$f\in\mathbf{P}^{\mathbb{A}}$	$\implies$	roots of $f \in (-\infty, \infty)$

Before we proceed, we should show that  $\mathbf{P}^{\mathbb{A}}$  is non-empty. Define

$$\langle \underline{\mathbf{x}; \mathbf{a}} \rangle_{\mathbf{n}}^{\mathbb{A}} == \prod_{i=0}^{\mathbf{n}-1} (\mathbb{A}^{i} \mathbf{x} - \mathbf{a})$$
 (8.1.1)

In case that  $\mathbb{A}(x) = x + 1$  we have

$$\langle \underline{\mathbf{x}}; \underline{\mathbf{0}} \rangle_{\mathbf{n}}^{\mathbb{A}} = (\mathbf{x})(\mathbf{x}+1)(\mathbf{x}+2)\cdots(\mathbf{x}+\mathbf{n}-1) = \langle \underline{\mathbf{x}} \rangle_{\mathbf{n}},$$
 (8.1.2)

if  $\mathbb{A}(x) = qx$  then

$$\left\langle \underline{x;-1} \right\rangle_{n}^{\mathbb{A}} = (1+x)(1+qx)\cdots(1+q^{n-1}x),$$
 (8.1.3)

and again if  $\mathbb{A}(x) = qx$  then

$$\langle \underline{\mathbf{x}}; \underline{\mathbf{0}} \rangle_{\mathbf{n}}^{\mathbb{A}} = q^{\binom{n}{2}} \mathbf{x}^{\mathbf{n}}.$$
 (8.1.4)

Since  $\mathbb{A}^{-1}\alpha < \alpha$  for  $\alpha \in \mathbb{A}^+$ , if  $a \in \mathbb{A}^+$  then the roots of  $\langle \underline{x}; \underline{a} \rangle_n^{\mathbb{A}}$  in increasing order are

$$\mathbb{A}^{-n+1}(\mathfrak{a}), \ldots, \mathbb{A}^{-2}(\mathfrak{a}), \mathbb{A}^{-1}(\mathfrak{a}), \mathfrak{a}$$

and the roots of  $\mathbb{A}\langle \underline{x}; \mathfrak{a} \rangle_n^{\mathbb{A}}$  are

$$\mathbb{A}^{-n}(\mathfrak{a}), \ldots, \mathbb{A}^{-3}(\mathfrak{a}), \mathbb{A}^{-2}(\mathfrak{a}), \mathbb{A}^{-1}(\mathfrak{a})$$

Consequently if  $a \in \mathbb{A}^+$  then  $\langle \underline{x}; a \rangle_n^{\mathbb{A}} \underline{\ll} \mathbb{A} \langle \underline{x}; a \rangle_n^{\mathbb{A}}$ .

**Lemma 8.1.** If  $f \in \mathbf{P}^{\mathbb{A}}$  then  $\mathbb{A}f \in \mathbf{P}^{\mathbb{A}}$ .

*Proof.* If  $f \in \mathbf{P}^{\mathbb{A}}$  then  $f \underline{\ll} \mathbb{A}f$ . Since  $\mathbb{A}$  preserves interlacing  $\mathbb{A}f \underline{\ll} \mathbb{A}^2f$  and hence  $\mathbb{A}f \in \mathbf{P}^{\mathbb{A}}$ .

Polynomials in  $\mathbf{P}^{\mathbb{A}}$  satisfy a hereditary property.

**Lemma 8.2.**  $P^{\mathbb{A}}$  is an order ideal. That is, if  $f \in P^{\mathbb{A}}$  and g divides f then  $g \in P^{\mathbb{A}}$ .

*Proof.* It suffices to show that if we remove one linear factor from f the resulting polynomial g is still in  $\mathbf{P}^{\mathbb{A}}$ . Since affine transformations preserve order, the effect of removing a root from f is to also remove the corresponding root in  $\mathbb{A}f$ . The remaining roots interlace, and so  $g \in \mathbf{P}^{\mathbb{A}}$ .

The next lemma gives a useful criterion for determining if a polynomial is in  $\mathbf{P}^{\mathbb{A}}$ .

Lemma 8.3. The following are equivalent

- 1.  $f \in \boldsymbol{P}^{\mathbb{A}}$
- 2. There is a g such that  $f \leftarrow g$  and  $Af \leftarrow g$ .
- *3. There is an* h *such that* f  $\leftarrow$  h *and* f  $\leftarrow$  Ah.

*Proof.* If (1) holds then  $f \leq Af$ . Choosing g = Af shows that (1) implies (2), and taking h = f shows that (1) implies (3). If (2) holds and  $f(x) = \prod (x - r_i)$  then  $A^{-1}r_i \leq r_i$ , so (1) follows from Lemma 1.4. If (3) holds then we know that  $f \leftarrow h$  and  $A^{-1}f \leftarrow h$ , so (2) implies that  $A^{-1}f \leq f$ . Applying A to each side shows that (1) holds.

## 8.2 The affine derivative

We define the affine derivative  $D_{\mathbb{A}}f$  of f as follows:

**Definition 8.4.** If f is a polynomial then  $D_{\mathbb{A}}f = \frac{\mathbb{A}f - f}{\mathbb{A}x - x}$ .

In case  $\mathbb{A}f = f(x+1)$  the affine derivative is the difference operator  $\Delta(f) = f(x+1) - f(x)$ . If  $\mathbb{A}f = f(qx)$  then  $D_{\mathbb{A}}f$  is the q-derivative:

$$\mathsf{D}_{q}(f) = \begin{cases} \frac{f(qx) - f(x)}{qx - x} & \text{if } x \neq 0\\ f'(x) & \text{if } x = 0 \end{cases}$$
(8.2.1)

The affine derivative is well defined since  $q \neq 1$  implies  $\mathbb{A}(x) \neq x$ . It is immediate from the Taylor series that  $D_{\mathbb{A}}$  is continuous. If we let  $\mathbb{A}$  approach the identity, then  $D_{\mathbb{A}}$  converges to the derivative. The degree of  $D_{\mathbb{A}}f$  is one less than the degree of f. As is to be expected for a derivative,  $D_{\mathbb{A}}x = 1$ , and there is a product rule

$$D_{\mathbb{A}}(fg) = \frac{\mathbb{A}f \cdot \mathbb{A}g - fg}{\mathbb{A}x - x} = \frac{\mathbb{A}f \cdot \mathbb{A}g - \mathbb{A}f \cdot g + \mathbb{A}f \cdot g - fg}{\mathbb{A}x - x} = \mathbb{A}f \cdot D_{\mathbb{A}}g + g D_{\mathbb{A}}f \quad (8.2.2)$$

In particular we have

$$\mathsf{D}_{\mathbb{A}}(\mathsf{x}\mathsf{f}) = \mathbb{A}\mathsf{f} + \mathsf{x}\;\mathsf{D}_{\mathbb{A}}\mathsf{f} = \mathsf{f} + (\mathbb{A}\mathsf{x})\;\mathsf{D}_{\mathbb{A}}\mathsf{f} \tag{8.2.3}$$

The formula for  $D_{\mathbb{A}}$  on the product of three terms is

$$\mathsf{D}_{\mathbb{A}}(f_1f_2f_3) = \ \mathsf{D}_{\mathbb{A}}f_1 \cdot (f_2f_3) + (\mathbb{A}f_1) \cdot \ \mathsf{D}_{\mathbb{A}}f_2 \cdot (f_3) + \mathbb{A}(f_1f_2) \cdot \ \mathsf{D}_{\mathbb{A}}f_3$$

and the general formula is

$$\mathsf{D}_{\mathbb{A}}(\mathsf{f}_{1}\cdots\mathsf{f}_{n}) = \sum_{k=1}^{n} \mathbb{A}(\mathsf{f}_{1}\cdots\mathsf{f}_{k-1}) \cdot \mathsf{D}_{\mathbb{A}}\mathsf{f}_{k} \cdot (\mathsf{f}_{k+1}\cdots\mathsf{f}_{n}) \tag{8.2.4}$$

The next lemma is the heart of the results about the affine derivative.

**Lemma 8.5.** If  $f \in \mathbf{P}^{\mathbb{A}}$  and  $\alpha$  is positive then  $f - \alpha \mathbb{A} f \in \mathbf{P}^{\mathbb{A}}$ .

*Proof.* We may suppose that f is monic of degree n, in which case the leading coefficient of Af is  $q^n$ . Since  $f \leq Af$  we may write  $Af = q^n f + h$  where  $f \leq h$  so that

$$f - \alpha \mathbb{A}f = (1 - \alpha q^n)f - \alpha h$$

Writing  $f = q^{-n} \mathbb{A} f - k$  where  $\mathbb{A} f \leq k$  yields

$$f - \alpha \mathbb{A}f = (q^{-n} - \alpha)\mathbb{A}f - k$$

There are two cases, depending in the size of  $\alpha$ .

**Case 1.** If  $0 < \alpha < q^{-n}$  then  $(1 - \alpha q^{-n})$  and  $(q^{-n} - \alpha)$  are positive, so by Corollary 1.30 we have

$$f - \alpha \mathbb{A}f = (1 - \alpha q^{-n})f - \alpha h \longleftarrow f$$
$$f - \alpha \mathbb{A}f = (q^{-n} - \alpha)\mathbb{A}f - k \longleftarrow \mathbb{A}f$$

**Case 2** If  $\alpha > q^{-n}$  then  $(1 - \alpha q^{-n})$  and  $(q^{-n} - \alpha)$  are negative, so by Corollary 1.30 we have

$$f - \alpha \mathbb{A}f = (1 - \alpha q^{-n})f - \alpha h \longrightarrow f$$
$$f - \alpha \mathbb{A}f = (q^{-n} - \alpha)\mathbb{A}f - k \longrightarrow \mathbb{A}f$$

In either case we can apply the affine criteria of Lemma 8.3 to conclude that  $f-\alpha \mathbb{A} f\in \textbf{P}^{\mathbb{A}}.$ 

*Remark* 8.6. The proof of the lemma did not use an analysis of the roots. If we look at the roots then we get another proof of the lemma, as well as information about the behavior when  $\alpha$  is positive.

If the roots of f are  $r_1 < \cdots < r_n$  then we know that

$$\mathbb{A}^{-1}\mathbf{r}_1 < \mathbf{r}_1 < \mathbb{A}^{-1}\mathbf{r}_2 < \mathbf{r}_2 < \dots < \mathbb{A}^{-1}\mathbf{r}_n < \mathbf{r}_n.$$

If  $\alpha$  is negative then the roots {s<sub>i</sub>} of f +  $\alpha A f$  lie in the intervals

$$(-\infty, \mathbb{A}^{-1}\mathbf{r}_1), (\mathbf{r}_1, \mathbb{A}^{-1}\mathbf{r}_2), \cdots, (\mathbf{r}_{n-1}, \mathbb{A}^{-1}\mathbf{r}_n)$$

and consequently  $s_i \leq \mathbb{A}^{-1}r_i < r_i \leq s_{i+1}$ . This shows that  $\mathbb{A}s_i < s_{i+1}$ , and hence  $f + \alpha \mathbb{A}f \in \mathbf{P}^{\mathbb{A}}$ .

If  $\alpha$  is positive the roots lie in the intervals

$$(\mathbb{A}^{-1}\mathbf{r}_1,\mathbf{r}_1), (\mathbb{A}^{-1}\mathbf{r}_2,\mathbf{r}_2), \cdots, (\mathbb{A}^{-1}\mathbf{r}_n,\mathbf{r}_n)$$

The roots  $\{s_i\}$  satisfy  $s_i \leq r_i \leq \mathbb{A}^{-1}r_{i+1} \leq s_{i+1}$ . We can not conclude (indeed it's not true) that  $\mathbb{A}s_i \leq s_{i+1}$  from these inequalities.

If we make stronger assumptions about the roots of f then we can draw conclusions about  $f + \alpha \mathbb{A}f$ . Suppose that  $f \ll \mathbb{A}^2 f$ . This means that the roots of f satisfy  $\mathbb{A}^2 r_i < r_{i+1}$ . The roots  $s_i$  now satisfy  $s_i \leqslant r_i, \mathbb{A}^{-2} r_{i+1} \leqslant s_{i+1}$ . Since the interlacing hypothesis implies that  $\mathbb{A}r_i \leqslant \mathbb{A}^{-1}r_{i+1}$  we conclude that  $\mathbb{A}s_i \leqslant s_{i+1}$ . A similar argument establishes the following result:

**Lemma 8.7.** If k is a positive integer greater than 1,  $f \ll \mathbb{A}^k f$  and  $\alpha$  is positive then  $g = f + \alpha \mathbb{A} f$  satisfies  $g \ll \mathbb{A}^{k-1} g$ .

The affine derivative has all the expected properties.

**Lemma 8.8.** Suppose  $f \in \mathbf{P}^{\mathbb{A}}$ .

- 1.  $D_{\mathbb{A}}f \in P^{\mathbb{A}}$ .
- 2. f  $\leq D_{\mathbb{A}}$ f.
- 3.  $\mathbb{A}f \leq D_{\mathbb{A}}f$
- 4. If  $f, g \in \mathbf{P}^{\mathbb{A}}$  satisfy  $f \leq g$  then  $D_{\mathbb{A}}f \leq D_{\mathbb{A}}g$ .

*Proof.* Lemma 8.5 implies that  $f - Af \in \mathbf{P}^A$ , so the first result follows from Lemma 8.2. Since  $f \leq f - Af$  and the smallest root of f - Af is the fixed point, we can divide out by x - Ax and conclude from Lemma 8.2 that  $f \leq D_A f$ . The third part is similar.

In order to show that  $D_{\mathbb{A}}$  preserves interlacing it suffices to show that  $D_{\mathbb{A}}(x+a)f \leq D_{\mathbb{A}}f$  where  $(x+a)f \in \mathbf{P}^{\mathbb{A}}$ . From the product rule (8.2.3) we find

$$\mathsf{D}_{\mathbb{A}}(x+a)\mathsf{f} = (x+a) \mathsf{D}_{\mathbb{A}}\mathsf{f} + \mathbb{A}\mathsf{f}$$

Since  $(x + a) D_{\mathbb{A}} f \leq D_{\mathbb{A}} f$  and  $\mathbb{A} f \leq D_{\mathbb{A}} f$  we can add these interlacings to conclude  $D_{\mathbb{A}}(x + a) f \leq D_{\mathbb{A}} f$ .

 $\mathbf{P}^{\mathbb{A}}$  is closed under ordinary differentiation.

**Lemma 8.9.** If  $f \in \mathbf{P}^{\mathbb{A}}$  then  $f' \in \mathbf{P}^{\mathbb{A}}$ 

*Proof.* Since  $f \in \mathbf{P}^{\mathbb{A}}$  we know that  $f + \alpha \mathbb{A}f \in \mathbf{P}$  for all  $\alpha$ . Since  $\frac{d}{dx}\mathbb{A}f = q\mathbb{A}(f')$  we have that  $f' + \alpha q\mathbb{A}f' \in \mathbf{P}$  for all  $\alpha$  and so  $f' \in \mathbf{P}^{\mathbb{A}}$ .

Only in certain special cases do  $\mathbb A$  and  $\ D_{\mathbb A}$  commute.

**Lemma 8.10.** For any positive integer n we have that  $q^n \mathbb{A}^n D_q(f) = D_q(\mathbb{A}^n f)$ . Consequently, if  $\mathbb{A} D_{\mathbb{A}} = D_{\mathbb{A}} \mathbb{A}$  then  $\mathbb{A} x = x + b$ .

Proof. First note that

$$\mathbb{A}(\mathbb{A}x-x) = \mathbb{A}(qx+b-x) = (q-1)(qx+b) + b = q(qx+b-x) = q(\mathbb{A}x-x).$$

This implies that for all positive n we have  $\mathbb{A}^n(\mathbb{A}x - x) = q^n(\mathbb{A}x - x)$ , and hence  $\mathbb{A}^{-n}(\mathbb{A}x - x) = q^{-n}(\mathbb{A}x - x)$ . We can now compute:

$$\begin{split} \mathsf{D}_{\mathsf{q}}(\mathbb{A}^{\mathsf{n}}\mathsf{f}) &= \frac{\mathbb{A}^{\mathsf{n}+1}\mathsf{f} - \mathbb{A}^{\mathsf{n}}\mathsf{f}}{\mathbb{A}\mathsf{x} - \mathsf{x}} \\ &= \mathbb{A}^{\mathsf{n}} \frac{\mathbb{A}\mathsf{f} - \mathsf{f}}{\mathbb{A}^{-\mathsf{n}}(\mathbb{A}\mathsf{x} - \mathsf{x})} \\ &= \mathsf{q}^{\mathsf{n}}\mathbb{A}^{\mathsf{n}} \mathsf{D}_{\mathsf{q}}\mathsf{f} \end{split}$$

We next consider properties of  $f(\mathbb{A})g$  and  $f(D_{\mathbb{A}})g$ . We first observe this immediate consequence of Lemma 8.5.

**Lemma 8.11.** *If*  $g \in P^{alt}$  *and*  $f \in P^{A}$  *then*  $g(A)f \in P^{A}$ .

**Corollary 8.12.** If  $g \in \mathbf{P}^{\text{pos}}(\mathbf{r})$  and  $f \ll \mathbb{A}^{r+1}f$  then  $g(\mathbb{A})f \in \mathbf{P}^{\mathbb{A}}$ . If  $f \ll \mathbb{A}^r$  then  $\sum_{0}^{r} {r \choose k} \mathbb{A}^k f \in \mathbf{P}$ .

*Proof.* Factor g, and apply Lemma inductively. For the second part, apply the first part to  $g = (1 + x)^r$ .

Next we have an additive closure property of the affine derivative.

**Lemma 8.13.** If  $g \in \mathbf{P}^{\mathbb{A}}$  and  $\alpha$  is positive then  $g - \alpha D_{\mathbb{A}}g \in \mathbf{P}^{\mathbb{A}}$ .

*Proof.* We know that  $g \leq D_{\mathbb{A}}g$  so  $g - \alpha D_{\mathbb{A}}g \leq g$ . Also,  $g \leq \mathbb{A}g \leq D_{\mathbb{A}}g$ , so  $g - \alpha D_{\mathbb{A}}g \ll \mathbb{A}g$ . Now apply Lemma 8.3.

**Lemma 8.14.** If  $g \in P^{alt}$  and  $f \in P^{A}$  then  $g(D_{A})f \in P^{A}$ .

*Proof.* Write  $g(x) = (x + a_1) \dots (x + a_n)$  where all  $a_i$  are negative. Since

$$g(\mathsf{D}_{\mathbb{A}})f = (\mathsf{D}_{\mathbb{A}} + \mathfrak{a}_1) \dots (\mathsf{D}_{\mathbb{A}} + \mathfrak{a}_n)f$$

we apply Lemma 8.13 inductively.

Here's a simple property about determinants.

**Lemma 8.15.** *Suppose*  $f, Af \in P^A \cap P^{pos}$ .

$$\begin{split} & If \begin{cases} \mathbb{A}x > x \text{ for all positive } x \\ \mathbb{A}x < x \text{ for all positive } x \end{cases} \quad then \begin{vmatrix} f & \mathbb{A}f \\ \mathbb{A}f & \mathbb{A}^2f \end{vmatrix} \text{ is } \begin{cases} \text{negative for positive } x. \\ \text{positive for positive } x. \end{cases} \\ & \text{The same inequalities hold for } \begin{vmatrix} f & D_{\mathbb{A}}f \\ D_{\mathbb{A}}f & D_{\mathbb{A}}^2f \end{vmatrix}. \end{split}$$

*Proof.* We begin with the first statement. We need to show that  $f \cdot \mathbb{A}^2 f < \mathbb{A} f \cdot \mathbb{A} f$ for positive x. Since Af is in  $P^{pos}$ , Af is positive for positive x, so it suffices to show that  $(\mathbb{A}f/\mathbb{A}^2f) > (f/\mathbb{A}f)$ .

Since  $f \in \mathbf{P}^{sep}$  we know that  $f \underline{\ll} \mathbb{A}f$ , and so there is a positive constant c and non-negative  $a_i$  such that

$$\mathbb{A} f = cf + \sum \mathfrak{a}_i \frac{f}{x + r_i}$$

where the roots of f are  $\{r_i\}$ . Thus

$$\begin{split} \frac{\mathbb{A}f}{f} &= c + \sum \frac{a_i}{x+r_i} \\ \left(\frac{\mathbb{A}f}{f}\right)' &= -\sum a_i \frac{1}{(x+r_i)^2} \end{split}$$

and consequently Af/f is decreasing. Since Ax > x, it follows that

$$\frac{f}{\mathbb{A}f}(x) < \frac{f}{\mathbb{A}f}(\mathbb{A}x) = \frac{\mathbb{A}f}{\mathbb{A}^2f}(x)$$

If Ax < x then the inequality is reversed. The last statement follows from the above since

$$\begin{vmatrix} f & \mathbb{A}f \\ \mathbb{A}f & \mathbb{A}^2f \end{vmatrix} = \frac{1}{(\mathbb{A}x - x)^2} \begin{vmatrix} f & \mathsf{D}_{\mathbb{A}}f \\ \mathsf{D}_{\mathbb{A}}f & \mathsf{D}_{\mathbb{A}}^2f \end{vmatrix}.$$

*Remark* 8.16.  $\langle \underline{x}; \underline{a} \rangle_{n}^{\mathbb{A}}$  in (8.1.1) is uniquely determined by the three conditions below.

- $\left< \underline{\mathbf{x}; \mathbf{a}} \right>_{\mathbf{0}}^{\mathbb{A}} = 1$ (1)
- $\left< \underline{a}; \underline{a} \right>_{n}^{\mathbb{A}} = 0$ (2)

(3)  $\mathsf{D}_{\mathbb{A}} \langle \underline{\mathbf{x}}; \underline{\mathbf{a}} \rangle_{n}^{\mathbb{A}} = [n] \mathbb{A} \langle \underline{\mathbf{x}}; \underline{\mathbf{a}} \rangle_{n-1}^{\mathbb{A}}$ This characterization of  $\langle \underline{\mathbf{x}}; \underline{\mathbf{a}} \rangle_{n}^{\mathbb{A}}$  is motivated a similar result by [98] in the case  $\mathbb{A}x = qx$ .

# 8.3 Sign interlacing for $P^{\mathbb{A}}$

Let's look at some graphs in order to better understand  $\mathbf{P}^{\mathbb{A}}$ . Suppose that  $f \leq \mathbb{A}f$ . The graph of  $\mathbb{A}f$  is the graph of f, with some shifting to the left and scaling. For instance, if  $\mathbb{A}x = x + 1$  and f = x(x - 1)(x - 3)(x - 5) then the graph of f and  $\mathbb{A}f$  is in Figure 8.3.



Figure 8.1: Understanding  $\mathbb{A}$ 

If  $f \leq g$  and  $Af \leq g$  then the roots of g must lie in the thickened segments of Figure 8.3. It's clear that the roots of g in the figure differ by at least one, as is guaranteed by Lemma 8.3.

If we translate the qualitative aspects of this graph, we get a version of sign interlacing. Assume that the roots of f are

$$r_1 < r_2 < \dots < r_n$$
 (8.3.1)

The roots of Af are

$$\mathbb{A}^{-1}\mathbf{r}_1 < \mathbb{A}^{-1}\mathbf{r}_2 < \dots < \mathbb{A}^{-1}\mathbf{r}_n \tag{8.3.2}$$

Since  $f \leq Af$  we can combine these orderings

$$\mathbb{A}^{-1}\mathbf{r}_1 < \mathbf{r}_1 \leqslant \mathbb{A}^{-1}\mathbf{r}_2 < \mathbf{r}_2 \leqslant \dots \leqslant \mathbb{A}^{-1}\mathbf{r}_n < \mathbf{r}_n \tag{8.3.3}$$

If  $f \leq g$  and  $\mathbb{A}f \leq g$  all have positive leading coefficients then it is clear from the graph that  $g(r_n) > 0$ ,  $g(\mathbb{A}^{-1}r_n) > 0$ ,  $g(r_{n-1} < 0, g(\mathbb{A}^{-1}r_{n-1}) < 0$ , and generally

$$(-1)^{n+i}g(r_i) > 0 \qquad (-1)^{n+i}g(\mathbb{A}^{-1}r_i) > 0 \tag{8.3.4}$$

This is sign interlacing for  $\mathbb{A}$ . We clearly have

**Lemma 8.17.** If g is a polynomial of degree n - 1 that satisfies (8.3.4) then  $g \in \mathbf{P}^{\mathbb{A}}$ .

The following polynomials play the role of  $f(x)/(x-r_{\rm i})$  in Lemma 1.20. For  $1\leqslant k\leqslant n$  define

$$f_{[k]} = \mathbb{A} \left[ (x - r_1) \cdots (x - r_{k-1}) \right] \cdot \left[ (x - r_{k+1}) \cdots (x - r_n) \right]$$
(8.3.5)

These are polynomials of degree n - 1 whose roots are

$$\mathbb{A}^{-1}r_1 < \mathbb{A}^{-1}r_2 \cdots \mathbb{A}^{-1}r_{k-1} < r_{k+1} < \cdots < r_n$$
(8.3.6)

Since the  $r_i$  are all distinct, it is easy to see that the  $f_{[k]}$  are independent. Thus, they form a basis for the polynomials of degree n - 1.

**Lemma 8.18.** Suppose that  $f \in \mathbf{P}^{\mathbb{A}}$  and  $g = \sum a_k f_{[k]}$  where all  $a_i$  are non-negative. Then  $g \in \mathbf{P}^{\mathbb{A}}$  and  $f \leq g$  and  $\mathbb{A}f \leq g$ .

*Proof.* The key point is that  $f \leq f_{[k]}$  and  $\mathbb{A}f \leq f_{[k]}$  - this follows from (8.3.1), (8.3.2), (8.3.3), (8.3.5). If all the  $a_i$  are non-negative then we can add these interlacings to finish the proof.

The fact that  $f \leq D_{\mathbb{A}}f$  follows from Lemma 8.18. From (8.2.4) we see that if  $f(x) = (x - r_1) \cdots (x - r_n)$  then

$$\begin{split} \mathsf{D}_{\mathbb{A}}(f) &= \ \mathsf{D}_{\mathbb{A}}(x - r_{1}) \cdots (x - r_{n}) \\ &= \sum_{k=1}^{n} \mathbb{A}(\ (x - r_{1}) \cdots (x - r_{k-1})\ ) \cdot 1 \cdot (\ (x - r_{k+1}) \cdots (x - r_{n})\ ) \\ &= f_{[1]} + \cdots + f_{[n]} \end{split}$$

Consequently, this is another proof that  $f \leq D_A f$ .

## 8.4 The affine exponential

We can formally construct an affine exponential  $E_{\mathbb{A}}$  function, but the series might not converge. We want two simple conditions

- (1)  $E_{\mathbb{A}}(0) = 1$
- (2)  $D_{\mathbb{A}}(E_{\mathbb{A}}) = \mathbb{A} E_{\mathbb{A}}$

The relation  $D_{\mathbb{A}}E_{\mathbb{A}} = \mathbb{A}E_{\mathbb{A}}$  determines  $E_{\mathbb{A}}$  up to a constant factor that is settled by assuming  $E_{\mathbb{A}}(0) = 1$ . If we write

$$\mathsf{E}_{\mathbb{A}}(\mathbf{x}) = 1 + \mathfrak{a}_1 \mathbf{x} + \mathfrak{a}_2 \mathbf{x}^2 + \cdots$$

and apply  $D_{\mathbb{A}}E_{\mathbb{A}} = \mathbb{A}E_{\mathbb{A}}$  we find

$$\mathfrak{a}_{n-1}\mathbb{A}(\mathfrak{x}^{n-1}) = \mathfrak{a}_n \mathsf{D}_{\mathbb{A}}(\mathfrak{x}^n)$$

and recursively solving for the coefficients yields

$$\mathsf{E}_{\mathbb{A}}(x) = \sum_{n=0}^{\infty} \prod_{k=1}^{n} \frac{\mathbb{A}^{k-1} x}{[k]_{\mathbb{A}}}$$

where

$$[k]_{\mathbb{A}} = (\mathbb{A}^k x - x)/(\mathbb{A} x - x) = [k] = q^{k-1} + q^{k-2} + \dots + q + 1.$$

and

$$\mathbb{A}^k \mathbf{x} = \mathbf{q}^k \mathbf{x} + \mathbf{b}[\mathbf{k} + 1]$$

Now this derivation is not valid if  $b \neq 0$ , but we still can use this definition since we have that

$$\mathsf{D}_{\mathbb{A}}\prod_{k=1}^{n}\frac{\mathbb{A}^{k-1}x}{[k]_{\mathbb{A}}} = \mathbb{A}\prod_{k=1}^{n-1}\frac{\mathbb{A}^{k-1}x}{[k]_{\mathbb{A}}}$$

and thus term by term differentiation establishes condition (2). In particular

$$\mathbb{A}x = x \qquad \qquad \mathsf{E}_{\mathbb{A}}(x) = \sum_{n=0}^{\infty} \frac{x^n}{n!} \qquad \qquad (8.4.1)$$

$$\mathbb{A}\mathbf{x} = q\mathbf{x} \qquad \qquad \mathsf{E}_{\mathbb{A}}(\mathbf{x}) = \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}}}{[n]!} \mathbf{x}^n \qquad \qquad (8.4.2)$$

$$\mathbb{A}x = q(x+1) \qquad \qquad \mathsf{E}_{\mathbb{A}}(x) = \sum_{n=0}^{\infty} \frac{\prod_{1}^{n} (q^{k-1}x + q[k])}{[n]!} \qquad \qquad (8.4.3)$$

$$Ax = x + 1 \qquad \qquad \mathsf{E}_{\mathbb{A}}(x) = \sum_{n=0}^{\infty} \frac{\langle \underline{x} \rangle_n}{n!} \qquad (8.4.4)$$

In the first case  $E_{\mathbb{A}}$  is the usual exponential. The second one is the q-exponential of (5.1.1) and has a representation as an infinite product. The last two have product representations for their finite sums:

Ax = qx 
$$\sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}}}{[n]!} x^n = \prod_{k=0}^{\infty} \left( 1 - q^k (1-q) x \right)$$
(8.4.5)

$$\mathbb{A}x = q(x+1) \qquad \sum_{n=0}^{N} \left( \prod_{k=1}^{n} \frac{\mathbb{A}^{k-1}x}{[k]} \right) = \prod_{k=1}^{N} \left( 1 + \frac{q^{k-1}x}{[k]} \right)$$
(8.4.6)

$$Ax = x + 1 \qquad \qquad \sum_{n=0}^{N} \frac{\langle \underline{x} \rangle_n}{n!} = \prod_{k=1}^{N} \left( 1 + \frac{x}{k} \right) = \frac{\langle \underline{x} + 1 \rangle_N}{N!} \qquad (8.4.7)$$

Unlike the partial sums of the exponential function, the partial sums of (8.4.4) and (8.4.3) have all real roots, as the product representation shows. Since the sum  $\sum q^{k-1}/[k]$  converges for |q| < 1 and diverges for |q| > 1, it follows that the product (8.4.6) converges for |q| < 1 and diverges for  $|q| \ge 1$ . Consequently, equation (8.4.3) is only valid for |q| < 1. The series in (8.4.4) does not converge.

The partial sums in the last two cases satisfy the exponential condition (1) and nearly satisfy (2). Let  $E_{A,N}$  denote the partial sums in (8.4.6) and (8.4.7). Then

$\mathbb{A}\mathbf{x} = \mathbf{q}(\mathbf{x} + 1)$	$D_{\mathbb{A}}(E_{\mathbb{A},N}) = \mathbb{A}E_{\mathbb{A},N-1}$
$\mathbb{A}\mathbf{x} = \mathbf{x} + 1$	$\Delta(E_{\mathbb{A},N}) = \mathbb{A} E_{\mathbb{A},N-1}$

## 8.5 A-interlacing and the symmetric derivative

We say that f and g  $\mathbb{A}$ -interlace if  $f + \alpha g \in \mathbf{P}^{\mathbb{A}}$  for all  $\alpha \in \mathbb{R}$ . If the degree of f is greater than the degree of g and f and g  $\mathbb{A}$ -interlace then we write  $f \leq^{\mathbb{A}} g$ . We give a criterion for affine interlacing, and use it to determine when the symmetric affine derivative is in  $\mathbf{P}^{\mathbb{A}}$ .

**Lemma 8.19.** If  $f \in \mathbf{P}^{\mathbb{A}}$  and  $f \leq \mathbb{A}^2 f$  then f and  $\mathbb{A} f \mathbb{A}$ -interlace.

*Proof.* From the interlacings  $f \leq Af$ ,  $f \leq A^2f$ ,  $Af \leq A^2f$  we conclude that if  $\alpha > 0$ 

$$f \leq f + \alpha \mathbb{A}f$$
$$f \ll \mathbb{A}f + \alpha \mathbb{A}^2f$$

and if  $\alpha < 0$ 

$$f \geq f + \alpha \mathbb{A}f$$
$$f \geq \mathbb{A}f + \alpha \mathbb{A}^2f$$

Thus  $f + \alpha \mathbb{A}f$  and  $\mathbb{A}(f + \alpha \mathbb{A}f)$  always have a common interlacing, so we find that  $f + \alpha \mathbb{A}f \in \mathbf{P}^{\mathbb{A}}$ .

Here is an example of polynomial Q that A-interlaces AQ. If A is increasing on  $\mathbb R$  then define

$$\mathbf{Q} = (\mathbf{x} + 1)(\mathbf{x} + \mathbb{A}^2 \mathbf{1})(\mathbf{x} + \mathbb{A}^4 \mathbf{1}) \cdots (\mathbf{x} + \mathbb{A}^{2n} \mathbf{1})$$

and if  $\mathbb{A}$  is decreasing define

$$\mathbf{Q} = (\mathbf{x} - 1)(\mathbf{x} - \mathbb{A}^2 \mathbf{1})(\mathbf{x} - \mathbb{A}^4 \mathbf{1}) \cdots (\mathbf{x} - \mathbb{A}^{2n} \mathbf{1})$$

If we apply  $\mathbb{A}^2$  to Q we have even powers from  $\mathbb{A}^2$  to  $\mathbb{A}^{2n+2}$ , so clearly  $Q \leq \mathbb{A}^2 Q$ .

Here is a variation on the lemma for two polynomials.

**Lemma 8.20.** If  $f \leq g$  and  $fg \in \mathbf{P}^{\mathbb{A}}$  then  $f \leq \mathbf{A}^{\mathbb{A}}$  g.

*Proof.* Suppose that the roots of f(x) are  $f_1 < \cdots$  and the roots of g(x) are  $g_1 < \cdots$ . The hypotheses imply that the roots appear in the order

$$\cdots g_i \leqslant \mathbb{A}g_i \leqslant f_i \leqslant \mathbb{A}f_i \leqslant g_{i+1} \leqslant \mathbb{A}g_{i+1} \leqslant \mathbb{A}f_1 \cdots$$

from which it follows that  $f \leq Ag$ . For positive  $\alpha$  (first column) and negative  $\alpha$  (second column) we have

$$f \underline{\ll} f + \alpha g \qquad \qquad f \underline{\gg} f + \alpha g \\ f \underline{\ll} A f + \alpha A g \qquad \qquad f \underline{\gg} A f + \alpha A g$$

Consequently,  $f + Ag \in \mathbf{P}^A$ .

Clearly  $f \leq ^{\mathbb{A}} g$  implies that  $f \leq g$  but  $\leq ^{\mathbb{A}}$  is a more restrictive condition than  $\leq$ , and the converse does not usually hold. We say that  $f < ^{\mathbb{A}} g$  if  $f \leq ^{\mathbb{A}} g$ , and f and g have no common factors. Using sign interlacing we can replace  $\leq$  and  $\leq ^{\mathbb{A}}$  in the lemma by < and  $<^{\mathbb{A}}$ . There is a simple criterion for such interlacing.

**Lemma 8.21.** Suppose that  $f \leq g$ . Then  $f \leq^{\mathbb{A}} g$  if and only if  $\begin{vmatrix} f & g \\ \mathbb{A}f & \mathbb{A}g \end{vmatrix}$  is never zero.

*Proof.* From Lemma 1.52 we have that  $f + \alpha g \ll \mathbb{A}f + \alpha \mathbb{A}g$  for all  $\alpha \in \mathbb{R}$ . This means that  $f + \alpha g \in \mathbf{P}^{\mathbb{A}}$  so the conclusion follows.

The symmetric derivative is

$$\mathsf{D}^{\text{sym}}_{\mathbb{A}}(\mathsf{f}) = \frac{\mathbb{A}(\mathsf{f}) - \mathbb{A}^{-1}(\mathsf{f})}{\mathbb{A}(\mathsf{x}) - \mathbb{A}^{-1}\mathsf{x}}$$

For Ax = x + 1 this is the symmetric difference

$$\mathsf{D}^{\texttt{sym}}_{\mathbb{A}} = \frac{\mathsf{f}(\mathsf{x}+1) - \mathsf{f}(\mathsf{x}-1)}{2}$$

and for Ax = qx it is

$$\mathsf{D}^{\texttt{sym}}_{\mathbb{A}}(\mathsf{f}) = \frac{\mathsf{f}(\mathsf{q} \mathsf{x}) - \mathsf{f}(\mathsf{x}/\mathsf{q})}{(\mathsf{q} - 1/\mathsf{q})\mathsf{x}}.$$

**Lemma 8.22.** If  $f \in \mathbf{P}^{\mathbb{A}}$  then  $D^{sym}_{\mathbb{A}} f \in \mathbf{P}$ . If in addition we have that  $f \leq \mathbb{A}^2 f$  then  $D^{sym}_{\mathbb{A}} f \in \mathbf{P}^{\mathbb{A}}$ .

*Proof.* Since  $f \in \mathbf{P}^{\mathbb{A}}$  we know that  $f \leq \mathbb{A}f$  and consequently  $f - \mathbb{A}f \in \mathbf{P}$ , so  $\mathsf{D}^{sym}_{\mathbb{A}} f \in \mathbf{P}$ . Next, assume that  $f \leq \mathbb{A}^2 f$ . We have the interlacings

$$\mathbb{A}^{-1} \mathbf{f} \underline{\ll} \mathbf{f} \underline{\ll} \mathbb{A} \mathbf{f}$$
$$\mathbb{A}^{-1} \mathbf{f} \underline{\ll} \mathbb{A}^{-1} \mathbf{f} \underline{\ll} \mathbb{A} \mathbf{f}$$

which imply that

$$\begin{aligned} \mathbf{f} &\leq \mathbb{A}\mathbf{f} - \mathbb{A}^{-1}\mathbf{f} \\ \mathbb{A}^{-1}\mathbf{f} &\leq \mathbb{A}\mathbf{f} - \mathbb{A}^{-1}\mathbf{f} \end{aligned}$$

The last two interlacings imply  $\mathbb{A}f - \mathbb{A}^{-1}f \in \mathbf{P}^{\mathbb{A}}$ , and so the conclusion follows.

Here's an example that shows that the symmetric derivative is not in  $\mathbf{P}^{\mathbb{A}}$  if the second condition is not satisfied. If  $\mathbb{A}x = x + 1$  then

$$\mathsf{D}^{sym}_{\mathbb{A}}(\underline{x})_{\mathfrak{n}} = \mathfrak{n}(x - \frac{\mathfrak{n} - 1}{2})(\underline{x - 1})_{\mathfrak{n} - 2}.$$

If n is odd then the symmetric derivative has a double root, and so is not in  $\mathbf{P}^{sep}$ .

## 8.6 Linear transformations

In this section we investigate transformations that map some subset of **P** to  $\mathbf{P}^{\mathbb{A}}$ . The key idea is the iteration of the linear transformation  $f \mapsto (x - \alpha)\mathbb{A}^{-1}f - \beta f$ .

**Lemma 8.23.** The linear transformation S:  $f \mapsto (x - \alpha)\mathbb{A}^{-1}f - \beta f$  where  $\beta \ge 0$  maps

$$\mathsf{S}: \boldsymbol{P}^{\mathbb{A}} \cap \boldsymbol{P}^{(\alpha,\infty)} \longrightarrow \boldsymbol{P}^{\mathbb{A}} \cap \boldsymbol{P}^{(\alpha,\infty)}$$

*Proof.* Since  $f \in \mathbf{P}^{\mathbb{A}}$  we know

$$\mathbb{A}^{-1}\mathbf{f} \leq \mathbf{f}$$

and using  $f\in I\!\!P^{(\,\alpha,\infty)}$  we get

$$(\mathbf{x} - \alpha) \mathbb{A}^{-1} \mathbf{f} \leq \mathbb{A}^{-1} \mathbf{f}.$$
(8.6.1)

$$(\mathbf{x} - \alpha) \mathbb{A}^{-1} \mathbf{f} \le \mathbf{f}. \tag{8.6.2}$$

By hypothesis  $\beta \ge 0$ , so we can apply Corollary 1.30 to (8.6.2) and Lemma 1.31 to (8.6.1) yielding

$$(\mathbf{x} - \alpha) \mathbb{A}^{-1} \mathbf{f} - \beta \mathbf{f} \leq \mathbf{f}$$
$$(\mathbf{x} - \alpha) \mathbb{A}^{-1} \mathbf{f} - \beta \mathbf{f} \leq \mathbb{A}^{-1} \mathbf{f}$$

which shows that  $S(f) \in \mathbf{P}^{\mathbb{A}}$ . Since  $\mathbb{A}^{-1}f \in \mathbf{P}^{(\alpha,\infty)}$  we can apply Corollary 1.30 to (8.6.2)

$$(\mathbf{x} - \boldsymbol{\alpha})\mathbb{A}^{-1}\mathbf{f} - \beta\mathbf{f} \underline{\ll} (\mathbf{x} - \boldsymbol{\alpha})\mathbb{A}^{-1}\mathbf{f}$$

and conclude that  $S(f) \in \mathbf{P}^{(\alpha,\infty)}$ .

Now, define polynomials by  $Q_n^{\alpha,\beta}(x) = ((x - \alpha)\mathbb{A}^{-1} - \beta)^n$  (1). Some particular examples are given in Table 8.1. They are immediate from their definitions, except for the Charlier polynomials, where we apply the recurrence (7.9.4).

Ax	α	β	$Q_n^{\alpha,\beta}$
x + 1	0	0	$(\underline{\mathbf{x}})_{\mathbf{n}}$
x + 1	0	β	$C_n^\beta(x)$
qx	0	0	$q^{-\binom{n}{2}}x^n$
qx	-1	0	$q^{-\binom{n}{2}}(-x;q)_n$

Table 8.1: Polynomials of the form  $Q_n^{\alpha,\beta}$ 

**Proposition 8.24.** The map  $T: x^n \mapsto Q_n^{\alpha,\beta}(x)$  is a linear transformation

$$P^{(-\beta,\infty)} \longrightarrow P^{(\alpha,\infty)} \cap P^{\mathbb{A}}$$

*Proof.* If we define  $S(f) = (x - \alpha)\mathbb{A}^{-1}f - \beta f$ , then from the definition of T we have that  $T(x^n) = S^n(1) = Q_n^{\alpha,\beta}$ , and so by linearity T(f) = f(S)(1). If we choose  $f = \prod (x - r_i)$  in  $\mathbf{P}^{(-\beta,\infty)}$  then each root  $r_i$  satisfies  $r_i \ge -\beta$ . Notice that

$$T(f) = \prod(S - r_i) = \prod((x - \alpha)\mathbb{A}^{-1} - \beta - r_i)(1)$$

Since  $\beta + r_i \ge 0$ , if follows from the lemma that  $T(f) \in \mathbf{P}^{(\alpha,\infty)} \cap \mathbf{P}^{\mathbb{A}}$ .

Here is a general criterion to tell if a transformation preserves  $\mathbf{P}^{\mathbb{A}}$ . See Lemma 8.56 for an application.

**Lemma 8.25.** If  $T: P \longrightarrow P$  preserves interlacing, and  $T\mathbb{A} = \mathbb{A}T$ , then  $T: P^{\mathbb{A}} \longrightarrow P^{\mathbb{A}}$ .

*Proof.* Assume that  $f \leq Af$ . Since T preserves interlacing we have  $Tf \leq TAf$ . Commutativity implies that TAf = ATf, and so  $T(f) \leq AT(f)$ , and hence  $T(f) \in \mathbf{P}^{\mathbb{A}}$ .

## **8.7** The case Ax = x + 1

In this section we study the affine operator<sup>1</sup>  $\mathbb{A}x = x + 1$ . The space of all polynomials f satisfying  $f(x) \leq f(x + 1)$  is denoted  $\mathbf{P}^{sep}$ , where *sep* stands for *separated*. Here are some necessary and sufficient conditions for a polynomial f to belong to  $\mathbf{P}^{sep}$ :

<sup>&</sup>lt;sup>1</sup>This operator is often denoted by E.

- 1.  $f(x) \leq f(x+1)$
- 2. The distance between any two roots of f is at least 1.
- 3. f(x) and f(x + 1) have a common interlacing.
- 4. For all positive  $\beta$  we have  $f(x) + \beta f(x+1) \in \mathbf{P}$ .

For example, both  $(\underline{x})_n$  and  $\langle \underline{x} \rangle_n$  are in  $\mathbf{P}^{sep}$ . The affine derivative is the difference operator  $\Delta(f) = f(x+1) - f(x)$ . The difference operator commutes with  $\mathbb{A}$ . Since the derivative maps  $\mathbf{P}^{sep}$  to itself, the roots of the derivative can't get too close:

**Lemma 8.26.** *If the distance between any two roots of a polynomial* f *in* P *is at least* 1, *then the distance between any two roots of the derivative* f' *is also at least* 1. *Similarly, the distance between any two roots of* f(x + 1) - f(x) *is at least one.* 

*Proof.* We know that  $f \in \mathbf{P}^{sep}$  implies that both f' and  $\Delta f$  are also in  $\mathbf{P}^{sep}$ .

If we express a polynomial in the basis  $(\underline{x})_n$ , and use the fact that  $\Delta(\underline{x})_n = n(\underline{x})_{n-1}$  we conclude

If  $\sum a_i(\underline{x})_i$  has roots that are separated by at least one, then the same is true for  $\sum ia_i(\underline{x})_{i-1}$ .

From Lemma 8.8 we find that differences preserve interlacing:

**Lemma 8.27.** If f,  $g \in \mathbf{P}^{sep}$  satisfy  $f \leq g$  then  $\Delta f \leq \Delta g$ .

**Corollary 8.28.** If  $g \in P^{alt}$  and  $f \in P^{sep}$  then  $g(A)f \in P^{sep}$  where A(f(x)) = f(x+1). Similarly, if  $g \in P^{alt}$  then  $g(\Delta)f \in P^{sep}$ .

*Proof.* Use Lemma 8.14 for the second part.

The second part of this corollary does not hold for all f. For instance, if f = x(x + 1) then  $f - \alpha \Delta f = (x + 1)(x - 2\alpha)$ , and the distance between roots is  $|1 + 2\alpha|$ . Thus  $f - \alpha \Delta f \in \mathbf{P}^{sep}$  if and only if  $\alpha \notin (-1, 0)$ . This is true generally.

**Lemma 8.29.** If  $f \in \mathbf{P}^{\mathbb{R} \setminus (-1,0)}$  then  $f(\Delta) : \mathbf{P}^{sep} \longrightarrow \mathbf{P}^{sep}$ .

*Proof.* It suffices to show that  $f - \alpha \Delta f \in \mathbf{P}^{sep}$  if  $f \in \mathbf{P}^{sep}$  and  $\alpha \notin (-1, 0)$ . Note that

$$f - \alpha \Delta f = -(\alpha + 1) \big( \frac{\alpha}{\alpha + 1} f(x + 1) - f(x) \big)$$

Since  $\alpha \notin (-1,0)$  implies  $\frac{\alpha}{\alpha+1} > 0$  we can apply Corollary 8.50 to finish the proof.

Here is a nice consequence of composition.

**Lemma 8.30.** Suppose that d is a positive integer. The linear transformation  $T: x^k \mapsto (\underline{x} + \underline{d} - \underline{k})_d$  maps  $P^{alt}$  to  $P^{sep}$ .

*Proof.* Let  $\mathbb{B}x = x - 1$ . The important observation is that  $\mathbb{B}^k(\underline{x + d})_d = T(x^k)$ . Thus,  $T(f) = f(\mathbb{B})(\underline{x + d})_d$ . Thus we only have to verify the conclusion for one factor. Since  $f \in \mathbf{P}^{a \ t}$ , we have to show that  $(\mathbb{B} - \alpha)$  maps  $\mathbf{P}^{sep}$  to itself if  $\alpha$  is positive. This follows from Lemma 8.5

Simple geometry forces a relationship between  $\Delta f$  and f'.

**Lemma 8.31.** If  $f \in P^{sep}$  then  $f' \ll \Delta f$ .

*Proof.* Consider Figure 8.2. The points a, b are consecutive zeros of f, and satisfy  $b - a \ge 1$ . Points d, e are on f, and the horizontal line between them has length exactly 1. Finally, c is a root of f'.

The x-coordinate of d is a root of f(x + 1) - f(x) = 0, and so is a root of  $\Delta f$ . Since the x coordinate of d is obviously less than c, it follows that  $f' \ll \Delta f$ .



Figure 8.2: The difference and the derivative

*Remark* 8.32. If  $f \in \mathbf{P}^{sep}$  then we know that  $f \leq f'$ ,  $f \leq \Delta f$ ,  $f' \in \mathbf{P}^{sep}$ , and  $\Delta f \in \mathbf{P}^{sep}$ . Even though  $f \leq f' + \Delta f$ , it is not necessarily the case that  $f' + \Delta f \in \mathbf{P}^{sep}$ . A simple counterexample is given by f = x(x - 1)(x - 2).

An important class of polynomials in  $\mathbf{P}^{sep}$  are the Charlier polynomials  $C_n^{\alpha}(x)$ . A simple consequence of the following result is that the Charlier polynomials are in  $\mathbf{P}^{sep}$ .

**Corollary 8.33.** *Assume*  $\alpha \leq 0$ .

- 1. The map  $x^i \mapsto C_n^{\alpha}$  maps  $P^{\alpha lt} \longrightarrow P^{\alpha lt} \cap P^{sep}$ .
- 2. The map  $x^{i} \mapsto (\underline{x})_{n}$  maps  $P^{alt} \longrightarrow P^{alt} \cap P^{sep}$ .
- 3. The map  $x^i \mapsto \langle \underline{x} \rangle_n$  maps  $P^{pos} \longrightarrow P^{pos} \cap P^{sep}$ .

*Proof.* Apply Proposition 8.24 to Table 8.1. Switch signs for the rising factorial in the result for the falling factorial.

Expressing this result in terms of coefficients yields

If  $\sum a_i x^i$  has all negative roots, then  $\sum a_i \langle \underline{x} \rangle_i$  has roots whose mutual distances are all at least one.

The following are consequences of Corollary 8.12.

**Corollary 8.34.** If  $g \in P^{pos}(n-1)$  and all roots of f are at least n apart then  $g(\mathbb{A})f \in P^{sep}$ .

**Corollary 8.35.** *If the roots of* f *are separated by at least* n + 1 *and*  $f \in P^{sep}(n)$  *then* 

$$\sum_{i=0}^{n} \binom{n}{i} f(n+i) \in P$$

If  $f \in \mathbf{P}^{sep}$  then

$$\sum_{i=0}^{n}(-1)^{i}\binom{n}{i}f(n+i)\in \textbf{P}$$

*Proof.* Apply the previous corollary to  $(1 + \mathbb{A})^n f$  and  $(\mathbb{A} - 1)^n f$ .

The next result is an interesting special case of Lemma 8.14.

**Corollary 8.36.** If  $g \in P^{sep}$  and m is a positive integer then

$$\sum_{i=0}^{m} (-1)^{i} \binom{m}{i} g(x+i) \in \boldsymbol{P}^{sep}.$$

*Proof.* Apply Lemma 8.14 with  $f(x) = x^m$ .

If we apply this result to  $(\underline{x})_n$  we get

$$\sum_{i=0}^{m} (-1)^{i} {m \choose i} (\underline{x+i})_{n} \in \mathbf{P}^{sep}.$$

The following corollary of Lemma 8.15 is a slight strengthening of Corollary 7.34.

**Lemma 8.37.** Suppose  $f \in P^{sep} \cap P^{pos}$ . For positive x

$$\left|\begin{smallmatrix}f(x+1)&f(x)\\f(x+2)&f(x+1)\end{smallmatrix}\right|>0$$

*Remark* 8.38. This does not necessarily hold for all  $x \in \mathbb{R}$ . If f = x(x + 1) then the determinant is 2(x + 1)(x + 2) which is negative for  $x \in (-2, -1)$ . However, if f = x(x + 2) then the determinant is positive for all x. More generally, it appears that if  $f(x) \leq f(x + 2)$  then the determinant is always positive. However, there are examples, e.g. x(x + 2)(x + 3.5), where f(x) and f(x + 2) don't interlace, yet the determinant is always positive.

**Example 8.39.** Here is a family of polynomials whose definition is similar to the Rodrigues definition of the Legendre polynomials (See Question 131). Define polynomials by the Rodrigues type formula

$$p_n = \Delta^n \prod_{i=1}^n (x^2 - i^2)$$

Since  $\prod_{i=1}^{n} (x^2 - i^2)$  is in  $\mathbf{P}^{sep}$ , the polynomial  $p_n$  is in  $\mathbf{P}^{sep}$ , and has degree n. Moreover, we claim that  $p_{n+1} \leq p_n$ . Since  $\Delta$  preserves interlacing in  $\mathbf{P}^{sep}$  it suffices to show that

$$\Delta \prod_{i=1}^{n+1} (x^2 - i^2) \le \prod_{i=1}^{n} (x^2 - i^2)$$

Now the roots of the left hand side are seen to be

$$-n, -n+1, \cdots, -2, -1, -.5, 1, 2, \cdots, n$$

and the roots of the right side are

$$-n, -n + 1, \cdots, -2, -1, 1, 2, \cdots, n$$

and so they interlace. The  $p_n$ 's are *not* orthogonal polynomials since they do not satisfy a three term recurrence. However, the  $p_n$  appear to satisfy a recurrence involving only three terms:

$$p_{n+2} = (n+2)^2 p_{n+1} + x(2n+3)p_{n+1} + (n+1)(n+2) \mathbb{A}p_{n+1} + x(2n+3) \mathbb{A}p_{n+1} - 2(2n+3)(n+1)^3 \mathbb{A}p_n$$

If we write a polynomial in the interpolation basis, then there is a simple estimate on the separation of roots. One consequence is that if  $\delta(f) > 0$  then  $\delta(f') > \delta(f)$ .

**Lemma 8.40.** Suppose that  $f = (x - a_1) \dots (x - a_n)$  where  $a_1 < \dots < a_n$  and that

$$g(x) = bf(x) + \sum_{i} b_{i} \frac{f(x)}{x - a_{i}}$$

where  $b \ge 0$ . If  $b_1 \ge b_2 \ge \cdots \ge b_n > 0$  then  $\delta(g) > \delta(f)$ .

*Proof.* Choose consecutive roots  $a_{j-1}, a_j, a_{j+1}$ , and points  $x_1, x_2$  satisfying  $a_{j-1} < x_1 < a_j < x_2 < a_{j+1}$ . We will show that  $\frac{g}{f}(x_2) - \frac{g}{f}(x_1) > 0$  if  $|x_1 - x_2| \leq \delta(f)$ . It follows that for  $|x_1 - x_2| \leq \delta(f)$ ,  $x_1$  and  $x_2$  can not be adjacent roots of g, so  $\delta(g) > \delta(f)$ .

We calculate

$$\begin{split} \frac{g}{f}(x_2) - \frac{g}{f}(x_1) &= \sum_{i=1}^n \frac{b_i}{x_2 - a_i} - \sum_{i=1}^n \frac{b_i}{x_1 - a_i} \\ &= \frac{b_1}{x_2 - a_1} + \frac{b_n}{a_n - x_1} + \sum_{i=1}^{n-1} \left( \frac{b_{i+1}}{x_2 - a_{i+1}} - \frac{b_i}{x_1 - a_i} \right) \end{split}$$

Now since  $b_i \ge b_{i+1}$ , we have

$$\geq \frac{b_1}{x_2 - a_1} + \frac{b_n}{a_n - x_1} + \sum_{i=1}^{n-1} b_{i+1} \left( \frac{1}{x_2 - a_{i+1}} - \frac{1}{x_1 - a_i} \right)$$
$$= \frac{b_1}{x_2 - a_1} + \frac{b_n}{a_n - x_1} + \sum_{i=1}^{n-1} b_{i+1} \frac{(a_{i+1} - a_i) - (x_2 - x_1)}{(x_2 - a_{i+1})(x_1 - a_i)}$$

The first two terms are positive; the numerator is positive since  $x_2 - x_1 \le \delta \le a_{i+1} - a_i$ , and the denominator is positive since  $x_1$  and  $x_2$  lie in consecutive intervals determined by the roots of f.

# 8.8 A multiplication in P<sup>sep</sup>

The product of two polynomials in  $\mathbf{P}^{sep}$  is not in  $\mathbf{P}^{sep}$ . We can remedy this situation by using a different multiplication. We define the bilinear transformation  $f \otimes g$  on a basis:

$$(\underline{\mathbf{x}})_{\mathbf{n}} \otimes (\underline{\mathbf{x}})_{\mathbf{m}} = (\underline{\mathbf{x}})_{\mathbf{n}+\mathbf{m}}$$

 $\otimes$  is a diamond product - see § 6.6.11.  $\Delta$  and  $\otimes$  interact as expected.

**Lemma 8.41.**  $\Delta(f \otimes g) = \Delta f \otimes g + f \otimes \Delta g$ 

*Proof.* We only need to check it on a basis.

$$\begin{array}{l} \Delta((\underline{x})_{n}\otimes(\underline{x})_{m}) = \Delta(\underline{x})_{n+m} = (n+m)(\underline{x})_{n+m-1} = \\ n(\underline{x})_{n-1}\otimes(\underline{x})_{m} + m(\underline{x})_{n}\otimes(\underline{x})_{m-1} \end{array}$$

_		
-	-	

It follows by induction that

$$\Delta(f_1\otimes\cdots\otimes f_n)=\sum_0^n f_1\otimes\cdots\otimes\Delta f_i\otimes\cdots\otimes f_n$$

and in particular the difference of a product of linear terms is

$$\Delta((\mathbf{x}-\mathbf{a}_1)\otimes\cdots\otimes(\mathbf{x}-\mathbf{a}_n))=\sum_{0}^{n}(\mathbf{x}-\mathbf{a}_1)\otimes\cdots\otimes(\widehat{\mathbf{x}-\mathbf{a}_i})\otimes\cdots\otimes(\mathbf{x}-\mathbf{a}_n)$$

where the hat means "omit".

Every polynomial in  $\mathbf{P}^{pos}$  can be factored in terms of  $\otimes$ .

**Lemma 8.42.** If  $f \in P^{pos}(n)$  is monic then there are positive  $a_1, \ldots, a_n$  such that

$$f(\mathbf{x}) = (\mathbf{x} + \mathbf{a}_1) \otimes (\mathbf{x} + \mathbf{a}_2) \otimes \cdots \otimes (\mathbf{x} + \mathbf{a}_n).$$

*Proof.* This follows from Proposition 6.56 since the linear transformation  $T: (\underline{x})_n \mapsto x^n$  maps  $\mathbf{P}^{pos}$  to itself.

*Remark* 8.43. Here's an example of the Lemma:

$$(x+3)(x+4) = (x+2) \otimes (x+6)$$

The Lemma shows that the product of certain terms with positive coefficients is in **P**; there are products of linear terms with positive coefficients that are not in **P**:

$$(\mathbf{x}+1)\otimes(\mathbf{x}+2)=(\mathbf{x}+1-\mathbf{i})(\mathbf{x}+1+\mathbf{i})$$

If  $f \in \mathbf{P}^{alt}$  then there might be no such representation. For example, the following is a unique representation:

$$(\mathbf{x} - 1 - \mathbf{i}) \otimes (\mathbf{x} - 1 + \mathbf{i}) = (\mathbf{x} - 1)(\mathbf{x} - 2)$$

**Lemma 8.44.** If  $f \in P^{sep} \cap P^{alt}$  and  $\beta \ge 0$  then

$$(\mathbf{x} - \mathbf{\beta}) \otimes \mathbf{f} \in \mathbf{P}^{sep} \cap \mathbf{P}^{alt}.$$

*Proof.* Let  $T(f) = (x - \beta) \otimes f$ . Since

$$\mathsf{T}((\underline{x})_{\mathfrak{n}}) = (\underline{x})_{\mathfrak{n}+1} - \beta(\underline{x})_{\mathfrak{n}} = (\mathfrak{x}\mathbb{A}^{-1} - \beta)(\underline{x})_{\mathfrak{n}}$$

we see that  $T(f) = (x \mathbb{A}^{-1} - \beta)f$ . Lemma 8.23 implies that  $T(f) \in \mathbf{P}^{sep} \cap \mathbf{P}^{alt}$ .  $\Box$ 

By iterating the Lemma, we get:

**Corollary 8.45.** If  $a_1, \ldots, a_n$  are positive then

$$(\mathbf{x} - \mathbf{a}_1) \otimes (\mathbf{x} - \mathbf{a}_2) \otimes \cdots \otimes (\mathbf{x} - \mathbf{a}_n) \in \mathbf{P}^{sep} \cap \mathbf{P}^{alt}.$$

We have seen that the following transformation preserved  $\mathbf{P}^{alt}$ .

**Corollary 8.46.** The transformation  $T: (\underline{x})_n \mapsto (\underline{x})_{n+1}$  maps

$$P^{sep} \cap P^{alt} \longrightarrow P^{sep} \cap P^{alt}.$$

*Proof.* Since  $T(f) = x \otimes f$  we can apply Lemma 8.44.

We would like to show that  $\otimes$  preserves  $\mathbf{P}^{sep} \cap \mathbf{P}^{alt}$ . See Question 90. This follows easily from Question 76, that the map  $x^n \longrightarrow \langle \underline{x} \rangle_n$  is a bijection between  $\mathbf{P}^{pos}$  and  $\mathbf{P}^{sep} \cap \mathbf{P}^{pos}$ .

Π

# **8.9** The analytic closure of **P**<sup>sep</sup>

The analytic closure  $\widehat{\mathbf{P}^{sep}}$  is the uniform closure on compact subsets of  $\mathbf{P}^{sep}$ . Obviously  $\widehat{\mathbf{P}^{sep}} \subsetneq \widehat{\mathbf{P}}$ . Since  $\mathbf{P}^{sep}$  is closed under differentiation and difference, so is  $\widehat{\mathbf{P}^{sep}}$ .

**Lemma 8.47.** The following functions are in  $\widehat{P^{sep}}$ :

- 1.  $\sin \pi x$ ,  $\cos \pi x$ .
- 2.  $e^{\alpha x}$
- 3.  $e^{\alpha x} f(x)$  for any  $f \in \mathbf{P}^{sep}$ .

 $e^{-\alpha x^2}$  is not in  $\widehat{P^{sep}}$  for any  $\alpha$ .

Proof. From the infinite product

$$\sin \pi x = \pi x \prod_{k=1}^{\infty} \left( 1 - \frac{x^2}{k^2} \right)$$

we see that the polynomials

$$\pi x \prod_{k=1}^n \left(1 - \frac{x^2}{k^2}\right)$$

converge to  $\sin \pi x$  and are in  $\mathbf{P}^{sep}$  since the roots are  $-n, -n+1, \dots, n-1, n$ . A similar argument applies to  $\cos \pi x$ .

Next, we show that for  $\alpha > 0$  we can find polynomials  $p_n$  such that

- 1.  $p_n \in \mathbf{P}^{sep}$
- 2. For every r there is an N such that n > N implies that the roots of  $p_n$  have absolute value at least r.
- 3.  $\lim p_n = e^{\alpha x}$

We start with the identity, valid for positive  $\beta$ :

$$(\beta+1)^{x} = \lim_{n \to \infty} \prod_{k=1}^{n} \left( 1 + \frac{\beta x}{n+\beta k} \right)$$
(8.9.1)

Let  $p_n$  be the product. The roots of  $p_n$  are  $-(n/\beta) - n, \ldots, -(n/\beta) - 1$  so  $p_n \in \mathbf{P}^{sep}$ . This shows that (1) and (2) are satisfied. If we choose positive  $\beta$  such that  $\beta + 1 = e^{\alpha}$  then  $p_n$  also satisfies (3). Thus,  $e^{\alpha x} \in \widehat{\mathbf{P}^{sep}}$  for positive  $\alpha$ . Since  $g(x) \in \mathbf{P}^{sep}$  if and only if  $g(-x) \in \mathbf{P}^{sep}$ , it follows that  $e^{\alpha x} \in \widehat{\mathbf{P}^{sep}}$  for all  $\alpha$ .

If  $f \in \mathbf{P}^{sep}$  then we can find an N such that n > N implies that  $p_n(x)f(x) \in \mathbf{P}^{sep}$ .  $\mathbf{P}^{sep}$ . Taking limits show that  $e^{\alpha x}f(x) \in \widehat{\mathbf{P}^{sep}}$ .

Finally, if  $e^{-\alpha x^2}$  belongs to  $\widehat{\mathbf{P}^{sep}}$ , then so do all its derivatives. The Rodrigues' formula for Hermite polynomials (7.8.1) implies that the Hermite polynomials  $H_n(x/\sqrt{\alpha})$  would be in  $\mathbf{P}^{sep}$ . This means that all roots of  $H_n(x)$  would be separated by at least  $\sqrt{\alpha}$ . However, from [168](6.31.21) we see that the minimum distance between roots of  $H_n$  goes to zero as n goes to infinity.

Alternatively, if  $e^{-\alpha x^2} \in \widehat{\mathbf{P}^{sep}}$  and  $f \in \mathbf{P}^{sep}$  then differentiating  $e^{-\alpha x^2} f$  and dividing out by the exponential factor would imply that  $f' - 2\alpha x f \in \mathbf{P}^{sep}$ . But if f = x(x+1)(x+2) then  $\delta(f' - 2xf) = .95$ .

The following consequence is due to Riesz [165].

**Corollary 8.48.** *If*  $f \in P^{sep}$  *and*  $\alpha \in \mathbb{R}$  *then*  $\alpha f + f' \in P^{sep}$ *.* 

*Proof.* Since  $e^{\alpha x} f(x) \in \mathbf{P}^{sep}$  we differentiate and find that  $(\alpha f + f')e^{\alpha x} \in \mathbf{P}^{sep}$ . The conclusion follows.

**Corollary 8.49.** If  $g(x) \in P$  and  $f \in P^{sep}$  then  $g(D)f \in P^{sep}$ .

The next result is not an independent proof of Lemma 8.5 since that Lemma was used to show that  $\mathbf{P}^{sep}$  is closed under differences.

**Corollary 8.50.** If  $f \in \mathbf{P}^{sep}$  and  $\beta > 0$  then  $\beta f(x+1) - f(x) \in \mathbf{P}^{sep}$ .

*Proof.* Note that  $\Delta(e^{\alpha x}f) = e^{\alpha x} (e^{\alpha}f(x+1) - f(x))$ .

As long as function in  $\widehat{\mathbf{P}}$  has no factor of  $e^{-\alpha x^2}$ , and all roots are at least one apart then f is in  $\widehat{\mathbf{P}^{sep}}$ .

**Lemma 8.51.** Assume that  $a_{n+1} - a_n \ge 1$  for n = 1, 2, ...

1. If 
$$f(x) = e^{\alpha x} \prod_{i=1}^{\infty} \left(1 - \frac{x}{a_i}\right) \in \widehat{P}$$
 then  $f \in \widehat{P^{sep}}$ .  
2. If  $f(x) = e^{\alpha x} \prod_{i=1}^{\infty} \left[ \left(1 - \frac{x}{a_i}\right) e^{-x/a_i} \right] \in \widehat{P}$  then  $f \in \widehat{P^{sep}}$ .  
3.  $1/\Gamma(x) \in \widehat{P^{sep}}$ .

*Proof.* The assumptions in (1) and (2) guarantee that the product converges. Part (1) follows from Lemma 8.47(3). From the proof of this lemma we let  $p(a, r) \in \mathbf{P}^{sep}$  be a polynomial whose roots have absolute value at least |r|, and  $\lim_{r\to\infty} p(a, r) = e^{ax}$ . Consider the polynomial

$$q_{n}(x) = p(\alpha, r_{0}) \prod_{i=1}^{n} \left( \left( 1 - \frac{x}{a_{i}} \right) p(-1/a_{i}, r_{i}) \right)$$

We can choose the  $r_i$  sequentially so that  $q_n \in \mathbf{P}^{sep}$ . Taking limits yields (2). The Weierstrass infinite product representation

$$\frac{1}{\Gamma(x)} = x e^{\gamma x} \prod_{n=1}^{\infty} \left[ \left( 1 + \frac{x}{n} \right) e^{-x/n} \right]$$

shows that  $\frac{1}{\Gamma(\mathbf{x})} \in \widehat{\mathbf{P}}$ . Since the roots are at negative integers, part (2) implies that  $1/\Gamma(\mathbf{x}) \in \widehat{\mathbf{P}^{sep}}$ .

Since  $\frac{1}{\Gamma(x)}$  is in  $\widehat{\mathbf{P}^{sep}}$ , so are all the differences  $\Delta^n \frac{1}{\Gamma(x)}$ . We can find a recursive formula for these differences. If we write

$$\Delta^{n} \frac{1}{\Gamma(\mathbf{x})} = \frac{f_{n}(\mathbf{x})}{\langle \underline{\mathbf{x}} \rangle_{n} \Gamma(\mathbf{x})}$$

then

$$\Delta^{n+1} \frac{1}{\Gamma(x)} = \frac{f_n(x+1)}{\langle \underline{x+1} \rangle_n \Gamma(x+1)} - \frac{f_n(x)}{\langle \underline{x} \rangle_n \Gamma(x)}$$
$$= \frac{f_n(x+1) - (x+n)f_n(x)}{\langle \underline{x} \rangle_{n+1} \Gamma(x)}$$

and consequently the numerators satisfy the recurrence

$$f_{n+1}(x) = f_n(x+1) - (x+n)f_n(x)$$

Since  $f_1(x) = 1 - x$ , the next lemma shows that all of the numerators are in  $\mathbf{P}^{sep}$ .

**Lemma 8.52.** Suppose that  $g_1(x) \in \mathbf{P}^{(0,\infty)} \cap \mathbf{P}^{sep}$  and the sequence  $\{g_n\}$  is defined by the recurrence

$$g_{n+1}(x) = g_n(x+1) - (x+n)g_n(x)$$

then  $g_n(\mathbf{x}) \in \mathbf{P}^{(-n+1,\infty)} \cap \mathbf{P}^{sep}$ .

*Proof.* The function  $\frac{1}{\langle \underline{x} \rangle_{n+1} \Gamma(x)}$  is in  $\widehat{\mathbf{P}^{sep}}$ , as can be seen by removing initial factors from the infinite product; its zeros are  $-n - 1, -n - 2, \cdots$ . Since  $\Delta^n \frac{g_1}{\Gamma(x)} = \frac{g_{n+1}}{\langle \underline{x} \rangle_{n+1} \Gamma(x)}$  is in  $\widehat{\mathbf{P}^{sep}}$ , the roots of  $g_{n+1}$  must lie in  $(-n, \infty)$ .

If we consider  $f/\Gamma$  we get

**Lemma 8.53.** If  $f \in P^{(1,\infty)} \cap P^{sep}$  then  $f(x+1) - xf(x) \in P^{(0,\infty)} \cap P^{sep}$ .

*Proof.* The largest root of  $1/\Gamma(x)$  is 0, so  $f(x)/\Gamma(x) \in \mathbf{P}^{sep}$ . Taking differences yields

$$\frac{f(x+1)}{\Gamma(x+1)} - \frac{f(x)}{\Gamma(x)}, = \frac{f(x+1) - xf(x)}{x\Gamma(x)} \in \widehat{\mathbf{P}^{sep}}$$

and the numerator is in  $\mathbf{P}^{sep}$ . The largest zero of  $x\Gamma(x)$  is -1, so the conclusion follows.

Here's a result that we get from using  $(\underline{x})_n$  in place of  $\frac{1}{\Gamma(x)}$ .

**Lemma 8.54.** If  $f \in P^{sep} \cap P^{(n,\infty)}$  then  $nf + (x+1)\Delta f \in P^{sep} \cap P^{(n-1,\infty)}$ .

*Proof.* Since the largest root of  $(\underline{x})_n$  is n - 1 it follows that  $(\underline{x})_n f \in \mathbf{P}^{sep}$ . Now taking the difference preserves  $\mathbf{P}^{sep}$ , so

$$\Delta(\underline{\mathbf{x}})_{n} \mathbf{f} = \mathbf{f} \Delta(\underline{\mathbf{x}})_{n} + (\Delta \mathbf{f}) \mathbb{A}(\underline{\mathbf{x}})_{n}$$
$$= (\underline{\mathbf{x}})_{n-1} (\mathbf{n}\mathbf{f} + (\mathbf{x}+1)\Delta \mathbf{f})$$

is in  $\mathbf{P}^{sep}$ . The result follows since factors of polynomials in  $\mathbf{P}^{sep}$  are in  $\mathbf{P}^{sep}$ .

Motivated by the analogy of  $\langle \underline{x} \rangle_n / n!$  and the exponential, we define a Laguerre like polynomial, and show that consecutive ones interlace.

**Lemma 8.55.** If  $f_n(x) = \Delta^n \langle \underline{x+1} \rangle_n (\underline{x-1})_n$  then  $f_n \in P^{sep}$  and  $f_n \leq f_{n-1}$ .

*Proof.* If we let  $p_n(x) = \langle \underline{x+1} \rangle_n (\underline{x-1})_n$  then

$$\Delta p_{n}(x) = n \langle \underline{x+2} \rangle_{n-1} (\underline{x-1})_{n-1} (1+2x)_{n-1} (1+2x)_{$$

Observe that that  $\Delta p_n \leq p_{n-1}$ . Since  $p_n \in \mathbf{P}^{sep}$  the conclusion follows by applying  $\Delta^{n-1}$ .

**Lemma 8.56.** If 
$$T(f) = f(x + \iota) + f(x - \iota)$$
 then  $T: \mathbb{P}^{sep} \longrightarrow \mathbb{P}^{sep}$ . [180]

*Proof.* Observe that  $T(f) = 2\cos(D)f$ , so that T maps  $\mathbf{P} \longrightarrow \mathbf{P}$  and preserves interlacing. Since T commutes with  $\mathbb{A}$ , the result follows from Lemma 8.25.

# 8.10 Integrating families of polynomials in $P^{sep}$ and $P^{\mathbb{A}}$

We now apply the integration results of §3.3.6 to  $\mathbf{P}^{sep}$ . Since polynomials in  $\mathbf{P}^{sep}$  are exactly those polynomials whose roots are separated by at least one,  $f \in \mathbf{P}^{sep}$  iff  $\delta(f) \ge 1$ . We will see more applications of integration when we consider the generalization of  $\mathbf{P}^{sep}$  to two variables.

One useful way of constructing locally interlacing families is to start with  $f(x) \in \mathbf{P}^{sep}$ , and form the family  $\{f(x + t)\}$ . The interval of interlacing is at least one.

Here is a simple application. Note that the hypothesis  $f' \in \mathbf{P}^{sep}$  doesn't imply  $f \in \mathbf{P}$ , yet we can conclude that  $\Delta f \in \mathbf{P}$ .

**Corollary 8.57.** Suppose that w(x) is positive on (0, 1). The mapping

$$f \mapsto \int_0^1 f(x+t)w(t) dt \qquad (8.10.1)$$

defines a linear transformation  $P^{\text{sep}} \longrightarrow P$ .

**Corollary 8.58.** If  $f' \in P^{sep}$  then  $\Delta f \in P$  where  $\Delta f = f(x + 1) - f(x)$ . *Proof.* Choose w(t) = 1, a = 0, b = 1, and  $f_t(x) = f'(x+t)$  in Proposition 3.37.

*Remark* 8.59. Integration is an example of a linear transformation  $T: \mathbf{P}^{sep} \longrightarrow \mathbf{P}$  that does not preserve interlacing. For example

$$f(x) = x(x+1) g(x) = x$$
  

$$T(f) = \int_0^1 f(x+t) dt T(g) = \int_0^1 g(x+t) dt$$
  

$$= (x+.59)(x+1.4) x+.5$$

Thus,  $f \leq g$  yet Tf and Tg don't interlace.

The following result is convolution of polynomials in **P**<sup>sep</sup>.

**Lemma 8.60.** *If*  $f, g \in P^{sep}$  *then* 

$$\int_0^1 f(x+t)g(x-t) \, dt \, \in \boldsymbol{P}$$

*Proof.* If we choose  $0 < t_1 < \cdots < t_n$  then let  $f_i(x) = f(x+t_i)$  and  $g_{n+1-i}(x) = g(x - t_i)$ . By Lemma 3.16 we know that  $\sum f_i g_{n+1-i} \in \mathbf{P}$ . Taking the limit shows the integral is in  $\mathbf{P}$ .

Note: the diagram in the proof of Lemma 3.16 is essentially the graph of f(x+t)g(x-t). Consideration of this diagram shows that we can find f, g so that if h is their convolution then  $\delta(h)$  is as small as desired.

Since we only need separation in one variable, we phrase the next result in terms of mutually interlacing families.

**Corollary 8.61.** *If*  $\{f_t\}$  *and*  $\{g_t\}$  *are locally interlacing families on*  $\mathbb{R}$  *with*  $\rho(f) \ge 1$  *and*  $\rho(g) \ge 1$ *, and we define* 

$$\Phi = \int_0^1 \int_0^{1-t} f_t(x)g_s(x) \, ds \, dt$$

*then*  $\Phi \in \mathbf{P}$ *.* 

*Proof.* If we define  $h_t = \int_0^t g_s(x) dt$  then by Lemma 3.43  $\{h_t\}$  is a locally interlacing family with  $\rho(h) \ge 1$ . Since  $\Phi = \int_0^1 f_t(x)h_{1-t}(x) dt$  we see that  $\Phi$  is a convolution of mutually interlacing polynomials. ( $\Phi$  is generally not in  $\mathbf{P}^{\text{sep}}$ .)

**Corollary 8.62.** If  $f \in P^{sep}$  then  $\int_0^1 f(x+t)e^{-t} dt \in P$ .

*Proof.* If we choose  $g(x) = e^{-x}$  which is in  $\widehat{\mathbf{P}^{sep}}$ , then we can apply the lemma to find that

$$\int_0^1 f(x+t)e^{x-t} dt = e^x \int_0^1 f(x+t)e^{-t} dt \in \mathbf{P}.$$

If we multiply by  $e^{-x}$  we get the conclusion.

We now to assume that  $\mathbb{A}x = qx$  where  $q \ge 1$ . We can integrate polynomials in  $\mathbf{P}^{\mathbb{A}}$  since they determine sequences of interlacing polynomials. Assume that  $f(x) \in \mathbf{P}^{\mathbb{A}}$ . It's clear that

$$f(x) \leq f(tx) \leq f(qx) \text{ if } 1 \leq t \leq q \tag{8.10.2}$$

Consequently, we have an interlacing family  $\{f(tx)\}$ . Applying the results of 3.3.6 we conclude that

**Corollary 8.63.** Suppose that w(x) is positive on (1, q). The mapping

$$f \mapsto \int_{1}^{q} f(tx)w(t) dt \qquad (8.10.3)$$

*defines a linear transformation*  $P^{\mathbb{A}} \longrightarrow P$ *.* 

**Corollary 8.64.** If  $f' \in \mathbf{P}^{\mathbb{A}}$  then  $D_q(f) \in \mathbf{P}$ .

Proof. Immediate from the computation

$$\int_{1}^{q} f'(tx) dt = (f(qx) - f(x))/x = (q-1) D_{q}(f)$$

We also have convolution. If  $f(x)\in {I\!\!P}^{\mathbb A}$  then replacing x by x/q in (8.10.2) shows

$$f(x/q) \leq f(x/t) \leq f(x) \text{ if } 1 \leq t \leq q \tag{8.10.4}$$

Equations (8.10.2) and (8.10.4) are reversed if 0 < q < 1. We can now apply Lemma 8.60 to conclude

**Lemma 8.65.** If  $f, g \in \mathbf{P}^{\mathbb{A}}$  where  $\mathbb{A}x = qx$ , and w(t) is positive on (1, q) if q > 1 and on (q, 1) if 0 < q < 1 then

$$\int_1^q w(t)f(xt)g(x/t)\,dt\in \boldsymbol{P}$$

# 8.11 Determinants and $P^{\mathbb{A}}$

In this section we look at connections between  $\mathbf{P}^{\mathbb{A}}$  and polynomials of the form  $|\mathbf{x}D + C|$  with symmetric C and diagonal D. We show that there is an analog of the differentiation formula (8.2.4) where the factors are replaced by determinants of principle submatrices. The rest of the section is concerned only with  $\mathbb{A}x = x + 1$ .

We assume that D is an n by n diagonal matrix with diagonal entries  $a_i$ , and  $C = (c_{ij})$  is a symmetric matrix. We are interested in the matrix M = xD + C. If M[i] is the ith principle submatrix, then

$$\frac{\mathrm{d}}{\mathrm{d}x} \det(\mathsf{M}) = a_1 \det(\mathsf{M}[1]) + \dots + a_n \det(\mathsf{M}[n]) \tag{8.11.1}$$

The formula for the affine derivative is similar:

**Lemma 8.66.** Assume all  $a_i$  are positive. If  $M_{[r]}$  is constructed from M[r] by applying A to the first r - 1 diagonal elements of M[r] then

$$D_{\mathbb{A}} \det(M) = a_1 \det(M_{[1]}) + \dots + a_n \det(M_{[n]})$$
(8.11.2)

For example,

$$M = \begin{pmatrix} a_{1}x + c_{11} & c_{12} & c_{13} \\ c_{12} & a_{2}x + c_{22} & c_{23} \\ c_{13} & c_{23} & a_{3}x + c_{33} \end{pmatrix}$$
$$M_{[1]} = \begin{pmatrix} a_{2}x + c_{22} & c_{23} \\ c_{23} & a_{3}x + c_{33} \end{pmatrix}$$
$$M_{[2]} = \begin{pmatrix} \mathbb{A}(a_{1}x + c_{11}) & c_{13} \\ c_{13} & a_{3}x + c_{33} \end{pmatrix}$$
$$M_{[3]} = \begin{pmatrix} \mathbb{A}(a_{1}x + c_{11}) & c_{12} \\ c_{12} & \mathbb{A}(a_{2}x + c_{22}) \end{pmatrix}$$

$$\mathsf{D}_{\mathbb{A}} \det(\mathsf{M}) = a_1 \det(\mathsf{M}_{[1]}) + a_2 \det(\mathsf{M}_{[2]}) + a_3 \det(\mathsf{M}_{[3]})$$

*Proof (of 8.66).* Note that the term  $a_i det(m_{[i]})$  is the determinant of the n by n matrix formed from M by setting all elements of the i-th row and column to zero, except for the main diagonal element, which is  $a_i$ . Since all terms of the equation are now n by n matrices, we can multiply by  $D^{-1/2}$  on each side, and so we may assume that all  $a_i$  are equal to 1. We now apply the following lemma, with y = Ax, to finish the proof.

**Lemma 8.67.** If C is an n by n symmetric matrix, and  $D_k$  is the n by n diagonal matrix whose first k diagonal elements are y and the remaining ones are x then

$$\frac{\det(C + xI) - \det(C + yI)}{x - y} = \\ \det(C + D_1)[1] + \det(C + D_2)[2] + \dots + \det(C + D_n)[n] \quad (8.11.3)$$
Notice that in the special case that M is a diagonal matrix the formula (8.11.2) reduces to (8.2.4). The limit of (8.11.3) as  $y \rightarrow x$  is (8.11.1).

For the remainder of this section we restrict ourselves to Ax = x + 1 and  $\mathbf{P}^{sep}$ . If  $f(x) = \prod (x - r_i)$  then the polynomials  $f/(x - r_i)$  are mutually interlacing. It is easy to find examples where the determinants of the principle submatrices of |xI + C| are not mutually interlacing. However, if |xI + C| is in  $\mathbf{P}^{sep}$  then we can find a family of mutually interlacing polynomials that generalize the  $f_{[k]}$  of (8.3.5).

Let 
$$M = xI + C$$
 and write  $M = \begin{vmatrix} x + c_{11} & v \\ v^{t} & M[1] \end{vmatrix}$ . Then

$$\begin{vmatrix} \mathbb{A}(\mathbf{x} + \mathbf{c}_{11}) & \mathbf{v} \\ \mathbf{v}^{\mathsf{t}} & \mathbf{M}[1] \end{vmatrix} = \begin{vmatrix} \mathbf{x} + \mathbf{c}_{11} & \mathbf{v} \\ \mathbf{v}^{\mathsf{t}} & \mathbf{M}[1] \end{vmatrix} + \begin{vmatrix} \mathbf{1} & \mathbf{0} \\ \mathbf{v}^{\mathsf{t}} & \mathbf{M}[1] \end{vmatrix}$$
$$= |\mathbf{M}| + |\mathbf{M}[1]|$$

Since  $|M| \leq |M[1]|$  we see that

$$|\mathsf{M}| \underline{\ll} \begin{vmatrix} \mathbb{A}(\mathsf{x} + \mathsf{c}_{11}) & \mathsf{v} \\ \mathsf{v}^{\mathsf{t}} & \mathsf{M}[1] \end{vmatrix}$$

If we let  $M_{[r]}$  be the result of applying  $\mathbb{A}$  to the first r diagonal elements of M then inductively we see that

$$|M| \underline{\ll} |M_{[1]}| \underline{\ll} |M_{[2]}| \underline{\ll} \cdots \underline{\ll} |M_{[n]}| = \mathbb{A} |M|$$

Thus, if  $|M| \in \mathbf{P}^{sep}$  then  $|M| \leq \mathbb{A}|M|$  and consequently the following polynomials are mutually interlacing

$$\{|\mathsf{M}|, |\mathsf{M}_{[1]}|, |\mathsf{M}_{[2]}|, \cdots, |\mathsf{M}_{[n-1]}|, \mathbb{A}|\mathsf{M}|\}$$

Every factor of a polynomial in  $\mathbf{P}^{sep}$  is in  $\mathbf{P}^{sep}$ , but the analogous property does not hold for principle submatrices. For instance, the matrix M below has the property that  $\delta(|\mathbf{M}[k]|) < \delta(|\mathbf{M}|)$  for k = 1, 2, 3, 4, 5.

$$\begin{pmatrix} x+2 & 4 & 6 & 6 & 5 \\ 4 & x+2 & 5 & 8 & 6 \\ 6 & 5 & x & 7 & 6 \\ 6 & 8 & 7 & x+2 & 4 \\ 5 & 6 & 6 & 4 & x \end{pmatrix}$$

However, the roots are always distinct.

**Lemma 8.68.** If M = xI + C is a matrix whose determinant has all distinct roots, then the determinants of all principle submatrices have all distinct roots.

*Proof.* The determinant of a principle submatrix interlaces the determinant of M.

# 8.12 The case Ax = qx with q > 1

In this section we consider the affine transformation  $\mathbb{A}f(x) = f(qx)$  where we first assume that that q > 1. We know that  $f \leq \mathbb{A}f$  implies that  $f \in \mathbf{P}^{\alpha lt}$ . There are simple conditions on the roots that characterize such polynomials. If the roots of f are  $\{a_i\}$  then the roots of  $\mathbb{A}f$  are  $\{a_i/q\}$ . If  $f \in \mathbf{P}^{\alpha lt}$  then since q > 1 the ordering of the roots is

$$\mathfrak{a}_1/\mathfrak{q} < \mathfrak{a}_1 < \mathfrak{a}_2/\mathfrak{q} < \mathfrak{a}_2 < \cdots$$

The ratio of consecutive roots is at least q. Conversely, if  $f \in \mathbf{P}^{alt}$  and the ratio of consecutive roots is at least q then  $f \underline{\ll} \mathbb{A}f$ .

The affine derivative is called the q-derivative, and is given in (8.2.1). The next lemma specializes the results in Section 8.2.

**Corollary 8.69.** Suppose Ax = qx where q > 1. The following are true:

- 1. If  $f \in \mathbf{P}^{\mathbb{A}}$  then  $f \in \mathbf{P}^{\operatorname{alt}}$ .
- 2. If  $f \in \mathbf{P}^{\mathbb{A}}$  then  $f' \in \mathbf{P}^{\mathbb{A}}$ .
- 3. If  $f \in \mathbf{P}^{\mathbb{A}}$  then  $D_q f \in \mathbf{P}^{\mathbb{A}}$ .
- 4. If  $f \leq g \in \mathbf{P}^{\mathbb{A}}$  then  $D_q f \leq D_q g$ .
- 5. If  $g \in \mathbf{P}^{alt}$  and  $f \in \mathbf{P}^{\mathbb{A}}$  then  $g(\mathbb{A})f \in \mathbf{P}^{\mathbb{A}}$ .
- 6. If  $g \in \mathbf{P}^{alt}$  and  $f \in \mathbf{P}^{\mathbb{A}}$  then  $g(\mathbf{D}_{q})f \in \mathbf{P}^{\mathbb{A}}$ .

For example, if we take  $g = (x-1)^n$  then the following polynomials are in  $\mathbf{P}^{\mathbb{A}}$ :

$$(\mathbb{A} - 1)^{n} f = \sum_{i=0}^{n} {n \choose i} f(q^{i}x)$$
$$(\mathsf{D}_{q} - 1)^{n} f = \sum_{i=0}^{n} {n \choose i} \mathsf{D}_{q}^{i} f(x)$$

In order to apply Lemma 8.14 we need the following formula for higher derivatives that is easily proved by induction:

$$\mathsf{D}_{q}^{\mathfrak{m}}\mathfrak{g}(x) = \frac{1}{(q-1)^{\mathfrak{m}}\mathfrak{q}^{\binom{\mathfrak{n}}{2}}\mathfrak{x}^{\mathfrak{m}}} \quad \sum_{\mathfrak{i}=0}^{\mathfrak{m}} (-1)^{\mathfrak{i}} \binom{\mathfrak{m}}{\mathfrak{i}}_{q} \mathfrak{q}^{\binom{\mathfrak{i}}{2}}\mathfrak{g}(\mathfrak{q}^{\mathfrak{i}}\mathfrak{x}).$$

 $\binom{n}{i}_{a}$  is the Gaussian binomial coefficient defined by

$$\binom{n}{i}_{q} = \frac{[n]!}{[k]! [n-k]!}$$

where as usual  $[n]! = [n][n-1] \cdots [1]$ .

**Corollary 8.70.** If  $g \in \mathbf{P}^{\mathbb{A}}$  and m is a positive integer then

$$\frac{1}{x^{\mathfrak{m}}} \quad \sum_{\mathfrak{i}=0}^{\mathfrak{m}} (-1)^{\mathfrak{i}} \binom{\mathfrak{m}}{\mathfrak{i}}_{\mathfrak{q}} \mathfrak{q}^{\binom{\mathfrak{i}}{2}} \mathfrak{g}(\mathfrak{q}^{\mathfrak{i}} x) \in \boldsymbol{P}^{\mathbb{A}}.$$

Lemma 8.15 implies that

**Lemma 8.71.** Suppose  $f \in \mathbf{P}^{\mathbb{A}} \cap \mathbf{P}^{\text{pos}}$  and q > 1. For all positive x

$$\left|\begin{array}{c}f(qx) & f(x)\\f(q^2x) & f(qx)\end{array}\right| > 0$$

8.13 The case Ax = qx with 0 < q < 1

The cases q > 1 and 0 < q < 1 are closely related. Let  $\mathbb{A}_q(x) = qx$ , and  $\mathbb{A}_{1/q}(x) = x/q$ . Consequently, if  $f \in \mathbf{P}$  then

$$f(x) \ll \mathbb{A}_q f(x) \iff f(-x) \ll \mathbb{A}_{1/q} f(-x)$$

The transformation  $x \mapsto -x$  is a bijection between  $\mathbf{P}^{\mathbb{A}}$  for q > 1 and  $\mathbf{P}^{\mathbb{A}}$  for 0 < 1/q < 1.

For the rest of this section we assume that 0 < q < 1. The fundamental polynomials in  ${\bf P}^{\mathbb{A}}$  are

$$(-x;q)_n = (1+x)(1+qx)(1+q^2x) + \cdots (1+q^{n-1}x)$$

which have roots

$$-q^{-(n-1)} < -q^{-(n-2)} < \dots < -q^{-1} < -1$$

Note that  $D_q(x;q)_n = [n](x;q)_{n-1}$ .

We can translate the results of Lemma 8.69:

**Corollary 8.72.** Suppose Ax = qx where 0 < q < 1. The following are true:

- 1. If  $f \in \mathbf{P}^{\mathbb{A}}$  then  $f \in \mathbf{P}^{\text{pos}}$ .
- 2. If  $f \in \mathbf{P}^{\mathbb{A}}$  then  $f' \in \mathbf{P}^{\mathbb{A}}$ .
- 3. If  $f \in \mathbf{P}^{\mathbb{A}}$  then  $D_{q}f \in \mathbf{P}^{\mathbb{A}}$ .
- 4. If  $f \leq g \in \mathbf{P}^{\mathbb{A}}$  then  $D_q f \leq D_q g$ .
- 5. If  $g \in \mathbf{P}^{pos}$  and  $f \in \mathbf{P}^{\mathbb{A}}$  then  $g(\mathbb{A})f \in \mathbf{P}^{\mathbb{A}}$ .
- 6. If  $g \in \mathbf{P}^{\text{pos}}$  and  $f \in \mathbf{P}^{\mathbb{A}}$  then  $g(\mathbf{D}_{q})f \in \mathbf{P}^{\mathbb{A}}$ .

Lemma 8.15 implies that

**Lemma 8.73.** Suppose  $f \in \mathbf{P}^{\mathbb{A}} \cap \mathbf{P}^{\text{pos}}$  and 0 < q < 1. For all positive x

$$\left|\begin{array}{c}f(qx) & f(x)\\f(q^2x) & f(qx)\end{array}\right| < 0$$

#### 8.14 The q-exponential

In this section we continue to assume that 0 < q < 1. We will use the q-exponential  $E_{\mathbb{A}}$  of (8.4.2) to derive properties of the q-Laguerre polynomial. We have already seen  $E_q(x)$  in (5.1.1).  $E_q(x)$  is an entire function if |q| < 1. Using the product representation it is easy to verify that  $D_q E_q(\alpha x) = \alpha \mathbb{A} E_q(\alpha x)$ .

From (5.1.1) we see that all the terms have positive coefficients, and the roots of the q-exponential  $\text{Exp}_q(x)$  are  $\{-q^{-n}\}_{n=1}^{\infty}$ , so clearly we have

$$E_q(x) \leq AE_q(x)$$

Thus,  $E_q$  is an entire function that is in the closure of  $\mathbf{P}^{\mathbb{A}}$ .

The Rodrigues' formula for Laguerre polynomials is

$$\mathcal{L}_{n}(\mathbf{x}) = \frac{1}{e^{-x}} \frac{1}{n!} \mathsf{D}^{n}(e^{-x} \mathbf{x}^{n})$$

We define the q-Laguerre polynomials by the formula

$$L_{n}^{\mathbb{A}}(x) = \frac{1}{\mathbb{A}^{n}\mathsf{E}_{q}(-x)} \mathsf{D}_{q}^{n}(\mathsf{E}_{q}(-x)(-x;q)_{n})$$

However, since  $E_q(-x) \notin \widehat{\mathbf{P}^{\mathbb{A}}}$ , it is by no means obvious that this is even a polynomial, let alone that it is in  $\mathbf{P}^{\mathbb{A}}$ . To this end, define

$$\mathbf{S}(\mathbf{n}) = \left\{ \mathbf{f} \cdot \mathbb{A}^{\mathbf{n}}(\mathsf{E}_{\mathbf{q}}(-\mathbf{x})) \mid \mathbf{f} \in \mathbf{P}^{\mathbb{A}} \right\}$$

We can control the action of  $D_q$  in S(n):

**Lemma 8.74.** For any non-negative integer n,  $D_q : S(n) \longrightarrow S(n+1)$ 

*Proof.* Apply the q-Leibnitz formula to the product:

$$\begin{split} \mathsf{D}_{\mathsf{q}}(\mathsf{f} \cdot \mathbb{A}^{\mathsf{n}}\mathsf{E}_{\mathsf{q}}(-\mathsf{x})) &= \ \mathsf{D}_{\mathsf{q}}\mathsf{f} \cdot \mathbb{A}(\mathbb{A}^{\mathsf{n}}\mathsf{E}_{\mathsf{q}}(-\mathsf{x})) + \mathsf{f} \cdot \ \mathsf{D}_{\mathsf{q}}(\mathbb{A}^{\mathsf{n}}\mathsf{E}_{\mathsf{q}}(-\mathsf{x})) \\ &= \ \mathsf{D}_{\mathsf{q}}\mathsf{f} \cdot \mathbb{A}^{\mathsf{n}+1}\mathsf{E}_{\mathsf{q}}(-\mathsf{x}) - \mathsf{q}^{\mathsf{n}}\mathsf{f} \cdot \mathbb{A}^{\mathsf{n}} \ \mathsf{D}_{\mathsf{q}}\mathsf{E}_{\mathsf{q}}(-\mathsf{x}) \\ &= (\ \mathsf{D}_{\mathsf{q}}\mathsf{f} - \mathsf{q}^{\mathsf{n}}\mathsf{f}) \cdot \mathbb{A}^{\mathsf{n}+1}\mathsf{E}_{\mathsf{q}}(-\mathsf{x}) \end{split}$$

Since  $D_q f - q^n f \in \mathbf{P}^{\mathbb{A}}$  by Lemma 8.5 it follows that  $D_q(f \cdot \mathbb{A}^n E_q(-x))$  is in S(n+1).

**Corollary 8.75.** The q-Laguerre polynomials are in  $P^{\mathbb{A}}$ .

*Proof.* Since  $(-x;q)_n \in \mathbf{P}^{\mathbb{A}}$ , it follows that  $(-x;q)_n \cdot E_q(-x)$  is in S(0). The Lemma above shows that  $\mathsf{D}_q^n([-x;q)_n \cdot E_q(-x)]$  is in S(n). This means that it has the form  $g \cdot \mathbb{A}^n E_q(-x)$  where g is in  $\mathbf{P}^{\mathbb{A}}$ , which proves the corollary.

244

There is a simple formula for the q-Laguerre polynomials.

$$L_n^{\mathbb{A}}(x) = \sum_{i=0}^n (-1)^i \, x^i \, \binom{n}{i} \, \frac{[n]!}{[i]!}$$

There are several q-analogs of the Hermite polynomials [103]. The variant that we use is motivated by the desire for a recursive formula similar to (7.8.6). We set  $H_0^q = 1$ ,  $H_1^q = x$  and

$$\mathsf{H}_{n+1}^{\mathsf{q}} = \mathsf{x} \, \mathsf{H}_{n}^{\mathsf{q}} - \mathsf{q}^{n} \, [n] \, \mathsf{H}_{n-1}^{\mathsf{q}}.$$

Since the coefficient of  $H_{n-1}^q$  is negative whenever q is positive, we see that  $H_n^q$  has all real roots for all positive q. From the definition it is easy to show by induction that

$$\mathsf{D}_{\mathsf{q}}\,\mathsf{H}_{\mathfrak{n}}^{\mathsf{q}} = [\mathfrak{n}]\,\mathsf{H}_{\mathfrak{n}-1}^{\mathsf{q}}$$

which is the desired analog of (7.8.6). If we combine these last two equations we find

$$H_n^q = (x - q^{n-1} D_q)(x - q^{n-2} D_q q) \dots (x - D_q)(1)$$

which is the q-analog of the identity  $H_n = (2x - D)^n(1)$ .

The q-hermite polynomials can not be in  $\mathbf{P}^{\mathbb{A}}$  since they have both positive and negative roots. However, they appear to be nearly so. See Question 83.

## 8.15 q-transformations

In this section we show that certain linear transformations associated to qseries map appropriate subsets of  $\mathbf{P}$  to  $\mathbf{P}$ . From earlier results specialized to the case at hand we have

**Corollary 8.76.** Let Ax = qx where q > 1.

- 1. The linear transformation  $x^n \mapsto q^{\binom{n}{2}} x^n$  maps  $P^{\text{alt}} \longrightarrow P^{\text{alt}} \cap P^{\mathbb{A}}$ .
- 2. The linear transformation  $x^n \mapsto (-x; q)_n$  maps  $P^{alt} \longrightarrow P^{(1,\infty)} \cap P^{\mathbb{A}}$ .

We next study the latter transformation and its inverse for all values of q. Define

$$T_{q}(x^{i}) = (x;q)_{i} = (1-x)(1-qx)\cdots(1-q^{i-1}x)$$
(8.15.1)

We will show the following:

**Theorem 8.77.** If  $T_q$  is defined in (8.15.1), then

- 1. If 0 < q < 1 then  $T_q : \boldsymbol{P}^{\mathbb{R} \setminus (0,1)} \longrightarrow \boldsymbol{P}^{\mathbb{R} \setminus (0,1)}$ .
- 2. If 0 < q < 1 then  $T_q^{-1} : \mathbf{P}^{(0,1)} \longrightarrow \mathbf{P}^{(0,1)}$ .
- 3. If 1 < q then  $T_q : \mathbf{P}^{(0,1)} \longrightarrow \mathbf{P}^{(0,1)}$ .

4. If 
$$1 < q$$
 then  $T_q^{-1} : \mathbf{P}^{\mathbb{R} \setminus (0,1)} \longrightarrow \mathbf{P}^{\mathbb{R} \setminus (0,1)}$ .

The idea is to discover the recursion relations satisfied by  $T_q$  and  $T_q^{-1}$ . We will show that  $T_{(1/q)} = T_q^{-1}$ , and this allows us to prove the results about  $T_q^{-1}$ . From the definition

$$\begin{split} \mathsf{T}_{\mathsf{q}}(x \cdot x^{n-1}) &= \mathsf{T}_{\mathsf{q}}(x^n) = (1-x)(1-\mathsf{q}x)(1-\mathsf{q}^2x) \cdots (1-\mathsf{q}^{n-1}x) \\ &= (1-x) \mathbb{A}\left((1-x)(1-\mathsf{q}x) \cdots (1-\mathsf{q}^{n-2})\right) \\ &= (1-x) \mathbb{A}\mathsf{T}_{\mathsf{q}}(x^{n-1}) \end{split}$$

which by linearity implies

$$T_{q}((x+b)f) = (1-x)AT_{q}(f) + bT_{q}(f)$$
(8.15.2)

Similarly,

$$\begin{split} x \, T_q(x^n) &= \frac{1}{q^n} \left( T_q(x^n) - T_q(x^{n+1}) \right) \; = \; T_q(\,(1-x)\mathbb{A}^{-1}\,x^n) \\ T_q^{-1} \left( x T_q(x^n) \right) &= (1-x)\mathbb{A}^{-1}x^n \\ T_q^{-1} \left( x T_q(g) \right) &= (1-x)\mathbb{A}^{-1}\,g \end{split}$$

Setting  $g = T_q^{-1}(f)$  yields

$$T_q^{-1}(xf) = (1-x)\mathbb{A}^{-1}T_q^{-1}(f)$$

Applying this last recursion to  $x^n$  yields

$$\mathsf{T}_{\mathsf{q}}^{-1}(\mathsf{x}^{\mathfrak{n}}) = \left(1 - \frac{\mathsf{x}}{1}\right) \left(1 - \frac{\mathsf{x}}{\mathsf{q}}\right) \cdots \left(1 - \frac{\mathsf{x}}{\mathsf{q}^{\mathfrak{n}-1}}\right) = \mathsf{T}_{1/\mathfrak{q}}(\mathsf{x}^{\mathfrak{n}})$$

Consequently, we have the equality  $T_q^{-1} = T_{1/q}$ . This relation shows that part 1 of Theorem 8.77 implies part 4, and part 3 implies part 2.

The next Lemma establishes properties of the recurrence (8.15.2).

**Lemma 8.78.** Suppose that 0 < q < 1, b is positive, A(x) = qx, and define

$$S(f) = (1-x)\mathbb{A}f + bf.$$

*Then if*  $f \in \mathbf{P}^{(1,\infty)}$  *and*  $\mathbb{A}(f) \underline{\ll} f$  *then*  $S(f) \in \mathbf{P}^{(1,\infty)}$ *, and*  $\mathbb{A}S(f) \underline{\ll} S(f)$ *.* 

*Proof.* Since  $\mathbb{A}(f) \leq f$  and all roots of f are greater than 1 it follows that  $(1 - x)\mathbb{A}(f) \leq f$ . By linearity, we know that  $(1 - x)\mathbb{A}(f) + bf + \gamma f \in \mathbf{P}$  for any  $\gamma$ , and consequently  $S(f) \leq f$ . This also shows that  $S(f) \in \mathbf{P}$ . Since  $f \in \mathbf{P}^{(1,\infty)}$  we know that S(f) has at most one root less than 1. It's easy to see that the coefficients of

S(f) alternate, so  $S(f) \in \mathbf{P}^{\alpha lt}$ . We need to show that S(f) has no roots in (0, 1). The leading coefficients of f and S(f) alternate in sign, and it's easy to check that f is positive as  $x \to -\infty$ . Since all roots of f are greater than 1, it follows that f is positive on (0, 1). Similarly,  $\mathbb{A}(f)$  is positive on (0, 1). Finally 1 - x is positive on (0, 1), and so S(f) is positive there. This implies that  $S(f) \in \mathbf{P}^{(1,\infty)}$ .

It remains to show that  $\mathbb{A}(S(f)) \leq S(f)$ , and we do this by analyzing the location of the roots of S(f). Suppose that the roots of f are  $r_1 < \cdots < r_n$ . The roots of  $\mathbb{A}f$  are  $\frac{r_1}{q} < \cdots < \frac{r_n}{q}$ , and since  $\mathbb{A}f \leq f$  we have the ordering

$$1 < r_1 < \frac{r_1}{q} < r_2 < \frac{r_2}{q} < \cdots < r_n < \frac{r_n}{q}$$

The signs of f and Af are constant on each of the intervals defined by consecutive terms of the above sequence. Now 1 - x is negative for x > 1, so the roots of S(f) lie in the intervals where f and Af have the same sign. Since both f and Af have the same sign on  $(\frac{r_n}{q}, \infty)$  the roots of S(f) are located in the intervals

$$(\frac{r_1}{q}, r_2), (\frac{r_2}{q}, r_3), \cdots (\frac{r_{n-1}}{q}, r_n), (\frac{r_n}{q}, \infty)$$

Since  $S(f) \leq f$  there is exactly one root in each interval. The smallest possible ratio between the first two roots of S(f) is found by taking the largest possible first root,  $r_2$ , and dividing it into the smallest possible second root,  $\frac{r_2}{q}$ . This ratio is  $\frac{1}{q}$ , and the same argument applies to all the possible ratios. Since all consecutive ratios of the roots of S(f) are at least  $\frac{1}{q}$ , we find that  $\mathbb{A}S(f) \leq S(f)$ .

*Proof of Theorem 8.77.* We begin with part 1, so assume that 0 < q < 1 and  $f \in \mathbf{P}^{(1,\infty)}$ . Now  $T_q((x + b)f) = S(T_q(f))$  by (8.15.2), so by induction it suffices to prove our result for f linear, which is trivial. The case for  $f \in \mathbf{P}^{pos}$  is similar and omitted.

The proof of part 3 is entirely similar to part 2 - we need a modified version of Lemma 8.78 - and is omitted. As observed above, the remaining parts follow from 1 and 3.  $\hfill \Box$ 

#### 8.16 Newton's inequalities for Ax = qx and Ax = x + 1

Suppose Ax = qx where q > 0. The coefficients of a polynomial in  $\mathbf{P}^{\mathbb{A}}$  satisfy constraints that are the q-generalizations of Newton's inequalities. The key observation is that  $\mathbf{P}^{\mathbb{A}}$  is closed under reversal.

**Lemma 8.79.** If  $\mathbb{A}x = qx$ , q > 0 and  $f \in \mathbf{P}^{\mathbb{A}}$  then  $f^{rev} \in \mathbf{P}^{\mathbb{A}}$ .

*Proof.* If the roots of f are

$$r_1 < r_2 < \cdots < r_n$$

then the roots of  $f^{rev}$  are

$$\frac{1}{r_n} < \frac{1}{r_{n-1}} < \dots < \frac{1}{r_1}$$

Since  $f\in P^{\mathbb{A}}$  the ratio of consecutive roots of f satisfies  $r_{i+1}/r_i \geqslant q$  for  $1\leqslant i < n$ . The ratio of consecutive roots of  $f^{re\nu}$  satisfies

$$\frac{1/r_k}{1/r_{k+1}} \geqslant \mathfrak{q}$$

and so  $f^{re\nu} \in \mathbf{P}^{\mathbb{A}}$ .

**Proposition 8.80.** If  $\mathbb{A}x = qx$  and we choose  $f(x) = a_0 + \dots + a_n x^n$  in  $\mathbf{P}^{\mathbb{A}}(n)$  then

$$\frac{a_{k+1}^2}{a_k a_{k+2}} \ge \frac{4}{(q+1)^2} \frac{[k+2][n-k]}{[k+1][n-k-1]}$$
(8.16.1)

In addition,

$$\frac{a_{k+1}^2}{a_k a_{k+2}} \ge \frac{1}{q} \frac{4}{(q+1)^2} \frac{[k+2][n-k]}{[k+1][n-k-1]}$$
(8.16.2)

*Proof.* We follow the usual proof of Newton's inequalities. Write

$$f(x) = \sum_{k=0}^{n} b_k \binom{n}{k}_{q} x^k$$

where  $\binom{n}{k}_{q} = [n]!/([k]![n-k]!)$ . The q-derivative of f satisfies

$$\mathsf{D}_{\mathsf{q}}\mathsf{f} = [\mathsf{n}]\sum_{k=1}^{\mathsf{n}} \mathfrak{b}_k \binom{\mathsf{n}-1}{k-1}_{\mathsf{q}} \mathsf{x}^{k-1}$$

Consequently, if we then apply the q-derivative k times, reverse, apply the q-derivative n - k - 2 times, and reverse again, the resulting polynomial is in  $\mathbf{P}^{\mathbb{A}}$ :

$$[n] \cdots [n-k+1] \left( b_k \binom{2}{0}_q + b_{k+1} \binom{2}{1}_q x + b_{k+2} \binom{2}{2}_q x^2 \right)$$
(8.16.3)

After removing the constant factor, the discriminant satisfies

$$(q+1)^2 b_{k+1}^2 - 4 b_k b_{k+2} \ge 0$$

Substituting  $b_k = a_k / {n \choose k}_q$  and simplifying yields the first conclusion.

Now suppose that 0 < q < 1. The roots of (8.16.3) are negative, and the coefficients are positive. Let

$$\begin{split} r_1 &= \frac{-(q+1)b_{k+1} - \sqrt{(q+1)^2 b_{k+1}^2 - 4 b_k b_{k+2}}}{2 b_{k+2}} \\ r_2 &= \frac{-(q+1)b_{k+1} + \sqrt{(q+1)^2 b_{k+1}^2 - 4 b_k b_{k+2}}}{2 b_{k+2}} \end{split}$$

Since (8.16.3) is in  ${I\!\!P}^{\mathbb A}$  we know  $r_1\leqslant q\,r_2$  so

$$0 \leqslant r_2 - q r_1 = (q+1) \left( (q-1)b_{k+1} + \sqrt{(q+1)^2 b_{k+1}^2 - 4b_k b_{k+2}} \right) / 2b_{k+2}$$

Simplifying yields  $qb_{k+1}^2 \ge b_k b_{k+2}$  which establishes the second part

Remark 8.81. The product

$$\prod_{k=0}^{n-1} (x+q^k) = \sum_{k=0}^n x^k \binom{n}{k}_q q^{\binom{n-k}{2}}$$
(8.16.4)

has Newton quotients

$$\frac{1}{q} \frac{[k+2][n-k]}{[k+1][n-k-1]}.$$

Numerical computation suggests that this is probably the best possible bound.

**Example 8.82.** If q = 1 then  $4/(q+1)^2 = 1$  and [n] = n, so (8.16.1) becomes the usual Newton inequalities (4.3.1). If we take  $f(x) = a_0 + a_1x + a_2x^2 + a_3x^3 \in \mathbf{P}^{\mathbb{A}}$  then

$$\begin{split} k &= 1 & \qquad \frac{a_1^2}{a_0 a_2} \geqslant \frac{4}{(q+1)^2}(1+q+q^2) \geqslant 3 \\ k &= 2 & \qquad \frac{a_2^2}{a_1 a_3} \geqslant \frac{4}{(q+1)^2}(1+q+q^2) \geqslant 3 \end{split}$$

The bound 3 is the same as the usual Newton's inequality bound, and is realized if q = 1. If  $q \neq 1$  then the bound is better. For example, if q = 2 then the ratio is at least  $3\frac{1}{9}$ .

Although the reverse of a polynomial in  $\mathbf{P}^{sep}$  generally isn't in  $\mathbf{P}^{sep}$ , there are also inequalities for the coefficients of polynomials in  $\mathbf{P}^{sep} \cap \mathbf{P}^{pos}$ . The idea is to apply Proposition 3.22 to the interlacing  $f \leq \Delta f$ . For example, if  $f = a_0 + \cdots + a_3 x^3 \in \mathbf{P}^{sep} \cap \mathbf{P}^{pos}$  then from the Proposition the following is totally positive:

$(a_1 + a_2 + a_3)$	$2a_2 + 3a_3$	$3a_3$	0		)
$\mathfrak{a}_0$	$a_1$	a <sub>2</sub>	$a_3$	0	
0	$a_1 + a_2 + a_3$	$2a_2 + 3a_3$	$3a_3$	0	
0	$a_0$	$a_1$	$a_2$	$a_3$	
:	:	:	:	:	·. ,
\ ·	•	•	·	•	• /

For example, from the submatrix  $\begin{pmatrix} a_2 & a_2 \\ 2a_2 + 3a_3 & 3a_3 \end{pmatrix}$  we get the inequality  $a_2 \ge 3a_3$ . The corresponding matrix for a polynomial of degree n is

$$\begin{pmatrix} a_{n-1} & a_n \\ (n-1)a_{n-1} + \binom{n}{2}a_n & na_n \end{pmatrix}$$

from which we conclude that  $a_{n-1} \ge {n \choose 2} a_n$ .

# 8.17 The case Ax = -x

In this section we look at the transformation  $\mathbb{A}(x) = -x$ . This is a degenerate case, but the associated linear transformation is interesting. We can characterize the elements of  $\mathbf{P}^{\mathbb{A}}$  in terms of pairs of interlacing polynomials.

**Lemma 8.83.** If  $\mathbb{A}(x) = -x$  and  $f(x) \leq f(-x)$  then there are  $f_0$ ,  $f_1$  in  $\mathbb{P}^{p \circ s}$  such that  $f_0 \leftarrow f_1$  and  $f = f_0(x) f_1(-x)$ . Conversely, given  $f_0$ ,  $f_1$  satisfying these conditions then  $f_0(x) f_1(-x)$  is in  $\mathbb{P}^{\mathbb{A}}$ .

*Proof.* Write  $f(x) = f_0(x)f_1(-x)$  where  $f_0(x)$  contains all the negative roots, and  $f_1(-x)$  all the positive roots. Since  $f(-x) = f_0(-x)f_1(x)$  the result follows easily from consideration of the interlacing of f(x) and f(-x).

The affine derivative of f(x) is (f(x) - f(-x))/2x. If we write

$$f(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots$$

then the affine derivative is

$$\mathsf{D}_{\mathbb{A}}(f) = \frac{1}{2} (a_1 + a_3 x^2 + a_5 x^4 + \cdots).$$

If we recall the *odd part* of f

$$f_o(z) = a_1 + a_3 z + a_5 z^2 + \cdots$$

then we can write

$$\mathsf{D}_{\mathbb{A}}(f) = \frac{1}{2} f_o(x^2).$$

An unexpected consequence is that

$$\mathsf{D}_{\mathbb{A}}(\mathsf{D}_{\mathbb{A}}(\mathsf{f})) = 0.$$

We saw the odd part (and the even part) of f in §7.7.15. Since the roots of  $D_{\mathbb{A}}(f)$  are the square roots of the roots of  $f_o$  we see that in order to have  $D_{\mathbb{A}}(f) \in \mathbf{P}$  we must have that the signs of  $D_{\mathbb{A}}(f)$  alternate. See Lemma 2.6.

One result of these computations is that if the degree of f is even then the degree of  $D_{\mathbb{A}}(f)$  is two less than the degree of f, and so does *not* interlace f. It is also easy to verify that we do not have interlacing if the degree is odd.

There is a linear transformation associated with  $\mathbb{A}$  that maps  $\mathbf{P}^{pos}$  to  $\mathbf{P}^{\mathbb{A}}$ . Define

$$\mathsf{T}(\mathsf{x}^{\mathsf{n}}) = \mathsf{x} \cdot \mathbb{A}\mathsf{x} \cdot \mathbb{A}^{2}\mathsf{x} \cdots \mathbb{A}^{\mathsf{n}-1}\mathsf{x} = (-1)^{\binom{\mathsf{n}}{2}}\mathsf{x}^{\mathsf{n}}$$

Unlike linear transformations  $\mathbf{P} \longrightarrow \mathbf{P}$  (Corollary 1.50), the leading coefficients of this transformation neither alternate, nor have the same sign.

**Lemma 8.84.** If  $T(x^n) = (-1)^{\binom{n}{2}} x^n$  then

- 1. T:  $P^{pos} \cup P^{pos} \longrightarrow P$
- 2. If  $f \in \mathbf{P}^{pos}(2n)$  then  $(Tf)(x) \leq (Tf)(-x)$ .
- 3. If  $f \in \mathbf{P}^{pos}(2n+1)$  then  $(Tf)(-x) \leq (Tf)(x)$ .

*Proof.* We first get the recurrence relation for T.

$$\mathsf{T}(x\,x^{n}) = (-1)^{\binom{n+1}{2}}x^{n+1} = x\,(-1)^{\binom{n}{2}}(-x)^{n} = x\,(\mathsf{T}x^{n})(-x)$$

and by linearity

$$\begin{split} T(xf) &= x(Tf)(-x) \\ T((x+\alpha)f) &= x\,(Tf)(-x) + \alpha\,(Tf)(x) \end{split}$$

We use this recursion to prove the lemma by induction on the degree of f. For n = 1 we chose a > 0. Now T(x+a) = x+a so we have that  $-x+a \ll x+a$  since a is positive.

We will do one of the cases; the others are similar. Suppose that  $g = (x + a)f \in \mathbf{P}^{pos}(2n)$ . Then  $f \in \mathbf{P}^{pos}(2n - 1)$  and by induction hypothesis  $(Tf)(-x) \leq (Tf)(x)$ . We want to show that  $(Tg)(x) \leq (Tg)(-x)$ . From the recurrence relation

$$\begin{pmatrix} \mathsf{Tg}(x) \\ \mathsf{A}(\mathsf{Tg}(x)) \end{pmatrix} = \begin{pmatrix} (\mathsf{Tg})(x) \\ (\mathsf{Tg})(-x) \end{pmatrix} = \begin{pmatrix} \mathsf{x}(\mathsf{Tf})(-x) + \mathsf{a}(\mathsf{Tf})(x) \\ -(\mathsf{x}(\mathsf{Tf})(x) + \mathsf{a}(\mathsf{Tf})(-x)) \end{pmatrix}$$
$$= \begin{pmatrix} \mathsf{x} & \mathsf{a} \\ \mathsf{a} & -\mathsf{x} \end{pmatrix} \begin{pmatrix} (\mathsf{Tf})(-\mathsf{x}) \\ (\mathsf{Tf})(\mathsf{x}) \end{pmatrix}$$

 $Tg \leq ATg$  follows from Corollary 3.54 since  $ATf \leq Tf$ .

# 8.18 The case Ax = 0

It is surprising that the degenerate case  $\mathbb{A}x = 0$  leads to interesting results. The class  $\mathbf{P}^{\mathbb{A}}$  is empty, since  $\mathbb{A}f = f(0)$  is a constant, and can not interlace f. However, there is a class of polynomials  $\mathbb{A}_0$  that are based on the affine derivative and its conjugate.

In this case the affine derivative is also called the *lower truncation operator* since the effect of  $D_{\mathbb{A}}$  is to remove the lowest term:

$$\mathsf{D}_{L}(f) = \frac{f(x) - f(0)}{x - 0}$$
$$\mathsf{D}_{L}(a_{0} + a_{1}x + \dots + a_{n}x^{n}) = a_{1} + a_{2}x + \dots + a_{n}x^{n-1}$$

The conjugate of the lower truncation operator by the reversal operator is the upper truncation operator

$$D_{U}(a_{0} + a_{1}x + \dots + a_{n}x^{n}) = a_{0} + a_{1}x + \dots + a_{n-1}x^{n-1}$$

Neither of these operators preserve roots (See p. 636.). This leads us to consider a set of polynomials for which both operators preserve roots.

**Definition 8.85.**  $\mathbb{A}_0$  is the largest set of polynomials in  $\mathbf{P}^{pos}$  with all distinct roots such that  $\mathsf{D}_{\mathsf{U}}(\mathbb{A}_0) \subset \mathbb{A}_0$  and  $\mathsf{D}_{\mathsf{L}}(\mathbb{A}_0) \subset \mathbb{A}_0$ .

It is unexpected that there is a characterization of the members of  $\mathbb{A}_0$  in terms of coefficients, and that this characterization is a kind of converse of Newton's inequalities (Theorem 4.8).

**Corollary 8.86.** A polynomial  $f = \sum a_i x^i$  in  $\mathbf{P}^{pos}$  with all distinct roots is in  $\mathbb{A}_0$  if and only if (5.4.1) holds.

*Proof.* If  $f \in A_0$  then by applying the right combinations of  $D_U$  and  $D_L$  we find that  $a_{i-1} + a_i x + a_{i+1} x^2$  is in  $A_0$ . Since all polynomials in  $A_0$  have all distinct roots the inequalities follow.

Conversely, suppose that the coefficients of f satisfy (5.4.1). If we apply either  $D_L$  or  $D_U$  to f then the resulting polynomial g has the same coefficients as f, except for either  $a_0$  or  $a_n$ . Consequently, g also satisfies the conditions of Theorem 5.19 and so has all real roots, all distinct.

The class  $\mathbb{A}_0$  satisfies the usual closure properties

**Corollary 8.87.**  $A_0$  *is closed under differentiation and the exponential operator.* 

*Proof.* If  $f = \sum a_i x^i$  is in  $\mathbb{A}_0$  then we only need check that the coefficients of f' and EXPf satisfy (5.4.1). This is immediate from (5.4.1) and

$$\begin{split} &i^2 a_i^2 \geqslant 4 \, (i-1)(i+1) \, a_{i-1} a_{i+1} \\ & \frac{a_i^2}{(i!)^2} \geqslant 4 \, \frac{a_{i-1}}{(i-1)!} \, \frac{a_{i+1}}{(i+1)!} \end{split}$$

It is not the case that  $\mathsf{D}_L(f)$  and f interlace. In fact, they are as far from sign interlacing as possible. If r is a non-zero root of  $f\in P^\pm$ , then  $\mathsf{D}_L(f)(r)=-f(0)/r$ , and so  $\mathsf{D}_L(f)$  has the same sign at all roots of f.

Part II

# Polynomials in several variables

#### CHAPTER



# **Polynomials in two variables**

In this chapter we generalize our results from polynomials in one variable to polynomials in two variables. Our goal is to generalize  $\mathbf{P}$  and  $\mathbf{P}^{\text{pos}}$ , to define interlacing even though there are no roots, and to use these facts to deduce properties of polynomials in one variable.

#### 9.1 The substitution property and determinants

All the polynomials that we will consider in this chapter satisfy a property called *substitution*. This is the analog of "all real roots" for polynomials in two variables. However, this alone is not sufficient; we will need another condition to generalize **P**.

**Definition 9.1.** If f is a polynomial in variables x, y then f satisfies xsubstitution if for every choice of a in  $\mathbb{R}$  the polynomial f(x, a) has all real roots, and the number of roots is the same for all a. We say that f satisfies substitution if it satisfies x-substitution and y-substitution. We let **Sub**<sub>2</sub> be the set of all polynomials in two variables that satisfy substitution.

A polynomial can satisfy x-substitution and not y-substitution. Figure 9.1 shows the graph f(x, y) = 0 of such a polynomial of total degree 2. Every vertical line meets the graph in two points, so f satisfies y-substitution. Since there are some horizontal lines that do not meet the graph, f does not satisfy x-substitution.

In the remainder of this section we discuss the substitution properties of polynomials that are defined by determinants.

**Example 9.2.** We start with an example of a polynomial that satisfies a different kind of substitution. Recall that the eigenvalues of a symmetric matrix are all real. Choose symmetric matrices A, B and consider the polynomial

$$f(x,y) = |I + xA + yB|$$
 (9.1.1)



Figure 9.1: The graph of a polynomial satisfying y but not x-substitution.

where I is the identity matrix. We claim that f(x, y) satisfies the property:

• For any  $\alpha$ ,  $\beta$  the polynomial  $f(\alpha z, \beta z)$  has all real roots.

Indeed, we see that

$$g(z) = f(\alpha z, \beta z) = |I + z(\alpha A + \beta B)|.$$

Since  $\alpha A + \beta B$  is symmetric, the roots of g(z) are given by  $-r^{-1}$  where r is an eigenvalue of  $\alpha A + \beta B$ .

It is not the case that |I + xA + yB| satisfies substitution. Here's a small example:

$$A = \begin{pmatrix} 2 & 4 \\ 4 & 4 \end{pmatrix} \qquad B = \begin{pmatrix} 2 & 4 \\ 4 & 2 \end{pmatrix}$$
$$f(x, y) = |I + xA + yB| = 1 + 6x - 8x^{2} + 4y - 20xy - 12y^{2}$$
$$f(x, 1) = -7 - 14x - 8x^{2}$$

and the latter polynomial has two complex roots. Figure 9.2 shows why every line through the origin will meet the graph, yet there are horizontal and vertical lines that miss it.

Example 9.3. We again start with equation (9.1.1). Now assume that

A, B,  $A^{-1}B$  are invertible and symmetric

We claim that

- For any  $\alpha$ ,  $\beta$  the polynomial  $f(\alpha z, \beta z)$  has all real roots.
- f(x, y) satisfies x and y substitution.



Figure 9.2: Meets all lines through the origin, fails substitution

Choose  $\alpha \in \mathbb{R}$  and observe

$$f(\mathbf{x}, \boldsymbol{\alpha}) = \left| A \right| \cdot \left| \mathbf{x} \mathbf{I} + (\boldsymbol{\alpha} A^{-1} \mathbf{B} + A^{-1}) \right|$$

Since  $A^{-1}B$  and A are symmetric by hypothesis,  $\alpha A^{-1}B + A^{-1}$  is symmetric. Thus the roots of  $f(x, \alpha)$  are all real since they are the negative of the eigenvalues of a symmetric matrix. For  $f(\alpha, x)$ , notice that  $B^{-1}A$  is symmetric, since it's the inverse of a symmetric matrix.

It is easy to find matrices that satisfy the assumptions of this example: take A and B to commute. For example, we take

$$\mathsf{A} = \begin{pmatrix} 1 & -1 & -1 & 1 \\ -1 & -1 & 0 & -3 \\ -1 & 0 & -2 & 1 \\ 1 & -3 & 1 & 1 \end{pmatrix}$$

and let  $B = A^2/10$ . Figure 9.3 shows the graph of the determinant, where the dot is the origin. The segments are not linear, even though they appear to be. It is clear that it satisfies substitution, and every line through the origin meets the graph in 4 points.

**Example 9.4.** Next, we assume that A, B are symmetric and positive definite n by n matrices. We claim that f(x, y) in (9.1.1) satisfies these properties:

- For any  $\alpha$ ,  $\beta$  the polynomial  $f(\alpha z, \beta z)$  has all real roots.
- f satisfies x and y substitution.
- All coefficients of f are positive.

The first part follows as before since A, B are symmetric. Next, A is positive definite, and so it has a positive definite square root. Let  $E^2 = A$  where E is positive definite. Factoring out |A| yields

$$f(x, y) = |A| \cdot |xI + y E^{-1}BE^{-1} + A^{-1}|$$



Figure 9.3: The graph of a determinant

This representation shows that  $f(x, \alpha)$  has all real roots. Factoring out B shows that  $f(\alpha, y)$  has all real roots. The fact that all the coefficients are positive follows from a more general fact [161].

**Lemma 9.5.** Suppose that  $A_1, ..., A_d$  are positive definite. All the coefficients of  $|x_1A_1 + \cdots + x_dA_d|$  are positive.

*Proof.* We prove it by induction, and it is immediate for d = 1. Assume all matrices are n by n. If I is a subset of  $\{1, 2, ..., n\}$ , and M is a matrix, then M[I] denotes the matrix composed from the rows and columns of M listed in I. Since  $A_1$  is positive definite, we may conjugate by  $A_1^{-1/2}$ , and so assume that  $A_1 = I$ , the identity. Upon expanding the determinant we see that

$$|x_1I + (x_2A_2 + \dots + x_dA_d)| = \sum_{I} x_1^{n-|I|} (x_2A_2 + \dots x_dA_d)[I]$$

Now any principle submatrix of a positive definite matrix is positive definite, so all the terms in the sum are positive definite. Thus, by the inductive hypothesis, all the summands have positive coefficients, and so the lemma is proved.  $\hfill \Box$ 

The following is a sort of converse to the above, in that it allows us to determine that the eigenvalues are positive from properties of the polynomials.

**Lemma 9.6.** If A is a matrix and  $|I+xA| \in P^{pos}$  then A has all positive eigenvalues.

*Proof.* If the eigenvalues of A are  $r_1, ..., r_n$  then  $|I+xA| = \prod (1-xr_i)$ . Since |I+xA| is in  $\mathbf{P}^{\text{pos}}$  it has all negative roots, which implies that all  $r_i$  are positive.  $\Box$ 

A consequence of this lemma is a kind of converse to Lemma 9.5.

**Corollary 9.7.** *If* A, B *are matrices, and* f(x, y) = |I + xA + yB| *is a polynomial that satisfies* 

- 1. All coefficients of f are positive.
- 2.  $f(x, 0) \in P$ .
- 3.  $f(0, x) \in P$ .

then A and B have positive eigenvalues.

**Example 9.8.** Here is another way of constructing matrices that satisfy substitution that we will study in detail later. We assume that A and C are symmetric n by n matrices, A is positive definite, and define

$$f(x,y) = det(xA + yI + C)$$
(9.1.2)

We claim that

- f(x, y) satisfies x and y substitution.
- The coefficients of terms of degree n are positive.

By assumption  $\alpha A + C$  is symmetric, so  $f(\alpha, y)$  has all real roots, since its roots are the negative eigenvalues of  $\alpha A + C$ . Let  $A = E^2$  where E is positive definite. Since

$$f(x, y) = |A| \cdot |xI + yA^{-1} + E^{-1}CE^{-1}|$$

and  $\mathsf{E}^{-1}\mathsf{C}\mathsf{E}^{-1}$  is symmetric, it follows that  $f(x,\alpha)$  has all real roots.

The polynomial  $f^{H}(x, y)$  formed by the terms of degree n is det(xA + yI), and the roots of  $f^{H}(1, y)$  are the negative eigenvalues of A. Thus, all roots of f(1, y) are negative since A is positive definite, and hence  $f^{H}$  has all positive coefficients.

We summarize the different matrices in Table 9.1.

A	В	$A^{-1}B$	С	Sub.	f <sup>H</sup> positive	Coefficients positive
sym. sym. posdef. posdef.	sym. sym. posdef I	sym.	I I I sym.	X X X	X X	x

Table 9.1: Properties of |Ax + By + C|

We can compute an explicit example of Lemma 9.5 that shows all terms are non-negative; this properly belongs in a later chapter since the matrices are only positive semi-definite.

**Lemma 9.9.** Suppose that  $X = \text{diag}(x_i)$ ,  $Y = \text{diag}(y_i)$  are d by d diagonal matrices, and  $v_1, \ldots, v_d$  are d-vectors. Let  $M = (v_1^t, \ldots, v_d^t)^t$ . Then

$$|X + y_1 v_1^{t} v + \dots + y_d v_d^{t} v_d| = \sum_{I \subset \{1, \dots, d\}} X_{I'} Y_I |M[I])|^2$$

*Proof.* By induction it<sup>1</sup> suffices to evaluate  $|y_1v_1^{t}v + \cdots + y_dv_d^{t}v_d|$ . But this can be written as  $|M^{t}YM| - |M^{t}||Y||M| - |y_1 \cdots + |M|^2$ 

$$|\mathsf{M}^{\mathsf{t}}\mathsf{Y}\mathsf{M}| = |\mathsf{M}^{\mathsf{t}}||\mathsf{Y}||\mathsf{M}| = \mathsf{y}_1 \cdots \mathsf{y}_d |\mathsf{M}|^2$$

**Corollary 9.10.** If A is symmetric,  $|I + xA| = \prod_{1}^{d} (1 + r_i x)$ ,  $v_1 \dots, v_d$  are d-vectors then

$$|I + xA + y_1 v_1^t v_1 + \dots + y_d v_d^t v_d| = \sum_{I \subset \{1, \dots, d\}} \frac{f(x)}{\prod_{i \in I} (1 + r_i x)} Y_I |M[I]|^2$$

*Proof.* Diagonalize A and apply the lemma.

If d = 2 we have

$$\begin{split} |I + xD + y_1 v_1 v_1^{t} + y_2 v_2 v_2^{t}| &= \\ &= f + y_1 \sum v_{1,i}^2 \frac{f}{1 + xr_i} + y_2 \sum v_{2,i}^2 \frac{f}{1 + xr_i} \\ &+ y_1 y_2 \sum_{i < j} \begin{vmatrix} v_{1,i} & v_{1,j} \\ v_{2,i} & v_{2,j} \end{vmatrix}^2 \frac{f}{(1 + xr_i)(1 + xr_j)} \quad (9.1.3) \end{split}$$

The statement for n = 3 is

$$\begin{split} |\mathbf{I} + \mathbf{x}\mathbf{D} + \mathbf{y}_{1}\mathbf{v}_{1}\mathbf{v}_{1}^{t} + \mathbf{y}_{2}\mathbf{v}_{2}\mathbf{v}_{2}^{t} + \mathbf{y}_{3}\mathbf{v}_{3}\mathbf{v}_{3}^{t}| \\ &= \mathbf{f} + \mathbf{y}_{1}\sum_{i$$

# 9.2 Interlacing in Sub<sub>2</sub>

Interlacing can be easily defined for polynomials satisfying substitution.

<sup>&</sup>lt;sup>1</sup>thanks to Neil White for this argument

**Definition 9.11.** If f and g are polynomials that satisfy substitution, then we say that f and g *interlace* if for every real  $\alpha$  the polynomial f +  $\alpha$ g satisfies substitution. Using the definition of substitution, we can restate this in terms of one variable polynomials:

If  $f, g \in \mathbf{Sub}_2$  then f and g interlace iff for all  $a \in \mathbb{R}$  we have that f(x, a) and g(x, a) interlace, as do f(a, y) and g(a, y).

We define  $\leq$  in terms of these one variable polynomials.

If  $f, g \in \mathbf{Sub}_2$  then  $f \leq g$  iff for all  $a \in \mathbb{R}$  we have  $f(x, a) \leq g(x, a)$ and  $f(a, y) \leq g(a, y)$ .

For homogeneous polynomials in two variables we can verify interlacing by reducing to a one variable problem.

**Lemma 9.12.** Suppose  $f, g \in Sub_2$  are homogeneous and that the coefficients of f are all positive. The following are equivalent

- 1. f<u>∢</u>g
- 2.  $f(1,y) \leq g(1,y)$
- 3.  $f(x,1) \leq g(x,1)$

*Proof.* Note that the first interlacing is in **Sub**<sub>2</sub>, and the last two in **P**. We can substitute x = 1 in the first interlacing to deduce the second, so assume  $f(1, y) \leq g(1, y)$ . By assumption  $f(1, y) \in \mathbf{P}^{\pm}$ . If the roots of f(1, y) are  $r_1, \ldots, r_n$  then the roots of f(a, y) are  $\{r_i/a\}$ , and the roots of f(x, b) are  $\{b/r_i\}$ . There are similar expressions for the roots of g. Since the roots are either all positive, we have  $f(a, y) \leq g(a, y)$  and  $f(x, a) \leq g(x, a)$ . Thus f and g interlace in **Sub**<sub>2</sub>. Similarly we can show that 1 and 3 are equivalent.

The relation of interlacing is preserved under limits.

**Lemma 9.13.** Suppose  $f_i, g_i$  are in  $\mathbf{Sub}_2$ , and  $f_i \rightarrow f, g_i \longrightarrow g$  where f, g are polynomials. If  $f_i$  and  $g_i$  interlace for all i then f and g interlace.

*Proof.* Since  $f_i + \alpha g_i$  is in **Sub**<sub>2</sub> for all i, and  $f_i + \alpha g_i$  converges to  $f + \alpha g$  it follows that  $f + \alpha g \in$  **Sub**<sub>2</sub>. (See Lemma 10.3.) Consequently f and g interlace.

Substitution is *not* preserved under the operations of differentiation<sup>2</sup>. Consequently, we must restrict ourselves to a subset of  $\mathbf{Sub}_2$ .

<sup>&</sup>lt;sup>2</sup> The partial derivative  $\frac{\partial f}{\partial x_1}$  of  $f = (x_1 + x_2 + 1)(x_1 + x_2)(x_1 - x_2)$  has imaginary roots for x = -1/4.

#### 9.3 Polynomials in two variables

We continue our journey toward the generalization of  $\mathbf{P}$  to polynomials in two variables. For polynomials of one variable it is often important to know the sign of the leading coefficient. For two variables the homogeneous part is the analog of the leading coefficient. The homogeneous part will determine the asymptotic behavior of the graph, which is central to our generalization.

If f is a polynomial in two variables, and we let x and y get large simultaneously then the behavior of f(x, y) is determined by the terms of highest total degree. If the maximum total degree is n then

$$f(x,y) = \sum_{i+j=n} c_{ij} x^{i} y^{j} + \sum_{i+j(9.3.1)$$

The homogeneous part  $f^H$  is the first sum in (9.3.1).

**Definition 9.14.** A homogeneous polynomial  $f(x, y) = c_0 x^n + \cdots + c_n y^n$  satisfies the *positivity condition* iff all  $c_i$  are positive.

Using the homogeneous part, we define the class  $P_2$  of polynomials that forms the 2-dimensional analog of "polynomials with all real roots".

**Definition 9.15.**  $P_2(n)$  consists of all polynomials f of degree n such that f satisfies x-substitution and y-substitution, and f<sup>H</sup> satisfies the positivity condition.

$$\mathbf{P}_2 = \mathbf{P}_2(1) \cup \mathbf{P}_2(2) \cup \mathbf{P}_2(3) \cup \dots$$

Sometimes the homogeneous part of a polynomial might be negative. For instance, if  $f(x, y) \in P_2(n)$ , then the homogeneous part of f(-x, -y) has all negative signs if n is odd. In this case,  $-f(-x, -y) \in P_2$ . If n is even then  $f(-x, -y) \in P_2$ . To simplify exposition, we just write  $f(-x, -y) \in \pm P_2$ , which is true for every n. We also express this as  $\pm f(-x, -y) \in P_2$ .

It's useful to note that f<sup>H</sup> is not arbitrary.

**Lemma 9.16.** *If*  $f \in \mathbf{P}_2$  *then*  $f^{\mathsf{H}} \in \mathbf{P}^{\mathsf{pos}}$ *.* 

*Proof.* See the proof of the more general result (Lemma 10.11).

Here are a few elementary facts about  $P_2$  and  $f^H$ .

#### Lemma 9.17.

- 1. If  $f(x, y) \in \mathbf{P}_2$  then  $f(y, x) \in \mathbf{P}_2$ .
- 2. If also  $g(x, y) \in P_2$ , then  $f(x, y)g(x, y) \in \mathbf{P}_2$ .
- 3. If  $a_1, \ldots, a_n$  are positive, and  $b_1, \ldots, b_n$  are arbitrary, then

$$(x + a_1y + b_1)(x + a_2y + b_2) \cdots (x + a_ny + b_n) \in P_2$$

- 4. If  $g \in \mathbf{P}_2$  then  $(fg)^H = f^H g^H$ .
- 5. If  $g \in \mathbf{P}_2$  has degree n 1 then  $(f + g)^H = f^H$ .
- 6.  $(f^{H})^{H} = f^{H}$ .
- 7.  $\left(\frac{\partial f}{\partial x}\right)^{H} = \frac{\partial}{\partial x} f^{H}$

*Proof.*  $f(y, x)^H$  is the reverse of  $f(x, y)^H$  and f(y, x) satisfies substitution since f(x, y) does. The second part follows from the  $(fg)^H = f^H \cdot g^H$ . Since each factor  $x + a_iy + b$  is easily seen to be in P<sub>2</sub>, the product is in **P**<sub>2</sub>. The rest follow easily from the definitions.

Let's explore the definition of  $P_2$ . We can write f(x, y) as a polynomial in either x or y:

$$f(x,y) = f_0(x) + f_1(x)y + \dots + f_n(x)y^n$$
(9.3.2)

$$= f^{0}(y) + f^{1}(y)x + \dots + f^{m}(y)x^{m}$$
(9.3.3)

$$= \sum_{i,j} c_{ij} x^i y^j \qquad (9.3.4)$$

The homogeneous part  $f^H$  is  $c_{0n}y^n + \cdots + c_{n0}x^n$ . Consequently, n = m, and  $f_n(x)$  and  $f^n(x)$  are non-zero constants. The coefficient polynomials  $f_i(x)$  have x-degree n - i since the total degree of f is n and the leading coefficient of  $f_i$  is  $c_{i,n-i}$  which is non-zero. We can summarize:

A polynomial  $f \in Sub_2$  given in (9.3.2),(9.3.3) is in  $P_2$  iff

- n = m
- The degree of  $f_i$  and  $f^i$  is n i.
- The leading coefficients of f<sub>0</sub>,..., f<sub>n</sub> (and f<sup>0</sup>,..., f<sup>n</sup>) are positive.

Of course the coefficient polynomials are highly interrelated. We will see shortly that consecutive coefficients interlace. A simple property is that if f in (9.3.2) is in  $P_2$  then since  $f(x, 1) \in P$  we have that

$$\mathsf{f}_0 + \mathsf{f}_1 + \dots + \mathsf{f}_n \in \mathbf{P}$$

If  $f_1 \underline{\ll} \cdots \underline{\ll} f_n$  is a sequence of mutually interlacing polynomials then we know that, from left to right, the roots appear in reverse order:  $f_n, \ldots, f_1$ . It follows that for any  $\alpha \in \mathbb{R}$  there is an i such that

$$f_1(\alpha), \dots, f_i(\alpha)$$
 all have the same sign  $\epsilon$   
 $f_{i+1}(\alpha), \dots, f_n(\alpha)$  all have the same sign  $-\epsilon$ .

We can use this observation to construct a family of polynomials that satisfy substitution and have all coefficients in  $\mathbf{P}$ , but are not in  $P_2$ . **Lemma 9.18.** Suppose  $f_1 \leq \cdots \leq f_n$  and  $g_1 \leq \cdots \leq g_n$  are two sequences of mutually interlacing polynomials. If

$$h(x,y) = \sum f_i(x)g_i(y) = \sum F_i(x)y^i = \sum G_i(y)x^i$$

then

- 1.  $h(x, \alpha)$  and  $h(\alpha, y)$  are in **P** for all  $\alpha \in \mathbb{R}$ .
- 2. All  $F_i$  and  $G_i$  are in P.
- 3. Consecutive  $F_i$ 's and  $G_i$ 's interlace.

*Proof.* Note that  $h(x, \alpha) = \sum f_i(x)g_i(\alpha)$ . The observation above shows that  $g_i(\alpha)$  has one sign change, so we can apply Lemma 1.23. This establishes the first part.

Since  $F_0 = h(x, 0)$  we see that  $F_0 \in \mathbf{P}$ . Now since  $\{g'(y)\}$  is a sequence of mutually interlacing polynomials we know that  $\sum f_i(x)g'_i(y)$  satisfies the first part, and its constant term is  $F_1$ . Continuing, all  $F_i$  are in  $\mathbf{P}$ .

Next, for any  $\beta \in \mathbb{R}$  the sequence  $\{(x + \beta)g_i(x)\}$  is mutually interlacing. By the second part the coefficient  $\beta F_i + F_{i-1}$  of  $y^i$  is in **P**. It follows that  $F_i$  and  $F_{i-1}$  interlace.

Although substitution holds for h, the degree condition does not. And, it is not true that  $h \in \overline{P}_2$  (take  $f_i = g_i$ ). However, by Lemma 21.77 we know that h is stable.

# 9.4 Inequalities for coefficients of a quadratic in P<sub>2</sub>

Let's look at the simplest non-trivial polynomials in P<sub>2</sub>. Consider a quadratic polynomial  $f(x, y) \in P_2$  where we write the terms in a grid



We have inequalities for the coefficients on each of the three outside lines, and the center square.

**Lemma 9.19.** *If*  $f \in P_2$  *is a quadratic with the above coefficients, then* 

$$\begin{array}{l} 0\leqslant a_{01}^2-4a_{02}a_{00}\\ 0\leqslant a_{10}^2-4a_{20}a_{00}\\ 0\leqslant a_{11}^2-4a_{02}a_{20}\\ 0\leqslant a_{01}a_{10}-a_{00}a_{11} \end{array}$$

*Proof.* Since  $f(x, 0) \in \mathbf{P}$ , the terms on the bottom row constitute the coefficients of a polynomial in  $\mathbf{P}$ , and the inequality is just Newton's inequality. Similarly for the left most terms. The diagonal terms are the coefficients of  $f^H$ , which is also a polynomial in  $\mathbf{P}$ .

If we solve the equation for x, then the discriminant is

 $\Delta_{x} = (a_{10} + a_{11} y)^{2} - 4 a_{20} (a_{00} + a_{01} y + a_{02} y^{2})$ 

The discriminant of  $\Delta_x$  as a function of y is

$$\Delta_{y} = 16 \left( a_{02} a_{10}^{2} - a_{01} a_{10} a_{11} + a_{00} a_{11}^{2} + a_{01}^{2} a_{20} - 4 a_{00} a_{02} a_{20} \right)$$

Since  $\Delta_x \ge 0$ , we know  $\Delta_y \le 0$ . Rewriting  $\Delta_y \le 0$  yields

$$a_{11}(a_{10}a_{01} - a_{00}a_{11}) \ge a_{02}(a_{10}^2 - 4a_{00}a_{20}) + a_{20}a_{01}^2$$

Now the all coefficients  $a_{20}$ ,  $a_{11}$ ,  $a_{02}$  are positive since  $f^{H} \in \mathbf{P}^{pos}$ , and by the above  $a_{10}^2 - 4a_{00} a_{20} \ge 0$ , so we conclude  $a_{10} a_{01} - a_{00} a_{11} \ge 0$ .

*Remark* 9.20. We can determine when each of the inequalities is strict. The first three are strict when the corresponding polynomials have all distinct roots. If  $a_{10}a_{01} - a_{00}a_{11} = 0$  then

$$0 = a_{02} \left( a_{10}^2 - 4 a_{00} a_{20} \right) + a_{20} a_{01}^2$$

and both summands are zero by the lemma. Since  $a_{20} > 0$  we see  $a_{01} = 0$ . Interchanging the roles of x and of y shows that  $a_{10} = 0$ . Finally,  $a_{10}^2 - 4a_{00} a_{22} = 0$  implies  $a_{00} = 0$ . We conclude that f(x, y) looks like



*Remark* 9.21. Polynomials of degree two in P<sub>2</sub> are more than three quadratic polynomials spliced together. Consider the two variable polynomial

The three polynomials on the boundary  $(x^2+2x+1, x^2+17x+16, x^2+8x+16)$  have all real roots, but the quadrilateral inequality is not satisfied, so the two variable polynomial is not in P<sub>2</sub>. This phenomena will be considered later (p. 322).

It is clear from the proof that if all four conditions of the lemma are satisfied then the polynomial is in  $P_2$ .

*Remark* 9.22. We will see later (p. 323) that the last inequality (the quadrilateral or rhombus inequality) is a consequence of interlacing of the coefficient polynomials. It can't be improved. For example, the diagram of coefficients of the polynomial  $(x + \epsilon y + 1)(\epsilon x + y + 1)$  is



and the ratio  $(a_{01}a_{10})/(a_{00}a_{11})$  equals

$$\frac{(1+\epsilon)^2}{1+\epsilon^2}$$

and this goes to 1 as  $\epsilon \rightarrow 0$ . If r, s > 0 and we let

$$f(x,y) = (x + \varepsilon(y+1))^r (y + \varepsilon(x+1))^s (\varepsilon x + y + 1)(x + \varepsilon y + 1)$$

then as  $\epsilon \to 0$  the ratio  $(a_{r,s+1}a_{r+1,s})/(a_{r,s}a_{r+1,s+1})$  converges to 1. Thus, there are no Newton inequalities with constant greater than 1.

The last inequality in Lemma 9.19 corresponds to the two adjacent triangles with a vertex in the lower left corner. The two other corners do not give the inequalities

$$a_{10}a_{11} - a_{01}a_{20} > 0$$
 (right corner) (9.4.1)  
 $a_{01}a_{11} - a_{10}a_{02} > 0$  (upper corner)

as we see by expanding  $(x + y - 1)^2$ :



We will see later (Proposition 10.41 that if the coefficients are positive then the inequalities (9.4.1) do hold.

**Example 9.23.** We can use these ideas to show that perturbations of products aren't necessarily in  $P_2$ . Let

$$f = e + (x + ay + b)(x + cy + d)$$

where  $a \neq c$  are positive. The discriminant  $\Delta_y$  is  $16(a - c)^2 e$ . Since this is positive for positive *e*, we conclude that there are perturbations of products that are not in P<sub>2</sub>.

For example, if f = (x + y)(x + 2y) + e, then  $f \in P_2$  if  $e \leq 0$ , and  $f \notin P_2$  if  $e \geq 0$ .

Although it is difficult to realize a set of numbers as the coefficients of a polynomial in  $\mathbf{P}_2^{\text{pos}}$ , the determinant is the only restriction for a rhombus.

**Lemma 9.24.** Suppose that  $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$  is a matrix with positive entries. The following are equivalent:

1. 
$$ad - bc > 0$$

 $\text{2. } \left(\begin{smallmatrix} a & b \\ c & d \end{smallmatrix}\right) = \left(\begin{smallmatrix} \alpha_{01} & \alpha_{11} \\ \alpha_{00} & \alpha_{10} \end{smallmatrix}\right) \textit{for some } \sum \alpha_{ij} \, x^i \, y^j \in I\!\!P_2^{\textit{pos}}.$ 

*Proof.* We may assume that c = 1. It suffices to consider the polynomial

 $f(x, y) = (1 + r_1 x + s_1 y)(1 + r_2 x + s_2 y).$ 

We want to find positive  $r_1$ ,  $s_1$ ,  $r_2$ ,  $s_2$  such that

$$\begin{pmatrix} a & b \\ 1 & d \end{pmatrix} = \begin{pmatrix} s_1 + s_2 & r_1 s_2 + r_2 s_1 \\ 1 & r_1 + r_2 \end{pmatrix}.$$

Now it is easy to see that

$$(0, ad) = \big\{ r_1 \, s_2 + r_2 \, s_1 \ | \ s_1 + s_2 = a, r_1 + r_2 = d, r_1, r_2, s_1, s_2 > 0 \big\}.$$

Since  $b \in (0, ad)$  by the determinant hypothesis we have found  $f \in \mathbf{P}_2^{\text{pos}}$ .  $\Box$ 

There are inequalities for the coefficients of  $f(x,y) \in P_2$  that come from Proposition 5.38

Lemma 9.25. If  $f(x,y) = \sum_0^n f_i(x) y^i \in {\tt P}_2$  and 0 < k < n then

$$\sum_0^k f_i(x)\,f_{k-i}(x)(-1)^{k-i} \geqslant 0$$

*Proof.* For any  $\alpha$  we apply Proposition 5.38 to  $f(\alpha, y)$ .

The inequality isn't strict. If k=0 then then the sum is  $f_0(\boldsymbol{x})^2$  which takes on the value 0.

#### 9.5 Solution curves

We have been discussing polynomials in  $P_2$  without actually looking at them. The geometric perspective gives important information about these polynomials. This is not surprising, since the condition that distinguishes  $P_2$  from **Sub**<sub>2</sub> concerns the structure of the homogeneous part, and the homogeneous part constrains the geometry of the graph. If f(x, y) is any polynomial, then the *graph* of f is defined to be

$$G_{f} = \{(x, y) | f(x, y) = 0\}$$

For example, if we choose a fifth degree polynomial

$$f = (x + y)(1 + x + 2y)(1 + x + 3y)(2 + x + 3y)(2 + x + 5y)$$
(9.5.1)

then the graph of  $g = 4f + 3 \frac{\partial f}{\partial x}$  (see Figure 9.4) is surprising linear outside of the central region. We will see later that g is in  $\mathbf{P}_2$ .



Figure 9.4: The graph of a polynomial in  $P_2$ 

We can decompose  $G_f$  into n curves, where n is the degree of f. Define  $r_i \colon \mathbb{R} \longrightarrow \mathbb{R}$  by setting  $r_i(y)$  to the i-th largest root of f(x, y) = 0. Since f satisfies x-substitution each  $r_i$  is well defined. The roots are continuous functions of the coefficients, and so each  $r_i$  is continuous. We call  $r_i(x)$  a *solution curve* of f. These are the analogs of the zeros of a polynomial in one variable. We can write

$$G_f = r_1(\mathbb{R}) \cup \cdots \cup r_n(\mathbb{R})$$

**Lemma 9.26.** If  $f \in \mathbf{P}_2$  then asymptotically the graph  $G_f$  is approximately a collection of infinite rays. The curve  $r_i(\mathbb{R})$  is asymptotically a ray with slope given by the *i*-th largest root of  $f^H(x, 1)$ . Consequently, for x, y sufficiently large  $G_f$  lies in the union of the upper left quadrant and the lower right quadrant.

*Proof.* If the degree of f is n the polynomial  $f^H(1, y)$  has n roots  $\beta_1, \ldots, \beta_n$  by Lemma 9.16. Consider

$$f(x,y) = f^H(x,y) + \sum_{i+j < n} c_{ij} x^i y^j$$

For x large and y approximately  $\beta_i x$  we see that

$$f(x,y) \approx f^{H}(x,y) + O(x^{n-1})$$

and hence for large |x| there is a root y of f(x, y) = 0 where y is close to  $\beta_i x$ .

Thus, for |x| large f(x, y) has n roots, approximately equal to  $\beta_1 x, \ldots, \beta_n x$ . Since  $f^H$  has all positive coefficients, the roots of  $f^H(1, y)$  are negative, and so all the  $\beta_i$  are negative. Each  $r_i(x)$  is in the upper left quadrant for x large, and in the lower right quadrant for x negative and |x| large.

The polynomial  $y^2 - x^2 + 1$  shows that x-substitution alone does not imply y-substitution. Theorem 9.27 remedies this situation by giving a condition that along with x-substitution implies y-substitution. This theorem is the best way to show that a polynomial is in **P**<sub>2</sub>.

**Theorem 9.27.** Suppose that f(x, y) is a polynomial that satisfies x-substitution, and  $f^H$  satisfies the positivity condition. Then f satisfies y-substitution, and is in  $P_2$ .

*Proof.* Since each solution curve  $r_i$  is continuous and and is asymptotically a ray that lies in the upper left quadrant or the lower right quadrant it follows that the graph of each  $r_i$  meets every horizontal line, so the equation  $r_i(x) = a$  has a solution for every a. In particular, for any y there are n solutions to f(x, y) = 0 since  $f(a, r_i(a)) = 0$  for each i. Since f(x, y) has degree n in x this implies that f(x, y) satisfies x-substitution.

*Remark* 9.28. It is important in the definition of substitution that the homogeneous part has all positive coefficients. For instance, choose  $g, h \in \mathbf{P}$ , and define f(x, y) = g(x)h(y). The homogeneous part of f is a monomial  $x^n y^m$ , and does not satisfy positivity, and so is not in P<sub>2</sub> The graph of f consists of n horizontal lines, and m vertical lines. We will see later that if  $g, h \in \mathbf{P}^{pos}$  then f is in the closure of P<sub>2</sub>.

Here is an important way of constructing polynomials in  $P_2$ . We will later show that these two conditions are equivalent.

Lemma 9.29. If A and C are symmetric matrices, and A is positive definite then

$$f(x, y) = \det(xA + yI + C) \in P_2.$$

*Proof.* This is Example 9.8

**Lemma 9.30.** If a solution curve of  $f \in P_2$  is horizontal or vertical at a point (a, b), then (a, b) is the intersection of at least two solution curves.

*Proof.* If the solution curve is horizontal at (a, b) then  $\frac{dy}{dx}(a, b) = 0$ . Since a solution curve is implicitly defined by f(x, y) = 0, we can differentiate to find  $f_x + \frac{dy}{dx} f_y = 0$ . This shows that  $f_x(a, b) = 0$ . Consequently, the point a is a double point of the function f(x, b), and so (a, b) lies on two solution curves. The vertical case is similar.

**Lemma 9.31.** The solution curves of a polynomial in  $P_2$  are non-increasing. They always go down and to the right.

*Proof.* If we ever had a solution curve that was increasing, then it would have a local minimum since it is asymptotic to a line of negative slope. A horizontal line slightly above this minimum would intersect it in two distinct points, contradicting the definition of solution curve. Similarly, considering the curve as a function of y we see it can never go left.

**Lemma 9.32.** If f is a polynomial in  $\mathbf{P}_2$  then  $f(x, x) \in \mathbf{P}$ . More generally, f(ax + b, cx + d) is in  $\mathbf{P}$  for any a, b, c, d such that a/c is either greater than the largest root of  $f^H$  or less than the smallest root of  $f^H$ .

*Proof.* Since each  $r_i(\mathbb{R})$  is connected and asymptotically is a line with negative slope, any line of positive slope intersects  $r_i(\mathbb{R})$ . See Figure 9.5. This proves the first part.

Under the hypothesis on a/c any solution curve either has asymptotes whose slopes are both larger (or both smaller) than a/c. Any line with slope a/c will intersect such a curve.



Figure 9.5: The intersection of f(x, y) = 0 and y = ax + b

The hypothesis on a/c are necessary. In Figure 9.5 it is easy to find lines that do not intersect the the solution curve whose ends are marked "\*".

The proof of the following theorem requires polynomials in four variables, and is proved in Theorem 10.60.

**Theorem 9.33.** *If*  $f \in \mathbf{P}_2$  *then for positive* a, b, c, d *we have* 

$$f(ax + by + u, cx + dy + v) \in \mathbf{P}_2$$

As an example, notice that if  $f \in \mathbf{P}_2$  then  $f(x, x + y) \in \mathbf{P}_2$ .

In addition to considering the intersection of the graph of f with lines we can consider the intersection of f with the hyperbola xy = -1. These intersections correspond to solutions of f(x, -1/x) = 0. Since this is not a polynomial, we multiply by  $x^n$  where n is the degree of f. The degree of  $x^n f(x, -1/x)$  is 2n.

For example, if  $f \in \mathbf{P}^{\text{pos}}(n)$  then  $F(x, y) = y^n f(x/y)$  is in  $\mathbf{P}_2$ . Obviously  $x^n F(x, -1/x) = (-1)^n f(-x^2)$ . Since all roots of f are negative, each root of f gives rise to 2 roots of  $x^n F(x, -1/x)$  and so all 2n roots are accounted for.

The reverse of one variable requires a negative sign. If we use 1/x instead of -1/x we get stable polynomials. (See p. 620.)

**Lemma 9.34.** *If*  $f \in \mathbf{P}_2(n)$  *then*  $x^n f(x, -1/x) \in \mathbf{P}(2n)$ .

*Proof.* Since all asymptotes of f have negative slope the graph of xy = -1 meets the graph of f in n points in the upper left quadrant, and n times in the lower right quadrant - see Figure 9.6. This gives 2n solutions to f(x, -1/x) = 0, and since this is the degree of  $x^n f(x, -1/x)$  the conclusion follows.



Figure 9.6: The intersection of f(x, y) = 0 and xy = -1

*Remark* 9.35. Harmonic functions are an important class of functions. A function is harmonic if it is the real part of an analytic function. As long as the degree is at least two then no polynomial in  $P_2$  is harmonic. A geometric explanation is that if f is a harmonic polynomial then the real part of f(x + iy) has asymptotes given by the 2n rays with angles  $\pi(2k + 1)/2n$ , k = 1, ..., 2n ([69]). Thus, the real part has asymptotes whose slopes are positive and negative, and hence is not in  $P_2$ .

**Example 9.36.** Recall (Example 9.2) that polynomials of the form f(x, y) = |I + xA + yB|, where A, B are n by n symmetric matrices, satisfy a strong line

intersection property: every line through the origin meets the graph of f in n points. This fails for polynomials in  $P_2$ . For instance, if we define

$$f(x,y) = \begin{vmatrix} 4+x+2y & -3\\ -3 & -2+x+y/2 \end{vmatrix} = -17 + 2x + x^2 - 2y + \frac{5xy}{2} + y^2$$

then  $f(x, y) \in P_2$ , and it is clear from the graph of f (Figure 9.7) that there is a range of lines through the origin that do not meet the graph of f.



Figure 9.7: Satisfies substitution, fails to meet all lines through the origin

# **9.6** Interlacing in P<sub>2</sub>

**Definition 9.37.** If  $f, g \in \mathbf{P}_2$  then f and g *interlace* iff  $f + \alpha g$  is in  $\pm \mathbf{P}_2$  for all  $\alpha$ . If f and g have the same degree then it is possible that  $(f + \alpha g)^H$  has negative coefficients. That's why we we require that  $f + \alpha g \in \pm \mathbf{P}_2$ , which we recall means that either  $f + \alpha g \in \mathbf{P}_2$ , or  $-(f + \alpha g) \in \mathbf{P}_2$ . If in addition the degree of f is one more than the degree of g then we say  $f \leq g$ . In this case  $(f + g)^H = f^H$ , and so the only condition we need to verify is substitution. This leads to an equivalent definition in terms of substitutions:

If f,  $g \in \mathbf{P}_2$  then  $f \leq g$  iff for all  $a \in \mathbb{R}$  we have  $f(x, a) \leq g(x, a)$  ( or for all  $a \in \mathbb{R}$  we have  $f(a, y) \leq g(a, y)$ ).

Figure 9.8 shows the graphs of the two interlacing polynomials

$$f = (x + y + 1)(x + 2y + 3)(x + 5y + 4)(x + 3y + 2)$$

and  $\frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}$ . From this we see that there really is a geometric interpretation to interlacing: if  $f \leq g$  then the solution curves of g interlace the solution curves of f.

We define strict interlacing  $f \leq g$  to mean that  $f \leq g$ , and the graphs of f and g are disjoint. It is easy to see that this is equivalent to saying that  $f(x, \alpha) \leq g(x, \alpha)$  for all  $\alpha \in \mathbb{R}$ .

The following lemma covers the fundamental properties of interlacing in  $\mathbf{P}_2$ .



Figure 9.8: The graphs of two interlacing polynomials in  $P_2$ 

**Theorem 9.38.** *If* f, g,  $h \in \mathbf{P}_2$  *and*  $f \leq g$ ,  $f \leq h$ ,  $\alpha$ ,  $\beta$  *positive then* 

- $\alpha g + \beta h \in P_2$
- $f \leq \alpha g + \beta h$

*Proof.* Lemma 1.10 shows that  $\alpha g + \beta h$  satisfies substitution. Since  $(\alpha g + \beta h)^H = \alpha g^H + \beta h^H$  it follows that  $\alpha g + \beta h$  satisfies the positivity condition.

**Theorem 9.39.**  $\mathbf{P}_2$  is closed under differentiation. If  $f \in \mathbf{P}_2$  then  $f \leq \frac{\partial f}{\partial x_i}$  for  $1 \leq i \leq 2$ .

*Proof.* Choose  $\alpha \in \mathbb{R}$ , and consider  $g = f + \alpha \frac{\partial f}{\partial x_1}$ . Since  $g^H = f^H$ , we will show that  $g \in P_2$  by showing g satisfies substitution. If we choose  $\beta \in \mathbb{R}$  then

$$g(x, \beta) = f(x, \beta) + \alpha \frac{\partial f}{\partial x_1}(x, \beta)$$

Since derivatives in one variable interlace, it follows from Theorem 9.27 that  $g \in P_2$ , and and that  $f \leq \frac{\partial f}{\partial x_1}$ .

One of the surprising things about this definition is that interlacing linear polynomials are highly constrained. In **P**, any two polynomials of degree one interlace. In **P**<sub>2</sub>, the only way that degree one polynomials interlace is for them to be essentially one-dimensional. Suppose that  $ax+by+c \leq sx+ty+u$  where a, b, s, t are positive. The requirement that  $(ax+by+c) + \alpha(sx+ty+u)$  lies in **P**<sub>2</sub>, implies that sx + ty is a multiple of ax + bx. If we define f(x) = x + c and  $g(x) = x+t/\alpha$  then we can express these two interlacing polynomials in terms of interlacing one variable polynomials. Namely, f(ax + by) = ax + by + c,  $\alpha g(ax + by) = sx + ty + u$ , and  $f \leq g$ . Thus there are no intrinsically two-dimensional interlacing linear polynomials. Geometrically, all this means is that the graph of interlacing linear polynomials must be parallel lines.

A consequence of this observation is that there appears to be no simple creation of *mutually interlacing polynomials*. We have seen how to create mutually interlacing polynomials using linear combinations of the products of n-1 factors from a polynomial with n factors. The fact that this set of polynomials is mutually interlacing follows from the fact that any n degree one polynomials in **P** are mutually interlacing.

Interlacing polynomials can be decomposed. This fact follows from from the assumption that the polynomials  $f + \alpha g$  all lie in  $\pm P_2$ , and hence the coefficients of the homogeneous part all have the same sign.

**Lemma 9.40.** Suppose  $f, g \in P_2$ , f and g have the same total degree, and f, g interlace. then

- 1.  $f^H$  and  $g^H$  are scalar multiples of each other.
- 2. There is an  $r \in \mathbf{P}_2$  and positive  $\alpha$  such that

r

a) 
$$f \leq r$$
  
b)  $g = \alpha f \pm dt$ 

*Proof.* First of all, we note that  $(f + \alpha g)^H = f^H + \alpha g^H$ , so  $f^H$  and  $g^H$  interlace. Since  $f + \alpha g \in \mathbf{P}_2$  for all  $\alpha$ , it follows that  $(f + \alpha g)^H \in \mathbf{P}^{pos}$  for all  $\alpha$ . From Lemma 2.14 there is a  $\gamma$  such that  $f + \gamma g$  is not in  $\mathbf{P}^{pos}$ , so the second case of Lemma 2.14 is not possible. Thus,  $g^H$  is a constant multiple of  $f^H$ .

By the previous lemma we can choose  $\alpha$  so that  $g^{H} = \alpha f^{H}$ . We define  $r = \pm (f - \alpha g)$  where the sign is chosen to make the coefficients of  $r^{H}$  positive. All linear combinations of f and g are in  $P_2$ , so r is in  $P_2$ . Moreover, the total degree of r is less than that of f since we removed all the highest degree terms. Also, f and r interlace since their linear span is the same as the linear span of f and g.

We use this definition to define  $\underline{\ll}$  for **P**<sub>2</sub>. Say that  $f \underline{\ll} g$  iff there is an  $r \in \mathbf{P}_2$  such that  $g = \alpha f + r$  for some positive  $\alpha$ , and  $f \underline{\lessdot} r$ .

#### **9.7** Linear transformations on **P**<sub>2</sub>

We study linear transformations on  $P_2$ . Just as in one variable, linear transformations preserving  $P_2$  preserve interlacing. The proof is immediate from the definition of interlacing.

**Theorem 9.41.** Suppose that  $T: P \longrightarrow P_2$  and  $S: P_2 \longrightarrow P_2$  are linear transformations. If f, g  $\in P$  interlace then Tf, Tg interlace. If f, g  $\in P_2$  interlace then Sf, Sg interlace.

**Corollary 9.42.** If  $f \leq g$  are in  $\mathbf{P}_2$ , and  $\mathfrak{a}$ ,  $\mathfrak{b}$  are positive then

$$a\frac{\partial f}{\partial x} + b\frac{\partial f}{\partial y} \le a\frac{\partial g}{\partial x} + b\frac{\partial g}{\partial y}$$

If  $f \lessdot g$  then  $\frac{\partial f}{\partial x_i} \lessdot \frac{\partial g}{\partial x_i}$ 

*Proof.* The map  $f \mapsto a \frac{\partial f}{\partial x} + b \frac{\partial f}{\partial y}$  maps  $P_2$  to itself, so it preserves interlacing. The second part follows form the one variable result.

**Lemma 9.43.** If  $f(x, y) \leq g(x, y)$  in  $P_2$  then  $f(x, x) \leq g(x, x)$ .

*Proof.* The map  $y \mapsto x$  is a linear transformation from  $P_2$  to P, and so it preserves interlacing.

A linear transformation  $\mathbf{P} \longrightarrow \mathbf{P}$  sometimes induces a linear transformation  $\mathbf{P}_2 \longrightarrow \mathbf{P}_2$ . Analyzing the geometry of the graph of the image of a polynomial under this induced transformation yields information about new transformations on  $\mathbf{P}$  or  $\mathbf{P}^{\text{pos}}$ .

**Theorem 9.44.** Suppose that  $T: P \longrightarrow P$  is a linear transformation that preserves degree, and maps polynomials with positive leading coefficients to polynomials with positive leading coefficients. The induced linear transformation

$$\mathsf{T}_*(\mathsf{x}^{\mathsf{i}}\mathsf{y}^{\mathsf{j}}) = \mathsf{T}(\mathsf{x}^{\mathsf{i}})\,\mathsf{y}^{\mathsf{j}}$$

defines a linear transformation from  $P_2$  to itself.

*Proof.* Suppose  $f = \sum f_i(x)y^i$  where  $f_i$  has degree n - i. Since T preserves degree, the degree of  $T(f_i)$  is n-i, and so  $(Tf)^H$  is a sum of terms  $x^{n-i}($  leading coefficient of  $Tf_i$ ). Since T preserves the sign of the leading coefficient,  $(Tf)^H$  has all positive terms.

By Theorem 9.27 it suffices to show that  $T_*(f)$  satisfies x-substitution. If we choose  $b\in\mathbb{R}$  then

$$(\mathsf{T}_* \mathsf{f})(\mathsf{x},\mathsf{b}) = \mathsf{T}(\mathsf{f}(\mathsf{x},\mathsf{b}))$$

Since  $f(x, b) \in \mathbf{P}$  we know T(f(x, b)) is in  $\mathbf{P}$ , and so  $T_*$  satisfies x-substitution.

We will revisit induced transformations in Chapter 11.11.12.

*Remark* 9.45. It is important to see that a linear transformation  $T: \mathbf{P} \longrightarrow \mathbf{P}$  does not in general induce a linear transformation  $\mathbb{P}_2 \longrightarrow \mathbf{P}_2$ . The assumptions of degree and positivity are essential. The proof shows that substitution will always be met. For instance, consider

$$T: g \mapsto g(\mathsf{D})x.$$

This satisfies T(1) = x, T(x) = 1 and  $T(x^k) = 0$  for k > 1. Since

$$T(x + y)^2 = T(x^2 + 2xy + y^2) = 2y + xy^2$$

we see that  $T(x + y)^2 \notin P_2$  since the homogeneous part is  $xy^2$ . See § 11.11.12.

**Corollary 9.46.** If  $T(x^iy^j) = H_i(x)H_j(y)$  then  $T: \mathbf{P}_2 \longrightarrow \mathbf{P}_2$ .

*Proof.* The maps  $x^i \mapsto H_i(x)$  and  $y^j \mapsto H_j(y)$  map **P** to itself by Corollary 7.44.

We can make use of a special function identity to verify Corollary 9.46 in a special case. We have

$$\begin{split} \mathsf{T}\left(x+y\right)^n &= \sum_{i=0}^n \binom{n}{k} \mathsf{T}(x^k y^{n-k}) \\ &= \sum_{i=0}^n \binom{n}{k} \mathsf{H}_k(x) \mathsf{H}_{n-k}(y) \\ &= 2^{n/2} \mathsf{H}_n\left(\frac{x+y}{\sqrt{2}}\right) \end{split}$$

and the last polynomial is in  $P_2$  since  $H_n$  is in P.

Theorem 9.44 is not true<sup>3</sup> if we only have that T:  $\mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}$ .

**Lemma 9.47.** Consider the maps  $T: f(x) \mapsto f(x + y)$  and  $S: f(x) \mapsto F(x, y)$ , where F(x, y) is the homogeneous polynomial corresponding to f. These are linear transformations  $T: \mathbf{P} \longrightarrow \mathbf{P}_2$  and  $S: \mathbf{P}^{\pm}(n) \longrightarrow \mathbf{P}_2(n)$ .

*Proof.* The map T is linear and preserves degree, and S is linear if we restrict to polynomials of the same degree.  $\Box$ 

**Corollary 9.48.** The linear transformation  $T(x^iy^j) = \frac{x^iy^j}{i!j!}$  maps  $P_2$  to  $P_2$ .

*Proof.* Apply Theorem 9.44 to the exponential map in each variable.

There are a few instances where we can describe all the coefficients of  $y^i$  in a polynomial belonging to  $P_2$ . Here are three examples - in each case the left sides are of the form f(x + y) so the right hand sides are in  $P_2$ .  $H_n$  is the Hermite polynomial, and the identity is a consequence of the Taylor series and the identity  $H'_n = 2nH_{n-1}$ .

$$\begin{split} (x+y+1)^n &= \sum_{i=0}^n \binom{n}{k} (x+1)^k y^{n-k} \\ H_n(x+y) &= \sum 2^k \binom{n}{k} H_{n-k} y^k \end{split}$$

If we expand f(x + y) in its Taylor series then for any  $f \in \mathbf{P}^{pos}$ 

$$f(x+y) = \sum_{i} f^{(i)}(x) \frac{y^{i}}{i!} \in \mathbf{P}_{2}$$
(9.7.1)

<sup>&</sup>lt;sup>3</sup> For example, we saw in §7.7.7 that  $T(x^n) = \langle \underline{x} \rangle_n$  is such a map. If we set f = (1 + x + y)(1 + 2x + y)(2 + 3x + y) then the coefficients of  $T_*f$  interlace but  $T_*f$  does not satisfy x-substitution since  $(T_*f)(3, y)$  has imaginary roots.
A linear transformation from P to P preserving  $P^{\text{pos}}$  determines a linear transformation from P to  $P_2$ .

**Lemma 9.49.** Suppose  $T: P \longrightarrow P$  is a linear transformation that preserves degree, and maps  $P^{pos}$  to itself. If  $T(x^i) = f_i$  and we define  $S(x^i) = y^i f_i(x/y)$ , then  $S: P^{pos} \longrightarrow P_2$ .

*Proof.* Suppose that  $g = \sum_{i=0}^{n} a_i x^i$  is in  $\mathbf{P}^{\text{pos}}$ . The homogeneous part of S(g) is  $a_n S(x^n) = a_n y^n f_n(x/y)$  whose homogenization is in  $\mathbf{P}^{\text{pos}}$  since T maps  $\mathbf{P}^{\text{pos}}$  to  $\mathbf{P}^{\text{pos}}$ . To verify substitution, choose  $\alpha$  and consider

$$S(g)(x, \alpha) = \sum a_i S(x^i)(x, \alpha) = \sum a_i \alpha^i f_i(x/\alpha)$$

This last polynomial is in  $\mathbf{P}^{pos}$ , as the diagram shows



**Example 9.50.** If we choose the affine map T(f) = f(x+1) then  $S(x^i) = (x+y)^i$ , and S(f) = f(x+y).

**Lemma 9.51.** *If* T:  $P \longrightarrow P$  and  $T(x^n) = p_n$ , then for any  $\alpha$  the linear transformation  $S(x^k) = x^{n-k} p_k(\alpha)$  maps  $P^{\pm}(n) \longrightarrow P(n)$ .

*Proof.* The proof follows from the diagram



Π

# 9.8 Applications to linear transformations on P

We now apply results about  $P_2$  to get information about linear transformations on **P**.

**Corollary 9.52.** Assume that the linear transformations  $T, S: P \longrightarrow P$  map  $x^n$  to a polynomial of degree n with positive leading coefficient, and maps  $P^{pos}$  to itself. . Define a transformation V on P(n) by  $V(x^i) = T(x^i) S(x^{n-i})$ . Then V maps  $P^{pos}(n)$  to itself.

Proof. V is the composition



where  $T_*$  and  $S_*$  are defined in Theorem 9.44. The conclusion follows form Lemma 9.32.

**Corollary 9.53.** If  $H_i(x)$  is the Hermite polynomial then the transformation  $x^i \mapsto H_i(x) H_{n-i}(x)$  maps  $P^{pos}(n)$  to P(n). If  $L_i(x)$  is the Laguerre polynomial then the transformation  $x^i \mapsto L_i(x) L_{n-i}(x)$  maps  $P^{pos}(n)$  to  $P^{pos}(n)$ . Also, the transformation  $x^i \mapsto L_i(x) H_{n-i}(x)$  maps  $P^{pos}(n)$  to P(n).

**Corollary 9.54.** Assume that the linear transformation  $T: P \longrightarrow P$  map  $x^n$  to a polynomial of degree n with positive leading coefficient. The linear transformation  $x^i \mapsto x^{n-i}T(x^i)$  maps  $P^{pos}(n)$  to itself.

*Proof.* Take S to be the identity in Corollary 9.52.

The next result follows from the corollary, but we will refine this result in Lemma 9.60.

**Lemma 9.55.** The transformation  $x^i \mapsto H_i(x)x^{n-i}$  maps  $P^{pos}(n)$  to P(n).

**Lemma 9.56.** *Suppose that*  $f \in P$ *, and*  $T: P \longrightarrow P$ *. Then* 

$$T_*(f(x+y))(x,x) \in P$$

*Proof.* We know that  $f(x+y) \in \mathbf{P}_2$ , and so  $T_*(f(x+y)) \in \mathbf{P}_2$ . Apply Lemma 9.32.

**Corollary 9.57.** Suppose  $T: P \longrightarrow P$  is a linear transformation that preserves degree and maps  $P^{\text{pos}}$  to itself. The Möbius transformation  $T_{1/z}$  satisfies  $T_{1/z}: P^{\text{pos}} \longrightarrow P^{\text{pos}}$ .

*Proof.* Let  $T(x^i) = f_i(x)$ , and recall that  $T_{1/z}(x^i) = f_i^{\text{REV}}$ . Let  $S(x^i) = y^i f_i(x/y)$  as in Lemma 9.49. Notice that  $S(x^n)(1,y) = y^n f_n(1/y) = T_{1/z}(y)$  and so  $S(f)(1,y) = T_{1/z}(f)$ . Now S maps  $P^{\text{pos}}$  to  $P_2$  by Lemma 9.49, and hence  $T_{1/z}(f) = S(f)(1,y) \in \mathbf{P}$ .

See (p. 512) for a different proof.

**Corollary 9.58.** The linear transformation  $x^i \mapsto L_i^{REV}(-x)$  maps  $P^{pos} \longrightarrow P^{pos}$ .

*Proof.* Apply Corollary 9.57 to Lemma 15.41.

The reason that we can substitute x = y in a polynomial in  $P_2$  is that the line x = y must intersect every solution curve. The line x + y = 1 does not necessarily meet every solution curve, but an easily met assumption about the homogeneous part will allow us to substitute 1 - x for y.

**Lemma 9.59.** Suppose  $T: P \longrightarrow P$  maps  $P^{pos}(n)$  to itself, and define  $S(x^i) = (1-x)^{n-i} T(x^i)$ . Assume that the leading coefficient of  $T(x^i)$  is  $c_i$ , and write  $f(x) = \sum a_i x^i$ . If  $f \in P^{pos}$  and either condition below is met then  $S(f) \in P$ .

- all roots of  $\sum a_i c_i x^i$  are greater than -1.
- all roots of  $\sum a_i c_i x^i$  are less than -1.

*Proof.* S is determined by the composition

 $P^{\text{pos}}(n) \xrightarrow{\text{homogenize}} P_2(n) \\
 s \downarrow \\
 P(n) \xleftarrow{(x,y)\mapsto(x,1-x)} P_2(n)$ 

If we choose  $f \in \mathbf{P}^{p \circ s}$  as in Corollary 9.52 we know  $T_*(F) \in \mathbf{P}_2$ , where F is f homogenized. The asymptotes of  $T_*(F)$  have slopes that are the roots of the homogeneous part of  $T_*(F)$ , which is  $\sum a_i c_i x^i$ . If either condition is met, then all solution curves meet the line x + y = 1.

The linear transformation T:  $x^k \mapsto H_k(x)x^{n-k}$  acting on **P**(n) becomes very simple if we make a transformation. We need the identity

$$T(x-2+\alpha)^n = H_n(\frac{\alpha x}{2})$$
(9.8.1)

If we substitute  $\alpha = 0$  then

$$\mathsf{T}(\mathsf{x}-2)^{\mathfrak{n}} = \mathsf{H}_{\mathfrak{n}}(0) = \begin{cases} 0 & \text{if $\mathfrak{n}$ is odd} \\ (-1)^{\mathfrak{n}/2}(\mathfrak{n}-1)! & \text{if $\mathfrak{n}$ is even} \end{cases}$$

This degenerate behavior at x = 2 is reflected in the following lemma.

**Lemma 9.60.** The linear transformation  $T: x^k \mapsto H_k(x)x^{n-k}$  acting on P(n) maps  $P^{(2,\infty)} \cup P^{(-\infty,2)}$  to P.

*Proof.* Consider the transformation S(f) = T(f(x - 2)) acting on P(n), and let  $W(x^k) = H_k(x)$ . The key observation (from (9.8.1)) is that

$$S(x^{k}) = T(x-2)^{k} = x^{n-k}H_{k}(0)$$
(9.8.2)

Therefore, if  $g = \sum a_i x^i$ , and the homogenization G is g, then

$$W_*(G) = \sum a_i y^{n-k} H_k(x)$$

so we get the basic relation

$$S(g)(y) = W_*(G)|_{x=0}$$

If  $g \in \mathbf{P}^{\pm}$ , then G is in  $\mathbb{P}_2$ , and  $W_*(G) \in \mathbb{P}_2$ , so  $S(g) \in \mathbf{P}$ . Therefore,  $T(g) \in \mathbf{P}$ .

*Remark* 9.61. We can use (9.8.1) to get precise information about the possible range of S in (9.8.2). Suppose that  $I_n$  is the smallest interval containing all the roots of  $H_n$ . The roots of  $H_n(\frac{\alpha x}{2})$  lie in the interval  $\frac{2}{\alpha}I_n$ . If we knew that T preserves interlacing then from Lemma 6.21 we conclude from (9.8.1) that if  $0 < \alpha < 1$  then

$$\mathsf{T}: \mathbf{P}^{(2+\alpha,\infty)}(\mathfrak{n}) \longrightarrow \mathbf{P}^{\frac{2}{\alpha}\mathbf{I}_{\mathfrak{n}}}$$

# **9.9 Properties of P**<sub>2</sub>

The main result of this section is that the coefficients of a polynomial in  $P_2$  are in P. This allows us to prove that a polynomial is in P by identifying it as a coefficient of a polynomial in  $P_2$ .

**Corollary 9.62.** If  $f \in \mathbf{P}_2$  then all coefficients  $f_i$  in (9.3.2) and all coefficients  $f^i$  in (9.3.3) are in  $\mathbf{P}$ . Moreover,  $f_i \leq f_{i+1}$  and  $f^i \leq f^{i+1}$  for  $0 \leq i < n$ .

*Proof.* If  $f \in \mathbf{P}_2$  then  $\left(\left(\frac{\partial}{\partial y}\right)^i f\right)(x,0) = i!f_i(x)$  and hence  $f_i \in \mathbf{P}$  by Theorem 9.39 and substitution.

Since  $f + \alpha \frac{\partial f}{\partial y}$  is in  $\mathbf{P}_2$  for any  $\alpha$  the coefficient of  $y^i$  is in  $\mathbf{P}$ . This coefficient is  $f_i + \alpha f_{i+1}$  and so we conclude that  $f_{i+1}$  and  $f_i$  interlace. Since the degree of  $f_i$  is greater than the degree of  $f_{i+1}$  we find  $f_i \leq f_{i+1}$ .

There is a simple condition that guarantees that all the coefficients strictly interlace.

**Lemma 9.63.** If  $f(x, y) \in P_2$  and f(x, 0) has all distinct roots, then

- 1. All  $f_i$  have distinct roots.
- 2.  $f_i \leq f_{i+1}$

*Proof.* Assume that  $f_0(r) = f_1(r) = 0$ . Let g(x, y) = f(x + r, y), and write  $g = \sum g_i(x)y^i = \sum g^j(y)x^j = \sum a_{ij}x^iy^j$ . Since  $g_0(0) = g_1(0) = 0$ , we find  $a_{0,0} = a_{0,1} = 0$ . Thus,  $g^0$  is divisible by  $x^2$ . Since  $g^1$  interlaces  $g_0, g^1$  is divisible by x. This implies that  $a_{1,0} = 0$ , and hence  $g_0$  has a double root at zero, but this contradicts the hypothesis that  $f_0$  has no repeated roots since  $g_0(x) = f_0(x+r)$ . Since  $f_0 < f_1$ , it follows that  $f_1$  has all distinct roots. Continuing, we see all  $f_i$  have distinct roots.

The polynomial f(x, y) = (1 + x + y)(1 + 2x + y) shows that f(x, 0) can have all distinct roots, yet f(0, y) can have repeated roots.

**Corollary 9.64.** *If*  $f \in \mathbf{P}_2$  *and*  $f_0 \ll f_1$  *then*  $f_i \ll f_{i+1}$  *for*  $0 \ll i < n$ .

**Lemma 9.65.** If f given in (9.3.2) satisfies y-substitution then the sequence of coefficients is log concave:  $\begin{vmatrix} f_i & f_{i+1} \\ f_{i+1} & f_{i+2} \end{vmatrix} \leq 0$  for  $0 \leq i \leq n-2$ . If  $f_0$  and  $f_1$  have no roots in common then  $\begin{vmatrix} f_i & f_{i+1} \\ f_{i+1} & f_{i+2} \end{vmatrix} < 0$ 

*Proof.* Since f has all real roots for any fixed x, we can apply Newton's inequalities (Corollary 4.11) to find that  $f_j^2(x) > f_{j-1}(x)f_{j+1}(x)$  unless  $f_0(r) = f_1(r) = 0$  for somer in which case we only have  $f_j^2(x) \ge f_{j-1}(x)f_{j+1}(x)$ .

#### Corollary 9.66. Set

$$f(x,y) = \prod_{i=1}^n (x+b_i+c_iy) = \sum_{i=0}^n f_j(x)y^j$$

Assume that the  $c_i$  are positive for  $1 \le i \le n$ . Then the coefficients  $f_j$  are polynomials of degree n - j, and

$$f_0 \underline{\lessdot} f_1 \underline{\lessdot} f_2 \underline{\lessdot} \dots$$

*If all* b<sub>i</sub> *are distinct then the interlacings are all strict.* 

*Proof.* Lemma 9.29 shows that  $f \in \mathbf{P}_2$ . By Corollary 9.64 it remains to show that  $f_0 < f_1$ . If we expand f we find (see Lemma 9.103)

$$f_1(x) = \sum_{j=1}^n c_j \frac{f_0(x)}{x + b_j}$$

Since all coefficients  $c_j$  are positive we can apply Lemma 1.20 to conclude that  $f_0 < f_1$  since  $f_0$  has all distinct roots.

**Corollary 9.67.** If  $f = \sum f_i(x)y^i$  and  $g = \sum g_i(x)y^i$  are both in  $\mathbf{P}_2(n)$  then

$$\sum_{i} f_{i}(x)g_{n-i}(x) \in P$$

*Proof.* The polynomial in question is the coefficient of  $y^n$  in the product fg.

Note that if we take  $f = \sum a_i x^{n-i} y^i$  and  $g = \sum g_i x^{n-i} y^i$  where  $\sum a_i x^i$  and  $\sum b_i x^i$  are both in  $\mathbf{P}^{pos}$  then we conclude that  $\sum a_{n-i} b_i x^{2i} \in \mathbf{P}$ . This is a simple modification of the Hadamard product.

*Remark* 9.68. Since the first two terms of the Taylor series (9.7.1) of f(x+y) are f and yf' it follows that  $f \leq f'$ . Thus, we can prove Rolle's theorem (Theorem 1.1) without analyzing the behavior of the graph of f and f'.

#### 9.10 The analog of $P^{pos}$

A polynomial in **P** with positive leading coefficient is also in  $\mathbf{P}^{\text{pos}}$  if and only all of its coefficients are positive. We define  $\mathbf{P}_2^{\text{pos}}$  in the same way:

 $\mathbf{P}_2^{\text{pos}} = \{ f \in \mathbf{P}_2 \mid \text{ all coefficients of f are positive} \}$ 

We start with a useful alternative criterion.

**Lemma 9.69.** Suppose  $f \in P_2$  and write  $f = \sum f_i(x)y^i$ . If  $f_0 \in \mathbf{P}^{pos}$  then  $f \in \mathbf{P}_2^{pos}$ .

*Proof.* Since  $f_i(x) \leq f_{i+1}(x)$  and since  $f_0(x) \in \mathbf{P}^{pos}$  it follows that all  $f_i$  are in  $\mathbf{P}^{pos}$ . Now the leading coefficients of all  $f_i$ 's are terms of  $f^H$  and are therefore positive. Consequently, all  $f_i$ 's have all positive coefficients, and thus  $f(x, y) \in \mathbf{P}_2^{pos}(n)$ .

We discuss the generalization of  $\mathbf{P}^{\text{pos}}$  to more than two variables in § 10.10.4, and more properties of  $\mathbf{P}_2^{\text{pos}}$  can be found there.

Our next result is that a simple translation takes a polynomial in  $P_2$  to one in  $\mathbf{P}_2^{\text{pos}}$ .

**Lemma 9.70.** *If*  $f(x, y) \in P_2(n)$ *, then there is an*  $\alpha$  *such that*  $f(x + \alpha, y) \in \mathbf{P}_2^{pos}(n)$ .

*Proof.* If  $f(x, y) = \sum f_i(x)y^i$ , then we can choose  $\alpha$  such that  $f_0(x + \alpha)$  is in  $\mathbf{P}^{pos}$ . It follows from Lemma 9.69 that  $f(x + \alpha, y) \in \mathbf{P}_2^{pos}$ .

Interlacing of polynomials in  $\mathbf{P}_2^{\text{pos}}$  stays in  $\mathbf{P}_2^{\text{pos}}$ .

**Lemma 9.71.** *If*  $f \in \mathbf{P}_2^{\text{pos}}$  and  $f \leq g$  then  $g \in \mathbf{P}_2^{\text{pos}}$ 

*Proof.* If we write  $f = \sum f_i(x)y^i$  and  $g = \sum g_i(x)y^i$  then we know that  $f_0 \leq g_0$ . Thus  $g_0 \in \mathbf{P}^{pos}$  and the conclusion follows from Lemma 9.69.

If f is in P<sub>2</sub> then  $f(x, \alpha x) \in \mathbf{P}$  for positive  $\alpha$ , but this can be false for negative  $\alpha$ . However, it is true for  $\mathbf{P}_2^{\text{pos}}$ .

**Proposition 9.72.** *If*  $f(x, y) \in \mathbf{P}_2^{pos}(n)$  *then*  $f(x, \alpha x) \in \mathbf{P}$  *for all*  $\alpha$ *.* 

*Proof.* If suffices to consider  $\alpha < 0$ . We will show that the line  $y = \alpha x$  meets the graph of f in n points. Without loss of generality we may assume that there are r roots of  $f^{H}(x, 1)$  greater than  $\alpha$ , and n - r less than  $\alpha$ , for some r, where  $0 \leq r \leq n$ .

Consider the upper left quadrant. Recall that the solution curves are asymptotic to lines whose slopes are the roots of  $f^{H}(x, 1)$ . Thus, there are n - r solution curves that are eventually above the line  $y = \alpha x$ . Since  $f \in \mathbf{P}_{2}^{\text{pos}}$ , each of these solution curves meets the x-axis in  $(-\infty, 0)$  which is below the line  $y = \alpha x$ . Thus, there are n - r intersection points in the upper left quadrant.

Similarly, there are r intersections in the lower right quadrant. We've found n intersections, and the proposition is proved.  $\Box$ 

Our next result is that all the homogeneous parts of a polynomial in  $P_2^{\text{pos}}$  are in  $P^{\text{pos}}$ . There are simple counterexamples that show that this is not true for arbitrary polynomials in  $P_2$ . The proof uses properties of homogeneous polynomials in three variables - see Lemma 10.37.

**Lemma 9.73.** Suppose  $f \in \mathbf{P}_2^{\text{pos}}(n)$  and  $g_r$  is the polynomial consisting of all terms of total degree r. If we set  $h_r(x) = g_r(x, 1)$  then  $h_r \in \mathbf{P}^{\text{pos}}$  for  $1 \leq r \leq n$  and we have interlacings

$$h_{\mathfrak{n}} \underline{\lessdot} h_{\mathfrak{n}-1} \underline{\lessdot} h_{\mathfrak{n}-2} \underline{\lessdot} \cdots$$

If  $f \in \mathbf{P}_2$  then the graphs of f and  $f^H$  have the same asymptotic behavior. Although these asymptotes do not interlace in general, they do if  $f \in \mathbf{P}_2^{\text{pos}}$ :

# **Lemma 9.74.** If $f \in \mathbf{P}_2^{\text{pos}}$ then $f(x, \alpha) \underline{\ll} f^H(x, \alpha)$ for $\alpha$ sufficiently large.

*Proof.* Assume  $f \in \mathbf{P}_2^{\text{pos}}(n)$  and let  $g_r$  be the homogeneous polynomial consisting of all terms of f of degree r. We can write

$$f(x, \alpha) = \sum_{i=0}^{n} g_i(x, \alpha)$$
(9.10.1)

We know from Lemma 9.73 that  $g_i(x, \alpha) \leq g_{i-1}(x, \alpha)$  for  $1 \leq i \leq n$ . We show that if  $\alpha$  is sufficiently large then  $f(x, \alpha)$  sign interlaces  $f^H(x, \alpha)$  and that the sign of  $f(x, \alpha)$  on the i-th largest root of  $f^H$  is  $(-1)^{n+i}$ .

If  $f^{H}(\beta, 1) = 0$  then  $f^{H}(\alpha\beta, \alpha) = g_{n}(\alpha\beta, \alpha) = 0$  since  $f^{H}$  is homogeneous. Upon computing  $f(\alpha\beta, \alpha)$  we find

$$f(\alpha\beta,\alpha) = 0 + g_{n-1}(\alpha\beta,\alpha) + O(\alpha^{n-2})$$

Now since  $g_n(x, \alpha) \leq g_{n-1}(x, \alpha)$  and the leading coefficients of both  $g_n(x, \alpha)$  and  $g_{n-1}(x, \alpha)$  are positive we know that the sign of  $g_{n-1}(\alpha\beta, \alpha)$  is  $(-1)^{n+i}$  where  $\beta$  is the i-th largest root of  $f^H(x, 1)$ . Since  $g_{n-1}(\alpha\beta, \alpha)$  is  $O(\alpha^{n-1})$  it follows that for  $\alpha$  sufficiently large

$$\operatorname{sgn} g_{n-1}(\alpha\beta, \alpha) = \operatorname{sgn} f(\alpha\beta, \alpha) = (-1)^{n+i}$$

and the conclusion follows.

#### 9.11 The representation of **P**<sub>2</sub> by determinants

The main result of this section is that there are simple determinant representations for polynomials in  $\mathbf{P}_2^{\text{pos}}$  and  $\mathbb{P}_2$ . Before we can prove that polynomials in  $\mathbf{P}_2^{\text{pos}}$  can be represented by determinants, we need the following result due to Vinnikov and Helton, see [79, 112, 174].

**Theorem 9.75.** Suppose that f(x,y) is a polynomial with the property that  $f(\alpha x, \beta x) \in \mathbf{P}$  for all  $\alpha, \beta \in \mathbb{R}$ . If f(0,0) = 1 then there are symmetric matrices B, C such that

$$f(x,y) = |I + xB + yC|.$$

**Theorem 9.76.** If  $f \in \mathbf{P}_2^{\text{pos}}$  and f(0,0) = 1 then there are positive definite symmetric matrices  $D_1, D_2$  so that

$$f(x,y) = |I + xD_1 + yD_2|.$$

*Proof.* Since Proposition 9.72 shows that f(x, y) satisfies the hypothesis of Theorem 9.75, Vinnikov's result shows that there are symmetric matrices B, C so that f(x, y) = |I + xB + yC|. The result now follows from Corollary 9.7.

**Corollary 9.77.** If  $f \in P_2$  then there is a symmetric matrix C, a positive definite diagonal matrix D, and a constant  $\beta$  so that

$$f(x, y) = \beta |xI + yD + C|.$$

*Proof.* There is an a so that  $f(x + a, y) \in \mathbf{P}_2^{\text{pos}}$ , so we can find positive definite matrices  $D_1, D_2$  and a constant  $\alpha = f(a, 0)$  so that

$$\begin{split} f(x+a,y) &= \alpha |I+xD_1+yD_2| \\ f(x,y) &= \alpha |I+(x-a)D_1+yD_2| \end{split}$$

Since  $D_1$  is positive definite, it has a positive definite square root  $D_1^{1/2}$ 

$$f(x,y) = \alpha |D_1| \cdot \left| D_1^{-1/2} (I - \alpha D_1) D_1^{-1/2} + xI + y D_1^{-1/2} D_2 D_1^{-1/2} \right|$$

If we let A be the symmetric matrix  $D_1^{-1/2}(I-aD_1)D_1^{-1/2}$ , B the positive definite matrix  $D_1^{-1/2}D_2D_1^{-1/2}$ , and  $\beta = \alpha |D_1|$  then

$$f(x, y) = \beta |xI + yA + B|$$

Now let  $ODO^{t} = A$  where O is orthogonal and D is diagonal

 $f(x, y) = \beta |xI + yD + O^{t}BO|$ 

Since O<sup>t</sup>BO is symmetric, the corollary is proved.

*Remark* 9.78. We can use the determinant representation of  $P_2$  to show that  $P_2$  is closed under differentiation without using any geometry. Recall that the characteristic polynomials of a symmetric matrix and any of its principle submatrices interlace (see Theorem 1.61).

Suppose that  $f(x, y) \in P_2(n)$ , represent it by |xI + yD + C| as above, and let M = xI + yD + C. Let  $\{d_i\}$  be the diagonal elements of D. Since the only occurrences of x and y in M are on the diagonal, it is easy to see that

$$\frac{\partial}{\partial x} |xI + yD + C| = |M[1]| + |M[2]| + \dots + |M[n]|$$
$$\frac{\partial}{\partial y} |xI + yD + C| = d_1 |M[1]| + d_2 |M[2]| + \dots + d_n |M[n]|$$

If we substitute  $\alpha$  for y we see that

$$|\mathbf{x}\mathbf{I} + \alpha \mathbf{D} + \mathbf{C}| \leq |\mathbf{M}[\mathbf{i}]|(\mathbf{x}, \alpha)$$

since principle submatrices interlace. All the polynomials |M[i]| have positive homogeneous part, so we simply add the interlacings to conclude that  $f(x, \alpha) \leq \frac{\partial}{\partial x} f(x, \alpha)$ . It follows that  $f \leq \frac{\partial f}{\partial x}$ , and in particular the derivative is in P<sub>2</sub>.

#### 9.12 When some coefficients are zero

If f(x) is a polynomial in **P**, then we know two facts about zero coefficients:

- 1. If the constant term is zero then f is divisible by x.
- 2. If two consecutive coefficients are zero, then all the coefficients of terms of smaller degree are zero.

We investigate the implications of zero coefficients for polynomials in P<sub>2</sub>. We will show that if a row (or column) has two consecutive zero coefficients then there is a large triangular region of zero coefficients. For example, the coefficient array of the polynomial  $f(x, y) = (x + y)^3(x + y + 1)^2$  has a triangular block of zeros:

The polynomial  $\partial f/\partial x$  has constant term zero, yet is irreducible.

**Lemma 9.79.** If  $f = \sum a_{i,j}x^iy^j \in P_2$  has a row (or column) with two consecutive zero coefficients, say  $a_{r,s-1} = a_{r,s} = 0$ , then then  $a_{i,j} = 0$  for all  $i + j \leq r + s$ .

*Proof.* We use the fact that if g(x) interlaces  $x^kh(x)$ , then g is divisible by  $x^{k-1}$ . If we write  $f = \sum f_i(x)y^i$  then the hypothesis says that  $f_r$  has two consecutive zeros, so  $f_r$  is divisible by  $x^s$ . We then know that  $f_{r+1}$  is divisible by  $x^{s-1}$ ,  $f_{r+2}$  is divisible by  $x^{s-2}$  and so on.

Next, write  $f = \sum F_j(y)x^j$ . The above paragraph shows that  $a_{r+1,k} = a_{r,k} = 0$  for  $0 \le k \le s-1$ , so  $F_k$  has two consecutive zeros, and hence is divisible by  $x^r$ . Continuing,  $F_{k+1}$  is divisible by  $x^{r-1}$ , and so on. Thus, we've found that all  $a_{ij}$  are zero if  $i + j \le r + s$ .

If we consider polynomials in  $\mathbf{P}_2^{\text{pos}}$  that have a quadrilateral where the strict rhombus inequality is not satisfied, then there is also a large triangular region with zeros.

**Lemma 9.80.** If  $f = \sum a_{i,j} x^i y^j \in \mathbf{P}_2^{pos}$  and there are r, s such that  $a_{r,s} a_{r+1,s+1} - a_{r+1,s} a_{r,s+1} = 0$  then  $a_{i,j} = 0$  for all  $i + j \leq r + s + 1$ .

*Proof.* Write  $f = \sum f_i(x)y^i$ . The expression  $a_{r,s}a_{r+1,s+1} - a_{r+1,s}a_{r,s+1}$  is the determinant of consecutive coefficients of two interlacing polynomials. If it is zero, then by Corollary 4.31 we have  $a_{r,s} = a_{r,s+1} = 0$ . The result now follows from Lemma 9.79.

**Corollary 9.81.** If  $f = \sum a_{i,j} x^i y^j \in \mathbf{P}_2^{pos}$  and  $f(0,0) \neq 0$  then all rhombus inequalities are strict. As with one variable, we find that agreement at two consecutive coefficients implies lots of agreement.

**Corollary 9.82.** Suppose that  $f \leq g$  in  $\mathbf{P}_2^{pos}$ ,  $f = \sum a_{i,j} x^i y^j$ , and  $g = \sum b_{i,j} x^i y^j$ . If there are r, s so that  $a_{r,s} = b_{r,s}$  and  $a_{r+1,s} = b_{r+1,s}$  then  $a_{i,j} = b_{i,j}$  for  $i + j \leq r + s + 1$ .

*Proof.* f - g is in  $\mathbf{P}_2^{\text{pos}}$ , and has two consecutive zeros.

We can replace the consecutive zero condition with equal minimum degree.

**Lemma 9.83.** Suppose that  $f = \sum f_i(x)y^i \in P_2$ , and let d be the minimum x-degree. If two consecutive  $f_i$  have degree d then so do all earlier ones.

*Proof.* We first differentiate d times with respect to f, and so assume that two consecutive coefficients are zero. By Lemma 4.10 all earlier coefficients are zero for all x, and so are identically zero.

#### 9.13 The Hadamard product

We can realize the Hadamard product as a coefficient in a polynomial in two variables. Begin with two polynomials of the same degree:

$$\begin{split} f(x) &= a_0 + a_1 x + a_2 x^2 + \dots + a_n x^n \\ g(x) &= b_0 + b_1 x + b_2 x^2 + \dots + b_n x^n \end{split}$$

Now reverse g and homogenize it.

$$G(x, y) = b_n x^n + b_{n-1} x^{n-1} y + b_{n-2} x^{n-2} y^2 + \dots + b_0 y^n.$$

If we multiply, and write as a series in y

$$f(y)G(x,y) = a_0b_nx^n + (a_1b_nx^n + a_0b_{n-1}x^{n-1})y + ... + (a_0b_0 + \dots + a_nb_nx^n)y^n + ... + a_nb_ny^{2n}$$

then the Hadamard product f \* g is the coefficient of  $y^n$ .

**Theorem 9.84.** The Hadamard product<sup>4</sup> of polynomials defines a bilinear map  $P(n) \times P^{\pm}(n) \longrightarrow P(n)$ . If  $f_1 \ll f_2$  then  $f_1 * g \ll f_2 * g$ .

<sup>&</sup>lt;sup>4</sup>*It does not define a map*  $P \times P \longrightarrow P$ . *For instance*  $(x^2 - 1) * (x^2 - 1) = x^2 + 1$  *is not in* P.

*Proof.* Without loss of generality we may we assume that  $g \in \mathbf{P}^{\text{pos}}$ . Since  $g \in \mathbf{P}^{\text{pos}}$  its reversal is in  $\mathbf{P}^{\text{pos}}$ , and the homogenization G(x, y) is in  $\mathbf{P}_2$ . Since f \* g is a coefficient of f(y)G(x, y) it is in  $\mathbf{P}$ . Interlacing follows since for fixed g the map  $f \mapsto f * g$  is a linear map  $\mathbf{P} \longrightarrow \mathbf{P}$ .

Actually, f(y) is not in  $P_2$ , but rather in the closure of  $P_2$  (see Chapter 11). However, the coefficients of f(y)G(x, y) are in **P**.

If we look more closely at the product f(y)G(x, y) we see that it equals

$$\cdots + (f * xg) y^{n-1} + (f * g) y^n + \frac{1}{x} (xf * g) y^{n+1} + \cdots$$

Consequently,

**Corollary 9.85.** *If*  $f, g, \in P^{pos}(n)$  *then* 

(1) 
$$(f * g)^2 \ge \left(\frac{n+1}{n}\right)^2 \frac{1}{x} (f * xg)(xf * g) \quad \text{for } x \ne 0$$
  
(2)  $\begin{vmatrix} f * g & xf * g \\ f * xg & xf * xg \end{vmatrix} \ge 0 \quad \text{for } x \ge 0$   
(3)  $f * xg - \frac{1}{x} (xf * g) \le f * g$ 

*Proof.* The first part is Newton's inequality (p. 106). The second part follows from the first and the identity xf \* xg = x(f \* g). Since f \* g and  $\frac{1}{x}(xf * g)$  interlace, and  $\frac{1}{x}(xf * g)$  has degree n - 1 we have the interlacings

$$f * xg \underline{\ll} f * g \underline{\ll} \frac{1}{x} (xf * g)$$

which implies the last part.

If we rewrite the last part in terms of coefficients we get

**Corollary 9.86.** If  $f = \sum a_i x^i$ ,  $g = \sum b_i x^i$  and  $f, g \in P^{\text{pos}}$  then

$$\sum a_i(b_{i+1}-b_i)x^i \in \boldsymbol{P}.$$

Surprising, we can do better than Theorem 9.84. We use a differential operator in  $P_2$ , and then evaluate it at 0 to get a polynomial in P. See Corollary 9.100 for an alternate proof. Define

$$\mathbf{x}^{i} *' \mathbf{x}^{j} = \begin{cases} i! \mathbf{x}^{i} & i = j \\ 0 & \text{otherwise} \end{cases}$$

Notice that  $f *' g = EXP^{-1} f * g$ . In terms of coefficients

$$f *' g = a_0b_0 + 1!a_1b_1x + 2!a_2b_2x^2 + \dots + n!a_nb_nx^n$$

**Theorem 9.87.** If  $f \in \mathbf{P}^{\pm}(n)$ ,  $g \in \mathbf{P}(n)$  then  $f *' g \in \mathbf{P}(n)$ .

*Proof.* Again assume that  $f \in \mathbf{P}^{pos}$ . Let G(x, y) be the homogeneous polynomial in  $\mathbf{P}_2$  corresponding to g. By Lemma 10.62 we know  $f(\frac{\partial}{\partial x})G \in \mathbf{P}_2$ , so evaluation at x = 0 gives a polynomial in  $\mathbf{P}$ . Consequently, the map  $f \times g \mapsto (f(\frac{\partial}{\partial x})G)(0, y)$  defines a linear transformation  $\mathbf{P}^{\pm}(n) \times \mathbf{P}(n) \longrightarrow \mathbf{P}(n)$ .

The polynomial f \*' g is the reverse of  $(f(\frac{\partial}{\partial x})G)(0,x)$ . To verify this, we check it for  $f = x^r$  and  $g = x^s$ :

$$\left(\frac{\partial}{\partial x}\right)^{r} \left(x^{s} y^{n-s}\right)(0,x) = \begin{cases} 0 & r \neq s \\ r! x^{n-r} & r = s \end{cases}$$

**Corollary 9.88.** If  $f \in \mathbf{P}^{\pm}(n)$ ,  $g_1 \underline{\ll} g_2 \in \mathbf{P}(n)$  then

$$f *' g_1 \underline{\ll} f *' g_2$$

The last lemma showed that  $\mathbf{P}^{\pm}(\mathfrak{n}) * \mathbf{P}(\mathfrak{n}) \subset EXP(\mathbf{P})$ . If we allow analytic functions then we get equality.

# Corollary 9.89. $\widehat{P^{\text{pos}}} * P = \text{EXP}(P)$

*Proof.* By taking limits we know that  $\widehat{\mathbf{P}^{\text{pos}}} * \mathbf{P} \subset \text{EXP}(\mathbf{P})$ . Since  $\text{EXP}(f) = e^x * f$  we see the the containment is an equality.

There is a partial converse to Theorem 9.87.

**Proposition 9.90.** Suppose that f is a polynomial. If for all  $g \in P^{\text{pos}}$  we have that  $f * g \in P$ , then  $\text{EXPf} \in P$ . If for all  $g \in P$  we have that  $f * g \in P$ , then  $\text{EXPf} \in \overline{P^{\pm}}$ .

*Proof.* Since  $(1 + x/n)^n \in \mathbf{P}^{pos}$  we know  $f * (1 + x/n)^n \in \mathbf{P}$ . Now  $(1 + x/n)^n$  converges uniformly to  $e^x$ , so  $f * e^x \in \mathbf{P}$ . The first part follows since  $f * e^x = \exp(f)$ , so it remains to show that  $f \in \mathbf{P}^{\pm}$  in the second case.

If  $f = \sum a_i x^i$  then  $f * x^i (x^2 - 1) = x^i (a_{i+2}x^2 - a_i)$  so it follows that  $a_i$ and  $a_{i+2}$  have the same sign. If  $a_{i+2}$  is non-zero, and  $a_{i+1} = 0$  then consider  $f * x^i (x + 1)^2 = x^i (a_{i+2}x^2 + a_i)$ . Since  $a_{i+2}$  and  $a_i$  have the same sign, it follows that  $a_i = 0$ . If we consider  $f * x^{i-j+2}(x+1)^j$  we see that all coefficients  $a_{i+1}, a_i, a_{i-1}, \ldots$  are zero. Thus,  $f = x^i f_1$  where  $f_1$  has all non-zero terms. Since the signs of the coefficients of  $f_1$  are either the same or are alternating, we get that  $f_1 \in \mathbf{P}^{\pm}$ .

As an application of Theorem 9.87 consider

**Lemma 9.91.** If  $f \in P$  then  $(1 - x)^{-n} * f$  is in P for n = 0, 1, 2, ...

*Proof.* From (7.10.6) we know that  $EXP(1-x)^{-n}$  is in  $\mathbf{P}^{pos}$  since  $e^x \in \mathbf{P}^{pos}$  and  $L_n(-x) \in \mathbf{P}^{pos}$ . The conclusion follows from Corollary 9.89.

The lemma explains the curious fact that  $f \mapsto (1 + x + x^2 + ...) * f$  sends **P** to itself, yet  $(1 + x + x^2 + ...) = 1/(1 - x)$  is not in  $\widehat{\mathbf{P}}$ .

If we re-express the lemma in terms of coefficients we find that the map

$$x^{i} \mapsto \frac{\langle \underline{n} \rangle_{i}}{i!} x^{i}$$
 (9.13.1)

which is just Theorem 7.1.

#### 9.14 Some differential operators

We use the fact that polynomials in **P** factor into linear factors to get differential operators on **P**<sub>2</sub>. We would like to know that if  $f \in \mathbf{P}_2$  then the differential operator  $f(\frac{\partial}{\partial x}, \frac{\partial}{\partial y})$  maps **P**<sub>2</sub> to itself, but the best we are able to prove is when f is a product of linear factors - see Lemma 10.62.

**Lemma 9.92.** If  $f \in \mathbf{P}$ ,  $g \in \mathbf{P}_2$  then  $f(\frac{\partial}{\partial x})g \in \mathbf{P}_2$ .

*Proof.* It suffices to show that  $g + \alpha \frac{\partial g}{\partial x}$  is in  $\mathbf{P}_2$ , but this is immediate from  $g \leq \frac{\partial g}{\partial x}$ .

Here is another differential operator that acts on  $P_2$ .

**Corollary 9.93.** If  $f \in P^{pos}$  and  $g \in P_2$  then  $f\left(-\frac{\partial^2}{\partial x \partial y}\right)g \in P_2$ .

*Proof.* If suffices to show that  $h(x, y) = \left(a - \frac{\partial^2}{\partial x \partial y}\right)g$  is in  $P_2$  for positive a. Since homogeneity is certainly satisfied since  $h^H = ag^H$  it suffices to check substitution. If  $\alpha \in \mathbb{R}$  then

$$\left(a - \frac{\partial^2}{\partial x \, \partial y}\right) g(x, \alpha) = ag(x, \alpha) - \frac{\partial}{\partial x} \left(\frac{\partial}{\partial y}g\right)(x, \alpha)$$

and the conclusion follows from  $g(x, \alpha) \leq \frac{\partial g}{\partial y}(x, \alpha)$  and Corollary 2.10.

**Example 9.94.** If we take  $f = (x - 1)^m$  and  $g = (x + y + 1)^n$  then

$$\left(\frac{\partial^2}{\partial x \, \partial y} - 1\right)^m (x + y + 1)^n = \sum_{i=0}^n (-1)^{m-i} \binom{m}{i} \left(\frac{\partial^2}{\partial x \, \partial y}\right)^i (x + y + 1)^n$$
$$= \sum_{2i \leqslant n} \binom{m}{i} (\underline{n})_{2i} (x + y + 1)^{n-2i}$$

and so we find that

$$\sum_{2i \leqslant n} (-1)^{m-i} \frac{(x+y+1)^{n-2i}}{i!(m-i)!(n-2i)!} \in \mathbf{P}_2$$

The differential operator  $g(x\frac{\partial}{\partial y})$  maps **P**<sub>2</sub> to itself.

**Lemma 9.95.** If  $g \in \mathbf{P}^{pos}$ ,  $f \in \mathbf{P}_2$  then  $g(x\frac{\partial}{\partial y}) f(x, y) \in \mathbf{P}_2$ .

*Proof.* Since all coefficients of g are positive the homogeneous part of  $g(x\frac{\partial}{\partial y}) f(x, y)$  has all positive coefficients. To show that  $g(\alpha \frac{\partial}{\partial y}) f(\alpha, y) \in \mathbf{P}$  it suffices to note that  $(b + \alpha \frac{\partial}{\partial y}) f(\alpha, y) \in \mathbf{P}$ .

**Example 9.96.** If we take  $g = (x + 1)^m$  and  $f = (x + y + 1)^n$  then

$$(x\frac{\partial}{\partial y}+1)^{\mathfrak{m}} (x+y+1)^{\mathfrak{n}} = \sum_{i=0}^{\mathfrak{m}} \binom{\mathfrak{m}}{i} (x\frac{\partial}{\partial y})^{i} (x+y+1)^{\mathfrak{n}}$$
$$= \sum_{i=0}^{\mathfrak{m}} \binom{\mathfrak{m}}{i} x^{i} (x+y+1)^{\mathfrak{n}-i} (\underline{\mathfrak{n}})_{i}$$

and so

$$\sum_{i=0}^{n} \binom{m}{i} x^{i} (x+y+1)^{n-i} (\underline{n})_{i} \in \mathbf{P}_{2}$$

**Corollary 9.97.** If  $f \in \mathbf{P}(n)$ ,  $g \in \mathbf{P}^{pos}(n)$ ,  $g = \sum b_i x^i$  then

$$h(x,y) = \sum b_i x^i f^{(i)}(y) \in \mathbf{P}_2$$

Proof. The homogeneous part of h is

$$a_n \sum b_i x^i(\underline{n})_i y^{n-i}$$

where  $a_n$  is the leading coefficient of f. Since  $h^H(x, 1) = a_n n! EXP(g)$  it follows that  $h^H$  is in **P**. Since all coefficients  $b_i$  are the same sign  $h^H(x, 1)$  is actually in  $\mathbf{P}^{pos}$ . We now apply Theorem 9.87 for fixed y, which shows that substitution holds.

Corollary 9.98 (Hermite-Poulin). If f, g are given in Corollary 9.97 then

$$\sum b_{i} f^{(i)}(y) \in \boldsymbol{P}^{pos}$$

Corollary 9.99 (Schur-Pólya). If f, g are given in Corollary 9.97 then

$$\sum b_{i} x^{i} f^{(i)}(x) \in \boldsymbol{P}$$

Here is a proof of a special case of Theorem 9.87.

**Corollary 9.100** (Schur). If f,  $g \in P^{pos}$ ,  $f = \sum a_i x^i g = \sum b_i x^i$ , then

$$\sum \mathfrak{i}!\,\mathfrak{a}_{\mathfrak{i}}\,\mathfrak{b}_{\mathfrak{i}}\,\mathfrak{x}^{\mathfrak{i}}\in P$$

*Proof.* Let G(x, y) be the homogenization of g. Since  $g \in \mathbf{P}^{p \circ s}$  we know that  $G(x, y) \in \mathbf{P}_2$ . Since  $f^{(i)}(0) = i!a_i$  it follows that

$$\sum i! a_i b_i x^i = f(x \frac{\partial}{\partial y}) G(x, y)|_{y=0}$$

and hence is in **P**.

It is possible for two different linear transformations  $P \longrightarrow P$  to agree on all polynomials up to degree n. For instance, suppose that f has degree n + 1 and all distinct roots. Choose  $g \in P$  such that  $f - g = \alpha x^{n+1}$  for some non-zero  $\alpha$ . If we define

$$S(h) = f(D)h$$
 and  $T(h) = g(D)h$ 

then  $(T - S)(h) = \alpha D^{n+1}h$ , and this is zero if the degree of h is at most n. Differential operators satisfy a uniqueness condition.

**Lemma 9.101.** Suppose that T(g) = f(D)g where  $f \in P$  has degree n. If S is a linear transformation  $P \longrightarrow P$  such that

$$S(x^i) = T(x^i)$$
 for  $0 \le i \le n+2$ 

then S = T.

*Proof.* Since  $(x - t)^{n+3} \leq (x - t)^{n+2}$  for any real t we have

$$S(x-t)^{n+3} \sim S(x-t)^{n+2} = T(x-t)^{n+2}$$

f has degree n so  $T(x-t)^{n+2}$  has a factor of  $(x-t)^2$  and therefore  $S(x-t)^{n+3}$  has t as a root. Thus

$$\begin{split} 0 &= S(x-t)^{n+3}(t) = S(x^{n+3})(t) + \sum_{0}^{n+2} \binom{n+3}{i} (-t)^{n+3-i} S(x^{i})(t) \\ &= S(x^{n+3})(t) + \sum_{0}^{n+2} \binom{n+3}{i} (-t)^{n+3-i} T(x^{i})(t) \\ &= S(x^{n+3})(t) + \left[ T(x-t)^{n+3}(t) - T(x^{n+3})(t) \right] \end{split}$$

Now t is a root of  $T(x - t)^{n+3}$  so we conclude that  $T(x^{n+3}) = S(x^{n+3})$ .

If we take f = 1 we get

**Corollary 9.102.** If T is a linear transformation  $P \longrightarrow P$  such that  $T(x^i) = x^i$  for i = 0, 1, 2 then T is the identity.

# 9.15 Realizing interlacing

There is a close relationship between interlacing and products: A pair of interlacing polynomials can be realized as the first two coefficients of a product. If the degrees are different the representation is essentially unique. A product is also determined by its first term and its homogeneous part.

**Lemma 9.103.** Assume that  $f_0 \in P(n)$  is monic,  $f_1$  has positive leading coefficient, and  $f_0 < f_1$ . There exist unique  $b_i, c_i$  such that (9.15.1) is satisfied, and all  $c_i$  are positive. The product is in  $P_2$ .

$$\prod_{i=1}^{n} (x + b_i + c_i y) = f_0(x) + f_1(x)y + \cdots$$
 (9.15.1)

*Proof.* Expanding the product, we find

$$\begin{split} f_0(x) &= \prod_{i=1}^n (x+b_i) \\ f_1(x) &= \sum_{j=1}^n \, c_j \prod_{i\neq j}^n (x+b_i) \end{split}$$

We can rewrite  $f_1$  in terms of  $f_0$ 

$$f_1(x) = \sum_{j=1}^n c_j \frac{f_0(x)}{x + b_j}$$

If we are given  $f_0$ ,  $f_1$  then we find the  $b_i$  as roots of  $f_0$ , and the  $c_i$  as the coefficients of  $f_1$  as in Lemma 1.20.

*Remark* 9.104. There is straightforward way to express  $f \leq g$  as a limit of strict interlacings. We can find non-negative  $a_i$  such that

$$\prod (x + a_i y + b_i) = f + y g + \cdots$$

It's enough to perturb the constants. For positive  $\epsilon$  and *distinct*  $\epsilon_i$  define

$$\prod (x + (a_i + \varepsilon) y + (b_i + \varepsilon_i)) = f_{\varepsilon} + y g_{\varepsilon} + \cdots$$

We claim that have that

- 1.  $f_{\varepsilon} < g_{\varepsilon}$  for all positive  $\varepsilon$  and sufficiently small  $\varepsilon_i$ .
- $\begin{array}{l} 2. \quad \lim_{\varepsilon, \varepsilon_{\mathfrak{i}} \to 0^{+}} f_{\varepsilon} = f \\ 3. \quad \lim_{\varepsilon, \varepsilon_{\mathfrak{i}} \to 0^{+}} g_{\varepsilon} = g \end{array} \end{array}$

We only need to check the first claim. Since the roots of  $f_{\varepsilon}$  are  $-(b_i + \varepsilon_i)$  and the  $\varepsilon_i$  are distinct, we see that if the  $\varepsilon_i$  are small enough then all roots of  $f_{\varepsilon}$  are distinct. Since  $a_i + \varepsilon$  is positive it follows that  $f_{\varepsilon} < g_{\varepsilon}$ .

We can also specify the first term and the homogeneous part.

**Lemma 9.105.** If  $h \in P^{pos}$  and  $f_0 \in P$  are monic of the same degree then there is a product

$$g(x,y) = \prod_{i} (a_iy + x + b_i) = f_0(x) + f_1(x)y + \cdots$$

such that  $g \in \mathbf{P}_2$  and  $g^H(x, 1) = h(x)$ .

*Proof.* If we write  $h(x) = \prod (x+a_i)$  and  $f_0(x) = \prod (x+b_i)$  then any ordering of the  $a_i$ 's and  $b_i$ 's gives a product. Since all  $a_i$  have the same sign, the product is in  $P_2$ .

As opposed to the unique representation of  $f \leq g$  as a product, we see from Lemma 9.106 that  $f \ll g$  are the first two coefficients of many products.

**Lemma 9.106.** Given  $f_1 \ll f_0$  there exist  $b_i$ ,  $c_i$ ,  $d_i$  such that (9.15.2) is satisfied, and the determinants  $\begin{vmatrix} 1 & b_i \\ c_i & d_i \end{vmatrix}$  are negative for all *i*. The product is in the closure of  $P_2$ .

$$F(x,y) = \prod_{i=1}^{n} (x + b_i + c_i y + d_i x y) = f_0(x) + f_1(x)y + \cdots$$
(9.15.2)

*Proof.* Define  $\alpha_i$ ,  $\beta_i$ ,  $\beta$  by

$$\begin{split} f_0(x) &= \prod_{i=1}^n (x+\alpha_i) \\ f_1(x) &= \beta f_0(x) + \sum_{i=1}^n \beta_i \frac{f_0(x)}{x+\alpha_i} \end{split}$$

Choose  $d_i$  so that  $\sum_{i=1}^n d_i = \beta.$  We see that  $f_0, f_1$  are the leading terms of the product

$$\prod_{i=1}^n (x+\alpha_i+d_ixy+(\beta_i+\alpha_id_i)y)$$

since the first term is obviously  $f_0$  and the second term is

$$\sum_{i} \frac{f_0}{x + \alpha_i} (d_i x + \beta_i + \alpha_i d_i) = \left(\sum_{i} d_i\right) f_0 + \sum_{i} \beta_i \frac{f_0}{x + \alpha_i}$$

Identify  $b_i = \alpha_i, c_i = \beta_i + \alpha_i d_i$  and note that the determinant is equal to  $\beta_i$  which is negative since  $f_1 \ll f_0$ . See see Lemma 11.28 and Corollary 11.7 for similar results.

Another consequence of looking at coefficients of products is the following lemma, which gives a "canonical" polynomial that interlaces two polynomials.

**Lemma 9.107.** *If*  $f = \prod (x+b_i)$  *and*  $c_i$ ,  $d_i$  *are non-negative, then we have a diagram of interlacing polynomials* 



*Proof.* Lemma 10.13 shows that Theorem 9.113 holds for arbitrary products, so the interlacings follow from the fact that the four terms in the diagram are the coefficients of 1, y, z, yz in the product

$$\prod_{i}(x+c_{i}y+d_{i}z+b_{i})$$

Another interpretation of this lemma is that if we are given f, g, h with positive leading coefficients, and f < g, h then there is a canonical choice of k such that f < g, k where all interlacings are <. In the case that g = h = f'

the canonical polynomial is  $\frac{f''}{2}$ . Since these polynomials are the coefficients of a polynomial in **P**<sub>3</sub> we will see that their determinant is negative. We can verify this directly:

$$\begin{vmatrix} f & \sum_{i} c_{i} \frac{f}{x+b_{i}} \\ \left| \sum_{j} d_{j} \frac{f}{x+b_{j}} & \sum_{i \neq j} c_{i} d_{j} \frac{f}{(x+b_{i})(x+b_{j})} \end{vmatrix} = \\ f^{2} \left( \sum_{i \neq j} \frac{c_{i} d_{j}}{(x+b_{i})(x+b_{j})} - \sum_{i,j} \frac{c_{i} d_{j}}{(x+b_{i})(x+b_{j})} \right) \\ = -f^{2} \left( \sum_{i} \frac{c_{i} d_{i}}{(x+b_{i})^{2}} \right) < 0$$

#### 9.16 Karlin's conjecture

We show that certain matrices formed from the coefficients of a polynomial in  $\mathbf{P}_2^{\text{pos}}$  are totally positive for positive values of x. A special case of this yields a solution to a version of Karlin's problem [41,47,100].

A Hankel matrix has equal entries on the minor diagonals. Given a sequence  $\mathbf{a} = (\alpha_1, \alpha_2, ...)$  and a positive integer d we can form the Hankel matrix  $H(\mathbf{a}; d)$ :

$$H(\boldsymbol{a};\boldsymbol{d}) = \begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \dots & \alpha_d \\ \alpha_2 & \alpha_3 & \alpha_4 & \dots & \alpha_{d+1} \\ \alpha_3 & \alpha_4 & \alpha_5 & \dots & \alpha_{d+2} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \alpha_d & \alpha_{d+1} & \alpha_{d+2} & \dots & \alpha_{2d} \end{pmatrix}$$

We are also interested in the Toeplitz matrix  $T(\mathbf{a}; d)$  associated to  $H(\mathbf{a}; d)$ . If we reverse all the columns of  $H(\mathbf{a}; d)$  we get  $T(\mathbf{a}; d)$ :

$$\mathsf{T}(\mathbf{a}; \mathbf{d}) = \begin{pmatrix} \alpha_{d} & \alpha_{d-1} & \alpha_{d-2} & \dots & \alpha_{1} \\ \alpha_{d+1} & \alpha_{d} & \alpha_{d-1} & \dots & \alpha_{2} \\ \alpha_{d+2} & \alpha_{d+1} & \alpha_{d} & \dots & \alpha_{3} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \alpha_{2d} & \alpha_{2d-1} & \alpha_{2d-2} & \dots & \alpha_{d} \end{pmatrix}$$

We can form  $T(\mathbf{a}; d)$  from  $H(\mathbf{a}; d)$  by shifting the rightmost column d - 1 places, the previous column d - 2 places, and so on. Consequently, we note the useful relation

$$\det (\mathsf{T}(\mathbf{a}; \mathbf{d})) = (-1)^{\binom{d}{2}} \det (\mathsf{H}(\mathbf{a}; \mathbf{d}))$$

The matrix  $\varphi(f)$  is a Toeplitz matrix. The first result picks out a submatrix of  $\varphi(f)$ .

**Lemma 9.108.** Suppose that  $f = a_0 + a_1x + a_2x^2 + \cdots \in P^{p \circ s}$ . For any positive integer the matrix  $T((a_0, a_1, \ldots); d)$  is totally positive.

*Proof.* The matrix  $T((a_0, a_1, ...); d)$ , with rows and columns reversed, is a submatrix of  $\varphi(f)$ .

We have an important consequence of this simple observation.

**Theorem 9.109.** Suppose that  $f(x, y) = \sum f_i(x)y^i \in \mathbf{P}_2^{pos}$ . The Toeplitz matrix  $T((f_0(x), f_1(x), f_2(x), ...); d)$  is totally positive for all non-negative x. In addition

$$(-1)^{\binom{d}{2}} \det H((f_0(x), f_1(x), f_2(x), \dots); d) \ge 0$$
 for  $x \ge 0$ 

**Corollary 9.110.** *If*  $f \in P^{pos}$  *then* 

(1) T(  $(f(x), \frac{f'(x)}{1!}, \frac{f^{(2)}(x)}{2!}, \frac{f^{(3)}(x)}{3!}, \dots)$  ); d) is totally positive for positive x.

(2) 
$$(-1)^{\binom{d}{2}} \det H((f(x), \frac{f'(x)}{1!}, \frac{f^{(2)}(x)}{2!}, \frac{f^{(3)}(x)}{3!}, \dots); d) \ge 0 \text{ for } x \ge 0.$$

*Proof.* Use the Taylor series for f(x + y) and apply Theorem 9.109.

We can restate the last part by saying that the sign of the determinant of the matrix whose ij entry is  $\frac{f^{(i+j)}}{(i+j)!}$  is  $(-1)^{\binom{d}{2}}$  whenever  $f \in \mathbf{P}^{pos}$ , and  $x \ge 0$ . Karlin's conjecture [41] is that

$$(-1)^{\binom{d}{2}} \det H((f(x), f'(x), f^{(2)}(x), f^{(3)}(x), \dots)); d) \ge 0 \text{ for } x \ge 0$$
(9.16.1)

It is easy to construct counterexamples to this.<sup>5</sup> Karlin's conjecture is true in special cases. Here's an example from [47].

**Corollary 9.111.** Define  $EXP_*^{-1}(x^iy^j) = j! x^iy^j$ , and let  $f \in \mathbf{P}^{pos}$ . If  $EXP_*^{-1}(f(x + y)) \in \mathbf{P}_2$  then Karlin's conjecture (9.16.1) holds.

We can also apply Theorem 4.23 to determinants of translates of a function. Suppose  $\mathbb{A}(x) = x + 1$  and let  $T(y^n) = (\underline{y})_n$ . Recall that

$$\sum_{i} i^n \frac{y^i}{i!} = e^y T^{-1}(y^n).$$

We want to determine the action of  $e^{Ay}$  on polynomials:

$$e^{\mathbb{A}y}x^{n} = \sum_{i} \frac{y^{i}}{i!}(x+i)^{n}$$
$$= \sum_{r,i} {n \choose r} x^{r} i^{n-r} \frac{y^{i}}{i!}$$
$$= \sum_{r} {n \choose r} x^{r} \sum_{i} i^{n-r} \frac{y^{i}}{i!}$$
$$= e^{y} \sum_{r} {n \choose r} x^{r} T^{-1} y^{n-r}$$
$$= e^{y} T_{*}^{-1} (x+y)^{n}$$

where  $T_*^{-1}$  is the induced map. By linearity

$$e^{\mathbb{A}y}f(x) = e^{y}T_{*}^{-1}f(x+y)$$
$$= \sum_{k=0}^{\infty} \frac{\mathbb{A}^{k}f(x)}{k!}y^{k}$$

 ${}^{5}f = (x + 1)(x + 2)(x + 3)$  and d = 3 is one.

Now we know that  $T^{-1}: \mathbf{P}^{pos} \longrightarrow \mathbf{P}^{pos}$  so for any positive value  $\alpha$  of x

$$\left[\mathsf{T}_*^{-1}\mathsf{f}(\mathsf{x}+\mathsf{y})\right](\alpha,\mathsf{y}) = \mathsf{T}^{-1}\mathsf{f}(\alpha+\mathsf{y}) \in \mathbf{P}^{\mathsf{pos}}$$

since  $f(\alpha + y) \in \mathbf{P}^{pos}$  and consequently  $e^{y}T_{*}^{-1}f(x+y) \in \widehat{\mathbf{P}^{pos}}$  for any positive x. We can now apply Theorem 4.23 to conclude

**Corollary 9.112.** If  $f \in P^{pos}$  then  $T((f, Af, A^2f/2!, A^3f/3!, ...); d)$  is totally positive for all  $x \ge 0$ .

# 9.17 Determinants of coefficients

In Lemma 4.34 we saw that the signs of certain two by two determinants of interlacing polynomials were always positive. Since any interlacing  $f \ll g$  can be viewed as the first two coefficients of a polynomial in  $P_2$  by Lemma 9.103, we can generalize these results to determinants of coefficients of a polynomial in  $P_2$ .

So, let's begin by considering the coefficients of  $x^i y^j$  in a product.

**Theorem 9.113.** Suppose  $f \in \mathbf{P}_2$ , f(x, 0) has all distinct roots, and set

$$f(x,y) = \sum_{i,j} d_{ij} x^{i} y^{j} = \sum_{i} f_{i}(x) y^{i}$$
(9.17.1)

*For*  $1 \leq i, j \leq n$  *we have* 

$$\begin{array}{c} (\text{rows}) & \left| \begin{array}{c} d_{ij} & d_{i+1,j} \\ d_{i+1,j} & d_{i+2,j} \end{array} \right| < 0 \\ (\text{columns}) & \left| \begin{array}{c} d_{ij} & d_{i,j+1} \\ d_{i,j+1} & d_{i,j+2} \end{array} \right| \leqslant 0 \\ (\text{squares}) & \left| \begin{array}{c} d_{ij} & d_{i+1,j} \\ d_{i,j+1} & d_{i+1,j+1} \end{array} \right| < 0 \\ \end{array}$$

If the roots of f(x, 0) aren't all distinct then all < signs are replaced with  $\leq$ .

*Proof.* We know that  $f_0 \leq f_1 \leq \cdots$ . The (rows) inequalities are Newton's inequalities since all  $f_i \in \mathbf{P}$ . The (columns) results follow upon writing the sum as  $\sum g_i(y)x^i$ . These inequalities are  $\leq$  since it necessarily true that f(0, y) has all distinct roots - consider (x + y + 1)(x + 2y + 1). The (squares) are negative since consecutive  $f_i$ 's interlace (Corollary 4.31).

*Remark* 9.114. We can also derive this from Lemma 9.19 in the same way that Newton's inequalities are derived from properties of the quadratic.

There are polynomials in P<sub>2</sub> for which all terms of odd degree are zero, For example, consider the coefficient array of  $((x + y)^2 - 1)^3$ .

In such a case we always have alternating signs.

**Lemma 9.115.** If  $f(x, y) \in P_2$  has the property that all the coefficients of odd degree are zero then the signs of the non-zero coefficients alternate.

*Proof.* If  $f(x, y) = \sum f_i(x)y^i$  then Lemma 2.5 implies that the signs of the non-zero coefficients of each  $f_i$  alternate. Writing  $f(x, y) = \sum g_i(y)x^i$  shows that the signs of each  $g_i$  alternate. Thus all the signs in the even rows and columns alternate, as do the signs in the odd rows and columns.

If  $f = \sum a_{ij}x^iy^j$  then the quadralateral inequality applied to the coefficients  $\{a_{00}, a_{10}, a_{01}, a_{11}\}$  yie;ds  $a_{00}a_{11} - a_{10}a_{01} < 0$ . Since  $a_{01} = a_{10} = 0$  we conclude that  $a_{00}$  and  $a_{11}$  have opposite signs. Thus, all non-zero signs alternate.

**Example 9.116.** When  $f \in \mathbf{P}_2$  is given by a determinant then the positivity of certain determinants in Theorem 9.113 is a consequence of a classical determinantal identity. Suppose f = |xD + yE + C| where D, E are diagonal matrices with positive entries, and C is symmetric. Assume the coefficients are given the expansion in (9.17.1). Let  $\Delta = \det(C)$ ,  $\Delta_i = \det(C[i])$ , and  $\Delta_{ij} = \det(C[i,j])$  where C[i] is C with the i-th row and column deleted. Then

$$\begin{split} d_{00} &= \Delta \\ d_{01} &= \sum d_i \Delta_i \\ d_{10} &= \sum e_i \Delta_i \\ d_{11} &= \sum_{i \neq j} d_i e_j \Delta_{ij} \\ \left| \begin{array}{c} d_{00} & d_{10} \\ d_{01} & d_{11} \end{array} \right| &= \sum_{i \neq j} \Delta \Delta_{ij} d_i e_j - \sum_{i,j} \Delta_i \Delta_j d_i e_j \\ &= \sum_{i \neq j} \left| \begin{array}{c} \Delta & \Delta_i \\ \Delta_j & \Delta_{ij} \end{array} \right| c_i d_j - \sum_i \Delta_i^2 d_i e_i \leqslant 0 \end{split}$$

since  $\begin{vmatrix} \Delta & \Delta_i \\ \Delta_j & \Delta_{ij} \end{vmatrix}$  is negative by Sylvester's identity for the minors of the adjugate.

Suppose that  $f = \sum a_{ij} x^i y^j$  is in  $P_2^{pos}(n)$ , and has all positive coefficients. We are interested in the matrices

$$\begin{pmatrix} a_{n,0} & \dots & a_{0,0} \\ \vdots & \ddots & \vdots \\ 0 & \dots & a_{0,n} \end{pmatrix} \text{ and } \begin{pmatrix} a_{0,n} & \dots & 0 \\ \vdots & \ddots & \vdots \\ a_{0,0} & \dots & a_{n,0} \end{pmatrix}$$
(9.17.2)

The determinant is  $\prod a_{i,n-i}$  which is positive since all the coefficients of f are positive. In addition, Theorem 9.113 implies that all 2 by 2 submatrices  $\begin{vmatrix} a_{r+1,s} & a_{r,s} \\ a_{r+1,s+1} & a_{r,s+1} \end{vmatrix}$  are positive. This is not terribly exciting, but there is a more interesting property of matrices to consider:

**Definition 9.117.** A polynomial f(x, y) has *totally positive coefficients* if either matrix in (9.17.2) is totally positive. Note that if one of the matrices is totally positive, so is the other.

Example 9.118. Consider a concrete example: choose

$$f = (1 + x + y) (2 + x + y) (1 + x + 2y).$$

If we expand f we get

$$2 + 5x + 4x^{2} + x^{3} + 7y + 11xy + 4x^{2}y + 7y^{2} + 5xy^{2} + 2y^{3}$$

and the corresponding matrix is

$$\begin{pmatrix} 2 & 7 & 7 & 2 \\ 0 & 5 & 11 & 5 \\ 0 & 0 & 4 & 4 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

The four by four determinant is 40. The matrices of the determinants of all three by three and all two by two submatrices are non-negative.

We are interested in polynomials in  $\mathbf{P}_2^{\text{pos}}$  with totally positive coefficients. There is a special case where we can explicitly compute some of these determinants:  $f = (x + y + 1)^n$ .

**Lemma 9.119.** *For* d = 0, ..., n

$$det \begin{pmatrix} \binom{n}{d,0} & \dots & \binom{n}{0,0} \\ \vdots & & \vdots \\ \binom{n}{d,d} & \dots & \binom{n}{0,d} \end{pmatrix} = \prod_{i=0}^{d} \binom{n}{i}$$

*Proof.* <sup>6</sup> The i, j entry of the matrix is  $\frac{n!}{(d-j)! \, i! \, (n-i+j-d)!}$ . If D is the determinant, then we can factor n! from all the entries, and (d-j)! from all the columns so that

$$\mathsf{D} = \frac{(n!)^{d+1}}{0! \cdots d!} \det\left(\frac{1}{i!(n-i+j-d)!}\right)$$

<sup>&</sup>lt;sup>6</sup>Thanks to Ira Gessel for this argument.

If we multiply each column by 1/(n - i + j - d)! then

$$\mathsf{D} = \frac{(\mathfrak{n}!)^{d+1}}{(0!\cdots d!)((\mathfrak{n}-d)!\cdots(\mathfrak{n})!)} \det\left(\binom{\mathfrak{n}-d+\mathfrak{j}}{\mathfrak{i}}\right)$$

The last determinant has entries that are polynomials in n. If we set  $p_i = \binom{n-d}{i}$  then the matrix is  $(p_i(n+j))$ . Using a generalization of the Vandermonde determinant [108] we can evaluate this determinant, and the result is exactly the desired formula.

**Lemma 9.120.** If f has totally positive coefficients then  $\frac{\partial f}{\partial x}$  has totally positive coefficients. If a is positive then (x + a)f has totally positive coefficients.

*Proof.* The argument is best described by an example, so assume that f has total degree 3. The determinants of the submatrices of the derivative matrix are just constant multiples of the determinants of the original matrix:

$$f = \begin{pmatrix} a_{3,0} & a_{2,0} & a_{1,0} & a_{0,0} \\ 0 & a_{2,1} & a_{1,1} & a_{0,1} \\ 0 & 0 & a_{1,2} & a_{0,2} \\ 0 & 0 & 0 & a_{0,3} \end{pmatrix}$$
$$\frac{\partial f}{\partial x} = \begin{pmatrix} a_{2,1} & a_{1,1} & a_{0,1} \\ 0 & 2a_{1,2} & 2a_{0,2} \\ 0 & 0 & 3a_{0,3} \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{pmatrix} \begin{pmatrix} a_{21} & a_{11} & a_{01} \\ 0 & a_{12} & a_{02} \\ 0 & 0 & a_{03} \end{pmatrix}$$

Next, assume that  $f\in P_2^{pos}$  has totally positive coefficients, and let  $\alpha$  be positive. The coefficient matrices of f and  $(x+\alpha)f$  are

$$\begin{pmatrix} a_{3,0} & a_{2,0} & a_{1,0} & a_{0,0} \\ 0 & a_{2,1} & a_{1,1} & a_{0,1} \\ 0 & 0 & a_{1,2} & a_{0,2} \\ 0 & 0 & 0 & a_{0,3} \end{pmatrix}$$

$$\begin{pmatrix} a_{3,0} & a_{2,0} + \alpha a_{3,0} & a_{1,0} + \alpha a_{2,0} & a_{0,0} + \alpha a_{1,0} & \alpha a_{0,0} \\ 0 & a_{2,1} & a_{1,1} + \alpha a_{2,1} & a_{0,1} + \alpha a_{1,1} & \alpha a_{1,0} \\ 0 & 0 & a_{1,2} & a_{0,2} + \alpha a_{1,2} & \alpha a_{0,2} \\ 0 & 0 & 0 & a_{0,3} & \alpha a_{0,3} \end{pmatrix}$$

We can write the coefficient matrix of (x + a)f as a product

$$\begin{pmatrix} a_{3,0} & a_{2,0} & a_{1,0} & a_{0,0} & 0 \\ 0 & a_{2,1} & a_{1,1} & a_{0,1} & 0 \\ 0 & 0 & a_{1,2} & a_{0,2} & 0 \\ 0 & 0 & 0 & a_{0,3} & 0 \end{pmatrix} \begin{pmatrix} 1 & \alpha & 0 & 0 & 0 \\ 0 & 1 & \alpha & 0 & 0 \\ 0 & 0 & 1 & \alpha & 0 \\ 0 & 0 & 0 & 1 & \alpha \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

1.

The first matrix of the product is totally positive by assumption, and the second is easily seen to be totally positive. Since the product of totally positive matrices is totally positive [100] it follows that the coefficient matrix of  $(x+\alpha)f$  is totally positive.

**Corollary 9.121.** *If*  $f, g \in P^{p \circ s}$  *have positive leading coefficients then* f(x)g(y) *has totally positive coefficients.* 

*Proof.* First of all, g(y) has totally positive coefficients since the matrix of  $g(y) = \sum_{0}^{n} b_i x^i$  is

$b_n$		$\mathfrak{b}_1$	$b_0$
0	•••	0	0
÷		÷	:
0		0	0/

If we write  $f(x) = \alpha(x + a_1) \dots (x + a_n)$  then we can inductively apply the lemma.

We conjecture (Question 102) that all polynomials that are product of linear terms in  $\mathbf{P}_2^{\text{pos}}$  have totally positive coefficients. It is not the case that all polynomials in  $\mathbf{P}_2^{\text{pos}}$  are totally positive. Consider the example where

	/1	0	0		/13	9	7		/5	7	8 \
M =	0	1	0	+x	9	7	5	+ y	7	11	12
	0	0	1/		7	5	4)		8/	12	14/

The determinant of M is in  $P_2^{\text{pos}}$  and equals

$$1 + 24x + 16x^{2} + 2x^{3} + 30y + 164xy + 62x^{2}y + 22y^{2} + 64xy^{2} + 4y^{3}$$

with coefficient matrix

$$\begin{pmatrix} 4 & 0 & 0 & 0 \\ 22 & 64 & 0 & 0 \\ 30 & 164 & 62 & 0 \\ 1 & 24 & 16 & 2 \end{pmatrix}$$

The determinant of the three by three matrix in the lower left corner is -1760. Of course, all the two by two submatrices have positive determinant.

We can't show that  $f \in \mathbf{P}_2^{\text{pos}}$  has totally positive coefficients, but we can show that the matrix of coefficients is TP<sub>2</sub>. This is a simple consequence of the rhombus inequalities, and reflects the geometric fact that the the graph of the coefficients is log-concave.

**Lemma 9.122.** If  $f \in \mathbf{P}_2^{pos}$  then the coefficient array (9.17.2) is  $TP_2$ .

*Proof.* We describe a special case; the general case is no different. Suppose  $f = \sum a_{i,j} x^i y^j$ , and we wish to show that  $\begin{vmatrix} a_{0,2} & a_{2,2} \\ a_{0,0} & a_{2,0} \end{vmatrix}$  is non-negative. Consider a portion of the coefficient array



From the rhombus inequalities we have

 $\begin{array}{l} a_{0,1} \; a_{1,0} \geqslant a_{0,0} \; a_{1,1} \\ a_{1,1} \; a_{2,0} \geqslant a_{1,0} \; a_{2,1} \\ a_{0,2} \; a_{1,1} \geqslant a_{0,1} \; a_{1,2} \\ a_{1,2} \; a_{2,1} \geqslant a_{1,1} \; a_{2,2} \end{array}$ 

Multiplying these inequalities and canceling yields the desired result:

 $a_{0,2} a_{2,0} \ge a_{0,0} a_{2,2}$ 

In general, the various vertices contribute as follows

- **corner** The corner vertices each occur once, on the correct sides of the inequality.
- **edge** The vertices on the interior of the boundary edges each occur twice, once on each side of the inequality.

interior The interior vertices occur twice on each side of the inequality.

Again, multiplying and canceling finishes the proof.

# 9.18 Generic polynomials and P<sub>2</sub><sup>sep</sup>

In one variable the interior of the space of polynomials consists of polynomials whose roots are all distinct. The analog in  $P_2$  is polynomials whose substitutions have all distinct roots. We also have the analog of  $P^{sep}$ , the set of all polynomials whose roots are at least one apart from one another.

**Definition 9.123.**  $P_2^{gen}$  consists of all polynomials f(x, y) in  $P_2$  such that

- 1.  $f(x, \alpha)$  has all distinct roots, for all  $\alpha \in \mathbb{R}$ .
- 2.  $f(\alpha, x)$  has all distinct roots, for all  $\alpha \in \mathbb{R}$ .

If we define  $\mathbb{A}_{x}f(x,y) = f(x+1,y)$  and  $\mathbb{A}_{y}f(x,y) = f(x,y+1)$  then  $\mathbf{P}_{2}^{sep}$  consists of all those polynomials in P<sub>2</sub> satisfying

•  $f \underline{\ll} \mathbb{A}_x f$ 

● f<u>≪</u>Ayf

For any fixed  $\alpha$  we see that= both  $f(x, \alpha)$  and  $f(\alpha, x)$  are in  $\mathbf{P}^{sep}$ , so  $f \in \mathbf{P}_2^{gen}$ .

The graph of a polynomial in  $\mathbf{P}_2^{\text{gen}}(n)$  is easy to describe: it's a set of n non-intersecting curves. If f is in  $\mathbf{P}_2^{\text{sep}}$  then all the curves are at least one apart on every vertical line, and they are also at least one apart on every horizontal line. In addition,

**Lemma 9.124.** If  $f \in \mathbf{P}_2^{gen}$  then the solution curves are always decreasing.

*Proof.* If the graph of f(x, y) = 0 had either horizontal or vertical points, then by Lemma 9.30 the graph would have double points. However, the solution curves are disjoint since  $f \in \mathbf{P}_2^{\text{gen}}$ . We conclude the the slope is always the same sign. Since the solution curves are asymptotic to lines with negative slope, the solution curves are everywhere decreasing.

**Example 9.125.** We can use polynomials in  $\mathbf{P}^{sep}$  to construct polynomials in  $\mathbf{P}_2^{sep}$ . If  $f(x) \in \mathbf{P}^{sep}$ , then f(x + y) is easily seen to be in  $\mathbf{P}_2^{sep}$ . The graph of f(x + y) is just a collection of parallel lines of slope 1 that are at least one apart in the x and y directions.

We can easily determine the minimum separation of the roots when the polynomial is a quadratic given by a determinant. We start with one variable. Define

$$f(\mathbf{x}) = \begin{vmatrix} a_1 \mathbf{x} + \alpha & \gamma \\ \gamma & a_2 \mathbf{x} + \beta \end{vmatrix}$$
(9.18.1)

The square of the distance between the roots of a polynomial  $ax^2 + bx + c$  is  $(b^2 - 4ac)/a^2$ . The squared difference between the roots of f is

$$\frac{1}{a_1^2 a_2^2}((\alpha + \gamma)^2 - 4a_1 a_2(\alpha \beta - \gamma^2)) = \frac{1}{a_1^2 a_2^2}(4a_1 a_2 \gamma^2 + (\alpha a_2 - \beta a_1)^2)$$

and hence the minimum squared distance is  $4\gamma^2/(a_1a_2)$ . To find the minimum of

$$f(x,y) = \begin{vmatrix} a_1x + b_1y + \alpha & \gamma \\ \gamma & a_2x + b_2y + \beta \end{vmatrix}$$
(9.18.2)

we note that if y is fixed then by the above the minimum distance is  $2|\gamma|/\sqrt{a_1a_2}$ . If x is fixed then the minimum distance is  $2|\gamma|/\sqrt{b_1b_2}$ . It follows that

**Lemma 9.126.** The polynomial f(x, y) of (9.18.2) is in  $\mathbf{P}_2^{sep}$  iff

$$|\gamma| \ge \frac{1}{2} \max\left\{\sqrt{a_1 a_2}, \sqrt{b_1 b_2}\right\}$$

*Remark* 9.127. The condition that the solution curves of a polynomial in  $\mathbf{P}_2^{\text{sep}}$  are at least one apart in the horizontal and vertical directions does not imply that they are one apart in all directions. From the above we know that  $f = (x + y + 1/2)(x + y - 1/2) \in \mathbf{P}_2^{\text{sep}}$ . The line x = y meets the graph, which is two parallel lines, in the points  $\pm (1/4, 1/4)$ . Thus, there are two points on different solution curves that are only  $\sqrt{2}/2 = .707$  apart.

**Example 9.128.** The polynomial xy - 1 is a limit of polynomials in  $\mathbf{P}_2^{\text{sep}}$ . We know that the limit

$$\lim_{\epsilon \to 0^+} \left| \begin{array}{c} x + \epsilon y & 1 \\ 1 & \epsilon x + y \end{array} \right| = xy - 1$$

expresses xy - 1 as a limit of polynomials in P<sub>2</sub>. In order to apply the lemma we need to multiply each side by  $\begin{vmatrix} 1 & 0 \\ 0 & e^{-1/2} \end{vmatrix}$ , which yields

$$\lim_{\varepsilon \to 0^+} \epsilon \left| \begin{array}{c} x + \epsilon y & \epsilon^{-1} \\ \epsilon^{-1} & x + \epsilon^{-1} y \end{array} \right| = xy - 1$$

Since  $2|c_{12}| = 2|c_{12}|/\sqrt{d_1d_2} = 2\varepsilon^{-1/2}$  we see that the roots are at least one apart when  $0 < \varepsilon < 1/4$ .

Polynomials in  $\mathbf{P}_2^{\text{sep}}$  satisfy the expected properties.

#### Lemma 9.129.

- 1.  $\mathbf{P}_{2}^{gen}$  and  $\mathbf{P}_{2}^{sep}$  are closed under differentiation.
- 2. The coefficients of any power of y (or of x) of a polynomial in  $\mathbf{P}_2^{\text{sep}}$  are in  $\mathbf{P}^{\text{sep}}$ .

*Proof.* The statement follows since differentiation preserves the properties of all distinct roots, and of interlacing. If we differentiate sufficiently many times, and substitute zero we can conclude that all coefficients are in  $\mathbf{P}^{\text{sep}}$ .

We can use properties of symmetric matrices to construct polynomials in  $\mathbf{P}_2^{\text{gen}}$ . We first show [134, page 124]

**Lemma 9.130.** *If* M *is a symmetric matrix whose superdiagonal entries are all non-zero, then the eigenvalues of* M *are distinct.* 

*Proof.* If we delete the first column and the last row of  $\alpha I + M$ , then the resulting matrix has all non-zero diagonal entries, since these entries come from the superdiagonal of M. Since the product of the diagonal entries is the product of the eigenvalues, the matrix is non-singular. Thus, if M is n by n, then  $\alpha I + M$  contains a submatrix of rank n - 1 for any choice of  $\alpha$ . If M had a repeated eigenvalue, then there would be an  $\alpha$  for which  $\alpha I + M$  had rank at most n - 2. Since  $\alpha I + M$  has rank at least n - 1, this contradiction establishes the lemma.

**Corollary 9.131.** *Suppose* C *is a symmetric matrix whose superdiagonal is all non-zero, and* D *is a diagonal matrix of positive diagonal entries. Then,* 

 $|\mathbf{x}\mathbf{I} + \mathbf{y}\mathbf{D} + \mathbf{C}| \in \mathbf{P}_2^{gen}$ 

*Moreover,*  $|xI + yD + \alpha C| \in \mathbf{P}_2^{sep}$  *for sufficiently large*  $\alpha$ *.* 

*Proof.* Let f(x, y) = |xI + yD + C|. The roots of  $f(x, \alpha)$  are the eigenvalues of  $\alpha D + C$ , and they are all distinct by the lemma. If  $f(\alpha, y) = 0$  then

$$\left|\alpha D^{-1} + yI + \sqrt{D^{-1}}C\sqrt{D^{-1}}\right| = 0$$

Thus, the roots of f(a, y) are the eigenvalues of  $\alpha D^{-1} + \sqrt{D^{-1}}C\sqrt{D^{-1}}$ . Since  $\sqrt{D^{-1}}C\sqrt{D^{-1}}$  satisfies the conditions of the lemma, the roots of  $f(\alpha, y)$  are all distinct. Thus,  $f(x, y) \in \mathbf{P}_2^{gen}$ .

The roots of  $|xI + yD + \alpha C|$  are the roots of  $f(\frac{x}{\alpha}, \frac{y}{\alpha})$ , and so the second part follows.

A similar argument shows the following.

**Corollary 9.132.** If  $D_1$  is a diagonal matrix with distinct positive entries, and  $D_2$  is a positive definite matrix whose superdiagonal is all non-zero, then

$$|I+xD_1+yD_2|\in \textbf{P}_2^{\text{gen}}\cap \textbf{P}_2^{\text{pos}}$$

**Corollary 9.133.** Every polynomial in  $P_2$  is the limit of polynomials in  $\mathbf{P}_2^{gen}$ .

*Proof.* By Corollary 9.77 if  $f(x, y) \in P_2$  then there is a diagonal matrix D and a symmetric matrix C such that f(x, y) = |xI + yD + C|. Let  $U_{\varepsilon}$  be the matrix that is all zero except for  $\varepsilon$ 's in the sub and super diagonal. The polynomial  $|xI + yD + C + U_{\varepsilon}|$  is in  $\mathbf{P}_2^{\text{gen}}$  for  $|\varepsilon|$  sufficiently small, and it converges to f(x, y) as  $\varepsilon \to 0$ .

**Example 9.134.** Here is an example of a cubic polynomial in  $P_2^{sep}$ . Let

$$f(x,y) = \begin{vmatrix} x+y & 1 & 0 \\ 1 & x+2y & 1 \\ 0 & 1 & x+3y \end{vmatrix} = -2x + x^3 - 4y + 6x^2y + 11xy^2 + 6y^3$$

The roots of  $f(x, \alpha)$  are

$$-2 \alpha, -2 \alpha - \sqrt{2 + \alpha^2}, -2 \alpha + \sqrt{2 + \alpha^2}$$

and the minimum distance between these roots is clearly  $\sqrt{2}$ . If  $f(\alpha, y) = 0$  then the roots are

$$\frac{-\alpha}{2}, \frac{-2\alpha - \sqrt{6+\alpha^2}}{3}, \frac{-2\alpha + \sqrt{6+\alpha^2}}{3}$$

and the minimum is seen to be  $1/\sqrt{2}$ . Consequently, replacing y by  $y/\sqrt{2}$ , x by  $\sqrt{2}x$ , and multiplying by  $\sqrt{2}$  yields

$$-4x + 4x^3 - 4y + 12x^2y + 11xy^2 + 3y^3 \in \mathbf{P}_2^{sep}$$

and is a polynomial satisfying  $\delta_x = \delta_y = 1$ .

The hyperbola xy - 1 is a good example of a polynomial whose graph has no intersections. This can be generalized.

**Lemma 9.135.** If D is a positive definite matrix with all distinct roots, and  $g \in P_2$  then  $|D - gI| \in \mathbf{P}_2^{gen}$ .

*Proof.* If D is an n by n matrix then the homogeneous part of |D - gI| is  $g^n$ . It suffices to show for each  $\alpha$  that  $|D - g(x, \alpha)I|$  has all distinct roots. There are a, b, c such that  $g(x, \alpha) = a(x + b)(x + c)$ . After translating and scaling we may assume that  $g(x, \alpha) = x^2 - 1$ . Now

$$|D - g(x, \alpha)I| = |(D + I) - x^2I|$$

Since D has all distinct eigenvalues, |D + I - xI| has all distinct positive roots. Taking square roots shows that  $|D + I - x^2I|$  has all distinct roots.

### 9.19 Integrating generic polynomials

We can integrate generic polynomials. To do this we first need to introduce some quantitative measures of root separation. We extend the one variable measure  $\delta(f)$  of separation to polynomials in P<sub>2</sub>:

$$\begin{split} \delta_{x}(f) &= \inf_{\alpha \in \mathbb{R}} \ \delta(f(x,\alpha)) \\ \delta_{y}(f) &= \inf_{\alpha \in \mathbb{R}} \ \delta(f(\alpha,y)) \end{split}$$

If the solution curves of f cross, then both  $\delta_y(f)$  and  $\delta_x(f)$  are zero. Thus,  $f \in \mathbf{P}_2^{\text{gen}}$  if either  $\delta_x(f) > 0$  or  $\delta_y(f) > 0$ . For an example, the polynomial

$$f(x,y) = (x + ay)(x + ay + b)$$

has  $\delta_x(f) = b$  and  $\delta_y(f) = b/a$ . This shows that  $(\delta_x(f), \delta_y(f))$  can take any pair of positive values.

Applying (3.6.1) to  $f \in P_2$  yields

$$0 \leqslant s \leqslant t \leqslant \delta_{y}(f) \implies (\forall \alpha \in \mathbb{R}) f(\alpha, y + s) \underline{\ll} f(\alpha, y + t)$$

This shows that  $f\in \textbf{P}_2^{sep}$  if and only if  $\delta_x(f)\geqslant 1$  and  $\delta_y(f)\geqslant 1.$ 

**Proposition 9.136.** Suppose that  $f \in \mathbf{P}_2^{gen}$ ,  $0 < \alpha \leq \delta_y(f)$ , and w(x) is a non-negative integrable function on  $\mathbb{R}$ . If we define

$$g(x,y) = \int_0^{\infty} f(x,y+t)w(t) dt$$

then  $g(x,y) \in \mathbf{P}_2^{gen}$ , and  $\delta_y(g) \ge \frac{1}{2}\alpha$ .

*Proof.* Since  $g^H = (\int_0^\alpha w(t) dt) f^H$ , we see that g satisfies the homogeneity condition. Proposition 3.37 implies that g satisfies substitution, and Proposition 3.42 shows that the separation number is as claimed. Since  $\delta_y(g) > 0$ , we know that  $g \in \mathbf{P}_2^{\text{gen}}$ .

*Remark* 9.137. If  $f \in \mathbf{P}_2^{sep}$  then  $\int_0^1 f(x, y + t) dt$  is in  $\mathbf{P}_2^{gen}$ , but even the simplest example fails to be in  $\mathbf{P}_2^{sep}$ . Consider

$$\begin{split} f(x,y) &= (x+y)(x+y+1) \\ \int_0^1 f(x,y+t) \, dt &= g(x,y) = 5/6 + (x+y)(x+y+2) \\ \delta_x(g) &= \delta_y(g) = \sqrt{2/3} < 1 \end{split}$$

Here's a curious corollary.

**Corollary 9.138.** Suppose that  $f \in \mathbf{P}_2^{gen}$ , and  $\delta_y(f) \ge 1$ . If  $\frac{\partial}{\partial y}h = f$ , then  $\Delta_y(h) \in \mathbf{P}_2^{gen}$ .

*Proof.* Take w = 1, and  $\alpha = 1$ . The corollary follows from Proposition 9.136 since

$$\int_0^1 f(x,y+t)dt = h(x,y+1) - h(x,y) = \Delta_y(h)$$

If  $f \in \mathbf{P}$ , then it follows from the definitions that

$$\delta(f) = \delta_x f(x+y) = \delta_y f(x+y)$$

Consequently,

**Corollary 9.139.** *If*  $f \in P$ , w(t) *positive, and*  $0 < \alpha \leq \delta(f)$  *then* 

$$\int_0^\alpha f(x+y+t)w(t)\,dt\in \mathbf{P}_2^{gen}$$

## 9.20 Recursive sequences of polynomials

In this section we show how to construct polynomials in  $P_2$  that are analogs of orthogonal polynomials. The next lemma (a special case of Lemma 10.112) is the key idea, for it allows us to construct a new polynomial in  $P_2$  from a pair of interlacing ones.

**Lemma 9.140.** If  $f \leq g$  are both in  $P_2$ , and a, b, d are positive then

$$(ax + by + c)f - dg < f$$

Two constructions are immediate consequences of the lemma. The first one shows that certain two-dimensional analogs of orthogonal polynomials are in  $P_2$ .

**Lemma 9.141.** Let  $a_n, b_n, d_n$  be positive, and define a sequence of polynomials by

$$P_{-1} = 0$$
  

$$P_{0} = 1$$
  

$$P_{n+1} = (a_{n}x + b_{n}y + c_{n})P_{n} - d_{n}P_{n-1}$$

Then all  $P_n$  are in  $P_2$  and

$$\mathsf{P}_0 \underline{\geq} \mathsf{P}_1 \underline{\geq} \mathsf{P}_2 \underline{\geq} \cdots$$

Note that  $P_n(x, \alpha)$  and  $P_n(\alpha, y)$  are orthogonal polynomials for any choice of  $\alpha$ . These polynomials also have a nice representation as determinants that shows that they are in **P**<sub>2</sub>. If we let  $\ell_i = a_i x + b_i y + c_i$  then we can construct a matrix

$\int \ell_1$	$\sqrt{d_2}$	0	0		0	0 \
$\sqrt{d_2}$	$\ell_2$	$\sqrt{d_3}$	0		0	0
0	$\sqrt{d_3}$	$\ell_3$	$\sqrt{d_4}$		0	0
0	0	$\sqrt{d_4}$	$\ell_4$		0	0
:	:	:	:	۰.	:	:
	•	•	•	•	·	· 1
$\setminus 0$	0	0	0		$\sqrt{d_n}$	$\ell_n$

whose determinant is  $P_n$ . If we set  $\ell_i = c_i$  then the matrix is called a *Jacobi matrix*, and its characteristic polynomial is an orthogonal polynomial in one variable.

**Lemma 9.142.** Let  $a_n, b_n, d_n, e_n$  be positive, and define a sequence of polynomials by

$$P_{-1} = 0$$

$$P_{0} = 1$$

$$P_{n+1} = (a_{n}x + b_{n}y + c_{n})P_{n} - \left(d_{n}\frac{\partial}{\partial y} + e_{n}\frac{\partial}{\partial x}\right)P_{n}$$

Then all  $P_n$  are in  $P_2$  and  $P_0 \ge P_1 \ge P_2 \ge \cdots$ 

### 9.21 Bezout matrices

Any two polynomials f, g of the same degree determine a matrix called the Bezout matrix. We give a simple argument that the Bezout matrix is positive definite if and only if f and g interlace. We also investigate properties of the Bezout matrix for polynomials in  $P_2$ .

**Definition 9.143.** If f(x), g(x) are polynomials of the same degree n the Bezout matrix of f and g is the symmetric matrix  $B(f, g) = (b_{ij})$  such that

$$\frac{f(x)g(y) - f(y)g(x)}{x - y} = \sum_{i,j=1}^{n} b_{ij} x^{i-1} y^{j-1}$$

In order to present the key result [152] we introduce the Vandermonde vector.

$$V_n(x) = (1, x, \dots, x^{n-1})^t$$

**Theorem 9.144** (Lander). If f, g are polynomials of degree n with no common zeros, and the zeros of f are  $r_1, \ldots, r_n$  then  $B(f,g) = (V^t)^{-1}DV^{-1}$  where

$$V = (V_n(r_1), \dots, V_n(r_n))$$
$$D = diag(g(r_1)f'(r_1), \dots, g(r_n)f'(r_n))$$

**Corollary 9.145.** *If*  $f \in P$  *and* f, g *have no common roots then* f *and* g *interlace if and only if the Bezout matrix is positive definite.* 

*Proof.* If the Bezout matrix is positive definite then g and f' have the same sign at the roots of f. Since f' sign interlaces f, so does g. The converse is similar.  $\Box$ 

This is a classical result. More interesting and recent is that using the Bezout matrix we can parametrize the solution curves of certain polynomials in  $P_2$ . We paraphrase Theorem 3.1 [152]

**Theorem 9.146.** Three polynomials  $f_0$ ,  $f_1$ ,  $f_2$  in one variable determine a map from the complex line  $\mathbb{C}$  to the complex plane  $\mathbb{C}^2$ :

$$z \mapsto \left(\frac{f_0(z)}{f_1(z)}, \frac{f_2(z)}{f_1(z)}\right) \tag{9.21.1}$$

The image is a rational curve defined by the polynomial

$$\Delta(\mathbf{x}, \mathbf{y}) = \det(B(f_0, f_2) + \mathbf{x}B(f_0, f_1) + \mathbf{y}B(f_2, f_1))$$
(9.21.2)

Choose mutually interlacing polynomials  $f_0$ ,  $f_1$ ,  $f_2$  in **P**(n). Each of the Bezout matrices B( $f_0$ ,  $f_1$ ), B( $f_0$ ,  $f_2$ ), B( $f_1$ ,  $f_2$ ) is positive definite since each pair is interlacing. It follows that  $\Delta(x, y)$  in (9.21.2) belongs to **P**<sub>2</sub><sup>pos</sup>. The solution curves are parametrized by (9.21.1).

Consider an example:

$$\begin{aligned} f_1 &= (2+x)(8+x)(12+x) \\ f_2 &= (3+x)(9+x)(13+x) \\ f_3 &= (7+x)(11+x)(15+x) \end{aligned} \quad \mathsf{B}(f_0,f_1) = \begin{pmatrix} 12600 & 2922 & 159 \\ 2922 & 785 & 47 \\ 159 & 47 & 3 \end{pmatrix} \\ \mathsf{B}(f_0,f_2) &= \begin{pmatrix} 90456 & 19074 & 963 \\ 19074 & 4109 & 211 \\ 963 & 211 & 11 \end{pmatrix} \qquad \mathsf{B}(f_1,f_2) = \begin{pmatrix} 89568 & 17292 & 804 \\ 17292 & 3440 & 164 \\ 804 & 164 & 8 \end{pmatrix} \end{aligned}$$

$$\begin{split} \Delta(x,y) &= (17325x^3 + 175200yx^2 + 155803x^2 + 242112y^2x + 465664yx + 191775x \\ &\quad + 73728y^3 + 240384y^2 + 223136y + 61425)/17325 \end{split}$$



Figure 9.9: The graph of a parametrized  $P_2$ 

All solutions to  $\Delta(x, y) = 0$  are of the form

x =	(t+2)(t+8)(t+12)	$u = \frac{(t+7)(t+11)(t+1)}{(t+1)(t+1)}$	5)
	(t+3)(t+9)(t+13)	$y = \frac{1}{(t+3)(t+9)(t+13)}$	3)

The graph of  $\Delta(x, y)$  is in Figure 9.9. The three curves are parametrized as follows:

$t\not\in(-13,-3)$
$t\in(-13,-9)$
$t\in(-9,-3)$

Now suppose that  $f \ll g \in P_2$ . The Bezout matrix with respect to x is

$$B_{x}(f,g) = \frac{f(x,y)g(z,y) - f(z,y)g(x,y)}{x-z}$$

Since  $f(x, \alpha)$  and  $g(x, \alpha)$  interlace for all  $\alpha$  it follows that  $B_x(f, g)$  is a matrix polynomial in one variable that is positive definite for all y. In particular,  $v^t B_x(f, g) v > 0$  for all non-zero vectors v. However,  $v^t B_x(x, y)v$  is generally not a stable polynomial.

The polynomial that determines the Bezout matrix has all coefficients in P.

**Lemma 9.147.** If  $f \leq g$  in **P** and

$$B(x,y) = \frac{1}{x-y} \begin{vmatrix} f(x) & f(y) \\ g(x) & g(y) \end{vmatrix}$$

then all coefficients of powers of x or of y are in **P**. If  $f, g \in P^{pos}$  then all coefficients of B are positive, and all coefficients are in  $P^{pos}$ .

*Proof.* Write  $g(x) = \sum a_i \frac{f}{x - r_i}$  where  $a_i \geqslant 0$ . Then

$$B(x,y) = \frac{1}{x-y} \left( f(x) \sum_{i=1}^{\infty} a_{i} \frac{f(y)}{y-r_{i}} - f(y) \sum_{i=1}^{\infty} a_{i} \frac{f(x)}{x-r_{i}} \right)$$
(9.21.3)

$$= \sum a_{i} \frac{r(x)}{x - r_{i}} \frac{r(y)}{y - r_{i}}$$
(9.21.4)

Since  $\{f(x)/(x - r_i)\}$  is mutually interlacing the result follows from Lemma 9.18. If  $f, g \in \mathbf{P}^{pos}$  then  $f(x)/(x - r_i)$  has all positive coefficients, so B has all positive coefficients.

*Remark* 9.148. It is not the case that  $B(x, y) \in \overline{P}_2$ . If it were then B(x, x) is in **P**, but  $B(x, x) = \begin{vmatrix} f(x) & f'(x) \\ g(x) & g'(x) \end{vmatrix}$  which is stable by Lemma 21.77.

CHAPTER



# **Polynomials in several variables**

#### **10.1** The substitution property

Just as for one variable there is a substitution property that all our polynomials will satisfy.

**Definition 10.1.** If f is a polynomial in variables  $x_1, \ldots, x_d$  then f satisfies  $x_i$ -substitution if for every choice of  $a_1, \ldots, a_d$  in  $\mathbb{R}$  the polynomial

$$f(a_1,\ldots,a_{i-1},x_i,a_{i+1},\ldots,a_d)$$

has all real roots. We say that f satisfies *substitution* if it satisfies  $x_i$ -substitution for  $1 \le i \le d$ .

It is clumsy to write out the variables so explicitly, so we introduce some notation. Let  $\mathbf{x} = (x_1, ..., x_d)$ ,  $\mathbf{a} = (a_1, ..., a_d)$ , and define

$$\mathbf{x}_{i}^{\mathbf{a}} = (a_{1}, \ldots, a_{i-1}, x_{i}, a_{i+1}, \ldots, a_{d})$$

The *degree* of a monomial  $x_1^{i_1} \cdots x_d^{i_d}$  is the sum  $i_1 + \cdots + i_d$  of the degrees. The *total degree* of a polynomial is the largest degree of a monomial. The  $x_i$ -*degree* of a polynomial is the largest exponent of  $x_i$ . An *index set* **I** is a sequence  $(i_1, i_2, \dots)$  of non-negative integers. A *monomial in* **x** can be written

$$\mathbf{x}^{\mathbf{I}} = \mathbf{x}_1^{\mathbf{i}_1} \mathbf{x}_2^{\mathbf{i}_2} \cdots$$

The size of an index set is  $|\mathbf{I}| = i_1 + i_2 + \cdots$ . The degree of  $\mathbf{x}^{\mathbf{I}}$  is  $|\mathbf{I}|$ .

We can write a polynomial f in terms of coefficients of monomials. If  $\mathbf{x} = (x_1, \dots, x_d)$  and  $\mathbf{y} = (y_1, \dots, y_e)$  then we write

$$f(\mathbf{x}, \mathbf{y}) = \sum_{\mathbf{I}} f_{\mathbf{I}}(\mathbf{x}) \mathbf{y}^{\mathbf{I}}.$$
 (10.1.1)

We can restate the definition of  $x_i$ -substitution:
•  $f(\mathbf{x})$  satisfies  $x_i$ -substitution iff  $f(\mathbf{x}_i^a) \in \mathbf{P}$  for all  $\mathbf{a} \in \mathbb{R}^d$ .

All the polynomials we consider will belong to this set:

 $Sub_d = \{all polynomials in d variables that satisfy substitution\}$ 

In the case d = 1 the set **Sub**<sub>1</sub> is just **P**. Consider these examples in several variables.

Example 10.2. All products of linear factors such as

$$(x_1 + 2x_2 - x_3 + 1)(x_1 - x_2)(x_2 + x_3)$$

satisfy substitution. More generally, if the factors of a product are linear in each variable separately then the product satisfies substitution. An example of such a polynomial is

$$(x_1x_2 + x_1 + 3)(x_1x_2x_3 + 2x_1x_4 + 1)$$

**Lemma 10.3.** If f is a polynomial, and there are  $f_j \in \mathbf{Sub}_d$  such that  $\lim f_j = f$  then f satisfies substitution.

*Proof.* Observe that for any  $\mathbf{a} \in \mathbb{R}^d$  and  $1 \leq i \leq d$  we have

$$f(\mathbf{x}_{i}^{\mathbf{a}}) = \lim_{i \to \infty} f_{j}(\mathbf{x}_{i}^{\mathbf{a}})$$

The result now follows from the corresponding result (Lemma 1.40) for one variable.  $\hfill \Box$ 

Substitution satisfies a factorization property.

**Lemma 10.4.** *If*  $fg \in \mathbf{Sub}_d$  *then*  $f, g \in \mathbf{Sub}_d$ .

*Proof.* The result is obvious for polynomials in one variable. The result follows from the fact that  $(fg)(\mathbf{x}_i^a) = f(\mathbf{x}_i^a) g(\mathbf{x}_i^a)$ .

The reversal transformation is one of the few operations that preserve the property of substitution.

**Proposition 10.5.** If f has  $x_i$ -degree  $k_i$  and  $f = \sum a_i x^I$  then the reverse of f is defined to be

$$\operatorname{rev}(f) = \sum a_{I} \mathbf{x}^{K-I}$$

where  $\mathbf{K} = (k_1, ...)$ . If  $f \in \mathbf{Sub}_d$  then  $rev(f) \in \mathbf{Sub}_d$ .

*Proof.* Let  $\mathbf{a} = (a_1, \dots, a_d)$ . If we substitute we see that

$$\begin{split} \mathsf{rev}(f)(\mathbf{x}_{i}^{\mathbf{a}}) &= \sum_{\mathbf{a}_{\mathbf{I}}} a_{\mathbf{I}} a_{1}^{k_{1}-i_{1}} \cdots x_{i}^{k_{i}-i_{i}} \cdots a_{d}^{k_{d}-i_{d}} \\ &= a_{1}^{k_{1}} \cdots a_{d}^{k_{d}} \sum_{\mathbf{a}_{\mathbf{I}}} a_{\mathbf{I}} a_{1}^{-i_{1}} \cdots x_{i}^{k_{i}-i_{i}} \cdots a_{d}^{-i_{d}} \\ &= a_{1}^{k_{1}} \dots a_{d}^{k_{d}} \mathsf{rev}(f(\mathbf{x}_{i}^{\mathbf{b}})) \end{split}$$

where  $\mathbf{b} = (1/a_1, \dots, 1/a_d)$ . Since reversal preserves roots in one variable, rev(f) satisfies substitution.

Interlacing can be easily defined for polynomials satisfying substitution.

**Definition 10.6.** If f and g are polynomials that satisfy substitution, then we say that f and g *interlace* if for every real  $\alpha$  the polynomial f +  $\alpha$ g satisfies substitution.

The relation of interlacing is preserved under limits.

**Lemma 10.7.** Suppose  $f_i, g_i$  are in  $\mathbf{Sub}_d$ , and  $\lim_{i\to\infty} f_i = f$ ,  $\lim_{i\to\infty} g_i = g$  where f, g are polynomials. If  $f_i$  and  $g_i$  interlace for all i then f and g interlace.

*Proof.* Since  $f_i + \alpha g_i$  is in  $\mathbf{Sub}_d$  for all i, and  $f_i + \alpha g_i$  converges to  $f + \alpha g$  it follows that  $f + \alpha g \in \mathbf{Sub}_d$ . Consequently f and g interlace.

#### **10.2** The class $P_d$ .

Before we can define  $P_d$  we need a few definitions that generalize the definitions for d = 2.

**Definition 10.8.** If f has degree n, then the *homogeneous part* of  $f = \sum a_I x^I$  is

$$f^{H} = \sum_{|I|=n} a_{I} \mathbf{x}^{I}$$
(10.2.1)

The homogeneous part controls the asymptotic behavior of the graph, just as the leading coefficient does in one variable.

**Definition 10.9.** A polynomial  $f(x_1, ..., x_d)$  of degree n satisfies the *positivity condition* if all coefficients of monomials of degree at most n are positive.

We now define

#### Definition 10.10.

 $P_d =$ all f in d variables such that

- f satisfies substitution
- f<sup>H</sup> satisfies the positivity condition (Definition 10.9).

 ${\tt P}_d$  is the analog of "polynomials with positive leading sign and all real roots".

We let  $P_d(n)$  denote the subset of  $P_d$  that consists of all polynomials of degree n.

The results in Lemma 9.16 also hold for polynomials in  $P_d$ . The homintogeneous part is always in  $U_d$ .

**Lemma 10.11.** If  $f \in P_d$  then  $f^H \in \mathbf{P}_d$ . If  $f \in \mathbf{Sub}_d$  then  $f^H \in \mathbf{Sub}_d$ .

*Proof.* Since  $f^H = (f^H)^H$  the positivity condition is satisfied. It remains to show that if  $f \in \mathbf{Sub}_d$  then  $f^H$  satisfies substitution. Assume that the degree of f is n, and write

$$\mathsf{f} = \sum_{|\mathbf{I}| = n} c_{\mathbf{I}} x^{\mathbf{I}} + \sum_{|\mathbf{I}| < n} c_{\mathbf{I}} x^{\mathbf{I}}.$$

Replace **x** by  $\mathbf{x}/\epsilon$  and multiply by  $\epsilon^{n}$ :

$$\varepsilon^{n}f(\mathbf{x}/\varepsilon) = \sum_{|\mathbf{I}|=n} c_{\mathbf{I}} \mathbf{x}^{\mathbf{I}} + \sum_{|\mathbf{I}| < n} \varepsilon^{n-|\mathbf{I}|} c_{\mathbf{I}} \mathbf{x}^{\mathbf{I}}.$$
 (10.2.2)

It follows that  $\epsilon^n f(\mathbf{x}/\epsilon)$  converges to  $f^H$  as  $\epsilon$  approaches 0. Lemma 10.3 shows that  $f^H$  satisfies substitution.

As is to be expected, substituting for several variables is a good linear transformation.

**Lemma 10.12.** Let  $\mathbf{x} = (x_1, \dots, x_d)$ ,  $\mathbf{y} = (y_1, \dots, y_e)$  and choose  $\mathbf{a} \in \mathbb{R}^e$ . The substitution map  $\mathbf{x}^I \mathbf{y}^J \mapsto \mathbf{x}^I \mathbf{a}_J$  is a linear transformation  $P_{d+e} \longrightarrow U_d$ .

*Proof.* Using induction it suffices to prove this in the case that e is 1. Let  $\mathbf{x}^{a} = (x_{1}, \ldots, x_{d}, a)$ . Choose  $f \in P_{d}$ , and let  $g = f(\mathbf{x}^{a})$ . Since f satisfies substitution so does g. The homogeneous part  $f(\mathbf{x}^{a})^{H}$  is found by deleting all the terms involving  $x_{d+1}$  in  $f^{H}$ . Since all coefficients of  $f^{H}$  are positive, so are those of  $f(\mathbf{x}^{a})^{H}$ . it follows that  $g \in P_{d}$ .

Unlike  $Sub_d$ , not all products of linear terms are in  $P_d$ . The lemma below shows that the signs must be coherent for this to occur. Note that there are no constraints on the signs of the constant terms.

Lemma 10.13. Suppose

$$f(\mathbf{x}) = \prod_{k=1}^{n} (b_k + a_{1k}x_1 + \dots + a_{dk}x_d).$$
(10.2.3)

Assume that all coefficients of  $x_1$  are positive. Then  $f \in P_d$  iff for each  $1 \leq i \leq d$  all coefficients of  $x_i$  are positive.

Proof. The homogeneous part of f is

$$f^{\mathsf{H}}(\mathbf{x}) = \prod_{k=1}^{n} (a_{1k}x_1 + \dots + a_{dk}x_d).$$

If the signs of the coefficients of  $x_i$  are positive then the coefficients of  $f^H(x_1, \ldots, x_d)$  are all positive.

Conversely, the terms in  $f^H$  involving only variables  $x_1$  and  $x_i$  are easily found. They factor into

$$\prod_{k=1}^{n} (a_{1k}x_1 + a_{ik}x_i)$$

If  $f^H$  satisfies the positivity property then the coefficients of this polynomial all have the same sign. Since the coefficients of  $x_1$  are all positive, the coefficients of  $x_i$  are positive. This is Lemma 2.1.

**Definition 10.14.** If  $f, g \in P_d$  then f and g *interlace* iff  $f + \alpha g$  is in  $\pm P_d$  for all  $\alpha$ . If in addition the degree of f is one more than the degree of g then we say  $f \leq g$ . In this case  $(f + g)^H = f^H$ , and so the only condition we need to verify is substitution. This leads to an equivalent definition in terms of substitutions:

If f, 
$$g \in \mathbb{P}_d$$
 then  $f \lessdot g$  iff  $f(\mathbf{x}_i^a) \lessdot g(\mathbf{x}_i^a)$  for all  $\mathbf{a} \in \mathbb{R}^d$  and  $1 \leq i \leq d$ .

The following important result is an immediate consequence of the definition. See the proof of Theorem 1.43.

**Theorem 10.15.** *If*  $T: P_d \longrightarrow P_d$  *is a linear transformation then* T *preserves inter-lacing.* 

**Lemma 10.16.** If f,  $g \in P_d$  satisfy  $f \leq g$  then  $f^H \leq g^H$ .

*Proof.* The transformation  $f \mapsto f^H$  maps  $P_d$  to itself.

Both  $\mathbf{Sub}_d$  and  $\mathbb{P}_d$  satisfies a loose form of closure. We will study the general question of closure in the next chapter.

**Lemma 10.17.** Assume f is a polynomial in d variables that satisfies the homogeneity condition. If there are polynomials  $f_i \in P_d$  such that  $\lim f_i = f$  then f is in  $P_d$ .

*Proof.* Applying Lemma 10.3 shows that f satisfies substitution. By hypothesis f satisfies the positivity condition, and hence  $f \in P_d$ .

#### Quadratics in $P_d$

There is an effective criterion to determine if a quadratic polynomial is in  $P_d$ . If we write  $f = \sum a_1 x^I$  then for all values of  $x_2, \ldots, x_d$  the polynomial f must have all real roots as a quadratic in  $x_1$ . This is true if and only if the discriminant is non-negative. If we write out the discriminant then we have a quadratic form that must be positive semi-definite.

For simplicity we only consider the case d = 3, but the general case presents no difficulties. Suppose that

$$\begin{split} \mathsf{f}(x,y,z) &= a_{000} + z\,a_{001} + z^2\,a_{002} + y\,a_{010} + y\,z\,a_{011} + y^2\,a_{020} + \\ &\quad x\,a_{100} + x\,z\,a_{101} + x\,y\,a_{110} + x^2\,a_{200} \end{split}$$

then the discriminant is

$$\begin{aligned} a_{100}^{\ 2} + 2\,z\,a_{100}\,a_{101} + z^2\,a_{101}^{\ 2} + 2\,y\,a_{100}\,a_{110} + 2\,y\,z\,a_{101}\,a_{110} \\ + y^2\,a_{110}^2 - 4\,a_{000}\,a_{200} - 4\,z\,a_{001}\,a_{200} - 4\,z^2\,a_{002}\,a_{200} \\ - 4\,y\,a_{010}\,a_{200} - 4\,y\,z\,a_{011}\,a_{200} - 4\,y^2\,a_{020}\,a_{200} \end{aligned}$$

Now we can homogenize this, and consider it as a quadratic q(y, z, w). Since the discriminant is non-negative, the quadratic must be non-negative for all values of y, z, w. Consequently, the matrix Q of the quadratic form q

$$\begin{pmatrix} a_{110}^2 - 4 a_{020} a_{200} & a_{101} a_{110} - 2 a_{011} a_{200} & a_{100} a_{110} - 2 a_{010} a_{200} \\ a_{101} a_{110} - 2 a_{011} a_{200} & a_{101}^2 - 4 a_{002} a_{200} & a_{100} a_{101} - 2 a_{001} a_{200} \\ a_{100} a_{110} - 2 a_{010} a_{200}, & a_{100} a_{101} - 2 a_{001} a_{200} & a_{100}^2 - 4 a_{000} a_{200} \end{pmatrix}$$

$$(10.2.4)$$

is positive semi-definite. Conversely, if Q is positive semi-definite then f(x, y, z) satisfies x-substitution. Note that the diagonal elements of Q are positive if and only if f(x, y, 0), f(x, 0, z), f(0, y, z) each have all real roots. We summarize for d = 3:

**Lemma 10.18.** *A* quadratic polynomial is in  $P_3$  if and only if it satisfies the positivity conditions and the matrix (10.2.4) is positive semi-definite.

We define  $\mathbf{P}_d^{sep}$  to be all those polynomials in  $\mathbb{P}_d$  such that all substitutions are in  $\mathbf{P}^{sep}$ . It is easy to construct quadratic polynomials in  $\mathbf{P}_d^{sep}$ .

Lemma 10.19. The polynomial

$$(a_1x_1 + \dots + a_dx_d)(b_1x_1 + \dots + b_dx_d) - c^2$$

 $\textit{is in } \mathbf{P}_d^{\textit{sep}} \textit{ if } \quad |c| \geqslant \frac{1}{2} \max_{1 \leqslant i < d} \sqrt{\alpha_i b_i}$ 

*Proof.* Notice that Lemma 9.126 does not depend on constants, so if we substitute for all but the i-th variable, the minimum distance between roots is  $2|c|/\sqrt{a_ib_i}$ .

## **10.3 Properties of P\_d**

In this section we establish the important fact that  $P_d$  is closed under differentiation. An important consequence of this will be results about the coefficients of polynomials in  $P_d$ .

As in  $P_2$  we only need to know that substitution holds in one variable to conclude that substitution holds in all variables. This is a very useful result.

**Theorem 10.20.** Suppose that f is a polynomial in d variables, and satisfies (see Definition 10.10)

- the positivity condition
- $x_i$ -substitution for some  $x_i$

then f satisfies substitution, and is in  $P_d$ .

*Proof.* It suffices to show that if f satisfies  $x_1$ -substitution then it satisfies  $x_2$ -substitution. For any  $\mathbf{a} = (a_1, \dots, a_d)$  let  $g(x_1, x_2) = f(x_1, x_2, a_3, \dots, a_d)$ . Since g satisfies  $x_1$ -substitution and  $g^H$  is a subset of  $f^H$ , we can apply Theorem 9.27 to conclude that g satisfies  $x_2$ -substitution.

**Theorem 10.21.** Suppose that  $T: P_d \longrightarrow P_d$  is a linear transformation that preserves degree, and for any monomial  $\mathbf{x}^I$  the coefficients of  $T(\mathbf{x}^I)^H$  are all positive. If  $\mathbf{y} = (\mathbf{y}_1, \dots, \mathbf{y}_e)$  then the linear transformation

$$\mathsf{T}_*(\mathbf{x}^I \mathbf{y}^J) = \mathsf{T}(\mathbf{x}^I) \, \mathbf{y}^J$$

*defines a linear transformation from*  $P_{d+e}$  *to itself.* 

*Proof.* The assumptions on the image of a monomial imply that  $T_*(f)$  satisfies the positivity condition for  $f \in P_{d+e}$ . By Theorem 10.20 it suffices to show that  $T_*(f)$  satisfies  $x_1$ -substitution. If we choose  $\mathbf{b} \in \mathbb{R}^e$  then

$$(\mathsf{T}_* \mathsf{f})(\mathbf{x}, \mathbf{b}) = \mathsf{T}(\mathsf{f}(\mathbf{x}, \mathbf{b}))$$

Since  $f(\mathbf{x}, \mathbf{b}) \in P_d$  by Lemma 10.12 we know  $T(f(\mathbf{x}, \mathbf{b}))$  is in  $P_d$ , and so satisfies  $x_1$ -substitution.

An important special case is when d is 1.

**Corollary 10.22.** Suppose  $T: P \longrightarrow P$  is a linear transformation that preserves degree, and maps polynomials with positive leading coefficient to polynomials with positive leading coefficient. We define a linear transformation on  $P_d$  by

$$\mathsf{T}_{i}(\mathsf{x}_{1}^{e_{1}}\cdots \mathsf{x}_{d}^{e_{d}}) = \mathsf{x}_{1}^{e_{1}}\cdots \mathsf{x}_{i-1}^{e_{i-1}} \mathsf{T}(\mathsf{x}_{i}^{e_{i}}) \, \mathsf{x}_{i+1}^{e_{i+1}}\cdots \mathsf{x}_{d}^{e_{d}}$$

 $T_i$  maps  $P_d$  to itself.

**Theorem 10.23.**  $P_d$  is closed under differentiation. If  $f \in \mathbf{P}_d$  then  $f \leq \frac{\partial f}{\partial x_i}$  for  $1 \leq i \leq d$ .

*Proof.* The linear transformation T of **P** defined by Tg = g + ag' maps **P** to itself for any a. Applying Theorem 10.21 shows that  $f + a \frac{\partial f}{\partial x_i}$  is in  $P_d$  for all a, and the conclusion follows.

**Corollary 10.24.** If  $f = \sum f_I(\mathbf{x})\mathbf{y}^I$  is in  $P_{d+e}$  and J is any index set then  $f_J(\mathbf{x})$  is in  $P_d(|J|)$ .

*Proof.* If  $\mathbf{J} = (j_1, \dots, j_e)$  then we can apply Theorem 10.23 since

$$f_{J}(\mathbf{x}) = \frac{1}{j_{1}!j_{2}!\cdots j_{e}!} \left( \frac{\partial^{j_{1}}}{\partial y_{1}} \cdots \frac{\partial^{j_{e}}}{\partial y_{e}} \right) f \Big|_{\mathbf{y}=0}$$

In  $P_2$  we know that consecutive coefficients interlace. The same is true in  $P_d$ , and the proof is the same.

**Lemma 10.25.** *Suppose that*  $f(\mathbf{x}, \mathbf{y}) \in P_{d+e}$ *, and we write* 

$$f(\mathbf{x}, \mathbf{y}) = \dots + f_{I}(\mathbf{x})\mathbf{y}^{I} + f_{I \cup y_{1}}(\mathbf{x})y_{1}\mathbf{y}^{I} + \dots$$

*then*  $f_I \leftarrow f_{I \cup y_1}$ .

**Theorem 10.26.** *If* f, g,  $h \in P_d$  and  $f \leq g$ ,  $f \leq h$  then

- $g + h \in P_d$
- $f \leq g + h$
- For any  $f, g \in P_d$  then  $fg \in \mathbf{P}_d$ .
- If  $g \leq k$  then  $f k \in P_d$ .

*Proof.* As we saw for two variables, Lemma 1.10 shows that g + h satisfies substitution. Since  $(g + h)^H = g^H + h^H$  it follows that g + h satisfies and positivity condition.

The product fg certainly satisfies substitution. Since  $(fg)^H = f^H g^H$  we see that the positivity condition is satisfied. The last one follows from the corresponding one variable result Lemma 1.31.

Two by two determinants of coefficients are non-positive. This is an immediate consequence of Theorem 9.113.

**Corollary 10.27.** Suppose that  $f(\mathbf{x}, \mathbf{y}, z) = \sum f_{ij}(\mathbf{x}) \mathbf{y}^i z^j$  is in  $P_{d+2}$ . Then

$$\begin{vmatrix} f_{i,j}(\mathbf{x}) & f_{i+1,j}(\mathbf{x}) \\ f_{i,j+1}(\mathbf{x}) & f_{i+1,j+1}(\mathbf{x}) \end{vmatrix} \leqslant 0$$

If we apply this corollary to the Taylor series for  $f(x_1 + y, x_2 + z, x_3, ..., x_d)$  we get

**Corollary 10.28.** If  $f(\mathbf{x}) \in P_d$ ,  $d \ge 2$ , then  $\begin{vmatrix} f & \frac{\partial}{\partial x_1} f \\ \frac{\partial}{\partial x_2} f & \frac{\partial}{\partial x_1} \frac{\partial}{\partial x_2} f \end{vmatrix} \leqslant 0$ .

*Remark* 10.29. If we are given a polynomial f that satisfies the above inequality for all pairs of distinct variables, then it does not imply that  $f \in P_d$ . However, this is true for multiaffine polynomials (See p. 389.).

## **10.4** The analog of P<sup>pos</sup>

 $\mathbf{P}_{d}^{pos}$  is the set of all polynomials in  $\mathbf{P}$  with all positive coefficients. The analog  $\mathbf{P}_{d}^{pos}$  of  $\mathbf{P}^{pos}$  is also defined in terms of the signs of the coefficients.  $\mathbf{P}_{d}^{pos}$  is closed under interlacing, and the homogeneous part of any polynomial in  $\mathbb{P}_{d}$  is always in  $\mathbf{P}_{d}^{pos}$ .

**Definition 10.30.**  $\mathbf{P}_d^{\text{pos}}(n)$  is the collection of all polynomials  $f \in P_d(n)$  such that all coefficients of monomials of degree at most n are positive.

That is, if  $f = \sum a_I x^I$  has total degree n, then  $a_I$  is positive for all  $|I| \leq n$ .

We can easily characterize products of linear terms that are in  $\mathbf{P}_d^{\text{pos}}$ . It follows from Lemma 10.13 that a product (10.2.3) is in  $\mathbf{P}_d^{\text{pos}}$  iff all  $a_{ik}$  are positive, and all  $b_k$  are positive.

#### Lemma 10.31.

- 1.  $\mathbf{P}_{d}^{pos}$  is closed under differentiation.
- 2. If  $f \in \mathbf{P}_d^{pos}$  then for any positive a we have  $f(a, x_2, \dots, x_d) \in \mathbf{P}_{d-1}^{pos}$ .
- 3. If  $f \in \mathbf{P}_{d+e}^{pos}$  then all coefficients  $f_I$  given in (10.1.1) are in  $\mathbf{P}_d^{pos}$ .

*Proof.* We only need to check positivity for the first statement, and this is immediate. The second statement is obvious, but it is false if we substitute a negative value.<sup>1</sup> Finally, differentiate sufficiently many times until  $f_I$  is the constant term, and then substitute 0 for y.

We can determine if a polynomial is in  $\mathbf{P}_d^{pos}$  just by looking at the constant terms with respect to each variable.

**Theorem 10.32.** Suppose that  $f \in P_d$  and for each  $1 \leq i \leq d$  the polynomial  $f(0, \ldots, 0, x_i, 0, \ldots, 0)$  is in  $\mathbf{P}^{pos}$ . Then, f is in  $\pm \mathbf{P}_d^{pos}$ .

*Proof.* We may assume that

$$f(0, ..., 0, x_i, 0, ..., 0)$$

has all positive, or all negative, signs. The proof now proceeds by induction on the number of variables. For d = 2 the hypothesis implies that in

$$f(x_1, x_2) = f_0(x_1) + \dots + f_n(x_1)x_2^n$$

the coefficient  $f_0$  is in  $\mathbf{P}^{\text{pos}}$ . We may assume that all coefficients of  $f_0$  are positive. Since  $f_0 \leq f_1$  we see that all coefficients of  $f_1$  have the same sign. Continuing, all the coefficients of  $f_i$  have sign  $\varepsilon_i$ . Now the hypothesis also implies that  $f(0, x_2)$  is in  $\mathbf{P}^{\text{pos}}$ . Since  $f(0, x_2)$  contains the constant terms of all  $f_i$  it follows that all  $\varepsilon_i$  have the same sign. Thus all coefficients of all the  $f_i$  are positive.

<sup>&</sup>lt;sup>1</sup> f =  $(1 + x_1 + x_2)(3 + x_1 + x_2)$  is in **P**<sup>pos</sup><sub>2</sub>, but f(-2, x<sub>2</sub>) =  $x_2^2 - 1$  is not in **P**<sup>±</sup>.

In general, if  $f \in P_d$  then by induction we know that all substitutions

$$f(x_1, ..., x_{i-1}, 0, x_{i+1}, ..., x_d)$$

are in  $\mathbf{P}_{d-1}^{\text{pos}}$ . We may assume that  $f(0, x_2, \dots, x_d)$  has all positive terms, and then use overlapping polynomials as above to conclude that all terms are positive.

 $\mathbf{P}_{d}^{\text{pos}}$  is closed under interlacing.

**Theorem 10.33.** *If*  $f \in \mathbf{P}_d^{\text{pos}}$ ,  $g \in P_d$  and  $f \leq g$  then g is in  $\mathbf{P}_d^{\text{pos}}$ .

*Proof.* The proof is by induction on the number of variables. For  $P^{pos}$  the result is true. Write

$$f(\mathbf{x}) = f_0(x_2, \dots, x_d) + f_1(x_2, \dots, x_d)x_1 + \dots + f_n(x_2, \dots, x_d)x_1^n.$$
  
$$g(\mathbf{x}) = g_0(x_2, \dots, x_d) + g_1(x_2, \dots, x_d)x_1 + \dots + g_{n-1}(x_2, \dots, x_d)x_1^{n-1}.$$

Since  $f_i \leq g_i$  for all i, and all  $g_i$  are in  $\mathbf{P}_{d-1}^{pos}$  by induction, it follows that the coefficients of  $g_i$  are either all positive, or all negative. Since  $g \in P_d$  the homogeneous part  $g^H$  has all positive coefficients, and so each  $g_i$  has some positive coefficient. This shows that  $g \in \mathbf{P}_d^{pos}$ .

The positivity of substitutions implies that a polynomial in  $P_d$  is in  $P_d^{\text{pos}}$ . Note that if f(x) is positive for all positive x then it is not necessarily the case that all coefficients are positive: consider  $1 + (x - 1)^2$ .

**Lemma 10.34.** If  $f(\mathbf{x}) \in P_d$  and  $f(\mathbf{a})$  is positive for all  $\mathbf{a} \in (\mathbb{R}^+)^d$  then  $f \in \mathbf{P}_d^{pos}$ .

*Proof.* We prove this by induction on d. If d = 1 and f(x) has positive leading coefficient then we can write  $f = \alpha \prod (x + r_i)$ . Since this is positive for all positive x it follows that no  $r_i$  can be negative, and therefore all coefficients are positive.

In general, write  $f(\mathbf{x}, \mathbf{y}) \in P_{d+1}$  as  $\sum f_i(\mathbf{x})\mathbf{y}^i$ . Now for any  $\mathbf{a} \in (\mathbb{R}^+)^d$  the hypothesis implies that  $f(\mathbf{a}, \mathbf{y})$  is positive for all positive  $\mathbf{y}$ . Thus all coefficients  $f_i(\mathbf{a})$  are positive, and so by induction all coefficients of  $f(\mathbf{x}, \mathbf{y})$  are positive.  $\Box$ 

The next two results show the relation between homogeneous polynomials and the positivity condition.

**Lemma 10.35.** If f is a homogeneous polynomial then  $f \in P_d$  iff  $f(1, x_2, ..., x_d) \in \mathbf{P}_{d-1}^{pos}$ .

*Proof.* The assumption that f is homogeneous is equivalent to the equality  $f = f^{H}$ . The result now follows from the definition of  $\mathbf{P}_{d}^{\text{pos}}$ .

**Lemma 10.36.** If  $f \in P_d(n)$  let F be the corresponding homogeneous polynomial. We construct F by replacing each monomial  $\mathbf{x}^I$  by  $\mathbf{x}^I \mathbf{x}_{d+1}^{n-|I|}$ . A necessary and sufficient condition that  $F \in P_{d+1}$  is that  $f \in \mathbf{P}_d^{pos}$ . *Proof.* All coefficients of  $F^H$  are positive. If  $F \in P_{d+1}$  then  $F^H = F$ . Since f is found by substituting 1 for  $x_{d+1}$  in F, all coefficients of f must be positive. Conversely, since the coefficients of F are the same as the coefficients of f we see that the assumption on f implies that F satisfies positivity.

We know that the terms of maximal total degree of a polynomial in  $P_d$  constitute a polynomial that is also in  $P_d$ . If  $f \in \mathbf{P}_d^{pos}$  then there is a stronger result:

**Lemma 10.37.** Suppose  $f \in \mathbf{P}_d^{\text{pos}}(n)$  and  $h_r$  is the polynomial consisting of all terms of total degree r. Then  $h_r \in \mathbf{P}_d^{\text{pos}}(r)$  for  $1 \leq r \leq n$ .

Proof. If we homogenize then

$$F(\mathbf{x}, \mathbf{z}) = g_n + zg_{n-1} + \dots + z^n g_0$$

Since  $g_r$  is the coefficient of  $z^{n-r}$  the conclusion follows.

Here is a condition for interlacing in  ${\bm P}_d^{\text{pos}}$  that only involves positive linear combinations.

**Lemma 10.38.** If  $f,g\in P^{\text{pos}}_d$  and deg(f)>deg(g) then  $f\underline{<}g$  if and only if these two conditions hold

(1) 
$$f + \alpha g \in \mathbf{P}_{d}^{pos}$$
 for all  $\alpha > 0$   
(2)  $f + \alpha g \mathbf{x}_{i} \in \mathbf{P}_{d}^{pos}$  for all  $\alpha > 0$  and some i

*Proof.* Assume that  $f \leq g$ . Both left hand sides are have homogeneous part equal to  $f^{H}$  plus perhaps a contribution for  $g^{H}$ , so they both satisfy the homogeneity conditions. We need to verify substitution. If we consider substituting for all but the  $x_i$  variable, and restrict **a** to be non-negative, then both  $f(\mathbf{x}^a)$  and  $g(\mathbf{x}^a)$  are in  $\mathbf{P}^{\text{pos}}$ . The conclusion now follows from Lemma 2.13.

Conversely, if (1) and (2) are satisfied then we can again use Lemma 2.13 to see that  $f \leq g$ .

Just as for one variable we can define a polar derivative. If  $f \in \mathbf{P}_d^{\text{pos}}$  then write  $f = \sum a_I \mathbf{x}^I$ . The corresponding homogeneous polynomial  $F = \sum a_I \mathbf{x}^I y^{n-|I|}$  is in  $\mathbf{P}_{d+1}$ , and if we differentiate with respect to y and then set y = 1 we get the polar derivative

$$\sum (n - |\mathbf{I}|) a_{\mathbf{I}} \mathbf{x}^{\mathbf{I}}$$
(10.4.1)

which can also be written in the more familiar fashion

$$nf - x_1 \frac{\partial f}{\partial x_1} - \dots - x_n \frac{\partial f}{\partial x_n}$$
 (10.4.2)

We will return to the polar derivative in Section 10.15.

Rather than forming the homogeneous polynomial with respect to all the variables, we can homogenize some of them.

**Lemma 10.39.** If  $f(\mathbf{x}, \mathbf{y}) = \sum_{i=1}^{n} a_i(\mathbf{x}) \mathbf{y}^i \in \mathbf{P}_{d+e}^{pos}(n)$  then  $g(\mathbf{x}, \mathbf{y}, z) = \sum_{i=1}^{n} a_i(\mathbf{x}) \mathbf{y}^i z^{n-|I|} \in \mathbf{P}_{d+e+1}(n)$ 

*Proof.*  $g^H = g$ , so g satisfies homogeneity condition. Since

$$g(\mathbf{x},\mathbf{y},z) = z^{n}f(\mathbf{x},\frac{\mathbf{y}}{z})$$

it follows that g satisfies substitution.

We have a more general form of polar derivative:

**Corollary 10.40.** If  $f(\mathbf{x}, \mathbf{y}) = \sum a_I(\mathbf{x}) \mathbf{y}^I \in \mathbf{P}_{d+e}^{pos}(n)$  then

$$f(\mathbf{x},\mathbf{y}) \leq \sum (n-|\mathbf{I}|)a_{\mathbf{I}}(\mathbf{x})\mathbf{y}^{\mathbf{I}}.$$

# **10.5** Coefficient inequalities for P<sub>2</sub><sup>pos</sup>, hives, and the Horn problem

Every polynomial in  $\mathbf{P}_2^{\text{pos}}$  determines a homogeneous polynomial in P<sub>3</sub>. If  $f(x, y, z) \in \mathbf{P}_3$  is such a homogeneous polynomial then the three polynomials f(x, 1, 0), f(0, x, 1), f(1, 0, x) have all real roots. We call these the *boundary polynomials* of f. We consider the converse:

#### The boundary polynomial problem:

Given three polynomials  $f_1(x)$ ,  $f_2(x)$ ,  $f_3(x)$  with all real roots, when is there a polynomial  $f(x, y, z) \in P_3$  whose boundary polynomials are  $f_1$ ,  $f_2$ ,  $f_3$ ?

Surprisingly, this is related to a famous problem involving matrices.

#### The additive Horn problem:

Fix an integer n, and let  $\alpha$ ,  $\beta$ ,  $\gamma \in \mathbb{R}^n$ . Is there a triple (A, B, C) of Hermitian matrices with A + B + C = 0, and eigenvalues  $\alpha$ ,  $\beta$ ,  $\gamma$ ? If so, we say *Horn's additive problem is solvable for*  $\alpha$ ,  $\beta$ ,  $\gamma$ .[161]

We begin by looking at a particular example, and then establishing inequalities for coefficients of polynomials in  $\mathbf{P}_2^{\text{pos}}$ . Taking logs of these inequalities gives a structure called a *hive*. We then recall the connection between hives and Horn's problem to get information about the boundary polynomial problem.

Suppose that we are given a polynomial of degree n in  $\mathbf{P}_3^{\text{pos}}$  that is homogeneous:

$$\begin{split} \mathsf{f}(x,y,z) &= (z+x+y)(4z+2x+y)(2z+3x+y)(3z+5x+y)\\ &= 24z^4 + 112\,xz^3 + 186\,x^2z^2 + 128\,x^3z + 30\,x^4 + 50\,yz^3 + 175\,xyz^2 + 190\,x^2yz + \\ &\quad 61\,x^3y + 35\,y^2z^2 + 80\,xy^2z + 41\,x^2y^2 + 10\,y^3z + 11\,xy^3 + y^4. \end{split}$$

We can represent the terms of f(x, y) in a triangular diagram.



The three boundary polynomials of f occur on the boundary of this diagram, and are

$$\begin{split} f(x,1,0) &= 30x^4 + 61x^3 + 41x^2 + 11x + 1\\ f(0,x,1) &= x^4 + 10x^3 + 35x^2 + 50x + 24\\ f(1,0,x) &= 24x^4 + 112x^3 + 186x^2 + 128x + 30 \end{split}$$

If we only list the coefficients then we get the triangular diagram



This array of numbers has the property that the coefficients in each of the three kinds of rhombus satisfy  $bc \ge ad$ :



**Proposition 10.41.** For any homogeneous polynomial in  $\mathbf{P}_3^{\text{pos}}$ , construct the triangular array of coefficients. In every rhombus formed by two adjacent triangles the product of the two numbers on the common edge is at least the product of the two remaining vertices.

*Proof.* The inequalities corresponding to the three different kinds of rhombus follow by applying the inequalities for polynomials in  $P_2$  (Theorem 9.113) to f(x, y, 1), f(x, 1, z), and f(1, y, z).

Suppose we choose  $f(x, y) = \sum a_{ij} x^i y^j$  in  $\mathbf{P}_2^{\text{pos}}$ . Since all coefficients are positive, we can define real numbers  $h_{ijk} = \log(a_{ij})$  where i + j + k = n. If we write the inequalities of Proposition 10.41 in terms of the  $h_{ij}$ 

$$\begin{aligned} a_{i+1,j} & a_{i,j+1} \geqslant a_{i,j} & a_{i+1,j+1} \\ a_{i+1,j-1} & a_{i,j-1} \geqslant a_{i,j} & a_{i+1,j-2} \\ a_{i-1,j+1} & a_{i-1,j} \geqslant a_{i,j} & a_{i-2,j+1} \end{aligned}$$

then they translate into inequalities about the  $h_{ijk}$ 's:

$$\begin{split} & h_{i+1,j,k-1} + h_{i,j+1,k-1} \geqslant h_{i,j,k} + h_{i+1,j+1,k-2} \\ & h_{i+1,j-1,k} + h_{i,j-1,k+1} \geqslant h_{i,j,k} + h_{i+1,j-2,k+1} \\ & h_{i-1,j+1,k} + h_{i-1,j,k+1} \geqslant h_{i,j,k} + h_{i-2,j+1,k+1} \end{split}$$

This is precisely the definition of a *hive* [161]. There is a wonderful theorem that relates hives to the Horn problem. The boundary of a hive consists of the three vectors ( $v_1$ ,  $v_2$ ,  $v_3$ ) formed by taking differences of consecutive terms on the boundary. In terms of the  $a_{ij}$ 's this is

$$\begin{split} \nu_1 &= (h_{n-1,1,0} - h_{n,0,0}, h_{n-2,2,0} - h_{n-1,1,0}, \cdots, h_{0,n,0} - h_{1,n-1,0}) \\ \nu_2 &= (h_{0,n-1,1} - h_{0,n,0}, h_{0,n-2,2} - h_{0,n-1,1}, \cdots, h_{0,0,n} - h_{0,1,n-1}) \\ \nu_3 &= (h_{1,0,n-1} - h_{0,0,n}, h_{2,0,n-2} - h_{1,0,n-1}, \cdots, h_{n,0,0} - h_{n-1,0,1}) \end{split}$$

then

**The Hive Theorem:** Horn's problem is solvable for  $(v_1, v_2, v_3)$  if and only if  $(v_1, v_2, v_3)$  is the boundary of a hive.[161]

We now determine the boundary of the hive corresponding to f.

$$\begin{split} \nu_1 &= \left( \log \ \frac{a_{n-1,1}}{a_{n,0}}, \log \ \frac{a_{n-2,2}}{a_{n-1,1}}, \dots, \right) \\ &= \text{ the logs of consecutive coefficients of the boundary polynomial } f(x,1,0) \\ \nu_2 &= \left( \log \ \frac{a_{0,n-1}}{a_{0,n}}, \log \ \frac{a_{0,n-2}}{a_{0,n-1}}, \dots, \right) \\ &= \text{ the logs of consecutive coefficients of the boundary polynomial } f(0,x,1) \\ \nu_3 &= \left( \log \ \frac{a_{1,0}}{a_{0,0}}, \log \ \frac{a_{2,0}}{a_{1,0}}, \dots, \right) \\ &= \text{ the logs of consecutive coefficients of the boundary polynomial } f(1,0,x) \end{split}$$

**Corollary 10.42.** If  $f(x, y, z) \in P_3$  is homogeneous, and  $v_1, v_2, v_3$  are the logs of the consecutive coefficients of the boundary polynomials, then Horn's problem is solvable for  $v_1, v_2, v_3$ .

Although polynomials in  $\mathbf{P}_2^{\text{pos}}$  determine hives, the converse is not true: here is an example of a hive that does not come from a polynomial. We don't write the hive, but rather the coefficients of polynomial; it's easy to check that all of the multiplicative rhombus inequalities are satisfied. If the array were determined by a polynomial in  $\mathbf{P}_2^{\text{pos}}$  then the polynomial corresponding to the coefficient of y,  $11 + 8x + 3x^2$ , would be in **P**, but it isn't.



Horn's problem is connected to the conjecture (now solved - see [64]) that specifies necessary and sufficient conditions for vectors  $\alpha$ ,  $\beta$ ,  $\gamma$  to satisfy Horn's problem. We can use this to get necessary conditions for the boundary polynomial problem. Here is one example. We first order  $\alpha$ ,  $\beta$ ,  $\gamma$ :

$$\begin{array}{l} \alpha: \alpha_1 \geqslant \alpha_2 \geqslant \cdots \geqslant \alpha_n \\ \beta: \beta_1 \geqslant \beta_2 \geqslant \cdots \geqslant \beta_n \\ \gamma: \gamma_1 \geqslant \gamma_2 \geqslant \cdots \geqslant \gamma_n \end{array}$$

Weyl's inequalities are

$$\alpha_i + \beta_j + \gamma_{n+2-i-j} \ge 0$$
 whenever  $i + j - 1 \le n$ 

We now apply this to our problem. For our particular polynomial we have that

$$\begin{aligned} \nu_1 &= \left(\log \frac{61}{30}, \log \frac{41}{61}, \log \frac{11}{41}, \log \frac{1}{11}\right) \\ \nu_2 &= \left(\log \frac{10}{1}, \log \frac{35}{10}, \log \frac{50}{35}, \log \frac{24}{50}\right) \\ \nu_3 &= \left(\log \frac{112}{24}, \log \frac{186}{112}, \log \frac{128}{186}, \log \frac{30}{128}\right) \end{aligned}$$

Notice that they are all decreasing - this is true in general, and is a consequence of Newton's inequalities. Since these quotients are decreasing, they are in the correct order to apply the Weyl inequalities. Using the values of  $v_1$ ,  $v_2$ ,  $v_3$  from above, the following is a necessary condition for the  $a_{i,j}$  to be the coefficients of boundary polynomials.

$$\log \frac{a_{n-i,i}}{a_{n-i+1,i-1}} + \log \frac{a_{0,n-i}}{a_{0,n-i+1}} + \log \frac{a_{n+2-i-j,0}}{a_{a+2-i-j+1}} \ge 0$$
$$\frac{a_{n-i,i} a_{0,n-i} a_{n+2-i-j,0}}{a_{n-i+1,i-1} a_{0,n-i+1} a_{a+2-i-j+1}} \ge 1$$

## **10.6** Log concavity in $P_2^{pos}$

We say earlier ((p. 118)) that the graph of the log of the coefficients of a polynomial in  $\mathbf{P}_2^{\text{pos}}$  is concave. We now prove a similar result for polynomials in  $\mathbf{P}_2^{\text{pos}}$ . Given a polynomial  $\sum a_{ij}x^iy^j$  in  $\mathbf{P}_2^{\text{pos}}$ , the set of points

$$\{(i, j, \log(a_{ij})) \mid a_{ij}x^iy^j \text{ is a term of f }\}$$

is called the graph of the log of the coefficients of f. We view this as the graph of a function defined on the points in the plane corresponding to the exponents appearing in f.

For example, the diagram below displays the log of the coefficients of a polynomial in  $\mathbf{P}_2^{\text{pos}}$ . The coordinates of the point in the lower left corner are (0,0), since 24 is the constant term. The coordinates of the rightmost point are (4,0), and the uppermost point's coordinates are (0,4).



**Proposition 10.43.** If  $f \in \mathbf{P}_2^{pos}$  then the graph of the log of the coefficients is concave. Every triangle in the planar diagram determines a triangle in the graph, and the graph lies on one side of the plane containing that triangle.

*Proof.* We will prove the second statement, for that is what we mean by saying that the graph is concave. Note that it suffices to show that the plane through a triangle lies above all of the three adjacent triangles. We now show that this is a consequence of the rhombus inequalities.

Consider the triangle T with solid lines



The plane in  $\mathbb{R}^3$  containing T is

$$\begin{aligned} (\log a_{r+1,s} - \log a_{r,s})x + (\log a_{r,s+1} - \log a_{r,s})y - z - \\ & \left(r \log a_{r+1,s} + s \, \log a_{r,s+1} - (r+s+1) \log a_{r,s}\right) = 0 \end{aligned}$$

The expression on the left side takes negative values for points above the plane, and positive values for points below the plane. If we evaluate it at the vertex  $(r + 1, s + 1, a_{r+1,s+1})$  of the dotted adjacent triangle we get

$$\log a_{r+1,s} + \log a_{r,s+1} - \log a_{r,s} - \log a_{r+1,s+1} = \\ \log \frac{a_{r+1,s} a_{r,s+1}}{a_{r,s} a_{r+1,s+1}} \ge 0$$

where the last inequality is the rhombus inequality. A similar computation shows that the plane lies above the other two adjacent triangles, so the graph is concave.

**Corollary 10.44.** Suppose that  $f = \sum a_{i,j}x^iy^j \in \mathbf{P}_2^{\text{pos}}$ . For all non-negative i, j, r, s for which  $a_{r,s} \neq 0$  we have

$$\left(\frac{a_{r+1,s}}{a_{r,s}}\right)^{i} \left(\frac{a_{r,s+1}}{a_{r,s}}\right)^{j} \geqslant \frac{a_{r+i,s+j}}{a_{r,s}}$$

*Proof.* Evaluate the plane through the triangle

$$\{(r, s, \log a_{r,s}), (r+1, s, \log a_{r+1,s}), (r, s+1, \log a_{r,s+1})\}$$

at the point  $(r + i, s + j, \log a_{r+i,s+j})$ . The result is positive, and simplifying yields the inequality.

*Remark* 10.45. If  $a_{0,0} = 1$ , then the inequality shows that the coefficients of f satisfy a simple bound:

$$a_{1,0}^{i} a_{0,1}^{j} \ge a_{i,j}.$$

### 10.7 Extending two polynomials

The boundary polynomial problem asks when can three polynomials be realized as the boundary polynomials of a polynomial in  $P_2^{\text{pos}}$ . We now ask the simpler question

Given two polynomials f(x), g(x) in  $\mathbf{P}^{\text{pos}}$ , describe the structure of all F(x, y) in  $\mathbf{P}_2^{\text{pos}}$  such that F(x, 0) = f(x) and F(0, y) = g(y). We say that F is an extension of f, g.

There is a trivial requirement that guarantees the existence of an extension. This result is similar to Lemmas 9.103 and 9.105.

**Lemma 10.46.** Assume that  $f, g \in \mathbf{P}^{pos}$ . There is an extension of f, g if and only if f(0) = g(0).

*Proof.* If there is an extension then F(0,0) = f(0) = g(0). Conversely, we may assume that f(0) = g(0) = 1. We can write  $f(x) = \prod(1 + r_i x)$  and  $g(x) = \prod(1 + s_i x)$ . The product below is the desired extension.

$$\prod (1 + r_i x + s_i y)$$

Suppose that F is an extension of f, g. We know that we can write  $F(x, y) = |I + xD_1 + yD_2|$  where  $D_1$  and  $D_2$  are positive definite. Moreover, since  $f(x) = |I + xD_1|$  we know the eigenvalues of  $D_1$ : they are the negative reciprocals of the roots of f. Similarly, we know the eigenvalues of  $D_2$ . We may assume that  $D_1$  is diagonal. Write  $D_2 = O \Lambda O^t$  where  $\Lambda$  is a diagonal matrix consisting of the eigenvalues of  $D_2$ , and O is an orthogonal matrix. Consequently,

There is an onto map from O(n), the group of orthogonal matrices, to the extensions of f, g.

Since the orthogonal matrices in O(2) are easy to describe we can completely determine the extensions for a quadratic.

**Lemma 10.47.** Suppose  $f = (1 + r_1 x)(1 + +r_2 x)$  and  $g = (1 + s_1 x)(1 + s_2 x)$ . All extensions of f, g are of the form below, where W lies in the interval determined by  $r_1s_1 + r_2s_2$  and  $r_1s_2 + r_2s_1$ .



*Proof.* If the orthogonal two by two matrix is a rotation matrix, the general extension of f, g is given by the determinant of

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \chi \begin{pmatrix} r_1 & 0 \\ 0 & r_2 \end{pmatrix} + \chi \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} s_1 & 0 \\ 0 & s_2 \end{pmatrix} \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$$

If we simplify the coefficient of xy in the determinant we find it to be

$$\frac{1}{2} [(r_1 + r_2)(s_1 + s_2) - (r_1 - r_2)(s_1 - s_2)\cos(2\theta)]$$

The two values at  $\cos 2\theta = \pm 1$  yield the bounds of the lemma. We get the same result if the orthogonal matrix does not preserve orientation.

The two ends of the interval for *W* are the values that arise from the two different products

$$\begin{aligned} &(1+r_1x+s_1y)(1+r_2x+s_2y)\\ &(1+r_1x+s_2y)(1+r_2x+s_1y). \end{aligned}$$

In general, if the polynomials are  $\prod (1 + r_i x)$  and  $\prod (1 + s_i y)$  then there are n! different products:

$$\prod (1 + r_i x + s_{\sigma i} y)$$

where  $\sigma$  is a permutation of 1,..., n. If n is odd then O(n) is connected, so the set of all extensions is connected. In the case n = 2 we saw that this set is convex, and the boundary consists of the two different products. This may be the case in general - see Question 123.

It follows from the above that if the quadratic f has all equal roots then there is a unique extension. This is true for any n.

**Lemma 10.48.** If  $g \in P^{pos}(n)$  then g(x) and  $(x + \alpha)^n$  have a unique extension.

*Proof.* Suppose f(x, y) is an extension, so that  $f(x, 0) = (x + \alpha)^n$  and f(0, y) = g(y). If we write  $f(x, y) = \sum f_i(x) y^i$  then  $f_0 = (x + \alpha)^n$  and  $f_0 \leq f_1$ , so there is a constant  $a_1$  such that  $f_1(x) = a_1(x + \alpha)^{n-1}$ . Continuing, we find constants  $a_i$  so that  $f_i(x) = a_i(x + \alpha)^{n-i}$ . Since  $f(0, y) = \sum a_i \alpha^{n-i} y^i$  we see that g(y) uniquely determines the  $a_i$ .

If  $f = \sum a_{ij} x^i y^j \in \mathbf{P}_2^{\text{pos}}$  is an extension of two polynomials, then  $\begin{vmatrix} a_{10} & a_{11} \\ a_{00} & a_{10} \end{vmatrix}$  only has one term that depends on the extension. We assume that f = |I + x D + y C| where D a diagonal matrix with entries  $\{d_i\}$ , and  $C = (c_{ij}) = \mathbb{O}E\mathbb{O}^t$  where E is diagonal with entries  $\{e_i\}$ .

Since every entry of y C has a factor of y, the only way we can get a term xy in the expansion of the determinant is to take x and y terms from the diagonal. Thus, the coefficient  $a_{11}$  of xy is

$$\sum_{i \neq j} d_i \, c_{jj} = \left( \sum d_i \right) \, \left( \sum c_{jj} \right) - \sum d_i c_{ii}$$

Now  $a_{00} = 1$ ,  $a_{10} = \sum d_i$  and  $a_{01} = \sum c_{jj}$ , so we conclude that

$$\begin{vmatrix} a_{10} & a_{11} \\ a_{00} & a_{10} \end{vmatrix} = \sum d_{i}c_{ii}$$

If we write  $\texttt{O}=(o_{\mathfrak{i}\mathfrak{j}})$  then  $c_{\mathfrak{i}\mathfrak{i}}=\sum e_{\mathfrak{j}}o_{\mathfrak{i}\mathfrak{j}}^2$  and consequently

$$\begin{vmatrix} a_{10} & a_{11} \\ a_{00} & a_{10} \end{vmatrix} = \sum d_i e_j o_{ij}^2$$

Now the maximum of  $\sum d_i e_j o_{ij}^2$  over all orthogonal matrices 0 is  $\sum d'_i e'_i$  where  $\{d'_i\}$  is  $\{d_i\}$  written in increasing order, and  $\{e'_i\}$  is  $\{e_i\}$  written in increasing order (Lemma 10.49). The minimum is  $\sum d'_i e''_i$  where  $\{e''_i\}$  is  $\{e_i\}$  written in decreasing order We thus can make a more precise inequality:

$$\sum d_i' e_i' \geqslant \begin{vmatrix} a_{10} & a_{11} \\ a_{00} & a_{10} \end{vmatrix} \geqslant \sum d_i' e_i''$$

When n = 2 we see that

$$\begin{vmatrix} a_{10} & a_{11} \\ a_{00} & a_{10} \end{vmatrix} = \cos^2(\theta) \left( d_1 e_1 + d_2 e_2 \right) + \sin^2(\theta) \left( d_1 e_2 + d_2 e_1 \right)$$

which shows that the determinant takes every value in the interval determined by the largest and smallest products.

We now prove the Lemma used above; it is a generalization of the wellknown rearrangement lemma.

**Lemma 10.49.** If  $a_1, \ldots, a_n$  and  $b_1, \ldots, b_n$  are increasing sequences of positive numbers then

$$\max_{\mathcal{O}} \sum_{i,j=1}^{n} o_{ij}^2 a_i b_j = \sum_i a_i b_i$$

where the maximum is over all n by n orthogonal matrices  $O = (o_{ij})$ .

*Proof.* If  $A = diag(a_1, ..., a_n)$  and  $B = diag(b_1, ..., b_n)$  then

$$\sum o_{ij}^2 a_i b_j = trace (O^t A O B)$$

Since orthogonal matrices are normal, we can apply Bourin's matrix Chebyshev inequality [21, page 3] to conclude that

trace 
$$(O^{t}AOB) \leq trace (O^{t}OAB) = trace (AB) = \sum a_{i}b_{i}$$
.

#### **10.8** Symmetric Polynomials

In this section we begin an investigation of the symmetric polynomials in  $\overline{\mathbf{P}}_d$ . We say that  $f(\mathbf{x})$  is symmetric if  $f(\mathbf{x}) = f(\sigma \mathbf{x})$  for all permutations  $\sigma$  of  $x_1, \ldots, x_d$ . For instance, the elementary symmetric functions are in  $\overline{\mathbf{P}}_d$ , and the product of any two symmetric functions in  $\overline{\mathbf{P}}_d$  is again a symmetric polynomial in  $\overline{\mathbf{P}}_d$ .

In two variables a homogeneous polynomial f(x, y) is symmetric if and only if it has the form

$$a_0y^n + a_1xy^{n-1} + a_2x^2y^{n-2} + \dots + a_2x^{n-2}y^2 + a_1x^{n-1}y + a_0x^n$$

The roots of f consist of paired roots  $\{r, 1/r\}$ , along with some  $\pm 1$ 's. In degree two such a polynomial is

$$y^2 + (r + \frac{1}{r})xy + x^2$$

It's immediate that the symmetric homogeneous polynomials in  $P_2(2)$  are of the form  $x^2 + \alpha xy + y^2$  where  $\alpha \ge 2$ . We can also characterize the symmetric homogeneous polynomials of degree two in three variables.

**Lemma 10.50.** The symmetric homogeneous polynomials in  $P_3(2)$  are of the form

$$x^2 + y^2 + z^2 + \alpha(xy + xz + yz)$$
 where  $\alpha \ge 2$ 

*Proof.* Setting z = 0 implies  $x^2 + \alpha x + 1 \in \mathbf{P}$ , and thus  $\alpha \ge 2$ . Conversely, if

$$D = \begin{pmatrix} r & 0 \\ 0 & 1/r \end{pmatrix} \qquad O = \frac{1}{r+1} \begin{pmatrix} \sqrt{r^2 + r + 1} & -\sqrt{r} \\ \sqrt{r} & \sqrt{r^2 + r + 1} \end{pmatrix}$$

then the determinant f of  $xI + yD + zODO^{t}$  equals

$$x^{2} + y^{2} + z^{2} + (r + 1/r)(xy + xz + yz)$$

Since D is positive definite and O is orthogonal it follows that  $f \in P_3$ . Finally, note that r + 1/r takes on all values  $\ge 2$ .

We can show that the symmetric homogeneous polynomials in  $P_d(2)$  follow the same pattern, but they don't always have a determinant representation.

**Lemma 10.51.** The symmetric homogeneous polynomials in  $P_d(2)$  are of the form

$$\sum_{1}^{d} x_{i}^{2} + \alpha \sum_{i < j} x_{i} y_{j} \qquad \textit{where } \alpha \geqslant 2$$

*Proof.* We use the definition of  $P_d$ . The homogeneous part is certainly positive, so we check substitution. Since we are considering a quadratic, we only need to check that the discriminant of  $x_d$  is non-negative. Subsituting  $a_i$  for  $x_i$  we get

$$\left(\alpha \sum_{1}^{d-1} \alpha_{i}\right)^{2} - 4\left(\sum_{1}^{d-1} \alpha_{i}^{2}\right)$$
$$= (\alpha - 2)\left((\alpha + 2)\sum_{1}^{d-1} \alpha_{i}^{2} + 2\alpha \sum_{i < j} \alpha_{i} \alpha_{j}\right)$$

The factor  $\alpha - 2$  is non-negative by hypothesis, and the second factor is a quadratic form in  $a_1, \ldots, a_{d-1}$  with matrix Q is

$$\begin{pmatrix} \alpha + 2 & \alpha & \alpha & \dots \\ \alpha & \alpha + 2 & \alpha & \dots \\ \alpha & \alpha & \alpha + 2 & \dots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix} = 2I + \alpha J$$

where J is the all one matrix. Now the eigenvalues of J are d - 1, 0, ..., 0, so the eigenvalues of Q are  $\alpha(d - 1) + 2, 2, ..., 2$ . Thus Q is positive definite, so the discriminant is non-negative, and our polynomial is in P<sub>d</sub>.

**Lemma 10.52.** *The polynomial* f(x, y, z, w) *given below has no representation of the form*  $|xI + yD_1 + zD_2 + wD_3|$  *where*  $D_1, D_2, D_3$  *are positive definite two by two matrices and*  $\alpha \ge 2$ .

$$f(x, y, z, w) = x^{2} + y^{2} + z^{2} + w^{2} + \alpha(xy + xz + xw + yz + yw + zw)$$

*Proof.* Since  $f(x, y, 0, 0) = x^2 + \alpha xy + y^2$  we see that we must have  $\alpha \ge 2$ , and that we can write f(x, y, 0, 0) = (x + r)(x + 1/r) where  $r \ge 1$ . We may assume that  $D_1$  is diagonal, with diagonal r, 1/r, and so  $\alpha = r + 1/r$ .

We proceed to determine D<sub>2</sub>. f is symmetric for all 24 permutations of the variables. Thus  $|I+xD_1| = |I+xD_2|$ , and therefore D<sub>2</sub> has the same eigenvalues as D<sub>1</sub>. We write

$$D_2 = \begin{pmatrix} \cos t & \sin t \\ -\sin t & \cos t \end{pmatrix} \begin{pmatrix} r & 0 \\ 0 & 1/r \end{pmatrix} \begin{pmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{pmatrix}$$

Since f(1, x, 0, 0) = f(0, x, 1, 0) we can equate the coefficients of x:

$$r + \frac{1}{r} = 2\cos^2(t) + \frac{(r^4 + 1)\sin^2(t)}{r^2}$$

Solving this equation yields

$$\label{eq:cost} \begin{split} \cos t &= \pm \frac{1}{1+r} \sqrt{1+r+r^2} \\ \sin t &= \pm \frac{1}{1+r} \sqrt{r} \end{split}$$

Consequently there are exactly four choices for  $D_1$  and for  $D_2$ :

$$\begin{split} D_1 &= \frac{1}{r+1} \begin{pmatrix} e_1 \sqrt{1+r+r^2} & e_2 \sqrt{r} \\ -e_2 \sqrt{t} & e_1 \sqrt{1+r+r^2} \end{pmatrix} \\ D_2 &= \frac{1}{r+1} \begin{pmatrix} f_1 \sqrt{1+r+r^2} & f_2 \sqrt{r} \\ -f_2 \sqrt{t} & f_1 \sqrt{1+r+r^2} \end{pmatrix} \end{split}$$

where  $e_1, e_2, f_1, f_2$  are  $\pm 1$ . If we equate the coefficients of x in  $|xD_1 + D_2| = |xD_1 + I|$  we get

$$\frac{2}{(1+r)^2} \left( e_2 f_2 r + e_1 f_1 (1+r+r^2) \right) = \frac{2e_1}{1+r} \sqrt{1+r+r^2}$$

It's easy to see that this equation has no solution for  $r \ge 1$ .

A symmetric homogeneous polynomial F(x, y, z) in  $P_3(3)$  has coefficient array 1

We first show that for any  $a \ge 3$  there is at least one b such that  $F(x, y, z) \in P_3$ . If we take  $a = \alpha + 1$  and  $b = 3\alpha$  then  $\alpha \ge 2$ . Now

$$(x + y + z)(x^{2} + y^{2} + z^{2} + \alpha(xy + xz + yz)) \in P_{3}$$

and has coefficient array

1

$$1$$

$$1 + \alpha \qquad 1 + \alpha$$

$$1 + \alpha \qquad 3\alpha \qquad 1 + \alpha$$

$$1 + \alpha \qquad 1 + \alpha$$

1

Newton's inequality applied to the first and second rows yield  $a \ge 3$  and  $b \ge 2a$ . However, we can get more precise results. As above, the roots of the bottom row are -r, -1/r, -1, so a = 1 + r + 1/r. We assume r > 1. Since the second row interlaces the first, we have that

$$-\mathfrak{r}\leqslant \frac{-\mathfrak{b}-\sqrt{\mathfrak{b}^2-4\mathfrak{a}^2}}{2\mathfrak{a}}\leqslant -1\leqslant \frac{-\mathfrak{b}+\sqrt{\mathfrak{b}^2-4\mathfrak{a}^2}}{2\mathfrak{a}}\leqslant -1/\mathfrak{r}$$

These inequalities yield

$$a(r+1/r) \ge b \ge 2a$$

Note that if r = 1 then a = 3 and b = 6. This is  $(x + y + z)^3$ , which is the unique extension of the bottom row and the left hand column. These are not the best bounds; empirically we find that the intervals for b appear to be

r=2	$\mathfrak{b} \in$	$(\frac{29}{4})$	,	$\frac{16}{2})$
r=3	$\mathfrak{b} \in$	$(\frac{90}{9})$	,	$\frac{\overline{38}}{3}$ )
r=5	$\mathfrak{b} \in$	$(\frac{326}{25})$	,	$\frac{142}{5})$

When is  $\sum a_i \sigma_i \in MA_d$ , where  $\sigma_i$  is the i'th elementary symmetric polynomial in  $x_1, \ldots, x_d$ ? If we set all  $x_i$  equal to x, then  $\sum a_i {d \choose i} x^i \in \mathbf{P}$ . We are not able to prove the converse; as usual we can't get the final factorial.

**Lemma 10.53.** *If*  $\sum a_i x^i \in P$  *then* 

- 1.  $\sum a_i \sigma_i \in MA_d$ .
- 2.  $\sum a_i i! \sigma_i \in MA_d$ .

*Proof.* The first sum is the coefficient of  $y^n$  in

$$\sum a_{i}y^{i}\left(\prod_{1}^{d}(y+x_{i})\right) = \sum a_{i}y^{i} \cdot \sum y^{n-i}\sigma_{i}$$

while the second equals

$$\left(\sum a_i D_y^i\right) \sum y^{n-i} \sigma_i \bigg|_{y=0}$$

**Example 10.54.** We give an example that supports the possibility that the converse is true. Consider the question

For which m is  $xyz - x - y - z + m = \sigma_3 - \sigma_1 + m \sigma_0 \in MA_3$ ?

We find

$$\begin{aligned} x^{3} - x + \mathfrak{m} \in \mathbf{P} & \implies & |\mathfrak{m}| \leqslant \sqrt{4/27} \\ x^{3}/6 - x + \mathfrak{m} \in \mathbf{P} & \implies & |\mathfrak{m}| \leqslant \sqrt{8/9} \\ \binom{3}{3}x^{3} - \binom{3}{1}x + \binom{3}{0}\mathfrak{m} \in \mathbf{P} & \implies & |\mathfrak{m}| \leqslant 2 \end{aligned}$$

We know from Example11.61 that the necessary and sufficient condition is that  $|m| \leq 2$ .

### **10.9** Linear transformations on P<sub>d</sub>

We have looked at linear transformations that map  $P_d$  to itself. In this section we look at mappings that either delete variables or introduce new variables.

The repeated application Theorem 10.21 with the linear transformation  $x \mapsto ax + b$  yields

**Corollary 10.55.** Choose  $a_1, b_1, ..., a_d, b_d \in \mathbb{R}$  where all of the  $a_i$ 's are positive. The linear transformation below maps  $\mathbf{P}_d$  to itself:

$$f(x_1,\ldots,x_d)\mapsto f(a_1x_1+b_1,\ldots,a_dx_d+b_d).$$

We can allow some of the  $a_i$  to be zero.

**Lemma 10.56.** *If*  $f(\mathbf{x}) \in \mathbf{P}_d$  *and exactly* r *terms of*  $a_1, \ldots, a_d$  *are non-zero, and they are positive, then*  $f(a_1x_1, \ldots, a_dx_d) \in \mathbf{P}_r$ .

We can also add new variables.

**Lemma 10.57.** *If*  $f \in P_d(n)$  *then*  $g = f(x_1 + y_1, x_2 + y_2, ..., x_d + y_d)$  *is in*  $P_{2d}(n)$ .

*Proof.* Certainly g satisfies substitution. Every term  $x_1^{i_1} \cdots x_d^{i_d}$  in f gives rise to  $(x_1 + y_1)^{i_1} \cdots (x_d + y_d)^{i_d}$  in g. This shows that

$$g^{H}(\mathbf{x}) = f^{H}(x_{1} + y_{1}, \dots, x_{d} + y_{d})$$

Since f satisfies the homogeneity conditions, so does g.

.. ..

**Lemma 10.58.** If  $f(\mathbf{x}) \in \mathbf{P}_d$  then the polynomial  $g = f(x, x, x_3, ..., x_d)$  is in  $\mathbf{P}_{d-1}$ . Also,  $f(x, x, ..., x) \in \mathbf{P}$ .

*Proof.* If d is two then this is Lemma 9.32. If d is larger than two then g satisfies  $x_d$ -substitution. Since  $f(x, x, x_3, ...)^H = f^H(x, x, x_3, ...)$  it follows that g satisfies the homogeneity condition. The second assertion follows by induction.

We can replace one variable by more complicated expressions.

**Lemma 10.59.** *If*  $f(\mathbf{x}) \in \mathbf{P}_d$ ,  $\beta \in \mathbb{R}$ , and  $a_1, \ldots, a_{d-1}$  are non-negative then

 $g(x_1, \dots, x_{d-1}) = f(x_1, x_2, \dots, x_{d-1}, a_1x_1 + \dots + a_{d-1}x_{d-1} + \beta) \in \mathbf{P}_{d-1}$ 

*Proof.* By induction on d. The case d = 2 is Lemma 10.58. Since g satisfies the positivity condition it remains to show that g satisfies substitution. If we substitute  $x_i = \alpha_i$  for all  $x_i$  except  $x_1$  and  $x_d$  then  $f(x_1, \alpha_2, ..., \alpha_{d-1}, x_d) \in \mathbf{P}_2$ . Consequently,  $f(x_1, \alpha_2, ..., \alpha_{d-1}, a_1x_1 + b)$  is in **P** for any choice of b. Choosing  $b = a_2\alpha_2 + \cdots + a_{d-1}\alpha_{d-1} + \beta$  and observing that

$$g(x_1, \alpha_2, ..., \alpha_{d-1}) = f(x_1, \alpha_2, ..., \alpha_{d-1}, a_1x_1 + a_2\alpha_2 + \cdots + a_{d-1}\alpha_{d-1} + \beta)$$

it follows that g satisfies substitution.

We now use these results to show how very general linear combinations preserve **P**.

**Theorem 10.60.** Suppose  $f(\mathbf{x}) \in \mathbf{P}_d$ . If M is a d by d matrix with positive entries then  $f(M\mathbf{x}) \in \mathbf{P}_d$ .

*Proof.* Let  $M = (m_{ij})$  and define a polynomial in d<sup>2</sup> variables  $x_{ij}$ 

$$g = f\left(\sum_{i} m_{i1}x_{i1}, \dots, \sum_{i} m_{id}x_{id}\right).$$

Repeated applications of Lemma 10.56 and Lemma 10.57 show that g is in  $P_{d^2}$ . We now use Lemma 10.58 to identify all pairs of variables  $x_{ij}$  and  $x_{ik}$ . The resulting polynomial is exactly f(Mx).

**Lemma 10.61.** If f,  $g \in \mathbf{P}_d$  and  $f = \sum_I a_I \mathbf{x}^I$  then for any index K

$$\sum_{I+J=K} \mathfrak{a}_I \frac{\mathfrak{g}^{(J)}}{J!} \in \mathbf{P}_{\mathsf{d}}.$$

*Proof.* Using the Taylor series for g we get

$$\mathsf{f}(\mathbf{y})\mathsf{g}(\mathbf{x}+\mathbf{y}) = \sum_{\mathbf{I},\mathbf{J}} \, \mathfrak{a}_{\mathbf{I}} \, \mathbf{x}^{\mathbf{I}} \, \frac{\mathbf{y}^{\mathbf{J}}}{\mathbf{J}!} \, \frac{\partial^{\mathbf{J}}}{\partial \mathbf{x}^{\mathbf{J}}} \, \mathfrak{g}(\mathbf{x})$$

The coefficient of  $\mathbf{y}^{\mathbf{K}}$  is the sum in the conclusion.

#### **10.10** The graph of polynomials in $P_d$

In P<sub>2</sub> we used the geometry of the graph of  $f(x, y) \in P_2$  to show that  $f(x, x) \in P$ . Now we use the fact that if  $f(x) \in P_d$  then  $f(x, x, ..., x) \in P$  to get information about the geometry of the graph of f.

For clarity we'll take d = 3. Choose  $f(x) \in P_d(n)$ . The graph  $G_f$  of f is  $\{(x, y, z) \mid f(x, y, z) = 0\}$ . Choose any point v = (a, b, c) on the plane  $x_1 + x_2 + x_3 = 0$ . Since f(x - v) = f(x - a, y - b, z - c) is in  $P_d$ , the equation f(x - a, x - b, x - c) = 0 has n solutions. Denote the i-th largest solution by  $s_i(v)$ , and let  $w_i(v) = (s_i(v) - a, s_i(v) - b, s_i(v) - c)$ . By construction we know that  $f(w_i(v)) = 0$ , so  $w_i(v)$  is a point on the graph of f. Consider the map  $v \mapsto w_i(v)$ . This is a function from the plane  $x_1 + x_2 + x_3 = 0$  to the graph of f that is 1 - 1, and continuous since solutions to equations are continuous functions of coefficients. Moreover, since the union of all lines of the form  $\{v + t(1, 1, 1)\}$  is a partition of  $\mathbb{R}^d$ , every point on the graph  $G_f$  is the union of n subsets  $G_{f_i}$  of  $\mathbb{R}^d$ , each homeomorphic to  $\mathbb{R}^{d-1}$ . These subsets might intersect.

In addition, since all coefficients of  $f^H$  are positive, there is an  $\alpha$  such if  $|\mathbf{x}| > \alpha$  and all coefficients of  $\mathbf{x}$  are positive then  $f(\mathbf{x}) > 0$ . Similarly, we may assume that if all coefficients are negative then  $f(\mathbf{x}) \neq 0$ .

Let  $\mathbb{R}^+$  be the set of all points with all positive coefficients,  $\mathbb{R}^-$  the set with all negative coefficients, and S be the ball of radius  $\alpha$ . The previous paragraph shows that the graph  $G_f$  separates  $\mathbb{R}^+ \setminus S$  from  $\mathbb{R}^- \setminus S$ .

We have seen that if a, b, c are positive and  $f \in P_3(n)$ , then  $f(ax + \alpha, bx + \beta, cx + \gamma)$  is in **P**(n). This is a special case of the following observation:

If a curve goes to infinity in all coordinates in  $\mathbb{R}^+$  and also in  $\mathbb{R}^-$  then it meets the graph of f in at least n points.

Here's a variation: suppose that the line  $\mathcal{L}$  is parallel to a coordinate axes. For instance,  $\mathcal{L}$  could be  $\{(a, b, c) + t(1, 0, 0) \mid t \in \mathbb{R}\}$ . Then,  $\mathcal{L}$  meets the graph in n points. To see this, notice that there is a sufficiently large T such that the line segment from (a + T, b, c) to (a + T, 0, 0) does not meet G<sub>f</sub>. Neither does the ray  $\{(a + T + t, 0, 0) \mid t \ge 0\}$ . Similarly analyzing negative values we can find a corresponding S and conclude that the piecewise linear curve consisting of the segments

$$\{(a + T + t, 0, 0) | t \ge 0\}$$
  
(a + T, b, c) - (a + T, 0, 0)  
(a + T, b, c) - (a - S, b, c)  
(a - S, b, c) - (a - S, 0, 0)  
{(a - S - t, 0, 0) | t \ge 0}

meets  $G_f$  in n points, and consequently the line  $\{(a+t,b,c) \mid t \in \mathbb{R}\}$  meets  $G_f$  in n points.

If  $f \in \mathbf{P}_d^{\text{pos}}$  then  $\mathbf{x} \in \mathbb{R}^+$  implies that  $f(\mathbf{x}) \neq 0$ . Any curve from a point in  $\mathbb{R}^+$  that goes to infinity in  $\mathbb{R}^-$  must intersect every  $\mathcal{G}_i$ .

#### **10.11** Differential operators

In this section we introduce differential operators on  $\mathbf{P}_d$  of the form

$$f(\frac{\partial}{\partial \mathbf{x}}) = f(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_d})$$

where f is a polynomial in d variables. To simplify notation, we will sometimes write  $\partial_x$  for  $\frac{\partial}{\partial x}$ . In Lemma 13.22 we will show that  $f(\partial_x)$  maps  $\mathbf{P}_d$  to itself. This requires the consideration of exponential functions. If we restrict ourselves to polynomials then

**Lemma 10.62.** Suppose that f is a product of linear terms

$$f = \prod_{i} (b_i + a_{1i}x_1 + \cdots + a_{di}x_d) = \prod_{i} (b_i + a_i \cdot x)$$

where all the coefficients  $a_{ji}$  are positive and  $\mathbf{a}_i = (a_{1i}, \dots, a_{di})$ . The map  $f(\frac{\partial}{\partial x})$  is a linear transformation from  $\mathbf{P}_d$  to itself.

*Proof.* It suffices to assume that f has a single factor. If g is in  $\mathbf{P}_d$ , then any derivative of g has total degree less than g. If the constant term of f is non-zero, then  $(f(\frac{\partial}{\partial x})g)^H$  is a constant multiple of  $g^H$ . If the constant term is zero then  $(f(\frac{\partial}{\partial x})g)^H$  is a positive linear combination of the partial derivatives of  $g^H$  which implies that all of its coefficients are positive. In either case  $(f(\frac{\partial}{\partial x})g)^H$  satisfies the positivity condition.

Substitution is also satisfied since since all the partial derivatives interlace g, and and are combined with the correct signs.

The next two lemmas are generalizations of Corollary 9.93 and Lemma 9.95; the proof are the same.

**Lemma 10.63.** If  $g \in P^{pos}$ ,  $f(\mathbf{x}) \in \mathbf{P}_d$  then  $g(-\partial x_1 \partial x_2) f(\mathbf{x}) \in \mathbf{P}_d$ .

**Lemma 10.64.** If  $g \in P^{\text{pos}}$ ,  $f(\mathbf{x}) \in \mathbf{P}_d$  then  $g(x_1 \frac{\partial}{\partial x_2}) f(\mathbf{x}) \in \mathbf{P}_d$ .

**Lemma 10.65.** Let  $\mathbf{c}^{i} = (\gamma_{1}^{i}, \dots, \gamma_{d}^{i})$  be vectors of all positive terms, and  $\mathbf{b}^{i} = (\beta_{1}^{i}, \dots, \beta_{d}^{i})$  vectors of all non-negative terms. The operator

$$\prod_{i} \left( \mathbf{c}^{i} \cdot \mathbf{x} - \mathbf{b}^{i} \cdot \left(\frac{\partial}{\partial \mathbf{x}}\right) \right)$$

maps  $\mathbf{P}_{d}$  to itself.

*Proof.* It suffices to verify it for one factor. Choose  $g \in \mathbf{P}_d$  and set

$$\mathbf{h} = \left( \mathbf{c} \cdot \mathbf{x} - \mathbf{b} \cdot \frac{\partial}{\partial \mathbf{x}} \right) \, \mathbf{g}.$$

Since  $h^H = (\mathbf{c} \cdot \mathbf{x}) g^H$  we see that h satisfies the homogeneity condition. Set  $k = (\mathbf{b} \cdot \frac{\partial}{\partial x}) g$ . Since derivatives interlace and all coefficients of  $\mathbf{b}$  are non-negative, we find that  $g \leq k$ . To verify substitution we must show that  $\mathbf{c} \cdot (\mathbf{x}_i^a) g(\mathbf{x}_i^a) - k(\mathbf{x}_i^a)$  is in  $\mathbf{P}$  for any  $\mathbf{a} \in \mathbb{R}^d$ . This follows from Lemma 1.20.

**Corollary 10.66.** Let  $\mathbf{c}^{i} = (\gamma_{1}^{i}, \dots, \gamma_{d}^{i})$  be vectors of all positive terms, and  $\mathbf{b}^{i} = (\beta_{1}^{i}, \dots, \beta_{d}^{i})$  vectors of all non-negative terms. The map

$$f \times g \mapsto f(\mathbf{c} \cdot \mathbf{x} - \mathbf{b} \cdot \frac{\partial}{\partial \mathbf{x}}) g(\mathbf{x})$$

*defines a map*  $\mathbf{P} \times \mathbf{P}_{d} \longrightarrow \mathbf{P}_{d}$ .

These last results were generalizations to d variables of the corresponding one variable results. The next result is strictly a property of more than one variable.

**Theorem 10.67.** If  $f(\mathbf{x}, \mathbf{y}) = \sum \alpha_I(\mathbf{x})\mathbf{y}^I$  has the property that  $f(\frac{\partial}{\partial \mathbf{x}}, \frac{\partial}{\partial \mathbf{y}})$  maps  $\mathbf{P}_{d+e}$  to itself, then all coefficients  $\alpha_I(\mathbf{x})$  determine operators  $\alpha_I(\frac{\partial}{\partial \mathbf{x}})$  that map  $\mathbf{P}_d$  to itself.

*Proof.* Choose  $g \in \mathbf{P}_d$ , a large integer N, and let  $\mathbf{y}^N = y_1^N \cdots y_e^N$ . The action of  $f(\frac{\partial}{\partial \mathbf{x}}, \frac{\partial}{\partial \mathbf{y}})$  on  $g(\mathbf{x})\mathbf{y}^N$  is

$$\begin{split} \mathsf{f}(\frac{\partial}{\partial x},\frac{\partial}{\partial y}) \mathsf{g}(x) y^{\mathsf{N}} &= \left(\sum \, \mathfrak{a}_{I}(\frac{\partial}{\partial x}) \, (\frac{\partial}{\partial y})^{I} \right) \mathsf{g}(x) y^{\mathsf{N}} \\ &= \sum \, \mathfrak{a}_{I}(\frac{\partial}{\partial x}) \mathsf{g}(x) \, \left(\frac{\partial^{I}}{\partial y^{I}} y^{\mathsf{N}}\right) \end{split}$$

Since N is large, all the y terms  $\frac{\partial^{1}}{\partial x^{1}}y^{N}$  are non-zero and distinct. Since the left hand side is in  $P_{d+e}$ , all its coefficients are in  $P_{d}$ , and hence  $a_{I}(\partial_{x})g(x)$  is in  $P_{d}$ . Thus  $a_{I}(\partial_{x}) \text{ maps } P_{d}$  to itself.

If we choose f to be a product of linear terms, then the coefficients are generally not products. This shows that there are polynomials h that are not products such that  $h(\frac{\partial}{\partial x})$  maps  $P_d$  to itself.

We now generalize Lemma 9.97, and find a Hadamard product.

Lemma 10.68. If 
$$\sum f_i(\mathbf{x})y^i \in P_{d+1}$$
 and  $\sum a_i x^i \in P^{pos}$  then  

$$\sum i! a_i f_i(\mathbf{x})y^i \in P_{d+1}$$
(10.11.1)

Proof. From Lemma 10.64 we know that

$$\sum \mathfrak{a}_{\mathfrak{i}}(\mathfrak{y}\mathfrak{d}_z)^{\mathfrak{i}}\sum \mathfrak{f}_{\mathfrak{j}}(\mathbf{x})z^{\mathfrak{j}}\in \overline{\mathbf{P}}_{d+2}$$

and the coefficient of  $z^0$  yields (10.11.1).

In the following lemma we would like to replace  $\exp_x f^{re\nu}$  with f, where we define  $\exp_x = \exp_{x_1} \cdots \exp_{x_d}$ .

**Lemma 10.69.** Suppose that  $f, g \in \mathbf{P}_d(n)$  and write  $f = \sum_{|I| \leq n} a_I \mathbf{x}^I$ . If  $f^{rev} = \sum a_I \mathbf{x}^{n-I}$  then

$$\left(\exp_{\mathbf{x}} f^{rev}\right)\left(\frac{\partial}{\partial \mathbf{x}}\right) g \in \mathbf{P}_{d}$$

*Proof.* The coefficient of  $y_1^n \cdots y_d^n$  in the product of f(y) and the Taylor series of g(x + y)

$$\left(\sum \mathfrak{a}_{I}\,\boldsymbol{y}^{I}\right)\left(\sum \mathfrak{g}^{(I)}(\boldsymbol{x})\frac{\boldsymbol{y}^{I}}{I!}\right)$$

equals

$$\sum \frac{a_{\mathbf{I}}}{(n-\mathbf{I})!} g^{(n-\mathbf{I})}(\mathbf{x}) = \left(\sum a_{\mathbf{I}} \frac{\mathbf{x}^{n-\mathbf{I}}}{(n-\mathbf{I})!}\right) \left(\frac{\partial}{\partial \mathbf{x}}\right) g$$

and the latter expression is exactly  $\exp_{x_1} \cdots \exp_{x_d} f^{re\nu}(\frac{\partial}{\partial x}) g$ .

Next we generalize Corollary 7.19 and (7.4.2).

**Lemma 10.70.** *If*  $g \in P$  *has no roots in* [0, n] *and*  $f \in P_d(n)$  *then* 

$$g(x_1 \frac{\partial}{\partial x_1}) f \in \mathbf{P}_d$$

*Proof.* It suffices to take  $g = -a + x_1$  where  $a \notin [0, n]$ . We need to show that  $h = -af + x_1 \frac{\partial f}{\partial x_1} \in \mathbf{P}_d$ . We first observe that Corollary 1.15 implies that h satisfies  $x_1$ -substitution. It remains to check the signs of the coefficients of  $h^H$ . If  $\mathbf{I} = (i_1, \ldots, i_d)$  and  $f = \sum a_i \mathbf{x}^I$  then the coefficient of  $\mathbf{x}^I$  in h is  $(i_1 - a)a_I$ . Since  $0 \leqslant i_1 \leqslant n$  it follows that if a < 0 the signs of all the coefficients of  $h^H$  are positive, and if a > n the signs are all negative. Thus, in either case  $h \in \pm \mathbf{P}_d$ .

**Corollary 10.71.** Suppose that  $f \in \mathbf{P}_d(n)$ , and  $g_1, \ldots, g_d$  have the property that none of them have a root in [0, n]. If  $f = \sum \alpha_I \mathbf{x}^I$  then

$$\sum_{I=(\mathfrak{i}_1,\ldots,\mathfrak{i}_d)} g_1(\mathfrak{i}_1)\cdots g_d(\mathfrak{i}_d) \, \mathfrak{a}_I \mathbf{x}^I \in \mathbf{P}_d \tag{10.11.2}$$

In particular, if  $f \in \mathbf{P}_d$  and  $g_1, \dots g_d \in \mathbf{P}^{pos}$  then (10.11.2) holds.

Proof. Use Lemma 10.70, equation (7.4.2), and the calculation

$$g(\mathbf{x}_1 \frac{\partial}{\partial \mathbf{x}_1}) = \sum_{\mathbf{I}} g(\mathbf{x}_1 \frac{\partial}{\partial \mathbf{x}_1}) a_{\mathbf{I}} \mathbf{x}^{\mathbf{I}}$$
$$= \sum_{\mathbf{I} = (i_1, \dots, i_d)} g(i_1) a_{\mathbf{I}} \mathbf{x}^{\mathbf{I}}$$

For example, if we take  $f = (x+y+1)^n$ ,  $g_1(x) = (x+1)^r$  and  $g_2(y) = (y+1)^s$  then

$$\sum_{0\leqslant i+j\leqslant n}(i+1)^r(j+1)^s\binom{n}{i,j}x^iy^j\ \in\ \textbf{P}_2$$

Here is a partial converse to Lemma 10.62.

**Proposition 10.72.** *If*  $f(\mathbf{x})$  *is a polynomial with the property that*  $f(\frac{\partial}{\partial \mathbf{x}})$  *maps*  $\mathbf{Sub}_d$  *to itself, then* f *is in*  $\mathbf{Sub}_d$ .

*Proof.* Let  $f = \sum a_I x^I$  have  $x_i$  degree  $k_i$ . Consider the action of  $f(\frac{\partial}{\partial x})$  on the polynomial  $g = x_1^{k_1 + N} \cdots x_d^{k_d + N}$  where N is large:

$$\begin{split} f(\frac{\partial}{\partial \mathbf{x}})g &= \sum \alpha_{I} \left(\frac{\partial}{\partial \mathbf{x}^{I}}\right)g \\ &= \sum \alpha_{I} (\underline{k_{1}+N})_{i_{1}} \cdots (\underline{k_{d}+N})_{i_{d}} \mathbf{x}_{1}^{k_{1}+N-i_{1}} \cdots \mathbf{x}_{d}^{k_{d}+N-i_{d}} \end{split}$$

where  $I = (i_1, ..., i_d)$ . If  $f(\frac{\partial}{\partial x})g$  is in  $Sub_d$  then so is its reversal(Proposition 10.5)

$$\sum a_{I}(\underline{k_{1}+N})_{i_{1}}\cdots(\underline{k_{d}+N})_{i_{d}}x_{1}^{i_{1}}\cdots x_{d}^{i_{d}}.$$

If we replace  $x_i$  by  $x_i/N$  we get that

$$\sum \mathfrak{a}_{I} \frac{(\underline{k_{1}+N})_{\mathfrak{i}_{1}}}{N^{\mathfrak{i}_{1}}} \cdots \frac{(\underline{k_{d}+N})_{\mathfrak{i}_{d}}}{N^{\mathfrak{i}_{d}}} x_{1}^{\mathfrak{i}_{1}} \cdots x_{d}^{\mathfrak{i}_{d}}.$$

is also in **Sub**<sub>d</sub>. Since this converges to f we conclude that  $f \in$ **Sub**<sub>d</sub>.

## 10.12 Higher order Newton inequalities

We show how the Newton inequalities (4.3.1) can be interpreted as statements about interlacing. This perspective leads to the Newton interlacing proposition.

If  $g(x) = b_0 + b_1 x + b_2 x^2 + \dots \in \mathbf{P}^{p \circ s}$  then Newton's inequalities state that  $b_k^2 \ge \frac{k+1}{k} b_{k-1} b_{k+1}$ . (If we know the degree of g we can be more precise.) We rewrite this as the sequence of inequalities

$$\frac{1 \cdot b_1}{b_0} \geqslant \frac{2 \cdot b_2}{b_1} \geqslant \frac{3 \cdot b_3}{b_2} \geqslant \cdots$$

Each fraction is the negative of the root of a linear polynomial, so we can reinterpret these inequalities as statements about the interlacing of a sequence of polynomials.

 $b_0 + 1 \cdot b_1 x \underline{\ll} b_1 + 2 \cdot b_2 x \underline{\ll} b_2 + 3 \cdot b_3 x \underline{\ll} \cdots$ 

These polynomials correspond to the double lines in the diagram below, where we write the coefficients of g' and g:



Now in this diagram g and g' can be considered to be the first two rows of a polynomial in  $P_2$ , as in Figure 10.1. If we simply take quadratic polynomials instead of linear polynomials, then we find that the resulting polynomials don't always have all real roots. The solution is to introduce binomial coefficients.

**Proposition 10.73.** Suppose that  $f \in \mathbf{P}_2^{\text{pos}}(n)$ . Choose a positive integer m. If  $f = \sum a_{i,j} x^i y^j$  then define the polynomials  $\mathfrak{F}_k$  for  $0 \leq k \leq n - m$  by

$$\mathfrak{F}_{k}(x) = \sum_{j=0}^{m} \mathfrak{a}_{k,j} \begin{pmatrix} k \\ j \end{pmatrix} x^{j}$$

All the  $\mathfrak{F}_k$  have all real roots and interlace:

$$\mathfrak{F}_0 \underline{\ll} \mathfrak{F}_1 \underline{\ll} \mathfrak{F}_2 \underline{\ll} \cdots \underline{\ll} \mathfrak{F}_{n-\mathfrak{m}}$$

Consider an example where m = 2 and n = 4. There are three polynomials, which are identified by the double lines in Figure 10.1.

$$\begin{split} \mathfrak{F}_0 &= a_{00} + 2a_{01}x + a_{02}x^2 \\ \mathfrak{F}_1 &= a_{10} + 2a_{11}x + a_{12}x^2 \\ \mathfrak{F}_2 &= a_{20} + 2a_{21}x + a_{22}x^2 \end{split}$$

The proposition asserts that  $\mathfrak{F}_0 \leq \mathfrak{F}_1 \leq \mathfrak{F}_2$ .

*Proof.* (of the proposition) If we write

$$f(x, y) = f_0(y) + f_1(y)x + f_2(y)x^2 + \cdots$$

then

$$\mathfrak{F}_{\mathbf{k}} = (1+\mathbf{y})^{\mathbf{m}} * \mathsf{f}_{\mathbf{k}}(\mathbf{y})$$

By Lemma 10.68 we know that  $(1 + y)^m * f(x, y) \in \overline{\mathbf{P}}_2$ . The conclusion of the proposition follows since consecutive coefficients of polynomials in  $P_2$  interlace.

#### 10.13 Negative subdefinite matrices

When does a quadratic form  $\mathbf{x}Q\mathbf{x}^t$  have the property that  $\mathbf{x}Q\mathbf{x}^t - c^2 \in \mathbf{P}_d$  for all c? The answer is simple: Q is negative subdefinite.

**Definition 10.74.** A real symmetric matrix A is *negative subdefinite* if it has exactly one positive eigenvalue, and all entries are positive.

Our goal is the following theorem, which will follow from Lemmas 10.86 and 10.87 in the next section.

**Theorem 10.75.** Suppose Q is a real symmetric matrix. The quadratic form  $\mathbf{x}Q\mathbf{x}^{t} - \mathbf{c}^{2}$  is in  $\mathbf{P}_{d}$  for all c if and only if Q is negative subdefinite.

In this section we establish various properties of negative subdefinite matrices. We should first observe that among all real symmetric matrices with all positive entries, the negative subdefinite matrices are those with the smallest possible number of positive eigenvalues. They can't all be negative, since Perron's theorem guarantees at least one positive eigenvalue.

We first note that subdefinite matrices are hereditary in the following sense:

**Lemma 10.76.** *If* A *is a negative subdefinite matrix then all the principle submatrices of* A *are also negative subdefinite.* 

*Proof.* It suffices to show that the matrix T formed by removing the last row and column is negative subdefinite. The eigenvalues of T interlace the eigenvalues of A by Theorem 1.61. Thus, since A has exactly n - 1 non-positive eigenvalues, T has at least n - 2 non-positive eigenvalues. Now the sum of the eigenvalues of T is the trace of T which is positive, so T must have at least one positive eigenvalue. Thus T has exactly one positive eigenvalue.



Figure 10.1: A polynomial in  $P_2(4)$ 

Here is an alternative characterization of negative subdefinite matrices. Recall that a symmetric matrix is positive definite if and only if the determinants of the principle submatrices are all positive. The k-th leading principle submatrix is the submatrix formed from the elements in the first k rows and columns.

**Lemma 10.77.** Suppose that the n by n matrix A has all positive entries, and all principle submatrices are invertible. Then, A is negative subdefinite if and only if the determinant of the k-th leading principle submatrix has sign  $(-1)^{k+1}$ , for  $1 \le k \le n$ .

*Proof.* If A is negative subdefinite, then the k-th principle submatrix is also negative subdefinite, and so it has k-1 negative eigenvalues, and one positive eigenvalue. Since the determinant is the product of the eigenvalues, the result follows.

Conversely, we prove by induction that the k-th leading principle submatrix is negative subdefinite. It is trivial for k = 1, so assume that the k-th principle submatrix is negative subdefinite. Since the eigenvalues of the (k + 1)-st leading principle submatrix P are interlaced by the eigenvalues of the k-th, it follows that P has at least k - 1 negative eigenvalues. Now, by hypothesis the determinant of P has sign  $(-1)^k$ , and so P must have exactly k negative eigenvalues, and therefore one positive one. Thus, P is negative subdefinite.

Note that this argument also shows that if Q is negative subdefinite and |Q| is non-zero, then all leading principle submatrices have non-zero determinant. Here is a useful criterion for extending negative subdefinite matrices.

**Lemma 10.78.** Suppose that  $Q = \begin{pmatrix} A & u \\ u^t & c \end{pmatrix}$  where all entries are positive, u is 1 by n, c is a scalar,  $|Q| \neq 0$ , and  $|A| \neq 0$ . Then, Q is negative subdefinite if and only if these two conditions are met:

- 1. A is negative subdefinite.
- 2.  $uA^{-1}u^{t} c > 0$

*Proof.* We use the Schur complement formula [85, page 22]:

$$\mathbf{Q}| = |\mathbf{A}| |\mathbf{c} - \mathbf{u}\mathbf{A}^{-1}\mathbf{u}^{\mathsf{t}}|$$

By hypothesis  $|c - uA^{-1}u^t|$  is negative, so |Q| and |A| have opposite signs, and the conclusion now follows from Lemmas 10.77 and 10.76.

**Example 10.79.** It is not difficult to find negative subdefinite matrices. Consider the matrix<sup>2</sup>  $aJ_n + bI_n$  where  $J_n$  is the n by n matrix of all 1's, and  $I_n$  is the identity matrix. For example,  $J_4 - I_4$  is

/0	1	1	1
1	0	1	1
1	1	0	1
$\backslash 1$	1	1	0/

<sup>&</sup>lt;sup>2</sup>In the statistics literature this is known, after scaling, as the equicorrelation matrix.

The eigenvalues of  $aJ_n + bI_n$  are na + b and n - 1 b's. If we choose b < 0, a > 0 and a + b > 0 then  $aJ_n + bI_n$  has exactly one positive eigenvalue and all positive entries. The matrices  $J_n - I_n$  are limits of negative subdefinite matrices.

We can use Lemma 10.78 to construct negative subdefinite matrices. For example, the following matrix

$$\begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & 2 & 3 & 4 & 5 \\ 3 & 3 & 3 & 4 & 5 \\ 4 & 4 & 4 & 4 & 5 \\ 5 & 5 & 5 & 5 & 5 \end{pmatrix}$$

is negative subdefinite since all entries are positive, and the determinants of the leading principle submatrices are 1, -2, 3, -4, 5.

We can embed negative subdefinite matrices in higher dimensional matrices.

**Lemma 10.80.** If Q is a d by d negative subdefinite matrix, then  $\begin{pmatrix} Q & 0 \\ 0 & 0 \end{pmatrix}$  is the limit of d + 1 by d + 1 negative subdefinite matrices.

*Proof.* Write  $Q = \begin{pmatrix} A & v \\ v^{t} & c \end{pmatrix}$  where A is d - 1 by d - 1. Consider the matrix

$$\mathsf{M} = \begin{pmatrix} \mathsf{A} & \mathsf{v} & \alpha \mathsf{v} \\ \mathsf{v}^{\mathsf{t}} & \mathsf{c} & \alpha \mathsf{c} \\ \alpha \mathsf{v}^{\mathsf{t}} & \alpha \mathsf{c} & e \end{pmatrix}$$

If we subtract  $\alpha$  times the dth row from the bottom row we see that

$$\begin{vmatrix} A & v & \alpha v \\ v^{t} & c & \alpha c \\ \alpha v^{t} & \alpha c & e \end{vmatrix} = \begin{vmatrix} A & v & \alpha v \\ v^{t} & c & \alpha c \\ 0 & 0 & e - \alpha^{2}c \end{vmatrix} = (e - \alpha^{2}c)|Q|$$

We choose positive  $\alpha$ , *e* so that  $e - \alpha^2 c < 0$ . Now M has all positive entries, and its eigenvalues are interlaced by the eigenvalues of Q. Thus, M has at least one positive eigenvalue, and at least d - 1 negative eigenvalues. Since the determinant of Q is the opposite sign from the determinant of M, it follows that M and Q have a different number of negative eigenvalues, so M had d negative eigenvalues, one positive one, and so is a negative subdefinite matrix. Taking the limit as *e* and  $\alpha$  go to zero gives the conclusion.

What do cones of negative subdefinite matrices look like? Equivalently, when are all positive linear combinations of a collection of negative subdefinite matrices also negative subdefinite? Here is an answer for two matrices.

#### **Lemma 10.81.** Suppose that

1.  $Q_1$  is negative subdefinite and invertible.

2. Q<sub>2</sub> has all non-negative entries.

3.  $Q_1^{-1}Q_2$  has no negative eigenvalues.

then

- 1.  $aQ_1 + bQ_2$  is negative subdefinite for all positive a, b.
- 2.  $Q_2$  is the limit of negative subdefinite matrices.

*Proof.* We claim that if a, b are positive then  $aQ_1 + bQ_2$  is never singular. If it were then there is a  $\nu$  such that

$$aQ_1\nu + bQ_2\nu = 0$$

which implies that

$$Q_1^{-1}Q_2\nu=-\frac{a}{b}\nu$$

But  $Q_1^{-1}Q_2$  has no negative eigenvalue, so this isn't possible. Thus, as a, b vary over positive reals the eigenvalues never change sign. Since a = 1 and b = 0 yields  $Q_1$  which has exactly one positive eigenvalue and no zero eigenvalues, all positive linear combinations  $aQ_1 + bQ_2$  have exactly one positive eigenvalue, and are invertible. Since all entries of  $Q_1$  are positive, and of  $Q_2$  are non-negative, the entries of  $aQ_1 + bQ_2$  are all positive. This establishes (1). The second part follows by taking  $a \to 0$  and b = 1.

**Corollary 10.82.** Suppose  $Q_1, Q_2, \ldots, Q_n$  are negative subdefinite invertible matrices.

- 1.  $aQ_1+bQ_2$  is negative subdefinite for all positive a, b iff  $Q_1^{-1}Q_2$  has no negative eigenvalues.
- 2. If  $Q_1, \ldots, Q_n$  lie in a cone of negative subdefinite matrices then  $Q_i^{-1}Q_j$  has no negative eigenvalues for  $1 \le i, j \le n$ .

*Remark* 10.83. If Q is an invertible negative subdefinite matrix, then the eigenvalues of  $Q^n$  are the nth power of the eigenvalues of Q. Thus, if n is an integer then  $Q^{2n+1}$  is also negative subdefinite. Moreover, if m is an integer then the eigenvalues of  $(Q^{2m+1})^{-1}(Q^{2n+1}) = Q^{2n-2m}$  are all positive. Thus,

$$aQ^{2n+1} + bQ^{2m+1}$$

is negative subdefinite.

Here is an example of a cone in dimension two. Define

$$Q_{a,b} = \begin{pmatrix} a & b \\ b & a \end{pmatrix}$$

As long as 0 < a < b,  $Q_{a,b}$  is negative subdefinite. Since 0 < a' < b' implies 0 < a + ta' < b + tb' for any positive t it follows that any positive linear combination of these matrices is still negative subdefinite.

The following property will be used in the next section. Recall that negative *semi*definite means that there are no positive eigenvalues.

**Theorem 10.84** ([133]). *Suppose that* A *is a real symmetric matrix that is not negative semidefinite. The following are equivalent:* 

- 1. A is negative subdefinite.
- 2. For every vector  $\mathbf{x}$ ,  $\mathbf{x}A\mathbf{x}^t > 0$  implies that  $A\mathbf{x}$  is a non-positive or non-negative vector.

The following lemma is used in the next section.

**Lemma 10.85.** If  $Q = \begin{pmatrix} a & v \\ v^t & C \end{pmatrix}$  is negative subdefinite, then  $v^t v - aC$  is positive semidefinite.

*Proof.* Assume that  $v^t v - aC$  is not positive semidefinite. Then, there is a *z* such that  $z(v^t v - aC)z^t < 0$ . This implies that  $zCz^t > \frac{1}{a}zv^tvz^t$ . Evaluate the quadratic form at  $w = (\alpha, z)$  where  $\alpha$  is to be determined.

$$(\alpha, z) \begin{pmatrix} a & v \\ v^{t} & C \end{pmatrix} \begin{pmatrix} \alpha \\ z^{t} \end{pmatrix} = \alpha^{2} a + \alpha (zv^{t} + vz^{t}) + zCz^{t}$$
$$> \alpha^{2} a + \alpha (zv^{t} + vz^{t}) + \frac{1}{a} zv^{t} vz^{t}$$
$$= \frac{1}{a} (\alpha a + zv^{t})^{2}$$

If  $\alpha a + zv^t \neq 0$  then  $wQw^t > 0$ , so by Theorem 10.84  $Qw^t$  has either all non-negative, or all non-positive, coefficients. But

$$Qw^{t} = (a\alpha + vz^{t}, \alpha v^{t} + Cz^{t}).$$

Since  $\alpha a + vz^t$  can be either positive or negative, as  $\alpha a + vz^t$  becomes zero, so must  $\alpha v^t + Cz^t$ . Thus, when  $\alpha = -\frac{1}{a}vz^t$  we have

$$(-\frac{1}{a}\nu z^{t})\nu^{t} + Cz^{t} = -\frac{1}{a}(\nu^{t}\nu - aC)z^{t} = 0$$

This contradicts our hypothesis that  $z(v^tv - aC)z^t < 0$ , and so the lemma is proved.

### 10.14 Quadratic Forms

We have seen substitution of some quadratic forms before. For instance, if  $f \in \mathbf{P}^{pos}$  then  $f(-x^2) \in \mathbf{P}$ . Also, if  $f \in \mathbf{P}^{pos}$ , then  $f(-xy) \in \overline{\mathbf{P}}_2$ . We will replace  $-x^2$  and -xy by quadratic forms determined by negative subdefinite matrices. We begin with substitution into the simplest polynomial:  $x + c^2$ :

When is  $-\mathbf{x}Q\mathbf{x}^{t} + c^{2}$  in  $\pm P_{2}$ ?

Let's first consider the case of a quadratic form in two variables. Take a matrix  $A = \begin{pmatrix} a_1 & a_2 \\ a_2 & a_3 \end{pmatrix}$  with corresponding quadratic form  $a_1x^2 + 2a_2xy + a_3y^2$ . We are interested when the polynomial

$$\begin{pmatrix} x & y \end{pmatrix} \begin{pmatrix} a_1 & a_2 \\ a_2 & a_3 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} - c^2 = a_1 x^2 + 2a_2 xy + a_3 y^2 - c^2$$

is in  $P_2$  for all c. We must have that all  $a_i$  are positive, and the discriminant should be non-negative. Thus

$$4a_2^2 - 4a_1a_3 + 4c^2 = -|a_1 a_2 a_3| + 4c^2 \ge 0$$

and hence  $|A| \leq 0$ . A has a positive eigenvalue, since the sum of the eigenvalues is the trace which is positive. Consequently,  $\mathbf{x}A\mathbf{x}^t - c^2 \in \mathbf{P}_2$  iff A is negative subdefinite.

**Lemma 10.86.** If Q is a d by d symmetric matrix such that  $xQx^t - c^2 \in \pm P_d$  for all c then

- 1. Q has all positive entries.
- 2. Q has exactly one positive eigenvalue.
- 3. *Q* is negative subdefinite.

*Proof.* Since  $\mathbf{x}Q\mathbf{x}^t - c^2 \in \pm \mathbf{P}_d$  the homogeneous part is in  $\mathbf{P}_d$ , and so the entries of  $\mathbf{x}Q\mathbf{x}^t$  are all the same sign, and are non-zero. If they are negative then letting  $Q = (a_{ij})$  and  $\mathbf{x} = (x, 0, ..., 0)$  yields  $\mathbf{x}Q\mathbf{x}^t - c^2 = a_{11}\mathbf{x}^2 - c^2$ . This has all real roots iff  $a_{11}$  is positive, so all entries of Q are positive.

For the second part we only need to assume that  $\mathbf{x}Q\mathbf{x}^t \in \overline{\mathbf{P}}_d$ . Recall Perron's theorem [85] that says that since all entries of Q are positive there is a unique largest positive eigenvalue  $\lambda$ , and the corresponding eigenvector  $v = (v_1, \ldots, v_d)$  has all positive entries. Suppose that  $\mu$  is an eigenvalue with eigenvector  $\mathbf{u} = (u_1, \ldots, u_d)$ . Since Q is symmetric and  $\lambda \neq \mu$  it follows that  $\mathbf{u}^t Q \mathbf{v} = 0$ . If we set  $\mathbf{f}(\mathbf{x}) = \mathbf{x}Q\mathbf{x}^t$  then we can replace each  $x_i$  by  $v_i \mathbf{x} + u_i$  and the resulting polynomial is in  $\mathbf{P}$  since all  $v_i$  are positive. Consequently,

$$f(xv + u) = (xv + u)Q(xv + u)^{t} = \lambda x^{2}|v|^{2} + \mu|u|^{2} \in \mathbf{P}$$

The only way that this can happen is if  $\mu$  is negative or 0, so there is exactly one positive eigenvalue.

**Lemma 10.87.** If Q is negative subdefinite then  $\mathbf{x}Q\mathbf{x}^t - \mathbf{c}^2 \in \mathbf{P}_d$  for all c.

*Proof.* The homogeneous part of  $\mathbf{x}Q\mathbf{x}^t - c^2$  is a polynomial with all positive coefficients since Q has all positive coefficients. Thus, we only have to check substitution.
Write  $Q = \begin{pmatrix} a & v \\ v^t & C \end{pmatrix}$  where v is  $1 \times (d-1)$  and C is  $(d-1) \times (d-1)$ , and let  $\mathbf{x} = (x_1, \mathbf{z})$  where  $\mathbf{z} = (x_2, \dots, x_d)$ . With this notation

$$\mathbf{x}\mathbf{Q}\mathbf{x}^{t} = (\mathbf{x}_{1}, \mathbf{z}) \begin{pmatrix} \mathbf{a} & \mathbf{v} \\ \mathbf{v}^{t} & \mathbf{C} \end{pmatrix} \begin{pmatrix} \mathbf{x} \\ \mathbf{z}^{t} \end{pmatrix}$$
$$= \mathbf{a}\mathbf{x}_{1}^{2} + \mathbf{x}_{1}(\mathbf{v}\mathbf{z}^{t} + \mathbf{z}\mathbf{v}^{t}) + \mathbf{z}\mathbf{C}\mathbf{z}^{t}$$

 $\mathbf{x}Q\mathbf{x}^t$  satisfies  $x_1$  substitution if and only if the discriminant is nonnegative. Now  $v\mathbf{z}^t = \mathbf{z}v^t$  since they are both scalars, so the discriminant condition is that

$$0 \leq 4(\mathbf{v}\mathbf{z}^{t})^{2} - 4\mathbf{a}\mathbf{z}C\mathbf{z}^{t} = 4\mathbf{z}(\mathbf{v}^{t}\mathbf{v} - \mathbf{a}C)\mathbf{z}^{t}$$

 $v^t v - aC$  is a symmetric matrix, so  $xQx^t$  satisfies  $x_1$ -substitution iff  $v^t v - aC$  is positive semidefinite. Now apply Lemma 10.85.

#### 10.15 The analog of x**D**

The differential operator  $\mathbf{x} \cdot \mathbf{\partial} = x_1 \frac{\partial}{\partial x_1} + \dots + x_d \frac{\partial}{\partial x_d}$  is the analog in  $\mathbf{P}_d^{\text{pos}}$  of the operator  $x \frac{d}{dx}$  on  $\mathbf{P}^{\text{pos}}$ . Here are some elementary properties of this operator.

- 1. If  $f(\mathbf{x})$  is homogeneous of degree n then  $(x_1 \frac{\partial}{\partial x_1} + \dots + x_d \frac{\partial}{\partial x_d})f = nf$ . This is the well-known Euler identity about homogeneous functions.
- 2. If  $f \in P_d(n)$  then  $\left( (x_1 \frac{\partial}{\partial x_1} + \dots + x_d \frac{\partial}{\partial x_d}) f \right)^H = n f^H$ . This is an immediate consequence of the previous property.
- 3. If  $g(\mathbf{x}) = \sum a_{\mathbf{I}} \mathbf{x}^{\mathbf{I}}$  then

$$f(x_1\frac{\partial}{\partial x_1} + \dots + x_d\frac{\partial}{\partial x_d})g = \sum a_I f(|I|)x^I$$

This is a simple consequence of linearity and the calculation

$$(\mathbf{x} \cdot \mathbf{\partial})\mathbf{x}^{\mathbf{I}} = (x_1 \frac{\partial}{\partial x_1} + \dots + x_d \frac{\partial}{\partial x_d})(x_1^{i_1} \cdots x_d^{i_d}) = (i_1 + \dots + i_d)x_1^{i_1} \cdots x_d^{i_d}$$

The following is the basic fact about  $\mathbf{x} \cdot \mathbf{\partial}$ .

**Proposition 10.88.** If  $f \in \mathbf{P}_d^{\text{pos}}(n)$  then  $(x_1 \frac{\partial}{\partial x_1} + \dots + x_d \frac{\partial}{\partial x_d}) f \in \mathbf{P}_d^{\text{pos}}$ . Moreover, if g is the polar derivative of f then

$$nf = (x_1 \frac{\partial}{\partial x_1} + \dots + x_d \frac{\partial}{\partial x_d})f + g.$$

We also have interlacings

$$f \underline{\ll} (x_1 \frac{\partial}{\partial x_1} + \dots + x_d \frac{\partial}{\partial x_d}) f \underline{\lessdot} g$$

*Proof.* These are immediate consequences of the properties of the polar derivative; see (10.4.1) and (10.4.2).

In the case that d is two we can derive them in a different way. Notice that

$$\frac{\partial}{\partial z} f(-xz, -yz) = -(xf_x(-xz, -yz) + yf_y(-xz, -yz))$$

Since the derivative is in  $\overline{\mathbf{P}}_3^{\text{pos}}$ , if we substitute z = -1 we get a polynomial in  $\overline{\mathbf{P}}_2^{\text{pos}}$ , which shows that  $xf_x + yf_y \in \overline{\mathbf{P}}_2^{\text{pos}}$ . Since  $xf_x + yf_y$  satisfies the homogeneity condition, it follows it is in  $\mathbf{P}_2^{\text{pos}}$ .

Finally, since the derivative of f(-xz, -yz) interlaces f(-xz, -yz), and is still true when we substitute z = -1, the interlacing part now follows. It's interesting to see that  $f(x\frac{\partial}{\partial x} + y\frac{\partial}{\partial y})g = f(z\partial_z)g(xz, yz)\Big|_{z=1}$ .

**Corollary 10.89.** If  $f \in P^{pos}$  and  $g = \sum a_I x^I \in P_d^{pos}$  then

$$\sum_{I} \mathbf{a}_{I} \mathbf{f}(|I|) \mathbf{x}^{I} \in \mathbf{P}_{d}^{pos}$$

*Proof.*  $\alpha g + \mathbf{x} \cdot \mathbf{\partial}(g) \in \mathbf{P}_d^{pos}$  by the above proposition, so  $(\mathbf{x} \cdot \mathbf{\partial} + \alpha)(g) \in \mathbf{P}_d^{pos}$  for positive  $\alpha$ . Now factor f, and use induction.

#### 10.16 Generalized Hadamard products

A generalized Hadamard product is a mapping of the form

monomial  $\times$  monomial  $\mapsto$  constant  $\times$  monomial

This generalizes multiplier transformations which have the form

monomial  $\mapsto$  constant  $\times$  monomial

We have seen two Hadamard type products

Theorem 9.84 
$$x^{i} * x^{j} \mapsto \begin{cases} x^{i} & i = j \\ 0 & i \neq j \end{cases}$$
  
Theorem 9.87  $x^{i} *' x^{j} \mapsto \begin{cases} i! x^{i} & i = j \\ 0 & i \neq j \end{cases}$ 

Both of these determine bilinear maps  $\mathbf{P}^{\text{pos}} \times \mathbf{P} \longrightarrow \mathbf{P}$ . In Proposition 15.51 we will determine sufficient conditions on constants  $c_i$  for a product of the form

$$x^{i} \circledast x^{j} \mapsto \begin{cases} c_{i} x^{i} & i = j \\ 0 & i \neq j \end{cases}$$

to determine a map  $P^{pos} \times P \longrightarrow P$ . We now study two products for  $P \times P_2 \longrightarrow P$ . Define

$$y^{i} * x^{j} y^{k} \mapsto \begin{cases} x^{j} & i = k \\ 0 & i \neq j \end{cases}$$
(10.16.1)

$$y^{i} *' x^{j} y^{k} \mapsto \begin{cases} i! x^{j} & i = k\\ 0 & i \neq k \end{cases}$$
(10.16.2)

**Proposition 10.90.** *The linear transformations in* (10.16.1) *and* (10.16.2) *determine mappings*  $P \times P_2 \longrightarrow P$ .

*Proof.* Choose  $f(x,y) = \sum f_i(x)y^i$  in  $P_2$ , and  $g = \sum_0^n a_i y^i$  in **P**. Since  $\sum a_{n-i}y^i \in \mathbf{P}$ , the product

$$\left(\sum f_{\mathfrak{i}}(x)y^{\mathfrak{i}}\right)\left(\sum \mathfrak{a}_{n-\mathfrak{j}}y^{\mathfrak{j}}\right)$$

is in  $\overline{\mathbf{P}}_2$ . The coefficient of  $y^n$  in the product equals  $\sum a_i f_i(x)$  and is in  $\mathbf{P}$ . This is exactly the product determined by (10.16.1).

The second product is a strengthening of the first since we can apply the exponential operator to (10.16.2) to obtain (10.16.1). Note that  $\left(g(\frac{\partial}{\partial y})y^{i}\right)(0) =$  i!a<sub>i</sub>. Since  $g(\frac{\partial}{\partial y})f$  is in P<sub>2</sub> by Lemma 10.62, the conclusion follows from the observation that

$$\left(g(\frac{\partial}{\partial y})f\right)(x,0) = \sum \frac{f_i(x)}{i!} \left(g(\frac{\partial}{\partial y})y^i\right)(0) = \sum f_i(x)a_i.$$

The same argument shows that

Corollary 10.91. If we define

$$\mathbf{y}^{\mathbf{i}} \ast' \mathbf{x}^{J} \mathbf{y}^{\mathbf{k}} \mapsto \begin{cases} \mathbf{i}! \, \mathbf{x}^{J} & \mathbf{i} = \mathbf{k} \\ 0 & \mathbf{i} \neq \mathbf{k} \end{cases}$$

then this determines a map  $P \times P_{d+1} \longrightarrow P_d$ .

**Corollary 10.92.** Suppose  $f(x, y) = \sum f_i(x)y^i$  is in  $P_{d+1}$ . The following two sums are in  $\overline{P}_d$ :

$$\sum (-1)^{\mathfrak{i}} \frac{\mathfrak{f}_{2\mathfrak{i}}(\mathbf{x})}{\mathfrak{i}!} \qquad \sum (-1)^{\mathfrak{i}} \mathfrak{f}_{2\mathfrak{i}}(\mathbf{x})$$

*Proof.* We can take limits to see that Corollary 10.91 determines a map  $\widehat{\mathbf{P}} \times \mathbb{P}_{d+1} \longrightarrow \mathbb{U}_d$ . These results now follow if we consider  $e^{-y^2} * f$  and  $e^{-y^2} *' f$ .  $\Box$ 

The reverse of a polynomial in  $P_d$  was defined in (11.11.1).

**Corollary 10.93.** If  $f = \sum f_i(\mathbf{x})y^i$  and  $g = \sum g_i(\mathbf{x})y^i$  are both in  $P_{d+1}(n)$  then

- 1.  $\sum f_i(\mathbf{x})g_{n-i}(\mathbf{x}) \in P_d$
- 2.  $\sum f_i(\mathbf{x})g_i^{re\nu}(\mathbf{x}) \in \overline{\mathbf{P}}_d$

*Proof.* For the first one we multiply and extract a coefficient as usual. For the second, apply (1) to f and  $g^{rev} = \sum g_i^{rev}(\mathbf{x})y^{n-i}$ .

## **10.17 THE BMV conjecture**

The Bessis-Moussa-Villani (BMV) conjecture, as reformulated by Lieb and Seiringer[114], is

**Conjecture 10.94** (BMV). *If* A, B *are positive definite matrices then*  $trace(I + (A + t B)^n)$  *has all positive coefficients for all positive integers* n.

We offer a stronger conjecture.

**Conjecture 10.95** (s-BMV). If  $A_i$ ,  $B_i$  are positive definite matrices,  $g_i \in P^{pos}$ , and  $f(x, y_1, ..., y_e) \in P_{1+e}$  then

- 1. det  $[f(x I + g_1(A_1 + B_1y) + \dots + g_e(A_e + B_ey))]$  has all positive coefficients.
- 2. If we write the determinant as  $\sum f_i(x)y^i$  then all  $f_i \in \mathbf{P}^{\text{pos}}$ .
- 3. In addition we have  $\cdots \leftarrow f_i \leftarrow f_{i+1} \leftarrow \cdots$ .

If C is a d × d matrix then the trace of C is the coefficient of  $x^{d-1}$  in the polynomial |xI + C|. Replacing x by x + 1 and applying this fact shows that the strong BMV conjecture applied to  $det(xI + I + (A + By)^n)$  implies the original BMV conjecture.

We can take limits and let the  $g_i$  belong to  $\mathbf{P}^{pos}$ . A case of particular interest is the exponential BMV conjecture:

**Conjecture 10.96** (e-BMV). *If* A<sub>i</sub>, B<sub>i</sub> *are positive definite matrices,* 

- 1. det  $[xI + e^{A_1 + B_1y} + \dots + e^{A_e + B_ey}]$  has all positive coefficients.
- 2. If we write the determinant as  $\sum f_i(x)y^i$  then all  $f_i \in \mathbf{P}^{pos}$ .
- 3. In addition  $\cdots \leftarrow f_i \leftarrow f_{i+1} \leftarrow \cdots$ .

It is not the case that  $|xI + (A + By)^n| \in P_2$  or that  $|xI + e^{A+By}| \in \widehat{\mathbf{P}}_2$ . For example, if we take A = B = I then the latter determinant is  $(x + e^{1+y})^d$  which is a polynomial in x, but an exponential polynomial in y.

The leading coefficient of  $|xI + g_1(A_1 + B_1y) + \cdots + g_e(A_e + B_ey)|$  is 1. We have partial information about the constant term.

**Lemma 10.97.** If A, B are positive definite and  $g \in P^{pos}$  then the constant term (with respect to x) of det [xI + g(A + By)] has all positive coefficients.

*Proof.* The constant term is det [g(A + By)]. If we write  $g(x) = \prod (x + a_i)$  where the  $a_i$  are positive then

$$det[g(A + By)] = \prod det[a_iI + A + By]$$

This has all positive coefficients since  $a_i I + A$  and B are positive definite.  $\Box$ 

There is one simple case where s-BMV(1) and e-BMV(1) hold.

**Lemma 10.98.** If  $A_1, B_1, \ldots$  are commuting positive definite matrices then *s*-BMV(1) and *e*-BMV(1) hold.

*Proof.* Let O simultaneously diagonalize all the  $A_i$ 's and  $B_i$ 's, so that  $OA_iO^t = diag(a_{i,j})$  and  $OB_iO^t = diag(b_{i,j})$  where the  $a_{i,j}$ 's and  $b_{i,j}$ 's are positive. Then

$$|\mathbf{x}\mathbf{I} + g_1(\mathbf{A}_1 + \mathbf{B}_1\mathbf{y}) + \dots + g_e(\mathbf{A}_e + \mathbf{B}_e\mathbf{y})| = \prod_j \left(\mathbf{x} + \sum_i g_i(a_{i,j} + b_{i,j}\mathbf{y})\right)$$

All coefficients are positive since  $g \in \mathbf{P}^{pos}$ , and  $a_{i,j}, b_{i,j} > 0$ . The exponential case follows by taking limits.

It is not even clear that s-BMV(2) and s-BMV(3) hold in this simple case. It would follow from the following

**Conjecture 10.99.** Suppose  $g \in P^{pos}$  and  $a_i, b_i$  are positive and n is a positive integer and write

$$\prod_{i=1}^{n} (x + g(a_i + b_i y)) = \sum f_i(x)y^i$$

All  $f_i \in \mathbf{P}^{pos}$  and  $f_i \longleftarrow f_{i+1}$ .

This is a special case of the composition conjecture, which we discuss in the next section.

#### **10.18** The composition conjecture

We investigate some special cases of the composition conjecture

**Conjecture 10.100.** Suppose that  $f(x, y_1, ..., y_e) \in \widehat{P}_{d+e}$ , and choose  $g_i(y) \in \widehat{P^{pos}}$ . If we write

$$F(\mathbf{x}, \mathbf{y}) = f(\mathbf{x}, g_1(\mathbf{y}), \dots, g_e(\mathbf{y})) = \sum f_i(\mathbf{x}) \mathbf{y}^i$$

then

- 1. All  $f_i$  are in  $\overline{\mathbf{P}}_d$ .
- 2.  $f_0 \longleftarrow f_1 \longleftarrow f_2 \longleftarrow \cdots$

*Remark* 10.101. It is not the case that  $F \in \widehat{\mathbf{P}}_{d+1}$ ; take f = x + y and  $g(y) = e^{y}$ . However, if  $\alpha \in \mathbb{R}$  then we clearly have  $F(\mathbf{x}, \alpha) \in \widehat{\mathbf{P}^{pos}}_{d}$ , since we are merely substituting  $g_i(\alpha)$  for  $y_i$ . Thus,  $F(\mathbf{x}, y)$  satisfies substitution for y, not substitution for x, and the coefficients of y interlace.

If  $f \in \widehat{\mathbf{P}^{\text{pos}}}_{d+e}$  and the composition conjecture holds then all coefficients of  $F(\mathbf{x}, \mathbf{y})$  are positive.

We first show if we have one exponential function then the composition conjecture holds. The proof is for d = 1, but an identical argument shows it for any d.

**Lemma 10.102.** Suppose  $f \in P_2$ , and  $g \in \widehat{P^{pos}}$ . If the coefficient  $h_n$  of  $y^n$  in  $e^{-xg(y)}$  is in P then the coefficient  $k_n$  of  $y^n$  in f(x, g(y)) is in P. If  $h_n \longleftarrow h_{n+1}$  then  $k_n \longleftarrow h_{n+1}$ .

*Proof.* Let  $T_g$  be the linear transformation

$$f \mapsto \text{coefficient of } y^n \text{ in } f(x, g(y)).$$

The generating function is

$$G(x, y, u, v) = \sum T_g(x^i y^j) \frac{(-u)^i (-v)^j}{i!j!}$$
  
= 
$$\sum \left[ \text{Coefficient of } y^n \text{ in } x^i g(y)^j \right] \frac{(-u)^i (-v)^j}{i!j!}$$
  
= 
$$e^{-ux} \left[ \text{Coefficient of } y^n \text{ in } e^{-vg(y)} \right].$$

By hypothesis this is in  $\widehat{\mathbf{P}}_3$ , so  $T_g$  maps  $\mathbb{P}_2$  to  $\mathbf{P}$ .

The second part is similar, and considers the map

$$T_{g,\alpha} \mapsto \left[ \text{coefficient of } y^n \text{ in } f(x,g(y)) \right] + \alpha \left[ \text{coefficient of } y^{n+1} \text{ in } f(x,g(y)) \right]$$

**Corollary 10.103.** If  $f \in P_2$  then the coefficient  $k_n$  of  $y^n$  in  $f(x, e^y)$  is in P. Moreover,  $k_n \longleftarrow k_{n+1}$ .

Proof. Recall the identity

$$e^{xe^{y}} = e^{x} \sum_{i=0}^{\infty} B_{i}(x) \frac{y^{i}}{i!}$$

where  $B_i$  is the Bell polynomial. Since the Bell polynomials are in **P** and consecutive ones interlace, the result follows from the preceding lemma.

The following conjecture arises naturally in the lemma following.

**Conjecture 10.104.** *Suppose that*  $\alpha_1, \ldots, \alpha_e$  *are positive and* n *is a positive integer. The mapping* 

$$\mathbf{x}^{I}\mathbf{y}_{1}^{\mathbf{t}_{1}}\cdots\mathbf{y}_{e}^{\mathbf{t}_{e}}\mapsto\mathbf{x}^{I}(\alpha_{1}\mathbf{t}_{1}+\cdots+\alpha_{e}\mathbf{t}_{e})^{n}$$

determines a map  $\mathbf{P}_{d}^{pos} \longrightarrow \mathbf{P}_{d+e}^{pos}$ .

**Lemma 10.105.** Choose positive  $a_i$  and let  $g_i(y) = e^{a_i y}$ . If Conjecture 10.104 is true then the composition conjecture(part 1) holds for these choices.

*Proof.* For simplicity of exposition we take e = 2. Write

$$f(\mathbf{x}, \mathbf{y}) = \sum a_{Irs} \mathbf{x}^{I} y_{1}^{r} y_{2}^{s}$$

Substituting for  $g_i$  yields

$$F(\mathbf{x}, \mathbf{y}) = \sum a_{\mathrm{Irs}} \mathbf{x}^{\mathrm{I}} e^{a_{1} \mathbf{y} \mathbf{r}} e^{a_{2} \mathbf{y} \mathbf{s}} = \sum a_{\mathrm{Irs}} \mathbf{x}^{\mathrm{I}} e^{(a_{1} \mathbf{r} + a_{2} \mathbf{s}) \mathbf{y}}$$

We find the coefficient  $g_n(\mathbf{x})$  of  $y^n$  by differentiating n times and letting y = 0

$$g_{n}(\mathbf{x}) = \sum a_{\mathrm{Irs}} \mathbf{x}^{\mathrm{I}} (a_{1}r + a_{2}s)^{n}$$

and this is in  $\boldsymbol{P}^{\text{pos}}_{d}$  by the conjecture.

Unfortunately, consider

**Example 10.106.** The operator  $a_1 y_1 \frac{\partial}{\partial y_1} + a_2 y_2 \frac{\partial}{\partial y_2}$  does not necessarily map  $\mathbf{P}_{d+e}^{pos}$  to itself. Consider

$$f = (x + y + z + 2)(x + y + 2z + 1)$$
  
$$g = yf_y + 2zf_z = [y(2y + 2x + 3)] + [6x + 9y + 10]z + 8z^2$$

The constant term of g is two lines that meet at (-3/2, 0). However, the coefficient of z is a line that does not pass through this point, so adjacent coefficients do not intersect, and hence  $g \notin P_3$ .

*Remark* 10.107. It is interesting to compute the generating function of the linear transformation

$$T_n: x^i y^r z^s \mapsto x^i (\alpha r + \beta s)^n$$

We know it does not define a map  $\mathtt{P}_3 \longrightarrow P$ , but it does appear to satisfy  $P_3^{pos} \longrightarrow P.$ 

$$\begin{split} G(x, y, z, u, v, w) &= \sum T_{n}(x^{i}y^{r}z^{s}) \frac{u^{i}v^{r}w^{s}(-1)^{i+r+s}}{i!r!s!} \\ &= \sum x^{i}(\alpha r + \beta s)^{n} \frac{u^{i}v^{r}w^{s}(-1)^{i+r+s}}{i!r!s!} \\ &= e^{-xu} \sum_{k=0}^{n} \alpha^{k} \beta^{n-k} \binom{n}{k} \sum_{r,s} \frac{r^{k}(-v)^{r} s^{n-k}(-w)^{s}}{r!s!} \\ &= e^{-xu} \sum_{k=0}^{n} \alpha^{k} \beta^{n-k} \binom{n}{k} (\sum \frac{r^{k}(-v)^{r}}{r!}) (\sum \frac{s^{n-k}(-v)^{r}}{s!}) \\ &= e^{-xu-v-w} \sum_{k=0}^{n} \alpha^{k} \beta^{n-k} \binom{n}{k} B_{k}(-v) B_{n-k}(-w) \end{split}$$

where  $B_k$  is the Bell polynomial. If  $\alpha = \beta = 1$  then this simplifies to

$$e^{-xu-v-w}B_n(-v-w)$$

which is in  $\widehat{\mathbf{P}}_4$ , but it does not appear to be in  $\widehat{\mathbf{P}}_4$  if  $\alpha \neq \beta$ .

## The derivative-substitution class

It appears that derivatives of compositions satisfy a a much stronger condition than interlacing.

**Definition 10.108.** A polynomial f(x,y) satisfies *derivative-substitution* if for  $m=0,1,\ldots$  and  $\alpha\in\mathbb{R}$ 

1. 
$$\frac{\partial^{m} f}{\partial y^{m}}(x, \alpha) \in \mathbf{P}$$
  
2.  $\frac{\partial^{m} f}{\partial y^{m}}(x, \alpha)$  and  $\frac{\partial^{m+1} f}{\partial y^{m+1}}(x, \alpha)$  interlace.

Clearly polynomials in  $P_2$  satisify derivative-substitution. The next lemma gives some of the properties of this class.

**Lemma 10.109.** If f(x,y) satisfies derivative-substitution and  $f(x,y) = \sum f_i(x)y^i$  then for m = 0, 1, ...

1.  $f_m \in \mathbf{P}$ .

- 2.  $f_m$  and  $f_{m+1}$  interlace.
- 3.  $f(x, y + \gamma)$  satisfies derivative-substitution for all  $\gamma \in \mathbb{R}$ .

*Proof.* Differentiating m times with respect to y and setting y = 0 shows part 1. From the second condition of the definition we have that for all  $\beta \in \mathbb{R}$ 

$$\frac{\partial^{\mathfrak{m}} f}{\partial y^{\mathfrak{m}}}(x,0) + \beta \frac{\partial^{\mathfrak{m}+1} f}{\partial y^{\mathfrak{m}+1}}(x,0) \in \mathbf{P}$$

which implies that  $f_i$  and  $f_{i+1}$  interlace.

The third part follows from the observation that

$$\frac{\partial^{\mathfrak{m}} f(x, y + \gamma)}{\partial y^{\mathfrak{m}}}(x, \alpha) = \frac{\partial^{\mathfrak{m}} f(x, y)}{\partial y^{\mathfrak{m}}}(x, \alpha + \gamma)$$

We have our final generalization of the BMV conjecture

#### Conjecture 10.110.

- 1. If  $f(x, y) \in P_2$  and  $g \in P$  then f(x, g(y)) satisfies derivative-substitution.
- 2. If A, B positive definite, and  $g \in P$  then |xI + g(A + By)| satisfies derivativesubstitution.

For the rest of this section we consider F(x, y) = f(x, g(y)). The Faà di Bruno formula is an expression for the m'th derivative of a composition. In our case we can write

$$\frac{\partial^{\mathfrak{m}}}{\partial y^{k}}f(x,\mathfrak{g}(y)) = \sum_{k} f^{(k)}(x,\mathfrak{g}(y)) A_{\mathfrak{m},k}(y)$$

Observe that this is the constant term of

$$\left(\sum_{k=0}^{m} A_{m,k}(y) \mathsf{D}_{y}^{k}\right) \cdot \left(\sum_{k=0}^{\infty} \mathsf{f}^{(k)}(x, \mathfrak{g}(y)) \frac{z^{k}}{k!}\right)$$

The right hand term is f(x, z + g(y)), and for  $\alpha \in \mathbb{R}$  we see that  $f(x, z + g(\alpha))$  is in P<sub>2</sub>. Define

$$A_{\mathfrak{m}}(\mathbf{x},\mathbf{y}) = \sum A_{\mathfrak{m},k}(\mathbf{x})\mathbf{y}^{k}$$

In order to show that F satisfies derivative-substitution we only need to show that  $A_m(\alpha, y) \in \mathbf{P}$ , and that  $A_m(\alpha, y) \leq A_{m+1}(\alpha, y)$ .

We can do a few special cases.

**Lemma 10.111.** If  $f \in P_2$  then  $f(x, e^y)$  satisfies derivative-substitution. If d is a positive integer then  $f(x, x^d)$  satisfies derivative-substitution.

*Proof.* These follow from Lemmas 1.77 and 1.78.

# **10.19** Recursive constructions

In this section we see how to construct new polynomials in  $\mathbf{P}_d$  from old ones. We can use these constructions to get infinite sequences of interlacing polynomials that are analogous to sequences of orthogonal polynomials.

**Lemma 10.112.** Let  $f \leq g$  be two polynomials in  $\mathbf{P}_d$ . Choose  $b \in \mathbb{R}$ , and a vector  $\mathbf{a} = (a_1, \dots, a_d)$  of positive real numbers. With these assumptions we can conclude that

- $1. \ (\mathfrak{b}+\mathbf{ax^t})\mathfrak{f}-\mathfrak{g} \ \in \mathbf{P}_d.$
- 2.  $(b + ax^t)f g \leq f$ .

*Proof.* Let  $h = (b + ax^t)f - g$ . Since  $h + \alpha f = (b + \alpha + ax^t)f - g$  it suffices to show that  $h \in \mathbf{P}_d$ . All terms of maximal total degree are obtained from f by multiplying by various coordinates of **a**, so they are all positive. If we substitute for all but one variable then we have to show that  $(a + cx)f - \tilde{g}$  is in **P**, where  $\tilde{f}$  and  $\tilde{g}$  are the results for substituting in f and g. Since  $\tilde{f} \leq \tilde{g}$  the result follows from Corollary 1.22.

If we iterate the construction of the lemma we can get an infinite sequence of interlacing polynomials.

**Corollary 10.113.** Choose constants  $b_1, b_2, ..., positive vectors <math>\mathbf{a}_1, \mathbf{a}_2, ..., and positive constants <math>c_1, c_2, ...$  Set  $f_{-1} = 0, f_0 = 1$  and define  $f_n$  recursively by

$$\mathbf{f}_{\mathbf{n}} = (\mathbf{b}_{\mathbf{n}} + \mathbf{a}_{\mathbf{n}}\mathbf{x}^{\mathsf{t}})\mathbf{f}_{\mathbf{n}-1} - \mathbf{c}_{\mathbf{n}}\mathbf{f}_{\mathbf{n}-2}.$$

All  $f_n$  are in  $\mathbf{P}_d$ , and we have interlacings

$$f_0 \ge f_1 \ge f_2 \ge$$

Since the derivative preserves interlacing, we can use derivatives to take the place of  $f_{n-2}$ .

**Corollary 10.114.** Choose constants  $b_1, b_2, ..., positive vectors <math>\mathbf{a}_1, \mathbf{a}_2, ..., and positive vectors <math>\mathbf{c}_1, \mathbf{c}_2, ...$  Set  $f_0 = 1$  and define  $f_n$  recursively by

$$\mathbf{f}_{n} = (\mathbf{b}_{n} + \mathbf{a}_{n}\mathbf{x}^{t})\mathbf{f}_{n-1} - \mathbf{c}_{n} \left(\frac{\partial}{\partial \mathbf{x}}\mathbf{f}_{n-1}\right)^{t}.$$

All  $f_n$  are in  $\mathbf{P}_d$ , and we have interlacings

$$f_0 \ge f_1 \ge f_2 \ge$$

**Example 10.115.** As an interesting special case, consider the recurrence  $f_0 = 1$  and

$$f_{n+1} = \mathbf{x}f_n - \partial \mathbf{x}f_n = (x_1 + \dots + x_d)f_n - (\frac{\partial}{\partial x_1} + \dots + \frac{\partial}{\partial x_d})f_n$$

If we define one variable polynomials  $h_0 = 1$  and  $h_n = xh_{n-1} - dh'_{n-1}$  then it is easy to verify that  $f_n = h_n(x_1 + \dots + x_d)$ . The  $h_n$  are orthogonal polynomials (essentially modified Hermite polynomials), and so we can view these  $f_n$  as generalized Hermite polynomials. In § 11.13.5 we will see a true generalization of Hermite polynomials to n-variables.

The linear combination  $(b + ax^t)g + f$  also has interlacing properties. This generalizes Corollary 1.14.

#### **Lemma 10.116.** If $f \leq g$ in $\mathbf{P}_d$ and $\mathbf{a}$ is positive then $(\mathbf{b} + \mathbf{a}\mathbf{x}^t)\mathbf{g} + \mathbf{f} \leq \mathbf{g}$ .

*Proof.* By linearity it suffices to show that the left hand side is in  $\mathbf{P}_d$ . The homogeneous part is  $(\mathbf{a}\mathbf{x}^t)g^H$ , the degrees are correct, and substitution follows from Corollary 1.14.

## 10.20 Two by two matrices preserving interlacing in $P_d$

In this section we generalize results from Chapter 3 to  $\overline{\mathbf{P}}_d$ . Many of the properties of one variable have immediate generalizations to d variables. However, we don't have a concept of "mutually interlacing" for more than one variable, so Theorems such as 3.7 are only possible for two by two matrices.

The following Lemma generalizes Corollary 3.8. The restrictive assumptions arise from the necessity to control the homogeneous part.

**Lemma 10.117.** Suppose that  $f_1 \ll f_2$  (or  $f_1 \ll f_2$ ) in  $P_d$  have positive leading coefficients, and  $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} f_1 \\ f_2 \end{pmatrix} = \begin{pmatrix} g_1 \\ g_2 \end{pmatrix}$ . If a, b, c, d are positive and if the determinant ad - bc is positive then  $g_1 \ll g_2$ , and if it is negative  $g_2 \ll g_1$ .

*Proof.* The positivity assumptions imply that  $g_1, g_2$  have positive homogeneous part. Corollary 3.8 implies that  $g_1, g_2 \in P_d$ , and that the interlacings hold as stated.

As long as all coefficients are positive then the matrices generalizing Corollary 3.54 preserve interlacing.

**Lemma 10.118.** *Suppose that*  $\alpha$ ,  $\beta$ , **a**, **b** *are vectors with all positive coordinates. The following matrices preserve interlacing in*  $P_d$ :

$$\begin{pmatrix} \alpha & \mathbf{a} \cdot \mathbf{x} \\ 0 & \beta \end{pmatrix} \quad \begin{pmatrix} \mathbf{a} \cdot \mathbf{x} & 0 \\ \alpha & \mathbf{b} \cdot \mathbf{x} \end{pmatrix} \quad \begin{pmatrix} \mathbf{a} \cdot \mathbf{x} & -\alpha \\ 0 & \mathbf{b} \cdot \mathbf{x} \end{pmatrix} \quad \begin{pmatrix} \mathbf{a} \cdot \mathbf{x} & -\alpha \\ \beta & \mathbf{b} \cdot \mathbf{x} \end{pmatrix}$$

*Proof.* All polynomials occurring in a product have positive homogeneous part. Substitution and interlacing follow from the one variable case.

Next, we generalize Proposition 3.56.

**Lemma 10.119.** Suppose that  $M = \begin{pmatrix} f & g \\ h & k \end{pmatrix}$  is a matrix of polynomials with in  $P_d$ . M preserves interlacing if

- 1.  $g \longrightarrow f, k \longrightarrow h$ , and both interlacings are strict.
- 2. The determinant  $\begin{vmatrix} f & g \\ h & k \end{vmatrix}$  is never zero.

*Proof.* If  $M({s \atop s}) = {u \atop v}$  then the homogeneous parts of u and v are positive. Interlacing and substitution follow from the one variable case.

How can we find such matrices? Suppose that  $f\in P_2^{gen}$ . Recall that this means that the graph of f has no intersection points. Choose  $g\in P_2^{gen}$  so that  $f \lessdot g$ . For instance, we could take  $g=\frac{\partial f}{\partial x}$ . Define

$$\mathsf{M} = \begin{pmatrix} \mathsf{f} & \mathsf{g} \\ \frac{\partial \mathsf{f}}{\partial \mathsf{x}} & \frac{\partial \mathsf{g}}{\partial \mathsf{x}} \end{pmatrix}$$

It follows from Lemma 1.55 that the determinant of M is never zero.

# **10.21** Polynomials from matrices

We can construct polynomials in several variables from symmetric matrices, and with some simple assumptions the resulting polynomial is in  $P_d$ . We begin with a construction for a polynomial in d variables that generalizes Lemma 9.29.

**Lemma 10.120.** *If* C *is symmetric and*  $A_2, \ldots, A_d$  *are pairwise commuting positive definite symmetric matrices, then* 

$$f(\mathbf{x}) = \det(x_1I + x_2A_2 + \dots + x_dA_d + C)$$

is in  $\mathbf{P}_{d}$ .

*Proof.* Since  $A_2, \ldots, A_d$  are pairwise commuting and symmetric, they are simultaneously diagonalizable. There is an orthogonal matrix P such that  $PA_kP^t = D_k$ , where  $D_k = (e_{ij}^k)$  is a diagonal matrix with all positive entries on the diagonal. We note

$$f(\mathbf{x}) = \det(x_1 I + x_2 A_2 + \dots x_d A_d + C)$$
  
= det(P<sup>t</sup>) det(x\_1 I + x\_2 D\_2 + \dots + x\_d D\_d + PCP<sup>t</sup>) det(P)

As before this satisfies  $x_1$ -substitution since PCP<sup>t</sup> is symmetric, and the homogeneous part is

$$f^{H}(\mathbf{x}) = \det(x_1 I + x_2 D_2 + \dots + x_d D_d)$$
  
=  $\prod_k (x_1 + x_2 e_{kk}^2 + \dots + x_d e_{kk}^d).$ 

which has all positive coefficients.

Remark 10.121. A variant of the lemma is to assume that

$$f(\mathbf{x}) = \det(x_1D_1 + \dots + x_dD_d + C)$$

where C is symmetric, all  $D_i$  are diagonal, and  $D_1$  has all positive diagonal entries. If  $\sqrt{D_1^{-1}}$  is the diagonal matrix whose entries are the inverses of the positive square roots of the diagonal entries of  $D_1$ , then

$$f(\mathbf{x}) = det(D_1) det\left(x_1I + x_2D_2D_1^{-1} + \dots + x_dD_dD_1^{-1} + \sqrt{D_1^{-1}} C \sqrt{D_1^{-1}}\right)$$

We can now apply Lemma 10.120 to see that  $f(\mathbf{x}) \in \mathbf{P}_d$ .

How can we find such commuting families of positive definite matrices? If f is a polynomial, and  $\lambda$  is an eigenvalue of a positive definite matrix A, then  $f(\lambda)$  is an eigenvalue of f(A). In particular, if  $f \in \mathbf{P}^{pos}$  then f(A) is positive definite. Since  $e^x$  is a limit of polynomials in  $\mathbf{P}^{pos}$  and any two polynomials in A commute, we have

**Corollary 10.122.** Suppose that  $f, f_2, ..., f_d$  are polynomials in  $P^{pos}$ , A is positive definite and  $\alpha, \alpha_2, ..., \alpha_d$  are positive. The following polynomials are in  $P_d$ .

$$det(x_1I + x_2f_2(A) + \dots + x_df_d(A) + f(A))$$
$$det(x_1I + x_2e^{\alpha_2A} + \dots + x_de^{\alpha_dA} + e^{\alpha A})$$

The polynomials that arise from these constructions can interlace.

**Lemma 10.123.** *If* D *is a diagonal matrix with all positive diagonal entries,* C *is symmetric, and* P *is a principle submatrix of* xI + yD + C*, then* 

$$|\mathbf{x}\mathbf{I} + \mathbf{y}\mathbf{D} + \mathbf{C}| \leq |\mathbf{P}| in \mathbf{P}_2$$

*Proof.* We may assume P is the result of deleting the first row and first column of |xI + yD + C|. If  $D = diag(d_i)$ ,  $C = (c_{ij})$ ,  $v = (c_{12}, \dots, c_{1n})$  then for any  $\alpha \in \mathbb{R}$  we can write

$$g_{\alpha}(\mathbf{x}, \mathbf{y}) = \begin{vmatrix} \mathbf{x} + d_1 \mathbf{y} + c_{11} + \alpha & \nu \\ \nu^{t} & P \end{vmatrix}$$
$$= \begin{vmatrix} \mathbf{x} + d_1 \mathbf{y} + c_{11} & \nu \\ \nu^{t} & P \end{vmatrix} + \begin{vmatrix} \alpha & 0 \\ 0 & P \end{vmatrix}$$
$$= |\mathbf{x}\mathbf{I} + \mathbf{y}\mathbf{D} + \mathbf{C}| + \alpha |\mathbf{P}|$$

Since  $g_{\alpha} \in \mathbf{P}_2$  for any  $\alpha \in \mathbb{R}$  it follows that the desired polynomials interlace.

*Remark* 10.124. If we take y = 0 then we recover the classical result that principle submatrices interlace. A similar result holds for P<sub>d</sub>. See Theorem 1.61.

If C is the all zero matrix and y = 1 then  $f(x) = |xI + yD + C| = \prod(x + d_i)$ and  $|P| = \frac{f(x)}{x+d_1}$ . These polynomials are the ones occurring in quantitative sign interlacing.

## **10.22** Extending three interlacing polynomials

Consider a polynomial in  $P_3$ 



If we are only given  $f_{00} \leq f_{01}$ ,  $f_{10}$  then we show that we can add terms to  $f_{00} + yf_{10} + zf_{01}$  so that the resulting polynomial F is in P<sub>3</sub>. (See p. 291.) We say that F is an *extension* of  $f_{00}$ ,  $f_{10}$ ,  $f_{01}$ . Given one extension, we show how to find many more with the same initial data.

The term  $f_{11}$  is not arbitrary, since it interlaces  $f_{01}$  and  $f_{10}$ , and satisfies  $f_{01}f_{10} - f_{00}f_{11} \ge 0$ . We show that if  $f_{00}$  has degree two then  $f_{10}f_{01} - f_{00}W \ge 0$  if and only if there is an extension of  $f_{00}$ ,  $f_{10}$ ,  $f_{01}$  such that  $W = f_{11}$ . We conjecture that this is true in general (Question 141).

**Lemma 10.125.** Suppose that  $f \leq g$  and  $f \leq h$ , where f, g, h have positive leading coefficients. Then there are  $F \in P_3$  such that if we write  $F(x, y, z) = \sum f_{ij}(x) y^i z^j$  then

$$f_{00} = f$$
,  $f_{10} = g$ ,  $f_{01} = h$ 

*Proof.* If we take three sequences  $a_i, b_i, c_i$ , a constant  $\alpha$ , and form the product

$$F = \alpha \prod (1 + a_i x + b_i y + c_i z)$$

then

$$F_{00} = \alpha \prod (1 + a_i x)$$

$$F_{10} = \sum b_i \frac{F_{00}}{1 + a_i x}$$

$$F_{01} = \sum c_i \frac{F_{00}}{1 + a_i x}$$

Given f, g, h we can, following the argument of Lemma 9.103, find  $\alpha$  and the sequences  $a_i$ ,  $b_i$ ,  $c_i$  so that  $F_{00} = f$ ,  $F_{10} = g$ ,  $F_{01} = h$ .

*Remark* 10.126. We can use a determinant identity to derive a different extension of three polynomials. Recall (9.1.3). If we let  $D = diag(d_i)$ , f = |I + xD|,  $V = (V_i)$  and  $W = (W_i)$  then

$$\begin{split} |I + xD + yVV^t + zWW^t| = \\ f + y\sum V_i^2 \frac{f}{1 + d_i x} + z\sum W_i^2 \frac{f}{1 + d_i x} + yz\sum_{i < j} \frac{f}{(1 + xd_i)(1 + xd_j)} \begin{vmatrix} V_i & V_j \\ W_i & W_j \end{vmatrix}^2 \end{split}$$

Since D, VV<sup>t</sup>, WW<sup>t</sup> are positive semi-definite, the right hand side is in  $\overline{\mathbf{P}}_3$ . (See Lemma 11.38.) It follows that we have an interlacing square



There are many determinants that will give the same initial data. This follows from the more general result that shows that the initial data only constrains the diagonals.

**Lemma 10.127.** Suppose that D is a diagonal matrix,  $D_1$  and  $E_1$  have the same diagonal, as do  $D_2$  and  $E_2$ . If we write

$$\begin{split} \left| I + xD + yD_1 + zD_2 \right| &= \sum U_{ij}(x)y^i z^j \\ \left| I + xD + yE_1 + zE_2 \right| &= \sum V_{ij}(x)y^i z^j \end{split}$$
 then 
$$U_{00} = V_{00}, \quad U_{10} = V_{10}, \quad U_{01} = V_{01}$$

*Proof.* Clearly  $U_{00} = V_{00}$ . Let  $D_1 = (d_{ij})$  and  $E_1 = (e_{ij})$ . Then

$$\begin{split} & U_{10} = \frac{\partial}{\partial y} \big| I + xD + yD_1 + 0D_2 \big|_{y=0} \\ &= \sum_{i,j} d_{ij} \big| (I + xD)[i,j] \big| \\ &= \sum_i d_{ii} \big| (I + xD)[i,i] \big| \qquad \text{since } I + xD \text{ is diagonal} \\ &= \sum_i e_{ii} \big| (I + xD)[i,i] \big| \qquad \text{since } e_{ii} = d_{ii} \\ &= V_{10} \end{split}$$

If we are given  $f_{00} \leftarrow f_{10}$ ,  $f_{01}$  in  $\mathbf{P}^{\text{pos}}$ , then by the first lemma we can realize  $f_{00}$ ,  $f_{10}$ ,  $f_{01}$  as the coefficients of  $|I + xD + yD_1 + zD_2|$  where D,  $D_1$ ,  $D_2$  are positive definite diagonal matrices. If we can find symmetric matrices  $S_1$ ,  $S_2$  with zero diagonal such that  $D_1 + S_1$  and  $D_2 + S_2$  are positive definite, then  $f_{00}$ ,  $f_{10}$ ,  $f_{01}$  are also realized as initial coefficients of  $|I + xD + y(D_1 + S_1) + z(D_2 + S_2)|$ .

**Example 10.128.** Consider the case  $f_{00} \in \mathbf{P}^{pos}(2)$ . Given the initial data  $f_{00} \leq f_{10}, f_{01}$  we first determine all  $\alpha$  that arise from Lemma 10.127. Consider

$$g(B,C) = \left| \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + x \begin{pmatrix} a_1 & 0 \\ 0 & a_2 \end{pmatrix} + y \begin{pmatrix} b_1 & B \\ B & b_2 \end{pmatrix} + z \begin{pmatrix} c_1 & C \\ C & c_1 \end{pmatrix} \right|$$

where we choose  $a_1, a_2, b_1, b_2, c_1, c_2$  so that

$$g(0,0) = f_{00} + yf_{10} + zf_{01} + \alpha yz.$$

This representation is possible by Lemma 10.125. Recall

If r, s are positive then  $|\begin{smallmatrix} r & t \\ t & s \end{smallmatrix}|$  is positive definite if and only if  $rs > t^2$ .

Thus,  $g(B, C) \in P_3(2)$  if and only if  $b_1b_2 > B^2$  and  $c_1c_2 > C^2$ . Now  $\alpha$  is the coefficient of yz which is

 $b_1c_2+b_2c_1+2BC.$ 

Using the constraints above, we find

$$\left(\sqrt{b_1c_2} - \sqrt{b_2c_1}\right)^2 \leqslant \alpha \leqslant \left(\sqrt{b_1c_2} + \sqrt{b_2c_1}\right)^2 \tag{10.22.1}$$

All  $f_{00}$ ,  $f_{01}$ ,  $f_{10}$  and  $\alpha$  satisfying this inequality have extensions to  $P_3$ .

Conversely, assume that  $f_{00}$ ,  $f_{10}$ ,  $f_{01}$  are determined by g(0,0). If we solve the inequality

 $f_{10}f_{01}-\alpha f_{00} \geqslant 0$ 

for  $\alpha$  we get (10.22.1).

## **10.23** Subsets of $P_d$

In one variable we had two different ways to talk about  $\mathbf{P}^{\text{pos}}$ . We could describe it in terms of coefficients (all the same sign), or in terms of roots (all negative). However,  $\mathbf{P}^{(-1,1)}$  has no definition in terms of coefficients, but is given only in terms of the location of its roots. In higher dimensions the graph corresponds to zeros in one dimension. We define

**Definition 10.129.** If  $K \subset \mathbb{R}^d$  then

$$\mathbf{P}_{d}^{\mathsf{K}} = \{ \mathsf{f} \in \mathtt{P}_{d} \mid \text{ the graph of } \mathsf{f} \text{ is disjoint from } \mathbf{K} \}$$
$$= \{ \mathsf{f} \in \mathtt{P}_{d} \mid \mathsf{f}(\mathbf{x}) = 0 \implies \mathbf{x} \notin \mathbf{K} \}$$

Consider some examples and properties.

- 1. If  $\mathbf{K} = \emptyset$  then  $\mathbf{P}_{d}^{\mathbf{K}} = \mathbb{P}_{d}$ , since there are no restrictions.
- 2. If  $\mathbf{K} \subset \mathbf{K}'$  then  $\mathbf{P}_d^{\mathbf{K}'} \subset \mathbf{P}_d^{\mathbf{K}}$ .
- 3. Let v, w be two points of  $\mathbb{R}_d$ , and let

$$\mathbf{K} = \{ \mathbf{x} \in \mathbb{R}^d \mid \mathbf{x} \ge \mathbf{w} \} \cup \{ \mathbf{x} \in \mathbb{R}^d \mid \mathbf{x} \le \mathbf{v} \}$$

Every solution variety of  $\mathbf{P}_d^{\mathbf{K}}$  must meet the segment  $\overline{vw}$ . This is an analog of  $\mathbf{P}^{(-1,1)}$ .

- 4. Let  $\mathbf{K} = \{(x, x, ..., x) \mid x \in \mathbb{R}\}$ . Since every  $f \in P_d$  satisfies  $f(x ..., x) \in \mathbf{P}$ , it follows  $\mathbf{P}_d^{\mathbf{K}} = \emptyset$ .
- 5. Suppose  $\mathbf{K} = \{ \mathbf{x} \in \mathbb{R}^2 \mid \mathbf{x} \ge 0 \text{ or } \mathbf{x} \le 0 \} \setminus (0,0)$ . Then,  $\mathbf{P}_2^{\mathbf{K}}$  consists of polynomials whose graphs are straight lines passing through the origin.
- 6. Suppose that  $\mathbf{K} = \{\mathbf{x} \in \mathbb{R}^n \mid \mathbf{x} \ge 0\}$ . We will show that  $\mathbf{P}_d^{\mathbf{K}} = \mathbf{P}_d^{\text{pos}}$ . Since all coefficients of  $\mathbf{P}_d^{\text{pos}}$  are positive, it follows that  $f \in \mathbf{P}_d^{\text{pos}}$  has no root in  $\mathbf{K}$ . We prove the converse by induction on d. For d = 1 we know that all the coefficients are the same sign if and only if no root is non-negative. Suppose that  $f \in P_d$  has no root  $\mathbf{x}$  with  $\mathbf{x} \ge 0$ . Write  $f(\mathbf{x}) = \sum f_i(x_1, \dots, x_{d-1}) \mathbf{x}_d^i$ . If we substitute  $x_d = 0$  then we see that  $f_0(x_1, \dots, x_{d-1})$  has no root with all coordinates non-negative, so by induction all coefficients of  $f_0(x_1, \dots, x_{d-1})$  have the same sign. Since all coefficients of  $f^H$  are positive, all coefficients of  $f_0(x_1, \dots, x_{d0-1})$  are positive.

Next, if  $f \in \mathbf{P}_d^K$ , then  $\frac{\partial f}{\partial x_d} \in \mathbf{P}_d^K$  since the roots of  $\frac{\partial f}{\partial x_d}$  lie between roots of f. Consequently, we conclude as above that  $f_1(x_1, \ldots, x_{d-1})$  has all positive coefficients. Continuing, we see f has all positive coefficients.

Many different **K** can determine the same set of polynomials. For example, the set  $\mathbf{K} = \{\mathbf{x} \in \mathbb{R}^d \mid \mathbf{x} \ge 0\}$  and  $\mathbf{K}' = \mathbf{K} \setminus p$  where p > 0 satisfy  $\mathbf{P}_d^{\mathbf{K}} = \mathbf{P}_d^{\mathbf{K}'}$ . However, if we choose p to be the origin, then  $\mathbf{P}_d^{\mathbf{K}} \subsetneq \mathbf{P}_d^{\mathbf{K}'}$  since  $x_1 + \dots + x_d$  is in  $\mathbf{P}_d^{\mathbf{K}'}$  but is not in  $\mathbf{P}_d^{\mathbf{K}}$ . We define

#### Definition 10.130.

$$\overline{K} = \bigcup \left\{ L \subset \mathbb{R}^d \mid P_d^K = P_d^L \right\}$$

In some cases we can determine  $\overline{\mathbf{K}}$ .

**Lemma 10.131.** Suppose that K is a closed convex region of  $\mathbb{R}^d$  whose supporting hyperplanes have an equation in  $P_d$ . Then,  $K = \overline{K}$ .

*Proof.* Since  $K \subset \overline{K}$  it suffices that any point v not in K is not in  $\overline{K}$ . Pick a supporting hyperplane H such that K is on one side of it, and v is in the interior of the other side. Choose a hyperplane H' parallel to H, and containing v. It follows that v lies on the graph of a polynomial that is disjoint from K, and hence  $v \notin \overline{K}$ .

*Remark* 10.132. In case d = 2 we see that if **K** is a closed convex region whose boundary is a non-increasing curve then  $\mathbf{K} = \overline{\mathbf{K}}$ .

We now ask when  $\mathbf{P}_d^{\mathbf{K}}$  is closed under differentiation. We say a set S is convex in the i-th coordinate if whenever *v*, *w* agree in all coordinates except the i-th, then the segment  $\overline{vw}$  is in S.

**Lemma 10.133.** If  $\mathbb{R}^d \setminus K$  is convex in the *i*-th coordinate then  $\mathbf{P}_d^K$  is closed under  $f \mapsto \frac{\partial f}{\partial x_i}$ .

*Proof.* To show that  $\frac{\partial f}{\partial x_i} \in \mathbf{P}_d^{\mathbf{K}}$  we fix all but the i-th coordinate and consider the roots of the resulting polynomial. Since the roots of  $\frac{\partial f}{\partial x_i}$  lie between the roots of f the convexity in the i-coordinate imply that they lie in  $\mathbb{R}^d \setminus \mathbf{K}$ .

# 10.24 Higher dimensional recurrences

In this section we investigate polynomials defined by two and three dimensional recurrences. Some of these polynomials are in one variable, and some in two. One of the interesting features is that we do not always have that consecutive terms interlace. We begin by considering polynomials  $f_{i,j}$  satisfying the similar recurrences

$$f_{i,j} = (x - a_i)f_{i-1,j} + b_{ij}f_{i,j-1}$$
(10.24.1)

$$f_{i,j} = (x - a_{ij})f_{i-1,j} + b_{ij}f_{i,j-1}$$
(10.24.2)

$$f_{i,j} = (x - a_i)f_{i-1,j} + b_i f_{i-1,j-1}$$
(10.24.3)

We can visualize these recursions using lattice diagrams:

We will see that recurrence (10.24.1) is a disguised product, that recurrence (10.24.2) fails unless the coefficients satisfy certain restrictions, and recurrence (10.24.3) comes from coefficients of products in  $P_2$ .

**Lemma 10.134.** Suppose  $f_{0,0} = 1$  and f satisfies recurrence (10.24.1). Then  $f_{i,j}$  is a constant multiple of  $\prod_{k=1}^{i} (x - a_k)$ . In particular, all  $f_{i,j}$  are in P, and consecutive  $f_{i,j}$  interlace.

*Proof.* Assume by induction of i, j that

$$f_{r,s} = \alpha_{r,s} \prod_{k=1}^{r} (x - a_k)$$

for all (r, s) < (i, j). We then have

$$\begin{split} f_{i,j} &= (x - a_i)f_{i-1,j} + b_{i,j}f_{i,j-1} \\ &= (x - a_i)\alpha_{i-1,j}\prod_{1}^{i-1}(x - a_k) + b_{i,j}\alpha_{i,j-1}\prod_{1}^{i}(x - a_k) \\ &= (\alpha_{i-1,j} + b_{i,j}\alpha_{i,j-1})\prod_{1}^{i}(x - a_k) \end{split}$$

Here is a simple recurrence whose proof that all members are in **P** only depends on the additivity of interlacing.

**Lemma 10.135.** Let  $h_n = \prod_{i=1}^{n} (x + a_i)$ , and let  $h_n \leftarrow g_n$  where all  $g_n$  have positive leading coefficients. If we define  $f_0 = 1$  and

$$f_n = (x + a_n)f_{n-1} + g_n$$

*then*  $h_n \leq f_n$ . *In particular, all*  $f_i$  *have all real roots.* 

*Proof.* We assume by induction that  $h_{n-1} \leq f_{n-1}$  Multiplying each side by by  $(x - a_n)$  yields  $h_n \leq (x - a_n)f_{n-1}$ . Since we also have  $h_n \leftarrow g_n$  we can add these interlacings to conclude that

$$h_n \underline{\ll} (x - a_n) f_{n-1} + g_n = f_n$$

We can apply this lemma to construct a particular sequence of polynomials whose members all have all real roots, but consecutive ones do not generally interlace.

**Corollary 10.136.** If  $f_0 = 1$  and  $f_n$  satisfies the recursion

$$f_n = (x + n)f_{n-1} + (x + n - 2)\prod_{1}^{n-1} (x + i)$$

then all  $f_n$  are in **P**.

*Proof.* Take 
$$h_n = \prod_{i=1}^{n} (x + i)$$
, and  $g_n = (x + n - 2)h_{n-1}$ .

**Lemma 10.137.** Suppose that  $f_{0,0} = 1$  and  $f_{i,j}$  satisfies (10.24.3) where all  $a_i$  have the same sign. Then all  $f_{i,j}$  are in **P**.

*Proof.* We claim that the  $f_{i,j}$  are the coefficients of the products

$$\prod_{j=1}^i(x+b_jy-a_j)=\sum_{j=0}^if_{i,j}(x)\,y^j$$

This is certainly true when i = 0. We only need to check that if we write

$$\prod_{j=1}^{i}(x+b_{j}y-a_{j})=\sum_{j=0}^{i}g_{i,j}(x)y^{j}$$

then the  $g_{i,j}$  satisfy the same recurrences as do the  $f_{i,j}$ :

$$\begin{split} \prod_{1}^{i} (x + b_{j}y - a_{j}) &= (x + b_{i}y - a_{i}) \sum_{j=0}^{i-1} g_{i-1,j}(x) y^{j} \\ &= \sum_{i} ((x - a_{i})g_{i-1,j}(x) - b_{i}g_{i-1,j-1}(x)) y^{j} \end{split}$$

and so we are done by induction.

We can generalize this argument to products in more variables - this leads to polynomials in x with more indices. For example, consider the recurrence

$$f_{i,j,k}(x) = (x - a_i)f_{i-1,j,k} + b_i f_{i,j-1,k} + c_i f_{i,j,k-1}$$
(10.24.4)

**Lemma 10.138.** If  $f_{0,0,0} = 1$  and  $f_{i,j,k}$  satisfies (10.24.4), all  $b_i$  have the same sign, and all  $c_i$  have the same sign then all  $f_{i,j,k}$  are in **P**.

*Proof.* Consider the product below, and proceed as in Lemma 10.137.

$$\prod_{h=1}^{i} (x - a_h + b_h y + c_h z) = \sum f_{i,j,k}(x) y^j z^k$$

As another product example, consider the recurrence

$$f_{i,j} = (a_i x + b_i) f_{i,j-1} + (c_i x + d_i) f_{i-1,j}$$
(10.24.5)

whose diagram is

$$f_{i-1,j} \xrightarrow{c_i x + d_i} f_{i,j} \begin{bmatrix} a_i x + b_1 \\ f_{i,j-1} \end{bmatrix}$$

**Lemma 10.139.** If  $f_{0,0} = 1$ ,  $f_i$ , j satisfies (10.24.5), and  $a_i d_i - c_i b_i > 0$  for all i then  $f_{i,j} \in P$ .

*Proof.* The assumptions imply that all the factors in the product below are in  $\overline{P}_{2}$ , so we can proceed as above

$$\prod_{j=1}^{i}((a_{j}x+b_{j})+(c_{j}+d_{j}x)y)=\sum_{j}f_{i,j}(x)y^{j}$$

In fact, we can take any polynomial in  $\overline{\mathbf{P}}_2$  that has some free parameters, take their product, and the coefficients  $f_{i,j}$  of  $y^j$  will satisfy a recurrence, and all the  $f_{i,j}$  belong to  $\mathbf{P}$ , and consequently will interlace.

Here is another disguised product

$$f_{i,j} = (x - b_i)f_{i-1,j} + (y - d_j)f_{i,j-1}$$
(10.24.6)

with diagram

$$f_{i-1,j} \xrightarrow{x-b_i} f_{i,j} \\ \begin{vmatrix} y - d_j \\ f_{i,j-1} \end{vmatrix}$$

The proof is a simple modification of the proof of Lemma 10.134.

**Lemma 10.140.** Suppose that  $f_{0,0} = 1$  and  $f_{i,j}$  satisfies (10.24.6). Then  $f_{i,j}$  is a constant multiple of  $\prod_{k=1}^{i} (x - b_k) \cdot \prod_{h=1}^{j} (y - d_h)$ .

We can get many recurrences from products. Here is a general approach. Let

$$g(\mathbf{x},\mathbf{y},\mathbf{z}) = \sum g_{\mathbf{r}}(\mathbf{x},\mathbf{z})\mathbf{y}^{\mathbf{r}}$$

where  $g \in \mathbf{P}_3$ , choose  $t_i \in \mathbb{R}$ , and consider the product

$$\prod_{j=1}^{\iota}g(x,y,t_j)=\sum_{i,j}f_{i,j}(x)y^j$$

We know that all  $f_{i,j} \in P$ , and consecutive ones interlace. The recurrence is easily found:

$$\begin{split} \prod_{j=1}^{i+1} g(x,y,t_j) &= g(x,y,t_{i+1}) \sum_{i,s} f_{i,s}(x) y^s \\ &= \left( \sum g_r(x,t_{i+1}) y^r \right) \sum_{i,s} f_{i,s}(x) y^s \end{split}$$

and comparing coefficients of y<sup>j</sup> yields

$$f_{i+1,j} = \sum_{r+s=j} g_r(x, t_{i+1}) f_{i,s}(x)$$

CHAPTER



# The polynomial closure of P<sub>d</sub>

In this chapter we study the polynomial closure of  $\mathbf{P}_d$ . For polynomials in one variable we know that  $\mathbf{P}(n)$  is exactly the set of polynomials in the closure of  $\mathbf{P}(n)$ , so the closure of  $\mathbf{P}$  leads to analytic functions. However, there are polynomials in the closure of  $\mathbf{P}_d(n)$  that are not in  $\mathbf{P}_d(n)$ , so we have two closures to consider: new polynomials and analytic functions. As expected, most properties of  $\mathbf{P}_d$  extend to these closures. We consider the analytic closure in the next chapter.

## **11.1** The polynomial closure of **P**<sub>2</sub>

In this section we investigate properties of the polynomial closure of  $P_2$ . We begin with the general definitions.

**Definition 11.1.**  $\overline{\mathbf{P}}_d(n)$  is the set of all polynomials that are limits of polynomials in  $\mathbb{P}_d(n)$ .  $\overline{\mathbf{P}}_d^{\text{pos}}(n)$  is defined similarly. We set

$$\overline{\mathbf{P}}_{d} = \overline{\mathbf{P}}_{d}(1) \cup \overline{\mathbf{P}}_{d}(2) \cup \dots$$

We say that f,  $g \in \overline{P}_d$  interlace if  $f + \alpha g \in \overline{P}_d$  for all real  $\alpha$ .

Although most properties of  $P_2$  extend to  $\overline{P}_2$ , there are some slight differences. The polynomial  $x^n y^m$  is in  $\overline{P}_2$  since

$$\lim_{\varepsilon \to 0^+} (x + \varepsilon y)^n (y + \varepsilon x)^m = x^n y^m$$

and  $(x + \epsilon y)^n (y + \epsilon x)^m$  is in P<sub>2</sub>. This shows that there are polynomials in  $\overline{P}_2$  where the x-degree is n, the y-degree is m, and the total degree is n + m. Consider this example that shows how the coefficients can interlace in complicated ways.

**Example 11.2.** The definition of f(x, y) as a limit of polynomials in P<sub>2</sub>

$$\begin{split} f(x,y) &= \lim_{\varepsilon \to 0^+} (\varepsilon x + y - 1)(\varepsilon x + y - 2)(\varepsilon x + y + 3)(x + y + 1)(x + 2y + 1) \\ &= (y - 1)(y - 2)(y + 3)(x + y + 1)(x + 2y + 1) \end{split}$$

shows that  $f \in \overline{\mathbf{P}}_2$ . If we expand f we get

$$\begin{split} &= (6+12x+6x^2) + (11+4x-7x^2)y + (-9-21x)y^2 + \\ &\quad (-13+2x+x^2)y^3 + (3+3x)y^4 + 2y^5 \\ &= f_0 + f_1\,y + f_2\,y^2 + f_3\,y^3 + f_4\,y^4 + f_5\,y^5 \end{split}$$

These coefficients interlace in ways that are not possible for polynomials in  $P_2$ , namely

$$f_0 \underline{\gg} f_1 \lessdot f_2 \triangleright f_3 \lessdot f_4.$$

We can use the idea of this construction to show that if  $g(x) \in \mathbf{P}$  then  $g(x) \in \overline{\mathbf{P}}_2$ . Simply note that  $g(x + \epsilon y + \epsilon)$  is in  $\mathbb{P}_2$  and converges to g(x) as positive  $\epsilon$  goes to 0. Similarly, if we let  $\epsilon$  go to zero through positive values we see that

$$\mathbf{P}^{\text{pos}} \subset \overline{\mathbf{P}}_{2}^{\text{pos}}$$

The mapping  $f \times g \mapsto f(x)g(y)$  defines a map  $\mathbf{P} \times \mathbf{P} \longrightarrow \overline{\mathbf{P}}_2$ , and  $\mathbf{P}^{\text{pos}} \times \mathbf{P}^{\text{pos}} \longrightarrow \overline{\mathbf{P}}_2^{\text{pos}}$ .

The two polynomials  $f = x^n y^m$  and  $g = x^{n-1} y^{m-1}$  do not interlace. However, they do satisfy  $f \stackrel{+}{\sim} g$  in the sense that  $f + \alpha g \in \mathbf{P}$  for all *positive*  $\alpha$ .

In order to see that polynomials in  $\overline{\mathbf{P}}_2$  interlace we can express them as limits of interlacings. For instance, the interlacing  $xy - 1 \leq x + y$  follows from

$$(xy-1) + \alpha(x+y) = \lim_{\varepsilon \to 0^+} \begin{vmatrix} \varepsilon x + y + \alpha & \sqrt{1+\alpha^2} \\ \sqrt{1+\alpha^2} & x + \varepsilon y + \alpha \end{vmatrix}$$

**Example 11.3.** Unlike in  $P_2$ , the signs of a polynomial in  $\overline{P}_2$  can be all negative. This follows from the observation that if  $f \in \overline{P}_2$  then  $-f \in \overline{P}_2$ . This is a consequence of the limit

$$\lim_{\varepsilon \to 0^+} (\varepsilon x + \varepsilon y - 1)f = -f$$

If  $f \in \overline{\mathbf{P}}_2(n)$  then we define the homogeneous part  $f^H$  to be all terms of total degree n. Unlike P<sub>2</sub>, the example above shows that all the coefficients of the homogeneous part can be negative. There are no other possibilities - see Lemma 11.15.

We can apply Lemma 9.32 to deduce

**Lemma 11.4.** If  $f \in \overline{\mathbf{P}}_2(n)$  then f(x, x) and  $x^n f(x, -1/x)$  are in  $\mathbf{P}$ .

This shows that xy + 1 isn't in  $\overline{P}_2$  even though it satisfies substitution. The graphs of polynomials in  $\overline{P}_2$  are more complicated than those in P<sub>2</sub>. Consider

$$f = (x + 4y + xy + 1)(x + y + 3xy + 2)(x + 5y + .5)(x + 2y + 1).$$

The graph of f in Figure 11.1 has two vertical asymptotes at x = -4 and x = -3. None the less, every vertical line except these two meets the graph in exactly four points.



Figure 11.1: The graph of of a polynomial in  $\overline{\mathbf{P}}_2$ 

We can use determinants to realize members of  $\overline{\mathbf{P}}_2$ .

**Lemma 11.5.** Suppose that C is a symmetric matrix, and D is a diagonal matrix where the i-th diagonal entry is  $d_{i1}x_1 + d_{i2}x_2$  where  $d_{ij}$  is non-negative. The polynomial det(C + D) is in  $\overline{P}_2$ .

*Proof.* Choose  $\epsilon > 0$  and let  $D_{\epsilon}$  have diagonal entries  $\sum (d_{ij} + \epsilon)x_i$ . A argument similar to Lemma 10.120 shows that  $det(C + D_{\epsilon}) \in P_2$ . Taking limits gives the result.

We now find some simple polynomials in  $\overline{\mathbf{P}}_2$  that are not in  $P_2$ .

**Lemma 11.6.** If  $f(x) \in P^{pos}$  then  $f(-xy) \in \pm \overline{P}_2$ .

*Proof.* Since f is a product of linear factors, it suffices to show that  $-xy + a^2 \in \overline{\mathbf{P}}_2$ . The determinant

$$\begin{array}{ccc} x + \epsilon y & a \\ a & y + \epsilon x \end{array}$$

is in P<sub>2</sub>, and converges to  $xy - a^2$ .

**Corollary 11.7.** Suppose a > 0. axy + bx + cy + d is in  $\pm \overline{\mathbf{P}}_2$  iff  $\begin{vmatrix} a & b \\ c & d \end{vmatrix} \leqslant 0$ .

*Proof.* Let f(x,y) = axy + bx + cy + d. Since  $(1/a)f(x - c/a, y - b/a) = xy - (bc - ad)/a^2$  we can scale and apply Lemma 11.6.

The lemma can be derived from Theorem 10.75. Also see Question 145. It is also a special case of Lemma 11.28. The necessity is the rhombus inequality (Proposition 10.41).

**Lemma 11.8.** *If*  $xy + bx + cy + d \in \overline{\mathbf{P}}_2$  *then there are*  $\alpha$ ,  $\beta$ ,  $\gamma$  *such that* 

$$f = egin{bmatrix} lpha + x & eta \ eta & \gamma + y \end{bmatrix}$$

*Proof.* If we choose  $\gamma = b$  and  $\alpha = c$  then we find that  $\beta^2 = bc - d$ . This expression is positive by the previous lemma.

**Example 11.9.** Since  $\begin{vmatrix} x+1 & 3 \\ y-2 \end{vmatrix} = -(2x + 11) + y(x + 1) \in \overline{\mathbf{P}}_2$  we see that a multiaffine polynomial  $f + yg \in \overline{\mathbf{P}}_2$  can have f, g with different leading signs. Of course we still have  $f \leq g$ .

**Example 11.10.** There are simple polynomials that are not in  $\overline{\mathbf{P}}_2$ . For instance,  $x^2 \pm y$  is not in  $\overline{\mathbf{P}}_2$ . Indeed,  $f(x, y) = x^2 - y$  doesn't satisfy substitution because  $f(x, -1) \notin \mathbf{P}$ . Similarly  $x^2 + y$  doesn't satisfy substitution.

We can extend Lemma 11.6 by replacing x by xy + bx + cy + d instead of by xy. The next result expresses this extension, but uses a more convenient replacement.

**Corollary 11.11.** *If*  $f \in P$  *then the polynomial*  $f((x + \alpha)(y + \beta) + \gamma)$  *is in*  $\overline{P}_2$  *if all roots of* f *are either greater than*  $\gamma$ *, or all roots of* f *are less than*  $\gamma$ *.* 

If  $f \in \mathbf{P}_2^{\text{pos}}$  then f(x, 0) and f(0, x) have all positive coefficients, and all negative roots. Thus the graph of f meets the x and y axis in negative values. If we intersect the graph with the line x = y then the intersection points are all negative. With this in mind, we establish

**Lemma 11.12.** If  $f \in \overline{\mathbf{P}}_2^{\text{pos}}$  then  $f(-x^2, -x^2) \in \mathbf{P}$ .

*Proof.* By continuity of roots it suffices to assume that  $f \in \mathbf{P}_2^{\text{pos}}$ . The roots of f(-x, -x) are all positive, so we can take their square roots.

This is a special case of Lemma 11.105.

*Remark* 11.13. If f(x, y) satisfies x substitution, and  $f^H$  is nearly homogeneous, f is not necessarily in P<sub>2</sub>. For instance, if f(x, y) = 1 + x(x + y) then f satisfies x substitution, and the homogeneous part has the form

 $x \times$  (homogeneous polynomial with positive coefficients).

However,  $f \notin \overline{\mathbf{P}}_2$  since  $f(\mathbf{x}, \mathbf{x}) \notin \mathbf{P}$ .

Here's a simple fact whose proof requires several closure properties of  $\overline{\mathbf{P}}_2$ .

**Lemma 11.14.** If  $ax + by + c \in \overline{\mathbf{P}}_2$  then a and b have the same sign.

*Proof.* After scaling and translating, it suffices to show that that x - y is not in  $\overline{\mathbf{P}}_2$ . If it were the case that  $x - y \in \overline{\mathbf{P}}_2$  then we also know that  $x - y + 3 \in \overline{\mathbf{P}}_2$ . Since  $\overline{\mathbf{P}}_2$  is closed under multiplication and differentiation, it follows that

$$f(x,y) = \frac{\partial}{\partial x}(x-y)(x-y+3)(x+y) = 6x + 3x^2 - 2xy - y^2 \in \overline{P}_2$$

However,  $f(-1, y) = -3 + 2y - y^2$  has two complex roots.

The homogeneous part of a polynomial in  $\overline{\mathbf{P}}_2$  is not too different from the homogeneous part of a polynomial in  $P_2$ .

**Lemma 11.15.** If  $f \in \overline{\mathbf{P}}_2$  then all non-zero coefficients of  $f^H$  have the same sign. There are non-negative integers n, m and  $g \in \mathbf{P}^{pos}$  so that  $f^H = \pm x^n y^m G$  where G is the homogenization of g.

*Proof.* If  $f^H$  only has one term then the result is true. If not all coefficients are the same sign then we can write  $f^H = \cdots + rx^ay^{n-a} + sx^by^{n-b} + \cdots$  where a < b, rs < 0, and all coefficients of  $x^ry^{n-r}$  are zero for a < r < b. Let  $g = (\partial_x)^a (\partial_y)^{n-b} f$  so that  $g^H(x,y) = cy^{b-a} + dx^{b-a}$  where cd < 0.

Substituting y = 1 shows that b - a is either 1 or 2. If b - a = 1 then g = cy + dx + e, but this is not possible by the previous lemma. If b - a = 2 then we can write  $g = \alpha x^2 + \beta xy + \gamma y^2 + \cdots$  where  $\alpha$  and  $\gamma$  have opposite signs. After scaling x and y and perhaps multiplying by -1 we can assume that  $g = x^2 + \beta xy - y^2 + \cdots$ . If  $\beta > 0$  then  $\partial_y g = \beta x - 2y + (constant)$  which is impossible by the lemma. If  $\beta < 0$  then  $\partial_x g = 2x + \beta y + (constant)$  which is also impossible.

The second statement follows immediately from the first since  $f^{H} \in \mathbf{P}$ .  $\Box$ 

It is a consequence of the lemma that the polynomial  $f^{H}(x, 1)$  has all non-positive roots if  $f \in \overline{\mathbf{P}}_2$ .

If we take a product of linear factors where the coefficients of x and y don't all have the same sign, then the resulting product isn't in  $P_2$ . However, if the constant terms are all the same sign then the coefficients are in **P**.

**Corollary 11.16.** Suppose that  $b_i \in \mathbb{R}$ , and all  $d_i$  have the same sign. If we write

$$\prod_{i=1}^n (y+b_ix+d_i) = \sum f_i(x)y^i$$

then

- (1)  $f_i(x) \in P$  for i = 0, ..., n 1.
- (2)  $f_i(x)$  and  $xf_{i+1}(x)$  interlace, for i = 0, ..., n-2.

*Proof.* By switching signs of x and y if necessary we may assume that all the  $d_i$  are negative. Replace y by xy:

$$\prod_{i=1}^{n}(xy+b_{i}x+d_{i})=\sum f_{i}(x)x^{i}y^{i}$$

Since the determinant of each factor is  $\begin{vmatrix} 1 & b_i \\ 0 & d_i \end{vmatrix} = d_i$  which is negative, we can apply Corollary 11.7 to conclude that the product is in  $\overline{\mathbf{P}}_2$ . Consequently, adjacent coefficients interlace, which finishes the proof.

If the roots of a polynomial are all positive or all negative, then the coefficients are all non-zero. We use this simple idea to constrain where the roots of coefficients lie.

**Corollary 11.17.** *Suppose that*  $f, g \in P^{pos}$  *and* s < r*. If we define* 

$$f(xy - ry)g(xy - sy) = \sum h_i(x)y^i$$

*then*  $h_i(x) \in \mathbf{P}^{[s,r]}$ .

*Proof.* We can apply Corollary 11.7 to see that  $h_i(x) \in \mathbf{P}$ . If  $\alpha > r$  then all coefficients of  $f((\alpha - r)y)$  and  $g((\alpha - s)y)$  are positive. Thus, all coefficients of  $(f(xy - ry)g(xy - sy))(\alpha, y)$  are positive, and hence  $h_i(\alpha)$  is not zero.

If  $\alpha < s$  then the roots of  $f((\alpha - r)y)$  and  $g((\alpha - s)y)$  are positive. Again, all coefficients of  $(f(xy - ry)g(xy - sy))(\alpha, y)$  are non-zero, and hence  $h_i(\alpha)$  is not zero.

If we define  $f_0(x) = J_0(2\sqrt{-x})$ , where  $J_0$  is the Bessel function, then we have the identity [89, (2.1.A)]:

$$f_0(xy+y)f_0(xy-y) = \sum_{n=0}^{\infty} P_n(x) \frac{(2y)^n}{n!n!}$$
(11.1.1)

where  $P_n(x)$  is the Legendre polynomial. Since  $f_0$  has all positive coefficients, it follows that the Legendre polynomials have roots in [-1, 1], as is well known. We will also use this identity to determine mapping properties of a Legendre transformation - see Lemma 15.48.

Any polynomial in **P** determines a mutually interlacing sequence (see Example 3.39). The following lemma shows that we can find a polynomial in  $\overline{\mathbf{P}}_2$  that interpolates two of these mutually interlacing sequences. We will show that given polynomials f(x), g(x) with roots  $\{r_i\}$ ,  $\{s_i\}$ , we can find F(x, y) in  $\overline{\mathbf{P}}_2(n)$  that satisfies

$$F(x, s_i) = g'(s_i) f_{n+1-i}(x)$$
  

$$F(r_i, y) = f'(r_i) g_{n+1-i}(y)$$

**Lemma 11.18.** Suppose that  $f(x) = \prod (x - r_i)$ ,  $g(x) = \prod (x - s_i)$  where  $r_1 \leq \cdots \leq r_n$  and  $s_1 \leq \cdots \leq s_n$ . Then

$$F(x,y) = \sum_{i=1}^{n} \frac{f(x)}{x - r_i} \frac{g(y)}{y - s_{n+1-i}} \in \overline{\mathbf{P}}_2$$
(11.1.2)

$$G(x,y) = \sum_{i=1}^{n} \frac{f(x+y)}{x+y-r_i} \frac{g(x+y)}{x+y-s_{n+1-i}} \in P_2$$
(11.1.3)

*Proof.* If we replace f(x, y) by  $f(x, \epsilon y)$  and g(x, y) by  $g(\epsilon x, y)$  where  $\epsilon$  is positive, then (11.1.3) implies that

$$\sum_{i=1}^{n} \frac{f(x + \varepsilon y)}{x + \varepsilon y - r_{i}} \frac{g(y + \varepsilon x)}{\varepsilon x + y - s_{n+1-i}}$$

and letting  $\epsilon$  go to zero establishes (11.1.2). The homogeneous part of G(x, y) is  $n(x+y)^{2n}$ . To check substitution, we choose  $\alpha \in \mathbb{R}$  and let  $f_{\alpha}(x) = f(x+\alpha)$ ,  $g_{\alpha}(x) = g(x+\alpha)$ . The result now follows from 3.16.

If  $\sum f_i(x)y^i \in P_2$  then  $f_i$  and  $f_{i+1}$  interlace. This is half true for  $f_i$  and  $f_{i+2}$ .

**Lemma 11.19.** If  $\sum f_i(x)y^i \in P_2$  then  $f_i \stackrel{+}{\sim} -f_{i+2}$ .

*Proof.* Since  $y^2 - \alpha \in \overline{\mathbf{P}}_2$  for positive  $\alpha$ , the product

$$(y^2 - \alpha) \sum f_i(x) y^i = \sum (f_{i+2}(x) - \alpha f_i(x)) y^i$$

is in  $\overline{\mathbf{P}}_2$ . Thus  $f_{\mathfrak{i}+2}(x) - \alpha f_{\mathfrak{i}}(x) \in \mathbf{P}$  for all positive  $\alpha$ .

A Hadamard product with a polynomial in  $\mathbf{P}_2^{\text{pos}}$  is effectively one with a polynomial in  $\mathbf{P}^{\text{pos}}$ .

**Lemma 11.20.** If  $f = \sum_{0}^{n} f_{i}(x)y^{i} \in \mathbf{P}_{1}^{pos}$  has the property that  $\sum_{0}^{n} g_{i}(x)f_{i}(x)y^{i} \in \mathbf{P}_{2}^{pos}$  for all  $\sum g_{i}(x)y^{i} \in \mathbf{P}_{2}^{pos}$  then  $f = F(x) \cdot \sum_{0}^{n} a_{i}y^{i}$  where  $F(x) \in \mathbf{P}^{pos}$  and  $\sum a_{i} \frac{y^{i}}{i!(n-i)!} \in \mathbf{P}^{pos}$ .

*Proof.* If we let  $g = (y + 1)^n$  then

$$\sum f_{i}g_{i}y^{i} = \sum {\binom{n}{i}}f_{i}(x)y^{i} \in \mathbf{P}_{2}^{pos}$$

Applying f again yields

$$\sum \binom{n}{i} f_i^2(x) y^i \in \mathbf{P}_2^{\text{pos}}$$

Since adjacent coefficients interlace, Lemma 1.34 implies that  $f_i = F(x)a_i$  for some constant  $a_i$ . Thus

$$\sum {\binom{n}{i}} f_i^2(x) y^i = n! F(x) \sum a_i \frac{y^i}{i!(n-i)!} \in \mathbf{P}_2^{\text{pos}}$$

which proves the theorem.

#### **11.2** Reversal and cancellation in **P**<sub>2</sub>

In this section we first consider two properties that are trivial for polynomials in one variable. First, the reversal of a polynomial in **P** is in **P**, and second, if  $xf \in \mathbf{P}$  then  $f \in \mathbf{P}$ . Neither is immediate in  $P_2$ , and we do not know the latter to be true if there are more than two variables. Finally, we discuss differential operators that preserve  $P_d$ .

The key to proving that  $xy - 1 \in \overline{\mathbf{P}}$  was the identity

$$xy - 1 = \begin{vmatrix} x & 1 \\ 1 & y \end{vmatrix}$$

which can be modified to give a sequence of polynomials in P<sub>2</sub> converging to xy - 1. Now xy - 1 is the reversal of x + y; we have a similar matrix argument that shows that the reversal is in  $\overline{\mathbf{P}}_2$ . If we use 1/x instead of -1/x (See p. 270.) we get stable polynomials (See p. 620.). The key is the following well known matrix identity for the determinant of a partitioned matrix:

$$\begin{vmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{vmatrix} = \begin{vmatrix} A_{11} \end{vmatrix} \begin{vmatrix} A_{22} - A_{21} A_{11}^{-1} A_{12} \end{vmatrix}$$
(11.2.1)

It follows easily that for matrices A, B of the same size

$$|\mathbf{x}\mathbf{B} - \mathbf{A}^2| = \begin{vmatrix} \mathbf{x}\mathbf{I} & \mathbf{A} \\ \mathbf{A} & \mathbf{B} \end{vmatrix}$$

**Lemma 11.21.** If  $f(x, y) \in P_2(n)$ , then  $y^n f(x, -1/y) \in \overline{\mathbf{P}}_2$ . Equivalently, if  $f = \sum f_i(x)y^i \in P_2(n)$ , then  $\sum f_i(x)(-y)^{n-i} \in \overline{\mathbf{P}}_2$ .

*Proof.* If we write  $f(x, y) = |I + xD_1 + yD_2|$  where  $D_1, D_2$  are positive definite then

$$\begin{aligned} x^{n}f(-1/x,y) &= |xI - D_{1} + xyD_{2}| \\ &= |x(I + yD_{2}) - D_{1}| \\ &= \begin{vmatrix} xI & \sqrt{D_{1}} \\ \sqrt{D_{1}} & I + yD_{2} \end{vmatrix} \\ &= \begin{vmatrix} \begin{pmatrix} 0 & 0 \\ 0 & I \end{pmatrix} + x \begin{pmatrix} I & 0 \\ 0 & 0 \end{pmatrix} + y \begin{pmatrix} 0 & 0 \\ 0 & D_{2} \end{pmatrix} + \begin{pmatrix} 0 & \sqrt{D_{1}} \\ \sqrt{D_{1}} & 0 \end{pmatrix} \end{aligned}$$

This represents f as a determinant of a matrix of size 2n that is clearly the limit of determinants of matrices that are in  $P_2$ .

Remark 11.22. Here is an alternative geometric argument. Define

$$f_{\varepsilon}(x, y) = (y + \varepsilon x)^{n} f(x + \varepsilon y, -1/(y + \varepsilon x)),$$

where  $\epsilon$  is positive. Since  $\lim_{\epsilon \to 0} f_{\epsilon} = f$ , we need to show that  $f_{\epsilon} \in P_2(2n)$ . The homogeneous part of  $f_{\epsilon}$  is clearly  $(x + \epsilon y)^n (y + \epsilon x)^n$ , and so all coefficients of  $f_{\epsilon}^H$  are positive. If we fix y, then the locus of  $\{(x + \epsilon y, -1/(y + \epsilon x))\}$  as x varies consists of two hyperbolas that open in the second and fourth quadrants. Consequently, every solution curve of f meets each of these hyperbolas - see Figure 9.6. Thus, for any y, there are 2n solutions, and substitution is satisfied.

**Proposition 11.23.** *The two variable reversal of a polynomial in*  $P_2(n)$  *is in*  $\overline{P}_2(2n)$ *. The two variable reversal of* f *is* f<sup>REV</sup> =  $x^n y^n f(1/x, 1/y)$ .

*Proof.* If we apply the lemma twice we see that  $x^n y^n f(-1/x, -1/y) \in \overline{\mathbf{P}}_2$ . Since the substitution of (-x, -y) for (x, y) preserves  $P_2$ , the result follows.

Note that the reversal of x + y is xy - 1, whereas the two variable reversal of x + y is x + y.

The reverse of a polynomial in  $P_2$  satisfies the assumptions of the following lemma.

**Lemma 11.24.** Suppose that  $f(x, y) = \sum f_i(x)y^i \in \widehat{P}_2$  or  $\overline{P}_2$ , and each  $f_i(x)$  is a polynomial of degree *i*. If the leading coefficients of the  $f_i$  alternate in sign then

$$g(\mathbf{x},\mathbf{y}) \in P_2(\mathbf{n}) \implies f(\mathbf{x},-\partial_{\mathbf{y}}) g(\mathbf{x},\mathbf{y}) \in P_2(\mathbf{n})$$

*Proof.* If we choose  $\alpha \in \mathbb{R}$  then  $f(\alpha, y)$  is in  $\widehat{\mathbf{P}}$ . It follows from Lemma 9.92 that  $f(\alpha, -\partial_y) g(\alpha, y) \in \mathbf{P}$ , and so  $f(x, -\partial_y) g(x, y)$  satisfies y-substitution.

Suppose the leading coefficient of  $f_i(x)$  is  $c_i$ . Since the sign of  $c_i$  alternates the homogeneous part of  $f_i(x)(-\partial_y)^i g(x,y)$  has degree n, and equals  $c_i(-x)^i(\partial_y^i g)^H$ . The homogeneous part of  $f(x, -\partial_y) g(x, y)$  has all positive signs (or all negative), and so the first part is proved. The second part is similar.

We need a lemma about truncating polynomials in **P** before we can proceed to canceling x in  $\overline{\mathbf{P}}_2$ .

**Lemma 11.25.** Suppose that  $f \in P(m-1)$  does not have multiple roots. If  $f_n \longrightarrow xf$  where all  $f_n \in P(m)$  then  $f_n(x) - f_n(0) \in P$  for n sufficiently large.

*Proof.* If  $r_1, \ldots, r_{m-1}$  are the roots of f', then none of  $f(r_1), \ldots, f(r_{m-1})$  are zero since f has no multiple roots. Let  $\alpha$  be the minimum value of  $|f(r_1)|, \ldots, |f(r_{m-1})|$ . Since  $f_n(0)$  converges to 0 there is an N such that if n > N we have  $|f_n(0)| < \alpha/2$ . And, if we choose N sufficiently large we can require that the minimum of  $\{|f_n(r)| | f'(r) = 0\}$  is at least  $\alpha/2$ . Notice that the graph of  $f_n(x) - f_n(0)$  is just the graph of  $f_n(x)$  shifted by less than  $\alpha/2$ . Since all the critical points of  $f_n(x)$  are more than  $\alpha/2$  away from the x axis,  $f_n(x)$  has all real roots.

**Proposition 11.26.** *If*  $xf \in \overline{\mathbf{P}}_2$  *then*  $f \in \overline{\mathbf{P}}_2$ *.* 

*Proof.* Since  $xf \in \overline{\mathbf{P}}_2(m)$  there are polynomials  $h_n(x, y) \in P_2(m)$  such that  $h_n \longrightarrow xf$ . We can write  $h_n(x, y) = x g_n(x, y) + k_n(y)$ . Since  $g_n \longrightarrow f$  it suffices to show that  $g_n \in P_2$ . Now  $h_n$  satisfies the homogeneity condition,

and therefore so does  $g_n$ . We need to show that  $g_n$  satisfies substitution. If we choose  $\alpha \in \mathbb{R}$  then  $xg_n(y, \alpha) + k_n(\alpha) \longrightarrow xf(y, \alpha)$ . The lemma above implies that  $g_n(y, \alpha) \in \mathbf{P}$  for  $n \ge N_{\alpha}$ . Now we can apply Lemma 11.118 to conclude that there is an N (the maximum of all the  $N_{\alpha}$ 's corresponding to roots of the resultant) such that n > N implies that  $g_n(y, \alpha) \in \mathbf{P}$  for all  $\alpha \in \mathbb{R}$ .

We would like to prove the converse of this.

**Lemma 11.27.** If  $f(\partial_x, \partial_y)g$  is in  $\overline{\mathbf{P}}_2$  for all  $g \in \overline{\mathbf{P}}_2$  then f is in  $\overline{\mathbf{P}}_2$ .

Proof. Consider the calculation

$$\begin{split} f(x,y) &= \sum_{0 \leqslant i,j \leqslant n} a_{ij} x^i y^j \\ f(\partial_x, \partial_y) x^R y^R &= \sum_{0 \leqslant i,j \leqslant n} a_{ij} (\underline{R})_i (\underline{R})_j x^{R-i} y^{R-j} \\ &= x^{R-n} y^{R-n} \sum_{0 \leqslant i,j \leqslant n} a_{ij} (\underline{R})_i (\underline{R})_j x^{n-i} y^{n-j} \end{split}$$

Replacing x by x/R and y by y/R yields

$$x^{R-n}y^{R-n}\sum_{0\leqslant i,j\leqslant n}a_{ij}\frac{(\underline{R})_i}{R^i}\frac{(\underline{R})_j}{R^j}x^{n-i}y^{n-j}\in\overline{P}_2$$

and so we can divide out the initial factors to conclude that

$$\sum_{0 \leqslant i,j \leqslant n} a_{ij} \frac{(\underline{R})_i}{R^i} \frac{(\underline{R})_j}{R^j} x^{n-i} y^{n-j} \in \overline{P}_2$$

Now letting  $R \to \infty$  shows that

$$\sum_{0\leqslant i,j\leqslant n}\mathfrak{a}_{ij}x^{n-i}y^{n-j}\in\overline{P}_2$$

Taking the reversal of the last expression and applying Proposition 11.23 finishes the proof.  $\hfill \Box$ 

## **11.3 Polynomials linear in** y

The polynomials of the form f(x)+yg(x) are well understood. First of all, since  $f + \alpha g$  is in **P** for all  $\alpha$ , it follows that f and g interlace, and their degrees differ by at most 1. If f and g have the same degree then we can write  $f = \beta g + h$  where  $g \leq h$ , so that  $f + yg = h + (y + \beta)g$ . Substituting  $y = y + \beta$  shows that we can always assume that the degrees differ by one. The next two lemmas show that the condition of interlacing is all we need to belong to  $\overline{\mathbf{P}}_2$ .

**Lemma 11.28.** *If*  $f \leftarrow g \in P$  *have positive leading coefficients then the polynomial* f + yg *lies in*  $\overline{P}_2$ .

*Proof.* If  $f \leftarrow g$  then we can write

$$f = (ax+b)g - \sum_{1}^{n} c_{i}^{2} \frac{g}{x+a_{i}}$$

where  $g = \prod (x + a_i)$  and a is non-negative. Since f + yg can be expressed as the determinant

y + ax + b	$c_1$	$c_2$	• • •	cn
$c_1$	$x + a_1$	0		0
c <sub>2</sub>	0	$x + a_2$		0
:	:		•.	:
c <sub>n</sub>	0			$x + a_n$

it follows that  $f + yg \in \overline{\mathbf{P}}_2$ .

**Lemma 11.29.** If  $f \leftarrow g \in P$  have positive leading coefficients then the polynomial yf - g lies in  $\overline{P}_2$ .

*Proof.* The reversal of f + yg is yg - f, so  $yg - f \in \overline{\mathbf{P}}_2$ .

*Remark* 11.30. If we choose  $f_0 \leq f_1$  and  $g_0 \leq g_1$  then both  $f_0 + yf_1$  and  $g_0 + yg_1$  are in  $\overline{\mathbf{P}}_2$ . Consequently their product

$$(f_0 + yf_1)(g_0 + yg_1) = f_0g_0 + (f_0g_1 + f_1g_0)y + f_1g_1y^2$$

is in  $\overline{\mathbf{P}}_2$ , and so the coefficients of the product interlace. This explains the Leibnitz property (Lemma 1.26):

$$f_0g_0 \underline{\lessdot} f_0g_1 + f_1g_0 \underline{\lessdot} f_1g_1.$$

The following lemma generalizes the Leibnitz property.

**Lemma 11.31.** Suppose  $f_i \longleftarrow g_i$  have positive leading coefficients for  $1 \le i \le n$  and define

$$h_k = \sum_{|I|=k} f_I g_{\{1,\dots,n\}\setminus I}$$

Then

 $h_1 \longleftarrow h_2 \longleftarrow \cdots \longleftarrow h_n$ 

*Proof.* The  $h_k$ 's are consecutive coefficients of  $\prod_i (f_i + yg_i)$ .

**Example 11.32.** In the case n = 3 the interlacings are

$$f_1f_2f_3 \longleftarrow f_1f_2g_3 + f_1g_2f_3 + g_1f_2f_3 \longleftarrow f_1g_2g_3 + g_1f_2g_3 + g_1g_2f_3 \longleftarrow g_1g_2g_3.$$

The first and last interlacings are easy, but the middle one is not so obvious.

379

Here is a similar result for common interlacings.

**Lemma 11.33.** If  $f_1, \ldots, f_n$  are in **P** and have a common interlacing then

$$\sum_{\mathfrak{i}_1 < \cdots < \mathfrak{i}_s} \mathfrak{f}_{\mathfrak{i}_1} \mathfrak{f}_{\mathfrak{i}_2} \cdots \mathfrak{f}_{\mathfrak{i}_s} \in \boldsymbol{P}$$

*Proof.* If  $g \leq f_i$  then  $g + yf_i \in \overline{\mathbf{P}}_2$  and therefore  $\prod_{i=1}^n (g + yf_i) \in \overline{\mathbf{P}}_2$ . The coefficient of  $y^s$  is in  $\mathbf{P}$  and is  $g^{n-s}$  times the desired polynomial.

**Corollary 11.34.** Suppose  $f = f_1(x) + yf_2(x)$  and  $g = g_1(x) + yg_2(x)$  are in  $\overline{\mathbf{P}}_2$  where  $f_1, f_2, g_1, g_2$  have positive leading coefficients. Then  $f \leq g$  iff

- 1.  $f_1 \lessdot g_1$
- 2.  $f_2 \lessdot g_2$

3. 
$$\left| \begin{array}{c} f_1 & f_2 \\ g_1 & g_2 \end{array} \right| < 0$$

Proof. This follows from Lemma 1.51 and Lemma 1.52.

There is no known criterion for a polynomial that is quadratic in y to belong to  $\overline{\mathbf{P}}_2$ . The following lemma gives a general construction for such polynomials, but it is not known if there are quadratics in y in  $\overline{\mathbf{P}}_2$  that are not given by this form.

**Lemma 11.35.** Suppose that  $f = (x + c_1) \cdots (x + c_n)$ ,  $f_i = \frac{f}{x + c_i}$ , and  $f_{ij} = \frac{f}{(x + c_i)(x + c_j)}$ . For any choice of  $a_i$ ,  $b_i$  the following polynomial is in  $\overline{\mathbf{P}}_2$ :

$$y^{2}f - y\left(\sum_{i} (a_{i}^{2} + b_{i}^{2})f_{i}\right) + \sum_{i < j} \left| \begin{array}{c} a_{i} & a_{j} \\ b_{i} & b_{j} \end{array} \right|^{2} f_{ij}$$

*Proof.* It suffices to show that the polynomial in question is the value of the determinant

$$\begin{vmatrix} y & 0 & a_1 & \dots & a_n \\ 0 & y & b_1 & \dots & b_n \\ a_1 & b_1 & x + c_1 & 0 \\ \vdots & \vdots & & \ddots & \vdots \\ a_n & b_n & 0 & \dots & x + c_n \end{vmatrix}$$
(11.3.1)

The coefficient of  $y^2$  is clearly f. To find the coefficient of y, consider the minor corresponding to the first row and column:

We have seen that this determinant has value  $yf - \sum b_i^2 f_i$ . The yf term comes from the y<sup>2</sup>f term in (11.3.1), and the remaining part is part of the coefficient of y. Similarly, the value of the minor corresponding to the y in the second row and column of (11.3.1) is  $yf - \sum a_i^2 f_i$ . Combining these gives the coefficient of y.

The constant term of the determinant (11.3.1) is the value of the determinant

0	0	$\mathfrak{a}_1$	• • •	a <sub>n</sub>
0	0	$\mathfrak{b}_1$		b <sub>n</sub>
<b>a</b> <sub>1</sub>	$\mathfrak{b}_1$	$\mathbf{x} + \mathbf{c}_1$		0
:	:		·	:
an	b <sub>n</sub>	0		$x + c_n$

Expanding using the first two rows (and some patience) gives the final result.

Here is another interpretation of the last result.

**Corollary 11.36.** Suppose  $f \in P(n)$ ,  $f \leq g$ ,  $f \leq h$ , and the leading coefficients of f, g, h are all positive. There is a  $k \in P(n-2)$  so that

$$\mathbf{k} - \mathbf{y}(\mathbf{g} + \mathbf{h}) + \mathbf{y}^2 \mathbf{f} \in \overline{\mathbf{P}}_2$$

*Proof.* Use Lemma 1.20 to write  $g = \sum a_i^2 f_i$  and  $h = \sum b_i^2 f_i$ , and now apply Lemma 11.35.

We can generalize this corollary to three variables.

**Lemma 11.37.** Suppose  $f \in P(n)$ ,  $f \leq g$ ,  $f \leq h$ , and the leading coefficients of f, g, h are all positive. There is a  $k \in P(n-2)$  so that

$$k-yg+zh+yzf\in \overline{P}_3$$

Proof. Following the evaluation in Lemma 11.35, we find that

$$F(x, y, z) = \begin{vmatrix} z & 0 & a_1 & \dots & a_n \\ 0 & y & b_1 & \dots & b_n \\ a_1 & b_1 & x + c_1 & 0 \\ \vdots & \vdots & & \ddots & \vdots \\ a_n & b_n & 0 & \dots & x + c_n \end{vmatrix}$$
$$= yz f - z h - y g + \sum_{i < j} \left| \begin{array}{c} a_i & a_j \\ b_i & b_j \end{array} \right|^2 f_{ij}$$

where f, g, h are as in the proof of Lemma 11.35. The polynomial  $k = \sum_{i < j} \left| \begin{array}{c} a_i & a_j \\ b_i & b_j \end{array} \right|^2 f_{ij}$  is in **P** since it is just F(x, 0, 0).

If we take g = h then k is zero, and so  $yz f - (y + z)g \in \overline{\mathbf{P}}_2$ . If we compute the determinant

an	an	an	0		$x + c_n$
:	÷	÷		·.	÷
<b>a</b> <sub>1</sub>	$\mathfrak{a}_1$	$\mathfrak{a}_1$	$\mathbf{x} + \mathbf{c}_1$		0
0	0	w	$\mathfrak{a}_1$		an
0	y	0	$\mathfrak{a}_1$		an
z	0	0	$a_1$		an

then we get yzwf - (yz - yw - zw)g which is therefore in  $\overline{\mathbf{P}}_4$ . More generally, if we compute the determinant of

$$\begin{vmatrix} \operatorname{diag}(y_i) & (A, \dots, A)^t \\ (A^t, \dots, A^t) & \operatorname{diag}(x + c_i) \end{vmatrix}$$

where  $A = (a_1, \ldots, a_n)$  then we find that

$$(\mathfrak{y}_1\cdots\mathfrak{y}_n)\,\mathsf{f}-\mathfrak{g}\left(\sum_{\mathfrak{j}}\frac{\mathfrak{y}_1\cdots\mathfrak{y}_n}{\mathfrak{y}_\mathfrak{j}}
ight)\in\overline{\mathbf{P}}_{n+1}.$$

One way to create polynomials of a fixed degree in y is to use positive semi-definite matrices.

**Lemma 11.38.** If  $D_1$  is an n by n positive definite matrix, and  $D_2$  is positive semidefinite of rank r, then  $|I + xD_1 + yD_2| \in \overline{P}_2$ , and has degree r in y.

*Proof.* If we diagonalize  $D_2$ , then there are exactly r non-zero terms on the diagonal, so the maximum degree is r. The coefficient of  $y^r$  is the determinant of the principle submatrix of  $I+xD_1$  corresponding to the zero diagonal entries of  $D_2$ , and is a polynomial in  $\mathbf{P}^{pos}$  of degree n - r.

**Example 11.39.** If  $v_1, \ldots, v_d$  are vectors then  $\left| \sum_{i}^{n} x_i v_i^{t} v_i \right|$  is linear in each variable since  $v_i^{t} v_i$  is a semi-definite rank 1 matrix.

**Example 11.40.** There is a simple formula if  $D_2$  has rank 1. If v is a vector of length n, A is n by n, and f = |I + xD| then

$$|I + vv^{t}| = 1 + v^{t}v$$
(11.3.2)  

$$|I + yvv^{t}| = 1 + yvv^{t}$$
  

$$|A + yvv^{t}| = |A|(1 + yv^{t}A^{-1}v)$$
  

$$|I + xD + yvv^{t}| = f(1 + yv^{t}(I + xD)^{-1}v)$$

If  $D = diag(d_i)$  then

$$|I+xD+y\nu\nu^t|=f+y\sum\nu_i^2\frac{f}{1+xd_i}$$

Since the coefficient of y interlaces the coefficient of f, we have another proof of part of Lemma 1.20. If f < g then  $g = \sum a_i f/(1 + xd_i)$  where  $a_i > 0$ . If we set  $w = (\sqrt{a_i})$  then

$$|\mathbf{I} + \mathbf{x}\mathbf{D}_1 + \mathbf{y}\mathbf{w}\mathbf{w}^{\mathsf{t}}| = \mathbf{f} + \mathbf{y}\mathbf{g}$$

This is a different determinant realization of f + yg (See p. 379.).

## 11.4 Constrained polynomials with zero coefficients

We look at some conditions on polynomials in  $\overline{\mathbf{P}}_2$  that imply they have a simple form. First of all, there are no non-trivial ways of adding one variable polynomials to get a polynomial in  $\overline{\mathbf{P}}_2$ .

**Lemma 11.41.** If  $f(x) + g(y) \in \overline{P}_2$  then at least one of the following holds

- 1. f or g is constant.
- 2. f and g have degree 1.

*Proof.* Assume that neither f nor g is constant. If f is linear then it takes arbitrarily large positive and negative values, so by Lemma 1.12 g is linear. If both f and g have degree at least 2 then the lemma implies that they have degree exactly two. Moreover, if f has positive leading coefficient then g has negative leading coefficient, but this is impossible for then  $(f(x) + g(y))^H$  does not have all coefficients of the same sign.

Linear combinations of the form f(x) + g(x)h(y) are trivial if deg(h) > 1.

**Lemma 11.42.** If  $f(x) + g(x)h(y) \in \overline{P}_2$  where deg(h) > 1 then g is a constant multiple of f.

*Proof.* If deg(h) > 2 the set  $\{r \in \mathbb{R} \mid h(y) + r \in P\}$  is bounded. Since  $h(y) + f(r)/g(r) \in P$  for all r such that  $g(r) \neq 0$  we have that

$$\left\{\frac{f(r)}{g(r)} \mid g(r) \neq 0\right\}$$

is bounded. Thus f/g extends to a bounded rational function on  $\mathbb{R}$  which is constant by Liouville's theorem.

If the degree of h is 2 then we may assume that f and g have no common factors. If g has a root then f/g takes on arbitrarily large positive and negative values, so by Lemma 1.12 h is linear. Thus g is constant, but this contradicts Lemma 11.41 since h has degree 2.
If  $\dots + a x^i + 0 x^{i+1} + b x^{i+2} + \dots \in \mathbf{P}$  then Newton's inequality implies that  $ab \leq 0$ . The analogous result in two variables is much stronger.

**Lemma 11.43.** If  $f(x, y) = \cdots + f_i(x) y^i + 0 y^{i+1} + f_{i+2}(x) y^{i+2} + \cdots \in \overline{P}_2$  and  $f_i$  and  $f_{i+2}$  are not zero then there are  $g, h \in P$  so that f(x, y) = g(x)h(y).

*Proof.* If we differentiate, reverse and differentiate with respect to y sufficiently many times we find that there are constants  $\alpha$ ,  $\beta$  so that  $\alpha f_i(x) + \beta f_{i+2}(x) y^2 \in P_2$ . By Lemma 11.42 we see that  $f_{i+2}$  is a constant multiple of  $f_i$ .

We can thus write

$$f(x, y) = \cdots + f_{i-1}(x) y^{i-1} + f_i(x) (a + b y^2) y^i + \cdots$$

for constants a, b. Differentiating and reversing yields constants  $c_i$  so that

 $c_{\mathfrak{i}-1}\,f_{\mathfrak{i}-1}(x)\,y^{\mathfrak{i}-1}+f_{\mathfrak{i}}(x)\,(c_{\mathfrak{i}}+c_{\mathfrak{i}+1}\,y^2)\,y^{\mathfrak{i}}\in\overline{P}_2$ 

Applying the lemma again shows that  $f_{i-1}$  is a multiple of  $f_i$ . Continuing, we see that we can write

$$f(x, y) = f_i(x) H(y) + f_{i+1}(x) y^{i+1} + \cdots$$

Reversing, we apply the same argument to finish the proof.

**Corollary 11.44.** *If*  $f \in \overline{P}_2$  *has all exponents of even degree, then* f(x, y) = g(x)h(y) *where*  $g, h \in P$ .

*Proof.* If we write  $f(x, y) = \sum f_i(x)y^i$  then if i is odd the hypotheses imply that  $f_i(x) = 0$ . The conclusion now follows from the lemma.

# **11.5** The polynomial closure of $P_d$

If V is the vector space of a polynomials in  $x_1, \ldots, x_d$  of total degree at most n, then  $P_d(n)$  is contained in V. The closure of  $P_d(n)$  in V is denoted  $\overline{P}_d(n)$ , and the closure of  $P_d^{pos}$  is  $\overline{P}_d^{pos}$ . We define

$$\overline{\mathbf{P}}_{d} = \overline{\mathbf{P}}_{d}(1) \cup \overline{\mathbf{P}}_{d}(2) \cup \dots$$

In this section we show that most properties of  $P_d$  apply to its closure. We also identify some specific polynomials in the closure that are not in  $P_d$ .

First of all, all monomials are in the closure.

**Lemma 11.45.** All monomials of at most d variables are in  $\overline{\mathbf{P}}_d$ .

*Proof.* The monomial  $x_1^{i_1} \cdots x_d^{i_d}$  is the limit of these polynomials in  $P_d$ :

$$(\mathbf{x}_1 + \boldsymbol{\varepsilon}(\mathbf{x}_2 + \cdots + \mathbf{x}_d))^{\boldsymbol{\iota}_1} \cdots (\boldsymbol{\varepsilon}(\mathbf{x}_1 + \cdots + \mathbf{x}_{d-1}) + \mathbf{x}_d)^{\boldsymbol{\iota}_d}.$$

Next, we have some important containments.

**Lemma 11.46.** If d < e then  $\overline{P}_d \subset \overline{P}_e$ . In the case d = 1 we have  $P \subset \overline{P}_2$ ,  $P^{\text{pos}} \subset \overline{P}_2^{\text{pos}}$ .

*Proof.* It suffices to assume e = d + 1. Choose  $f \in \overline{P}_d(n)$ . Since f is in the closure of  $P_d(n)$  there is a sequence  $f_1, f_2, \ldots$  of polynomials in  $P_d(n)$  that converge to f. For any positive  $\epsilon$  we know that  $f_i(x_1 + \epsilon x_{d+1}, x_2, \ldots, x_d)$  is in  $P_{d+1}$ . Since  $\lim_{\epsilon \to 0} f_i(x_1 + \epsilon x_{d+1}, x_2, \ldots, x_d) = f_i$  we see that f is the limit of polynomials in  $P_{d+1}$ . It is clear that if the homogeneous part of f has all positive coefficients, then so does the homogeneous part of

 $f(x_1 + \varepsilon x_{d+1}, x_2, \dots, x_d)$ . This implies that  $\overline{\mathbf{P}}_d \subset \overline{\mathbf{P}}_e$ .

If  $f \in \mathbf{P}(n)$  then  $f(x + \epsilon y)$  has homogeneous part  $c(x + \epsilon y)^n$  and so all  $f(x + \epsilon y)$  are in P<sub>2</sub>. If  $f \in \mathbf{P}^{pos}$  then all its coefficients are positive, and so the same is true of  $f(x + \epsilon y)$ .

We can use determinants to realize members of  $\overline{\mathbf{P}}_d$ . The proof is the same as Lemma 11.5.

**Lemma 11.47.** Suppose that C is a symmetric matrix, and D is a diagonal matrix where the i-th diagonal entry is  $\sum d_{ij}x_i$  where  $d_{ij}$  is non-negative. The polynomial det(C + D) is in  $\overline{P}_d$ .

Before we go too far, let's see that there are polynomials that are *not* in  $\overline{\mathbf{P}}_d$ . Consider this simple consequence of Lemma 10.59 :

**Lemma 11.48.** *If* f(x, ..., x) *does not have all real roots then*  $f \notin \overline{\mathbf{P}}_d$ .

This implies that xyz + 1 is not in  $\overline{\mathbf{P}}_3$ , because  $\pm x^3 + 1$  does not have all real roots. We can slightly generalize the lemma to say that if we replace some of the variables by  $\pm x$  and the resulting polynomial is not in  $\overline{\mathbf{P}}_d$  then neither is f. This shows that xy + z is not in  $\overline{\mathbf{P}}_3$  since  $\pm x^2 + z$  is not in  $\overline{\mathbf{P}}_2$  by Example 11.10.

There are simple polynomials in  $\overline{\mathbf{P}}_3$ . Since

$$\begin{vmatrix} x & 1 & 0 \\ 1 & y & 1 \\ 0 & 1 & z \end{vmatrix} = xyz - x - z$$

it follows from Lemma 11.47 that  $xyz - x - z \in \overline{\mathbf{P}}_3$ .

**Lemma 11.49.** The elementary symmetric functions are in  $\overline{\mathbf{P}}$ .

*Proof.* The i-th elementary symmetric function of  $x_1, \ldots, x_d$  is the coefficient of  $y^i$  in the product

$$(\mathbf{y} + \mathbf{x}_1)(\mathbf{y} + \mathbf{x}_2) \cdots (\mathbf{y} + \mathbf{x}_d)$$

Since each of the factors is in  $\overline{\mathbf{P}}_{d+1}$ , the product is in  $\overline{\mathbf{P}}_{d+1}$ , and so the coefficients of y are in  $\overline{\mathbf{P}}_d$ .

As long as there is no obvious restriction, the interior of  $\mathbf{Sub}_d$  is contained in  $\overline{\mathbf{P}}_d$ .

**Lemma 11.50.** *If*  $f \in \text{int } \mathbf{Sub}_d$  *and*  $f^H$  *has all non-negative terms then*  $f \in \overline{\mathbf{P}}_d$ .

*Proof.* With these assumptions on f the reason that f might not be in  $P_d$  is that  $f^H$  might have some zero coefficients. If f has total degree n then we write

$$\mathsf{f} = \sum_{I} c_{I} x^{I} \qquad \qquad \mathsf{g} = \sum_{|I| = n, c_{I} = 0} x^{I}$$

For any  $\alpha > 0$  the polynomial  $f + \alpha g$  satisfies the homogeneity condition. Let 0 be an open neighborhood in  $\mathbf{Sub}_d(n)$  that contains f. For sufficiently small  $\alpha$  we have that  $f + \alpha g$  is in 0, so for these  $\alpha$  it follows that  $f + \alpha g$  satisfies substitution. Since the homogeneous part of  $f + \alpha g$  has all positive coefficients  $f + \alpha g$  is in  $\mathbf{P}_d(n)$ , and converge to f.

Recall the definition of interlacing:

f, g  $\in \overline{\mathbf{P}}_d$  interlace iff f +  $\alpha$ g is in  $\overline{\mathbf{P}}_d$  for every  $\alpha$ .

The following are all consequences of the corresponding results for  $P_d$ .

**Theorem 11.51.** Suppose  $f \in \overline{\mathbf{P}}_d$ .

- 1. If  $g \in \overline{\mathbf{P}}_e$  then  $fg \in \overline{\mathbf{P}}_{d+e}$ .
- 2. If  $f \leq g$ , h and g,  $h \in \overline{\mathbf{P}}_d$  then  $f \leq g + h$  and  $g + h \in \overline{\mathbf{P}}_d$ .
- 3. f and  $\frac{\partial f}{\partial x_i}$  interlace.
- 4. Linear transformations extend from  $P \longrightarrow P$  to  $\overline{P}_d \longrightarrow \overline{P}_d$ .
- 5. Polynomial limits of polynomials in  $\overline{\mathbf{P}}_{d}$  are in  $\overline{\mathbf{P}}_{d}$ .
- 6. If  $f = \sum a_I(\mathbf{x})\mathbf{y}$  then  $a_I(\mathbf{x}) \in \overline{\mathbf{P}}_d$

We can generalize Lemma 11.28 to  $\mathbf{P}_d$ , but we no longer have a nice determinant representation.

**Lemma 11.52.** If  $f \leq g \in \mathbf{P}_d$  then  $f + x_{d+1}g \in \overline{\mathbf{P}}_{d+1}$ .

*Proof.* To simplify notation we assume that d = 3. Define a polynomial in four variables

$$F_{\varepsilon} = f(x + \varepsilon u, y + \varepsilon u, z + \varepsilon u) + ug(x + \varepsilon u, y + \varepsilon u, z + \varepsilon u).$$

Since  $\lim_{\varepsilon\to 0}F_\varepsilon=f+ug$  it suffices to verify that  $F_\varepsilon\in P_4.$  The homogeneous part satisfies

$$F_{\epsilon}^{H} = f^{H}(x + \epsilon u, y + \epsilon u, z + \epsilon u) + ug^{H}(x + \epsilon u, y + \epsilon u, z + \epsilon u).$$

Since both  $f^H$  and  $g^H$  have all positive terms, it follows that  $F_{\epsilon}^H$  also does. It remains to check substitution. Setting  $y = \alpha, z = \beta, u = \gamma$  we need to show that

$$F_{\varepsilon}(x, \alpha, \beta, \gamma) = f(x + \varepsilon \gamma, \alpha + \varepsilon \gamma, \beta + \varepsilon \gamma) + \gamma g(x + \varepsilon \gamma, \alpha + \varepsilon \gamma, \beta + \varepsilon \gamma)$$

is in **P**. Since  $f \leq g$  in **P**<sub>3</sub> it follows that

$$f(w, \alpha + \epsilon \gamma, \beta + \epsilon \gamma) \leq g(w, \alpha + \epsilon \gamma, \beta + \epsilon \gamma)$$

Since any linear combination of these two polynomials is in **P** we add  $\gamma$  times the second one to the first and replace *w* by  $x + \epsilon \gamma$  to finish the proof.

If we don't introduce a new variable then the result is easier.

**Corollary 11.53.** If  $f \leq g$  in  $\mathbf{P}_d$  then  $x_ig + f \in \overline{\mathbf{P}}_d$  for  $1 \leq i \leq d$ .

Proof. Use Lemma 10.116

The polynomial f+yg+zh+yzk is in  $\overline{\mathbf{P}}_3$  if f, g, h, k satisfy certain interlacing assumptions. Here's a more general result.

**Lemma 11.54.** If  $\mathbf{y} = (y_1, \dots, y_d)$  and  $\mathbf{F} = \sum_I f_I(\mathbf{x}) g_I(\mathbf{y})$  satisfies

- 1. x-substitution
- 2. deg  $(f_{00\dots 0}(\mathbf{x})) \ge deg (f_I(\mathbf{x})) + deg (g_I(\mathbf{y}))$  for all I
- 3. all  $f_I$  and  $g_I$  have all non-negative coefficients

*then*  $F \in \overline{\mathbf{P}}_{d+1}$ 

*Proof.* The notation  $f_{0...0}$  refers to the terms that contain no  $y_i$ 's. Let  $G_{\varepsilon} = F(x + \varepsilon y, y)$  where  $y = y_1 + \cdots + y_d$  and  $\varepsilon > 0$ . If  $n = deg(f_{0...0}(x))$  then condition two implies that  $n \ge deg(f_1(x + \varepsilon y)g_I(y))$  so the degree of  $G_{\varepsilon}$  is n. The homogeneous part of  $G_{\varepsilon}$  includes a term  $(x + \varepsilon y)^n$ , and the third condition guarantees that all other contributions are non-negative, so  $G_{\varepsilon}$  has all positive terms. If we let  $\mathbf{a} = (a_1, \ldots, a_d)$  then  $G_{\varepsilon}(x, \mathbf{a}) = F(x + \varepsilon a, \mathbf{a})$  where  $a = a_1 + \cdots + a_d$ . Now  $F(x, \mathbf{a})$  has all real roots by the first condition, and so the translate by  $\varepsilon a$  is also in  $\mathbf{P}$ , and so  $G_{\varepsilon}(x, \mathbf{a})$  satisfies x-substitution. Since  $\lim_{\varepsilon \to 0} G_{\varepsilon} = F$ , it follows that  $F \in \overline{\mathbf{P}}_{d+1}$ .

Corollary 11.55. Suppose that

$$F(x, y, z) = f_{00}(x) + yf_{10}(x) + zf_{01}(x) + yzf_{11}(x)$$

satisfies

- 1.  $F(x, a, b) \in P$  for all  $a, b \in \mathbb{R}$ .
- 2.  $f_{00} \in P^{pos}$ .
- 3.  $\deg(f_{00}) \ge \deg(f_{10}) + 1$
- 4.  $deg(f_{00}) \ge deg(f_{01}) + 1$
- 5.  $\deg(f_{00}) \ge \deg(f_{11}) + 2$ .
- 6. All f<sub>ij</sub> have non-negative coefficients

*then*  $F \in \overline{\mathbf{P}}_3$ .

*Remark* 11.56. We can multiply the Taylor series and extract coefficients to derive some properties of derivatives (Theorem 9.39, Corollary 9.93). Starting with  $f \in P_2$  and writing

$$\begin{split} f(x+u,y+\nu) &= f(x,y) + f_x(x,y)u + f_y(x,y)\nu + \\ &\quad f_{xx}(x,y)\frac{u^2}{2} + f_{xy}(x,y)u\nu + f_{yy}(x,y)\frac{y^2}{2} + .\,. \end{split}$$

we first multiply by  $u + \alpha$ . The coefficient of u is  $\alpha f_x + f$ . Since this is in  $\overline{\mathbf{P}}_2$  for all  $\alpha$  we conclude  $f \leq f_x$ .

Next, choose  $\alpha > 0$  and multiply by  $u^2 - 2\alpha \in \overline{\mathbf{P}}_4$ . The coefficient of  $u^2$  is  $\alpha f_{xx} - f$ , so we conclude that  $f - \alpha f_{xx} \in \mathbf{P}_2$  for all positive  $\alpha$ . If instead we multiply by  $1 - \alpha uv \in \mathbf{P}_2$  we find that the coefficient  $f - \alpha f_{xy}$  of uv is in  $\mathbf{P}_2$ .

Since (u + 1)(v + 1) - 1 = uv + u + v is in  $\overline{\mathbf{P}}_2$ , the coefficient  $f_x + f_y + f_{xy}$  of uv is in  $\mathbf{P}_2$ .

In general, if  $g(u, v) = \sum a_{rs} u^r v^s \in \overline{\mathbf{P}}_2$  then

$$f(x + u, y + v) = \sum D_x^i D_y^j f \frac{u^i v^j}{i! j!}$$
$$f(x + u, y + v)g(u, v) = \sum a_{rs} D_x^i D_y^j f \frac{u^{i+r}}{i! j!}$$

The coefficient of  $v^n u^m$  is

$$\sum \frac{a_{n-i,m-j}}{i!j!}\mathsf{D}_x^i\mathsf{D}_y^j\mathsf{f}$$

and is in  $P_2$ .

**Lemma 11.57.** Suppose that  $f(x_1, ..., x_d) = \sum \mathbf{a}_I \mathbf{x}^I$  is a non-zero polynomial, and *define* 

$$g(\mathbf{x}) = f(\varepsilon_{10} + \sum \varepsilon_{1j} x_j, \dots, \varepsilon_{d0} + \sum \varepsilon_{dj} x_j)$$

If **I** is an index for which  $\mathbf{a}_{I} = 0$  then the coefficient of  $\mathbf{x}^{I}$  in  $g(\mathbf{x})$  is a non-zero polynomial in the  $\varepsilon_{ij}$ .

*Proof.* A monomial  $\mathbf{a}_{\mathbf{I}}\mathbf{x}^{\mathbf{J}}$  in f contributes a product

$$\mathbf{a}_{J} \prod_{k=1}^{d} \left( \epsilon_{k0} + \sum_{i} \epsilon_{ki} \mathbf{x}_{i} \right)^{j_{k}}$$
(11.5.1)

to  $g(\mathbf{x})$ . Every term in this expansion has the form  $\varepsilon_{1?}^{j_1} \cdots \varepsilon_{d?}^{j_d}$ . where each "?" represents an index in 0, ..., d. Consequently the coefficient of  $\mathbf{x}^I$  in  $g(\mathbf{x})$  is a sum of terms of the form

(coefficient of  $x^J$ ) × (at most one term arising from  $x^J$ )

and so there is no cancellation. Note that (11.5.1) has terms of every index of degree at most deg(f).

**Lemma 11.58.** If  $f(\mathbf{x}) \in \overline{\mathbf{P}}_d$  has non-negative coefficients then it is a limit of polynomials in  $\mathbf{P}_d^{\text{pos}}$ .

Proof. Define

$$f_{\varepsilon}(\textbf{x}) = f\big(\varepsilon_{10} + \sum \varepsilon_{1j} x_j, \ldots, \varepsilon_{d0} + \sum \varepsilon_{dj} x_j\big).$$

By Lemma 11.57 the terms in  $f_{\epsilon}$  with coefficient zero in f are a sum of polynomials in the  $\epsilon_{ij}$  with positive coefficients in  $g(\mathbf{x})$ . Observe that f is the limit of  $f_{\epsilon}$  through positive values of  $\epsilon_{ii} \rightarrow 1$  and  $\epsilon_{i \neq j} \rightarrow 0$ . All the  $f_{\epsilon}$  are in  $P_d$ , have all positive coefficients for all indexes of degree at most deg(g), and so f is a limit of polynomials in  $\mathbf{P}_d^{\text{pos}}$ .

Interlacings in  $\overline{\mathbf{P}}_d$  can be approximated by interlacings in  $\mathbb{P}_d$ .

**Corollary 11.59.** If  $f \leftarrow g$  in  $\overline{\mathbf{P}}_d$  then there are  $f_n, g_n$  such that

- $\lim f_n = f$
- $\lim g_n = g$
- $f_n \longleftarrow g_n$

*Proof.* Since  $f + yg \in \overline{P}_{d+1}$  we let  $h_{\varepsilon}$  be the polynomial determined by the lemma above that converges to f + yg. If  $h_{\varepsilon} = f_{\varepsilon}(\mathbf{x}) + yg_{\varepsilon}(\mathbf{x}) + \cdots$  then  $f_{\varepsilon}, g_{\varepsilon}$  are the desired polynomials.

# 11.6 Multiaffine polynomials

A *multiaffine polynomial* is a polynomial that is linear in each variable. We look at a few properties of such polynomials; for a more detailed discussion see [bbs].

### Definition 11.60.

 $\mathsf{MA}_{d} = \{ \mathsf{f} \in \overline{\mathbf{P}}_{d} \mid \text{all variables have degree } 1 \}$ 

We can write a multiaffine polynomial as

$$\sum_{I \subset \{1, \dots, d\}} a_I x_I$$

A multiaffine polynomial in  $MA_d$  can be visualized as an assignment of real numbers to the  $2^d$  vertices of the d-cube.

There is no simple way of determining if a polynomial f(x) is multiaffine. Brändén[23] proved that a necessary and sufficient condition is that

$$\frac{\partial f}{\partial x_i}\frac{\partial f}{\partial x_j} - f\frac{\partial^2 f}{\partial x_i\partial x_j} \geqslant 0 \quad \text{ for all } 1\leqslant i,j\leqslant n.$$

In the case d = 2 this is the simple criterion

 $a + bx + cy + dxy \in MA_2 \Leftrightarrow ad - bc \leqslant 0$ 

**Example 11.61.** Here is an application for a polynomial of total degree 3. Consider the question:

For which m is  $f(x, y, z) = xyz - x - y - z + m \in MA_3$ ?

Using the criterion we find

$$\frac{\partial f}{\partial x}\frac{\partial f}{\partial y} - f \cdot \frac{\partial^2 f}{\partial x \partial y} = z^2 - zm + 1$$

This is non-negative if and only if  $|m| \leq 2$ . A similar calculation shows

xyz – 
$$ax - by - cz + m \in MA_3$$
 iff a, b, c  $\geq 0$  and  $m^2 \leq 4abc$ 

Here are three examples of general multiaffine polynomials in  $MA_d$ ; the first two are trivial to show multiaffine, but the last is the Grace polynomial Grace<sub>d</sub> and does not have an elementary proof that it is in  $MA_d$ .

$$\begin{split} & \prod_{1}^{d} (x_i + y_i) \\ & \prod_{1}^{d} (x_i y_i - 1) \\ & \sum_{\sigma \in \text{sym}(d)} \prod_{1}^{d} (x_i + y_{\sigma i}) \end{split}$$

It's easy to construct multiaffine polynomials using this lemma:

**Lemma 11.62.** If  $f \in \overline{P}_d$  then the multiaffine part of f is in  $MA_d$ .

*Proof.* The usual reversal, differentiation, reversal shows that if  $\sum f_i(\mathbf{x})y^i \in \overline{\mathbf{P}}_d$  then  $f_0(\mathbf{x}) + f_1(\mathbf{x})y \in \overline{\mathbf{P}}_d$ . Repeating this for each variable shows that the multiaffine part of f is in  $\overline{\mathbf{P}}_d$ .

**Example 11.63.** If we let  $f = (1 + x + y + z)^n$  then the multiaffine part is

1 + n(x + y + z) + n(n-1)(xy + xz + yz) + n(n-1)(n-2)xyz

and is in MA<sub>3</sub>.

This last example is an example of a multiaffine polynomial that is symmetric in its variables. See Lemma 10.53.

Multiaffine polynomials are closed under multiplication in the following sense

**Corollary 11.64.** If f,  $g \in MA_d$  the the multiaffine part of fg is also in  $MA_d$ .

We can also construct multiaffine polynomials using determinants. If S is symmetric and  $D_1, \ldots, D_d$  are positive definite then the multiaffine part of

$$|S + x_1D_1 + \cdots + x_dD_d|$$

is multiaffine. We don't always have to take the multiaffine part. If  $\nu_1,\cdots,\nu_d$  are vectors then

$$|\mathbf{S} + \mathbf{x}_1 \mathbf{v}_1^{\mathrm{t}} \mathbf{v}_1 + \dots + \mathbf{x}_d \mathbf{v}_d^{\mathrm{t}} \mathbf{v}_d|$$

is multiaffine.

There are no interesting multiaffine bilinear forms.

**Lemma 11.65.** If  $f = \sum_{1}^{n} \alpha_{ij} x_i y_j \in MA_{2d}$  then we can write

$$f = \left(\sum_{1}^{n} \alpha_{i} x_{i}\right) \cdot \left(\sum_{1}^{n} \beta_{i} y_{i}\right)$$

where all  $\alpha_i$  have the same sign, and all  $\beta_i$  have the same sign.

*Proof.* Write  $(a_{ij}) = (v_1, \dots, v_d)^t$  so that

$$f = (x_1, \dots, x_d) \begin{pmatrix} v_1 \\ \vdots \\ v_d \end{pmatrix} \begin{pmatrix} y_1 \\ \vdots \\ y_d \end{pmatrix}$$

We see  $\frac{\partial^2 f}{\partial x_1 \partial x_j} = 0$ , and hence  $\frac{\partial f}{\partial x_1} \frac{\partial f}{\partial x_j} \ge 0$ . If  $\mathbf{y} = (y_1, \dots, y_d)^t$  then  $\frac{\partial f}{\partial x_1} = v_1 \cdot y$  and  $\frac{\partial f}{\partial x_i} = v_j \cdot y$ . Consequently,

$$(\mathbf{v}_1 \cdot \mathbf{y})(\mathbf{v}_j \cdot \mathbf{y}) \ge 0$$

If the product of two linear forms is never negative then they must be multiples of one another. Thus we can write  $v_j = \alpha_j v_1$  for some constant  $\alpha_j$  which leads to the the desired form

$$\mathbf{f} = \begin{bmatrix} \alpha_1 \mathbf{x}_1 + \dots + \alpha_d \mathbf{x}_d \end{bmatrix} \cdot \begin{bmatrix} \nu_1 \mathbf{y} \end{bmatrix}$$

Each of the factors is in  $MA_d$  so all the coefficients of a factor have the same sign.  $\Box$ 

The following might be true for arbitrary multiaffine polynomials with positive coefficients.

**Lemma 11.66.** Polynomials with positive coefficients in  $MA_2$  are closed under Hadamard product.

*Proof.* We know a+bx+cy+dxy is in MA<sub>2</sub> iff  $\begin{vmatrix} b & a \\ d & c \end{vmatrix} \ge 0$ . If a, b, c, d are positive then  $\begin{vmatrix} b & a \\ d & c \end{vmatrix}$  is positive semi-definite. Since positive semi-definite matrices are closed under Hadamard product, the conclusion follows.

#### **11.7** Multiaffine polynomials with complex coefficients

Multiaffine polynomials with complex coefficients are much more complicated than those with real coefficients. A multiaffine polynomial with real coefficents determines polynomials with complex coefficients, but not all of them. We give this simple construction, and then restrict ourselves to multiaffine polynomials with two variables, where we will see an equivalence with certain Möbius transformations.

Suppose that  $f(x_1, ..., x_d) \in MA_d$ . If  $\sigma_1, ..., \sigma_d$  lie in the upper half plane then  $f(x_1 + \sigma_1, ..., x_d + \sigma_d)$  is multiaffine, has complex coefficients, and is non-vanishing for  $\mathbf{x} \in UHP$ .

So, assume that  $f(x, y) = \alpha x + \beta + \gamma y x + \delta y$ . Solve f(x, y) = 0 for y:

$$y = -\frac{\alpha x + \beta}{\gamma x + \delta}.$$

If M is the Möbius transformation with matrix  $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$  then y = -M(x). If  $x \in UHP$  and  $f \in HP_2$  then  $y \in \mathbb{C}-UHP$ , for otherwise we have a solution with both x, y in the upper half plane. Thus,  $x \in UHP$  if and only if  $M(x) \in \overline{UHP}$ . Consequently,

f is complex multiaffine  $\iff M : UHP \longrightarrow \overline{UHP}$ .

For the rest of this section we determine conditions on a Möbius transformation to map UHP  $\longrightarrow$  UHP.

If M maps the upper half plane to itself then the image of the upper half plane is either a half plane parallel to the real line, or the interior of a circle lying in the upper half plane.

Suppose that  $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$  is a matrix with complex entries that maps the upper half plane to itself. Since our entries are complex, we can divide by the square root of the determinant, and so without loss of generality we may assume that **the determinant is one**. In the case the image of the upper half plane is a half plane contained in the upper half plane we have

$$\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} = \begin{pmatrix} e & f \\ g & h \end{pmatrix} + \iota t \begin{pmatrix} g & f \\ 0 & 0 \end{pmatrix}$$

where  $\begin{pmatrix} e & f \\ a & h \end{pmatrix}$  is a matrix with real entries and determinant 1, and  $t \ge 0$ .

The more interesting case is when the image of the real line is a circle constrained in the upper half plane. The image of the upper half plane is either the interior or the exterior of the circle. We first find the center of the circle<sup>1</sup>

center = 
$$\frac{\beta\overline{\gamma} - \alpha\overline{\delta}}{\delta\overline{\gamma} - \gamma\overline{\delta}}$$

<sup>&</sup>lt;sup>1</sup>Thanks to David Wright for this information.

The image of  $\infty$  is  $\alpha/\gamma$ , so the radius is the distance from the center to  $\alpha/\gamma$ :

radius = 
$$\frac{1}{|2\Im(\gamma\overline{\delta})|}$$

The image of the circle is entirely in the upper half plane if the center lies in the upper half plane, and the radius is less than the imaginary part of the center. This gives the middle condition below. If this holds, then the image of the upper half plane lies in the circle if and only if 0 is not the image of a point in the upper half plane. Since  $-\beta/\alpha$  maps to zero, we have the three conditions

$$\Im(\text{center}) > 0 \qquad \qquad \Re\left( \begin{vmatrix} \overline{\alpha} & \beta \\ \overline{\gamma} & \delta \end{vmatrix} \right) \geqslant 1 \qquad \qquad \Im(\beta/\alpha) \geqslant 0$$

It is more interesting to write these conditions in terms of the real and imaginary parts. Write

$\alpha = a_1 + \iota a_2$	$\beta = b_1 + \iota b_2$		
$\gamma = c_1 + \iota c_2$	$\delta = d_1 + \iota d_2$		

Note that if we want to show  $\Im(\sigma/\tau) > 0$  it is enough to show  $\Im(\sigma\overline{\tau}) > 0$ . The three conditions are

$$\begin{array}{ll} 2\left(b_{1}c_{1}+b_{2}c_{2}-a_{1}d_{1}-a_{2}d_{2}\right)\left(-c_{1}d_{2}+c_{2}d_{1}\right)>0 & (\text{center})\\ \\ -b_{1}c_{1}-b_{2}c_{2}+a_{1}d_{1}+a_{2}d_{2}\geqslant1 & (\text{radius})\\ \\ a_{1}b_{2}-a_{2}b_{1}>0 & (\text{interior}) \end{array}$$

Now the determinant of M is one, which gives us two equations

(imaginary part)	$-b_2c_1 - b_1c_2 + a_2d_1 + a_1d_2 = 0$
(real part)	$-b_1c_1 + b_2c_2 + a_1d_1 - a_2d_2 = 1$

If we reexpress these in terms of determinants we get

$$\begin{pmatrix} \begin{vmatrix} a_1 & b_1 \\ c_1 & d_1 \end{vmatrix} + \begin{vmatrix} a_2 & b_2 \\ c_2 & d_2 \end{vmatrix} \begin{pmatrix} c_1 & c_2 \\ d_1 & d_2 \end{vmatrix} > 0$$
 (center)  
$$\begin{vmatrix} a_1 & b_1 \\ c_1 & d_1 \end{vmatrix} + \begin{vmatrix} a_2 & b_2 \\ c_2 & d_2 \end{vmatrix} \ge 1$$
 (radius)  
$$\begin{vmatrix} a_1 & b_1 \\ c_1 & d_1 \end{vmatrix} - \begin{vmatrix} a_2 & b_2 \\ c_2 & d_2 \end{vmatrix} = 1$$
 (real part)

and these simplify to four conditions

$$\begin{vmatrix} a_1 & b_1 \\ c_1 & d_1 \end{vmatrix} > 1 \qquad \begin{vmatrix} a_2 & b_2 \\ c_2 & d_2 \end{vmatrix} \leqslant 0 \qquad \begin{vmatrix} a_1 & a_2 \\ b_1 & b_2 \end{vmatrix} > 0 \qquad \begin{vmatrix} c_1 & c_2 \\ d_1 & d_2 \end{vmatrix} > 0$$

Expressed in terms of the original coefficients this is

 $|\Re(\mathsf{M})| \ge 1$   $|\Im(\mathsf{M})| \le 0$   $\Im(\alpha/\beta) \le 0$   $\Im(\gamma/\delta) \le 0$ 

**Example 11.67.** The Möbius transformation with matrix  $M = \begin{pmatrix} 1+\iota & \iota \\ 2+\iota & 1+\iota \end{pmatrix}$  maps the real line to the circle with center  $1 + \iota/2$  and radius 1/2. Since the solution to Mx = 0 is  $\iota/(1 + \iota) = -1/2(1 + \iota)$ , the upper half plane maps to the interior of the circle. The corresponding complex multiaffine polynomial is

 $\mathbf{i} + (1 + \mathbf{i})\mathbf{x} + (1 + \mathbf{i})\mathbf{y} + (2 + \mathbf{i})\mathbf{x}\mathbf{y}$ 

The four conditions are

 $|\mathfrak{R}(\mathsf{M})| = 1$   $|\mathfrak{I}(\mathsf{M})| = 0$   $\operatorname{Im}(\alpha/\beta) = -1$   $\mathfrak{I}(\gamma/\delta) = -1/2.$ 

# 11.8 Nearly multiaffine polynomials

A *nearly multiaffine polynomial* is a polynomial that is degree one in all but one variable.

**Lemma 11.68.** If  $yf(x) + zg(x) \in \overline{\mathbf{P}}_3$  then f is a constant multiple of g.

*Proof.* We first remove all common factors of f and g so that it suffices to show that f and g are constant. Since  $yf + zg \in \overline{\mathbf{P}}_3$  it follows that f and g interlace. We consider the possibilities:

First assume  $f \leq g$ ; the case  $g \leq f$  is the same. If we let y = x then  $(xf) + zg \in \overline{\mathbf{P}}_2$ . But this is impossible, since this implies that g and xf interlace yet deg(xf) = 2 + deg(g).

We next assume  $f \ll g$ . All roots of g are smaller than the largest root s of f. If r > s then  $(x - r)f + zg \in \overline{P}_2$ , and hence  $(x - r)f \leq g$ . But this implies that g has a root in [s, r], which is a contradiction.

Thus, f and g must be constant, which proves the lemma.

**Corollary 11.69.** If  $f(x, y) = \sum_{1}^{d} y_i f_i(x) \in \overline{P}_{d+1}$  then there are positive  $\alpha_i$  and  $g \in P$  such that

$$f(\mathbf{x}, \mathbf{y}) = g(\mathbf{x}) \left( \alpha_1 \mathbf{y}_1 + \cdots + \alpha_d \mathbf{y}_d \right)$$

We next show that there are no interesting matrices M(x) such that  $yMz^t \in \overline{\mathbf{P}}_d$  for vectors  $\mathbf{y}, \mathbf{z}$  of variables. The only such M have the form  $g(x)v^t \cdot w$  where v, w are vectors of positive constants.

**Lemma 11.70.** If  $f(x, y, z) = \sum_{1}^{d} y_i z_j f_{ij}(x) \in \overline{P}_{2d+1}$  then there are  $g \in P$  and positive  $\alpha_i, \beta_i$  so that

$$f(x, y, z) = g(x) \left( \sum \alpha_i y_i \right) \left( \sum \beta_i z_i \right).$$

*Proof.* Consider the matrix  $M(x) = (f_{ij}(x))$ . For any  $\alpha \in \mathbb{R}$  we know from Lemma 11.65 that  $M(\alpha)$  has rank 1, so for all distinct i, j, k, l

$$f_{ij}(\alpha)f_{kl}(\alpha) = f_{il}(\alpha)f_{kj}(\alpha)$$

and hence  $f_{ij}f_{kl} = f_{il}f_{kj}$  as polynomials. Now

$$f_{11} \cdot M = (f_{11}f_{ij}) = \begin{pmatrix} f_{11} \\ \vdots \\ f_{1d} \end{pmatrix} \begin{pmatrix} f_{11} & \dots & f_{d1} \end{pmatrix}$$

since  $f_{11}f_{ij} = f_{1j}f_{j1}$ . It follows that

$$f_{11} \cdot f(\mathbf{x}, \mathbf{y}, \mathbf{z}) = \left(\sum y_i f_{1i}\right) \left(\sum z_i f_{i1}\right).$$

Since each factor is in  $\overline{\mathbf{P}}_{2d+1}$  we can use the corollary to write

$$f_{11} \cdot f(x, \boldsymbol{y}, \boldsymbol{z}) = g(x) \big( \sum \alpha_i y_i \big) \big( \sum \beta_i z_i \big).$$

Clearly  $f_{11}$  divides g, giving the result.

We can apply the characterization of complex multiaffine polynomials to get properties of nearly multiaffine polynomials.

**Lemma 11.71.** Suppose  $f(x) + g(x)y + h(x)z + k(x)yz \in \overline{P}_3$ . If  $\sigma \in UHP$ ,  $\delta^2 = \left| \begin{array}{c} f(\sigma) & g(\sigma) \\ h(\sigma) & k(\sigma) \end{array} \right|$  and  $\left| \begin{array}{c} \alpha & \beta \\ \gamma & \delta \end{array} \right| = \left| \begin{array}{c} f(\sigma)/\delta & g(\sigma)/\delta \\ h(\sigma)/\delta & k(\sigma)/\delta \end{array} \right|$  then

$$\begin{vmatrix} \mathfrak{R}(lpha) & \mathfrak{R}(eta) \\ \mathfrak{R}(\gamma) & \mathfrak{R}(\delta) \end{vmatrix} \geqslant 1 \qquad \begin{vmatrix} \mathfrak{I}(lpha) & \mathfrak{I}(eta) \\ \mathfrak{I}(\gamma) & \mathfrak{I}(\delta) \end{vmatrix} \leqslant 0$$

# 11.9 The polar derivative and the Schur-Szegö map

The polar derivative was defined for polynomials in  $\mathbf{P}_d^{\text{pos}}$  (See p. 321.) but its properties still hold for  $\overline{\mathbf{P}}_d$  as well if we restrict ourselves to a single variable. If  $f(\mathbf{x}, y) \in \mathbb{P}_{d+1}$  then the polar derivative with respect to y, which we shall denote as  $\partial_y^{\text{polar}}$ , is defined as the composition

 $\xrightarrow{\text{reverse y}}, \xrightarrow{\text{differentiate}}, \xrightarrow{\text{reverse y}}$ 

 $\partial_y^{polar}$  defines a map from  $\overline{\mathbf{P}}_{d+1}$  to itself since reversal maps  $\overline{\mathbf{P}}_{d+1}$  to itself. As in one variable we have the formula

$$\partial_{u}^{\text{polar}} f = nf - y\partial_{y} f$$

where n is the degree of y in f. The key fact about polar derivatives is

**Lemma 11.72.** *If*  $f(\mathbf{x}, \mathbf{y}) \in \overline{\mathbf{P}}_{d+1}$  *then* 

- 1.  $\partial_y$  and  $\partial_y^{polar}$  commute
- 2.  $\partial_{y}f$  and  $\partial_{y}^{polar}f$  interlace.

Proof. Commutativity is the calculation

$$\partial_{y}\partial_{u}^{\text{polar}}f = nf' - f' - yf'' = (n-1)f' - yf'' = \partial_{u}^{\text{polar}}\partial_{y}$$

Choose  $\alpha \in \mathbb{R}$  and let  $g(\mathbf{x}, y) = f(\mathbf{x}, y - \alpha)$ . Since  $g \in \overline{\mathbf{P}}_{d+1}$  we know that its polar derivative is in  $\overline{\mathbf{P}}_{d+1}$  and therefore  $ng - y\partial_y g \in \overline{\mathbf{P}}_{d+1}$ . Substituting  $y + \alpha$  for y yields

$$nf(\mathbf{x},\mathbf{y}) - (\mathbf{y} + \alpha)f(\mathbf{x},\mathbf{y}) = \partial_{\mathbf{y}}^{\text{polar}}f - \alpha\partial_{\mathbf{y}}f \in \overline{\mathbf{P}}_{d+1}$$

which proves the lemma.

We can apply this to derive an important theorem. We call the transformation the *Schur-Szegö map*.

**Theorem 11.73.** If  $f = \sum_{0}^{n} a_{i}y^{i} \in P(n)$  and  $\sum g_{i}(x)y^{i} \in \overline{P}_{d+1}(n)$  then

$$\sum a_i i! (n-i)! g_i(\mathbf{x}) \in \mathbf{P}_d$$

Equivalently, the transformation  $y^{i} \times f(x)y^{j} \mapsto \begin{cases} 0 & i \neq j \\ i! (n-i)! f(x) & i = j \end{cases}$ defines a map  $P(n) \times \overline{P}_{d+1}(n) \longrightarrow \overline{P}_{d}(n).$ 

*Proof.* By the lemma we see that if  $f(x) \in \textbf{P}$  and F(x,y) is the homogenization of f then

$$F(\partial_y, \partial_y^{polar}) \colon \overline{\mathbf{P}}_{d+1} \longrightarrow \overline{\mathbf{P}}_{d+1}$$

since  $\partial_y + \alpha \partial_y^{\text{polar}} : \overline{\mathbf{P}}_{d+1} \longrightarrow \overline{\mathbf{P}}_{d+1}$  and the regular and polar derivatives commute. It's easy to see that

$$\vartheta_{y}^{i} \left( \vartheta_{y}^{\text{polar}} \right)^{n-i} y^{j} = \begin{cases} i!(n-i)! & i=j \\ 0 & i\neq j \end{cases}$$

so we have

$$F(\partial_{y}, \partial_{y}^{\text{polar}}) \sum_{i} g_{i}(\mathbf{x}) y^{i} = \sum_{i,j} a_{i} \partial_{y}^{i} \left( \partial_{y}^{\text{polar}} \right)^{n-i} y^{j} g_{j}(\mathbf{x})$$
$$= \sum_{i} a_{i} i! (n-i)! g_{i}(\mathbf{x}) \in \overline{\mathbf{P}}_{d} \quad (11.9.1)$$

There are two important consequences that previously were proved using Grace's theorem.

**Corollary 11.74** (Schur-Szegö). *If*  $\sum a_i x^i \in P$  and  $\sum b_i x^i \in P^{pos}$  then

 $\sum a_i b_i i! (n-i)! x^i \in \mathbf{P}.$ 

*Proof.* Since  $\sum b_i x^i \in \mathbf{P}^{pos}$  we know that  $\sum b_i (-xy)^i \in \overline{\mathbf{P}}_2$ . Now apply the theorem.

**Corollary 11.75** (Borceau, Branden, Shapiro). *If*  $\sum a_i x^i \in P$  *and*  $g(x) \in P$  *then* 

$$\sum (n-i)! a_{n-i} g^{(i)}(x) \in \boldsymbol{P}.$$

*Proof.* Apply the theorem to

$$g(x+y) = \sum g^{(i)}(x) \frac{y^i}{i!} \in P_2.$$

More generally, if  $T: \mathbf{P} \longrightarrow P_2$  is a map (See p. 512.) then we get a map



# 11.10 Determining the direction of interlacing

If all linear combinations of f and g are in  $\overline{\mathbf{P}}_d$  then either f  $\leftarrow$  g or g  $\leftarrow$  f. Determining the direction of interlacing is easy if the degrees are different; here are some ways of determining the direction when the degrees are the same. First of all,

**Lemma 11.76.** *If*  $f \sim g$  *in*  $P_d$  *have the same degree then either*  $f \leq g$  *or*  $g \leq f$ .

*Proof.* The homogeneous part of  $\alpha f + g$  is positive for large  $\alpha$ , and negative for very negative  $\alpha$ . Thus, there is an  $\alpha$  for which one of the coefficients of  $(\alpha f + g)^H$  is zero, but this is only possible if  $(\alpha f + g)^H = 0$ . For this  $\alpha$  define  $r = \alpha f + g$ . Now  $f \sim r$ , and since r has smaller degree than f we have  $f \leq r$ . Thus  $g = \alpha f + r$ , from which the conclusion follows.

**Lemma 11.77.** *If*  $f, g \in P_d$ ,  $f \leq g$ , and  $g \leq f$  then g is a scalar multiple of f.

*Proof.*  $f \leq g$  implies that there are  $\alpha > 0$  and r such that  $f \leq r$ ,  $f = \alpha g + r$  and r has positive leading coefficient. But  $g \leq f$  implies  $f = (1/\alpha)g - (1/\alpha)r$  where -r has positive leading coefficient. It follows that r = 0.

**Lemma 11.78.** Suppose that  $f \sim g$  in  $P_d$ . If either

$$\forall a_2, \ldots, a_d \in \mathbb{R} \quad f(x, a_2, \ldots, a_d) \longleftarrow g(x, a_2, \ldots, a_d)$$

or

$$\exists a_2, \ldots, a_d \in \mathbb{R} \quad f(x, a_2, \ldots, a_d) \ll g(x, a_2, \ldots, a_d)$$

*then* f  $\leftarrow$  g.

*Proof.* We may assume that both f and g have degree n. If  $g \leq f$  then we know that

 $g(x, a_2, \ldots, a_d) \leq f(x, a_2, \ldots, a_d)$ 

The hypothesis gives the other direction, so

$$g(x, a_2, \ldots, a_d) = \alpha f(x, \alpha_2, \ldots, \alpha_d)$$

Since this holds for all x and  $\alpha_i \in \mathbb{R}$  this implies that  $g = \alpha f$ , so  $f \longleftarrow g$  still holds.

If we have strict interlacing for one choice of  $a_i$  then we can't also have  $g(x, a_2, ..., a_d) \leq f(x, a_2, ..., a_d)$ .

We only need interlacing in one variable to determine interlacing for  $P_2$ .

**Lemma 11.79.** If  $f, g \in P_2$  and  $f(x, a) \underline{\ll} g(x, a)$  for all  $a \in \mathbb{R}$  then  $f \underline{\ll} g$ .

*Proof.* It suffices to show that  $f^H$  and  $g^H$  are scalar multiples. Geometrically the hypotheses say that the solution curves for g interlace the solution curves for f. For large positive x we therefore have  $f^H(x) \leq g^H(x)$  and for small negative x we have  $f^H(-x) \leq g^H(-x)$  which implies  $f^H$  is a scalar multiple of  $g^H$ .

Substitution for  $P_d$  preserves the direction of interlacing but this is not true for  $\overline{P}_d$ . For example

$$\begin{vmatrix} x & 1 & 1 \\ 1 & y & 1 \\ 1 & 1 & z \end{vmatrix} = 2 - x - y + z(xy - 1)$$

so  $2 - x - y \leq xy - 1$ . However, substitution of y = 2 and y = -2 gives interlacings in different directions:

$$\begin{array}{rrrr} -x &= (2-x-y)(x,2) & \underline{\gg} & (xy-1)(x,2) &= 2x-1 \\ -x+4 &= (2-x-y)(x,-2) & \underline{\ll} & (xy-1)(x,-2) &= -2x-1 \end{array}$$

In the following we let  $\mathbf{a} = (a_1, \dots, a_d) \in \mathbb{R}^d$  and  $\mathbf{b} = (b_1, \dots, b_d) \in (\mathbb{R}^+)^d$ . From Lemma 10.59 we know that if  $f(\mathbf{x}) \in \overline{\mathbf{P}}_d$  then  $f(\mathbf{a} + \mathbf{b}t) \in \mathbf{P}$ . The converse is also true, see [18]. **Lemma 11.80.** If  $f \sim g$  in  $\overline{P}_d$  and  $f(T) \leftarrow g(T)$  for all  $\mathbf{a} \in \mathbb{R}^d$  and  $\mathbf{b} \in (\mathbb{R}^+)^d$ where  $T = \mathbf{a} + \mathbf{b}t$  then  $f \leftarrow g$ .

*Proof.* We first assume f,  $g \in P_d(n)$ . If we take  $\mathbf{a} = (0, a_2, ..., a_d)$  and  $\mathbf{b}_{\varepsilon} = (1, \varepsilon, ..., \varepsilon)$  then  $f(\mathbf{a} + t\mathbf{b}_{\varepsilon}) \leq g(\mathbf{a} + t\mathbf{b}_{\varepsilon})$ . Since  $f(\mathbf{a} + t\mathbf{b}_{\varepsilon})$  and  $g(\mathbf{a} + t\mathbf{b}_{\varepsilon})$  also have degree n it follows that  $f(\mathbf{a} + t\mathbf{b}) \leq g(\mathbf{a} + t\mathbf{b})$ . Thus

$$\forall a_i \in \mathbb{R}$$
  $f(t, a_2, \dots, a_d) \leq g(t, a_2, \dots, a_d)$ 

so f  $\leq g$ .

Now assume that  $f, g \in \overline{\mathbf{P}}_d$  have the same degree. Let  $X = x_1 + \cdots + x_d$ ,  $f_{\varepsilon}(\mathbf{x}) = f(\mathbf{x} + \varepsilon X)$ , and  $g_{\varepsilon}(\mathbf{x}) = g(\mathbf{x} + \varepsilon X)$ . We claim

- 1.  $f_{\varepsilon}, g_{\varepsilon} \in P_d$
- 2.  $\lim_{\varepsilon \to 0} f_{\varepsilon} = f$ ,  $\lim_{\varepsilon \to 0} g_{\varepsilon} = g$ .
- 3.  $f_{\varepsilon}(T) \leftarrow g_{\varepsilon}(T)$ .

Given 1,2,3 then by the first part we have that  $f_{\varepsilon}(\mathbf{x}) \leftarrow g_{\varepsilon}(\mathbf{x})$ . All terms have the same degree, so using Lemma 1.42  $f(\mathbf{x}) \leq g(\mathbf{x})$ .

Now 1) follows from Lemma 10.59, 2) is clear, and 3) follows from the first part as follows. If

$$S = \mathbf{a} + t\mathbf{b} + \varepsilon \left[\sum \alpha_i + t \sum b_i\right](1, 1, ..., 1)$$

then  $f_{\varepsilon}(T) = f(S)$  and  $g_{\varepsilon}(T) = g(S)$ 

Interlacing in  $\overline{\mathbf{P}}_d$  is the limit of interlacing in  $\mathbb{P}_d$ .

**Corollary 11.81.** If  $f \leftarrow g$  in  $\overline{\mathbf{P}}_d$  then there are  $f_n, g_n$  such that

- 1.  $f_n, g_n \in P_d$ .
- 2.  $f_n \leftarrow g_n$ .
- 3.  $\lim_{n\to\infty} f_n = f$  and  $\lim_{n\to\infty} g_n = g$ .

*Proof.* First assume  $f \leq g$ . Since  $f + x_{d+1} \in \overline{\mathbf{P}}_{d+1}$  there are  $F_n \in P_{d+1}$  such that  $\lim F_n = f + x_{d+1}g$ . If we write  $F_n = f_n + x_{d+1}g_n + \cdots$  then  $f_n$ ,  $g_n$  satisfy the three conclusions.

If  $f \leftarrow g$  then we can write  $g = \alpha f + r$  where  $f \leq r$ . Now apply the first paragraph to f and r.

**Corollary 11.82.** If  $f \leq g$  in  $\overline{\mathbf{P}}_d$  then  $\begin{vmatrix} f & g \\ \frac{\partial f}{\partial x_i} & \frac{\partial g}{\partial x_i} \end{vmatrix} \leqslant 0$  for i = 1, ..., d.

*Proof.* We approximate and reduce the problem to the corresponding result for  $\mathbb{P}_d$ . If suffices to assume that  $f \leq g$  and i = 1. Since  $f + x_{d+1}g \in \overline{\mathbf{P}}_{d+1}$  there are  $F_n = f_n + x_{d+1}g_n + \cdots \in \mathbb{P}_{d+1}$  such that  $\lim_{n \to \infty} F_n = f + x_{d+1}g$ . Now by Taylor's theorem

$$F_{n}(x_{1}+t, x_{2}, \dots, x_{d+1}) = f_{n} + x_{d+1}g + t\left(\frac{\partial f_{n}}{\partial x_{1}} + x_{d+1}\frac{\partial g_{n}}{\partial x_{1}}\right) + \cdots$$

so Corollary 10.27 tells us that

$$\begin{vmatrix} f_n & g_n \\ \frac{\partial f_n}{\partial x_1} & \frac{\partial g_n}{\partial x_1} \end{vmatrix} \leqslant 0$$

and taking limits finishes the proof.

**Corollary 11.83.** If  $f \sim g \in \overline{P}_d$  and  $\begin{vmatrix} f & g \\ \frac{\partial f}{\partial x_i} & \frac{\partial g}{\partial x_i} \end{vmatrix} < 0$  for some  $\mathbf{x} \in \mathbb{R}^d$  and some i where  $1 \leq i \leq d$  then  $f \leq g$ .

*Proof.* We know  $f \leq g$  or  $g \leq f$ . If the latter holds then

$$\begin{vmatrix} f & g \\ \frac{\partial f}{\partial x_{i}} & \frac{\partial g}{\partial x_{i}} \end{vmatrix} \geqslant 0$$

but this contradicts the hypothesis.

#### **11.11** Reversal in P<sub>d</sub>

We use the properties of the graph of a polynomial in  $\mathtt{P}_d$  to prove that the reversal is in  $\overline{P}_d.$ 

**Lemma 11.84.** If  $f(\mathbf{x}) \in P_d(n)$  then the reversal  $f^{rev}$  is in  $\overline{\mathbf{P}}_d(nd)$ , where

$$f^{rev}(\mathbf{x}) = (x_1 \cdots x_d)^n f\left(\frac{1}{x_1}, \dots, \frac{1}{x_d}\right)$$
(11.11.1)

Proof. Define

$$g_{\varepsilon}(\mathbf{x}) = \left(\prod_{i=1}^{d} (x_i + \varepsilon \sum_{i \neq j} x_j)\right)^n f\left(\frac{1}{x_1 + \varepsilon \sum_{1 \neq j} x_j}, \dots, \frac{1}{x_d + \varepsilon \sum_{d \neq j} x_j}\right)$$

Since  $\lim_{\varepsilon \to 0} g_{\varepsilon}(\mathbf{x}) = f^{rev}(\mathbf{x})$ , it suffices to show that  $g_{\varepsilon} \in P_d(nd)$ . Now  $g_{\varepsilon}$  clearly satisfies homogeneity, so we just have to check substitution. Let  $x_1 = t, x_2 = a_2, \dots, x_d = a_d$ , and set  $a = a_2 + \cdots a_d$ . It suffices to show that the second factor, which is

$$f\left(\frac{1}{t+\varepsilon \mathfrak{a}},\frac{1}{\varepsilon t+\varepsilon \mathfrak{a}+(1-\varepsilon)\mathfrak{a}_{2}},\ldots,\frac{1}{\varepsilon t+\varepsilon \mathfrak{a}+(1-\varepsilon)\mathfrak{a}_{d}}\right),$$

has exactly nd roots. So, consider the curve

$$\mathfrak{C} = \left(\frac{1}{\mathfrak{t} + \mathfrak{ea}}, \frac{1}{\mathfrak{et} + \mathfrak{ea} + (1 - \mathfrak{e})\mathfrak{a}_2}, \dots, \frac{1}{\mathfrak{et} + \mathfrak{ea} + (1 - \mathfrak{e})\mathfrak{a}_d}\right)$$

The curve C has singularities at

$$-\epsilon a, -a - \frac{1-\epsilon}{\epsilon}a_2, \ldots, -a - \frac{1-\epsilon}{\epsilon}a_d$$

As t goes to  $\pm \infty$  the curve goes to 0, so there are d components in the closure of the image of C. A component has two coordinates that are unbounded; the other coordinates only vary by a bounded amount. The curve goes to  $+\infty$  in one coordinate, and is bounded in the others. In the other direction, it goes to  $-\infty$  in the other coordinate, and is bounded in the remaining ones. Consequently, the curve is eventually above the graph in one direction, and below in the other, so each component meets the graph in n points. Since there are d components of C we have nd intersection points, and so substitution is satisfied.

Using reversal, an argument entirely similar to Lemma 11.27 shows that

**Lemma 11.85.** If  $f(\partial_x)g$  is in  $\overline{\mathbf{P}}_d$  for all  $g \in \overline{\mathbf{P}}_d$  then f is in  $\overline{\mathbf{P}}_d$ .

**Lemma 11.86.** Suppose that  $f(\mathbf{x}, \mathbf{y}) = \sum f_i(\mathbf{x}) \mathbf{y}^i \in \widehat{\mathbf{P}}_{d+1}$  or  $\overline{\mathbf{P}}_{d+1}$ , and the degree of  $f_i(\mathbf{x})$  is i.

If the homogeneous parts of the  $f_i$  alternate in sign then

$$g(\mathbf{x},\mathbf{y},\mathbf{z}) \in P_{d+e+1}(\mathbf{n}) \implies f(\mathbf{x},-\partial_{\mathbf{y}}) g(\mathbf{x},\mathbf{y},\mathbf{z}) \in P_{d+e+1}(\mathbf{n}).$$

*Proof.* If e > 0 so that there are some **z** variables, then we replace  $f(\mathbf{x}, \mathbf{y})$  by  $f(\mathbf{x} + \epsilon \mathbf{z}, \mathbf{y})$ , which still satisfies the degree hypothesis. We now follow the proof of Lemma 11.24

# 11.12 Induced transformations

A linear transformation T on polynomials in one variable determines an induced transformation defined on polynomials in two variables:

$$\mathsf{T}_*(\mathsf{x}^{\mathsf{i}}\mathsf{y}^{\mathsf{j}}) = \mathsf{T}(\mathsf{x}^{\mathsf{i}})\,\mathsf{y}^{\mathsf{j}}.$$

Conversely, we will see that polynomials in  $\overline{\mathbf{P}}_2$  can determine linear transformations on  $\mathbf{P}$ . We have seen examples of induced transformations (See p. 317.) in  $\mathbb{P}_d$  where the results are easier.

We first make the important observation that if  $T: \mathbf{P} \longrightarrow \mathbf{P}$  then T does not necessarily determine a linear transformation  $T_*: \overline{\mathbf{P}}_2 \longrightarrow \overline{\mathbf{P}}_2$ . As in Remark 9.45, we take T(g) = g(D)x and note that (but see Lemma 11.91)

$$\mathsf{T}(\mathsf{x}\mathsf{y}-1)=\mathsf{y}-\mathsf{x}\not\in\overline{\mathbf{P}}_2$$

However, if  $T: \mathbf{P} \longrightarrow \mathbf{P}$  then the induced transformation always satisfies x-substitution, as the calculation below shows:

$$(\mathsf{T}_*\mathsf{f})(\mathsf{x},\alpha) = \mathsf{T}(\mathsf{f}(\mathsf{x},\alpha)) \in \mathbf{P}$$
 since  $\mathsf{f}(\mathsf{x},\alpha) \in \mathbf{P}$ .

**Definition 11.87.** If T is a linear transformation on polynomials in one variable then we say that T *satisfies induction* if the induced transformation  $T_*$  determines a map  $T_* : P_2 \longrightarrow \overline{P}_2$ . If  $T_* : P_2^{pos} \longrightarrow \overline{P}_2$  then we say that T *satisfies induction on*  $\mathbf{P}_2^{pos}$ .

Note that if T satisfies induction then T maps **P** to itself, since  $f \in \mathbf{P}$  implies  $T(f) = T_*(f) \in \overline{\mathbf{P}}_2$ . Also, if T and S satisfy induction, then so does their composition ST. Here are two assumptions that guarantee that induction is satisfied.

**Lemma 11.88.** Suppose that  $T: P \longrightarrow P$ . If either of the these two conditions hold then T satisfies induction.

- 1. There is an integer r such that if  $T(x^i) \neq 0$  then the degree of  $T(x^i)$  is i + r, and  $T(x^i)$  has positive leading coefficients.
- 2. T(g) = f \* g where  $f \in \mathbf{P}^{pos}$ .

*Proof.* The first part is a slightly more general statement that Theorem 9.44; the proof is similar. In this case we have  $T_* : P_2 \longrightarrow P_2$ .

In the second case, suppose that  $h(x,y) \in P_2(n)$ , and  $f \in \mathbf{P}^{pos}(r)$ . If r is at least n then the homogeneous part of T(h) has all positive terms. However, if r < n then  $T(h)^H$  has some zero coefficients. Replace f by  $f_{\varepsilon} = f(x)(1 + \varepsilon x)^n$  where  $\varepsilon$  is positive. The corresponding transformation  $S(g) = f_{\varepsilon}(x) * g$  satisfies  $S_*(h) \in P_2$  since  $f_{\varepsilon} \in \mathbf{P}^{pos}(n + r)$ . Taking limits as  $\varepsilon$  goes to zero shows that  $T_*(h) \in \overline{\mathbf{P}}_2$ .

We can resolve the problem with the transformation  $g \mapsto g(D)f$  by introducing a negative factor

**Corollary 11.89.** If f(x) is in P then the linear transformation  $T: g \mapsto g(-D)f$  satisfies induction.

Proof. We calculate

$$\sum T(x^i)\frac{(-y)^i}{i!} = \sum (\mathsf{D}^i f)\frac{y^i}{i!} = f(x+y).$$

Since  $f(x + y) \in P_2$ , we can apply Lemma 11.91.

**Lemma 11.90.** If  $F(x, y) = \sum f_i(x)(-y)^i$  is in  $\overline{\mathbf{P}}_2$  then

1. The linear transformation  $T: x^i \mapsto f_i$  maps  $P \longrightarrow P$ .

2. T satisfies induction.

Proof. By Lemma 11.21 the reverse of F satisfies

$$\begin{split} F_{rev}(x,y) &= \sum \, f_i(x) y^{n-i} \in \overline{P}_2 \\ Choose \quad h(x,y) &= \sum \, h_i(x) \, y^i \in \mathtt{P}_2 \\ T(h_i(x)) &= \text{coefficient of } y^n \text{ in } h_i(y) F_{rev}(x,y) \\ T_*(h(x,y)) &= \text{coefficient of } z^n \text{ in } h(z,y) F_{rev}(x,z) \end{split}$$

Since h(z, y) and  $F_{rev}(x, z) \in \overline{\mathbf{P}}_3$ , it follows that  $T_*(h) \in \overline{\mathbf{P}}_2$ . Finally, (2) holds since T satisfies induction.

As is often the case, if we replace multiplication by a differential operator and add a constraint, we can introduce a factorial.

**Lemma 11.91.** Suppose that  $f(x, y) = \sum \frac{f_i(x)}{i!} (-y)^i$  is in  $\overline{P}_2$ , and each  $f_i$  is a polynomial of degree *i* with positive leading coefficient.

- 1. The linear transformation  $T: x^i \mapsto f_i \text{ maps } P \longrightarrow P$ .
- 2. T satisfies induction.

*Proof.* Choose  $g(x, y) = \sum g_i(y)x^i$  in P<sub>2</sub>. By Lemma 11.86 we know that  $f(x, -\partial_z)g(y, z) \in \overline{\mathbf{P}}_3$ . The coefficient of  $z^0$  is in  $\overline{\mathbf{P}}_2$  and equals

$$\sum \frac{f_i(x)}{i!} (\partial_z)^i g_i(y) z^i = \sum f_i(x) g_i(y) = (T_*g)$$

We can show a linear transformation preserves all real roots by applying it to test functions.

**Lemma 11.92.** If T is a linear transformation on polynomials and  $T_*(1-xy)^n \in \overline{P}_2$  for n = 1, 2, ... then  $T : P \longrightarrow P$ .

*Proof.* Let  $g = \sum_{0}^{r} a_{i} x^{i} \in \mathbf{P}$  and calculate

$$g(-\mathsf{D}_{y})\mathsf{T}_{*}\left(1-\frac{xy}{n}\right)^{n}\Big|_{y=0} = \sum_{i=0}^{r} \alpha_{i}(-1)^{i}\mathsf{T}_{*}\sum_{i=0}^{n} \left(\frac{-xy}{n}\right)^{k} \binom{n}{k}\Big|_{y=0}$$
$$= \sum_{i=0}^{r} \alpha_{i}\frac{(\underline{n})_{i}}{n^{i}}\mathsf{T}(x^{i})$$

Since  $g(-D_y)$  maps  $\overline{P}_2$  to itself, and evaluation yields a polynomial in **P**, we see the above polynomial is in **P**. Finally, taking the limit of polynomials yields

$$\lim_{n\to\infty}\sum_{i=0}^{r}\alpha_{i}\frac{(\underline{n})_{i}}{n^{i}}T(x^{i})=\sum\alpha_{i}T(x^{i})=T(g(x))$$

## 11.13 Linear Transformations

One of the ways that we use to show that a polynomial has all real roots is to identify it as a coefficient of a polynomial in  $\overline{P}_2$ . We use this idea to show that some linear transformations preserve **P**.

**Theorem 11.93.** Suppose f is a polynomial in  $\mathbf{P}_2$  of degree n with coefficients given in (9.3.2). The linear transformation defined by  $T: x^i \mapsto f_{n-i}$  maps  $\mathbf{P}(n)$  to itself. Similarly the transformation  $x^i \mapsto f_i$  maps  $\mathbf{P}$  to itself.

*Proof.* Choose a polynomial  $g \in \mathbf{P}$ . Since  $g(y) \in \overline{\mathbf{P}}_2$  the result of multiplying g(y)f(x, y) is in  $\overline{\mathbf{P}}_2$ , and hence the coefficient of  $y^n$  is in  $\mathbf{P}$ . The coefficient of  $y^n$  is exactly Tg since

$$\begin{split} g(y)f(x,y) &= \left(\sum a_{j}y^{j}\right) \ \left(\sum f_{j}(x)y^{j}\right) \\ &= \sum_{k} \left(y^{k}\sum_{i+j=k}a_{j}f_{i}(x)\right) \end{split}$$

If we replace g by its reverse then the transformation is  $x^i \mapsto f_i$ .  $\Box$  **Corollary 11.94.** *If*  $f \in P$  *then the map*  $T: x^i \mapsto \frac{f^{(i)}}{i!}$  *preserves roots and interlacing. Proof.* Apply the Theorem to the Taylor series

$$f(x + y) = f(x) + f'(x)y + \cdots$$

An alternative argument follows from the observation that T(g) = g(D)EXP(f). Or, we could apply EXP to Theorem 11.93.

**Lemma 11.95.** If F(x, y) is the homogenization of f then define a map T:  $f \times g \mapsto F(-\frac{\partial}{\partial x}, y)$  g. This defines a linear transformation

$$P^{\text{pos}} \times \overline{\mathbf{P}}_2 \longrightarrow \overline{\mathbf{P}}_2.$$

*Proof.* Since  $f \in \mathbf{P}^{pos}$  we can write  $F = \prod(\alpha_i x + y)$  where all  $\alpha_i$  are positive. From Lemma 10.65 we know that

$$\prod_{i} (\epsilon x + y - \alpha_{i} \frac{\partial}{\partial x})$$

maps  $\mathbf{P}_2$  to itself. The result follows by letting  $\epsilon$  go to zero.

A similar application of Lemma 10.65 shows

**Lemma 11.96.** If  $f \in P$  and  $g \in \overline{P}_2$  then  $f(y - D_x)g(x, y) \in \overline{P}_2$ .

*Proof.* If suffices to show that  $(y + \epsilon x - D_x)h(x, y) \in \mathbf{P}_2$  if  $h \in \overline{\mathbf{P}}_2$  and  $\epsilon$  is positive. This follows from Lemma 10.65.

There is an interesting identity related to this last result:

$$f(y + D_x) = \sum_i f^{(i)}(D_x)g \frac{y^i}{i!}$$

Since this is linear in f only need to verify it for  $f = x^k$ 

$$\begin{split} \sum_{i} (x^k)^{(i)} (\mathsf{D}_x) g \, \frac{y^i}{i!} &= \sum_{i} \frac{(\underline{k})_i}{i!} \mathsf{D}_x^{k-i} g \, y^i \\ &= \sum_{i} \binom{k}{i} \mathsf{D}_x^{k-i} y^i \, g \\ &= (y + \mathsf{D}_x)^k \, g \end{split}$$

The next result constructs a transformation by looking at coefficients.

**Theorem 11.97.** Choose  $f(\mathbf{x}, \mathbf{y}) \in \mathbf{P}_{2d}$ , write f in terms of coefficients  $f = \sum_{I} \alpha_{I}(\mathbf{x}) \mathbf{y}^{I}$ , and let K be an index set. The linear transformation defined by

$$\mathbf{x}^{J} \mapsto \mathfrak{a}_{K-J}(\mathbf{x})$$

maps  $\overline{\mathbf{P}}_{d}$  to itself.

*Proof.* If  $g(x) \in P_d$  then  $g(x) \in \overline{P}_{2d}$ , and hence  $g(y)f(x,y) \in \overline{P}_{2d}$ . If  $g(y) = \sum_I b_J y^J$  then we can write

$$g(\mathbf{y})f(\mathbf{x},\mathbf{y}) = \sum_{I,J} \, b_J \, a_I(\mathbf{x}) \mathbf{y}^{I+J}.$$

The coefficient of  $\mathbf{y}^{\mathbf{K}}$  in this product is in  $\overline{\mathbf{P}}_{d}$ , and equals

$$\sum_{I+J=K} \, b_J \, \mathfrak{a}_I(x)$$

This is exactly T(g) since

$$\mathsf{T}(\mathfrak{g}(\mathbf{x})) = \mathsf{T}\left(\sum_{\mathbf{J}} \mathfrak{b}_{\mathbf{J}} \, \mathbf{x}^{\mathbf{J}}\right) = \sum_{\mathbf{J}} \mathfrak{b}_{\mathbf{J}} \mathfrak{a}_{\mathbf{K}-\mathbf{J}}(\mathbf{x})$$

**Corollary 11.98.** *If*  $f \in P_d$  *and* K *is an index set then the map* 

$$\mathbf{x}^{K-I} \mapsto rac{1}{I!} rac{\partial^I}{\partial \mathbf{x}^I} f(\mathbf{x})$$

*defines a map*  $\mathbf{P}_d \mapsto \mathbf{P}_d$ .

*Proof.* Since  $f(\mathbf{x}) \in \mathbf{P}_d$  we know that  $f(\mathbf{x}+\mathbf{y}) \in \mathbf{P}_{2d}$ . The Taylor series of  $f(\mathbf{x}+\mathbf{y})$  is

$$f(\mathbf{x} + \mathbf{y}) = \sum_{\mathbf{I}} \frac{\mathbf{y}^{\mathbf{I}}}{\mathbf{I}!} \frac{\partial^{\mathbf{I}}}{\partial \mathbf{x}^{\mathbf{I}}} f(\mathbf{x}).$$
(11.13.1)

The corollary now follows from the proof of Theorem 11.97.

# 11.14 The diamond product

Recall (§ 6.6.11) the diamond product  $\binom{x}{r} \diamond \binom{x}{s} = \binom{x}{r+s}$ . We show that  $f \diamond g$  has all real roots for certain sets of f and g by identifying it as a diagonal of a polynomial in  $\overline{\mathbf{P}}_3$ . In order to do this we need an identity [178]:

$$f \Diamond g = \sum_{i} \frac{(x+1)^{i} x^{i}}{i! i!} \mathsf{D}^{i} f \mathsf{D}^{i} g$$

It turns out that the signed version of this identity also gives a bilinear product that often preserves roots, so we will begin with it.

Lemma 11.99. Define a bilinear map

$$S(f,g) = \sum_{i} (-1)^{i} \frac{(x+1)^{i} x^{i}}{i! i!} D^{i} f D^{i} g$$

*If* (f, g) *has roots in any one of the four quarter planes below* <sup>2</sup> (see Figure 11.2) *then*  $S(f, g) \in \mathbf{P}$ .

- (1)  $(-\infty, 0) \times (-\infty, -1)$
- (2)  $(-\infty, -1) \times (-\infty, 0)$
- (3)  $(0,\infty) \times (-1,\infty)$
- (4)  $(-1,\infty) \times (0,\infty)$

*Proof.* We begin with two consequences of Taylor's theorem

$$\begin{split} f(x+xy) &= \sum_{i} \frac{x^{i} D^{i} f}{i!} y^{i} \\ g(x+z+xz) &= \sum_{j} \frac{(x+1)^{j} D^{j} g}{j!} z^{j} \end{split}$$

<sup>2</sup> There are examples where  $S(f,g) \notin P$  if f, g have roots in any one of the shaded regions of Figure 11.2



Figure 11.2: Domains of the signed diamond product

The signed diamond product is the signed diagonal in the product

$$f(x+xy)g(x+z+xz) = \left(\sum_{i} \frac{x^{i}D^{i}f}{i!}y^{i}\right) \cdot \left(\sum_{j} \frac{(x+1)^{j}D^{j}g}{j!}z^{j}\right) \quad (11.14.1)$$

For easy of exposition we define  $f(x, y) \in \overline{\mathbf{P}^{alt}}_2$  if and only if  $f(x, -y) \in \overline{\mathbf{P}}_2$ . The four domains of the lemma lead to the four cases:

 $\begin{array}{ll} (a) & f(\,x(y+1)\,)\in\overline{\textbf{P}^{alt}}_2 & g(\,(x+1)(y+1)-1\,)\in\overline{\textbf{P}^{alt}}_2 \\ (b) & g(\,x(y+1)\,)\in\overline{\textbf{P}^{alt}}_2 & f(\,(x+1)(y+1)-1\,)\in\overline{\textbf{P}^{alt}}_2 \\ (c) & f(\,x(y+1)\,)\in\overline{\textbf{P}}_2 & g(\,(x+1)(y+1)-1\,)\in\overline{\textbf{P}}_2 \\ (d) & g(\,x(y+1)\,)\in\overline{\textbf{P}}_2 & f(\,(x+1)(y+1)-1\,)\in\overline{\textbf{P}}_2 \end{array}$ 

For instance, suppose that (1) holds. If  $f \in \mathbf{P}^{(-\infty,0)}$  then  $f(x+xy) = f(x(y+1)) \in \overline{\mathbf{P}^{alt}}_2$ . If  $g(x) \in \mathbf{P}^{(-\infty,-1)}$  then  $g(x-1) \in \mathbf{P}^{(-\infty,0)}$ , and hence  $g(x+z+xz) = g((x+1)(z+1)-1) \in \overline{\mathbf{P}^{alt}}_2$ . Consequently, case (a) above holds. Both of the factors of the left hand side of (11.14.1) are in  $\overline{\mathbf{P}^{alt}}_2$ , so the product is as well. The signed diagonal of the right hand side of (11.14.1) is in **P** by Lemma 13.10.

**Proposition 11.100.** If  $f \times g \in (P^{(-1,\infty)} \times P^{pos}) \cup (P^{pos} \times P^{(-1,\infty)})$ then  $f \Diamond g \in P$ .

*Proof.* The cases are similar to Lemma 11.99. Suppose that  $f \in \mathbf{P}^{\text{pos}}$  and  $g \in \mathbf{P}^{(-1,\infty)}$ . Then we see that  $f(-x(y+1)) \in \overline{\mathbf{P}}_2$ . As before  $g(x + z + xz) \in \overline{\mathbf{P}}_2$ , so

$$h(x, y, z) = f(-x(y+1)) g(x+z+xz) \in \overline{\mathbf{P}}_2.$$

The signed diagonal of h(x, y, z) is the diamond product of f and g.

We now consider a different map  $m : \mathbf{P} \times \mathbf{P} \longrightarrow \mathbf{P}$ .

**Proposition 11.101.** Suppose  $\mathfrak{m}(f, g) = f(D)g$ , and  $T(x^n) = H_n$ . Then  $f \bigotimes_T^m g$ :  $P \times P \longrightarrow P$ . *Proof.* We will show that  $\oint_{T}^{m} g = (Tf)(2D)g(x)$  which implies the conclusion using Corollary 7.15 and Corollary 7.44. First of all, from  $(H_n)' = 2nH_{n-1}$  it follows that

$$\mathsf{D}^k \mathsf{H}_n = 2^k (\underline{n})_k \mathsf{H}_{n-k}$$

and consequently

$$\mathsf{T}^{-1}(\mathsf{D}^k \mathsf{H}_n) = 2^k (\underline{n})_k x^{n-k} = 2^k \mathsf{D}^k x^n$$

By linearity in  $H_n$  and  $x^n$ 

$$\mathsf{T}^{-1}(\mathsf{D}^{\mathsf{k}}\mathsf{T}\mathfrak{g}) = (2\mathsf{D})^{\mathsf{k}}\mathfrak{g}$$

By linearity in  $D^k$  it follows that for any polynomial h

$$\mathsf{T}^{-1}(\mathsf{h}(\mathsf{D})\mathsf{T}\mathfrak{g}) = \mathsf{h}(2\mathsf{D})\mathfrak{g}$$

and hence choosing h = Tf yields

$$\Gamma^{-1}(\mathsf{Tf}(\mathsf{D})\mathsf{Tg}) = (\mathsf{Tf})(\mathsf{2D})\mathsf{g}$$

If we use the difference operator  $\Delta(f) = f(x + 1) - f(x)$  instead of the derivative, then we have a similar result, except that the Hermite polynomials are replaced with the Charlier polynomials.

**Proposition 11.102.** Suppose  $\mathfrak{m}(f, g) = f(\Delta)g$ , and  $T(x^n) = C_n^{\alpha}$ . Then  $f \bigotimes_T^{\alpha} g$ :  $P^{\operatorname{alt}} \times P \longrightarrow P$ .

*Proof.* We will show that  $f \bigotimes_{T}^{\alpha} g = (Tf)(\Delta)g(x)$  which implies the conclusion using Corollary 7.15 and Corollary 7.45. Since  $\Delta(C_{n}^{\alpha}) = nC_{n-1}^{\alpha}$  it follows that

$$\Delta^k C_n^{\alpha} = (\underline{n})_k C_{n-k}^{\alpha}$$

and consequently

$$\mathsf{T}^{-1}(\Delta^k \, \mathsf{C}^{\alpha}_n) = (\underline{n})_k \mathsf{x}^{n-k} = \mathsf{D}^k \mathsf{x}^n$$

By linearity

$$\mathsf{T}^{-1}(\mathsf{h}(\Delta)\mathsf{T}\mathfrak{g}) = \mathsf{h}(\mathsf{D})\mathfrak{g}$$

and hence choosing h = Tf yields

$$\mathsf{T}^{-1}\mathsf{T}\mathsf{f}(\Delta)\mathsf{T}\mathsf{g} = (\mathsf{T}\mathsf{f})(\mathsf{D})\mathsf{g}$$

A diamond-type product associated with falling and rising factorials has elementary proofs of its properties. (But see Question 45.)

**Lemma 11.103.** If  $T: x^n \mapsto (\underline{x})_n$  and  $S: x^n \mapsto \langle \underline{x} \rangle_n$  then  $\diamondsuit_{TS} : P^{alt} \times P^{alt} \longrightarrow P^{alt}$  where  $\diamondsuit_{TS} (f \times g) = S^{-1}(Tf \times Tg).$ 

*Proof.* The proof follows from the diagram



## 11.15 Substituting into quadratic forms

We have seen that if  $f \in \mathbf{P}^{\text{pos}}$  then  $f(-xy) \in \pm \overline{\mathbf{P}}_2$ . Now xy is the quadratic form corresponding to the matrix  $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  which is a limit of negative subdefinite matrices. Here is a generalization:

**Lemma 11.104.** Suppose that the d by d matrix Q is the limit of negative subdefinite matrices. If  $f \in \mathbf{P}^{pos}$  then  $f(-\mathbf{x}Q\mathbf{x}^t) \in \pm \overline{\mathbf{P}}_d$ .

*Proof.* If we write  $f = \prod (x + c_i)$  where all  $c_i$  are positive, then each factor  $-xQx^t + c_i$  of  $f(-xQx^t)$  is in  $\pm \overline{P}_d$ , so their product is in it as well.

For instance, if we take  $Q = \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}$  with associated quadratic form xy + xy + zy then Q is a limit of negative subdefinite matrices since  $Q = J_3 - I_3$ . Consequently, if f(x) has all negative roots then  $f(-(xy + xy + yz)) \in \pm \overline{P}_3$ .

We have seen that if  $f(x) \in \mathbf{P}^{p \circ s}$  then  $f(-x^2) \in \mathbf{P}$ . Here's a different generalization to  $\mathbf{P}_d$ .

**Lemma 11.105.** Suppose that Q is an e by e negative subdefinite matrix. If  $f(\mathbf{x}) \in \mathbf{P}_d^{\text{pos}}(n)$  then  $f(-\mathbf{x}Q\mathbf{x}^t, \dots, -\mathbf{x}Q\mathbf{x}^t) \in \pm \mathbf{P}_e(2n)$ .

*Proof.* If  $f = \sum a_I x^I$  then

$$\mathfrak{g}(\mathbf{x}) = \mathfrak{f}(-\mathbf{x}Q\mathbf{x}^{\mathsf{t}},\ldots,-\mathbf{x}Q\mathbf{x}^{\mathsf{t}}) = \sum \mathfrak{a}_{I}(-\mathbf{x}Q\mathbf{x}^{\mathsf{t}})^{|I|}.$$

Since  $f \in \mathbf{P}_d^{\text{pos}}$  we know that f(x, ..., x) has all negative roots. For every c0 the polynomial  $-\mathbf{x}Q\mathbf{x}^t + c^2$  satisfies substitution. Consequently, for every root of f(x, ..., x) = 0 we substitute for all but one variable, and find two roots. This accounts for all 2n roots of g. Since the degree and positivity conditions are clearly satisfied for  $(-1)^n g$ , we see  $(-1)^n g \in \mathbf{P}_e(2n)$ .

The only bilinear forms in  $\overline{\mathbf{P}}_{2d}$  are products.

**Lemma 11.106.** If  $f = \sum_{i,j=1}^{d} a_{ij} x_i y_j \in \overline{P}_{2d}$  then there are non-negative  $b_i$ ,  $c_i$  so that

$$\sum_{i,j=1}^{d} a_{ij} x_i y_j = \pm (b_1 x_1 + \dots + b_d x_d) (c_1 y_1 + \dots + c_d y_d)$$

*Proof.* We will show that all two by determinants  $\begin{vmatrix} a_{i,k} & a_{i,l} \\ a_{j,k} & a_{j,l} \end{vmatrix}$  are zero. This implies that all rows are multiples of one another, and the conclusion follows. It suffices to show that  $\begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} = 0$ . If we set  $x_j = y_j = 0$  for j > 2 then

$$a_{11} x_1 y_1 + a_{12} x_1 y_2 + a_{21} x_2 y_1 + a_{22} x_2 y_2 \in \mathbf{P}_4.$$

If all of  $a_{11}$ ,  $a_{12}$ ,  $a_{21}$ ,  $a_{22}$  are zero then the determinant is zero, so we may assume that  $a_{11} > 0$ . By Lemma 11.15 we see that  $a_{12}$ ,  $a_{21}$ ,  $a_{22}$  are non-negative. Substituting  $x_2 = y_2 = 1$  and applying Corollary 11.7 shows that  $a_{11}a_{22} - a_{12}a_{21} \leq 0$ . If  $a_{12}$  is non-zero then substituting  $x_2 = y_1 = 1$  and using the lemma we conclude that  $a_{21}a_{12} - a_{11}a_{22} \leq 0$ . We get the same inequality if  $a_{21}$  is non-zero. Consequently, the determinant is zero.

#### 11.16 Simultaneous negative pairs of matrices

In Lemma 11.105 we substituted the *same* quadratic from into a polynomial. When can we substitute two different quadratic forms into a polynomial in  $P_2$  and have the resulting polynomial be in  $P_d$ ? We need a condition on pairs of quadratic forms.

**Definition 11.107.** If p(x), q(x) are polynomials with positive leading coefficients, then we say that they are a *simultaneous negative pair* if there is some value x such that  $p(x) \leq 0$  and  $q(x) \leq 0$ . If  $Q_1$  and  $Q_2$  are two matrices, we say that they are a *simultaneous negative pair of matrices* if for all vectors of constants **a** the polynomials  $\mathbf{x}_1^{\mathbf{a}}Q_1\mathbf{x}_1^{\mathbf{a}\,\mathbf{t}}$  and  $\mathbf{x}_1^{\mathbf{a}}Q_2\mathbf{x}_1^{\mathbf{a}\,\mathbf{t}}$  are a simultaneous negative pair.

The first question to address is when does  $\{Q, Q\}$  form a simultaneous negative pair?

**Lemma 11.108.** Suppose that Q has all positive entries.  $\{Q, Q\}$  is a simultaneous negative pair iff Q is negative subdefinite.

*Proof.* Assume that {Q, Q} is a simultaneous negative pair. We show that  $xQx' - c^2 \in P_d$ , which implies that Q is negative subdefinite. Since  $xQx' - c^2$  has positive homogeneous part, we need to verify substitution. Choose a vector **a** and consider  $x^aQx^{a'}$ . For large values of x this goes to infinity. Since there is a value of x making  $x^aQx^{a'}$  non-positive, it follows that for fixed **a**,  $x^aQx^{a'}$  takes on all positive values. Consequently  $x^aQx^{a'} - c^2$  has two real zeros, and so substitution is satisfied.

The converse is similar.

**Proposition 11.109.** Suppose that  $f \in \mathbf{P}_2^{pos}(n)$ . If  $Q_1$  and  $Q_2$  form a simultaneous negative pair of matrices then  $g(\mathbf{x}) = f(-\mathbf{x}Q_1\mathbf{x}^t, -\mathbf{x}Q_2\mathbf{x}^t)$  is in  $\pm P_d(2n)$ .

*Proof.* Choose a vector of constants **a** and consider the parametrized curve in the plane given by

$$\mathcal{C} = \left\{ (-\mathbf{x}_{1}^{\mathbf{a}} Q_{1} \mathbf{x}_{1}^{\mathbf{a} \, \mathsf{t}}, -\mathbf{x}_{1}^{\mathbf{a}} Q_{2} \mathbf{x}_{1}^{\mathbf{a} \, \mathsf{t}}) \mid \mathbf{x}_{1} \in \mathbb{R} \right\}.$$

Since  $Q_i$  has all positive coefficients, the limit of  $-\mathbf{x}_1^a Q_i \mathbf{x}_1^{a^t}$  as  $\mathbf{x}_1 \longrightarrow \infty$  is  $-\infty$ . Thus, the curve  $\mathcal{C}$  is eventually in the lower left quadrant. Since  $Q_1, Q_2$  are a simultaneous negative pair of matrices we know that  $\mathcal{C}$  meets the upper right quadrant. It then follows that  $\mathcal{C}$  meets each of the solution curves of f, since the graph of f meets the  $\mathbf{x}$  axis to the left of the origin, and the  $\mathbf{y}$  axis below the origin. Each solution curve yields two intersections, so we find all 2n solutions. As before,  $\pm g(\mathbf{x})$  satisfies the positivity condition, so  $\pm g \in \mathbb{P}_d(2n)$ .

**Example 11.110.** What are conditions on a pair of matrices that make them a simultaneous negative pair? Suppose  $Q_1 = \begin{pmatrix} c & v \\ v^t & C \end{pmatrix}$  and  $Q_2 = \begin{pmatrix} b & w \\ w^t & B \end{pmatrix}$  are negative subdefinite d by d matrices.

Choose a vector **a** of constants. We will find the value  $V_1$  of  $\mathbf{x}_1^a Q_2 \mathbf{x}_1^{a\,t}$  evaluated at the smallest solution to  $\mathbf{x}_1^a Q_1 \mathbf{x}_1^{a\,t} = 0$ , and the value  $V_2$  of  $\mathbf{x}_1^a Q_1 \mathbf{x}_1^{a\,t}$  evaluated at the smallest solution to  $\mathbf{x}_1^a Q_2 \mathbf{x}_1^{a\,t} = 0$ . If for every **a** either  $V_1$  or  $V_2$  is negative then  $Q_1, Q_2$  form a simultaneous negative pair. This will be satisfied if we can show that  $V_1 + V_2$  or  $V_1 V_2$  is never positive.

Without loss of generality we may scale by positive values, and thus assume that b = c = 1.

$$\begin{split} \boldsymbol{x}_{1}^{a}Q_{1}\boldsymbol{x}_{1}^{a\,t} &= (x_{1},\boldsymbol{a})\left(\begin{smallmatrix}1 & \nu\\ \nu^{t} & C\end{smallmatrix}\right)\left(\begin{smallmatrix}x_{1}\\a^{t}\end{smallmatrix}\right) \\ &= x_{1}^{2} + 2\boldsymbol{a}\nu^{t}x_{1} + \boldsymbol{a}C\boldsymbol{a}^{t} \\ \boldsymbol{x}_{1}^{a}Q_{2}\boldsymbol{x}_{1}^{a\,t} &= (x_{1},\boldsymbol{a})\left(\begin{smallmatrix}1 & \nu\\ \nu^{t} & B\end{smallmatrix}\right)\left(\begin{smallmatrix}x_{1}\\a^{t}\end{smallmatrix}\right) \\ &= x_{1}^{2} + 2\boldsymbol{a}\boldsymbol{w}^{t}x_{1} + \boldsymbol{a}B\boldsymbol{a}^{t} \end{split}$$

To find  $V_1$ , we solve for  $x_1$  and take the smaller root

$$\mathbf{x}_1 = -\mathbf{a}\mathbf{v}^{\mathsf{t}} - \sqrt{(\mathbf{a}\mathbf{v}^{\mathsf{t}})^2 - \mathbf{a}C\mathbf{a}^{\mathsf{t}}}$$

Since  $Q_1$  is negative subdefinite we know  $x_1$  is real. We then substitute into  $x_1^a Q_2 x_1^{at}$ . We then do the same for  $V_2$ . Simplification occurs, and

$$V_1 + V_2 = 2(\mathbf{a}\nu^t - \mathbf{a}w^t) \left( (\mathbf{a}\nu^t - \mathbf{a}w^t) + \sqrt{\mathbf{a}(\nu\nu^t - C)\mathbf{a}^t} - \sqrt{\mathbf{a}(ww^t - B)\mathbf{a}^t} \right)$$

Notice that if v = w then  $V_1 + V_2 = 0$ , so  $Q_1, Q_2$  form a simultaneous negative pair in this case. Reconsidering our earlier normalization, this shows that if two negative subdefinite matrices have an equal row then they form a simultaneous negative pair.

**Example 11.111.** In this example we will show that xy and yz are limits of simultaneous negative pairs. Let  $Q_1$  and  $Q_2$  be the quadratic forms corresponding to the matrices  $M_1$  and  $M_2$ :

$$M_{1} = \begin{pmatrix} e & 1 & e \\ 1 & e & e^{2} \\ e & e^{2} & e^{4} \end{pmatrix}$$
$$M_{2} = \begin{pmatrix} e^{4} & e^{2} & e \\ e^{2} & e & 1 \\ e & 1 & e \end{pmatrix}$$

We will show that  $M_1$  and  $M_2$  are a simultaneous negative pair for 0 < e < 1. When e = 0 the quadratic forms are 2xy and 2yz. The determinant of  $M_1$  and of  $M_2$  is  $(e-1)^2 e^3 (1+e)$ , which is positive for 0 < e < 1. Since the eigenvalues for e = .1 are (1.1, -0.9, -0.0008), we see that  $M_i$  is negative subdefinite for 0 < e < 1. Notice that  $Q_1(x, y, z) = Q_2(z, y, x)$ .

If we solve for y in  $Q_1$ , and substitute into  $Q_2$  we get

$$V_{1}(x,z) = \frac{(1-e)(x-z)}{e} \times \left[ (2+2e-e^{2}-e^{3}-e^{4}) x + (e^{2}+e^{3}-e^{4}) z + 2 (1+e) \sqrt{(1-e^{2})x^{2}+(e^{4}-e^{5})z^{2}} \right]$$

We next determine the sign of  $V_1$  in the xz plane. First, note that  $V_1(x, z) = 0$  has solution x = z. The remaining factor has the form  $ax + bz + (cx^2 + dz^2)^{1/2} = 0$ , where a, b, c, d are positive. The solution to such an equation is found to be of the form  $x = \alpha z$  and  $x = \beta z$ , . In our case, it is found that  $\alpha$  and  $\beta$  are positive,  $\alpha > \beta$ , and  $\alpha\beta = 1$ . The solution is only valid for negative z. Thus, the sign of  $V_1(x, z)$  is as follows. Consider the four rays in clockwise order around the origin:

- A:  $(0,0) (\infty,\infty)$
- B:  $(0,0) (-\infty, -\alpha\infty)$
- C:  $(0,0) (-\infty, -\infty)$
- D:  $(0,0) (-\infty, -\beta\infty)$

Then  $V_1$  is positive in the sectors AB and CD, and negative in the sectors BC and DA. By symmetry,  $V_2(x, z) = V_1(z, x)$ , and since  $\alpha\beta = 1$  the roots of the complicated factor yield lines with the same slope. Thus  $V_2(x, z)$  is negative in the sectors AB and CD, and positive in the sectors BC and DA. Consequently,  $V_1V_2$  is never positive.

We conclude that for every value of x, z we can find a value of y for which both quadratic forms are non-positive, so  $M_1$  and  $M_2$  form a negative simultaneous pair.

**Corollary 11.112.** *If*  $f(x,y) \in \mathbf{P}_2^{\text{pos}}(n)$  *then*  $f(-xz, -yz) \in \pm \overline{\mathbf{P}}_3(2n)$ .

**Corollary 11.113.** Suppose that the linear transformation  $x^i \mapsto \alpha_i x^i$  maps  $P^{pos}$  to itself. The linear transformation

$$x^i y^j \mapsto \alpha_{i+j} x^i y^j$$

maps  $\mathbf{P}_2^{pos} \longrightarrow P_2$ .

*Proof.* If  $f(x, y) = \sum a_{ij}x^iy^j$  then by Corollary 11.112 we know

$$\sum a_{ij}(-1)^{i+j} x^i y^j z^{i+j} \in \pm \overline{\mathbf{P}}_3$$

Applying the induced transformation  $z^k \mapsto \alpha_k z^k$  yields that

$$\sum a_{ij}(-1)^{i+j} x^i y^j \alpha_{i+j} z^{i+j} \in \pm \overline{\mathbf{P}}_3$$

Substituting z = -1 shows that

$$\sum a_{ij} x^i y^j \alpha_{i+j} \in \mathtt{P}_2$$

**Corollary 11.114.** The linear transformation  $x^iy^j \mapsto \frac{x^iy^j}{(i+j)!}$  maps  $\mathbf{P}_2^{\text{pos}}$  to itself.

Denote by T the transformation of Corollary 11.114. Here are some consequences:

1. If we take  $f(x, y) = (x + y + 1)^n$  then applying T yields

$$\sum_{0 \leqslant i,j \leqslant n} \frac{n!}{i!j!(i+j)!} x^i y^j \in \mathbf{P}_2^{\text{pos}}$$

2. If we take  $g\in I\!\!P^{pos}$  then  $g(-xy)\in \pm I\!\!P_2^{pos}.$  If  $g(-x)=\sum b_i x^i$  then

$$\sum b_i \frac{x^i y^i}{(2i)!} \in \mathbf{P}_2^{\text{pos}}$$

#### **11.17** The interior of $P_d$

We have seen (§ 1.1.6) that the polynomials in one variable with all distinct roots can be characterized as the interior of **P**. We can extend these ideas to  $\mathbf{P}_{d}$ .

The set of all polynomials of degree at most n in  $P_d$  is a vector space V of finite dimension, and so has the standard topology on it. A polynomial f of degree n is in the interior of  $P_d$ , written  $f \in \text{int } P_d$ , if there is an open

neighborhood 0 of f in V such that all polynomials in 0 are also in  $P_d$ . The boundary of  $P_d$  is defined to be  $P_d \setminus \text{int } P_d$ .

We say that  $f \leq g$  if there are open neighborhoods of f, g such that if  $\tilde{f}$  is in the open neighborhood of f, and  $\tilde{g}$  is in the open neighborhood of g then  $\tilde{f} \leq \tilde{g}$ . Both f and g are necessarily in the interior of  $\mathbf{P}_d$ .

Surprisingly, products of linear factors are on the boundary.

**Lemma 11.115.** If n > 1 and  $f = \prod_{k=1}^{n} (b_k + x_1 + a_{2k}x_2 + \dots + a_{dk}x_d)$  is in  $P_d$  then f is in the boundary of  $P_d$ .

*Proof.* We first make a small perturbation so that all the coefficients  $a_{ij}$  are distinct. Consider the perturbation

$$h = (\epsilon + (b_1 + x_1 + \dots)(b_2 + x_1 + \dots)) \prod_{k=3}^{n} (b_k + x_1 + a_{2k}x_2 + \dots + a_{dk}x_d)$$

If we set all but one of  $x_2, ..., x_d$  to zero, then Example 9.23 shows that we can choose  $\epsilon$  such that the first factor does not satisfy substitution.

There is an effective way of determining if a polynomial is in the interior of  $P_2$ . It is only practical for polynomials of small degree. In order to do this, we must introduce the resultant.

#### Definition 11.116. If

$$f(x) = a_0 + a_1x + \dots + a_nx^n$$
  
$$g(x) = b_0 + b_1x + \dots + b_mx^m$$

then the resultant of f and g is defined to be the determinant of the n + m by n + m matrix

The resultant  $R_f$  of f(x, y) and  $\frac{\partial f}{\partial y}(x, y)$  is a polynomial in one variable. The resultant has the property that  $R_f(a) = 0$  if and only if f(a, y) has a double root.

We can use the resultant to show that a polynomial satisfies substitution.

**Lemma 11.117.** If f(x, y) satisfies the following two conditions then f satisfies *y*-substitution:

- $f(\alpha, y)$  is in **P** for some  $\alpha$ .
- R<sub>f</sub> has no real roots.

*Proof.* If there is an a for which f(a, y) does not have all real roots, then as t goes from  $\alpha$  to a there is a largest value of t for which f(t, y) has all real roots. For this value we must have that f(t, y) has a double root, but this is not possible since the resultant  $R_f$  has no real roots.

We can extend this lemma to the case where  $R_f$  has some real roots. The proof is similar.

**Lemma 11.118.** Suppose f(x, y) has resultant  $R_f$ , and that the distinct roots of  $R_f$  are  $r_1 < r_2 < \cdots < r_s$ . If f satisfies the condition below then f satisfies y-substitution.

• Set  $r_0 = -\infty$  and  $r_{s+1} = \infty$ . For every  $0 \le i \le s$  there is an  $\alpha_i \in (r_i, r_{i+1})$  such that  $f(\alpha_i, y) \in \mathbf{P}$ .

**Lemma 11.119.** *If*  $f \in P_2$  *has the property that the resultant*  $\text{Res}(f, f_y)$  *has only simple roots then* f *is in the interior of*  $P_2$ .

*Proof.* The resultant is a continuous function of the coefficients, so if g is close to f then  $R_g$  is close to  $R_f$ . Consequently we can find an open neighborhood  $\bigcirc$  of f so that all polynomials in  $\bigcirc$  have a resultant with simple roots, and exactly as many roots as  $R_f$ . Moreover, we may choose  $\bigcirc$  small enough that each interval satisfies the condition of the previous lemma. We now apply the lemma to conclude that every function in  $\bigcirc$  satisfies y-substitution. We can also choose  $\bigcirc$  small enough to insure that all its members satisfy the homogeneity conditions, so that  $\bigcirc \subset P_2$ .

**Example 11.120.** Of course, the resultant must fail to show that linear products are in the interior, since they are on the boundary. The resultant of

$$(x + a_1 + y b_1) (x + a_2 + y b_2) (x + a_3 + y b_3)$$

is

$$-b_{1} b_{2} b_{3} (a_{1} - a_{2} + y b_{1} - y b_{2})^{2} (a_{1} - a_{3} + y b_{1} - y b_{3})^{2} (a_{2} - a_{3} + y b_{2} - y b_{3})^{2}$$

and in general

Resultant 
$$\left(\prod_{i=1}^{n} x + a_i + b_i y\right) = b_1 \cdots b_n \prod_{i < j} (a_j - a_i + y(b_j - b_i))^2$$

There are many values of y for which the resultant is 0, and they are all double roots, so the previous lemma doesn't apply.

## 11.18 The cone of interlacing

Lemma 1.20 in Chapter 1 explicitly describes all the polynomials that interlace a given polynomial of one variable. In this section we study the collection of all the polynomials that interlace a given polynomial in  $\overline{\mathbf{P}}_d$ . This set forms a cone, and in some cases we are able to determine its dimension.

**Definition 11.121.** Suppose that  $f \in \overline{\mathbf{P}}_d$ . The *interlacing cone of* f is defined to be

$$\mathbf{C}$$
 ONE $(\mathbf{f}) = \left\{ \mathbf{g} \in \overline{\mathbf{P}}_d \,|\, \mathbf{f} \leq \mathbf{g} \right\}.$ 

We should first note that if  $g_1, g_2 \in CONE(f)$  and a, b > 0 then  $g = ag_1 + bg_2$  satisfies  $f \leq g$  and  $g \in \overline{P}_d$ ., so CONE(f) is closed under positive linear combinations. Thus, CONE(f) is indeed a cone. In some cases we can explicitly determine the cone of interlacing. See Corollary 11.123.

The dimension of a cone is the least number of elements whose non-negative linear combinations span the cone. From Lemma 11.122 we see that dim  $C ONE(x^n y^m) = 1$ .

The first lemma is an analog of the one variable fact that all polynomials that interlace  $x^n$  are of the form  $cx^m$ .

**Lemma 11.122.** Suppose that  $f \in \overline{\mathbf{P}}_2$  has the property that f and  $x^n y^m$  interlace. If f has x-degree at most n, and y-degree at most m then there are a, b, c, d such that

$$f = x^{n-1}y^{m-1}(a + bx + cy + dxy).$$

*Proof.* For any  $\alpha \in \mathbb{R}$  we substitute to find that  $x^n \alpha^m$  and  $f(x, \alpha)$  interlace. Since the x-degree of f is at most n we know that there are constants  $r(\alpha)$  and  $s(\alpha)$  depending on  $\alpha$  such that

$$f(x, \alpha) = r(\alpha)x^n + s(\alpha)x^{n-1}.$$

However, we do not yet know that  $r(\alpha)$  and  $s(\alpha)$  are polynomials in  $\alpha$ . If we solve  $f(1, \alpha) = f(2, \alpha) = 0$  for  $r(\alpha)$  and  $s(\alpha)$  then we see that  $r(\alpha)$  and  $s(\alpha)$  are rational functions of  $\alpha$ . Thus,  $f(x, y) = x^{n-1}(r(y) + s(y)x)$  and so we can conclude that  $x^{n-1}$  divides f. Similarly,  $y^{m-1}$  divides f. Since  $x^{n-1}y^{m-1}$  divides f, the conclusion follows from the degree assumption.

#### Corollary 11.123.

$$\boldsymbol{C} ONE(\mathbf{x}^{n}\mathbf{y}^{m}) = \left\{ a \mathbf{x}^{n-1} \mathbf{y}^{m-1} w here \ a > 0 \right\}.$$

*Proof.* The x and y degrees of a polynomial g such that  $x^n y^m \leq g$  must be n-1 and m-1 respectively. Apply Lemma 11.122.

The next result is obvious in **P**. It's worth noting however that the positivity condition is not true for polynomials in one variable that do not have all real roots.<sup>3</sup>

<sup>&</sup>lt;sup>3</sup> The product of two polynomials with positive and negative coefficients can have all positive coefficients:  $(2x^2 + 3x + 2)(x^2 - x + 2) = 2x^4 + x^3 + 3x^2 + 4x + 4$ .

**Theorem 11.124.** *If*  $fg \in \mathbf{P}_2$  *then either*  $f, g \in \mathbf{P}_2$  *or*  $-f, -g \in \mathbf{P}_2$ .

*Proof.* From Lemma 10.4 we know that f and g satisfy substitution. It remains to show that  $\pm f^H$  has all positive coefficients. We know that  $(fg)^H = f^H g^H$ , and since  $fg \in \mathbf{P}_2$  we can factor

$$f^{H}g^{H} = (fg)^{H} = \prod_{i} (a_{i}x + b_{i}y)$$

where all  $a_i$ ,  $b_i$  are positive. It follows that  $f^H$  and  $g^H$  are products of some of the terms  $a_ix + b_iy$ , and the theorem is proved.

The following lemma is harder than it seems. See the short proof in [31].

**Lemma 11.125.** If f(x, y) is a continuous function defined for all x, y with the property that for any  $\alpha \in \mathbb{R}$  both  $f(x, \alpha)$  and  $f(\alpha, y)$  are polynomials then f is itself a polynomial.

We use the last lemma and the factorization property (Theorem 11.124) to determine what polynomials interlace  $f^n$ .

**Lemma 11.126.** If  $f, g \in \mathbf{P}_2$  and  $f^n \leq g$  then there is a  $g_1 \in \mathbf{P}_2$  such  $f \leq g_1$  and  $g = f^{n-1}g_1$ .

*Proof.* If we substitute  $y = \alpha$  then  $f(x, \alpha)^n \leq g(x, \alpha)$  so from Lemma 1.20 we conclude

$$g(\mathbf{x}, \alpha) = f(\mathbf{x}, \alpha)^{n-1} g_1(\mathbf{x}, \alpha)$$

where we only know that g(x, y) is a polynomial in x for any value of y. Similarly we get

$$g(\beta, y) = f(\beta, y)^{n-1}g_2(\beta, y)$$

where  $g_2(x, y)$  is a polynomial in y for any fixed x. Now

$$\frac{g(x,y)}{f(x,y)^{n-1}} = g_1(x,y) = g_2(x,y)$$

so  $g_1$  satisfies the hypothesis of Lemma 11.125 and hence is a polynomial in x and y. Finally, to see that  $f \leq g_1$  note that since  $f^n \leq f^{n-1}g_1$  we know that

$$f^{n-1}(f + \alpha g_1) \in \mathbf{P}_2$$

We now apply Theorem 11.124 to conclude that  $f + \alpha g_1 \in \mathbf{P}_2$ , and so  $f \leq g_1$ .  $\Box$ 

**Corollary 11.127.** C ONE $((x + y)^n) = \{ \alpha (x + y)^{n-1} \mid \alpha \in \mathbb{R} \}$ 

**Corollary 11.128.** If  $f \in \mathbf{P}_2$  then dim  $C ONE(f) = dim C ONE(f^k)$  for any  $k \ge 1$ .

Next, we have an analog of quantitative sign interlacing.

**Theorem 11.129.** *If*  $h \in \overline{P}_2$ ,  $f(x) = (x-a_1) \dots (x-a_n)$  and  $g(y) = (y-b_1) \dots (y-b_m)$  satisfy  $f(x)g(y) \leq h$  then

$$h(x,y) = \sum_{i,j} c_{ij} \frac{f(x)}{x - a_i} \frac{g(y)}{y - b_j}$$
(11.18.1)

where all  $c_{ij}$  are non-negative. Conversely, if h is given by (11.18.1) where all  $c_{ij}$  have the same sign or are zero then  $f(x)g(y) \leq h(x,y)$ .

*Proof.* If we fix y then from quantitative sign interlacing (Lemma 1.20) we can write

$$h(x,y) = \sum \alpha_i(y) \frac{f(x)}{x - a_i}$$

and by holding x fixed

$$h(x,y) = \sum \beta_j(x) \, \frac{g(y)}{y - b_j}$$

Substituting  $y = b_j$  yields

$$\begin{split} h(x,b_j) &= \beta_j(x)g'(b_j) \\ &= \sum \alpha_i(b_j) \, \frac{f(x)}{x-a_i} \end{split}$$

so  $\beta_j(x)$  is a polynomial in x. Next substituting  $x = a_i$  and then  $y = b_j$  gives

$$h(a_i, y) = \alpha_i(y)f'(a_i)$$
$$h(a_i, b_i) = \alpha_i(b_i)f'(a_i)$$

and so

$$\begin{split} h(x,y) &= \sum_{j} \frac{h(x,b_{j})}{g'(b_{j})} \frac{g(y)}{y-b_{j}} \\ &= \sum_{i,j} \frac{\alpha_{i}(b_{j})}{g'(b_{j})} \frac{f(x)}{x-a_{i}} \frac{g(y)}{y-b_{j}} \\ &= \sum_{i,j} \frac{h(a_{i},b_{j})}{f'(a_{i})g'(b_{j})} \frac{f(x)}{x-a_{i}} \frac{g(y)}{y-b_{j}} \end{split}$$

We now determine the sign of  $c_{ij}$ . The sign of  $h(a_i, b_j)$  isn't arbitrary, for we must have that  $f(x) \leq h(x, \alpha)$  for any  $\alpha$ , and thus all the coefficients

$$s_{i}(\alpha) = \sum_{j} \frac{h(a_{i}, b_{j})}{f'(a_{i})g'(b_{j})} \frac{g(\alpha)}{\alpha - b_{j}}$$

of  $\frac{f(x)}{x-\alpha_i}$  must have the same sign. If we evaluate  $s_i(x)$  at a root  $b_k$  of g we get

$$s_{i}(b_{k}) = \frac{h(a_{i}, b_{k})}{f'(a_{i})g'(b_{k})}g'(b_{k}) = \frac{h(a_{i}, b_{k})}{f'(a_{i})}$$

Consequently  $sgn s_i(b_k) = (-1)^{n+i}sgn(h(a_i, b_k))$  since the sign of  $f'(a_i)$  is  $(-1)^{n+i}$ . Thus the sign of  $h(a_i, b_k)$  is  $c_k(-1)^i$  where  $c_k$  depends on k. A similar argument shows that the sign is also  $d_i(-1)^k$ , so we conclude that the sign is  $e(-1)^{i+j}$  where e is a constant. This implies that the sign of

$$\frac{h(a_i, b_j)}{f'(a_i)g'(b_j)}$$

is constant, as desired. The converse follows easily.

**Corollary 11.130.** *If all*  $a_i$ ,  $b_j$  *are distinct*,  $f(x) = (x - a_1) \dots (x - a_n)$ , and  $g(y) = (y - b_1) \dots (y - b_m)$  then dim C ONE $(f(x)g(y)) \leq nm$ .

*Proof.* The basis for the cone is all the products  $\frac{f(x)}{x-a_i} \frac{g(y)}{y-b_i}$ .

# 11.19 Products of linear factors

In this section we look at the interlacing properties of products of linear factors. We construct polynomials in  $\overline{\mathbf{P}}_{2}^{\text{pos}}$  whose coefficients are mutually interlacing polynomials.

**Theorem 11.131.** Let  $p_1, \ldots, p_n$  and  $q_1, \ldots, q_n$  be two sequences of polynomials with positive leading coefficients, and define

$$F=\prod_1^n(p_i+q_iy)=f_0(x)+f_1(x)y+\cdots+f_n(x)y^n.$$

- 1. If  $p_i \leq q_i$  for i = 1, ..., n then  $f_i \leq f_{i+1}$  for i = 1, ..., n.
- 2. If  $p_i \leq q_i$  in  $P^{pos}$  for i = 1, ..., n then  $f_i \leq f_{i+1}$  for i = 1, ..., n.

*Proof.* If  $p_i \leq q_i$  then  $p_i + yq_i$  is in  $\overline{\mathbf{P}}_2$ , and so the product F is in  $\overline{\mathbf{P}}_2$ . The coefficients of F are in **P**, and hence adjacent coefficients interlace.

In the second case, we know that consecutive coefficients interlace - the problem is determining which direction. We prove this by induction. We need only to look at the two leading coefficients, so let

 $p = cx^{r} + dx^{r-1} + \cdots$   $q = \alpha x^{r} + \beta x^{r-1} + \cdots$   $f_{i} = a_{i}x^{n} + b_{i}x^{n-1} + \cdots$   $f_{i+1} = a_{i+1}x^{n} + b_{i+1}x^{n-1} + \cdots$
Consecutive terms of the product (p + yq)F are

$$(\alpha a_{i-1} + c a_i) x^{n+r} + (\beta a_{i-1} + d a_i + \alpha b_{i-1} + c b_i) x^{n+r+1} + \cdots$$
$$(\alpha a_i + c a_{i+1}) x^{n+r} + (\beta a_i + d a_{i+1} + \alpha b_i + c b_{i+1}) x^{n+r+1} + \cdots$$

The determinant of the first two coefficients is

 $\begin{vmatrix} \alpha a_{i-1} + c a_i & \beta a_{i-1} + d a_i + \alpha b_{i-1} + c b_i \\ \alpha a_i + c a_{i+1} & \beta a_i + d a_{i+1} + \alpha b_i + c b_{i+1} \end{vmatrix}$ 

which can be written as

$$\begin{vmatrix} c & \alpha \\ d & \beta \end{vmatrix} \begin{vmatrix} a_i & a_{i+1} \\ a_{i-1} & a_i \end{vmatrix} + \alpha^2 \begin{vmatrix} a_{i-1} & a_i \\ b_{i-1} & b_i \end{vmatrix} + c^2 \begin{vmatrix} a_i & a_{i+1} \\ b_i & b_{i+1} \end{vmatrix} + \alpha c \begin{vmatrix} a_{i-1} & a_{i+1} \\ b_{i-1} & b_{i+1} \end{vmatrix}$$

The first determinant is positive since  $p \leq q$ , the second is Newton's inequality, and the last three follow from  $f_i \leq f_{i+1}$ .

**Corollary 11.132.** Suppose that constants  $a_i, b_i, c_i, d_i$  have positive determinants  $\begin{vmatrix} b_i & a_i \\ c_i & d_i \end{vmatrix}$  and that  $a_i d_i$  is positive for  $1 \le i \le n$ . If

$$\prod_{i=1}^{n} (a_{i}x + b_{i} + (d_{i}x + c_{i})y) = f_{0} + f_{1}y + \dots + f_{n}y^{n}$$

then

$$f_0 \underline{\ll} f_1 \underline{\ll} \cdots \underline{\ll} f_n.$$

*Proof.* If we set  $p_i = d_i x + c_i$  and  $q_i = a_i x + b_i$  then the relationship  $p_i \leq q_i$  is equivalent to  $-c_i/d_i > -b_i/a_i$ . Since  $a_i d_i$  is positive the condition is equivalent to the positivity of the determinant. Now apply Theorem 11.131.

**Corollary 11.133.** Suppose that constants  $c_i$  and  $d_i$  satisfy

$$0 < d_1 < c_1 < d_2 < c_2 < \cdots < d_n < c_n$$

If

$$f = \prod_{i}^{n} (xy + x + c_iy + d_i) = f_0(x) + \dots + f_n(x)y^n$$

*then*  $(f_0, \ldots, f_n)$  *is mutually interlacing.* 

*Proof.* The determinants of the factors are  $\begin{vmatrix} 1 & 1 \\ c_i & d_i \end{vmatrix} = d_i - c_i > 0$ , so we can apply Corollary 11.132 to conclude that

$$f_0 \underline{\ll} f_1 \underline{\ll} \cdots \underline{\ll} f_n.$$

The hypothesis are that  $f_0 = \prod (x + d_i) \leq \prod (x + c_i) = f_n$ . It follows that this sequence of polynomials is mutually interlacing.

*Remark* 11.134. The interlacings in the last three results are strict if we assume that the interlacings in the hypothesis are strict.

What happens in Corollary 9.66 if some of the coefficients of x are zero? Theorem 11.135 shows that we still have interlacing, but some interlacings change from  $\ll$  to  $\ll$ .

**Theorem 11.135.** *Suppose that*  $f \in P$  *is a polynomial of degree* r *and let* 

$$f(y)\prod_{i=1}^{n}(y+a_{i}x+b_{i})=\sum_{i=0}^{n+r}p_{i}(x)y^{i}$$

Assume that  $\prod (a_i x + b_i)$  has all distinct roots, all  $a_i$  have the same sign  $\varepsilon$ , and write  $f(y) = \sum c_i y^i$ .

• If  $c_0c_1\varepsilon > 0$  then

$$p_0 \ll p_1 \ll \ldots \ll p_n \lessdot p_{n+1} \lessdot \ldots \lessdot p_{n+r-1}$$

• If  $c_0c_1 \varepsilon < 0$  then

$$\mathfrak{p}_0 \gg \mathfrak{p}_1 \gg \ldots \gg \mathfrak{p}_n \lessdot \mathfrak{p}_{n+1} \lessdot \ldots \lessdot \mathfrak{p}_{n+r-1}$$

*Proof.* Assume  $\epsilon > 0$ . Both f(y) and the product are in  $\overline{\mathbf{P}}_2$ , so all the coefficients interlace. It remains to see how they interlace. The first two coefficients are

$$\begin{split} p_0 &= c_0 \prod (a_i x + b_i) \\ p_1 &= c_1 p_0 + c_0 \sum_j \prod_{i \neq j} (a_i x + b_i) \\ &= c_1 p_0 + c_0 \sum_j \frac{p_0}{a_i x + b} \\ &= c_1 p_0 + c_0 \sum_j \frac{1}{a_i} \frac{p_0}{x + b/a_i}. \end{split}$$

Assume that  $c_0c_1 \epsilon > 0$ , so  $c_0, c_1 \neq 0$ . By hypothesis  $p_0$  has all distinct roots. The coefficient  $c_0/a_i$  has the same sign as  $c_1$ , so  $p_1 \gg p_0$ . Now apply Corollary 9.64. The remaining cases are similar.

#### 11.20 The interlacing of products

In one variable, every product  $\prod (x + a_i)$  is interlaced by many different products. In two variables, the only products  $\prod (x + a_iy + b_i)$  that are interlaced by products are images of one variable polynomials. In particular, most products in two variables are not interlaced by any other products. This is actually a result about lines in the plane, and the proof requires the fundamental result due to Sylvester and Gallai [1]:

In any configuration of n points in the plane, not all on a line, there is a line which contains exactly two of them.

Since this is a property of points, lines, and incidence, we can dualize:

In any configuration of n lines in the plane, not all parallel nor all coincident, there is a point which lies on exactly two of them.

**Theorem 11.136.** Suppose that  $f, g \in P_2(n)$  are products of linear factors, fg has no repeated factors, and f, g interlace. Then there are interlacing polynomials  $h, k \in P$  with homogenizations H, K, and constants  $\alpha, \beta$  such that either

$f(x,y) = h(x + \alpha y + \beta)$	$g(x,y) = k(x + \alpha y + \beta) \text{ or }$
$f(x, y) = H(x, \alpha y + \beta)$	$g(x, y) = K(x, \alpha y + \beta)$

*Proof.* The graph of f consists of n lines of negative slope in the plane, as does the graph of g. By assumption, all these lines are distinct. If the lines are all parallel then we get the first case of the conclusion, and if they are all coincident then we have the second case.

If we substitute any value for x then the resulting polynomials interlace, so we see that the intersections of the graph G of fg with any vertical line almost always consist of 2n points, and these points alternate between being roots of f, or roots of g. The same is true for horizontal lines.

Consider a point (u, v) of the plane that lies on an even number of lines of G. By Sylvester-Gallai we know that there is at least one such point. Slightly to the left of v a horizontal line meets the lines through (u, v) in points that (from the left) are in f, g, f, g, ... (say), and slightly to the right of v a horizontal line meets the lines through (u, v) in g, f, g, f, .... Since a horizontal line meets all the lines of G, we get the important conclusion that on the horizontal line through (u, v) all the lines of G must intersect, and all such intersections have even degree. A similar statement holds for the vertical lines. Figure 11.3 shows an example of such intersections, where solid lines are factors of f, and dashed lines are factors of g.



Figure 11.3: The graph of fg meeting a horizontal line

Now, take the leftmost such vertical line: this is the line through the intersection point of G that has an even number of lines through it, and has smallest x coordinate. On this line, take the point of intersecting lines with the largest y coordinate. Label the intersection (s, t). A line of G has negative slope, and can not meet the line y = t to the right of (s, t), and also the line x = s below (s, t). This implies that all lines pass through (s, t) and this contradiction establishes the theorem.

*Remark* 11.137. Although the lemma shows that there are no interesting interlacing products of linear terms, there are interesting *positive* interlacing products. Suppose that  $f = \prod_{1}^{n} (x + a_{i}y + b_{i})$  is a product in P<sub>2</sub>. We claim that

$$\frac{f(x,y)}{x+a_1+b_1} \stackrel{+}{\sim} \frac{f(x,y)}{x+a_2+b_2} \stackrel{+}{\sim} \cdots \stackrel{+}{\sim} \frac{f(x,y)}{x+a_n+b_n}$$

Indeed, each of them interlaces f, so any positive linear combination does as well. More generally, if  $M = I + xD_1 + yD_2$  where  $D_1$ ,  $D_2$  are positive definite and M[i] is the ith principle submatrix then

$$|\mathcal{M}[1]| \stackrel{+}{\sim} |\mathcal{M}[2]| \stackrel{+}{\sim} \cdots \stackrel{+}{\sim} |\mathcal{M}[n]|.$$

#### 11.21 Characterizing transformations

Just as in the one variable case, we can characterize linear transformations T of  $P_2$  that satisfy  $f \leq Tf$ . As expected, T is a linear combination of derivatives.

**Lemma 11.138.** If  $T : P_2 \longrightarrow P_2$  satisfies  $f \leq Tf$  for all  $f \in P_2$  then there are nonnegative c, d such that  $Tf = (c \frac{\partial}{\partial x} + d \frac{\partial}{\partial y})f$ .

*Proof.* It suffices to show it holds for polynomials of degree n. Let  $t_1, t_2$  be positive. Since

$$(t_1x+t_2y)^n \underline{\lessdot} T(t_1x+t_2y)^n$$

there is a constant  $\alpha_t$  depending on  $t_1$ ,  $t_2$  such that

$$T(t_1x + t_2y)^n = \alpha_t(t_1x + t_2y)^{n-1}.$$
 (11.21.1)

We will determine T by equating coefficients, but first we need to find  $\alpha_t$ . Since T decreases degree we can write

$$\mathsf{T}(x^iy^{n-i}) = \mathfrak{a}_i x^i y^{n-1-i} + \mathfrak{b}_i x^{i-1} y^{n-i}$$

Thus

$$\begin{aligned} \mathsf{T}(\mathsf{t}_{1}\mathsf{x} + \mathsf{t}_{2}\mathsf{y})^{\mathsf{n}} &= \mathsf{t}_{1}^{\mathsf{n}}\mathsf{T}(\mathsf{x}^{\mathsf{n}}) + \mathsf{n}\mathsf{t}_{1}^{\mathsf{n}-1}\mathsf{t}_{2}\,\mathsf{T}(\mathsf{x}^{\mathsf{n}-1}\mathsf{y}) + \cdots \\ &= \mathsf{t}_{1}\,\mathsf{b}_{\mathsf{n}}\mathsf{x}^{\mathsf{n}-1} + \mathsf{n}\mathsf{t}_{1}^{\mathsf{n}-1}\mathsf{t}_{2}\big(\mathfrak{a}_{\mathsf{n}-1}\mathsf{x}^{\mathsf{n}-1} + \mathfrak{b}_{\mathsf{n}-1}\mathsf{x}^{\mathsf{n}-2}\mathsf{y}\big) + \cdots \\ &= \alpha_{\mathsf{t}}\big(\mathsf{t}_{1}^{\mathsf{n}-1}\mathsf{x}^{\mathsf{n}-1} + (\mathsf{n}-1)\mathsf{t}_{1}^{\mathsf{n}-2}\mathsf{t}_{2}\,\mathsf{x}^{\mathsf{n}-2}\mathsf{y} + \cdots\big) \end{aligned}$$

Equating coefficients of  $x^{n-1}$  in (11.21.1) yields

$$\begin{split} \sum \binom{n}{i} t_1^i t_2^{n-i} \mathsf{T}(x^i y^{n-i}) \\ &= (t_1 b_n + \mathfrak{n} b_{n-1} t_2) \sum \binom{n-1}{j} t_1^j t_2^{n-1-j} \mathsf{T}(x^j y^{n-1-j}) \end{split}$$

and conbined with (11.21.2) yields

$$t_1b_n + nb_{n-1}t_2 = \alpha_t$$

Substituting this into (11.21.1) and equating coefficients of  $t_1^i t_2^{n-i}$  yields

$$\begin{split} \mathsf{T}(x^{i}y^{n-i}) &= \mathfrak{b}_{n}\,\mathfrak{i}\,x^{i-1}y^{n-i} + \mathfrak{a}_{n-1}\,(n-i)x^{i}y^{n-1-i} \\ &= \left(\mathfrak{b}_{n}\frac{\partial}{\partial x} + \mathfrak{a}_{n-1}\frac{\partial}{\partial y}\right)x^{i}y^{n-i} \end{split}$$

The conclusion now follows by linearity.

This holds more generally; we sketch the proof.

**Lemma 11.139.** If  $T : P_d \longrightarrow P_d$  satisfies  $f \leq Tf$  for all  $f \in P_d$  then there are non-negative  $c_i$  such that  $Tf = \left(\sum c_i \frac{\partial}{\partial x_i}\right) f$ 

Proof. As above,

$$\left(\sum_{i=1}^{n} t_{i}x_{i}\right)^{n} \leq T \left(\sum_{i=1}^{n} t_{i}x_{i}\right)^{n}$$
$$T \left(\sum_{i=1}^{n} t_{i}x_{i}\right)^{n} = \alpha_{t} \left(\sum_{i=1}^{n} t_{i}x_{i}\right)^{n-1}$$

Write

$$\mathsf{T}(\mathsf{x}_1^{i_1}\cdots \mathsf{x}_d^{i_d}) = \sum \mathfrak{a}_{\mathfrak{i}_1,\ldots,\mathfrak{i}_d;\mathfrak{j}} \mathsf{x}_1^{i_1}\cdots \mathsf{x}_{\mathfrak{j}}^{i_{\mathfrak{j}}-1}\cdots \mathsf{x}_d^{i_d}$$

Equating coefficients of  $x_1^{n-1}$  yields

$$\alpha_t = t_1 \mathfrak{a}_{n0\cdots 0} + n \sum_{j>1} t_j \mathfrak{a}_{n-10\cdots 1\cdots 0}$$

Finally, equating coefficients uniquely determines T on monomials, which completes the proof.  $\hfill \Box$ 

#### **11.22** Orthogonal polynomials in P<sub>d</sub>

There are many different definitions for orthogonal polynomials in several variables (see [53, 166]). We know two infinite families of orthogonal polynomials in more than one dimension that are also in  $P_d$ : the Hermite polynomials (§ 13.13.5) and Appell polynomials (this section).

Knowing that an orthogonal polynomial is in  $P_2$  allows us to recover some of the one dimensional properties concerning roots. Although such a polynomial does not factor into linear factors, all of its coefficients (of powers of either x or of y) factor into linear factors, and adjacent coefficients interlace.

We begin by studying Rodrigues' formulas in  $P_d$ . In one variable a Rodrigues' formula represents a family of polynomials by an expression of the

form  $ch^{-1}D^{n}(hp^{n})$  where h is a function, c is a constant, and p is a polynomial. Here are three classic examples:

Laguerre 
$$L_n^{(\alpha)}(x) = \frac{1}{n!} \frac{1}{e^{-x} x^{\alpha}} \left(\frac{d}{dx}\right)^n (e^{-x} x^{\alpha} x^n)$$
  
Hermite  $H_n(x) = (-1)^n \frac{1}{e^{-x^2}} \left(\frac{d}{dx}\right)^n \left(e^{-x^2}\right)$   
Jacobi  $P_n^{\alpha,\beta}(x) = \frac{(-1)^n}{2^n n!} \frac{1}{(1-x)^{\alpha}(1+x)^{\beta}} \times \left(\frac{d}{dx}\right)^n \left((1-x)^{\alpha}(1+x)^{\beta}(1-x^2)^n\right)$ 

The next proposition generalizes the Rodrigues' formula for Laguerre polynomials to define polynomials in d variables.

**Proposition 11.140.** Suppose that  $\mathbf{b} = (\beta_1, \dots, \beta_d)$  and  $\mathbf{I} = (\mathfrak{i}_1, \dots, \mathfrak{i}_d)$  are all non-negative, and all entries of  $\mathbf{a} = (\alpha_1, \dots, \alpha_d)$  are at least -1. If  $f \in \overline{\mathbf{P}}_d$  then the following is a polynomial in  $\overline{\mathbf{P}}_d$ :

$$\frac{1}{e^{-\mathbf{x}\cdot\mathbf{b}}\mathbf{x}^{\mathbf{a}}}\,\frac{\partial^{|I|}}{\partial\mathbf{x}^{I}}\,\left(e^{-\mathbf{x}\cdot\mathbf{b}}\mathbf{x}^{\mathbf{a}}\mathbf{f}\right)$$

*Proof.* To reduce the necessary notation, let's assume d = 2. We can not just apply results about differentiation of polynomials in  $\mathbf{\overline{P}}_2$  since the objects we are differentiating are not polynomials when the  $\alpha_i$  are not integers and the  $\beta_i$  aren't all zero. So, we enlarge our class of objects:

$$\mathsf{P}(\mathsf{r},\mathsf{s}) = \left\{ e^{-\beta_1 \varkappa - \beta_2 \vartheta} \mathsf{x}^{\alpha_1} \mathsf{x}^{\mathsf{r}} \mathsf{y}^{\alpha_2} \mathfrak{y}^{\mathsf{s}} \mathsf{f} \mid \mathsf{f} \in \overline{\mathbf{P}}_2 \right\}$$

We then are going to show that

$$\frac{\partial}{\partial x}: P(r,s) \longrightarrow P(r-1,s)$$
(11.22.1)  
$$\frac{\partial}{\partial y}: P(r,s) \longrightarrow P(r,s-1)$$
(11.22.2)

Since

$$e^{-\beta_1 x - \beta_2 y} x^{\alpha_1} x^r y^{\alpha_2} y^s f \in P(r, s)$$

it will follow that

$$\frac{\partial^{n+m}}{\partial x^n \partial y^m} \left( e^{-\beta_1 x - \beta_2 y} x^{\alpha_1} x^r y^{\alpha_2} y^s f \right) \in \mathsf{P}(0,0)$$

which implies the conclusion.

It suffices only to prove (11.22.1); the proof of (11.22.2) is similar. Differentiating,

$$\begin{split} \frac{\partial}{\partial x} \left( e^{-\beta_1 x - \beta_2 y} x^{\alpha_1} x^r y^{\alpha_2} y^s f \right) &= e^{-\beta_1 x - \beta_2 y} x^{\alpha_1} x^{r-1} y^{\alpha_2} y^s \quad \times \\ \left( -\beta_1 x f + (\alpha_1 + r) f + x \frac{\partial f}{\partial x} \right) \end{split}$$

Since r is at least 1,  $\alpha_1 + r$  is non-negative, so we know that that  $-\beta_1 x f + (\alpha_1 + r)f + x \frac{\partial f}{\partial x} \in \mathbf{P}_2$ , and hence (11.22.1) holds.

There are several varieties of Appell polynomials. We begin with the simple, no parameter version, and then introduce more complexity. The version below is in [166] for d = 2. If  $I = (i_1, ..., i_d)$  and  $x = (x_1, ..., x_d)$  then define

$$\begin{aligned} A_{\mathbf{I}}(\mathbf{x}) &= \frac{1}{\mathbf{I}!} \frac{\partial^{|\mathbf{I}|}}{\partial \mathbf{x}^{\mathbf{I}}} \left( (1 - \Sigma \mathbf{x})^{|\mathbf{I}|} \mathbf{x}^{\mathbf{I}} \right) \\ &= \frac{1}{i_1! \cdots i_d!} \frac{\partial^{i_1 + \cdots + i_d}}{\partial x_1^{i_1} \cdots \partial x_d^{i_d}} \left( (1 - x_1 - \cdots - x_d)^{i_1 + \cdots + i_d} x_1^{i_1} \cdots x_d^{i_d} \right) \end{aligned}$$

**Proposition 11.141.**  $A_I(\mathbf{x})$  is in  $\overline{\mathbf{P}}_d$ .

*Proof.* Since  $(1 - \Sigma \mathbf{x})^{|\mathbf{I}|}$  is in  $\mathbf{P}_d$ , the product with  $x_1^{i_1} \cdots x_d^{i_d}$  is in  $\overline{\mathbf{P}}_d$ . Since  $\overline{\mathbf{P}}_d$  is closed under differentiation,  $A_{\mathbf{I}}(\mathbf{x}) \in \overline{\mathbf{P}}_d$ .

Proposition 11.142. Adjacent Appell polynomials interlace.

*Proof.* Without loss of generality, we have to show that

$$A_{\mathfrak{i}_1,\ldots,\mathfrak{i}_d}(\mathbf{x}) \leq A_{\mathfrak{i}_1,\ldots,\mathfrak{i}_d-1}(\mathbf{x}).$$

Since we can ignore constant factors, we need to show that

$$\frac{\partial^{|\mathbf{I}|}}{\partial x_1^{i_1} \cdots \partial x_d^{i_d}} \left( (1 - \Sigma \mathbf{x})^{|\mathbf{I}|} \, x_1^{i_1} \cdots x_d^{i_d} \right) \leq \\ \frac{\partial^{|\mathbf{I}| - 1}}{\partial x_1^{i_1} \cdots \partial x_d^{i_d - 1}} \left( (1 - \Sigma \mathbf{x})^{|\mathbf{I}| - 1} \, x_1^{i_1} \cdots x_d^{i_d - 1} \right)$$

Since differentiation preserves interlacing, we need to show that

$$\frac{\partial}{\partial x_d} \left( \left. \left( 1 - \Sigma \boldsymbol{x} \right)^{|\boldsymbol{I}|} \boldsymbol{x}_1^{i_1} \cdots \boldsymbol{x}_d^{i_d} \right. \right) \underline{\ll} \left( \left. \left( 1 - \Sigma \boldsymbol{x} \right)^{|\boldsymbol{I}| - 1} \boldsymbol{x}_1^{i_1} \cdots \boldsymbol{x}_d^{i_d - 1} \right. \right)$$

Differentiating and factoring the left side yields

$$(1 - \Sigma \mathbf{x})^{|\mathbf{I}| - 1} \mathbf{x}_1^{\mathbf{i}_1} \dots \mathbf{x}_d^{\mathbf{i}_d - 1} \left( -|\mathbf{I}| \mathbf{x}_d + \mathbf{i}_d (1 - \Sigma \mathbf{x}) \right) \leq \\ \left( \left( 1 - \Sigma \mathbf{x} \right)^{|\mathbf{I}| - 1} \mathbf{x}_1^{\mathbf{i}_1} \dots \mathbf{x}_d^{\mathbf{i}_d - 1} \right)$$

Since the left side is a multiple of the right side by a linear term that is in  $P_{d}$ , the last interlacing holds, and so the proposition is proved.

The polynomials for d = 1, 2 are

$$\begin{aligned} A_{n}(x) &= \frac{1}{n!} \frac{d^{n}}{dx^{n}} \left( (1-x)^{n} x^{n} \right) \\ A_{n,m}(x_{1}, x_{2}) &= \frac{1}{n!m!} \frac{\partial^{n+m}}{\partial x_{1}^{n} \partial x_{2}^{m}} \left( (1-x_{1}-x_{2})^{n+m} x_{1}^{n} x_{2}^{m} \right) \end{aligned}$$

The one dimensional polynomials are a modified Legendre polynomial. The first few of the one and two dimensional polynomials are listed in Chapter 24.9 The Appell polynomials of different dimensions are related since  $A_{i_1,...,i_d}(x_1,...,x_d) = A_{i_1,...,i_d,0}(x_1,...,x_d,0)$ .

Next we consider Appell polynomials with extra parameters ([166]). Let  $\mathbf{a} = (\alpha_1, \dots, \alpha_d)$ , and define

$$\begin{split} \mathsf{A}_{I}^{a}(x) &= \frac{x^{-a}}{I!} \frac{\partial^{|I|}}{\partial x^{I}} \left( (1 - \Sigma x)^{|I|} x^{I+a} \right) \\ &= \frac{x_{1}^{-\alpha_{1}} \cdots x_{d}^{-\alpha_{d}}}{i_{1}! \cdots i_{d}!} \frac{\partial^{i_{1} + \cdots + i_{d}}}{\partial x_{1}^{i_{1}} \cdots \partial x_{d}^{i_{d}}} \\ &\qquad \qquad \left( (1 - x_{1} - \cdots - x_{d})^{i_{1} + \cdots + i_{d}} x_{1}^{i_{1} + \alpha_{1}} \cdots x_{d}^{i_{d} + \alpha_{d}} \right) \end{split}$$

**Proposition 11.143.** *If*  $\alpha_i \ge -1$  *for*  $1 \le i \le d$  *then*  $A_I^a(\mathbf{x}) \in \mathbf{P}_d$ 

*Proof.* Since  $(1 - \Sigma I)^{|I|} \mathbf{x}^{I} \in \overline{\mathbf{P}}_{d}$ , the result follows from Proposition 11.140 with b = 0.

If we attempt to generalize the Rodrigues' formula for the Laguerre polynomials with the definition

$$\frac{1}{e^{-x\cdot b}x^a}\,\frac{\partial^{|I|}}{\partial x^I}\,\left(e^{-b\cdot x}x^ax^I\right)$$

then it is easy to see that this factors into a product of one dimensional Laguerre polynomials. We use  $\Sigma x$  to force the  $x_i$ 's to be dependent. The following polynomials are in  $\overline{P}_d$ .

$$\frac{1}{e^{-\mathbf{x}\cdot \mathbf{b}} x^a} \; \frac{\partial^{|\mathbf{I}|}}{\partial x^{\mathbf{I}}} \; \left( e^{-\mathbf{b}\cdot \mathbf{x}} x^a (\boldsymbol{\Sigma} \mathbf{x})^{|\mathbf{I}|} \right)$$

Here is a two dimensional example that is (28) from [166, Page 159]. Since  $xy - 1 \in \overline{\mathbf{P}}_2$ , the following is in  $\overline{\mathbf{P}}_2$  for b > 0:

$$\frac{1}{e^{-b(x+y)}} \frac{\partial^{n+m}}{\partial x^n \partial y^m} \left( e^{-b(x+y)} (xy-1)^{n+m} \right)$$

The Rodrigues' formula for the Legendre polynomials is

$$\mathsf{P}_{\mathsf{n}}(\mathsf{x}) = \frac{1}{2^{\mathsf{n}} \mathsf{n}!} \left(\frac{\mathsf{d}}{\mathsf{d} \mathsf{x}}\right)^{\mathsf{n}} (\mathsf{x}^2 - 1)^{\mathsf{n}}$$

We can give a Rodrigues' formula for an analog of Legendre polynomials, but the resulting family does not appear to have any recursive properties. Let Q be a negative subdefinite symmetric d by d matrix, and let  $\mathbf{x} = (x_1, \dots, x_d)$ . For any index set I we define

$$\mathsf{P}_{\mathbf{I}}(\mathbf{x}) = \frac{\partial^{|\mathbf{I}|}}{\partial \mathbf{x}^{\mathbf{I}}} (\mathbf{x} Q \mathbf{x}^{\mathsf{t}} - 1)^{|\mathbf{I}|}$$

Since Q is negative subdefinite, the quadratic form  $\mathbf{x}Q\mathbf{x}^t - 1$  is in  $\mathbf{P}_d$ , and since multiplication and differentiation preserve  $\mathbf{P}_d$ , it follows that  $\mathbf{P}_I(\mathbf{x}) \in \mathbf{P}_d$ . *Remark* 11.144. There is a different definition of Appell polynomials in [53] that does not lead to polynomials in  $\overline{\mathbf{P}}_d$ . In the case that d = 2 these polynomials are, up to a constant factor, defined to be

$$U_{n,m}(x,y) = (1 - x^2 - y^2)^{-\mu + 1/2} \frac{\partial^{n+m}}{\partial x^n \, \partial y^m} \left(1 - x^2 - y^2\right)^{n+m+\mu - 1/2}$$

where  $\mu$  is a parameter. Since

$$U_{2,0}(x,y) = -\left((3+2\mu) \left(1 - (2+2\mu)x^2 - y^2\right)\right)$$

it is clear that  $U_{2,0}(x, y)$  does not satisfy substitution, and so is not in **P**<sub>2</sub>.

When we generalize the classical orthogonal polynomials, there are many choices and no one way of doing it. We can notice that these generalized Legendre polynomials are similar to the definition of Appell polynomial in [53], except that he uses a positive definite matrix (the identity) leading to the quadratic form (in two variables)  $x^2 + y^2 - 1$ . Replacing the identity by a negative subdefinite matrix leads to a definition of generalized Appell polynomials as

$$\frac{1}{(\mathbf{x}Q\mathbf{x}^{t}-1)^{\mu}}\partial^{I} (\mathbf{x}Q\mathbf{x}^{t}-1)^{|I|+\mu}$$

where  $\mu$  is a parameter. That this polynomial is in  $\mathbf{P}_d$  follows from the following theorem, whose proof is similar to Proposition 11.140 and omitted.

**Proposition 11.145.** *If*  $f \in \mathbf{P}_d$ ,  $\mu$  *any parameter,* I *any index set, then* 

$$\frac{1}{\mathsf{f}^{\mu}} \frac{\partial^{|I|}}{\partial x^{I}} \mathsf{f}^{|I|+\mu} \in \mathbf{P}_{\mathsf{d}}$$

#### **11.23** Arrays of interlacing polynomials

The polynomials determined by Rodrigues' type formulas form interlacing arrays. For instance, consider the polynomials

$$f_{n,m} = \frac{\partial^{n+m}}{\partial x^n \, \partial y^m} \, g^{n+m}$$

where  $g \in P_2$  is of degree 2. Since  $\frac{\partial g}{\partial x}$  is a polynomial of degree 1 in  $P_2$  we know that

$$g^{n+m-1}\frac{\partial g}{\partial x} \leq g^{n+m-1}$$

and therefore

$$\frac{\partial^{n+m-1}}{\partial x^{n-1} \partial y^m} \frac{\partial}{\partial x} g^{n+m} \leq \frac{\partial^{n+m-1}}{\partial x^{n-1} \partial y^m} g^{n+m-1}$$

which implies that  $f_{n,m} \leq f_{n-1,m}$ . These interlacing take the following form, where  $h \leftarrow k$  stands for  $h \leq k$ :



We consider two examples for g that are of the form  $\mathbf{x}Q\mathbf{x}^t - 1$  where Q is negative subdefinite.

**Example 11.146.** First, take g = xy - 1. In this case we have an explicit formula for  $f_{n,m}$ :

$$\begin{split} f_{n,m} &= \frac{\partial^{n+m}}{\partial x^n \, \partial y^m} \, (xy-1)^{n+m} \\ &= \sum \frac{\partial^{n+m}}{\partial x^n \, \partial y^m} \, \binom{n+m}{i} x^i y^i (-1)^{n-i} \\ &= \sum_{i=max(n,m)}^{n+m} \, \binom{n+m}{i} (\underline{n})_i (\underline{m})_i x^{i-n} y^{i-m} \\ &= (n+m)! \sum_{i=max(n,m)}^{n+m} \, \binom{i}{i-n,i-m} x^{i-n} y^{i-m} \end{split}$$

Using this formula we can verify that  $f_{n,m}$  satisfies a simple two term recurrence relation

$$f_{n,m} = \frac{1}{n} \left( (n+m)^2 y f_{n-1,m} - m(n+m)(n+m-1) f_{n-1,m-1} \right)$$

If we substitute y = 1 we get an array of one variable polynomials such that every polynomial interlaces its neighbors.

x	$2 x^2$	$6 x^3$
-2+4x	$-12  \mathrm{x} + 18  \mathrm{x}^2$	$-72 x^2 + 96 x^3$
-12 + 18  x	$24 - 144  x + 144  x^2$	$360 \mathrm{x} - 1440 \mathrm{x}^2 + 1200 \mathrm{x}^3$
$-72 + 96  \mathrm{x}$	$360 - 1440 \mathrm{x} + 1200 \mathrm{x}^2$	$-720 + 8640 \text{ x} - 21600 \text{ x}^2 + 14400 \text{ x}^3$

**Example 11.147.** Next, we consider the quadratic form corresponding to the negative subdefinite matrix  $\begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$ . If we define

$$f_{n,m} = \frac{\partial^{n+m}}{\partial x^n \, \partial y^m} \, (x^2 + y^2 + 4xy - 1)^{n+m}$$

then we again get an array of interlacing polynomials. The degree of  $f_{n,m}$  is n + m. In this case if we substitute y = 0 then we get an array of interlacing one variable polynomials, but there does not appear to be any nice recurrence relation.

1	4 x	$-4 + 36 x^2$
2 x	$-8 + 24 x^2$	$-216 \mathrm{x} + 408 \mathrm{x}^3$
$-4 + 12  x^2$	$-144 \mathrm{x} + 240 \mathrm{x}^3$	$432 - 4896  x^2 + 6000  x^4$
$-72 x + 120 x^3$	$288 - 2880  x^2 + 3360  x^4$	$24480  x - 120000  x^3 + 110880  x^5$

CHAPTER



# Polynomials satisfying partial substitution

In this chapter we are interested in polynomials in two kinds of variables that satisfy substitution only for positive values of the variables of the second kind. Such polynomials naturally arise from transformations that map  $P^{pos}$  to P.

## **12.1** Polynomials satisfying partial substitution in two variables

We introduce the class  $\mathcal{P}_{1,1}$  of polynomials f(x, y) that satisfy substitution for positive y.

**Definition 12.1.** The class  $\mathscr{P}_{1,1}$  consists of all polynomials f(x, y) satisfying

- 1. The homogeneous part  $f^{H}$  is in  $P^{pos}$ .
- 2.  $f(x, \beta) \in \mathbf{P}$  for all  $\beta \ge 0$ .

If all the coefficients are non-zero, then  $f \in \mathcal{P}_{1,1}^{pos}$ . The polynomials in the closure of  $\mathcal{P}_{1,1}$  are  $\overline{\mathcal{P}}_{1,1}$ , and those in the closure of  $\mathcal{P}_{1,1}^{pos}$  are  $\overline{\mathcal{P}}_{1,1}^{pos}$ .

We have the inclusion  $P_2 \subsetneq \mathcal{P}_{1,1}$ . The next example shows that  $\mathcal{P}_{1,1}$  is not closed under differentiation, and so the containment is strict.

**Example 12.2.**  $\mathscr{P}_{1,1}$  is not closed under differentiation with respect to y. However, if  $f \in \mathscr{P}_{1,1}(n)$  then there is an  $\alpha \ge 0$  such that  $\frac{\partial f}{\partial y}(x, y + \alpha) \in \mathscr{P}_{1,1}(n)$ .

The calculations below show that  $f \in \mathcal{P}_{1,1}$ , but that  $\frac{\partial f}{\partial y} \notin \mathcal{P}_{1,1}$ . The calculations do show that  $\frac{\partial f}{\partial y}(x, y + \alpha) \in \mathcal{P}_{1,1}$  for  $\alpha \ge 1/8$ .

$$\begin{split} \mathsf{f} &= (-3 + x + y)^2 \, \left( 18 - 19 \, x + 5 \, x^2 - 22 \, y + 10 \, x \, y + 5 \, y^2 \right) \\ \mathsf{f}_y &= 2 \, \left( -3 + x + y \right) \, \left( 51 - 45 \, x + 10 \, x^2 - 48 \, y + 20 \, x \, y + 10 \, y^2 \right) \\ \text{roots of f} : & \left\{ 3 - y, 3 - y, \frac{19 - 10 \, y \pm \sqrt{1 + 60 \, y}}{10} \right\} \\ \text{roots of f}_y : & \left\{ 3 - y, \frac{45 - 20 \, y \pm \sqrt{15} \, \sqrt{-1 + 8 \, y}}{20} \right\} \end{split}$$

Every vertical line  $x = \beta$  to the left of the smallest root of f(x, 0) has n - 1 roots of  $\frac{\partial f}{\partial y}(\beta, y)$ . Since these n - 1 solution curves are asymptotic to lines whose slopes lie between the slopes of the asymptotes of f, every sufficiently positive horizontal line will meet all n - 1 curves.

What do the graphs of polynomials in  $\mathcal{P}_{1,1}$  look like? The solution curves do not necessarily go from the upper left quadrant to the lower right quadrant as is the case for polynomials is  $\mathbf{P}_2$ , but they can loop back to the upper left quadrant.



Figure 12.1:  $L_5(x; y)$  and its derivative (dashed) with respect to y

In general, since  $f^H$  has all negative roots, the asymptotes of f are lines in the upper left quadrant with negative slopes. As  $x \to -\infty$ , the solution curves eventually have positive y-coordinates. Thus, there is an  $\alpha_1$  such that  $f(\alpha, y) \in \mathbf{P}^{\alpha lt}$  for all  $\alpha < \alpha_1$ , and also an  $\alpha_0$  such that  $f(\alpha, y) \in \mathbf{P}$  for all  $\alpha < \alpha_0$ . In Figure 12.1  $\alpha_0$  is about .1, and  $\alpha_1$  is about -12.6. A formal proof of this general fact follows the argument of Theorem 9.27.

#### 12.2 Examples

Polynomials satisfying partial substitution arise naturally in several ways.

**Example 12.3.** The generalized Laguerre polynomials  $L_n(x; y)$  are in  $\mathcal{P}_{1,1}$  since their homogeneous part is  $\frac{1}{n!}(x + y)^n$ , and they they satisfy the three term recurrence [168, (5.1.10)]

$$\begin{split} nL_n(x;y) &= (-x+2n+y-1)L_{n-1}(x;y) - (n+y-1)L_{n-2}(x;y) \\ L_0(x;y) &= 1 \quad L_1(x;y) = x+y+1 \end{split}$$

The following lemma generalizes the fact that three term recurrences determine orthogonal polynomials.

**Lemma 12.4.** Suppose that  $a_i, b_i, d_i, e_i$  are positive for  $i \ge 0$ . If

$$\begin{split} f_n(x;y) &= (a_n x + b_n y + c_n) f_{n-1}(x;y) - (d_n y + e_n) f_{n-2}(x;y) \\ f_0(x;y) &= 1 \quad f_{-1}(x;y) = 0 \end{split}$$

*then*  $f_n(x; y) \in \mathcal{P}_{1,1}$  *for*  $n = 0, 1, \ldots$ 

*Proof.* The construction yields polynomials that satisfy the degree requirements, and the homogeneous part is  $\prod_i (a_ix + b_iy)$ . If y is positive then the recurrence defines a three term recurrence for orthogonal polynomials, so they are in **P**.

Note that if  $\alpha$  is sufficiently negative, then  $f_n(x; \alpha)$  satisfies a recurrence that does not define a sequence of orthogonal polynomials, so we do not expect any polynomials from this construction to lie in P<sub>2</sub>.

**Example 12.5.** We can construct two variable Charlier polynomials from the one variable polynomials by

$$C_n(x;y) = (-1)^n C_n^y(-x).$$

They also satisfy a recurrence relation  $C_{-1}(x; y) = 0$ ,  $C_0(x; y) = 1$  and

$$C_{n+1}(x;y) = (x + y + n) C_n(x;y) - ny C_{n-1}(x;y).$$

It follows from the Lemma that  $C_n(x; y) \in \mathfrak{B}_{1,1}$ . Figure 12.2 shows  $C_5(x; y)$ .

If  $f \in \mathbf{P}^{pos}$  then  $f(-xy) \in \overline{\mathbf{P}}_2$ , and consequently is in  $\overline{\mathcal{P}}_{1,1}$ . However, more is true:

**Lemma 12.6.** If  $f \in \mathbf{P}$ , then  $f(xy) \in \overline{\mathcal{P}}_{1,1}$ . If  $f \in \mathbf{P}^{pos}$ , then  $f(xy) \in \overline{\mathcal{P}}_{1,1}^{pos}$ .



Figure 12.2: The Charlier polynomial  $C_5(x; y)$ 

*Proof.* Since  $\mathscr{P}_{1,1}$  is closed under multiplication, the comment above shows that it suffices to show that  $1 + xy \in \overline{\mathscr{P}}_{1,1}$ . Consider  $f_{\varepsilon}(x, y) = 1 + (x + \varepsilon y)(y + \varepsilon x)$ . For positive  $\varepsilon$  the graph of  $f_{\varepsilon}$  is a hyperbola, and the minimum positive value is found to be  $d_{\varepsilon} = \frac{2\sqrt{\varepsilon}}{|1-\varepsilon^2|}$ . Since the asymptotes have slope  $-\varepsilon$  and  $-1/\varepsilon$ , the polynomial  $f_{\varepsilon} - d_{\varepsilon}$  meets every horizontal line above the x-axis in two points, and the degree and homogeneity conditions are met, so  $f_{\varepsilon} - d_{\varepsilon} \in \mathscr{P}_{1,1}$ . Since  $\lim_{\varepsilon \to 0} (f_{\varepsilon} - d_{\varepsilon}) = f$  it follows that  $f \in \overline{\mathscr{P}}_{1,1}$ .

**Example 12.7.** We can construct polynomials that satisfy partial substitution but not the homogeneity condition in a manner that is similar to the construction of polynomials in  $\mathbf{P}_2^{\text{pos}}$ . Suppose that A is a positive definite n by n matrix, and B is a symmetric matrix that has positive and negative eigenvalues. Define f(x, y) = |I + yA + xB|. If y is positive then I + yA is positive definite, and so all roots of f(x, y) are real. The homogeneous part of f(x, y) is |yA + xB|. Since A is positive definite, the roots of  $f^H(x, 1)$  are the roots of  $|I + xA^{-1/2}BA^{-1/2}|$ . Since B has positive and negative eigenvalues, Sylvester's law of inertia shows that  $A^{-1/2}BA^{-1/2}$  does also. Consequently, the roots of  $f^H$  are not all the same sign, and thus  $f \notin \mathcal{P}_{1,1}$ .

Consider a concrete example. A was chosen to be a random positive definite matrix, C was a random symmetric matrix, D was the diagonal matrix with diagonal (1, 1, 1, -1), and B = CDC. If f(x, y) = |I + yA + xB|, then Figure 12.3 shows that  $f(x, y) \notin P_2$ . (The dashed line is the x-axis.)

$$\begin{split} f(x,y) &= 1 + 11.42 \, x + 4.84 \, x^2 - 7.13 \, x^3 - 3.67 \, x^4 + 15.48 \, y \\ &\quad + 42.56 \, x \, y - 9.60 \, x^2 \, y - 21.88 \, x^3 \, y + 12.05 \, y^2 + 13.80 \, x \, y^2 \\ &\quad - 5.94 \, x^2 \, y^2 + 1.99 \, y^3 + 0.96 \, x \, y^3 + 0.07 \, y^4 \end{split}$$



Figure 12.3: A polynomial of the form |I + yA + xB|

We summarize these observations in a lemma. The last two parts are proved in the same way as the results for  $P_2$ .

**Lemma 12.8.** *If* A *is positive definite, and* B *is symmetric with positive and negative eigenvalues then* f(x, y) = |I + yA + xB| *satisfies* 

- 1. f(x, y) has asymptotes of positive and negative slope.
- 2.  $f(\mathbf{x}, \alpha) \in \mathbf{P}$  for  $\alpha \ge 0$ .
- 3.  $f(\alpha, y) \in \mathbf{P}$  for all  $\alpha$ .
- 4.  $f(\alpha x, x) \in P$  for all  $\alpha$ .

#### 12.3 Induced transformations

We now look at induced linear transformations on  $\mathbf{P}^{pos}$ .

**Lemma 12.9.** If  $T: \mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}^{\text{pos}}$ , T preserves degree and the sign of the leading coefficient, then  $T_*: \mathcal{B}_{1,1}^{\text{pos}} \longrightarrow \mathcal{B}_{1,1}^{\text{pos}}$ . If  $T: \mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}$  then  $T_*: \mathcal{B}_{1,1}^{\text{pos}} \longrightarrow \mathcal{B}_{1,1}$ .

*Proof.* The hypotheses imply that  $T_*f$  satisfies homogeneity assumption. We next verify substitution for y. For  $\beta \ge 0$  we have

$$(\mathsf{T}_*\mathsf{f}(\mathsf{x},\mathsf{y}))(\mathsf{x},\mathsf{\beta}) = \mathsf{T}(\mathsf{f}(\mathsf{x},\mathsf{\beta})) \in \mathbf{P}^{\mathsf{pos}}$$

since  $f(x, \beta) \in \mathbf{P}^{pos}$ , and consequently  $T_*(f) \in \mathscr{O}_{1,1}^{pos}$ . The second part is similar.

We can easily determine the slopes of the asymptotes of  $T_*f$ . Suppose that  $f = \sum a_{i,j}x^iy^j$ , and  $c_i$  is the leading coefficient of  $T(x^i)$ . Then

$$(\mathsf{T}_*\mathsf{f})^\mathsf{H}(x,1) = \left(\sum a_{i,j}\mathsf{T}(x^i)y^j\right)^\mathsf{H}(x,1) = \sum a_{i,n-i}\,c_ix^i$$

The slopes of the asymptotes are the roots of this last polynomial.

**Corollary 12.10.** Suppose  $T: \mathbf{P}^{pos} \longrightarrow \mathbf{P}^{pos}$ , T preserves degree and the sign of the leading coefficient. If  $f \in \mathbf{P}^{pos}$  then  $T_*f(x + y) \in \mathcal{B}_{1,1}^{pos}$ .

*Proof.* Since 
$$f(x + y) \in \mathcal{P}_{1,1}^{pos}$$
 we can apply Lemma 12.9.

**Lemma 12.11.** Choose a positive integer n, and define the linear transformation  $S: x^i \mapsto T(x^i)y^{n-i}$ . If  $T: \mathbf{P}^{pos} \longrightarrow \mathbf{P}^{pos}$  preserves degree and the sign of the leading coefficient then  $S: \mathbf{P}^{pos} \longrightarrow \mathcal{O}_{1,1}^{pos}$ .

*Proof.* If  $f \in \mathbf{P}^{pos}$  then the homogenization F of f is in  $\mathcal{P}_{1,1}$ . Since  $S(f) = T_*(F)$  we can apply Lemma 12.9 to conclude that  $T_*(F) \in \mathcal{P}_{1,1}^{pos}$ .

**Corollary 12.12.** If  $f \in P^{pos}(n)$  and  $S(x^i) = \langle \underline{x} \rangle_i y^{n-i}$  then  $S(f) \in \mathscr{B}_{1,1}^{pos}$ .

*Proof.* The map  $x^i \mapsto \langle \underline{x} \rangle_i$  satisfies the hypotheses of Lemma 12.14.

**Example 12.13.** If we choose  $f = (x - 1)^n$  then from the corollary

$$S((x+1)^n) = \sum_{i=0}^n \langle \underline{x} \rangle_i y^{n-i} \binom{n}{i} = n! \sum_{i=0}^n \binom{x+i-1}{i} \frac{y^{n-i}}{(n-i)!}$$

The last polynomial is in  $\mathscr{P}_{1,1}^{pos}$ .

**Lemma 12.14.** If  $T: \mathbf{P}^{pos} \longrightarrow \mathbf{P}^{pos}$  and we define a linear transformation  $S(x^n) = y^n [T(x^n)(x/y)]$  then  $S: \mathbf{P}^{pos} \longrightarrow \mathcal{P}_{1,1}$ .

*Proof.* if  $f(x) = \sum a_i x^i$  is in  $\mathbf{P}^{\text{pos}}$  then

$$S(f)(x, \alpha) = \sum_{k=0}^{n} \alpha_{i} \alpha^{i} [T(x^{i})(x/\alpha)]$$
$$= T(f(\alpha x))(x/\alpha)$$

The last expression is in  $\mathbf{P}^{\text{pos}}$  for positive  $\alpha$ . Finally,  $S(f)^{\text{H}}$  is the homogenization of  $T(x^n)$ , and has all positive coefficients since T maps to  $\mathbf{P}^{\text{pos}}$ .

#### **12.4** Applications to transformations

The graph of a polynomial in  $\mathcal{P}_{1,1}$  goes up and to the left. If we intersect it with an appropriate curve we will find sufficiently many intersections that determine a polynomial in **P**. Using these techniques we can establish mapping properties of transformations.

**Lemma 12.15.** If  $f \in \mathcal{P}_{1,1}$ ,  $f(\alpha, y) \in P$  for  $\alpha \leq 0$ , and  $f(0, y) \in P^{pos}(n)$  then  $f(x, x) \in P^{pos}$ .

*Proof.* We will show that the graph of f intersects the line y = x in n points that have negative x coordinate. We know that the graph of f has asymptotes that have negative slope, so f is eventually in the third quadrant. However, the assumption that  $f(x, 0) \in \mathbf{P}^{p \circ s}$  means that all solution curves begin on the negative y axis. Consequently, all the curves must intersect the line y = x. See Figure 12.4.



Figure 12.4: The intersection of f(x, y) = 0 and y = x.

**Lemma 12.16.** Suppose that  $T: \mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}^{\text{pos}}$ , where T preserves degree, and  $T(x^i)$  has leading coefficient  $c_i$ . Choose  $f \in \mathbf{P}^{\text{pos}}(n)$ , and define  $S(x^i) = T(x^i)x^{n-i}$ . If  $\sum c_i a_i x^i \in \mathbf{P}^{(-\infty,-1)}$  where  $f = \sum a_i x^i$  then  $S(f) \in \mathbf{P}^{\text{pos}}$ .

*Proof.* Since  $f \in \mathbf{P}^{pos}$  we know that the homogenization F of f is in  $\mathcal{P}_{1,1}$ . From Lemma 12.9 the induced map  $T_*$  satisfies  $T_*(F) \in \mathcal{P}_{1,1}$ . By hypothesis, the asymptotes all have slope less than -1. Each solution curve meets the x-axis in  $(-\infty, 0)$  since  $(T_*(F))(x, 0) = T(x^n) \in \mathbf{P}^{pos}$ . Consequently, the graph of  $T_*F$  meets the line x = -y in n points, and these are the roots of S(f). See Figure 12.5.

**Corollary 12.17.** The linear transformation,  $V : x^i \mapsto \langle \underline{x} \rangle_i x^{n-i}$  maps  $P^{(0,1)}$  to  $P^{\text{pos}}$ . Similarly,  $U : x^i \mapsto (\underline{x})_i x^{n-i}$  maps  $P^{(0,1)}(n)$  to  $P^{\text{alt}}$ .



Figure 12.5: The intersection of  $T_*(F)$  and y = -x.

*Proof.* If we define  $T(x^i) = \langle \underline{x} \rangle_i$  and F is the homogenization of f then  $T_*(F)(x, x) = V(f)$ . If  $f \in \mathbf{P}^{(0,1)}(n)$  then  $F \in \mathbf{P}_2(n)$  and so from Corollary 12.12 we know that  $T_*(F) \in \mathcal{P}_{1,1}$ . Since  $(T_*(F))(x, 0) = \langle \underline{x} \rangle_n$  the graph of  $T_*(F)$  meets the y axis below the line x = -y. Since the leading coefficient of  $\langle \underline{x} \rangle_i$  is 1 the asymptotes of  $T_*(F)$  are the roots of f, which are between 0 and 1. Thus  $T_*(F)$  is asymptotic to lines that lie below x = -y, and so every solution curve of  $T_*(F)$  meets x = -y above the x axis. See Figure 12.6.



Figure 12.6: The intersection of  $T_*(F)$  and y = -x.

**Lemma 12.18.** If  $T: \mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}$  preserves degree,  $T(x^n)$  is monic, a > 0, and we define  $V(x^n) = x^n [T(x^n)(-a/x)]$  then  $V: \mathbf{P}^{(-a,0)} \longrightarrow \mathbf{P}$ .

*Proof.* If we define  $S(x^n) = y^n [(Tx^n)(x/y)]$  then V(f) = (Sf)(-a, x). Since  $T(x^n)$  is monic we find that (Sf)(x, 0) = f(x), so the roots of (Sf)(x, 0) are exactly the roots of f. Consequently, the vertical line through -a is to the left of the intersection of the graph of Tf with the x-axis, and so meets all the solution curves since  $Sf \in \mathcal{P}_{1,1}$  by Lemma 12.14.

**Corollary 12.19.** If  $V : x^n \mapsto -(x-1)(2x-1)\cdots((n-1)x-1)$  then  $V : \mathbf{P}^{(-1,0)} \longrightarrow \mathbf{P}^{(0,1)}$ .

The transformation  $x^n \mapsto (x+1)(2x+1)\cdots((n-1)x+1)$  maps  $P^{(0,1)} \longrightarrow P^{(-1,0)}$ .

*Proof.* We know  $V : \mathbf{P}^{(-1,0)} \longrightarrow \mathbf{P}$  by applying the Lemma to  $T : x^n \mapsto \langle \underline{x} \rangle_n$ . To see that V has range  $\mathbf{P}^{(0,1)}$  we consider the intersection of the graph with the line x = -y. If  $f = \sum a_i x^i$  and S is as in the Lemma then

$$S(x^n) = (x + y)(x + 2y) \cdots (x + ny)$$

so  $(Sf)(x, -x) = a_0 + a_1x$ . Since  $f \in \mathbf{P}^{p \circ s}$ ,  $a_0$  and  $a_1$  are positive, and so the graph of Vf does meets the line x = -y once in the upper left quadrant.

The homogeneous part of S(f) is  $x(x + y)(x + 2y) \cdots (x + (n - 1)y)$ , so the graph of S(f) has asymptotes with slopes  $1, 1/2, \cdots, 1/(n-1)$ , and one vertical asymptote. For positive y the graph of S(f) consists of n curves starting at the roots of f. The vertical asymptote begins below y = -x and so meets the line y = -x. Thus, the other curves lie entirely below y = -x, and consequently the line x = -1 meets these solution curves beneath the line y = -x, and hence all the roots of V are in (0, 1).

Replace x by -x for the second statement.

For example, consider Figure 12.7, the graph of Sf, where f(x) is the polynomial  $(x + .3)^3(x + .9)^3$ . The two dots are are at (-.9, 0) and (-.3, 0) which correspond to the roots of f. Except for the vertical asymptote, the graph of Sf lies under the line x = -y. The dashed line x = -1 meets the graph in points whose y coordinates lie in (0, 1).



Figure 12.7: An example for Corollary 12.19

#### 12.5 Möbius transformations

We use properties of the graph of  $T_*f(xy)$  to establish properties of Möbius transformations acting on linear transformations. We know from Corollary 9.57 that if T maps  $\mathbf{P}^{\text{pos}}$  to itself then so does  $T_{1/z}$ . It is more complicated when the domain is not  $\mathbf{P}^{\text{pos}}$ .

The graph of  $T_*f(xy)$  is different from the graphs of Figure 12.1 and Figure 12.4 since there are vertical and horizontal asymptotes. Figure 12.8 is the graph of  $T_*(f(xy))$  where  $T(x^i) = \langle \underline{x+1} \rangle_i$  and  $f \in \mathbf{P}^{pos}$  has degree 3.



Figure 12.8: A graph of  $T_*(f(xy))$  where  $f \in \mathbf{P}^{pos}$ 

**Lemma 12.20.** Let  $T: \mathbf{P}^{pos} \longrightarrow \mathbf{P}^{pos}$  where T preserves degree, and  $c_i$  is the leading coefficient of  $T(x^i)$ . For any positive integer n and  $f = \sum a_i x^i$  in  $\mathbf{P}^{pos}(n)$ 

- 1.  $(\mathsf{T}_*\mathsf{f}(\mathsf{x}\mathsf{y}))(\mathsf{x},\beta) \in \boldsymbol{P}^{\mathsf{pos}}$  for all  $\beta > 0$ .
- 2.  $T_*f(xy) \in \overline{\mathcal{P}}_{1,1}$
- 3. The vertical asymptotes of  $T_*f(xy)$  occur at the roots of  $T(x^n)$ .
- 4. The horizontal asymptotes are of the form xy = s, where s is a root of  $\sum a_i c_i x^i$ .

*Proof.* The first part follows from

$$T_*(f(xy))(x,\beta) = \sum a_i T(x^i)\beta^i = \sum a_i T(\beta^i x^i) = T(f(\beta x)).$$

As the coefficient of  $y^n$  is  $T(x^n)$  there are vertical asymptotes at the roots of  $T(x^n)$ . We know that  $f(xy) \in \mathscr{P}_{1,1}^{pos}$  so  $T_*f(x,y) \in \mathscr{P}_{1,1}^{pos}$ . The horizontal asymptotes are governed by the terms with equal x and y degrees. The polynomial so determined is  $\sum a_i c_i x^i y^i$ , which establishes the last part. Lemma 2.4 shows that all the  $c_i$  have the same sign, so s is negative.

If we assume that  $f \in \mathbf{P}^{alt}$  then Figure 12.8 is reflected around the y axis. Figure 12.9 shows the hyperbola and the intersection points with the graph that give the roots in the next lemma.

**Corollary 12.21.** Let  $T: \mathbf{P}^{\text{alt}} \longrightarrow \mathbf{P}^{\text{alt}}$  where T preserves degree, and  $c_i$  is the leading coefficient of  $T(x^i)$ . For any  $f = \sum a_i x^i$  in  $\mathbf{P}^{\text{alt}}$  such that  $\sum c_i a_i x^i$  is in  $\mathbf{P}^{(0,1)}$  the polynomial  $T_{1/2}(f)$  is in  $\mathbf{P}^{\text{alt}}$ .

*Proof.* As usual let  $T_*(x^iy^j) = T(x^i)y^j$ . If we define  $T(x^i) = p_i$  then

$$T_{1/z}(f) = \sum a_i x^i p_i(1/x) = \sum a_i [T_*(x^i y^i)(1/x, x)] = T_*(f(xy))(1/x, x)$$

Thus, the roots of  $T_{1/z}$  are found at the intersection of  $T_*(f(xy))$  and the hyperbola xy = 1. As long as the solution curves of  $T_*(f(xy))$  go to zero more rapidly than 1/x then xy = 1 will intersect every solution curve. By Lemma 12.20 the horizontal asymptotes are of the form xy = s where s is a root of  $\sum c_i a_i x^i$ . By assumption 0 < s < 1, and hence  $T_{1/z}(f)$  has all real roots.



Figure 12.9: The roots of  $T_{1/z}(f)$  for  $f \in \mathbf{P}^{alt}$ 

Similarly, we have

**Corollary 12.22.** Let  $T: \mathbf{P}^{pos} \longrightarrow \mathbf{P}^{pos}$  where T preserves degree, and  $c_i$  is the leading coefficient of  $T(x^i)$ . For any  $f = \sum a_i x^i$  in  $\mathbf{P}^{pos}$  such that  $\sum c_i a_i x^i$  is in  $\mathbf{P}^{(-1,0)}$  the polynomial  $T_{-1/z}(f)$  is in  $\mathbf{P}^{pos}$ .

Here is another consequence of changing signs.

**Corollary 12.23.** Let  $T: \mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}^{\text{alt}}$  where T preserves degree, and  $c_i$  is the leading coefficient of  $T(x^i)$ . For any  $f = \sum_{i=1}^{n} a_i x^i$  in  $\mathbf{P}^{\text{alt}}$  such that  $\sum_{i=1}^{n} c_i a_i (-x)^i$  is in  $\mathbf{P}^{(0,1)}$  the polynomial  $T_{1/z}(f)$  is in  $\mathbf{P}^{\text{alt}}$ .

*Proof.* If we let S(x) = T(-x) then S satisfies the hypothesis of Corollary 12.22. The conclusion follows since  $S_{-1/z} = T_{1/z}$ .

We can apply Corollary 12.21 to the monic Laguerre and Charlier polynomials (Corollary 7.47 and Corollary 7.45) since all the  $c_i$  are 1.

**Corollary 12.24.** If  $\tilde{L}_{n}^{(\alpha)}$  is the monic Laguerre polynomial then the transformation  $x^{i} \mapsto \tilde{L}_{i}(\alpha; x)^{\text{REV}}$  maps  $P^{(0,1)}$  to P.

**Corollary 12.25.** If  $C_n^{\alpha}$  is the Charlier polynomial then the transformation  $x^i \mapsto C_i^{\alpha \text{ REV}}$  maps  $P^{(0,1)}$  to P.

In the falling factorial the constant terms of the factors changed. Now we let the coefficient of x change. Consider the linear transformation

$$T(x^{n}) = (1-x)(1-2x)(1-3x)\cdots(1-nx)$$
(12.5.1)

**Lemma 12.26.** The linear transformation (12.5.1) maps  $P^{(0,1)}$  to P.

*Proof.* If  $S(x^n) = (\underline{x-1})_n$  then  $T = S_{1/z}$ . The result now follows from Corollary 12.19.

Next we consider the Möbius transformation  $z \longrightarrow (1 - z)$ .

**Lemma 12.27.** Suppose that  $T: \mathbf{P}^{pos} \longrightarrow \mathbf{P}$ , and let  $f(x) = \sum a_i x^i \in \mathbf{P}^{pos}(n)$ . Define  $c_i$  to be the leading coefficient of  $T(x^i)$ . We can conclude that  $\sum a_i(1 - x)^{n-i}T(x^i) \in \mathbf{P}$  if any of the following hold:

- 1.  $T(\mathbf{x}^n) \in \mathbf{P}^{(-\infty,1)}$  and  $\sum a_i c_i \mathbf{x}^i \in \mathbf{P}^{(-\infty,-1)}$ .
- 2.  $T(x^n) \in \mathbf{P}^{(1,\infty)}$  and  $\sum a_i c_i x^i \in \mathbf{P}^{(-1,0)}$ .
- 3.  $T(x^n) \in \mathbf{P}^{(-\infty,1)}$  and  $\sum a_i c_i x^i \in \mathbf{P}^{\text{alt}}$ .

*Proof.* We need to show that each solution curve of  $T_*(f)$  meets the line x + y = 1. Since  $T: \mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}$  there are solution curves for all positive y. The asymptotes of  $T_*(f)$  are the roots of  $\sum a_i c_i x^i$ , and  $T_*(f)(x, 0) = T(x^n)$  so the geometry of the solution curves is given in Figure 12.10. It is clear that each solution curve must meet the line x + y = 1.

#### 12.6 Partial substitution in several variables

We now generalize  $\mathscr{P}_{1,1}$  to more variables. In short, we consider polynomials in two kinds of variables such that if we substitute arbitrary values for all but one of the x variables, and only non-negative values for all the y variables, then the resulting polynomial is in **P**. In more detail,

**Definition 12.28.** The class  $\mathscr{P}_{d,e}$  consists of all polynomials  $f(\mathbf{x}; \mathbf{y})$  where  $\mathbf{x} = (x_1, \dots, x_d)$  and  $\mathbf{y} = (y_1, \dots, y_e)$  satisfying



Figure 12.10: The geometry of the solution curves of Lemma 12.27

- 1. The total degree equals the degree of each variable.
- 2. The homogeneous part  $f^H$  is in  $P_{d+e}$ .
- 3.  $f(\mathbf{x}; \beta) \in P_d$  where all  $\beta = (\beta_1, \dots, \beta_e)$  are non-negative.

We define  $\overline{\mathcal{P}}_{d,e}$  to be the set of all polynomials that are the limit of polynomials in  $\mathcal{P}_{d,e}$ . The set of all polynomials in  $\mathcal{P}_{d,e}$  with all positive coefficients is  $\mathcal{P}_{d,e}^{\text{pos}}$ .

Notice that  $\mathscr{P}_{d,0}^{pos}$  consists of polynomials in  $\mathbb{P}_d$  with all non-negative coefficients, and is exactly  $\mathbf{P}_{d+e}^{pos}$ . We have the inclusions

$$\mathbf{P}_{d+e}^{\text{pos}} = \mathcal{P}_{d+e,0}^{\text{pos}} \subseteq \mathcal{P}_{d+e-1,1}^{\text{pos}} \subseteq \cdots \subseteq \mathcal{P}_{1,d+e-1}^{\text{pos}}$$

If d, e > 1 then substitution of positive values determines two maps

$$\mathcal{P}_{d,e}^{\text{pos}} \longrightarrow \mathcal{P}_{d-1,e}^{\text{pos}}$$
 (substituting for an x value)  
 $\mathcal{P}_{d,e}^{\text{pos}} \longrightarrow \mathcal{P}_{d,e-1}^{\text{pos}}$  (substituting for a y value)

 $\mathcal{P}_{d,e}$  and  $\mathcal{P}_{d,e}^{pos}$  are closed under multiplication. We can not define interlacing in  $\mathcal{P}_{d,e}^{pos}$  as closed under all linear combinations, since we don't allow negative coefficients. If f,  $g \in \mathcal{P}_{d,e}^{pos}$  we define  $f \stackrel{+}{\sim} g$  to mean that  $f + \alpha g \in \mathcal{P}_{d,e}^{pos}$ for all non-negative  $\alpha$ . For example, if  $\ell$  is linear and in  $\mathcal{P}_{d,e}^{pos}$ , and  $f \in \mathcal{P}_{d,e}^{pos}$ , then  $\ell f \stackrel{+}{\sim} f$ .

A linear polynomial  $c+\sum a_i x_i + \sum b_i y_i$  is in  $\mathcal{P}_{d,e}$  if and only if it is in  $\mathbb{P}_{d+e}$ . The quadratic case is much more interesting, and is considered in Section 12.7.

It is easy to construct polynomials in  $\mathcal{P}_{d,e}$  by generalizing the inductive construction of Lemma 12.4. The proof is straightforward and omitted.

**Lemma 12.29.** Suppose that  $\mathbf{a}_i, \mathbf{b}_i, \mathbf{d}_i, \mathbf{e}_i$  are vectors of positive constants for  $i \ge 0$ , and that  $\mathbf{c}_i$  can be be positive or negative. If

$$\begin{split} f_n(\mathbf{x};\mathbf{y}) &= (\mathbf{a}_n \cdot \mathbf{x} + \mathbf{b}_n \cdot \mathbf{y} + \mathbf{c}_n) f_{n-1}(\mathbf{x};\mathbf{y}) - (\mathbf{d}_n \cdot \mathbf{y} + \mathbf{e}_n) f_{n-2}(\mathbf{x};\mathbf{y}) \\ f_0(\mathbf{x};\mathbf{y}) &= 1 \quad f_1(\mathbf{x};\mathbf{y}) = \mathbf{a}_1 \cdot \mathbf{x} + \mathbf{b}_1 \cdot \mathbf{y} + \mathbf{c}_1 \end{split}$$

then  $f_n(\mathbf{x}; \mathbf{y}) \in \mathcal{P}_{d,e}$  for  $n = 0, 1, \ldots$ 

#### 12.7 Quadratic forms

Our goal in this section is to determine conditions for the quadratic form  $\begin{pmatrix} \mathbf{x} & \mathbf{y} \end{pmatrix} Q \begin{pmatrix} \mathbf{x} & \mathbf{y} \end{pmatrix}^{t}$  to belong to  $\mathcal{P}_{d,e}$ . We first consider the case when d = 1, and then when e = 1.

A matrix Q is called *copositive* if  $xQx^t \ge 0$  for every vector x with all nonnegative entries. Unlike positive definite matrices, there is no good characterization of copositive matrices.

**Lemma 12.30.** Suppose that  $Q = \begin{pmatrix} a & v \\ v^{t} & C \end{pmatrix}$ , where C is symmetric and a is positive. The quadratic form  $f(x; y) = (x, y)Q(x, y)^{t}$  has all real roots for all non-negative y if the matrix  $v^{t}v - aC$  is copositive.

*Proof.* Expanding the quadratic form yields

$$(\mathbf{x}, \mathbf{y}) \begin{pmatrix} \mathbf{a} & \mathbf{v} \\ \mathbf{v}^{t} & \mathbf{C} \end{pmatrix} \begin{pmatrix} \mathbf{x} \\ \mathbf{y}^{t} \end{pmatrix} = \mathbf{a}\mathbf{x}^{2} + \mathbf{x}(\mathbf{y}\mathbf{v}^{t} + \mathbf{v}\mathbf{y}^{t}) + \mathbf{y}\mathbf{C}\mathbf{y}^{t}$$

Since  $\mathbf{y}\mathbf{v}^{t} = \mathbf{v}\mathbf{y}^{t}$ , the discriminant of the equation is  $\mathbf{y}(\mathbf{v}^{t}\mathbf{v} - \mathbf{a}C)\mathbf{y}^{t}$ .

**Lemma 12.31.** Suppose that  $Q = \begin{pmatrix} A & u \\ u^{t} & c \end{pmatrix}$  where all entries are positive, A is a d by d symmetric matrix, and c is a scalar. The quadratic form  $f(\mathbf{x}; \mathbf{y}) = (\mathbf{x} \ \mathbf{y}) Q (\mathbf{x} \ \mathbf{y})^{t}$  is in  $\mathcal{P}_{d,1}$  iff Q is negative subdefinite.

*Proof.* If Q is negative subdefinite then  $f(x, y) \in P_{d+1}$ , and since all terms are positive it is in  $\mathcal{P}_{d,1}$ .

For the converse, we only need to show that  $f(\mathbf{x}; \mathbf{y}) \in P_{d+1}$ , since this implies that Q is negative subdefinite. Since all entries of Q are positive,  $f(\mathbf{x}, \mathbf{y})$  satisfies the degree condition. Since  $f(\mathbf{x}, \mathbf{y})^H = f(\mathbf{x}, \mathbf{y})$ , it suffices to show that  $f(\mathbf{x}, \mathbf{y})$  satisfies substitution for any choice of values for  $\mathbf{x}$  and  $\mathbf{y}$ . By hypothesis,  $f(\mathbf{x}, 1) \in P_d$ . Now since  $f(\mathbf{x}; z) = z^2 f(\frac{\mathbf{x}}{z}, 1)$  and  $f(\frac{\mathbf{x}}{z}; 1)$  satisfies substitution, it follows that  $f(\mathbf{x}; z)$  satisfies substitution.

**Proposition 12.32.** Suppose that  $Q = \begin{pmatrix} A & u \\ u^t & C \end{pmatrix}$  where all entries are positive, A is a d by d symmetric matrix, and C is an e by e symmetric matrix. The quadratic form

$$\begin{pmatrix} \mathbf{x} & \mathbf{y} \end{pmatrix} Q \begin{pmatrix} \mathbf{x}^{t} \\ \mathbf{y}^{t} \end{pmatrix}$$

is in  $\mathcal{P}_{d,e}$  iff the two conditions are met:

- 1. A is negative subdefinite.
- 2.  $uA^{-1}u^t C$  is copositive.

*Proof.* If the quadratic form is in  $\mathcal{P}_{d,e}$  then substituting  $\mathbf{y} = 0$  shows that  $\mathbf{x}A\mathbf{x}^t \in P_d$ , so A is negative subdefinite.

If we substitute non-negative values  $\beta$  for **y**, expand and regroup the quadratic form, we get

$$\begin{pmatrix} \mathbf{x} & \beta \end{pmatrix} \begin{pmatrix} \mathbf{A} & \mathbf{u} \\ \mathbf{u}^{t} & \mathbf{C} \end{pmatrix} \begin{pmatrix} \mathbf{x} \\ \beta \end{pmatrix} = \mathbf{x} \mathbf{A} \mathbf{x}^{t} + \beta \mathbf{u}^{t} \mathbf{x}^{t} + \mathbf{x} \mathbf{u} \beta^{t} + \beta \mathbf{C} \beta^{t}$$
$$= \begin{pmatrix} \mathbf{x} & 1 \end{pmatrix} \begin{pmatrix} \mathbf{A} & \mathbf{u} \beta^{t} \\ \beta \mathbf{u}^{t} & \beta \mathbf{C} \beta^{t} \end{pmatrix} \begin{pmatrix} \mathbf{x} \\ 1 \end{pmatrix}$$
(12.7.1)

By Lemma 12.31,  $f(\mathbf{x}, z)$  is in  $P_{d+1}$  iff  $\begin{pmatrix} A & u\beta^t \\ \beta u^t & \beta C\beta^t \end{pmatrix}$  is negative subdefinite. By Lemma 10.78 this is the case iff

$$0 \leq (\mathfrak{u}\beta^{t})(A^{-1})(\beta\mathfrak{u}^{t}) - (\beta C\beta^{t}) = \beta(\mathfrak{u}A^{-1}\mathfrak{u}^{t} - C)\beta^{t}$$

which finishes the proof.

*Remark* 12.33. Here is an alternative derivation in the case that A is 2 by 2. We only need to determine when the quadratic form satisfies substitution. This will be the case iff the discriminant, with respect to the variable  $x_1$ , is positive. This discriminant is a quadratic form, and its discriminant, with respect to  $x_2$ , equals

$$a |A| \mathbf{y}(\mathbf{u}A^{-1}\mathbf{u}^{t} - C)\mathbf{y}$$

The condition that this is non-positive is equivalent to the positivity condition since A is negative subdefinite, and therefore |A| is negative.

*Remark* 12.34. Let's work out an example in light of this theory. We start with

$$g(x;y) = x^2 + 2xy + y^2 + 4x + 3y + 2$$

and the homogenized form is

$$G(x; y, z) = x^{2} + 2xy + y^{2} + 4xz + 3yz + 2z^{2}$$
$$= (x, y, z) \begin{pmatrix} 1 & 1 & 2\\ 1 & 1 & 3/2\\ 2 & 3/2 & 2 \end{pmatrix} (x, y, z)^{t}$$

To show that this is in  $\mathcal{P}_{1,2}$  we note A = (1), u = (1,2),  $C = \begin{pmatrix} 1 & 3/2 \\ 3/2 & 2 \end{pmatrix}$ 

$$\mathfrak{u} A^{-1} \mathfrak{u}^{t} - C = \begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix} - \begin{pmatrix} 1 & 3/2 \\ 3/2 & 2 \end{pmatrix} = \begin{pmatrix} 0 & 1/2 \\ 1/2 & 2 \end{pmatrix}$$

which is copositive. Substituting z = 1 shows  $g(x; y) \in \mathcal{P}_{1,1}$ .

*Remark* 12.35. A quadratic form in  $\mathscr{P}_{1,1}$  is in  $\mathbb{P}_2$ . To see this, let  $Q = \begin{pmatrix} a & b \\ b & c \end{pmatrix}$ , and assume  $(x y)Q(x y)^t \in \mathscr{P}_{1,1}$ . Applying Proposition 12.32 we see that  $b^2/a - c > 0$  or equivalently  $\begin{vmatrix} a & b \\ b & c \end{vmatrix} < 0$ , and hence Q is negative subdefinite. The example in the previous remark is a quadratic polynomial in  $\mathscr{P}_{1,1} \setminus \mathbb{P}_2$ , but it comes from a quadratic from three variables.

**Lemma 12.36.**  $x_1y_1 + \cdots x_dy_d \in \overline{\mathcal{P}}_{d,d}$ 

*Proof.* We will give the computations for d = 2, and describe the general construction. If we define

$$Q_{\epsilon} = \begin{pmatrix} \epsilon^{3} & \epsilon & 1 & \epsilon \\ \epsilon & \epsilon^{3} & \epsilon & 1 \\ 1 & \epsilon & \alpha & \alpha \\ \epsilon & 1 & \alpha & \alpha \end{pmatrix}$$

then  $(x_1, x_2, y_1, y_2)Q_{\varepsilon}(x_1, x_2, y_1, y_2)^t$  converges to  $x_1y_1 + x_2y_2$ . It suffices to show that  $u_{\varepsilon}A_{\varepsilon}u_{\varepsilon}$  has positive entries, for we just choose  $\alpha$  small enough so that  $u_{\varepsilon}A_{\varepsilon}u_{\varepsilon} - C_{\alpha}$  is positive.

$$\begin{split} \mathfrak{u}_{\varepsilon} A_{\varepsilon}^{-1} \mathfrak{u}_{\varepsilon} &= \begin{pmatrix} 1 & \varepsilon \\ \varepsilon & 1 \end{pmatrix} \begin{pmatrix} \varepsilon^{3} & \varepsilon \\ \varepsilon & \varepsilon^{3} \end{pmatrix}^{-1} \begin{pmatrix} 1 & \varepsilon \\ \varepsilon & 1 \end{pmatrix} \\ &= \frac{1}{\varepsilon (1+\varepsilon)(1+\varepsilon^{2})} \begin{pmatrix} 2+\varepsilon+\varepsilon^{2} & 1+\varepsilon+2\varepsilon^{2} \\ 1+\varepsilon+2\varepsilon^{2} & 2+\varepsilon+\varepsilon^{2} \end{pmatrix} \end{split}$$

In general, we define J(a, b) to be the d by d matrix whose diagonal is a, and remaining elements are b. We let  $A_{\varepsilon} = J(\varepsilon, \varepsilon^3)$ ,  $U_{\varepsilon} = J(1, \varepsilon)$  and  $C_{\varepsilon} = J(\alpha, \alpha)$ . With some work it can be verified that  $U_{\varepsilon}A_e^{-1}U_{\varepsilon}$  has all positive entries.

Note that this gives an element of  $\overline{\mathcal{P}}_{2,2}$  that is not in  $\overline{\mathbf{P}}_4$ . If it were, we could substitute  $x_2 = y_2 = 1$ , but  $x_1y_1 + 1$  is not in  $\overline{\mathbf{P}}_2$ .

*Remark* 12.37. We can construct polynomials of higher degree in  $\mathcal{P}_{1,1}$  by multiplying quadratics. For instance,  $\begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & \varepsilon \\ 1 & \varepsilon & c \end{pmatrix}$  satisfies the conditions of Proposition 12.32 for 0 < c < 1. If we choose  $0 < c_i < 1$  and let  $\varepsilon \longrightarrow 0^+$ , then the following polynomial is in  $\overline{\mathcal{P}}_{1,1}$ :

$$\prod_{i}(x^2+2xy+y^2+2x+c_i)$$

CHAPTER



### The analytic closure of P<sub>d</sub>

 $\mathbf{P}_d$  is the closure of  $\mathbf{P}_d$  under uniform convergence on compact domains. The uniform closure of  $\mathbf{P}_d^{\text{pos}}$  is  $\widehat{\mathbf{P}^{\text{pos}}}_d$ . Many results about these closures follow by taking limits of results about polynomials. For instance, we can conclude from Lemma 9.32 that if  $f(x, y) \in \widehat{\mathbf{P}}_2$  then  $f(x, x) \in \widehat{\mathbf{P}}_2$ . Similarly, we can also extend definitions from polynomials to these closures. As an example, we say that f,  $g \in \widehat{\mathbf{P}}_d$  interlace if  $f + \alpha g \in \widehat{\mathbf{P}}_d$  for all real  $\alpha$ .

#### 13.1 The analytic closure

The properties of  $\hat{\mathbf{P}}_d$  are similar to those of  $\hat{\mathbf{P}}$ , except that we do not have a characterization of the members as we do in one variable. To start off,

**Lemma 13.1.**  $\hat{P}_d$  consists of analytic functions whose domain is  $\mathbb{C}^d$ .

*Proof.* If  $f(\mathbf{x})$  is in  $\widehat{\mathbf{P}}_d$  then it is a limit of polynomials  $f_n(\mathbf{x})$  in  $\mathbf{P}_d$ . If we choose  $\mathbf{a} \in \mathbb{R}^d$  then  $f(\mathbf{x}^a_i)$  is the uniform limit of polynomials  $f_n(\mathbf{x}^a_i)$  in  $\mathbf{P}$  and hence  $f(\mathbf{x}^a_i)$  is analytic. Consequently, f is analytic in each variable separately, and so we can apply Hartog's Theorem [83] to conclude that  $f(\mathbf{x})$  is analytic.

From Section 11.5 we know that  $\hat{\mathbf{P}}_e \subset \hat{\mathbf{P}}_d$  if e < d, and  $\hat{\mathbf{P}^{\text{pos}}} \subset \hat{\mathbf{P}^{\text{pos}}}_2$ . The most important non-polynomial function in  $\hat{\mathbf{P}}$  is  $e^x$ . Thus  $e^x$  is in all  $\hat{\mathbf{P}}_d$ . There is another exponential function that is in  $\hat{\mathbf{P}}_d$ .

**Lemma 13.2.** If  $d \ge 2$  then  $e^{-xy}$  is in  $\widehat{P}_d$ . Also,  $e^{xy} \notin \widehat{P}_d$ .

*Proof.* From Corollary 11.7 we know that n - xy is in  $\overline{\mathbf{P}}_2$ . Writing  $e^{-xy}$  as a limit

$$e^{-xy} = \lim_{n \to \infty} n^{-n} (n - xy)^n$$
 (13.1.1)

it follows that all the factors of the product are in  $\overline{\mathbf{P}}_2$ , and hence  $e^{-xy}$  is in  $\widehat{\mathbf{P}}_2$ . If  $e^{xy} \in \widehat{\mathbf{P}}_2$  then substituting x = y yields  $e^{x^2} \in \widehat{\mathbf{P}}$  which is false. Since  $f(x, y) = e^{-xy}$  is in  $\widehat{\mathbf{P}^{\text{pos}}}_2$  we can apply Lemma 11.4 to find that  $f(x, x) = e^{-x^2}$  is in  $\widehat{\mathbf{P}}$ , as we saw earlier. Since  $\widehat{\mathbf{P}} \subset \widehat{\mathbf{P}^{\text{pos}}}_d$  we see  $e^{-x^2}$  is in  $\widehat{\mathbf{P}^{\text{pos}}}_d$ . We can combine these members of  $\widehat{\mathbf{P}^{\text{pos}}}_d$  to conclude

**Proposition 13.3.** If H is a d by d matrix with non-negative entries then  $e^{-xHx^{t}}$  is in  $\widehat{P^{pos}}_{d}$ .

*Proof.* It suffices to show that  $e^{-\sum a_{ij}x_ix_j}$  is in  $\widehat{\mathbf{P}^{pos}}_d$  for all non-negative  $a_{ij}$ . Since

$$e^{-\sum a_{ij}x_ix_j} = \prod e^{-a_{ij}x_ix_j}$$

and all the factors of the product are in  $\widehat{\mathbf{P}^{\text{pos}}}_d$  it follows that the product itself is in  $\widehat{\mathbf{P}^{\text{pos}}}_d$ , and so is the left hand side.

The non-negativity of the coefficients of H is important. For instance, if H = (-1) then  $e^{-xHx} = e^{x^2}$  and this is not in  $\widehat{\mathbf{P}}$  by (5.2.2).

*Remark* 13.4. We can easily construct functions in  $\widehat{\mathbf{P}}_2$ . If  $\sum |a_i| < \infty$  then the infinite product  $\prod_{i=1}^{\infty} (1 + a_i z)$  is an entire function. Suppose we choose sequences of positive real numbers  $\{a_i\}, \{b_i\}$  so that  $\sum a_i$  and  $\sum b_i$  are finite. The product

$$f(x,y) = \prod_{1}^{\infty} (1 + a_i x + b_i y)$$

is an entire function since

$$\sum |a_i x + b_i y| \leqslant |x| \sum |a_i| + |y| \sum |b_i| < \infty$$

Since f is the limit of the partial products which are in  $\mathbf{P}_2$ , f(x, y) is in  $\widehat{\mathbf{P}}_2$ .

**Example 13.5.** We will show that  $sin(x) + y cos(x) \in \hat{\mathbf{P}}_2$ , which is an example of a member of  $\hat{\mathbf{P}}_2$  that is not a product. We start with the two product approximations to sin and cos:

$$f_{n}(x) = x \prod_{k=1}^{n} \left( 1 - \frac{x^{2}}{k^{2}\pi^{2}} \right)$$
$$g_{n}(x) = \prod_{k=0}^{n} \left( 1 - \frac{4x^{2}}{(2k+1)^{2}\pi^{2}} \right)$$

and notice that  $f_{n+1} \leq g_n$ . Lemma 11.28 implies that  $f_{n+1}(x) + yg_n(x) \in \overline{\mathbf{P}}_2$ . Taking the limit, we find that  $\sin(x) + y\cos(x) \in \widehat{\mathbf{P}}_2$ .

The same limit argument used in Lemma 13.2 can be applied to Corollary 9.93: **Lemma 13.6.** If  $f \in \widehat{P^{pos}}$  and  $g \in \overline{P}_d$  then  $f(-\partial_x \partial_y)g \in \overline{P}_d$ .

This has a very important corollary:

**Corollary 13.7.** If  $f \in \overline{P}_d$  then  $e^{-\partial x_1 \partial x_2} f \in \overline{P}_d$ . More generally,  $e^{-\partial x \cdot \partial y} : \overline{P}_{2d} \longrightarrow \overline{P}_{2d}$ .

*Proof.* Since  $e^{-xy} \in \widehat{\mathbf{P}}_2$  we can apply limits to Lemma 13.6. The second part follows from

$$e^{-\partial \mathbf{x} \cdot \partial \mathbf{y}} = e^{-\partial \mathbf{x}_1 \partial \mathbf{y}_1} \cdots e^{-\partial \mathbf{x}_d \partial \mathbf{y}_d}$$

**Corollary 13.8.** The map  $x^k \mapsto L_k(xy)$  maps  $P^{pos} \longrightarrow \overline{P}_2$ .

Proof. From (7.10.3) the following diagram commutes



and consequently this commutes:

e



Lemma 13.6 has another important consequence:

**Theorem 13.9.** If  $f = \sum a_{ij} x^i y^j$  in  $\mathbf{P}_2^{\text{pos}}$  then the diagonal polynomials  $\text{diag}(f) = \sum a_{ii} x^i$  and  $\text{diag}_1(f) = \sum a_{ii} i! x^i$  are in  $\mathbf{P}^{\text{pos}}$ .

*Proof.* Since f is in  $\mathbf{P}_2^{\text{pos}}(n)$  the homogenization  $F = \sum a_{ij} x^i y^j z^{n-i-j}$  is in  $\mathbf{P}_3$ . The linear transformation  $g \mapsto e^{-\partial_x \partial_y} g$  maps  $\overline{\mathbf{P}}_2$  to itself, and by Theorem 10.21 it extends to a linear transformation mapping  $\overline{\mathbf{P}}_3$  to itself. We compute

$$\begin{aligned} {}^{-\partial_{x}\partial_{y}}\mathbf{F} &= e^{-\partial_{x}\partial_{y}}\sum_{\mathbf{i},\mathbf{j}}a_{\mathbf{i}\mathbf{j}}x^{\mathbf{i}}y^{\mathbf{j}}z^{\mathbf{n}-\mathbf{i}-\mathbf{j}} \\ &= \sum_{\mathbf{i},\mathbf{j},\mathbf{k}}a_{\mathbf{i}\mathbf{j}}\frac{(-1)^{\mathbf{k}}}{\mathbf{k}!}\left(\frac{\partial}{\partial x}\right)^{\mathbf{k}}\left(\frac{\partial}{\partial y}\right)^{\mathbf{k}}x^{\mathbf{i}}y^{\mathbf{j}}z^{\mathbf{n}-\mathbf{i}-\mathbf{j}} \\ &= \sum_{\mathbf{i},\mathbf{j},\mathbf{k}}a_{\mathbf{i}\mathbf{j}}\frac{(-1)^{\mathbf{k}}}{\mathbf{k}!}(\mathbf{i})_{\mathbf{k}}(\mathbf{j})_{\mathbf{k}}x^{\mathbf{i}-\mathbf{k}}y^{\mathbf{j}-\mathbf{k}}z^{\mathbf{n}-\mathbf{i}-\mathbf{j}} \end{aligned}$$
(13.1.2)

Substituting x = y = 0 gives a polynomial in **P** 

$$e^{-\partial_{x}\partial_{y}}\mathsf{F}\big|_{x=y=0} = \sum_{k} a_{kk}(-1)^{k}k! z^{n-2k}$$

The reverse of this last polynomial is in **P**, so  $\sum a_{kk}(-1)^k k! z^{2k}$  is in **P**. Now all  $a_{kk}$  are positive since  $f \in \mathbf{P}_2^{\text{pos}}$ , and so  $\sum a_{kk}(-1)^k k! x^k$  is in **P**. Replacing x by -x, and an application of the exponential map shows that  $\sum a_{ii} x^i$  is also in **P**.

A similar proof establishes a similar result in higher dimensions, but we are not able to eliminate the negative factors.

**Lemma 13.10.** If 
$$f(\mathbf{x}, \mathbf{y}, z) = \sum_{ij} f_{ij}(\mathbf{x}) y^i z^j$$
 is in  $\mathbf{P}_{d+2}(n)$  then  $\sum_{ij} (-1)^k \mathbf{k}! f_{kk}(\mathbf{x})$  and  $\sum_{ij} (-1)^k f_{kk}(\mathbf{x})$  are in  $\mathbf{P}_d$ .

*Proof.* We follow the proof above and apply the operator  $e^{-\partial_y \partial_z}$ , but we do not need to homogenize. For the second part, apply the first part to exp(f) where the exponential map operates on the y variable.

*Remark* 13.11. Here is an alternative proof that the Hadamard product maps  $\mathbf{P}^{\text{pos}} \times \mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}^{\text{pos}}$ . It does not show that the domain could be enlarged to  $\mathbf{P} \times \mathbf{P}^{\text{pos}}$ .

If  $f, g \in \mathbf{P}^{\text{pos}}$  then  $f(x)g(y) \in \overline{\mathbf{P}}_2^{\text{pos}}$ . Applying Theorem 13.9 shows that the diagonal of f(x)g(y) is in  $\mathbf{P}^{\text{pos}}$ . But the diagonal is Hadamard product: if  $f(x) = \sum a_i x^i, g(y) = \sum b_i y^i$  then the diagonal of  $\sum a_i b_j x^i y^j$  is  $\sum a_i b_i x^i$ .

**Corollary 13.12.** If  $\sum f_i(x)y^i$  and  $\sum g_i(x)y^i$  are both in  $P_{d+1}$  then

$$\sum (-1)^{i} f_{i}(\mathbf{x}) g_{i}(\mathbf{x}) \in \mathbf{P}_{d}$$

*Proof.* The product  $(\sum f_i(\mathbf{x})y^i) (\sum g_i(\mathbf{x})z^i)$  is in  $\overline{\mathbf{P}}_{d+2}$ . Now apply Lemma 13.10.

This is false without the factor  $(-1)^i$ . Just take f = g; the sum  $\sum f_I(\mathbf{x})g_I(\mathbf{x})$  is positive if the constant term of f is non-zero. We note some similar results:

**Corollary 13.13.** If  $f = \sum a_{ij} x^i y^j$  is in  $\mathbf{P}_2^{\text{pos}}$  then for any non-negative integers r, s the polynomial

$$\sum_{i} a_{i+r,i+s} \frac{(i+r)!(j+s)!}{r!s!i!} x^{i} \in P$$

*Proof.* Instead of substituting x = y = 0 in (13.1.2) take the coefficient of  $x^r y^s$  and apply a similar argument.

Corollary 13.14. If  $f=\sum \alpha_{ij}x^iy^j$  is in  $P_2^{\text{pos}}(n)$  then define

$$s_k(x) = \sum_{i-j=k} a_{i,j} x^{j}$$

For -n < k < n we have that  $s_k \in P$ , and  $s_k, s_{k-1}$  have a common interlacing. The map  $x^k \mapsto s_k$  sends  $P^{p \circ s}(n)$  to itself.

*Proof.* For any positive  $\alpha$  the polynomial  $g = (x^k + \alpha x^{k-1})f$  is in the closure of  $\mathbf{P}_2^{\text{pos}}$ , and diag  $g = s_k + \alpha s_{k-1}$ . By Proposition 1.35  $s_k$  and  $s_{k-1}$  have a common interlacing. If  $h = \sum b_k y^k$  is in  $\mathbf{P}^{\text{pos}}(n)$  then h(y) is in the closure of  $\mathbf{P}_2^{\text{pos}}(n)$ , and the following calculations show that T(h) is in  $\mathbf{P}$ .

$$diagh(y)f(x,y) = diag\sum_{i,j,k} b_k a_{ij} x^i y^{j+k}$$
$$= \sum_k b_k diag\sum_{i,j} a_{ij} x^i y^{j+k}$$
$$= \sum_k b_k \sum_{i=j+k} a_{ij} x^i$$
$$= \sum b_k s_k = T(h)$$

**Corollary 13.15.** *For any* n *these polynomials have all real roots:* 

$$\sum_{i} \binom{n}{i,i} x^{i} \qquad \sum_{i} \frac{x^{i}}{i!(n-2i)!}$$

*Proof.* The two polynomials in question are the diagonal polynomials of a polynomial in  $\mathbf{P}_2$ , namely  $(x + y + 1)^n$ .

*Remark* 13.16. Here's a variation on the even part (§ 6.7.15). If  $f = \sum a_i x^i$  is in  $\mathbf{P}^{alt}$  then f(x+y) is in  $\mathbf{P}_2^{pos}$ . Now  $diag(x+y)^k$  is either 0 if k is odd, or  $\binom{2m}{m}x^m$  if k = 2m is even. Consequently,  $diag_1 f(x+y) = \sum \frac{(2i)!}{i!} a_{2i}x^i$  is in  $\mathbf{P}$ .

In the remark we considered f(x+y). If we consider f((x+1)(y+1)) then we get a quite different transformation:

**Corollary 13.17.** The linear transformation  $x^r \mapsto \sum_i {\binom{r}{i}}^2 x^i$  maps  $P^{alt}$  to itself.

*Proof.* If  $f \in \mathbf{P}^{alt}$  then  $f(xy) \in \overline{\mathbf{P}}_2$ , and so after substituting we have that  $f((x+1)(y+1)) \in \mathbf{P}_2$ . If we write  $f = \sum a_i x^i$  then

$$\begin{split} f(\,(x+1)(y+1)\,) &= \sum_{i} \, \alpha_{i}(x+1)^{i}(y+1)^{i} \\ diag \; f(\,(x+1)(y+1)\,) &= \sum_{i} \, \alpha_{i} \sum_{j} \, \binom{i}{j}^{2} \, x^{j} \end{split}$$

which establishes the result.

In §1.1.16 we saw that the partial sums of the function  $e^x$  in  $\widehat{\mathbf{P}}$  do not have all real roots, but adding another factorial lands in  $\mathbf{P}$ .

**Lemma 13.18.** If  $f = \sum a_i \frac{x^i}{i!}$  is in  $\widehat{P}$  then for any n the polynomial  $\sum_{i=0}^n \binom{n}{i} a_i x^i = n! \sum_{i=0}^n \frac{a_i}{i!(n-i)!} x^i$  has all real roots.

*Proof.* The operator  $f(\partial_x)$  maps  $P_2$  to itself, so consider

$$\begin{split} f(\partial_x)(x+y)^n \Big|_{x=0} &= \sum_{i,j} \alpha_i y^{n-j} \binom{n}{j} \mathsf{D}^i x^j \Big|_{x=0} \\ &= \sum_k \alpha_k y^{n-k} \binom{n}{k} \end{split}$$

Taking the reverse establishes the lemma.

We can take limits of the Hadamard product to conclude from Theorem 9.87

**Corollary 13.19.** If  $f \in \widehat{P^{pos}}$  and  $g \in \widehat{P}$  then  $EXP^{-1}f * g$  is in  $\widehat{P}$ .

Here's a representation of  $\widehat{\mathbf{P}}$  as the image of one exponential.

Lemma 13.20.

$$\widehat{\boldsymbol{P}} = \{\mathsf{T}(\boldsymbol{e}^{\mathsf{x}}) \mid \mathsf{T}: \boldsymbol{P}^{\mathsf{pos}} \longrightarrow \boldsymbol{P}\}$$
(13.1.3)

$$\widehat{P^{\text{pos}}} = \{\mathsf{T}(e^{x}) \mid \mathsf{T}: P^{\text{pos}} \longrightarrow P^{\text{pos}}\}$$
(13.1.4)

*Proof.* Since  $T: \mathbf{P}^{pos} \longrightarrow \mathbf{P}$  extends to  $T: \widehat{\mathbf{P}^{pos}} \longrightarrow \widehat{\mathbf{P}}$ , we know that  $T(e^x) \in \widehat{\mathbf{P}}$ . Conversely, choose  $f \in \widehat{\mathbf{P}}$  and define

$$\mathsf{T}_{\mathsf{f}}(\mathsf{g}) = \mathsf{f} \ast' \mathsf{g} = \mathsf{f} \ast \mathsf{EXP}^{-1}(\mathsf{g})$$

We know that  $T_f : \widehat{\mathbf{P}^{pos}} \longrightarrow \widehat{\mathbf{P}}$ . Since  $T_f(e^x) = f$ , we have equality. The second case is similar.

Unlike **P**, factors of elements in  $\widehat{\mathbf{P}}_2$  aren't necessarily in  $\widehat{\mathbf{P}}_2$ . A simple example is  $\mathbf{x} = (e^{xy})(xe^{-xy})$ . Note that  $xe^{-xy} \in \widehat{\mathbf{P}}_2$ , yet  $e^{xy} \notin \widehat{\mathbf{P}}_2$ . However, we do have

**Lemma 13.21.** If  $f(x, y)e^{-xy} \in \widehat{P}_2$ , then  $f(x, \alpha)$  and  $f(\alpha, x)$  are in  $\widehat{P}$  for any choice of  $\alpha \in \mathbb{R}$ .

*Proof.* If  $f(x, y)e^{-xy} \in \widehat{\mathbf{P}}_2$ , then  $f(x, \alpha)e^{-\alpha x} \in \widehat{\mathbf{P}}$ . Since  $e^{\alpha x} \in \widehat{\mathbf{P}}$  it follows that  $f(x, \alpha) \in \widehat{\mathbf{P}}$ .

452

#### 13.2 Differential operators

If we substitute derivatives for some variables, the result maps  $\overline{\mathbf{P}}_d$  to itself. This generalization of the corresponding property for  $f(\mathsf{D})$  acting on  $\mathbf{P}$  has many important uses.

Lemma 13.22. If  $f(\mathbf{x}) \in \overline{\mathbf{P}}_d$  then  $f(-\partial_{\mathbf{x}}) \colon \overline{\mathbf{P}}_d \longrightarrow \overline{\mathbf{P}}_d$ .

*Proof.* Since  $e^{-\partial_x \partial_y}$  maps  $\overline{\mathbf{P}}_{2d}$  to itself the lemma follows from the identity

$$\left. e^{-\partial_{\mathbf{x}}\partial_{\mathbf{y}}} g(\mathbf{x}) f(\mathbf{y}) \right|_{\mathbf{y}=0} = f(-\partial_{\mathbf{x}})g(\mathbf{x}).$$

By linearity we only need to check it for monomials:

$$e^{-\partial_x\partial_y}x^Iy^J\bigg|_{y=0} = \left.\frac{(-\partial_x)^J}{J!}x^I \cdot (\partial_y)^Jy^J\bigg|_{y=0} = (-\partial_x)^Jx^I$$

We use the curious equality below to derive a corollary due to Sokal and Lieb [115].

**Lemma 13.23.** If f(x, y) is any polynomial then

$$f(x, y - \partial_x) = e^{-\partial x \partial y} f(x, y)$$

*Proof.* By linearity we may assume that  $f(x, y) = x^{s}y^{r}$ . The left hand side is

$$(\mathbf{y} - \mathbf{d}_{\mathbf{x}})^{r} \mathbf{x}^{s} = \sum_{i=0}^{r} \mathbf{y}^{r-i} {\binom{r}{i}} (-1)^{i} \mathbf{d}_{\mathbf{x}}^{i} \mathbf{x}^{s}$$
$$= \sum_{i=0}^{r} \mathbf{y}^{r-i} \mathbf{x}^{s-i} \frac{(-1)^{i}}{i!} \frac{r!}{(r-i)!} \frac{s!}{(s-i)!}$$
$$= \sum_{i=0}^{\infty} \frac{(-\mathbf{d}_{\mathbf{x}} \mathbf{d}_{\mathbf{y}})^{i}}{i!} \mathbf{x}^{s} \mathbf{y}^{r}$$

which is exactly the right hand side.

**Corollary 13.24.** *If*  $f(\mathbf{x}, \mathbf{y}) \in P_{2d}$  *then* 

- 1.  $f(\mathbf{x}, \mathbf{y} \partial_{\mathbf{x}}) \in \overline{\mathbf{P}}_{2d}$ .
- 2.  $f(\mathbf{x}, -\partial_{\mathbf{x}}) \in \overline{\mathbf{P}}_d$ .

Proof. If we iterate Lemma 13.23 we get

$$f(x_1, y_1 - \partial x_1, \dots, x_d, y_d - \partial x_d) = e^{-\partial x_1 \partial y_2} \cdots e^{-\partial x_d \partial y_d} f(\mathbf{x}, \mathbf{y})$$

The first part now follows from Corollary 13.7. If we set  $\mathbf{y} = 0$  we get the second part.

_	

If we substitute  $\partial x_i$  for  $x_i$  then the result is much easier to prove.

**Lemma 13.25.** If  $f = \sum f(x_1, ..., x_{d-1}) x_d^k$  is in  $P_d$  and  $f(0) \neq 0$  then the linear transformation  $g \mapsto \sum f_k(x_1, ..., x_{d-1}) \frac{\partial^k}{\partial x_d^k}(g)$  maps  $P_d$  to itself.

*Proof.* The homogeneous part of T(g) is  $f(0)g^H$ , so T satisfies the homogeneity condition. If we choose  $a_1, \ldots, a_{d-1} \in \mathbb{R}$  then

$$\mathsf{T}(\mathfrak{g})(\mathfrak{a}_1,\ldots,\mathfrak{a}_{d-1},\mathfrak{x}_d)=\mathsf{f}(\mathfrak{a}_1,\ldots,\mathfrak{a}_{d-1},\mathfrak{d}_{\mathfrak{x}_d})\mathfrak{g}(\mathfrak{a}_1,\ldots,\mathfrak{a}_{d-1},\mathfrak{x}_d)$$

and this is in **P** since  $f(a_1, \ldots, a_{d-1}, x)$  and  $g(a_1, \ldots, a_{d-1}, x)$  are in **P**.

Next is a Hadamard product result.

**Lemma 13.26.** If  $\sum f_I(\mathbf{x})\mathbf{y}^I$  and  $\sum g_I(\mathbf{x})\mathbf{y}^I$  are in  $\overline{\mathbf{P}}_{2d}$  then

$$\sum_{I} (-1)^{I} I! f_{I}(\mathbf{x}) g_{I}(\mathbf{x}) \in \overline{\mathbf{P}}_{d}$$

*Proof.* A calculation shows:

$$\left(\sum f_{I}(\mathbf{x})(-\partial \mathbf{y})^{I}\right)\left(\sum g_{I}(\mathbf{x})\mathbf{y}^{I}\right)\Big|_{\mathbf{y}=\mathbf{0}} = \sum_{I} (-1)^{I} \mathbf{I}! f_{I}(\mathbf{x}) g_{I}(\mathbf{x})$$

**Lemma 13.27.** If  $Q = (q_{ij})$  is a symmetric d by d matrix, then  $e^{-xQx^t} \in \widehat{P}_d$  if and only if all elements of Q are non-negtive.

*Proof.* If all elements are non-negative then it is easy to see that  $e^{-xQx^t} \in \widehat{\mathbf{P}}_d$ . Since  $\alpha - xy \in \overline{\mathbf{P}}_2$  for  $\alpha > 0$ , compute

$$e^{-(\alpha x^2+2bxy+cy^2)}(\partial_x,\partial_y)(\alpha-xy) = \alpha - xy + 2b$$

Since this is in  $\overline{\mathbf{P}}_2$ , it follows that  $\alpha + 2b \ge 0$  and so b is non-negative.

#### 13.3 Limits and transformations

If f and g are entire functions and T is a linear transformation then we define f(T)g to be the *formal* series

$$\sum_{i=0}^{\infty} \sum_{j=0}^{\infty} a_i b_j T^i(x^j)$$
 (13.3.1)

where  $f = \sum_{0}^{\infty} a_i x^i$  and  $g = \sum_{0}^{\infty} b_i x^i$ . This is only a formal sum since there is no apriori reason why f(T)g should exist. For a full discussion of the complexities when  $T = aD + xD^2$  see [106].

We are interested in the case where  $f, g \in \widehat{\mathbf{P}}$  and  $T: \mathbf{P} \longrightarrow \mathbf{P}$ . We assume that  $f_n \rightarrow f$  and  $g_n \rightarrow g$  where convergence is uniform convergence on compact subsets. The problem is then to determine when

 $\lim_{n \to \infty} f_n(T)g_n \text{ exists and equals } f(T)g.$ 

The simplest case is when all limits involve polynomials.

#### Lemma 13.28. If

- 1. T decreases degree.
- 2.  $h(T): \mathbf{P} \longrightarrow \mathbf{P}$  for all  $h \in \mathbf{P}$ .
- 3. f  $\in \widehat{P}$
- *then*  $f(T): P \longrightarrow P$ .

*Proof.* If  $f_n \to g$  and  $g \in \mathbf{P}(m)$  then both  $f_n(T)g$  and f(T)g are polynomials since  $T^k(g) = 0$  for  $k \ge m$ . Since the coefficients of  $f_n$  converge to the coefficients of f we see that  $f_n(T)g \to f(T)g$ . By hypothesis all  $f_n(T)g \in \mathbf{P}$ , so  $f(T)g \in \mathbf{P}$ .

**Example 13.29.** For example, since f(D) maps **P** to itself it follows that  $e^{D}$ : **P**  $\longrightarrow$  **P**. Of course, this is trivial since  $e^{D}f(x) = f(x + 1)$ , but the same argument applies to  $e^{-\partial_{x}\partial_{y}}$  (see Corollary 13.7).

In order to work with entire functions we recall Montel's theorem:

Any locally bounded sequence of holomorphic functions  $f_n$  defined on an open set D has a subsequence which converges uniformly on compact subsets to a holomorphic function f.

It follows that if T is an operator that takes bounded sequences to bounded sequences and if  $f_n \longrightarrow f$  then  $Tf_n \longrightarrow Tf$ . For instance, the k'th derivative is a bounded operator by Cauchy's integral formula:

$$f^{(k)}(\mathfrak{a}) = \frac{k!}{2\pi\mathfrak{i}} \oint_{\mathcal{C}} \frac{f(z)}{(z-\mathfrak{a})^{k+1}} \, \mathrm{d}z$$

where C is a small circle centered at a. More generally, if  $f(x, y) = \sum f_i(x)y^i$  is a polynomial in two variables then the operator

$$T \colon \mathfrak{g} \mapsto \sum f_{\mathfrak{i}}(x) D^{\mathfrak{i}}\mathfrak{g}$$

is a bounded operator since it is a sum of products of locally bounded functions  $f_i$  times bounded operators. We therefore have

**Lemma 13.30.** If T is a differential operator as above then  $\lim_{n\to\infty} T(1-\frac{xy}{n})^n$  exists, equals  $T(e^{-xy})$ , and is an entire function.
In order to prove that sequences are uniformly bounded we use the bounds on the size of a polynomial.

Lemma 13.31. Suppose that T is a linear transformation and

- 1.  $T_* : \overline{\mathbf{P}}_2 \longrightarrow \overline{\mathbf{P}}_2$
- 2.  $T(1) \in \mathbb{R} \setminus 0$  and  $f(0) \neq 0$ .
- 3.  $f_n \to f$  in  $\widehat{\mathbf{P}}_2$  where  $f_n \in \overline{\mathbf{P}}_2$ .

then  $T_*(f_n) \longrightarrow T_*(f)$ .

*Proof.* Write  $f_n = \sum_i F_{n,i}(x)y^i$  and  $f = \sum_i F_i(x)y^i$ . Now  $Tf_n = \sum T(F_{n,i})y^i$  is in  $\overline{\mathbf{P}}_2$  by hypothesis, so for any  $\alpha \in \mathbb{R}$  we have  $(Tf_n)(\alpha, y) \in \mathbf{P}$  and therefore we can apply Lemma 4.16:

$$\sup_{|y|\leqslant r} (\mathsf{T}\,\mathsf{f}_n)(\alpha,y) \leqslant \mathsf{T}(\mathsf{F}_{n,0}) \exp\left(r\frac{\mathsf{T}\,\mathsf{F}_{n,1}(\alpha)}{\mathsf{T}(\mathsf{F}_{n,0})} + 2r^2\frac{(\mathsf{T}\,\mathsf{F}_{n,1}(\alpha))^2}{(\mathsf{T}\mathsf{F}_{n,0})^2} + r^2\frac{\mathsf{T}\,\mathsf{F}_{n,2}(\alpha)}{\mathsf{T}(\mathsf{F}_{n,0})}\right)$$

If we define

$$a = \inf_{n} T(F_{n,0}) \qquad A = \sup_{n} T(F_{n,0})$$
$$B_{r} = \sup_{n,|\alpha| \leq r} T(F_{n,1}(\alpha)) \qquad C_{r} = \sup_{n,|\alpha| \leq r} T(F_{n,2}(\alpha))$$

then A,  $B_r$ ,  $C_r$  are finite and a is non-zero by hypothesis so

$$\sup_{|\alpha| \leq r, |y| \leq r} (\mathsf{T} f_n)(\alpha, y) \leq A \exp\left(r\frac{B}{a} + 2r^2\frac{B^2}{a^2} + r^2\frac{C}{a}\right)$$

Thus  $T(f_n)$  is uniformly bounded and therefore converges to T(f).

#### **13.4** Totally positive functions

We raise some questions about totally positive functions and  $\mathbf{P}_2^{\text{pos}}$ .

**Definition 13.32.** A function f(x, y) is *strictly totally positive* if for every positive integer n and sequences  $x_1 < x_2 < \cdots < x_n$  and  $y_1 < y_2 < \cdots < y_n$  the determinant

$$\begin{vmatrix} f(x_1, y_1) & f(x_1, y_2) & \dots & f(x_1, y_n) \\ f(x_2, y_1) & & \vdots \\ \vdots \\ f(x_n, y_1) & \dots & f(x_n, y_n) \end{vmatrix}$$
(13.4.1)

is positive. If the sequences  $\{x_i\}$  and  $\{y_i\}$  are restricted to positive values then we say that f(x, y) is a strictly totally positive function on  $\mathbb{R}^2_+$ .

If we take n = 1 we see that a necessary condition for f(x, y) to be strictly totally positive on  $\mathbb{R}^2_+$  is that f(x, y) is positive for x and y positive. A natural question (106) is to determine conditions on polynomials g(x, y) so that 1/g is strictly totally positive on  $\mathbb{R}^2_+$ .

We know three functions for which this holds. It is well known that  $e^{xy}$  is strictly totally positive, and  $e^{xy}$  is of the form 1/g where  $g = e^{-xy} \in \widehat{\mathbf{P}}_2$ . The simplest function in  $\mathbf{P}_2^{\text{pos}}$  is x + y; this is strictly totally positive on  $\mathbb{R}_+^2$ . That this is so follows from a famous evaluation. Recall Cauchy's double alternant:

$$\det_{1 \leqslant i,j \leqslant n} \left( \frac{1}{x_i + y_j} \right) = \frac{\prod_{1 \leqslant i < j \leqslant n} (x_i - x_j)(y_i - y_j)}{\prod_{1 \leqslant i,j \leqslant n} (x_i + y_j)}$$

The numerator is positive for any pair of strictly increasing sequences; the denominator is positive since all  $x_i$  and  $y_i$  are positive.

For the last example, there is a quadratic analog (13.4.2) of Cauchy's formula due to Borchardt[159] where perm is the permanent. Since the permanent of a matrix with all positive entries is positive, it follows that  $(x + y)^{-2}$  is strictly totally positive on  $\mathbb{R}^2_+$ .

$$\det_{1 \leqslant i, j \leqslant n} \left( \frac{1}{(x_i + y_j)^2} \right) = \det_{1 \leqslant i, j \leqslant n} \left( \frac{1}{x_i + y_j} \right) \operatorname{perm}_{1 \leqslant i, j \leqslant n} \left( \frac{1}{x_i + y_j} \right)$$
(13.4.2)

If h(x) is any function that is positive for positive x, and f(x, y) is totally positive on  $\mathbb{R}^2_+$  then h(x)f(x, y) is strictly totally positive on  $\mathbb{R}^2_+$ . This follows from simple properties of the determinant:

$$\det_{1\leqslant i,j\leqslant n} \left( h(x_i)f(x_i,y_j) \right) = h(x_1)\cdots h(x_n) \ \det_{1\leqslant i,j\leqslant n} \left( f(x_i,y_j) > 0 \right)$$

Consequently, if  $g \in \widehat{\mathbf{P}}_2$  satisfies the property that 1/g is strictly totally positive on  $\mathbb{R}^2_+$  then so does  $e^{-x^2}g$ .

It is not true that 1/g is positive semi-definite if  $g \in \mathbf{P}_2^{\text{pos}}$  - see [12]. We need a few properties of positive definite matrices [95].

**Definition 13.33.** Let A, B be Hermitian matrices. We write  $B \prec A$  if the matrix A - B is positive definite. If A - B is positive semidefinite we write  $B \preceq A$ . The relation  $\preceq$  makes the set of all Hermitian matrices into a partially ordered

Here are some properties of  $\prec$  that we will use.

- 1.  $A \prec B$  implies det(A) < det(B).
- 2.  $A \prec B$  implies  $B^{-1} \prec A^{-1}$ .

set.

3. A  $\prec$  B and C positive definite implies A + C  $\prec$  B + C.

4. A  $\prec$  B and C positive definite implies CAC  $\prec$  CBC.

If f(x) is a positive definite matrix for all x in an interval on the real line, then we say that f is *increasing* if a < b implies  $f(a) \prec f(b)$ . f is *decreasing* if -f is increasing.

**Lemma 13.34.** If A and B are positive definite matrices, and  $0 < x_1 < x_2$ ,  $0 < y_1 < y_2$  then

$$\begin{vmatrix} \frac{1}{|I+x_1A+y_1B|} & \frac{1}{|I+x_2A+y_1B|} \\ \frac{1}{|I+x_1A+y_2B|} & \frac{1}{|I+x_2A+y_2B|} \end{vmatrix} > 0$$

*Proof.* If we let  $C = x_2A$ ,  $D = y_2B$ ,  $x = x_1/x_2$ ,  $y = y_1/y_2$  then 0 < x, y < 1 and C, D are positive definite. We need to show that

$$\begin{vmatrix} \frac{1}{|I + xC + yD|} & \frac{1}{|I + C + yD|} \\ \frac{1}{|I + xC + D|} & \frac{1}{|I + C + D|} \end{vmatrix} > 0$$

Since all the matrices are positive definite the determinants of positive linear combinations are positive. Thus we need to show that  $^{\rm 1}$ 

$$|I + xC + D||I + C + yD| > |I + xC + yD||I + C + D|$$

which is equivalent to

$$\frac{|I + xC + yD|}{|I + xC + D|} < \frac{|I + C + yD|}{|I + C + D|}$$

If we denote the left hand side by F(x) then we will show that F(x) is an increasing function.

$$\begin{split} I+xC+D \text{ is increasing}\\ (I+xC+D)^{-1} \text{ is decreasing}\\ D^{1/2}(I+xC+D)^{-1}D^{1/2} \text{ is decreasing}\\ (y-1)D^{1/2}(I+xC+D)^{-1}D^{1/2} \text{ is increasing}\\ I+(y-1)D^{1/2}(I+xC+D)^{-1}D^{1/2} \text{ is increasing} \end{split}$$

The determinant of the last expression is F(x) - this follows from the identity

$$|(\alpha A + B)B^{-1}| = |I + \alpha A^{1/2}B^{-1}A^{1/2}|$$

<sup>&</sup>lt;sup>1</sup>Thanks to Han Engler for this argument.

# 13.5 Hermite polynomials

We use generating functions to define Hermite polynomials in d variables. The main result is that they are in  $P_d$ .

Let S be a  $n \times n$  Hermitian matrix,  $\mathbf{y} = (y_1, \dots, y_n)$  and  $\mathbf{x} = (x_1, \dots, x_n)$ . The Hermite polynomials determined by S are defined to be the coefficients of the exponential generating function

$$f_{S}(\mathbf{x}, \mathbf{y}) = \exp\left(-\mathbf{y}S\mathbf{y}^{*} - 2\mathbf{y}S\mathbf{x}^{*}\right) = \sum_{\mathbf{I}} H_{\mathbf{I}}(\mathbf{x})\frac{(-\mathbf{y})^{\mathbf{I}}}{\mathbf{I}!}$$
(13.5.1)

where  $I = (i_1, ..., i_n)$ ,  $I! = i_1! \cdots i_n!$ ,  $y^I = y_1^{i_1} \cdots y_n^{i_n}$ , and  $z^*$  is the conjugate transpose of z. The factor  $2ySx^*$  has a negative sign, so we can use Proposition 13.3 to conclude that  $f_S \in \hat{P}_{2d}$ .

If we take n = 1 and S = (1) then the exponential is  $e^{-y_1^2 - 2x_1y_1}$  which is exactly the generating function of the one variable Hermite polynomials.

If we multiply (13.5.1) by  $exp(-xSx^{t})$  then

$$\exp(-xSx^{t}) \sum H_{I} \frac{(-y)^{I}}{I!} = \exp(-(y+x)S(y+x)^{t}).$$

We use this to derive the Rodrigues' formula for the Hermite polynomials. The Taylor series of  $g(\mathbf{x})$  asserts

$$g(\mathbf{x} + \mathbf{y}) = \sum_{\mathbf{I}} \left(\frac{\partial}{\partial \mathbf{x}}\right)^{\mathbf{I}} g(\mathbf{x}) \frac{\mathbf{y}^{\mathbf{I}}}{\mathbf{I}!}$$

so if we choose  $g(\mathbf{x}) = \exp(-\mathbf{x}S\mathbf{x}^t)$  then

$$\begin{split} \exp(-(\mathbf{y}+\mathbf{x})S(\mathbf{y}+\mathbf{x})^{\mathrm{t}}) &= \sum \left(\frac{\partial}{\partial \mathbf{x}}\right)^{\mathrm{I}} e^{-\mathbf{x}S\mathbf{x}^{\mathrm{t}}} \frac{\mathbf{y}^{\mathrm{I}}}{\mathrm{I}!} \\ &= \sum e^{-\mathbf{x}S\mathbf{x}^{\mathrm{t}}} H_{\mathrm{I}}(\mathbf{x}) \frac{(-\mathbf{y})^{\mathrm{I}}}{\mathrm{I}!}. \end{split}$$

Equating coefficients of y yields the Rodrigues' formula

$$\mathsf{H}_{I}(x) = (-1)^{|I|} \, e^{xSx^{t}} \, \left(\frac{\partial}{\partial x}\right)^{I} \, e^{-xSx^{t}}.$$

If we rewrite this as

$$e^{-xSx^{t}}\;\mathsf{H}_{I}(x)=(-1)^{|I|}\left(\frac{\partial}{\partial x}\right)^{I}\;e^{-xSx^{t}}$$

then it follows that  $H_I(\mathbf{x})$  satisfies substitution.

In order to show that  $H_I$  is in  $P_n$  it suffices to determine the homogeneous part of  $H_I$ , and we use the Rodrigues formula for that. We first note that if  $e_i$  is the coordinate vector with a 1 in the i-th place, and 0 elsewhere then

$$\frac{\partial}{\partial x_i} \mathbf{x} \mathbf{S} \mathbf{x}^{\mathsf{t}} = (e_i \mathbf{S} + \mathbf{S} e_i^{\mathsf{t}}) \mathbf{x}.$$

Consequently, if we differentiate  $g(\mathbf{x}) \exp(-\mathbf{x} S \mathbf{x}^t)$  where g is some polynomial then

$$\frac{\partial}{\partial x_i} g(\mathbf{x}) \exp(-\mathbf{x} S \mathbf{x}^t) = k(\mathbf{x}) \exp(-\mathbf{x} S \mathbf{x}^t)$$

where  $k(\mathbf{x})$  is a polynomial, and moreover

$$\mathbf{k}(\mathbf{x})^{\mathsf{H}} = ((\mathbf{e}_{\mathsf{i}}\mathsf{S} + \mathsf{S}\mathbf{e}_{\mathsf{i}}^{\mathsf{t}})\mathbf{x})\mathsf{g}(\mathbf{x})^{\mathsf{H}}.$$

Upon writing

$$\left(\frac{\partial}{\partial x_{i}}\right)^{1} \exp(-\mathbf{x}S\mathbf{x}^{t}) = \mathbf{j}(\mathbf{x}) \exp(-\mathbf{x}S\mathbf{x}^{t})$$

we conclude that the homogeneous part of j is

$$\mathfrak{j}(\mathbf{x})^{\mathsf{H}} = \prod \left( \left( e_{\mathfrak{i}} S + S e_{\mathfrak{i}}^{\mathsf{t}} \right) \mathbf{x} \right).$$

By assumption all coefficients of S are positive, so all the coefficients of  $j^H$  are positive, and by the above it follows that the homogeneous part of  $H_I$  has all positive coefficients. This proves

**Theorem 13.35.** If S is a symmetric matrix with all positive coefficients then the Hermite polynomials determined by (13.5.1) are in  $P_d$ .

#### 13.6 Hermite polynomials in two variables

The results of this section exist simply to explain properties that can be observed in a particular graph of a Hermite polynomial in two variables. If we take  $Q = \begin{pmatrix} 1 & 3 \\ 3 & 2 \end{pmatrix}$  and I = (2, 3) then

$$\begin{split} \mathsf{H}_{2,3} = & -16 \left( 108 \, \text{x} - 207 \, \text{x}^3 + 54 \, \text{x}^5 + 114 \, \text{y} - 876 \, \text{x}^2 \, \text{y} + 432 \, \text{x}^4 \, \text{y} - 990 \, \text{x} \, \text{y}^2 \right. \\ & + & 1206 \, \text{x}^3 \, \text{y}^2 - 332 \, \text{y}^3 + 1420 \, \text{x}^2 \, \text{y}^3 + 744 \, \text{x} \, \text{y}^4 + 144 \, \text{y}^5 ) \end{split}$$

The homogeneous part factors into  $-32(3 + x)^2(2 + 3x)^3$ , which explains the asymptotes in Figure 13.1. The rest of the graph of H<sub>2,3</sub> is quite surprising:

The following result shows that all the  $H_{i,j}$  are in  $\mathbf{P}_2^{gen}$ , but doesn't explain the striking regularity.



Figure 13.1: The graph of  $H_{2,3}$ 

**Lemma 13.36.** If S is not a diagonal matrix then  $H_{i,j} \in \mathbf{P}_2^{gen}$ .

*Proof.* Let  $S = \begin{pmatrix} a & b \\ b & c \end{pmatrix}$  where  $b \neq 0$ , and set  $Q = ax^2 + 2bxy + cy^2$ . Suppose that  $H_{i,j}$  has intersecting solution curves. If so, then  $H_{i,j}$  and  $\frac{\partial}{\partial x}H_{i,j}$  have a common zero. Now since

$$\begin{split} \frac{\partial}{\partial x} \mathbf{H}_{ij} &= \frac{\partial}{\partial x} \left( e^{\mathbf{Q}} \, \partial_x^i \, \partial_y^j \, e^{-\mathbf{Q}} \right) \\ &= e^{\mathbf{Q}} \, \partial_x^{i+1} \, \partial_y^j \, e^{-\mathbf{Q}} + \mathbf{Q}_x e^{\mathbf{Q}} \, \partial_x^i \, \partial_y^j \, e^{\mathbf{Q}} \\ &= \mathbf{H}_{i+1,j} + \mathbf{Q}_x \mathbf{H}_{ij} \end{split}$$

it follows that  $H_{i,j}$  and  $H_{i+1,j}$  have a common zero. Since  $e^Q$  has no zeros, both  $\partial_x^i \partial_y^j e^Q$  and  $\partial_x^{i+1} \partial_y^j e^Q$  have a common root. Since  $\partial_x^i \partial_y^j e^Q \leq \partial_x^{i+1} \partial_y^j e^Q$ , we conclude from Corollary 9.42 that  $\partial_x^{i-1} \partial_y^j e^Q$  and  $\partial_x^i \partial_y^j e^Q$  have a common root. Continuing, we see  $\partial_y^j e^Q$  and  $\partial_x \partial_y^j e^Q$  have a common root. Eliminating derivatives of y, we conclude that  $\partial_y e^Q$  and  $\partial_x \partial_y e^Q$  have a common root. Because  $H_{1,0} = -2bx - 2cy$ , we can solve for x and substitute into

$$H_{1,1} = 2 \left( -b + 2 a b x^{2} + 2 b^{2} x y + 2 a c x y + 2 b c y^{2} \right)$$

which yields -2b. Since  $b \neq 0$ , there are no common roots.

We follow Nunemacher in finding the exact equations of the asymptotes. Let

$$Q = ax^{2} + 2bxy + cy^{2}$$

$$H_{n,m}(x, y) = -e^{ax^{2} + 2bxy + cy^{2}} \left(\frac{\partial}{\partial x}\right)^{n} \left(\frac{\partial}{\partial y}\right)^{m} e^{-(ax^{2} + 2bxy + cy^{2})}$$

$$H_{n,m+1} = -2(bx + cy)H_{n,m} + \partial_{y}H_{n,m}$$

$$H_{n+1,m} = -2(ax + by)H_{n,m} + \partial_{x}H_{n,m}$$

From the recursions it easily follows that the homogeneous part of  $H_{n,m}$  is  $(-2)^{n+m}(ax + by)^n(bx + cy)^m$ . Consequently, there are n asymptotes in one direction, and m in another. We proceed to find the precise equations of these asymptotes. It's easy to see that we can write

$$\begin{split} H_{n,m} &= (ax + by)^n Q_{n,m}(x,y) + (ax + by)^{n-2} Q_{n-2,m}(x,y) + \cdots \\ H_{n,m+1} &= (ax + by)^n Q_{n,m+1}(x,y) + (ax + by)^{n-2} Q_{n-2,m+1}(x,y) + \cdots \\ &= -2(bx + cy) \left[ (ax + by)^n Q_{n,m}(x,y) + (ax + by)^{n-2} Q_{n-2,m}(x,y) + \cdots \right] \\ &+ \partial_y \left[ (ax + by)^n Q_{n,m}(x,y) + (ax + by)^{n-2} Q_{n-2,m}(x,y) + \cdots \right] \end{split}$$

Now when we substitute x = b, y = -a, all terms with a factor of (ax + by) vanish, so equating terms of like degree yields

$$Q_{n,m+1}(b,-a) = -2(bx + cy)Q_{n,m}(b,-a)$$
  

$$Q_{n-2,m+1}(b,-a) = -2(bx + cy)Q_{n-2,m}(b,-a)$$
  
...

Thus we see that

$$Q_{n-2r,m}(b,-a) = (-2(bx+cy))^m Q_{n-2r,0}(b,-a)$$

The asymptotes [132] are  $ax + by = \alpha$  where  $\alpha$  is the root of

$$\begin{split} t^{n}Q_{n,m}(b,-a) + t^{n-2}Q_{n-2,m}(b,-a) + \cdots \\ &= (-2(bx+cy))^{m} \left[ t^{n}Q_{n,0}(b,-a) + t^{n-2}Q_{n-2,0}(b,-a) + \cdots \right] \end{split}$$

and so the asymptotes of  $H_{n,m}$  and  $H_{n,0}$  are the same. But,

$$H_{n,0}(x,y) = -e^{\alpha x^2 + 2bxy} \left(\frac{\partial}{\partial x}\right)^n e^{-(\alpha x^2 + 2bxy)}$$

consists of parallel lines, so we can substitute y = 0 to see that the asymptotes  $ax + by = \alpha$  to  $H_{n,m}$  satisfy  $H_{n,0}(\alpha, 0) = 0$ . Since

$$H_{n,0}(x,0) = e^{\alpha x^2} \left(\frac{\partial}{\partial x}\right)^n e^{-\alpha x^2}$$

is the Hermite polynomial  $H_n(x/\sqrt{a})$  we conclude:

**Proposition 13.37.** The asymptotes of  $H_{n,m}$  are of the form

$$ax + by = \alpha$$
  $bx + cy = \beta$ 

where  $H_n(\alpha/\sqrt{a}) = 0$  and  $H_m(\beta/\sqrt{c}) = 0$ .

The coefficients of some Hermite polynomials obey a recursion similar to the three term recursion for orthogonal polynomials. Choose a positive symmetric matrix S, fix a positive integer n, and define

$$H_{n}(\mathbf{x}) = e^{\mathbf{x}S\mathbf{x}'} \left(\frac{\partial}{\partial x_{1}}\right)^{n} e^{-\mathbf{x}S\mathbf{x}'}$$
$$= \sum_{i=0}^{n} f_{i}(x_{1}, \dots, x_{d-1}) x_{d}^{i}$$

If we let  $v = (v_1, \dots, v_d)$  be the first row of S, then it easy to see that

$$H_{n+1} = -2(v_1x_1 + \dots + v_dx_d)H_n - 2nv_1H_{n-1}$$

We can use this recursion to prove by induction on n that

$$f_k = \left(\sum_{i=1}^{d-1} \frac{n+1}{n-k} \frac{\nu_i}{\nu_d} x_i\right) f_{k+1} - \left(\frac{(n+1)(n+2)}{2(n-k)} \frac{\nu_1}{\nu_d^2}\right) f_{k+2}$$

If d=2 , and  $S=\left(\begin{smallmatrix}a&b\\b&c\end{smallmatrix}\right)$  then

$$f_k(x) = \left(\frac{k+1}{n-k} \frac{a}{b}\right) x f_{k+1}(x) - \left(\frac{(k+1)(k+2)}{2(n-k)} \frac{a}{b^2}\right) f_{k+2}(x)$$

We get a three term recursion if we consider  $H_{n,1}$  instead of  $H_{n,0}$ , but the coefficients of  $H_{n,2}$  have a five term recursion that includes  $x^2 f_{n+2}$  and  $x f_{n+3}$ .

CHAPTER



# **Extending** P<sub>d</sub>

The goal in this Chapter is to investigate polynomials that only satisfy substitution for some values of y. There are several different approaches.

# **14.1** Taking the image of **P**<sub>2</sub>

We define the set  $\mathfrak{P}_2$  of polynomials as images of polynomials in  $\mathbf{P}_2$ . Lemma 13.20 provides the motivation for our analogues, as it shows that we can define  $\widehat{\mathbf{P}}$  in terms of linear transformations. Recall its results:

$$\widehat{\mathbf{P}} = \{\mathsf{T}(e^{x}) \mid \mathsf{T} \colon \mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}\}$$
$$\widehat{\mathbf{P}^{\text{pos}}} = \{\mathsf{T}(e^{x}) \mid \mathsf{T} \colon \mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}^{\text{pos}}\}$$

We now consider two variables.

#### Definition 14.1.

$$\begin{split} \mathfrak{P}_2 &= \left\{\mathsf{T}_*\left(\mathsf{f}\right) \mid \mathsf{f} \in \mathbf{P}_2^{\text{pos}} \wedge \mathsf{T} \colon \mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P} \wedge \mathsf{T} \text{ satisfies induction} \right\} \\ \mathfrak{P}_2^{\text{pos}} &= \left\{\mathsf{T}_*\left(\mathsf{f}\right) \mid \mathsf{f} \in \mathbf{P}_2^{\text{pos}} \wedge \mathsf{T} \colon \mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}^{\text{pos}} \wedge \mathsf{T} \text{ satisfies induction} \right\} \\ \widehat{\mathfrak{P}}_2 &= \text{ closure of } \mathfrak{P}_2 \\ \widehat{\mathfrak{P}}_2^{\text{pos}} &= \text{ closure of } \widehat{\mathfrak{P}}_2 \end{split}$$

The following lemma enumerates basic properties of  $\mathfrak{P}_2$ .

**Lemma 14.2.** Suppose  $g = T_* f$  is in  $\mathfrak{P}_2$ . Then

$$\begin{split} & 1. \ \mathbf{P}_{2}^{\text{pos}} \subsetneq \mathfrak{P}_{2}^{\text{pos}} \subsetneq \mathfrak{P}_{2} \subsetneq \mathfrak{P}_{1,1}. \\ & 2. \ g^{H}(x,1) \in \mathbf{P} \\ & 3. \ g(x,\alpha) \in \mathbf{P}^{\text{pos}} \text{ for } \alpha \geqslant 0. \end{split}$$

- 4.  $\frac{\partial g}{\partial x} \in \mathfrak{P}_2$
- 5.  $\frac{\partial g}{\partial u} \in \mathfrak{P}_2$
- 6. The coefficient of any power of y is in  $P^{\text{pos}}$ .
- 7. If  $h(x) \in P^{pos}(n)$  the the homogenization  $H = y^n h(x/y)$  is in  $\mathfrak{P}_2$ .
- 8. If  $S: \mathbb{P}^{pos} \longrightarrow \mathbb{P}^{pos}$  then  $S_*: \mathfrak{P}_2 \longrightarrow \mathfrak{P}_2$ .

If we assume that  $g \in \mathfrak{P}_2^{pos}$  then parts (4), (5), and (8) hold with  $\mathfrak{P}_2$  replaced by  $\mathfrak{P}_2^{pos}$  and in (2) we have that  $g^H(x, 1) \in \mathbf{P}^{pos}$ .

*Proof.* If we take T to be the identity then  $\mathbf{P}_2^{\text{pos}} \subseteq \mathfrak{P}_2$ . The example below shows that this is a proper containment. Since  $g^H = T_*(f^H)$  we see  $g^H(x,1) \in \mathbf{P}$ . If  $\alpha \ge 0$  then  $f(x, \alpha) \in \mathbf{P}^{\text{pos}}$ , and so  $g(x, \alpha) = T_*f(x, \alpha) \in \mathbf{P}^{\text{pos}}$ . If we define  $S(h) = \frac{\partial}{\partial x} T(h)$  then  $S: \mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}^{\text{pos}}$ , and so  $\frac{\partial g}{\partial x} \in \mathfrak{P}_2$ . Next,  $\frac{\partial g}{\partial y} = T_*(\frac{\partial f}{\partial y})$ , and so  $\frac{\partial g}{\partial y} \in \mathfrak{P}_2$  since  $\frac{\partial f}{\partial y} \in \mathbf{P}_2^{\text{pos}}$ . Since we can differentiate with respect to y, and substitute y = 0, we see that all coefficients of powers of y are in  $\mathbf{P}^{\text{pos}}$ . The coefficients of H are all positive, and substitution is clear. The last one is immediate from the definition:  $S_*(T_*f) = (ST)_*f$ .

**Example 14.3.** Members of  $\mathfrak{P}_2^{pos}$  do not necessarily satisfy substitution for negative  $\alpha$ . Consider T:  $x^n \mapsto \langle \underline{x} \rangle_n$  and  $f = (x+y+1)^2 \in \mathbf{P}_2^{pos}$ . If  $g(x,y) = T_* f$  then  $g \in \mathfrak{P}_2$ , yet  $g(x,-2) = x^2 - x + 1$  has complex roots.

#### **Constructing elements in** $\mathfrak{P}_2$

We can construct elements of  $\mathfrak{P}_2$  by explicitly exhibiting them in the form  $T_*f$  as in the definition, or by a general construction.

**Example 14.4.** The generalized Laguerre polynomials  $L_n(x; y)$  are in  $\mathfrak{P}_2^{\text{pos}}$ . This is a consequence of the surprising identity (14.1.1). If we define T:  $x^n \mapsto (\underline{x})_n$  then

$$T_{*}^{-1} \left< \underline{x + y + 1} \right>_{n} = n! L_{n}(x; y)$$
(14.1.1)

1

Now T<sup>-1</sup> maps  $\mathbf{P}^{\text{pos}}$  to itself and  $\langle \underline{x} + \underline{y} + \underline{1} \rangle_n \in \mathbf{P}_2^{\text{pos}}$ . It follows from (14.1.1) that  $L_n(x; \underline{y}) \in \mathfrak{P}_2^{\text{pos}}$ . I don't know a good proof for (14.1.1); there is an uninteresting inductive proof.

**Example 14.5.** We claim  $xy - 1 \in \mathfrak{P}_2$ . First of all, the transformation T:  $x^n \mapsto H_n(x/2)$  maps **P** to itself, and satisfies T(1) = 1, T(x) = x,  $T(x^2) = x^2 - 2$ . Now choose a positive integer m, and define  $S(f) = T^m(f)$ . Then S: **P**  $\longrightarrow$  **P** and

$$S(1) = 1$$
  $S(x) = x$   $S(x^2) = x^2 - 2m$ 

Set  $\varepsilon = 1/(2m)$ , and define

$$f_{m} = S_{*}(x + \varepsilon y)(y + \varepsilon x)$$
$$= xy(1 + \varepsilon^{2}) + \varepsilon y^{2} + \varepsilon x^{2} - \varepsilon y^{2}$$

 $\lim_{\mathfrak{m}\to\infty}\mathfrak{f}_\mathfrak{m}=xy-1$ 

A similar argument shows that  $xy - \alpha \in \mathfrak{P}_2$  for any positive  $\alpha$ . Notice that we are using a limit of transformations to show that an element is in  $\mathfrak{P}_2$ , not a sequence of elements.

**Example 14.6.** The map  $T: x^i \mapsto \langle \underline{x} \rangle_i$  satisfies  $\mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}^{\text{pos}}$ . If  $f(x) \in \mathbf{P}^{\text{pos}}$ , then the homogenization of f(x) is in  $\mathbf{P}_2$ . Consequently, if we define  $S(x^i) = \langle \underline{x} \rangle_i y^{n-i}$  then  $S(f) \in \mathfrak{P}_2^{\text{pos}}$ .

**Example 14.7.** We can use transformations to get elements of  $\mathfrak{P}_2$  or  $\mathfrak{P}_2^{\text{pos}}$  from elements in  $\mathbf{P}^{\text{pos}}$ . Suppose that T is a linear transformation satisfying T:  $\mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}^{\text{pos}}$ , and S is a linear transformation then the composition below gives elements of  $\mathfrak{P}_2$  from elements of  $\mathbf{P}^{\text{pos}}$ .

$$\mathbf{P}^{\text{pos}} \xrightarrow{S} \mathbf{P}_2 \xrightarrow{T_*} \mathfrak{P}_2$$

Useful choices of S are

$$f(x) \mapsto f(x + y)$$
  
 $f(x) \mapsto$  homogenization of  $f(x)$ 

In the latter case the composition is  $x^i \mapsto T(x^i)y^{n-i}$  where n is the degree of f.

#### The graphs of polynomials in $\mathfrak{P}_2$

The graphs of polynomials in  $\mathfrak{P}_2$  resemble polynomials in  $\mathbb{P}_2$  in the upper left quadrant, since they satisfy substitution for positive y, and their homogeneous part has all negative roots. Here are two examples. Figure 12.1 shows the graph of  $L_5(x;y)$ . In general, the Laguerre polynomials  $L_n(x;y)$  have a series expansion (7.10.2) that shows that  $L_n(x;y)$  is a polynomial of degree n in both x and y, with all positive terms. The leading term of the coefficient of  $x^k$  is

$$\frac{y^{n-k}}{(n-k)!k!}$$

and hence the homogeneous part of  $L_n(x; y)$  is  $\frac{1}{n!}(x + y)^n$ . Consequently, the solution curves are all asymptotic to lines with slope -1. Also,  $L_n(0; y) = \binom{n+y}{n}$  implies that the graph of  $L_n(x; y)$  meets the y-axis in -1, -2, ..., -n. All lines  $y = \alpha x$  where  $\alpha$  is positive will meet the curve in negative y values, so  $L_n(x; \alpha x) \in \mathbf{P}^{\text{pos}}$ .  $\mathfrak{P}_2$  is closed under differentiation with respect to y, as can also be seen in Figure 12.1.

For the next example, we take T:  $x^n \mapsto \langle \underline{x} \rangle_n$ , and pick (at random)

$$f = (x + y + 1)(x + 2y + 2)(x + 3y + 1)(x + 5y + 2).$$

then



Figure 14.1: The graph of T<sub>\*</sub>f

Figure 14.1 shows the graph of  $T_*f$ . In Figure 12.1 the solution curves all meet the y axis, but this is not true in Figure 14.1. Also, the line y = x does not intersect all the solution curves, and so  $f(x, x) \notin \mathbf{P}$ .

If we pick our slopes appropriately, then we can intersect all solution curves.

**Lemma 14.8.** Suppose that  $f(x, y) \in \mathfrak{P}_2^{\text{pos}}$ , and that  $\alpha$  is the largest root of  $f^{H}$ . If  $0 > \beta > \alpha$  then  $f(x, \beta x) \in \mathbf{P}^{\text{pos}}$ .

*Proof.* The line  $y = \beta x$  meets the x-axis to the right of intersection points of the solution curves, and is eventually less than every solution curve, so it must meet every one.

#### A partial Pólya-Schur result

Next, we look at two actions on  $\mathfrak{P}_2$  by polynomials.

**Lemma 14.9.** Suppose that  $g \in \mathfrak{P}_2$  and  $h \in \mathbf{P}^{pos}$ . Then

1.  $h(y)g(x,y) \in \mathfrak{P}_2$ .

2. 
$$h(\frac{\partial}{\partial x})g(x,y) \in \mathfrak{P}_2$$
.

3.  $h(\frac{\partial}{\partial u})g(x,y) \in \mathfrak{P}_2$ .

If  $g \in \mathfrak{P}_2^{pos}$  then all the statements are true with  $\mathfrak{P}_2$  replaced by  $\mathfrak{P}_2^{pos}$ .

*Proof.* Assume that  $g = T_*f$  as in the definition. If  $h(y) \in \mathbf{P}^{pos}$ , then  $h(y) \in \widehat{\mathbf{P}^{pos}}_2$ , so  $h(y)f(x,y) \in \widehat{\mathbf{P}^{pos}}_2$ . The first part now follows from

$$\mathsf{T}_*\mathsf{h}(\mathsf{y})\mathsf{f}(\mathsf{x},\mathsf{y}) = \mathsf{h}(\mathsf{y})\mathsf{T}_*\mathsf{f}(\mathsf{x},\mathsf{y}) = \mathsf{h}(\mathsf{y})\mathfrak{g}(\mathsf{x},\mathsf{y})$$

For the second part, it suffices to show that  $g + \alpha g' \in \mathfrak{P}_2$  if  $\alpha \ge 0$ . This follows from the fact that  $k \mapsto k + \alpha k'$  maps  $\mathbf{P}^{\text{pos}}$  to itself. To verify the third part, if suffices to show that  $g + \alpha g_y \in \mathfrak{P}_2$  for positive  $\alpha$ . But  $T_*(f + \alpha f_y) = g + \alpha g_y$ since  $T_*$  commutes with  $\frac{\partial}{\partial y}$ , and  $f + \alpha f_y \in \mathbf{P}_2$ .

We can use this result to go from an element of  $\mathfrak{P}_2$  to a transformation.

**Proposition 14.10.** If  $T_*(e^{xy}) \in \widehat{\mathfrak{P}}_2^{\operatorname{pos}}$  then  $T: P^{\operatorname{pos}} \longrightarrow P^{\operatorname{pos}}$ . In other words, suppose  $g(x,y) = \sum g_i(x) \frac{y^i}{i!} \in \widehat{\mathfrak{P}}_2^{\operatorname{pos}}$ . The linear transformation  $T: x^i \mapsto g_i$  defines a map  $P^{\operatorname{pos}} \longrightarrow P^{\operatorname{pos}}$ .

*Proof.* If we choose  $h(x) = \sum_{0}^{n} a_{i} x^{i} \in \mathbf{P}^{pos}$  then The coefficient of  $y^{n}$  in  $h^{rev}(\frac{\partial}{\partial u})g(x,y)$  is

$$\sum a_i g_i(x) = T(h)$$

Since  $h^{rev}(\frac{\partial}{\partial y})g(x,y) \in \widehat{\mathfrak{P}}_2$ , we know that all coefficients are in  $\mathbf{P}^{pos}$ .

# 14.2 Polynomials in several variables with positive coefficients

In this section we look at the set  $Q_{d,e}$  of certain polynomials in d + e variables that have non-negative coefficients. The motivation for this set comes from considering these three properties of  $P_2$ . Assume that  $f(x, y) \in P_2$ .

- 1.  $f(x, \alpha) \in \mathbf{P}$  for all  $\alpha$ .
- 2.  $f(\alpha, x) \in \mathbf{P}$  for all  $\alpha$ .
- 3.  $f(x, \alpha x) \in \mathbf{P}$  for all positive  $\alpha$ .

 $\Omega_{d,e}$  is constructed by starting with  $\mathbf{P}_d^{\text{pos}}$  and adding *e* new variables so that the resulting polynomials satisfy certain substitution conditions. We next define interlacing, and verify that it obeys the expected properties. The graphs of polynomials in  $\Omega_{0,2}$  are more complicated than those in  $\mathbf{P}_2^{\text{pos}}$ , but their behavior in each quadrant is explained by their defining conditions. We find some general constructions for polynomials in  $\Omega_{d,e}$  that are not necessarily in  $\mathbf{P}_{d+e}^{\text{pos}}$ . We then look at a special class of polynomials in  $\Omega_{d,e}$  - those that are linear in the last *e* variables. They can be identified with collections of polynomials that are defined on the corners of a cube. The three variable case of these polynomials leads to matrices that preserve interlacing.

#### Definitions

In order to better motivate our definition, we begin by restating the definition for interlacing mentioned above. Let F(x, y) = f(x) + yg(x). If  $F(x, \alpha)$  and  $F(x, \alpha x)$  are in  $\mathbf{P}^{\text{pos}}$  for all non-negative  $\alpha$  then f and g interlace. This suggests our first definition:

**Definition 14.11.**  $\Omega_{1,1}$  is the set of all polynomials F in x, y with non-negative coefficients that satisfy the two conditions

# $F(x, \alpha) \in \mathbf{P}^{pos}$ for all non-negative $\alpha$ $F(x, \alpha x) \in \mathbf{P}^{pos}$ for all non-negative $\alpha$

The general definition extends polynomials in  $\mathbf{P}_d^{\text{pos}}$  rather than polynomials in  $\mathbf{P}^{\text{pos}}_d$ . We let  $\mathbf{x} = (x_1, \dots, x_d)$ ,  $\mathbf{y} = (y_1, \dots, y_e)$ , and  $\mathbf{y}_b^a = (\alpha_1 \mathbf{x}_{b_1}^{c_1}, \dots, \alpha_e \mathbf{x}_{b_e}^{c_e})$ .

**Definition 14.12.**  $\Omega_{d,e}$  is the set of all polynomials  $f(\mathbf{x}, \mathbf{y})$  in d+e variables with non-negative coefficients such that for all non-negative  $\alpha_1, \ldots, \alpha_e$ , all  $b_i$  which satisfy  $b_i \in \{1, 2, \ldots, d\}$  and all  $c_1, \ldots, c_e$  which are 0 or 1 we have that  $f(\mathbf{x}, \mathbf{y}_b^a) \in \mathbf{P}_d^{\text{pos}}$ .

In other words, if we substitute either a non-negative constant or a non-negative constant multiple of some x variable for each y variable then the resulting polynomial is in  $\mathbf{P}_d^{\text{pos}}$ . If d is zero, then we substitute either a non-negative constant or a non-negative multiple of x for each y variable.

Notice that the conditions for a polynomial to be in  $\Omega_{1,2}$  are a subset of those that define  $\Omega_{0,3}$ . Consequently,  $\Omega_{0,3} \subset \Omega_{1,2}$ . More generally we have

**Lemma 14.13.** If  $d \ge 2$  and  $e \ge 0$  then  $Q_{d,e} \subset Q_{d-1,e+1}$  and  $Q_{0,e+1} \subset Q_{1,e}$ .

*Proof.* Since a polynomial in  $\Omega_{0,e+1}$  satisfies all the conditions of  $\Omega_{1,e}$  plus a few more involving substitutions for the first variable, we have  $\Omega_{0,e+1} \subset \Omega_{1,e}$ . If  $d \ge 2$  then choose  $f \in \Omega_{d,e}$  and consider the substitution conditions that must be met for f to be in  $\Omega_{d-1,e+1}$ :

$$f(x_1, \dots, x_{d-1}, \alpha_1, \alpha_2 x_{b_2}^{c_2}, \dots, \alpha_{e+1} x_{b_{e+1}}^{c_{e+1}}) \in \mathbf{P}_{d-1}^{\text{pos}}$$
(14.2.1)

$$f(x_1, \dots, x_{d-1}, \alpha_1 x_{b_1}, \alpha_2 x_{b_2}^{c_2}, \dots, \alpha_{e+1} x_{b_{e+1}}^{c_{e+1}}) \in \mathbf{P}_{d-1}^{\text{pos}}$$
(14.2.2)

(14.2.1) holds since

$$f(x_1,\ldots,x_{d-1},x_d,\alpha_2x_{b_2}^{c_2},\ldots,\alpha_{e+1}x_{b_{e+1}}^{c_{e+1}}) \in \mathbf{P}_d^{\text{pos}}$$

and substitution of a constant for  $x_d$  gives a polynomial in  $\mathbf{P}_{d-1}^{\text{pos}}$ . (14.2.6) holds because we can substitute any positive multiple of a variable for a variable in  $\mathbf{P}_d^{\text{pos}}$ , and the result is in  $\mathbf{P}_{d-1}^{\text{pos}}$ .

 $\Omega_{d,e}$  is not empty since  $\mathbf{P}_{d+e}^{\text{pos}} \subset \Omega_{d,e}$ . We allow some coefficients to be zero, so  $\Omega_{d,e} \subset \Omega_{d,e+1}$ , and therefore we have the containments

$$\begin{split} \mathbf{P}^{\text{pos}} &= \mathcal{Q}_{1,0} &= & \mathcal{Q}_{0,1} \\ & & \subset & & \subset \\ \mathbf{P}^{\text{pos}}_2 &= \mathcal{Q}_{2,0} &\subset & \mathcal{Q}_{1,1} & \supset & \mathcal{Q}_{0,2} \\ & & \subset & & \subset \\ \mathbf{P}^{\text{pos}}_3 &= \mathcal{Q}_{3,0} &\subset & \mathcal{Q}_{2,1} &\subset & \mathcal{Q}_{1,2} & \supset & \mathcal{Q}_{0,3} \end{split}$$

**Example 14.14.** If we choose  $f_i \leq g_i \in \mathbf{P}^{pos}$  then it follows from the results in the next section that

$$\prod_{i=1}^{n} (f_i + yg_i) \in Q_{1,1}$$

The polynomial  $f(x, y) = x + y^2$  is in  $\Omega_{1,1}$  but is not in  $\Omega_{0,2}$  since  $f(1, y) \notin \mathbf{P}$ . The polynomial

$$f = 2x + xy + (1 + 3x)z + yz$$
(14.2.3)

is in  $\Omega_{1,2}$  but is not in  $\Omega_{0,3}$ . To verify this we just make the four substitutions that define  $\Omega_{1,2}$ , and check that in each case the polynomials have all real roots. It is not in  $\Omega_{0,3}$  since f(.01, x, .01x) has imaginary roots.

**Example 14.15.** Suppose that  $f \leq g$  where  $f, g \in \mathbf{P}^{pos}$  have positive leading coefficients. We know that (xg, f, g) is totally interlacing. If  $\alpha$  is positive then the matrix below on the left is totally non-negative

$$\begin{pmatrix} 1 & \alpha & 0 \\ 0 & 1 & \alpha \end{pmatrix} \begin{pmatrix} xg \\ f \\ g \end{pmatrix} = \begin{pmatrix} xg + \alpha f \\ f + \alpha g \end{pmatrix}$$

and hence

$$xg + \alpha f \longleftarrow f + \alpha g$$
 (14.2.4)

If we define

$$h(x, y, z) = xg + zf + y(f + zg)$$
 (14.2.5)

then (14.2.4) shows that  $h(x, y, \alpha) \in Q_{1,1}$  for positive  $\alpha$ . Also, replacing  $\alpha$  by  $1/\alpha$  and multiplying by  $\alpha$  yields  $\alpha xg + f \leftarrow \alpha f + g$ . Multiplying the right by x shows that

$$xg + (\alpha x)f \longleftarrow f + (\alpha x)g$$

which implies  $h(x, y, \alpha x) \in Q_{1,1}$ . Consequently,  $h(x, y, z) \in Q_{1,2}$ .

#### **Elementary properties**

We define interlacing, and then show that  $Q_{d,e}$  satisfies the expected closure properties. If we consider the definition of  $Q_{1,1}$ , we see that we can define interlacing in  $\mathbf{P}^{\text{pos}}$  as follows:

 $f \underline{\lessdot} g \text{ in } \mathbf{P}^{p \, o \, s} \text{ if and only if } deg(f) > deg(g) \text{ and } f + yg \in \mathbb{Q}_{1,1}.$ 

If f, g  $\in Q_{d,e}$  then we say that  $f \leq g$  if and only if the total x-degree of f is greater than that of g, and  $f + y_{e+1}g \in Q_{d,e+1}$ . All this means is that  $f \leq g$  if and only if

$$\begin{split} & \text{total } x\text{-degree}(f) > \text{total } x\text{-degree}(g) \\ & \text{f}(x,y) + \alpha g(x,y) \in \mathfrak{Q}_{d,e} \\ & \text{f}(x,y) + \alpha x_{i}g(x,y) \in \mathfrak{Q}_{d,e} \text{ (for any } 1 \leqslant i \leqslant d) \end{split}$$

We use  $\leq$  to define  $\leftarrow$ :

 $f \leftarrow g$  iff we can write  $g = \alpha f + h$  where  $\alpha \ge 0$  and  $f \le h$ .

In order to show that  $f \leq g$  all we need to do is to show that all substitutions interlace.

**Lemma 14.16.** Suppose that  $f, g \in Q_{d,e}$ , and that the x-degree of f is greater than the x-degree of g. The following are equivalent

- 1.  $f(\mathbf{x}, \mathbf{y}_{\mathbf{b}}^{\mathbf{a}}) \leq g(\mathbf{x}, \mathbf{y}_{\mathbf{b}}^{\mathbf{a}})$  in  $\mathbf{P}_{d}^{pos}$  for all appropriate  $\mathbf{a}$  and  $\mathbf{b}$
- 2.  $f(\mathbf{x}, \mathbf{y}) + y_{e+1}g(\mathbf{x}, \mathbf{y}) \in Q_{d,e+1}$
- 3.  $f(\mathbf{x}, \mathbf{y}) \leq g(\mathbf{x}, \mathbf{y})$  in  $Q_{d,e}$ .

*Proof.* The second and third are equivalent since they are the definition of interlacing. To show that (3) implies (1) note that interlacing implies that  $f + y_{e+1}g \in Q_{d,e+1}$ . We can substitute for  $y_{e+1}$  and conclude that

$$\begin{split} \mathsf{f}(\mathbf{x}, \mathbf{y}_{\mathbf{b}}^{\mathbf{a}}) + \alpha \, \mathsf{g}(\mathbf{x}, \mathbf{y}_{\mathbf{b}}^{\mathbf{a}}) \in \mathbf{P}_{\mathrm{d}}^{\mathrm{pos}} \\ \mathsf{f}(\mathbf{x}, \mathbf{y}_{\mathbf{b}}^{\mathbf{a}}) + \alpha \mathsf{x}_{\mathrm{i}} \, \mathsf{g}(\mathbf{x}, \mathbf{y}_{\mathbf{b}}^{\mathbf{a}}) \in \mathbf{P}_{\mathrm{d}}^{\mathrm{pos}} \end{split}$$

and the conclusion follows from Lemma 10.38. That (1) implies (3) is similar.  $\Box$ 

**Lemma 14.17.** Assume that  $f, g, h \in Q_{d,e}$ .

- 1. fg  $\in Q_{d,e}$ .
- 2. If  $f \leftarrow g$  then  $hf \leftarrow hg$ .
- 3. If  $f \leftarrow g$  and  $f \leftarrow h$  then  $g + h \in Q_{d,e}$  and  $f \leftarrow g + h$ .
- 4. If  $g \leftarrow f$  and  $h \leftarrow f$  then  $g + h \in Q_{d,e}$  and  $g + h \leftarrow f$ .
- 5. If  $\alpha_1, \ldots, \alpha_d$  and  $\beta_1, \ldots, \beta_e$  are non-negative then  $f(\alpha_1 x_1, \ldots, \alpha_d x_d, \beta_1 y_1, \ldots, \beta_e y_e) \in Q_{d,e}$ .

*Proof.* We first consider the case where all "—" are " $\leq$ ". The first one is immediate from the observation that  $fg(x, y_b^a) = f(x, y_b^a) g(x, y_b^a)$  The second follows from the first since  $hf + y_{e+1}hg = h(f + y_{e+1}g)$ , and  $f + y_{e+1}g \in Q_{d,e+1}$  by definition of interlacing.

For assertion (3), in order to verify that  $f + y_{e+1}(g+h) \in Q_{d,e+1}$  we use the properties of interlacing in  $\mathbf{P}_d^{\text{pos}}$ . Since  $f \leq g$ , we know that by Lemma 14.16

$$f(\mathbf{x}, \mathbf{y}_b^{\mathbf{a}}) \leq g(\mathbf{x}, \mathbf{y}_b^{\mathbf{a}})$$
  
$$f(\mathbf{x}, \mathbf{y}_b^{\mathbf{a}}) \leq h(\mathbf{x}, \mathbf{y}_b^{\mathbf{a}})$$

and adding these interlacings together gives

$$f(\mathbf{x}, \mathbf{y}_{\mathbf{b}}^{\mathbf{a}}) \leq g(\mathbf{x}, \mathbf{y}_{\mathbf{b}}^{\mathbf{a}}) + h(\mathbf{x}, \mathbf{y}_{\mathbf{b}}^{\mathbf{a}})$$

It follows that  $f + y_{e+1}(g + h) \in Q_{d,e+1}$ . The proof of the next statement is similar. The last statement is immediate from the definition.

In the general case  $f \leftarrow g$  we use the representation  $g = \alpha f + k$  where  $\alpha \ge 0$  and  $f \le h$ . For instance, to establish (2), we know that  $f \le k$  implies  $hf \le hk$ , and therefore  $hg = \alpha hf + hk$  implies  $hf \leftarrow hg$ .

Factors of polynomials in  $Q_{d,e}$  are generally in  $Q_{d,e}$ .

**Lemma 14.18.** If  $fg \in Q_{1,e}$ , and f has only non-negative coefficients then  $f \in Q_{1,e}$ . If  $fg \in Q_{d,e}$ , and f has only non-negative coefficients and  $f(\mathbf{x}, \mathbf{y}_{\mathbf{b}}^{a}) \in \mathbf{P}_{d}^{\text{pos}}$  for all appropriate  $\mathbf{y}_{\mathbf{b}}^{a}$  then  $f \in Q_{d,e}$ .

*Proof.* If some substitution resulted in a complex root for f, then it would be a complex root for fg as well. Since f has non-negative coefficients, f is in  $Q_{d,e}$ .

#### Constructing polynomials in $Q_{1,2}$

We can easily construct polynomials in  $\Omega_{1,2}$  using products. If  $f \leq g$ ,  $h \leq k$  are polynomials in  $\mathbf{P}^{\text{pos}}$  then  $f + yg \in \Omega_{1,2}$ ,  $h + zk \in \Omega_{1,2}$ , and therefore  $(f + yg)(h + zk) \in \Omega_{1,2}$ . However, these polynomials are also in  $\overline{\mathbf{P}}_3^{\text{pos}}$ . We do not know if  $\Omega_{1,2}$  is different from  $\overline{\mathbf{P}}^{\text{pos}}_{4_3}$ .

**Lemma 14.19.** Suppose that f, g, h,  $k \in P^{pos}$  have all positive coefficients. If

f k where all interlacings are "
$$\ll$$
" or all are " $\ll$ " and the determinant  $\begin{vmatrix} f & g \\ h & k \end{vmatrix}$ 

has no negative roots then

$$f + yg + zh + yzk \in Q_{1,2}$$

*Proof.* It suffices to show that  $f + \alpha g < k + \alpha h$  and  $f + \alpha xg < k + \alpha xh$  for all positive  $\alpha$ . The interlacing assumptions imply that all four linear combinations are in  $\mathbf{P}^{\text{pos}}$ . We now follow the proof of Lemma 1.52 to show that the first interlacing holds since the determinant in the conclusion is never zero for negative x. Again by following the proof of Lemma 1.52, the second holds since  $\left| \frac{f \times g}{h \times k} \right| = x \left| \frac{f g}{h \times k} \right|$ .

Note that Example 14.15 is a special case of the Lemma. Take

 $xg \int_{f}^{f} g$ . Since  $\begin{vmatrix} xg & f \\ g & f \end{vmatrix} = xg^2 - f^2$  clearly has no negative roots, the

Lemma applies.

It is easy to find polynomials in  $Q_{1,2}$  using Lemma 14.19. This is a special case of a more general construction in the next section.

**Lemma 14.20.** Choose  $w(x, y, z) \in \mathbf{P}_3^{pos}$ , and write

$$w(x, y, z) = f(x) + g(x)y + h(x)z + k(x)yz + \cdots$$

Then  $f(x) + g(x)y + h(x)z + k(x)yz \in Q_{1,2}$ .

*Proof.* f, g, h, k are all in  $P_2$ , and Theorem 9.113 shows that the determinant is never zero.

For instance, if  $v(x) \in \mathbf{P}^{pos}$ , then the Taylor series expansion of v(x+y+z) shows that

$$\nu(\mathbf{x}) + \nu'(\mathbf{x}) \cdot (\mathbf{y} + z) + \frac{\nu''(\mathbf{x})}{2} \mathbf{y} \mathbf{z} \in \mathcal{Q}_{1,2}$$

#### **Polynomials linear in the y variables**

An interesting class of polynomials in  $Q_{d,e}$  consists of those polynomials that are linear in the y variables. In general, if f is a polynomial in  $Q_{d,e}$  that is linear in  $y_1, \ldots, y_d$  then we can represent f by labeling the vertices of a d-cube with appropriate coefficients of f.

We can find such polynomials in  $\Omega_{d,e}$  by taking initial terms of polynomials in  $\mathbf{P}_{d+e}^{\text{pos}}$ .

**Lemma 14.21.** Suppose  $f(\mathbf{x}, \mathbf{y}) \in \mathbf{P}_{d+e}^{pos}$ , and let  $g(\mathbf{x}, \mathbf{y})$  consist of all terms of f that have degree at most one in every y variable. Then  $g \in Q_{d,e}$ .

*Proof.* For simplicity we illustrate the proof with e = 2; the general proof is the same. If we write

$$f(\mathbf{x}, \mathbf{y}, z) = f_{00}(\mathbf{x}) + f_{10}(\mathbf{x})\mathbf{y} + f_{01}(\mathbf{x})z + f_{11}(\mathbf{x})\mathbf{y}z + \cdots$$
$$g(\mathbf{x}, \mathbf{y}, z) = f_{00}(\mathbf{x}) + f_{10}(\mathbf{x})\mathbf{y} + f_{01}(\mathbf{x})z + f_{11}(\mathbf{x})\mathbf{y}z$$

then the coefficient of yz in  $(y + x_i^r \alpha)(z + x_i^s \beta)f(x, y, z)$  is

$$g(\mathbf{x}, \mathbf{x}_{i}^{r} \boldsymbol{\alpha}, \mathbf{x}_{i}^{s} \boldsymbol{\beta}) = f_{00}(\mathbf{x}) + f_{10}(\mathbf{x})\mathbf{x}_{i}^{r} \boldsymbol{\alpha} + f_{01}(\mathbf{x})\mathbf{x}_{i}^{s} \boldsymbol{\beta} + f_{11}(\mathbf{x})\mathbf{x}_{i}^{r} \mathbf{x}_{i}^{s} \boldsymbol{\alpha} \boldsymbol{\beta}$$

Consequently, we see that since  $\alpha$ ,  $\beta$  are positive,

$$g(\mathbf{x}, \alpha, \beta), g(\mathbf{x}, x_i\alpha, \beta), g(\mathbf{x}, \alpha, x_i\beta), g(\mathbf{x}, x_i\alpha, x_i\beta)$$

are all in  $\mathbf{P}_{d}^{\text{pos}}$ , and so  $g(\mathbf{x}, \mathbf{y}, z) \in \mathcal{Q}_{d,e}$ .

**Lemma 14.22.** Suppose  $f(x, y) \in Q_{1,e}$  and has x degree r and  $y_i$  degree  $s_i$ . The reverse polynomial below is in  $Q_{1,e}$ :

$$\mathbf{x}^{\mathbf{r}}\mathbf{y}_{1}^{s_{1}}\cdots\mathbf{y}_{e}^{s_{e}} f(\frac{1}{\mathbf{x}},\frac{1}{\mathbf{y}_{1}},\ldots,\frac{1}{\mathbf{y}_{e}})$$

*Proof.* Any substitution for  $y_i$  by a constant yields a polynomial in some  $Q_{1,e'}$  where e' < e. We can also scale each  $y_i$ , so we only need to consider the substitution  $y_i \rightarrow x$ . Thus we need to verify that

$$x^{r+s_1+\dots+s_e} f(\frac{1}{x},\dots,\frac{1}{x}) \in \mathbf{P}^{pos}$$
 (14.2.6)

If a monomial in f(x, y) is  $a_I x^i y^I$  with total degree i + |I|, then in (14.2.6) the degree is n - (i + |I|) where  $n = r + s_1 + \dots + s_e$ . Consequently, (14.2.6) is the usual reverse of  $f(x, x, \dots, x)$  which we know to be in  $\mathbf{P}^{\text{pos}}$ .

For instance, if d = 1 then Lemma 14.22 shows that if

$$f_{11}(x) + y f_{21}(x) + z f_{12}(x) + y z f_{22}(x) \in Q_{1,2}$$

then the reverse is also in  $Q_{1,2}$ 

$$yz f_{11}(x) + z f_{21}(x) + y f_{12}(x) + f_{22}(x) \in Q_{1,2}$$

## Matrices preserving interlacing

Matrices determined by polynomials linear in y and z are closed under multiplication. We can use this fact to show that these matrices also preserve interlacing. This is a special case of Theorem 24.11.

**Lemma 14.23.** Suppose that  $f, g \in Q_{d,2}$  are linear in y and z and write

$$f = f_{11} + yf_{12} + zf_{21} + yzf_{22}$$
$$g = g_{11} + yg_{12} + zg_{21} + yzg_{22}$$

*Represent* f, g by matrices in the following way

$$F = \begin{pmatrix} f_{12} & f_{22} \\ f_{11} & f_{21} \end{pmatrix}$$
$$G = \begin{pmatrix} g_{12} & g_{22} \\ g_{11} & g_{21} \end{pmatrix}$$

If the product is  $FG = \begin{pmatrix} h_{12} & h_{22} \\ h_{11} & h_{21} \end{pmatrix}$ 

then  $h_{11}+yh_{12}+zh_{21}+yzh_{22}\in \mathbb{Q}_{d,2}$ 

*Proof.* The proof is an application of the Leibnitz rule in disguise. We know that

$$f_{11} + yf_{12} \leq f_{21} + yf_{22}$$
$$g_{11} + zg_{21} \leq g_{12} + zg_{22}$$

and multiplying by the right hand sides

$$(f_{11} + yf_{12})(g_{12} + zg_{22}) \leq (f_{21} + yf_{22})(g_{12} + zg_{22}) (f_{21} + yf_{22})(g_{11} + zg_{21}) \leq (f_{21} + yf_{22})(g_{12} + zg_{22})$$

and adding them together yields

$$(f_{11} + yf_{12})(g_{12} + zg_{22}) + (f_{21} + yf_{22})(g_{11} + zg_{21}) \leq (f_{21} + yf_{22})(g_{12} + zg_{22})$$

The matrix representation of the left hand polynomial is exactly FG.

**Corollary 14.24.** If  $f \in Q_{1,2}$  is given as in the lemma,  $g_1 \longleftarrow g_2$  and we set

$$\begin{pmatrix} f_{12} & f_{22} \\ f_{11} & f_{21} \end{pmatrix} \begin{pmatrix} g_2 \\ g_1 \end{pmatrix} = \begin{pmatrix} h_2 \\ h_1 \end{pmatrix}$$

*then*  $h_1 \leftarrow h_2$ .

*Proof.* We check that  $g = (g_1 + yg_2)(1 + z)$  is in  $\Omega_{1,2}$ . Since  $g_1 \leftarrow g_2$  we know that  $g_1 + yg_2 \in \Omega_{1,1}$ . Multiplying by 1 + z shows  $g \in \Omega_{1,2}$ .

The matrix of g is  $\begin{pmatrix} g_2 & g_2 \\ g_1 & g_1 \end{pmatrix}$ . Applying the lemma and considering only the first column of the product gives the conclusion.

We can use the determinant condition of Lemma 14.19 to find such matrices.

**Corollary 14.25.** Suppose that f, g, h,  $k \in P^{pos}$  have all positive coefficients. If

$$f \left[ \begin{array}{c} k \\ h \end{array} \right] k$$
 where all interlacings are < and the determinant  $\left| \begin{array}{c} f & g \\ h & k \end{array} \right|$  has no negative

roots then the matrix  $\begin{pmatrix} h & k \\ f & q \end{pmatrix}$  preserves interlacing for polynomials in  $P^{pos}$ .

**Example 14.26.** The determinant of the polynomials in the interlacing square below has only positive roots, so the corresponding matrix preserves interlacing. If we multiply this matrix by  $\begin{pmatrix} 0 & x \\ 1 & 0 \end{pmatrix}$  which preserves interlacing, we find that the matrix of polynomials below preserves interlacing for polynomials in  $\mathbf{P}^{\text{pos}}$ .

$$4 + 88x \xrightarrow{>} 8 + 260x + 216x^{2}$$

$$\ll \uparrow \qquad \qquad \uparrow \checkmark$$

$$32 + 64x \xrightarrow{>} 64 + 416x$$

$$\begin{pmatrix} 32x + 64x^{2} & 64x + 416x^{2} \\ 4 + 88x & 8 + 260x + 216x^{2} \end{pmatrix}$$

We can not give a characterization of the matrices that preserve interlacing, but we can find a few restrictions. We begin with the linear case.

**Lemma 14.27.** Suppose  $f, g \in Q_{1,2}$  have the property that for all  $p \leftarrow q \in Q_{1,2}$  it holds that  $fp + gq \in Q_{1,2}$ . Then  $g \leftarrow f$ .

*Proof.* If we choose  $p = \alpha q$  for positive  $\alpha$  then  $p \leq q$ , and so  $p(\alpha f + g) \in Q_{1,2}$ . Since factors of polynomials in  $Q_{1,2}$  are in  $Q_{1,2}$ , we conclude that  $\alpha f + g \in Q_{1,2}$ . Next, if we choose  $p = \alpha x q$  then again  $p \leq q$ , and so  $xf + g \in Q$ . It follows that  $g \leq f$ .

Lemma 14.28. Suppose that

$$\begin{pmatrix} f_{12} & f_{22} \\ f_{11} & f_{21} \end{pmatrix} \begin{pmatrix} g_2 \\ g_1 \end{pmatrix} = \begin{pmatrix} h_2 \\ h_1 \end{pmatrix}$$

and the  $f_{ij}$  have the property that whenever  $g_1 \leq g_2$  in  $Q_{1,2}$  then it holds that  $h_1 \leq h_2$  in  $Q_{1,2}$ . Then, the  $f_{ij}$  are in  $Q_{1,2}$  and form an interlacing square

$$\begin{array}{ccc} f_{12} \twoheadleftarrow f_{22} \\ \downarrow & \downarrow \\ f_{11} \twoheadleftarrow f_{21} \end{array}$$

*Proof.* The horizontal arrows follow from the previous lemma. If we take a limit, we may assume that  $g_1$  is zero. Thus, we have that  $f_{11}g_2 \leq f_{12}g_2$  which implies that  $f_{11} \leq f_{12}$  and that  $f_{11}$ ,  $f_{12}$  are in  $Q_{1,2}$ . Taking  $g_2$  to be zero gives the properties of the second column.

#### 14.3 Epsilon substitution

We have looked at properties of polynomials f(x, y) where  $f(x, \alpha)$  is guaranteed to be in **P** for only positive  $\alpha$ . We now reduce the range of acceptable  $\alpha$ .

**Definition 14.29.**  $\mathcal{P}_d^{\epsilon}$  consists of those polynomials  $f(x; y_1, \ldots, y_d)$  with the property that there is an  $\epsilon$ , depending on f, such that

 $|\alpha_1| \leq \varepsilon, \ldots, |\alpha_d| \leq \varepsilon \implies f(x; \alpha_1, \ldots, \alpha_d) \in \mathbf{P}$ 

It's easy to tell from the graph if a polynomial is in  $\mathscr{P}_1^{\varepsilon}$ . For instance, Figure 14.3 is the graph of a polynomial of degree 12. We can see that every horizontal line  $y = \alpha$  intersects the graph in 12 points if  $|\alpha|$  is sufficiently small, and consequently  $f \in \mathscr{P}_1^{\varepsilon}$ .

Here are a few simple facts that need no proof.

- 1. If f,  $g \in \mathcal{P}_d^{\epsilon}$  then  $fg \in \mathcal{P}_d^{\epsilon}$ .
- 2. If  $f \in \mathcal{P}_d^{\epsilon}$  then  $\frac{\partial f}{\partial x} \in \mathcal{P}_d^{\epsilon}$ .
- 3. If  $f \in \mathcal{P}_d^{\epsilon}$  then  $f(x; 0, \ldots, 0) \in \mathbf{P}$ .

Let's consider two examples.

**Example 14.30.** We will show that  $\sum_{i=0}^{n} P_i(x) {n \choose i} y^{n-i} \in \mathfrak{P}_1^{\epsilon}$  where  $P_i$  is the Legendre polynomial. We make use of the formula [168]

$$\sum_{i=0}^{n} {n \choose i} P_i(x) y^{n-i} = (1 + 2xy + y^2)^{n/2} P_n((x+y)(1 + 2xy + y^2)^{-1/2})$$
(14.3.1)

We show that we can take  $\varepsilon=1.$  If  $\mathsf{P}_n(\beta)=0,$  then we know that  $|\beta|<1.$  Solving

$$\beta = (x + y)(1 + 2xy + y^2)^{-1/2}$$

for x yields

$$x=-y+y\ \beta^2-\sqrt{\beta^2-y^2\ \beta^2+y^2\ \beta^4}$$

Now if |y| < 1, then |x| < 1. In this case not only is (14.3.1) in  $\mathcal{P}_1^{\epsilon}$ , but we have the stronger result that

$$|\alpha| \leqslant 1 \quad \Longrightarrow \quad \sum_{i=0}^{n} P_{i}(x) {n \choose i} \alpha^{n-i} \in \mathbf{P}^{(-1,1)}$$

Example 14.31. Define the polynomials of degree n

$$\begin{split} f_n(x,y) &= \frac{\partial^n}{\partial y^n} \left( x^2 + y^2 - 1 \right)^n \\ h_n(y) &= \left( \frac{d}{dy} \right)^n (y^2 - 1)^n \end{split}$$

Figure 14.2 is the graph of  $f_6$ . We have the relationship

$$f_n(x,y) = 0$$
 iff  $h_n\left(\frac{y}{\sqrt{1-x^2}}\right) = 0$ 

Note that h is, up a factor, the Legendre polynomial, and so has the same roots. Since  $f_n(0, y) = h_n(y)$  we see that the intersection of the graph of  $f_n$  with the y axis is at the roots of the Legendre polynomial. As  $n \to \infty$ , these roots become dense in (-1, 1).

The relationship between  $f_n$  and  $h_n$  shows that the graph of  $f_n$  consists of ovals through the roots of  $h_n$  on the y axis that all meet at (1,0) and (-1,0). Consequently, any horizontal line close enough to the x-axis will meet the graph in n points, and so  $f_n \in \mathcal{P}_1^{\varepsilon}$ .

Since the smallest positive root of the Legendre polynomial becomes arbitrarily small, we see that the  $\varepsilon$  for which the interval  $-\varepsilon \leqslant \alpha \leqslant \varepsilon$  implies  $f_n(x, \alpha) \in \mathbf{P}$  becomes arbitrarily small. None the less, all  $f_n$  are in  $\mathscr{P}_1^{\varepsilon}$ .



Figure 14.2: The graph of  $\partial_{u}^{6}(x^{2} + y^{2} - 1)^{6} = 0$ 

In the case d = 1, this last observation says that the constant term (with respect to y) of a polynomial in  $\mathcal{P}_1^{\epsilon}$  is in **P**. If all derivatives with respect to y are in  $\mathcal{P}_1^{\epsilon}$  then all the coefficients are in **P**.

We now show that 2-variable analogs (14.3.2) of the Legendre polynomials are in  $\mathscr{P}_1^e$ , and consequently all their coefficients are in **P**. Figure 14.3 is the graph of  $\mathfrak{p}_{6,6,12}$ .

$$p_{n,m,r}(x,y) = \left(\frac{\partial}{\partial x}\right)^n \left(\frac{\partial}{\partial y}\right)^m \left(x^2 + y^2 - 1\right)^r$$
(14.3.2)



Figure 14.3: The graph of  $p_{6,6,12}$ 

Since differentiation with respect to x preserves  $\mathscr{P}_1^{\varepsilon}$ , we may assume that n = 0. We need to consider how differentiation with respect to y affects the graph of  $p_{0,m,r}$ . In Figure 14.4 we have the graph of  $p_{0,6,6}$  and  $p_{0,7,6}$ . The light lines are in  $p_{0,6,6}$  (see Figure 14.2) and the dark lines are in  $p_{0,7,6}$ . It is easy to

see that



Figure 14.4: The graph of  $\vartheta_y^6(x^2+y^2-1)^6=0$  and its y derivative

We determine a point (a, b) in  $p_{0,7,6}$  by considering the line x = a, and finding the intersection of the line with the graph. If -1 < a < 1 then there are 6 intersection points. If (a, b) is in the graph of  $p_{0,7,6}$  then b is a root of  $\frac{\partial}{\partial y}p_{0,6,6}(a, y)$ , and so there are five possible b's, and they interlace the six intersection points.

Thus, the graph consists of five curves joining (-1,0) and (1,0), each one passing between a different pair of adjacent curves of  $p_{0,6,6}$ , as can be seen in the figure.

This is the general situation:  $h_{0,0,r}$  consists of r circles  $x^2 + y^2 = 1$ . Each successive y derivative introduces a curve between existing curves, each joining (-1) and (1,0).

**Proposition 14.32.** All  $p_{n,m,r}$  in (14.3.2) lie in  $\mathscr{P}_1^{\varepsilon}$ . All coefficients of powers of y in  $p_{n,m,r}$  lie in  $\mathbf{P}^{[-1,1]}$ .

*Proof.* The only part that remains to be proved is that the roots of the coefficients lie in [-1, 1], but this is clear from the graph.

We can show that  $p_{0,m,r}$  only meets the axis in  $\pm 1$ . We claim that

$$\left(\frac{\partial}{\partial y}\right)^{n} \left(x^{2} + y^{2} - 1\right)^{r} (x, 0) = \begin{cases} n! \binom{r}{n/2} (x^{2} - 1)^{r - n/2} & \text{r even, } r \ge n\\ 0 & \text{otherwise} \end{cases}$$
(14.3.3)

$$\sum_{n=1}^{\infty} \left[ \left( \frac{\partial}{\partial z} \right)^n (x^2 + y^2 - 1)^r \right] (x, 0) \frac{z^n}{z^n} = (x^2 + z^2)^n$$

This is an easy consequence of Taylor series:

$$\sum_{n=0} \left\lfloor \left( \frac{\partial}{\partial y} \right) \quad \left( x^2 + y^2 - 1 \right)^r \right\rfloor \ (x,0) \ \frac{z^n}{n!} = (x^2 + z^2 - 1)^r$$

so the left hand side of (14.3.3) is the coefficient of  $z^n$  in  $(x^2 + z^2 - 1)^n$  which yields the right hand side of (14.3.3).

CHAPTER



# Generating functions of linear transformations

The goal of this chapter is to explain the use of generating functions to prove that linear transformations preserve particular spaces of polynomials.

# 15.1 The idea of generating function

Suppose T is a linear transformation and S is a set of polynomials. Generating functions are a method for proving that T maps S to S. At its simplest, there is a function G, the *base generating function*, and a set of functions S' such that

$$\mathsf{T}(\mathsf{G}) \in \mathsf{S}' \implies \mathsf{T}: \mathsf{S} \longrightarrow \mathsf{S}. \tag{15.1.1}$$

We say that we have a *Pólya-Schur* result if in addition

$$\mathsf{T}:\mathsf{S}\longrightarrow\mathsf{S}\implies\mathsf{T}(\mathsf{G})\in\mathsf{S}'.$$
(15.1.2)

The general method relies on expressing T(f) in terms of T(G), where  $f \in S$ . T(G) is the induced map on G, defined by  $T(\mathbf{x}^{I} \mathbf{y}^{J}) = T(\mathbf{x}^{I}) \mathbf{y}^{J}$ . We need the following data, where  $\mathbf{x} = (x_1, \dots, x_d)$  and  $\mathbf{y} = (y_1, \dots, y_d)$ :

S	A set of polynomials in <b>x</b> .
S′	A set of functions in <b>x</b> and <b>y</b> satisfying
	$f(\mathbf{x}, \mathbf{y}) \in S' \implies f(\mathbf{x}, 0) \in S.$
G	The base generating function.
W	A map from S to maps $S' \rightarrow S'$ .

These combine in the fundamental identity for  $f \in S$ :

$$(Wf) T(G)|_{y=0} = T(f)$$
 (15.1.3)

Let's verify that the assumptions above imply (15.1.1). If  $f \in S$  and  $T(G) \in S'$  then W(f)T(G) lies in S'. Evaluation at  $\mathbf{y} = 0$  gives an element of S, so  $T(f) \in S$ .

S	S'	G	u	Comments
Р	$\widehat{\mathbf{P}}_2$	e <sup>-xy</sup>	-Dy	
Ρ <sub>d</sub>	$\widehat{\mathbf{P}}_{2d}$	$e^{-x \cdot y}$	$-D_{\mathbf{y}}$	
$\mathcal{H}_1$	$\widehat{\mathcal{H}_2}$	e <sup>xy</sup>	Dy	
$MA_d$	$MA_{2d}$	$\prod_{1}^{d}(x_{i} + y_{i})$	$-D_{\mathbf{y}}$	
P <sup>sep</sup>	?	$(1 + x)^y$	$\Delta$	
Р	$\widehat{\mathbf{P}^{\text{pos}}}$	e <sup>x</sup>	Dy	multiplier transformation
$\mathcal{H}_1$	$\widehat{\mathcal{H}_1}$	e <sup>x</sup>	Dy	multiplier transformation
$\mathbf{P}(\mathbf{n})$	$P_2$	$(x + y)^n$	$-D_y$	bounded degree

Table 15.1: Generating function data

How do we find G and W? All we need is a dimension decreasing transformation  $U^{I}$ . We define W(f) = f(U), construct dual polynomials  $p_{I}(\mathbf{y})$  satisfying

$$\mathbf{U}^{\mathbf{I}}\mathbf{p}_{\mathbf{J}}(\mathbf{y})\big|_{\mathbf{y}=\mathbf{0}} = \begin{cases} \mathbf{0} & \mathbf{I} \neq \mathbf{J} \\ \mathbf{1} & \mathbf{I} = \mathbf{J} \end{cases}$$

and define

$$\mathsf{G}(\textbf{x},\textbf{y}) = \sum_{\textbf{I}} \textbf{x}^{\textbf{I}} \textbf{p}_{\textbf{I}}(\textbf{y}).$$

The fundamental identity is formally trivial, since we only need to check on monomials:

$$|\mathbf{U}^{I} \mathsf{T}(\mathsf{G})|_{\mathbf{y}=0} = |\mathbf{U}^{I} \sum \mathsf{T}(\mathbf{x}^{J}) \mathsf{p}_{J}(\mathbf{y})|_{\mathbf{y}=0} = \sum \mathsf{T}(\mathbf{x}^{J}) |\mathbf{U}^{I} \mathsf{p}_{J}(\mathbf{y})|_{\mathbf{y}=0} = \mathsf{T}(\mathbf{x}^{I}).$$

The only difficult part is the verification of the mapping requirement

$$f \times g \mapsto f(U)g$$
 satisfies  $S \times S' \longrightarrow S'$ . (15.1.4)

In order to use the method of generating functions effectively we must be able to compute T(G), and have some knowledge of the members of S'. This isn't always easy.

Proving a Pólya-Schur result is harder, and often requires some analytic information in order to show that  $T(G) \in S'$ . For instance, if we take T to be the identity it isn't even clear why T(G) = G is in S'.

We now discuss, with simple examples, each of the cases in Table 15.1.

**Example 15.1.** *Transformations*  $P \longrightarrow P$ 

We take S = P,  $S' = \hat{P}_2$ , and U = -D, with dual polynomials  $p_i(y) = (-y)^i/i!$ . The base generating function is

$$e^{-xy} = \sum_0^\infty x^i \frac{(-y)^i}{i!}$$

and is in  $\widehat{\mathbf{P}}_2$ . We need to verify the mapping condition (15.1.4). Let  $f(x) \in \mathbf{P}$  and  $g(x, y) \in \overline{\mathbf{P}}_2$ . We know that  $f(-\mathsf{D})g \in \widehat{\mathbf{P}}_2$ , and therefore we can apply the method of generating functions to prove that transformations preserve all real roots.

**Lemma 15.2.** The Hadamard product  $x^{i} \circledast x^{j} \mapsto \begin{cases} 0 & i \neq j \\ x^{i} & i = j \end{cases}$ determines a map  $P^{pos} \times P \longrightarrow P$ .

*Proof.* Choose  $f \in \mathbf{P}^{pos}$  and define  $T_f(g) = f \circledast g$ . If  $f = x^n$  then it's easy to see that  $T_f(e^{-xy}) = (-xy)^n$ . It follows that in general  $T(e^{-xy}) = f(-xy)$ . Since  $f \in \mathbf{P}^{pos}$  we know that  $f(-xy) \in \overline{\mathbf{P}}_2$ , which is contained in  $\widehat{\mathbf{P}}_2$ .

It is interesting that the base generating function is equivalent to base generating functions for each positive integer.

**Lemma 15.3.** If 
$$T_*(1 - xy)^n \in \mathbf{P}_2$$
 for  $n = 1, 2, ...$  then  $T: \mathbf{P} \longrightarrow \mathbf{P}_2$ 

*Proof.* Since  $f(-D_y)$  preserves  $\overline{P}_2$  we know that  $f(-D_y)T_*(1 - xy/n)^n$  is in  $\overline{P}_2$ . The result follows from the claim that

$$\lim_{n \to \infty} f(-\mathsf{D}_{y}) \mathsf{T}\left(1 - \frac{xy}{n}\right)^{n} \Big|_{y=0} = \mathsf{T}(f)$$

By linearity it suffices to show this for  $f(x) = x^k$  in which case

$$\begin{split} (-\mathsf{D}_{\mathfrak{y}})^{k}\mathsf{T}_{*}\bigg(1-\frac{x\mathfrak{y}}{\mathfrak{n}}\bigg)^{\mathfrak{n}}\bigg|_{\mathfrak{y}=0} &= \mathsf{D}^{k}\sum \binom{\mathfrak{n}}{\mathfrak{i}}\bigg(\frac{\mathfrak{y}}{\mathfrak{n}}\bigg)^{\mathfrak{i}}\mathsf{T}(\mathfrak{x}^{\mathfrak{i}})\bigg|_{\mathfrak{y}=0} \\ &= \binom{\mathfrak{n}}{k}\mathfrak{n}^{-k}\mathsf{T}(\mathfrak{x}^{k}) \longrightarrow \mathsf{T}(\mathfrak{x}^{k}). \end{split}$$

Alternatively, see Corollary 11.89

Of course, the two approaches are equivalent:

**Lemma 15.4.** If  $T_*(e^{-xy}) \in \overline{\mathbf{P}}_2$  then  $T_*(1-xy)^n) \in \overline{\mathbf{P}}_2$  for n = 1, 2, ...Conversely, if  $T(1) \neq 0$  and  $T_*(1-xy)^n \in \overline{\mathbf{P}}_2$  for all n then  $T_*(e^{-xy}) \in \widehat{\mathbf{P}}_2$ .

*Proof.* Since  $1 - yz \in \overline{\mathbf{P}}_2$  the operator  $1 + z\partial_y$  preserves  $\widehat{\mathbf{P}}_3$ . We have

$$(1+z\partial y)^{n} T_{*}(e^{-xy})\Big|_{y=0} = \sum (-1)^{k} z^{k} T(x^{k}) \binom{n}{k} = T_{*}(1-xz)^{n}$$

Since  $(1 - xy/n)^n \rightarrow e^{-xy}$  the second part follows from Lemma 13.31.  $\Box$ 

**Example 15.5.** Transformation  $\overline{\mathbf{P}}_{d} \longrightarrow \overline{\mathbf{P}}_{d}$ 

The argument for  $\overline{\mathbf{P}}_d$  is exactly the same as for **P**. If  $\mathbf{I} = (i_1, \dots, i_d)$  we let

$$U^{I} = \left(\frac{\partial}{\partial y_{1}}\right)^{i_{1}} \cdots \left(\frac{\partial}{\partial y_{d}}\right)^{i}$$

The dual polynomials are

$$p_{I} = \frac{y_{1}^{i_{1}} \cdots y_{d}^{i_{d}} (-1)^{i_{1} + \dots + i_{d}}}{i_{1}! \cdots i_{d}!}$$

For an example, here is a Hadamard product result.

**Lemma 15.6.** Suppose that  $f_1, \ldots, f_d$  are in  $P^{\text{pos}}$ . The linear transformation  $g \mapsto (f_1(x_1) \cdots f_d(x_d)) \circledast g$  defines a map  $\overline{P}_d \longrightarrow \overline{P}_d$ .

*Proof.* The generating function of the map is  $f_1(-x_1y_1) \cdots f_d(-x_dy_d)$  which is in  $\overline{P}_{2d}$  since all  $f_i \in \mathbf{P}^{pos}$ .

**Example 15.7.** *Transformations*  $\mathcal{H}_1 \longrightarrow \mathcal{H}_1$ 

Since  $f \times g \mapsto f(D)g$  maps  $\mathcal{H}_1 \times \mathcal{H}_1 \longrightarrow \mathcal{H}_1$  we know that  $T(e^{xy}) \in \widehat{\mathcal{H}}_2$ implies  $T : \mathcal{H}_1 \longrightarrow \mathcal{H}_1$ .

We start wwith a Hadamard product result.

**Lemma 15.8.** The map  $f \times g \mapsto f * g$  determines a map  $P^{pos} \times \mathcal{H}_1 \longrightarrow \mathcal{H}_1$ .

*Proof.* If we fix  $f \in \mathbf{P}^{pos}$  then the generating function of  $g \mapsto g * f$  is exp f(xy). Since exp  $f \in \mathbf{P}^{pos}$  we know that exp  $f(xy) \in \mathcal{H}_2$ .

*Remark* 15.9. If T does not map  $\mathcal{H}_1 \longrightarrow \mathcal{H}_1$  but rather maps  $\mathcal{H}_1 \longrightarrow \mathcal{H}_1 \cup 0$  then there might not be a Pólya-Schur type result.

Consider the Hadamard product T:  $f \mapsto (x^2 + 1) \circledast f$ . In coordinates this is

$$\mathsf{T}(\sum \mathfrak{a}_{\mathfrak{i}} x^{\mathfrak{i}}) = \mathfrak{a}_0 + \mathfrak{a}_2 x^2.$$

Note that T(x) = 0, and since all  $a_i$  are non-negative,  $T : \mathcal{H}_1 \longrightarrow \mathcal{H}_1 \cup 0$ .

The generating function of T is  $(xy)^2/2+1$  which is *not* in  $\mathcal{H}_2$ , so there is no Pólya-Schur type result. In addition the induced map fails to map  $\mathcal{H}_2 \longrightarrow \mathcal{H}_2$ :

$$\mathsf{T}(x+y)^2 = x^2 + y^2 \not\in \mathcal{H}_2$$

In general, the map  $g \mapsto f \times g$  has generating function  $\exp f(xy)$ , and this is in  $\mathcal{H}_2$  if and only if  $\exp f \in \mathbf{P}^{pos}$ .

**Corollary 15.10.** *The following linear transformations map*  $\mathcal{H}_1$  *to*  $\mathcal{H}_1$ *.* 

Exponential 
$$x^k \mapsto \frac{x^k}{k!} x^k$$
  
Binomial  $x^k \mapsto \binom{n}{k} x^k$   
Laguerre  $x^k \mapsto L_k(-x)$   
Laguerre<sup>REV</sup>  $x^k \mapsto L_k^{REV}(-x)$ 

*Proof.* See Table 15.2.

Next we have an inner product result. If  $f=\sum f_i(x)y^i$  and  $g=\sum g_i(x)y^i$  then we define

$$\langle f,g \rangle = \sum i!f_ig_i$$

This is clearly an inner product. The generating function of the map

$$T_f: g \mapsto \langle f, g \rangle$$

is

$$\sum T_f(x^iy^j)\frac{u^i\nu^j}{i!j!} = \sum \left\langle f, x^iy^j \right\rangle \frac{u^i\nu^j}{i!j!} = \sum j!f_jx^i\frac{u^i\nu^j}{i!j!} = e^{xu}f(x,\nu)$$

Since this is in  $\mathcal{H}_2$  if  $f \in \mathcal{H}_2$  we conclude

**Lemma 15.11.** If  $f \in \mathcal{H}_2$  then the map  $g \mapsto \langle f, g \rangle$  satisfies  $\mathcal{H}_2 \times \mathcal{H}_2 \longrightarrow \mathcal{H}_1$ .

Sums of squares of polynomials in  $P^{pos}$  are not necessarily stable, but the sum of the squares of the coefficients of a polynomial in  $P_2^{pos}$  is stable. Since  $P_2^{pos} \subset \mathcal{H}_2$  we can apply the inner product.

Corollary 15.12. If  $\sum f_i(x)y^i \in P_2^{\text{pos}}$  then  $\sum f_i^2$  is in  $\mathcal{H}_1$ .

For example, using the Taylor series  $\sum \frac{f^{(i)}}{i!} y^i$  yields that if  $f \in \mathbf{P}^{pos}$  then

$$\sum \left(\frac{f^{(\mathfrak{i})}}{\mathfrak{i}!}\right)^2 \in \mathfrak{H}_1$$

**Corollary 15.13.** *If*  $T: P \longrightarrow P$ ,  $T: P^{\text{pos}} \longrightarrow P^{\text{pos}}$ , and T preserves degree then  $\sum \left(\frac{T(x^i)}{i!}\right)^2 \in \widehat{\mathcal{H}_1}$ .

For instance, if  $L_n$  is the Laguerre polynomial then the transformation  $x^n \mapsto L_n(-x)$  satisfies the hypotheses of the corollary, so

$$\sum_{n} \frac{L_{n}(-x)^{2}}{n!n!} \in \widehat{\mathcal{H}_{1}}.$$

#### **Example 15.14.** *Multiaffine polynomials*

The unusual aspect of multiaffine polynomials is that the generating function is a polynomial. There are two choices for U, and they lead to base generating functions that are the reverse of one another. In each case the mapping properties follow from the properties for  $\overline{\mathbf{P}}_d$ .

If we take  $U^{I} = (-1)^{|I|} \hat{D}_{y_{I}}$  then the dual polynomials are  $(-1)^{|I|} y_{I}$  and the generating function is

$$\sum_I x_I (-1)^{|I|} y_I = \prod_1^d (1-x_i y_1)$$

We can also take  $U^I=\mathsf{D}_{y_{\{1,\dots,d\}\setminus I}}$  with dual polynomials  $y_{\{1,\dots,d\}\setminus I}.$  The generating function is

$$\sum_{\mathbf{I}} \mathbf{x}_{\mathbf{I}} \mathbf{y}_{\{1,\dots,d\}\setminus \mathbf{I}} = \prod_{1}^{u} (\mathbf{x}_{i} + \mathbf{y}_{i})$$

**Example 15.15.** Transformations  $P^{sep} \longrightarrow P^{sep}$ 

We do not yet understand this case. We have  $S = \mathbf{P}^{sep}$ ,  $U = \Delta_y$ , with dual polynomials  $(y)_i/i!$ . The base generating function is

$$\sum_{i=0}^{\infty} x^{i} \frac{(\underline{y})_{i}}{i!} = (1+x)^{y}$$

Note that the base generating function is not entire, but is analytic for |x| < 1 and all y. It is not clear what S' is, and we do not know how to prove the mapping property.

#### **Example 15.16.** *Polynomials with separated roots*

We start with the fact that  $\Delta$  maps  $\mathbf{P}^{sep}$  to itself. The dual polynomials are  $(y)_n/n!$ . Thus, the base generating function is

$$\sum x^{i} \frac{(\underline{y})_{i}}{i!} = (1+x)^{y}$$

However, in this case we do not know the correct space of functions S' that contain the base generating function.

#### Example 15.17. Multiplier transformations for P

Let  $T : x^i \mapsto a_i x^i$  be a multiplier transformation. If we treat this a map on polynomials then we know that if  $T(e^{-xy}) \in \widehat{\mathbf{P}}_2$  then  $T : \mathbf{P} \longrightarrow \mathbf{P}$ . However, since T is a multiplier transformation  $T(e^{-xy}) = T(e^x)(-xy)$ . It follows that if  $T(e^x) \in \widehat{\mathbf{P}^{pos}}$  then  $T(e^{-xy}) \in \widehat{\mathbf{P}}_2$ , and hence  $T : \mathbf{P} \longrightarrow \mathbf{P}$ .

## **Example 15.18.** *Multiplier transformations for* $\mathcal{H}_1$

Let  $T : x^i \mapsto a_i x^i$  be a multiplier transformation. As above we know that if  $T(e^{xy}) \in \widehat{\mathcal{H}_2}$  then  $T : \mathcal{H}_1 \longrightarrow \mathcal{H}_1$ . Now  $T(e^{xy}) = T(e^x)(xy)$ , and the only way for f(xy) to belong to  $\mathcal{H}_2$  is if  $f \in \widehat{\mathbf{P}^{pos}}$ .

**Lemma 15.19.** If the multiplier transformation T satisfies  $T(e^x) \in \widehat{P^{pos}}$  then

$$1. \ \mathsf{T}: \boldsymbol{P} \longrightarrow \boldsymbol{P}$$

2. 
$$T : \mathcal{H}_1 \longrightarrow \mathcal{H}_1$$

**Example 15.20.** *Polynomials of bounded degree* 

In this case we have that  $T(G) \in S'$  implies that  $V : S \longrightarrow S$  where V is a transformation depending on T.

We take  $U = -D_u$  and  $G = (x + y)^n$ .

$$\mathbf{U}^{k}\mathsf{T}(\mathsf{G})\big|_{\mathfrak{y}=0} = \sum (-\mathsf{D}_{\mathfrak{y}})^{k}\mathsf{T}(\mathfrak{x}^{n-k})\mathfrak{y}^{k}\binom{\mathfrak{n}}{k}\big|_{\mathfrak{y}=0} = (-1)^{k}\mathsf{T}(\mathfrak{x}^{n-k})(\underline{\mathfrak{n}})_{k}$$

If we define  $V(x^k) = T(x^k)(\underline{n})_k$  then by taking the revese and negating x we see that V:  $P(n) \longrightarrow P(n)$ .

In this case even the identity yields an interesting result:

**Lemma 15.21.** The mapping  $x^k \mapsto x^k(\underline{n})_k$  maps  $P(n) \longrightarrow P(n)$ .

#### **15.2** Multiplier transformations

The main results of this section could be derived as special cases of the previous section, but the proofs are easy. We are concerned with the generating functions of multiplier transformations  $x^i \mapsto a_i x^i$ . If  $T(x^i) = a_i x^i$  then the generating function of T is

$$T_{*}(e^{-xy}) = \sum_{i=0}^{\infty} T(x^{i}) \frac{(-y)^{i}}{i!} = \sum_{i=0}^{\infty} \alpha_{i} \frac{(-xy)^{i}}{i!} = F(-xy)$$
(15.2.1)

where

$$F(\mathbf{x}) = \sum a_{i} \frac{\mathbf{x}^{i}}{i!}$$

We will consider the generating function to be a function  $T(e^{-x}) = F(x)$  of one variable. Our goal is to prove that (A)  $\implies$  (C) for multiplier transformations.

Theorem 15.22 (Pólya-Schur).

- 1.  $\mathbf{P}^{\text{pos}}$  is precisely the set of generating functions of linear transformations  $x^i \mapsto a_i x^i$  that map  $\mathbf{P}^{\text{pos}}$  to  $\mathbf{P}^{\text{pos}}$ .
- 2.  $\widehat{\mathbf{P}}$  is precisely the set of generating functions of linear transformations  $x^i \mapsto a_i x^i$  that map  $\mathbf{P}^{\text{pos}}$  to  $\mathbf{P}$ .

*Proof.* If  $T: \mathbf{P}^{pos} \longrightarrow \mathbf{P}^{pos}$  then

$$T(e^{x}) = \lim_{n \to \infty} T\left(1 + \frac{x}{n}\right)^{n}.$$

Sine  $(1 + x/n)^n \in \mathbf{P}^{pos}$  we conclude from Lemma 4.16 that the polynomials in the limit are in  $\mathbf{P}^{pos}$  by assumption on T. Thus, the generating function is in  $\widehat{\mathbf{P}^{pos}}$  by Lemma 13.31. A similar observation shows that the generating function in the second case is in  $\widehat{\mathbf{P}}$ .

Conversely, assume that the generating function f is in  $\widehat{\mathbf{P}^{\text{pos}}}$  and choose g(x) in  $\mathbf{P}^{\text{pos}}$ . By Theorem 9.87 the Hadamard-type product \*' satisfies

$$\mathsf{T}(\mathfrak{g}) = \mathsf{F} \ast' \mathfrak{g} \in \mathbf{P} \tag{15.2.2}$$

and therefore  $T(g) \in \mathbf{P}^{pos}$  since all coefficients are positive. In the second case  $f \in \widehat{\mathbf{P}}$  and  $g \in \mathbf{P}^{pos}$ , so we only get that  $T(g) \in \mathbf{P}$ .

*Remark* 15.23. This result is known as the Pólya-Schur Theorem [137]. Part (1) is sometimes stated in terms of maps  $\mathbf{P} \longrightarrow \mathbf{P}$ , but this can be deduced from the following lemma.

**Lemma 15.24.** If  $T: P^{\text{pos}} \longrightarrow P^{\text{pos}}$  is a multiplier transformation, then T extends to a linear transformation  $P \longrightarrow P$ .

*Proof.* Since the generating function F of T is in  $\widehat{\mathbf{P}^{\text{pos}}}$ , the result follows from (15.2.2) and Theorem 9.87.

Here's another consequence that can be found in [137].

**Lemma 15.25.** If  $f \in \widehat{P}$  has all positive coefficients then  $f \in \widehat{P^{pos}}$ .

*Proof.* If  $f \in \widehat{\mathbf{P}}$  then consider the linear transformation given by (15.2.2). If  $g \in \mathbf{P}^{\text{pos}}$  then  $\mathsf{T}g \in \mathbf{P}^{\text{pos}}$  since f has all positive coefficients. Thus,  $f \in \widehat{\mathbf{P}^{\text{pos}}}$ .

$$(B) \Longrightarrow (C)$$

**Lemma 15.26.** If  $T(x^i) = a_i x^i$  and  $T((1 + x)^n) \in P$  for infinitely many n then  $T(e^x) \in \widehat{P}$ .

*Proof.* If we substitute x/n for x then we see that

$$\lim_{n \to \infty} \mathsf{T}\left(1 + \frac{x}{n}\right)^n = \mathsf{T}(e^x) \in \widehat{\mathbf{P}}$$

By Theorem 15.22 we are done.

If T is as in the last lemma, then since  $(x + 1)^n \leq (x + 1)^{n-1}$ , we know that  $T(x + 1)^n \leq T(x + 1)^{n-1}$ . In [42] it is shown that the interlacing is strict.

## **15.3** Linear transformations $P \longrightarrow P$

Table 15.2 lists some examples of generating functions of linear transformations that map **P** to **P**. The first four are elementary, and the rest are standard formulas (e.g. [72]). In the Hadamard product and f(xD) we must choose  $f \in \mathbf{P}^{\pm}$ . In the table  $I_0(z)$  is the modified Bessel function of the first kind and  $J_0(z)$  is the Bessel function. In the *Multiplier* entry we are given a linear transformation of the form  $T(x^i) = a_i x^i$  which maps **P** to itself. The generating function of T in the sense of the previous section is  $f(x) = \sum a^i \frac{x^i}{i!}$  and is in  $\widehat{\mathbf{P}}$ .

**Example 15.27.** If T:  $\mathbf{P} \longrightarrow \mathbf{P}$  doesn't preserve degree and the sign of the leading coefficient then the generating function might not be in  $\widehat{\mathbf{P}}_2$ . Consider  $T(g) = G(\mathsf{D})f(x)$ . We have seen that this linear transformation doesn't extend to a map  $\overline{\mathbf{P}}_2 \longrightarrow \overline{\mathbf{P}}_2$  when f(x) = x. An easy calculation shows that the generating function is  $(x - y)e^{-xy}$ , which isn't in  $\widehat{\mathbf{P}}_2$ .

Name					$T_*(e^{-xy})$
Affine	$\sum_{i=0}^{\infty}$	$(ax+b)^i$	$\frac{(-y)^{i}}{i!}$	=	$e^{-(ax+b)y}$
Derivative	$\sum_{i=0}^{t=0}$	ix <sup>i-1</sup>	$\frac{(-y)^{i}}{i!}$	=	-ye <sup>-xy</sup>
Derivative	$\sum_{i=0}^{\infty}$	$f(D)x^{\mathfrak{i}}$	$\frac{(-y)^{i}}{i!}$	=	$f(-y)e^{-xy}$
Derivative	$\sum_{i=0}^{\infty}$	$f(xD)x^{i}$	$\frac{(-y)^{i}}{i!}$	=	$f(-xy)e^{-xy}$
Exponential	$\sum_{i=0}^{i=0}$	$\frac{x^{i}}{i!}$	$\frac{(-y)^{i}}{i!}$	=	$J_0(2\sqrt{-xy})$
Hadamard	$\sum_{i=0}^{\infty}$	$f(x) * x^i$	$\frac{(-y)^{i}}{i!}$	=	(EXPf)(-xy)
Hadamard'	$\sum_{i=0}^{\infty}$	$[x^r * x^r = r!x^r] * x^i$	$\frac{(-y)^{i}}{i!}$	=	f(-xy)
Hermite	$\sum_{i=0}^{i=0}$	H <sub>i</sub>	$\frac{(-y)^{i}}{i!}$	=	$e^{-2xy-y^2}$
Laguerre	$\sum_{i=0}^{t=0}$	$L_i(-\mathbf{x})$	$\frac{(-y)^{i}}{i!}$	=	$e^{-y}J_0(2\sqrt{xy})$
Laguerre REV	$\sum_{i=0}^{i=0}$	$L_{i}^{\text{ Rev}}(-x)$	$\frac{(-y)^{i}}{i!}$	=	$e^{-xy}J_0(2\sqrt{y})$
Laguerre	$\sum_{i=0}^{i=0}$	Lin	$\frac{(-y)^{i}}{i!}$	=	$e^{-y}L_n(x+y)$
Multiplier	$\sum_{i=0}^{\iota=0}$	$a_i x^i$	$\frac{(-y)^{i}}{i!}$	=	f(-xy)

Table 15.2: Generating Functions for transformations  $P \longrightarrow P$ 

*Remark* 15.28. The generating function of the linear transformation  $f \mapsto f + \alpha f'$  is  $(1 + \alpha y)e^{-xy}$ . Since this is in  $\hat{\mathbf{P}}_2$  it follows that  $f + \alpha f'$  is in  $\mathbf{P}$  for all  $f \in \mathbf{P}$ . This is another proof of Rolle's theorem (Theorem 1.1).

**Example 15.29.** What happens if T doesn't map **P** to itself? The generating functions should be functions that are not in  $\hat{\mathbf{P}}_2$ . Let's look at some examples. The Hermite transformation  $T(x^i) = H_i$  maps **P** to itself. Since this is not an affine map we know that the inverse does not map **P** to itself. Using identities in [146] we can find the generating function of  $T^{-1}$ :

$$T^{-1}(x^{n}) = \left(\frac{i}{2}\right)^{n} H_{n}\left(\frac{-ix}{2}\right)$$
$$T_{*}(e^{-xy}) = \sum_{n=0}^{\infty} \left(\frac{i}{2}\right)^{n} H_{n}\left(\frac{-ix}{2}\right) \frac{(-y)^{n}}{n!}$$
$$= \sum_{n=0}^{\infty} H_{n}\left(\frac{-ix}{2}\right) \frac{\left(\frac{(-iy)}{2}\right)^{n}}{n!}$$
$$= e^{\frac{-2xy+y^{2}}{4}}$$

This is not in  $\widehat{\mathbf{P}}_2$  since substituting x = 0 gives  $e^{y^2/4}$  which is not in  $\widehat{\mathbf{P}}$ .

Next, consider  $T(x^i) = \langle \underline{x} \rangle_i$  which maps  $P^{\text{pos}}$  to itself. We can compute

$$T_*(e^{-xy}) = \sum_{i=0}^{\infty} T(x^i) \frac{(-y)^i}{i!} = \sum_{i=0}^{\infty} x(x+1) \cdots (x+i-1) \frac{(-y)^i}{i!} = (1+y)^{-x}$$

Although the generating function has a simple form it isn't in  $\hat{\mathbf{P}}_2$ . This is because it is not an entire function:  $T_*(e^{-xy})(1,y) = (1+y)^{-1}$  has a pole at y = -1.

An example of a different sort is given by the generating function of the q-derivative. Recall if  $\mathbb{A}(x) = qx$  then the q-derivative is  $\frac{f(qx) - f(x)}{x(q-1)}$ . The generating function is

$$\frac{e^{-qxy}-e^{-xy}}{x(q-1)}$$

which is just the q-derivative of  $e^{-xy}$ . It's not in  $\widehat{\mathbf{P}}$  since substituting y = 1 gives a function that has complex roots.

**Example 15.30.** If  $T(x^i) = H_i$ , then using the generating function we can show that  $T(\mathbf{P}) \not\subset EXP(\mathbf{P})$  without exhibiting any particular polynomial in  $EXP(\mathbf{P})$  that is not in  $T(\mathbf{P})$ . If  $T(\mathbf{P}) \subset EXP(\mathbf{P})$  then  $EXP^{-1}T(\mathbf{P}) \subset \mathbf{P}$ . We will show that

the generating function of  $EXP^{-1}T$  isn't in  $\widehat{\mathbf{P}}_2$ . We compute

$$\sum_{i} EXP^{-1} T(x^{i}) \frac{(-y)^{i}}{i!} = EXP_{*}^{-1} \left( \sum_{i} T(x^{i}) \frac{(-y)^{i}}{i!} \right)$$
$$= EXP_{*}^{-1} e^{-2xy-y^{2}}$$
$$= e^{-y^{2}} \sum_{i} i! \frac{(-2xy)^{i}}{i!}$$
$$= \frac{e^{-y^{2}}}{1+2xy}$$

This last function is not entire.

# **15.4** Linear transformations $P \longrightarrow P$ - general properties

In this section we establish general properties of generating functions. In the next section we will apply some of the results to show that various linear transformations preserve  $\mathbf{P}$ .

If a linear transformation T has generating function F(x, y), then it is easy to compute the generating function of the Möbius transformation  $T_M$  (§ 6.6.7), where  $M : z \mapsto \frac{az+b}{cz+d}$ . To begin with, there is a very useful relationship between the generating functions of a linear transformation T and the transformation  $T_{1/z}$ . Recall (§ 6.6.7) that if  $T(x^n) = p_n(x)$  then  $T_{1/z}(x^n) = p_n^{\text{REV}}$ . The generating function of  $T_{1/z}$  is

$$\sum_{n=0}^{\infty} T_{1/z}(x^n) \frac{(-y)^n}{n!} = \sum_{n=0}^{\infty} x^n p_n (1/x) \frac{(-y)^n}{n!} = \sum_{n=0}^{\infty} p_n (1/x) \frac{(-xy)^n}{n!} = F(1/x, xy)$$

More generally, if M is a Möbius transformation  $M(z) = \frac{az+b}{cz+d}$  then we define  $T_M(x^n) = (cx + d)^n T(x^n) \left(\frac{ax+b}{cx+d}\right)$ , and the generating function is

$$\sum_{i} (cx+d)^{i} T\left(\frac{ax+b}{cx+d}\right) \frac{(-y)^{i}}{i!} = F\left(\frac{ax+b}{cx+d}, (cx+d)y\right)$$

If we first compose with an affine transformation then the generating function has an exponential factor.

**Lemma 15.31.** Suppose that T is a linear transformation with generating function F(x, y). If S(f) = T(f(ax + b)) then the generating function of S is  $e^{-by}F(x, ay)$ .
*Proof.* A computation:

$$\sum_{n} S(x^{n}) \frac{(-y)^{n}}{n!} = \sum_{n} T((ax+b)^{n}) \frac{(-y)^{n}}{n!}$$
$$= \sum_{n} \sum_{k} a^{k} T(x^{k}) b^{n-k} {n \choose k} \frac{(-y)^{n}}{n!}$$
$$= \sum_{k} a^{k} T(x^{k}) \frac{(-y)^{k}}{k!} \sum_{n \ge k} \frac{(by)^{n-k}}{(n-k)!}$$
$$= \sum_{k} T(x^{k}) \frac{(-ay)^{k}}{k!} e^{-by}$$
$$= e^{-by} F(x, ay)$$

The next result shows that the coefficients of the polynomials defining a root preserving transformation are highly constrained.

**Lemma 15.32.** Suppose that  $T: \mathbf{P} \longrightarrow \mathbf{P}$ . Choose a non-negative integer r and let  $d_i$  be the coefficient of  $x^r$  in  $T(x^i)$ . Then, the series  $\sum_{i=0}^{\infty} d_i \frac{x^i}{i!}$  is in  $\widehat{\mathbf{P}}$ , and  $x^i \mapsto d_i x^i$  maps  $\mathbf{P}^{pos} \longrightarrow \mathbf{P}$ .

In addition, for any n the polynomial  $\sum {\binom{n}{i}} d_i x^i$  is in **P**.

*Proof.* The first sum is the coefficient of  $x^r$  in the generating function of T. The second follows from Lemma 13.18.

In particular, the generating function of the constant terms of a root preserving linear transformation is in  $\hat{\mathbf{P}}$ .

The leading coefficients are similarly constrained.

**Lemma 15.33.** Suppose  $T: \mathbf{P} \longrightarrow \mathbf{P}, T: \mathbf{P}^{pos} \longrightarrow \mathbf{P}^{pos}$ , T preserves degree, and the leading coefficient of  $T(x^i)$  is  $c_i$ . Then  $\sum c_i \frac{x^i}{i!} \in \widehat{\mathbf{P}}$ , and  $x^i \mapsto c_i x^i$  maps  $\mathbf{P}^{pos}$  to itself.

*Proof.* Corollary 9.57 shows that  $T_{1/z} : \mathbf{P} \longrightarrow \mathbf{P}$ . The constant term of  $T_{1/z}(\mathbf{x}^i)$  is  $c_i$ , so we can apply the previous lemma.

We have seen (Corollary 1.50) that if  $T: x^n \mapsto f_n(x)$  maps  $\mathbf{P} \longrightarrow \mathbf{P}$ , then the map  $x^n \mapsto c_n x^n$ , where  $c_n$  is the leading coefficient of  $f_n$ , also maps  $\mathbf{P}$  to itself. Evaluation has similar properties.

**Corollary 15.34.** Suppose that  $T: x^n \mapsto f_n(x)$  preserves degree and maps  $P \longrightarrow P$ , then for any  $\alpha \in \mathbb{R}$  the multiplier transformation  $x^n \mapsto f_n(\alpha)x^n$  maps  $P^{pos} \longrightarrow P$ .

*Proof.* If the generating function  $T_*(e^{-xy} \in \widehat{\mathbf{P}}_2$ , then evaluation is in  $\widehat{\mathbf{P}}$ , and so

$$\sum f_n(\alpha) \frac{(-y)^n}{n!} \in \widehat{P}$$

from which the conclusion follows.

If we have two linear transformations S, T with generating functions in  $\widehat{\mathbf{P}}_2$  then we can multiply their generating functions by  $\sum a_i y^{n-i} z^{n-i}$ :

$$\left(\sum_{i=0}^{\infty} S(x^{i}) \frac{(-y)^{i}}{i!}\right) \left(\sum_{i=0}^{\infty} T(x^{i}) \frac{(-z)^{i}}{i!}\right) \left(\sum_{i=0}^{n} a_{i} y^{n-i} z^{n-i}\right)$$

If  $\sum_{0}^{n} a_{i} x^{i} \in \mathbf{P}^{pos}$  then the third factor is in  $\widehat{\mathbf{P}^{pos}}_{2}$  so all the factors are in  $\widehat{\mathbf{P}}_{3}$ . The coefficient of  $y^{n} z^{n}$  is  $\sum_{i \in S(x^{i})T(x^{i}) \atop i!i!} \frac{a_{i}S(x^{i})T(x^{i})}{i!i!}$ . Consequently, the linear transformation  $x^{n} \mapsto \sum_{i} a_{i} \frac{S(x^{i})T(x^{i})}{i!i!}$  maps  $\mathbf{P}^{pos}$  to  $\mathbf{P}$ . We can remove the factorials.

**Lemma 15.35.** Suppose S, T are linear transformations with generating functions in  $\widehat{P}_2$ . The map  $x^i \mapsto (-1)^i S(x^i) T(x^i)$  maps  $P^{pos} \longrightarrow P$ .

*Proof.* If  $\sum a_i x^i \in \mathbf{P}^{alt}$  then we can apply Lemma 10.63. Simply observe (as above) that since

$$\begin{split} \left(\sum_{i=0}^{n} \alpha_{i}(-1)^{i} \partial_{y}^{i} \partial_{z}^{i}\right) \left(\sum_{i=0}^{\infty} S(x^{i}) \frac{(-y)^{i}}{i!}\right) \left(\sum_{i=0}^{\infty} T(x^{i}) \frac{(-z)^{i}}{i!}\right) \Big|_{y=z=0} \\ &= \sum_{i=0}^{n} \alpha_{i}(-1)^{i} S(x^{i}) T(x^{i}) \end{split}$$

and the differential operator maps  $\widehat{\mathbf{P}}_3$  to itself so the right hand side is in **P**.

We can use the relationship between linear transformations  $\mathbf{P} \longrightarrow \mathbf{P}$  and elements of  $\hat{\mathbf{P}}_2$  to construct new linear transformations from old. We start with a linear transformation, determine the corresponding element of  $\hat{\mathbf{P}}_2$ , manipulate it to create a new element of  $\hat{\mathbf{P}}_2$ , and then convert back to a linear transformation. Unfortunately, we don't know many ways to get new elements of  $\hat{\mathbf{P}}_2$  from old ones. Differentiation in  $\hat{\mathbf{P}}_2$  yields some simple transformations. Suppose that

$$\begin{split} f(x,y) &= \sum \mathsf{T}(x_i) \frac{(-y)^i}{i!} \\ &\frac{\partial f}{\partial x} = \sum \frac{d}{dx} \mathsf{T}(x^i) \frac{(-y)^i}{i!} \\ &\text{ and so } \frac{\partial}{\partial x} \text{ corresponds to } g \mapsto \frac{d}{dx}(\mathsf{T}g) \\ &\frac{\partial f}{\partial y} = -\sum \mathsf{T}(x^i) \frac{(-y)^{i-1}}{(i-1)} \\ &\text{ and } \frac{\partial}{\partial y} \text{ corresponds to } g \mapsto -\mathsf{T}(xg) \end{split}$$

Multiplication in  $\widehat{\mathbf{P}}_2$  gives a convolution of linear transformations.

**Lemma 15.36.** Let S, T:  $P \longrightarrow P$  preserve degree and the sign of the leading coefficient. Then the linear transformation below also maps  $P \longrightarrow P$ .

$$x^{n} \mapsto \sum_{i=0}^{n} {n \choose i} T(x^{i}) S(x^{n-i})$$

*Proof.* Multiplying two elements in  $\hat{\mathbf{P}}_2$  yields an element of  $\hat{\mathbf{P}}_2$ :

$$\left(\sum \mathsf{T}(x^i)\frac{(-y)^i}{i!}\right)\left(\sum \mathsf{S}(x^j)\frac{(-y)^j}{j!}\right) = \sum_{n=0}^{\infty} \left(\sum_{i=0}^n \binom{n}{i} \mathsf{T}(x^i)\mathsf{S}(x^{n-i})\right)\frac{(-y)^n}{n!}$$

which establishes the lemma.

*Remark* 15.37. This really is a property of linear transformations that satisfy induction. Indeed, if S, T satisfy induction, then since  $W(f) = (S_*T_*f(x + y))(x, x)$  we see that  $W(f) \in \mathbf{P}$ . The example  $S(g) = g(\mathsf{D})x$ , T(f) = f yields  $W(1) = 1 + x^2$ . This is not a counterexample, since S does not satisfy induction.

*Remark* 15.38. We revisit the characterization results of § 6.6.1. If T is a degree preserving linear transformation such that Tf and f interlace for all  $f \in \mathbf{P}$ , then the linear transformation  $T_{\alpha}(f) = T(f) + \alpha f$  maps  $\mathbf{P}$  to itself. Thus, for all  $\alpha$ , the generating function of  $T_{\alpha}$  is in  $\widehat{\mathbf{P}}_2$ . If G is the generating function of T, then the generating function of  $T_{\alpha}$  is  $G + \alpha e^{-xy}$ . Consequently, G and  $e^{-xy}$  interlace.

We know that such linear transformations are given by Tf = axf + bf' + cf where a and b have the same sign. The generating function of this transformation is  $(ax + by + c)e^{-xy}$ . This makes it clear why we must have the sign condition. See Question 168.

Since we do not have a characterization of functions in  $\hat{\mathbf{P}}_2$ , this point of view does not give a *proof* of the characterization theorems, just an understanding of them.

**Lemma 15.39.** If T is an onto linear transformation defined on polynomials in one variable and the linear transformation S:  $T(x^i)y^i \mapsto T(x^j)y^i$  maps  $P_2$  to itself, then T is an affine transformation.

*Proof.* Since  $S(T(x^i)y^j) = T(x^j)y^i$  by linearity we see  $S(T(f(x))y^j) = T(x^j)f(y)$  for any polynomial f. Choosing  $f = T^{-1}(x^i)$  shows that  $S(x^iy^j) = T(x^j)T^{-1}(y^i)$ . The compositions

$$\mathbf{P} \longrightarrow \overline{\mathbf{P}}_2 \longrightarrow \overline{\mathbf{P}}_2 \longrightarrow \mathbf{P}$$

$$f(x) \xrightarrow{x \mapsto x} f(x) \xrightarrow{y=1} T(f)$$

$$f(x) \stackrel{x \mapsto y}{\longmapsto} f(y) \longmapsto T^{-1}(f) \stackrel{x=1}{\longmapsto} T^{-1}(f)$$

show that both T and  $T^{-1}$  map **P** to itself, and consequently T is affine.

# **15.5** Applications: Linear transformations $P \longrightarrow P$

We now have a powerful tool for showing that a linear transformation maps **P** to itself. We use the generating function to show that many linear transformations preserve roots.

**Example 15.40.** If we start with an element of  $\widehat{\mathbf{P}}_2$  then we get a linear transformation. For instance, if  $f \in \mathbf{P}$ , then  $f(x + y) \in \mathbf{P}_2 \subset \widehat{\mathbf{P}}_2$ . The Taylor series of f is

$$f(x+y) = \sum f^{(i)}(x) \frac{y^{i}}{i!} = \sum (-1)^{i} f^{(i)}(x) \frac{(-y)^{i}}{i!}$$

and consequently the linear transformation  $x^i \mapsto (-1)^i f^{(i)}$  is a map  $P \longrightarrow P$ . Precomposing with  $x \mapsto -x$  shows that  $x^i \mapsto f^{(i)}$  maps  $P \longrightarrow P$ .

In Corollary 7.47 we showed that the Laguerre transformation maps  $\mathbf{P}^{\text{alt}}$  to itself. Now we show that it actually maps **P** to itself.

**Lemma 15.41.** The mapping  $x^n \mapsto L_n(-x)$  (the n-th Laguerre polynomial) maps P to itself.

*Proof.* It suffices to know that the generating function of the transformation is in  $\hat{\mathbf{P}}_2$ . From Table 15.2 the generating function is  $J_0(2\sqrt{xy})$  where the Bessel function  $J_0(z)$  is given by the series

$$J_0(z) = \sum_{k=0}^{\infty} \frac{(-1)^k}{2^{2k} k! k!} z^{2k}$$
$$J_0(2\sqrt{xy}) = \sum_{k=0}^{\infty} \frac{(-1)^k}{k! k!} x^k y^k$$

The Bessel function has a product formula

$$J_0(z) = \prod_{k=1}^{\infty} \left(1 - \frac{z^2}{r_i^2}\right)$$

and hence

$$J_0(2\sqrt{xy}) = \prod_{k=1}^\infty \left(1 - \frac{2xy}{r_i^2}\right)$$

Since the factors of this last product are all in  $\overline{\mathbf{P}}_2$ , it follows that  $J_0(2\sqrt{xy})$  is in  $\widehat{\mathbf{P}^{\text{pos}}}_2$ , and hence the generating function given of  $x^n \mapsto L_n(-x)$  is in  $\widehat{\mathbf{P}^{\text{pos}}}_2$ .

Since the linear transformation T:  $x^n \mapsto L_n$  maps **P** to **P**, we know that  $T^{-1}$  does not map **P** to itself. It's easy to verify that

$$\mathsf{T}^{-1} = \mathsf{E}\mathsf{X}\mathsf{P}^{-1} \circ \mathsf{T} \circ \mathsf{E}\mathsf{X}\mathsf{P}^{-1}.$$

Since  $EXP^{-1}$  doesn't map **P** to itself, it isn't surprising that  $T^{-1}$  doesn't either.

**Lemma 15.42.** If n is a positive integer, then the linear transformation  $x^i \mapsto L_n^i(-x)$  maps P to itself.

*Proof.* From Table 15.2 the generating function F(x, y) of the transformation is  $e^{y}L_{n}(x + y)$ . Since  $F(x, y) \in \widehat{\mathbf{P}}_{2}$ , the linear transformation maps **P** to itself.  $\Box$ 

We can apply Möbius transformations to show that the reverse of the Laguerre transformation also maps **P** to itself.

**Lemma 15.43.** The linear transformation  $S(x^n) = L_n^{REV}(-x)$  maps P to P.

*Proof.* The generating function of  $T(x^n) = L_n(-x)$  is  $e^y J_0(2\sqrt{xy})$ . Since  $S = T_{1/z}$  we know the generating function G(x, y) of S is  $F(1/x, xy) = e^{-xy} J_0(2\sqrt{y})$ . Now  $e^{-xy} \in \widehat{\mathbf{P}}_2$ , and  $J_0(2\sqrt{y}) \in \widehat{\mathbf{P}}$ , and hence  $G(x, y) \in \widehat{\mathbf{P}}_2$ .

The next result is an immediate consequence of Lemma 15.35.

**Corollary 15.44.** If  $H_n$  is the Hermite polynomial, and  $L_n$  is the Laguerre polynomial, then the following linear transformations map  $P^{pos}$  to P.

- $x^n \mapsto H_n(x)^2$
- $x^n \mapsto L_n(-x)^2$
- $x^n \mapsto H_n(x)L_n(-x)$

Lemma 6.28 characterizes the linear transformations of the form  $T(x_n) = \prod_{i=1}^n (x + a_i)$  that map **P** to itself. We conclude

Lemma 15.45. The only choice of constants ai satisfying

$$\sum_{n=0}^{\infty} \left( \prod_{i=0}^{n} (x + a_i) \frac{y^n}{n!} \right) \in \widehat{P}_2$$

*is*  $a_1 = a_2 = a_3 \cdots$ .

We can use generating functions to get results converting complex roots by applying appropriate affine transformations. See [26].

Lemma 15.46. Consider the linear transformations

$$T: f(x) \mapsto x f(x+1) + (x-1) f(x-1)$$
$$S: f(x) \mapsto (\iota x + \frac{1}{2}) f(x-\iota) + (\iota x - \frac{1}{2}) f(x+\iota)$$

*The following diagram commutes* 



*Proof.* It is easy to verify that the diagram commutes at the element level, so it suffices to show that S maps **P** to itself. We compute the generating function of S:

$$\sum_{n=0}^{\infty} \left( (\imath x + \frac{1}{2})(x - \imath)^n + (\imath x - \frac{1}{2})(x + \imath)^n \right) \frac{(-y)^n}{n!}$$
  
=  $\frac{1}{2} \left( -e^{\imath (-1 + \imath x)y} + e^{\imath (1 + \imath x)y} + (2\imath) e^{\imath (-1 + \imath x)y} x + (2\imath) e^{\imath (1 + \imath x)y} x \right)$   
=  $\imath (2x \cos(y) + \sin(y)) e^{-xy}$ 

Now Example 13.5 shows that  $2x \cos(y) + \sin(y) \in \widehat{\mathbf{P}}_2$ , so all factors are in  $\widehat{\mathbf{P}}_2$ , and hence  $\widehat{S}$  maps  $\mathbf{P}$  to itself.

We can determine more transformations that are sums of shifted arguments using Lemma 5.10.

Lemma 15.47. If  $f(x) = \sum a_i x^i$  is in  $P^{(-1,1)}$  and we define  $T(g) = \sum_k a_k \left(g(x + k\iota) + g(x - k\iota)\right)$ 

then T maps **P** to itself.

*Proof.* It suffices to compute the generating function of T. Notice that the generating function of the transformation  $x^n \mapsto (x + \alpha)^n$  is  $e^{-xy - \alpha y}$ . Consequently, the generating function of T is

$$2e^{-xy} \sum_{k} a_k \left(\frac{e^{-ky\iota} + e^{ky\iota}}{2}\right) = 2e^{-xy} \sum_{k} a_k \cos(ky)$$

Since  $f(x) \in \mathbf{P}^{\Delta}$  we know that  $\sum a_k \cos(ky) \in \widehat{\mathbf{P}}$  - see the proof of Lemma 20.76. Thus the generating function is in  $\widehat{\mathbf{P}}_2$ .

For example, we can take  $f = (x+1)^n$ . The lemma shows that if  $g \in \mathbf{P}$  then

$$\sum_{k=0}^{n} {n \choose k} \left( g(x+k\iota) + g(x-k\iota) \right) \in \mathbf{P}.$$

Now we have properties of the Legendre and Jacobi polynomials.

**Lemma 15.48.** The linear transformation  $x^k \mapsto P_k(x)/k!$  defines a map  $P \longrightarrow P$ .

*Proof.* Since  $J_0(x) \in \widehat{\mathbf{P}^{pos}}$ , it follows from (11.1.1) that  $f_0(xy+y)f_0(xy-y) \in \mathbb{P}_2$ . Consequently the generating function for the transformation is in  $\widehat{\mathbf{P}}_2$ .

We can generalize this to Jacobi polynomials. We use the *Mathematica* definition of  $P_n^{\lambda,\mu}(x)$ .

**Lemma 15.49.** For  $\lambda, \mu > -1$  the linear transformation

$$x^{n} \mapsto \frac{n!}{\langle \underline{\lambda+1} \rangle_{n} \langle \underline{\mu+1} \rangle_{n}} P_{n}^{\lambda,\mu}(x)$$

*defines a map*  $P \longrightarrow P$ *.* 

Proof. From [89, (2.1.A)] we let

$$\mathsf{f}_{\lambda}(z) = \sum_{k=0}^{\infty} \frac{1}{k!} \frac{\Gamma(\lambda+1)}{\Gamma(\lambda+k+1)} z^{k}$$

then we have that

$$f_{\lambda}(xz-z)f_{\mu}(xz+z) = \sum_{n=0}^{\infty} \frac{2^{n}}{\langle \underline{\lambda+1} \rangle_{n} \langle \underline{\mu+1} \rangle_{n}} P_{n}^{\lambda,\mu}(x) z^{n}$$

Now  $f_{\lambda}$  is a modified Bessel function and is in  $\widehat{\mathbf{P}^{pos}}$ , so we follow the argument of the previous lemma.

#### 15.6 Applications: the diamond product

We can show that a bilinear mapping T satisfies T:  $\mathbf{P}^{I} \times \mathbf{P} \longrightarrow \mathbf{P}$  by showing that for every  $f \in \mathbf{P}^{I}$  the linear transformation  $T_{f}(g) = T(f, g)$  maps **P** to itself.

We now show that the diamond product based on the Hermite polynomials maps  $P \times P \longrightarrow P$ . The proof shows that the generating function of the transformation is in  $\widehat{P^{\mathfrak{alt}}}_2$  by using some special function identities for Hermite polynomials.

Recall (§ 7.11.14) that if T is a linear transformation then the diamond product is given by  $f \bigotimes_{T} g = T^{-1}(T(f) T(g))$ .

**Lemma 15.50.** Suppose  $T(H_n) = x^n$  where  $H_n$  is the Hermite polynomial. The diamond product  $\diamondsuit_T$  defines a mapping  $P \times P \longrightarrow P$ .

*Proof.* We fix  $f \in \mathbf{P}$ , and show that the generating function of  $g \mapsto f \bigotimes_{T} g$  is in  $\widehat{\mathbf{P}}_2$ . We recall two identities that can be found in [146]

$$T(x^{n}) = \left(\frac{i}{2}\right)^{n} H_{n}\left(\frac{-ix}{2}\right)$$
(15.6.1)  
$$H_{n}(x) H_{k}(x) = \sum_{j=0}^{\min(n,k)} 2^{j} \binom{n}{j} \binom{k}{j} j! H_{n+k-2j}(x)$$

We simplify matters by initially setting  $f = x^n$ . The generating function of  $g \mapsto x^n \mathop{\Diamond}_T g$  is

$$\begin{split} \mathsf{F}(\mathbf{x},\mathbf{y}) &= \sum_{k=0}^{\infty} \mathsf{T}^{-1}(\mathsf{T}(\mathbf{x}^{n})\,\mathsf{T}(\mathbf{x}^{k}))\,\frac{(-\mathbf{y})^{k}}{k!} \\ &= \sum_{k=0}^{\infty} \mathsf{T}^{-1}\left(\left(\frac{\mathbf{i}}{2}\right)^{n}\,\mathsf{H}_{n}\left(\frac{-\mathbf{i}\mathbf{x}}{2}\right)\,\left(\frac{\mathbf{i}}{2}\right)^{k}\,\mathsf{H}_{k}\left(\frac{-\mathbf{i}\mathbf{x}}{2}\right)\right)\,\frac{(-\mathbf{y})^{k}}{k!} \\ &= \sum_{k=0}^{\infty} \mathsf{T}^{-1}\left((\mathsf{H}_{n}\,\mathsf{H}_{k})\left(\frac{-\mathbf{i}\mathbf{x}}{2}\right)\right)\,\left(\frac{\mathbf{i}}{2}\right)^{n+k}\,\frac{(-\mathbf{y})^{k}}{k!} \\ &= \sum_{k=0}^{\infty} \mathsf{T}^{-1}\left(\sum_{j=0}^{\min(n,k)} 2^{j}\binom{n}{j}\binom{k}{j}j!\,\mathsf{H}_{n+k-2j}\left(\frac{-\mathbf{i}\mathbf{x}}{2}\right)\right)\,\left(\frac{\mathbf{i}}{2}\right)^{n+k}\,\frac{(-\mathbf{y})^{k}}{k!} \end{split}$$

Applying the identity (15.6.1) yields

$$\begin{aligned} H_{n+k-2j}\left(\frac{-ix}{2}\right) &= \left(\frac{i}{2}\right)^{-n-k+2j} \mathsf{T}(x^{n+k-2j}) \quad \text{and so} \\ \mathsf{F}(x,y) &= \sum_{k=0}^{\infty} \sum_{j=0}^{\min(n,k)} 2^{j} \binom{n}{j} \binom{k}{j} j! \left(\frac{i}{2}\right)^{-n-k+2j} x^{n+k-2j} \left(\frac{i}{2}\right)^{n+k} \frac{(-y)^{k}}{k!} \\ &= e^{-xy} \left(\frac{2x+y}{2}\right)^{n} \end{aligned}$$

By linearity, we conclude that the generating function G(x,y) for  $g\mapsto f\mathop{\Diamond}_T g$  is

$$e^{-xy}f\left(\frac{2x+y}{2}\right)$$

Since  $G(x, y) \in \hat{P}_2$ , and so the diamond product maps  $P \times P \longrightarrow P$ .

The Hermite diamond product has a simple description using the Hermite basis, and using this description we can find some interesting linear transformations. The diamond product is simply  $H_n \bigotimes_T H_m = H_{n+m}$ . Upon fixing m = 1 the linear transformation  $f \mapsto f \bigotimes_T H_1$  defines a map  $\mathbf{P} \longrightarrow \mathbf{P}$ . Using the definition of the diamond product above, this shows that the linear transformation  $H_n \mapsto H_{n+1}$  determines a map  $\mathbf{P} \longrightarrow \mathbf{P}$ . Since  $H_n \bigotimes_T 4x^2 = H_n \bigotimes_T (H_2 + 2H_0)$ , the map  $H_i \mapsto H_{i+2} + 2H_i$  maps  $\mathbf{P} \longrightarrow \mathbf{P}$ .

#### 15.7 Applications: generalized Hadamard products

In this section we characterize several generalized Hadamard products. We have seen that the two Hadamard products  $x^i * x^i = x^i$  and  $x^i *' x^i = i!x^i$  map  $P \times P^{pos} \longrightarrow P$ . The next result characterizes such general Hadamard products.

**Proposition 15.51.** Suppose that  $g(x) = \sum_{i=0}^{\infty} a_i \frac{x^i}{i!i!}$ . The generalized Hadamard

product

$$x^{i} \circledast x^{j} = \begin{cases} a_{i}x^{i} & i = j \\ 0 & otherwise \end{cases}$$

defines a map  $P \circledast P^{pos} \longrightarrow P$  if and only if  $g(x) \in \widehat{P^{pos}}$ .

*Proof.* Choose  $f = \sum b_i x^i \in \mathbf{P}^{pos}$ , and consider the map  $T_f : g \mapsto f \circledast g$ . We will show that the generating function of  $T_f$  is in  $\widehat{\mathbf{P}}_2$ . Compute:

$$\sum T_{f}(x^{n})\frac{(-y)^{n}}{n!} = \sum (x^{n} \circledast f)\frac{(-y)^{n}}{n!}$$
$$= \sum b_{n}a_{n}\frac{x^{n}(-y)^{n}}{n!}$$
$$= (f *'g)(-xy)$$

Since  $f \in \mathbf{P}^{pos}$ , and  $g \in \widehat{\mathbf{P}^{pos}}$ , we know that  $f *' g \in \mathbf{P}^{pos}$ , and consequently  $(f *' g)(-xy) \in \widehat{\mathbf{P}^{pos}}_2$ .

Conversely, since the Hadamard product extends to  $\widehat{\mathbf{P}}$  the following is in  $\widehat{\mathbf{P}^{\text{pos}}}$ :  $e^x \circledast e^x = \sum a_i \frac{x^i}{i!i!}$ 

We can generalize this to products defined by

$$\mathbf{x}^{i_1} imes \mathbf{x}^{i_2} imes \cdots imes \mathbf{x}^{i_d} \mapsto egin{cases} \mathbf{a}_i \mathbf{x}^i & i = i_1 = i_2 \cdots = i_d \ \mathbf{0} & \text{otherwise} \end{cases}$$

where all  $a_i$  are positive. A similar argument shows that a necessary condition that this product determines a linear transformation  $(\mathbf{P}^{pos})^d \longrightarrow \mathbf{P}^{pos}$  is

$$\sum_{i=0}^{\infty} \frac{a_i}{(i!)^d} x^i \in \widehat{{\textbf{P}}^{\text{pos}}}$$

Next we consider maps  $P_2^{\text{pos}} \longrightarrow P^{\text{pos}}$  that have the form

$$x^{i}y^{j} \mapsto \begin{cases} a_{i}x^{i} & i = j\\ 0 & \text{otherwise} \end{cases}$$
(15.7.1)

If  $a_i = 1$  this was called the diagonal map diag, and if  $a_i = i!$  this was called diag<sub>1</sub>. See Theorem 13.9.

**Lemma 15.52.** The map (15.7.1) defines a map  $T: \mathbf{P}_2^{\text{pos}} \longrightarrow \mathbf{P}^{\text{pos}}$  iff  $G = \sum a_i \frac{x^i}{i!i!} \in \widehat{\mathbf{P}^{\text{pos}}}$ .

*Proof.* Since  $T(e^x e^y) = \sum a_i \frac{x^i}{i!i!}$  the condition  $G \in \widehat{\mathbf{P}^{pos}}$  is necessary. From Theorem 13.9 we see that  $a_i = i!$  determines a transformation that satisfies the conclusions of the theorem. Consequently, if  $G \in \widehat{\mathbf{P}^{pos}}$  then we can express T as a composition



The map  $x^i \mapsto \frac{a_i}{i!} x^i$  defines a map  $\mathbf{P}^{pos} \longrightarrow \mathbf{P}^{pos}$  since  $\sum \frac{a_i}{i!} \frac{x^i}{i!} \in \widehat{\mathbf{P}}$  by hypothesis.

Note that  $diag_1(f(x)g(y)) = f*'g$ . Consequently, Theorem 9.87, in the case both polynomials are in  $\mathbf{P}^{\text{pos}}$ , follows from this lemma, since we can use the embedding  $\mathbf{P}^{\text{pos}} \times \mathbf{P}^{\text{pos}} \longrightarrow \overline{\mathbf{P}}_2^{\text{pos}}$  given by  $f \times g \mapsto f(x)g(y)$ .

We next show that the only products of the form

$$x^{i} \times x^{j} \mapsto a_{i+j} x^{i+j} \tag{15.7.2}$$

that determine maps  $\mathbf{P} \times \mathbf{P} \longrightarrow \mathbf{P}$  are compositions.

**Lemma 15.53.** A product (15.7.2) determines a map  $T: P \times P \longrightarrow P$  iff  $\sum a_n \frac{x^n}{n!} \in \widehat{P}$ . This is equivalent to saying that the product factors through P:

$$\begin{array}{c|c} P \times P & & 1 \\ \hline P \times P & & P \\ \hline multiplication & & \\ P & & \\ \end{array}$$

*Proof.* It suffices to evaluate  $T(e^x, e^x)$ :

$$\Gamma(\mathbf{x}^{\mathbf{x}}, \mathbf{e}^{\mathbf{x}}) = \sum_{i,j=0}^{\infty} \frac{a_{i+j}}{i!j!} \mathbf{x}^{i+j}$$
$$= \sum_{n=0}^{\infty} a_n \frac{\mathbf{x}^n}{n!} \sum_{i+j=n} \frac{n!}{i!j!}$$
$$= \sum_{n=0}^{\infty} a_n \frac{(2\mathbf{x})^n}{n!}$$

We can only state necessary conditions in the more general cases. The proofs are similar to the above, and are omitted.

**Lemma 15.54.** If the map (15.7.3) induces  $P_{2d} \longrightarrow U_d$  then  $\sum \mathbf{a}_I \frac{\mathbf{x}^I}{(I!)^2} \in \widehat{\mathbf{P}}_d$ . If the map (15.7.4) induces  $P_{2d} \longrightarrow U_d$  then  $\sum \mathbf{a}_I \frac{\mathbf{x}^I}{I!} \in \widehat{\mathbf{P}}_d$ .

$$\mathbf{x}^{I}\mathbf{y}^{J} \mapsto \begin{cases} \mathbf{a}_{I}\mathbf{x}^{I} & I = J\\ 0 & otherwise \end{cases}$$
(15.7.3)

$$\mathbf{x}^{I}\mathbf{y}^{J} \mapsto \mathbf{a}_{I+J}\mathbf{x}^{I+J} \tag{15.7.4}$$

#### **15.8** Generating functions on P(n)

The utility of generating functions is that if a certain function (the generating function) computed from a linear transformation is in some space, then the linear transformation has some nice mapping properties. In this section we look at linear transformations such as  $x^r \mapsto x^{n-r}$  that are only defined on  $\mathbf{P}(n)$ . The following proposition does not characterize such generating functions, since there is a necessary factorial.

**Proposition 15.55.** Suppose that T is a linear transformation of polynomials of degree at most n such that  $T_*(x+y)^n \in \overline{P}_2$ . Then, the linear transformation  $x^i \mapsto \frac{T(x^i)}{i!}$  maps P(n) to itself.

*Proof.* Choose  $g(x) = \sum a_i x^i \in \mathbf{P}(n)$ . Since  $g^{rev}(\frac{\partial}{\partial y})$  maps  $\overline{\mathbf{P}}_2$  to itself, we see that

$$\begin{split} g^{rev}(\frac{\partial}{\partial y})T_*(x+y)^n \bigg|_{y=0} &= g^{rev}(\frac{\partial}{\partial y})\sum_{i=0}^n \binom{n}{i}T(x^i)y^{n-i}\bigg|_{y=0} \\ &= \sum_{i=0}^n \binom{n}{i}T(x^i)\left(g^{rev}(\frac{\partial}{\partial y})y^{n-i}\bigg|_{y=0}\right) \\ &= \sum_{i=0}^n \binom{n}{i}T(x^i)(n-i)!a_i \\ &= n! T \circ \text{EXP}(g) \end{split}$$

and hence  $T \circ EXP(g) \in \mathbf{P}(n)$ .

The converse needs an extra hypothesis.

**Proposition 15.56.** Suppose  $T: P(n) \longrightarrow P(n)$ . If  $T_*(x + y)^n$  satisfies the homogeneity condition then  $T_*(x + y)^n \in P_2(n)$ .

*Proof.* We only need to check substitution, and if we substitute  $\alpha$  for y then  $T_*(x+y)^n(x,\alpha) = T(x+\alpha)^n \in \mathbf{P}$ .

**Example 15.57.** Here are a few examples.

1. Möbius transformations. Suppose  $T(x^i) = (ax + b)^i(cx + d)^{n-i}$ . Then

$$T_{*}(x+y)^{n} = \sum_{i=0}^{n} {n \choose i} (ax+b)^{i} (cx+d)^{n-i} y^{n-i} = ((ax+b)+y(cx+d))^{n-i} y^{n-i} = (ax+b)^{n-i} y^{n-i} = (ax+b)^{$$

2. Polar derivative. If  $T(x^i) = (n - i)x^i$  then

$$\mathsf{T}_*(x+y)^n = \sum_{i=0}^n \binom{n}{i} (n-i)x^i y^{n-i} = ny(x+y)^n$$

3. Reversal (with a negative sign). If  $T(x^i) = (-x)^{n-i}$  then

$$T_{*}(x+y)^{n} = \sum_{i=0}^{n} {n \choose i} (-1)^{n-i} x^{n-i} y^{n-i} = (1-xy)^{n}$$

4. Hermite. Let  $T(x^n) = H_n$ . From (6.8.2) we know that

$$\mathsf{T}_*(x+y)^n = \mathsf{T}(x^n)(x+2y) = \mathsf{H}_n(x+2y) \in \mathtt{P}_2.$$

This implies that  $x^n \mapsto H_n/n!$  maps P(n) to itself, but in this case we know T maps P(n) to itself.

5. In Lemma 8.30 we saw that the linear transformation T:  $x^k \mapsto (\underline{x + n - k})_n$  maps  $\mathbf{P}^{alt}(n) \longrightarrow \mathbf{P}(n)$ . The generating function

$$T_*(x+y)^n = \sum_{i=0}^n \binom{n}{i} (\underline{x+n-k})_n y^{n-i} = (y+1)^n \langle \underline{x} \rangle_n$$

is in  $\overline{\mathbf{P}}_2(2n)$ . Proposition 15.55 only allows us to conclude that

$$x^{k} \mapsto \frac{(x+n-k)_{n}}{k!} \operatorname{maps} \mathbf{P}(n) \longrightarrow \mathbf{P}(n).$$
 (15.8.1)

Since it is not true that  $x^k \mapsto (\underline{x + n - k})_n$  maps  $\mathbf{P}(n)$  to itself, we see that the factorial in Proposition 15.55 is necessary.

#### 15.9 Higher dimensional generating functions

If  $T: \mathbf{P}_d \longrightarrow \mathbf{P}_d$  is a linear transformation, then its generating function is a function of 2d variables:

$$\mathsf{T}_*(e^{-xy}) = \sum_{\mathbf{I}} \mathsf{T}(x^{\mathbf{I}}) \frac{(-y)^{\mathbf{I}}}{\mathbf{I}!}$$

Here are some examples.

The generating function of the identity transformation is  $e^{-x \cdot y}$ .

Suppose that  $T: \mathbf{P} \longrightarrow \mathbf{P}$  and let F(x, y) be the generating function of T. If we define  $T_*(x^i y^j) = T(x^i)y^j$  then the generating function  $F_*$  of  $T_*$  is

$$\begin{split} F_*(x,y,u,\nu) &= \sum_{i,j=0}^{\infty} T(x^i) y^j \frac{(-u)^i}{i!} \frac{(-\nu)^j}{j!} \\ &= \left(\sum_{i=0}^{\infty} T(x^i) \frac{(-u)^i}{i!}\right) \left(\sum_{j=0}^{\infty} y^j \frac{(-\nu)^j}{j!}\right) = F(x,u) e^{-y\nu} \end{split}$$

More generally, if we have another linear transformation  $S: \mathbf{P} \longrightarrow \mathbf{P}$  with generating function G(x, y) then the generating function of  $x^i y^j \mapsto T(x^i)S(y^j)$  is F(x, u)G(y, v).

If T:  $P_2 \longrightarrow P_2$  is a linear transformation and a, b are positive, then  $S(f) = a \frac{\partial}{\partial x}T(f) + b \frac{\partial}{\partial x}T(f)$  satisfies  $T(f) \le S(f)$ . If T has generating function F, then the generating function of S is  $aF_x + bF_y$ .

The generating function of the linear transformation  $g \mapsto f(\partial_x, \partial_y)g$  has generating function  $f(u, v)e^{-xu-yv}$ . This follows using linearity from the calculation

$$\sum_{i,j} \partial_x^r \partial_y^s (x^i y^j) \frac{(-u)^i}{i!} \frac{(-\nu)^j}{j!} = \left(\sum_i \partial_x^r x^i \frac{(-u)^i}{i!}\right) \left(\sum_j \partial_y^r y^j \frac{(-\nu)^j}{j!}\right)$$
$$= \left(u^r e^{-xu}\right) \left(\nu^s e^{-y\nu}\right)$$

In general, the generating function of  $g \mapsto f(\partial_x)g$  is  $f(x)e^{-x \cdot y}$ . Similarly, the generating function of  $f \mapsto f(\partial_x)g$  is g(x + y).

**Proposition 15.58.** *If* T *is a linear transformation on*  $\overline{\mathbf{P}}_d$  *and*  $T_*(e^{-\mathbf{x}\cdot\mathbf{y}}) \in \widehat{\mathbf{P}}_{2d}$  *then*  $T: \overline{\mathbf{P}}_d \longrightarrow \overline{\mathbf{P}}_d$ .

*Proof.* Since  $f(\partial_x)$  maps  $\widehat{P}_{2d}$  to itself the proof is the same as the case d = 1.  $\Box$ 

**Corollary 15.59.** If S is a d by d positive symmetric matrix, and  $H_I(\mathbf{x})$  is the corresponding Hermite polynomial then the linear transformation  $\mathbf{x}^I \mapsto H_I(\mathbf{x})$  maps  $\overline{\mathbf{P}}_d$  to  $\overline{\mathbf{P}}_d$ .

*Proof.* The generating function of the Hermite polynomials (13.5.1) is  $\exp(-ySy^* - 2ySx^*)$  which is in  $\widehat{\mathbf{P}}_{2d}$ . We can now apply the Proposition.

The proposition implies that differential operators preserve  $P_d$ .

**Lemma 15.60.** *If*  $f \in P_d$  *then the linear transformation* 

$$T: g \mapsto f(\partial_x)g$$

maps  $P_d$  to itself.

*Proof.* The generating function of T is  $f(\mathbf{y})e^{-\mathbf{x}\cdot\mathbf{y}}$  which is in  $\widehat{\mathbf{P}}_{2d}$ .

Table 15.3 lists some generating functions in higher dimensions. Note that these transformations do not all map  $P_d$  to itself since not all of the generating functions are in  $\hat{P}_{2d}$ . The operator  $EXP_x$  is  $EXP_{x_1} \cdots EXP_{x_d}$ , and the generalized Hurwitz transformation is defined as

$$\mathsf{T}(\mathbf{x}^{\mathbf{I}}) = \begin{cases} \mathbf{x}^{\mathbf{I}/2} & \text{if all coordinates of } \mathbf{I} \text{ are even} \\ 0 & \text{otherwise} \end{cases}$$
(15.9.1)

#### 15.10 Higher dimensional multiplier transformations

The Pólya-Schur theorem does not generalize to two variables. Although we will see that all multiplier transformations are products of one dimensional multipliers, there are examples of transformations that do not map  $P_2$  to itself, yet have a two variable generating function in  $\overline{P}_2$ .

[18] observed that all higher dimensional multiplier transformations are just products of one dimensional maps:

Name					$T_*(e^{-xy})$
differentiation	$\sum_{\mathbf{I}}$	$f(\vartheta_x)x^I$	$\frac{(-\mathbf{y})^{\mathbf{I}}}{\mathbf{I}!}$	=	$f(\mathbf{y})e^{-\mathbf{x}\cdot\mathbf{y}}$
Hermite	$\sum_{I}^{I}$	$H_{I}(x)$	$\frac{(-\mathbf{y})^{\mathbf{I}}}{\mathbf{I}!}$	=	$e^{-ySy'-2ySx'}$
Hadamard	$\sum_{I}$	$f(\mathbf{x}) \ast \mathbf{x}^{\mathrm{I}}$	$\frac{(-\mathbf{y})^{\mathbf{I}}}{\mathbf{I}!}$	=	$EXP_{\mathbf{x}}f(-x_1y_1,\ldots,-x_dy_d)$
Hurwitz	$\sum_{I}$	$\mathbf{x}^{\mathrm{I}/2}$	$\frac{(-\mathbf{y})^{\mathbf{I}}}{\mathbf{I}!}$	=	$\cosh(y_1\sqrt{x_1})\cdots\cosh(y_d\sqrt{x_d})$
identity	$\sum_{I}^{I}$	x <sup>I</sup>	$\frac{(-\mathbf{y})^{\mathbf{I}}}{\mathbf{I}!}$	=	$e^{-\mathbf{x}\cdot\mathbf{y}}$

Table 15.3: Generating Functions in higher dimensions

Lemma 15.61. The following are equivalent:

- 1.  $T: \mathbf{x}^I \longrightarrow \alpha_I \mathbf{x}^I$  maps  $\overline{\mathbf{P}}_d$  to itself.
- 2. T is a product of one-dimensional transformations.

*Proof.* We first assume that d = 2, so assume that  $T(x^iy^j) = \alpha_{ij}$ . Following [18] we apply T to two test functions

$$\begin{split} & \mathsf{T} x^i y^j (1+x) (1+y) = x^i y^j \big( \alpha_{i,j} + \alpha_{i+1,j} x + \alpha_{i,j+1} y + \alpha_{i+1,j+1} x y \big) \\ & \mathsf{T} x^i y^j (1+x) (1-y) = x^i y^j \big( \alpha_{i,j} + \alpha_{i+1,j} x - \alpha_{i,j+1} y - \alpha_{i+1,j+1} x y \big). \end{split}$$

By Proposition 11.26 or Lemma 20.1 both of the factors are in  $\overline{\mathbf{P}}_2$ , and by Theorem 9.113

$$\alpha_{i,j}\alpha_{i+1,j+1} - \alpha_{i+1,j}\alpha_{i,j+1} \ge 0$$
$$-\alpha_{i,j}\alpha_{i+1,j+1} + \alpha_{i+1,j}\alpha_{i,j+1} \ge 0$$

and therefore

$$\alpha_{i,j}\alpha_{i+1,j+1} = \alpha_{i+1,j}\alpha_{i,j+1} \tag{15.10.1}$$

We now consider the *support* S of T - that is, the set of all (i, j) such that  $\alpha_{i,j} \neq 0$ . If r is a non-negative integer then the composition

$$\mathbf{P} \xrightarrow{\mathbf{x}^n \mapsto \mathbf{x}^n \mathbf{y}^r} \overline{\mathbf{P}}_2 \xrightarrow{\mathsf{T}} \overline{\mathbf{P}}_2 \xrightarrow{\mathbf{x}^n \mathbf{y}^r \mapsto \mathbf{x}^n} \mathbf{P}$$

determines a map  $\mathbf{P} \longrightarrow \mathbf{P}$ . Since this map is a multiplier transformation in one variable we know that the non-zero coefficients have no gaps, and so form an interval. Thus, all intersections of the support S with a horizontal or

vertical line are intervals. The identity (15.10.1) shows that we can not have the configurations

$$\begin{array}{c|c} \neq 0 & 0 \\ \neq 0 & \neq 0 \end{array} \qquad \qquad \begin{array}{c} \neq 0 & \neq 0 \\ \hline 0 & \neq 0 \end{array}$$

in the support, so it follows that the support is a rectangle.

We may assume that  $\alpha_{0,0} = 1$ . Using the recursion (15.10.1) and induction yields  $\alpha_{ij} = \alpha_{i,0} \alpha_{0,j}$ . It follows that T is a product of the one dimensional transformations

$$T_r: x^n \mapsto \alpha_{n,0} x^n$$
  $T_c: y^n \mapsto \alpha_{0,n} y^r$ 

The case for general d is no different; we use the fact that the intersection with every d - 1 dimensional face is a product to conclude the support is a product.

**Example 15.62.** Here is an example of a multiplier transformation that does not map  $P_2 \longrightarrow P_2$  but whose two variable generating function is in  $\widehat{P}_2$ . The two variable generating function of  $T: x^i y^j \mapsto (i+j)^2 x^i y^j$  is

$$\sum_{i,j=0}^{\infty} \frac{(i+j)^2}{i!j!} x^i y^j = e^{x+y} (x+y)(x+y+1)$$

which is in  $\widehat{\mathbf{P}}_2$ . We know T maps  $\mathbf{P}_2^{\text{pos}}$  to itself, but T does not map  $P_2$  to itself. If

$$f = -193 + 43 x + 21 x^{2} + x^{3} + 262 y + 196 x y + 14 x^{2} y + 448 y^{2} + 64 x y^{2} + 96 y^{3} + 100 y^{2} 

then Tf(-2, y) has complex roots. Note that the usual generating function is

$$\sum_{i,j=0}^{\infty} (i+j)^2 x^i y^j \frac{(-u)^i (-\nu)^j}{i! \, j!} = e^{-ux - \nu y} (ux + \nu y) (ux + \nu y - 1)$$

and the latter expression is not in  $\hat{\mathbf{P}}_4$ .

*Remark* 15.63. In Corollary 11.114 we saw that the linear transformation  $x^i y^j \mapsto \frac{x^i y^j}{(i+j)!}$  maps  $\mathbf{P}_2^{\text{pos}}$  to itself, yet it does not map  $\mathbf{P}_2$  to itself. The generating function is

$$\sum_{i,j} \frac{x^i y^j}{(i+j)!} = \frac{xe^x - ye^y}{x - y}$$

This can be seen as follows. If the generating function is s, then the terms of  $\frac{x}{y}(s-1)$  largely cancel, leaving only  $\frac{x}{y}e^x - e^y$ . This generating function is not in  $\widehat{\mathbf{P}}_2$  since it is not an entire function.

Just as in one variable (Lemma 1.49) a linear transformation  $T: \overline{\mathbf{P}}_2 \longrightarrow \overline{\mathbf{P}}_2$  determines a simpler linear transformation.

**Lemma 15.64.** *If*  $T: \overline{P}_2 \longrightarrow \overline{P}_2$ , and we can write

$$T(x^{i}y^{j}) = g_{ij}(x, y) = c_{ij}x^{i}y^{j} + \text{ terms of lower degree}$$

then the linear transformation  $S(x^iy^j) = c_{ij}x^iy^j$  also maps  $\overline{P}_2 \longrightarrow \overline{P}_2$ .

*Proof.* Choose  $f = \sum a_{ij} x^i y^j$  in  $\overline{\mathbf{P}}_2$ . Substitute  $x/\alpha$  for x,  $y/\beta$  for y and apply T. Next, substitute  $\alpha x$  for x and  $\beta y$  for y. The result is that the following polynomial is in  $\overline{\mathbf{P}}_2$ :

$$\sum_{i,j} a_{ij} \alpha^i \beta^j g_{ij}(x/\alpha, y/\beta)$$

Next, using the fact that

$$\lim_{\alpha \to 0^+} \lim_{\beta \to 0^+} \alpha^{i} \beta^{j} g_{ij}(x/\alpha, y/\beta) = c_{ij} x^{i} y^{j}$$

we see that

$$\lim_{\alpha \to 0^+} \lim_{\beta \to 0^+} \sum_{i,j} a_{ij} \alpha^i \beta^j g_{ij}(x/\alpha, y/\beta) = \sum_{i,j} a_{ij} c_{ij} x^i y^j = S(f) \in \overline{P}_2$$

**Example 15.65.** The two variable generating function of  $x^r y^s \mapsto f(r+s)x^r y^s$  can be easily found.

$$\sum (i+j)^{n} \frac{x^{i}y^{j}}{i!j!} = e^{x+y} T_{**}^{-1} (x+y)^{n}$$

where  $T\colon x^n=(\underline{x})_n$ , and  $T_{**}^{-1}(x^ry^s)=T^{-1}(x^r)T^{-1}(y^s).$  By linearity we find the generating function

$$\sum f(x+y)\frac{x^{i}y^{j}}{i!j!} = e^{x+y}T_{**}^{-1}f(x+y) = (e^{x}T^{-1}f)(x+y)$$

where the last equality follows from Example 6.53. It follows that if  $f \in \mathbf{P}^{\text{pos}}$  then this generating function is in  $\widehat{\mathbf{P}}_2$ . We saw that the linear transformation corresponding to  $f(x) = x^2$  does not map  $P_2$  to itself.

#### 15.11 Differential operators

We continue the investigation of differential operators. We first compute the generating functions of differential operators acting on **P**, extract some properties, show some eigenpolynomials are in **P**, and then extend some of the results to  $P_d$ .

Consider the linear transformation T defined on polynomials of one variable

$$T(g) = \sum_{i=0}^n f_i(x) g^{(i)}(x)$$

We define  $f(x,y) = \sum f_i(x)y^i,$  and write  $\mathsf{T}(g) = f(x,\mathsf{D})g.$  The generating function of T is

$$\begin{split} \sum_{i=0}^{\infty} \mathsf{T}(x^{j}) \frac{(-y)^{j}}{j!} &= \sum_{j=0}^{\infty} \sum_{i=0}^{n} \mathsf{f}_{i}(x) \mathsf{D}^{i} x^{j} \frac{(-y)^{j}}{j!} \\ &= \sum_{i=0}^{n} \mathsf{f}_{i}(x) \mathsf{D}^{i} \sum_{j=0}^{\infty} x^{j} \frac{(-y)^{j}}{j!} \\ &= \sum_{i=0}^{n} \mathsf{f}_{i}(x) \mathsf{D}^{i} e^{-xy} \\ &= \sum_{i=0}^{n} \mathsf{f}_{i}(x) (-y)^{i} e^{-xy} \\ &= \mathsf{f}(x, -y) e^{-xy} \end{split}$$

Thus, we have

**Proposition 15.66.** *If*  $f \in \overline{P}_2$  *then* f(x, -D) *maps* P *to* P*. Conversely, if* 

- 1. f(x, -D) maps **P** to **P**.
- 2. The coefficients of the homogeneous part of f(x, y) are all positive.

then  $f \in P_2$ .

*Proof.* The generating function is  $f(x, y)e^{-xy} \in \widehat{\mathbf{P}}_2$ . In Lemma 7.14 we saw that  $f(x, \alpha) \in \mathbf{P}$  for all  $\alpha \in \mathbb{R}$ , so  $f \in P_2$ .

*Remark* 15.67. For example, if we take F = x + y then the linear transformation is  $f \mapsto xf - f'$ . We know that this is in **P** since  $xf \leq f'$  implies  $xf - f' \leq f$ .

**Corollary 15.68.** Suppose that  $T: P \longrightarrow P$ . The linear transformation

$$\mathfrak{g}\mapsto \sum \mathsf{T}(x^k)\frac{\mathfrak{g}^{(k)}(x)}{k!}(-1)^k$$

maps  $P \longrightarrow P$ .

*Proof.* If f(x, y) is the the generating function of T, then  $f(x, y) \in P_2$ . The map is  $g(x) \mapsto f(x, -D)g(x)$ .

In [10] they show that for n sufficiently large there is an eigenpolynomial f(x, D) of degree n. We now show that the eigenpolynomials are in **P** if  $f \in \overline{\mathbf{P}}_2$ , and f satisfies a degree condition.

**Lemma 15.69.** *Choose*  $f(x, y) \in \overline{\mathbf{P}}_2(d)$ 

$$f(x, y) = f_0(x) + f_1(x)y + f_2(x)y^2 + \dots + f_d(x)y^d$$

where the degree of  $f_i$  is i. If n is sufficiently large then there is a polynomial  $p \in P(n)$ and constant  $\lambda$  such that  $f(x, D)p = \lambda p$ .

*Proof.* Since f(x, D) maps **P** to itself, and preserves degree, we will apply Lemma 7.85. It suffices to show that there is a dominant eigenvalue, which is the same as finding a dominant leading coefficient.

The r-th diagonal element of M is the coefficient  $c_r$  of  $x^r$  in  $f(x, D)x^r$ . Denoting the leading coefficient of  $f_i$  by  $C_i$ ,

$$c_d = \sum_{i=0}^d c_i \left(\underline{r}\right)_i = C_d(\underline{r})_d + O(r^{d-1}).$$

For n sufficiently large,  $c_n$  is the largest eigenvalue of M.

*Remark* 15.70. The identity  $e^{-(\partial_x + \partial_y)^2}(x + y)^n = (x + y)^n$  shows that the operator  $e^{-(\partial_x + \partial_y)^2}$  has polynomial eigenvalues of every degree.

We can generalize some of the above to more variables. Suppose that  $f(\pmb{x},\pmb{y})\in \mathtt{P}_{d+e}$  and we define a differential operator

$$T(g(\mathbf{x})) = \sum_{I} f_{I}(\mathbf{x}) D^{I}g(\mathbf{x})$$
(15.11.1)

It is easy to see that the generating function of T is simply  $f(x, y)e^{-x \cdot y}$ .

Proposition 15.71. Suppose

- 1.  $f(\mathbf{x}, -D)$  maps  $P_d$  to  $P_d$ .
- 2. The coefficients of the homogeneous part of  $f(\mathbf{x}, \mathbf{y})$  are all positive.

then  $f \in P_{d+e}$ . Conversely, if  $f(\mathbf{x}, \mathbf{y}) \in \overline{\mathbf{P}}_{2d}$  then  $f(\mathbf{x}, -D)$  maps  $\overline{\mathbf{P}}_d$  to itself.

*Proof.* If suffices to show that  $f(\mathbf{x}, \mathbf{y})$  satisfies substitution. The proof is the same as Lemma 7.14: apply  $f(\mathbf{x}, -D)$  to  $e^{-\mathbf{a}\cdot\mathbf{x}}$  to conclude that  $f(\mathbf{x}, \mathbf{a})e^{-\mathbf{a}\cdot\mathbf{x}} \in P_d$ . Multiplying by  $e^{\mathbf{a}\cdot\mathbf{x}}$  finishes the proof.

The second part follows form the proposition.

Which polynomials f(x, y) determine operators f(x, D) that map  $P^{pos} \longrightarrow P$ , and don't map  $P \longrightarrow P$ ? Here's a necessary condition.

**Lemma 15.72.** Suppose that f(x, y) determine an operator f(x, D) that maps  $P^{\text{pos}} \longrightarrow P$ . If f(x, -y) satisfies degree and positivity then  $f(x, -y) \in \mathscr{P}_{1,1}$ .

*Proof.* Let  $f(x,y) = \sum a_{ij}x^iy^j$ . Since  $e^{-\alpha x} \in \mathbf{P}^{pos}$  for positive  $\alpha$ , we see that

$$\left(\sum a_{ij}x^{i}\mathsf{D}^{j}\right)e^{-\alpha x} = \left(\sum a_{ij}x^{i}(-\alpha)^{j}\right)e^{-\alpha x}$$

is in  $\widehat{\mathbf{P}}$ , and hence  $f(x, -\alpha) \in \mathbf{P}$  for all positive  $\alpha$ . Since f(x, -y) satisfies degree and positivity,  $f(x, -y) \in \mathcal{P}_{1,1}$ .

## **15.12** Generating functions for maps $P_2 \leftrightarrow P$

If  $T: \mathbf{P}_2 \longrightarrow \mathbf{P}$  then the generating function of T lies in  $\widehat{\mathbf{P}}_3$ . We compute the generating functions for many of these maps.

The most basic map of all is evaluation. The generating function of  $f(x, y) \mapsto f(z, \alpha)$  is

$$\sum T(x^{i}y^{j})\frac{(-u)^{i}(-\nu)^{j}}{i!j!} = \sum z^{i}\alpha^{j}\frac{(-u)^{i}(-\nu)^{j}}{i!j!} = e^{-uz-\alpha\nu}$$

Next, consider the diagonal map  $f(x, y) \mapsto f(z, z)$ . The generating function is

$$\sum T(x^{i}y^{j})\frac{(-u)^{i}(-\nu)^{j}}{i!j!} = \sum z^{i+j}\frac{(-u)^{i}(-\nu)^{j}}{i!j!} = e^{-(u+\nu)z}$$

The map that extracts the coefficient of a fixed monomial also maps  $P_2 \longrightarrow \mathbf{P}$ . If  $T(x^iy^j) = z^i$  if j = k and 0 otherwise then the generating function is

$$\sum T(x^{i}y^{j})\frac{(-u)^{i}(-v)^{j}}{i!j!} = \sum_{i} z^{i}\frac{(-u)^{i}(-v)^{k}}{i!k!} = \frac{(-v)^{k}}{k!}e^{-uz}$$

Finally, we consider the map  $f(x, y) \mapsto f(D, x)$ . More precisely, this is the map  $x^i y^j \mapsto \frac{d^i}{dz^i} z^j = (\underline{j})_i z^{j-i}$ .

$$\begin{split} \sum_{i\leqslant j} (\underline{j})_i z^{j-i} \frac{(-u)^i (-\nu)^j}{i!j!} &= \sum_{j=0}^\infty \frac{(-z\nu)^j}{j!} \sum_{i=0}^j {j \choose i} \left(\frac{-u}{z}\right)^i \\ &= \sum_{j=0}^\infty \frac{(-z\nu)^j}{j!} \left(1 + \frac{-u}{z}\right)^j = e^{-z\nu + u\nu} \end{split}$$

We summarize these few results in Table 15.4

We can construct transformations  $P \longrightarrow {\tt P}_2$  by homogenizing transformations  $P \longrightarrow P.$ 

Name				$T_*(e^{-xy})$
evaluation at $\alpha$	f(x, y)	$\mapsto$	$f(z, \alpha)$	$e^{-uz-\alpha v}$
diagonal	f(x, y)	$\mapsto$	f( <i>z</i> , <i>z</i> )	$e^{-z(u+v)}$
coefficient	f(x, y)	$\mapsto$	coef. of y <sup>k</sup>	$\frac{v^k}{k!}e^{-uz}$
derivative	f(x, y)	$\mapsto$	f(-D, z)	$e^{-\nu(u+z)}$

Table 15.4: Generating Functions for  $\mathbf{P}_2 \longrightarrow \mathbf{P}$ 

**Lemma 15.73.** Suppose  $T: \mathbf{P} \longrightarrow \mathbf{P}$  and  $T(x^n) \in \mathbf{P}^{pos}(n)$ . If  $T(x^i) = f_i(x)$  and  $T_H(x^i) = f_i(x/y)y^i$  then  $T_H: \mathbf{P} \longrightarrow P_2$ .

*Proof.* If  $g = \sum a_i x^i \in \mathbf{P}(n)$  then  $T_H(g) = \sum a_i f_i(x/y) y^i$ . The homogeneous part of  $T_H(g)$  is  $f_n(x/y) y^n$  which has all positive terms since  $T(x^n) \in \mathbf{P}^{pos}$ . Substituting  $\alpha$  for y yields

$$T_{H}(g)(x,\alpha) = \sum a_{i}f_{i}(x/\alpha)\alpha^{i} = \sum (\alpha^{i}a_{i})f_{i}(x/\alpha) = T(g(\alpha x))(x/\alpha)$$

which is in **P**.

If the generating function of T is G(x, y) then the generating function of  $T_H$  is G(x/y, yz).

Note that evaluation at x = 1 shows that  $T: x^i \longrightarrow f_i^{rev}$  maps  $P \longrightarrow P$ . This gives another proof of Corollary 9.57.

**Example 15.74.** If T is the affine transformation  $x \mapsto x+1$  then  $T_H(g) = g(x+y)$ . For a more complicated example, if  $T(x^i) = L_i(-x)$  is the Laguerre transformation then

$$T_{H}(x^{n}) = \sum_{0}^{n} \frac{1}{k!} {n \choose k} x^{k} y^{n-k}$$

# **15.13** Linear transformations $P^{pos} \longrightarrow P$

In this section we consider maps  $\mathbf{P}^{pos} \longrightarrow \mathbf{P}^{pos}$ . If we have a linear transformation  $T: \mathbf{P}^{alt} \longrightarrow \mathbf{P}^{alt}$  then we can construct a map  $S: \mathbf{P}^{pos} \longrightarrow \mathbf{P}^{pos}$  by S(f)(x) = T(f(-x))(-x). The two generating functions satisfy

$$f(x,y) = \sum T(x^{n}) \frac{(-y)^{n}}{n!}$$
  
$$g(x,y) = \sum S(x^{n}) \frac{(-y)^{n}}{n!} = \sum (-1)^{n} T(x^{n}) (-x) \frac{y^{n}}{n!}$$

and so g(x, y) = f(-x, -y).

**Theorem 15.75.** Suppose  $T: P^{pos} \longrightarrow P^{pos}$ , T preserves degree and the sign of the leading coefficient. If

$$F(x,y) = T_*(e^{-xy}) = \sum_{i=0}^{\infty} T(x^i) \frac{(-y)^i}{i!}$$
(15.13.1)

 $\textit{then } F(x,y) \in \widehat{\mathcal{P}}_{1,1}.\textit{ If } T \colon \textit{P}^{alt} \longrightarrow \textit{P}^{alt}\textit{ then } F(x,-y) \in \widehat{\mathcal{P}}_{1,1}.$ 

*Proof.* It suffices to assume that  $T: \mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}^{\text{pos}}$ . Since  $e^{-xy} \in \widehat{\mathbf{P}}_2 \subset \widehat{\mathbf{P}}_{1,1}$ , we know that  $F(x, y) = T_*(e^{-xy}) \in \widehat{\mathbf{P}}_{1,1}$ .

**Theorem 15.76.** Suppose that  $F(x, y) = \sum f_i(x) \frac{(-y)^i}{i!}$  is in  $\widehat{\mathcal{P}}_{1,1}$ . If each  $f_i$  is a polynomial and we define  $T(x^i) = f_i/i!$  then  $T: \mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}$ .

*Proof.* We follow the proof for  $\mathbf{P}_d$ , but since  $\mathcal{P}_{1,1}$  is not closed under differentiation we need to multiply. This is why we have the factorial appearing.

Since  $\mathcal{P}_{1,1}$  is closed under multiplication, so is  $\widehat{\mathcal{P}}_{1,1}$ . Look at Table 15.5. We know that  $e^{-\alpha x} \in \widehat{\mathbf{P}}_2$ , and  $(1 + y)^x \in \widehat{\mathcal{P}}_{1,1}$  since the latter is the generating function of a linear transformation  $\mathbf{P}^{\alpha lt} \longrightarrow \mathbf{P}^{\alpha lt}$ . Their product is in  $\widehat{\mathcal{P}}_{1,1}$ , and is the generating function for the Charlier transformation.

Table 15.5 lists the generating functions of some linear transformations  $\mathbf{P}^{\text{alt}} \longrightarrow \mathbf{P}^{\text{alt}}$ . Table 15.13 lists the generating functions of linear transformations  $\mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}$  whose image is not  $\mathbf{P}^{\text{pos}}$ . Notice that most of these generating functions are not in  $\hat{\mathbf{P}}_2$  - indeed, they are not analytic for all real values of the parameters, but only positive y.

**Example 15.77.** We can determine the closed formula for the generating function F(x, y) of T:  $x^i \mapsto (-1)^{\binom{i}{2}} x^i$  given in Table 15.5 by considering four sums determined by the index mod 4. The result is

$$F(x,y) = \sum_{i=0}^{\infty} (-1)^{\binom{i}{2}} x^{i} \frac{y^{i}}{i!} = \sqrt{2} \cos(xy - \pi/4).$$

The cosine has a representation as an infinite product

$$\cos(x) = \prod_{k=0}^{\infty} \left( 1 - \frac{4x^2}{(2k+1)^2 \pi^2} \right)$$

so the generating function can be expressed as the infinite product

$$F(x,y) = \prod_{k=0}^{\infty} \left( 1 - \frac{4(xy - \frac{\pi}{4})^2}{(2k+1)^2 \pi^2} \right).$$

Name					$T_*(e^{-xy})$
Binomial	$\sum_{i=0}^{\infty}$	$\begin{pmatrix} x \\ i \end{pmatrix}$	$\frac{(-y)^{\mathfrak{i}}}{\mathfrak{i}!}$	=	$_{1}F_{1}(-x, 1, -y)$
Charlier	$\sum_{i=0}^{\infty}$	$C_n^{\alpha}$	$\frac{(-y)^{i}}{i!}$	=	$e^{-\alpha y}(1+y)^{\chi}$
Falling Factorial	$\sum_{i=0}^{\infty}$	$(\underline{x})_i$	$\frac{(-y)^{i}}{i!}$	=	$(1-y)^x$
Hurwitz	$\sum_{i=0}^{\infty}$	x <sup>i</sup>	$\frac{y^{2i}}{(2i)!}$	=	$\cosh(\sqrt{xy^2})$
Hurwitz	$\sum_{i=0}^{\infty}$	x <sup>i</sup>	$\frac{\mathtt{y}^{4\mathtt{i}}}{(4\mathtt{i})!}$	=	$\tfrac{1}{2}\left(cos(x^{1/4}y)+cosh(x^{1/4}y)\right)$
q-series	$\sum_{i=0}^{\infty}$	$(-1)^{\binom{i}{2}} x^i$	$\frac{(-y)^{i}}{i!}$	=	$\sqrt{2}\cos(xy-\pi/4)$
Rising Factorial	$\sum_{i=0}^{\infty}$	$\left( \left< \underline{x} \right>_i \mapsto x^i \right)$	$\frac{(-y)^{i}}{i!}$	=	$e^{\mathbf{x}(1-e^{-y})}$

Table 15.5: Generating functions for  $P^{\texttt{alt}} \longrightarrow P^{\texttt{alt}}$ 

Name		$T_*(e^{-xy})$
Hermite	$\sum_{i=0}^{\infty} H_{n}^{re\nu} \frac{(-y)^{i}}{i!}$	$= e^{2y-x^2y^2}$
Hermite	$\sum_{i=0}^{\infty} x^n H_n \frac{(-y)^i}{i!}$	$= e^{2x^2y-x^2y^2}$
Laguerre	$\sum_{i=0}^{i=\infty} L_n^{rev}  \frac{(-y)^i}{i!}$	$= e^{xy}J_0(2\sqrt{y})$

Table 15.6: Generating functions for  $\mathbf{P}^{pos} \longrightarrow \mathbf{P}$ 

The product is not in  $\widehat{\mathbf{P}}_2$  as the factorization shows

$$\begin{pmatrix} 1 - \frac{4(xy - \frac{\pi}{4})^2}{(2k+1)^2 \pi^2} \end{pmatrix} = \left( 1 - \frac{2(xy - \frac{\pi}{4})}{(2k+1)\pi} \right) \left( 1 + \frac{2(xy - \frac{\pi}{4})}{(2k+1)\pi} \right)$$
$$= F_1(x, y)F_2(x, y)$$

 $F_1(x,y) \in \widehat{P}_2$ , but  $F_2(x,-y) \in \widehat{P}_2$ . This reflects the fact that the linear transformation  $x^i \mapsto (-1)^{\binom{i}{2}} x^i$  maps  $P^{pos}$  to P, and does not map  $P \longrightarrow P$ .

**Example 15.78.** One way to find elements in  $\mathcal{P}_{1,1}$  is to compute the generating function of simple linear transformations. The generating function of  $U(x^n) = 0$  if n odd, and  $x^n$  if n even is

$$\sum_{n=0}^{\infty} x^{2n} \frac{y^{2n}}{(2n)!} = \cosh(xy)$$

while the generating function of  $V(x^n) = 0$  if n even, and  $x^n$  if n odd is

$$\sum_{n=0}^{\infty} x^{2n+1} \frac{y^{2n+1}}{(2n)!} = \sinh(xy)$$

Let  $T_e(x^n) = 0$  if n is odd, and  $x^{n/2}$  if n is even. We know that  $T_e : \mathbf{P}^{pos} \longrightarrow \mathbf{P}^{pos}$ . Consequently, the map  $S(f) = T_e(f)(1 - x^2)$  maps  $\mathbf{P}^{pos} \longrightarrow \mathbf{P}$ . The generating function of  $T_e$  is for positive x

$$\sum_{n=0}^{\infty} x^n \frac{y^{2n}}{(2n)!} = \cosh(y\sqrt{x})$$

and the generating function of S is

$$\sum_{n=0}^{\infty} (1-x^2)^n \frac{y^{2n}}{(2n)!} = \cos(y\sqrt{1-x^2})$$

Finally, let  $T_o(x) = 0$  if n even, and  $x^{(n-1)/2}$  if n odd. The generating function of  $T_o$  for positive x is

$$\sum_{n=0}^{\infty} x^n \frac{y^{2n+1}}{(2n+1)!} = \sqrt{x} \sinh(y\sqrt{x})$$

#### **15.14** More generating functions

In this section we list in Table 15.14 a few generating functions for linear transformations that map  $\mathbf{P}^{\text{finite}} \longrightarrow \mathbf{P}^{J}$ . These functions are not entire - in order to converge we must assume that  $|\mathbf{x}|, |\mathbf{y}| < 1$ .

Name			$T_*(e^{-x\mathfrak{y}})$				
Chebyshev	$\sum_{i=0}^{\infty}$	T <sub>i</sub>	$\frac{(-y)^{i}}{i!}$	=	$e^{xy} \cosh(\sqrt{y^2(x^2-1)})$		
Legendre	$\sum_{i=0}^{\infty}$	Pi	$\frac{(-y)^{i}}{i!}$	=	$e^{xy} {}_0F_1(;1;\frac{1}{4}(y^2-1)x^2)$		
Euler	$\sum_{i=0}^{i=0}$	A <sub>i</sub>	$\frac{(-y)^{i}}{i!}$	=	$\frac{e^{xy} - xe^{xy}}{e^{xy} - xe^{y}}$		
Factorial	$\sum_{\mathfrak{i}=0}^\infty$	${\textstyle\prod}_{k=1}^{i}(1-kx)$	$\frac{(-y)^{i}}{i!}$	=	$(1+xy)^{-1+1/x}$		

Table 15.7: Miscellaneous generating functions

Since the method of finding bivariate generating functions may not be familiar, I'll describe how to find the generating function for the Euler transformation. We compute the generating function of  $x^n \mapsto A_n$  by first computing the generating function of  $x^n \mapsto B_n = x^n A_n(\frac{x+1}{x})$ . We use the recurrence for  $B_n$  to derive a partial differential equation for the generating function, which we use the method of characteristics to solve. So recall (Lemma 7.60) that  $B_n$  satisfies the recursion  $B_{n+1} = (x + 1)(B_n + xB'_n)$ . Substituting this into the generating function for T

$$F(x,y) = \sum_{n=0}^{\infty} B_n \frac{y^n}{n!}$$
$$= \sum_{n=0}^{\infty} (x+1)(B_n + xB'_n) \frac{y^n}{n!}$$

with some manipulation yields the partial differential equation

$$(x+1)F(x,y) = (-x^2 - x)F_x(x,y) + F_y(x,y)$$
(15.14.1)

The method of characteristics assumes that x, y are functions of s, t, where we consider t a variable, and s a parameter. We choose these functions to make each side of (15.14.1) an exact differential. Thus, we have to solve the

three equations

$$\frac{d}{dt}x(s,t) = -x^2 - x \tag{15.14.2}$$

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathbf{y}(\mathbf{s},\mathbf{t}) = 1 \tag{15.14.3}$$

$$\frac{d}{dt}F(s,t) = (x+1)F(s,t)$$
(15.14.4)

with the initial conditions

$$x(s,0) = s$$
  
 $y(s,0) = 0$   
 $F(s,0) = 1$ 

We solve (15.14.2) and (15.14.3) for x, y, and then use these two solutions to express (15.14.4) entirely in terms of s, t. We solve that equation, solve for s, t in terms of x, y, substitute into our last solution, and we are done! Of course, a computer algebra system is indispensable for these calculations.

a computer algebra system is indispensable for these calculations. The solution is  $F(x, y) = \frac{e^y}{1 + x - xe^y}$ . We then apply the first Möbius transformation  $z \mapsto 1/z$  which yields the generating function F(1/x, xy), and then  $z \mapsto z - 1$  which gives F(1/(x - 1), (x - 1)y) which simplifies to our result in the table. CHAPTER



# Recurrences with polynomial coefficients

How can we construct sequences of real-rooted polynomials  $\{f_i\}_{i=0}^{\infty}$  such that for all positive n the  $f_i$  satisfy the recurrence

$$f_{n}(x)g_{0}(x) + f_{n+1}(x)g_{1}(x) + \dots + f_{n+d}(x)g_{d}(x) = 0$$
(16.0.1)

where the  $g_i$  are polynomials in x? We say that the sequence  $\{f_i\}$  satisfies a *recursion with polynomial coefficients*. We will show several ways of doing constructing such sequences. Unfortunately, we don't know how to reverse the process. That is, given a sequence that satisfies a recursion with polynomial coefficients, prove that all the terms have all real roots.

**Example 16.1.** Here are two simple examples. If  $f_i = x^i$  then we have the recurrence

$$f_n(x) \cdot x - f_{n+1}(x) = 0.$$

A less trivial example is given by the Chebyshev polynomials  $U_k$  of the second kind. They satisfy the recurrence

$$U_{n}(x) - U_{n+1}(x) \cdot 2x + U_{n+2} = 0$$

In each of these cases the consecutive terms interlace, but this will not be true in general.

The Hermite polynomial  $H_n$  satisfies the recursion

$$H_{n+1} = 2xH_n - 2nH_{n-1}$$

This is *not* a recursion with polynomial coefficients since the factor (2n) depends on the index. However, they do satisfy a differential recursion with polynomial coefficients

$$\mathsf{H}_{\mathsf{n}} = 2\mathsf{x}\mathsf{H}_{\mathsf{n}-1} - \mathsf{H}_{\mathsf{n}-1}'.$$

Every sequence that satisfies a recursion with polynomial coefficients also satisfies a determinantal identity. This has nothing to do with the property of having real roots or not. Given a sequence as in (16.0.1) we have the identity

$$(g_0 \quad g_1 \quad \cdots \quad g_d) \begin{pmatrix} f_n & f_{n+1} & \cdots & f_{n+d} \\ f_{n+1} & f_{n+2} & \cdots & f_{n+d+1} \\ \vdots & & \vdots \\ f_{n+d} & f_{n+d+1} & \cdots & f_{n+2d} \end{pmatrix} = 0$$

If the  $g_i$  are not identically zero the determinant is zero:

$$\begin{vmatrix} f_{n} & f_{n+1} & \cdots & f_{n+d} \\ f_{n+1} & f_{n+2} & \cdots & f_{n+d+1} \\ \vdots & & \vdots \\ f_{n+d} & f_{n+d+1} & \cdots & f_{n+2d} \end{vmatrix} = 0$$

#### 16.1 The general construction

Many of our recursions arise from repeatedly multiplying by a polynomial. Here's a trivial example:

$$f_0 = f$$
$$f_n = g f_{n-1}$$

where f,  $g \in \mathbf{P}$ . More generally, we turn polynomial multiplication into matrix multiplication, and the characteristic polynomial becomes the recursion.

So, our construction requires

- 1. A d by d matrix M that might have polynomial entries, or perhaps even linear transformations as entries.
- 2. A vector v of polynomials.
- 3. For i = 0, 1, ..., all the entries of  $M^i v$  have all real roots.

If  $\sum_{0}^{d} \alpha_{i} \lambda^{i}$  is the characteristic polynomial of M and  $n = 0, 1, \dots$ , then

$$\sum_{i=0}^d \mathfrak{a}_i \left( M^{i+n} \nu \right) = 0.$$

Thus if we write

$$M^{i} v = (p_{i,1}, \ldots, p_{i,d})^{\dagger}$$

then each sequence of polynomials  $\{p_{i,k}\}_{i=1}^\infty$  has all real roots, and satisfies the same recursion

$$\sum_{i=0}^{d} a_i p_{i+n,k} = 0.$$

The vector v of polynomials can be

- 1. A collection of mutually interlacing polynomials.
- 2. All coefficients of a polynomial in  $\overline{\mathbf{P}}_2$ .
- 3. The initial coefficients of a function in  $\widehat{\mathbf{P}}_2$ .

# **16.2** Recursions from f(y) and g(x, y)

In our first construction we always get the same recursion; this is due to the fact that our matrix is lower triangular. We start with the following data

$$\begin{split} f(y) &= \sum_{0}^{d} \alpha_{i} y^{i} & \in P \\ g(x,y) &= \sum_{i} g_{i}(x) y^{i} & \in \overline{P}_{2} \\ f(y)g(x,y) &= \sum_{i} h_{i}(x) y^{i} \end{split}$$

and we have the relation

$$\begin{pmatrix} a_0 & 0 & 0 & \dots & 0 \\ a_1 & a_0 & 0 & \dots & 0 \\ a_2 & a_1 & a_0 & \dots & 0 \\ \vdots & & & \ddots & \\ a_d & a_{d-1} & a_{d-2} & \dots & a_0 \end{pmatrix} \begin{pmatrix} g_0 \\ g_1 \\ g_2 \\ \vdots \\ g_d \end{pmatrix} = \begin{pmatrix} h_0 \\ h_1 \\ h_2 \\ \vdots \\ h_d \end{pmatrix}$$

Since  $f(y)g(x, y) \in \overline{\mathbf{P}}_2$  it follows that all the coefficients  $h_0, \ldots, h_d$  have all real roots. We follow the outline in the previous section. The characteristic polynomial is  $(\lambda - a_0)^d$ . If we write

$$M^{\mathfrak{i}} \mathfrak{v} = (\mathfrak{p}_{\mathfrak{i},1},\ldots,\mathfrak{p}_{\mathfrak{i},d})^{\mathfrak{t}}$$

then all  $\{p_{i,k}\}$  satisfy

$$p_{n,k} - {d \choose 1} a_0 p_{n-1,k} + \dots + (-1)^d a_0^d p_{n-d,k} = 0$$

Note that the recursion is effectively independent of f if  $a_0 \neq 0$ , since we can rescale to make  $a_0 = 1$ . Here's an example where d = 3:

$$f = (y+1)^3 \qquad g = (x+2y+1)(x+3y+2)$$
$$M = \begin{pmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 & 0 \\ 3 & 3 & 1 & 0 \\ 1 & 3 & 3 & 1 \end{pmatrix}$$

$$\nu = \begin{pmatrix} x^2 + 3x + 2 \\ 5x + 7 \\ 6 \\ 0 \end{pmatrix} \qquad \qquad M\nu = \begin{pmatrix} x^2 + 3x + 2 \\ 3x^2 + 14x + 13 \\ 3x^2 + 24x + 33 \\ x^2 + 18x + 41 \end{pmatrix}$$
$$M^2\nu = \begin{pmatrix} x^2 + 3x + 2 \\ 6x^2 + 23x + 19 \\ 15x^2 + 75x + 78 \\ 20x^2 + 135x + 181 \end{pmatrix} \qquad M^3\nu = \begin{pmatrix} x^2 + 3x + 2 \\ 9x^2 + 32x + 25 \\ 36x^2 + 153x + 141 \\ 84x^2 + 432x + 474 \end{pmatrix}$$

The four sequences

$x^2 + 3x + 2$	$x^2 + 3x + 2$	$x^2 + 3x + 2$	$x^2 + 3x + 2$	
5x + 7	$3x^2 + 14x + 13$	$6x^2 + 23x + 19$	$9x^2 + 32x + 25$	
6	$3x^2 + 24x + 33$	$15x^2 + 75x + 78$	$36x^2 + 153x + 141$	
0	$x^2 + 18x + 41$	$20x^2 + 135x + 181$	$84x^2 + 432x + 474$	

have all real roots and satisfy the recurrence

$$p_n - 3p_{n-1} + 3p_{n-2} - p_{n-3} = 0.$$

#### **Recursions from** f(y, z) **and** g(x, y, z)16.3

Given a polynomial in  $P_3$  and  $\nu$  the vector of the first k coefficients of a polynomial in  $P_2$  we can construct a matrix that preserves such initial sequences. This gives us recursions of real rooted polynomials. We give the general construction, and then give some examples.

**Construction 16.2.** We are given  $f = \sum f_{i,j}(x) y^i z^j$  in  $\overline{P}_3$ , and g(x,y) = $\sum_{0}^{r} g_{i}(x)y^{i}$  is the first r + 1 terms of a function in  $\widehat{\mathbf{P}}_{2}$ .

We define

$$M = (f_{r-i,j})_{0 \leq i,j \leq r}$$
$$v_0 = (g_0, \dots, g_r)$$

By Theorem 24.11 we know that all polynomials

$$\mathsf{M}^{\mathsf{k}}\mathsf{v}_0 = \mathsf{v}_{\mathsf{k}} = (\mathsf{v}_{0,\mathsf{k}},\ldots,\mathsf{v}_{\mathsf{r},\mathsf{k}})$$

have all real roots. Suppose that M satisfies

$$\sum_{0}^{s} \alpha_{i} M^{i} = 0$$

Then the sequences

$$g_0 = v_{0,0}, v_{0,1}, v_{0,2}, \dots$$
  

$$g_1 = v_{1,0}, v_{1,1}, v_{1,2}, \dots$$
  

$$\dots$$
  

$$g_r = v_{r,0}, v_{r,1}, v_{r,2}, \dots$$

all satisfy the same recurrence

$$\sum_{i=0}^s \alpha_i p_i = 0$$

#### Example 16.3. We start with

$$\begin{vmatrix} x+1 & 1 & 1 \\ 1 & y & 1 \\ 1 & 1 & z \end{vmatrix} = yz(1+x) - y - z + (1-x) \in \overline{\mathbf{P}}_3$$

The matrix

$$\mathsf{M} = \begin{pmatrix} -1 & 1-x \\ 1+x & -1 \end{pmatrix}$$

preserves initial sequences of length two. M satisfies its characteristic polynomial

$$\lambda^2 + 2\lambda + x^2 = 0$$

If  $g_0(x) + y g_1(x) \in \overline{\mathbf{P}}_2$  then the polynomial sequences

$$\begin{pmatrix} -1 & 1-x \\ 1+x & -1 \end{pmatrix}^n \begin{pmatrix} g_0 \\ f_0 \end{pmatrix} = \begin{pmatrix} g_n \\ f_n \end{pmatrix}$$

are in  ${\bf P}$  and satisfy

$$g_n + 2 g_{n-1} + x^2 g_{n-2} = 0$$
  
$$f_n + 2 f_{n-1} + x^2 f_{n-2} = 0$$

Now we choose  $g_0 = -1$  and  $g_1 = x$ . Since  $g_0 + yg_1 = xy - 1 \in \overline{\mathbf{P}}_2$  we can apply the above construction

$$\nu = \begin{pmatrix} -1 \\ x \end{pmatrix} \qquad \mathsf{M}\nu = \begin{pmatrix} -x^2 + x + 1 \\ -2x - 1 \end{pmatrix} \qquad \mathsf{M}^2\nu = \begin{pmatrix} 3x^2 - 2x - 2 \\ -x^3 + 4x + 2 \end{pmatrix} \qquad ..$$

giving us the two sequences of polynomials

Consecutive  $g_i$ 's or  $f_i$ 's do not necessarily interlace, but they all are in **P**. However,  $g_i$  and  $f_i$  do interlace.

**Example 16.4.** The polynomial xy + xz + yz - 1 is in  $\overline{\mathbf{P}}_3$  since  $\begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}$  is negative subdefinite. For a particular choice of v we have nice formulas for the two

polynomial sequences.

$$\begin{split} \mathsf{M} &= \begin{pmatrix} x & -1 \\ 1 & x \end{pmatrix} \\ \mathsf{CP}(\mathsf{M}) &= y^2 - 2xy + x^2 + 1 \\ \mathsf{p}_n &= 2x\mathsf{p}_{n-1} - (x^2 + 1)\mathsf{p}_{n-2} \\ \upsilon &= (x, 1) \\ \mathsf{M}^k \upsilon &= (\mathsf{f}_k, \mathsf{g}_k) \\ \mathsf{f}_k &= \sum_i x^{n-2i} \binom{\mathsf{n}}{2i} (-1)^i \\ \mathsf{g}_k &= \sum_i x^{n-2i+1} \binom{\mathsf{n}}{2i-1} (-1)^{i+1} \end{split}$$

We know that  $f_k$  and  $g_k$  have all real roots since they are derived from the even and odd parts of  $(x-1)^n$ 

**Example 16.5.** In this example we construct a sequence of polynomials in two variables that satisfy a three term recursion. We begin with the Grace polynomial that is in  $\overline{\mathbf{P}}_4$ 

$$Grace_2 = 2x_1x_2 + (x_1 + x_2)(y_1 + y_2) + 2y_1y_2$$

Assume that  $f = f_0(x_1, x_2) + f_1(x_1, x_2)y_1 \in \overline{P}_3$ . Consider the coefficients of  $y_1$  in  $f \cdot Grace_2$ 

 $\begin{array}{ll} \text{Coefficient of } y_1^0: & [2x_1x_2+y_2(x_1+x_2)]f_0 \\ \text{Coefficient of } y_1^1: & [2x_1x_2+y_2(x_1+x_2)]f_1+[2y_2+x_1+x_2]f_0 \\ \text{Coefficient of } y_1^2: & [2y_2+x_1+x_2]f_1 \end{array}$ 

The coefficient of  $y_1^1$  is a polynomial in  $y_2$ , so the coefficients of  $y_2^0$  and  $y_2^1$  interlace:

Coefficient of 
$$y_2^0$$
:  $(x_1 + x_2)f_0 + 2x_1x_2f_1$   
Coefficient of  $y_2^1$ :  $2f_0 + (x_1 + x_2)f_1$ 

It follows that the matrix

$$\mathsf{M} = \begin{pmatrix} \mathsf{x}_1 + \mathsf{x}_2 & 2\mathsf{x}_1\mathsf{x}_2 \\ 2 & \mathsf{x}_1 + \mathsf{x}_2 \end{pmatrix}$$

preserves interlacing pairs of polynomials in two variables. As expected, the determinant,  $(x_1 - x_2)^2$ , is non-negative.

The sequence of vectors

$$\begin{pmatrix} f_0 \\ f_1 \end{pmatrix}$$
,  $M \begin{pmatrix} f_0 \\ f_1 \end{pmatrix}$ ,  $M^2 \begin{pmatrix} f_0 \\ f_1 \end{pmatrix}$ ,  $\cdots$ 

consists of polynomials in  $\overline{\mathbf{P}}_2$ . Write  $M^i \begin{pmatrix} f_0 \\ f_1 \end{pmatrix} = \begin{pmatrix} g_i \\ h_i \end{pmatrix}$ . The characteristic polynomial of M is

$$z^2 - 2(x_1 + x_2)z + (x_1 - x_2)^2$$

and since M satisfies its characteristic polynomial we have

$$M^{n+2} \cdot \nu - 2(x_1 + x_2)M^{n+1} \cdot \nu + (x_1 - x_2)^2M^n \cdot \nu = 0$$

for any vector v. Therefore, we have the recurrences

$$\begin{split} g_{n+2} &- 2(x_1+x_2)g_{n+1} + (x_1-x_2)^2 g_n = 0 \\ h_{n+2} &- 2(x_1+x_2)h_{n+1} + (x_1-x_2)^2 h_n = 0 \end{split}$$

If we want to construct examples then we must begin with  $g_0 \leftarrow h_0$ , and not with  $g_0, g_1$ . For example, if we take a degenerate case  $g_0 = 1$ ,  $h_0 = 0$  then the first few pairs are

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \begin{pmatrix} x_1 + x_2 \\ 2 \end{pmatrix}, \quad \begin{pmatrix} x_1^2 + 6x_1x_2 + x_2^2 \\ 4x_1 + 4x_2 \end{pmatrix}, \quad \begin{pmatrix} x_1^3 + 15x_1^2x_2 + 15x_1x_2^2 + x_2^3 \\ 6x_1^2 + 20x_1x_2 + 6x_2^2 \end{pmatrix},$$

It's not hard to see that we have an explicit formula

$$g_n(x_1, x_2) = \sum_{i=0}^n \binom{2n}{2i} x_1^i x_2^{n-i}$$

and the  $g_i$  satisfy the recurrence above with initial conditions  $g_0 = 1$  and  $g_1 = x_1 + x_2$ .

### **16.4** Recursions from $f(x, -D_y)$ and g(x, y)

We know that if  $f(x, y) \in \overline{P}_2$  then  $f(x, -D_y)$  maps  $\overline{P}_2 \longrightarrow \overline{P}_2$ . This guarantees that our recursions will consist of polynomials with all real roots.

We assume

$$\begin{split} f(x,y) &= \sum_{i} f_{i}(x)y^{i} & \in \widehat{P}_{2} \\ f(0,0) &\neq 0 \\ g(x,y) &= \sum_{i=0}^{d} g_{i}(x)y^{i} & \in \mathtt{P}_{2} \end{split}$$

The action of  $f(x, -D_y)$  never decreases y-degree, so it determines a d + 1 by d + 1 matrix acting on the coefficients of polynomials in  $P_2(d)$ .

#### Example 16.6. We assume

$$f(x,y) = (x+y)^3$$
  $g(x,y) = g_0(x) + g_1(x)y + g_2(x)y^2$ 

The action of  $f(x, -D_y)$  on g is given by the matrix

$$\begin{pmatrix} x^3 & -3x^2 & 6x \\ 0 & x^3 & -6x^2 \\ 0 & 0 & x^3 \end{pmatrix}$$

This gives recurrences of the form

$$p_n - 3x^3p_{n-1} + 3x^6p_{n-2} - x^9p_{n-3} = 0$$

**Example 16.7.** In this case we let  $f = e^{-xy}$ , and  $g = \sum_{i=1}^{4} g_i(x)y^i$ . The matrix is

/1	$-\mathbf{x}$	$\chi^2$	$-\chi^3$	$x^4$
0	1	-2x	$3x^2$	$-4x^3$
0	0	1	-3x	6x <sup>2</sup>
0	0	0	1	-4x
0	0	0	0	1 /

The characteristic polynomial is  $(\lambda - 1)^5$ , so these recurrences are of the form

$$p_n - 5p_{n-1} + 10p_{n-2} - 10p_{n-3} + 5p_{n-4} - p_{n-5} = 0.$$

# **16.5** Recurrences from $f(-\mathbf{D}_x, y)$ and g(x, y)

If we use  $f(-\mathsf{D}_x,y)$  instead of  $f(x,-\mathsf{D}_y)$  then we get recurrences involving derivatives. We assume

$$\begin{split} f(x,y) &= \sum_{i} f_{i}(x)y^{i} & \in \widehat{P}_{2} \\ g(x,y) &= \sum_{i=0}^{d} g_{i}(x)y^{i} & \in P_{2} \end{split}$$

We give a very simple example - note that the degree of the recurrence comes from the number of coefficients we consider, and not from the degree of f.

#### Example 16.8. We let

$$f(x,y) = y - D_x$$
  $g = \sum g_i(x)y^i$ 

We consider only the first four coefficients of g, so the action of  $y - D_x$  is

$$-g_0' + y(g_0 - g_1') + y^2(g_1 - g_2') + y^3(g_2 - g_3').$$

As a matrix this is

$$\begin{pmatrix} -\mathsf{D}_{\mathsf{x}} & 0 & 0 & 0\\ 1 & -\mathsf{D}_{\mathsf{x}} & 0 & 0\\ 0 & 1 & -\mathsf{D}_{\mathsf{x}} & 0\\ 0 & 0 & 1 & -\mathsf{D}_{\mathsf{x}} \end{pmatrix} \begin{pmatrix} \mathsf{g}_{0}\\ \mathsf{g}_{1}\\ \mathsf{g}_{2}\\ \mathsf{g}_{3} \end{pmatrix}$$

The characteristic polynomial is  $(\lambda + D_x)^4$ , so the recurrence is

$$p_n + 4p'_{n-1} + 6p^{(2)}_{n-2} + 4p^{(3)}_{n-3} + p^{(4)}_{n-4} = 0.$$

#### 16.6 Recurrences from mutually interlacing polynomials

If a matrix preserves mutually interlacing polynomials then we get recursive sequences of polynomials in **P**.

Example 16.9. The matrix and initial vector

$$M = \begin{pmatrix} 1 & x+1 & x+1 \\ 1 & 1 & x+1 \\ 1 & 1 & 1 \end{pmatrix} \qquad \qquad \nu = \begin{pmatrix} x-1 \\ x \\ x+1 \end{pmatrix}$$

determine three polynomial sequences  $M^i\nu=(f_i,g_i,h_i)$  that satisfy the recurrence

$$p_n = 3 p_{n-1} + 3 x p_{n-2} + x^2 p_{n-3}.$$

To find the initial terms we need to compute Mv and  $M^2v$ .

$$M\nu = \begin{pmatrix} 2x^2 + 4x \\ x^2 + 4x \\ 3x \end{pmatrix} \qquad M^2\nu = \begin{pmatrix} x^3 + 10x^2 + 11x \\ 6x^2 + 11x \\ 3x^2 + 11x \end{pmatrix}$$

It follows that the following sequence of polynomials has all real roots. It also appears that  $f_i \leq f_{i-1}$  but that doesn't follow from the general construction, and I don't know how to prove it.

$$\begin{split} h_0 &= x + 1 \\ h_1 &= 3x \\ h_2 &= 3x^2 + 11x \\ h_n &= 3h_{n-1} + 3xh_{n-2} + x^2h_{n-3} \end{split}$$

**Example 16.10.** In this example we start with the recurrence, and find a matrix that preserves mutually interlacing polynomials. Consider the sequence

$$\begin{aligned} q_0 &= q_1 = q_2 = \dots = q_d = 1 \\ q_k &= q_{k-1} + xq_{k-d} \text{ for } k > d \\ q_{d+1} &= 1 + x, \quad q_{d+2} = 1 + 2x, \quad \dots \quad q_{2d+1} = 1 + (d+1)x \end{aligned}$$

Rewrite the recurrence in matrix terms:

/1	0	0	0		x)	$(q_{k-1})$		$(q_k)$	
1	0	0	0		0	$q_{k-2}$		$q_{k-1}$	
0	1	0	0		0		_	:	
:		•.	۰.		:	•		•	
·		•	•		·				
$\setminus 0$	0			1	0/	$\langle q_{k-d} \rangle$		$\langle q_{k-d+1} \rangle$	

The d+1 by d+1 matrix preserves mutually interlacing polynomials. Since the vector  $(q_{d+1}, \ldots, q_{2d+1})$  is mutually interlacing, we can repeatedly apply the Lemma to conclude that  $(q_k, \ldots, q_{k+d})$  is a vector of mutually interlacing vectors. In particular, all  $q_n$  have all real roots.

If a two by two matrix preserves interlacing polynomials then we get a recurrence. This is really just a special case of a general matrix preserving the first k coefficients of a polynomial, and also a special case of a matrix preserving mutual interlacing.

**Example 16.11.** The matrix  $\begin{pmatrix} 1 & x \\ 1 & 0 \end{pmatrix}$  preserves interlacing. If we let  $\begin{pmatrix} 1 & x \\ 1 & 0 \end{pmatrix}^n \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} p_n \\ p_{n-1} \end{pmatrix}$ 

then all  $p_n \in \mathbf{P}$ . The characteristic polynomial is  $\lambda^2 - \lambda - x$ , so we have the recurrence

$$p_n = p_{n-1} + x p_{n-2}$$

#### **16.7** Alternative approaches

There are many specialized ways to prove facts about recurrences. [54] has some very interesting results about recurrences with polynomial coefficients.

Of course, there is a 1-1 correspondence between recurrences with polynomial coefficients and rational functions. Here we start with a rational function.

If we write

$$\frac{1}{1 - (x + y)^n} = \sum_{j=0}^{\infty} \frac{h_{n,j}(x)}{(1 - x^n)^{j+1}} y^j$$

then we will show that the roots of the coefficients  $h_{n,j}$  lie on n equally spaced rays through the origin. For example, if n = 2 we have

$$\frac{1}{1-(x+y)^2} = \frac{1}{1-x^2} + \frac{2x}{(1-x^2)^2}y + \frac{3x^2+1}{(1-x^2)^3}y^2 + \frac{4(x^3+x)}{(1-x^2)^4}y^3 + \cdots$$

and all the roots lie on the imaginary axis. There is a simple expression for these coefficients

$$\frac{1}{1-(x+y)^n} = \sum_{i} (x+y)^{ni} = \sum_{j} y^j \cdot \sum_{i=0}^{\infty} \binom{ni}{j} x^{ni-j}.$$
The coefficient of  $y^j$  is  $x^{-j}g(x^n)$  where

$$g_{n,j}(x) = \sum_{i=0}^{\infty} \binom{ni}{j} x^i = \frac{f_{n,j}(x)}{(1-x)^{j+1}}.$$

The series converges for |x| < 1. If x is positive then the series is positive or divergent. Thus,  $g_{n,j}$  has no positive roots. Here are some values of the sum

$$\begin{array}{c|ccccc} n=1 & n=2 & n=3 \\ \hline j=0 & \frac{1}{1-x} & \frac{1}{1-x} & \frac{1}{1-x} \\ j=1 & \frac{x}{(1-x)^2} & \frac{2x}{(1-x)^2} & \frac{3x}{(1-x)^2} \\ j=2 & \frac{x^2}{(1-x)^3} & \frac{3x^2+x}{(1-x)^3} & \frac{6x^2+3x}{(1-x)^3} \end{array}$$

We will show that  $f_{n,j} \in \mathbf{P}$ , which establishes the claim. We write  $g_{nj}(x)$  in terms of a differential operator. First, let  $\binom{nx}{j} = \sum a_k x^k$ .

$$\begin{split} g_{n,j}(x) &= \sum_{i} \binom{ni}{j} x^{i} = \sum_{i,k} \alpha_{k} i^{k} x^{i} \\ &= \sum_{i,j} \alpha_{k} (x\mathsf{D})^{k} x^{i} = \bigl(\sum_{i} (x\mathsf{D})^{k} \alpha_{k}\bigr) \frac{1}{1-x} = \binom{nx\mathsf{D}}{j} \frac{1}{1-x}. \end{split}$$

It simplifies the argument if we eliminate the denominator, so we will show that  $F_j \in \textbf{P}$  where

$$(\mathbf{n}\mathbf{x}\mathbf{D})(\mathbf{n}\mathbf{x}\mathbf{D}-1)\cdots(\mathbf{n}\mathbf{x}\mathbf{D}-\mathbf{j}+1)\frac{1}{1-\mathbf{x}}=\frac{\mathbf{F}_{\mathbf{j}}}{(1-\mathbf{x})^{\mathbf{j}+1}}$$

This yields the recurrence

$$F_{n,j+1} = \left[ \left( (j+1)n + j \right) x + j \right] F_{n,j} - x(x-1)F'_{n,j}$$

This recurrence is of the form

$$p_n = (ax+b)p_{n-1} - x(x-1)p'_{n-1}$$

Since x(x - 1) is positive at the roots of  $p_n$  it follows that all  $f_j$  have all real roots and all positive coefficients.

CHAPTER



# Matrices

The theme of this chapter is that there are several ways to represent sequences of interlacing polynomials by Hermitian matrices. There is a 1-1 correspondence between Hermitian matrices and interlacing sequences of monic polynomials. Since orthogonal polynomials can be realized as the characteristic polynomials of certain Hermitian matrices, many results about orthogonal polynomials can be generalized to Hermitian matrices.

# 17.1 Basics

We first recall some basic facts about matrices. A\* denotes the conjugate transpose of A. The matrix A is *Hermitian* if  $A = A^*$ . Now we are only considering real polynomials, so we could restrict ourselves to symmetric matrices, but it is no harder to consider Hermitian matrices. A matrix U is unitary if  $UU^* = I$ , where I is the identity matrix. A Hermitian matrix has all real eigenvalues. The spectral theorem says that there is a unitary matrix U and a diagonal matrix  $\Lambda$  consisting of the eigenvalues of A such that  $A = U\Lambda U^*$ . If all entries of A are real, then U can be chosen to be real, so that it is an orthogonal matrix.

The characteristic polynomial of A is |xI - A|, and is written CP(A). The set of roots of the characteristic polynomial of A is written  $\lambda(A)$ . We say that  $A \ll B$  if CP(A)  $\ll$  CP(B),  $A \ll B$  iff CP(A)  $\ll$  CP(B), and so on. A Hermitian matrix is *positive definite* if all of its eigenvalues are positive, and is *positive semidefinite* if the eigenvalues are all non-negative.

If the matrix A is invertible then interlacing properties of A determine interlacing properties of  $A^{-1}$ .

- If A is invertible and  $\lambda(A) = (a_1, \dots, a_n)$ , then  $\lambda(A^{-1}) = (\frac{1}{a_1}, \dots, \frac{1}{a_n})$ .
- If A is positive definite and  $A \ll B$  then  $B^{-1} \ll A^{-1}$ .
- If A is positive definite and  $A \leq B$  then  $A^{-1} \leq B^{-1}$ .

• The characteristic polynomial of A<sup>-1</sup> is

$$|xI - A^{-1}| = |-A^{-1}| |I - xA| = |-A^{-1}| |(1/x)I - A|x^{n}$$

where A is n by n. Since the reverse of a polynomial f is  $f^{REV} = x^n f(1/x)$ , we can write

$$CP(A^{-1}) = \frac{1}{|-A|} (CP(A))^{REV}$$

# 17.2 Principle submatrices

In this section we establish the important result that Hermitian matrices correspond to sequences of interlacing polynomials.

**Definition 17.1.** For n by n matrix M , and index sets  $\alpha, \beta \subset \{1, ..., n\}$ , the (sub)matrix that lies in the rows indexed by  $\alpha$  and columns indexed by  $\beta$  is denoted M{ $\alpha, \beta$ }. If  $\alpha = \beta$  then we write M{ $\alpha$ }; such a submatrix is called a *principle* submatrix (§ 10.10.14). Sometimes we want to refer to a submatrix via the deletion of rows; in this case we let M[ $\alpha$ ] be the submatrix resulting from the deletion of the rows and columns not listed in  $\alpha$ . The size  $|\alpha|$  of an index set  $\alpha$  is its cardinality.

We know that if A is a principle submatrix of B then  $B \leq A$ .

Suppose that f is a polynomial with roots  $roots(f) = (a_1, ..., a_n)$ . Set  $\Lambda$  equal to the diagonal matrix whose elements are  $(a_1, ..., a_n)$ , and let  $y^* = (y_1, ..., y_n)$ . The following calculation shows the relation between the characteristic polynomial of  $\Lambda$  and an extension of  $\Lambda$ . Suppose that

$$A = \begin{pmatrix} \Lambda & \vdots & y \\ \dots & \dots \\ y^* & \vdots & a \end{pmatrix}$$

The characteristic polynomial of A is

$$det(xI - A) = det \begin{pmatrix} xI - A & \vdots & -y \\ \dots & \dots & \dots \\ -y^* & \vdots & x - a \end{pmatrix}$$

$$= det \begin{pmatrix} I & \vdots & 0 \\ \dots & \dots & \dots \\ [(xI - A)^{-1}y]^* & \vdots & 1 \end{pmatrix} \begin{pmatrix} xI - A & \vdots & -y \\ \dots & \dots & \dots \\ -y^* & \vdots & x - a \end{pmatrix} \times$$

$$\begin{pmatrix} I & \vdots & (xI - A)^{-1}y \\ \dots & \dots & \dots \\ 0 & \vdots & I \end{pmatrix}$$

$$= det \begin{pmatrix} xI - A & \vdots & 0 \\ \dots & \dots & \dots \\ 0 & \vdots & (x - a) - y^*(xI - A)^{-1}y \end{pmatrix}$$

$$= [(x - a) - y^*(xI - A)^{-1}y] det(xI - A)$$

$$= (x - a)f(x) - \sum_{i=1}^{n} y_i^2 \frac{f(x)}{x - a_i}$$

If  $p_i$  is the characteristic polynomial of  $A\{1, ..., i\}$ , then from repeated applications of Theorem 1.61 we have a sequence of interlacing polynomials

$$\mathfrak{p}_{\mathfrak{n}} \underline{\lessdot} \mathfrak{p}_{\mathfrak{i}-1} \underline{\lessdot} \dots \underline{\lessdot} \mathfrak{p}_0$$

Conversely, we can use the above calculation to show that given the  $p_i$ 's, we can find a Hermitian matrix determining them.

**Theorem 17.2.** Suppose that  $p_0, p_1, \ldots, p_n$  is a sequence of polynomials such that the degree of  $p_i$  is i, and  $p_i \leq p_{i-1}$  for  $1 \leq i \leq n$ . There is a Hermitian matrix A such that the characteristic polynomial of  $A\{1, \ldots, i\}$  is a multiple of  $p_i$ , for  $0 \leq i \leq n$ .

*Proof.* We prove this by induction on n. The case n = 1 is clear, so assume that A is an n by n Hermitian matrix such that the characteristic polynomial of  $A\{1, ..., i\}$  is a multiple of  $p_i$ , for  $0 \le i \le n$ . Choose an orthogonal matrix U so that UAU\* is the diagonal matrix  $\Lambda$  whose diagonal consists of the roots of  $p_n$ . We need to find a number b and vector  $z^* = (z_1, ..., z_n)$  so that the characteristic polynomial of

$$\begin{pmatrix} A & \vdots & z \\ \dots & \dots \\ z^* & \vdots & b \end{pmatrix}$$
(17.2.1)

is equal to a multiple of  $p_{n+1}$ . Write

$$B = \begin{pmatrix} U & \vdots & 0 \\ \cdots & \cdots & 0 \\ 0 & \vdots & 1 \end{pmatrix} \begin{pmatrix} \Lambda & \vdots & z \\ \cdots & \cdots & z^* & \vdots & b \end{pmatrix} \begin{pmatrix} U^* & \vdots & 0 \\ \cdots & \cdots & 0 \\ 0 & \vdots & 1 \end{pmatrix} = \begin{pmatrix} \Lambda & \vdots & y \\ \cdots & \cdots & y^* \\ y^* & \vdots & a \end{pmatrix}$$
(17.2.2)

The characteristic polynomial of A is  $cp_n$  for some constant c. The characteristic polynomial of (17.2.1) is the same as the characteristic polynomial B in (17.2.2). Since  $p_{n+1} \leq p_n$ , we can apply Lemma 1.20 and find an a and  $y_i^2$  such that

$$p_{n+1} = (x-a)p_n - \sum y_i^2 \frac{p_n(x)}{x-a_i}$$

where  $roots(p_n) = (a_1, ..., a_n)$ . The desired matrix is  $U^*BU$ .

If  $p_{i+1} < p_i$  for all i then all of the coefficients  $y_i^2$  are non-zero. In this case there are  $2^n$  choices of  $y_i$ 's, and so there are  $2^n$  different A's.

**Example 17.3.** The construction of a matrix from a sequence of characteristic polynomials involves solving many equations. Here are a few examples of n by n matrices M where we are given a polynomial f and the determinant of the first i rows and columns is a constant multiple of  $f^{(i)}$ .

1. If 
$$f = (x - a)(x - b)$$
 then  $M = \begin{pmatrix} \frac{a+b}{2} & \frac{a-b}{2} \\ \frac{a-b}{2} & \frac{a+b}{2} \end{pmatrix}$   
2. If  $f = (x - 1)(x - 2)(x - 3)$  then  $M = \begin{pmatrix} 2 & \frac{-1}{\sqrt{3}} & 0 \\ \frac{-1}{\sqrt{3}} & 2 & \frac{-2}{\sqrt{3}} \\ 0 & \frac{-2}{\sqrt{3}} & 2 \end{pmatrix}$   
3. If  $f = x^2(x - 1)$  then  $M = \frac{1}{3} \begin{pmatrix} 1 & -1 & -1 \\ -1 & 1 & 1 \\ -1 & 1 & 1 \end{pmatrix}$ 

**Example 17.4.** If a Hermitian matrix H is partitioned into diagonal blocks, and the polynomials determined by each of these blocks strictly interlace, we do not have strict interlacing for the polynomials determined by H. Assume that A is an r by r Hermitian matrix determining the polynomial sequence  $p_r < \cdots < p_1$ , and B is an s by s Hermitian matrix determining the polynomial sequence g<sub>s</sub>  $< \cdots < q_1$ . The partitioned matrix

$$\begin{pmatrix} A & \vdots & 0 \\ \cdots & \cdots & 0 \\ 0 & \vdots & B \end{pmatrix}$$

、

,

determines the polynomial sequence

$$p_1 \! \geqslant \! p_2 \! \geqslant \cdots p_r \underline{\geqslant} p_r \, q_1 \! \ge \! p_r \, q_2 \! \ge \cdots \underline{\geqslant} p_r \, q_s$$

If a Hermitian matrix M is diagonal with diagonal entries  $(a_1, ..., a_n)$ , then the characteristic polynomial is  $f(x) = (x - a_1) \cdots (x - a_n)$ . The characteristic polynomial of a single row/column deletion M[i] equals  $f(x)/(x - a_i)$ . Since these are the polynomials occurring in quantitative sign interlacing it is not surprising that the characteristic polynomials of single deletions are also important. For instance, an immediate consequence of Theorem 1.61 is that for any choice of positive  $b_i$  the polynomials

$$\sum_{i=1}^{n} b_i \ CP(M[i])$$

have all real roots.

Analogous to the formula

$$\frac{\mathrm{d}}{\mathrm{d}x}f(x) = \sum_{i=1}^{n} \frac{f(x)}{x - a_{i}}$$

we have

$$\frac{d}{dx}\operatorname{CP}(M) = \sum_{i=1}^n \operatorname{CP}(M[i])$$

This can be established by considering the derivative of only the terms in the expansion of the determinant that have diagonal entries in them.

**Example 17.5.** It is not the case that the characteristic polynomials of the single deletion submatrices of an n by n matrix span an n-dimensional space. The matrix  $M = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  has eigenvalues 1, -1 but M[1] = M[2] = (0), so they have the same characteristic polynomial, x. More generally, for any symmetric C with distinct eigenvalues none of which are 1 or -1, the matrix M below has M[1] = M[2], so the characteristic polynomials do not span an n-dimensional space.

$$\begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & C \end{pmatrix}$$

An n by n matrix M has  $2^n$  principle submatrices. Their characteristic polynomials can be combined to form polynomials that are in  $\overline{P}_{n+1}$ .

**Lemma 17.6.** Let M be an n by n symmetric matrix. For each  $\alpha \subset \{1, ..., n\}$  let  $\mathbf{x}^{\alpha} = \prod_{i \in \alpha} x_i$ . Then the polynomials below are in  $\overline{\mathbf{P}}_{n+1}$ .

$$\sum_{\substack{\alpha \subset \{1, \dots, n\} \\ \substack{\alpha \subset \{1, \dots, n\}}} \operatorname{CP}(\mathsf{M}\{\alpha\}) \mathbf{x}^{\alpha}$$

*Proof.* The polynomial f(x, x) in question is nothing other than the characteristic polynomial of M + D where D is the diagonal matrix whose diagonal is  $(x_1, \ldots, x_n)$ . It remains to see that  $f \in \overline{\mathbf{P}}_{n+1}$ . Consider  $g_{\varepsilon} = CP(M + D + \varepsilon zI)$ where  $z = x_1 + \cdots + x_n$ . The homogeneous part of  $g_{\varepsilon}$  is  $\prod_i (x + x_i + \varepsilon (x_1 + \cdots + x_n))$  and for  $\varepsilon > 0$  this is in  $\mathbf{P}_{n+1}$ . If we substitute values for  $x_1, \ldots, x_n$ then the resulting polynomial is the characteristic polynomial of a symmetric matrix and so is in  $\mathbf{P}$ . Since  $\lim_{\varepsilon \to 0^+} g_{\varepsilon} = f$  it follows that f is in the closure of  $\mathbf{P}_{n+1}$ .

The second assertion follows from the first by setting x = 0.

For example, if  $M = \begin{pmatrix} a & b \\ b & c \end{pmatrix}$  then the two polynomials are

$$det \begin{vmatrix} a-x-x_1 & b \\ b & c-x-x_2 \end{vmatrix} = (x-a)(x-c) - b^2 + (x-a)x_2 + (x-c)x_1 + x_1x_2 det \begin{vmatrix} a-x_1 & b \\ b & c-x_2 \end{vmatrix} = (ac-b^2) - ax_2 - cx_1 + x_1x_2$$

# 17.3 The Schur complement

The Schur complement of a matrix plays a role that is similar to contraction for a graph. Although the initial definition is unintuitive, there is a simple interpretation in terms of deletion in the inverse that is easy to understand.

**Definition 17.7.** For an n by n matrix M , and index sets  $\alpha$ ,  $\beta \subset \{1, ..., n\}$  where a and  $\beta$  have the same size, the Schur complement of the (sub)matrix that lies in the rows indexed by  $\alpha$  and columns indexed by  $\beta$  is

$$A/[\alpha,\beta] = A\{\alpha',\beta'\} - A\{\alpha',\beta\}A\{\alpha,\beta\}^{-1}A\{\alpha,\beta'\}$$

If  $\alpha = \beta$  then we write  $M/[\alpha]$ . Notice that if  $\alpha$  has size k, then the Schur complement has dimension n - k.

**Example 17.8.** If M is diagonal with diagonal entries  $(a_1, ..., a_n)$ , and  $\alpha \subset \{1, ..., n\}$ , then the Schur complement  $M/[\alpha]$  is the diagonal matrix  $M\{\alpha'\}$ .

The Schur complement may be computed by taking the inverse, deleting appropriate rows and columns, and taking the inverse of the resulting matrix.

**Theorem 17.9 ([5]).** *If* A *is invertible, then*  $A/[\alpha, \beta] = (A^{-1}\{\beta', \alpha'\})^{-1}$ 

An interesting property is that two disjoint deletions commute.

**Theorem 17.10 ([5]).** *If*  $\alpha_1, \alpha_2, \beta_1, \beta_2$  *are index sets with*  $|\alpha_1| = |\beta_1|, |\alpha_2| = |\beta_2|$ , and  $\alpha_1 \cap \alpha_2 = \beta_1 \cap \beta_2 = \emptyset$  then

$$\left(M/[\alpha_1,\beta_1]\right)/[\alpha_2,\beta_2] = \left(M/[\alpha_2,\beta_2]\right)/[\alpha_1,\beta_1] = M/[\alpha_1\cup\alpha_2,\beta_1\cup\beta_2]$$

We can now show that Schur complements preserve interlacing.

**Theorem 17.11** ([160]). *If* A *is a definite (positive or negative) Hermitian matrix, and if*  $|\alpha| = |\beta| = 1$ *, then* A  $\leq A/[\alpha, \beta]$ *.* 

*Proof.* Since  $A^{-1}$  is Hermitian and its principle submatrices interlace, we get  $A^{-1} \leq A^{-1}\{\beta', \alpha'\}$ . Since all eigenvalues of A have the same sign, when we take inverses interlacing is preserved:

$$\mathsf{A} = \left(\mathsf{A}^{-1}\right)^{-1} \leq \left(\mathsf{A}^{-1}\{\beta',\alpha'\}\right)^{-1} = \mathsf{A}/[\alpha,\beta]$$

**Corollary 17.12.** If M is non-singular definite Hermitian, and we set  $p_k = CP(M/[\{1..k\}])$  then

$$\mathfrak{p}_{\mathfrak{n}} \underline{\lessdot} \mathfrak{p}_{k-1} \underline{\lessdot} \dots \underline{\lessdot} \mathfrak{p}_{1}$$

*Proof.* Since  $(M/[\{1..k\}])/[\{k+1\}] = M/[\{1..k+1\}]$  by Theorem 17.10, we can apply Theorem 17.11.

Another simple consequence is that we can realize polynomial sequences by characteristic polynomials of Schur complements.

**Theorem 17.13.** Suppose that  $p_0, p_1, \ldots, p_n$  is a sequence of polynomials such that the degree of  $p_i$  is i, and  $p_i \leq p_{i-1}$  for  $1 \leq i \leq n$ . There is a Hermitian matrix A such that the characteristic polynomial of  $A/[\{1, \ldots, i\}]$  is a constant multiple of  $p_i$ , for  $0 \leq i \leq n$ .

*Proof.* Let B be a Hermitian matrix such that  $CP(B\{\{1..k\}\}) = (p_i)^{REV}$ .  $B^{-1}$  is the desired matrix.

## **17.4** Families of matrices

We consider properties of the eigenvalues of continuous families of Hermitian matrices. We begin with a definition of *a locally interlacing family* of matrices.

**Definition 17.14.** If  $\{H_t\}$  is a continuous family of Hermitian matrices such that the family of their characteristic polynomials is a locally interlacing family, then we say that  $\{H_t\}$  is a *locally interlacing family*.

If the entries of a Hermitian matrix are functions of a parameter, then we can determine the derivatives of the eigenvalues with respect to this parameter. This result is known as the Hellmann-Feynman theorem:

**Theorem 17.15.** Let H(x) be an n by n Hermitian matrix whose entries have continuous first derivatives with respect to a parameter x for  $x \in (a, b)$ . Let  $\lambda(H(x)) = (\lambda_1(x), \ldots, \lambda_n(x))$  be the eigenvalues of H(x), with corresponding eigenvectors  $u_1, \ldots, u_n$ . If H' is the matrix formed by the derivatives of the entries of H(x), then for  $j = 1, \ldots, n$ 

$$\frac{d\lambda_i}{dx} = \frac{u_i^* H' u_i}{u_i^* u_i}$$

*Proof.* More generally, we consider an inner product for which  $(H_t u, v) = (u, H_t v), (u, v) = (v, u)$ , and (v, v) > 0 if  $v \neq 0$ . From the fact that the  $u_i$  are eigenvectors, we find that for any  $s \neq t$ 

$$\begin{aligned} (H_{t}u_{i}(t), u_{i}(s)) &= \lambda_{i}(t) (u_{i}(t), u_{i}(s)) \\ (H_{s}u_{i}(t), u_{i}(s)) &= (H_{s}u_{i}(s), u_{i}(t)) \\ &= \lambda_{i}(s) (u_{i}(t), u_{i}(s)) \end{aligned}$$

Subtracting and dividing by t - s gives the difference quotient

$$\left(\frac{\mathsf{H}_{\mathsf{t}}-\mathsf{H}_{\mathsf{s}}}{\mathsf{t}-\mathsf{s}}\mathsf{u}_{\mathsf{i}}(\mathsf{t}),\mathsf{u}_{\mathsf{i}}(\mathsf{s})\right) = \frac{\lambda_{\mathsf{t}}-\lambda_{\mathsf{s}}}{\mathsf{t}-\mathsf{s}}(\mathsf{u}_{\mathsf{i}}(\mathsf{t}),\mathsf{u}_{\mathsf{i}}(\mathsf{s}))$$

The result follows by taking the limit as s goes to t.

**Corollary 17.16.** If H(x) is an n by n Hermitian matrix whose entries have continuous first derivatives with respect to a parameter x for  $x \in (a, b)$ , and H'(x) is positive definite for all x, then  $\{H_t\}$  is a locally interlacing family.

**Example 17.17.** Suppose that H is Hermitian. Consider the family  $\{e^{tH}\}$ . The derivative of this family is

$$\frac{\mathrm{d}}{\mathrm{dt}}\left(e^{\mathrm{tH}}\right) = \mathrm{H}e^{\mathrm{tH}}$$

If v is an eigenvector of H with eigenvalue a, then  $He^{tH}v = H(e^{at}v) = ae^{at}v$ and so v is an eigenvector of H' with eigenvalue  $ae^{at}$ . These eigenvalues are increasing if and only if a is positive, so we conclude that  $\{e^{tH}\}$  is locally interlacing if and only if H is positive definite.

**Corollary 17.18.** If  $\{H_t\}$  is a family of Hermitian matrices such that  $\frac{d}{dt}H_t$  is strictly totally positive, then the families of characteristic polynomials of any principle submatrix are locally interlacing.

*Proof.* Since every principle submatrix of H' is positive definite, the result follows from Corollary 17.16.

The next result is known as the *monotonicity theorem*, since it states that adding a positive definite matrix increases all the eigenvalues.

**Corollary 17.19.** *If* A, B *are Hermitian matrices, and* B *is positive definite, then the family*  $\{A + tB\}$  *is locally interlacing.* 

*Proof.* The derivative  $\frac{d}{dt}(A + tB)$  is simply B.

**Example 17.20.** Consider the family  $\{H + tI\}$ , where H is Hermitian. Since I is positive definite, it is an locally interlacing family. If the characteristic polynomial of H is f(x), then

$$||xI - (H + tI)| = |(x - t)I - H| = f(x - t)$$

Another elementary family is given by {tH}. Since |xI - tH| = f(x/t) it is locally interlacing.

We can relate interlacing of polynomial sequences with their Hermitian matrices, but first we need to recall the *positive definite ordering* (p. 458) of Hermitian matrices.

**Lemma 17.21.** Suppose that A, B are Hermitian. They determine polynomial sequences  $p_i = ||xI - A\{1..i\}|$  and  $q_i = ||xI - B\{1..i\}|$ . If  $B \prec A$  then  $p_i \prec q_i$  for  $1 \leqslant i \leqslant n$ .

*Proof.* Consider the family  $M_t = tA + (1-t)B$ . The derivative is A - B which is positive definite, so it is a locally interlacing family. Since  $M_0 = B$  and  $M_1 = A$  it follows that the roots of  $M_t$  are increasing as a function of t, so  $p_i \prec q_i$ .

# 17.5 Permanents

If  $A = (a_{ij})$  is a matrix then the *permanent* of A, written per(A), is the determinant without alternating signs:

$$per(A) = \sum_{\sigma \in S_n} \prod_{i=1}^n a_{i,\sigma(i)}$$

Unlike the determinant, polynomials such as per(xI+C) where C is symmetric almost never have all real roots. If we define J to be the all 1 matrix, then there is a beautiful conjecture [76]

**Conjecture 17.22.** If  $A = (a_{ij})$  is a real n by n matrix with non-negative entries which are weakly increasing down columns then the permanent of A + xJ has all real roots.

The permanents of principle submatrices of A + xJ do not interlace the permanents of A + xJ, but we do have common interlacing:

**Lemma 17.23.** Suppose that A satisfies the conditions above, and the conjecture is true. Then per(A + xJ) and per((A + xJ)[1]) have a common interlacing.

*Proof.* If we let M = A + xJ so that  $M = M_0$  where

$$M_{\alpha} = \begin{vmatrix} a_{11} + x + \alpha & v + x \\ w + x & A[1] + xJ[1] \end{vmatrix}$$

By linearity of the permanent we have

$$per(M_{\alpha}) = per \begin{vmatrix} a_{11} + x & v + x \\ w + x & A[1] + xJ[1] \end{vmatrix} + \alpha per(A[1] + xJ[1])$$

If we choose  $\alpha$  to be negative then the conjecture implies that  $per(M_{\alpha})$  has all real roots. Consequently,  $per(M) + \alpha per(M[1])$  has all real roots for all negative  $\alpha$ . This implies that per(M) and per(M[1]) have a common interlacing.

*Remark* 17.24. The proof can be easily modified to show that if we remove the first row and any column from A then the permanents have a common interlacing. Empirically, it appears that all the submatrices have a common interlacing.

#### **17.6** Matrices from polynomials

If A is a constant matrix and f(x) is a polynomial then f(xA) is a matrix whose entries are polynomials in x. We look for conditions on A or f that imply that all entries of f(xA) are in **P**.

A simple case is when A is a diagonal matrix  $diag(a_1, ..., a_n)$ . The resulting matrix f(xA) is a diagonal matrix. To check this, write  $f(x) = \sum b_i x^i$ :

$$\begin{split} f(xA) &= \sum b_i x^i diag(a_1^i, \dots, a_n^i)) \\ &= diag\left(\sum b_i (xa_1)^i, \dots, \sum b_i (xa_n)^i\right) \\ &= diag(f(a_1x), \dots, f(a_nx)) \end{split}$$

If A is diagonalizable then we can write  $A = SAS^{-1}$  where  $A = diag(d_1, \dots, d_n)$ . The entries of f(xA) are linear combinations of  $f(d_1x), \dots, f(d_nx)$  since

$$\begin{split} f(xA) &= \sum b_i x^i S \Lambda^i S^{-1} \\ &= S \left( \sum b_i x^i \Lambda^i \right) S^{-1} \\ &= S \operatorname{diag}(f(d_1 x), \dots, f(d_n x)) S^{-1} \end{split}$$

For example, if  $A = \begin{pmatrix} 1 & 1 \\ 0 & a \end{pmatrix}$  then A is diagonalizable, and  $A^n = \begin{pmatrix} 1 & \frac{a^n - 1}{a - 1} \\ 0 & a^n \end{pmatrix}$  so that

$$f(xA) = \begin{pmatrix} f(x) & \frac{1}{\alpha-1}(f(\alpha x) - f(x)) \\ 0 & f(\alpha x) \end{pmatrix}$$

It is easy to find polynomials f for which the entries of the above matrix are not in **P**.

A different kind of example is given by  $A = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$  which is not diagonalizable, yet f(xA) has an especially simple form. Since  $A^n = \begin{pmatrix} 1 & 0 \\ n & 1 \end{pmatrix}$  it follows that

$$f(xA) = \sum b_{i}x^{i} \begin{pmatrix} 1 & 0 \\ i & 1 \end{pmatrix} = \begin{pmatrix} f & 0 \\ xf' & f \end{pmatrix}$$

This example can be generalized to the matrices whose entries above the diagonal are 0, and the rest are 1. For instance, when n = 4

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 \end{pmatrix}^{n} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ \binom{n}{1} & 1 & 0 & 0 \\ \binom{n+1}{2} & \binom{n}{1} & 1 & 0 \\ \binom{n+2}{3} & \binom{n+1}{2} & \binom{n}{1} & 1 \end{pmatrix}$$

It follows easily that

$$f(xA) = \begin{pmatrix} f & 0 & 0 & 0 \\ xD(f) & f & 0 & 0 \\ xD^2(xf) & xD(f) & f & 0 \\ xD^3(x^2f) & xD^2(xf) & xD(f) & f \end{pmatrix}$$

Thus for these matrices f(xA) has all its entries in **P** for any choice of f. It is easy to characterize those polynomials for which f(xA) is a matrix whose entries are all in **P** for every choice of A.

**Lemma 17.25.** *If* f *is a polynomial then the following are equivalent:* 

- 1. f(xA) has all entries in **P** for every matrix A.
- 2.  $f(x) = cx^n(ax + b)$  for constants a, b, c.

*Proof.* If (2) holds then

$$f(xA) = cx^{n}(axA^{n+1} + bA^{n}).$$

Consequently, all entries of f(xA) are of the form  $x^n$  times a linear term, and so all entries are in **P**.

Conversely, assume f(xA) has all entries in **P** for any choice of A. Consider the matrix  $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$  and its powers:

$$\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}^{4k} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \qquad \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}^{4k+1} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$
$$\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}^{4k+2} = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \qquad \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}^{4k+3} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

If 
$$f(x) = \sum_{i=1}^{n} a_i x^i$$
 then

$$\begin{split} f(xA) &= \begin{pmatrix} \sum_{i} a_{4i} x^{4i} - a_{4i+2} x^{4i+2} & \sum_{i} a_{4i+1} x^{4i+1} - a_{4i+3} x^{4i+3} \\ - \left( \sum_{i} a_{4i+1} x^{4i+1} - a_{4i+3} x^{4i+3} \right) & \sum_{i} a_{4i} x^{4i} - a_{4i+2} x^{4i+2} \end{pmatrix} \\ &= \begin{pmatrix} f_e(-x^2) & x f_o(-x^2) \\ -x f_o(-x^2) & f_e(-x^2) \end{pmatrix} \end{split}$$

where  $f_e$  and  $f_o$  are the even and odd parts of f. If we consider  $B = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  then a similar computation shows that

$$f(\mathbf{x}B) = \begin{pmatrix} f_e(\mathbf{x}^2) & \mathbf{x}f_o(\mathbf{x}^2) \\ -\mathbf{x}f_o(\mathbf{x}^2) & f_e(\mathbf{x}^2) \end{pmatrix}$$

Now  $f_e(x^2) \in \mathbf{P}$  iff  $f_e \in \overline{\mathbf{P}^{alt}}$ , and  $f_e(-x^2) \in \mathbf{P}$  iff  $f_e \in \overline{\mathbf{P}^{pos}}$ , and hence  $f_e(x) = ax^r$  for some a, r. Similarly  $f_o(x) = bx^s$  for some b, s, and thus  $f(x) = ax^{2r} + bx^{2s+1}$ . If we choose A = (1) we see that  $f \in \mathbf{P}$ , and hence 2r and 2s + 1 must be consecutive integers. This concludes the proof of the lemma.

CHAPTER



# **Matrix Polynomials**

A *matrix polynomial* is a polynomial whose coefficients are matrices. Equivalently, it is a matrix whose entries are polynomials. We assume all matrices are  $\mu$  by  $\mu$ . We write

$$f(\mathbf{x}) = A_0 + \dots + A_n \mathbf{x}^r$$

The leading coefficient of f(x) is the matrix  $A_n$ . We are interested in the class of *hyperbolic* matrix polynomials that are a nice generalization of **P**.

**Definition 18.1.** The *hyperbolic matrix polynomials in* d *variables of degree* n, denoted  $Hyp_d(n)$ , consists of all matrix polynomials  $f(\mathbf{x}) = \sum A_I \mathbf{x}^I$  such that

- 1.  $\langle f \cdot v, v \rangle = v^t f(\mathbf{x}) v \in \mathbb{P}_d$  for all  $v \in \mathbb{R}^{\mu}$ .
- 2.  $f(\mathbf{x})$  has degree n.
- 3. If  $|\mathbf{I}| = n$  then  $A_{\mathbf{I}}$  is positive definite.

 $\mathsf{Hyp}_d^{\mathrm{pos}}(n)$  is the subset of  $\mathsf{Hyp}_d(n)$  where all coefficients are positive definite. If we don't wish to specify the degree we write  $\mathsf{Hyp}_d$  and  $\mathsf{Hyp}_d^{\mathrm{pos}}$ .

A hyperbolic matrix polynomial of degree zero is a constant matrix, and must be positive definite.

If  $g \in P_d$ , and P is a positive definite matrix then  $g P \in Hyp_d$ . Although this is a trivial construction, the embedding  $P_d \longrightarrow Hyp_d$  is important.

#### 18.1 Introduction

In this section we establish a few simple properties about hyperbolic matrix polynomials. The role of positive coefficients in  $\mathbf{P}$  is played by positive definite polynomials in  $\mathsf{Hyp}_d$ . For instance, a matrix polynomial of degree 1 in  $\mathsf{Hyp}_d$  has the form

$$A_1x_1 + \cdots + A_dx_d + S$$

where the  $A_i$  are positive definite, and S is symmetric. If S is positive definite then the polynomial is in Hyp<sub>d</sub><sup>pos</sup>.

The analog of multiplying by a positive constant is replacing f by  $A^{t}fA$ . The leading coefficients are positive definite, and if w = Av then

$$< A^{t}fA \cdot v, v >= v^{t}A^{t}fAv = < f \cdot w, w > \in P_{d}.$$

In the case that d is 1 then we can multiply on the left and right by the square root of the inverse of the leading coefficient, and so we may assume that the coefficient of  $x^n$  is the identity matrix.

**Lemma 18.2.** Any principle submatrix of a matrix in  $Hyp_d$  is also in  $Hyp_d$ . In particular, all diagonal entries are in  $P_d$ .

*Proof.* This is immediate, for we just need to set the i-th coordinate of v equal to zero to show the hyperbolic property for the i-th principle submatrix. The leading coefficients are positive definite since all principle submatrices of a positive definite matrix are positive definite.

**Lemma 18.3.** Suppose that f(x) is a diagonal matrix  $diag(f_1(x), \ldots, f_{\mu}(x))$ . The following are equivalent:

1.  $f \in Hyp_1$ .

2.  $f_1, \ldots, f_{\mu}$  have a common interlacing.

*Proof.* If  $\mu = 2$  then

$$(\mathbf{a},\mathbf{b})\left(\begin{smallmatrix}f_1&0\\0&f_2\end{smallmatrix}
ight)(\mathbf{a},\mathbf{b})^{\mathbf{t}}=\mathbf{a}^2\mathbf{f}_1+\mathbf{b}^2\mathbf{f}_2\in\mathbf{P}$$

Thus all positive linear combinations of  $f_1$  and  $f_2$  lie in **P**, and so  $f_1$  and  $f_2$  have a common interlacing. It follows that any two of the  $f_i$  have a common interlacing, and so they all have a common interlacing.

Conversely, if they have a common interlacing then all positive linear combinations are in **P**, and so

$$(\mathfrak{a}_1,\ldots,\mathfrak{a}_\mu)\operatorname{diag}(\mathfrak{f}_1,\ldots,\mathfrak{f}_\mu)(\mathfrak{a}_1,\ldots,\mathfrak{a}_\mu)^{\operatorname{t}} = \mathfrak{a}_1^2\mathfrak{f}_1 + \cdots \mathfrak{a}_\mu^2\mathfrak{f}_\mu \in \mathbf{P}.$$

*Remark* 18.4. We can use this result to show that the tensor (Kronecker) product does not preserve hyperbolic matrix polynomials. If  $f = \begin{pmatrix} x & 0 \\ 0 & x-1 \end{pmatrix}$  then  $f \otimes f \otimes f$  is an 8 by 8 diagonal matrix whose diagonal contains  $x^3$  and  $(x-1)^3$ . Since these two do not have a common interlacing,  $f \otimes f \otimes f$  is not in Hyp<sub>1</sub>.

Replacing x by Ax in a polynomial with all real roots does not usually yield a hyperbolic matrix polynomial.

**Corollary 18.5.** Suppose that A is a positive definite matrix with eigenvalues  $d_1, \ldots, d_{\mu}$ , and  $f \in P$ . Then  $f(xA) \in Hyp_1$  if and only if  $\{f(d_1x), \ldots, f(d_{\mu}x)\}$  have a common interlacing.

*Proof.* Let O be orthogonal, and D diagonal so that  $A = O^{t}DO$ . Then

$$f(xA) = O^{t} f(xD) O = O^{t} diag(f(xd_{i})) O$$

Thus,  $f(xA) \in Hyp_1$  if and only if  $diag(f(xd_i)) \in Hyp_1$ . Lemma 18.3 completes the proof.

*Remark* 18.6. There is a simple criterion for  $\{f(d_ix)\}_i$  to satisfy the corollary. Suppose that  $roots(f) = (r_i)$ , and  $\alpha > 1$ . The roots of  $f(\alpha x)$  are  $r_1/\alpha < \cdots < r_n/\alpha$ . If they have a common interlacing then

$$r_1/\alpha < r_1 < r_2/\alpha < r_2 \cdots$$

and thus  $\alpha < \min_{i>j} r_i/r_j$ . If  $0 < \alpha < \beta$  then  $f(\alpha x)$  and  $f(\beta x)$  have a common interlacing if and only if f(x) and  $f(\beta x/\alpha)$  do. We conclude that  $\{f(d_i x)\}$  has a common interlacing if and only if

$$\max_{i,j} \frac{d_i}{d_j} \leqslant \min_{i>j} \frac{r_i}{r_j}$$

Note that if this condition is satisfied then they are actually mutually interlacing.

If we add off-diagonal entries that are all equal, then we can determine when the matrix polynomial is in  $Hyp_1$ . Define

$$\sigma_{n} = \max_{a_{1},\dots,a_{n}} \frac{a_{1}a_{2} + a_{2}a_{3} + a_{3}a_{4} + \dots + a_{n-1}a_{n}}{a_{1}^{2} + \dots + a_{n}^{2}}$$

Some values of  $\sigma_n$  are

$$\sigma_2 = \frac{1}{2}$$
  $\sigma_3 = \frac{\sqrt{2}}{2}$   $\sigma_4 = \frac{1+\sqrt{5}}{4}$   $\sigma_5 = \frac{\sqrt{3}}{2}$ 

**Lemma 18.7.** The n by n tridiagonal matrix below is in  $Hyp_1$  if and only if  $f+2\alpha g \in P$  for  $|\alpha| \leq \sigma_n$ .

$$\mathfrak{m}(\mathbf{x}) = \begin{pmatrix} \mathbf{f} & \mathbf{g} & \mathbf{0} & \dots & \\ \mathbf{g} & \mathbf{f} & \mathbf{g} & \mathbf{0} & \dots & \\ \mathbf{0} & \mathbf{g} & \mathbf{f} & \mathbf{g} & \mathbf{0} & \dots \\ \mathbf{0} & \mathbf{0} & \ddots & \ddots & \ddots & \ddots \end{pmatrix}$$

*Proof.* If  $v = (a_1, \ldots, a_n)^t$ , then

$$v^{t}mv = (a_{1}^{2} + \dots + a_{n}^{2})f + 2(a_{1}a_{2} + a_{2}a_{3} + a_{3}a_{4} + \dots + a_{n-1}a_{n})g$$

Dividing by the first factor, we see that this is in **P** if and only if  $f + 2\alpha g \in \mathbf{P}$  for  $|\alpha| \leq \sigma_n$ .

The off-diagonal element can even be a constant if it is small enough.

**Corollary 18.8.** *Suppose that*  $f \in P$ *. The following are equivalent:* 

$$\begin{split} & 1. \ \left(\begin{smallmatrix} f & c \\ c & f \end{smallmatrix}\right) \in \ \textit{Hyp}_1 \\ & 2. \ |c| \leqslant \min_{f'(\alpha)=0} |f(\alpha)| \\ \end{split}$$

*Proof.* Geometrically, the second condition means that the largest value of |c| satisfies  $f + b \in \mathbf{P}$  for all  $|b| \leq |c|$ . Since  $2\sigma_2 = 1$ , the result now follows from the lemma.

The off diagonal entries are not necessarily in  $P_d$ . This can be deduced from Lemma 18.7 by taking g small relative to f. For example,

$$\begin{pmatrix} x^2-1 & \varepsilon(x^2+1) \\ \varepsilon(x^2+1) & x^2-1 \end{pmatrix} \in \ \text{Hyp}_1 \Longleftrightarrow |\varepsilon| \leqslant 1$$

The reverse preserves  $Hyp_1$ .

**Lemma 18.9.** If  $f(x) = \sum_{i=0}^{n} A_i x^i \in Hyp_1$ , and  $A_0$  is positive definite, then the reverse  $f^{rev}(x) = \sum_{i=0}^{n} A_i x^{n-i}$  is also in  $Hyp_1$ .

*Proof.* Since  $A_0$  is positive definite, the leading coefficient of  $f^{rev}$  is positive definite. The conclusion follows from the calculation

$$< \nu f^{re\nu}, \nu > = < \nu x^n f(1/x), \nu > = x^n < \nu f(1/x), \nu > = < \nu f, \nu >^{re\nu}.$$

The product of hyperbolic matrix polynomials is generally not hyperbolic since the product of the leading coefficients, which are positive definite matrices, is not usually positive definite. We do have

**Lemma 18.10.** If f, 
$$g \in Hyp_1$$
 then  $f^t g f \in Hyp_1$ .

*Proof.* If A is the leading coefficient of f and B is the leading coefficient of g, then the leading coefficient of  $f^t g f$  is  $A^t b A$  which is positive definite. If  $v \in \mathbb{R}^{\mu}$  and w = fv then

$$v^t f^t g f m v = w^t g w \in \mathbf{P}$$

#### 18.2 Interlacing

There are two equivalent ways to define interlacing in  $Hyp_d$ : the usual linearity definition, or reduction to one variable. We start with the latter, since it is more precise.

**Definition 18.11.** Suppose that  $f, g \in \mathsf{Hyp}_d$  have degree n and n - 1 respectively. We define

 $\begin{array}{ll} f \leq g & \text{if and only if} \quad \nu^t f \nu \leq \nu^t g \nu & \text{for all } \nu \in \mathbb{R}^{\mu} \setminus 0 \\ f < g & \text{if and only if} \quad \nu^t f \nu < \nu^t g \nu & \text{for all } \nu \in \mathbb{R}^{\mu} \setminus 0 \end{array}$ 

 $\ll$  and  $\underline{\ll}$  are defined similarly.

**Lemma 18.12.** Suppose that  $f, g \in Hyp_d$  have degree n and n-1 respectively. The following are equivalent:

1.  $f + \alpha g \in Hyp_d$  for all  $\alpha \in \mathbb{R}$ .

2. f<u>≪</u>g

Proof. By hypothesis we know that

$$v^{t}(f + \alpha g)v = v^{t}fv + \alpha v^{t}gv \in P_{d}$$
 for all  $\alpha \in \mathbb{R}$ 

and consequently we have that  $v^t f v \leq v^t g v$ . The converse is similar.

We now have some easy consequences of the definition:

**Lemma 18.13.** Assume that  $f, g, h \in Hyp_d$ .

- 1. If  $f \leq g$  and  $f \leq h$  then  $f \leq g + h$ . In particular,  $g + h \in Hyp_d$ .
- 2. If  $f \leq g \leq h$  then  $f h \leq g$ .
- 3. If  $k \in P_d$  then  $fk \in Hyp_d$ .
- 4. If all  $a_i$  are positive and  $f \in Hyp_d$  then

$$(a_1x_1 + \cdots + a_dx_d + b)f \leq f$$

- 5. If the i-th diagonal element of f is  $f_i$ , and of g is  $g_i$  then  $f \leq g$  implies  $f_i \leq g_i$ .
- 6. If  $F \leq G$  in  $P_d$  and  $M_1$ ,  $M_2$  are positive definite matrices then  $FM_1 \leq GM_2$  in  $Hyp_d$ .
- 7. If  $g_1, \ldots, g_r$  have a common interlacing in  $P_d$ , and  $M_1, \ldots, M_r$  are positive definite matrices then  $g_1 M_1 + \cdots + g_r M_r \in Hyp_d$ .

*Proof.* By hypothesis we know that  $v^t f v \leq v^t g v$  and  $v^t f v \leq v^t h v$ . Since f, g, h  $\in$  Hyp<sub>d</sub> we know that the leading coefficients of these three terms are positive, so we can add interlacings in P<sub>d</sub>:

$$v^{t} f v \leq v^{t} g v + v^{t} h v = v^{t} (g + h) v$$

The proof of the second one is similar. The third one follows since  $P_d$  is closed under multiplication, and k is a scalar:

$$\langle \mathsf{fk} \cdot \mathsf{v}, \mathsf{v} \rangle = \mathsf{k} \langle \mathsf{f} \cdot \mathsf{v}, \mathsf{v} \rangle \in \mathsf{P}_{\mathsf{d}}$$

The fourth follows from the third one. For the next one, note that if  $e_i$  is the vector with 1 in the i-th place and zeros elsewhere, then  $e_i^t f e_i$  is the i-th diagonal element of f.

If  $F \leq G$  then  $v^t M_1 v$  and  $v^t M_2 v$  are positive so  $v^t F M_1 v \leq v^t G M_2 v$ . If  $G \leq g_i$  for  $1 \leq i \leq r$  then  $G I \leq g_i M_i$  by the previous result, and the conclusion follows by adding interlacings.

The interlacing of even linear polynomials leads to some complexity.

Lemma 18.14. Suppose that A, C are positive definite, and B, D are symmetric.

- 1.  $xI B \leq xI D$  iff  $B \ge D$  (i.e. B D is positive definite).
- 2.  $xA B \leq xC D$  iff

$$\frac{v^{t}Bv}{v^{t}Av} \ge \frac{v^{t}Dv}{v^{t}Cv} \quad \text{for all } v \tag{18.2.1}$$

*Proof.* If  $xI - B \leq xI - D$  then for all non-zero v we have

$$v^{t}(xI - B)v \underline{\ll} v^{t}(xI - D)v$$
$$(v^{t}v)x - v^{t}Bv \underline{\ll} (v^{t}v)x - v^{t}Dv$$

These are linear polynomials with positive leading coefficients, and they interlace iff  $\nu^t B\nu \ge \nu^t D\nu$ . Thus  $\nu^t (B - D)\nu \ge 0$ , so  $B \ge D$ .

In the second case

$$v^{t}Av x - v^{t}Bv \ll v^{t}Cv x - v^{t}Dv$$

for all non-zero v. Solving each equation for x and using the first part yields the conclusion.

The condition (18.2.1) is not well understood. Let  $\lambda_{\min}(W)$  and  $\lambda_{\max}(W)$  be the smallest and largest eigenvalues of a matrix W. Suppose that matrices A, B, C, D are as in the lemma and (18.2.1) is satisfied for all non-zero v. If W is any symmetric matrix then the Rayleigh-Ritz inequality [85] says that

$$\lambda_{\min}(W) \leqslant rac{
u^t W 
u}{
u^t 
u} \leqslant \lambda_{\max}(W) \quad ext{for all } 
u 
eq 0$$

We may assume that  $\nu$  has norm 1. The values of  $\nu^{t} W \nu$  lie in the interval  $(\lambda_{\min}(W), \lambda_{\max}(W))$ . If (18.2.1) holds it follows that

$$(\nu^{t}B\nu) (\nu^{t}D\nu) \ge (\nu^{t}A\nu) (\nu^{t}C\nu) \quad \text{for all } \nu \lambda_{\max}(B)\lambda_{\max}(D) \le \lambda_{\min}(A)\lambda_{\min}(C)$$
 (18.2.2)

The values of  $\frac{\nu^t B \nu}{\nu^t A \nu}$  lie in the interval  $[\lambda_{min}(BA^{-1}), \lambda_{max}(BA^{-1})]$  and therefore

$$\lambda_{\min}(BA^{-1}) \leq \lambda_{\max}(DC^{-1}) \quad \text{and} \quad (18.2.3)$$
$$\lambda_{\min}(BA^{-1})\lambda_{\min}(CD^{-1}) \leq 1$$

Conversely, if we know that the stronger condition

 $\lambda_{\text{max}}(BA^{-1}) < \lambda_{\text{min}}(CD^{-1})$ 

holds then we can conclude that (18.2.1) holds.

Hyp<sub>d</sub> behaves well under induced linear transformations.

**Lemma 18.15.** Suppose that  $T: P_d \longrightarrow P_d$ , and if d is 1 then T preserves the sign of the leading coefficient. Define a map on  $Hyp_d$  by  $T(\sum A_I \mathbf{x}^I) = \sum A_I T(\mathbf{x}^I)$ . Then  $T: Hyp_d \longrightarrow Hyp_d$ .

*Proof.* If  $f \in \mathsf{Hyp}_d$  then the leading coefficient of T(f) is positive definite. The result now follows from

$$T < f \cdot v, v > = < T(f) \cdot v, v >$$

The derivative behaves as expected. The proofs follow from the previous lemma, and the arguments for **P**.

**Lemma 18.16.** Suppose that  $f, g \in Hyp_d$  and  $h \in P^{pos}$ .

- 1.  $\frac{\partial f}{\partial x_i} \in Hyp_d$ . 2.  $f \leq \frac{\partial f}{\partial x_i}$
- 3. If  $f \leq g$  then  $\frac{\partial f}{\partial x_i} \leq \frac{\partial g}{\partial x_i}$
- 4.  $h(\partial/\partial x_i)f \in Hyp_d$

**Example 18.17.** What does the graph of a diagonal matrix polynomial look like? Consider  $f = \begin{pmatrix} (x-1)(x-3) & 0 \\ 0 & (x-2)(x-4) \end{pmatrix}$ . We have the interlacings

$$(x-1)(x-2)I \leq f \leq (x-3)(x-4)I$$

If we graph  $(1, \nu)^{t} f(1, \nu)$  then Figure 18.1 shows that it stays in the regions determined by (x - 1)(x - 2) = 0 and (x - 3)(x - 4) = 0. The small dashed lines are the graph of (x - 1)(x - 3) = 0, and the large dashed lines are the roots of (x - 2)(x - 4) = 0. This is a general phenomenon – see page 556.



Figure 18.1: The graph of a diagonal matrix polynomial

## 18.3 Common interlacings

If  $f \in Hyp_1$  and  $g \in \mathbf{P}$ , then  $f \leq gI$  if and only if  $v^t f v \leq g$  for all  $v \in \mathbb{R}^{\mu}$ . If either condition holds we say that f *has a common interlacing*.

The usual definition[122] of hyperbolic polynomials allows the vectors v to be complex. This is equivalent to the existence of a common interlacing. Recall that  $w^*$  is the conjugate transpose.

**Lemma 18.18.** *Suppose that* f *is a matrix polynomial whose leading coefficient is the identity. The following are equivalent.* 

**1**  $w^* f w \in \mathbf{P}$  for all  $w \in \mathbb{C}^{\mu}$ .

**2a** { $v^t f v \mid v \in \mathbb{R}^{\mu}$ } has a common interlacing.

**2b** f has a common interlacing.

*Proof.* 2a and 2b are equivalent by definition, so assume that 1 holds. Write w = u + iv where u, v are real vectors. Then

$$w^* fw = u^* fu + u^* f(w) + (w)^* fu + (w) fw$$
$$= u^t fu + v^t fv$$

Replacing v by  $\alpha v$  shows that  $u^t f u + \alpha^2 v^t f v \in \mathbf{P}$  for all  $\alpha$ , so  $u^t f u$  and  $v^t f v$  have a common interlacing. Conversely, if all  $u^t f u$  and  $v^t f v$  have a common interlacing then  $w^* f w \in \mathbf{P}$ .

Do all hyperbolic polynomials in  $Hyp_1$  have a common interlacing? This is an unsolved question, but it is true for two by two matrices.

**Lemma 18.19.** Any two by two matrix  $F \in Hyp_1$  has a common interlacing.

*Proof.* If we write  $F = \begin{pmatrix} f & h \\ h & g \end{pmatrix}$  and let  $v = (\beta, 1)$  then

$$v^{t} f v = \beta^{2} f + 2\beta h + g$$

Lemma 3.50 guarantees the existence of a common interlacing.

There is a simple condition that implies a common interlacing.

**Lemma 18.20.** Assume  $f(x) = x^2I + Ax + B$  where A, B are symmetric. If there is an  $\alpha$  such that  $f(\alpha)$  is negative definite then f has a common interlacing.

*Proof.* Since  $\nu^t f(\alpha)\nu < 0$  and  $\nu^t f(\beta)\nu > 0$  for  $\beta$  sufficiently large, it follows that  $f \leq (x - \alpha)I$ .

We have seen that the polynomials of a diagonal hyperbolic matrix polynomial have a common interlacing. This is true in general.

**Corollary 18.21.** *The diagonal elements of a polynomial in*  $Hyp_1$  *have a common interlacing.* 

*Proof.* Any two by two principle submatrix is in  $Hyp_1$ , and by Lemma 18.19 the diagonal elements have a common interlacing. Since any two diagonal elements have a common interlacing, they all have a common interlacing.  $\Box$ 

**Corollary 18.22.** *If*  $f \in Hyp_1$  *has a common interlacing then the determinant and trace of* f *are in* P*.* 

*Proof.* If det( $f(\lambda)$ ) = 0 then there is a  $w \in \mathbb{C}$  so that  $f(\lambda)w = 0$ , and thus  $w^*f(\lambda)w = 0$ . By Lemma 18.18 we know that  $w^*fw \in \mathbf{P}$  for all complex w, so  $\lambda$  is real.

Since the diagonal elements have a common interlacing, their sum is in  $\mathbf{P}$ .

**Lemma 18.23.** If  $f \leq g$  in  $Hyp_1$  then trace(f)  $\leq$  trace(g) in **P**.

*Proof.* For any  $\alpha \in \mathbb{R}$  we know that  $f + \alpha g \in \mathsf{Hyp}_1$ . Now the trace is a linear map  $\mathsf{Hyp}_1 \longrightarrow \mathbf{P}$ , so trace $(f) + \alpha \operatorname{trace}(g) \in \mathbf{P}$ , which finishes the proof.  $\Box$ 

**Example 18.24.** Here is a simple example of a polynomial with a common interlacing. If v = (a, b) and

$$\mathbf{f} = \mathbf{x} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

then

$$v^{t} f v = (a^{2} + b^{2})x + 2ab$$

Consequently, the roots of  $v^t f v$  lie in the closed interval [-1, 1] and so

$$(\mathbf{x}-1)(\mathbf{x}+1)\mathbf{I} \leq \mathbf{f}.$$

The determinant of f is  $x^2 - 1$ , which is in **P**, as the corollary tells us.

Empirical investigations suggest that the adjugate of a matrix that determines a polynomial in  $\mathbf{P}_d^{\text{pos}}$  is in  $\text{Hyp}_d$ . We prove this for d = 1, and show that there is a common interlacing for all d.

Assume that  $D_1, \ldots, D_d$  are positive definite  $\mu$  by  $\mu$  matrices, and consider the matrix

$$M = Id + x_1D_1 + \cdots + x_dD_d.$$

Recall that M[i, j] is the matrix formed by removing the i'th row and j'th column of M. The adjugate of M is

$$\operatorname{Adj}[M] = \left( \left| (-1)^{i+j} M[i,j] \right| \right).$$

The adjugate satisfies

$$\mathrm{Adj}[\mathrm{M}] \cdot \mathrm{M} = |\mathrm{M}|\mathrm{Id}$$

and our conjecture is that  $Adj(M) \in Hyp_d$ . First of all, we show that Adj(M) has a common interlacing.

**Lemma 18.25.** If M is as above and  $v \in \mathbb{R}^{\mu}$  then  $|\mathsf{M}| \leq v^{t} \operatorname{Adj}(\mathsf{M}) v$ . In particular,  $v^{t} \operatorname{Adj}(\mathsf{M}) v \in \overline{\mathbf{P}}_{d}^{pos}$ .

*Proof.* Choose  $\nu=(\nu_i)\in\mathbb{R}^\mu$  and define  $u=((-1)^i\nu_i).$  Introducing a new variable y, define

$$N = M + y \cdot uu^t$$
.

Now  $N \in \overline{\mathbf{P}}_{d+1}^{pos}$  since  $uu^t$  is positive semi-definite. Also,  $uu^t$  has rank one, so N is linear in y. Expanding |N| yields

$$|\mathbf{N}| = |\mathbf{M}| + y \sum_{i,j} (-1)^{i+j} v_i v_j |\mathbf{M}[i,j]|$$
$$= |\mathbf{M}| + y v^t \operatorname{Adj}(\mathbf{M}) v$$

We conclude that  $|M| \leq v^t \operatorname{Adj}(M) v$  since consecutive coefficients interlace.

In order to show that  $Adj(M) \in Hyp_d$  we need to show that the coefficients of the homogeneous part are positive definite. It's easy for d = 1.

**Lemma 18.26.** If D is positive definite then  $Adj(Id + xD) \in Hyp_1$ .

*Proof.* Let M = Id + xD. The degree of |M[i, j]| is at most  $\mu - 1$ , so the degree of Adj(M) is at most  $(\mu - 1)\mu$ . The coefficient of  $x^{\mu(\mu-1)}$  is  $((-1)^{i+j} |D[i, j]|) = |D|D^{-1}$ . Since  $D^{-1}$  is positive definite, we see that  $Adj(M) \in Hyp_1$ .

We can sometimes substitute matrices for constants, and we get hyperbolic matrices with a common interlacing. Recall that the norm ||M|| is the maximum eigenvalue of a positive definite matrix M and satisfies

$$\|\mathbf{M}\| = \sup_{\mathbf{v}\neq 0} \frac{\mathbf{v}^{\mathsf{t}} \, \mathbf{M} \, \mathbf{v}}{\mathbf{v}^{\mathsf{t}} \, \mathbf{v}}.$$

**Lemma 18.27.** *Suppose that* M *is positive definite and* deg(f) > deg(g)*.* 

- 1. If  $f + \alpha g \in P$  for  $0 \leq \alpha \leq ||M||$  then  $fI + gM \in Hyp_1$ .
- 2.  $fI \leq fI + gM$

*Proof.* The leading coefficient of fI + gM is I. To check the inner product condition, compute

$$v^{t} (fI + gM) v = v^{t} v \left( f + \frac{v^{t} M v}{v^{t} v} g \right)$$

Since  $\frac{v^t M v}{v^t v} \leq ||M||$  the inner product condition holds. Next,

$$fI + \alpha(fI + gM) = (1 + \alpha) \left( fI + \frac{\alpha}{1 + \alpha} gM \right)$$

and the latter polynomial is in  $\text{Hyp}_1$  since  $\left\|\frac{\alpha}{1+\alpha}M\right\| \leq \|M\|$  for all positive  $\alpha$ .

**Lemma 18.28.** If  $f \in Hyp_d$  has a common interlacing then  $|f| \in P_d$ .

*Proof.* If we substitute for all but one of the variables then we get a polynomial in  $Hyp_1$  with a common interlacing. We know that its determinant is in **P**, so substitution is satisfied.

Assume that f has degree n. If we write  $f = \sum_{I} a_{I} x^{I}$  then the homogeneous part of |f| is  $|\sum_{|I|=n} a_{I} x^{I}|$ . If we replace each monomial  $x^{I}$  by a new variable  $y_{I}$  then we know that  $|\sum_{|I|=n} a_{I} y_{I}|$  has all positive coefficients since all the  $a_{I}$  are positive definite. Replacing the  $y_{I}$  by  $y^{I}$  shows that  $|f|^{H}$  has all positive coefficients.

**Lemma 18.29.** If  $f \leq g$  in  $Hyp_1$  then  $f + yg \in \overline{Hyp_2}$ .

Proof. It suffices to show that

$$F_{\varepsilon}(x,y) = f(x + \varepsilon y) + (y + \varepsilon x)g(x + \varepsilon y) \in Hyp_2$$

for positive  $\epsilon$ . Now the homogeneous part is

$$F_{\epsilon}^{H} = f^{H}(x + \epsilon y) + (y + \epsilon x)g^{H}(x + \epsilon y)$$

and  $f^H$ ,  $g^H$  are matrix polynomials with positive definite coefficients it follows that  $F_{\epsilon}^H$  is a sum of polynomials with positive definite coefficients, and so all its coefficients are positive definite.

Next, we verify substitution. Fix  $v \in \mathbb{R}^n$  and let  $G(x, y) = v^t f v + y v^t g v$ . By hypothesis  $v^t f v \leq v^t g v$ , so we know that  $G(x + \varepsilon y, y + \varepsilon x) \in P_2$ . Thus,  $F_{\varepsilon} \in Hyp_2$ .

**Lemma 18.30.** *If*  $gI \leq f$  *in*  $Hyp_1$  *then* 

- 1.  $|gI + yf| \in \overline{\mathbf{P}}_2$
- 2. If we write  $|I + yf| = \sum f_i(x)y^i$  then  $f_i \in \mathbf{P}$ .
- 3.  $f_i \leq gf_{i+1}$ .

*Proof.* The first part follows from the previous lemma. If we expand the determinant we get

$$|gI + yf| = \sum g^{\mu - i} f_i(x) y^i$$

Since this is in  $\mathtt{P}_2$  we see that  $g^{\mu-i}f_i \lessdot g^{\mu-(i-1)}f_{i+1}$  which implies that  $f_i \underline{\lessdot} gf_{i+1}.$ 

The next corollary generalizes the fact the the determinant and trace are in **P**.

**Corollary 18.31.** *If* 
$$f \in Hyp_1$$
 *has a common interlacing then*  $\sum_{|S|=k} |f[S]| \in P$ .

*Proof.* We have  $S \subset \{1, ..., n\}$  and f[S] is the submatrix of f only using the rows and columns in S. The sum in question is the coefficient of  $y^{n-k}$  in |ygI + f|, and this is in **P** by the last lemma.

*Remark* 18.32. If we drop the assumption that matrices are positive definite then we can find a 2 × 2 matrix M that satisfies  $v^t M v \in \mathbf{P}$  for all v, yet has no common interlacing. Choose

$$F(x, y, z) = f + y g + z h + y z k \in \overline{\mathbf{P}}_3$$

Choose  $\alpha, \beta \in \mathbb{R}$  and  $\alpha \neq 0$ . Since

$$\begin{pmatrix} \alpha & \beta \end{pmatrix} \begin{pmatrix} f & g \\ h & k \end{pmatrix} \begin{pmatrix} \alpha \\ \beta \end{pmatrix} = \alpha^2 \left[ f + \frac{\beta}{\alpha} (g + h) + \left( \frac{\beta}{\alpha} \right)^2 k \right]$$

we see it is in **P** since it equals  $F(x, \beta/\alpha, \beta/\alpha)$ . For a particular example, consider

$$\begin{aligned} (x+y+z+1)(x+2y+3z+5) \\ &= 5+6x+x^2+y(7+3x)+z(8+4x)+5yz+2y^2+3z^2 \end{aligned}$$

Taking coefficients yields the matrix

$$\mathsf{M} = \begin{pmatrix} 5+6\mathbf{x}+\mathbf{x}^2 & 8+4\mathbf{x} \\ 7+3\mathbf{x} & 5 \end{pmatrix}$$

The table below shows that there is no common interlacing.

b	roots o	$f(1b)M(1b)^t$
-5	-35.1	-5.8
5	2.0	26.9

#### 18.4 Orthogonal-type recurrences

The fact that  $f \leq g \leq h \implies f-h < g$  is all we need to establish orthogonal type recurrences in Hyp<sub>d</sub>.

Lemma 18.33. Define a sequence of matrix polynomials by

$$\begin{split} f_0 &= U \\ f_1 &= A_1 x_1 + \dots + A_d x_d + S \\ f_k &= (a_{k,1} x_1 + \dots + a_{k,d} x_d + b_k) f_{k-1} - c_k f_{k-2} \end{split}$$

where U and the  $A_i$  are positive definite, S is symmetric, all  $c_k$  are positive, and all  $a_{i,i}$  are positive.

Then all  $f_k$  are in  $Hyp_d$  and  $f_k \leq f_{k-1}$ .

*Proof.* We note that  $f_0$  and  $f_1$  are in Hyp<sub>d</sub>. By induction we have that

$$(a_{k,1}x_1 + \cdots + a_{k,d}x_d + b_k)f_{k-1} \leq f_{k-1} \leq f_{k-2}$$

and the observation above finishes the proof.

The off-diagonal entries of a matrix polynomial in  $\text{Hyp}_1$  might not have all real roots. Here's an explicit example. The coefficient of x in  $f_1$  below is positive definite, as is  $f_0$ , so  $f_2 \in \text{Hyp}_1$ . However, the off diagonal entry of  $f_2$  has two complex roots. The diagonal elements have roots (-2.4, .4) and (-2.5, 1), so they have a common interlacing.

In general, all the diagonal elements of the sequence  $f_0, f_1, \ldots$  form an orthogonal polynomial sequence. The off diagonal elements obey the three term recursion, but the degree 0 term might not be positive, and so they might not have all real roots.

$$f_{0} = \begin{pmatrix} 2 & 3 \\ 3 & 6 \end{pmatrix}$$

$$f_{1} = \begin{pmatrix} 1 & -1 \\ -1 & 2 \end{pmatrix} \mathbf{x} + \mathbf{I}$$

$$f_{2} = (\mathbf{x} + 1)f_{1} - f_{0}$$

$$= \begin{pmatrix} \mathbf{x}^{2} + 2\mathbf{x} - 1 & -\mathbf{x}^{2} - \mathbf{x} - 3 \\ -\mathbf{x}^{2} - \mathbf{x} - 3 & 2\mathbf{x}^{2} + 3\mathbf{x} - 5 \end{pmatrix}$$

**Example 18.34.** Next, an example in two variables. The coefficients of x and y in  $f_1$  below are positive definite, but not all entries of  $f_2$  are in  $P_d$  since the homogeneous part has positive and negative coefficients.

$$\begin{split} f_0 &= \begin{pmatrix} 2 & 3 \\ 3 & 6 \end{pmatrix} \\ f_1 &= \begin{pmatrix} 1 & -1 \\ -1 & 2 \end{pmatrix} x + \begin{pmatrix} 3 & 1 \\ 1 & 2 \end{pmatrix} y + I \\ f_2 &= (x + 2y + 1)f_1 - f_0 \\ &= \begin{pmatrix} x^2 + 5yx + 2x + 6y^2 + 5y - 1 & -x^2 - yx - x + 2y^2 + y - 3 \\ -x^2 - yx - x + 2y^2 + y - 3 & 2x^2 + 6yx + 3x + 4y^2 + 4y - 5 \end{pmatrix} \end{split}$$

#### **18.5 Problems with products**

It's reasonable to think that a product such as

$$(xI - S_1)(xI - S_2) \cdots (xI - S_n)$$

should be in Hyp<sub>1</sub>, but this fails, even in the diagonal case. If we take each  $S_i$  to be a diagonal matrix, then the product is a diagonal matrix, and we know that a necessary condition is that there is a common interlacing. To try to satisfy this condition, we can require that all eigenvalues of  $S_1$  are less than all eigenvalues of  $S_2$ , which are less than all eigenvalues of  $S_3$ , and so on. However, this does not always produce hyperbolic matrix polynomials. If we perturb a diagonal matrix that gives a hyperbolic polynomial by conjugating some of the terms, then we can lose hyperbolicity.

If  $e^{\iota \tau} = \begin{pmatrix} \cos \tau & \sin \tau \\ -\sin \tau & \cos \tau \end{pmatrix}$  then the graph of the inner product of the polynomial

$$f(\mathbf{x}) = \left(\mathbf{x}\mathbf{I} - e^{-2\iota} \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix} e^{2\iota}\right) \left(\mathbf{x}\mathbf{I} - e^{-\iota} \begin{pmatrix} 3 & 0 \\ 0 & 4 \end{pmatrix} e^{\iota}\right)$$

is given in Figure 18.2, and is clearly not in  $Hyp_1$ .



Figure 18.2: The graph of a matrix product not in Hyp<sub>1</sub>

Here is an example of a product that is in  $Hyp_1$ .

$$\begin{pmatrix} xI - \begin{pmatrix} 1.24 & -0.42 \\ -0.42 & 1.75 \end{pmatrix} \end{pmatrix} \begin{pmatrix} xI - \begin{pmatrix} 3.97 & -0.15 \\ -0.15 & 3.02 \end{pmatrix} \end{pmatrix} \begin{pmatrix} xI - \begin{pmatrix} 6.99 & -0.04 \\ -0.04 & 6.00 \end{pmatrix} \end{pmatrix}$$

where the eigenvalues of the matrices are (1, 2), (3, 4), (6, 7). The graph of  $(1, y)^t f(1, y)$  is in Figure 18.3. The three dashed lines are the asymptotes, and the dots are the roots of the determinant, drawn on the x-axis. The graph shows that  $f \in Hyp_1$ , since each horizontal line meets the graph in exactly three points.



Figure 18.3: The graph of a hyperbolic matrix polynomial

Some constructions that you would expect to give hyperbolic polynomials give stable matrix polynomials. For instance, if A is positive definite then  $(xI + A)^2$  is generally not hyperbolic, but is a stable matrix polynomial. (See p. 657.)

# 18.6 Sign Interlacing

Recall that f sign interlaces g if the sign alternates as we evaluate f at the roots of g. There is a similar definition for  $f \in Hyp_1$ , where the role of alternating signs is replaced by alternating positive and negative definite matrices.

**Lemma 18.35.** Suppose that  $g \in P$  has roots  $r_1 < \cdots < r_n$ . If  $f \in Hyp_1$  satisfies gI < f then

 $(-1)^{n+i}f(r_i)$  is positive definite for  $1 \leq i \leq n$ 

*Proof.* The hypotheses imply that  $v^t(gI)v = (v^tv)g < v^tfv$ , so  $v^tfv$  sign interlaces g. Thus,  $(-1)^{n+i}v^tf(r_i)v > 0$  for all non-zero v, and hence  $(-1)^{n+i}f(r_i)$  is positive definite.

The converse can be used to show that a matrix polynomial is in  $Hyp_1$ .

**Lemma 18.36.** Suppose that  $g \in P$  has roots  $r_1 < \cdots < r_{n+1}$ . If

- 1.  $(-1)^{n+i}f(r_i)$  is positive definite for  $1 \leq i \leq n+1$
- 2. f has degree n.

3. The leading coefficient of f is positive definite.

*then*  $f \in Hyp_1$  *and*  $gI \leq f$ .

*Proof.* From the first condition we get that  $v^t f v$  sign interlaces g. Since the degree of g is one more than the degree of  $v^t f v$ , we conclude that  $g < v^t f v$ . This implies that  $v^t f v \in \mathbf{P}$ , and that gI < f.

**Construction 18.37.** We can use sign interlacing to construct elements of  $Hyp_1$ . We start with the data

 $\begin{array}{ll} \mbox{real numbers} & p_1 < p_2 < \cdots < p_{n+1} \\ \mbox{positive definite matrices} & m_1, \cdots, m_{n+1} \end{array}$ 

We claim that the following is in Hyp<sub>1</sub>:

$$f(x) = \sum_{k=1}^{n+1} m_k \prod_{i \neq k} (x - p_i)$$
(18.6.1)

In order to verify that  $f \in Hyp_1$  we see that

$$f(p_k) = \prod_{i \neq k} (p_k - p_i) m_k$$

and the sign of the coefficient of  $m_k$  is  $(-1)^{n+k+1}$  since the  $p_i$  are increasing. Since the leading coefficient of f(x) is  $\sum_k m_k$  which is positive definite we can now apply the lemma.

Note that if  $\mu = 1$  this is exactly Lemma 1.20

**Construction 18.38.** There is a similar construction for polynomials in  $Hyp_d$ . The idea is that we construct a polynomial by adding together lots of interlacing polynomials, multiplied by positive definite matrices. So, we start with

positive definite d by d matrices  $D_1, \ldots, D_d$ positive definite d by d matrices  $M_1, \ldots, M_d$ 

Let

$$A = I + x_1 D_1 + \dots + x_d D_d$$
  
f(x) = |A|  
f<sub>i</sub>(x) = determinant of i'th principle submatrix of A  
g(x) = f\_1(x)M\_1 + \dots + f\_d(x)M\_d

Since  $f \leq f_i$  it follows that  $fI \leq f_iM_i$ . Addition yields our conclusion:  $fI \leq g$  in  $Hyp_d$ .

The Hadamard product of two matrix polynomials  $f = \sum A_i x^i$  and  $g = \sum B_i x^i$  is  $f * g = \sum A_i * B_i x^i$  where  $A_i * B_i$  is the usual Hadamard product of matrices. In certain circumstances Hyp<sub>1</sub> is closed under the Hadamard product \* with a positive definite matrix.

**Lemma 18.39.** Suppose that  $f \leq gI$  where  $f \in Hyp_1$  and  $g \in P$ . If P is a positive definite matrix then  $f * P \leq gI$ . In particular,  $f * P \in Hyp_1$ .

*Proof.* If the roots of g are  $r_1 < \cdots < r_n$  then we know that  $(-1)^{n+i} f(r_i)$  is positive definite. Now the Hadamard product of two positive definite matrices is again positive definite [85], so

$$(-1)^{n+i}(f * P)(r_i) = (-1)^{n+i}f(r_i) * P$$

is positive definite, as is the leading coefficient of f \* P. The conclusion follows.  $\Box$ 

The Hadamard product of three hyperbolic matrix polynomials can be in  $Hyp_1$ .

**Lemma 18.40.** If f, g,  $h \in Hyp_1$  have the same common interlacing then  $f * g * h \in Hyp_1$ .

*Proof.* Let  $r_1, \ldots, r_{n+1}$  be the roots of the common interlacing. We know that

$$(-1)^{n+i}f(r_i), \quad (-1)^{n+i}g(r_i), \quad (-1)^{n+i}h(r_i)$$

are all positive definite, and so  $(-1)^{n+i}(f * g * h)(r_i)$  is positive definite. We now apply Lemma 18.36.

#### 18.7 Root zones

Hyperbolic matrix polynomials in one variable are generalizations of polynomials with all real roots. There are three different notions of the zeros of a hyperbolic matrix polynomial f(x), and all reduce to the same idea if the matrix dimension is 1:

- 1. The roots of the diagonal elements.
- 2. The roots of the determinant of f.
- 3. The roots of all the quadratic forms  $v^t f v$ .

What does the graph of a hyperbolic matrix polynomial  $f \in Hyp_1$  look like? When  $\mu = 2$  we can graph  $(1, y)f(1, y)^t$  – Figure 18.3 is a typical picture.

The first point to note is that the graph does not contain large x values. This can be seen as follows. It suffices to consider v satisfying |v| = 1, and that the leading coefficient of f is the identity. In this case  $v^t f v$  is monic and the coefficients of  $v^t f v$  are all bounded (by compactness), and so the roots are bounded.

If a polynomial  $f \in Hyp_1$  has a common interlacing g, then if we graph f and the vertical lines corresponding to the roots of g, then the strip between any two consecutive such vertical lines contains exactly one solution curve of f.

The usual terminology is as follows. If  $f \in Hyp_1$  define  $\Delta_k$  to be the set of all x such that there is a v where x is the k'th largest root of  $v^t f v$ .  $\Delta_k$  is called the k'th *root zone*.

Interlacing of root zones implies interlacing.

**Lemma 18.41.** *If the root zones of two hyperbolic matrix polynomials alternate then the polynomials interlace.* 

*Proof.* Let  $r_1, \ldots, r_n$  be the root zones of r, and  $s_1, \ldots, s_n$  the root zones of s. The hypotheses mean that the zones are arranged as in the diagram

$$\underbrace{\qquad}_{r_1} f_1 \underbrace{\qquad}_{s_1} g_1 \underbrace{\qquad}_{r_2} f_2 \underbrace{\qquad}_{s_2} g_2 \cdots \cdots g_{n-1} \underbrace{\qquad}_{r_n} f_n \underbrace{\qquad}_{s_n} g_n$$

The  $f_i$ 's and the  $g_i$ 's are values that separate the various root zones. Let  $f = \prod(x - f_i)$  and  $g = \prod(x - g_i)$ . The interlacings

easily imply that  $r \leq s$ .

**Example 18.42.** The converse of this lemma is not true – it fails even for diagonal hyperbolic matrix polynomials. Let

$$f = \begin{pmatrix} (x+32)(x+36) & 0\\ 0 & (x+14)(x+35) \end{pmatrix} \qquad f' = \begin{pmatrix} 2(x+34) & 0\\ 0 & 2(x+24.5) \end{pmatrix}$$

We know that  $f \leq f'$  yet their root zones overlap (Figure 18.4).



Figure 18.4: Overlapping root zones

**Lemma 18.43.** The diagonal elements of a hyperbolic polynomial have a common interlacing.

*Proof.* If  $f \in Hyp_1$  then each two by two principle submatrix is in  $Hyp_1$ , and the two diagonal elements have a common interlacing. Consequently, every two diagonal polynomials have a common interlacing, so they all have one.

#### CHAPTER 18. MATRIX POLYNOMIALS

If the degree of a matrix polynomial is greater than two then we do not have nice pictures such as Figure 18.3. We can make another graph by drawing all of the graphs  $v^t f v$  for all non-zero v. The result looks like Figure 18.5. The points on the x axis show that all the curves  $v^t f v$  have a common interlacing. If  $\lambda_{\min}(M)$  and  $\lambda_{\max}(M)$  are the smallest and largest eigenvalues of a symmetric matrix M then we have the inequality

$$\lambda_{\min}(f(x)) \leqslant \frac{\nu^t f \nu}{\nu^t \nu} \leqslant \lambda_{\max}(f(x))$$

The dashed boundary curves in the figure are the graphs of the maximum and minimum eigenvalues of f(x) as a function of x.



Figure 18.5: The graphs of  $v^{t} f v$  as v varies

The Hadamard product preserves  $Hyp_1$  if one of the terms is a diagonal polynomial.

**Lemma 18.44.** If f,  $g \in Hyp_1$  and f is diagonal then  $f * g \in Hyp_1$ .

*Proof.* Since the diagonal elements of f and of g have common interlacings F and G, the result follows from the fact that  $f_{ii} \leq F$  and  $g_{ii} \leq G$  implies (Lemma 7.9)

$$f_{ii} * g_{ii} \leq F * G$$

The roots of the diagonal fall in the root zones. Indeed, if the diagonal elements of f are  $f_i$ , and  $e_i$  is a coordinate vector, then  $e_i^t f e_i = f_i$ .

The location of the roots of the determinant of a hyperbolic matrix polynomial fit nicely into the root zone picture.

**Lemma 18.45.** Suppose that  $f \in Hyp_1(n)$  has a common interlacing. Then, det(f) has  $n\mu$  roots, all real. In the interior of each root zone all f(x) are positive definite or negative definite. There are  $\mu$  zeros of the determinant in each root zone.

*Proof.* This follows from Lemma 18.35, since we know how many positive eigenvalues ( $\mu$  or 0) there are in each region between the root zones. As  $\alpha$ 

procedes from one root of the common interlacing to the next, there is a change from all eigenvalues of one sign to all eigenvalues of the opposite sign. Each time an eigenvalue changes sign, the determinant must be zero.

**Example 18.46.** Suppose that  $\mu = n = 3$ , and that the roots of det(f) are  $r_1, \ldots, r_9$ . Then

interval	number of positive roots
$(-\infty, \mathbf{r}_1)$	0
$(r_1, r_2)$	1
$(r_2, r_3)$	2
$(r_3, r_4)$	3
$(r_4, r_5)$	2
$(r_5, r_6)$	1
$(r_6, r_7)$	0
$(r_7, r_8)$	1
$(r_8, r_9)$	2
$(\mathbf{r}_9, \infty)$	3

We can show that a polynomial is in  $Hyp_1$  by checking if all roots of the determinant are real, and that the number of positive eigenvalues in each interval follows the patterns described above.

#### 18.8 Quadratics and Newton inequalities

Quadratic matrix polynomials have been widely studied [111]. We derive a simple inequality for eigenvalues of the quadratic, and use it in the usual way to get Newton inequalities for hyperbolic matrix polynomials.

There are necessary and sufficient conditions for a quadratic matrix polynomials to belong to Hyp<sub>1</sub>, but they are not as elementary as for polynomials.

**Lemma 18.47.** Suppose  $f(x) = Ax^2 + Bx + C$  where A and C are non-negative definite.

1. If 
$$f \in Hyp_1$$
 then  $\lambda_{max}(BA^{-1})\lambda_{max}(BC^{-1}) \ge 4$ 

2. If  $\lambda_{\min}(BA^{-1}) \lambda_{\min}(BC^{-1}) \ge 4$  then  $f \in Hyp_1$ .

*Proof.* If  $f \in Hyp_1$  then

$$v^{t}Av x^{2} + v^{t}Bv x + v^{t}Cv \in \mathbf{P}$$

Thus the discriminant must be non-negative, and so

$$(\mathbf{v}^{\mathsf{t}}\mathsf{B}\mathbf{v})^2 \ge 4(\mathbf{v}^{\mathsf{t}}\mathsf{A}\mathbf{v})(\mathbf{v}^{\mathsf{t}}\mathsf{C}\mathbf{v})$$

Now  $v^t A v$  and  $v^t C v$  are non-negative, so this is equivalent to

$$\frac{\nu^{t}B\nu}{\nu^{t}A\nu} \ge 4\frac{\nu^{t}C\nu}{\nu^{t}B\nu}$$

We now apply (18.2.3).

**Corollary 18.48.** Suppose that  $f = \sum A_i x^i \in Hyp_1^{pos}(n)$ . For k=1,...,n-1 the following inequality holds:

$$\lambda_{\max}(A_k A_{k+1}^{-1}) \lambda_{\max}(A_k A_{k-1}^{-1}) \ge (1 + \frac{1}{k})(1 + \frac{1}{n-k})$$

*Proof.* Since  $Hyp_1^{pos}$  is closed under differentiation and reversal, we can follow the usual proof of Newton's inequalities.

The coefficients of the associated quadratic form are not arbitrary. Suppose that  $f(x) = \sum A_i x^i$  is a matrix polynomial in  $Hyp_1^{pos}(n)$ . We know that for all  $v \neq 0$  and  $0 \leq i \leq n$ 

$$\lambda_{i}^{\min} \leqslant \frac{\nu^{t} A \nu}{\nu^{t} \nu} \leqslant \lambda_{i}^{\max}$$

Define the two vectors

$$\Lambda_{\min} = (\lambda_0^{\min}, \dots, \lambda_n^{\min})$$
$$\Lambda_{\max} = (\lambda_0^{\max}, \dots, \lambda_n^{\max})$$

The inequality above shows that for  $|\nu| = 1$  all the polynomials  $\nu^t f \nu$  are polynomials of the form poly(c) where  $0 \leq \Lambda_{min} \leq c \leq \Lambda_{max}$ . (See p. 39.).

# **18.9** Newton inequalities in Hyp<sub>2</sub>

The coefficients of a quadratic in  $Hyp_2$  satisfy inequalities, and they lead to inequalities for general polynomials in  $Hyp_2$ . We begin with an example. Consider the polynomial f(x, y) (p. 552):



If we compute  $v^t f v$  then we get a polynomial in P<sub>2</sub>. For instance, if we take v = (2,3) we get



This polynomial satisfies the rhombus inequalities, so we can use this information to get inequalities for the original polynomial. In general, we start with  $f(x, y) = \sum A_{i,j} x^i y^j$  in  $Hyp_2^{pos}$ . If v is any vector then

$$\nu^t f \nu = \sum \left( \nu^t A_{i,j} \nu \right) x^i y^j \in \mathbf{P}_2$$

The rhombus inequalities (Proposition 10.41) imply that the following holds for all vectors v

$$(v^{t} A_{i,j+1}v) (v^{t} A_{i+1,j}v) > (v^{t} A_{i,j}v) (v^{t} A_{i+1,j+1}v)$$

Using (18.2.3) this implies rhombus inequalities in  $Hyp_2$ :

$$\lambda_{\max}(A_{i,j+1}A_{i,j}^{-1}) \geqslant \lambda_{\min}(A_{i+1,j+1}A_{i+1,j}^{-1})$$

## 18.10 Singular polynomials

We now look at some examples of matrix polynomials f for which the leading coefficient of a matrix polynomial is positive semi-definite and singular, and  $v^{t} f v \in \mathbf{P}$  for all  $v \in \mathbb{R}^{\mu}$ . Unlike hyperbolic matrix polynomials, the root zones can be infinite, or all of them can even be equal.

First, we consider the matrix  $F = \begin{pmatrix} f & h \\ h & 0 \end{pmatrix}$  where deg(f) > deg(h). If f is monic then the leading coefficient of F is  $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$ , which has eigenvalues 0 and 1, and is positive semi-definite. The quadratic form is  $\alpha^2 f + 2\alpha\beta h$  which is in **P** for all  $\alpha$ ,  $\beta$  if and only if f < h. Figure 18.6 is the graph of such a polynomial.



Figure 18.6: Positive semi-definite leading coefficient

Assume that  $f \leq h$ . If  $\varepsilon > 0$  and  $F_{\varepsilon} = \begin{pmatrix} f & h \\ h & \varepsilon f \end{pmatrix}$  then  $F_{\varepsilon} \in Hyp_1$  since its leading coefficient is  $\begin{pmatrix} 1 & 0 \\ 0 & \varepsilon \end{pmatrix}$ , and the quadratic form is a linear combination of f and h. Consequently, F is the limit of polynomials in  $Hyp_1$ . Note that F has a common interlacing, but the root zones can be unbounded.

Next, we have an example with no common interlacing. Let

$$F = \begin{pmatrix} (x+1)(x+3) & 3x+5\\ 3x+5 & 8 \end{pmatrix}.$$

The leading coefficient is again  $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$  and the quadratic form factors:

$$(\alpha, \beta) F(\alpha, \beta)^{t} = (\alpha(x+1) + 2\beta)(\alpha(x+3) + 4\beta).$$

Consequently, for every  $x_0$  there are two v such that the quadratic form  $v^t F(x_0)v$  is zero. Thus, the two root zones are both equal to  $\mathbb{R}$ , there is no common interlacing, and so F is not a limit of polynomials in Hyp<sub>1</sub>.

# **18.11** Matrix polynomials with determinants in P<sub>d</sub>

In this last section we consider the properties the set of matrix polynomials with determinants in  $\overline{\mathbf{P}}_d$ . Define

$$\mathcal{M} = \{ \text{matrix polynomial } M \mid |M| \in \overline{\mathbf{P}}_d \text{ for some } d \}$$

If y is a variable not occuring in A or B then we say that  $A \leftrightarrow B$  if and only if  $A + yB \in M$ .

If the matrices are 1 by 1 then this is just the usual definition of interlacing. One interesting question is what properties of  $\overline{\mathbf{P}}_d$  extend to  $\mathcal{M}$ ? Clearly  $\mathcal{M}$  is closed under substitution of real values for variables, and of  $x_i + a$  for  $x_i$ .

Here are some basic interlacing properties:

#### **Lemma 18.49.** *If* $A \leftrightarrow B$ *then*

- 1. B ↔ −A
- 2.  $A + \alpha B \leftrightarrow B$  for  $\alpha \in \mathbb{R}$ .
- 3.  $A \leftrightarrow B + \alpha A$  for  $\alpha \in \mathbb{R}$ .
- 4. If  $C \in M$  then  $AC \Leftrightarrow BC$  and  $CA \Leftrightarrow CB$ .

*Proof.* If  $|A + yB| \in \overline{\mathbf{P}}_d$  then reversing just y yields  $|yA - B| \in \overline{\mathbf{P}}_d$ . Next, replacing y by  $y + \alpha$  shows that  $|A + (y + \alpha)B| \in \overline{\mathbf{P}}_d$ . From  $B \leftrightarrow -A$  we get that  $B + \alpha(-A) \leftrightarrow -A$ , and applying the first part again gives  $A \leftrightarrow B + (-\alpha)A$ . For the next one, note that |AC + yBC| = |C||A + yB|.

Some constructions:

#### Lemma 18.50.

1. If A is symmetric, all  $D_i$  and B are positive semi-definite, then

$$A + \sum x_i D_i \nleftrightarrow B.$$

- 2. If  $Z = diag(z_1, ..., z_d)$  and A is positive semi-definite then
  - a)  $Z + A \in \mathcal{M}$ b)  $Z - A \in \mathcal{M}$

c) 
$$AZ + I \in M$$
  
d)  $ZA + I \in M$   
e) If  $W = diag(w_i)$  and O is orthogonal then  $OZ + WO \in M$ ,

*Proof.* The first one is the statement that  $|A + yB + \sum x_iD_i| \in \overline{\mathbf{P}}_d$ . For the second, reverse each variable separately, and then reverse all at once. The next follows from the fact that  $A + Z = A + \sum z_iD_i$  where  $D_i$  is the positive semi-definite matrix that is all zero except for a 1 in location (i, i). If we reverse all variables we have

$$z_1 \cdots z_d |A + Z^{-1}| = |AZ + I| = |ZA + I|.$$

For the last one,  $Z + O^{-1}WO$  is a sum of variables times positive semi-definite matrices.

If a < b then  $x + a \ll x + b$ . The corresponding result for  $\mathcal{M}$  is

**Lemma 18.51.** If A < B are positive definite and  $Z = diag(z_i)$  then

$$A + Z \leftrightarrow B + Z$$
.

*Proof.* Since B – A is positive definite, Lemma 18.50 implies that

 $A + Z \leftrightarrow B - A.$ 

Applying Lemma 18.49 yields the conclusion

$$A + Z \leftrightarrow B - A + (A + Z) = B + Z.$$

**Corollary 18.52.** *With the same hypotheses,*  $BZ + I \leftrightarrow AZ + I$ .

Proof. Reverse all variables.

As [16] observed, it is an interesting question if A + Z  $\leftrightarrow$  B + Z implies A  $\leqslant$  B.
Part III

## Polynomials with complex roots

CHAPTER



# Polynomials non-vanishing in a region

In this chapter we introduce polynomials that are non-vanishing (NV) in a region  $\mathcal{D}$ . We show how geometric properties of  $\mathcal{D}$  imply properties of the sets of polynomials such as closure under differentiation, or under reversal.

**Definition 19.1.** Suppose that  $\mathcal{D} \subsetneq \mathbb{C}$ . Define

 $\begin{aligned} \mathsf{NV}_{d}(\mathcal{D}) &= \big\{ \mathsf{f} \mid \mathsf{f} \text{ has complex coefficients and } \mathsf{f}(\sigma) \neq 0 \text{ for all } \sigma \in \mathcal{D} \big\} \\ \mathsf{rNV}_{d}(\mathcal{D}) &= \big\{ \mathsf{f} \mid \mathsf{f} \text{ has real coefficients and } \mathsf{f}(\sigma) \neq 0 \text{ for all } \sigma \in \mathcal{D} \big\} \end{aligned}$ 

Here are some examples of the spaces we will consider.

**Example 19.2.** If  $\mathcal{D}$  is the product of d open upper half planes, then a polynomial in NV<sub>d</sub>( $\mathcal{D}$ ) is called a *upper half plane polynomial*. It is a *real upper half-plane polynomial* if it is in rNV<sub>d</sub>( $\mathcal{D}$ ). The spaces of such polynomials are denoted HP<sub>d</sub> and rHP<sub>d</sub>. Many results about these polynomials can be found in [23], [18] and Chapter 20.

**Example 19.3.** If  $\mathcal{D}$  is a product of d open right half planes then a polynomial in NV<sub>d</sub>( $\mathcal{D}$ ) is called a *right half-plane polynomial*, or a *Hurwitz polynomial*. NV<sub>d</sub>( $\mathcal{D}$ ) is denoted  $\mathcal{H}_d(\mathbb{C})$  and rNV<sub>d</sub>( $\mathcal{D}$ ) is denoted  $\mathcal{H}_d$ . In the  $\mathcal{H}_1(\mathbb{C})$  case, a right half-plane polynomial has all roots in the closed left half plane. Such polynomials are usually called stable, or Hurwitz-stable. More information about these polynomials can be found in [33] and Chapter 21.

**Example 19.4.** If S is a vertical strip such as  $\{z \mid -1 < \Re(z) < 1\}$  then we consider  $\mathcal{D} = \mathbb{C} \setminus S$ . The set  $\mathsf{NV}_1(\mathbb{C} \setminus S)$  consists of polynomials whose zeros all lie in the strip S. More information about them can be found in Section 19.10.

**Example 19.5.** The open unit disk  $\Delta$  in  $\mathbb{C}$  has two generalizations to  $\mathbb{C}^d$ . The first is the open ball  $\Delta_d$  of radius 1, and the second is the product  $\Delta \times \cdots \times \Delta$ . The spaces are considered in Section 19.11.

**Example 19.6.** Define the sector  $S_n = \{z \mid -\pi/n < \arg(z) < \pi/n\}$ . For instance,  $S_1$  is  $\mathbb{C} \setminus (-\infty, 0)$ , and  $S_2$  is the open right half plane. We consider properties of product of sectors  $\mathsf{NV}_d(\prod_i (S_{n_i}))$  in various examples in this chapter and Section 19.12.

#### **19.1** Fundamental properties

In order to derive interesting properties we we usually assume that our domain is an open cone  $\mathcal{C}$  in  $\mathbb{C}^d$ . This means  $\mathcal{C}$  is closed under addition and multiplication by positive constants. We have a few properties that hold for general domains. The proofs are immediate from the definition.

#### Lemma 19.7.

- 1.  $f(\mathbf{x})g(\mathbf{x}) \in NV_d(\mathcal{D})$  if and only if  $f(\mathbf{x}) \in NV_d(\mathcal{D})$  and  $g(\mathbf{x}) \in NV_d(\mathcal{D})$ .
- 2. Suppose  $f(\mathbf{x}, \mathbf{y}) \in \mathsf{NV}_{d+e}(\mathcal{D} \times \mathcal{E})$  where  $\mathcal{D} \subset \mathbb{C}^d$  and  $\mathcal{E} \subset \mathbb{C}^e$ . If  $\alpha \in \mathcal{E}$  then  $f(\mathbf{x}, \alpha) \in \mathsf{NV}_d(\mathcal{D})$ .

The following classical theorem of Hurwitz also holds for analytic functions in d variables.

**Theorem 19.8** (Hurwitz). Let  $(f_n)$  be a sequence of functions which are all analytic and without zeros in a region  $\Omega$ . Suppose, in addition, that  $f_n(z)$  tends to f(z), uniformly on every compact subset of  $\Omega$ . Then f(z) is either identically zero or never equal to zero in  $\Omega$ .

The next lemma is an immediate consequence of the general Hurwitz's theorem:

**Lemma 19.9** (Polynomial closure). *If*  $\mathcal{D}$  *is open and* { $f_i$ } *is a sequence of polynomials in*  $NV_d(\mathcal{D})$  *that converge uniformly on compact subsets to a polynomial* g *then either* g *is zero or is in*  $NV_d(\mathcal{D})$ .

The next lemma gives several simple ways to construct new polynomials in  $NV_d(\mathcal{C})$  from old ones. The proofs easily follow from the definition.

**Lemma 19.10.** Assume that C is a cone.

- 1. If  $f(\mathbf{x}) \in NV_d(\mathbb{C})$  then  $f(\mathbf{x} + \mathbf{y}) \in NV_{2d}(\mathbb{C} \times \mathbb{C})$  where  $\mathbf{x} = (x_1, \dots, x_d)$  and  $\mathbf{y} = (y_1, \dots, y_d)$ .
- 2. If a is a positive constant and  $f(\mathbf{x}) \in NV_d(\mathcal{C})$  then  $f(a x_1, ..., a x_d) \in NV_d(\mathcal{C})$ .
- 3. If  $a_i$  are positive constants,  $C_i$  are cones, and  $f(\mathbf{x}) \in NV_d(\prod C_i)$  then  $f(a_1 x_1, \ldots, a_d x_d) \in NV_d(\prod C_i)$ .

C	$\mathcal{C}^{-1}$	ρ	$\mathcal{C}/\mathcal{C}$
UHP	-UHP	-1	$\mathbb{C}\setminus(-\infty,0)$
RHP	RHP	1	$\mathbb{C}\setminus(-\infty,0)$
$Q_{\mathrm{I}}$	$Q_{\mathrm{IV}}$	l	RHP
Sn	Sn	1	S <sub>2n</sub>

Table 19.1: Properties of cones

The homogeneous part  $f^H$  of a polynomial consists of the terms of maximum total degree.

**Corollary 19.11.** *If*  $f \in NV_d(H)$   $\mathcal{C}$  *then*  $f^H \in NV_d(\mathcal{C})$ .

*Proof.* This follows from the above, the fact that f<sup>H</sup> is never zero and

$$f^{\mathsf{H}}(x) = \lim_{r \to \infty} \frac{f(rx)}{r^n}$$

where n is the degree of  $f^{H}$ .

The next result refines the earlier substitution result.

**Lemma 19.12** (Substitution). Suppose that  $\mathbb{D}$  is open and  $\overline{\mathbb{D}}$  is the closure of  $\mathbb{D}$ . If  $f(\mathbf{x}, \mathbf{y}) \in \mathsf{NV}_{d+e}(\mathbb{D} \times \mathcal{E})$  and  $\alpha \in \overline{\mathbb{D}}$  then  $f(\alpha, \mathbf{y}) \in \mathsf{NV}_{e}(\mathcal{E})$  or  $f(\alpha, \mathbf{y}) = 0$ .

*Proof.* If  $\alpha \in \mathcal{D}$  then  $f(\alpha, \mathbf{y}) \in \mathsf{NV}_d(\mathcal{E})$ . If  $\alpha \in \overline{\mathcal{D}}$  then we approach  $f(\alpha, \mathbf{y})$  by sequences in  $\mathsf{NV}_d(\mathcal{D} \times \mathcal{E})$ , so we can apply Lemma 19.9.

The constant in the next lemma only depends on  $\mathbb{C}$  and is called the *reversal constant*.

**Lemma 19.13** (Reversal). Assume  $\mathbb{C} \subset \mathbb{C}$  is a cone and suppose that  $\sum x^i f_i(\mathbf{y}) \in NV_{d+1}(\mathbb{C} \times D)$ . There is a unique  $\rho \in \mathbb{C}$  with  $|\rho| = 1$  such that

$$\sum (\rho x)^{n-i} f_i(\mathbf{y}) \in \textit{NV}_{d+1}(\mathcal{C} \times \mathcal{D}).$$

*Proof.* If  $\mathcal{C}^{-1} = \{z^{-1} \mid z \in \mathcal{C}_1\}$  then  $\mathcal{C}^{-1}$  is also a cone, whose elements are the conjugates of  $\mathcal{C}$ . Since  $\mathcal{C}$  and  $\mathcal{C}^{-1}$  only differ by a rotation there is a unique rotation  $\rho \in \mathbb{C}$  so that  $\rho \mathcal{C}^{-1} = \mathcal{C}$ . If  $(\sigma_1, \ldots, \sigma_{d+1}) \in \mathcal{C} \times \mathcal{D}$  then

$$\sum (\rho \sigma_1)^{n-i} f_i(\sigma_2, \dots, \sigma_{d+1}) = (\rho \sigma_1)^n \sum \left(\frac{\rho}{\sigma_1}\right)^i f_i(\sigma_2, \dots, \sigma_{d+1})$$

and this is non-zero since  $\rho/\sigma_1 \in \mathbb{C}$ . See Table 19.1.

We now consider when a linear polynomial is in  $\mathsf{NV}_d(\mathcal{D})$ . The following are elementary.

**Lemma 19.14.** *The following are equivalent, where*  $C_i \subset \mathbb{C}$  *are cones.* 

- 1.  $\sum_{1}^{d} a_i x_i \in NV_d(\prod C_i).$
- 2.  $0 \notin \sum_{1}^{d} a_i \mathcal{C}_i$
- 3.  $\sum_{1}^{d} a_i C_i$  is contained in a half plane.

**Example 19.15.** Suppose  $\mathbb{C} = \mathbb{S}_2 \times \mathbb{S}_4$  and consider  $x + \alpha y$ . In order for  $\mathbb{S}_2 + \alpha \mathbb{S}_4$  to remain in a half plane, we must have that  $\alpha \mathbb{S}_4 \subset \mathbb{S}_2$ . Thus we conclude that

$$x + \alpha y \in \mathsf{NV}_2(\mathbb{S}_2 \times \mathbb{S}_4) \Longleftrightarrow -\pi/4 \leqslant \alpha \leqslant \pi/4$$

If r > 0 and  $\mathcal{C} \subset \mathbb{C}$  then  $x + ry \in \mathsf{NV}_2(\mathcal{C}^2)$ , and hence  $\rho xy + r$  and  $xy + r/\rho$  are in  $\mathsf{NV}_2(\mathcal{C}^2)$ .

There's a simple condition if all the planes are half planes.

**Lemma 19.16.** *If all cones*  $C_i$  *are the same half plane* C*, then the following are equivalent:* 

- 1.  $\sum_{1}^{d} a_i x_i \in NV_d(\mathbb{C}^d)$ .
- 2. All non-zero  $a_i$  have the same argument.

*Proof.* If  $\sum \alpha_i x_i \in \mathsf{NV}_d(\mathbb{C}^d)$  then we may assume that  $\alpha_1$  is non-zero, so we may divide through and assume  $\alpha_1 = 1$ . We now need to show that all non-zero  $\alpha_i$  are positive. If some  $\alpha_i$  is not positive, then  $\mathbb{C}$  and  $\alpha \mathbb{C}$  are different, and so their sum equals  $\mathbb{C}$ . The converse is immediate.

The corollary is simple and the proof is omitted.

**Corollary 19.17.** *If* C *is a half plane then*  $\beta + x_1 + \alpha_2 x_2 + \cdots + \alpha_d x_d \in NV_d(C^d)$  *if and only if* 

- 1. All  $\alpha_i$  are non-negative.
- 2.  $\beta$  is in the closure of C.
- Following [18,33] we have

**Lemma 19.18.** Suppose  $\mathbb{C} \subset \mathbb{C}$  and  $f(\mathbf{x})$  is homogeneous. The following are equivalent

- 1.  $f(\mathbf{x}) \in NV_d(\mathbb{C}^d)$
- 2.  $f(\mathbf{a} + t\mathbf{b}) \in NV_1(\mathbb{C}/\mathbb{C})$  where  $\mathbf{a}, \mathbf{b} > 0$ .

*Proof.* C/C denotes the set { $\alpha/\beta \mid \alpha, \beta \in C$ }. If  $\alpha/\beta \in C/C$  where  $\alpha, \beta \in C$  and f has degree n then by homogeneity

$$f(\mathbf{a} + (\alpha/\beta)\mathbf{b}) = \beta^{n} f(\beta \mathbf{a} + \alpha \mathbf{b}) \neq 0$$

since  $\beta \mathbf{a} + \alpha \mathbf{b} \in \mathbb{C}^d$ . Conversely, let  $\sigma_1, \ldots, \sigma_d \in \mathbb{C}$ . Since  $\mathbb{C}$  is a cone there are  $\alpha, \beta \in \mathbb{C}$  and  $\mathbf{a}, \mathbf{b} > 0$  such that  $\beta \mathbf{a} + \alpha \mathbf{b} = (\sigma_1, \ldots, \sigma_d)$ . Thus

$$f(\sigma_1,\ldots,\sigma_d) = f(\beta \mathbf{a} + \alpha \mathbf{b}) = \beta^n f(\mathbf{a} + \mathbf{b}(\alpha/\beta)) \neq 0$$

#### 19.2 Differentiation

We can not differentiate with respect to a variable and stay in the same class, but we can if the corresponding factor is a half plane.

**Lemma 19.19.** If  $f(x, y) \in NV_{d+1}(\mathbb{C} \times \mathbb{D})$  and  $\mathbb{C}$  is a half plane then  $\frac{\partial}{\partial x}f(x, y) \in NV_{d+1}(\mathbb{C} \times \mathbb{D})$ .

*Proof.* Suppose that  $(\alpha_1, \ldots, \alpha_{d+1}) \in \mathbb{C} \times \mathcal{D}$ . Since

$$\left(\frac{\partial f}{\partial x}\right)(\alpha_1,\ldots,\alpha_{d+1}) = \left[\frac{\partial}{\partial x}(x,\alpha_2,\ldots,\alpha_{d+1})\right](\alpha_1)$$

it suffices to assume that d = 1. So, if  $g \in NV_1(\mathcal{C})$  then all roots lie in the complement of  $\mathcal{C}$ . Now  $\mathcal{C}$  is a half plane by hypothesis, so its complement is a closed half plane, and in particular is convex. By Gauss-Lucas the roots of g' lie in the convex hull of the roots of g, and hence lie in the complement of  $\mathcal{C}$ . Consequently,  $g' \in NV_1(\mathcal{C})$ .

**Lemma 19.20.** If C is a half plane,  $f(x, y) = \sum x^i f_i(y) \in NV_{d+1}(C \times D)$  then either  $f_i(y) = 0$  or

$$f_i(\mathbf{y}) \in NV_d(\mathcal{D})$$
 for  $i = 0, 1, ...$ 

*Proof.* Differentiate with respect to x until the desired coefficient is the constant term, and then substitute zero for x.  $\Box$ 

**Example 19.21.** The assumption that C is a half plane is essential. We give an example of a polynomial in NV<sub>2</sub>(S<sub>4</sub>) whose coefficients are not all in NV<sub>1</sub>(S<sub>4</sub>). Let

$$f(x) = (x - 1)^2 + 4 = (x - 1 - 2\iota)(x - 1 + 2\iota).$$

Clearly  $f(x) \in \mathsf{NV}_1(S_4)$ , but  $f'(x) = 2(x-1) \notin \mathsf{NV}_1(S_4)$ . Now consider

$$f(x + y) = (x - 1)^2 + 4 + 2y(x - 1) + y^2$$

We know  $f(x + y) \in NV_2(S_4)$ , and the coefficient of  $y^0$  in f(x + y) is in  $NV_1(S_4)$  but the coefficient of  $y^1$  is not.

**Lemma 19.22.** *If*  $\sum x^i f_i(\mathbf{y}) \in NV_{d+1}(\mathbb{C} \times \mathbb{D})$  and  $\mathbb{C}$  is a half plane then

$$f_i(\mathbf{y}) + x f_{i+1}(\mathbf{y}) \in NV_{d+1}(\mathcal{C} \times \mathcal{D})$$
 for  $i = 0, 1, ...$ 

*Proof.* Because C is a half plane we can differentiate and reverse, so all the polynomials below are in  $NV_{d+1}(C \times D)$ .

$$\begin{array}{ll} \mbox{reversing} & \sum (\rho x)^{n-i} f_i(\mathbf{y}) \\ n-k-1 \mbox{ differentiations} & \sum (\rho)^{n-i} x^{k+1-i} (\underline{n-i})_{n-k-1} f_i(\mathbf{y}) \\ \mbox{ reverse} & \sum (\rho)^{n-i} (\rho x)^i (\underline{n-i})_{n-k-1} f_i(\mathbf{y}) \\ \mbox{ k differentiations} & \sum (\rho)^{n-i} (\rho x)^{i-k} (\underline{n-i})_{n-k-1} (\underline{i})_k f_i(\mathbf{y}) \\ \mbox{ which equals} & \rho^n \big( f_k(\mathbf{y}) (n-k)! k! + x f_{i+1}(\mathbf{y}) (n-k-1)! (k+1)! \big) \end{array}$$

Dividing and rescaling yields the result.

**Corollary 19.23.** *If*  $f(\mathbf{x}) \in NV_{d+1}(\mathbb{C} \times \mathbb{D})$  *and*  $\mathbb{C}$  *is a half plane then* 

$$f(\mathbf{x}) + y \frac{\partial f}{\partial x_1} \in NV_{d+2}(\mathcal{C} \times \mathcal{C} \times \mathcal{D})$$
  
$$\rho y f(\mathbf{x}) + \frac{\partial f}{\partial x_1} \in NV_{d+2}(\mathcal{C} \times \mathcal{C} \times \mathcal{D})$$

where  $\rho$  is the reversal constant.

*Proof.* Since  $f(x_1 + y, x_2, ..., x_d) \in \mathsf{NV}_{d+2}(\mathbb{C} \times \mathbb{C} \times \mathbb{D})$  we can apply the lemma to the Taylor series

$$f(x_1 + y, \cdots, x_d) = f(x) + \frac{\partial f}{\partial x_1}y + \cdots$$

The second statement follows by reversing the first.

We have seen that coefficients of the homogeneous part of a polynomial in  $\overline{\mathbf{P}}_d$  all have the same sign. A similar result holds for  $\mathsf{NV}_d(\mathcal{C})$ : if we multiply by the appropriate scalar then all the coefficients are positive:

**Lemma 19.24.** If C is a half plane and  $f \in NV_d(C^d)$  then all non-zero coefficients of  $f^H$  have the same argument.

*Proof.* We first assume that f is homogeneous and use induction on d. For d = 2 we note that  $x + ay \in NV_2(\mathbb{C}^2)$  if and only if  $a \ge 0$ . Thus if  $f \in NV_2(\mathbb{C}^2)$  is homogeneous then

$$f = \alpha \prod_i (x + \alpha_i y)$$
 where all  $\alpha_i \geqslant 0$ 

so all non-zero coefficients have argument  $arg(\alpha)$ .

If  $f(\mathbf{x}, \mathbf{y}) \in \mathsf{NV}_{d+1}(\mathbb{C}^{d+1})$  is homogeneous then we can write

$$f(\mathbf{x},\mathbf{y}) = \sum_{i} y^{i} f_{i}(\mathbf{x})$$

where all  $f_i \in NV_d(\mathbb{C}^d)$  and are homogeneous. By induction all coefficients of  $f_i$  have the same argument.

By Lemma 19.18 with a = (1, ..., 1, 0) and b = (0, ..., 0, 1) we have that

$$f(1\ldots,1,t) = \sum t^i f_i(1,\ldots,1) \in \mathbf{P}^{\text{pos}}$$

since  $\mathcal{C}/\mathcal{C} = \mathbb{C} \setminus (-\infty, 0)$ . Now the argument of  $f_i(1, \ldots, 1)$  equals the argument of the coefficients of  $f_i$ , and since  $f(1, \ldots, 1, t) \in \mathbf{P}^{\text{pos}}$  all coefficients have the same argument.

If f is not homogeneous and has degree n then the conclusion follows from Corollary 19.11.  $\hfill \Box$ 

#### 19.3 Interlacing

Interlacing was initially defined by the positions of roots, and then by linearity. Now we define it algebraically.

**Definition 19.25.** Suppose  $f, g \in NV_d(\mathcal{D})$ . We say that g *interlaces* f *in*  $\mathcal{C}$ , written  $f \nleftrightarrow g$ , if and only if

$$f(\mathbf{x}) + yg(\mathbf{x}) \in \mathsf{NV}_{d+1}(\mathcal{D} \times \mathcal{C})$$

*Remark* 19.26. Note that if  $C' \subset C$  then interlacing in C implies interlacing in C', but the converse is false. For example, we claim that

$$x^2 + y(ax + b) \in \mathsf{NV}_2(S_4 \times S_4)$$
 if  $a, b > 0$ 

This is clear, since both the image of  $S_4$  by  $x^2$  and y(ax+b) lies in  $S_2$ . However, we also have that  $f(x, y) = x^2 + y(x+1)$  is not in  $NV_2(S_2 \times S_4)$ . To see this, we need to exhibit one zero in  $S_2 \times S_4$ :

f( 
$$(1+2\iota)/8$$
 ,  $(19-42\iota)/680$  ) = 0

However, it's not had to check that  $x^2 + y(x + \iota) \in NV_2(S_2 \times S_4)$ , and therefore is also in  $NV_2(S_4 \times S_4)$ 

We first have some properties of interlacing that do not require one of the cones to be a half plane.

**Lemma 19.27.** Assume that f, g,  $h \in NV_d(\mathcal{D})$ , and  $\mathcal{C}$  is a cone.

- 1. fg <~ fh in C iff g <~ h in C.
- 2. If  $-1 \notin \mathbb{C}$  then  $f \nleftrightarrow f$  in  $\mathbb{C}$ .
- 3. If  $\rho$  is the reversing constant for C then f  $\leftrightarrow \rho$  g in C implies  $\rho g \leftrightarrow \rho$  f in C.
- 4. If  $0 \notin C_1 + C_2$  and  $f \in NV_2(C_1 \times C_2)$  then  $xf \nleftrightarrow f$  in  $C_2$ .

*Proof.* The first is trivial. To check if  $f \Leftrightarrow f$  we need to see that  $f(\mathbf{x}) + y f(\mathbf{x}) = f(\mathbf{x}) (y+1)$  is non-zero. Now  $f(\mathbf{x}) \neq 0$ , and  $y+1 \neq 0$  by hypothesis. For the next one,  $f + yg \in \mathsf{NV}_{d+1}(\mathbb{C} \times \mathbb{C}_0)$  implies that  $\rho g + f \in \mathsf{NV}_{d+1}(\mathbb{C} \times \mathbb{C}_0)$ .

Finally, to check  $xf \Leftrightarrow f$  we check that  $xf + yf \in \mathsf{NV}_2(\mathfrak{C}_1 \times \mathfrak{C}_2)$ . Simply note that  $(x + y)f \neq 0$  by hypothesis.

Most interesting results about interlacing require a half plane. As always, we expect adjacent coefficients as well as the derivative to interlace. The corollary is a consequence of Lemma 19.22 and Corollary 19.23.

**Corollary 19.28.** Suppose  $\mathcal{C}$  is a half plane and  $f = \sum x^i f_i(\mathbf{y}) \in NV_d(\mathcal{C} \times \mathcal{D})$ .

- 1.  $f_i \nleftrightarrow f_{i+1}$  for i = 0, 1, ...
- 2. f  $\leftrightarrow \frac{\partial f}{\partial x}$ .

**Lemma 19.29.** *If* C *is a half plane and*  $f, g, h, k \in NV_d(D)$  *then (all interlacings in* C)

- 1. f  $\leftrightarrow$  g and f  $\leftrightarrow$  h implies f  $\leftrightarrow$  ag + bh for a, b > 0.
- 2.  $f \leftrightarrow g$  and  $h \leftrightarrow k$  implies  $fh \leftrightarrow fk + gh \leftrightarrow gk$ .
- 3.  $g \nleftrightarrow f$  and  $h \nleftrightarrow f$  implies  $ag + bh \nleftrightarrow f$  for a, b > 0.
- 4. If  $f \nleftrightarrow g \nleftrightarrow h$  then  $f + \rho h \nleftrightarrow g$ .

*Proof.* All these results follow using the interlacing of adjacent coefficients. For the first we have

$$(f + yg)(f + yh) = f2 + yf(g + h) + y2gh$$

so  $f^2 \nleftrightarrow f(g + h)$  and therefore  $f \nleftrightarrow g + h$ . The second is similar. For the third we use

$$(g + y\rho f)(g + y\rho h) = g^2 + gy\rho(f + h)$$

so  $g \nleftrightarrow \rho(f + h)$  and reversing again yields  $f + h \nleftrightarrow g$ . The last follows from the third since  $\rho h \nleftrightarrow g$ .

Next is the recurrence for orthogonal-type polynomials.

**Lemma 19.30.** Assume C is a half plane and  $-1 \notin C$ . If  $f_0, f_1 \in NV_d(C \times D)$  then

$$\begin{split} &f_1 \nleftrightarrow f_0 \\ &f_n = (a_n x_1 + b_n) f_{n-1} + \rho \, c_n \, f_{n-2} \quad \textit{where } a_n, b_n, c_n > 0. \end{split}$$

*then*  $f_n \leftrightarrow f_{n-1}$  *in*  $\mathcal{C}$ .

*Proof.* By induction we may assume that  $f_{n-1} \nleftrightarrow f_{n-2}$ . Since  $-1 \notin \mathcal{C}$  we know  $b_n f_{n-1} \nleftrightarrow f_{n-1}$ . Since  $\mathcal{C} + \mathcal{C}$  doesn't contain zero we know  $x_1 f_{n-1} \nleftrightarrow f_{n-1}$ , and thus  $(a_n x_1 + b_n) f_{n-1} \twoheadleftarrow f_{n-1}$ . By reversal of  $f_{n-1} \twoheadleftarrow f_{n-2}$  we have  $\rho f_{n-2} \twoheadleftarrow f_{n-1}$ , so addition gives the conclusion.

**Example 19.31.** We can use differentiation to derive properties of  $NV_1(S_4)$  even though  $NV_1(S_4)$  isn't closed under differentiation. Since  $x+y^2$  and  $x+(y+1)^2$  are in  $NV_2(S_2 \times S_4)$  the coefficients of powers of x in  $[x+y^2]^2 [x+(y+1)^2]^2$  interlace. Thus all the following polynomials are in  $NV_1(S_4)$  and interlace in  $S_2$ .

$$y^4(y+1)^4 \leftrightarrow 2y^2(y+1)^2 (2y^2+2y+1) \leftrightarrow 6y^4+12y^3+10y^2+4y+1 \leftrightarrow 2 (2y^2+2y+1) \leftrightarrow 1$$

We can use the example above to give a general construction.

**Lemma 19.32.** *If*  $f, g \in NV_1(S_4)$  *and*  $f \nleftrightarrow g$  *in*  $S_2$  *then* 

$$y^2 f(y) \nleftrightarrow f(y) + y^2 g(y) \nleftrightarrow g(y)$$
 in  $S_2$ 

*Proof.* Since  $x + y^2$  and f(y) + xg(y) are in  $NV_2(S_2 \times S_4)$  their product is also. Thus

$$y^2 f(y) + x [f(y) + y^2 g(y)] + x^2 [g(y)] \in \mathsf{NV}_2(\mathfrak{S}_2 \times \mathfrak{S}_4).$$

Remark 19.33. It follows from the lemma that the two matrices

$$\begin{pmatrix} 1 & y^2 \\ 0 & 1 \end{pmatrix} \qquad \qquad \begin{pmatrix} y^2 & 0 \\ 1 & y^2 \end{pmatrix}$$

map pairs of polynomials in  $NV_1(S_4)$  that interlace in  $S_2$  to another pair of polynomials in  $NV_1(S_4)$  that interlace in  $S_2$ 

#### **19.4** Some determinant properties

In this section we establish some properties of determinants we need to construct polynomials.

If A and B are anti-diagonal matrices with anti-diagonal entries  $(\mathfrak{a}_i)$  and  $(\mathfrak{b}_i)$  then

$$\begin{vmatrix} A & I \\ -I & B \end{vmatrix} = (a_1b_n + 1)\cdots(a_nb_1 + 1)$$
(19.4.1)

The proof is an easy induction. For instance, if n = 4 then

 $\begin{vmatrix} \cdot & \cdot & \cdot & a_1 & 1 & \cdot & \cdot & \cdot \\ \cdot & \cdot & a_2 & \cdot & \cdot & 1 & \cdot & \cdot \\ \cdot & a_3 & \cdot & \cdot & \cdot & \cdot & 1 & \cdot \\ a_4 & \cdot & \cdot & \cdot & \cdot & \cdot & 1 \\ -1 & \cdot & \cdot & \cdot & \cdot & \cdot & b_1 \\ \cdot & -1 & \cdot & \cdot & \cdot & b_2 & \cdot \\ \cdot & \cdot & -1 & \cdot & b_3 & \cdot & \cdot \\ \cdot & \cdot & \cdot & -1 & b_4 & \cdot & \cdot & \cdot \end{vmatrix} = (a_1b_4+1)(a_2b_3+1)(a_3b_2+1)(a_4b_1+1)$ 

where the dots "." are zeros. A similar argument shows that if I is the n by n identity matrix then  $\begin{vmatrix} I & \iota I \\ \iota I & I \end{vmatrix} = 2^n$ .

**Lemma 19.34.** If A is a skew symmetric matrix, B is positive definite, and S is symmetric then

$$|\mathbf{A} + \mathbf{B} + \mathbf{i}\mathbf{S}| \neq 0$$

*Proof.*  $A + B + \iota S \neq 0$  since B is positive definite. If  $|A + B + \iota S| = 0$  then there are non-zero real u, v such that

$$(\mathbf{A} + \mathbf{B} + \mathbf{i}\mathbf{S})(\mathbf{u} + \mathbf{i}\mathbf{v}) = \mathbf{0}$$

Now A, B, S, u, v are real, so we can separate real and imaginary parts:

$$Au + Bu - Sv = 0$$
$$Av + Bv + Su = 0$$

The inner product < r, s >=  $r^ts$  satisfies < Ar, r >= 0 since A is skew-symmetric. Thus

$$<$$
 Bu, u > -  $<$  Sv, u > = 0  
 $<$  Bv, v > +  $<$  Su, v > = 0

S is symmetric, so adding the last two equations yields

$$<$$
 Bu, u  $>$  +  $<$  Bv, v  $>$  = 0

However, B is positive definite and  $\langle Br, r \rangle$  is positive unless r = 0. Thus u = v = 0, a contradiction.

**Corollary 19.35.** *If* S, B *are symmetric and either one is positive definite then*  $|S+\iota B|$  *is not zero.* 

*Proof.* Take S = 0. If necessary, multiply by  $\iota$ .

Lemma 19.36.

$$\begin{vmatrix} A & B \\ -B & A \end{vmatrix} = |A + \iota B| \cdot |A - \iota B|$$

If A is a skew-symmetric matrix then  $\begin{vmatrix} -A & B \\ -B & -A \end{vmatrix} = \begin{vmatrix} A & B \\ -B & A \end{vmatrix}$ 

*Proof.* Assume all matrices are n by n. We compute

$$2^{2n} \begin{vmatrix} A & B \\ -B & A \end{vmatrix} = \begin{vmatrix} (I & -\iota I \\ -\iota I & I \end{pmatrix} \begin{pmatrix} A & B \\ -B & A \end{pmatrix} \begin{pmatrix} I & \iota I \\ \iota I & I \end{pmatrix} \end{vmatrix}$$
$$= \begin{vmatrix} 2(A + \iota B) & 0 \\ 0 & 2(A - \iota B) \end{vmatrix}$$

The second part follows from the first.

**Corollary 19.37.** *If* A *is a skew-symmetric matrix,* S *is symmetric, and* B *is positive definite then* 

$$\begin{vmatrix} A & S + \iota B \\ -S - \iota B & A \end{vmatrix} \neq 0$$

*Proof.* The determinant equals  $|2(A - B + \iota S)| \cdot |2(A + B - \iota S)|$ , and is non-zero by Lemma19.34.

In the next result we determine properties of eigenvalues.

**Lemma 19.38.** Suppose that A, B are positive definite and  $\alpha + \beta \iota$  is an eigenvalue of  $A + B\iota$ .

1.  $0 < \alpha < \rho(A), 0 < \beta < \rho(B), \beta/\alpha \leq \rho(BA^{-1}).$ 

- 2. If A is only symmetric then  $\beta$  is positive.
- 3. If B is only symmetric then  $\alpha$  is positive.

*Proof.* Let u + iv be an eigenvector of  $\alpha + \beta i$ , where u, v are real. Then<sup>1</sup>

 $(A + B\iota)(u + \iota v) = (\alpha + \beta\iota)(u + \iota v)$ 

Separating real and imaginary parts yields

$$Au - Bv = \alpha u - \beta v$$
$$Bu + Av = \alpha v + \beta u$$

 $\langle x, y \rangle = x^{t}y$  is the usual inner product. The symmetry of A and B yields:

$$< u, Au > - < u, Bv > = \alpha < u, u > -\beta < u, v >$$
$$< u, Bv > + < v, Av > = \alpha < v, v > +\beta < u, v >$$

We can now solve for  $\alpha$ , and a similar calculation yields  $\beta$ 

$$\alpha = \frac{\langle u, Au \rangle + \langle v, Av \rangle}{\langle u, u \rangle + \langle v, v \rangle} \leq \rho(A)$$
  
$$\beta = \frac{\langle u, Bu \rangle + \langle v, Bv \rangle}{\langle u, u \rangle + \langle v, v \rangle} \leq \rho(B)$$
  
$$\frac{\beta}{\alpha} = \frac{\langle u, Bu \rangle + \langle v, Bv \rangle}{\langle u, Au \rangle + \langle v, Av \rangle} \leq \rho(BA^{-1})$$

We use these formulas to establish the various parts of the lemma.

#### **19.5** Determinants

We can construct elements of various  $NV_d(\mathcal{C})$  using determinants.

**Corollary 19.39.** If  $\gamma \in \mathbb{C} \setminus 0$ , H is the upper half plane,  $D_i$  are  $n \times n$  positive definite matrices and S is symmetric then

$$\left|\gamma S + \sum_{1}^{d} x_{k} D_{k}\right| \in \mathit{NV}_{d}((\gamma H)^{d})$$

<sup>&</sup>lt;sup>1</sup>Thanks to math.sci.research and Tony O'Conner for this argument.



Figure 19.1: Location of roots of  $|-xI + A + B\iota|$  if B > A.

*Proof.* If  $\gamma \sigma_1, \ldots, \gamma \sigma_d \in \gamma H$  where  $\sigma_k = \alpha_k + \iota \beta_k$  then

$$\begin{split} \Big| \gamma \, S + \sum_{1}^{d} x_k \, D_k \Big| (\gamma \sigma_1, \dots, \gamma \sigma_d) &= \gamma^n \big| S + \sum \sigma_k \, D_k \big| \\ &= \gamma^n \big| \big( S + \sum \alpha_k \, D_k \big) + \iota \big( \sum \beta_k \, D_k \big) \big| \end{split}$$

and this is non-zero by the lemma since  $\beta_k > 0$  and so  $\sum \beta_k D_k$  is positive definite.

Next we construct polynomials in  $NV_d(\mathbb{C}^d)$  where  $\mathbb{C}$  is not a half plane.

**Lemma 19.40.** If  $A_k$ ,  $B_k$  are positive definite and C is the first quadrant then

$$\left|\sum_{1}^{d} x_{k}(A_{k} + \iota B_{k})\right| \in \textit{NV}_{d}(\mathbb{C}^{d})$$

*Proof.* If  $\alpha_k + \iota \beta_k \in \mathbb{C}$  then  $\alpha_k$  and  $\beta_k$  are positive, and

$$\left|\sum_{1}^{d} (\alpha_{k} + \iota \beta_{k})(A_{k} + \iota B_{k})\right| = \left|\sum_{1}^{d} (\alpha_{k}A_{k} - \beta_{k}B_{k}) + \iota \sum_{1}^{d} (\alpha_{k}B_{k} + \beta_{k}A_{k})\right|$$
(19.5.1)

Since  $\sum_{1}^{d} (\alpha_k B_k + \beta_k A_k)$  is positive definite the result follows from Lemma 19.35.

**Lemma 19.41.** If  $A_k$  is positive definite,  $A_k > B_k$  and  $\mathcal{C} = \{a + b\iota | a > b > 0\}$  then

$$\left|\sum_{1}^{u} x_{k}(A_{k} + \iota B_{k})\right| \in NV_{d}(\mathcal{C}^{d})$$

*Proof.* The hypothesis imply that

$$\alpha_k A_k - \beta_k B_k = A_k(\alpha_k - \beta_k) + \beta_k(A_k - B_k)$$

is positive definite, so the result follows from Lemma 19.35 and (19.5.1).  $\Box$ 

**Lemma 19.42.** Suppose A and B are symmetric and  $f(x) = |-xI + A + B\iota|$ . If B > A > 0 then  $f \in NV_1(\mathcal{D})$  where  $\mathcal{D} = \{a + b\iota \mid a > b > 0\}$ .

*Proof.* Since B > A we know that  $\rho(BA^{-1}) > 1$ . From the lemma the eigenvalues  $\alpha + \beta \iota$  satisfy  $\beta > \alpha > 0$ . The roots of f are the negative eigenvalues of  $A + B\iota$ , so they lie in the region of the first quadrant bounded by x = y and x = 0. See Figure 19.1.

#### **19.6** Linear transformations

Properties of the Hadamard product in **P** depended on the identification of the Hadamard product as a coefficient:

If  $f(x) \in \mathbf{P}^{pos}$  and  $g \in \mathbf{P}$  then f(-xy) and  $g(y) \in \overline{\mathbf{P}}_2$ , and the Hadamard product is a coefficient of f(-xy)g(y).

The reason why  $f(-xy) \in \overline{\mathbf{P}}_2$  is that  $\alpha - xy \in \overline{\mathbf{P}}_2$  if and only if  $\alpha > 0$ , and this follows by reversing  $x + \alpha y$ . A similar argument works for half planes.

**Lemma 19.43.** Suppose that C is a half plane with reversal constant  $\rho$ , and  $\ell$  is the ray  $\rho^{-1}\mathbb{R}^+$ . The Hadamard product

$$\sum a_{\mathfrak{i}} x^{\mathfrak{i}} \times \sum g_{\mathfrak{i}}(\mathbf{y}) x^{\mathfrak{i}} \mapsto \sum a_{\mathfrak{i}} g_{\mathfrak{n}-\mathfrak{i}}(\mathbf{y}) x^{\mathfrak{i}}$$

determines a map

$$\boldsymbol{P}^{\ell} \times \boldsymbol{NV}_{d}(\mathbb{C}^{d}) \longrightarrow \boldsymbol{NV}_{d}(\mathbb{C}^{d})$$

*Proof.* If r > 0 then  $x + ry \in NV_2(\mathbb{C}^2)$  so  $\rho xy + r \in NV_2(\mathbb{C}^2)$  and hence  $xy - (-r/\rho) \in NV_2(\mathbb{C}^2)$ . Thus, if  $f(x) \in \mathbf{P}^\ell(n)$  then  $f(xy) \in NV_2(\mathbb{C}^2)$ . Write  $f = \sum a_i x^i$  and  $g = \sum g_i(y) x^i$ . The coefficient of  $z^n$  in

$$f(xz)g(z) = \sum_{i,j} a_i x^i z^i g_j(y) z^j$$

is in  $\mathsf{NV}_d(\mathfrak{C})$  and equals  $\sum \mathfrak{a}_i \mathfrak{g}_{n-i}(y) x^i.$ 

If we consider f(x + y) we get

**Lemma 19.44.** If C is a half plane then  $f(x) \times \sum b_i y^i \mapsto \sum f^{(i)}(x) b_{n-i}/i!$  determines a map

$$NV_2(\mathbb{C}^2) \times NV_1(\mathbb{C}) \longrightarrow NV_2(\mathbb{C}^2)$$

*Proof.* Write  $f(x+y) = \sum f^{(i)}(x) y^{i}/i!$ . The coefficient of  $y^{n}$  in  $f(x+y) \sum b_{i}y^{i}$  is  $f^{(i)}(x)$ 

$$\sum \frac{f^{(\iota)}(x)}{\mathfrak{i}!} \mathfrak{b}_{\mathfrak{n}-\mathfrak{i}}.$$

-		
F		
I		
I		
L		

**Lemma 19.45.** Suppose C is a half plane with reversal constant  $\rho$ .

- 1. If  $f \in NV_d(\mathbb{C}^d)$  then  $f + \rho \partial_{x_1} \partial_{x_2} f \in NV_d(\mathbb{C}^d)$ .
- 2. The linear transformation  $g \times f \mapsto g(\rho \partial_{x_1} \partial_{x_2}) f$  determines a map

$$P^{\text{pos}} \times NV_{d}(\mathbb{C}^{d}) \longrightarrow NV_{d}(\mathbb{C}^{d})$$

*Proof.* The first part follows from Lemma 19.29 and  $f \nleftrightarrow \partial_{x_2} f \nleftrightarrow \partial_{x_1}(\partial_{x_2} f)$ . The second part follows by induction.

#### **19.7** Analytic functions

We define  $\widehat{NV_d}(\mathcal{C})$  to be the uniform closure on compact subsets of  $NV_d(\mathcal{C})$ . Here are some exponential functions in  $NV_d(\mathcal{C})$ .

**Lemma 19.46.** *Suppose*  $\mathcal{C} \subset \mathbb{C}$  *is a cone.* 

- 1. If  $\beta \in \overline{\mathbb{C}} \setminus 0$  then  $e^{x/\beta} \in \widehat{NV_1}(\mathbb{C})$ .
- 2. If  $\rho$  is the reversal constant then  $e^{\rho \times y} \in \widehat{NV_2}(\mathcal{C})$ .

*Proof.* Since  $\beta + x \in \mathsf{NV}_1(\mathfrak{C})$  we have  $\left(1 + \frac{x/\beta}{n}\right)^n$  is in  $\mathsf{NV}_1(\mathfrak{C})$ , and hence  $e^{x/\beta} \in \mathsf{NV}_1(\mathfrak{C})$ . For the second part we know  $x + y \in \mathsf{NV}_2(\mathfrak{C}^2)$ , so  $1 + \rho xy \in \mathsf{NV}_2(\mathfrak{C}^2)$ . Taking limits and following the first part finishes the proof.

**Proposition 19.47.** If C is a half plane with reversal constant  $\rho$  then

1.  $f \mapsto e^{\rho \partial_x \cdot \partial_y} f$  determines a map

$$NV_{d}(\mathbb{C}^{d}) \longrightarrow NV_{d}(\mathbb{C}^{d})$$

2.  $f(\mathbf{x}) \times g(\mathbf{x}) \mapsto f(\rho \partial \mathbf{x})g(\mathbf{x})$  determines a map

$$NV_{d}(\mathbb{C}^{d}) \times NV_{d}(\mathbb{C}^{d}) \longrightarrow NV_{d}(\mathbb{C}^{d}).$$

*Proof.* We know that  $g + \rho \partial_{x_1} \partial_{y_1} g \in \mathsf{NV}_d(\mathbb{C}^d)$ . Thus  $g \mapsto \left(1 + \frac{\rho \partial_{x_1} \partial_{y_1}}{n}\right)^n g$  maps  $\mathsf{NV}_d(\mathbb{C}^d)$  to itself, since the homogeneous part of  $\left(1 + \frac{\rho \partial_{x_1} \partial_{y_1}}{n}\right)^n g$  equals  $g^H$ . Consequently, taking limits shows that the homogeneous part of  $e^{\rho \partial_{x_1} \partial_{y_1}}g$  equals  $g^H$ , and in particular is not zero. The result now follows from Lemma 19.9 and the observation that

$$\rho \, \partial_{\mathbf{x}} \cdot \partial_{\mathbf{y}} = \rho \partial_{\mathbf{x}_1} \partial_{\mathbf{y}_1} + \dots + \rho \partial_{\mathbf{x}_d} \partial_{\mathbf{y}_d}$$

The second part relies on the identity

$$\left. e^{\rho \partial_{\mathbf{x}} \partial_{\mathbf{y}}} \, g(\mathbf{x}) \, f(\mathbf{y}) \right|_{\mathbf{y}=\mathbf{0}} = \left. f(\rho \partial_{\mathbf{x}}) g(\mathbf{x}) \right.$$

By linearity we only need to check it for monomials:

$$e^{\rho \partial_{\mathbf{x}} \partial_{\mathbf{y}}} \mathbf{x}^{\mathbf{I}} \mathbf{y}^{\mathbf{J}} \Big|_{\mathbf{y}=0} = \frac{(\rho \partial_{\mathbf{x}})^{\mathbf{J}}}{\mathbf{J}!} \mathbf{x}^{\mathbf{I}} \cdot (\partial_{\mathbf{y}})^{\mathbf{J}} \mathbf{y}^{\mathbf{J}} \Big|_{\mathbf{y}=0} = (\rho \partial_{\mathbf{x}})^{\mathbf{J}} \mathbf{x}^{\mathbf{I}}$$

**Example 19.48.** The infinite product  $\cos(x) = \prod_{k=0}^{\infty} \left(1 - \frac{4x^2}{(2k+1)^2\pi^2}\right)$  shows that  $\cos(\gamma x) \in \mathsf{NV}_1(\gamma \mathsf{H})$ . Taking derivatives shows

 $\cos(\gamma x) \nleftrightarrow -\gamma \sin(\gamma x)$  and  $\sin(\gamma x) \bigstar \gamma \cos(\gamma x)$ 

This gives us elements in  $NV_2(\gamma H)$ :

 $\cos(\gamma x) - \gamma y \sin(\gamma x)$  and  $\sin(\gamma x) + \gamma y \cos(\gamma x)$ 

#### 19.8 Homogeneous polynomials

Define  $\mathsf{homog}_d$  to be the set of homogeneous polynomials contained in  $\mathsf{HP}_d$ . If we don't want to specify d we write  $\mathsf{homog}_*$ . By Lemma 19.24 we can write  $f(\mathbf{x}) \in \mathsf{homog}_d$  as  $\alpha \cdot g(\mathbf{x})$  where  $g(\mathbf{x})$  has all positive coefficients. If d = 2 then  $f(x, y) \in \mathsf{homog}_d$  is of the form  $\alpha \cdot G(x, y)$  where G is the homogenization of a polynomial in  $\mathbf{P}^{\mathsf{pos}}$ . In general,  $f \in \mathsf{homog}_d$  is a multiple of the homogenization of a polynomial in  $\mathbf{P}^{\mathsf{pos}}_{d-1}$ , since  $\overline{\mathbf{P}}_{d-1} = \mathsf{rHP}_{d-1}$  (Theorem 20.3).

The first lemma shows that there is nothing special about the upper half plane in the definition of  $homog_d$ ; any half plane through the origin will do.

Lemma 19.49.  $homog_d = \bigcap_{\gamma \in \mathbb{C} \setminus 0} NV_d((\gamma H)^d)$ 

*Proof.* If  $f \in \mathsf{homog}_d$  has degree n and  $\sigma \in \mathsf{H}^d$  then  $f(\gamma \sigma) = \gamma^n f(\sigma) \neq 0$  so  $f \in \mathsf{NV}_d((\gamma \mathsf{H})^d)$ .

Conversely, if  $f(\mathbf{x})$  has degree n and is in the intersection,  $\sigma \in H^d$  and  $\gamma \neq 0$  then  $f(\gamma \sigma) \neq 0$ . But  $f(\gamma \sigma)$  is a polynomial in  $\gamma$ , so the solutions to  $f(\gamma \sigma) = 0$  must all be  $\gamma = 0$ . Thus  $f(\gamma \sigma)$  is divisible by  $\gamma^n$ , and  $f(\gamma \sigma) = \gamma^n f(\sigma)$ . Since this holds for all  $\sigma \in H^d$  we see that  $f(\gamma \mathbf{x}) = \gamma^n f(\mathbf{x})$ , so f is homogeneous.  $\Box$ 

There is a simple determinant construction for polynomials in  $homog_d$ , but not all such polynomials can be so realized.

**Corollary 19.50.** *If all* 
$$D_i$$
 *are positive definite then*  $\left|\sum_{1}^{d} x_k D_k\right| \in homog_d$ . *Proof.* This follows from Corollary 19.39.

Next, we construct homogeneous polynomials from elementary symmetric functions. Fix a positive integer n, and let  $\sigma_i(y_1, \ldots, y_n)$  be the i'th elementary symmetric function of  $y_1, \ldots, y_n$ . If **I** is an index set then

$$\sigma_{I}(\mathbf{x}) = \prod_{i \in I} \sigma_{i}(x_{i,1}, \dots, x_{i,n})$$

579

**Lemma 19.51.** For positive integers n and m we have that  $\sum_{|I|=m} \sigma_I(x) \in homog_*$ .

*Proof.* Since  $uz - 1 \in HP_2$ 

$$\prod_{i=1}^{n} (x_{i,1}z - 1) = \sum_{k} \sigma_{k}(x_{i,1}, \dots, x_{i,n})(-z)^{n-k} \in \mathsf{HP}_{*}$$

and thus

$$\prod_{j=1}^d \left(\sum_k \sigma_k(x_{j,1},\ldots,x_{j,n})(-z)^{n-k}\right) \in \mathsf{HP}_*$$

The coefficient of  $z^m$  is  $\pm \sum_{|I|=m} \sigma_I(\mathbf{x})$ . This is in homog<sub>\*</sub> since all monomials have degree m.

Now we have the Schur-Szegö theorem for homogeneous polynomials.

**Lemma 19.52.** If  $f = \sum a_I \mathbf{x}^I$ ,  $g = \sum b_I \mathbf{x}^I$ , deg(f) = deg(g), and  $f, g \in homog_d$  then

$$\sum a_I b_I I! \mathbf{x}^I \in homog_d$$

*Proof.* If f has degree n then  $f(\partial x)g(x + y)$  is in homog<sub>\*</sub> and equals

$$\sum_{|\mathbf{I}|=|\mathbf{J}|=n} a_{\mathbf{I}} \, \partial \mathbf{x}^{\mathbf{I}} \, b_{\mathbf{J}} \, (\mathbf{x}+\mathbf{y})^{\mathbf{J}} = \sum a_{\mathbf{I}} \, b_{\mathbf{I}} \, \mathbf{I}! \, \mathbf{x}^{\mathbf{I}}$$

since  $(\partial \mathbf{x}^{I})(\mathbf{x} + \mathbf{y})^{J} = \begin{cases} 0 & I \neq J \\ I! \, \mathbf{y}^{I} & I = J \end{cases}$ 

**Lemma 19.53.** If  $\sum \alpha_I \mathbf{x}^I \in HP_d$  then  $\sum \alpha_I \sigma_I(\mathbf{x}) I! \in HP_d$ .

Proof. Note that

$$(\partial \mathbf{x})^{i} [(\mathbf{x} + \mathbf{x}_{1}) \cdots (\mathbf{x} + \mathbf{x}_{n})] \Big|_{\mathbf{x} = 0} = \sigma_{i}(\mathbf{x}) i!$$

and thus

$$\left(\sum a_{I} \partial x^{I}\right) \left(\prod_{i=1}^{d} \prod_{k=1}^{n} (x_{i} + x_{i,k})\right) \bigg|_{x=0} = \sum a_{I} \sigma_{I}(x) I!.$$

**Corollary 19.54.** *The linear transformation*  $\mathbf{x}^{I} \mapsto \sigma_{I}(\mathbf{x}) I!$  *maps*  $HP_{*} \longrightarrow HP_{*}$  *and*  $homog_{*} \longrightarrow homog_{*}$ .

#### **19.9** Möbius transformations

Suppose  $Mz = \frac{az+b}{cz+d}$  is a Möbius transformation where  $|M| = ad - bc \neq 0$ . If  $f(x_1, ..., x_r)$  is homogeneous of degree n then we say that f is *M*-invariant if  $(Mf)(\mathbf{x}) = |M|^n f(\mathbf{x})$  for all Möbius transformations M where  $|M| \neq 0$  and

$$(\mathsf{Mf})(\mathsf{x}_1,\ldots,\mathsf{x}_r) = \left[(\mathsf{c}\mathsf{x}_1+\mathsf{d})\cdots(\mathsf{c}\mathsf{x}_r+\mathsf{d})\right]^n \cdot f\left(\frac{\mathsf{a}\mathsf{x}_1+\mathsf{b}}{\mathsf{c}\mathsf{x}_1+\mathsf{d}},\ldots,\frac{\mathsf{a}\mathsf{x}_r+\mathsf{b}}{\mathsf{c}\mathsf{x}_r+\mathsf{d}}\right)$$

We will see (Lemma 19.55) that there are no M-invariant polynomials in  $homog_d$ , but we can find polynomials  $f(\mathbf{x}, \mathbf{y})$  such that

1. f is M-invariant.

2.  $f(\mathbf{x}, -\mathbf{y}) \in homog_d$ 

We will call such polynomials *Ruelle polynomials* [150]. It is easy to show that a polynomial is M-invariant; the difficulty is showing that it is  $homog_d$ .

**Lemma 19.55.** If  $f \in HP_n$  then there is a Möbius transformation M so that Mf  $\notin HP_n$ .

*Proof.* Choose  $\alpha_i$  so that  $f(\alpha_1, ..., \alpha_n) = 0$ , and choose  $\beta$  so that  $\beta + \alpha_i$  is in the upper half plane for all i. If  $Mz = z - \beta$  then the roots of

$$Mf(x_1, \ldots, x_n) = f(x_1 - \beta, \ldots, x_n - \beta)$$

are  $\beta + \alpha_1, \dots, \beta + \alpha_n$  which are all in the upper half plane, so Mf  $\notin HP_n$ .  $\Box$ 

We now give several constructions of M-invariant polynomials. They all depend on the fact that x - y is M-invariant:

$$M(x-y) = (cx+d)(cy+d)\left[\frac{ax+b}{cx+d} - \frac{ay+b}{cy+d}\right] = |M|(x-y)$$

**Construction 19.56.** If  $(a_{ij}) = A$  is an r by r matrix and  $X = diag(x_1, ..., x_r)$ ,  $Y = diag(y_1, ..., y_r)$  then we define

$$f(\mathbf{x}, \mathbf{y}) = \det(AX - YA) = \det(a_{ij}(x_i - y_j)).$$

 $f(\mathbf{x}, \mathbf{y})$  is M-invariant since

$$Mf = det(a_{ij}M(x_i - y_j)) = det(|M|a_{ij}(x_i - y_j)) = |M|^r f(X, Y).$$

Construction 19.57. With the same setup as above we let

$$g(\mathbf{x}, \mathbf{y}) = per(AX - YA)$$

where per is the permanent. The same argument shows  $g(\mathbf{x}, \mathbf{y})$  is M-invariant.

If we expand the determinant or the permanent we find that these two constructions are special cases of

**Construction 19.58.** If  $\alpha_{\sigma} \in \mathbb{C}$  for each permutation  $\sigma$  then define

$$h(\mathbf{x},\mathbf{y}) = \sum_{\sigma \in \mathfrak{sym}(r)} \alpha_{\sigma} \prod_{i=1}^{r} (x_i - y_{\sigma i}).$$

This is clearly M-invariant. Ruelle [150] proved that all Ruelle polynomials can be written in this form, but the  $\alpha_{\sigma}$  are not unique.

**Example 19.59.** If J is the r by r all 1 matrix then Cauchy's determinant formula [108] gives

$$\begin{aligned} f(\mathbf{x}, \mathbf{y}) &= |JX - YJ| = |(x_i - y_j)| = \prod_{i < j} (x_i - x_j) \cdot \prod_{i < j} (y_i - y_j) \\ &= \sum_{\sigma \in sym(r)} sign(\sigma) \prod_{i=1}^{r} (x_i - y_{\sigma i}) \end{aligned}$$

Note that f(X, -Y) is not in  $HP_{2r}$  since it contains  $x_i - x_j$  factors. However, we will see later that

$$per(XJ - JY) = \sum_{\sigma \in sym(r)} \prod_{1}^{r} (x_i - y_{\sigma i})$$

is a Ruelle polynomial. We call this the Grace polynomial, denoted  $Grace_r$ . For instance,

$$\begin{aligned} & \text{Grace}_2 = (x_1 - y_1)(x_2 - y_2) + (x_1 - y_2)(x_2 - y_1) \\ &= 2x_1x_2 - (x_1 + x_2)(y_1 + y_2) + 2y_1y_2 \\ &= 2\sigma_2(x_1, x_2)\sigma_0(y_1, y_2) - \sigma_1(x_1, x_2)\sigma_1(y_1, y_2) + 2\sigma_0(x_1, x_2)\sigma_2(y_1, y_2) \end{aligned}$$

The next lemma is Ruelle's original definition of a Ruelle polynomial.

**Lemma 19.60** (Ruelle).  $f(\mathbf{x}, \mathbf{y})$  *is a Ruelle polynomial iff whenever there is a circle separating*  $\sigma = (\sigma_1, \ldots, \sigma_n)$  *from*  $\tau = (\tau_1, \ldots, \tau_m)$  *then*  $f(\sigma, \tau) \neq 0$ .

*Proof.* Choose a Möbius transformation such that  $M\sigma$  is in the upper half plane and  $M\tau$  is in the lower half plane. Since f(Mx, -My) is in  $homog_d$  we have that  $f(M\sigma, -M\tau) \neq 0$ . The converse is harder, and can be found in [150].

**Lemma 19.61** (Ruelle). *If* A *is unitary then*  $f(\mathbf{x}, \mathbf{y}) = |AX - YA|$  *is a Ruelle polynomial.* 

*Proof.* We have seen that f is M-invariant. If A is unitary then  $A^{-1} = A^*$  so

$$f(x, y) = |AX - YA| = |A| |X - A^*YA|$$

If the rows of A are  $A_1, \ldots, A_r$  then

$$f(\mathbf{x}, \mathbf{y}) = |A| \det\left(X + \sum y_i A_i^* A_i\right)$$

Since all  $A_i^*A_i$  are positive semi-definite the determinant is in HP<sub>2r</sub>.

**Example 19.62.** In this example we give the Ruelle polynomial that is constructed using the general three by three orthogonal matrix [11] where  $\beta = a^2 + b^2 + c^2 + d^2$ .

$$A = \frac{1}{\beta} \begin{pmatrix} a^2 + b^2 - c^2 - d^2 & 2(bc + ad) & 2(bd - ac) \\ 2(bc - ad) & a^2 - b^2 + c^2 - d^2 & 2(ab + cd) \\ 2(ac + bd) & 2(cd - ab) & a^2 - b^2 - c^2 + d^2 \end{pmatrix}$$

If we expand the determinant we get the following permutation representation:

$$\begin{array}{rl} (y_3+x_1)(y_2+x_2)(y_1+x_3)+\\ 4(ac+bd)^2&(y_2+x_1)(y_3+x_2)(y_1+x_3)+\\ 4(ac-bd)^2&(y_3+x_1)(y_1+x_2)(y_2+x_3)+\\ 4(b^2-c^2)(a^2-d^2)&(y_1+x_1)(y_3+x_2)(y_2+x_3)+\\ 4(a^2-b^2)(d^2-c^2)&(y_2+x_1)(y_1+x_2)(y_3+x_3)+\\ a^2-b^2+c^2-d^2)^2&(y_1+x_1)(y_2+x_2)(y_3+x_3)\end{array}$$

In order to show that the Grace polynomial is a Ruelle polynomial we need to recall Grace's theorem:

**Theorem 19.63** (Grace). If  $f = \sum_{0}^{n} a_{i}x^{i}$ ,  $g = \sum_{0}^{n} b_{i}x^{i}$ , and the roots of f are separated by a circle from the roots of g then

$$\sum_{0}^{n} a_{i} b_{n-i} (-1)^{i} i! (n-i)! \neq 0$$

Corollary 19.64. The Grace polynomial is a Ruelle polynomial.

*Proof.* We know the Grace polynomial is M-invariant; we now show it satisfies the second condition. Suppose that  $x_1, \ldots, x_n$  and  $y_1, \ldots, y_n$  are in the upper half plane. Grace's theorem applied to  $\prod (x - x_i)$  and  $\prod (x + y_i)$  shows that

$$\begin{split} 0 &\neq \sum (-1)^i \sigma_i(x_1, \dots, x_n) \cdot \sigma_{n-i}(y_1, \dots, y_n) (-1)^i \, i! \, (n-i)! \\ &= \sum \sigma_i(x_1, \dots, x_n) \sigma_{n-i}(y_1, \dots, y_n) \, i! \, (n-i)! \end{split}$$

Comparing monomials shows that this is exactly  $\text{Grace}_r(\mathbf{x}, -\mathbf{y})$ , and it is non-zero for substitutions in the upper half plane.

Here's a simple one-variable corollary.

**Corollary 19.65.** *If*  $a_i, b_i \in \mathbb{R}$  *then* 

$$\sum_{\sigma \in \text{sym}(n)} \prod_i (x + a_i + b_{\sigma i}) \in \textbf{P}$$

*Proof.* Replace  $y_i$  by  $b_i$  and  $x_i$  by  $x + a_i$  in the Grace polynomial.

The conjecture below generalizes Grace's theorem:

**Conjecture 19.66.** If n is a positive integer,  $\sigma_i(x_k)$  the i'th elementary symmetric polynomial on  $x_{k,1}, \ldots, x_{k,n}$  and  $\sigma_I(\mathbf{x}) = \prod_k \sigma_{i_k}(x_k)$  then

$$\sum_{|I|=n}\sigma_{I}(x)\,I!\in\textit{homog}_{*}$$

#### 19.10 Properties of polynomials with zeros in a strip

Assume that S is the strip  $\{z \mid -1 < \text{Re}(z) < 1\}$ . We will consider properties of  $NV_1(\mathbb{C} \setminus S)$ . If  $f \in NV_1(\mathbb{C} \setminus S)$  then all roots of f lie in S. Since S is convex we have that

 $\mathsf{NV}_1(\mathbb{C} \setminus \mathbb{S})$  is closed under differentiation.

The reverse of  $x - \alpha$  has root  $1/\alpha$ , so

 $\mathsf{NV}_1(\mathbb{C} \setminus \mathbb{S})$  is not closed under reversal.

A non-trivial property is this interlacing result:

**Lemma 19.67.** If S is a vertical strip and  $f \in NV_1(\mathbb{C} \setminus S)$  then for all real  $\alpha$ 

 $f + \iota \alpha f' \in \mathit{NV}_1(\mathbb{C} \setminus \mathbb{S})$ 

The proof follows from the next lemma.

**Lemma 19.68.** Suppose that  $f = \prod (x - r_k)$ , and define

$$g = f + \iota \alpha \sum_{k} a_{k} \frac{f(x)}{x - r_{k}}$$

where  $\alpha \in \mathbb{R}$  and all  $a_k$  are non-negative. If a + bi is a root of g then

$$\min_{k} \Re(\mathbf{r}_{k}) \leqslant \mathfrak{a} \leqslant \max_{k} \Re(\mathbf{r}_{k})$$

*Proof.* Let  $r_k = s_k + \iota t_k$ . Dividing by f yields

$$1 + \iota \alpha \sum_{k} a_{k} \frac{1}{a + b\iota - s_{k} - \iota t_{k}} = 0$$
$$\sum_{k} a_{k} \frac{a - \iota b - s_{k} + \iota t_{k}}{|a + b\iota - s_{k} - \iota t_{k}|^{2}} = \iota/\alpha$$

Taking the real part yields

$$\sum_k \mathfrak{a}_k \frac{\mathfrak{a} - s_k}{|\mathfrak{a} + \mathfrak{b}\mathfrak{l} - s_k - \mathfrak{l}\mathfrak{t}_k|^2} = 0$$

Thus a can't be less than all  $s_k$ , nor greater than all  $s_k$ .

**Corollary 19.69.** If  $f \in P$  then the map  $g \mapsto f(\iota D)g$  determines a linear transformation

$$NV_1(\mathbb{C}\setminus\mathbb{S})\longrightarrow NV_1(\mathbb{C}\setminus\mathbb{S})$$

**Corollary 19.70.** *If*  $g \in NV_1(\mathbb{C} \setminus S)$  *then* 

$$g+g''+g^{(4)}/2!+g^{(6)}/3!+\cdots\in\textit{NV}_1(\mathbb{C}\setminus\texttt{S})$$

*Proof.* Apply the previous corollary to  $f = e^{-x^2}$ .

The polynomial xy - 1 is in  $NV_2(\mathbb{C} \setminus S)$  since |x| > 1 if  $x \notin S$ . If  $x, y \notin S$  then |xy| > 1, and hence  $xy - 1 \neq 0$ . This observation will be generalized in the next section

#### 19.11 Polynomials non-vanishing on the open unit ball

Suppose that  $\Delta_d$  is the open unit ball in  $\mathbb{C}^d$ . That is,

$$\Delta_{\mathbf{d}} = \{ \mathbf{z} \in \mathbb{C}^{\mathbf{d}} \mid |\mathbf{z}| < 1 \}$$

We will construct some simple non-trivial elements of  $\mathsf{NV}_d(\mathfrak{D})$  where  $\mathfrak{D} \subset \mathbb{C}$ , and use them to show that  $\mathsf{NV}_d(\Delta^d)$  has non-trivial elements. Recall that  $\mathsf{NV}_d(\mathfrak{D})$  contains trivial products of the form

$$\prod_{i=1}^d \prod_j (x_i - \alpha_{ij})$$

where  $\alpha_{ij} \notin D$ . The next lemma constructs elements that are not products.

**Lemma 19.71.** Suppose that the ball  $B = \{z \mid |z - \sigma| < r\}$  is contained in  $\mathbb{C}^d \setminus \mathcal{D}$ . If  $\sigma = (\sigma_1, \dots, \sigma_d)$  and  $0 < b < r^d$  then

$$(\mathbf{x} - \sigma_1) \cdots (\mathbf{x} - \sigma_d) - \mathbf{b} \in \mathit{NV}_d(\mathcal{D})$$

*Proof.* If  $(\tau_1, \ldots, \tau_d) \notin B$  then  $|\tau_i - \sigma_i| \ge r$ , so  $|(\tau_1 - \sigma_1) \cdots (\tau_d - \sigma_d)| \ge r^d$ . Since  $0 < b < r^d$  it follows that

$$(\tau_1 - \sigma_1) \cdots (\tau_d - \sigma_d) - b \in \mathsf{NV}_d(\mathcal{D})$$

**Example 19.72.** If H is the upper half plane, then the ball  $\{z \mid |z + \iota| < 1\}$  is contained in  $\mathbb{C}^d \setminus H^d$ . It follows that

$$(\mathbf{x}_1 + \mathbf{i}) \cdots (\mathbf{x}_d + \mathbf{i}) - 1 \in \mathsf{HP}_d$$

A similar argument shows that

$$(\mathbf{x}_1+1)\cdots(\mathbf{x}_d+1)-1\in\mathcal{H}_d$$

Here's a small extension to  $NV_d(\Delta_d)$ .

**Lemma 19.73.** If f(x) has all its zeros in the open unit disk then

$$f(x_1 \cdots x_d) \in \mathcal{NV}_d(\Delta_d)$$

*Proof.* Use the lemma with  $B = \Delta_d$ , and multiply many of the polynomials together.

It is easy to see that  $xy - 1/2 \in NV_2(\mathbb{C}^2 \setminus \Delta \times \Delta)$  since if neither x nor y is in  $\Delta \times \Delta$  then both have absolute value at least one, so their product has absolute value at least one. There is a simple geometric condition for a multiaffine polynomial to belong to  $NV_2(\mathbb{C}^2 \setminus \Delta^2)$ .

**Lemma 19.74.** Suppose that f(x, y) = a + bx + cy + dxy, and let  $M = \begin{pmatrix} a & c \\ b & d \end{pmatrix}$ .

- 1.  $f \in NV_2(\mathbb{C}^2 \setminus \Delta^2) \iff M \colon \mathbb{C}^2 \setminus \Delta^2 \longrightarrow \Delta^2$
- $2. \ f \in \textit{NV}_2(\mathbb{C}^2 \setminus \overline{\Delta}^2) \Longleftrightarrow M \colon \mathbb{C}^2 \setminus \overline{\Delta}^2 \longrightarrow \overline{\Delta}^2$

*Proof.* Assume that  $|y| \ge 1$ . If f(x, y) = 0 we must show that |x| < 1. Solving for x

$$\mathbf{x} = -\frac{\mathbf{a} + \mathbf{c}\mathbf{y}}{\mathbf{b} + \mathbf{d}\mathbf{y}} = -\mathbf{M}(\mathbf{y})$$

Thus |x| < 1 if and only if |M(y)| < 1. The second case is similar.

Recall the result (p. 616) that if  $0 < a_1 < \cdots < a_n$  then  $\sum a_i x^i$  has all its roots in the open unit ball. Here's a conjecture in two variables

**Conjecture 19.75.** *If*  $0 < a_1 < \cdots < a_n$  *then* 

$$\sum (a_i + a_j) x^i y^j \in \textit{NV}_2(\mathbb{C}^2 \setminus \overline{\Delta}^2)$$

We can establish the multiaffine case of this conjecture.



Figure 19.2: The image of the circle under a Möbius transformation

**Lemma 19.76.** If 
$$0 < a < b$$
 then  $2a + (a + b)(x + y) + 2bxy \in NV_2(\mathbb{C}^2 \setminus \overline{\Delta}^2)$ .

*Proof.* This is a question about Möbius transformations. We take three points on the unit circle, and find the unique circle containing their image. We check that  $M(\infty)$  lies in this circle, and that this circle lies in the unit circle.

$$M(1) = \frac{3a+b}{3b+a} \qquad M(-1) = 1$$
$$M(1) = \frac{(2+i)a+ib}{a+(1+2i)b} \qquad M(\infty) = \frac{a+b}{2b}$$
$$center = \frac{2(a+b)}{a+3b} \qquad radius = \frac{b-a}{a+3b}$$

Since the center is on the real line, it follows that the image is the circle with diameter M(1) M(-1). Thus the image lies in the closed unit circle.

The following lemma constructs a stable polynomial from a polynomial in  $NV_1(\mathbb{C} \setminus \Delta)$ .

**Lemma 19.77.** If  $0 < a_0 < \cdots < a_n$  then  $\sum_0^n a_i(x+1)^i(1-x)^{n-i} \in \mathcal{H}_1$ .

*Proof.* The Möbius transformation  $z \mapsto \frac{1+z}{1-z}$  maps the right half plane to  $\mathbb{C} \setminus \Delta$ . It follows that if  $f(x) \in \mathsf{NV}_1(\mathbb{C} \setminus \Delta)$  then  $f((1+x)(1-x)) \in \mathcal{H}_1$ . If we let  $f(x) = \sum a_i x^i$  then  $f \in \mathsf{NV}_1(\mathbb{C} \setminus \Delta)$ . Multiplying by  $(1-x)^n$  yields the result.  $\Box$ 

#### 19.12 Polynomials non-vanishing in a sector

We begin with a simple construction of polynomials with no roots in a sector. We claim that if f and g have positive leading coefficient then

$$f \in \mathbf{P}^{\text{pos}}(n) \& g \in \mathbf{P}^{\text{pos}}(m) \implies f(x) + g(y) \in \mathsf{rNV}_2(\mathbb{S}_{2n} \times \mathbb{S}_{2m})$$

To see this, we first claim that the image of  $S_{2n}$  under f is the open right half plane. If  $f = \prod (x + r_i)$  where  $r_i \ge 0$ , and  $\sigma \in S_n$ , then  $r_i + \sigma \in S_{2n}$ , so  $f(\sigma)$  is the product of n points whose arguments are all less than  $\pi/(2n)$ , Similarly, if  $\tau \in S_{2m}$  then  $g(\tau)$  is also in the right half plane, so their sum is as well. In particular, their sum is not zero.

For example,

$$\mathbf{x}^{n} + \mathbf{y}^{m} \in \mathsf{rNV}_{2}(\mathbb{S}_{2n} \times \mathbb{S}_{2m})$$

This construction easily generalizes.

**Lemma 19.78.** Suppose that  $n_1, \ldots, n_d$  are positive integers. If all  $f_i(x) \in \overline{P^{pos}}(n_i)$  have positive leading coefficient then

$$f_1(x_1) + \dots + f_d(x_d) \in \textit{rNV}_d(\$_{2n_1} \times \dots \times \$_{2n_d})$$

CHAPTER



### Upper half plane polynomials

There is a natural subset of polynomials with complex coefficients that has properties similar to **P**: those whose roots lie in the lower half plane. However, we call them *upper half plane polynomials* because the generalization to more variables uses the fact that they are non-zero in the upper half plane. For a quick summary with proofs see [59].

We first recount properties that follow from the general theory of Chapter 19, and then investigate polynomials in one variable.

#### 20.1 Properties of upper half plane polynomials

We apply the general results of Chapter 19 to derive results about polynomials non-vanishing in the upper half plane. If UHP is the upper half plane then  $U_d(\mathbb{C}) = \mathsf{NV}_d(\mathsf{UHP}^d)$ . The reversal constant is -1, and  $\mathsf{UHP}/\mathsf{UHP} = \mathbb{C} \setminus (-\infty, 0)$ . If f,  $g \in U_d(\mathbb{C})$  then we write  $f \xleftarrow{u} g$  if  $f + zg \in U_{d+1}(\mathbb{C})$ . From Chapter 19 we have

#### Lemma 20.1.

- 1.  $U_d(\mathbb{C})$  is closed under differentiation.
- 2. If  $f \in U_d(\mathbb{C})$  then  $f \xleftarrow{u} \partial f / \partial x_i$
- 3. If  $\sum x^i f_i(\mathbf{y}) \in U_{d+1}(\mathbb{C})$  then  $f_i(\mathbf{y}) \in U_d(\mathbb{C})$  and  $f_i \xleftarrow{U} f_{i+1}$ .
- 4.  $fg \in U_d(\mathbb{C})$  if and only if  $f \in U_d(\mathbb{C})$  and  $g \in U_d(\mathbb{C})$ .
- 5. If  $f \xleftarrow{u} g$  then  $g \xleftarrow{u} -f$ .
- 6. If  $f \xleftarrow{u} g \xleftarrow{u} h$  then  $f h \xleftarrow{u} g$ .
- 7. If S is symmetric and all  $D_i$  are positive definite then  $|S + \sum_1^d x_i D_i| \in U_d(\mathbb{C})$ .

- 8. The Hadamard product  $\sum a_i x^i \times \sum f_i(\mathbf{y}) x^i \mapsto \sum a_i f_{n-i}(\mathbf{y}) x^i$  determines a map  $\mathbf{P}^{pos} \times U_d(\mathbb{C}) \longrightarrow U_d(\mathbb{C})$ .
- 9.  $f(\mathbf{x}) \times g(\mathbf{x}) \mapsto f(-\partial \mathbf{x})g(\mathbf{x})$  determines a map

$$U_{d}(\mathbb{C}) \times U_{d}(\mathbb{C}) \longrightarrow U_{d}(\mathbb{C}).$$

10. The following are equivalent

a) 
$$f(\mathbf{x}) \in U_d(\mathbb{C})$$

b)  $f(\mathbf{a} + t\mathbf{b}) \in U_1(\mathbb{C})$  for  $\mathbf{a} \in \mathbb{R}^d$ ,  $\mathbf{b} > 0$ .

*Proof.* The only modification is that we do not need homogeneity in the last one. Following the proof of Lemma 19.18 we see (1) implies (2). Conversely, consider  $\sigma_1, \ldots, \sigma_d$  in UHP. Since 1 is in the closure of UHP we can find an  $\alpha$  and positive **a**, **b** such  $(\sigma_1, \ldots, \sigma_d) = \mathbf{a} + \alpha \mathbf{b}$ . Thus,  $f(\sigma_1, \ldots, \sigma_d) = f(\mathbf{a} + \mathbf{b}\alpha) \neq 0$ .

There are two reversals in  $U_d(\mathbb{C})$ , a single variable reversal with a minus sign, and a full reversal with no signs.

#### Lemma 20.2.

- 1. If  $\sum x^i f_i(\mathbf{y}) \in U_d(\mathbb{C})$  has degree n then  $\sum (-x)^{n-i} f_i(\mathbf{y}) \in U_d(\mathbb{C})$ .
- 2. If  $x_i$  has degree  $e_i$  then  $x_1^{e_1} \cdots x_d^{e_d} f(1/x_1, \dots, 1/x_d) \in U_d(\mathbb{C})$ .

*Proof.* Only the second one requires proof. If  $\sigma_i \in UHP$  then  $1/\overline{\sigma_i} \in UHP$ . Thus

$$f(1/\sigma_1,\ldots,1/\sigma_d) = f(1/\overline{\sigma_1},\ldots,1/\overline{\sigma_d}) \neq 0.$$

1: need real and im parts in Pd

#### Theorem 20.3. $U_d = \overline{\mathbf{P}}_d$ .

*Proof.* We first show  $U_d \subset \overline{P}_d$ . If  $f(\mathbf{x}) \in U_d$  then define  $g_{\varepsilon}(\mathbf{x}) = f(\mathbf{x} + \varepsilon(x_1 + \cdots + x_d))$  where  $\varepsilon > 0$ . We may assume that  $f^H$  has all non-negative terms, so  $g_{\varepsilon}^H$  has all positive terms. Define  $\mathbf{a} = (0, r_2, \ldots, r_d)$  and  $\mathbf{b} = (t, 0, \ldots, 0)$ . Since  $U_d$  is closed  $g_{\varepsilon}(\mathbf{a} + \mathbf{b}t) \in U_1$ . Since  $U_1 = \mathbf{P}$ , we see that  $g_{\varepsilon}$  satisfies substitution, and therefore  $g_{\varepsilon} \in U_d$ . The conclusion follows since  $\lim_{\varepsilon \to 0^+} g_{\varepsilon} = f$ .

Next we show  $U_d \subset U_d$  which implies  $\overline{\mathbf{P}}_d = U_d$ . If  $f \in U_d$  and is homogeneous then  $f(\mathbf{a} + \mathbf{bt}) \in \mathbf{P}^{\text{pos}}$  for all  $\mathbf{a}, \mathbf{b} > 0$ , so  $f \in U_d$  by Lemma 20.1. If f is not homogeneous then choose  $\mathbf{c} \in \mathbb{R}^d$  so that  $f(\mathbf{x} + \mathbf{c})$  has all positive coefficients. Since  $f(\mathbf{x} + \mathbf{c}) \in \mathbf{P}_d$  we can homogenize it  $y^n f(\mathbf{x}/y) = F(\mathbf{x}, y)$ . Since  $F(\mathbf{x}, y) \in U_{d+1}$  and is homogeneous we know from the previous paragraph that  $F(\mathbf{x}, y) \in U_{d+1}$ , and thus  $f(\mathbf{x} + \mathbf{c}) = F(\mathbf{x}, 1) \in U_d$ . Now  $\mathbf{c}$  is real so it follows that  $f(\mathbf{x}) \in U_d$ .

**Lemma 20.4.** If  $f(\mathbf{x}, \mathbf{y}^2) \in U_d$  then  $f(\mathbf{x}, \mathbf{y}) \in \mathbf{P}_d$ .

*Proof.* If  $\sigma, \sigma_i \in \mathsf{UHP}$  then we can write  $\sigma = \tau^2$  where  $\tau \in \mathsf{UHP}$ . Thus

$$f(\sigma_1,\ldots,\sigma_d,\sigma) = f(\sigma_1,\ldots,\sigma_d,\tau^2) \neq 0$$

which proves the lemma.

Of course the converse is false – consider x + 1 and  $x^2 + 1$ .

#### 20.2 Constructions from determinants

We can construct polynomials in HP<sub>d</sub> using skew-symmetric matrices.

Lemma 20.5. If A is skew symmetric and B is positive definite then

$$\begin{vmatrix} \mathsf{x} \, \mathsf{A} & \mathsf{B} \\ -\mathsf{B} & \mathsf{y} \, \mathsf{A} \end{vmatrix} \in \overline{\mathbf{P}}_2$$

*Proof.* If  $b = B^{-1/2} = b^t$  then

$$\begin{pmatrix} b & 0 \\ 0 & b \end{pmatrix} \begin{pmatrix} xA & B \\ -B & yA \end{pmatrix} \begin{pmatrix} b & 0 \\ 0 & b \end{pmatrix} = \begin{pmatrix} xbAb & I \\ -I & ybAb \end{pmatrix} = \begin{pmatrix} xC & I \\ -I & yC \end{pmatrix}$$

where bAb = C is skew symmetric. We can find an orthogonal matrix 0 and D an anti-diagonal matrix with positive anti-diagonal  $(d_i)$  such that  $OCO^t = \begin{pmatrix} 0 & D \\ -D & 0 \end{pmatrix}$ . Clearly

$$\begin{pmatrix} 0 & . \\ . & 0 \end{pmatrix} \begin{pmatrix} x C & I \\ -I & x C \end{pmatrix} \begin{pmatrix} 0^{t} & . \\ . & 0^{t} \end{pmatrix} = \begin{pmatrix} . & xD & I & . \\ -xD & . & . & I \\ -I & . & . & yD \\ . & -I & -yD & . \end{pmatrix}$$

where the dot is a zero matrix. By (19.4.1) this determinant is  $\prod_{1}^{n} (d_{i}d_{n-i}xy - 1)$  which is in  $\overline{P}_{2}$ .

**Example 20.6.** If A is the skew-symmetric n by n matrix that is all 1 above the diagonal, and all -1 below, then

$$\begin{vmatrix} \mathbf{x} & \mathbf{A} & \mathbf{I} \\ -\mathbf{I} & \mathbf{A} \end{vmatrix} = \left[ \sum (-\mathbf{x})^k \binom{\mathbf{n}}{2k} \right]^2$$

We already knew that this is in **P** since it's just the even part of  $(x - 1)^n$ .

The following corollary was proved by a different method in [62].

**Corollary 20.7.** Suppose A is skew-symmetric and B is positive definite. The determinant and pfaffian of  $\begin{pmatrix} xA & B \\ -B & A \end{pmatrix}$  have alternating coefficients.

*Proof.* Substituting y = 1 in Lemma 20.5 shows that all the roots are positive, and hence the coefficients alternate. Since the pfaffian is the square root of the determinant all the roots are negative, and so the coefficients again alternate.

We can construct polynomials in d variables of even total degree.

**Corollary 20.8.** If A is skew-symmetric, and  $B_1, \ldots, B_d$  are positive definite then

$$f(\mathbf{x}) = \left| \begin{pmatrix} A & 0 \\ 0 & A \end{pmatrix} + x_1 \begin{pmatrix} 0 & B_1 \\ -B_1 & 0 \end{pmatrix} + \dots + x_d \begin{pmatrix} 0 & B_d \\ -B_d & 0 \end{pmatrix} \right| \in \textit{HP}_d$$

 $f(\mathbf{x})$  is a square, and all non-zero monomials of  $f(\mathbf{x})$  have even total degree.

*Proof.* If we substitute  $\alpha_k + \beta_k \iota$  for  $x_k$  where all  $\beta_k$  are positive, then the determinant equals

$$\left| \begin{pmatrix} A & \sum \alpha_k B_k \\ -\sum \alpha_k B_k & A \end{pmatrix} + \iota \begin{pmatrix} 0 & \sum \beta_k B_k \\ -\sum \beta_k B_k & 0 \end{pmatrix} \right|$$

and this matrix is non-zero by Lemma 19.34 since  $\sum \beta_k B_k$  is positive definite. Since the matrix is anti-symmetric  $f(\mathbf{x})$  is a square. Finally, using Lemma 19.36,

$$f(-\mathbf{x}) = \begin{pmatrix} A & -\sum x_k B_k \\ \sum x_k B_k & A \end{pmatrix} = \begin{pmatrix} -A & -\sum x_k B_k \\ \sum x_k B_k & -A \end{pmatrix} = f(\mathbf{x})$$

**Example 20.9.** If d = 1,  $B_1$  is the identity, and A is the matrix of the last example then

$$\begin{split} \left| \begin{pmatrix} A & 0 \\ 0 & A \end{pmatrix} + x \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix} \right| &= \left[ \sum_{i=0}^{n} (-1)^{i} x^{n-2i} \binom{n}{2i} \right]^{2} \\ |A + \iota x I| &= \sum_{i=0}^{n} (-1)^{i} x^{n-2i} \binom{n}{2i} \end{split}$$

In the last corollary the matrix had real coefficients and the polynomial was a square. If we allow complex coefficients then we can take the square root:

**Corollary 20.10.** If A is skew-symmetric and all  $B_k$  are positive definite then

$$g(\mathbf{x}) = \left| A + \iota \sum_{1}^{d} x_{\iota} B_{\iota} \right| \in \textit{rHP}_{d}$$

and all monomials have even degree.

Proof. The determinant in Corollary 20.8 equals

$$|\mathbf{A} + \mathbf{i} \sum_{1}^{d} \mathbf{x}_{i} \mathbf{B}_{i}| |\mathbf{A} - \mathbf{i} \sum_{1}^{d} \mathbf{x}_{i} \mathbf{B}_{i}|$$

If C denotes the sum then

$$|\mathbf{A} + \mathbf{i}\mathbf{C}| = |(\mathbf{A} + \mathbf{i}\mathbf{C})^{\mathsf{t}}| = |-\mathbf{A} + \mathbf{i}\mathbf{C}| = |\mathbf{A} - \mathbf{i}\mathbf{C}|$$

Thus both factors are equal and since the conjugate of  $|A + \iota C|$  is  $|A - \iota C|$  they both have real coefficients. This also shows that  $g(-\mathbf{x}) = g(\mathbf{x})$ , and there fore all terms have even degree.

**Lemma 20.11.** Suppose that  $D_1$ ,  $D_2$ ,  $D_3$  are positive definite matrices, and S is symmetric. If

$$f(x, y) = |x D_1 + y D_2 + S + \iota D_3|$$

then  $f(x, y) \in U_2$ .

*Proof.* The homogeneous part of f is  $|xD_1 + yD_2|$  which has all positive coefficients by Lemma 9.5. For any  $\alpha \in \mathbb{R}$ 

$$f(x, \alpha) = |x D_1 + (\alpha D_2 + S) + \iota D_3|$$

and by this is in  $\Im$  since  $\alpha D_2 + S$  is symmetric.

**20.3**  $SL_2(\mathbb{R}[x])$ 

In this section we characterize the set of matrices of  $\text{SL}_2(\mathbb{R}[x])$  whose entries interlace. We define

#### Definition 20.12.

$$SL_2(\mathbb{R}[x])_P = \left\{ \begin{pmatrix} g & f \\ k & h \end{pmatrix} \mid gh - fk = 1 \text{ and } f \xleftarrow{u} g \xleftarrow{u} k, f \xleftarrow{u} h \xleftarrow{u} k \right\}$$

We begin with some elementary properties.

**Lemma 20.13.** Suppose  $\begin{pmatrix} g & f \\ k & h \end{pmatrix} \in SL_2(\mathbb{R}[x])_P$  is not constant.

- 1. deg(g) + deg(h) = deg(f) + deg(k).
- 2. If sign(r) is the sign of the leading coefficient of r then sign(g) · sign(h) = sign(f) · sign(k).
- 3.  $\left(\begin{smallmatrix} g & f \\ k & h \end{smallmatrix}\right)^{-1} \notin SL_2(\mathbb{R}[\mathbf{x}])_{\mathsf{P}}.$

*Proof.* The first two follow from the fact that gh - fk is a constant, so the leading coefficients of gh and fk must cancel. The determinant of the inverse is 1, but all the interlacings go in the opposite direction.

*Remark* 20.14. Consider some necessary conditions for the matrix  $M = \begin{pmatrix} g & f \\ k & h \end{pmatrix}$  to belong to  $SL_2(\mathbb{R}[x])_P$ .

- 1. M has determinant one.
- 2. Degrees of adjacent entries differ by at most one.
- 3. All entries are in **P**.
- 4. Either  $f \xleftarrow{u} g$  or  $f \xleftarrow{u} h$  or  $g \xleftarrow{u} k$  or  $h \xleftarrow{u} k$ .

The matrix  $\begin{pmatrix} 1 & x \\ x & x^{2}-1 \end{pmatrix}$  satisfies all conditions except (1). If (2) fails then not all adjacent polynomials can interlace. It is possible for only condition (3) to fail – consider  $\begin{pmatrix} 1 & x \\ x & x^{2}+1 \end{pmatrix}$ . In addition, only condition (4) can fail

$$\left(\begin{array}{ccc} x^2+3x & x^2+3x+2 \\ x^2+3x-\frac{1}{2} & x^2+3x+\frac{3}{2} \end{array}\right).$$

All the entries are in **P**, but no two entries interlace.

**Lemma 20.15.** *If*  $M = \begin{pmatrix} g & f \\ k & h \end{pmatrix}$  *has determinant* 1 *and three out four interlacings then*  $M \in SL_2(\mathbb{R}[x])_P$ .

*Proof.* Using the first part of the next lemma we can multiply by certain of the first three matrices of the lemma to assure that all leading coefficients are positive. The interlacings were preserved, so we know that the degrees must be  $\binom{n+b}{n+a+b}$  where a, b  $\ge 0$ .

Suppose that we do not know that  $g \longrightarrow f$ . Since  $h \longrightarrow f$  we know that h sign interlaces f. Now fk - gh = 1, so g has the same sign at the roots of f as h does, so g sign interlaces f. Since  $deg(g) \leq deg(f)$  it follows that  $g \longrightarrow f$ . The remaining cases are similar.

We now show that certain matrices preserve  $SL_2(\mathbb{R}[x])_P$ .

**Lemma 20.16.** Suppose a > 0. Multiplying on the right or left by any of these matrices maps  $SL_2(\mathbb{R}[x])_P$  to itself:

 $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \quad \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \quad \begin{pmatrix} 1 & ax+b \\ 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 \\ -ax-b & 1 \end{pmatrix}$ 

*Proof.* The first three follow easily from the fact that

 $\mathsf{f} \xleftarrow{u} \mathsf{g} \quad \Leftrightarrow \quad -\mathsf{f} \xleftarrow{u} -\mathsf{g} \quad \Leftrightarrow \quad -\mathsf{g} \xleftarrow{u} \mathsf{f} \quad \Leftrightarrow \quad \mathsf{g} \xleftarrow{u} -\mathsf{f}.$ 

If  $\begin{pmatrix} g & f \\ k & h \end{pmatrix} \in SL_2(\mathbb{R}[x])_P$  then

$$\begin{pmatrix} g & f \\ k & h \end{pmatrix} \begin{pmatrix} 1 & ax+b \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} g & (ax+b)g+f \\ k & (ax+b)k+h \end{pmatrix}$$

Now if we have  $r \xleftarrow{U}{\leftarrow} s$  then  $r + ys \in U_2$ . Since  $a \ge 0$  we may substitute

$$r + (y + ax + b)s = r + (ax + b)s + ys \implies r + (ax + b)s \xleftarrow{u} s.$$

This shows that we have three out of the four interlacings, and since the determinant is one, the conclusion follows from Lemma 20.16. Multiplying on the other side is similar.

The last one follows from

$$\begin{pmatrix} 1 & 0 \\ -ax-b & 1 \end{pmatrix} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & ax+b \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

**Proposition 20.17.**  $SL_2(\mathbb{R}[x])_P$  is generated by

$$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \qquad \begin{pmatrix} c & 0 \\ 0 & 1/c \end{pmatrix} \qquad \begin{pmatrix} 1 & ax+b \\ 0 & 1 \end{pmatrix}$$

where  $a \ge 0, b \in \mathbb{R}, c \neq 0$ .

*Proof.* The degree of  $M = \begin{pmatrix} g & f \\ k & h \end{pmatrix}$  is deg(g) + deg(h). If M has degree at least two then we show that there are matrices  $A, B, M_1$  such that  $M = AM_1B$  where A, B are certain products of the first two generators, and  $M_1 \in SL_2(\mathbb{R}[x])_P$  has lower degree.

We can multiply on the left or right by the first two generators so that the leading coefficients in the top row are positive. If the matrix is  $\begin{pmatrix} g & f \\ k & h \end{pmatrix}$  then  $f \xleftarrow{U} g$ , and since they have positive leading coefficients  $f \longleftarrow g$ . Thus we can write f = (ax + b)g - s where  $a \ge 0$ , g < s, and s has positive leading coefficient. We write

$$\begin{pmatrix} g & f \\ k & h \end{pmatrix} = \begin{pmatrix} g & (ax+b)g-s \\ k & h \end{pmatrix} = \begin{pmatrix} g & -s \\ k & -h_1 \end{pmatrix} \begin{pmatrix} 1 & ax+b \\ 0 & 1 \end{pmatrix}$$

where  $h_1$  has positive leading coefficient, and  $h = (ax + b)k - h_1$ . We show that  $\begin{pmatrix} g & -s \\ k & -h_1 \end{pmatrix} \in SL_2(\mathbb{R}[x])_P$ , and by Lemma 20.15 we need to show that  $k \leq h_1$ .

Now  $deg(g) + deg(h_1) = deg(k) + deg(s)$ , and deg(g) = deg(s) + 1 so we have  $deg(h_1) = deg(k) - 1$ . From  $-gh_1 + sk = 1$  we see that  $h_1$  and g have the same signs on the roots of k. Since g alternates sign on the roots of k so does  $h_1$ . From  $deg(h_1) < deg(k)$  it follows that  $k < h_1$ .

If we reduce a matrix to one of degree one, then after multiplying by appropriate generators it has the form  $\begin{pmatrix} c & ax+b \\ 0 & 1/c \end{pmatrix}$ , which is easily seen to be a product of the generators.

**Corollary 20.18.**  $SL_2(\mathbb{R}[x])_P$  is closed under multiplication.

**Corollary 20.19.** If  $\begin{pmatrix} g & f \\ k & h \end{pmatrix} \in SL_2(\mathbb{R}[x])_P$  then  $f + y g + z h + yz k \in \overline{\mathbf{P}}_3$ .

*Proof.* It is clear that each of the first two generators of  $SL_2(\mathbb{R}[x])_P$  determines a polynomial in  $\overline{P}_3$ . The last generator determines the polynomial ax+b+y+z which in is  $\overline{P}_3$  since a is non-negative.

To complete the proof we verify that multiplication by each of the generators preserves the property of being in  $\overline{\mathbf{P}}_3$ . We start with a polynomial  $F = f + yg + zh + yzk \in \overline{\mathbf{P}}_3$ . There are three cases:

 $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$  This matrix transforms  $\begin{pmatrix} g & f \\ k & h \end{pmatrix}$  to  $\begin{pmatrix} -k & -h \\ g & f \end{pmatrix}$  which corresponds to

$$-k - yh + zf + yzg = -(k + yh + f(-z) + y(-z)g)$$

which equals  $-zF(x, y, -1/z) \in \overline{\mathbf{P}}_3$ .

 $\begin{pmatrix} c & 0 \\ 0 & 1/c \end{pmatrix}$  This transforms F to  $c(f + yg + (z/c^2)h + y(z/c^2)k) = F(x, y, z/c^2)$ which is in  $\overline{\mathbf{P}}_3$ .

 $\begin{pmatrix} 1 & ax+b \\ 0 & 1 \end{pmatrix}$  This transforms F to

$$(ax + b)g + f + yg + z[(ax + b)k + h] + yzk = F(x, ax + b + y, z)$$

which is in  $\overline{\mathbf{P}}_3$ .

**Corollary 20.20.**  $SL_2(\mathbb{R}[x])_P$  acts on polynomials  $f(x) + g(x)y + h(x)z + k(x)yz \in \overline{P}_3$  by matrix multiplication.

#### 20.4 Matrices of nearly quadratic polynomials

In this section we consider matrices formed from polynomials in  $U_3$  that have degree at most 2 in y and z. We show how to construct such matrices of arbitrarily large degree and constant determinant.

**Definition 20.21.**  $M_3 = \left\{ (f_{ij})_{0 \leqslant i, j \leqslant 2} \mid \sum_{i,j=0}^2 f_{ij}(x) \, y^i z^j \in U_3 \right\}$ 

Many operations on  $F(x, y, z) = \sum f_{ij}y^i z^j$  corespond to matrix multiplication. We only consider  $U_3$ , but the arguments apply to all  $U_k$ . Let

$$\mathsf{M} = \begin{pmatrix} \mathsf{f}_{00} & \mathsf{f}_{10} & \mathsf{f}_{20} \\ \mathsf{f}_{01} & \mathsf{f}_{11} & \mathsf{f}_{21} \\ \mathsf{f}_{02} & \mathsf{f}_{12} & \mathsf{f}_{22} \end{pmatrix}$$

**Example 20.22.** Suppose we replace y by cy, where c > 0. Since  $F(x, cy, z) \in U_3$  the new matrix is in  $M_3$  and equals

$$\begin{pmatrix} f_{00} & c f_{10} & c^2 f_{20} \\ f_{01} & c f_{11} & c^2 f_{21} \\ f_{02} & c f_{12} & c^2 f_{22} \end{pmatrix} = M \begin{pmatrix} 1 & . & . \\ . & c & . \\ . & . & c^2 \end{pmatrix}$$

If we replace *z* by *cz* then we multiply on the left

$$\begin{pmatrix} f_{00} & f_{10} & f_{20} \\ c f_{01} & c f_{11} & c f_{21} \\ c^2 f_{02} & c^2 f_{12} & c^2 f_{22} \end{pmatrix} = \begin{pmatrix} 1 & \cdot & \cdot \\ \cdot & c & \cdot \\ \cdot & \cdot & c^2 \end{pmatrix} M$$

**Example 20.23.** Next we consider F(x, y + a, z) which is in  $U_3$  for all a. The corresponding matrix equals

$$\begin{pmatrix} f_{00} + af_{10} + a^2f_{20} & f_{10} + 2af_{20} & f_{20} \\ f_{01} + af_{11} + a^2f_{21} & f_{11} + 2af_{21} & f_{21} \\ f_{02} + af_{12} + a^2f_{22} & f_{12} + 2af_{22} & f_{22} \end{pmatrix} = M \begin{pmatrix} 1 & . & . \\ a & 1 & . \\ a^2 & 2a & 1 \end{pmatrix}$$

If we consider F(x, y, z + a) then we multiply on the left by:

$$\begin{pmatrix} 1 & a & a^2 \\ . & 1 & 2a \\ . & . & 1 \end{pmatrix} M$$

**Example 20.24.** If a is positive then  $F(x, y + ax, z) \in U_3$  and  $F(x, y, z + ax) \in U_3$ . These correspond to multiplying on the right by

$$\begin{pmatrix} 1 & \cdot & \cdot \\ ax & 1 & \cdot \\ a^2x^2 & 2ax & 1 \end{pmatrix}$$

or on the left by

$$\begin{pmatrix} 1 & ax & a^2x^2 \\ . & 1 & 2ax \\ . & . & 1 \end{pmatrix}$$

Example 20.25. If we reverse with respect to y then we get

$$\begin{pmatrix} f_{20} & -f_{10} & f_{00} \\ f_{21} & -f_{11} & f_{01} \\ f_{22} & -f_{12} & f_{01} \end{pmatrix} = M \begin{pmatrix} \cdot & \cdot & 1 \\ \cdot & -1 & \cdot \\ 1 & \cdot & \cdot \end{pmatrix}$$

If we reverse with respect to *z* then we multiply on the left.

**Example 20.26.** In order to get started we need some simple matrices. Since  $(y + z)^2 \in U_3$  the following matrix is in  $M_3$ 

$$\begin{pmatrix} \cdot & \cdot & 1 \\ \cdot & 2 & \cdot \\ 1 & \cdot & \cdot \end{pmatrix}$$

and has determinant -2.

**Example 20.27.** Finally, we show how to combine these matrices to make matrices in  $M_3$  with determinant -2. Let

$$C = \begin{pmatrix} 1 & . & . \\ x & 1 & . \\ x^2 & 2x & 1 \end{pmatrix} \qquad A = \begin{pmatrix} . & . & 1 \\ . & -1 & . \\ 1 & . & . \end{pmatrix} \qquad B = \begin{pmatrix} . & . & 1 \\ . & 2 & . \\ 1 & . & . \end{pmatrix}$$

The matrix  $BC(AC)^n$  has degree 2n + 2. For instance,  $BC(AC)^1$  equals

$$\left(\begin{array}{rrrr} x^4-2x^2+1 & 2x^3-2x & x^2\\ 2x^3-2x & 4x^2-2 & 2x\\ x^2 & 2x & 1 \end{array}\right)$$

#### **20.5** $\mathfrak{I}$ , a subset of $U_1(\mathbb{C})$

We now consider polynomials in  $U_1(\mathbb{C})$  with positive leading coefficients and strict interlacing.

#### Definition 20.28.

$$\begin{split} \mathfrak{I} &= \{ \mathsf{f} + \mathsf{g} \iota \mid \mathsf{f}, \mathsf{g} \in \mathbf{P} \text{ have positive leading coefficients and } \mathsf{f} \lessdot \mathsf{g} \} \\ \mathfrak{I}^{\mathrm{pos}} &= \{ \mathsf{f} + \mathsf{g} \iota \mid \mathsf{f}, \mathsf{g} \in \mathbf{P}^{\mathrm{pos}} \text{ have positive leading coefficients and } \mathsf{f} \lessdot \mathsf{g} \} \\ \overline{\mathfrak{I}} &= \{ \mathsf{f} + \mathsf{g} \iota \mid \mathsf{f}, \mathsf{g} \in \mathbf{P} \text{ have positive leading coefficients and } \mathsf{f} \lessdot \mathsf{g} \} \\ \widehat{\mathfrak{I}} &= \{ \mathsf{f} + \mathsf{g} \iota \mid \mathsf{f}, \mathsf{g} \in \mathbf{P} \text{ have positive leading coefficients and } \mathsf{f} \lessdot \mathsf{g} \} \\ \widehat{\mathfrak{I}} &= \text{ the uniform closure of } \mathfrak{I} \end{split}$$

If  $h(x) = \sum a_i x^i$  then we define  $h_{\Re} = \sum \Re(a_i) x^i$  and  $h_{\Im} = \sum \Im(a_i) x^i$ . With this notation, the interlacing condition of the definition of  $\Im$  is that  $h_{\Re} < h_{\Im}$  where  $h = f + g\iota$ .

*Remark* 20.29. There are two reasons we only consider <, and not also  $\ll$ . The first is that if we consider a product  $\prod(x + \sigma_i) = f(x) + \iota g(x)$  where  $\sigma_i \in \mathbb{C}$  then the degree of f(x) is greater than the degree of g(x).

Second, if the degrees of the real and imaginary polynomials are equal then the leading coefficient isn't positive nor even real; it's complex. However, this is not a serious problem, for we can multiply it to have the correct form:

*Lemma* 20.30. If  $f \ll g$ , a > 0 is the leading coefficient of f and b > 0 is the leading coefficient of g then

$$(\mathfrak{a} - \mathfrak{bl})(\mathfrak{f} + \mathfrak{lg}) \in \mathfrak{I}$$

*Proof.* The leading coefficient of (a - bi)(f + ig) is  $a^2 + b^2$ . Now

$$\begin{aligned} (a - b\iota)(f + \iota g) &= af + bg + \iota(ag - bf) \\ \begin{pmatrix} a & b \\ -b & a \end{pmatrix} \begin{pmatrix} f \\ g \end{pmatrix} &= \begin{pmatrix} af + bg \\ ag - bf \end{pmatrix} \end{aligned}$$

The determinant is  $a^2 + b^2$ , and a is positive, so the interlacing on the right hand side is strict (Corollary 3.8).

It remains to determine the sign of the leading coefficient of ag - bf. Write

$$\begin{split} f &= c_0 + \dots + c_{n-1} x^{n-1} + a \, x^n \\ g &= d_0 + \dots + d_{n-1} x^{n-1} + b \, x^n \end{split}$$

Since  $f \ll g$  we know that  $\left| \begin{array}{c} d_{n-1} & b \\ c_{n-1} & a \end{array} \right| > 0$  and this is the leading coefficient of ag - bf.

*Remark* 20.31. If  $F = f + \iota g$  then F has no real roots. If  $r \in \mathbb{R}$  were a root of F then  $f(r) + \iota g(r) = 0$ , which implies that f(r) = g(r) = 0. This contradicts the hypothesis that f and g have no roots in common.

If  $f \in \mathbf{P}^{pos}$  then there are further restrictions on the location of the roots.

**Lemma 20.32.** If  $f, g \in P^{pos}$ ,  $f \leq g$ , and f, g have positive leading coefficients then the roots of  $f + \iota g$  have negative real part and negative imaginary part.

*Proof.* We know the imaginary part is negative. If  $f = \prod_k (x - r_k)$  and  $f(\alpha) + \iota g(\alpha) = 0$  then there are positive  $a_k$  such that

$$\begin{split} \sum_{j} a_{j} \left( f/(x-r_{j}) \right) &= g(x) \\ \prod_{k} (\alpha-r_{k}) + \iota \sum_{j} a_{j} \frac{1}{\alpha-r_{j}} \prod_{k} (\alpha-r_{k}) = 0 \\ 1 + \iota \sum_{j} a_{j}/(\alpha-r_{j}) &= 0 \end{split}$$

Taking the imaginary part yields

$$\Re(\alpha)\sum_{j}\frac{a_{j}}{|\alpha-r_{j}|^{2}}=\sum_{j}a_{j}\frac{r_{j}}{|\alpha-r_{j}|^{2}}$$

Since the  $a_j$ 's are positive and the  $r_j$ 's are negative, it follows that the real part is negative.

It's easy to construct polynomials in  $\Im$  from polynomials in **P**:

**Lemma 20.33.** Suppose that  $f \in P$ ,  $\sigma$  is in the upper half plane and  $\alpha$  is positive. Then

- 1.  $f(x + \sigma) \in \mathfrak{I}$ . 2.  $\int_{0}^{1} f(x + \alpha \iota t) dt \in \mathfrak{I}$ . 3.  $\int_{0}^{1} f(x + \alpha \iota t) dt \in \mathfrak{I}$ . 4.  $\iota^{n} f(-\iota x) \in \mathfrak{I}$  if  $f \in P^{pos}(n)$ . 5. If  $g(x) \in \mathfrak{I}$  then  $g(\alpha x) \in \mathfrak{I}$  for positive  $\alpha$ .
- 6. If  $f \in P$  has all distinct roots then  $f(x) + \iota f'(x) \in \mathfrak{I}$ .

*Proof.* The roots of  $f(x + \sigma)$  all have imaginary part  $-\Im(\sigma)$  which is negative. Corollary 22.30 shows that all the roots of the integral have imaginary part equal to  $-\alpha/2$ , and so the integral is in  $\Im$ .

If  $f = x^n + ax^{n-1} + \cdots$  then  $\iota^n f(-\iota x) = x^n + \iota ax^{n-1} + \cdots$  Since  $f(x) \in \mathbf{P}^{pos}$ , all the roots of  $f(-\iota x)$  lie in the lower half plane. Thus the roots are in the correct location, and the leading coefficients are positive.

The remaining ones are obvious.
A useful property of  $\mathfrak{I}$  is the following, which follows from the fact that  $\mathfrak{I} \subset U_1(\mathbb{C})$ , and the leading coefficient of the product is positive.

## **Lemma 20.34.** $\Im$ and $\widehat{\Im}$ are closed under multiplication.

*Remark* 20.35. We can also prove that  $\Im$  is closed under multiplication using properties of matrices. It suffices to show that if  $f + \mathfrak{gl}$  and  $h + k\mathfrak{l}$  are in  $\Im$ , then so is their product  $(\mathfrak{fh} - \mathfrak{gk}) + (\mathfrak{fk} + \mathfrak{gh})\mathfrak{l}$ . Notice that both  $\mathfrak{fh} - \mathfrak{gk}$  and  $\mathfrak{fk} + \mathfrak{gh}$  have positive leading coefficients, and that the degree of  $\mathfrak{fh} - \mathfrak{gk}$  is one more than the degree of  $\mathfrak{fk} + \mathfrak{gh}$ . If we write these terms as a matrix product

$$\begin{pmatrix} fh - gk \\ fk + gh \end{pmatrix} = \begin{pmatrix} h & -k \\ k & h \end{pmatrix} \begin{pmatrix} f \\ g \end{pmatrix}$$

then the conclusion follows from Lemma 3.57.

Here is a simple consequence.

**Corollary 20.36.** *If* f < g *and* n *is a positive integer then* 

$$\begin{split} & If \, n = 2m \\ & \sum_{k=0}^m f^{2k} g^{2m-2k} \binom{2m}{2k} (-1)^k < \quad \sum_{k=0}^m f^{2k+1} g^{2m-2k-1} \binom{2m}{2k+1} (-1)^k \\ & If \, n = 2m+1 \\ & \sum_{k=0}^m f^{2k} g^{2m+1-2k} \binom{2m+1}{2k} (-1)^k > \quad \sum_{k=0}^m f^{2k+1} g^{2m-2k} \binom{2m+1}{2k+1} (-1)^k \end{split}$$

*Proof.* Expand  $(f + \iota g)^n$ .

The following corollary is useful.

**Corollary 20.37.** If  $T: P \longrightarrow P$  preserves the degree and sign of the leading coefficient then  $T: \overline{\mathfrak{I}} \longrightarrow \overline{\mathfrak{I}}$ . If T also preserves strict interlacing then  $T: \mathfrak{I} \longrightarrow \mathfrak{I}$ .

*Proof.* Since T is a transformation that only involves real coefficients, we know that

$$\mathfrak{R}(\mathsf{T}(\mathsf{f})) = \mathsf{T}(\mathfrak{R}(\mathsf{f})) \text{ and } \mathfrak{I}(\mathsf{T}(\mathsf{f})) = \mathsf{T}(\mathfrak{I}(\mathsf{f})).$$

**Corollary 20.38.** These linear transformations map  $\Im$  to itself.

- 1.  $f \mapsto f + \alpha f'$  where  $\alpha \in \mathbb{R}$ .
- 2.  $f \mapsto g(D)f$  where  $g(x) \in P$
- 3.  $f \mapsto EXP(f)$ .

4.  $f \mapsto H_n$ .

*Proof.* The only observation required is that the second one follows from the first.  $\Box$ 

*Remark* 20.39. Note that the first transformation is equivalent to the statement that if  $g \leq h$  then  $g + \alpha g' \leq h + \alpha h'$ . This is Lemma 1.55.

### **Corollary 20.40.** *If* $f \in \mathfrak{I}$ *then*

- 1.  $f(x + i) + f(x i) \in \overline{\mathfrak{I}}$
- 2.  $(1/\iota) (f(x + \iota) f(x \iota)) \in \overline{\mathfrak{I}}$
- *Proof.* The first one is  $2\cos(D)f$ , and the second is  $2\sin(D)f$ .

**Lemma 20.41.** Suppose that  $S_1, S_2 : P \longrightarrow P$  are linear transformations that preserve  $\leq$  and the sign of the leading coefficient, and satisfy  $S_1(f) \leq S_2(f)$  for all  $f \in P$ . Define  $T(f) = S_1(f) + \iota S_2(g)$ . Then,

- $T: P \longrightarrow \overline{\mathfrak{I}}$
- $T: \overline{\mathfrak{I}} \longrightarrow \overline{\mathfrak{I}}$

*Proof.* The first is immediate from the hypotheses of the Lemma. Since  $T(f + \iota g) = S_1(f) - S_2(g) + \iota(S_1(g) + S_2(f))$  the degrees and the signs of the leading coefficients are correct. We have

$$S_{1}(f) - S_{2}(g) \leq S_{1}(g) \text{ since } S_{1}(f) \leq S_{1}(g) \leq S_{2}(g)$$
$$\leq S_{2}(f) \text{ since } S_{1}(f) \leq S_{2}(f) \leq S_{2}(g)$$

Adding these two interlacings gives the desired interlacing.

Here are some examples of linear transformations satisfying the hypothesis of the lemma. Suppose that  $U : P \longrightarrow P$  preserves  $\leq$  and the sign of the leading coefficient.

$$\begin{array}{ll} S_1(f) = U(xf) & S_2(f) = U(f) \\ S_1(f) = U(f) & S_2(f) = U(f)' \\ S_1(f) = U(f) & S_2(f) = U(f') \end{array}$$

*Remark* 20.42. Here is a different proof of part of the Hermite-Biehler theorem. We prove that if  $f \leq g$  and the leading coefficients of f and g are positive, then  $h(x) = f(x) + \iota g(x)$  has all roots in -H. First of all, notice that h(z) can not have any real roots, since if  $h(\alpha) = 0$  and  $\alpha$  is real then  $f(\alpha) = g(\alpha) = 0$  which contradicts  $f \leq g$ . Next, it follows from from Example 3.59 that  $(x + \iota)^n = f_{\mathfrak{R}} + g_{\mathfrak{I}}\iota$  is in  $\mathfrak{I}$  since  $f_{\mathfrak{R}} \leq g_{\mathfrak{I}}$ , and  $(x + \iota)^n \in U_1(\mathbb{C})$ . Since any pair of strictly interlacing polynomials can be reached by a path of strictly interlacing polynomials with positive leading coefficients must have all roots in the lower half plane, since the roots can never be real.

П



Figure 20.1: Trajectories for  $f + t \iota g$ 

The real parts of the roots of a polynomial in  $\Im$  are constrained by the location of the roots of the real part of the polynomial. See Question 95.

**Lemma 20.43.** Suppose that  $f = f_{\Re}(x) + \iota f_{\Im}(x) \in \Im$ , and assume that all the roots of  $f_{\Re}$  lie in an interval I. Then, the real parts of the roots of f also lie in I.

*Proof.* We may assume  $f_{\Re}$  is monic, and write  $f_{\Re}(x) = \prod(x - r_k)$ ,  $f_{\Im}(x) = \sum a_k f_{\Re}/(x - r_k)$  where the  $a_k$  are non-negative. We show that if  $\Re(\sigma)$  is greater than all  $r_k$  then  $f(\sigma) \neq 0$ . The case where  $\Re \sigma$  is less than all the roots is similar. Dividing by  $f_{\Re}$ 

$$0 = 1 + \iota \sum_{k} a_{k} \frac{1}{\sigma - r_{k}}$$

Since  $\Re \sigma > r_k$ , all terms in the sum have positive real part, and so  $\sigma$  is not a zero of f.

Suppose that  $f \leq g$  and consider the plot (Figure 20.1) of the roots of  $F_t = (1-t)f + \iota t g$  for  $0 \leq t \leq 1$ . From the lemma we know that the real part of the roots of  $F_t$  lies in the interval determined by the roots of f. At t = 0 the roots are the roots of f, and as t increases the roots move to the roots of g, except for one root whose imaginary part goes to  $-\infty$ .

Here's a variation of the lemma where we start with polynomials of equal degree.

**Lemma 20.44.** Suppose that  $g \blacktriangleleft f$  where f and g have the same degree. If  $f + \iota g = \prod (x - r_k - \iota s_k)$ , f has roots  $a_1 \leqslant \cdots \leqslant a_n$ , g has roots  $b_1 \leqslant \cdots \leqslant b_n$  then

$$\mathbf{r}_{k} \in \left[\frac{a_{1}+b_{1}}{2}, \frac{a_{n}+b_{n}}{2}\right]$$

*Proof.* Suppose that  $\alpha + \iota\beta$  is a root of  $f + \iota g$ . Then

$$\prod (\alpha + \iota\beta - a_k) + \iota \prod (\alpha + \iota\beta - b_k) = 0$$
$$\prod \frac{\alpha + \iota\beta - b_k}{\alpha + \iota\beta - a_k} = \iota$$
$$\prod \frac{(\alpha - b_k)^2 + \beta^2}{(\alpha - a_k)^2 + \beta^2} = 1$$

Since  $g \blacktriangleleft f$  we have that  $b_k \le a_k$ . Thus, if  $b_k < a_k$  and  $\alpha < (a_k + b_k)/2$  then the kth term is less than one, and if  $\alpha > (a_k + b_k)/2$  then the kth term is greater than 1. This implies the result.

### 20.6 Some simple consequences

**Corollary 20.45.** If  $f(x, y) \in U_2$  and  $\sigma$  has positive imaginary part then  $f(x, \sigma) \in \mathfrak{I}$ .

*Proof.* Since  $U_2 \subset U_2(\mathbb{C})$  substitution of  $\sigma$  yields a polynomial in  $U_1(\mathbb{C})$ , and this is in  $\mathfrak{I}$  since the coefficient of  $x^n$  is positive.

**Corollary 20.46.** If  $f(x, y) \in U_2$  and

$$f(x, y) = f_0(x) + f_1(x)y + f_2(x)y^2 + \cdots$$

then

$$f_0 - f_2 + f_4 - \cdots \leq f_1 - f_3 + f_5 - \cdots$$

*Proof.* This is just a restatement of Corollary 20.45, where  $f_{\Re}$  is the left hand side, and  $f_{\Im}$  is the right hand side.

**Lemma 20.47.** If  $f = \sum_{ij} a_{ij} x^i y^j \in \mathbf{P}_2(2n)$  then  $\sum_{i \equiv j \pmod{2}} (-1)^{(n+i+j)/2} a_{ij} x^i y^j$  is in  $\mathbf{P}_2$ 

*Proof.* Since  $g(x, y) = \iota^n f(-\iota x, -\iota y) \in U_2(\mathbb{C})$ , the expression of the conclusion is the real part of g.

**Corollary 20.48.** If f < g in  $U_2$  then the matrix  $\begin{pmatrix} f & -g \\ g & f \end{pmatrix}$  maps a pair of interlacing polynomials in  $U_2$  to a pair of interlacing polynomials in  $P_2$ .

*Proof.* The action of the matrix on  $\binom{h}{k}$  is the same as the multiplication of  $h + \iota k$  by  $f + \iota g$ .

**Corollary 20.49.** *Suppose that*  $f(x, y) \in U_2$ *. Then* 

1. 
$$\frac{1}{\iota} (f(x, y + \iota) - f(x, y - \iota)) \in U_2(\mathbb{C})$$
  
2. 
$$f(x, y + \iota) + f(x, y - \iota) \in U_2(\mathbb{C})$$
  
3. 
$$f(x + \iota, y - \iota) + f(x - \iota, y + \iota) - f(x + \iota, y + \iota) - f(x - \iota, y - \iota) \in U_2(\mathbb{C})$$

*Proof.* The first one is sin(D) and the second is cos(D) applied to the y variable. If we apply sin to (1) using differentiation with respect to x, then we get (3).

**Lemma 20.50.** If 
$$f(x, y) \in U_2$$
 then  $\int_0^1 f(x, y + \iota t) dt \in rHP_2$ .

*Proof.* If we write the integral as  $g + \iota h$  where g, h have real coefficients, then

$$\int_0^1 f(\alpha, y + \iota t) dt = g(\alpha, y) + \iota h(\alpha, y)$$

Since the left hand side is in  $\mathfrak{I}$ , we see that  $\mathfrak{g}(\alpha, \mathfrak{y}) \leq \mathfrak{h}(\alpha, \mathfrak{y})$  for all  $\mathfrak{y}$ . It's clear that the homogeneous parts of  $\mathfrak{g}$  and  $\mathfrak{h}$  have all positive coefficients, and so they are in  $U_2$  and interlace.

*Remark* 20.51. If  $f \leq g$  in  $\mathfrak{I}$  then we can write g in terms of the roots of f, but the coefficients might be complex. For instance, suppose

$$D_{1} = \begin{pmatrix} 95 & 40 & 78 & 61 \\ 40 & 35 & 39 & 48 \\ 78 & 39 & 70 & 60 \\ 61 & 48 & 60 & 70 \end{pmatrix} \qquad D_{2} = \begin{pmatrix} 90 & 65 & 73 & 77 \\ 65 & 57 & 65 & 68 \\ 73 & 65 & 83 & 73 \\ 77 & 68 & 73 & 90 \end{pmatrix}$$
$$S = \begin{pmatrix} 3 & 6 & 3 & 3 \\ 6 & 1 & 2 & 7 \\ 3 & 2 & 6 & 5 \\ 3 & 7 & 5 & 6 \end{pmatrix} \qquad f = |xI + yD_{1} + S + iD_{2}|$$

Since  $D_1, D_2$  are positive definite, and S is symmetric, we know that  $f \in rHP_2$ . If  $f = f_0(x) + f_1(x)y + \cdots$  then  $f_0 \leq f_1$ . If  $roots(f_0) = (r_i)$  then we can write

$$\begin{split} f_1 &= \sum_{1}^{4} a_i \frac{f_0}{x - r_i} & \text{where} \\ a_1 &= -17. - 292. \mathfrak{l} & a_2 &= -1.38 - 13.2 \mathfrak{l} \\ a_3 &= -0.775 - 12.3 \mathfrak{l} & a_4 &= 3.17 - 2.45 \mathfrak{l} \end{split}$$

## **20.7** Interlacing in $\Im$

We make the usual definition of interlacing in  $\Im$  (closure under linear combinations), and it turns out that we have already seen the definition! We then gather some properties of interlacing in  $\Im$ . We begin with the case of unequal degrees.

**Definition 20.52.** If f,  $g \in \mathfrak{I}$ , then  $f \lessdot g$  if and only if  $f + \alpha g \in \mathfrak{I}$  for all  $\alpha \in \mathbb{R}$ .

Suppose that  $f = f_{\mathfrak{R}} + \iota f_{\mathfrak{I}}$  and  $G = g_{\mathfrak{R}} + \iota g_{\mathfrak{I}}$ . f < g is equivalent to  $f_{\mathfrak{R}} + \iota f_{\mathfrak{I}} + \alpha(g_{\mathfrak{R}} + \iota g_{\mathfrak{I}}) \in \mathfrak{I}$ , which, if we express this in terms of the definition of  $\mathfrak{I}$  is

$$f_{\mathfrak{R}} + \alpha g_{\mathfrak{R}} < f_{\mathfrak{I}} + \alpha g_{\mathfrak{I}}.$$

It follows from Lemmas 1.51 and 1.52 that we have

**Lemma 20.53.** Suppose that  $f = f_{\mathfrak{R}} + \iota f_{\mathfrak{I}}$  and  $g = g_{\mathfrak{R}} + \iota g_{\mathfrak{I}}$  are in  $\mathfrak{I}$ . Then f < g if and only if

- 1.  $f_{\mathfrak{R}} \lessdot g_{\mathfrak{R}}$
- 2.  $f_{\mathfrak{I}} \lessdot g_{\mathfrak{I}}$
- 3.  $\begin{vmatrix} f_{\mathfrak{R}} & f_{\mathfrak{I}} \\ g_{\mathfrak{R}} & g_{\mathfrak{I}} \end{vmatrix} < 0$

Such polynomials exist. For instance,

 $(8 + 14x + 7x^2 + x^3) + \iota(22 + 25x + 6x^2) < (16 + 17x + 4x^2) + \iota(29 + 15x)$ 

These polynomials arise from a general construction - see Lemma 20.57.

Here are a few simple properties of  $\leq$  for  $\mathfrak{I}$ . These are the  $\mathfrak{I}$ -analogs of the usual interlacing properties in **P**. Note that multiplication preserves strict interlacing. Many of these can also be easily proved using the non-vanishing definition of  $\mathfrak{I}$ , but it interesting to see that we can prove them just using properties of interlacing polynomials.

**Lemma 20.54.** *Assume*  $f, g, h \in \mathfrak{I}$ .

- $1. \ f \lessdot f'.$
- 2. If  $f \leq g$  and  $f \leq h$  then  $f \leq \alpha g + \beta h$  for positive  $\alpha$ ,  $\beta$ . In particular,  $\alpha f + \beta g \in \mathfrak{I}$ .
- 3. If  $f \leq g \leq h$  then  $f h \leq g$ .
- 4. If  $f \leq g$  and  $\mathfrak{I}(\sigma) > 0$  then  $f + \sigma g \in \mathfrak{I}$ .
- 5. If  $f \leq g$  then  $fh \leq gh$ .
- 6. If  $f \in \mathfrak{I}^{pos}$  then  $f(-D)g \in \mathfrak{I}$ .
- 7. If  $f \in \mathfrak{I}^{pos}$  and  $f^{rev} = f^{rev}_{\mathfrak{R}} + \mathfrak{l} f^{rev}_{\mathfrak{I}}$  then  $f^{rev} \in \mathfrak{I}^{pos}$ .
- 8. Suppose that f factors as  $\prod (x \sigma_k)$ .
  - a)  $\frac{f}{x \sigma_k} \in \Im$ b)  $f < \sum a_k \frac{f}{x - \sigma_k}$  for any non-negative  $a_k$ .

*Proof.* If  $f = f_{\Re} + \iota f_{\Im}$  then  $f' = f'_{\Re} + \iota f'_{\Im}$ . Since  $f_{\Re} < f_{\Im}$ , the interlacing conditions hold. The determinant condition is Lemma 1.55.

In order to see that  $f \leq \alpha g + \beta h$  we have to check the three conditions, and that  $\alpha g + \beta h \in \mathfrak{I}$ . The first two follow from the additivity of interlacing, and the third one follows from the linearity of the determinant:

$$\begin{vmatrix} f_{\mathfrak{R}} & f_{\mathfrak{I}} \\ \alpha g_{\mathfrak{R}} + \beta h_{\mathfrak{R}} & \alpha g_{\mathfrak{I}} + \beta h_{\mathfrak{I}} \end{vmatrix} = \alpha \begin{vmatrix} f_{\mathfrak{R}} & f_{\mathfrak{I}} \\ g_{\mathfrak{R}} & g_{\mathfrak{I}} \end{vmatrix} + \beta \begin{vmatrix} f_{\mathfrak{R}} & f_{\mathfrak{I}} \\ h_{\mathfrak{R}} & h_{\mathfrak{I}} \end{vmatrix}$$

Since the real parts of f and  $\alpha g + \beta h$  interlace, as do the imaginary parts, and the determinant is negative, we can apply Lemma 1.59. This shows that the real and imaginary parts of  $\alpha g + \beta h$  interlace, and hence it is is in  $\Im$ .

If  $f \lt g \lt h$  then the argument is nearly the same as the previous one.

If we write  $\sigma = a + bi$  then  $f + \sigma g = (f + ag) + i(bg)$ . Since f + ag < bg it suffices to consider f + ig. Expanding into real and imaginary parts

$$f + \iota g = (f_{\mathfrak{R}} - g_{\mathfrak{I}}) + \iota(f_{\mathfrak{I}} + g_{\mathfrak{R}})$$

and using the interlacings  $f_{\mathfrak{R}} \ll f_{\mathfrak{I}} + g_{\mathfrak{R}} \ll g_{\mathfrak{I}}$  yields

$$\Re(f + \iota g) = f_{\mathfrak{R}} - g_{\mathfrak{I}} \lt f_{\mathfrak{I}} + g_{\mathfrak{R}} = \mathfrak{I}(f + \iota g)$$

Since  $f \in \mathfrak{I}^{pos}$  both  $f_{\mathfrak{R}}$  and  $f_{\mathfrak{I}}$  have all negative roots, and consequently  $f_{\mathfrak{R}}(-x)f_{\mathfrak{I}}(-x)$  has negative leading coefficient, and positive constant term. Consequently, Proposition 7.17 implies that we have the interlacing square



It follows that

$$f_{\mathfrak{R}}(-\mathsf{D})g_{\mathfrak{R}} - f_{\mathfrak{I}}(-\mathsf{D})g_{\mathfrak{I}} \lessdot f_{\mathfrak{R}}(-\mathsf{D})g_{\mathfrak{I}} + f_{\mathfrak{I}}(-\mathsf{D})g_{\mathfrak{R}}$$

Since all the roots of the real and imaginary part are positive, reversal preserves interlacing.

Since f < g we know that  $f + \alpha g \in \mathfrak{I}$  for all real  $\alpha$ . Thus  $hf + \alpha hg = h(f + \alpha g) \in \mathfrak{I}$  since  $\mathfrak{I}$  is closed under multiplication.

Since interlacing is preserved by addition it will suffice to show that  $f < f/(x - \sigma_k)$ . If  $\sigma_k = r_k + \iota s_k$  then we compute

$$f + \alpha \frac{f}{x - r_k - \iota s_k} = \frac{f}{x - r_k - \iota s_k} \cdot (x - r_k - \iota s_k + \alpha).$$

Since both factors are in  $\Im$ , so is their product, and thus we have shown interlacing.

There are two choices for the Hadamard product of polynomials in  $\mathfrak{I}$ . Suppose that  $f = \sum \alpha_i x^i$  and  $g = \sum \beta_i x^i$ .

- 1. Same definition as before:  $f * g = \sum \alpha_i \beta_i x^i$ .
- 2. Separate the real and complex parts:  $f *^{\iota} g = f_{\mathfrak{R}} * g_{\mathfrak{R}} + \iota f_{\mathfrak{I}} * g_{\mathfrak{I}}$ .

Surprisingly, each one preserves  $\Im$ .

**Corollary 20.55.** *If*  $f \in \mathfrak{I}$ ,  $g \in \mathfrak{I}^{pos}$  *then*  $f *^{\iota} g \in \mathfrak{I}^{pos}$ .

*Proof.* This is Lemma 7.9.

**Corollary 20.56.** *If*  $f, g \in \mathfrak{I}^{pos}$  *then*  $f * g \in \mathfrak{I}$ .<sup>1</sup>

<sup>&</sup>lt;sup>1</sup>*Unlike* P, *it's not true that the Hadamard product \* maps*  $\Im \times \Im^{pos} \longrightarrow \Im$ 

*Proof.* Write in terms of real and imaginary parts:

$$f * g = (f_{\mathfrak{R}} * g_{\mathfrak{R}} - f_{\mathfrak{I}} * g_{\mathfrak{I}}) + \iota(f_{\mathfrak{R}} * g_{\mathfrak{I}} + f_{\mathfrak{I}} * g_{\mathfrak{R}})$$

Since Hadamard product preserves interlacing in  $\mathbf{P}^{\text{pos}}$ , the interlacing of the two sides follows from the interlacings

$$f_{\mathfrak{R}} * g_{\mathfrak{R}} \lessdot f_{\mathfrak{R}} * g_{\mathfrak{I}} + f_{\mathfrak{I}} * g_{\mathfrak{R}} \ll f_{\mathfrak{I}} * g_{\mathfrak{I}}$$

The result is only in  $\Im$  since  $f_{\Re} * g_{\Re} - f_{\Im} * g_{\Im}$  might have negative signs.  $\Box$ 

We can construct interlacing polynomials in  $\Im$  from polynomials in  $P_2^{\text{pos}}$  and  $P_3^{\text{pos}}$ .

Lemma 20.57.

- 1. If  $f_0(x) + f_1(x)y + \cdots \in \mathbf{P}_2^{pos}$  and  $f_0$  has all distinct roots then  $f_i + \iota f_{i+1} \leqslant f_{i+1} + \iota f_{i+2}$ .
- 2. If  $\sum f_{i,j}(x)y^iz^j \in \mathbf{P}_3^{pos}$  and  $f_{0,0}$  has all distinct roots then

$$f_{i,j} + \iota f_{i+1,j} \leqslant f_{i,j+1} + \iota f_{i+1,j+1}$$
  
$$f_{i,j} + \iota f_{i,j+1} \leqslant f_{i+1,j} + \iota f_{i+1,j+1}$$

*Proof.* Since  $f_0 \leq f_1$  by Lemma 9.63, we know from Lemma 9.65 that  $\begin{vmatrix} f_i & f_{i+1} \\ f_{i+1} & f_{i+2} \end{vmatrix} < 0$ . From Corollary 9.64 we see that  $f_i \leq f_{i+1} \leq f_{i+2}$ . This establishes the first part.

For the second part, since  $f_{0,0}$  has all distinct roots, it follows that all interlacings are strict. Thus, in order to verify the interlacing in  $\Im$  we need to show that the determinant  $\begin{vmatrix} f_{i,j+1} & f_{i+1,j+1} \\ f_{i,j+1} & f_{i+1,j+1} \end{vmatrix}$  is positive for all x. If we substitute  $\alpha$  for x, then Corollary 9.81 says that all quadrilateral inequalities are strict if  $f(\alpha, 0, 0) \neq 0$ . Moreover, if they aren't strict, then  $f(\alpha, y, z)$  satisfies the hypothesis of Lemma 9.80, and so there is a triangular region of zeros. It follows that  $f_{0,0}(x)$  and  $f_{1,0}(x)$  have a common root, which is a contradiction.

Example 20.58. Consider an example. Write

$$(1 + x + y + z)(2 + x + 2y + z)(4 + x + 3y + 2z)$$

as any array with the usual horizontal and vertical interlacings.



Each vertical and horizontal arrow determines a polynomial in  $\mathfrak{I}$ . If we only consider the horizontal arrows then we have the following interlacings in  $\mathfrak{I}$ :

**Lemma 20.59.** If  $f(x) \in \Im$  then  $\int_0^1 f(x + \iota t) dt \in \overline{\Im}$ .

*Proof.* If we write  $f = f_{\Re} + \iota f_{\Im}$  then we know

$$F_{\mathfrak{R}} = \int_0^1 f_{\mathfrak{R}}(x + \iota t) \, dt \in \mathfrak{I} \qquad \qquad F_{\mathfrak{I}} = \int_0^1 f_{\mathfrak{I}}(x + \iota t) \, dt \in \mathfrak{I}$$

Since  $f_{\mathfrak{R}} + \alpha f_{\mathfrak{I}} \in \mathbf{P}$  for all  $\alpha$  it follows that  $F_{\mathfrak{R}} + \alpha F_{\mathfrak{I}} \in \mathfrak{I}$  for all  $\alpha \in \mathbb{R}$ . Thus,  $F_{\mathfrak{R}} \leq F_{\mathfrak{I}}$  in  $\overline{\mathfrak{I}}$ , and therefore  $\int_{0}^{1} f(x + \iota t) = F_{\mathfrak{R}} + \iota F_{\mathfrak{I}} \in \overline{\mathfrak{I}}$ .

### 20.8 Particular interlacings

It is easy to describe all the polynomials that interlace a given polynomial in  $\mathbf{P}$ . This is an unsolved problem in  $\mathfrak{I}$ . We have a number of examples and simple observations.

**Example 20.60.** In **P**,  $(x - a)^2 \leq x - b$  if and only if a = b. The first surprise is that there are many polynomials interlacing  $(x + \iota)^2$ . We show that  $\sigma$  satisfies  $(x + \iota)^2 \leq x - \sigma$  in  $\Im$  if and only if  $\sigma$  lies in the disk of radius 1 centered at  $-\iota$ .

To see this, write  $\sigma = u + w$ . The interlacing requirements are met if |u| < 1 and v > 0. The determinant requirement is that

$$0 > \begin{vmatrix} x^2 - 1 & 2x \\ x + u & \nu \end{vmatrix} = x^2(\nu - 2) - 2xu - \nu$$

This means that the discriminant is non-positive, and thus  $u^2 + (v + 1)^2 \leq 1$ . The general quadratic appears to also have a simply stated answer - see Question 184.

Example 20.61. Another difference from interlacing in P is that

$$(x + \iota)^2 \ll (x + \iota + \alpha)^2$$
 if and only if  $0 < \alpha < 2$ .

Note that

$$(\mathbf{x} + \mathbf{i} + \alpha)^2 = (\mathbf{x} + \mathbf{i})^2 + 2\alpha(\mathbf{x} + \alpha/2 + \mathbf{i})$$

In order to have interlacing we need that  $\alpha > 0$  and that  $(x+\iota)^2 < (x+\alpha/2+\iota)$ . The latter happens exactly when  $|\alpha| < 2$  by the first example

More generally, if  $x + \iota$ )<sup>2</sup>  $\ll (x + \alpha + \beta \iota)^2$ , and  $\beta$  is not zero, then

$$(\mathbf{x} + \mathbf{i} + \alpha)^2 - (\mathbf{x} + \mathbf{i})^2 = 2(\alpha \mathbf{i} - \mathbf{i} + \beta)\mathbf{x} + \cdots$$

If  $\alpha \neq 1$  then the difference is not in  $\mathfrak{I}$ .

**Example 20.62.** If we replace x by  $\beta x$  in the first two examples, and observe that interlacing is preserved under  $x \mapsto x + r$  where  $r \in \mathbb{R}$ , then we see that

1.  $(x + \sigma)^2 \ll (x + \tau)$  iff  $|\sigma - \tau| < 2\Im(\sigma)$ . 2.  $(x + \sigma)^2 \ll (x + \sigma + \alpha)^2$  iff  $0 < \alpha < 2\Im(\sigma)$ .

As  $\sigma$  approaches the real line, both interlacings converge to the usual interlacings in **P**.

**Example 20.63.** Here are some of the properties of the polynomials interlacing  $(x+\iota)^n$ . If  $(x+\iota)^n \le f(x)$  and we differentiate n-2 times then  $(x+\iota)^2 < f^{(n-2)}$ . Now the single root of  $f^{(n-2)}$  is the average of the roots of f. Consequently we conclude:

The average of the roots of f lies in the unit disk centered at 1.

Next, we claim that if  $(x + \iota)^n \leq (x + r\iota)^{n-1}$  where r is real, then  $0 < r < \frac{n}{n-1}$ . To see this, let  $f = (x + \iota)^n$  and  $g = (x + r\iota)^{n-1}$ , and note that

$$\begin{split} f_{\mathfrak{R}} &= x^n + \cdots & f_{\mathfrak{I}} = n x^{n-1} + \cdots \\ g_{\mathfrak{R}} &= x^{n-1} + \cdots & f_{\mathfrak{I}} = (n-1) x^{n-2} r + \cdots \end{split}$$

so coefficient of  $x^{2n-2}$  in  $f_{\Re}g_{\Im} - f_{\Im}g_{\Re}$  is (n-1)r - n. Since the determinant is negative, the leading coefficient must be negative, and so  $r \leq n/(n-1)$ .

If  $(x + \sigma)^n \ll (x + \tau)^n$  then differentiating n - 2 times shows that  $(x + \sigma)^2 \ll (x + \tau)^2$ , and so  $0 < \tau - \sigma < 2$ .

**Example 20.64.** Given n, we can determine an r so that if  $s \in (r, n/(n-1))$  then  $(x + \iota)^n \leq (x + s \iota)^{n-1}$ . To do this, we compute the discriminant of the determinant. This is a polynomial in s. The intervals between consecutive distinct roots of the discriminant have the property that the determinant has no multiple roots for s in the interval. If all roots of the determinant are complex, and the leading coefficient is negative then the determinant is everywhere negative.

Once we have these intervals then it is easy to verify that the real and complex parts interlace for values of s in these intervals. Table 20.1 lists a few of these intervals.

n	interval			
2	(0,	2)		
3	(1/9,	3/2)		
4	(.23,	4/3)		
5	(.34,	5/4)		
6	(.42,	6/5)		

Table 20.1: Intervals where  $(x + \iota)^n \leq (x + s \iota)^{n-1}$ 

### **20.9** The geometry of interlacing in $\Im$

There is a geometric condition for the interlacing of polynomials in  $\mathfrak{I}$ . Although it is in general hard to verify it does help to understand why polynomials in  $\mathfrak{I}$  fail to interlace.

Suppose that f,  $g \in \mathfrak{I}$  satisfy deg(f) = deg(g) + 1 and have no common factors. If they don't interlace then there is an  $\alpha \in \mathbb{R}$  and  $\sigma$  in the closed upper half plane such that  $(f + \alpha g)(\sigma) = 0$ . Now if  $\alpha = 0$  all the roots are in the lower half plane, and the roots are continuous functions of  $\alpha$ , so we see that

If f and g don't interlace then there are  $\alpha$ ,  $t \in \mathbb{R}$  such that  $(f + \alpha g)(t) = 0$ . If they do interlace then  $(f + \alpha g)(t)$  is never zero.

Now if  $f(t) + \alpha g(t) = 0$  then  $\frac{f}{g}(t) \in \mathbb{R}$ , so  $\arg(f/g)$  is a multiple of  $\pi$ . This gives us a way to check interlacing. Let

$$f = \prod (x - r_i) \qquad \qquad g = \prod (x - s_i)$$

Thus, if deg(f) = n then f and g interlace iff the *arg sum* 

$$\arg \frac{f}{g} = \sum_{1}^{n} \arg(x - r_i) - \sum_{1}^{n-1} \arg(x - s_i)$$
(20.9.1)

is not a multiple of  $\pi$  for any real x. Sometimes we can use geometric arguments to show that this is indeed the case.

**Example 20.65.** Suppose that the roots of f and g all have the same imaginary part, and their real parts alternate. (Figure 20.2) Now

$$\sum_1^n arg(x-r_\mathfrak{i}) - \sum_1^{n-1} arg(x-s_\mathfrak{i}) = arg(x-r_n) + \sum_1^{n-1} \big(arg(x-r_\mathfrak{i}) - arg(x-s_\mathfrak{i})\big).$$

Since the roots alternate all of these differences have the same sign. The arg sum is the sum of the shaded angles in the figure. It's clear that their sum is between  $\pi$  and 0 for all x, and so they interlace.



Figure 20.2: Interlacing example

**Example 20.66.** Now we look at the behavior at infinity. If deg(f) = deg(g)+1 and x goes to positive infinity then the angle between any of the 2n - 1 points and x is close to 0, so the arg sum converges to 0. As x goes to  $-\infty$  the arg sum goes to  $\pi$ . See Figure 20.3 for an example of a plot of the arg sum of two interlacing polynomials.



Figure 20.3: An argument plot for interlacing polynomials

**Example 20.67.** The polynomials whose roots are given in Figure 20.4 don't interlace. This can be seen from the plot of the arg sum in Figure 20.5 since the curve crosses the line y = 0. The failure to interlace can also be checked using Lemma 20.53.



**Example 20.68.** In this example we give four roots, for which no choice of a fifth root leads to interlacing. Consider Figure 20.6. The arg sum is

$$\angle f_1 0g_1 + \angle f_2 0g_2 + \angle 10f_2 = \pi/2 + \pi/2 + \angle 10f_3 > \pi$$



Figure 20.5: An argument plot for non-interlacing polynomials

so no choice of  $f_3$  can give interlacing polynomials.



Figure 20.6: Roots of non-interlacing polynomials

**Example 20.69.** We revisit the problem of finding all  $\beta$  for which  $(x + \iota)^n$  and  $(x + \beta \iota)^{n-1}$  interlace. The arg sum is

$$A(x, \beta) = n \arctan(1/x) - (n-1) \arctan(\beta/x)$$

The unique solution to  $A(x, \beta) = 0$  is  $\beta = n/(n-1)$  as we saw before. The solution to  $A(x, \beta) = \pi$  determines the lower bound. We find the smallest possible  $\beta$  by solving

$$A(\mathbf{x}, \boldsymbol{\beta}) = 0$$
$$\frac{\partial}{\partial \mathbf{x}} A(\mathbf{x}, \boldsymbol{\beta}) = 0$$

Using the addition formula for arc tangents

$$\arctan x + \arctan y = \arctan \frac{x+y}{1-xy}$$

we see that these are polynomial equations. The first few solutions are in Table 20.1.

# 20.10 Orthogonal type recurrences

We can define orthogonal type recurrences whose members all interlace in  $\Im$ ; the only new condition is that the imaginary part of the constant term is positive.

**Lemma 20.70.** Choose constants  $a_k > 0$ ,  $\Im(b_k) > 0$ ,  $c_k > 0$ . If  $p_{-1} = 0$ ,  $p_0 = 1$  and

$$p_{n+1} = (a_n x + b_n) p_n - c_n p_{n-1}$$

*then*  $p_n \in \mathfrak{I}$  *and we have interlacings*  $p_1 \gg p_2 \gg p_3 \gg \cdots$ .

*Proof.* We prove that  $p_{n+1} \leq p_n$  by induction on n. We first note that  $p_1 \in \mathfrak{I}$  since its only root has negative imaginary part by the hypotheses on  $\mathfrak{a}_0$  and  $\mathfrak{b}_0$ . Thus,  $p_1 \leq p_0$  since all linear combinations are in  $\mathfrak{I}$ .

Next, assume that  $p_n < p_{n-1}$ . All the leading coefficients of the polynomials in the interlacing

$$(\mathfrak{a}_n \mathfrak{x} + \mathfrak{b}_n) \mathfrak{p}_n \lessdot \mathfrak{p}_n \lessdot \mathfrak{c}_n \mathfrak{p}_{n-1}$$

are positive, and so it follows from Lemma 20.54 that

$$(\mathfrak{a}_n \mathbf{x} + \mathfrak{b}_n) \mathfrak{p}_n - \mathfrak{c}_n \mathfrak{p}_{n-1} \lessdot \mathfrak{p}_n$$

which finishes the proof.

**Example 20.71.** We get two interrelated recurrences if we separate into real and imaginary parts. Consider the simple example

$$\mathfrak{p}_{\mathfrak{n}+1} = (\mathfrak{x} + \mathfrak{i}) \mathfrak{p}_{\mathfrak{n}} - \mathfrak{p}_{\mathfrak{n}-1}$$

If we write  $p_n = r_n + \iota s_n$  then

$$r_{n+1} = x r_n - r_{n-1} - s_n$$
  
 $s_{n+1} = x s_n - s_{n-1} + r_n$ 

The recurrences for just the r's and s's are the same:

$$\begin{split} r_{n+1} &= 2x\,r_n - (3+x^2)\,r_{n-1} + 2x\,r_{n-2} - r_{n-3} \\ s_{n+1} &= 2x\,s_n - (3+x^2)\,s_{n-1} + 2x\,s_{n-2} - s_{n-3} \end{split}$$

It is hard to see how one would ever show that the r's and s's interlace if we only had these recurrences.

**Example 20.72.** Consider the Chebyshev recurrence  $p_{n+1} = 2x p_n - p_{n-1}$ , except we start with  $p_1 = x + \iota$ . An easy induction shows that

$$p_n = T_n + \iota U_{n-1}$$

We can see directly that this is in  $\Im$  since the Chebyshev polynomials  $T_n$  and  $U_n$  have positive leading coefficients and satisfy  $T_n \triangleleft U_{n-1}$ .

<b>Rising Factorial</b>	$p_n =$	$\langle \underline{\mathbf{x}} \rangle_{\mathbf{n}} + \mathbf{i} \langle \underline{\mathbf{x}} \rangle_{\mathbf{n}-1}$
	$p_n =$	$\lfloor n-1-\iota+x \rfloor p_{n-1}+\iota \lfloor n-3+x \rfloor p_{n-2}$
Hermite	$p_n =$	$H_n + \iota H_{n-1}$
	$p_n =$	$2xp_{n-1}-p'_{n-1}$
Bell	$p_n =$	$B_n + \iota B_{n-1}$
	$p_n =$	$x(p_{n-1} + p'_{n-1})$
Euler	$p_n =$	$B_n + \iota B_{n-1}$
	$p_n =$	$(nx - \iota)p_{n-1} + (n-2)\iota x p_{n-2} + (x - x^2)(p'_{n-1} + \iota p'_{n-2})$
Laguerre	$p_n =$	$L_n(-x) + \iota L_{n-1}(-x)$
-	$n p_n =$	$\big[x+n-\iota(n-2)\big]p_{n-1}+\iota(n-2)p_{n-2}+(1+\iota)p_{n-1}'$
Legendre	$p_n =$	$P_n + \iota P_{n-1}$
-	$p_n =$	$(x+\iota)p_{n-1}-(n-2)p_{n-2}+2x^2p_{n-1}'-2xp_{n-2}'$
Gegenbauer	$p_n =$	$G_n^{(2)} + \iota G_n^{(2)}$
0	$n(n-1)p_n =$	$2(n+2)(n-1) x p_{n-1} - (n+1)(n+2)p_{n-2} + 2(1-x^2)p'_{n-1}$

Table 20.2: Recurrences for  $f_n + \iota f_{n-1}$ 

Here's a simple generalization of that fact.

**Lemma 20.73.** Suppose orthogonal polynomial sequences  $\{p_i\}$  and  $\{q_i\}$  both satisfy the recurrence

$$r_{n+1}(x) = (a_n x + b_n)r_n(x) - c_n r_{n-1}(x)$$

where all  $a_n$  and  $c_n$  are positive. If

$$p_0 = 1$$
,  $p_1 = x$ ,  $q_0 = 0$ ,  $q_1 = 1$ 

then

$$p_n < q_n$$
 and  $\begin{vmatrix} p_n & p_{n+1} \\ q_n & q_{n+1} \end{vmatrix} > 0.$ 

*Proof.* Let  $w_i$  satisfy the same recurrence with  $w_0 = 1$  and  $w_1 = x + \iota$ . By induction  $w_n = p_n + \iota q_n$ , and  $w_n$  satisfies the conditions of Lemma 20.70. Thus,  $w_n < w_{n+1}$  which gives the conclusions of the lemma.

If we have a sequence of interlacing polynomials  $f_k \leq f_{k-1}$  and we let  $p_n = f_n + \iota f_{n-1}$  then all  $p_n$  are in  $\overline{\mathfrak{I}}$ . Table 20.10 lists recurrence relations for the  $p_n$  for several polynomial sequences.

## **20.11** The transformation $x \mapsto \exp(-\iota x)$

We have seen (Lemma 5.10) that  $\sin(x) - \alpha \in \widehat{\mathbf{P}}$  when  $-1 \leq \alpha \leq 1$ . The next lemma extends this to complex exponentials. We then use this result to determine mapping properties of the Chebyshev polynomials.

**Lemma 20.74.** *If*  $|\sigma| \leq 1$  *then*  $e^{-\iota x} + \sigma \in \widehat{\mathfrak{I}}$ .

*Proof.* If we write x = a + bi then

$$e^{-\iota x} + \sigma = e^{-\iota \iota} e^{b} + \sigma$$

Now  $|e^{-\alpha \iota}| = 1$ , and  $e^b > 1$  since b > 0, so the sum is non-zero since  $|\sigma| \leq 1$ .

**Proposition 20.75.** If  $f \in \mathbf{P}^{\Delta}$  then  $f(e^{-\iota x}) \in \widehat{\mathfrak{I}}$ .

*Proof.* This is an immediate consequence of Lemmas 20.34 and 20.74. This is essentially due to Pólya: see [89, page 50].

**Corollary 20.76.** If  $T_n$  and  $U_n$  are the Chebyshev polynomials then  $x^n \mapsto T_n(x)$ and  $x^n \mapsto U_n(x)$  map  $P^{(-1,1)} \longrightarrow P^{(-1,1)}$ .

*Proof.* We only consider  $T_n$ ; the case of  $U_n$  is similar.  $T_n(x)$  is defined by  $T_n(\cos x) = \cos(nx)$ , so the diagram below commutes at the element level. Proposition 20.75 implies that the top arrow maps as claimed, and hence  $x^n \mapsto \cos nx$  maps as in the diagram. Since  $g(\cos x) \in \widehat{\mathbf{P}}$  if and only if  $g(x) \in \mathbf{P}^{(-1,1)}$ , the proof is complete.



**Example 20.77.**  $\frac{1-e^{-\imath y}}{y}$  is another function in  $\hat{\jmath}$ . To see that this is so, note that

$$\frac{1 - e^{-y\iota}}{y} = \frac{1 - \cos y}{y} + \iota \frac{\sin y}{y}$$
$$= \frac{2 \sin^2(y/2)}{y} + \iota \frac{\sin y}{y}$$

The infinite product expansion shows that  $\sin^2(y/2)/y$  and  $\sin(y)/y$  are in  $\widehat{\mathbf{P}}$ . Moreover, if we approximate them by their partial products then they interlace ( $\underline{<}$ ). For instance, for zeros up to  $\pm 2\pi$  the roots are

$$\frac{\sin^2(y/2)/y:}{\sin(y)/y:} -2\pi, -2\pi, 0, 0, 2\pi, 2\pi$$

If we know that a polynomial is in  $\mathbf{P}^{\Delta}$  then we can use Proposition 20.75 to get information about trigonometric and exponential polynomials. We use a result due to Eneström-Kakeya [20] to create a polynomial in  $\mathbf{P}^{\Delta}$ .

**Theorem 20.78.** *If*  $f(x) = a_0 + a_1x + \cdots + a_nx^n$  *and* 

$$a_n > a_{n-1} > \cdots > a_0 > 0$$
 (20.11.1)

then  $f \in \mathbf{P}^{\Delta}$ .

**Corollary 20.79.** *If the real numbers*  $\{a_k\}$  *satisfy* (20.11.1) *then* 

- 1.  $\sum_{k=0}^{n} a_k \sin(kx)$  has all real roots.
- 2.  $\sum_{k=0}^{n} a_k \cos(kx)$  has all real roots.
- 3. The roots of  $\sum_{k=0}^{n} a_k \sin(kx)$  and  $\sum_{k=0}^{n} a_k \cos(kx)$  interlace.

*Proof.* Since  $f(x) = a_0 + a_1x + \cdots + a_nx^n$  is in  $\mathbf{P}^{\Delta}$ , we can apply Proposition 20.75 to conclude that  $\sum a_k e^{-\iota kx} \in \widehat{\mathfrak{I}}$ . This implies (1), (2), and (3).

If we take limits we get [139, III 205]

**Corollary 20.80.** If  $\phi(x)$  is a positive increasing function on (0,1) with  $\int_0^1 \phi(x) dx$  finite then

$$\int_0^1 \phi(x) e^{-\iota t x} \, dx \in \widehat{\mathfrak{I}}.$$

*Proof.* Since the integral of  $\phi$  is finite we have

$$\int_0^1 \phi(x) e^{-\iota tx} dx = \lim_{n \to \infty} \sum_{k=1}^n \frac{1}{n} \phi(k/n) e^{-\iota tk/n}$$

Since  $\phi$  is increasing the coefficients  $(1/n)\phi(k/n)$  are positive and increasing. It follows from Proposition 20.75 that the sum is in  $\hat{\mathcal{I}}$ , and hence the limit is as well.

### 20.12 Newton inequalities

We have seen mutually interlacing polynomials evaluated at points in the upper half plane (Figure 3.4). We now give an example of the ratio of the coefficients of a polynomial in  $\mathbf{P}_2^{\text{pos}}$  evaluated at a point in  $\mathbf{Q}_I$ . If  $f = \sum f_i(x)y^i$  then Figure 20.7 shows the ratios of a fourth degree polynomial evaluated at a point  $\sigma \in \mathbf{Q}_I$ .

There are two striking facts in this picture:



Figure 20.7: Coefficient ratios of  $f \in \mathbf{P}_2^{\text{pos}}$  and  $\sigma \in \mathsf{Q}_I$ 

1. The ratios are all in the first quadrant.

2. The real and imaginary parts are increasing.

The first part is easy; the second part is contained in the following lemma. We can differentiate, so we only need to establish it for  $f_0$  and  $f_1$ . Since  $f_0 \leq f_1$ , we can write

$$\frac{f_0}{f_1} = \left(\sum \frac{a_i}{x + r_i}\right)^{-1}$$

where the  $a_i$  and the  $r_i$  are positive. If  $\sigma \in Q_I$  then  $\sigma + r_i \in Q_I$ , and so  $\sum (\sigma + r_i)^{-1} \in Q_{IV}$ , which establishes the first observation above.

The lemma below is another generalization of Newton's inequality. If we let  $f = \sum a_i x^i \in \mathbf{P}^{pos}$  then  $f(x + y) \in \mathbf{P}_2^{pos}$ . Choosing  $\sigma = 0$  in the Proposition gives that

$$0 \leqslant \mathfrak{R}\frac{\mathfrak{a}_{\mathfrak{i}}}{\mathfrak{a}_{\mathfrak{i}+1}} \leqslant \mathfrak{R}\frac{\mathfrak{a}_{\mathfrak{i}+1}}{\mathfrak{a}_{\mathfrak{i}+2}}$$

since  $f^{(i)}(0)/i! = a_i$ . The imaginary part is vacuous.

**Lemma 20.81.** Suppose that  $\sum f_i(x) y^i \in P_2^{\text{pos}}(n)$ . For all  $\sigma \in Q_I$  and  $0 \leqslant i \leqslant n-2$ 

$$\begin{array}{rcl} 0 & \leqslant & \mathfrak{R} \; \frac{f_{\mathfrak{i}}(\sigma)}{f_{\mathfrak{i}+1}(\sigma)} & \leqslant & \mathfrak{R} \; \frac{f_{\mathfrak{i}+1}(\sigma)}{f_{\mathfrak{i}+2}(\sigma)} \\ \\ 0 & \leqslant & \mathfrak{I} \; \frac{f_{\mathfrak{i}}(\sigma)}{f_{\mathfrak{i}+1}(\sigma)} & \leqslant & \mathfrak{I} \; \frac{f_{\mathfrak{i}+1}(\sigma)}{f_{\mathfrak{i}+2}(\sigma)} \end{array}$$

*Proof.* Since  $\sigma \in Q_I$  we know that  $f(\sigma, y) \in \mathfrak{I}$ . The inequality now follows from Lemma 20.82.

If all the roots of a polynomial lie in a quadrant then there are Newton inequalities for the real parts, imaginary parts, and magnitudes of the coefficients.

**Lemma 20.82.** Suppose that  $\sum_{0}^{n} a_i x^i$  has all roots in a quadrant. The table below summarizes the properties of the sequence  $\frac{a_0}{a_1}$ ,  $\frac{a_1}{a_2}$ ,  $\frac{a_2}{a_3}$ ,... where "neg,  $\downarrow$ " means it is a decreasing sequence of negative terms, and so on.

	$Q_{\mathrm{I}}$	$Q_{II}$	$Q_{\rm III}$	$Q_{\rm IV}$
Real parts	neg,↓	pos,↑	pos,↑	neg,↓
Imaginary parts	neg,↓	neg,↓	pos,↑	pos,↑
Magnitude	$\uparrow$	$\uparrow$	Ŷ	$\uparrow$

*Proof.* As usual, we differentiate, reverse and differentiate; the result is a quadratic with all roots in the original quadrant. We will work with the third quadrant since we can use properties of  $\Im$ . The upper two quadrants follow using conjugation, and the fourth quadrant is similar.

Thus, we may assume that our polynomial is

$$(x + a)(x + b) + \iota c(x + d)$$
  $0 < a < d < b$ ,  $0 < c$ 

We simply need to compute the real parts, imaginary parts and magnitudes, and show that the inequalities are satisfied. If we write the polynomial as  $a_0 + a_1x + a_2x^2$  then

$$\begin{aligned} a_0 &= ab + icd & a_1 &= a + b + ic \\ a_2 &= 1 & \\ \Re \frac{a_0}{a_1} &= \frac{ba^2 + b^2a + c^2d}{a^2 + 2ba + b^2 + c^2} & \\ \Re \frac{a_0}{a_1} &= \frac{c(-ab + db + ad)}{a^2 + 2ba + b^2 + c^2} & \\ \Im \frac{a_0}{a_1} &= \frac{a^2b^2 + c^2d^2}{(a + b)^2 + c^2} & \\ \left| \frac{a_1}{a_2} \right|^2 &= (a + b)^2 + c^2 \end{aligned}$$

All six terms are clearly positive, and it is easy to check that all three inequalities are satisfied.

The bound for absolute values is stronger that the bound we get from considering sectors - see 677.

CHAPTER



# Stable polynomials

A one variable polynomial is stable (sometimes called Hurwitz stable) if all its roots lie in the closed left half plane. This is a well studied class of polynomials with many applications - e.g. [34]. There are two classes of stable polynomials in one variable:

 $\mathcal{H}_1$  = real coefficients and roots in the closed left half plane  $\mathcal{H}_1(\mathbb{C}) =$  complex coefficients and roots in the closed left half plane

Since polynomials whose roots are in the closed left half plane are non-zero in the right half plane, we can apply the results of Chapter 19 to get properties of stable polynomials in d variables.

### Definition 21.1.

 $\mathcal{H}_{d}(\mathbb{C}) = \begin{cases} \text{All polynomials } f(x_{1}, \dots, x_{d}) \text{ with complex coefficients} \\ \text{such that } f(\sigma_{1}, \dots, \sigma_{d}) \neq 0 \text{ for all } \sigma_{1}, \dots, \sigma_{d} \text{ in the right} \\ \text{half plane. If we don't need to specify d we simply write} \\ \mathcal{H}(\mathbb{C}). \end{cases}$ We

call such polynomials stable polynomials. In one variable they are often called Hurwitz stable.

After establishing the basic properties of stable polynomials in d variables, we focus our attention on stable polynomials with real coefficients in one variable. We will see that whenever a polynomial is shown to be strictly positive, it's likely that it's stable. There are three kinds of interlacing for stable polynomials, and *positive interlacing* is especially useful for showing that a polynomial is stable.

For a short summary of the properties of stable polynomials, see [58].

### **21.1** Stable polynomials in d variables

We apply the general results of Chapter 19 to derive results about polynomials non-vanishing in the open right half plane (RHP). The reversal constant is 1, and RHP/RHP =  $\mathbb{C} \setminus (-\infty, 0)$ . If f, g  $\in \mathcal{H}_d(\mathbb{C})$  then we write f  $\xleftarrow{H}$  g if  $f + zg \in \mathcal{H}_{d+1}(\mathbb{C})$ . From Chapter 19 we have

**Lemma 21.2.** *Suppose that*  $f \in \mathcal{H}_d(\mathbb{C})$ *.* 

- 1. If  $\sigma_i \in RHP$  and  $r_i > 0$  then  $f(r_1x_1 + \sigma_1, \dots, r_dx_d + \sigma_d) \in \mathcal{H}_d(\mathbb{C})$ .
- 2.  $f(x_1 + y, x_2, \dots, x_d) \in \mathcal{H}_{d+1}(\mathbb{C}).$
- 3. If  $\sigma \in \mathbb{R}$  then  $f(\iota \sigma, x_2, \ldots, x_d) \in \mathcal{H}_{d-1}(\mathbb{C}) \cup \{0\}$ .
- 4. If  $f \xleftarrow{H} g$  then  $g \xleftarrow{H} f$ .
- 5. If  $f \xleftarrow{H} g \xleftarrow{H} h$  then  $f + h \xleftarrow{H} g$ .
- 6.  $\mathfrak{H}_{d}(\mathbb{C})$  is closed under differentiation. That is, if  $g \in \mathfrak{H}_{d}(\mathbb{C})$  then  $\frac{\partial}{\partial x_{i}} f \in \mathfrak{H}_{d}(\mathbb{C}) \cup \{0\}$ .
- 7. f  $\leftarrow^{H} \partial f / \partial x_i$
- 8. If  $\sum x^i f_i(\mathbf{y}) \in \mathcal{H}_{d+1}(\mathbb{C})$  then  $f_i(\mathbf{y}) \in \mathcal{H}_d(\mathbb{C}) \cup \{0\}$  and  $f_i \xleftarrow{\mathsf{H}} f_{i+1}$ .
- 9.  $f(\mathbf{x}) \times g(\mathbf{x}) \mapsto f(\partial \mathbf{x})g(\mathbf{x})$  determines a map

$$\mathcal{H}_{d}(\mathbb{C}) \times \mathcal{H}_{d}(\mathbb{C}) \longrightarrow \mathcal{H}_{d}(\mathbb{C}).$$

- 10. If  $\sum x^i f_i(\mathbf{y}) \in \mathcal{H}_d(\mathbb{C})$  has degree n then  $\sum x^{n-i} f_i(\mathbf{y}) \in \mathcal{H}_d(\mathbb{C})$ .
- 11. If  $x_i$  has degree  $e_i$  then  $x_1^{e_1} \cdots x_d^{e_d} f(1/x_1, \dots, 1/x_d) \in \mathcal{H}_d(\mathbb{C})$ .
- 12. The Hadamard product  $\sum a_i x^i \times \sum f_i(\mathbf{y}) x^i \mapsto \sum a_i f_{n-i}(\mathbf{y}) x^i$  determines a map  $P^{\text{pos}} \times \mathcal{H}_d(\mathbb{C}) \longrightarrow \mathcal{H}_d(\mathbb{C})$ .
- 13. If S is skew-symmetric and all  $D_i$  are positive definite then  $|S + \sum_{i=1}^{d} x_i D_i| \in \mathcal{H}_d$ .
- 14. The following are equivalent
  - a)  $f(\mathbf{x}) \in \mathcal{H}_d$
  - b)  $f(\mathbf{a} + t\mathbf{b}) \in \mathfrak{H}_1$  for  $\mathbf{a}, \mathbf{b} > 0$ .

*Proof.* Part (1) shows a) implies b). Conversely, consider  $\sigma_1, \ldots, \sigma_d$  in RHP. Since 1 is in the closure of RHP we can find an  $\alpha$  and positive  $\mathbf{a}, \mathbf{b}$  such  $(\sigma_1, \ldots, \sigma_d) = \mathbf{a} + \alpha \mathbf{b}$ . Thus,  $f(\sigma_1, \ldots, \sigma_d) = f(\mathbf{a} + \mathbf{b}\alpha) \neq 0$ .

### 21.2 Real stable polynomials

We now consider properties that depend on real coefficients. It is immediate that  $\mathbf{P}^{\text{pos}} \subset \mathcal{H}_1$ ; this is still true for more variables.  $\mathbf{P}_d^{\text{pos}}$  was defined in terms of  $U_d$  and the positivity of the coefficients. We can also express this in terms of non-vanishing.

**Lemma 21.3.**  $f(\mathbf{x}) \in \mathbf{P}_d^{pos}$  if and only if  $f(\frac{\sigma_1}{\sigma_0}, \dots, \frac{\sigma_d}{\sigma_0}) \neq 0$  for all  $\sigma_i$  in the upper half plane.

*Proof.* We know that  $f(\mathbf{x}) \in \mathbf{P}_d^{\text{pos}}$  if and only if its homogenization is in  $U_{d+1}$ . Now the homogenization equals  $x^n f(\frac{x_1}{x}, \cdots, \frac{x_d}{x})$  where n is the degree of f, so the lemma is now clear.

**Lemma 21.4.** 
$$\overline{\mathbf{P}}_{d}^{pos} = \mathcal{H}_{d} \cap \overline{\mathbf{P}}_{d}$$

*Proof.* We first show that  $\mathbf{P}_d^{\text{pos}} \subset \mathcal{H}_d \cap U_d$ . Suppose that  $f(\mathbf{x}) \in \mathbf{P}_d^{\text{pos}}$ , and  $\tau_1, \ldots, \tau_d$  are in the right half plane. Now  $\iota \tau_1, \ldots, \iota \tau_d$  lie in the upper half plane, and so by Lemma 21.3

$$0 \neq f\big(\frac{\iota\tau_1}{\iota}, \dots, \frac{\iota\tau_d}{\iota}\big) = f(\tau_1, \dots, \tau_d)$$

Thus,  $f(\mathbf{x}) \in \mathcal{H}_d$ . Next, if  $f \in \mathcal{H}_d \cap \overline{\mathbf{P}}_d$  then  $f \in \overline{\mathbf{P}}_d$ , and f has all non-negative coefficients, so  $f \in \overline{\mathbf{P}}_d^{\text{pos}}$  by Lemma 11.58

**Corollary 21.5.** The unsigned reversal of a polynomial in  $\mathbf{P}_{d}^{pos}$  is in  $\mathcal{H}_{d}$ .

*Proof.* The only point is that  $\mathbf{P}_d^{\text{pos}} \subset \mathcal{H}_d$ .

**Lemma 21.6.** If  $f \in H_d$  then all coefficients have the same sign.

*Proof.* By Lemma 11.57 and following the proof of Lemma 11.58 we can write f as a limit of  $f_{\varepsilon} \in \mathcal{H}_d$  where all coefficients of  $f_{\varepsilon}$  are non-zero. Since the coefficients of a polynomial in  $\mathcal{H}_1$  with all non-zero coefficients have the same sign, an easy induction shows that all the coefficients of  $f_{\varepsilon}$  have the same sign. Thus, the non-zero coefficients have the same sign.

**Lemma 21.7.** If  $f(\mathbf{x}) \in \mathfrak{H}_d$  and the homogenization  $\mathsf{F}$  of  $\mathsf{f}$  is in  $\mathfrak{H}_{d+1}$  then  $\mathsf{f} \in \overline{\mathbf{P}}_d^{\text{pos}}$ .

*Proof.* If F is in  $\mathcal{H}_{d+1}$  then for  $\tau_0, \ldots, \tau_d \in \mathsf{RHP}$  we know that

$$f\big(\frac{\tau_1}{\tau_0},\ldots,\frac{\tau_d}{\tau_0}\big)\neq 0$$

If  $\sigma_1, \ldots, \sigma_d \in \mathsf{UHP}$  then we can choose  $\tau_0 \in \mathsf{RHP}$  so that  $\tau_0^{-1}$  rotates all of the  $\sigma_i$  to the right half plane. Thus  $\sigma_i = \tau_i / \tau_0$  where  $\tau_i \in \mathsf{RHP}$ , and so f is non-vanishing on the upper half plane. Thus  $f \in \overline{\mathbf{P}}_d$ , and by Lemma 21.6 all coefficients of f have the same sign. Consequently, f is in  $\overline{\mathbf{P}}_d^{pos}$ .

*Remark* 21.8. Here is a particular example of a polynomial in  $\mathcal{H}_2$  whose homogenization is not in  $\mathcal{H}_3$ . Consider  $f(x, y) = 1 + (x+y) + (x+y)^2$  with homogenization F(x, y, z). F has a root in the right half plane:  $F(1, -.23 - 1.86\iota, 2\iota) = 0$ .

Multiplying each variable by a new variable is similar to homogenizing.

**Lemma 21.9.**  $f(y x) \in \mathcal{H}_{d+1}$  *if and only if*  $f(x) \in P_d^+$ .

*Proof.* If  $f(y\mathbf{x}) \in \mathcal{H}_{d+1}$  then substituting 1 shows  $f(\mathbf{x}) \in \mathcal{H}_d$ . If  $\sigma_1, \ldots, \sigma_d \in$  UHP then choose  $\tau \in \mathsf{RHP}$  so that  $\tau \sigma_1, \ldots, \tau \sigma_d \in \mathsf{RHP}$ . Since  $\tau^{-1} \in \mathsf{RHP}$ 

$$f(\sigma_1,\ldots,\sigma_d) = f(\tau^{-1}(\tau\sigma_1,\ldots,\tau\sigma_d)) \neq 0$$

Thus  $f \in U_d$ , so  $f \in P_d^+$ . Conversely, assume that  $f \in P_d^+$ . We know that the homogenization of f is in  $P_{d+1}^+$ . Writing  $f = \sum a_I x^I$  and taking the reverse yields

$$\sum a_I x^I \, \boldsymbol{y}^{n-|I|} \implies \sum a_I x^I \, \boldsymbol{y}^{|I|} = \boldsymbol{\mathsf{f}}(\boldsymbol{y} \cdot \boldsymbol{x}) \in \boldsymbol{\mathfrak{H}}_{d+1}$$

so the proof is complete.

**Lemma 21.10.** *If*  $f(\mathbf{x}, \mathbf{y}^2) \in \mathcal{H}_d$  *then*  $f(\mathbf{x}, \mathbf{y}) \in \mathbf{P}^{pos}d$ .

*Proof.* See the proof of the corresponding result for  $U_d$ , Lemma 20.4.

**Lemma 21.11.** Suppose that  $f(x) = \sum_{0}^{n} a_{i}x^{i} \in \mathcal{H}_{1}$ . If some internal coefficient is zero then  $f(x) = x^{r}g(x^{2})$  where  $g \in \mathbf{P}^{pos}$ .

*Proof.* We can write

$$f(x) = x^r \times \prod_1^s (x + a_i) \times \prod_1^t ((x + b_i)^2 + c_i)$$

where  $a_i$  is positive,  $b_i$ ,  $c_i \ge 0$  and  $b_i c_i \ne 0$ . If the second factor is non-empty then there are no internal zeros. If any  $b_i$  is non-zero there are no internal zeros, so  $f(x) = x^r \prod (x^2 + c_i)$  from which the conclusion follows.

**Lemma 21.12** (Hermite-Biehler). Suppose that  $f(\mathbf{x})$  is a polynomial, and write  $f(\mathbf{x}) = f_{\mathsf{E}}(\mathbf{x}) + f_{\mathsf{O}}(\mathbf{x})$  where  $f_{\mathsf{E}}(\mathbf{x})$  (resp.  $f_{\mathsf{O}}(\mathbf{x})$ ) consists of all terms of even (resp. odd) degree. Then

 $f \in \mathcal{H}(\mathbb{C})$  if and only if  $f_E \xleftarrow{H} f_O$ .

*Proof.* Since  $f(\mathbf{x}) = f_E(\mathbf{x}) + f_O(\mathbf{x}) \in \mathcal{H}$  we know  $f(-\mathbf{i}\mathbf{x}) = f_E(-\mathbf{i}\mathbf{x}) + f_O(-\mathbf{i}\mathbf{x}) \in U(\mathbb{C})$ . Now  $f_E(-\mathbf{i}\mathbf{x})$  has all real coefficients, and  $f_O(-\mathbf{i}\mathbf{x})$  has all imaginary coefficients, so by Hermite-Biehler for  $U(\mathbb{C})$  we know that

$$f_{E}(-\iota x) + (y/\iota)f_{O}(-\iota x) \in U(\mathbb{C})$$

Returning to  $\mathcal{H}(\mathbb{C})$  yields  $f_{\mathsf{E}}(\mathbf{x}) + yf_{\mathsf{O}}(\mathbf{x}) \in \mathcal{H}(\mathbb{C})$ .

### **21.3** Polynomials (not) in $\mathcal{H}_d$

We give some examples of polynomials that are (not) in  $\mathcal{H}_d$ .

**Example 21.13.** Since the map  $z \mapsto z^2$  maps  $\mathsf{RHP} \longrightarrow \mathbb{C} \setminus (-\infty, 0)$  it follows that the equation  $x^2 + y^2 = 0$  has solutions in the right half plane, so  $x^2 + y^2 \notin \mathcal{H}_2$ .

**Example 21.14.** We claim  $xy + a^2 \in \mathcal{H}_2$  if  $a \in \mathbb{R}$ . As above, the image of RHP under xy is  $\mathbb{C} \setminus (-\infty, 0)$ , so adding a positive amount will not make xy equal to zero.

**Example 21.15.** If  $\sigma$  is complex (and not negative real) then  $x^2 - \sigma \notin \mathcal{H}_1$ . Indeed, one of  $\sigma^{1/2}$ ,  $-\sigma^{1/2}$  lies in the right half plane. Consequently,  $xy - \sigma \notin \mathcal{H}_2$ .

These last two examples imply that

**Lemma 21.16.** If f is a polynomial with positive coefficients then  $f(xy) \in \mathcal{H}_2$  if and only if  $f \in \mathbf{P}^{pos}$ .

The stable polynomials of degree 1 in  $\mathcal{H}_2$  are clearly ax + by + c where a, b are positive and c is non-negative. If a, b, c are positive then we know that

$$ax^2 + bx + c \in egin{cases} \mathbf{P}^{\mathrm{pos}} & ext{if } b^2 - 4ac \geqslant 0 \ \mathcal{H}_1 & ext{if } b^2 - 4ac \leqslant 0 \end{cases}$$

**Lemma 21.17.** *If* a, b, c, d *are positive then*  $axy + bx + cy + d \in \mathcal{H}_2$ .

*Proof.* If a + bx + cy + dxy = 0 then  $y = -\frac{a+bx}{c+dx}$ . By the following lemma the Möbius transformation  $z \mapsto \frac{a+bz}{c+dz}$  maps the right hand plane to itself. Thus, if  $x \in \mathsf{RHP}$  then  $y \notin \mathsf{RHP}$ , and so  $a + bx + cy + dxy \in \mathcal{H}_2$ .

This can also be derived from the corresponding criterion for  $U_2(\mathbb{C})$ .

*Remark* 21.18. It is not the case that a multiaffine polynomial with all positive coefficients is in  $\mathcal{H}_3$ . If

$$f(x, y, z) = (135/4) + x + y + z + xy + xz + yz + 24xyz$$

then  $f(1/2 + \iota, 1/2 + \iota, 1/2 + \iota) = 0$ , so  $f \notin \mathcal{H}_3$ .

**Lemma 21.19.** If the Möbius transformation M satisfies  $Mz = \frac{a+bz}{c+dz}$  then the following are equivalent:

- 1. M maps the right half plane to itself.
- 2. At least one of the following holds
  - a) a, b, c, d have the same sign (or some are zero).
  - b) b, d have the same sign and ad = bc. In this case the image is the single point b/d.

*Proof.* If  $r + \iota s \in \mathsf{RHP}$  then we need to show that  $\mathsf{M}(r + \iota s) \in \mathsf{RHP}$ . Equivalently, we show that if r > 0 then  $\Re \mathsf{M}(r + \iota s) > 0$ . Now

$$M(r+\iota s) = \frac{bds^2 + (a+br)(c+dr)}{(c+dr)^2 + d^2s^2} + \iota \frac{(bc-ad)s}{(c+dr)^2 + d^2s^2}$$

and we can ignore the positive denominator, so we need to show that

$$bds^{2} + (a + br)(c + dr) > 0$$
 (21.3.1)

but this is clear if a, b, c, d, r are positive. If ad = bc then Mz = b/d which is in the right half plane since b/d > 0.

Conversely, assume (21.3.1) holds. Assume that  $abcd \neq 0$ ; the remaining cases are easily taken care of. Taking s large shows that bd is positive, so b and d have the same sign. Taking s = 0 and r close to zero shows that ac is positive, so a and c have the same sign. Take s = 0. If ab < 0 then the factor (a + br) has a positive root, and so (a + br)(c + dr) will be negative for some r close to that root, unless the root is also a root of the other factor. In this case a + bz is a multiple of c + dz, and so ad = bc.

There are many quadratic forms in  $\mathcal{H}_d$ . The next lemma generalizes the fact that  $x^2 - 1 \in \mathbf{P}$ , and  $x^2 + 1 \in \mathcal{H}_1$ .

Lemma 21.20. If Q is a d by d negative subdefinite matrix then

$$\mathbf{x}^{t}Q\mathbf{x} + \boldsymbol{\alpha} \in \begin{cases} \boldsymbol{\mathcal{H}}_{d} & \textit{if } \boldsymbol{\alpha} \geqslant \boldsymbol{0} \\ \overline{\mathbf{P}}_{d} & \textit{if } \boldsymbol{\alpha} \leqslant \boldsymbol{0} \end{cases}$$

*Proof.* If  $\alpha < 0$  then  $\mathbf{x}^t Q \mathbf{x} + \alpha \in \overline{\mathbf{P}}_d$  by Lemma 10.87. Now if  $f(x_1, \dots, x_d) \in H\mathbf{P}_d$  then  $f(\mathfrak{u}_1, \dots, \mathfrak{u}_d) \in \mathcal{H}_d$ . Since  $\overline{\mathbf{P}}_d \subset H\mathbf{P}_d$  it follows that

$$-(\mathbf{x}^{\mathsf{t}} Q \mathbf{x} - \alpha) = (\mathbf{\iota} \mathbf{x})^{\mathsf{t}} Q(\mathbf{\iota} \mathbf{x}) + \alpha \in \mathcal{H}_{\mathsf{d}}.$$

The graphs of stable polynomials (Figure 21.1 is a sketch of one) are different from the graphs of polynomials in  $U_2$ .

The following famous theorem of Heilman-Lieb determines a multiaffine stable polynomial.

**Theorem 21.21.** *Let* G = (V, E) *be a graph,*  $V = \{1, ..., n\}$ *. To each edge*  $e = ij \in E$  *assign a non-negative real number*  $\lambda_{ij}$ *. Then the polynomial* 

$$M_{G}(z) = \sum_{M \text{ is a matching } ij \in M} \prod_{ij \in M} \lambda_{ij} z_{i} z_{j}$$

is stable.



Figure 21.1: The graph of a stable polynomial in two variables

### 21.4 Stable polynomials with complex coefficients

In this section we consider polynomials with complex coefficients in one variable. Since a 90° rotation moves the right half plane to the upper half plane, we can easily transfer properties from  $\mathfrak{I}$  to  $\mathfrak{H}_1(\mathbb{C})$ . If f has degree n define  $\phi(f) = \mathfrak{1}^n f(-\mathfrak{lx})$ . Note that the factor of  $\mathfrak{l}^n$  implies that  $\phi$  is *not* a linear transformation.

**Lemma 21.22.** *If* f *has positive leading coefficient then*  $f \in \mathcal{H}_1(\mathbb{C})$  *iff*  $\varphi(f) \in \mathfrak{I}$ *.* 

*Proof.* We may assume f is monic. All roots of f(-tx) lie in the lower half plane, and the leading coefficient is  $(-t)^n$ . Thus,  $\phi(f)$  is monic, and all roots are in the lower half plane.

Interlacing in  $\mathcal{H}_1(\mathbb{C})$  is different form **P**. Any definition of interlacing should satisfy  $f \leq f'$ , and this should mean that all linear combinations of f and f' are stable. However, this is false. If f = x + 1 then f - 2f' = x - 1 is not stable.

 $\mathfrak{I}$  is invariant under the transformation  $x \mapsto x + \alpha$ , where  $\alpha \in \mathbb{R}$ , but  $\mathfrak{H}_1(\mathbb{C})$  is preserved by  $x \mapsto x + \iota \alpha$ . In addition,  $\mathfrak{H}_1(\mathbb{C})$  is preserved by  $x \mapsto x + \alpha$  if  $\alpha > 0$ . Consequently, there are two new kinds of interlacing along with  $\xleftarrow{\mathsf{H}}$ .

Assume that f,  $g \in \mathcal{H}_1(\mathbb{C})$ 

$f \stackrel{\iota}{\sim} g$	iff	$\forall lpha \in \mathbb{R}$	$f + \alpha \iota g \in \mathfrak{H}_1(\mathbb{C})$
$f \overset{H}{\sim} g$	iff	$\forall \alpha \in \mathbb{R} > 0$	$f+\alphag\in {\mathcal H}_1({\mathbb C})$
$f \stackrel{H}{\longleftarrow} g$	iff	$\forall \alpha \in RHP$	$f+\alpha g\in {\mathcal H}_1({\mathbb C})$

We will see later that  $f \stackrel{\iota}{\sim} f'$  and  $f \stackrel{H}{\sim} f''$ . Taking limits shows that

 $\textbf{Lemma 21.23. } \textit{If } f,g \in \mathfrak{H}_1(\mathbb{C}) \textit{ then } f \xleftarrow{H} g \implies f \stackrel{\iota}{\sim} g \textit{ and } f \stackrel{H}{\sim} g$ 

*Remark* 21.24. It is not true that  $f \stackrel{H}{\sim} g \implies f \stackrel{\iota}{\sim} g$ . If  $f = x^2$  and g = 1 then clearly  $f \stackrel{H}{\sim} g$ , but  $x^2 + \iota$  has roots with positive and negative real parts.

The next result allows us to transfer interlacing results from  $\mathfrak{I}$  to  $\mathcal{H}_1(\mathbb{C})$ .

**Lemma 21.25.**  $f \xleftarrow{H} g$  in  $\mathcal{H}_1(\mathbb{C})$  if and only if  $\phi(f) \stackrel{\iota}{\sim} \phi(g)$  in  $\mathfrak{I}$ .

*Proof.* Since  $\phi$  is a bijection between  $\mathcal{H}_1(\mathbb{C})$  and  $\mathfrak{I}$ , the computation below establishes the lemma.

$$\begin{aligned} \varphi(f + \alpha \iota g) &= \iota^n (f(-\iota x) + \alpha \iota g(-\iota x)) \\ &= \varphi(f) - \alpha \iota^{n-1} g(-\iota x) \\ &= \varphi(f) - \alpha \varphi(g) \end{aligned}$$

Most properties of interlacing in **P** carry over to  $\stackrel{\iota}{\sim}$  in  $\mathcal{H}_1(\mathbb{C})$ , except for some occasional sign changes.

**Lemma 21.26.** *Assume* f, g,  $h \in \mathcal{H}_1(\mathbb{C})$ .

- 1. If  $f \stackrel{\iota}{\sim} g$  then  $fh \stackrel{\iota}{\sim} gh$ .
- 2. If  $\sigma \in RHP$  then  $(x + \sigma)f \stackrel{\iota}{\sim} f$ .
- 3.  $f \stackrel{\iota}{\sim} f'$ .
- 4. If  $f \in \mathbf{P}^{pos}$  then  $f(\iota \mathbf{D})g \in \mathcal{H}_1(\mathbb{C})$ .
- 5. If  $f \stackrel{\iota}{\sim} g$  and  $f \stackrel{\iota}{\sim} h$  then  $f \stackrel{\iota}{\sim} \alpha g + \beta h$  for positive  $\alpha, \beta$ . In particular,  $\alpha f + \beta g \in \mathfrak{H}_1(\mathbb{C})$ .
- 6. If  $f \stackrel{\iota}{\sim} g \stackrel{\iota}{\sim} h$  in  $\mathcal{H}_1(\mathbb{C})$  then  $f + h \stackrel{\iota}{\sim} g$ .
- 7.  $\mathbf{x} \mathbf{\sigma} \in \mathfrak{H}_1(\mathbb{C})$  iff  $\mathbf{\sigma} \in \mathbf{RHP}$ .
- 8.  $x \stackrel{\mathsf{H}}{\longleftarrow} 1$

Proof. The first is immediate. The second is a consequence of

$$(\mathbf{x} + \sigma)\mathbf{f} + \alpha \mathbf{i}\mathbf{f} = (\mathbf{x} + \sigma + \alpha \mathbf{i})\mathbf{f}$$

and  $x + \sigma + \alpha \iota \in \mathcal{H}_1(\mathbb{C})$ . The chain rule shows that  $\phi(f') = \phi(f)'$ . Since  $f \in \mathcal{H}_1(\mathbb{C})$  implies that  $\phi(f) \in \mathfrak{I}$ , we know that  $\phi(f) \leq \phi(f)' = \phi(f')$ .

We only need to check the next one for f = x + a where a > 0. We see that  $f(\iota D)g = \iota g' + ag$  which is in  $\mathcal{H}_1(\mathbb{C})$  by the previous part.

The next part follows from the corresponding result (Lemma 20.54) for  $\Im$ . We have interlacings

$$\mathfrak{l}^{n}\mathfrak{f}(\mathfrak{l} x) \underline{\lessdot} \mathfrak{l}^{n-1}\mathfrak{g}(\mathfrak{l} x) \underline{\lessdot} \mathfrak{l}^{n-2}\mathfrak{h}(\mathfrak{l} x)$$

in  $\mathfrak I$  where f has degree n. In  $\mathfrak I$  we have

$$\iota^{n}(f+h)(\iota x) = \iota^{n}f(\iota x) - \iota^{n-2}h(\iota x) \leq \iota^{n-1}g(\iota x)$$

and the result follows. The last two are trivial.

We have orthogonal type recurrences in  $\mathcal{H}_1(\mathbb{C})$ , except the last sign is reversed.

**Corollary 21.27.** *If*  $a_i, c_i > 0$ ,  $\Re(\beta_i) \ge 0$  *then define* 

$$\begin{split} p_{-1} &= 0 \\ p_{0} &= 1 \\ p_{n} &= (a_{n}x + \beta_{n}) \, p_{n-1} + c_{n} \, p_{n-2} \end{split}$$

All  $p_n$  are stable, and

$$\cdots p_3 \xleftarrow{H} p_2 \xleftarrow{H} p_1$$

*Proof.* Using induction, these follow from the last part of Lemma 21.26 and the interlacings

$$(x + \beta_n)p_{n-1} \xleftarrow{H} p_{n-1} \xleftarrow{H} p_{n-2}$$

**Lemma 21.28.** Suppose that  $f(x) = \prod (x - \sigma_k)$  is in  $\mathcal{H}_1(\mathbb{C})$ . If all  $\alpha_k$  are positive and

$$g(x) = \sum_{k} \alpha_{k} \frac{f(x)}{x - \sigma_{k}}$$
(21.4.1)

then  $g \in \mathfrak{H}_1(\mathbb{C})$  and  $f \xleftarrow{\mathsf{H}} g$ .

*Proof.* By additivity of interlacing for  $\mathcal{H}_1(\mathbb{C})$  it suffices to show that  $f \xleftarrow{H} f/(x - \sigma_k)$ , or  $(x - \sigma_k)h \xleftarrow{H} h$  where  $h = f/(x - \sigma_k)$ . This interlacing holds since  $x - \sigma_k \xleftarrow{H} 1$ .

Here are some lemmas that use the geometry of  $\mathbb{C}$  and are useful in proving that polynomials are stable. The first lemma gives useful information about the mappings determined by quotients.

**Lemma 21.29.** Assume that r, s > 0 and  $\sigma \in Q_I$ .

$$\begin{array}{ll} \displaystyle \frac{1}{r+\sigma} \in \mathcal{Q}_{IV} & \displaystyle \frac{1}{(r+\sigma)(s+\sigma)} \in \textit{lower half plane} \\ \displaystyle \frac{\sigma}{r+\sigma} \in \mathcal{Q}_{I} & \displaystyle \frac{\sigma}{(r+\sigma)(s+\sigma)} \in \textit{right half plane} \end{array}$$

*Proof.* Suppose  $\sigma = a + bi$  where a, b > 0. The left two follow from

$$\frac{1}{r+\sigma} = \frac{a+r-\imath b}{|r+\sigma|^2} \qquad \qquad \frac{\sigma}{r+\sigma} = \frac{a^2+b^2+ar+\imath br}{|r+\sigma|^2}$$

Since  $1/(r + \sigma)$  and  $1/(s + \sigma)$  are in the fourth quadrant, their product is in the lower half plane. Since  $\sigma/(r + \sigma)$  is in the first quadrant, and  $1/(s + \sigma)$  is in the fourth quadrant, their product is in the right half plane.

In **P** we know that f and f'' don't interlace, but  $f - \alpha f'' \in \mathbf{P}$  for all positive  $\alpha$ . There is a similar result for  $\mathcal{H}_1(\mathbb{C})$ .

**Lemma 21.30.** Suppose that  $f = \prod (x - \sigma_i)$  is in  $\mathcal{H}_1(\mathbb{C})$ . If all  $\alpha_{ij}$  are non-negative then

$$f \stackrel{H}{\sim} \sum_{i \neq j} \alpha_{ij} \frac{f}{(x - \sigma_i)(x - \sigma_j)}$$
(21.4.2)

*Proof.* If  $\tau$  is a root of the sum of the right and left hand sides of (21.4.2) then, after dividing by  $f(\tau)$ , we know that

$$1 + \sum_{i \neq j} \alpha_{ij} \frac{1}{(\tau - \sigma_i)(\tau - \sigma_j)} = 0$$

By Lemma 21.29 the sum lies in the lower half plane. Thus  $\tau$  does not satisfy (21.4.2), so  $\tau$  must lie in the open left half plane. Similarly for the lower right quadrant.

If we restrict all the coefficients to be equal, the lemma gives an alternate proof that that  $f \stackrel{H}{\sim} f''$ .

**Lemma 21.31.** *If*  $f \in \mathcal{H}_1(\mathbb{C})$ ,  $\mathfrak{R}(\alpha) \ge 0$ ,  $\beta \ge 0$  *then* 

$$\alpha f + \beta f' \in \mathcal{H}_1(\mathbb{C}).$$
  
 $\alpha f + \beta f'' \in \mathcal{H}_1(\mathbb{C})$ 

*Proof.* Assume that  $f = \prod (x - \sigma_i)$  where  $\Re \sigma_i < 0$ . Factoring out f(x), let  $\tau$  be a solution of

$$\alpha + \beta \sum \frac{1}{x - \sigma_i} = 0$$

If  $\Re(\tau) \ge 0$  then  $\Re 1/(\tau - \sigma_i) > 0$ , so the sum has positive real part, and can't equal  $-\alpha/\beta$ . The second follows from Lemma 21.30.

We know the following corollary holds more genearally in  $\mathcal{H}_d(\mathbb{C})$ , but in  $\mathcal{H}_1(\mathbb{C})$  we can prove it by induction.

**Corollary 21.32.** *If*  $f, g \in \mathcal{H}_1(\mathbb{C})$  *then*  $f(D)g \in \mathcal{H}_1(\mathbb{C})$ *.* 

*Proof.* Factoring f, it suffices to show that  $g + \alpha g' \in \mathcal{H}_1(\mathbb{C})$  for  $\mathfrak{R}(\alpha) > 0$ . This follows from the lemma.

Linear transformations on **P** induce linear transformations on  $\mathcal{H}_1(\mathbb{C})$ .

**Lemma 21.33.** Suppose  $T: P \longrightarrow P$  is a linear transformation that preserves degree, the sign of the leading coefficient, and strict interlacing. The linear transformation

$$\mathbf{x}^k \mapsto \mathbf{\iota}^k \, \mathsf{T}(\mathbf{x}^k)(-\mathbf{\iota} \, \mathbf{x})$$

*maps*  $\mathcal{H}_1(\mathbb{C})$  *to itself.* 

*Proof.* Since T induces a linear transformation in  $\Im$  (p. 600), the conclusion follows from the diagram

**Corollary 21.34.** The transformations  $x^k \mapsto \iota^k H_k(-\iota x)$  and  $x^k \mapsto \iota^k L_k(\iota x)$  map  $\mathcal{H}_1(\mathbb{C})$  to  $\mathcal{H}_1(\mathbb{C})$ , and  $\mathcal{H}_1$  to  $\mathcal{H}_1$ .

*Proof.* The maps  $x^n \mapsto H_n(x)$  and  $x^n \mapsto L_n(-x)$  satisfy the hypotheses of the lemma. It follows from the above that they map  $\mathcal{H}_1(\mathbb{C})$  to itself. Since  $\iota^k H_k(-\iota x)$  and  $\iota^k L_k(\iota x)$  are real polynomials, the maps also send  $\mathcal{H}_1$  to itself.

If all roots of two polynomials lie in given quadrants then we know the location of the roots of their Hadamard product. In the table below the notation  $-\mathcal{H}_1(\mathbb{C})$  means all roots in the right half plane, and  $-\mathfrak{I}$  means all roots in the upper half plane. Recall that  $\sum a_i x^i * \sum b_i x^i = \sum a_i b_i x^i$ .

**Lemma 21.35.** (*The location of* f \* g)

		Quadrant			
		1	2	3	4
	1	J	$-\mathcal{H}_1(\mathbb{C})$	$-\Im$	$\mathfrak{H}_1(\mathbb{C})$
	2	$-\mathcal{H}_1(\mathbb{C})$	$-\Im$	$\mathcal{H}_1(\mathbb{C})$	I
Quadrant	3	$-\Im$	$\mathcal{H}_1(\mathbb{C})$	I	$-\mathcal{H}_1(\mathbb{C})$
	4	$\mathcal{H}_1(\mathbb{C})$	J	$-\mathfrak{H}_1(\mathbb{C})$	$-\Im$

*Proof.* If f and g are in the third quadrant then they are in  $\mathfrak{I}^{pos}$ , and so their Hadamard product is in  $\mathfrak{I}$ . The Lemma now follows from the identity

$$f(\alpha x) * g(\beta x) = |f * g|(\alpha \beta x)$$

which holds for all complex  $\alpha$ ,  $\beta$ .

Note that this shows that the Hadamard product of stable polynomials with complex coefficients is not necessarily stable. For example

 $(x + 1 + \iota) * (x + 1 + 2\iota) = x - 1 + 3\iota.$ 

### 21.5 The analytic closure

We define the analytic closure of  $\mathcal{H}_1$  and  $\mathcal{H}_1(\mathbb{C})$  as usual:

$$\label{eq:Hamiltonian} \begin{split} \widehat{\mathcal{H}_d} = \ \text{the uniform closure of } \mathcal{H}_d \\ \widehat{\mathcal{H}_d}(\mathbb{C}) = \ \text{the uniform closure of } \mathcal{H}_d(\mathbb{C}) \end{split}$$

We can determine the exponentials that are in  $\mathcal{H}_1(\mathbb{C})$ ,

Lemma 21.36.

- 1.  $e^{\alpha x}$ ,  $e^{\alpha x^2}$  are in  $\widehat{\mathcal{H}}_1$  for all positive  $\alpha$ .
- 2.  $e^{\alpha \iota x}$  is in  $\mathcal{H}_1(\mathbb{C})$  for all  $\alpha \in \mathbb{R}$ .
- 3.  $\sinh(\alpha x)/x$  and  $\cosh(\alpha x)$  are in  $\widehat{\mathcal{H}_1}$  for  $\alpha \in \mathbb{R}$ .
- 4.  $e^{-x}$ ,  $e^{-x^2}$  are not in  $\mathcal{H}_1(\mathbb{C})$ .

Proof. Consider the limits

$$e^{\alpha x} = \lim_{n \to \infty} \left(1 + \frac{\alpha x}{n}\right)^n$$
$$e^{\alpha x^2} = \lim_{n \to \infty} \left(1 + \frac{\alpha (x + 1/n)^2}{n}\right)^n$$
$$e^{\alpha \iota x} = \lim_{n \to \infty} \left(1 + \frac{\alpha \iota x}{n}\right)^n$$

In the first two cases the part being exponentiated is in  $\mathcal{H}_1$ , and in  $\mathcal{H}_1(\mathbb{C})$  in the third, so the limits are as claimed. The infinite product formulas

$$\sinh(x)/x = \prod_{n=1}^{\infty} \left(1 + \frac{x^2}{n^2 \pi^2}\right) \qquad \cosh(x) = \prod_{n=1}^{\infty} \left(1 + \frac{4x^2}{(2n-1)^2 \pi^2}\right)$$

show that  $\sinh(x)/x$  and  $\cosh(x)$  are in  $\widehat{\mathcal{H}_1}$ .

To show non-existence we use Corollary 21.32. If  $e^{-x} \in \widehat{\mathcal{H}}_1$  then  $e^{-\mathsf{D}} \mathsf{f} \in \mathcal{H}_1$ , but this fails for  $\mathsf{f} = 2x + 1$ . If  $e^{-x^2} \in \widehat{\mathcal{H}}_1$  then  $e^{-\mathsf{D}^2} \mathsf{f} \in \mathcal{H}_1$ , and this fails for  $\mathsf{f} = (2x + 1)^2 + 1$ .

### 21.6 Real stable polynomials in one variable

The class  $\mathcal{H}_1$  shares many properties with  $\mathbf{P}^{pos}$ , as the first result shows.

#### Lemma 21.37.

- 1. If f, g have real coefficients then  $fg \in \mathcal{H}_1$  iff f and g are in  $\mathcal{H}_1$ .
- 2. If  $f \in \mathcal{H}_1$  then all coefficients of f have the same sign.
- 3. If  $f \in \mathcal{H}_1$  then  $f^{rev} \in \mathcal{H}_1$ .

*Proof.* The first is obvious. Factor monic  $f \in \mathcal{H}_1$  as

$$\prod(x-r_j) \cdot \prod(x-\sigma_k)(s-\overline{\sigma}_k)$$

where the  $r_j$  are negative, and the  $\sigma_k$  have negative real part. Expanding  $(x - \sigma_k)(s - \overline{\sigma}_k) = x^2 - 2\Re(\sigma_k)x + |\sigma_k|^2$  shows that all the factors have positive coefficients, and so the coefficients of f are all positive.

The roots of  $f^{rev}$  are the inverses of the roots of f. Since the roots of f lie in the open left half plane, so do the roots of  $f^{rev}$ .

The converse of 2 is false:

$$(x-1-2i)(x-1+2i)(x+2-i)(x+2+i) = x^4 + 2x^3 + 2x^2 + 10x + 25$$

There are simple conditions for the stability of polynomials of small degree.

Lemma 21.38. Assume all these polynomials have real coefficients.

- 1. A quadratic  $ax^2 + bx + c$  is stable if and only if all coefficients have the same sign.
- 2. A cubic  $x^3 + ax^2 + bx + c$  is stable if and only if all coefficients are positive and ab > c.
- 3. A quartic  $x^4 + ax^3 + bx^2 + cx + d$  is stable if and only if all coefficients are positive and  $abc > c^2 + a^2d$ .

*Proof.* We know all coefficients must be positive. The first one is trivial. For the remaining two, we convert to  $\Im$ , and the interlacing condition becomes the inequalities.

$$\begin{aligned} x^3 + ax^2 + bx + c & \text{ is stable iff} \\ x^3 - bx + \iota(ax^2 - c) & \text{ is in } \Im \text{ iff} \\ x^3 - bx &\leq ax^2 - c & \text{ iff} \\ \sqrt{b} > \sqrt{c/a} & \text{ which is equivalent to } ba > c \end{aligned}$$

$$\begin{aligned} x^4 + ax^3 + bx^2 + cx + d & \text{ is stable iff} \\ x^4 - bx^2 + d + \iota(ax^3 - cx) & \text{ is in } \Im \text{ iff} \\ x^4 - bx^2 + d &\leq ax^3 - cx & \text{ iff} \\ \frac{1}{2} \left( b - \sqrt{b^2 - 4d} \right) &\leq \frac{c}{a} &\leq \frac{1}{2} \left( b + \sqrt{b^2 - 4d} \right) & \text{ iff} \\ c^2 + a^2d &< abc \end{aligned}$$

In ove variable  $F_E$  is a sum of even powers, and  $f_O$  is a sum of odd powers so we can refine the previous result. Recall that  $f_e$  and  $f_o$  are the even and odd parts of f.

**Lemma 21.39.** Assume that f is a polynomial of degree n with all positive coefficients.

$$f \in \mathfrak{H}_1 \text{ if and only if } \begin{cases} f_e \leqslant f_o & n \text{ even} \\ f_e \ll f_o & n \text{ odd} \end{cases}$$

*Proof.* We first assume that the degree n of  $f\in {\mathcal H}_1$  is even. If we write  $f(x)=f_e(x^2)+xf_o(x^2)$  then

$$\phi(f) = \iota^n \left[ f_e(-x^2) - \iota x f_o(-x^2) \right]$$

Since n is even  $f_e(-x^2) \ll x f_o(-x^2)$  and hence  $f_e(x) \ll f_o(x)$ , which implies  $\psi(f) \in \mathfrak{I}^{pos}$ . The converse direction follows by reversing the argument.

If n = 2m + 1 then

$$\varphi(f)=(-1)^{\mathfrak{m}}\left(xf_{o}(-x^{2})+\iota f_{e}(-x^{2})\right)$$

which implies that  $xf_o(-x^2) \leqslant f_e(-x^2)$ , and hence  $f_e \ll f_o$ .

It follows from the lemma that if the degree of  $f \in \mathcal{H}_1$  is even then  $f_e + \iota f_o \in \mathfrak{I}$ . If the degree is odd, then  $(1 - \iota)(f_e + \iota f_o) \in \mathfrak{I}$ .

If  $\tau$  is a root of  $f \in \mathcal{H}_1$  then  $f/(x - \tau)$  is not necessarily in  $\mathcal{H}_1$ . If we pair the complex roots then we get interlacing:

**Lemma 21.40.** *Suppose that*  $f(x) \in \mathcal{H}_1$ *, and write* 

$$f(x) = \prod_{j=1}^{n} (x - r_j) \cdot \prod_{k=1}^{m} (x - \sigma_k)(s - \overline{\sigma}_k)$$

*If all*  $\alpha_k$ ,  $\beta_j$  *are non-negative and*  $\Re(\sigma_k) < 0$  *then* 

$$\mathsf{f} \xleftarrow{\mathsf{H}} \sum_{j=1}^n \beta_j \frac{\mathsf{f}(x)}{x-r_j} + \sum_{k=1}^m \alpha_k \frac{x-\mathfrak{R}(\sigma_k)}{(x-\sigma_k)(x-\overline{\sigma}_k)} \, \mathsf{f}(x)$$

Proof. The conclusion follows from Lemma 21.28 and the observation that

$$\frac{\mathbf{x} - \Re(\sigma_{k})}{(\mathbf{x} - \sigma_{k})(\mathbf{x} - \overline{\sigma}_{k})} = \frac{1}{2} \left( \frac{1}{\mathbf{x} - \sigma_{k}} + \frac{1}{\mathbf{x} - \overline{\sigma}_{k}} \right)$$

**Example 21.41.** It is not easy to determine all polynomials interlacing a given polynomial. For example, we will find all linear polynomials interlacing  $(x + 1)^2$  by translating to  $\mathfrak{I}$ . Assume that  $(x + 1)^2 \leq \sigma x + \tau$ . Then

$$(\mathbf{x}+1)^2 + lpha \mathbf{i}(\sigma \mathbf{x}+ au) \in \mathfrak{H}_1(\mathbb{C})$$

for all real  $\alpha$ . Since the sum of the roots is  $-2 - \alpha \iota \sigma$  and lies in the left half plane for all  $\alpha$  it follows that  $\Im(\sigma) = 0$ . After scaling, we may assume that

$$(\mathbf{x}+1)^2 \leq \mathbf{x}+\mathbf{a}+\mathbf{b}\mathbf{i}.$$

We now translate this interlacing to  $\Im$ :

$$x^2 - 1 + 2\iota x \lessdot x + a\iota - b$$

The determinant condition is

$$0 > \begin{vmatrix} x^2 - 1 & 2x \\ x - b & a \end{vmatrix} = (a - 2)x^2 + 2xb - a$$

Solving the equations yields

- 1.  $(x + 1)^2 \leq x + \tau$  in  $\mathcal{H}_1(\mathbb{C})$  if and only if  $\tau$  lies in a circle of radius one centered at 1.
- 2.  $(x+1)^2 \leq x+t$  in  $\mathcal{H}_1$  if and only 0 < t < 2.

If 0 < a < 1 then we can give an explicit polynomial in  $\mathcal{H}_2$  that shows that  $(x+1)^2 \lessdot x + a.$ 

$$\begin{vmatrix} y + x \left(\alpha^2 + \sqrt{\alpha^2 + 1}\alpha + 1\right) + 1 & \alpha \\ -\alpha & y + x \left(\alpha^2 - \sqrt{\alpha^2 + 1}\alpha + 1\right) + 1 \\ = (1 + \alpha^2)(1 + x)^2 + 2(1 + \alpha^2)\left(x + \frac{1}{1 + \alpha^2}\right)y + y^2 \end{vmatrix}$$

We have seen the first kind of region in  $\mathfrak{I}$ . Restricted to real polynomials, we get an interval in  $\mathfrak{H}_1$ , but just a single point in  $\mathbf{P}^{\text{pos}}$ .

A similar calculation shows the following:

$$(x+1)(x+2) \le (x+1) + a(x+2)$$
 in  $\mathcal{H}_1$  iff  $a > -1/2$ 

If  $f = \prod(x - w_i) \in \mathbf{P}$  then in Lemma 1.20 and Lemma 1.23 we gave conditions on constants  $c_i$  such that  $\sum c_i f/(x - w_i)$  is in  $\mathbf{P}$ . If  $f \in \mathbf{P}^{\text{pos}}$  we now ask what choice of constants determines a stable interlacing polynomial. The example above shows that some constants can be negative.

**Lemma 21.42.** Suppose that  $f = (x + r_1) \cdots (x + r_n)$  where  $0 < r_1 < \cdots < r_n$ . If

$$a_k + a_{k+1} + \cdots + a_n \ge 0$$
 for  $k=1,\ldots,n$ 

and at least one sum is positive, then  $f \xleftarrow{H} \sum a_k \frac{f}{x+r_k}$  in  $\mathfrak{H}_1.$ 

*Proof.* We need to verify that the quotient

$$\frac{1}{f}\sum a_k \frac{f}{x+r_k} = \sum \frac{a_k}{x+r_k}$$

maps  $\Re(\sigma) > 0$  to the right half plane. Let  $\sigma = \alpha + \beta \iota$  with  $\alpha > 0$ . Then,

$$\begin{split} \mathfrak{R} & \sum \frac{a_{k}}{\sigma + r_{k}} = \mathfrak{R} \sum a_{k} \frac{r_{k} + \alpha - \beta \iota}{(r_{k} + \alpha)^{2} + \beta^{2}} \\ & = \sum a_{k} \frac{r_{k} + \alpha}{(r_{k} + \alpha)^{2} + \beta^{2}} \\ & \geqslant \frac{1}{(r_{n} + \alpha)^{2} + \beta^{2}} \sum a_{k}(\alpha + r_{k}) \end{split}$$

This is positive since

$$\sum a_k(\alpha + r_k) = (\alpha + r_1)(a_1 + \dots + a_n) + (r_2 - r_1)(a_2 + \dots + a_n) + (r_3 - r_2)(a_3 + \dots + a_n) + \dots + (r_n - r_{n-1})a_n > 0$$

by hypothesis.

**Example 21.43.** If  $0 < r_1 < r_2$  and we choose  $a_1 + a_2 > 0$ ,  $a_2 > 0$  then  $f = (x + r_1)(x + r_2) < a_1(x + r_2) + a_2(x + r_1)$  in  $\mathcal{H}_1$ . Scaling so that  $a_1 + a_2 = 1$ , this is  $f \leq x + r_1 + (r_2 - r_1)a_1$ . Since  $1 \geq a_1$ , it follows that

$$(x + r_1)(x + r_2) \le x + t \text{ in } \mathcal{H}_1 \text{ if } 0 < t < r_2$$

We can determine all  $t \in \mathbb{R}$  for which  $(x + r_1)(x + r_2) \leq x + t$  in  $\mathcal{H}_1$ . Converting to  $\Im$  yields  $x^2 + (r_1 + r_2)x\mathbf{i} - r_1r_2 \leq x + \mathbf{i}t$ . The interlacing conditions are satisfied, and the determinant condition yields

$$(x + r_1)(x + r_2) \le x + t \text{ in } \mathcal{H}_1 \text{ iff } 0 < t < r_1 + r_2$$

Although the example shows that the hypotheses of the lemma do not determine all interlacing polynomials, there is a sense in which they are the best possible.

**Lemma 21.44.** If  $a_1, \ldots, a_n$  have the property that for all  $0 < r_1 < \cdots < r_n$  we have

$$f \lessdot \sum_{k=1}^n \alpha_k \frac{f}{x+r_k} \text{ in } \mathcal{H}_1$$

where  $f = (x + r_1) \cdots (x + r_n)$  then

$$a_k + a_{k+1} + \dots + a_n \ge 0$$
 for  $k=1,\dots,n$ 

*Proof.* Assume that  $a_k + \cdots + a_n < 0$ . Take  $r = r_1 = \cdots = r_{k-1}$  and  $s = r_k = \cdots = r_n$ . Then the real part of the quotient, where  $\sigma = \alpha + \beta \iota$ , is equal to

$$\begin{split} &\sum_{i=1}^{k-1} a_i \frac{r+\alpha}{|\sigma+r|^2} + \sum_{i=k}^n a_i \frac{s+\alpha}{|\sigma+s|^2} = \\ &\frac{r+\alpha}{|\sigma+r|^2} \sum_{i=1}^{k-1} a_i + \frac{s+\alpha}{|\sigma+s|^2} \sum_{i=k}^n a_i \end{split}$$

Since the second sum is negative this expression will be negative if we can choose constants so that

$$\frac{r+\alpha}{|\sigma+r|^2} \bigg/ \frac{s+\alpha}{|\sigma+s|^2} < \epsilon$$

for appropriately small positive  $\epsilon$ . This is the case if we pick  $|\sigma| \gg s \gg r \gg \alpha$ .

The next lemma follows from the corresponding facts for  $\mathcal{H}_1(\mathbb{C})$ .

**Lemma 21.45.** If f,  $g \in \mathcal{H}_1$  then  $f(D)g \in \mathcal{H}_1$ . If  $f \in P^{\text{pos}}$  then  $f(D^2)g \in \mathcal{H}_1$ .

Combining the positive and negative cases, we get

**Corollary 21.46.** *If*  $f \in P^{pos}$  *then* 

$$f + \alpha f'' \in \begin{cases} P & \alpha \leqslant 0 \\ \mathfrak{H}_1 & \alpha > 0 \end{cases}$$

We describe some interactions among  $\mathbf{P}^{pos}$ ,  $\mathcal{H}_1$ ,  $\mathcal{H}_1(\mathbb{C})$  and  $\mathfrak{I}$ .

**Lemma 21.47.** *If*  $F \in \mathfrak{I}$  *has all coefficients in the first quadrant then*  $F \in \mathcal{H}_1(\mathbb{C})$ *.* 

*Proof.* Write  $F = f + \iota g$ . If  $f = \prod (x - r_k)$  and  $g/f = \sum \frac{a_k}{x - r_k}$  where all  $a_k$  are positive then  $(f + \iota g)(\sigma) = 0$  implies

$$1 + \iota \sum \frac{a_i}{|\sigma - r_i|^2} (\overline{\sigma} - r_i) = 0$$

It follows that the real part of  $\sigma$  lies in the interval  $(\min r_k, \max r_k)$ . If all the coefficients are positive then all roots of f are negative, so the real parts of  $\sigma$  are negative, and  $F \in \mathcal{H}_1(\mathbb{C})$ .
The intersection of  $\mathcal{H}_1(\mathbb{C})$  and  $\mathfrak{I}$  contains polynomials that do not have all positive coefficients. If  $f + \iota g$  is

$$(x - .08)(x + .7)(x + 1)(x + 1.7)(x + 2) + \iota(x + .3)(x + .9)(x + 1.7)(x + 1.8)$$

then the roots are all are in the third quadrant

$$(-1.9 - 0.1\iota, -1.7 - 0.003\iota, -0.9 - 0.04\iota, -0.5 - 0.5\iota, -0.2 - 0.2\iota)$$

**Lemma 21.48.** Assume that  $f, g \in \mathbf{P}^{pos}$ . If  $f \xleftarrow{P} g$  in  $\mathbf{P}$  then  $f \xleftarrow{H} g$  in  $\mathcal{H}_1$ .

*Proof.* If 
$$f \xleftarrow{P} g$$
 then  $f + yg \in \mathbf{P}_2^{\text{pos}} \subset \mathcal{H}_2$ , so  $f \xleftarrow{H} g$ .

The converse is false.

*Remark* 21.49. The image of  $\mathcal{H}_1$  in  $\mathfrak{I}$  is easily described. If  $f = \sum^n a_k x^k$  then

$$\Phi(f) = \iota^{n} \sum_{k \equiv n}^{n} x^{k} a_{k} (-\iota x)^{k}$$
  
= 
$$\sum_{k \equiv n \pmod{2}} x^{k} a^{k} (-1)^{(n-k)/2} + \iota \sum_{k \not\equiv n \pmod{2}} x^{k} a^{k} (-1)^{(n-k-1)/2}$$

Thus the real part consists of the terms of even degree with alternating signs and the odd part consists of terms of odd degree with alternating signs, or vice versa. This is equivalent to  $\overline{g(x)} = g(-x)$ . Since  $\phi$  is just a rotation, the roots of  $\phi(f)$  are all in the lower half plane, and are symmetric around the imaginary axis. Note that the image of  $\mathcal{H}_1$  isn't  $\mathfrak{I}^{\text{pos}}$ .

**Lemma 21.50.** If  $f \in P^{pos}$ , and  $T(x^i) = x^{i-1}$  for  $i \ge 1$  and T(1) = 0 then  $T: P^{pos} \longrightarrow \mathcal{H}_1$ .

*Proof.* Note that T(f) = (f(x) - f(0))/x. If  $f = \prod(x + a_i)$  where all  $a_i$  are positive then  $\alpha$  is a root of T(f) implies that  $f(\alpha) = f(0)$ , and therefore

$$\prod \left(\frac{\mathbf{x}}{\mathbf{a}_{i}}+1\right)=1$$

If x has positive real part then  $x/a_i$  also has positive real part, so  $|1+x/a_i| > 1$ . Consequently the product above can't be 1, and so no roots are in the right half plane.

## **21.7** Coefficient inequalities in $\mathcal{H}_1$

Newton's inequalities do not hold for  $\mathcal{H}_1$ . This can be seen easily in the quadratic case. If  $f = a_0 + a_1x + a_2x^2 \in \mathcal{H}_1$  then

$$\frac{a_1^2}{a_0 a_2} \quad \begin{cases} \ge 4 & \text{if the roots are real} \\ \le 4 & \text{if the roots are complex} \end{cases}$$

Moreover, the quotient can take on any value in  $(0, \infty)$ .

However, there are many inequalities involving the coefficients. For instance, if  $f = \sum a_i x^i \in \mathcal{H}_1$  then we know that

$$a_0 - a_2 x + \cdots < a_1 - a_3 x + \cdots$$

and therefore  $| {a_1 \ a_2 \ a_2} | > 0$  which is equivalent to

$$\frac{a_1 a_2}{a_0 a_3} > 1 \tag{21.7.1}$$

A much more general result is that the Hurwitz matrix is totally positive.(See p. 649.) This means that

/a1	a <sub>3</sub>	$a_5$		)
a <sub>0</sub>	$a_2$	$a_4$		
0	$\mathfrak{a}_1$	$a_3$	$a_5$	
0	$a_0$	$a_2$	$\mathfrak{a}_4$	
(:	÷	÷	÷	·)

is totally positive. For example, if we choose  $f \in \mathcal{H}_1(4)$  then

$$0 \leqslant \begin{vmatrix} a_1 & a_3 & 0 \\ a_0 & a_2 & a_4 \\ 0 & a_1 & a_3 \end{vmatrix} = a_1 a_2 a_3 - a_1^2 a_4 - a_3^2 a_0$$

This inequality is the necessary and sufficient condition (Lemma 21.38) for a positive quartic to be stable.

Here's a simple yet surprising conclusion about initial segments of stable polynomials.

**Lemma 21.51.** *If*  $\sum a_i x^i \in P^{pos}$  *then* 

$$\sum_{k=0}^1 a_i x^i \stackrel{+}{\sim} \sum_{k=0}^2 a_i x^i \stackrel{+}{\sim} \sum_{k=0}^3 a_i x^i \stackrel{+}{\sim} \sum_{k=0}^4 a_i x^i \quad \text{in } \mathcal{H}_1$$

In particular, each of these partial sums is stable.

*Proof.* We begin with the middle interlacing; the first is trivial. We need to verify that

 $\alpha(a_0+a_1x+a_2x^2)+(a_0+a_1x+a_2x^2+a_3x^3)=(1+\alpha)(a_0+a_1x+a_2x^2)+a_3x^3$ 

is stable. From (21.7.1) we know that  $a_1a_2 > a_0a_3$ , so

$$(1 + \alpha)a_1(1 + \alpha)a_2 > (1 + \alpha)a_0a_3$$

and therefore we have positive interlacing. For the last interlacing, we need to show that

$$(1 + \alpha)(a_0 + a_1x + a_2x^2 + a_3x^3) + a_4x^4$$

is stable for positive  $\alpha$ . To do this we need to show that

$$(1+\alpha)^3a_1a_2a_3>a_4(1+\alpha)^2a_1^2+(1+\alpha)^2a_3^2a_0$$

The Newton inequalities give

$$a_1^2 \geqslant \frac{8}{3}a_0a_2 \qquad \qquad a_2^2 \geqslant \frac{9}{4}a_1a_3 \qquad \qquad a_3^2 \geqslant \frac{8}{3}a_2a_4$$

and as a consequence

$$a_2a_3 \ge 6a_1a_4 \qquad \qquad a_1a_2 \ge 6a_0a_3$$

and so

$$\begin{aligned} &(1+\alpha)^3 a_1 a_2 a_3 \geqslant 6(1+\alpha)^2 a_1^2 a_4 \\ &(1+\alpha)^3 a_1 a_2 a_3 \geqslant 6(1+\alpha)^2 a_3^2 a_0 \end{aligned}$$

Addition yields the desired result.

Note that the first two only require that f is stable, but the last interlacing uses  $f \in \mathbf{P}^{pos}$ .

*Remark* 21.52. It is not true that the higher partial sums are stable. For  $k \ge 5$  the partial sum  $1 + x + \cdots + x^k/k!$  of the exponential series is not stable [187]. The sum of the first k+1 coefficients of  $(1+x/n)^n$  converges to  $1+x+\cdots+x^k/k!$ , and  $(1+x/n)^n$  is in  $\mathbf{P}^{\mathrm{pos}}$ . Thus, for n sufficiently large the partial sums aren't stable. For instance, the sum of the first six terms of  $(1+x/n)^n$  is stable for for  $n \le 23$  and not stable for n > 23.

## 21.8 Positive interlacing

We study positive interlacings  $f \stackrel{H}{\sim} g$ . If f, g in **P**, then one way of showing  $f \leq g$  is sign interlacing. This can be viewed as describing the behavior of the map

$$\frac{\mathsf{g}}{\mathsf{f}}\colon \mathbb{R}\longrightarrow \mathbb{R}\setminus\infty.$$

In order to prove that  $f \stackrel{H}{\sim} g$  in  $\mathcal{H}_1$ , we look at the image of the first quadrant rather than the real line. The next lemma gives criteria for  $\stackrel{+}{\sim}$  and  $\stackrel{H}{\sim}$ . The proof is omitted.

Lemma 21.53.

- 1.  $f \stackrel{+}{\sim} g iff \frac{f}{g} \colon Q_{I} \longrightarrow RHP.$
- 2.  $f \stackrel{\mathsf{H}}{\sim} g \operatorname{iff} \frac{f}{g} \colon Q_{\mathrm{I}} \longrightarrow \mathbb{C} \setminus (-\infty, 0).$

If  $f,g\in P$  then the implication  $f\underline{\lessdot} g\implies f\overset{\mathsf{H}}{\sim} g$  is trivial. It's also easy in  $\mathcal{H}_1.$ 

**Lemma 21.54.** If f, 
$$g \in \mathcal{H}_1$$
 and  $f \leq g$  implies  $f \stackrel{\sqcap}{\sim} g$ 

*Proof.* We know that the image of the closed right plane under  $x \mapsto \frac{f}{g}(x)$  lies in the open right half plane. Thus,  $f \stackrel{H}{\sim} g$ .

We have seen (See p. 54.) that if  $f \stackrel{H}{\sim} g$ , h in  $\mathbf{P}^{pos}$  then g + h might not have all real roots. However,

**Lemma 21.55.** If  $f \stackrel{H}{\sim} g$ , h are in  $P^{pos}$  then g + h is stable.

*Proof.* From Lemma 21.53 we know that evaluation of g/f and h/f at a point in the closed first quadrant gives a point in the right half plane, and so their sum lies there too.

**Lemma 21.56.** If  $f \xleftarrow{H} g$  then  $f \stackrel{H}{\sim} x g$ 

*Proof.* We assume  $f, g \in \mathcal{H}_1$ ; the case  $f, g \in \mathcal{H}_1(\mathbb{C})$  is similar. Since  $f \triangleleft g$  we know that  $f/g: \overline{\mathbf{Q}_I} \longrightarrow \mathsf{RHP}$ . To show  $f \stackrel{\mathsf{H}}{\sim} xg$  we choose  $\alpha \ge 0$  and prove that  $f + \alpha xg \in \mathcal{H}_1$ . We do this by establishing that  $f + \alpha xg \stackrel{\mathsf{H}}{\sim} f$ . If  $\sigma \in \overline{\mathbf{Q}_I}$  then

$$\frac{f + \alpha x g}{f}(\sigma) = 1 + \alpha \sigma \frac{f(\sigma)}{g(\sigma)}$$

Now  $(g/f)(\sigma) \in \mathsf{RHP}$  since f < g and  $\alpha \sigma \in \overline{\mathsf{RHP}}$  so  $\frac{f + \alpha x g}{f}(\sigma)$  is in  $\mathsf{RHP} + \overline{\mathsf{RHP}} \times \mathsf{RHP} = \mathbb{C} \setminus (0, \infty)$ .

Next is a Leibniz-like rule for second derivatives.

**Lemma 21.57.** *If*  $f, g \in P^{pos}$  *then*  $fg'' + f''g \in \mathcal{H}_1$ .

*Proof.* Recall Lemma 21.53. If we evaluate f/f'' and g/g'' at a point in the closed first quadrant then we get a point in the lower half plane. Thus f/f'' + g/g'' is stable.

Note that fg'' + f''g might not lie in  $\mathbf{P}^{pos}$ . For example, if  $f = (x + 1)^2$  and  $g = (x + 2)^2$  then fg'' + f''g is always positive, and so has no real zeros. Positive interlacings can be multiplied.

**Lemma 21.58.** If  $f \leq g$  and  $h \leq k$  in  $P^{pos}$ 

1. fh  $\stackrel{\text{H}}{\sim}$  gk *in*  $\mathcal{H}_1$ .

- 2. fk  $\stackrel{\text{H}}{\sim}$  gh *in*  $\mathcal{H}_1$ .
- 3.  $f^2 \stackrel{H}{\sim} g^2$  in  $\mathcal{H}_1$ .

*Proof.* We know that f/g and h/k both map the first quadrant to itself, so (fh)/(gk) maps the first quadrant to the upper half plane, so fh  $\stackrel{H}{\sim}$  gk. The last is a special case of the first. Since k/h maps Q<sub>I</sub> to Q<sub>IV</sub>, we see that (fh)/(gk) maps to Q<sub>IV</sub>  $\cup$  Q<sub>I</sub>  $\cup$  Q<sub>II</sub>  $\setminus$  0, from which the second statement follows.

**Corollary 21.59.** *If*  $a_i$ ,  $b_i$ ,  $c_i > 0$ , *then the sequence defined by* 

$$p_{-1} = 0$$
  

$$p_{0} = 1$$
  

$$p_{n} = (a_{n}x + b_{n}) p_{n-1} + c_{n} p_{n-2}$$

then all  $p_n$  are stable, and

$$\dots p_3 \stackrel{H}{\sim} p_2 \stackrel{H}{\sim} p_1$$

*Proof.* From Corollary 21.27 we know that  $p_n \leq p_{n-1}$ , and so  $p_n \stackrel{H}{\sim} p_{n-1}$  by Lemma 21.54.

The preceding lemma can be considered as a construction of stable orthogonal polynomials. There is a simple case where the roots are all known explicitly [80]. If  $f_1 = 1$ ,  $f_2 = x$  and

$$f_{n+1}(x) = x f_n(x) + f_{n-1}(x)$$

then the polynomials are known as the Fibonacci polynomials and have purely imaginary roots:

$$2\iota \cos(k\pi/n)$$
,  $1 \leq k \leq n-1$ .

*Remark* 21.60. If we take two stable polynomials

$$f = x^4 + 6x^3 + 14x^2 + 10x + 4$$
  $g = x^2 + 2x + 2$ 

then how do we go about showing that  $f \stackrel{H}{\sim} g$ ? If we use the definition then we need to show that  $f + t g \in \mathcal{H}_1$  for all positive t. Since this is a quartic we could apply the criterion (Lemma 21.38) for a quartic to be stable. A more general technique uses the fact (Lemma 21.53) that f/g must map the closed first quadrant to  $\mathbb{C} \setminus (-\infty, 0]$ . If  $\sigma \in \overline{Q_I}$  then this would follow from  $\Im(f(\sigma)/g(\sigma)) > 0$ . To show this we just need to show that the polynomial  $\Im(f(a + b\iota)g(a - b\iota))$  is positive for  $a, b \ge 0$ . If we are lucky, this polynomial will have no minus signs, as in this case:

$$f(a + bi)g(a - bi) = 2b(a^2 + b^2)^3 \times (a^5 + 6a^4 + 2b^2a^3 + 16a^3 + 8b^2a^2 + 27a^2 + b^4a + 8b^2a + 24a + 2b^4 + 3b^2 + 6)$$

This approach doesn't quite work here, since a = b = 0 has imaginary part 0. However, f(0)/g(0) = 2 which is not on the negative axis, so we conclude that  $f \stackrel{H}{\sim} g$ .

The Bezout polynomial is stable.

**Lemma 21.61.** The Bezout polynomial  $\frac{1}{x-y} \begin{vmatrix} f(x) & f(y) \\ g(x) & g(y) \end{vmatrix}$  is stable if  $f \xleftarrow{P} g \in P^{pos}$ .

Proof. Recall (9.21.3)

$$B(x,y) = \sum a_i \frac{f(x)}{x - r_i} \frac{f(y)}{y - r_i}$$
$$\frac{B(x,y)}{f(x)f(y)} = \sum a_i \frac{1}{x - r_i} \frac{1}{y - r_i}$$

If  $f, g \in \mathbf{P}^{pos}$ ,  $\sigma \in Q_1, \tau \in Q_1$  then  $\frac{1}{\sigma - r_i} \frac{1}{\tau - r_i}$  is in the lower half plane, and so the sum misses  $(-\infty, 0)$ . If  $\sigma \in Q_1, \tau \in Q_4$  then  $\frac{1}{\sigma - r_i} \frac{1}{\tau - r_i}$  is in the right half plane, and so the sum again misses  $(-\infty, 0)$ . The remaining cases are similar, so we conclude  $f(x)f(y) \stackrel{H}{\sim} B(x, y)$ .

When a polynomial is always positive then it is not necessarily stable. Here's a familiar case where it is.

**Lemma 21.62.** If f < g in  $P^{pos}$  then  $\begin{vmatrix} f & g \\ f' & g' \end{vmatrix}$  is stable. If f has all distinct roots then  $\begin{vmatrix} f & f' \\ f' & g' \end{vmatrix}$  is stable.

*Proof.* If we write  $f = \prod (x + r_i)$  and  $g = \sum a_i f/(x + r_i)$  then

$$f'g-fg'=f^2\,\sum\frac{a_i}{(x+r_i)^2}$$

and Lemma 21.53 applies. Use f < f' for the second one.

**Example 21.63.** Here's a nice example of the lemma. Assume that  $f_0, f_1, f_2, ...$  is an orthogonal polynomial sequence with strict interlacing:  $f_{k+1} < f_k$  for k = 1, ... The Christoffel-Darboux formula [168] states that

$$f_0(x)f_0(y) + f_1(x)f_1(y) + \dots + f_n(x)f_n(y) = \frac{k_n}{k_{n+1}} \frac{1}{x-y} \begin{vmatrix} f_n(y) & f_{n+1}(y) \\ f_n(x) & f_{n+1}(x) \end{vmatrix}$$

for positive constants  $k_n$ . If we let x = y then

$$f_0(x)^2 + f_1(x)^2 + \dots + f_n(x)^2 = \frac{k_n}{k_{n+1}} \begin{vmatrix} f_n(x) & f_{n+1}(x) \\ f'_n(x) & f'_{n+1}(x) \end{vmatrix}$$

It follows from the Lemma that

$$f_0(x)^2 + f_1(x)^2 + \dots + f_n(x)^2$$
 is stable

We can also state Lemma 21.62 in terms of Wronskians: if f < g in  $\mathbf{P}^{pos}$  then the Wronskian W(f,g) = fg' - f'g is stable.

More generally we have

**Lemma 21.64.** The square determined by  $|I + xD_1 + yD_2 + zD_3|$ , where all  $D_i$  are positive definite,  $D_1$  is diagonal, and  $D_2$ ,  $D_3$  have all positive entries, yields a stable polynomial.

*Proof.* The assumptions mean that we are to show that  $\begin{vmatrix} f & g \\ h & k \end{vmatrix}$  is stable, where

$$f + gy + hz + kyz + \cdots = |I + xD_1 + yD_2 + zD_3|$$

and all matrices are positive definite. Assume that  $D_1$  is diagonal  $(r_i)$ ,  $D_2 = (a_{ij})$ ,  $D_3 = (b_{ij})$ . We have

$$\begin{split} f &= \prod (1+r_{i}x) \\ g &= \sum_{i} a_{ii}f/(1+r_{i}x) \\ h &= \sum_{j} b_{jj}f/(1+r_{j}x) \\ k &= \sum_{i \neq j} \begin{vmatrix} a_{ii} & b_{ij} \\ a_{ij} & b_{jj} \end{vmatrix} \frac{f}{(1+r_{i}x)(1+r_{j}x)} \\ gh &- fk = f^{2} \left( \sum_{i} \frac{a_{ii}b_{ii}}{(1+r_{i}x)^{2}} + \sum_{i \neq j} \frac{a_{ij}b_{ij}}{(1+r_{i}x)(1+r_{j}x)} \right) \end{split}$$

Since D<sub>1</sub>, D<sub>2</sub>, D<sub>3</sub> are positive definite, their diagonals are positive, so  $(gh - fk)/f^2$  lies in the lower half plane by Lemma 21.29. Lemma 21.53 implies that  $f^2 \stackrel{H}{\sim} gh - fk$ , and therefore gh - fk is stable.

**Corollary 21.65.** *If* f, g, h, k *are as above, and*  $0 < \alpha \leq 1$  *then*  $\begin{vmatrix} \alpha f & g \\ h & k \end{vmatrix}$  *is stable. Proof.* The above computation shows that

$$gh-\alpha fk=gh-fk+(1-\alpha)fk=f^2\left(\sum \frac{a_ib_i}{(1+r_ix)^2}+(1-\alpha)\frac{k}{f}\right)$$

From Lemma 21.53 we know that if we evaluate at a point in the first quadrant then all the terms lie in the lower half plane.  $\Box$ 

We can slightly improve Lemma 21.62.

**Corollary 21.66.** If  $f \in \mathbf{P}^{\text{pos}}$  and  $0 < \alpha \leq 1$  then  $\begin{vmatrix} \alpha f & f' \\ f' & f'' \end{vmatrix}$  is stable.

Proof. Applying the calculations of the lemma to

$$f(x + y + z) = f(x) + f'(x)y + f'(x)z + f''(x)yz/2 + \dots \in \mathbf{P}_{3}^{\text{pos}}$$

shows that  $(\alpha/2)ff'' - f'f'$  is stable.

Newton's inequality says that a certain determinant is positive. If we introduce a second variable, this becomes a statement that a polynomial is stable. The first part follows from Lemma 21.64.

**Lemma 21.67.** If  $f = \sum f_i(x)y^i$  is in  $\mathbf{P}_2^{pos}$  then

$f_0^2$	$\stackrel{\rm H}{\sim}$	$f_1^2 - f_0 f_2$	in $\mathcal{H}_1$
$f_0 f_1$	н ~	$f_1^2 - f_0 f_2$	in $\mathcal{H}_1$
$f_0^2$	$\stackrel{\rm H}{\sim}$	$x(f_1^2-f_0f_2)$	in $\mathcal{H}_1$
$f_0 f_1$	$\stackrel{\rm H}{\sim}$	$x(f_1^2-f_0f_2)$	in $\mathcal{H}_1$

In particular,  $f_1^2 - f_0 f_2$  is (weakly) stable, and all coefficients are non-negative.

*Proof.* We compute  $f_1^2 - f_0 f_2$  using the determinant representation for f. So, assume that  $f = |I + xD_1 + yD_2|$ . We may assume that  $D_1$  is diagonal,  $D_1 = diag(r_i)$ , and that  $D_2 = (d_{ij})$ . Then

$$\begin{split} f_0(x) &= \prod_i (1+r_i x) \\ f_1(x) &= \sum_i d_{ii} \frac{f_0}{1+r_i x} \\ f_2(x) &= \sum_{i < j} \left| \begin{array}{c} d_{ii} & d_{ij} \\ d_{ij} & d_{jj} \end{array} \right| \frac{f_0}{(1+r_i x)(1+r_j x)} \\ f_1^2 - f_0 f_2 &= \sum_i d_{ii}^2 \frac{f_0^2}{(1+xr_i)^2} + \sum_{i < j} (d_{ij}^2 + d_{ii} d_{jj}) \frac{f_0^2}{(1+xr_i)(1+xr_j)} \\ \frac{f_1^2 - f_0 f_2}{f_0^2} &= \sum_i \frac{d_{ii}^2}{(1+xr_i)^2} + \sum_{i < j} \frac{(d_{ij}^2 + d_{ii} d_{jj})}{(1+xr_i)(1+xr_j)} \end{split}$$

Now the diagonal entries of a positive definite matrix are positive, so all the coefficients of the expansion of  $f_1^2 - f_0 f_2$  are positive. Consequently, evaluation at a point in the closed first quadrant yields a point in the open lower half plane, which establishes the first part.

For the second interlacing we have

$$\frac{f_1^2 - f_0 f_2}{f_0 f_1} = \frac{f_0}{f_1} \left( \sum_{i} \frac{d_{ii}^2}{(1 + xr_i)^2} + \sum_{i < j} \frac{(d_{ij}^2 + d_{ii} d_{jj})}{(1 + xr_i)(1 + xr_j)} \right)$$

When we evaluate a point in the closed first quadrant,  $f_0/f_1$  is in the first quadrant, and the part in the parentheses is in the lower half plane. Consequently, their product lies in an open half plane P (the rotation of the lower half plane by the argument of the point) which misses the negative real axis.

Since  $f_1^2 - f_0 f_2$  is stable, all of its coefficients have the same sign. If we let x = 0 then this is positive, by Newton's inequality for f(0, y). Since the constant term is positive, all terms are positive.

When we introduce the factor of x, then the sum in parentheses for  $x(f_1^2 - f_0f_2)/f_0^2$  lies in the right half plane by lemma 21.29. The case where the denominator is  $f_0f_1$  yields a point in the product of the first quadrant and right half plane, and so again misses the negative real axis.

If we choose  $\beta > 0$  then  $f_0(\beta)$ ,  $f_1(\beta)$ ,  $f_2(\beta)$  are the first three terms of a polynomial in  $\mathbf{P}^{\text{pos}}$ , so we know that  $f_1(\beta)^2 > 2f_0(\beta) f_2(\beta)$ . This suggests that we can improve the result above.

**Corollary 21.68.** If  $0 < \alpha < 2$  and f,  $f_i$  are as above then

1. 
$$f_1^2 - \alpha f_0 f_2$$
 is stable.  
2.  $\begin{vmatrix} f_{k+1} & f_{k+2} \\ f_k & f_{k+1} \end{vmatrix}$  is stable.

*Proof.* For the second, we differentiate k times and apply the first part. Using the notation of the lemma, the first part follows from the identity

$$\begin{split} \frac{f_1^2 - \alpha f_0 f_2}{f_0^2} = \\ & \sum_i \frac{d_{ii}^2}{(1 + x r_i)^2} + \alpha \sum_{i < j} \frac{d_{ij}^2}{(1 + r_i x)(1 + r_j x)} + (2 - \alpha) \sum_{i < j} \frac{d_{ii} d_{jj}}{(1 + r_i x)(1 + r_j x)} \end{split}$$

We can use the lemma to construct stable polynomials from polynomials in  $\mathbf{P}^{\text{pos}}$ .

**Corollary 21.69.** If  $f \in P^{pos}$  and a, b, c > 0 then  $af^2 + bff' + cx(ff'' - (f')^2)$  is stable.

*Proof.* From Lemma 21.67 we know that each of the three terms positive interlace  $f^2$ , so any positive combination is stable

The next lemma arises from considerations involving the Euler polynomials. Unlike the last lemma, we must look carefully at the terms of the quotient.

**Lemma 21.70.** *If*  $f \in P^{pos}(n)$  *then* 

$$Q = x \left( nf^2 + (x-1)^2 \left[ ff' + x(ff'' - f'^2) \right] \right) \in \mathcal{H}_1$$

*Proof.* We show that Q and f<sup>2</sup> interlace positively. First,

$$Q/f^2 = x \left( n + (x - 1)(f'/f) + x [ff'' - f'^2]/f^2 \right).$$

If  $f = \prod (x + r_i)$  where all  $r_i$  are positive then

$$\frac{f'}{f} = \sum_{i=1}^{n} \frac{1}{x + r_i} \qquad \qquad \frac{f'^2 - ff''}{f^2} = \sum_{i=1}^{n} \frac{1}{(x + r_i)^2}$$

Thus the quotient satisfies

$$\begin{split} Q/f^2 &= x \bigg( n + (x-1) \sum \frac{1}{x+r_i} - x \sum \frac{1}{(x+r_i)^2} \bigg) \\ &= x \sum \bigg( 1 + \frac{x-1}{x+r_i} - \frac{x}{(x+r_i)^2} \bigg) \\ &= \sum \frac{xr_i(1+r_i+x)}{(x+r_i)^2} \end{split}$$

If we substitute  $\alpha + \beta \iota$  for x then

$$\begin{aligned} \frac{xr_{i}(1+r_{i}+x)}{(x+r_{i})^{2}} = \\ \frac{r_{i}\left(\beta^{4}+\left(2\alpha^{2}+3r_{i}\alpha+\alpha+r_{i}(r_{i}+2)\right)\beta^{2}+\alpha(r_{i}+\alpha)^{2}(r_{i}+\alpha+1)\right)}{|r_{i}+\alpha|^{2}} \end{aligned}$$

Thus, the sum is in the first quadrant if  $\alpha + \beta \iota$  is in the first quadrant.

## 21.9 Linear transformations

Linear transformations preserve linearity, so the following is immediate.

**Lemma 21.71.** If  $T: \mathcal{H}_1 \longrightarrow \mathcal{H}_1$  preserves degree and the sign of the leading coefficient then T preserves interlacing and positive interlacing.

**Lemma 21.72.** *If*  $f, g \in \mathcal{H}_1$  *then*  $f * g \in \mathcal{H}_1$ .

*Proof.* If  $f \in \mathbf{P}^{pos}$  then the method of generating functions shows that the Hadamard product maps  $\mathbf{P}^{pos} \times \mathcal{H}_1 \longrightarrow \mathcal{H}_1$ . However, we need a different approach when  $f \notin \mathbf{P}^{pos}$ . There does not appear to be a direct way to do this. We follow [66]: convert to  $U_1$ , take Hadamard products there, and use the fact that  $f \leq g$  and  $h \leq k$  implies  $f * h \leq g * k$ .

**Lemma 21.73.** If T:  $x^n \longrightarrow \langle \underline{x} \rangle_n$  then T maps  $\mathcal{H}_1$  to  $\mathcal{H}_1$  and  $\mathcal{H}_1(\mathbb{C})$  to  $\mathcal{H}_1(\mathbb{C})$ .

*Proof.* It suffices to show that T maps  $\mathcal{H}_1(\mathbb{C})$  to itself; we proceed by induction. So assume that  $f \in \mathcal{H}_1(\mathbb{C})$  and  $T(f) \in \mathcal{H}_1(\mathbb{C})$ . If  $a \in \mathsf{RHP}$  then will show that  $T(x+a)f \stackrel{H}{\sim} T(f)$  which implies that  $T(x+a)f \in \mathcal{H}_1(\mathbb{C})$ . From (7.7.1) we know that

$$T(x + a)f = (x + a)T(f) + T(xf').$$

If  $\sigma \in \overline{\mathsf{Q}_{\mathrm{I}}}$  then

$$\frac{\Gamma(\mathbf{x} + \mathbf{a})f}{\Gamma(f)}(\sigma) = (\sigma + \mathbf{a}) + \frac{T(\mathbf{x}f')}{T(f)}(\sigma)$$

Now  $T(xf') \stackrel{H}{\sim} T(f)$  so  $\frac{T(xf')}{T(f)}(\sigma)$  is in  $\mathbb{C} \setminus (0, \infty)$ . Thus  $\frac{T(x+a)f}{T(f)}(\sigma)$  is in  $\mathbb{C} \setminus (0, \infty)$  and the lemma is proved.

**Lemma 21.74.** If  $T: x^n \longrightarrow (\underline{x})_n$  then  $T^{-1}$  maps  $\mathfrak{H}_1$  to  $\mathfrak{H}_1$  and  $\mathfrak{H}_1(\mathbb{C})$  to  $\mathfrak{H}_1(\mathbb{C})$ .

*Proof.* We proceed as above. Let  $S = T^{-1}$ . From (7.6.1) we have

$$S(x + a)f = (x + a) S(f) + x(S(f))$$

To show  $S(x+\alpha)f \stackrel{H}{\sim} S(f)$  we choose  $\sigma \in \overline{\mathsf{Q}_{\mathrm{I}}}$  and then

$$\frac{S(x+a)f}{S(f)}(\sigma) = (\sigma+a) + \sigma \frac{(S(f))'}{S(f)}(\sigma)$$
(21.9.1)

Since  $S(f) \in \mathcal{H}_1(\mathbb{C})$  by induction and S(f) < (S(f))' we have  $(S(f))'/S(f)(\sigma) \in RHP$  and therefor (21.9.1)  $\in RHP + RHP \times \overline{RHP} = \mathbb{C} \setminus (0, \infty)$ .

**Lemma 21.75.** If 
$$T: x^n \mapsto \begin{cases} x^{n/2} & n \text{ even} \\ 0 & n \text{ odd} \end{cases}$$
 then  $T: \mathcal{H}_1 \longrightarrow P^{\text{pos}}$ .

*Proof.* If  $f \in \mathcal{H}_1$  has degree n and we write  $f(x) = f_e(x^2) + xf_o(x^2)$  then

$$\mathfrak{l}^{\mathfrak{n}}(\mathfrak{f}_{\mathfrak{e}}(-\mathbf{x}^2) + (-\mathfrak{l}\mathbf{x})\mathfrak{f}_{\mathfrak{o}}(-\mathbf{x}^2)) \in \mathfrak{I}$$

Thus,  $f_e$  and  $f_o$  are in **P**, and so  $T(f) = f_e$  since all coefficients are positive.  $\Box$ 

The Charlier polynomials are defined by

$$C_{-1}^{\alpha} = 0 \qquad C_0^{\alpha} = 1$$
  
$$C_{n+1}^{\alpha}(x) = (x - n - \alpha)C_n^{\alpha}(x) - \alpha nC_{n-1}^{\alpha}(x)$$

**Lemma 21.76.** If  $\alpha \in RHP$  and  $T: x^n \mapsto C_n^{\alpha}$  then  $T^{-1}$  maps  $\mathfrak{H}_1(\mathbb{C})$  to  $\mathfrak{H}_1(\mathbb{C})$ , and  $\mathfrak{H}_1$  to  $\mathfrak{H}_1$ .

*Proof.* We proceed as with the rising and falling factorials. We show by induction that for  $a \in \mathsf{RHP}$  we have that  $T^{-1}(x+a)f \stackrel{H}{\sim} T^{-1}(f)$ . From the recurrence

$$T^{-1}(x + a)f = (x + \alpha + a)T^{-1}f + (x + \alpha)(T^{-1}(f))$$

we choose  $\sigma \in \mathsf{RHP}$  and consider the quotient

$$\frac{\mathsf{T}^{-1}(\mathsf{x}+\mathfrak{a})\mathsf{f}}{\mathsf{T}^{-1}(\mathsf{f})}(\sigma) = (\sigma+\alpha+\mathfrak{a}) + (\sigma+\mathfrak{a})\frac{\left(\mathsf{T}^{-1}(\mathsf{f})\right)'}{\mathsf{T}^{-1}(\mathsf{f})}(\sigma)$$

We see that the quotient lies in  $\mathsf{RHP} + \mathsf{RHP} \times \mathsf{RHP} \subset C \setminus (0, \infty)$  since  $\mathfrak{a}, \alpha$  and  $\sigma$  are all in  $\mathsf{RHP}$ .

If we have two mutually interlacing sequences then the sum of products in reverse order is in  $\mathbf{P}$  (See p. 66.). If we use the original order we get a stable polynomial.

**Lemma 21.77.** Suppose that  $f_1, f_2, \dots, f_n$  and  $g_1, g_2, \dots, g_n$  are two sequences of mutually interlacing polynomials in  $\mathbf{P}^{pos}$ . Then

$$f_1g_1 + f_2g_2 + \dots + f_ng_n \in \mathcal{H}_1$$

*Proof.* If  $1 \le k \le n$  then  $f_1 \le f_k$  and  $g_1 \le g_k$ . If  $\sigma$  is in the closed first quadrant then  $\frac{f_k}{f_1}(\sigma)$  and  $\frac{g_k}{g_1}(\sigma)$  are in the open fourth quadrant. Thus  $\frac{f_k g_k}{f_1 g_1}(\sigma)$  are in the open lower half plane. Addition shows that

$$\frac{1}{f_1g_1}\sum_{k=1}^n f_kg_k \quad \text{is in the open lower half plane}$$

which finishes the proof.

We can apply this lemma to integrate families of interlacing interlacing polynomials. We also view these results as linear transformations on  $\mathbf{P}^{sep}$ .

**Lemma 21.78.** If f, 
$$g \in \mathbf{P}^{sep}$$
 then  $\int_0^1 f(x+t)g(x+t) dt$  is stable.

*Proof.* The proof is the same as Lemma 8.60, except we use the preceding lemma instead of Lemma 3.16.  $\Box$ 

**Corollary 21.79.** If  $f \in \mathbf{P}^{sep}$  then  $\int_0^1 e^t f(x+t) dt$  is stable.

*Proof.* Apply the lemma to  $e^x$  and f, and then factor out  $e^x$  from the result.  $\Box$ 

**Example 21.80.** If we take  $f \in \mathbf{P}^{sep}$  then we see that  $\int_0^1 f(x+t)^2 dt$  is stable. In particular,

$$\int_0^1 \left( \langle \underline{\mathbf{x}} \rangle_n \right)^2 d\mathbf{t} \quad \text{is stable.}$$

**Lemma 21.81.** If  $f \in \mathfrak{H}_1$  then  $\int_0^1 f(x+t) dt \in \mathfrak{H}_1$ .

*Proof.* Suppose deg(f) = n, let  $g(x) = \iota^n f(-\iota x)$ , and  $h(x) = \int_0^1 f(x+t) dt$ . Since  $g(x) \in \Im$  we have from Lemma 20.59

$$\int_0^1 g(x+\iota t)\,dt\in\mathfrak{I}\qquad\text{ and therefore }\qquad\int_0^1\iota^nf(-\iota x+t)\,dt\in\mathfrak{I}.$$

The last integral is  $\iota^n h(-\iota x)$ , so  $h \in \mathcal{H}_1$ .

It is surprising that we do not need to make any other assumptions about f in order that the difference is stable.

**Corollary 21.82.** If  $f \in \mathcal{H}_1$  then  $\Delta(f) = f(x+1) - f(x) \in \mathcal{H}_1$ .

*Proof.* Since  $f' \in \mathcal{H}_1$  we have

$$f(x+1)-f(x)=\int_0^1 f(x+t)'\,dt\in \mathfrak{H}_1.$$

**Example 21.83.** Here's a simple example. We know that  $x^n \in \mathcal{H}_1$ , and the roots of  $\Delta x^n = (x+1)^n - x^n$  are

$$-\frac{1}{2} - \iota \frac{\sin \frac{2\pi k}{n}}{-2 + 2\cos \frac{2\pi k}{n}} \qquad k = 1, \dots, n$$

They all lie in the open left half plane, so  $\Delta x^n \in \mathcal{H}_1$ .

# 21.10 The totally stable matrix conjecture

Recall that if  $f(x) = \sum \alpha_i x^i$  is in  $I\!\!P^{\text{pos}}$  then the matrix

$$\begin{pmatrix} a_0 & a_1 & a_2 & a_3 & a_4 & a_5 & a_6 & \dots \\ 0 & a_0 & a_1 & a_2 & a_3 & a_4 & a_5 & \dots \\ 0 & 0 & a_0 & a_1 & a_2 & a_3 & a_4 & \dots \\ 0 & 0 & 0 & a_0 & a_1 & a_2 & a_3 & \dots \\ \vdots & \vdots & & & \ddots & \ddots & \ddots \end{pmatrix}$$
(21.10.1)

is totally positive. This means that all minors are non-negative. Now, assume that  $f(x, y) = \sum f_i(x)y^i$  and form the matrix

$$\varphi(f)(\mathbf{x},\mathbf{y}) = \begin{pmatrix} f_0 & f_1 & f_2 & f_3 & f_4 & f_5 & f_6 & \dots \\ 0 & f_0 & f_1 & f_2 & f_3 & f_4 & f_5 & \dots \\ 0 & 0 & f_0 & f_1 & f_2 & f_3 & f_4 & \dots \\ 0 & 0 & 0 & f_0 & f_1 & f_2 & f_3 & \dots \\ \vdots & \vdots & & \ddots & \ddots & \ddots \end{pmatrix}.$$
(21.10.2)

If  $f \in \mathbf{P}_2^{\text{pos}}$  and  $\alpha > 0$  we know that  $f(\alpha, y) \in \mathbf{P}^{\text{pos}}$ , so all minors of  $\varphi(f)(\alpha, y)$  are positive. Now, whenever we have a polynomial that is positive for x > 0 we should consider whether it is stable. Empirical evidence suggests the following:

**Conjecture 21.84** (The totally stable matrix conjecture). *If*  $f(x, y) \in \mathbf{P}_2^{pos}$  *then all minors of the matrix* (21.10.2) *are stable.* 

Here's a simple consequence:

**Lemma 21.85.** If  $f \in \mathbf{P}_2^{\text{pos}}$  and all minors of  $\phi(f)$  are stable, then all minors have non-negative coefficients.

*Proof.* If g(x) is a minor then since g is stable we know that all coefficients have the same sign. If we set  $x = \alpha \ge 0$  then the matrix  $\varphi(f(\alpha, y))$  is just the matrix of a one variable polynomial in  $\mathbf{P}^{\text{pos}}$  and we know that all of its minors are non-negative. Thus,  $g(\alpha)$  is non-negative for all positive  $\alpha$ . Since all coefficients are the same sign they are positive.

**Example 21.86.** For a simple example, consider  $\varphi(1 + x + y)^4 =$ 

$$\begin{pmatrix} (1+x)^4 & 4(1+x)^3 & 6(1+x)^2 & 4(1+x) & 1 & 0 & \dots \\ 0 & (1+x)^4 & 4(1+x)^3 & 6(1+x)^2 & 4(1+x) & 1 & \dots \\ 0 & 0 & (1+x)^4 & 4(1+x)^3 & 6(1+x)^2 & 4(1+x) & \dots \\ 0 & 0 & 0 & (1+x)^4 & 4(1+x)^3 & 6(1+x)^2 & \dots \\ \vdots & \vdots & & \ddots & \ddots & \ddots \end{pmatrix}.$$

Every minor is a power of (1 + x) times a minor of the coefficients, which we know to be positive, so all minors are stable.

**Example 21.87.** Lemma 21.67 shows that the 2 by 2 determinants  $\begin{vmatrix} f_k & f_{k+1} \\ f_{k+1} & f_{k+2} \end{vmatrix}$  are stable.

**Example 21.88.** If  $f(x) \in \mathbf{P}^{pos}$  then  $f(x + y) \in \mathbf{P}_2^{pos}$ , and the entries of the matrix come from the Taylor series.

$$\phi(f(x+a)) = \begin{pmatrix} f & f^{(1)} & f^{(2)}/2! & f^{(3)}/3! & f^{(4)}/4! & f^{(5)}/5! & f^{(6)}/6! & \dots \\ 0 & f & f^{(1)}/1! & f^{(2)}/2! & f^{(3)}/3! & f^{(4)}/4! & f^{(5)}/5! & \dots \\ 0 & 0 & f & f^{(1)}/1! & f^{(2)}/2! & f^{(3)}/3! & f^{(4)}/4! & \dots \\ 0 & 0 & 0 & f & f^{(1)}/1! & f^{(2)}/2! & f^{(3)}/3! & \dots \\ \vdots & \vdots & & \ddots & \ddots & \ddots \end{pmatrix}.$$

We know that some of the two by two determinants are stable. For example,

$$\begin{vmatrix} f^{(3)}/3! & f^{(4)}/4! \\ f^{(4)}/4! & f^{(5)}/5! \end{vmatrix} = \frac{1}{4!4!} \left(\frac{4}{5}gg'' - g'g'\right)$$

where  $g = f^{(3)}$ . Since 4/5 < 1 Corollary 21.66 applies to show that it is stable.

*Remark* 21.89. If  $f = \sum a_i x^i$  is a stable polynomial then  $f_e \leq f_o$ . Proposition 3.22 therefore implies that

$$\begin{pmatrix} a_0 & a_2 & a_4 & \cdots & & \\ 0 & a_1 & a_3 & a_5 & \cdots & \\ 0 & 0 & a_0 & a_2 & a_4 & \cdots & \\ 0 & 0 & 0 & a_1 & a_3 & a_5 & \cdots & \\ & & \ddots & \ddots & \ddots & \end{pmatrix}$$

is totally positive. This matrix is known as the Hurwitz matrix [7].

Earlier (See p. 300.) we saw that the matrix of coefficients of a polynomial in  $\mathbf{P}_2^{\text{pos}}$  is totally positive<sub>2</sub>, but there are examples where some three by three determinants are negative. However, there is a surprising conjecture that is not fazed by this fact.

**Conjecture 21.90.** If  $D_1$ ,  $D_2$ ,  $D_3$  are positive definite matrices and M is the matrix of coefficients of y, z of  $|I + x D_1 + y D_2 + z D_3|$  then all minors of M are stable polynomials in x.

For example, if we use the same example with a particular choice for  $D_1$ 

$$M = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} + x \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} + y \begin{pmatrix} 13 & 9 & 7 \\ 9 & 7 & 5 \\ 7 & 5 & 4 \end{pmatrix} + z \begin{pmatrix} 5 & 7 & 8 \\ 7 & 11 & 12 \\ 8 & 12 & 14 \end{pmatrix}$$

then the coefficient matrix is a 4 by 4 matrix, and the non-constant 3 by 3 minors are

$$\begin{pmatrix} -1620x^3 - 7560x^2 - 4884x + 1760 & -1080x^2 - 3512x - 4544 & -768x - 4096 \\ -1080x^2 - 3496x - 3376 & -720x - 1312 \\ -744x - 2728 \end{pmatrix}$$

All entries are stable polynomials. Note that if we substitute zero for x we get the example on page 300.

## 21.11 Determinants of Hankel matrices

If  $p_1, p_2, \ldots$  is a sequence of polynomials then the d by d Hankel matrix (See p. 294.) of this sequence is

$$H({p_i}; d) = \begin{pmatrix} p_1 & p_2 & p_3 & \dots & p_d \\ p_2 & p_3 & p_4 & \dots & p_{d+1} \\ p_3 & p_4 & p_5 & \dots & p_{d+2} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ p_d & p_{d+1} & p_{d+2} & \dots & p_{2d} \end{pmatrix}$$

In this section we study properties of Hankel matrices. This includes the kind of polynomial (all real roots, stable, etc.), the sign of the coefficients, and interlacing properties.

A polynomial sequence  $\{p_n(x)\}_{n \ge 0}$  is called q*-log-concave* if all coefficients of

$$p_{n} p_{n+2} - p_{n+1}^{2} = \begin{vmatrix} p_{n} & p_{n+1} \\ p_{n+1} & p_{n+2} \end{vmatrix} = H[\{p_{n}, p_{n+1}, p_{n+2}\}; 2]$$

are positive for  $n \ge 0$ . If all coefficients are negative it is q-*log-convex* [118]. A stronger condition is that the polynomial  $p_{n+1}^2 - p_n p_{n+2}$  is stable. For instance, Lemma 21.67 shows that the coefficients of a polynomial in  $\mathbf{P}_2^{\text{pos}}$  are q-log-convex.

**Lemma 21.91.** If  $B_n$  is the Bell polynomial then  $B_{n+2}B_n - B_{n+1}^2$  is weakly stable with positive leading coefficient. In particular, it is q-log-convex.

*Proof.* The Bell polynomials satisfy the recurrence  $B_{n+1} = x(B_n + B'_n)$ , and therefore

$$B_{n+2}B_n - B_{n+1}^2 = x \left( B_n^2 + B_n B_n' + x (B_n B_n'' - B_n'^2) \right)$$

Since  $B_n \in \overline{\mathbf{P}^{pos}}$  it follows from Corollary 21.69 that this is in  $\mathcal{H}_1$ .

In addition, we have shown that  $B_n^2 \stackrel{H}{\sim} B_n B_{n+2} - B_{n+1}^2$ .

**Lemma 21.92.** If  $A_n$  is the Euler polynomial then  $A_{n+2}A_n - A_{n+1}^2$  is weakly stable with positive coefficients.

Proof. The Euler polynomial satisfies the recurrence

$$A_{n+1} = x((n+1)A_n + (1-x)A'_n)$$

and so substitution yields

$$A_{n+2}A_n - A_{n+1}^2 = x \left( (1+n)A_n^2 + (x-1)^2 \left[ A_n A_n' + x (A_n A_n'' - (A_n')^2 \right] \right)$$

This sum is the same as the sum in Lemma 21.70, except that it has n + 1 in place of n. This only adds a point in the first quadrant to the quotient, so we conclude that  $A_{n+2}A_n - A_{n+1}^2$  is stable. The value at 0 is known to be positive [118], so all terms are positive. Since  $A_n$  has 0 for a root, we only get weakly stable.

The following general result is easy, since everything factors. The work is finding the constant.

**Lemma 21.93.** If n and d are positive then  $|H[\{\langle \underline{x} \rangle_n, \langle \underline{x} \rangle_{n+1}, \dots \}; d]|$  is in **P**.

*Proof.* We first prove the result for d = 3. If n = 1 and d = 3 then the matrix is

$$\begin{pmatrix} x & x(x+1) & x(x+1)(x+2) \\ x(x+1) & x(x+1)(x+2) & x(x+1)(x+2)(x+3) \\ x(x+1)(x+2) & x(x+1)(x+2)(x+3) & x(x+1)(x+2)(x+3)(x+4) \end{pmatrix}$$

with determinant

$$2x^{3}(x+1)^{2}(x+2)$$

If we let  $f_n(x)$  be the determinant of  $H[\{\langle \underline{x} \rangle_n, \langle \underline{x} \rangle_{n+1}, \ldots\}; 3]$  then it is easy to see that

$$\begin{split} f_n(x) &= \langle \underline{x} \rangle_n \langle \underline{x} \rangle_{n+1} \langle \underline{x} \rangle_{n+2} \times \\ & \begin{vmatrix} 1 & 1 & 1 \\ x+n & x+n+1 & x+n+2 \\ (x+n)(x+n+1) & (x+n+1)(x+n+2) & (x+n+2)(x+n+3) \end{vmatrix} \\ &= 2 \langle \underline{x} \rangle_n \langle \underline{x} \rangle_{n+1} \langle \underline{x} \rangle_{n+2} \end{split}$$

In general, the answer is

$$(d-1)!(d-2)!\cdots 1!$$
  $\langle \underline{x} \rangle_n \langle \underline{x} \rangle_{n+1} \cdots \langle \underline{x} \rangle_{n+d-1}$ 

It's easy to find the polynomial factors. If we remove them and replace x + n by y we are left with

$$\begin{vmatrix} 1 & \dots & 1 \\ \langle \underline{y} \rangle_1 & & \langle \underline{y} + d - 1 \rangle_1 \\ \vdots & & \vdots \\ \langle \underline{y} \rangle_{d-1} & \dots & \langle \underline{y} + d - 1 \rangle_{d-1} \end{vmatrix}$$

From [108] we see that this equals the Vandermonde determinant

$$\begin{vmatrix} 1 & \dots & 1 \\ y & y+d-1 \\ \vdots & \vdots \\ y^d & \dots & (y+d-1)^{d-1} \end{vmatrix}$$

which equals

$$\prod_{0 < i < j < d} \left( [y+j] - [y+i] \right) = \prod_{0 < i < j < d} (j-i)$$

and this is precisely  $(d - 1)! \cdots 1!$ .

**Example 21.94.** The Chebyshev T polynomials have trivial Hankel determinants. First of all,

$$\begin{vmatrix} T_{n} & T_{n+1} \\ T_{n+1} & T_{n+2} \end{vmatrix} = \begin{vmatrix} T_{n} & T_{n+1} \\ 2xT_{n} - T_{n-1} & 2xT_{n+1} - T_{n} \end{vmatrix} = \begin{vmatrix} T_{n-1} & T_{n} \\ T_{n} & T_{n+1} \end{vmatrix}$$

Continuing, we find they are all equal to  $x^2 - 1$ . Thus, all the Hankel matrices  $H[\{T_n\}; 2]$  are equal to  $x^2 - 1$ , so they all have all real roots.

Since  $T_{n+1} = 2xT_n - T_{n-1}$  has coefficients that do not depend on n, all the higher Hankel determinants are zero.

We end this section with some empirical results about the Hankel determinants of the form  $H(\{p_n\}; d]$ . "cpx" means roots in all quadrants, and  $\checkmark$  means it has been proved.

Family	d	Kind of polynomial	Sign of coefficients	
Bell	2	stable $\checkmark$	positive $\checkmark$	
	≥3	cpx	positive	
Euler	2	stable $\checkmark$	positive $\checkmark$	
	≥3	cpx	positive	
Narayana	2	stable	positive	
	3	stable	positive	
	≥3	cpx	positive	
Laguerre	2	stable	negative	
	3	stable	negative	
	4	stable	positive	
	n	stable	$(-1)^{\binom{n}{2}}$	

# 21.12 Constructing stable polynomials

We construct matrices whose determinant is a stable polynomial. Recall that a matrix A is called *skew-symmetric* if  $A^T = -A$ . If A is skew-symmetric then

- 1. The eigenvalues of A are purely imaginary.
- 2. If D is positive definite, then the eigenvalues of A D have negative real part.

Using the latter property, and the fact that  $f(x)\in {\mathcal H}_1$  if and only if  $\iota^nf(-\iota x)\in {\mathfrak I}(n)$  we have we have

**Lemma 21.95.** If  $D_1, \ldots$  are positive definite, S is symmetric, and A is skew-symmetric then

- 1.  $|I + x_1 D_1 + \dots + x_d D_d + \iota S| \in \mathcal{H}_1(\mathbb{C})$
- 2.  $|\mathbf{I} + \mathbf{x} \mathbf{D}_1 + \dots + \mathbf{x}_d \mathbf{D}_d + \mathbf{A}| \in \mathcal{H}_d$

For example, the determinant of

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + x \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix} + y \begin{pmatrix} 3 & 1 \\ 1 & 2 \end{pmatrix} + \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

is in  $\mathcal{H}_2$  and equals

$$2x^2 + 8yx + 3x + 5y^2 + 5y + 2$$
.

*Remark* 21.96. It's easy to construct a determinant representation  $\alpha |I + xD + A|$  of  $f \in \mathcal{H}_1$ , where  $\alpha > 0$ . Factoring f, it suffices to show this for polynomials of degree one and two - the general case follows by taking direct sums.

$$\begin{aligned} \mathbf{x} + \mathbf{a} &= \mathbf{a} \left| (1) + \mathbf{x} \left( 1/a \right) + (0) \right| \\ (\mathbf{x} + \mathbf{a})^2 + \mathbf{b}^2 &= \mathbf{a}^2 \left| \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \mathbf{x} \begin{pmatrix} 1/a & 0 \\ 0 & 1/a \end{pmatrix} + \begin{pmatrix} 0 & \mathbf{b}/a \\ -\mathbf{b}/a & 0 \end{pmatrix} \right| \end{aligned}$$

**Example 21.97.** A special case of the matrix xI + D + A is the Jacobi-like matrix

(	$x + a_1$	$\mathfrak{b}_1$	0	0	0	0	)
	$-b_1$	$x + a_1$	<b>s</b> <sub>1</sub>	0	0	0	]
	0	$-s_1$	$x + a_2$	$\mathfrak{b}_2$	0	0	
	0	0	$-b_2$	$x + a_2$	s <sub>2</sub>	0	
	0	0	0	$-s_2$	$x + a_3$	$b_3$	
	0	0	0	0	$-b_3$	$x + a_3$	
	:	:	:	:		•.	. ]
	:	:	:	:		•	••• /

whose determinant is stable. The determinant of the submatrix consisting of the first n rows and columns is an n'th degree polynomial  $p_n$ . The recurrence relations for  $p_n$  are

$$p_{2k} = (x + a_k) p_{2k-1} + b_k^2 p_{2k-2}$$
$$p_{2k+1} = (x + a_{k+1}) p_{2k} + s_k^2 p_{2k-1}$$

If all the  $a_i$  are positive then we know from the Lemma above that all  $p_k$  are stable ; this also follows from the recurrence of Corollary 21.27. Moreover, the corollary shows that they interlace.

If a polynomial f(x) is in  $\mathbf{P}^{pos}$  then all its roots are negative, so  $f(x^2)$  has purely imaginary roots. There is a similar result for  $\mathbf{P}_2^{pos}$ .

**Lemma 21.98.** If  $f(x, y) \in \mathbf{P}_2^{pos}$  then  $f(x, x^2) \in \mathcal{H}_1$ .

*Proof.* Since f has real coefficients it suffices to show that  $f(\sigma, \sigma^2) \neq 0$  where  $\sigma$  is in the first quadrant. If so, then  $\sigma^2$  is in the upper half plane, and hence  $f(\sigma, \sigma^2) \neq 0$  since  $f \in U_2$ .

Alternatively, this is a consequence of the next lemma.

**Lemma 21.99.** If  $D_1$  and  $D_2$  are positive definite then  $|I + xD_1 + x^2D_2|$  is stable.

*Proof.* If  $\sigma = a + bi$  is in the first quadrant then

 $|I + \sigma D_1 + \sigma^2 D_2| = |(I + aD_1 + (a^2 - b^2)D_2) + \iota(bD_1 + 2abD_2)|$ 

The coefficient of *i* is positive definite, so the determinant is non-zero.

#### 21.13 Stable matrix polynomials

A hyperbolic matrix polynomial M is determined by the properties of the family of polynomials  $v^t M v$ . If we posit that all the quadratic forms are stable we get stable matrix polynomials.

**Definition 21.100.** An n by n matrix polynomial M is *stable* if  $v^t M v$  is stable for all non-zero  $v \in \mathbb{R}^n$ , and the leading coefficient is positive definite.

Most properties of hyperbolic matrix polynomials carry over to stable matrix polynomials. For instance, we define

The following lemma is easily proved using properties of stable polynomials.

**Lemma 21.101.** *Suppose that* f, g, h *are stable matrix polynomials, and*  $k \in \mathcal{H}_1$ .

- 1. All elements of  $Hyp_1^{pos}$  are stable matrix polynomials.
- 2.  $f \leftarrow H g$  iff  $f + \sigma g$  is a stable matrix polynomial for all  $\sigma \in RHP$ .
- 3. If  $f \xleftarrow{H} g$  and  $f \xleftarrow{H} h$  then  $f \xleftarrow{H} g + h$ . In particular, g + h is a stable matrix polynomial.
- 4. If  $f \xleftarrow{H} g \xleftarrow{H} h$  then  $f + h \xleftarrow{H} g$ .
- 5. fk is a stable matrix polynomial.
- 6. If all  $a_i$  are positive then  $(a_1x_1 + \cdots a_dx_d + b)f \xleftarrow{H} f$ .
- 7. If the i-th diagonal element of f is  $f_i$ , and of g is  $g_i$  then  $f \leq g$  implies  $f_i \leq g_i$ .
- 8.  $\frac{df}{dx}$  is a stable matrix polynomial.
- 9. f  $\leftarrow \frac{H}{dx}$
- 10. If  $f \xleftarrow{H} g$  then  $\frac{df}{dx} \xleftarrow{H} \frac{dg}{dx}$

We can construct stable matrix polynomials using positive interlacing. We start with the data

positive real numbers
$$a_1, b_1, \dots, a_n, b_n$$
positive definite matrices $m_1, \dots, m_n$ 

We claim that

$$M(x) = \sum_{k=1}^{n} m_k \prod_{i \neq k} (x + a_i)^2 + b_i^2$$

is a stable matrix polynomial. In order to verify this we first define

$$g(x) = \prod_i (x + a_i)^2 + b_i^2$$

and we know that

$$g(x) \xleftarrow{H} \frac{g(x)}{(x+a_k)^2 + b_k^2} = \prod_{i \neq k} (x+a_i)^2 + b_i^2.$$

Next,

$$u^{t} M \nu = \sum_{k} (\nu^{t} m_{k} \nu) \frac{g(x)}{(x + a_{k})^{2} + b_{k}^{2}}$$

Since  $v^t m_k v$  is positive, it follows that  $g \stackrel{H}{\leftarrow} v^t M v$  since each of the terms positively interlaces g(x), and positive interlacing is closed under positive linear combinations.

More generally, we can also construct stable matrix polynomials using common interlacing.

**Lemma 21.102.** If  $f_1, \ldots, f_r$  have a common interlacing in  $\mathcal{H}_1$  and  $m_1, \ldots, m_r$  are positive definite matrices then

 $f_1 m_1 + \cdots + f_r m_r$  is a stable matrix polynomial.

*Proof.* The leading coefficient is a positive linear combination of positive definite matrices and so is positive definite. If  $g \stackrel{H}{\leftarrow} f_i$  then

$$v^{t} g v = (v^{t} v)g \xleftarrow{H} (v^{t} m_{i} v)f_{i} = v^{t} f_{i} m_{i} v$$

and therefore

$$v^{t} g v \xleftarrow{\mathsf{H}} v^{t} (f_{1}m_{1} + \dots + f_{r}m_{r}) v.$$

A quadratic polynomial is stable if all its coefficients are positive. The analogous result holds for matrix polynomials.

**Lemma 21.103.** If  $M_1$ ,  $M_2$ ,  $M_3$  are positive definite matrices then the matrix polynomial  $M_1 + x M_2 + x^2 M_3$  is a stable matrix polynomial, and its determinant is stable.

*Proof.* We must verify that  $v^t(M_1 + x M_2 + x^2 M_3)v$  is stable for all non-zero  $v \in \mathbb{R}^{\mu}$ . Now all the coefficients of  $(v^t M_1 v) + x(v^t M_2 v) + x^2(v^t M_3 v)$  are positive since the  $M_i$  are positive definite, and , so this is a stable polynomial.

If  $\lambda$  is a root of  $|M_1 + x M_2 + x^2 M_3|$  then  $\lambda \neq 0$  since  $M_1$  has positive determinant. There is a non-zero vector v in  $\mathbb{C}^{\mu}$  such that

$$(\mathbf{M}_1 + \lambda \,\mathbf{M}_2 + \lambda^2 \,\mathbf{M}_3)\mathbf{v} = 0$$

Recall that if  $v^*$  is the conjugate transpose, then  $v^* M v > 0$  for non-zero v and positive definite M. Multiplying on the left by  $v^*$  yields

$$(v^* M_1 v) + \lambda (v^* M_2 v) + \lambda^2 (v^* M_3 v) = 0$$

Since this is a quadratic with positive coefficients, it is a stable polynomial, and so  $\lambda$  lies in the left half plane.

Surprisingly, powers such as  $(xI+A)^d$  are generally not stable matrix polynomials. We do have

**Lemma 21.104.** *If* A *is positive definite, and* B *is positive semi-definite then*  $(xI + A)^2 + B$  *is a stable matrix polynomial, and generally not hyperbolic.* 

*Proof.* First take B = 0. If  $v \in \mathbb{R}^{\mu}$  is non-zero then all coefficients of

$$x^{2}(v^{t}v) + x(2v^{t}Av) + (v^{t}A^{2}v)$$

are positive, so  $(xI + A)^2$  is a stable matrix polynomial. Next, we show that the discriminant  $4((\nu^t A \nu)^2 - (\nu^t \nu)(\nu^t A^2 \nu))$  is never positive. Define an inner product  $< \nu, w >= \nu^t w$ . Then

$$(v^{\mathsf{t}} \mathsf{A} v)^2 - (v^{\mathsf{t}} v)(v^{\mathsf{t}} \mathsf{A}^2 v) = < v, \mathsf{A} v >^2 - < v, v > < \mathsf{A} v, \mathsf{A} v > \leqslant 0$$

by the Cauchy-Schwartz inequality. If B is positive semi-definite then the discriminant is

$$4((v^{t} A v)^{2} - (v^{t} v)(v^{t} A^{2} v)) - 4(v^{t} v)(v^{t} B v)$$

which is still non-positive.

There is a similar result for cubics, but we need a condition on the eigenvalues to get stability.

**Lemma 21.105.** Suppose A is a positive definite matrix with minimum and maximum eigenvalues  $\lambda_{\min}$  and  $\lambda_{\max}$ . If  $\frac{\lambda_{\max}}{\lambda_{\min}} \leq 9$  then  $(xI + A)^3$  is a stable matrix polynomial.

*Proof.* We need to show that if v is any non-zero vector then

$$x^{3}(v^{t}v) + x^{2}(3v^{t}Av) + x(3v^{t}A^{2}v) + (v^{t}A^{3}v)$$

is stable. Since all the coefficients are positive, by Lemma 21.38 this is the case precisely when

$$\frac{(3\nu^{\mathrm{t}} A \nu)(3\nu^{\mathrm{t}} A^{2} \nu)}{(\nu^{\mathrm{t}} \nu)(\nu^{\mathrm{t}} A^{3} \nu)} \ge 1$$

Now this is equivalent to

$$\frac{(v^{\mathrm{t}} A v)}{(v^{\mathrm{t}} v)} \times \frac{(v^{\mathrm{t}} A^{2} v)}{(v^{\mathrm{t}} A^{3} v)} \ge \frac{1}{9}$$

The first factor lies in  $[\lambda_{\min}, \lambda_{\max}]$ , and the second lies in  $[1/\lambda_{\max}, 1/\lambda_{\min}]$ , so their product lies in  $\left[\frac{\lambda_{\min}}{\lambda_{\max}}, \frac{\lambda_{\max}}{\lambda_{\min}}\right]$ . By hypothesis  $\lambda_{\min}/\lambda_{\max} \ge 1/9$ , so the proof is complete.

The ratio  $\lambda_{max}/\lambda_{min}$  is known as the *condition number* of A, so we could restate the hypothesis to say that the condition number is at most 9.

CHAPTER



# Transformations in the complex plane

In this chapter we are interested in polynomials whose roots might be complex. If S is a region in the complex plane, then  $\mathbf{P}^{S}$  means all polynomials whose roots all lie in S. We will give some examples of linear transformations T and regions I, J such that T:  $\mathbf{P}^{I} \longrightarrow \mathbf{P}^{J}$ .

It is difficult to describe the behavior of a transformation on all polynomials, so we sometimes restrict ourselves to a special subset of polynomials. Define the function  $\phi : \mathbb{C} \mapsto \mathbf{P}^{\mathbb{C}}(n)$  by  $z \mapsto (x - z)^n$ . Given a transformation T we are interested in determining the image T  $\phi$ . In the best case we can find a a simple function  $\phi'$  so that the diagram commutes:



Given a set  $\mathcal{P}$  of polynomials we are also interested in all the roots of polynomials in  $\mathcal{P}$ . We extend the meaning of roots(f) to a set of polynomials. Define

$$\mathsf{roots}(\mathcal{P}) = \{ \zeta \in \mathbb{C} \mid \exists f \in \mathcal{P} \land f(\zeta) = 0 \}$$

If S is a region in  $\mathbb C,$  and T is a transformation, then since  $\varphi S \subset {I\!\!P}^S$  we know that

$$\operatorname{roots}(\mathsf{T}(\varphi S)) \subseteq \operatorname{roots}(\mathsf{T}(\mathbf{P}^S)).$$
(22.0.2)

We are interested in situations when (22.0.2) is an equality. See Question 205. We generalize the Hermite-Biehler theorem that characterizes polynomials whose roots are in the lower half plane in terms of interlacing properties of their real and imaginary parts.

The closed disk of radius r centered at  $\sigma$  is  $\Delta(\sigma, r)$ ; the unit disk is  $\Delta$ , and the boundary of the unit disk is the unit circle  $\partial \Delta$ .

## 22.1 The derivative and interlacing

We recall the Gauss-Lucas theorem and prove a converse.

The fundamental result about the derivative is the the Gauss-Lucas theorem which states that the roots of the derivative of f lie in the convex hull of the roots of f. It follows that

**Theorem 22.1.** If S is a convex subset of  $\mathbb{C}$  then the linear transformation  $f \mapsto f'$  maps  $\mathbf{P}^{S}$  to itself.

The converse is also true.

**Proposition 22.2.** If T is a linear transformation with the property that T:  $P^{S} \longrightarrow P^{S}$  for all convex regions S, then Tf is a multiple of some derivative.

*Proof.* We first choose S to be the single point 0. It follows that  $T(x^n) = a_n x^{e_n}$ , since  $T(x^n)$  can only have 0 as a root. Ignoring the trivial case, there is some non-zero  $a_k$ . Let s be the smallest integer so that  $a_s \neq 0$ . Next, consider  $S = \{-1\}$ . Since  $(x + 1)^s \in \mathbf{P}^S$  and  $T(x + 1)^s = a_s x^{e_s}$ , it follows that  $e_s = 0$ .

Consider  $\$ = \{-\lambda\}$ . The only members of  $\mathbf{P}^{\$}$  are multiples of powers of  $(x + \lambda)$ . We compute  $T(x + \lambda)^r$  in two ways:

$$T(x+\lambda)^{r} = \beta_{r}(x+\lambda)^{m_{r}} = \beta_{r} \sum_{j=0}^{m_{r}} {m_{r} \choose j} x^{j} \lambda^{m_{r}-j}$$
$$= \sum_{k=0}^{r} {r \choose k} T(x^{k}) \lambda^{r-k} = \sum_{k=0}^{r} {r \choose k} a_{k} x^{e_{k}} \lambda^{r-k}$$

Since these are identities in  $\lambda$  that hold for infinitely many values of  $\lambda$ , the coefficients of powers of  $\lambda$  must be equal. From consideration of the coefficient of  $\lambda^{r-k}$  we find

$$\binom{r}{k}a_{k}x^{e_{k}} = \beta_{r}\binom{m_{r}}{m_{r}-r+k}x^{m_{r}-r+k}$$
(22.1.1)

Choosing k = s shows that  $\beta_r \neq 0$  for  $r \ge s$ . Consequently,  $a_k \neq 0$  for  $k \ge s$ . Substituting r = k shows that  $\beta_r = a_r$ . If we take  $r \ge s$  then comparing exponents shows that  $e_k = m_r - r + k$ . It follows from this that  $e_k = k - c$  for some constant c, and  $k \ge s$ . Since we know that  $e_s = 0$ , it follows that  $T(x^k) = a_k x^{k-s}$  for  $k \ge s$  and is 0 for k < s. In addition,  $m_r = r - s$ . Substituting this into (22.1.1) yields

$$a_r = \begin{cases} \frac{a_s}{s!} \frac{r!}{(r-s)!} & r \ge s\\ 0 & r < s \end{cases}$$

which implies that  $T(x^r) = a_s D^s x^r$ , as promised.

Every point of S is a root of a derivative of a polynomial supported on the boundary.

**Lemma 22.3.** If S is a convex region of  $\mathbb{C}$ , and D is the derivative, then

roots( 
$$D(P^{\partial S})$$
) = S.

*Proof.* If  $\sigma, \tau \in \partial S$  then  $f(x) = (x - \sigma)(x - \tau) \in \mathbf{P}^{\partial S}$ . Since  $f' = 2x - (\sigma + \tau)$ , we see that  $\frac{1}{2}(\sigma + \tau) \in \text{roots}(\mathsf{D}\mathbf{P}^{\partial S})$ . Since S is convex, the set of all midpoints of pairs of vertices on the boundary equals all of S.

# 22.2 Properties of algebraic Interlacing

In this section we consider some questions about the location of the roots of algebraic interlacing polynomials.

**Definition 22.4.** Suppose  $f(z) = \sigma(z - r_1) \cdots (z - r_n)$ . We say that  $f \preccurlyeq g$  if and only if we can write

$$g(z) = f(z)\left(\frac{\rho_1}{x-r_1}+\cdots+\frac{\rho_1}{x-r_1}\right)$$

where  $\rho_1, \ldots, \rho_n$  are all real and non-negative.

It is immediate from this definition that if  $f, g \in \mathbf{P}$  then  $f \leq g$  iff  $f \leq g$ . For any f we have that  $f \leq f'$ , since we know

$$f'(x) = f(x) \left(\frac{1}{x-r_1} + \dots + \frac{1}{x-r_n}\right).$$

The Gauss-Lucas property holds for this definition of interlacing, and the proof is nearly the same, see [121, Theorem 6.1'].

**Lemma 22.5.** Suppose that S is a convex region, and  $f \in P^{\delta S}$ . If  $f \prec g$  then  $g \in P^{S}$ .

We now extend the Gauss-Lucas Theorem by showing that every possible point is a root of some interlacing polynomial. Let  $Hull(\zeta_1, ..., \zeta_n)$  denote the convex hull of the points  $(\zeta_1, ..., \zeta_n)$ . If these points are the roots of f, then we write  $Hull(f) = Hull(\zeta_1, ..., \zeta_n)$ .

**Proposition 22.6.** *If* f *is any polynomial, then every point in* Hull(f) *is a root of some* g *where*  $f \preccurlyeq g$ . *Consequently,* 

$$Hull(f) = roots(\{g \mid f \preccurlyeq g\})$$

*Proof.* lemma 22.5 shows that  $roots(\{g \mid f \preccurlyeq g\}) \subseteq Hull(f)$ . Let  $\zeta = \{\zeta_1, \ldots, \zeta_n\}$  be the roots of f. We first show that if  $0 \in Hull(f)$  then  $0 \in roots(\{g \mid f \preccurlyeq g\})$ . For  $\beta = (\beta_1, \ldots, \beta_n)$  we define

$$f_{\beta} = \sum_{k=1}^{n} \beta_k \frac{f}{x - \zeta_k}$$
(22.2.1)

The requirement that  $f_{\beta}(0) = 0$  implies that

$$0 = \pm \zeta_1 \cdots \zeta_n \, \left( \frac{\beta_1}{\zeta_1} + \cdots + \frac{\beta_n}{\zeta_n} \right)$$

Thus, it suffices to see that 0 is in the convex hull of  $\{1/\zeta_1, ..., 1/\zeta_n\}$ , which is clear, since 0 is in the convex hull of  $\{\zeta_1, ..., \zeta_n\}$ .

Choose  $\alpha$  in the convex hull of f. Define the Blaschke (see last section) transformation

$$\mathsf{T}\colon z\mapsto \frac{z-\alpha}{1-\overline{\alpha}z}$$

Consider  $f_{\beta}(z)$  and the transformed polynomial  $g_{\beta}(z) = (1 - \overline{\alpha}z)^n f_{\beta}(Tz)$ . If  $\beta$  is chosen so that  $g_{\beta}(0) = 0$  then  $f_{\beta}(\alpha) = 0$ . So, we need to know that if  $\alpha$  is in the convex hull of the  $\zeta_i$  then 0 is in the convex hull of  $T\zeta_i$ . But since  $\alpha \in Hull(f)$  there are  $\gamma_i$  so that

$$\alpha = \sum \gamma_i \zeta_i$$

and so

$$0 = \mathsf{T}(\alpha) = \sum \gamma_{i} \mathsf{T}(z_{i})$$

Thus, 0 is in the hull, and we are done.

This last proposition showed that there are no restrictions on the locations of roots of polynomials interlacing a given polynomial. However, there are restrictions on the simultaneous location of *all* the roots of a given polynomial interlacing f. Write  $f(x) = (x - r_1) \cdots (x - r_n)$ , and define

$$f_{\beta}(z) = \sum \beta_{i} \frac{f(z)}{z - r_{i}}$$

Letting  $s_1, \ldots, s_{n-1}$  be the roots of  $f_\beta$ , and substituting  $z = r_k$  we have

$$\prod_{\mathfrak{i}}(r_k-s_{\mathfrak{i}})=\beta_k\prod_{j\neq k}(r_k-r_j)$$

and since  $\beta_k$  is positive we have

$$\sum_{i} arg(r_k - s_i) = \sum_{j \neq k} arg(r_k - r_j)$$

This says that the sum of the angles (mod  $2\pi$ ) formed at  $r_k$  by the roots of  $f_\beta$  is a fixed quantity. We can therefore derive restrictions such as

The roots of  $f_{\beta}$  can not satisfy

$$0 \leqslant arg(r_k - s_i) < \frac{1}{n-1} \sum_{j \neq k} arg(r_j - r_k) \qquad \text{for } 1 \leqslant i \leqslant n-1$$

662

For example, if n = 3 this restriction becomes:

The angles at a root of f formed by the two roots can not both be less than half the angle at the vertex.

In the picture below, it follows that  $f_{\beta}$  has one root in each triangle marked a (or one each in b, or one each in c). The interior lines are the angle bisectors.



Figure 22.1: Where the roots are.

# **22.3** The map $\phi$

When we have a formula for  $T(x - z)^n$  we can use this information to determine a map  $\phi'$  so that (22.0.1) is satisfied. We begin with multiplier transformations.

**Example 22.7.** Multiplier transformations have a very simple action in the complex plane. Suppose that T is a multiplier transformation. We have a fundamental commutativity diagram that is immediate from the definition of multiplier:



**Lemma 22.8.** Suppose that T is a multiplier transformation  $x^k \mapsto a_k x^k$  and  $r_1, \ldots, r_n$  are the roots of  $T(x-1)^n$ . The definition of  $\varphi'$  below makes (22.0.1) commute.

$$\phi': z \mapsto \prod_{k} (x - r_k z)$$

*Proof.* If  $z \in \mathbb{C}$ , then  $T(x - z)^n(x) = 0$  implies that  $T(\frac{x}{z} - 1)^n(x) = 0$ . Since T is a multiplier transformation this is equivalent to  $T(x - 1)^n(\frac{x}{z}) = 0$ , which implies the result.

**Example 22.9.** If we use the formula (7.10.4) for Laguerre polynomials with the transformation  $T(x^k) = L_k(x)$  we get

$$T(x+y)^{n} = (y+1)^{n}T(x^{n})(\frac{x}{y+1})$$

$$T\left(\frac{x+y}{y+1}\right)^{n} = T(x^{n})(\frac{x}{y+1})$$
(22.3.1)

and by linearity yields

$$T\left(f\left(\frac{x+y}{y+1}\right)\right) = T(f)(\frac{x}{y+1})$$

If we substitute  $\alpha(y + 1)$  for x in (22.3.1) then we see that

$$T(x + y)^{n} (\alpha(y + 1)) = (y + 1)^{n} T(x^{n})(\alpha) = (y + 1)^{n} L_{n}(\alpha)$$

and that therefore the roots of  $T((x - y)^n)$  are given by  $\alpha(1 - y)$ , where  $\alpha$  is a root of  $L_n$ . In particular, it follows that

**Lemma 22.10.** Suppose  $T(x^n) = L_n(x)$  has roots  $r_1, \ldots, r_n$ . The following definition of  $\phi'$  makes (22.0.1) commute:

$$\phi': z \mapsto \prod_{k=1}^{n} (x - r_{i}(1+z))$$

For instance, if  $\alpha$  is positive then the roots of  $T(x + \alpha \iota)^n$  lie on vertical rays emanating from the roots of  $L_n$ .

**Example 22.11.** We use the addition formula (22.4.2) for Hermite polynomials to determine  $\phi'$ .

**Lemma 22.12.** Suppose  $T(x^n) = H_n(x)$  has roots  $r_1, \ldots, r_n$ . The following definition of  $\phi'$  makes (22.0.1) commute:

$$\phi': z \mapsto \prod_{k=1}^{n} (x - (r_i + (1/2)(1+z)))$$

*Proof.* From the addition formula (22.4.2) with  $f = (x + 1)^n$  we have

$$\mathsf{T}(\mathsf{x}+1)^{\mathsf{n}}\big|_{\mathsf{x}\mapsto\mathsf{x}+\mathsf{y}} = \mathsf{T}_{*}(\mathsf{x}+2\mathsf{y}+1)^{\mathsf{n}}$$

If we let z = 2y + 1 then

$$T(x+1)^{n} (x-1/2+z/2) = T_{*}(x+z)^{n}$$

If r is a root of  $T(x + 1)^n$ , then x - 1/2 + z/2 = r yields a root of the right hand side. Thus x = r + 1/2 + z/2.

**Example 22.13.** Suppose that  $T(x^n) = x^n H_n(1/x) = H_n^{rev}(x)$ . We have the following identity that can be derived from the similar identity for the Hermite polynomial transformation.

$$T(x+2u-2)^n = u^n T(x^n) (\frac{x}{u})$$
 (22.3.2)

**Lemma 22.14.** Suppose  $T(x^n) = x^n H_n(1/x)$  has roots  $r_1, \ldots, r_n$ . The following definition  $\phi'$  makes (22.0.1) commute:

$$\varphi':z\mapsto \prod_k\ (x-r_k(1+z/2)\,)$$

*Proof.* We must show that  $T(x - z)^n = \prod_k (x - r_k(1 + z/2))$ . The conclusion follows as before.

#### 22.4 Lines, rays, and sectors

We show that some transformations preserve lines or rays or sectors in the complex plane.

**Lemma 22.15.** Suppose  $T: x^n \mapsto a_n x^n$  is a multiplier transformation that maps  $P^{pos}$  to itself. For any  $\sigma \in \mathbb{C}$ , we have  $T: P^{\sigma \mathbb{R}} \longrightarrow P^{\sigma \mathbb{R}}$ .

*Proof.* Suppose  $f(x) = \prod_{i=1}^{n} (x - \sigma r_i)$  where  $r_i \in \mathbb{R}$ , and let  $g(x) = \prod (x - r_i)$ . Then since T is a multiplier transformation

$$T f(x) = T(f(\sigma x))(x/\sigma) = T(\sigma^n g(x))(x/\sigma) = \sigma^n T(g(x))(x/\sigma)$$

Since T:  $\mathbf{P} \longrightarrow \mathbf{P}$  we know that  $\mathsf{T}(\mathfrak{g}(\mathbf{x})) \in \mathbf{P}$ , and so  $\mathsf{TF} \in \mathbf{P}^{\sigma \mathbb{R}}$ .

Similarly, the roots of  $(x - \iota - \alpha)^n$  lie on the horizontal lines  $\{\iota r_k + \mathbb{R}\}$ .

Here is a very general way of finding linear transformations that preserve lines.

**Proposition 22.16.** Suppose  $f \in \widehat{P}$ . The linear transformation  $g \mapsto f(\sigma D)g$  maps  $P^{\xi+\sigma\mathbb{R}}$  to itself for any  $\xi \in \mathbb{R}$  and  $\sigma \in \mathbb{C}$ .

*Proof.* Since f is a limit of polynomials in **P**, it suffices to prove the result for linear polynomials. So, choose  $g \in \mathbf{P}^{\xi+\sigma\mathbb{R}}$ ; it suffices to show that  $(\sigma D + \beta)g = \sigma g' + \beta g$  is also in  $\mathbf{P}^{\xi+\sigma\mathbb{R}}$  for any  $\beta \in \mathbb{R}$ . Since  $\frac{d}{dx}g(x+\gamma) = g'(x+\gamma)$ , we take  $\xi = 0$ .

If  $g \in \mathbf{P}^{S}$  and we write  $g(x) = \prod_{i=1}^{n} (x - \sigma r_{k})$  with  $r_{i} \in \mathbb{R}$ , then

$$\mathfrak{g}(\sigma x)=\sigma^n\prod(x-r_k).$$

If we define  $h(x) = \sigma^{-n}g(\sigma x)$  then  $h \in \mathbf{P}$ . Note also that  $h'(x) = \sigma^{-n+1}g'(\sigma x)$ . When  $k(x) = \sigma g'(x) + \beta g(x)$  then

$$\mathbf{k}(\sigma \mathbf{x}) = \sigma \mathbf{g}'(\sigma \mathbf{x}) + \beta \mathbf{g}(\sigma \mathbf{x}) = \sigma^{\mathbf{n}}(\mathbf{h}'(\mathbf{x}) + \beta \mathbf{h}(\mathbf{x})).$$

Since  $h \in \mathbf{P}$ , it follows that  $\sigma^n k(\sigma x) \in \mathbf{P}$ , and hence the roots of k(x) are real multiples of  $\sigma$ .

**Corollary 22.17.** If  $f(x) \in \mathbf{P}^{\alpha+\iota\mathbb{R}}$  then  $f(x+1) + f(x-1) \in \mathbf{P}^{\alpha+\iota\mathbb{R}}$ .

*Proof.* Simply note that

$$f(x + 1) + f(x - 1) = (e^{D} + e^{-D})f = 2\cos(\iota D)f$$

and  $\cos(x) \in \widehat{\mathbf{P}}$ .

*Remark* 22.18. This transformation appears in [26], where it is used to show that certain polynomials have roots whose real part is 1/2.

In the the next two examples the regions are rays and horizontal lines.

**Lemma 22.19.** Consider  $T: x^k \mapsto H_k(x)x^{n-k}$  acting on P(n). Every polynomial with roots on a ray in the complex plane based at 2 is mapped by T to a line through the origin with the same slope.

*Proof.* Assume that f has all its roots on the line  $2 + \alpha \sigma$ , where  $\sigma \in \mathbb{C}$  is fixed, and  $\alpha \ge 0$ . If we define S(f) = T(f(x - 2)), then it suffices to determine the action of S on polynomials with roots on the ray through  $\sigma$ . Since  $\sigma^{-n}f(\sigma x)$ ) has all positive roots, the result follows from Lemma 9.60

**Lemma 22.20.** If  $T(x^k) = H_k(x)$  then  $\mathbf{P}^{\mathbb{R}+\alpha \iota} \longrightarrow \mathbf{P}^{\mathbb{R}+\alpha \iota/2}$ . In other words, if f(x) is a polynomial whose roots all have imaginary part  $\alpha \iota$ , then the roots of T(f) all have imaginary part  $\frac{\alpha \iota}{2}$ .

Proof. The addition formula for Hermite polynomials (7.8.4) implies that

$$\mathsf{T}(\mathsf{x}^{\mathsf{n}})(\mathsf{x}+\mathsf{y}) = \mathsf{T}_*(\mathsf{x}+2\mathsf{y})^{\mathsf{n}}$$

and by linearity

$$T(f)(x + y) = T_* f(x + 2y)$$
(22.4.1)

Replacing x + y by x in this equation yields

$$T(f) = (T_*(f(x+2y))(x-y))$$
(22.4.2)

Equation (22.4.2) implies the following diagram commutes:



Multiplier transformations preserve sectors because rotation commutes with such transformations.

**Lemma 22.21.** Suppose  $T: x^n \mapsto a_n x^n$  maps **P** to itself and has all positive coefficients. If S is a sector of angle at most  $\pi$  then T maps **P**<sup>S</sup> to itself.

*Proof.* Since T preserves interlacing and the sign of the leading coefficient it maps  $f + \iota g \in \mathfrak{I}$  to  $Tf + \iota Tg \in \mathfrak{I}$ . If  $|\sigma| = 1$  is a rotation then T maps  $\mathbf{P}^{-\sigma H}$  to itself. Consequently.

$$\mathsf{T} \colon \mathbf{P}^{-\sigma\mathsf{H}\cap-\mathsf{H}} = \mathbf{P}^{-\sigma\mathsf{H}} \cap \mathbf{P}^{-\mathsf{H}} \longrightarrow \mathbf{P}^{-\sigma\mathsf{H}} \cap \mathbf{P}^{-\mathsf{H}} = \mathbf{P}^{-\sigma\mathsf{H}\cap-\mathsf{H}}$$

We can choose  $\sigma$  so that  $-\sigma H \cap -H$  is a sector of any desired opening of at most  $\pi$ . Another rotation to S finishes the proof.

## 22.5 The set of roots

In this section we look at properties of the set of roots. If we are in the fortunate situation of (22.0.1) then for any subset S of  $\mathbb{C}$  we know that

$$roots(T\varphi S) = roots(\varphi'S)$$

In general we can not describe the behavior so well. We next note that we can refine Lemma 6.21 as follows:

**Lemma 22.22.** Suppose that  $T: P^{(a,b)} \longrightarrow P$  preserves interlacing. Then

$$roots(T(\phi(a, b))) = roots(T(P^{(a, b)})).$$

*(* **1**)

*Proof.* The proof follows from the proof of Lemma 6.21 and using the continuity of roots.

If  $\ell_n$  is the largest root of  $L_n$ , then from Lemma 22.10

```
roots(T(\phi\Delta)) = \int \{roots \text{ of } T(x-z)^n \mid |z| \leq 1\}
```

consists of all points in the circle of radius  $\ell_n$  centered at  $\ell_n$ . See Figure 22.2 which shows  $roots(T(\phi(\partial \Delta)))$  when n is 5. Since  $\ell_n$  goes to infinity as n goes to infinity, the union of all these circles covers the right half plane. This proves

**Proposition 22.23.** *If*  $T(x^k) = L_k$ , *then*  $\{\Re(\zeta) \ge 0\} \subseteq \text{$ *roots* $}(T(\varphi\Delta))$ .

In other words, if  $\Re(\xi) \ge 0$  then there is a *z* and positive integer n where  $|z| \le 1$  such that  $\xi$  is a root of  $T(x - z)^n$ .

## **22.6** Complex Transformations involving f(x + i)

We introduce some transformations that involve complex numbers, and show that they preserve P because they are other known transformations in disguise.

To begin, define the transformation T(f) = f(x+i) + f(x-i). If f has all real coefficients, then T(f) has all real coefficients, since Tf is equal to its conjugate.



Figure 22.2: Image of the unit circle under Laguerre transformation

**Lemma 22.24.** The transformation  $T: f(x) \mapsto f(x+i) + f(x-i)$  preserves roots.

*Proof.* Since  $e^{\alpha D} = f(x + a)$  it follows that  $\cos(D)f = \frac{1}{2}Tf$ . Since  $\cos(x)$  is in  $\widehat{\mathbf{P}}$ ,  $\cos(D)$  preserves roots, and thus so does T.

*Remark* 22.25. More generally, if we use the fact that  $\cos(ax + b) \in \widehat{\mathbf{P}}$  we see that

$$\cos(a\mathsf{D}+b)\mathsf{f}(x) = \frac{1}{2} \big( e^{\imath b} \mathsf{f}(x+\imath a) + e^{-\imath b} \mathsf{f}(x-\imath a) \big)$$

is in **P**. [27]

We can iterate the construction of the following lemma. If we let the number of terms go to zero then the sum becomes a convolution. See [29].

**Lemma 22.26.** *If*  $a_1, \ldots, a_r, b_1, \ldots, b_r$  *are real numbers, and*  $f \in P$ *, then* 

$$\frac{1}{2^{r}}\sum f(x \pm a_{1}i \pm a_{2}i \pm \cdots \pm a_{r}i)e^{i(\pm b_{1} \pm b_{2} \cdots \pm b_{r})} \in P$$

where the sum is over all  $2^{r}$  choices of  $\pm$  signs.

*Proof.* This is  $\cos(a_1 D + b_1) \cdots \cos(a_r D + b_r) f$ .

We get a similar result using sin instead of cos. The proof is omitted.

**Lemma 22.27.** The map  $f(x) \mapsto \frac{f(x+i)-f(x-i)}{2i}$  maps **P** to itself.

Next we look at a variation of the above.

**Lemma 22.28.** If n is a positive integer and  $f(x) \in P$  then

$$f(x) + \sum_{k=-n}^{n} f(x+ki) \in P$$

*Proof.* We compute that

$$f(x) + \sum_{k=-n}^{n} f(x+ki) = 2(1+\cos(D)+\cos(2D)+\dots+\cos(nD)) f(x)$$
$$= 2\left[\frac{\sin((n+1)x/2)}{\sin(x/2)}\cos(nx/2)\right] (D) f(x)$$

Now  $\cos(nx/2)$  is in  $\widehat{\mathbf{P}}$ , and so is the first factor, since we can write an infinite product expansion for it, and the factors on the bottom are canceled out by terms on the top.

**Corollary 22.29.** If 
$$f(x) \in P$$
 then  $\int_{-1}^{1} f(x + \iota t) dt \in P$ .

*Proof.* Approximate the integral by the sum

$$\frac{1}{2n}\left(f(x)+\sum_{k=-n}^n f(x+\iota k/n)\right).$$

This sum is in **P**, for we can scale f(x) in the Lemma.

Here is an alternative proof not using sums. If  $\mathsf{T}(\mathsf{f})$  is the integral in the Corollary, then

$$T(x^{n}) = \int_{-1}^{1} (x + \iota t)^{n} dt = \frac{1}{\iota} \left( \frac{(x + \iota)^{n+1}}{n+1} - \frac{(x - \iota)^{n+1}}{n+1} \right)$$
$$= 2 \frac{e^{\iota D} - e^{-\iota D}}{2\iota} \frac{x^{n+1}}{n+1} = 2 \sin(D) \int x^{n}$$

Thus  $T(f) = 2 \sin(D) \int f$ . This in in **P** since

$$\sin(\mathsf{D}) \int \mathsf{f} = \left(\frac{\sin(x)}{x}(\mathsf{D})\right) \mathsf{D}(\int \mathsf{f}) = \left(\frac{\sin(x)}{x}(\mathsf{D})\right) \mathsf{f}$$
(22.6.1)

and  $\frac{\sin(x)}{x}$  is in  $\widehat{\mathbf{P}}$ .

Translating and scaling this result gives the following corollary, which will prove useful when we look at polynomials with complex coefficients (Chapter 20).

**Corollary 22.30.** If  $\alpha > 0$  and  $f(x) \in P$  then the all the roots of  $\int_0^1 f(x + \alpha t) dt$  have imaginary part  $-\alpha/2$ .

*Proof.* Observe that

$$\int_{0}^{1} f(x + \alpha \iota t) dt = \frac{1}{2} \int_{0}^{2} f(f + \alpha \iota t/2) dt = \frac{1}{2} \int_{-1}^{1} f([x - \alpha \iota/2] + \alpha \iota t/2) dt$$

Since the last integral has all real roots, the roots of the first integral have imaginary part  $-\alpha/2$ .

### 22.7 The Hermite-Biehler problem

This section is concerned with determining how many roots of a polynomial with complex coefficients lie in the upper half plane. If a polynomial h has  $\alpha$  roots in the Upper half plane, and  $\beta$  roots in the Lower half plane, then we define  $\mathcal{UL}(h) = (\alpha, \beta)$ .

**The Hermite-Biehler problem:** Suppose that f, g have all real coefficients and no common real roots. Express  $U\mathcal{L}(f + \iota g)$  in terms of the arrangement of the roots of f and g.

Suppose f and g are two polynomials with real coefficients, with perhaps some complex roots. The polynomial  $f + \iota g$  has the following fundamental property which follows from the fact that f and g have all real coefficients:

If  $f + \iota g$  has a real root, then f and g have that root in common.

Thus, if f and g have no real roots in common, then all the roots of  $f + \iota g$  are complex. Our approach is based on this simple consequence:

If  $f_t$  and  $g_t$  are a continuous family of real polynomials where  $f_t$ and  $g_t$  have no common root for all  $0 \le t \le 1$ , then  $\mathcal{UL}(f_0 + \iota g_0) =$  $\mathcal{UL}(f_1 + \iota g_1)$  since the roots can never cross the real axis.

We begin by describing three transformations of  $f + \iota g$  that preserve the number of roots in each half plane. We assume that f and g have no common real roots. If  $UL(f + \iota g) = (\alpha, \beta)$  then  $UL(-f - \iota g) = (\alpha, \beta)$ . Taking conjugates shows that

$$\mathcal{UL}(f - \iota g) = \mathcal{UL}(-f + \iota g) = (\beta, \alpha)$$

Consequently, we will always assume that f and g have positive leading coefficients.

#### 1. Join adjacent roots

Suppose that u, v are adjacent roots of g that are not separated by a root of f, and let  $u_t = tu + (1 - t)v$ . The transformation below doesn't ever create a  $g_t$  with a root in common with f since  $u_t$  lies between u and v. Thus  $\mathcal{UL}(h) = \mathcal{UL}(h_1)$  since  $h_0 = h$ .

$$h_t = f + \iota \frac{g}{x - \nu} \cdot (x - u_t)$$

## 2. Replace double roots with $x^2 + 1$

Suppose that u is a double root of g. The following transformation moves a pair of roots of g into the complex plane, missing all real roots of f, and lands at  $\pm \iota$ . Consequently, we get a factor of  $x^2 + 1$ . Again,  $\mathcal{UL}(h) = \mathcal{UL}(h_1)$ .

$$h_t=f+\iota\frac{g}{(x-u)^2}\,\left((x-(1-t)u)^2+t\right)$$

## 3. **Replace complex roots with** $x^2 + 1$

If a + bi is a root of g then since g has all real coefficients g has a factor of  $(x - a)^2 + b^2$ . The transformation below converts this factor to  $x^2 + 1$ .

$$h_t = f + \iota \frac{g}{(x-a)^2 + b^2} \left( (x - (1-t)a)^2 + (1-t)b^2 + t \right)$$

We can also do the above constructions to f. If we do them as many times as possible, then we have found a polynomial that we call the **HB**-simplification of h:

$$H = (x^2 + 1)^r F + \iota (x^2 + 1)^s G$$

The constructions above guarantee that  $\mathcal{UL}(h) = \mathcal{UL}(H)$ . In addition, F and G interlace, since we have removed all consecutive roots belonging to only one of f, g. We can now state our result:

**Proposition 22.31.** Suppose f and g are polynomials with real coefficients, positive leading coefficients, and no common real roots. Let  $H = (x^2 + 1)^r F + \iota(x^2 + 1)^s G$  be the HB-simplification of  $h = f + \iota g$ , and set m = max(r, s). Then

$$\mathcal{UL}(h) = \begin{cases} (m, m) + (0, \deg(F)) & \text{if } F \longleftarrow G\\ (m, m) + (\deg(G), 0) & \text{if } G \longleftarrow F \end{cases}$$

*Proof.* If F and G have the same degree then without loss of generality we may assume that  $F \ll G$ . Suppose that r > s. Consider

$$h_t = (x^2 + 1)^r f + \iota (tx^2 + 1)(x^2 + 1)^s g$$

Since deg(h<sub>t</sub>) = deg(h),  $\mathcal{UL}(h_1) = \mathcal{UL}(h)$ . We can do a similar transformation if r < s, so we may assume that both exponents are equal to  $m = \max(r, s)$ . If write  $(x^2 + 1)^m H_1 = H$  then we see that  $\mathcal{UL}(h) = \mathcal{UL}(h_1) + (m, m)$ . Since  $H_1 = F + \iota G$ , we can apply the classical Hermite-Biehler theorem to conclude that  $\mathcal{UL}(H_1) = (0, deg(F))$ . If F < G then we consider  $G_{\varepsilon} = \varepsilon F + G$  and let  $\varepsilon$  go to zero. The proposition is now proved.

For completeness, here's a quick proof of the Hermite-Biehler theorem.

Suppose  $h = f + \iota g$  with  $f \ll g$ , and the roots of f are  $\nu_1, \ldots, \nu_n$  then there is a positive  $\sigma$  and positive  $a_i$  so that

$$g = \sigma f + \sum a_i \frac{f}{x - v_i}$$

A root  $\rho$  of f +  $\iota$ g satisfies

$$\begin{split} \sigma + \sum \frac{a_i}{\rho - \nu_i} &= \mathfrak{l} \\ \sigma + \sum \frac{a_i(\overline{\rho} - \nu_i)}{|\rho - \nu_i|^2} &= \mathfrak{l} \end{split}$$
from which it follows that  $\rho$  has negative imaginary part.

**Example 22.32.** Here is an example. Suppose that

$$f = (x^{2} + 2)(x^{2} + 3)(x - 2)(x - 4)(x - 5)(x - 8)(x - 9)(x - 10)$$
  
$$g = (x^{2} + 5)(x - 1)(x - 3)(x - 6)(x - 7)$$

We write the roots of f and g in order, and successively remove pairs of adjacent roots of f (or of g)

gfgffggfff gfgggfff gfgfff gfgf

Thus  $F \leq G$ . Since f has degree 10 and g has degree 6, we see that r = (10-2)/2 = 4, and s = (6-2)/2 = 2. Thus m = 4 and

$$\mathcal{UL}(f + \iota g) = (4, 6)$$

The image of the upper half plane H under a counterclockwise rotation of angle  $\alpha$  is  $e^{i\alpha}$ H. A polynomial f(x) has all its roots in this half plane if and only if  $f(e^{\alpha i}) \in \mathbf{P}^{H}$ , and this is the case exactly when

$$\Re(f(e^{\alpha \iota})) \triangleleft \Im(f(e^{\alpha \iota}))$$

and the leading coefficients have opposite signs. However, it is difficult in general to express the real and imaginary parts of  $f(e^{i\alpha})$  in terms of coefficients of f. When  $e^{i\alpha}$  is a root of unity then there is an explicit formula. We describe the general case, and then look at three special cases.

Choose a positive integer n and let  $\zeta = e^{2\pi i/n}$ . Write  $f = \sum a_i x^i$  and define

$$f_k(x) = \sum_{s \equiv k \pmod{n}} a_s x^s$$

We can express  $f(\zeta x)$  in terms of the  $f_k$ :

$$f(\zeta x) = \sum_{k=0}^{n-1} f_k(\zeta x) = \sum_{k=0}^{n-1} \zeta^k f_k(x)$$

If we define

$$\begin{split} F_0(x) &= \sum_{k=0}^{n-1} \cos(2k\pi/n) \, \Re(f_k(x)) - \sin(2k\pi/n) \, \Im(f_k(x)) \\ F_1(x) &= \sum_{k=0}^{n-1} \sin(2k\pi/n) \, \Re(f_k(x)) + \cos(2k\pi/n) \, \Im(f_k(x)) \end{split}$$

then since  $\zeta = \cos(2k\pi/n) + \iota \sin(2k\pi/n)$ 

$$f(\zeta x) = F_0(x) + \iota F_1(x)$$

and  $F_0$  and  $F_1$  have all real coefficients. Consequently,  $f \in \mathbf{P}^{\zeta H}$  iff  $F_0 \ll F_1$  and the leading coefficients have opposite signs.

If  $f(\boldsymbol{x})$  has all real coefficients then the imaginary parts are zero and we have

$$\begin{split} F_0(x) &= \sum_{k=0}^{n-1} \cos(2k\pi/n)\,f_k(x) \\ F_1(x) &= \sum_{k=0}^{n-1} \sin(2k\pi/n)\,f_k(x) \end{split}$$

We now look at three special cases.

**Example 22.33.** If n = 3 then  $\zeta = \frac{-1}{2} + \frac{\sqrt{3}}{2}i$ , and  $\zeta^2 = \frac{-1}{2} - \frac{\sqrt{3}}{2}i$ . We define

$$F_{0}(x) = \Re(f_{0}) + \frac{-1}{2}\Re(f_{1}) - \frac{\sqrt{3}}{2}\Im(f_{1}) + \frac{1}{2}\Re(f_{2}) - \frac{\sqrt{3}}{2}\Im(f_{2})$$
(22.7.1)

$$F_{1}(x) = \Im(f_{0}) + \frac{\sqrt{3}}{2}\Re(f_{1}) + \frac{-1}{2}\Im(f_{1}) + \frac{\sqrt{3}}{2}\Re(f_{2}) - \frac{1}{2}\Im(f_{2})$$
(22.7.2)

If the coefficients of f are real then

$$F_0(x) = f_0 + \frac{-1}{2}f_1 + \frac{1}{2}f_2$$
$$F_1(x) = \frac{\sqrt{3}}{2}f_1 + \frac{\sqrt{3}}{2}f_2$$

We conclude:

**Lemma 22.34.** All roots of f have argument between  $2\pi/3$  and  $5\pi/3$  if and only if  $F_0$  and  $F_1$  in (22.7.1),(22.7.2) satisfy  $F_0 \ll F_1$  and their leading coefficients have opposite signs. If f has all real coefficients then the interlacing criterion is

$$2f_0 - f_1 + f_2 \leqslant f_1 + f_2.$$

**Example 22.35.** We now take n = 4. Since  $e^{2\pi \iota/4} = \iota$ , if we follow the calculation above, then

$$\begin{split} F_0 &= \mathfrak{R}(f_0) - \mathfrak{I}(f_1) - \mathfrak{R}(f_2) + \mathfrak{I}(f_3) \\ F_1 &= \mathfrak{I}(f_0) + \mathfrak{R}(f_1) - \mathfrak{I}(f_2) - \mathfrak{R}(f_3) \end{split}$$

and consequently f has roots with only negative real parts if and only if  $F_0 \ll F_1$ . This can be further simplified. If we assume that f(x) has all real coefficients then

$$F_0 = f_0 - f_2$$
  
$$F_1 = f_1 - f_3$$

If we write  $f(x) = p(x^2) + xq(x^2)$  then  $F_0(x) = p(-x^2)$  and  $F_1(x) = xq(-x^2)$ . In this case, the result is also known as the Hermite-Biehler [81] theorem:

**Theorem 22.36** (Hermite-Biehler). Let  $f(x) = p(x^2) + xq(x^2)$  have all real coefficients. The following are equivalent:

- 1. All roots of f have negative real part.
- 2.  $p(-x^2)$  and  $xq(-x^2)$  have simple interlacing roots, and the leading coefficients are the same sign.

**Example 22.37.** We next take n = 8, where  $\zeta = \frac{1+1}{\sqrt{2}}$ . We assume that all coefficients of f are real. Define

$$F_0(\mathbf{x}) = f_0 + \frac{f_1}{\sqrt{2}} - \frac{f_3}{\sqrt{2}} - f_4 - \frac{f_5}{\sqrt{2}} + \frac{f_7}{\sqrt{2}}$$
(22.7.3)

$$F_1(x) = \frac{f_1}{\sqrt{2}} + f_2 + \frac{f_3}{\sqrt{2}} - \frac{f_5}{\sqrt{2}} - f_6 - \frac{f_7}{\sqrt{2}}$$
(22.7.4)

**Lemma 22.38.** If f(x) has all real coefficients, then  $\Re(r) < \Im(r)$  for every root r of f(x) if and only if  $F_0$ ,  $F_1$  in (22.7.3),(22.7.4) have leading coefficients of opposite signs, and satisfy  $F_0 < F_1$ .

Suppose we are given a line  $\ell$  in the complex plane such that the angle the line makes with the real line is a rational multiple of  $2\pi$ . We can find an interlacing restriction such that a polynomial has all its roots in a particular half plane determined by  $\ell$  if and only if the interlacing condition is met, along with agreement or disagreement of the signs of the leading coefficients.

We can combine these interlacing and sign conditions, provided the relevant angles are rational. We can summarize this as

**Lemma 22.39.** Suppose that S is a convex region determined by n lines, and each line makes a rational angle with the real axis. Then, we can find n interlacing and sign restrictions such that a polynomial has all its roots in S if and only if these n restrictions are satisfied.

#### 22.8 The Bôcher-Walsh problem

The essence of the Bôcher-Walsh theorem can be stated [155] in the following form

If all roots of f lie in the upper half plane and all roots of g lie in

the lower half plane then  $\begin{vmatrix} f & f' \\ g & g' \end{vmatrix}$  has no real zeros.

This is reminiscent of the Gauss-Lucas theorem, which in general follows from the fact that if f has all its roots in the upper half plane then so does f'. Unlike the Hermite-Biehler theorem, it's easy to determine the location of the roots of the determinant.

**Lemma 22.40.** Suppose that  $f \blacktriangleleft f_1$  has all roots in the upper half plane, and  $g \blacktriangleleft g_1$  has all its roots in the lower half plane. If deg(f) = n > m = deg(g) then

f	$f_1  _{has}$	∫n-1	roots in the upper half plane
g	$g_1   mas$	m	roots in the lower half plane

*Proof.* If the determinant is zero then  $fg_1 - gf_1 = 0$ , so  $f_1/f = g_1/g$ . Because of algebraic interlacing, we can find non-negative  $a_i$ ,  $b_i$  so that

$$\sum a_i \frac{1}{x - r_i} = \sum b_i \frac{1}{x - s_i}$$

where  $roots(f) = (r_i)$  and  $roots(g) = (b_i)$ . If we evaluate the equation at any real number the left hand side is in the upper half plane, and the right hand side is in the lower half plane, so there are no real roots.

Thus, as f and g vary among polynomials satisfying the hypothesis the roots of the determinant never cross the real axis. We may therefore choose  $f = (x - \iota)^n$ ,  $f_1 = f'$ ,  $g = (x + \iota)^m$  and  $g_1 = g'$ , in which case

$$\begin{vmatrix} f & f_1 \\ g & g_1 \end{vmatrix} = \begin{vmatrix} (x-\iota)^n & n(x-\iota)^{n-1} \\ (x-\iota)^m & m(x-\iota)^{m-1} \end{vmatrix} = (x-\iota)^{n-1}(x+\iota)^{m-1}((n-m)x+(n+m))$$

This determinant has n - 1 roots in the upper half plane, and m in the lower half plane, so the lemma is proved.

If n = m then the proof shows that there are n - 1 roots in each half plane.

#### 22.9 Newton's inequalities in the complex plane

If we constrain the location of the roots of a polynomial to a region S of the complex plane, then geometric properties of S lead to restrictions on the Newton quotients. It turns out that the inequalities only depend on the behavior of the Newton quotients on quadratics, so we define

$$NQ(S) = \inf_{f \in \mathbf{P}^{S}(2)} \left| \frac{a_{1}^{2}}{a_{0}a_{2}} \right|$$

where  $f(x) = a_0 + a_1x + a_2x^2$ . Consider some examples. The usual argument shows that  $NQ(\mathbb{R}^+) = 4$ . Similarly,  $NQ(\sigma\mathbb{R}^+) = 4$  for any  $\sigma \in \mathbb{C} \setminus 0$ . Since  $(x - \sigma)(x + \sigma)$  has Newton quotient zero, we see that  $NQ(\sigma\mathbb{R}) = 0$  for any  $\sigma \in \mathbb{C} \setminus 0$ . The Newton quotient of  $(x - \sigma)^2$  is 4, and therefore  $NQ(S) \leq 4$  for any S.

**Proposition 22.41.** Suppose that S is a region of  $\mathbb{C}$  such that S and S<sup>-1</sup> are convex. If  $f(x) = \sum a_i x_i$  is in  $P^{S}(n)$  then

$$\left|\frac{a_k^2}{a_{k-1}a_{k+1}}\right| \ge \frac{NQ(8)}{4} \left(\frac{k+1}{k} \cdot \frac{n-k+1}{n-k}\right)$$
(22.9.1)

Proof. We follow the usual proof of Newton's inequalities. Write

$$f(x) = \sum_{k=0}^{n} b_k \binom{n}{k} x^k$$

where  $\binom{n}{i}b_i = a_i$ . The derivative of f satisfies

$$f'(x) = n \sum_{k=1}^{n} b_k \binom{n-1}{k-1} x^{k-1}$$

By Gauss-Lucas, the derivative is in  $\mathbf{P}^{\$}$  since \$ is convex. If we form the reverse of the polynomial,  $x^n f(1/x) = \sum {n \choose i} b_{n-i} x^i$  then the roots of the reverse are the inverses of the roots of f, and thus lie in  $\$^{-1}$ . Since  $\$^{-1}$  is convex, any derivatives of the reverse are in  $\mathbf{P}^{\$^{-1}}$ . Consequently, if we apply the derivative k times, reverse, apply the derivative n - k - 2 times, and reverse again, the resulting polynomial is in  $\mathbf{P}^{\$}$ :

$$\mathbf{n}\cdots(\mathbf{n}-\mathbf{k}+1)\left(\mathbf{b}_{\mathbf{k}}\binom{2}{0}+\mathbf{b}_{\mathbf{k}+1}\binom{2}{1}\mathbf{x}+\mathbf{b}_{\mathbf{k}+2}\binom{2}{2}\mathbf{x}^{2}\right)\in\mathbf{P}^{\mathbb{S}}$$
(22.9.2)

The Newton quotient of this quadratic equation satisfies

$$\frac{4 b_{k+1}^2}{b_k b_{k+2}} \geqslant NQ(S)$$

Substituting  $b_k = a_k / {n \choose k}$  and simplifying yields the conclusion.

In general it is hard to determine NQ(S). The next lemma gives a good lower bound for the Newton quotient that is often enough to determine NQ(S).

**Lemma 22.42.** If  $0 < \alpha < \beta < 2\pi$  and r, s > 0 then the Newton quotient for  $(x - re^{\iota\alpha})(x - se^{\iota\beta})$  is at least  $4\cos^2(\frac{\alpha - \beta}{2})$ .

*Proof.* The Newton quotient is invariant under multiplication by any complex number, so we multiply by  $(1/r)e^{-(\alpha+\beta)/2}$ . Thus we may assume that  $f = (x - e^{i\gamma})(x - te^{-i\gamma})$  where  $\gamma = (\beta - \alpha)/2$ . If t > 0 then the Newton quotient is

$$\left|\frac{(-e^{\iota\gamma}-t\,e^{-\iota\gamma})^2}{t}\right| = \frac{1}{t} + t + 2\cos(2\gamma)$$

This is minimized for t = 1, and has value  $4 \cos^2(\gamma)$ .

676

Sectors satisfy the conditions of the proposition.

**Corollary 22.43.** *If*  $0 < \alpha < \beta < \pi$  *and*  $\S$  *is the sector*  $\alpha \leq \arg z \leq \beta$  *then* 

$$\left|\frac{a_{k}^{2}}{a_{k-1}a_{k+1}}\right| \ge \cos^{2}(\frac{\alpha-\beta}{2})\left(\frac{k+1}{k} \cdot \frac{n-k+1}{n-k}\right)$$

*Proof.* If u, v are any two points in the sector, then the difference of their arguments is at most  $\beta - \alpha$ . From the lemma the Newton quotient is minimized when  $\alpha$  and  $\beta$  are as different as possible, so the minimum is attained at  $(\alpha - \beta)/2$ .

**Corollary 22.44.** If  $f(x) = \sum a_i x^i$  is a polynomial of degree n satisfying the condition that the real parts of all the roots have the same sign, and the imaginary parts of the roots all have the same sign then

$$\left|\frac{a_k^2}{a_{k-1}a_{k+1}}\right| \ge \frac{1}{2} \, \left(\frac{k+1}{k} \cdot \frac{n-k+1}{n-k}\right)$$

*Proof.* The region \$ is a quadrant. Thus  $\alpha - \beta = \pi/2$ , and  $4\cos^2(\frac{\alpha-\beta}{2}) = 2$ .

**Corollary 22.45.** If  $f(x) = \sum a_i x^i$  is a polynomial of degree n whose roots all have the same argument then

$$\left|\frac{a_{k}^{2}}{a_{k-1}a_{k+1}}\right| \geqslant \left(\frac{k+1}{k} \cdot \frac{n-k+1}{n-k}\right)$$

We can give a lower bound for more general regions.

**Corollary 22.46.** If S and S<sup>-1</sup> are convex regions, and if S is contained within a sector of angle width  $\alpha$  then for  $f \in \mathbf{P}^{S}$ 

$$\left|\frac{a_{k}^{2}}{a_{k-1}a_{k+1}}\right| \ge \cos^{2}(\alpha/2) \, \left(\frac{k+1}{k} \cdot \frac{n-k+1}{n-k}\right)$$

For an example of such a region, consider a circle of radius 1 around  $\sigma$  where  $|\sigma| > 1$ . Simple trigonometry shows that the cosine of one half the angle between the two tangents to the circle is  $\sqrt{|\sigma|^2 - 1}/|\sigma|$ . We conclude that for all f with roots in this circle we have

$$\left|\frac{a_k^2}{a_{k-1}a_{k+1}}\right| \geqslant \frac{|\sigma|^2 - 1}{|\sigma|^2} \, \left(\frac{k+1}{k} \, \frac{n-k+1}{n-k}\right)$$

If we restrict f to have all real coefficients then the results are simpler.

**Corollary 22.47.** Suppose that S and S<sup>-1</sup> are convex regions of  $\mathbb{C}$ . If  $f(x) = \sum a_i x_i$  is in  $P^S(n)$  then

$$\left|\frac{a_{k}^{2}}{a_{k-1}a_{k+1}}\right| \ge \inf_{z \in S \cap \overline{S}} \cos^{2}(\arg(z)) \left(\frac{k+1}{k} \frac{n-k+1}{n-k}\right)$$
(22.9.3)

**Corollary 22.48.** *If*  $0 \le \alpha \le \pi$  *and* \$ *is one of the two sectors* 

- 1.  $-\alpha \leqslant \arg z \leqslant \alpha$
- 2.  $\pi \alpha \leq \arg z \leq \pi + \alpha$

and f is a polynomial with all real coefficients in  ${\textbf P}^S(n)$  then

$$\left|\frac{a_{k}^{2}}{a_{k-1}a_{k+1}}\right| \geqslant \cos^{2}(\alpha) \ \left(\frac{k+1}{k} \ \frac{n-k+1}{n-k}\right)$$

CHAPTER



## **Equivariant Polynomials**

In this chapter we consider polynomials that are fixed under the actions of a cyclic group.

#### 23.1 Basic definitions and properties

A polynomial is *equivariant* if it is fixed under the action of a group. For polynomials in one variable the group we consider is  $\mathbb{Z}/n\mathbb{Z}$ , the integers mod n, for which we take the multiplicative generator  $\zeta = e^{2\pi i/n}$ . The action of a group element  $\zeta^k$  takes a polynomial f(z) to  $f(\zeta^k z)$ . Since  $\mathbb{Z}/n\mathbb{Z}$  is cyclic we only need to verify that  $f(z) = f(\zeta z)$  in order to know that f is equivariant.

When n is 1 then all polynomials are equivariant and we are left with the theory of polynomials in **P** and  $\mathbf{P}^{\pm}$ . In general, we want to consider equivariant polynomials that have the largest possible number of real roots. Unfortunately, this is not quite the definition because we must treat zero roots specially:

**Definition 23.1.** The set  $\mathbf{P}/\mathbf{Z}_n$  consists of all polynomials with real coefficients fixed under the action of  $\mathbb{Z}/n\mathbb{Z}$  such that if we write  $f(z) = z^s g(z)$  where  $g(0) \neq 0$  then g has degree nk and k real roots. We also define (where g is as in the preceding sentence)

 $\mathbf{P}^+ / \mathbf{Z}_n = \{ f \in \mathbf{P} / \mathbf{Z}_n | all real roots of g are positive \}$  $\mathbf{P}^- / \mathbf{Z}_n = \{ f \in \mathbf{P} / \mathbf{Z}_n | all real roots of g are negative \}$ 

We should make a few observations about this definition. First, if f(z) is an equivariant polynomial with a term  $cz^k$  then  $f(z) = f(\zeta z)$  implies that  $cz^k = c(\zeta z)^k$  which requires that  $\zeta^k = 1$ . Since  $\zeta$  is a primitive n-th root of unity, the non-zero terms of an equivariant polynomial have degree divisible by n. This shows that s in the definition is also divisible by n. Moreover, if g

in the definition has a real root r then it also has roots  $r\zeta, r\zeta^2, \ldots, r\zeta^{n-1}$ . If n is odd then these are all complex. Thus when n is odd the number of non-zero real roots of any equivariant polynomial is at most the degree of g divided by n. If n is even then the number of *positive* roots is at most the degree divided by n. In addition, if n is even then g has positive and negative roots, so both  $P^+/Z_n$  and  $P^-/Z_n$  are empty.

From the remarks above we see that if monic  $f \in \mathbf{P}/\mathbf{Z}_n$  has real roots (or positive roots if n is even)  $r_1, \ldots, r_k$  then we can write

$$\begin{split} f(z) &= z^{ns} \prod_{k} (z - r_k) (z - \zeta r_k) \cdots (z - \zeta^{n-1} r_k) \\ &= \tilde{f}(z^n) \end{split}$$

where

$$\tilde{\mathsf{f}}(\mathsf{x}) = z^{\mathsf{s}} \prod_{\mathsf{k}} (\mathsf{x} - \mathsf{r}^{\mathsf{n}}_{\mathsf{k}})$$

This gives us a useful representation for polynomials in  $\mathbf{P}/\mathbf{Z}_n$ . A polynomial f(z) is in  $\mathbf{P}/\mathbf{Z}_n$  iff there is a polynomial  $\tilde{f} \in \mathbf{P}$  (or  $\mathbf{P}^{alt}$  if n even) such that  $f(z) = \tilde{f}(z^n)$ . Since this correspondence is linear, we have a linear transformation

$$\widetilde{}: \mathbf{P} / \mathbf{Z}_n \longrightarrow \begin{cases} \mathbf{P}^{alt} & n \text{ even} \\ \mathbf{P} & n \text{ odd} \end{cases}$$

that is a bijection.

**Example 23.2.** Consider the case n = 2. Since  $P/\mathbb{Z}_2 = \{g(x^2) \mid g \in \overline{P^{alt}}\}$  if g has non-zero roots  $r_1, \ldots, r_m$  then  $g(x^2)$  has non-zero roots  $\pm \sqrt{r_i}$  and hence  $P/\mathbb{Z}_2 \subset P$ . The Hermite polynomials  $H_n$  are invariant under  $\mathbb{Z}_2$ . Corollary 7.44 shows that the map  $x^k \mapsto H_k(x)$  defines a linear transformation  $P/\mathbb{Z}_2 \longrightarrow P/\mathbb{Z}_2$ .

If n > 2 then  $P/Z_n$  is not contained in P.

It is at first surprising that the derivative does not preserve  $P/Z_n$ :

**Lemma 23.3.** If n is odd and  $f \in P/\mathbb{Z}_n$  then  $f' \notin P/\mathbb{Z}_n$  but  $f^{(n)} \in P/\mathbb{Z}_n$ . For any  $k \ge 0$  we have that  $z^k f^{(k)}(z)$  is in  $P/\mathbb{Z}_n$ .

*Proof.* From the definition

$$f'(\zeta z) = \lim_{\zeta \to 0} \frac{f(\zeta(z+h)) - f(\zeta z)}{\zeta h} = \zeta^{-1} f'(z)$$

This shows that f(z) is not equivariant, but that zf'(z) is equivariant. Similarly  $f^{(k)}(\zeta z) = \zeta^{-k}f^{(k)}(z)$  so that  $z^k f^{(k)}(z)$  is equivariant. It remains to show that f' has the right number of real roots.

From Rolle's theorem, if a polynomial p has s real roots, then the derivative has at least s - 1 real roots, and they interlace the real roots of p. It might be the case that p' has more roots than p, but we always have that if p has a roots, and p' has b roots then  $a \equiv b + 1 \pmod{2}$ .

Without loss of generality we may assume that  $\tilde{f}$  has all distinct roots, all non-zero. Assume that n is odd. If f has degree nm then  $\tilde{f}$  and f have m non-zero real roots. The derivative f' has either m - 1 or m - 2 non-zero real roots, and an n - 1 fold root at 0. The total number of roots of f' has opposite parity from m, so f' has exactly m - 1 non-zero roots. Since zf' is equivariant, we can write  $zf'(z) = g(z^n)$  where  $g \in \mathbf{P}$  and hence  $zf'(z) \in \mathbf{P}/\mathbf{Z}_n$ .

Next,  $f^{(2)}$  has at least m - 2 non-zero roots, and an n - 2 fold root at 0. The same argument as above shows that  $z^2 f^{(2)}$  has an n fold root at 0, and exactly m - 1 non-zero roots. Consequently  $z^2 f^{(2)}$  is in  $\mathbf{P}/\mathbf{Z}_n$ . Continuing,  $f^{(i)}$  has an n - i fold root at 0 and exactly m - 1 non-zero roots. As above,  $z^i f^{(i)}$  is in  $\mathbf{P}/\mathbf{Z}_n$ .

Since  $z^n f^{(n)}$  is in  $\mathbf{P} / \mathbf{Z}_n$  we can divide by  $z^n$  and thus  $f^{(n)}$  has exactly m-1 non-zero roots, and is also in  $\mathbf{P} / \mathbf{Z}_n$ .

Notice that although there is a bijection between  $P/Z_n$  and P some results are more natural in  $P/Z_n$ . For example, if  $f(x) = g(x^5)$  then  $f^{(5)}$  is in  $P/Z_5$ . In terms of g this says that if  $g \in P$  then

 $120 g'(x) + 15000 x g''(x) + 45000 x^2 g^{(3)}(x) + 25000 x^3 g^{(4)}(x) + 3125 x^4 g^{(5)}(x)$ 

is also in **P**.

There is a general principle at work here: properties of **P** extend to **P**/**Z**<sub>n</sub>, but we often have to replace x with  $z^n$ , or f' with  $f^{(n)}$ .

**Corollary 23.4.** If n is odd and k is a positive integer then the linear transformation  $T: x^j \mapsto (\underline{jn})_k x^j$  maps P to itself. In addition,  $x^j \mapsto {\binom{jn}{k}} x^j$  also maps P to itself. If n is even the domain of the maps is  $P^{\text{alt}}$ .

*Proof.* The map T is the composition

$$\mathbf{P} \xrightarrow{\tilde{\mathbf{x}}} \mathbf{P} / \mathbf{Z}_n \xrightarrow{z^k f^{(k)}} \mathbf{P} / \mathbf{Z}_n \xrightarrow{\tilde{\mathbf{x}}} \mathbf{P}$$

where  $\tilde{\star}$  is the map  $g(z) \mapsto g(z^n)$ . To verify this claim, we calculate the image of a monomial  $z^j$ :

$$z^{j} \xrightarrow{\tilde{x}} z^{jn} \xrightarrow{z^{k}f^{(k)}} z^{k} (\underline{jn})_{k} z^{jn-k} \xrightarrow{\tilde{z}} (\underline{jn})_{k} z^{j}$$

To derive the second part, divide by the constant k!.

#### 23.2 Interlacing

We make the usual definition of linear interlacing:

**Definition 23.5.** Two polynomials  $f, g \in P/\mathbb{Z}_n$  *interlace* if and only if  $f + \alpha g$  is in  $P/\mathbb{Z}_n$  for all real  $\alpha$ .

It is very easy to determine interlacing in  $\mathbf{P}/\mathbf{Z}_n$ .

**Lemma 23.6.** Suppose f, g are in  $P/Z_n$ . If f and g interlace then  $\tilde{f}$  and  $\tilde{g}$  interlace. The converse is true if n is odd.

*Proof.* Since the map  $f \mapsto \tilde{f}$  is linear it follows that if f, g interlace then  $\tilde{f}$ ,  $\tilde{g}$  interlace. Conversely, if  $\tilde{f}$  and  $\tilde{g}$  interlace then  $\tilde{f} + \alpha \tilde{g}$  is in **P**, and hence  $f + \alpha g = (\tilde{f} + \alpha \tilde{g})(z^n)$  is in **P**/**Z**<sub>n</sub> if n is odd.

If n is even then the converse is not true. For instance, if  $\tilde{f} = x - 1$  and  $\tilde{g} = x - 2$  then  $\tilde{f}(x^2)$  and  $\tilde{g}(x^2)$  do not interlace. However, the *positive* roots of  $\tilde{f}(x^2)$  and  $\tilde{g}(x^2)$  interlace.

**Corollary 23.7.** *If* f,  $g \in P/Z_n$  *and* f, g *interlace then the degrees of* f *and* g *differ by* -n, 0, *or* n.

*Proof.* The degrees of  $\tilde{f}$  and  $\tilde{g}$  differ by at most one.

**Corollary 23.8.** If  $f \in \mathbf{P}^+ / \mathbf{Z}_n$  then  $f \leq zf' \leq z^2 f^{(2)} \leq \cdots$ .

*Proof.* If we write  $z^{i}f^{(i)} = g_{i}(z^{n})$  then the proof of Lemma 23.3 shows that  $g_{i} \leq g_{i+1}$  and the Corollary follows.

Every linear transformation on **P** (or  $\mathbf{P}^{alt}$  is n even) induces a linear transformation on  $\mathbf{P}/\mathbf{Z}_n$ , and vice versa. So suppose that  $T: \mathbf{P} \longrightarrow \mathbf{P}$ . We define a linear transformation  $T_*$  by the composition



We can also use the diagram to define T if we are given  $T_*$ , so linear transformations on  $P/Z_n$  are not very interesting. For instance, we have

**Lemma 23.9.** If f,  $g \in P/Z_n$  then the Hadamard product f \* g is in  $P/Z_n$ .

*Proof.* Observe that  $(f * g)(z) = (\tilde{f} * \tilde{g})(z^n)$ .

## 23.3 Creating equivariant polynomials and the Hurwitz theorem

Whenever a group acts on functions we can construct equivariant functions by averaging over the group. In our case, we define the average of a polynomial over  $\mathbb{Z}/n\mathbb{Z}$  to be

$$\hat{f}(z) = \frac{1}{n} \sum_{k=0}^{n-1} f(\zeta^k z)$$
(23.3.1)

The average is obviously equivariant, and has a very simple interpretation in terms of coefficients. If  $f = \sum a_k x^k$  then

$$\hat{f}(z) = \frac{1}{n} \sum_{k} a_{k} \left( \sum_{j=0}^{n-1} \zeta^{jk} \right) z^{k}$$

and since  $\zeta$  is a root of unity

$$\sum_{j=0}^{n-1} \zeta^{jk} = \begin{cases} 0 & k \nmid n \\ n & k \mid n \end{cases}$$

This gives us our fundamental formula:

$$\hat{f}(z) = \sum_{k \equiv 0 \pmod{n}} a_k x^k \tag{23.3.2}$$

Since the even part of a polynomial is just the average of f over  $\mathbb{Z}/2\mathbb{Z}$  the Hurwitz theorem Theorem 7.64 can be restated:

The average over  $\mathbb{Z}/2\mathbb{Z}$  of a polynomial in  $\mathbf{P}^{\text{pos}}$  is in  $\mathbf{P}^{\text{alt}}$ .

More generally we have

**Theorem 23.10.** If n is odd and  $f \in P^{alt}$  then  $\hat{f} \in P^+ / Z_n$ .

*Proof.* A stronger result has been proved earlier. Here we give an alternate analytic proof of a part of Theorem 7.65. The key idea is to apply (7.4.2) since we can also express  $\hat{f}$  as g(xD)f where

$$g(z) = \frac{1}{n} \frac{\sin(\pi z)}{\sin(\pi z/n)}.$$

In order to see this, notice that the zeros of g are at the integers not divisible by n, and if k is an integer then g(kn) = 1. (If n is even then  $g(kn) = (-1)^k$ .)

If  $f \in \mathbf{P}^{alt}(m)$  then g(z) has  $m - \lfloor m/n \rfloor$  zeros in (0, m). By Corollary 1.15 it follows that g(xD)f has at least  $\lfloor m/n \rfloor$  zeros. However, if  $f = \sum a_i x^i$  then

$$g(x\mathsf{D})f = \sum g(i)a_ix^i = \sum_{n\mid i}a_ix^i = \hat{f}$$

Now  $\hat{f}$  has degree  $n\lfloor m/n \rfloor$  and hence has at most  $\lfloor m/n \rfloor$  real roots. Consequently  $\hat{f}$  has exactly  $\lfloor m/n \rfloor$  real roots, and is in  $P^+/Z_n$ .

**Corollary 23.11.** If  $f = \sum a_k x^k$  is in  $P^{\text{alt}}$  (or  $\widehat{P^{\text{alt}}}$ ) then  $\sum a_{nk} x^k$  is in  $P^{\text{alt}}$  (or  $\widehat{P^{\text{alt}}}$ ).

**Corollary 23.12.** If n is odd then  $\sum_{k=0}^{\infty} \frac{z^k}{(nk)!}$  is in  $\widehat{P^{\alpha lt}}$ .

**Corollary 23.13.** *If*  $f \leq g$  *in* P *then*  $\hat{f} \stackrel{+}{\sim} \hat{g}$ *. Proof.* The average is a linear transformation  $P^{alt} \longrightarrow P/Z_n$ .

#### 23.4 Hermite polynomials

The Hermite polynomials have analogs in  $\mathbf{P}/\mathbf{Z}_n$ . If we replace  $x^2$  by  $x^n$  in (7.8.1) we get polynomials  $H_k^n$  that are nearly equivariant.

$$\mathsf{H}^{n}_{k}(z) = (-1)^{k} e^{z^{n}} \left(\frac{\mathrm{d}}{\mathrm{d}z}\right)^{k} e^{-z^{n}}$$

From the definition we find the recurrence

$$\mathsf{H}_{k+1}^{\mathfrak{n}}(z) = \mathfrak{n} z^{\mathfrak{n}-1} \mathsf{H}_{k}^{\mathfrak{n}}(z) - \mathsf{H}_{k}^{\mathfrak{n}}(z)'$$

which shows that the degree of  $H_k^n$  is k(n-1). If we write  $z^k H_k^n(z) = g_k^n(z^n)$  then the degree of  $g_k^n$  is k and

$$g_{k+1}^n(x) = (nx+k)g_k^n(x) - nxg'(x)$$

From the recurrence we see that  $H_k^n$  satisfies  $H_k^n(\zeta z) = \zeta^{-k} H_k^n(z)$ . The recurrence also shows that all  $g_k^n$  are in  $\mathbf{P}^{alt}$ , and hence

**Lemma 23.14.** The polynomials  $z^k H_k^n$  are in  $P^+/Z_n$  (or  $P/Z_n$  if n is even). In particular,  $H_{nk}^n(z)$  is in  $P/Z_n$ .

We can also express  $H_k^n$  as a composition:

$$H_{k}^{n}(z) = (nz^{n-1} - D)^{k}(1)$$

CHAPTER



## Polynomials modulo an ideal

If S is a set of polynomials in variables  $\mathbf{x}$ , and  $\mathcal{I}$  is an ideal in  $\mathbb{R}[\mathbf{x}]$  (or  $\mathbb{C}[\mathbf{x}]$ ) then in this chapter we study S/J. There are two cases that we are particularly interested in:

- 1. **P**/{ $x^n$ }
- 2.  $U_2/\{y^n\}$

We want to find intrinsic descriptions of S/J as well as to resolve the usual questions of interlacing, coefficient inequalities, and linear transformations.

#### 24.1 Initial segments of P

An element of  $\mathbf{P}/\{x^n\}$  can be viewed as a polynomial  $f = \sum_{k=0}^{n-1} a_i x^i$  such that there is an extension to  $\mathbf{P}$ . By this we mean a polynomial  $\tilde{f}$  in  $\mathbf{P}$  such that the first n coefficients of  $\tilde{f}$  and f are equal. For instance,  $1 + 7x + 28x^2 \in \mathbf{P}/\{x^3\}$  since there is a polynomial, namely  $(1 + x)^7$ , that begins  $1 + 7x + 28x^2 + \cdots$ . We sometimes say that elements of  $\mathbf{P}/\{x^{n+1}\}$  are n-initial segments of  $\mathbf{P}$ . Note that  $\mathbf{P}(n-1) \subset \mathbf{P}/\{x^n\}$  since  $f \in \mathbf{P}(n-1)$  is its own extension.

Although  $\mathbf{P}/\{x^2\} = \{a + bx \mid a, b \in \mathbb{R}\}$  is trivial, even the quadratic case for  $\mathbf{P}^{pos}$  takes some effort.

**Lemma 24.1.**  $P^{\text{pos}}/\{x^3\} = \{a + bx + cx^2 \mid a, b, c > 0 \& b^2 > 2ac\}$ 

*Proof.* If  $f = \sum a_i x^i \in P^{pos}(n)$  then Newton's inequalities yield

$$\frac{a_1^2}{a_0a_2} \geqslant \frac{2}{1}\frac{n}{n-1} > 2$$

Conversely, assume that  $b^2 > 2ac$ . If we can find  $f = a + bx + cx^2 + \cdots \in \mathbf{P}^{pos}$  then if we divide by a, and replace x by xb/a we can find  $1 + x + Cx^2 + \cdots \in \mathbf{P}$ 

where 0 < C < 1/2. It therefore suffices to show that if 0 < C < 1/2 then we can find  $f = 1 + x + Cx^2 + \cdots \in \mathbf{P}^{pos}$ .

The quadratic equation shows that

$$1 + x + tx^2 \in \mathbf{P}$$
 for all  $0 < t < 1/4$ 

Squaring yields that

$$(1 + x + tx^2)^2 = 1 + 2x + (2t + 1)x^2 + \dots \in \mathbf{P}$$
 for all  $0 < t < 1/4$ 

and replacing 2x by x yields

$$1 + x + (1/4 + t/2)x^2 + \dots \in \mathbf{P}$$
 for all  $0 < t < 1/4$ 

If we let s = 1/4 + t/2 we have

$$1 + x + sx^2 + \dots \in \mathbf{P}$$
 for all  $0 < s < 3/8$ 

Continuing by induction, if we expand  $(1 + x + tx^2)^{2^n}$ , we get that

$$1 + x + sx^2 + \cdots \in \mathbf{P}$$
 for all  $0 < s < 1/2 - 2^{-n}$ 

Consequently, if we choose n so that  $2^{-n} < 1/2 - C$  then we can realize  $1 + x + Cx^2$  as a 2-initial segment.

The argument in the lemma shows that powers of  $1 + x + ax^2$  converge, after renormalization, to  $1 + x + x^2/2$ . Here's a more general result that shows why the exponential function appears.

**Lemma 24.2.** If  $f = 1 + a_1x + \dots + a_{n-1}x^{n-1} + x^n$  then

$$\lim_{n\to\infty} f(x/n)^n = e^{a_1 x}$$

*Proof.* If we write  $f = \prod (x + r_i)$  then  $\prod r_i = 1$  and  $\sum 1/r_i = a_1$ .

$$\lim_{m \to \infty} f(x/m)^m = \lim_{x \to \infty} \prod (x/m + r_i)^m$$
$$= \prod_1^n \lim_{m \to \infty} \left(\frac{x}{mr_i} + 1\right)^m = \prod_1^n e^{x/r_i} = e^{\alpha_1 x}$$

Not only do the powers converge to an exponential function, but the coefficients are bounded by the coefficients of  $e^x$ . In addition, Lemma 24.2 shows that there are polynomials in  $\mathbf{P}^{\text{pos}}/\{x^{n+1}\}$  whose coefficients are arbitrarily close to 1/i!.

**Lemma 24.3.** If 
$$1 + x + a_2x + \cdots + a_nx^n \in P^{pos}$$
 then  $0 < a_i < 1/i!$ .

*Proof.* Newton's inequalities give that  $1/2 > a_2$ . Assuming that the Lemma is true up to n - 1, we see that

$$\begin{split} a_{n-1}^2 &> \frac{n}{n-1} a_{n-2} a_n & \text{Newton} \\ a_n &< \frac{1}{(n-1)!^2} \frac{n-1}{n} \frac{1}{a_{n-2}} & \text{induction} \\ a_n &< 1/n! & \text{induction} \end{split}$$

If  $f = b_0 + b_1 x + \dots \in \mathbf{P}^{pos}/\{x^{n+1}\}$  then

$$\frac{1}{b_0} f\left(\frac{b_0 x}{b_1}\right) = 1 + x + \cdots$$

is also in  $\mathbf{P}^{\text{pos}}/\{x^{n+1}\}$  Thus, in order to describe  $\mathbf{P}^{\text{pos}}/\{x^{n+1}\}$  it suffices to determine which polynomials  $1 + x + a_2x^2 + \cdots + a_nx^n$  are in  $\mathbf{P}^{\text{pos}}/\{x^{n+1}\}$ . The lemma shows that the set of all such  $(a_2, \dots, a_n)$  is bounded. If  $1 + x + a_2x^2 + a_3x^3 \in \mathbf{P}^{\text{pos}}/\{x^4\}$  then we know several bounds:

$1/2 > \mathfrak{a}_2 > 0$	
$1/6 > \mathfrak{a}_3 > 0$	
$\mathfrak{a}_2^2 > (3/2)\mathfrak{a}_3$	Newton
$1 + a_3 \ge 2a_2$	(24.3.1)

Empirical evidence suggests that these are also sufficient.

#### **24.2** Initial segments of **P**<sub>2</sub>

We now introduce  $\mathbf{P}_2^{\text{pos}}/\{y^{n+1}\}$ . Its elements can be considered to be sums  $f_0(x) + \cdots + f_n(x)y^n$  arising from a polynomial in P<sub>2</sub>. Here are two examples of elements in  $\mathbb{P}_2/\{y^n\}$ .

• Let  $f \in \mathbf{P}^{pos}$ . The Taylor series for f(x + y) shows

$$f + y \frac{f'}{1!} + y^2 \frac{f''}{2!} + \dots + y^n \frac{f^{(n)}}{n!} \in \mathbb{P}_2/\{y^{n+1}\}$$

• If we choose  $f = \prod (x + a_i + b_i y)$  and set  $g = \prod (x + a_i)$  then

$$g+y \sum b_j \frac{g}{x+a_j} + y^2 \sum_{j \neq k} b_j b_k \frac{g}{(x+a_j)(x+a_k)} \quad \in \mathbb{P}_2/\{y^3\}$$

The case n = 2 is very familiar.

$$\begin{split} \mathbb{P}_2 / \{y^2\} &= \{f + y g \mid f \leq g \text{ in } \mathbf{P} \} \\ \mathbf{P}_2^{\text{pos}} / \{y^2\} &= \{f + y g \mid f \leq g \text{ in } \mathbf{P}^{\text{pos}} \} \end{split}$$

We think that the characterization for  $\mathbf{P}_2^{\mathrm{pos}}/\{y^3\}$  is similar to the result for  $\mathbf{P}_2^{\mathrm{pos}}/\{x^3\}$ . Namely,  $f_0+y$   $f_1+y^2$   $f_2$  is a 2-initial segment if and only if  $f_0 \leq f_1 \leq f_2$  and  $f_1^2 - 2f_0f_2 \geq 0$  for positive x. Necessity is easy since we just substitute for  $\alpha$  and apply the result for  $\mathbf{P}^{\mathrm{pos}}/\{x\}$ . The problem is existence.

#### 24.3 Coefficient inequalities

There are Newton's inequalities for  $\mathbf{P}^{\text{pos}}/\{x^n\}$ . If we choose  $f = \sum_0^{n-1} a_i x^i$  in  $\mathbf{P}^{\text{pos}}/\{x^n\}$  and f has an extension of degree r then Newton's inequalities tell us that

$$\frac{a_{k+1}^2}{a_k a_{k+2}} \geqslant \frac{k+1}{k} \cdot \frac{r-k+1}{r-k} \qquad \qquad 0 \leqslant k < n-2$$

Since we don't know r all we can say is that

$$\frac{a_{k+1}^2}{a_k a_{k+2}} > \frac{k+1}{k} \qquad \qquad 0 \leqslant k < n-2$$

In addition to these Newton inequalities, we have determinant inequalities for  $\mathbf{P}^{\text{pos}}/\{x^n\}$ . Any subdeterminant of (4.5.2) that only involves  $a_0$  up to  $a_{n-1}$  must be non-negative. For instance, if  $a_0 + a_1x + a_2x^2 + a_3x^3 \in \mathbf{P}^{\text{pos}}/\{x^4\}$  then

$$\begin{vmatrix} a_1 & a_2 & a_3 \\ a_0 & a_1 & a_2 \\ 0 & a_0 & a_1 \end{vmatrix} \geqslant 0$$

.

which gives the inequality

$$a_1^3 + a_0^2 a_3 \ge 2a_0 a_1 a_2 \tag{24.3.1}$$

If we are given  $a_0 + \cdots + a_n x^n \in \mathbf{P}^{pos}/\{x^{n+1}\}$  then there is an  $f \in \mathbf{P}^{pos}$  with  $f = \sum a_i x_i$ , but we don't know the degree m of f. We can also use Newton's inequalities to get lower bounds on m. First of all,

$$\frac{a_k^2}{a_{k-1}a_{k+1}} \geqslant \frac{k+1}{k} \left( 1 + \frac{1}{m-k} \right) \qquad \qquad \text{for } 1 \leqslant k < n$$

Solving for m yields

$$\mathfrak{m} \ge k + \frac{\mathfrak{m} + 1}{\mathfrak{m}} \left( \frac{\mathfrak{a}_k^2}{\mathfrak{a}_{k-1} \mathfrak{a}_{k+1}} - \frac{k+1}{k} \right)^{-1}$$

These inequalities hold for  $1 \le k < n$ . For example, if k = 1 the inequality is

$$m \ge 1 + 2\left(\frac{a_1^2}{a_0 a_2} - 2\right)^{-1}$$
 (24.3.2)

We see m increases as the Newton quotient gets closer to its minimum value. It's not surprising that the exponential series is the unique series where the Newton quotient equals its lower bound for all indices.

For an example, we claim that the smallest extension of  $1 + nx + {\binom{x}{2}}x^2$  has degree n. Indeed, if we compute the bound (24.3.2) we get exactly n.

*Remark* 24.4.  $\mathbf{P}/\{x^n\}$  isn't closed. If we have a sequence  $f_k \in \mathbf{P}/\{x^n\}$  such that  $f_k \to f$  then the extensions might not converge to a polynomial. For instance, (24.3.2) shows that  $1 + x + x^2/2 \notin \mathbf{P}/\{x^3\}$ . It is, however, the limit of  $(1 + x/n)^n$  in  $\mathbf{P}/\{x^3\}$ .

#### 24.4 There are many extensions

We show that polynomials in  $\mathbf{P}^{pos}$  and  $\mathbf{P}^{pos}_2$  have extensions of arbitrary degree.

**Lemma 24.5.** If  $f \in \mathbf{P}^{pos}/\{x^n\}$  has an extension of degree m, then f has extensions of degree m + k for k = 1, 2, ...

*Proof.* We observed earlier (p. 128) that if  $f \in \mathbf{P}(n)$  then there is a positive  $\alpha$  such that  $f + \alpha x^{n+1} \in \mathbf{P}$ . Using this fact, we establish the lemma by induction.

**Lemma 24.6.** If  $f \in \mathbf{P}_2^{pos}/\{y^n\}$  has an extension of degree m, then f has extensions of degree m + k for k = 1, 2, ...

*Proof.* If  $f \in \mathbf{P}_2^{\text{pos}}(n)$ , and g(x, y) consists of the terms of degree n then we will show that there is an  $\epsilon$  such that

$$F_{\epsilon}(x,y) = f(x,y) + \epsilon(x+y)g(x,y)$$

is in  $\mathbf{P}_2^{\text{pos}}(n+1)$ . Since the asymptotes of the graph of f are also asymptotes of the graph of  $F_{\varepsilon}$  for any  $\varepsilon$ , we see that there is an N such that if  $|\alpha| > N$  then  $F_{\varepsilon}(\alpha, y) \in \mathbf{P}$ . Also, for any  $\alpha$  we can, by the observation in the proof above, find an  $\varepsilon$  such that  $F_{\varepsilon}(\alpha, y) \in \mathbf{P}$ . Since [-N, N] is compact, we can find an  $\varepsilon$  that works for all points in this interval, and hence it works for all of  $\mathbb{R}$ .

A polynomial in  $\mathbf{P}(n)$  determines an element in  $\mathbf{P}/\{x^r\}$  for  $1 \leq r < n$ . However, it is not true that an element of  $\widehat{\mathbf{P}}$  determines a member of  $\mathbf{P}/\{x^r\}$ . For instance,  $e^x \in \widehat{\mathbf{P}}$ , but it is known [187] that no k-initial segment is in  $\mathbf{P}$  except for k = 1. Corollary 5.20 is an example of the other extreme. The polynomial

$$\sum_1^\infty x^k 2^{-k^2}$$

has the property that *all* initial segments are in  $\mathbf{P}^{pos}$ .

#### 24.5 Interlacing

We can define interlacing as usual. We say that f and g interlace, written  $f \sim g$ , in  $\mathbf{P}/\mathbb{J}$  if  $f + \alpha g \in \mathbf{P}/\mathbb{J}$  for all  $\alpha \in \mathbb{R}$ . If we restrict  $\alpha$  to positive values we write  $f \stackrel{+}{\sim} g$ .

Here is a simple result.

**Lemma 24.7.** Suppose  $a + bx + cx^2$  and  $A + Bx + Cx^2$  are in  $P^{\text{pos}}/\{x^3\}$ . Then,  $a + bx + cx^2 \stackrel{+}{\sim} A + Bx + Cx^2$  if and only if  $bB \ge aC + cA$ .

*Proof.*  $a + bx + cx^2 + \alpha(A + Bx + Cx^2)$  is in  $\mathbf{P}^{\text{pos}}/\{x^3\}$  if and only if

$$(b + \alpha B)^2 > 2(a + \alpha A)(c + \alpha C)$$

and this holds for all positive  $\alpha$  if and only if  $bB \ge aC + Ac$  holds.

It is not easy to show that two polynomials do or don't interlace. For example we show that x and  $(x + 1)^2$  do not interlace in  $\mathbf{P}/\{x^3\}$ . If they did then  $(x + 1)^2 + \alpha x \in \mathbf{P}/\{x^3\}$  for all  $\alpha$ . Newton's inequality says that

$$(2+\alpha)^2 \ge 2$$

and this doesn't hold for all  $\alpha$ , so they don't interlace. It *is* the case that for any positive t

$$x \stackrel{+}{\sim} (x + t)^2$$
 in **P**/{ $x^3$ }

since  $(2t + \alpha)^2 > 2t^2$  for  $\alpha \ge 0$ .

Since there is no concept of degree in  $\mathbf{P}/\{x^{n+1}\}$  there is no idea of the direction of interlacing, so we can not expect simple inequalities between interlacing polynomials. We do have

**Lemma 24.8.** Suppose  $\sum a_i x^i$  and  $\sum b_i x^i$  are interlacing polynomials in  $P/\{x^{n+1}\}$ . The following inequality holds for  $0 \le k < n-1$ :

$$\begin{split} \left( (k+2)a_{k}b_{k+2} + (k+2)a_{k+2}b_{k} - 2(k+1) \right) a_{k+1}b_{k+1} \\ \\ \leqslant 4(k+1)^{2} \bigg( a_{k+1}^{2} - \frac{k+2}{k+1}a_{k}a_{k+2} \bigg) \bigg( b_{k+1}^{2} - \frac{k+2}{k+1}b_{k}b_{k+2} \bigg) \end{split}$$

*Proof.* Apply Newton's inequality to  $\sum (a_k + \alpha b_k)x^k$ . The result is a quadratic in  $\alpha$ ; the inequality is the statement that the discriminant is non-positive.

#### 24.6 Linear transformations

A linear transformation  $\mathbf{P} \longrightarrow \mathbf{P}$  does not in general determine a linear transformation  $\mathbf{P}/\{x^n\} \longrightarrow \mathbf{P}/\{x^n\}$  since T applied to different extensions can give different results. We will see that Hadamard products do preserve  $\mathbf{P}/\{x^n\}$ , and there are some other interesting preserving transformations arising from  $\mathbf{P}_3^{\text{pos}}$ .

We recall an elementary result about ideals.

**Lemma 24.9.** If  $\mathfrak{I},\mathfrak{J}$  are ideals of  $\mathbb{R}[x]$  then a linear transformation  $\mathsf{T}$  on  $\mathbb{R}[x]$  such that  $\mathsf{T}(\mathfrak{I}) \subset \mathfrak{J}$  determines a map  $\mathsf{T}^* : \mathbf{P}/\mathfrak{I} \longrightarrow \mathbf{P}/\mathfrak{J}$ .

*Proof.* The point is that the map involves a choice, and the hypothesis makes the map well defined. We define  $T^*$  by



For example, the derivative map  $D : f \mapsto f'$  maps the ideal generated by  $x^n$  to the ideal generated by  $x^{n-1}$ . We therefore have

$$\mathsf{D}\colon \mathbf{P}/\{\mathbf{x}^n\}\longrightarrow \mathbf{P}/\{\mathbf{x}^{n-1}\}$$

More generally, we have

**Lemma 24.10.** If  $f \in \mathbf{P}(r)$  then  $f(\mathbf{D}): \mathbf{P}/\{x^n\} \longrightarrow \mathbf{P}/\{x^{n-r}\}$ .

P is closed under products, and  $\mathbb{J}\times\mathbb{J}\subset\mathbb{J}$  so the product induces a linear transformation

$$\mathbf{P}/\mathbb{J} \times \mathbf{P}/\mathbb{J} \longrightarrow \mathbf{P}/\mathbb{J}$$

If  $f = \sum a_i x^i$  and  $g = \sum b_i x^i$  then the matrix representation of

$$\mathbf{P}/\{\mathbf{x}^n\} \times \mathbf{P}/\{\mathbf{x}^n\} \longrightarrow \mathbf{P}/\{\mathbf{x}^n\}$$

is given by

$$\begin{pmatrix} b_0 \\ b_1 \\ \vdots \\ b_{n-1} \end{pmatrix} \mapsto \begin{pmatrix} a_0 & & \\ a_1 & a_0 & \\ \vdots & \vdots & \ddots & \\ a_{n-1} & a_{k-2} & \dots & a_0 \end{pmatrix} \begin{pmatrix} b_0 \\ b_1 \\ \vdots \\ b_{n-1} \end{pmatrix}$$

We get more interesting matrices if we use two variables. Consider

$$\overline{P}_2^{\text{pos}} \times P \xrightarrow{\text{multiplication}} \overline{P}_2^{\text{pos}} \xrightarrow{\text{extract coefficient}} P$$

If we take the coefficient of  $y^n$  then we get a map

$$\overline{\mathbf{P}}_{2}^{pos} \times \mathbf{P}/\{x^{n+1}\} \xrightarrow{\text{multiplication}} \overline{\mathbf{P}}_{2}^{pos}/\{x^{n+1}\} \xrightarrow{\text{coefficient of } y^{n}} \mathbf{P}/\{x^{n+1}\}$$

For example, take  $g(y) = \sum b_i y^i \in \mathbf{P}$  and  $f(x, y) = \sum a_{ij} x^i y^j \in \overline{\mathbf{P}}_2$ . Then  $fg \in \overline{\mathbf{P}}_2$ , and the coefficient of  $y^n$ , is in  $\mathbf{P}$ . The coefficient of  $x^j y^n$  is  $\sum_j b_i a_{j,n-i}$ .

This gives a matrix that preserves  $\mathbf{P}/\{x^{n+1}\}$ . For instance, if we consider 2-initial segments and take n = 2 then the mapping

$$\begin{pmatrix} b_0 \\ b_1 \\ b_2 \end{pmatrix} \mapsto \begin{pmatrix} a_{02} & a_{01} & a_{00} \\ a_{12} & a_{11} & a_{10} \\ a_{22} & a_{21} & a_{20} \end{pmatrix} \begin{pmatrix} b_0 \\ b_1 \\ b_2 \end{pmatrix}$$

maps  $\mathbf{P}/\{x^3\}$  to itself. Similarly, we can find matrices that preserve  $P_2/\{y^n\}$ . If we take  $f(x, y, z) \in P_3$  and  $g(x, z) \in \mathbf{P}_2$  then

**Theorem 24.11.** Suppose  $f \in \overline{P}_3$  where  $f = \sum f_{i,j}(x) y^i z^j$ . If we define

$$\mathsf{M} = (\mathsf{f}_{\mathsf{n}-\mathsf{i}-1,\mathsf{j}})_{0 \leqslant \mathsf{i},\mathsf{j} \leqslant \mathsf{n}-1}$$

then M maps  $P_2/\{y^n\}$  to itself.

Consider an example. Choose  $g\in {I\!\!P}^{pos}$  , and consider g(x+y+z) in Theorem 24.11. The matrix M is

$\left(\underline{g^{(2)}}\right)$	g′	<u>g</u>
2!0!	1!0!	0!0!
g <sup>(3)</sup>	<u>g</u> <sup>(2)</sup>	<u>g'</u>
2!1!	1!1!	0!1!
<u>g</u> <sup>(4)</sup>	$g^{(3)}$	<u>g</u> <sup>(2)</sup>
\2!2!	1!2!	0!2!/

We can use the theorem to find matrices M and vectors v such that  $M^k v$  consists of interlacing polynomials for k = 1, 2, ... For example, take

$$\begin{split} & \mathsf{f} = (\mathsf{x} + \mathsf{y} + 1)(\mathsf{x} + 2\mathsf{y} + 2)(2\mathsf{x} + \mathsf{y} + 1) \\ & = (2\mathsf{x}^3 + 7\mathsf{x}^2 + 7\mathsf{x} + 2) + (7\mathsf{x}^2 + 14\mathsf{x} + 6)\mathsf{y} + (7\mathsf{x} + 6)\mathsf{y}^2 + 2\mathsf{y}^3 \\ & \mathsf{v} = (2\mathsf{x}^3 + 7\mathsf{x}^2 + 7\mathsf{x} + 2, 7\mathsf{x}^2 + 14\mathsf{x} + 6, 7\mathsf{x} + 6, 2) \\ & \mathsf{g} = (\mathsf{x} + \mathsf{y} + \mathsf{z} + 1)^2(\mathsf{x} + 2\mathsf{y} + 3\mathsf{z} + 4)^2 \\ & \mathsf{M} = \begin{pmatrix} 6(4\mathsf{x} + 7) & 22\mathsf{x}^2 + 86\mathsf{x} + 73 & 2(\mathsf{x} + 1)(\mathsf{x} + 4)(4\mathsf{x} + 7) & (\mathsf{x} + 1)^2(\mathsf{x} + 4)^2 \\ 30 & 2(29\mathsf{x} + 53) & 2(17\mathsf{x}^2 + 70\mathsf{x} + 62) & 6(\mathsf{x} + 1)(\mathsf{x} + 2)(\mathsf{x} + 4) \\ 0 & 37 & 2(23\mathsf{x} + 44) & 13\mathsf{x}^2 + 56\mathsf{x} + 52 \\ 0 & 0 & 20 & 12(\mathsf{x} + 2) \end{pmatrix} \end{split}$$

Since  $M^k \nu \in \mathbb{P}_2/\{y^4\}$  for k = 1, ..., it follows that all vectors  $M^k \nu$  consist of interlacing polynomials. If we take  $g = (x + y + z + 1)^6$  then the matrix M is composed of multinomial coefficients:

$$\begin{pmatrix} \binom{6}{3,0} (x+1)^3 & \binom{6}{2,0} (x+1)^4 & \binom{6}{1,0} (x+1)^5 & \binom{6}{0,0} (x+1)^6 \\ \binom{6}{3,1} (x+1)^2 & \binom{6}{2,1} (x+1)^3 & \binom{6}{1,1} (x+1)^4 & \binom{6}{0,1} (x+1)^5 \\ \binom{6}{3,2} (x+1) & \binom{6}{2,2} (x+1)^2 & \binom{6}{1,2} (x+1)^3 & \binom{6}{0,2} (x+1)^4 \\ \binom{6}{3,3} & \binom{6}{2,3} (x+1) & \binom{6}{1,3} (x+1)^2 & \binom{6}{0,3} (x+1)^3 \end{pmatrix}$$

The Hadamard product defines a map  $P^{pos} \times P \longrightarrow P$ . Since  $f \times \cdot$  maps  $\{x^n\}$  to itself, we have a map

$$\mathbf{P}^{\text{pos}}/\{x^n\} \times \mathbf{P}/\{x^n\} \longrightarrow \mathbf{P}/\{x^n\}$$

The converse is slightly more complicated.

**Lemma 24.12.** Suppose that the Hadamard product with f maps the closure of  $P^{\text{pos}}/\{x^n\}$  to itself. Then  $\exp(f) \in P^{\text{pos}}/\{x^n\}$ .

*Proof.*  $e^x$  is in the closure of  $\mathbf{P}^{\text{pos}}/\{x^n\}$ .

Here is an interesting construction of matrices that preserve interlacing in  $\mathbf{P}^{\text{pos}}$ . Recall CP(M) is the characteristic polynomial of M, and M[1,2] is M with the first two rows and columns removed.

**Lemma 24.13.** If M is positive definite then the matrix below preserves interlacing in  $P^{\text{pos}}$ .

$$\begin{pmatrix} CP(M[1]) & CP(M) \\ CP(M[1,2]) & CP(M[2]) \end{pmatrix}$$

Proof. The determinant of

$$-xI + M + \begin{pmatrix} y & 0 & \dots \\ 0 & z & 0 & \dots \\ \vdots & \vdots & 0 \end{pmatrix}$$

equals

$$CP(M) + y CP(M[1]) + z CP(M[2]) + y z CP(M[1,2])$$

Now apply Theorem 24.11.

#### 24.7 Stable polynomials

Some stable cases are easy.

#### Lemma 24.14.

$$\begin{aligned} 1. \ \ &\mathcal{H}_{1}\!/\{x^{2}\} = \left\{a_{0} + a_{1}x + a_{2}x^{2} \mid a_{0}, a_{1}, a_{2} > 0\right\}\\ 2. \ \ &\mathcal{H}_{1}\!/\{x^{4}\} = \left\{a_{0} + a_{1}x + a_{2}x^{2} + a_{3}x^{3} \mid a_{0}, a_{1}, a_{2}, a_{3} > 0 \ \& \ a_{1}a_{2} > a_{0}a_{3}\right\}\\ 3. \ \ &\mathcal{H}_{1}\!/\{x^{5}\} \supset \left\{a_{0} + \dots + a_{4}x^{4} \mid a_{0}, \dots, a_{4} \geqslant 0 \ \& \ a_{1}a_{2}a_{3} \geqslant a_{1}^{2}a_{4} + a_{3}^{2}a_{0}\right\}\end{aligned}$$

*Proof.* Lemma 21.38 shows that if the inequalities are strict then the polynomials are already in  $\mathcal{H}_1$ . The quadratic is trivial, and we know from (21.7.1) that  $a_1a_2 > a_0a_3$  characterizes cubic stable polynomials. If  $f \in \mathcal{H}_1$  then by Lemma 21.38 we know that its 4-initial segment is in  $\mathcal{H}_1$ .

Example 24.15. We conjecture that

$$\mathcal{H}_{2}/\{y^{2}\} \stackrel{?}{=} \{f \underline{\lessdot} g \mid f, g \in \mathcal{H}_{1}\}$$

The left hand side is certainly contained in the right hand side; we give an example to show that certain elements of the right hand side are in the left.

We know that  $(x + 1)^2 \leq x + a$  in  $\mathcal{H}_1$  if and only if 0 < a < 2. Choose positive  $\alpha$  and consider the matrix

$$M = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + y \begin{pmatrix} e_1 & 0 \\ 0 & e_2 \end{pmatrix} + \begin{pmatrix} 0 & -\alpha \\ \alpha & 0 \end{pmatrix} + x \begin{pmatrix} 1 + \alpha^2 - \alpha\sqrt{1 + \alpha^2} & 0 \\ 0 & 1 + \alpha^2 + \alpha\sqrt{1 + \alpha^2} \end{pmatrix}$$

The determinant of M is

$$(1+\alpha^2)(1+x)^2 + y\left[(e_1+e_2) + x\left((1+\alpha^2)(e_1+e_2) + (e_1-e_2)\alpha\sqrt{1+\alpha^2}\right)\right] + \cdots$$

Thus, the first term is a constant multiple of  $(x + 1)^2$ , so we need to determine what the coefficient of y is. If we divide the coefficient of x in this coefficient by the constant term  $e_1 + e_2$  we get

$$1 + \alpha^2 + \frac{e_1 - e_2}{e_1 + e_2} \alpha \sqrt{1 + \alpha^2}$$

Now as  $\alpha$ ,  $e_1$ ,  $e_2$  range over the positive reals, the expression above takes all values in  $(1/2, \infty)$ , and therefore all linear terms x + a where 0 < a < 2 are realized by a determinant. Thus,  $(x + 1)^2 + y(x + a)$  lies in  $\mathcal{H}_1/\{y^2\}$ .

The next result is the 2  $\times$  2 analog of Theorem 24.11; the general case is similar.

**Lemma 24.16.** Suppose  $h(x, y) = \sum h_i(x)y^i$  is in  $\mathcal{H}_2$ . For appropriate n the matrix  $\begin{pmatrix} h_{n-1} & h_{n-2} \\ h_n & h_{n-1} \end{pmatrix}$  maps  $\mathcal{H}_2 / \{y^2\}$  to itself.

*Proof.* We can write  $F = f + gy + \cdots$  where  $F \in \mathcal{H}_2$ . The coefficients of  $y^n$  and  $y^{n-1}$  in hF are  $h_n f + h_{n-1}g$  and  $h_{n-1}f + h_{n-2}g$ . The result follows since adjacent coefficients interlace in  $\mathcal{H}_2$ .

#### 24.8 Other ideals

A representative element of  $\mathbf{P}_2^{\text{pos}}/\{x^2, y^2\}$  is a polynomial a + bx + cy + dxy. The characterization is very simple.

**Lemma 24.17.** 
$$\mathbf{P}_{2}^{pos}/\{x^{2}, y^{2}\} = \{a + bx + cy + dxy \mid a, b, c, d > 0 \& | a b c d | < 0\}$$

*Proof.* If  $f = a + bx + cy + dxy + \cdots$  is in P<sub>2</sub> then Theorem9.113 says that  $\begin{vmatrix} a & b \\ c & d \end{vmatrix} < 0$ . Conversely, assume the determinant is negative. By dividing by a and scaling x and y it suffices to show that  $1 + x + y + Dxy \in \mathbf{P}_2^{\text{pos}}/\{x^2, y^2\}$  where 0 < D < 1. Consider the product

$$\begin{aligned} (1+\alpha x+\beta y)(1+(1-\alpha)x+(1-\beta)y)\\ &=1+x+y+(\alpha(1-\beta)+\beta(1-\alpha))xy+\cdots \end{aligned}$$

Clearly since 0 < D < 1 we can find  $) < \alpha, \beta < 1$  so that  $(\alpha(1-\beta)+\beta(1-\alpha)) = D$ .

Although  $2xy - 1 \notin P_2$ , the product

$$(x + y + 1)(x + y - 1) = -1 + 2xy + \cdots$$

shows that  $2xy - 1 \in \mathbb{P}_2/\{x^2, y^2\}$ . A similar argument shows that

**Lemma 24.18.**  $P_2/\{x^2, y^2\} = \{a + bx + cy + dxy \mid \begin{vmatrix} a & b \\ c & d \end{vmatrix} \le 0\}$ 

The elements of  $\mathbf{P}_3^{\text{pos}}/\{y^2,z^2\}$  are interlacing squares with non-negative determinant

$$\begin{array}{cccc} \mathsf{f}_{00} & \underline{<} & \mathsf{f}_{10} \\ | \wedge & | \wedge \\ \mathsf{f}_{01} & \underline{<} & \mathsf{f}_{11} \end{array}$$

arising from  $\sum f_{ij}(x)y^i z^j \in \mathbf{P}_3^{pos}$ . See the next section.

**Lemma 24.19.** If  $f \leq g$ , h then  $f + yg + zh \in P_3/\{y^2, yz, z^2\}$ .

Proof. The argument similar to Lemma 9.103. If we write

$$\begin{split} f &= \prod (1+r_i\,x) \\ g &= \sum a_i\,f/(1+r_i\,x) \\ h &= \sum b_i\,f/(1+r_i\,x) \end{split}$$

then

$$\prod (1 + r_i x + a_i y + b_i z) = f + g y + h z + \sum_{i \neq j} a_i b_j \frac{f}{(1 + r_i x)(1 + r_j x)} yz + \cdots$$

**Lemma 24.20.**  $P_2/\{x - y\} \approx P$ 

*Proof.* A representative of  $\mathbb{P}_2/\{x - y\}$  is a polynomial of the form f(x, x) where  $f(x, y) \in \mathbb{P}_2$ . Now we know that  $f(x, x) \in \mathbf{P}$  so this gives a map  $\mathbb{P}_2/\{x - y\} \longrightarrow \mathbf{P}$ . Conversely, if  $g \in \mathbf{P}$  has positive leading coefficients and we define f(x, y) = g((x + y)/2) then  $f(x, y) \in \mathbb{P}_2$  and f(x, x) = g.

If we make a divisibility assumption then we get induced maps.

Lemma 24.21. Suppose that the linear transformation T satisfies

1. 
$$T(x^n)$$
 divides  $T(x^{n+1})$  for  $n \ge 1$ .  
2.  $T: \mathbf{P}^{\mathcal{R}} \longrightarrow \mathbf{P}^{\mathcal{R}}$   
3.  $T^{-1}: \mathbf{P}^{\mathcal{S}} \longrightarrow \mathbf{P}^{\mathcal{S}}$ 

then

$$\begin{array}{rcl} \mathsf{T} & : & \boldsymbol{P}^{\mathcal{R}}\!/\{\mathbf{x}^n\} & \longrightarrow & \boldsymbol{P}^{\mathcal{R}}\!/\{\mathsf{T}(\mathbf{x}^n)\}\\ \mathsf{T}^{-1} & : & \boldsymbol{P}^{\mathcal{S}}\!/\{\mathsf{T}(\mathbf{x}^n)\} & \longrightarrow & \boldsymbol{P}^{\mathcal{S}}\!/\{\mathbf{x}^n\} \end{array}$$

*Proof.* If  $g = \sum a_i x^i$  is any polynomial then we claim that there exist  $\alpha_i \in \mathbb{R}$  such that

$$T(gx^n) \in \{T(x^n)\}$$
 (24.8.1)

$$g T(x^n) = \sum \alpha_i T(x^{n+i})$$
(24.8.2)

For (24.8.1) we have

$$T(gx^n) = T(x^n) \sum a_i \frac{T(x^{n+i})}{T(x^n)}$$

which is of the form  $h(x) T(x^n)$  by the first hypothesis. For (24.8.2), using induction it suffices to show that it is true for g = x. If we write  $T(x^{n+1}) = (\alpha + \beta x)T(x^n)$  then

$$\mathbf{x} \mathsf{T}(\mathbf{x}^n) = \frac{1}{\beta} \mathsf{T}(\mathbf{x}^{n+1}) - \frac{\alpha}{\beta} \mathsf{T}(\mathbf{x}^n)$$

If  $f_0 \in \mathbf{P}^{\mathcal{R}}/\{x^n\}$  then write  $f_0 = f + g x^n$  where  $f \in \mathbf{P}^{\mathcal{R}}$ . Applying T

$$\mathsf{T}(\mathsf{f}_0) = \mathsf{T}(\mathsf{f}) + \mathsf{T}(\mathfrak{g}\,\mathsf{x}^n)$$

and using the facts that  $T(f) \in \mathbf{P}^{\mathcal{R}}$  and  $T(gx^n) \in \{T(x^n)\}$  shows that  $T(f_0)$  is in  $\mathbf{P}^{\mathcal{R}}/\{T(x^n)\}$ . If  $f_0 \in \mathbf{P}^{\mathcal{S}}/\{T(x^n)\}$  then write  $f_0 = f + gT(x^n)$  where  $f \in \mathbf{P}^{\mathcal{S}}$ . Applying  $T^{-1}$  yields

$$\mathsf{T}^{-1}(\mathsf{f}_0) = \mathsf{T}^{-1}(\mathsf{f}) + \mathsf{T}^{-1}(\mathsf{g}\,\mathsf{T}(\mathsf{x}^n))$$

Since  $T^{-1}(g T(x^n)) = x^n \sum \alpha_i x^i$  and  $T^{-1}(f) \in \mathbf{P}^{\delta}$  we see that  $T^{-1}(f_0)$  is in  $\mathbf{P}^{\delta} / \{x^n\}$ .

**Corollary 24.22.** Let  $T(x^n) = (\underline{x})_n$ . Then

$$\begin{array}{rcl} \mathsf{T} & : & \boldsymbol{P}^{\operatorname{alt}}\!/\{\mathbf{x}^{\mathfrak{n}}\} & \longrightarrow & \boldsymbol{P}^{\operatorname{alt}}\!/\{\mathsf{T}(\mathbf{x}^{\mathfrak{n}})\}\\ \mathsf{T}^{-1} & : & \boldsymbol{P}^{\operatorname{pos}}\!/\{\mathsf{T}(\mathbf{x}^{\mathfrak{n}})\} & \longrightarrow & \boldsymbol{P}^{\operatorname{pos}}\!/\{\mathbf{x}^{\mathfrak{n}}\} \end{array}$$

**Corollary 24.23.** *If*  $a + bx + cx^2 \in P^{pos}/\{x(x-1)(x-2)\}$  *then*  $(b+c)^2 > 2ac$ . *Proof.* If  $a + bx + cx^2 \in P^{pos}/\{x(x-1)(x-2)\}$  then and  $T(x^n) = (\underline{x})_n$  then

$$T^{-1}(a + bx + cx^2) = a + bx + c(x^2 + x) \in \mathbf{P}^{\text{pos}}/\{x^3\}$$

This implies that  $(b + c)^2 > 2ac$ .

#### 24.9 Coefficients of extensions

In this section we choose  $f\in \mathbb{S}/\mathbb{J}$  and a monomial  $x^I,$  and consider properties of the set

$$\left\{ \alpha \mid \alpha \text{ is the coefficient of } \mathbf{x}^{I} \text{ in } F \in S \text{ where } F \equiv f \pmod{\mathfrak{I}} \right\}$$

Our first elementary example is a linear polynomial. We take  $S = P^{pos}$ ,  $J = \{x^3\}$ ,  $x^I = x^3$ , and f = a + bx.

**Lemma 24.24.** *If* a, b > 0 *then* 

$$\left\{ \alpha \mid a + bx + \alpha x^2 \in \mathbf{P}^{\text{pos}}/\{x^3\} \right\} = \left( 0, b^2/(2\alpha) \right)$$

*Proof.* Use the fact that  $a + bx + \alpha x^2 \in \mathbf{P}^{pos}/\{x^3\}$  if and only if  $b^2 > 2\alpha\alpha$ .  $\Box$ 

The corresponding problem for  $P_2^{pos}$  is unsolved. We believe that if  $f+yg \in P_2^{pos}/\{y^2\}$  (equivalently  $f \leq g$ ) then

$$\left\{ h \left| f + yg + hy^2 \in \mathbf{P}_2^{\text{pos}} / \{y^3\} \right\} \stackrel{?}{=} \left\{ h \left| g \leq h \& \left| \frac{f}{g} \frac{g}{h} \right| \leqslant 0 \right\} \right\}$$

The conjectured solution for  $\mathbf{P}_3^{\text{pos}}$  is similar. If  $f + gy + hz \in \mathbf{P}_3^{\text{pos}}/\{y^2, z^2\}$  then

$$\left\{ k \left| f + yg + hz + kyz \in \mathbf{P}_{3}^{\text{pos}} / \{y^{2}, z^{2}\} \right\} \stackrel{?}{=} \left\{ k \left| g, h \leq k \& \left| f \atop h k \right| \leqslant 0 \right\}$$
(24.9.1)

We don't know how to approach these problems, but we can establish properties of the interlacing squares of the last question. Using these, we can establish (24.9.1) when the degree of f is two. So, we consider the new question

What are the extensions of three interlacing polynomials to an interlacing square with a positive determinant?

We are given three polynomials f, g, h in  $\mathbf{P}^{\text{pos}}$  with positive leading coefficients that satisfy  $f \leq g$ ,  $f \leq h$ , and consider the set  $\mathcal{K}$  of all polynomials k with positive leading coefficient satisfying

$$\begin{array}{l} f < g \\ \land \quad \land \\ h < k \end{array}$$
 (24.9.2)

$$\left| \begin{smallmatrix} f & g \\ h & k \end{smallmatrix} \right| > 0 \qquad \text{for } x \in \mathbb{R} \tag{24.9.3}$$

First of all,

**Lemma 24.25.** *X* is a non-empty bounded convex set of polynomials.

*Proof.* Lemma 24.19 shows  $\mathcal{K}$  is non-empty. Suppose that  $k_1$  and  $k_2$  satisfy (24.9.2) and (24.9.3). If  $0 < \alpha < 1$  then  $\alpha$  and  $1 - \alpha$  are positive, so

 $\begin{array}{cccc} f & < & g \\ \land & & \land \\ h & < & \alpha \, k_1 + (1 - \alpha) \, k_2 \end{array}$ 

From the determinant hypothesis:

$$\begin{split} & gh-fk_1>0\\ & gh-fk_2>0. \end{split}$$

Multiplying and adding yields

$$gh - f(\alpha k_1 + (1 - \alpha)k_2) > 0$$

Thus  $\alpha k_1 + (1 - \alpha)k_2$  satisfies the two conditions, and is in  $\mathcal{K}$ .

Since f(1) > 0 we see that g(1)h(1)/f(1) > k(1). Consequently, all coefficients of k are bounded by g(1)h(1)/f(1).

**Example 24.26.** If  $f = (x + 1)^n$ ,  $g = h = (x + 1)^{n-1}$  then the interlacing requirement implies that  $k = \alpha(x+1)^{n-2}$ . The determinant requirement implies that  $0 < \alpha < 1$ , and so

$$\mathcal{K} \subseteq \left\{ \alpha (x+1)^{n-2} \,\mid\, 0 < \alpha < 1 \right\}$$

In order to see that this is an equality, we proceed as follows. If  $a_i, b_i$  are non-negative then  $0 \leq (\sum a_i b_i) \leq (\sum a_i) (\sum b_j)$ . So, if  $0 \leq \alpha \leq 1$  choose  $a_i, b_i$  so that  $\sum a_i = \sum b_i = 1$  and  $\sum a_i b_i = 1 - \alpha$ . We have

$$\begin{split} & \prod_{1}^{n} (x+1+a_{i}y+b_{i}z) \\ & = (x+1)^{n} + y \, (x+1)^{n-1} + z \, (x+1)^{n-1} + y z (x+1)^{n-2} \biggl( \sum_{i\neq j} a_{i}b_{j} \biggr) \end{split}$$

This is the desired representation since

$$\sum_{i\neq j} a_i b_j = \left(\sum a_i\right) \left(\sum b_j\right) - \sum a_i b_i = 1 - (1 - \alpha) = \alpha.$$

**Example 24.27.** If  $F(x, y, z) \in \mathbf{P}_3^{pos}$  and  $F = \sum f_{ij}(x)y^i z^j$  then  $f_{11}$  is an extension of  $f_{00}, f_{10}, f_{01}$ . Thus, the polynomial  $f_{11} \in \mathcal{K}$ . In particular, if  $f(x) \in \mathbf{P}^{pos}$ , then  $f(x + y + z) \in \mathbf{P}_3^{pos}$ , so f''(x)/2 is an extension of f, f', f'.

In the case that f has degree 2,  $\mathcal{K}$  is an interval of positive numbers. We may assume that the polynomials are as given below, where all constants are positive:

$$\begin{array}{rcl} (x+r_1)(x+r_2) & < & t_1(x+r_1)+t_2(x+r_2) \\ & & & \land \\ s_1(x+r_1)+s_2(x+r_2) & < & & k \end{array} \tag{24.9.4}$$

The construction of the lemma suggests we consider the polynomial

 $(x + r_1 + s_1y + t_1z)(x + r_2 + s_2y + t_2z)$ 

It provides an extension where k is  $s_1t_2 + s_2t_1$ . Note that this value does not depend on the roots of f. Using these polynomials, we have

**Lemma 24.28.** The set  $\mathcal{K}$  of extensions of (24.9.4) is the interval

$$(s_1t_2 + s_2t_1 - 2\sqrt{s_1s_2t_1t_2}, s_1t_2 + s_2t_1 + 2\sqrt{s_1s_2t_1t_2}).$$

*Proof.* The polynomial gh - kf is a polynomial in x, and for it to be positive its discriminant must be negative. The discriminant is a quadratic in k whose roots are  $s_1t_2 + s_2t_1 \pm 2\sqrt{s_1s_2t_1t_2}$ .

An explanation of these conditions is that when n = 2 (24.9.2) and (24.9.3) are exactly what we need to extend to a polynomial in  $\mathbf{P}_3^{\text{pos}}$ .

**Lemma 24.29.** Assume that deg(f) = 2. The polynomials f, g, h, k in  $P^{pos}$  satisfy the conditions (24.9.2) and (24.9.3) above if and only if

$$f + gy + hz + yzk \in \mathbf{P}_3^{pos}/\{y^2, z^2\}$$

*Equivalently, if*  $f + gy + hz \in \mathbf{P}_3^{pos}/\{y^2, z^2\}$  and f has degree 2 then (24.9.1) holds.

*Proof.* We only need to show that if we are given f, g, h, k satisfying the conditions then there is such a polynomial in  $P_3$ . We start with the matrix

$$M = \begin{pmatrix} (r_1 + x + s_2 y + t_2 z)/r_1 & d_2 y + e_2 z \\ d_2 y + e_2 z & (r_2 + x + s_1 y + t_1 z)/r_2 \end{pmatrix}$$

and notice that  $r_1r_2 |M|$  is in P<sub>3</sub> and equals

 $f + gy + hz + (-2d_2e_2r_1r_2 + s_2t_1 + s_1t_2)yz + \cdots$ 

From the Lemma above we know that we can write

$$k = s_1 t_2 + s_2 t_1 + 2\beta$$
 where  $|\beta| \leq \sqrt{s_1 s_2 t_1 t_2}$ 

Since the matrix M is of the form  $I + xD_1 + yD_2 + zD_3$  we know that we must choose  $d_2$ ,  $e_2$  so that  $D_1$ ,  $D_2$ ,  $D_3$  are positive definite matrices. Now  $D_2 = \begin{pmatrix} s_2/r_1 & d_2 \\ d_2 & s_1/r_2 \end{pmatrix}$  and  $D_3 = \begin{pmatrix} t_2/r_1 & e_2 \\ e_2 & t_1/r_2 \end{pmatrix}$ . the determinant must be positive  $d_2$  and  $e_2$  must satisfy

$$s_1 s_2/(r_1 r_2) > d_2^2$$
  
 $t_1 t_2/(r_1 r_2) > e_2^2$ 

Notice that if d<sub>2</sub>, e<sub>2</sub> satisfy these inequalities then

$$\sqrt{s_1s_2t_1t_2} \geqslant r_1r_2d_2e_2$$

Consequently we can choose  $d_2$ ,  $e_2$  satisfying  $d_2e_2r_1r_2 = \beta$ .

**Corollary 24.30.** If f, g, h, k satisfy (24.9.2) and (24.9.3) and f has degree 2 then  $\begin{vmatrix} f & g \\ h & k \end{vmatrix}$  is a stable polynomial.

*Proof.* This follows from the lemma and Lemma 21.64.

*Remark* 24.31. If the determinant is always negative (rather than positive) then the corollary fails. The following polynomials form an interlacing square with everywhere negative determinant that is not stable.

Appendix

# Glossary of polynomials, spaces, and interlacing

#### Polynomials

An	Euler polynomial
C <sub>n</sub> <sup>α</sup>	Charlier polynomial with parameter $\alpha$
$C_n(x;y)$	$(-1)^{n}C_{n}^{y}(-x) = (-1)^{n}n!L_{n}^{-x-n}(y) = (-1)^{n}n!L_{n}(-y;-x-n)$
$G_n^{(\alpha)}$	Gegenbauer polynomial with parameter $\alpha$
H <sub>i</sub>	Hermite polynomial
HI	Hermite polynomial in d variables
L <sub>n</sub>	Laguerre polynomial
Ĺn	monic Laguerre polynomial: n!L <sub>n</sub>
L <sub>n</sub> <sup>α</sup>	Laguerre polynomial with parameter $\alpha$
$L_n(x;y)$	$L_n^y(-x)$
P <sub>n</sub>	Legendre polynomial
$Q_n^{\alpha,\beta}$	$\left(\left.\left(x-lpha ight)\mathbb{A}^{-1}-eta ight. ight)^{n}\left(1 ight) ight.$
T <sub>n</sub>	Chebyshev polynomial
$(\underline{\mathbf{x}})_{\mathbf{n}}$	falling factorial $x(x-1)\cdots(x-n+1)$
$\langle \underline{\mathbf{x}} \rangle_{\mathbf{n}}$	rising factorial $x(x + 1) \cdots (x + n - 1)$
$(x;q)_n$	$(1-x)(1-qx)\cdots(1-q^{n-1}x)$

Р	All polynomials in one variable with all real roots.
Ŷ	The analytic closure of <b>P</b> .
<b>P</b> <sup>pos</sup>	All polynomials in <b>P</b> with all negative roots and all positive (or all negative) coefficients: $P^{(-\infty,0)}$
$\widehat{\mathbf{P}^{pos}}$	The analytic closure of $\mathbf{P}^{pos}$ .
<b>P</b> <sup>alt</sup>	All polynomials in <b>P</b> with all positive roots and alternating coefficients: $\mathbf{P}^{(0,\infty)}$
$\widehat{\mathbf{P}^{alt}}$	The analytic closure of $\mathbf{P}^{alt}$ .
$\mathbf{P}^{\pm}$	$\mathbf{P}^{alt} \cup \mathbf{P}^{pos}$
$\mathbf{P}^{\mathrm{I}}$	All polynomials in <b>P</b> with all roots in an interval I.
P <sup>sep</sup>	Polynomials in <b>P</b> whose roots are all at least one apart.
$\widehat{\mathbf{P}^{sep}}$	The analytic closure of $\mathbf{P}^{sep}$ .
I	Polynomials $f + \iota g$ where $f < g$ .
$\overline{\mathfrak{I}}$	Polynomials $f + \iota g$ where $f \leq g$ .
$\widehat{\mathfrak{I}}$	Analytic closure of J.
$\mathcal{H}_1$	Stable polynomials with real coefficients
Н°	Interior of $\mathcal{H}_1$
$\widehat{\mathcal{H}}$	Analytic closure of $\mathcal{H}_1$
$\mathcal{H}_1(\mathbb{C})$	Stable polynomials with complex coefficients
$\widehat{\mathcal{H}_1}(\mathbb{C})$	Analytic closure of $\mathcal{H}_1(\mathbb{C})$

Sets of polynomials in one variable

### Sets of polynomials in several variables

U <sub>d</sub>	Polynomials in d variables non-vanishing in the upper half plane					
$\mathrm{U}_{d}(\mathbb{C})$	Polynomials in d variables and complex coefficients non- vanishing in the upper half plane					
$\mathcal{H}_d$	Polynomials in d variables non-vanishing in the right half plane					
${\mathfrak H}_d({\mathbb C})$	Polynomials in d variables with complex coefficients non- vanishing in the right half plane					
$P_d^+$	$\mathbb{U}_{d}(\mathbb{C})\cap\widetilde{\mathbb{H}}_{d}$					
P <sub>d</sub>	Polynomials in d variables satisfying degree conditions, substitution, and homogeneous part in $\mathbf{P}_{d-1}$					
<b>P</b> <sub>d</sub>	Polynomials in the closure of $P_d$					
$\widehat{\mathbf{P}}_{d}$	The analytic closure of P <sub>d</sub>					
$\mathbf{P}_{d}^{\text{pos}}$	Polynomials in $\mathbf{P}_{d}$ with all positive coefficients.					
$\overline{\mathbf{P}}_{d}^{\text{pos}}$	Polynomials in the closure of $\mathbf{P}_{d}^{pos}$					
$\widehat{\mathbf{P}^{\text{pos}}}_d$	The analytic closure of $\mathbf{P}_{d}^{\text{pos}}$					
$\mathcal{P}_{d,e}$	Polynomials satisfying substitution for $\mathbf{x} \in \mathbb{R}$ , $\mathbf{y} \ge 0$ , the degree condition, and homogeneous part in $\mathbf{P}_{d+e}$ .					
Ø <sub>d,e</sub>	Polynomials in the closure of $\mathcal{P}_{d,e}$ .					
$\widehat{\wp}_{\mathrm{d},e}$	The analytic closure of $\mathcal{P}_{d,e}$ .					
$Q_{d,e}$	Extension of $\mathbf{P}_{d}^{pos}$ by <i>e</i> variables subject to restricted substitution					
$\mathcal{H}_d$	stable polynomials with real coefficients in d variables					
$\widehat{\mathcal{H}_d}$	analytic closure of $\mathcal{H}_d$ in d variables					
$\mathbf{P}_2^{\text{gen}}$	Polynomials in $P_2$ whose solution curves are all distinct					
$\mathbf{P}_2^{\text{sep}}$	Polynomials in $P_2$ whose substitutions are all in $\mathbf{P}^{sep}$ .					

#### Kinds of interlacing

$f \xleftarrow{u} g$	$f + yg \in U_d(\mathbb{C})$
$f \stackrel{H}{\longleftarrow} g$	$f+yg\in\mathfrak{H}_d(\mathbb{C})$
$f \stackrel{P}{\longleftarrow} g$	$f + yg \in P_d^+$
$f\stackrel{+}{\sim} g$	$f + \alpha g \in \mathbf{P}$ for $\alpha \ge 0$
$f \stackrel{H}{\sim} g$	$f+\alpha g\in \mathfrak{H}_d \text{ for } \alpha \geqslant 0$
$f \stackrel{P}{\sim} g$	$f + \alpha g \in P_d^+ \text{ for } \alpha \geqslant 0$
f <u>∢</u> g	The roots of f and g alternate, and $deg(g) + 1 = deg(f)$
f <u>≪</u> g	The roots of f and g alternate, $deg(g) = deg(f)$ , and the largest root belongs to f
$f \longleftarrow g$	$f \leq g \text{ or } g \ll g \text{ in } \mathbf{P}$

## **Tables of transformations**

	f	T(f)	Domain	Range		
Multiplier transformations $\mathbf{P} \longrightarrow \mathbf{P}$						
Binomial	xn	$\binom{nd}{k} \chi^n$	Р	Р	681	d odd, $k = 1, 2$
Binomial	xn	$(\underline{nd})_{k} x^{n}$	Р	Р	681	d  odd, k = 1, 2
Derivative	xn	$nx^{n-1}$	Р	Р	25	
Exponential	xn	$x^n/n!$	Р	Р	192	
Exponential	x <sup>n</sup>	$x^n n!/(kn)!$	Р	Р	192	k = 1, 2,
Exponential	χn	$x^n/(kn)!$	Р	Р	192	k = 1, 2,
q-Exponential	xn	$x^n/[n]!$	Р	Р	170	q > 1
q-series	xn	$q^{\binom{n}{2}}x^n$	Р	Р	170	q  < 1
Rising factorial	x <sup>n</sup>	$\frac{\langle \underline{n} \rangle_i}{i!} x^i$	Р	Р	288	
Transformations	$\mathbf{P} \longrightarrow \mathbf{P}$	Р				
Derivative	f	f′	Р	Р	495	
Falling Factorial	$(\underline{\mathbf{x}})_{\mathbf{n}}$	$(\underline{\mathbf{x}})_{n+1}$	Р	Р	185	
Hermite	x <sup>n</sup>	H <sub>n</sub>	Р	Р	188	
Hermite	Hn	$H_{n+1}$	Р	Р	188	
Laguerre	x <sup>n</sup>	Ln	Р	Р	191	
Laguerre	x <sup>n</sup>	$L_n^{REV}$	Р	Р	496	
Laguerre	χn	L <sub>k</sub> <sup>n</sup>	Р	Р	496	$k = 1, 2 \dots$
Legendre	x <sup>n</sup>	$P_n/n!$	Р	Р	498	
Rising Factorial	$\left< \underline{x} \right>_n$	$\left< \underline{\mathbf{x}} \right>_{n+1}$	Р	Р	185	
Transformations	s P <sup>pos</sup> –	$\rightarrow \mathbf{P}$				
Affine	x <sup>n</sup>	$(-1)^{\binom{n}{2}}x^n$	<b>P</b> <sup>pos</sup>	Р	250	
Chebyshev	xn	$T_n^{REV}$	P <sup>pos</sup>	Р	203	
Laguerre	x <sup>n</sup>	$L_n^{\text{ Rev}}(-x)$	<b>P</b> <sup>pos</sup>	<b>P</b> <sup>pos</sup>	277	
Transformations	B P <sup>pos</sup> –	$\rightarrow \mathbf{P}^{pos}$				
Charlier	C <sub>n</sub> <sup>α</sup>	x <sup>n</sup>	<b>P</b> <sup>pos</sup>	<b>P</b> <sup>pos</sup>	189	?
Falling factorial	$(\underline{\mathbf{x}})_{\mathbf{n}}$	x <sup>n</sup>	<b>P</b> <sup>pos</sup>	<b>P</b> <sup>pos</sup>	182	
Falling factorial	$(\underline{x})_n$	$\langle \underline{\mathbf{x}} \rangle_{\mathbf{n}}$	<b>P</b> <sup>pos</sup>	$\mathbf{P}^{pos}$	183	
Hurwitz	χn	$\begin{cases} x^{n/2} & n \text{ even} \\ 0 & n \text{ odd} \end{cases}$	<b>P</b> <sup>pos</sup>	<b>P</b> <sup>pos</sup>	199	
<b>Rising</b> factorial	x <sup>n</sup>	$\left\langle \underline{\mathbf{x}} \right\rangle_{\mathrm{n}}$	<b>P</b> <sup>pos</sup>	<b>P</b> <sup>pos</sup>	183	

Rising Factorial  $x^n$   $\langle x \rangle /n!$   $\mathbf{P}^{pos}$   $\mathbf{P}^{pos}$  185
	f	T(f)	Domain	Range		
Transformations	$\mathbf{P}^{alt} \longrightarrow$	Р				
Affine	x <sup>n</sup>	$(-1)^{\binom{n}{2}}x^n$	$\mathbf{P}^{alt}$	Р	250	
Chebyshev	x <sup>n</sup>	$T_n^{REV}$	$\mathbf{P}^{alt}$	Р	203	
Hermite	x <sup>n</sup>	$H_nL_n$	$\mathbf{P}^{alt}$	Р	496	
Hermite	x <sup>n</sup>	$(H_n)^2$	$\mathbf{P}^{alt}$	Р	496	
Laguerre	x <sup>n</sup>	$(L_n)^2$	$\mathbf{P}^{alt}$	Р	496	
Transformations	$\mathbf{P}^{alt} \longrightarrow$	<b>P</b> <sup>alt</sup>				
Binomial	x <sup>n</sup>	$\binom{x}{n}$	$\mathbf{P}^{alt}$	$\mathbf{P}^{alt}$	194	
Binomial	xn	$\binom{nd}{k} x^n$	$\mathbf{P}^{alt}$	$\mathbf{P}^{alt}$	681	d even, $k = 1, 2 \dots$
Binomial	x <sup>n</sup>	$(\underline{nd})_k x^n$	$\mathbf{P}^{alt}$	$\mathbf{P}^{alt}$	681	d even, $k = 1, 2 \dots$
Charlier	xn	C <sub>n</sub> <sup>α</sup>	$\mathbf{P}^{alt}$	$\mathbf{P}^{alt}$	189	?
Falling factorial	xn	$(\underline{\mathbf{x}})_{\mathbf{n}}$	$\mathbf{P}^{alt}$	$\mathbf{P}^{alt}$	182	
Legendre	xn	$P_{\mathcal{M}}(x)/n!$	$\mathbf{P}^{alt}$	$\mathbf{P}^{alt}$	202	
Rising factorial	$\left< \underline{\mathbf{x}} \right>_{\mathbf{n}}$	x <sup>n</sup>	$\mathbf{P}^{alt}$	$\mathbf{P}^{alt}$	183	
Transformations	$\mathbf{S} \mathbf{P}^{\mathrm{I}} \longrightarrow \mathbf{P}$	Ι				
Charlier	x <sup>n</sup>	$^{\text{REV}}C_{n}^{\alpha}$	$\mathbf{P}^{(-1,0)}$	Р	442	
Chebyshev	x <sup>n</sup>	Tn	$\mathbf{P}^{(-1,1)}$	$\mathbf{P}^{(-1,1)}$	615	
Euler	x <sup>n</sup>	An	$\mathbf{P}^{(-1,0)}$	$\mathbf{P}^{alt}$	197	
Euler-Frobenius	x <sup>n</sup>	Pn	$\mathbf{P}^{(-1,1)}$	$\mathbf{P}^{(-1,1)}$	198	
Factorial	x <sup>n</sup>	$\prod_{1}^{n}(1-kx)$	$P^{(0,1)}$	$\mathbf{P}^{(0,1)}$	442	
Falling Factorial	x <sup>i</sup>	$(\underline{\mathbf{x}})_{i}^{Rev}$	$P^{(0,1)}$	$\mathbf{P}^{(0,1)}$	442	
Laguerre	x <sup>n</sup>	$_{\rm REV} \tilde{L}_n^{(\alpha)}$	$\mathbf{P}^{(0,1)}$	Р	442	$\alpha \geqslant -1$ ,
Q-series	x <sup>n</sup>	$(\mathbf{x};\mathbf{q})_{\mathbf{n}}$	$\mathbf{P}^{\mathbb{R}\setminus(0,1)}$	$\mathbf{P}^{\mathbb{R}\setminus(0,1)}$	244	0 < q < 1,
Q-series	x <sup>n</sup>	$(\mathbf{x};\mathbf{q})_{\mathbf{n}}$	$\mathbf{P}^{(0,1)}$	$\mathbf{P}^{(0,1)}$	244	1 < q,
Q-series	$(x;q)_n$	x <sup>n</sup>	$\mathbf{P}^{\mathbb{R}\setminus(0,1)}$	$\mathbf{P}^{\mathbb{R}\setminus(0,1)}$	244	1 < q,
Q-series	$(x;q)_n$	x <sup>n</sup>	$\mathbf{P}^{(0,1)}$	$\mathbf{P}^{(0,1)}$	244	0 < q < 1,
Rising Factorial	x <sup>n</sup>	$\left< \underline{x} \right>_n / n!$	${\bf P}^{(1,\infty)}$	$\mathbf{P}^{(1,\infty)}$	185	
Transformations	$\mathbf{P}(n)$ —	$\rightarrow \mathbf{P}(n)$				
Polar derivative	x <sup>i</sup>	$(n-i)x^i$	$\mathbf{P}(n)$	$\mathbf{P}(n-1)$	104	
Reversal	χ <sup>i</sup>	$x^{n-i}$	$\mathbf{P}(n)$	$\mathbf{P}(n)$	173	
Möbius	x <sup>i</sup>	$x^i(1-x)^{n-i}$	$\mathbf{P}(n)$	$\mathbf{P}(\mathbf{n})$	146	

	f	T(f)	Domain	Range							
Transformations	$\mathbf{P}^{\mathrm{I}}(\mathfrak{n})$ –	$\rightarrow \mathbf{P}^{\mathrm{I}}(\mathfrak{n})$									
Binomial	x <sup>i</sup>	$\binom{\alpha}{i} x^{i}$	$\mathbf{P}^{pos}(n)$	$\mathbf{P}^{pos}(n)$	171	$\alpha > n-2 \geqslant 0$					
Binomial	x <sup>i</sup>	$\begin{pmatrix} x+n-i \\ n \end{pmatrix}$	$\mathbf{P}^{(0,1)}(n)$	$\mathbf{P}^{alt}(n)$	194						
Chebyshev	x <sup>i</sup>	$(T_r)^i$	$\mathbf{P}^{(-1,1)}(\mathbf{n})$	$\mathbf{P}^{(-1,1)}(\mathbf{nr})$	168						
Factorials	$(\underline{\mathbf{x}})_{\mathbf{k}}$	$\langle \underline{\mathbf{x}} \rangle_{\mathbf{n}-\mathbf{k}}$	<b>P</b> <sup>pos</sup>	<b>P</b> <sup>pos</sup>	184						
Falling factorial	x <sup>i</sup>	$(\underline{\alpha})_{i} x^{i}$	$\mathbf{P}^{pos}(n)$	$\mathbf{P}^{pos}(n)$	171	$\alpha > n-2 \geqslant 0$					
Falling factorial	x <sup>i</sup>	$(\underline{x})_i x^{n-i}$	$\mathbf{P}^{pos}(n)$	$\mathbf{P}^{pos}(n)$	184						
Falling factorial	x <sup>i</sup>	$(\underline{\mathbf{x}+\mathbf{n}-\mathbf{i}})_{\mathbf{n}}\frac{\mathbf{x}^{\mathbf{i}}}{\mathbf{i}!}$	$\mathbf{P}^{pos}(n)$	$\mathbf{P}^{pos}(n)$	184						
Hermite	xi	$H_i H_{n-i}$	$\mathbf{P}^{pos}(n)$	$\mathbf{P}(n)$	277						
Hermite	x <sup>i</sup>	$x^{n-i}H_i$	$\mathbf{P}^{(2,\infty)}(\mathfrak{n})$	$\mathbf{P}(n)$	277						
Hermite	xi	$x^{n-i}H_i$	$\mathbf{P}^{pos}(n)$	$\mathbf{P}(n)$	277						
Laguerre	x <sup>i</sup>	$x^{n-i}L_i$	$\mathbf{P}^{pos}(n)$	$\mathbf{P}(n)$	277						
Laguerre	x <sup>i</sup>	$L_i \ L_{n-i}$	$\mathbf{P}^{pos}(n)$	$\mathbf{P}(n)$	277						
<b>Rising Factorial</b>	xi	$\langle \underline{x} \rangle_i x^{n-i}$	$\mathbf{P}^{(0,1)}(n)$	$\mathbf{P}^{pos}(n)$	437						
Rising factorial	x <sup>i</sup>	$\left< \underline{\mathbf{x}} \right>_{n-i}$	$\mathbf{P}^{pos}(n)$	$\mathbf{P}^{pos}(n)$	184						
Affine transformations with $Ax = x + 1$											
Binomial	x <sup>n</sup>	$(x+d-n)_d$	$\mathbf{P}^{alt}$	P <sup>sep</sup>	227	d = 1, 2,					
Charlier	x <sup>n</sup>	$C_n^{\alpha}$	$\mathbf{P}^{(-lpha,\infty)}$	$\mathbf{P}^{(-lpha,\infty)} \cap \mathbf{P}^{sep}$	228						
Derivative	f	f′	P <sup>sep</sup>	P <sup>sep</sup>	227						
Difference	f	$\Delta f$	P <sup>sep</sup>	P <sup>sep</sup>	227						
Falling Factorial	x <sup>n</sup>	$(\underline{\mathbf{x}})_{\mathbf{n}}$	$\mathbf{P}^{alt}$	$\mathbf{P}^{alt} \cap \mathbf{P}^{sep}$	228						
<b>Rising Factorial</b>	xn	$\langle \underline{\mathbf{x}} \rangle_{\mathbf{n}}$	<b>P</b> <sup>pos</sup>	$\mathbf{P}^{pos} \cap \mathbf{P}^{sep}$	228						
Transformations	of stabl	le polynomials									
Charlier	xn	C <sub>n</sub> <sup>α</sup>	$\mathcal{H}_1(\mathbb{C})$	$\mathcal{H}_1(\mathbb{C})$	646	$\alpha \in RHP$					
Charlier	x <sup>n</sup>	C <sub>n</sub> <sup>α</sup>	$\mathcal{H}_1$	$\mathcal{H}_1$	646	$\alpha \in RHP$					
Difference	f	f(x+1)-f(x)	$\mathcal{H}_1$	$\mathcal{H}_1$	648						
Hermite	x <sup>n</sup>	$\iota^n H_n(-\iota x)$	$\mathcal{H}_1$	$\mathcal{H}_1$	629						
Laguerre	xn	$\iota^n L_n(\iota x)$	$\mathcal{H}_1$	$\mathcal{H}_1$	629						
Rising factorial	x <sup>n</sup>	$\langle \underline{\mathbf{x}} \rangle_{\mathbf{n}}$	$\mathcal{H}_1(\mathbb{C})$	$\mathcal{H}_1(\mathbb{C})$	645						
Rising factorial	x <sup>n</sup>	$\langle \underline{\mathbf{x}} \rangle_{\mathbf{n}}$	$\mathcal{H}_1$	$\mathcal{H}_1$	645						
Falling factorial	$(\underline{\mathbf{x}})_{\mathbf{n}}$	x <sup>n</sup>	$\mathfrak{H}_1(\mathbb{C})$	$\mathfrak{H}_1(\mathbb{C})$	646						
Falling factorial	$(\underline{\mathbf{x}})_{\mathbf{n}}$	x <sup>n</sup>	$\mathcal{H}_1$	$\mathcal{H}_1$	646						
Hurwitz	x <sup>n</sup>	$\begin{cases} x^{n/2} & n \text{ even} \\ 0 & n \text{ odd} \end{cases}$	$\mathcal{H}_1$	<b>P</b> <sup>pos</sup>	646						

#### Transformations in two or more variables

Exponential	x <sup>i</sup> y <sup>j</sup>	$\frac{x^{i}y^{j}}{(i+j)!}$	$\mathbf{P}_2^{\mathrm{pos}}$	$\mathbf{P}_2^{\mathrm{pos}}$	413
Evaluation	$f\times x^I$	$f( I )x^{I}$	$\mathbf{P}^{pos} \times \mathbf{P}_{d}^{pos}$	$\mathbf{P}_{d}^{\text{pos}}$	349
Polar derivative	f	$(\mathbf{x} \cdot \mathbf{\partial})\mathbf{f}$	$\mathbf{P}_{d}^{\mathrm{pos}}$	$\mathbf{P}_{d}^{\text{pos}}$	348

# **Empirical tables of transformations**

In the table on the next page we list the results of some empirical testing. We constructed ten transformations for each polynomial family. We use the abbreviations The transformation did not appear to map any interval to **P**.

•			The transformation did not appear to map any interva
Х			The transformation appears to map $\mathbf{P}^{X} \longrightarrow \mathbf{P}$ .
$\mathbb{R}$			The transformation appears to map $\mathbf{P}^{(-\infty,\infty)} \longrightarrow \mathbf{P}$ .
$\mathbb{R}^+$			The transformation appears to map $\mathbf{P}^{(0,\infty)} \longrightarrow \mathbf{P}$ .
$\mathbb{R}^{-}$			The transformation appears to map $\mathbf{P}^{(-\infty,0)} \longrightarrow \mathbf{P}$ .
I			The transformation appears to map $\mathbf{P}^{(-1,1)} \longrightarrow \mathbf{P}$ .
$\mathbb{I}^+$			The transformation appears to map $\mathbf{P}^{(0,1)} \longrightarrow \mathbf{P}$ .
$\mathbb{I}^{-}$			The transformation appears to map $\mathbf{P}^{(-1,0)} \longrightarrow \mathbf{P}$ .
	1	1.	

An underlined entry such as (-1,1) means that the transformation is known to map  $\mathbf{P}^{(-1,1)} \longrightarrow \mathbf{P}$ . The other entries are unsolved questions.

The polynomial families in the first column are in the glossary, except for the last three. The Laurent-X polynomials are defined by  $p_{-1} = 0$ ,  $p_0 = 1$  and

Laurent	-1		$p_{n+1}$	$= \mathbf{x}(\mathbf{p}_{\mathbf{n}} - \mathbf{p}_{\mathbf{n}-1})$				
Laurent	-2		$\mathfrak{p}_{\mathfrak{n}+1}$	$= \mathbf{x}(\mathbf{p}_{\mathbf{n}} - \mathbf{n}\mathbf{p}_{\mathbf{n}-1})$				
Laurent	-3		$\mathfrak{p}_{\mathfrak{n}+1}$	$= (\mathbf{x} - \mathbf{n})\mathbf{p}_{\mathbf{n}} - \mathbf{n}\mathbf{x}\mathbf{p}_{\mathbf{n}-1}$				
There are some cavaets on a few entries.								
$P_n$	χn	$\mapsto$	$p_n x^{d-n}$	proved for $(-\infty, 0)$				
$\binom{x}{n}$	x <sup>n</sup>	$\mapsto$	$p_n$	only proved for each half				
(x; 1/2)	χn	$\mapsto$	$p_n/n!$	perhaps it's $(-\infty, 0) \cup (0, \infty)$				

The second table lists the actions of just a single transformation  $x^n \mapsto$  polynomial on various subsets of the complex plane.

EMPIRICAL TABLES OF TRANSFORMATIONS

	xn	pn	x <sup>n</sup>	xn	xn	pn	x <sup>n</sup>	x <sup>n</sup>	x <sup>n</sup>	xn
	$\downarrow \\ p_n$	$\downarrow_{x^n}$	$\downarrow p_n^{rev}$	$\downarrow p_n^2$	$\downarrow p_n x^n$	$\downarrow p_{d-n}$	$\downarrow p_n p_{d-n}$	$\downarrow p_n x^{d-n}$	$\downarrow p_n/n!$	↓ n!p <sub>n</sub>
H <sub>n</sub>	$\mathbb{R}_{\sim}$		$(-\infty, 1)$	$\mathbb{R}^+_{\!\!\!\sim\!\sim}$	$\mathbb{R}^+$	$\mathbb{R}^+$	$\mathbb{R}_{\sim}^{\pm}$	(	$\mathbb{R}$	I
Ln	$\mathbb{R}_{\sim}$		$\mathbb{R}_{\sim}$	$\mathbb{R}^+_{\!\!\!\sim\!\sim\!\!\!\sim}$	$\mathbb{R}^{-}$		$\mathbb{R}_{\sim}^{\pm}$	$\underline{\mathbb{R}^{-}} \cup \mathbb{R}^{+}$	$\mathbb{R}_{\sim}$	$(-\infty, 1)$
Pn	I		$\mathbb{R}^{\pm}$	$\mathbb{I}^+$	$\mathbb{I}^+$	$\mathbb{R}\setminus\mathbb{I}$	$\mathbb{R}^{\pm}$	$\mathbb{R}_{\times}^{\pm}$	$\mathbb{R}_{\sim}$	I
T <sub>n</sub>	$\mathbb{I}_{\!\!\!\!}$		$\mathbb{R}^{\pm}$	$\mathbb{I}^+$	$\mathbb{I}^+$	$\mathbb{R}^+ \cup \mathbb{R}^-$	$\mathbb{R}^{\pm}$	$\mathbb{R}^{\pm}$	$\overset{\mathbb{I}}{\sim}$	I
An	$\mathbb{I}_{\!\!\sim}^-$	$(1,\infty)$	I				$\mathbb{R}$	I	$\mathbb{R}^{-}$	$\mathbb{I}^-$
$C_n^0$	$\mathbb{R}^+_{\!$	$\mathbb{R}^{\sim}$	$\mathbb{I}_{\!\!\sim}^+$			$(1,\infty)$	$\mathbb{R}^+$	$\mathbb{I}^+$	$(-\infty, -1)$ $\cup \mathbb{R}^+$	$\mathbb{I}^+$
$G_n^{(2)}$	I		$(-\infty,1)$	$\mathbb{I}^+$	$\mathbb{I}^+$	$\mathbb{R}$	$\mathbb{R}^{-}$	$(-\infty, 1)$	$\mathbb{R}$	I
$(\underline{x})_n$	$\overset{\mathbb{R}^+}{\sim}$	$\mathbb{R}^{\sim}$	$\mathbb{I}_{\!\!\sim\!}^+$			$(1,\infty)$	$\mathbb{R}^+$	$\mathbb{I}^+_{\!\!\sim}$	$\underbrace{(-\infty,-1)}_{\cup \mathbb{R}^+}$	$\mathbb{I}^+$
$\langle \underline{x} \rangle_n$	$\mathbb{R}^{-}_{\sim}$	$\overset{\mathbb{R}^+}{\longrightarrow}$	$\mathbb{I}^+$			$(-\infty, -1)$	$\mathbb{R}^+$	$\mathbb{I}_{\!\!\sim}^+$	$\mathbb{R}^{-}_{(1,\infty)}$	$\mathbb{I}^-$
$\binom{x}{n}$	$\mathbb{R} \setminus \mathbb{I}^-$	$\mathbb{I}^-$	•	$(1,\infty)$		$\mathbb{I}^-$	$\mathbb{R}^{\pm}$	•	$(-\infty, -1)$	$\overset{\mathbb{R}^+}{\underset{\sim}{\longrightarrow}}$
									$\cup \mathbb{R}^+_{\!$	
(x;2)	$\mathbb{I}_{\!\!\sim}^+$	$\operatorname{Right}^+$	$\mathbb{R}^+$			$\mathbb{R}^{-}$	$\mathbb{R}^+$	$\mathbb{R}^+$		$\mathbb{I}^+$
(x;1/2)	$\mathbb{R} \setminus \mathbb{I}^+_{}$	$\overset{\mathbb{I}^+}{\sim}$		$(1,\infty)$			$\mathbb{R}^{\pm}$		$\mathbb{R}-\mathbb{I}^+$	$\mathbb{R}^{-}$
Exp	$\mathbb{R}_{\sim}$			$\mathbb{R}^+_{\!\!\!\sim\!\!\!\sim\!\!}$	$\overset{\mathbb{R}^+}{\sim}$				$\mathbb{R}_{\sim}$	$\mathbb{R}_{\sim}$
$2^{\binom{n}{2}}x^n$		$\mathbb{R}_{\sim}$				$\mathbb{R}$				
$(1/4)^{\binom{n}{2}} x^n$	$\mathbb{R}_{\sim}$			$\mathbb{R}^+$	$\mathbb{R}^+$				$\mathbb{R}_{\sim}$	$\mathbb{R}_{\sim}$
$(1/2)^{\binom{n}{2}}x^n$	$\mathbb{R}_{\sim}$			$\mathbb{R}^+_{\!$	$\overset{\mathbb{R}^+}{\sim}$				$\mathbb{R}$	
$(-1)^{\binom{n}{2}}x^{n}$ Laurent-1	$\overset{\mathbb{R}^{\pm}}{\mathbb{I}^{+}}$	$\overset{\mathbb{R}^{\pm}}{\underset{\cdot}{\overset{\cdot}{\overset{\cdot}}}}$	$\cdot$ $\mathbb{R} - \mathbb{I}^+$	$\mathbb{R}^+_{\sim}$		$\mathbb{R}$ .	$\cdot \\ \mathbb{R}^+ \cup \mathbb{R}^-$	$\mathbf{\mathbb{R}}-\mathbb{I}^+$	$\frac{\mathbb{R}^{\pm}}{\mathbb{R}^{+}}$	
Laurent-2	$\mathbb{R}^+_{\!\!\sim\!\!\sim}$	•	$\underline{\mathbb{R}^+} \cup \mathbb{R}^-$	•			$\mathbb{R}^+ \cup \mathbb{R}^-$	$\mathbb{R}^+ \cup \mathbb{R}^-$	$\mathbb{R}^+$	•
Laurent-3	$\mathbb{R}^+_{\sim}$		I			$\mathbb{I}^-$	$\mathbb{R}^+ \cup \mathbb{R}^-$	I	$\mathbb{R}^+$	$.\mathbb{I}^+$

$\chi^n\mapsto$	QI	$Q_{\mathrm{II}}$	$Q_{\mathrm{III}}$	$Q_{\mathrm{IV}}$	$\mathcal{H}_1$	$\mathcal{H}_1(\mathbb{C})$	I	Р	ιP
H <sub>n</sub>	-J	$-\Im$	I	I	Р		I	$\mathop{P}_{\sim}$	
L <sub>n</sub>	$Q_{\mathrm{II}}$	$Q_{\mathrm{II}}$	$Q_{\mathrm{III}}$	$Q_{\mathrm{III}}$	$\mathcal{H}_1$	$\mathfrak{H}_1(\mathbb{C})$	$Q_{\mathrm{III}}$	$\underset{\sim}{P}$	$\mathcal{H}_1(\mathbb{C})$
P <sub>n</sub>	$-\Im$	$-\Im$	I	I	•		I		
T <sub>n</sub>	$-\Im$	$-\Im$	I	I			I	•	
An	•				$\mathcal{H}_1$		•	•	
$G_n^{(2)}$	$-\Im$	$-\Im$	I	I			I		
$(\underline{\mathbf{x}})_{\mathbf{n}}$				•				•	•
$\langle \underline{\mathbf{x}} \rangle_{\mathbf{n}}$	•				$\mathcal{H}_1$		•	•	
$\binom{x}{n}$	•						•	•	
(x;2)	•				$-\mathcal{H}_1$		•	•	
(x;1/2)	•	$-\mathcal{H}_1(\mathbb{C})$	$-\mathcal{H}_1(\mathbb{C})$		$-\mathcal{H}_1$	$-\mathcal{H}_1(\mathbb{C})$	•	•	$-\mathfrak{H}_1(\mathbb{C})$
Exp	$\widetilde{\textbf{Q}}_{L}$	$\widecheck{\textbf{Q}}_{II}$	$\underline{Q}_{III}$	$\widetilde{Q}_{IV}$	$\underbrace{\mathcal{H}}_{1}$	$\underbrace{\mathcal{H}_1(\mathbb{C})}$	$\widetilde{\mathfrak{I}}$	$\mathop{\underline{P}}_{\sim}$	<b>₽</b>
$2^{\binom{n}{2}}x^n$									
$(1/4)^{\binom{n}{2}}x^n$	$Q_{\mathrm{I}}$	$Q_{\mathrm{II}}$	$Q_{\mathrm{III}}$	$Q_{\mathrm{I}V}$	$\mathcal{H}_1$	${\mathfrak H}_1({\mathbb C})$	I	Р	ιP
$(-1)^{\binom{n}{2}} x^n$					Р				
Laurent — 1	$-\Im$			I			•		
Laurent – 2	$-\Im$			I					
Laurent – 3	•	•	•			•			
	1								

# Tables of determinants and integrals

We assume that all polynomials in one variable have positive leading coefficients.

$egin{array}{c} \mathfrak{a}_k \ \mathfrak{a}_{k-1} \end{array}$	$\begin{vmatrix} a_{k+1} \\ a_k \end{vmatrix}$	< 0	where $\sum_{i=1}^{n} a_i x_i \in \mathbf{P}$ has all distinct roots. If they are not all distinct, then it is still < unless $a_k = 0$	106
a <sub>i</sub> b <sub>i</sub>	$\begin{vmatrix} a_{i+1} \\ b_{i+1} \end{vmatrix}$	< 0	and $a_{k-1}a_{k+1} = 0$ . where $f = \sum a_i x^i$ , $g = \sum b_i x^i$ , and $f \ll g$ .	115
f   f'	g g'	< 0	where $f < g$ or $f \ll g$ .	29
f  h	g k	< 0	where $deg(f) = deg(g) + 1 = deg(h) + 1 = deg(h) + 2$ and for all all $\alpha$ , $\beta$ , not both zero, we have that $\alpha f + \beta g < \alpha h + \beta k$	27
f  Af		< 0	for positive $x$ if $\mathbb{A}x > x$ and $f \in {I\!\!P}^{pos} \cap {I\!\!P}^{\mathbb{A}}$	219
f Af		> 0	for positive $x$ if $\mathbb{A} x < x$ and $f \in {I\!\!P}^{\text{pos}} \cap {I\!\!P}^{\mathbb{A}}$	219
$\begin{vmatrix} f \\ \Delta f \end{vmatrix}$	$\left. \begin{array}{c} \Delta f \\ \Delta^2 f \end{array} \right $	< 0	for positive x if $f\in P^{\text{pos}}\cap P^{sep}$	219
$\begin{vmatrix} f(x) \\ f(x+1) \end{vmatrix}$	$\left. \begin{array}{c} f(x+1) \\ f(x+2) \end{array} \right $	< 0	for positive x and $f \in \mathbf{P}^{pos} \cap \mathbf{P}^{sep}$	229
$\begin{vmatrix} d_{ij} \\ d_{i,j+1} \end{vmatrix}$	$ \begin{vmatrix} d_{i+1,j} \\ d_{i+1,j+1} \end{vmatrix} $	< 0	$f(x,y) = \sum d_{ij}x^iy^j$ is in P <sub>2</sub> and $f(x,0)$ has all distinct roots.	296
f   f <sub>y</sub>	$\left. \begin{array}{c} f_x \\ f_{xy} \end{array} \right $	≤ 0	$f \in P_d, d \ge 2.$	318
$\begin{vmatrix} f_{00} \\ f_{01} \end{vmatrix}$	$ \left. \begin{array}{c} f_{10} \\ f_{11} \end{array} \right  $	≪ 0	$\begin{split} &f\in \mathtt{P}_{d+2} \text{ and } f(\textbf{x}, y, z) = f_{00}(\textbf{x}) \!+\! f_{10}(\textbf{x}) y \!+\! f_{01}(\textbf{x}) z \!+\! f_{11}(\textbf{x}) y z \!+\! \cdots \end{split}$	318
$\begin{vmatrix} 1/f(x_1,y_1) \\ 1/f(x_2,y_1) \end{vmatrix}$	$\frac{1/f(x_2, y_1)}{1/f(x_2, y_2)}$	> 0	$f\in P_2$	458

Next, we have examples of three by three determinants, all of which are special cases of n by n determinants. All these matrices are totally positive for non-negative x.

$$\begin{vmatrix} a_{2} & a_{1} & a_{0} \\ a_{3} & a_{2} & a_{1} \\ a_{4} & a_{3} & a_{2} \end{vmatrix} \qquad \sum a_{i}x^{i} \in \mathbf{P}^{pos} \quad 113$$

$$\begin{vmatrix} f_{2} & f_{1} & f_{0} \\ f_{3} & f_{2} & f_{1} \\ f_{4} & f_{3} & f_{2} \end{vmatrix}$$

$$\begin{vmatrix} \frac{1}{2!}f'' & f' & f \\ \frac{1}{3!}f^{(3)} & \frac{1}{2!}f'' & f' \\ \frac{1}{4!}f^{(4)} & \frac{1}{3!}f^{(3)} & \frac{1}{2!}f'' \end{vmatrix} \qquad f \in \mathbf{P}^{pos} \quad 294$$

$$\begin{vmatrix} \frac{1}{2!}f(x+2) & f(x+1) & f(x) \\ \frac{1}{3!}f(x+3) & \frac{1}{2!}f(x+2) & f(x+1) \\ \frac{1}{4!}f(x+4) & \frac{1}{3!}f(x+3) & \frac{1}{2!}f(x+2) \end{vmatrix}$$

$$f \in \mathbf{P}^{pos} \quad 114$$

$$\frac{1}{2!}(x+2)^{n} & (x+1)^{n} & x^{n} \\ \frac{1}{3!}(x+3)^{n} & \frac{1}{2!}(x+2)^{n} & (x+1)^{n} \\ \frac{1}{4!}(x+4)^{n} & \frac{1}{3!}(x+3)^{n} & \frac{1}{2!}(x+2)^{n} \end{vmatrix}$$

$$114$$

$$\int_{-1}^{1} f(x + it) dt \in \mathbf{P}. \qquad f \in \mathbf{P} \qquad 669$$

$$\int_{0}^{1} f(x + it) dt \in \Im \qquad f \in \mathbf{P} \text{ and } \alpha > 0 \qquad 599$$

$$\int_{0}^{1} f(x + it) dt \in \Im \qquad f \in \mathbf{P} \text{ and } \alpha > 0 \qquad 608$$

$$\int_{0}^{1} f(x + it) dt \in \Im \qquad f \in \mathbf{P} \qquad 608$$

$$\int_{0}^{1} f(x + it) dt \in \mathcal{H}_{1} \qquad f \in \mathcal{H}_{1} \qquad 647$$

$$\int_{0}^{1} \phi(x) e^{itx} dx \in \widehat{\Im}. \qquad \phi(x) \uparrow \text{ and positive on } (0, 1) \qquad 616$$

$$\int_{0}^{1} f(x, y + it) dt \in \Im_{2} \qquad f \in \mathbf{P}_{2} \qquad 603$$

$$\int_{0}^{1} f(x, t) dt \in \mathbf{P} \qquad f \in \mathbf{P}_{2}^{sep} \qquad 75$$

$$\int_{0}^{1} f(x, t) g(x, -t) dt \in \mathbf{P} \qquad f, g \in \mathbf{P}_{2}^{sep} \qquad 237$$

# **Tables of Polynomials**

## Appell polynomials in one variable

 $A_n(x)$ 

 $\begin{array}{ll} A_0(x) &= 1 \\ A_1(x) &= 1-2\,x \\ A_2(x) &= 1-6\,x+6\,x^2 \\ A_3(x) &= 1-12\,x+30\,x^2-20\,x^3 \\ A_4(x) &= 1-20\,x+90\,x^2-140\,x^3+70\,x^4 \\ A_5(x) &= 1-30\,x+210\,x^2-560\,x^3+630\,x^4-252\,x^5 \end{array}$ 

# Appell polynomials in two variables

 $A_{n,m}(x)$ 

 $\begin{array}{lll} A_{0,0}(x,y) &= 1 \\ A_{1,0}(x,y) &= 1-2\,x-y \\ A_{0,1}(x,y) &= 1-x-2\,y \\ A_{2,0}(x,y) &= 1-6\,x+6\,x^2-2\,y+6\,x\,y+y^2 \\ A_{1,1}(x,y) &= 1-4\,x+3\,x^2-4\,y+8\,x\,y+3\,y^2 \\ A_{0,2}(x,y) &= 1-2\,x+x^2-6\,y+6\,x\,y+6\,y^2 \end{array}$ 

Charlier with parameter 1

 $C_n^1(\mathbf{x})$ 

 $\begin{array}{rll} C_0(x) = & 1 \\ C_1(x) = & -1+x \\ C_2(x) = & 1-3x+x^2 \\ C_3(x) = & -1+8x-6x^2+x^3 \\ C_4(x) = & 1-24x+29x^2-10x^3+x^4 \\ C_5(x) = & -1+89x-145x^2+75x^3-15x^4+x^5 \end{array}$ 

Charlier in two variables

$$C_n(x;y) = (-1)^n C_n^y(-x)$$

 $\begin{array}{rll} C_0(x;y) = & 1 \\ C_1(x;y) = & x+y \\ C_2(x;y) = & x+x^2+2xy+y^2 \\ C_3(x;y) = & 2x+3x^2+x^3+3xy+3x^2y+3xy^2+y^3 \\ C_4(x;y) = & 6x+11x^2+6x^3+x^4+8xy+12x^2y+4x^3y+6xy^2+6x^2y^2+4xy^3+y^4 \\ C_5(x;y) = & 24x+50x^2+35x^3+10x^4+x^5+30xy+55x^2y+30x^3y+5x^4y+ \\ & 20xy^2+30x^2y^2+10x^3y^2+10xy^3+10x^2y^3+5xy^4+y^5 \end{array}$ 

#### Chebyshev

 $T_n(\mathbf{x})$ 

 $\begin{array}{rcl} T_0(x) = & 1 \\ T_1(x) = & x \\ T_2(x) = & -1+2x^2 \\ T_3(x) = & -3x+4x^3 \\ T_4(x) = & 1-8x^2+8x^4 \\ T_5(x) = & 5x-20x^3+16x^5 \end{array}$ 

# Euler

 $A_n(x)$ 

 $\begin{array}{rll} A_0(x) = & 1 \\ A_1(x) = & x \\ A_2(x) = & x + x^2 \\ A_3(x) = & x + 4x^2 + x^3 \\ A_4(x) = & x + 11x^2 + 11x^3 + x^4 \\ A_5(x) = & x + 26x^2 + 66x^3 + 26x^4 + x^5 \end{array}$ 

#### **Falling factorial**

 $(\underline{\mathbf{x}})_{\mathbf{n}}$ 

 $\begin{array}{ll} (\underline{x})_0 &= 1 \\ (\underline{x})_1 &= x \\ (\underline{x})_2 &= -x + x^2 \\ (\underline{x})_3 &= 2 \ x - 3 \ x^2 + x^3 \\ (\underline{x})_4 &= -6 \ x + 11 \ x^2 - 6 \ x^3 + x^4 \\ (\underline{x})_5 &= 24 \ x - 50 \ x^2 + 35 \ x^3 - 10 \ x^4 + x^5 \end{array}$ 

Falling factorial inverse (Exponential or Bell polynomials)

 $B_n(\mathbf{x})$ 

 $\begin{array}{ll} B_0(x) &= 1 \\ B_1(x) &= x \\ B_2(x) &= x+x^2 \\ B_3(x) &= x+3 \ x^2+x^3 \\ B_4(x) &= x+7 \ x^2+6 \ x^3+x^4 \\ B_5(x) &= x+15 \ x^2+25 \ x^3+10 \ x^4+x^5 \end{array}$ 

## Gegenbauer

 $G_n(x)$ 

 $\begin{array}{ll} G_0(x) &= 1 \\ G_1(x) &= 4x \\ G_2(X) &= 12x^2-2 \\ G_3(x) &= 32x^3-12x \\ G_4(X) &= 80x^4-48x^2+3 \\ G_5(x) &= 192x^5-160x^3+24x \end{array}$ 

## Hermite

 $\mathsf{H}_{\mathfrak{n}}(x)$ 

 $\begin{array}{rll} H_0(x) = & 1 \\ H_1(x) = & 2x \\ H_2(x) = & -2 + 4x^2 \\ H_3(x) = & -12x + 8x^3 \\ H_4(x) = & 12 - 48x^2 + 16x^4 \\ H_5(x) = & 120x - 160x^3 + 32x^5 \end{array}$ 

#### Laguerre

 $L_{n}(\boldsymbol{x})$ 

$L_0(\mathbf{x}) =$	1
$L_1(\mathbf{x}) =$	1-x
$L_2(\boldsymbol{x}) =$	$\frac{1}{2}\left(2-4x+x^2 ight)$
$L_3(x) = \\$	$\tfrac{1}{6}\left(6-18x+9x^2-x^3\right)$
$L_4(x) = \\$	$\frac{1}{24}\left(24 - 96x + 72x^2 - 16x^3 + x^4\right)$
$L_5(x) = \\$	$\frac{1}{120} \left(120 - 600 x + 600 x^2 - 200 x^3 + 25 x^4 - x^5\right)$

# Q-Laguerre

 $L_n^{\mathbb{A}}(x)$ 

 $\begin{array}{rcl} L_0^{\mathbb{A}}(x) = & 1 \\ L_1^{\mathbb{A}}(x) = & -x+1 \\ L_2^{\mathbb{A}}(x) = & x^2-2[2]x+[2] \\ L_3^{\mathbb{A}}(x) = & -x^3+3[3]x^2-3[2][3]x+[2][3] \\ L_4^{\mathbb{A}}(x) = & x^4-4[4]x^3+6[3][4]x^2-4[2][3][4]x+[2][3][4] \end{array}$ 

#### Laguerre in two variables

$$\begin{split} L_n(x;y) &= L_n^y(-x) \\ L_0(x) &= 1 \\ L_1(x;y) &= 1+x+y \\ L_2(x;y) &= \frac{1}{2}\big(2+4\,x+x^2+3\,y+2\,x\,y+y^2\big) \\ L_3(x;y) &= \frac{1}{6}\big(6+18\,x+9\,x^2+x^3+11\,y+15\,x\,y+3\,x^2\,y+6\,y^2+3\,x\,y^2+y^3\big) \\ L_4(x;y) &= \frac{1}{24}\big(24+96\,x+72\,x^2+16\,x^3+x^4+50\,y+104\,x\,y+42\,x^2\,y+4\,x^3\,y+35\,y^2+36\,x\,y^2+6\,x^2\,y^2+10\,y^3+4\,x\,y^3+y^4\big) \\ L_5(x;y) &= \frac{1}{120}\big(120+600\,x+600\,x^2+200\,x^3+25\,x^4+x^5+274\,y+770\,x\,y+470\,x^2\,y+90\,x^3\,y+5\,x^4\,y+225\,y^2+355\,x\,y^2+120\,x^2\,y^2+10\,x^3\,y^2+85\,y^3+70\,x\,y^3+10\,x^2\,y^3+15\,y^4+5\,x\,y^4+y^5\big) \end{split}$$

# Legendre

```
P_n(x)
```

 $\begin{array}{rll} P_0(x) = & 1 \\ P_1(x) = & x \\ P_2(x) = & \frac{1}{2} \left( -1 + 3 x^2 \right) \\ P_3(x) = & \frac{1}{2} \left( -3 x + 5 \, x^3 \right) \\ P_4(x) = & \frac{1}{8} \left( 3 - 30 \, x^2 + 35 x^4 \right) \\ P_5(x) = & \frac{1}{8} \left( 15 x - 70 x^3 + 63 x^5 \right) \end{array}$ 

## Narayana

$$\begin{split} N_n(x) &= \ 1 \\ N_1(x) &= \ x \\ N_2(x) &= \ x^2 + x \\ N_3(x) &= \ x^3 + 3x^2 + x \\ N_4(x) &= \ x^4 + 6x^3 + 6x^2 + x \\ N_5(x) &= \ x^5 + 10x^4 + 20x^3 + 10x^2 + x \end{split}$$

# **Rising factorial**

 $\left< \underline{\mathbf{x}} \right>_{\mathbf{n}}$ 

 $\begin{array}{ll} \langle \underline{x} \rangle_{0} &= 1 \\ \langle \underline{x} \rangle_{1} &= x \\ \langle \underline{x} \rangle_{2} &= x + x^{2} \\ \langle \underline{x} \rangle_{3} &= 2 \ x + 3 \ x^{2} + x^{3} \\ \langle \underline{x} \rangle_{4} &= 6 \ x + 11 \ x^{2} + 6 \ x^{3} + x^{4} \\ \langle \underline{x} \rangle_{5} &= 24 \ x + 50 \ x^{2} + 35 \ x^{3} + 10 \ x^{4} + x^{5} \end{array}$ 

# **Rising factorial inverse**

$$\langle \underline{x} \rangle_n^{-1}$$

$$\begin{array}{ll} \langle \underline{x} \rangle_{0}^{-1} &= 1 \\ \langle \underline{x} \rangle_{1}^{-1} &= x \\ \langle \underline{x} \rangle_{2}^{-1} &= -x + x^{2} \\ \langle \underline{x} \rangle_{3}^{-1} &= x - 3 \ x^{2} + x^{3} \\ \langle \underline{x} \rangle_{4}^{-1} &= -x + 7 \ x^{2} - 6 \ x^{3} + x^{4} \\ \langle \underline{x} \rangle_{5}^{-1} &= x - 15 \ x^{2} + 25 \ x^{3} - 10 \ x^{4} + x^{5} \end{array}$$

# Questions

There are many unsolved problems about polynomials.

#### Polynomials with all real roots

*Question* 1. Is the minimum of the roots of the Hadamard product of polynomials in  $\mathbf{P}^{(0,1)}(n)$  attained at  $f = g = (x + 1)^n$ ?

*Question* 2. Show that for all n the falling factorial  $(\underline{x})_n$  is the derivative of a polynomial with all real roots. (See 164)

*Question* 3. Find all polynomials h(x,y) such that for all  $f \leq g$  we have  $h(f,g) \in \mathbf{P}$ .

*Question* 4. Show that  $f_n, g_n$  defined below satisfy  $f_n \leftarrow f_{n-1}$  and that  $g_n \leftarrow g_{n-1}$ , where  $\binom{n}{i,i}$  and  $\binom{n}{i,i,i}$  are multinomial coefficients. In addition,  $f_n < f_{n-2}$  and  $g_n < g_{n-3}$ . (There is the obvious generalization.) Corollary 13.15 shows that  $f_n \in \mathbf{P}$ .

$$f_n = \sum_{i=0}^n \binom{n}{i,i} x^i \qquad \qquad g_n = \sum_{i=0}^n \binom{n}{i,i,i} x^i$$

*Question* 5. Suppose n is a positive integer. Show the following interlacings, where B, L, N, E are the Bell, Laguerre, Narayana and Euler polynomials.

$$\begin{split} & exp^{-1} \, B_{n+1} \longleftarrow exp^{-1} \, B_n \\ & exp^{-1} \, L_{n+1} \longleftarrow exp^{-1} \, L_n \\ & exp^{-1} \, N_{n+1} \longleftarrow exp^{-1} \, N_n \\ & exp^{-1} \, E_{n+1} \longleftarrow exp^{-1} \, E_n \end{split}$$

*Question* 6. Under what conditions does  $0 \le a \le b$  imply there is a g such that  $poly(c) \le g$  for all  $a \le c \le b$ ?

*Question* 7. (C. Johnson) Given  $f, g \in \mathbf{P}$ , not necessarily interlacing, what is the maximum number of complex roots of linear combinations of f and g? Can this number be computed from the arrangement of the roots of f and g?

*Question* 8 (The Faà di Bruno problem - (See p. 39.)). Suppose that  $g \in \widehat{\mathbf{P}}$ . For any  $\alpha \in \mathbb{R}$  and m = 0, 1, ...,

- 1.  $F_m(\alpha, y)$  has all real roots.
- 2.  $F_{m+1}(\alpha, y) \leq F_m(\alpha, y)$ .

#### Polynomials with all positive coefficients

*Question* 9. If  $T: \mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}$ , then what constraints are there on the signs of  $T(x^i)$ ? The example  $x^i \mapsto (-1)^{\binom{i}{2}} x^i$  shows that the pattern -++ is possible. In addition to all signs the same, and all signs alternating, are there any other possibilities?

*Question* 10. If  $\sum a_i x^i \in \mathbf{P}^{pos}$  and  $\alpha > 1$  show that  $\sum a_i^{\alpha} x^i \in \mathbf{P}^{pos}$ . See [74] for a questionable argument. Compare Question 132.

*Question* 11. If  $f \in \mathbf{P}^{\text{pos}}$ , consider the set  $S = \{g \mid f \stackrel{+}{\sim} g\}$ . Unlike the case where  $\stackrel{+}{\sim}$  is replaced by  $\leq$ , the set S is not a cone. However, it is a union of a collection of cones. What is a usefful description of S?

#### Matrices that preserve interlacing

*Question* 12. If  $M = (f_{ij})$  is a matrix of polynomials with positive leading coefficients that preserves mutually interlacing polynomials then does  $(f'_{ij})$  also preserve mutual interlacing? If so, the following would be true:

Write  $M = \sum M_i x^i$  where each  $M_i$  is a matrix of constants. Each  $M_i$  is totally positive<sub>2</sub>.

*Question* 13. If  $(f_1, ..., f_n)$  is a sequence of mutually interlacing polynomials, then is  $\{(a_i) \in \mathbb{R}^n | \sum a_i f_i \in \mathbf{P}\}$  a convex set?

*Question* 14. Describe all matrices M of polynomials with the property that if v is a vector of mutually interlacing polynomials then the vector Mv also consists of mutually interlacing polynomials.

*Question* 15. A question related to the previous one. If v is fixed, describe all matrices M such that  $M^i v$  is mutually interlacing, for all positive integers i.

A special case. Find all  $M = \begin{pmatrix} f & g \\ h & k \end{pmatrix}$  such that  $M^i \begin{pmatrix} 1 \\ 1 \end{pmatrix}$  consists of interlacing polynomials for all positive integers i.

*Question* 16. If  $f \leq g$  and fF - gG = 1 where deg(F) = deg(G) - 1 = deg(f) - 2 then are F and G unique?

*Question* 17. Find all d by d matrices that are **P**-positive semi-definite. That is, if  $f \in \mathbf{P}$  is  $\sum a_i x^i$ , and  $A = (a_0, \dots, a_{d-1})$ , then  $AQA^t \ge 0$ .

Are all **P**-positive semi-definite matrices the sum of a positive semidefinite matrix and a **P**-positive semi-definite matrix whose non-zero entries are only on the anti-diagonal?

#### **Homogeneous Polynomials**

*Question* 18. Given  $f \in \mathbf{P}$ , are there  $a, b, c \in \mathbb{R}$  and  $g \in \mathbf{P}$  such that  $\left(a + b\frac{\partial}{\partial x} + c\frac{\partial}{\partial u}\right)g = f$ ?

*Question* 19. If T:  $\mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}^{\text{pos}}$  then when is  $\varphi(\mathsf{T})$  totally positive? It appears to be so for the Laguerre transformation.

#### Analytic functions

*Question* 20. Which hypergeometric functions are in  $\mathbf{P}$ ?

*Question* 21. Which polynomials f(x, y) have the property that  $f(x, \sin x) \in \widehat{\mathbf{P}}$ ? How about  $f(x, e^x) \in \widehat{\mathbf{P}}$ ?

*Question* 22. Is there a q-analog  $\Gamma_q$  of the gamma function for which  $\Gamma_q(z + 1)/\Gamma_q(kz + 1)$  is entire, and only has negative zeros?

*Question* 23. If  $f \in \widehat{\mathbf{P}}$ , and for all  $a \in \mathbb{R}$  we know that  $f + a \in \widehat{\mathbf{P}}$  then show that f = ax + b. This is trivial when f is a polynomial.

#### Linear Transformations of polynomials

*Question* 24. Characterize all linear transformations T (see (6.3.1)) defined by  $T(x^n) = \prod_{i=1}^{n} (x + a_i)$  that map  $\mathbf{P}^{\text{pos}}$  to  $\mathbf{P}$ .

Question 25. Can we characterize transformations T that satisfy

$$\begin{array}{rccc} \mathsf{T} \colon & \mathbf{P}^{\text{pos}} & \longrightarrow & \mathbf{P}^{\text{pos}} \\ \mathsf{T}^{-1} \colon & \mathbf{P}^{\text{alt}} & \longrightarrow & \mathbf{P}^{\text{alt}} \end{array}$$

*Question* 26. Can we characterize transformations T that satisfy  $T: \mathbf{P}^{alt} \longrightarrow \mathbf{P}^{pos}$  and  $T^2$  is the identity? The Laguerre transformation is such an example.

*Question* 27. Characterize all linear transformations  $T: \mathbf{P} \longrightarrow \mathbf{P}$  that commute with differentiation ( $\mathsf{DT} = \mathsf{TD}$ ).

*Question* 28. Suppose that V is the infinite dimensional vector space of all linear transformations from polynomials to polynomials, and W is the subset of V consisting of all linear transformations that map  $\mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}$ . Then W contains no open sets.

*Question* 29. If  $T(x^n) = d_n x^n$  satisfies  $T(\mathbf{P}^{pos}) \subset \mathbf{P}^{pos}$  then does T preserve interlacing?

*Question* 30. If  $T: \mathbf{P} \longrightarrow \mathbf{P}$  is a linear transformation then describe the possible sets  $S = \{ deg(T(x)), deg(T(x^2)), deg(T(x^3)), \cdots \}$ . Since  $T(x^n)$  and  $T(x^{n+1})$  interlace, we know that adjacent terms of S differ by at most 1. The example  $f \mapsto f(D)g$  shows that S can be bounded.

Question 31. Can we find linear transformations T such that

- 1. T:  $\mathbf{P} \longrightarrow \mathbf{P}$ .
- 2.  $T(x^i) = x^i$  for  $0 \leq i < r$
- 3.  $T(x^r) \neq x^r$

 $x^n \mapsto H_n(x)$  works for r = 2.

#### Linear transformations that preserve roots

Question 32. Show that

1. 
$$T(x^i) = \langle \underline{x} \rangle_i \langle \underline{x} \rangle_{n-i}$$
 maps  $\mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}^{\text{pos}}$ .

2.  $T(x^i) = \langle \underline{x} \rangle_i (\underline{x})_{n-i} \text{ maps } \mathbf{P}^{\text{pos}} \longrightarrow \mathbf{P}.$ 

Question 33. Lie and Wang [117] showed that

$$\begin{vmatrix} B_{n+1} & B_{n+2} \\ B_n & B_{n+1} \end{vmatrix}$$

has all positive coefficients where  $B_n$  are the Bell polynomials (p. 189). Show that the determinant is actually stable.

*Question* 34. Show that the linear transformation  $x^i \mapsto H_i(x)x^{n-i}$  maps  $\mathbf{P}^{(-\infty,2)}(n) \longrightarrow \mathbf{P}$ .

*Question* 35. If  $T(H_n) = x^n$ , then T does not map **P**,  $\mathbf{P}^{alt}$  or  $\mathbf{P}^{pos}$  to **P**. Show that EXP  $\circ$  T and T  $\circ$  EXP both map **P** to itself.

*Question* 36. Show that the transformation  $x^k \mapsto H_n^q$  where  $H_n^q$  is the q-Hermite polynomial maps  $\mathbf{P}^{(-1,1)} \longrightarrow \mathbf{P}$  for q > 1.

*Question* 37. Suppose  $r_i > 1 + r_{i-1}$  are positive constants. Show that the transformation  $x^n \mapsto \prod_{i=1}^n (x - r_i)$  maps  $\mathbf{P}^{(0,1)} \longrightarrow \mathbf{P}$ .

*Question* 38. Show that the transformation  $x^i \mapsto P_i^{REV}$ , the reverse of the Legendre polynomial, maps  $\mathbf{P}^{alt} \cup \mathbf{P}^{pos} \longrightarrow \mathbf{P}^{pos}$ . The transformation satisfies

$$\mathsf{T}(x^{n+1}) = \frac{2n+1}{n+1}\mathsf{T}(x^n) - \frac{x^2}{n+1}\mathsf{T}((x^n)')$$

*Question* 39. Show that the transformation  $x^i \mapsto P_i$ , where  $P_i(x)$  is the Legendre polynomial, maps  $\mathbf{P}^{(-1,1)} \longrightarrow \mathbf{P}^{(-1,1)}$ .

*Question* 40. For which  $\alpha$ ,  $\beta$  is it the case that  $T: \mathbf{P} \longrightarrow \mathbf{P}$ , where  $T: x^i \mapsto \frac{P_i^{\alpha,\beta}}{i!}$ , and  $P_i^{\alpha,\beta}(x)$  is the Jacobi polynomial?

*Question* 41. If T is the linear transformation  $x^n \mapsto T_n$  and  $Mz = \frac{z+1}{z-1}$  then  $T_M : \mathbf{P}^{(-1,1)} \to \mathbf{P}^{pos}$ . See Lemma 7.69.

*Question* 42. Show that the transformation  $E_i \mapsto x^i$ , where  $E_i$  is the Euler polynomial, maps  $\mathbf{P}^{(1,\infty)} \longrightarrow \mathbf{P}^{alt}$ .

*Question* 43. Show that the exponential diamond product  $f \bigotimes_{EXP} g = EXP^{-1}(EXPf \times EXPg)$  is a bilinear map  $\mathbf{P}^{pos} \times \mathbf{P}^{alt} \longrightarrow \mathbf{P}$ . This map is equivalent to  $x^i \bigotimes_{EXP} x^j = {i+j \choose i} x^{i+j}$ .

*Question* 44. Show that if  $T(x^i) = L_i(x)$ , the Laguerre polynomial, then the Laguerre diamond product  $f \bigotimes_T g = T^{-1}(T(f) \times T(g))$  is a bilinear map  $\mathbf{P}^{(1,\infty)} \times \mathbf{P}^{(-\infty,1)} \longrightarrow \mathbf{P}$ .

*Question* 45. If T:  $x^n \mapsto (\underline{x})_n$  then the diamond product  $f \bigotimes_T g = T^{-1}(T(f) \times T(g))$  maps  $\mathbf{P} \times \mathbf{P}^{pos} \longrightarrow \mathbf{P}$ . See Lemma 11.103.

*Question* 46. We can modify the definition of diamond product by replacing multiplication by the Hadamard product. Given a linear transformation T, define S by

$$\begin{array}{c} f \times g \longmapsto \xrightarrow{T \times T} & Tf \times Tg \\ s \overbrace{\downarrow}^{\overline{I}} & & & \downarrow^{\overline{I}} \\ T^{-1}(Tf * Tg) \xleftarrow{T^{-1}} & Tf * Tg \end{array}$$

- (A) Let  $T(x^n) = (\underline{x})_n$ . Show that  $S: \mathbf{P}^{\pm} \times \mathbf{P}^{\pm} \longrightarrow \mathbf{P}$ .
- (B) Show that the above result holds if we use the modified Hadamard product  $x^r *' x^r = r! x^r$ .
- (C) If  $T(H_i) = x^i$ , then show that

S: 
$$(\mathbf{P}^{(\infty,-1)} \cup \mathbf{P}^{(1,\infty)}) \times (\mathbf{P}^{(\infty,-1)} \cup \mathbf{P}^{(1,\infty)}) \longrightarrow \mathbf{P}$$

*Question* 47. If  $T(x^n) = P_m^{\alpha,\beta}$ , the n-th Jacobi polynomial, for which  $\alpha, \beta$  does T map  $\mathbf{P}^{-1,1}$  to itself? Some computations show that there are examples that do not map  $\mathbf{P}^{-1,1}$  to itself, but they are all where  $\alpha \neq \beta$  and  $-1 < \alpha, \beta < 0$ .

*Question* 48. Show that the transformation  $x^i \mapsto \prod (1 + (-1)^i x) = (x; -1)_i$  maps  $P^{\text{pos}} \cup P^{\alpha \text{lt}} \longrightarrow P$ .

*Question* 49. For which a, b does the transformation  $(x; a) \mapsto (x; b) \text{ map } \mathbf{P} \longrightarrow \mathbf{P}$ ?

*Question* 50. What assumptions on S,T do we need to make the following true?

If S, T are linear transformations that map  $\mathbf{P}^{\text{pos}}$  to itself, then define the transformation U on  $\mathbf{P}^{\text{pos}}(n)$  by  $x^i \mapsto T(x^i)S(x^{n-i})$ . If S, T map  $\mathbf{P}^{\text{alt}}$  to itself then U maps  $\mathbf{P}^{\text{alt}}(n)$  to itself and preserves interlacing.

*Question* 51. Determine all orthogonal polynomial families  $\{p_n\}$  so that  $x^n \mapsto p_n$  maps  $\mathbf{P}^{\mathbf{I}}$  to  $\mathbf{P}$  for some interval  $\mathbf{I}$ .

*Question* 52. Identify a polynomial f in P(n) with  $\alpha$ f in projective n + 1 space. What is

$$\frac{\text{Vol} DP(n+1)}{\text{Vol} P(n)}$$

*Question* 53. If T:  $\mathbf{P} \longrightarrow \mathbf{P}$ , then is it possible to compute  $\frac{Vol(T(\mathbf{P}))}{Vol(\mathbf{P})}$ ? What are upper and lower bounds for this ratio? Should we restrict ourself to  $\mathbf{P}(n)$ ?

*Question* 54. Let all denote all polynomials. If  $T: \mathbf{P} \longrightarrow \mathbf{P}$  and  $\frac{Vol(T(\mathbf{P}))}{Vol(\mathbf{P})}$  is small then  $\frac{Vol(T(all) \cap \mathbf{P})}{Vol(\mathbf{P})}$  is large.

*Question* 55. Suppose that T:  $\mathbf{P}(n) \longrightarrow \mathbf{P}(n)$ . What are the obstructions to extending T to a linear transformation that maps  $\mathbf{P}(n + 1)$  to itself?

*Question* 56. Show that if  $T(x^i) = (\underline{\alpha})_i x^i$  where  $\alpha < n - 2$  is not an integer then  $T((x + 1)^n) \notin \mathbf{P}$ . See Lemma 7.2.

*Question* 57. When is it true that if  $f \in \mathbf{P}^{\text{pos}}$ , then for all positive integers i

$$f(i)^2 > f(i-1)f(i+1).$$

Corollary 7.34 shows that the inequality with  $\ge$  in place of > is true. *Question* 58. Suppose that  $p_i$  is a sequence of polynomials defined by

$$p_0 = 1$$
  

$$p_1 = x$$
  

$$p_{n+1} = xp_n - c_n p_{n-1}$$

where the  $c_n$  are constants. If the linear transformation  $x^n \mapsto p_n$  maps **P** to itself then are the  $p_i$  essentially the Hermite polynomials?

*Question* 59. If  $T(x^k) = \frac{\langle \underline{x} \rangle_k}{k!}$  then show that  $T: \mathbf{P}^{\mathbb{R} \setminus \{0,1\}} \longrightarrow \mathbf{P}^{\mathbb{R} \setminus \{0,1\}}$ .

*Question* 60. If  $T(x^k) = \frac{\langle x \rangle_k}{k!}$  then show that for any  $\alpha \in (0, 1)$  there is an n such that  $T(x - \alpha)^n$  does not have all real roots.

*Question* 61. If T is a linear transformation, and  $T(x - r)^n \in \mathbf{P}$  for all positive integers n and all r satisfying  $a \leq r \leq b$ , then is it true that  $T: \mathbf{P}^{(a,b)} \longrightarrow \mathbf{P}$ ?

*Question* 62. Find examples of transformations T and constant a for which  $T(x + a)^n$  is a scalar, for all n. This is satisfied by  $T(x^i) = (x + n - i)_n$ 

*Question* 63. If T:  $x^n \mapsto \langle \underline{x} \rangle_n / n!$  then show that T maps polynomials whose roots all have real part 1/2 to polynomials with real part 1/2. [142]

*Question* 64. In Example 1.85 we saw that the commutator of the two transformations  $x^n \mapsto N_n$  and  $x_n \mapsto Dx^n$  maps the sequence  $\{x^n\}$  to an interlacing sequence. It appears that more is true. An entry in Table 64, such as **P** at the intersection of the row  $A_n$  and column  $B_n$  means that empirical evidence suggests that

$$T_1: x^n \mapsto A_n$$
  $T_2: x^n \mapsto B_n$   $\Longrightarrow$   $T_1T_2 - T_2T_1: \mathbf{P}^{pos} \longrightarrow \mathbf{P}$ 

#### **Affine Transformations**

*Question* 65. Show that if  $f(x) \leq f(x+2)$  then

$$\left| \begin{array}{c} f(x+1) & f(x) \\ f(x+2) & f(x+1) \end{array} \right| > 0$$

	$N_n$	An	Bn	D	Ln	$\langle \underline{\mathbf{x}} \rangle_{\mathbf{n}}$
Nn		Р	Р	$\mathbf{P}^{pos}$	<b>P</b> <sup>pos</sup>	
An			Р	$\mathbf{P}^{pos}$	P <sup>pos</sup>	•
Bn				${\mathcal H}$	$\mathbf{P}^{pos}$	Р
D					$\mathbf{P}^{(-\infty,0]}$	P <sup>pos</sup>
$L_n$						$\mathbf{P}^{pos}$

Table 1: Purported range of **P**<sup>pos</sup> under the commutator map

*Question* 66. If  $T(x^n)$  is  $H_n(x), x^n/n!$ , or  $L_n(x)$  then show that the minimum separation is achieved at the falling factorial. That is, if  $f \in \mathbf{P}^{sep}$  show that

$$\delta T(f) \ge \delta T(\underline{x})_n$$

*Question* 67. Is  $(\underline{x})_i \mapsto (\underline{x})_{n-i}$  a bijection on  $\mathbf{P}^{sep}(n)$ ?

*Question* 68. Show that  $x^n \mapsto L_n$  ( $L_n$  is the Laguerre polynomial) maps  $\mathbf{P}^{sep}$  to itself.

*Question* 69. Show that  $\langle \underline{x} \rangle_n \mapsto x^n$  maps  $\mathbf{P}^{\text{alt}}$  to  $\mathbf{P}^{\text{sep}}$ .

*Question* 70. Suppose all r<sub>i</sub> are positive and define

$$\mathsf{T}(x^n) = \prod_{i=1}^n (x - r_i)$$

If  $T(x^n) \in \mathbf{P}^{sep}$  then  $T: \mathbf{P}^{(0,1)} \longrightarrow \mathbf{P}^{(1,\infty)} \cap \mathbf{P}^{sep}$ .

*Question* 71. If T:  $\mathbf{P}^{sep} \longrightarrow \mathbf{P}^{sep}$  satisfies  $f \leq Tf$  then is  $Tf = \alpha f' + \beta \Delta(f)$ ? *Question* 72. If  $\delta(f) > 1$  then is there a g with all real roots such that  $\Delta(g) = f$ ? *Question* 73. Since  $\prod_{i=0}^{2n-1} (x - 2i)$  is a polynomial of degree 2n in  $\mathbf{P}^{sep}$ , we know that

$$p_{n} = \Delta^{n} \prod_{i=0}^{2n-1} (x-2i) \in \mathbf{P}^{sep}(n)$$

Although it is not the case that  $p_n \leq p_{n-1}$ , it appears to be nearly true. Show that if  $g_n = p_n(x)/(x - \alpha)$  where  $\alpha$  is the largest root of  $p_n$  then  $p_{n-1} \leq g_n$ . *Question* 74. Which series  $f(x) = \sum_{0}^{\infty} a_i x^i$  have the property that f \* g is in  $\mathbf{P}^{\mathbb{A}}$  if  $g \in \mathbf{P}^{\mathbb{A}}$ ? This is an affine analog of Theorem 15.22.

*Question* 75. If f has all its roots in (-1, 0) and T:  $x^n \longrightarrow \langle \underline{x} \rangle_n$  then the roots of Tf satisfy: the largest is larger than all roots of f, and the rest are all less than -1.

*Question* 76. Show that the map  $x^n \longrightarrow \langle \underline{x} \rangle_n$  is a bijection between  $\mathbf{P}^{pos}$  and  $\mathbf{P}^{sep} \cap \mathbf{P}^{pos}$ .

*Question* 77. For any A is there an infinite family of polynomials  $h_n$  so that

- $D_{\mathbb{A}}h_n = (const)h_{n-1}$
- $h_n \in \mathbf{P}^{\mathbb{A}}$

*Question* 78. If  $\{a_i\}$  is a sequence such that  $g \leq \sum a_i g^{(i)}$  whenever g is  $(\underline{x})_n$  then do we have interlacing for all  $g \in \mathbf{P}^{sep}$ ?

*Question* 79. If f is a polynomial such that  $f(D_{\mathbb{A}})\mathbf{P}^{\mathbb{A}} \subset \mathbf{P}^{\mathbb{A}}$  then is  $f \in \mathbf{P}^{\mathbb{A}}$ ?

*Question* 80. Does  $Tf = x f(x/2) - \Delta(f)$  preserve roots in some appropriate space?

Question 81. If  $f(x) \gg g(x)$ ,  $f(x) \gg f(x + 1)$ ,  $g(x) \gg g(x + 1)$  then is  $\begin{vmatrix} f(x) & g(x) \\ f(x+1) & g(x+1) \end{vmatrix} \neq 0$ ?

*Question* 82. What are conditions on f that guarantee that  $\begin{vmatrix} f & \Delta f \\ \Delta f & \Delta^2 f \end{vmatrix}$  is never zero? This determinant is also equal to  $\begin{vmatrix} f(x) & f(x+1) \\ f(x+1) & f(x+2) \end{vmatrix}$ .

*Question* 83. Show that the positive part of the q-Hermite polynomials is in  $\mathbf{P}^{\mathbb{A}}$ .

*Question* 84. For which polynomials does the q-Hermite transformation  $x^n \mapsto H_n^q$  (see § 8.8.12) preserve real roots?

*Question* 85. Recall  $\delta(f)$  is the minimum distance between roots of f. What is  $\inf_{f} \frac{\delta(exp(f))}{\delta(f)}$ ? It is 2 for quadratic and 1 for linear.

*Question* 86. What is  $\inf_{f} \frac{\delta(f*g)}{\delta(f)\delta(g)}$ ?

*Question* 87. For which polynomials F(x, z) is  $F(x, \Delta) : \mathbf{P}^{sep} \longrightarrow \mathbf{P}^{sep}$ ? How about  $F(x, A) : \mathbf{P}^{sep} \longrightarrow \mathbf{P}^{sep}$ ?

Question 88. Consider the recurrence

 $p_{n+2} = (a_n x + b_n + (c_n x + d_n) \mathbb{A}) p_{n+1} - e_n \mathbb{A} p_n$ 

What conditions on the coefficients do we need so that all  $p_n \in \mathbf{P}^{sep}$ , and  $p_{n+1} \leq p_n$  for all n?

*Question* 89. Show that the real parts of the roots of  $\Delta^k [\langle \underline{x} \rangle_n \langle \underline{x} \rangle_n]$  satisfy these two properties.

- 1. The real parts are all integers, or integers plus 1/2.
- 2. If k = n + r where  $r \ge -1$  then the real parts are all equal, and have value -(2n + r 1)/2.

*Question* 90. Show the following:

The map  $f \times g \mapsto f \otimes g$  satisfies  $\mathbf{P}^{sep} \cap \mathbf{P}^{alt} \times \mathbf{P}^{sep} \cap \mathbf{P}^{alt} \longrightarrow \mathbf{P}^{sep} \cap \mathbf{P}^{alt}$ . (See p. 231.) *Question* 91. Find an explicit formula for the q-Laguerre polynomials. It is probably similar to the definition of the Laguerre polynomials:

$$L_n(x) = \frac{1}{n!} \sum_{i=0}^n \binom{n}{i} \frac{n!}{i!} (-x)^n$$

*Question* 92. Suppose that  $f_t, g_t, h_t$  are locally interlacing families with  $\rho(f) \ge 1$ ,  $\rho(g) \ge 1$ ,  $\rho(h) \ge 1$ . Show that the triple convolution below is in **P**.

$$\int_{0}^{1} \int_{0}^{1-t} f_{t}(x) g_{s}(x) h_{1-s-t}(x) \, ds \, dt$$

*Question* 93. Suppose that  $f \in \mathbf{P}^{\mathbb{A}}$  where  $\mathbb{A}x = qx$  with q > 0. Show that if  $f = \sum a_i x^i$  then (see 247)

$$\frac{a_{k+1}^2}{a_k a_{k+2}} \ge \frac{1}{q} \frac{[k+2][n-k]}{[k+1][n-k-1]}.$$

*Question* 94. Suppose that  $\mathbb{A}x = qx$  where q > 1,  $\mathbb{B}x = q^{1/2}x$  and  $\mathsf{D}^{sym}_{\mathbb{A}}f(x) = (\mathbb{A}f - \mathbb{A}^{-1}f)/(\mathbb{A}x - \mathbb{A}^{-1}x)$ . Show that if  $f \in \mathbf{P}^{\mathbb{A}}$  then  $\mathsf{D}^{sym}_{\mathbb{A}}f \in \mathbf{P}^{\mathbb{B}}$ .

*Question* 95. Suppose that  $f \leq \mathbb{A}^2 f$  and the polynomial  $f + \mathfrak{l} \mathbb{A} f$  has roots  $r_1 + \mathfrak{l} \mathfrak{s}_1, \ldots, \mathfrak{r}_n + \mathfrak{l} \mathfrak{s}_n$ , and  $g = (\mathfrak{x} - \mathfrak{r}_1) \cdots (\mathfrak{x} - \mathfrak{r}_n)$ . Show that

$$f \underline{\ll} g \underline{\ll} \mathbb{A}f.$$

(It then follows that  $\mathbb{A}f \underline{\ll} \mathbb{A}g \underline{\ll} \mathbb{A}^2 f$ . Since  $f \underline{\ll} \mathbb{A}^2 f$  the sequence  $f, g, \mathbb{A}f, \mathbb{A}g, \mathbb{A}^2 f$  is mutually interlacing, and hence  $g \in \mathbf{P}^{\mathbb{A}}$ .)

Question 96. Define the q-Charlier polynomials [125] by

$$C_{n+1}(x, a; q) = (x - aq^n - [n])C_n(x, a; q) - a[n]q^{n-1}C_{n-1}(x, a; q),$$

where  $C_{-1}(x, a; q) = 0$  and  $C_0(x, a; q) = 1$ . If q > 1 and a > 1 then show that  $C_n(x, a; q)$  is in  $\mathbf{P}^{\mathbb{A}}$ , where  $\mathbb{A}x = qx$ .

#### **Polynomials in P**<sub>2</sub>

*Question* 97. If  $f \in \mathbf{P}_2^{sep}$  then show that  $\Delta_x(f) = f(x + 1, y) - f(x, y)$  is in  $\mathbf{P}_2^{sep}$ . *Question* 98. Show that the linear transformation  $x^i y^j \mapsto \langle \underline{x} \rangle_i \langle \underline{y} \rangle_j$  maps  $P_2$  to itself.

*Question* 99. Suppose that we have three polynomials  $f, g, h \in P_2$  where  $af + bg + ch \in \pm P_2$  for all  $a, b, c \in \mathbb{R}$ . Are f, g, h necessarily linearly dependent?

*Question* 100. If f(x,y) is a product of linear factors, then for which linear transformations T is Tf ever a product of linear factors?

*Question* 101. Suppose that  $f \in \mathbf{P}_2$  and write

$$f = f_0 + f_1 y + f_2 y^2 + \dots + f_n y^n$$

Consider the  $d \times d$  determinant

$$det_{d} = \begin{vmatrix} f_{0} & f_{1} & \dots & f_{d-1} \\ f_{1} & f_{2} & \dots & f_{d} \\ f_{2} & f_{3} & \dots & f_{d+1} \\ \vdots & & & \\ f_{d-1} & f_{d} & \dots & f_{2d-2} \end{vmatrix}$$

Empirical evidence suggests that

- If d = 2r is even and d < n then  $det_d$  always has the same sign  $(-1)^r$ .
- If d is odd then the determinant has positive and negative values.

This is similar to results for matrices of orthogonal polynomials found in [99]. The case where f is in  $\mathbf{P}_2^{\text{pos}}$  is discussed in § 4.4.5.

*Question* 102. (The total positivity conjecture) Suppose that  $f \in \mathbf{P}_2^{\text{pos}}(n)$  is a product of linear terms, and write  $f = \sum a_{ij} x^i y^j$ . When is the matrix below totally positive? (Lemma 9.120 is a special case.)

$$\begin{array}{cccc} a_{0d} & \dots & a_{00} \\ \vdots & & \vdots \\ a_{dd} & \dots & a_{d0} \end{array}$$

*Question* 103. We can form polynomials from the coefficients of a polynomial in **P**<sub>2</sub>. Suppose that  $f = \sum a_{i,j} x^i y^j$ . Define M(i, j; d) to be the determinant of the d by d matrix  $(a_{i+r,j+s})$ , where  $0 \le r, s < d$ . For example,  $M(i, j, 1) = a_{i,j}$ , and  $M(i, j; 2) = a_{i,j}a_{i+1,j+1} - a_{i,j+1}a_{i+1,j}$ .

When is  $\sum M(i, j; d)x^iy^j \in \pm \mathbf{P}_2$ ?

*Question* 104. Show that the product of two polynomials in  $\mathbf{P}_2^{\text{pos}}$  with totally positive coefficients has totally positive coefficients.

*Question* 105. Suppose  $f = \sum a_i x^i$  where  $f \in \mathbf{P}^{pos}(n)$ . The polynomial f(x+y) is in  $\mathbf{P}_2$  and

$$\begin{split} f(x+y) &= \sum_{0}^{n} \alpha_{i}(x+y)^{i} \\ &= \sum_{i=0}^{n} \sum_{j=0}^{i} \binom{i}{j} \alpha_{i} x^{i-j} y^{j} \end{split}$$

Show that the following matrix is totally positive

$$\begin{pmatrix} \dots & a_3 & a_2 & a_1 & a_0 \\ & \dots & 3a_3 & 2a_2 & a_1 \\ & & \dots & 3a_3 & a_2 \\ & & & \dots & a_3 \\ & & & & \dots \end{pmatrix}$$

This is a stronger statement than Theorem 4.23.

*Question* 106. Suppose that  $f(x, y) \in \widehat{\mathbf{P}}_2$  is positive whenever x and y are positive. This is the case if  $f \in \mathbf{P}_2^{\text{pos}}$ ; it also holds for  $e^{-xy-x^2}$ . When is 1/f is totally positive on  $\mathbb{R}^2_+$ ?

*Question* 107. The conjecture in [76] is that if  $A = (a_{ij})$  is a real n by n matrix with non-negative entries which are weakly increasing down columns then the permanent of A + xJ has all real roots.

Show that the permanents of all the submatrices of A + xJ obtained by deleting any row and column have a common interlacing with per(A + xJ).

*Question* 108. If a linear transformation is defined on  $P_2$ , and T maps all products  $\prod(y + a_ix + b_i)$  to  $P_2$ , then does T map  $P_2$  to itself?

*Question* 109. Which products have coefficients satisfying a recurrence? That is, given a product  $\prod (a_i + b_i y + c_i) = \sum f_i(x)y^i$  for which  $a_i, b_i, c_i$  do we have  $f_i = x\alpha_i f_{i-1} + \beta_i f_{i-2}$  for constants  $\alpha_i, \beta_i$ .

*Question* 110. Given  $f \in \mathbf{P}(n)$  for which r can we find a product of n + r terms ax + by + c so that the coefficient of  $y^r$  is f? We can always do it for r = 1, since all we need do is choose g < f.

*Question* 111. If  $f_0 < f_1 < f_2$  then is there a polynomial  $f_0 + f_1(x)y + f_2(x)y^2 + \cdots + f_ny^n$  that satisfies x-substitution?

*Question* 112. If  $f \in \mathbf{P}_2$  and  $\prod (a_i x + b_i y + c_i) \leq f$  then is  $f = \sum d_i \frac{f}{a_x + b_i y + c_i}$ ?

*Question* 113. The Legendre polynomials can be defined by  $P_n(x) = \frac{(-1)^n}{2^n n!} D^n (x^2 - 1)^n$ . This leads to the guess that the map

$$x^i y^j \mapsto \alpha_{n,m} D^i (x^2 - 1)^j$$

might map  $P_2$  to P for some constants  $\alpha_{n,m}$ . This is not true when all  $\alpha_{n,m}$  are equal to 1. For instance, the image of (x + y + 16)(x + 4y + 25) under this map has complex roots. Is there a choice of constants for which it is true?

*Question* 114. What are the different isotopy types of graphs of polynomials in  $P_2(n)$ ?.

*Question* 115. If  $f \in P_2$  then let  $p_n$  be an eigenpolynomial. This means that there exists a  $\lambda_n$  such that  $f(x, D)p_n = \lambda_n p_n$ . See Lemma 15.69.

1. When do they exist?

- 2. When is  $p_n \in \mathbf{P}$ ?
- 3. When does  $p_{n+1} \leq p_n$ ?

*Question* 116. For which diagonal matrices D and symmetric matrices C is the integral below in  $P_2$ ?

$$\int_0^1 |\mathbf{x}\mathbf{I} + \mathbf{y}\mathbf{D} + e^{\mathbf{t}\mathbf{C}}| \, \mathrm{d}\mathbf{t}$$

*Question* 117. Let C be a function from the reals to symmetric matrices. Define

$$f(x,t) = |xI + C(t)|.$$

For every t we kare given that  $f(x, t) \in \mathbf{P}$ . What assumptions on C must we make so that  $f \in \widehat{\mathbf{P}}_2$ ? Consider  $C(t) = e^{tE}$ , where E is symmetric. The roots of f(x, t) are the negative eigenvalues of  $e^{tE}$ , and they are  $\{e^{t\lambda_i}\}$  where the eigenvalues of E are  $\{\lambda_i\}$ . Is this in  $\widehat{\mathbf{P}}_2$ ?

*Question* 118. If  $f(x, y) \in P_2$ , and  $\sigma$  has positive real and imaginary parts then we know that the roots of  $f(x, \sigma)$  all lie in the lower half plane. Show that the argument of a root  $\rho$  of  $f(x, \sigma)$  satisfies

$$|\pi - \arg(\rho)| \leq 2 \arg(\sigma).$$

Question 119. If  $f \in \mathbf{P}_2^{\text{pos}}$ ,  $f = \sum f_i(x)y^i$ ,  $f_0 = \sum_k f_{2k}(x)$ ,  $f_1(x) = \sum_k f_{2k+1}(x)$  then all the roots of  $f_0$  and of  $f_1$  have negative real part (they may be complex).

*Question* 120. Can we remove the factorials from Corollary 4.28? Suppose  $f \in \mathbf{P}^{pos}$ , d is a positive integer, and  $\alpha > 0$ . Then show that

$f(\alpha + d)$	$f(\alpha + d - 1)$	 $f(\alpha)$	
•		:	
:		:	$\geq 0$
$f(\alpha + 2d)$	$f(\alpha + 2d - 1)$	 $f(\alpha + d)$	

*Question* 121. If d is odd show that the derivative of N(f) (see (4.5.4)) is positive for all x

*Question* 122. If d is even show that N(f) (see (4.5.4)) is positive for all x. (Not just for positive x.)

*Question* 123. Is the set of all extensions of two polynomials in  $\mathbf{P}^{\text{pos}}$  a convex set? More precisely, is it the convex hull of the n! polynomials arising from products? (p. 327)

*Question* 124. (*The analog of common interlacing for*  $P_2$ ) If f,  $g \in P_2$  and  $\alpha f + \beta g \in P_2$  for all non-negative  $\alpha$ ,  $\beta$  then is there an  $h \in P_2$  such that  $f \leq h$  and  $g \leq h$ ?

Using the one variable result, it is easy to see that there is a continuous function h such that the graph of h interlaces the graphs of f and g. Can we choose such an h to be in  $P_2$ ?

*Question* 125. If  $f = \sum a_{i,j} x^i y^j \in \mathbf{P}_2^{\text{pos}}$  then the n - 2 by n - 2 matrix

$$\left(\frac{\mathfrak{a}_{\mathfrak{i},\mathfrak{j}}\,\mathfrak{a}_{\mathfrak{i}+1,\mathfrak{j}+1}}{\mathfrak{a}_{\mathfrak{i}+1,\mathfrak{j}}\,\mathfrak{a}_{\mathfrak{i},\mathfrak{j}+1}}\right)_{0\leqslant\mathfrak{i},\mathfrak{j}\leqslant\mathfrak{n}-2}$$

has  $\lfloor \frac{n-1}{2} \rfloor$  negative eigenvalues and  $\lfloor \frac{n}{2} \rfloor$  positive eigenvalues.

*Question* 126. (A generalization of Kurtz's theorem(p. 128)) Suppose that  $f = \sum_{i+j \leq n} a_{i,j} x^i y^j$  has all positive coefficients. If for all relevant i, j the following inequalities hold then  $f \in P_2$ .

$$\frac{a_{i,j}a_{i,j+1}}{a_{i-1,j}a_{i+1,j}} \ge 2, \quad \frac{a_{i,j+1}a_{i+1,j}}{a_{i,j}a_{i+1,j+1}} \ge 2, \quad \frac{a_{i,j}a_{i+1,j+1}}{a_{i,j+1}a_{i+1,j+1}} \ge 2$$

In other words, if all rhombus inequalities hold with constant 2 then  $f \in P_2$ . It is a consequence of these inequalities that  $a_{i,j}^2 \ge 4a_{i,j-1}a_{i,j+1}$  and so by Kurtz's theorem we know that all  $f_i \in \mathbf{P}^{pos}$  where  $f = \sum f_i(x)y_i$ .

#### Polynomials in several variables

*Question* 127. Given f < g in  $\mathbf{P}^{pos}$  is there  $\sum h_{ij}(x)y^i z^j \in \mathbf{P}_3^{pos}$  such that

$$h_{00} = f$$
  $h_{10} = g$   $\begin{vmatrix} h_{00} & h_{10} \\ h_{01} & h_{11} \end{vmatrix} = 1$ 

*Question* 128. For a fixed polynomial f in  $\mathbf{P}_d$  what are the possible dimensions of the cone of interlacing of f?

*Question* 129. Can every polynomial is  $\mathbf{P}_d$  be realized as a coefficient of a product of linear terms with a large number of variables?

*Question* 130. If  $f = \sum a_I(x)y^I$ ,  $g = \sum b_I(x)y^I$ ,  $f \in P_{d+e}^{pos}(n)$  and  $g \in P_{d+e}^{pos}(n)$ , then when is the "Hadamard Product" below in  $P_{d+e}(n)$ ?

$$\sum a_I(x) b_I(x) y^I$$

It appears to hold for d = 0, e = 2, or d = e = 1. If it held for d = e = 1 then we could easily derive a consequence due to [66]: If  $f_0 < f_1$  and  $g_0 < g_1$  are all in  $\mathbf{P}^{\text{pos}}$  then  $f_0 * g_0 < f_1 * g_1$ . The proof goes like this: We can find  $f, g \in \mathbf{P}_2^{\text{pos}}$  such that

$$f(x,y) = f_0(x) + f_1(x)y + \cdots$$
  
g(x,y) = g\_0(x) + g\_1(x)y + \cdots

By Question 130

 $f_0*g_0+f_1*g_1y+\cdots\in \textbf{P}_2$ 

Since the coefficients interlace,  $f_0 * g_o \leq f_1 * g_1$ .

Question 131. The "generalized Legendre" polynomials are of the form

$$\left(\frac{\partial}{\partial x_1}\right)^{n_1} \dots \left(\frac{\partial}{\partial x_r}\right)^{n_r} (x_1^2 + x_2^2 + \dots + x_r^2 - 1)^{n_1 + \dots + n_r}$$

They do not satisfy substitution for  $x_i$ 's in the unit ball.

- 1. Show that the one variable polynomials that are coefficients of monomials of the form  $x_2^{e_2}x_3^{e_3}\cdots x_r^{e_r}$  are in **P**, where the  $e_i$  are positive integers.
- 2. If the case r = 2 show that if

$$f_{2i,2j}(x^2,y^2) = \left(\frac{\partial}{\partial x}\right)^{2i} \left(\frac{\partial}{\partial y}\right)^{2j} (x^2 + y^2 - 1)^{2i+2j}$$

then  $f_{2i,2j}(x, y) \in \mathbf{P}_2$ .

*Question* 132. If e > 1 and  $\sum a_I \mathbf{x}^I \in \mathbf{P}_d^{\text{pos}}$ , then  $\sum a_I^e \mathbf{x}^I \in \mathbf{P}_d^{\text{pos}}$ .

*Question* 133. Suppose that  $Q_1, \ldots, Q_d$  are negative subdefinite matrices with the property that for all positive  $\alpha_1, \ldots, \alpha_d$  the matrix  $\alpha_1 Q_1 + \cdots + \alpha_d Q_d$  is negative subdefinite. (Find a general construction for such matrices.) Show that if  $f(\mathbf{x}) \in \mathbf{P}_d^{\text{pos}}(n)$  then

$$(-1)^{n} f(-\mathbf{x}Q_1\mathbf{x}^t,\ldots,-\mathbf{x}Q_d\mathbf{x}^t) \in \mathbf{P}_d(2n)$$

*Question* 134. Suppose that T:  $P_d \longrightarrow P$  is a linear transformation with the property that monomials map to monomials. Can we characterize such T? Are they constructed out of maps from  $P_2 \longrightarrow P$ ? And, are these of the form  $x^i y^j \mapsto z^{i+j}, x^i y^j \mapsto \alpha^i z^j$ , coefficient extraction, and  $x^i y^j \mapsto (\underline{j})_i z^{j-i}$ ? For instance, if it's true that  $x^i y^j \mapsto (\underline{j})_i z^{j-i}$  maps  $P_2 \longrightarrow P$ , then we have a map

$$\mathbb{P}_3 \xrightarrow{(x,y,v)\mapsto(x,x,v)} \mathbb{P}_2 \xrightarrow{f(x,U)} \mathbb{P}_2$$

$$x^{i}y^{j}v^{k} \longmapsto x^{i+j}v^{k} \longmapsto (\underline{i+j})_{k}z^{k-i-j}$$

More generally, can we describe all *multiplier transformations*? These are linear transformations  $P_d \longrightarrow P_e$  that map monomials to monomials.

*Question* 135. Prove the following. Note that the only difficulty is to replace the zero entries of Q with small positive values so that there is still exactly one positive eigenvalue. There is only a problem if the determinant is zero.

Lemma 24.32. If Q is a matrix with

1. all non-negative entries

2. exactly one positive eigenvalue

then Q is a limit of negative subdefinite matrices.

#### QUESTIONS

*Question* 136. Show that the Hurwitz transformation (15.9.1) maps  $\mathbf{P}_{d}^{pos}$  to itself.

*Question* 137. Show that the linear transformation  $x_1^{i_1} \cdots x_d^{i_d} \mapsto \frac{x_1^{i_1} \cdots x_d^{i_d}}{(i_1 + \cdots + i_d)!}$ maps  $\mathbf{P}_d^{\text{pos}}$  to itself. (It does not map  $\mathbf{P}_2$  to itself.)

*Question* 138. Suppose that  $f_{0,0} = 1$  and  $f_{i,j}$  satisfies (10.24.2) where the constants satisfy  $b_{i,j} \ge 0$  and

$$a_{0,0} \leq a_{0,j} \leq a_{1,0} \leq a_{1,j} \leq a_{2,0} \leq a_{2,j} \leq a_{3,0} \cdots$$
 for  $j = 1, 2, \dots$ 

Then we have  $f_{i,j} \leq f_{i,0}$  for all i, j. Consecutive  $f_{i,j}$  do not necessarily interlace. *Question* 139. For which regions  $D \subset \mathbb{R}^2$  is it the case that for all  $f \in \mathbf{P}_3^{sep}$  the integral below is in **P**?

$$\int_D f(x,s,t)\,ds\,dt$$

*Question* 140. For which regions  $D \subset \mathbb{R}^2$  is it the case that for all f,  $g \in \mathbf{P}_2^{sep}$  the integral below is in **P**?

$$\int_D f(x,s)g(x,t)\,ds\,dt$$

In this case, the conclusion holds if D is the product  $(0, 1) \times (0, 1)$  or if D is the segment  $0 \leq s, t$  and s + t = 1 or if D is the triangle  $0 \leq s, t$  and  $s + t \leq 1$ .

*Question* 141. Suppose that  $f_{00} \leq f_{01}$ ,  $f_{10}$ . Show that  $f_{10}f_{01} - f_{00}W \ge 0$  if and only if there is an extension of  $f_{00}$ ,  $f_{10}$ ,  $f_{01}$  such that  $W = f_{11}$ . (See p. 361.)

*Question* 142. If f(x, y) has the property that  $f(\alpha, y) \ll g(\alpha, y + 1)$  and  $f(x, \alpha) \ll f(x + 1, \alpha)$  for all real  $\alpha$  does it follow that  $f \in \mathbf{P}^{sep}$ ?

#### The polynomial closure of P<sub>d</sub>

*Question* 143. We know that a + bx + cy + dxy is in  $\overline{\mathbf{P}}_2$  for any choice of a, b, c, d satisfying  $\begin{vmatrix} a & b \\ c & d \end{vmatrix} < 0$ . For which choices of constants  $a_I$  do we have

$$\sum_{I} \mathfrak{a}_{I} x^{I} \in \overline{P}_{d}$$

where the sum is over all  $2^d$  monomials I? For instance, if d = 3 a determinant such as

$$\begin{array}{ccccc} x + a & d_1 & d_2 \\ d_1 & y + b & d_3 \\ d_2 & d_3 & z + c \end{array}$$

has the desired form. Brändén's criterion generally leads to infeasible problems

*Question* 144. Suppose that L is a lattice in  $\mathbb{Z}^d$  with basis vectors  $v = \{v_1, \ldots, v_e\}$ . If  $\mathbf{I} = (i_1, \ldots, i_e)$  is an index set then set  $\mathbf{I} \cdot v = i_1 v_1 + \cdots + i_e v_e$ . When is it true that

If 
$$f(\mathbf{x}) = \sum a_{\mathbf{I}} \mathbf{x}^{\mathbf{I}}$$
 is in  $\mathbf{P}_{d}^{\text{pos}}$  then  $\sum a_{\mathbf{I} \cdot \mathbf{v}} \mathbf{x}^{\mathbf{I}} \in \overline{\mathbf{P}}_{e}^{\text{pos}}$ 

Notice that by taking  $v_1 = (2)$  the Hurwitz theorem on even and odd parts (Theorem 7.64) is a consequence of this question. Empirically, the question holds for  $v_1 = (3,3)$  or (4,4) or (2,4), and fails for  $v_1 = (3,2)$  or (2,1). The question fails if we replace  $\mathbf{P}_2^{\text{pos}}$  by  $\mathbf{P}_2$ .

The diagonal (Theorem 13.9) of a polynomial in **P**<sub>2</sub> is found by taking  $v_1 = (1,1)$ . For another example, take  $f = (x + y + z + 1)^n$ ,  $v_1 = (1,1,0)$  and  $v_2 = (0,0,1)$ . The resulting polynomial is in **P**<sub>2</sub>, and the coefficient of  $x_2^k$  is  $\sum {k \choose i,i} x_1^i$ .

*Question* 145. If  $f \in \overline{\mathbf{P}}_d$  has degree 1 in each variable then is there a d by d matrix A and a diagonal matrix of the form  $(d_i x_i)$  such that det(A + D) = f? In particular, can xyz - x - y - z + 1 be so represented?

*Question* 146. Suppose  $f = \sum a_{i,j}x^iy^j \in P_2$ , and define  $f_{rs} = \sum a_{2i+r,2j+s}x^iy^j$  where  $r, s \in \{0, 1\}$ . Show that

$$\mathsf{f}_{00} + \mathsf{u}\mathsf{f}_{10} + \mathsf{u}\mathsf{f}_{01} + \mathsf{u}\mathsf{v}\mathsf{f}_{11} \in \overline{\mathbf{P}}_4.$$

Is this true in the special case that f is a product of linear factors?

Question 147. Suppose that  $g \in \overline{P}_2^{\text{pos}}(n)$  and write  $g(x, y) = \sum_{i=0}^{n} g_i(x)y^i$ . Define

$$f(x) = \sum_{i=0}^{n} g_i(x^{n+1})x^i = g(x^{n+1}, x)$$

then the absolute value of the argument of any root of f is at least  $\frac{\pi}{n+1}$ .

Note that if n = 1 then this states that  $g_0(x^2) + xg_1(x^2)$  has negative real part for  $g_0 \leftarrow g_1$  - the classic Hurwitz stability result.

*Question* 148. Suppose that  $f_1, \ldots, f_n$  are mutually interlacing. Are there positive  $a_i$  so that

$$\sum_{i=1}^n a_i f_i(x) \, y^i \in \overline{\textbf{P}}_2?$$

A special case of this: suppose the roots of f are  $r_1\leqslant \cdots \leqslant r_n.$  Are there positive  $a_i$  so that

$$\sum_{i=1}^{n} a_{i} \frac{f(x)}{x - r_{i}} y^{i} \in \overline{\mathbf{P}}_{2}?$$

*Question* 149. Let  $T(x^n) = P_n$ , the Legendre polynomial. Show that if  $f \in \overline{P}_2$ , then  $T_*(f)(\alpha, y) \in \mathbf{P}$  for  $|\alpha| < 1$ . The case  $f = (-1 + xy)^n$  is [168, page 387, problem 69] and (14.3.1).

*Question* 150. Recall the Hermite polynomials  $H_{r,s}$  in Theorem 13.35. We know that we can represent  $H_{r,s}$  as the determinant of a matrix. Show that we have the following representation of  $H_{r,1}$  as the determinant of the tridiagonal matrix

$$(-1)^{r+1}e^{ax^2+2bx+cy^2}\frac{\partial^{r+1}}{\partial x^r\partial y}e^{-(ax^2+2bxy+cy^2)} =$$

$$\begin{vmatrix} 2(ax+by) & \sqrt{2a} & 0 & 0 & \dots & \dots & 0 \\ \sqrt{2a} & 2(ax+by) & \sqrt{4a} & 0 & \dots & \dots & 0 \\ 0 & \sqrt{4a} & 2(ax+by) & \sqrt{6a} & \dots & \dots & 0 \\ & & \ddots & \ddots & & & \\ 0 & \dots & \dots & \sqrt{2(r-2)a} & 2(ax+by) & \sqrt{2(r-1)a} & 0 \\ 0 & \dots & \dots & 0 & \sqrt{2(r-1)a} & 2(ax+by) & \sqrt{2rb} \\ 0 & \dots & \dots & 0 & 0 & \sqrt{2rb} & 2(bx+cy) \end{vmatrix}$$

*Question* 151. For which symmetric matrices Q is  $xQy^t \in \overline{P}_{2d}$ ? Setting x = y shows that  $xQx^t \in \overline{P}_d$ , so Q is negative subdefinite.

*Question* 152. Suppose that f, g, h,  $k \in \mathbf{P}$  satisfy

1. Interlacing: 
$$f \begin{pmatrix} g \\ h \end{pmatrix} k$$

2. Determinant: 
$$\begin{vmatrix} f & g \\ h & k \end{vmatrix} \leq 0$$

Show that  $f + y g + z h + y z k \in \overline{\mathbf{P}}_3$ .

#### The analytic closure of P<sub>d</sub>

*Question* 153. If L<sub>n</sub> is the Legendre polynomial then [168, page 69]

$$f(x,y) = \sum_{i=0}^{\infty} L_n(x)y^n = (1 - 2xy + y^2)^{-1/2}$$

If  $g \in \mathbf{P}^{(-\infty,1)}(n)$  then show that the coefficients of  $y^n, \dots, y^{2n}$  in g(y)f(x,y) are in **P**. The coefficients of  $y^2, \dots, y^{n-1}$  are not necessarily in **P**.

*Question* 154. When do all the coefficients of  $y^i$  in a product of terms of the form  $(a + y)^{bx+c}(d + y)^{rx+s}$  have all real roots? The coefficient of  $y^3$  in  $(1 + y)^x(2+y)^x$  is not in **P**. It seems that all coefficients are in **P** for  $(1+y)^x(2-y)^x$ . *Question* 155. Show that the linear transformation  $x^I \mapsto H_I$  where  $H_I$  is the d-dimensional Hermite polynomial, maps  $P_d$  to itself.

*Question* 156. Suppose that  $\{p_n\}$  is an orthogonal polynomial system. Can we find constants  $a_i$  so that

$$\sum_{i=0}^\infty \alpha_i p_i(x) y^i \in \widehat{P}_2$$

#### Extending P<sub>d</sub>

*Question* 157. Some questions about  $\mathfrak{P}_2$ .

- 1. Is  $\mathfrak{P}_2$  closed under multiplication?
- 2. If  $f \in \mathbf{P}^{\text{pos}}$  then is  $f(-xy) \in \mathfrak{P}_2$ ?
- 3. Is  $e^{-xy} \in \mathfrak{P}_2$ ? This would follow from the above question.
- 4. Does Lemma 12.4 hold for  $\mathfrak{P}_2$ .

*Question* 158. Consider the map  $x^n \mapsto L_n(x; y)$ . Show that it maps  $\mathbf{P} \longrightarrow \mathcal{P}_{1,1}$ . *Question* 159. Is  $y\sqrt{x}$  in the closure of  $\mathcal{P}_{1,1}$ ?

*Question* 160. Which functions analytic in Re(z) > 0 are uniform limits (in Re(z) > 0) of polynomials with all real roots?

*Question* 161. If  $f \in \mathcal{P}_{1,1}$  then show that f(x, y) determine an operator f(x, D) that maps  $\mathbf{P}^{pos} \longrightarrow \mathbf{P}$ 

*Question* 162. Is  $Q_{d,e}$  closed under differentiation? Are coefficients of  $Q_{d,e}$  well-behaved?? Are there interesting transformations that preserve  $Q_{d,e}$ ?

*Question* 163. Suppose f,  $g \in Q_{d,2}$  have the property that for all  $p \leftarrow q \in Q_{d,2}$  it holds that  $fp + gq \in Q_{d,2}$ . Then  $g \leftarrow f$ .

#### **Generating Functions**

*Question* 164. If  $\frac{d}{dx}f_i(x) = (\underline{x})_i$  then the generating function satisfies

$$\frac{\partial}{\partial x} \left( \sum f_i(x) \frac{y^i}{i!} \right) = \sum (\underline{x})_i \frac{y^i}{i!}$$
$$= (1+y)^x$$

Integration with respect to x shows

$$\sum f_i(x) \frac{y^i}{i!} = \frac{(1+y)^x}{\log(1+y)} + c(y)$$

If we choose c(y) = -1/y and write

$$\sum g_{\mathfrak{i}}(x)\frac{y^{\mathfrak{i}}}{\mathfrak{i}!} = \frac{(1+y)^{x}}{\log(1+y)} - \frac{1}{y}$$

then show that  $g_i(x)$  interlaces  $(\underline{x})_i$  and hence is in **P**. This solves Question 2.

Question 165. Can we use the identity

$$\sum (\Delta^{i} f) \frac{(\underline{y})_{i}}{i!} = f(x + y)$$

to get information about linear combinations of  $\Delta^{i} f$ ? *Question* 166. Is the following true?

Suppose that  $a_1..., a_r$  and  $b_1,..., b_s$  are positive and satisfy  $(\sum b_i) - (\sum a_i) > -1$ . Then the Hadamard-type product

$$x^{\mathfrak{m}} * '' x^{\mathfrak{n}} \mapsto \begin{cases} \prod_{\substack{i=1\\s}}^{r} (a_{i}\mathfrak{n})! \\ \prod_{j=1}^{s} (b_{j}\mathfrak{n})! \\ 0 & \text{otherwise} \end{cases}$$

maps  $\mathbf{P} \times \mathbf{P}^{pos} \longrightarrow \mathbf{P}$ . The assumptions on the a's and b's guarantee that the genus is 0. The problem is to show that all the roots are real.

*Question* 167. In Lemma 15.69 we saw that  $f(x, \partial)$  had an eigenpolynomial  $p_n$  of degree n for n sufficiently large. Show that there is an eigenpolynomial for all n, and that  $p_n \ll p_{n-1}$ .

*Question* 168. Show that if  $G \in \widehat{\mathbf{P}}_2$  interlaces  $e^{-xy}$  then  $G = (ax + by + c)e^{-xy}$  where a and b have the same sign.

More generally, if  $f \in \overline{\mathbf{P}}_2$ , determine all G such that  $G \in \widehat{\mathbf{P}}_2$  interlaces  $f(x,y)e^{-xy}$ .

#### **Recurrences with polynomial coefficients**

*Question* 169. What are necessary and sufficient conditions for  $g_0(x), \ldots, g_d(x)$  so that there exists a sequence  $f_0, f_1, \ldots$  satisfying

- 1.  $f_i \in \mathbf{P}, i = 0, 1, ...$
- 2.  $\sum_{i=0}^{d} g_i f_{i+n} = 0$  for  $n = 0, 1, \ldots$

#### Matrices

*Question* 170. If  $f = \prod(x + b_iy + c_i) = \sum p_i(x)y^i$  then can we explicitly construct a matrix A such that |A[1, ..., i]| is a multiple of  $p_i$ ?

*Question* 171. If A is a d by d matrix of non-negative constants then the sequence

$$\label{eq:h0} \begin{split} h_0 &= I \\ h_{n+1} &= xAh_n - h_n' \end{split}$$

defines a sequence of d by d matrices whose entries are all in **P**.

*Question* 172. Suppose that A, A' are diagonal matrices with positive diagonal entries, and C, C' are symmetric matrices. If |xI + yA + C| = |xI + yA' + C'| then what is the relationship between A, A', and C, C'?

*Question* 173. Interlacing was defined for matrices in terms of their characteristic polynomials. Is there a notion of "interlacing" for general linear transformations? For example, if S and T are compact self-adjoint linear operators on a Hilbert space, then it is clear what it means to say that the discrete spectrums of S and T interlace. However, what does it mean to say that the continuous spectrums interlace?

*Question* 174. Suppose M = xI+yD+C where C is symmetric, and  $D = (d_{ij})$  is a diagonal matrix with positive diagonal. If f(x, y) = |M|, and g(x, y) = |M[1]| then define  $h = f - (x + d_{11}y)g$ . Show that  $h \in \mathbf{P}_2$  and  $f \leq h$ .

*Question* 175. Suppose that  $P_0 = 0$ ,  $P_1$  is the n-dimensional identity matrix, and  $P_{n+1} = x P_n - JP_{n-1}$  where J is the  $n \times n$  matrix of all 1's. The diagonals of  $P_n$  are all equal to a polynomial  $f_n$ , and the off diagonals are all equal to  $g_n$ . It isn't true that  $f_n \in \mathbf{P}$ , but show that  $g_n \in \mathbf{P}$  and  $g_{n+1} \leq g_n$ .

#### Matrix polynomials

*Question* 176. Do all matrix polynomials in Hyp<sub>1</sub> have a common interlacing?

*Question* 177. Is Hyp<sub>1</sub> closed under Hadamard products?

*Question* 178. If  $f \in Hyp_1$  has degree n then does the function

 $x \mapsto$  maximum eigenvalue of f(x)

have exactly n zeros?

*Question* 179. Suppose that  $D_1, \ldots$  and  $E_1, \ldots$  are positive definite matrices. Show that the determinant of

$$(D_1 + x E_1) * (D_2 + x E_2) * \cdots * (D_n + x E_n)$$

is a stable polynomial, where \* is the Hadamard product.

More generally, if A, B are matrices with polynomial entries, and |A| and B| are stable polynomials then is it true that |A \* B| is stable?

Note that all the polynomials in question are matrix polynomials in Hyp<sub>1</sub>.

#### Polynomials non-vanishing in a region

*Question* 180. Determine all the exponential functions in  $\widehat{NV}_1(\mathbb{C})$  where  $\mathbb{C}$  is a cone in  $\mathbb{C}$ .

#### Upper half plane polynomials

*Question* 181. If A is skew-symmetric and  $f(\mathbf{x}) \in U_n$  then is  $f(A\mathbf{x}) \in U_n$ ? If n = 2 and  $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$  then  $f(A\mathbf{x}) = f(y, -x)$  which is in  $U_2$  if f is.



Figure 1: The imaginary part of a polynomial in  $\Im_2$ 

*Question* 182. Describe all the polynomials that interlace a given polynomial in  $U_1(\mathbb{C})$ .

*Question* 183. If  $f \leq g$  in  $U_1(\mathbb{C})$ , and all coefficients are positive then show that  $f \cdot \overline{g} = h(x) + \iota k(x)$  where h and k are stable.

*Question* 184. If  $(x - \sigma)(x - \tau) \in U_1(\mathbb{C})$  then show that  $(x - \sigma)(x - \tau) < x - \nu$  if and only if  $\nu$  lies in the ellipse whose foci are  $\sigma, \tau$ , and is tangent to the x axis.

*Question* 185. If  $f \in \mathbf{P}^{pos}$  then show that  $\int_0^1 f(-\iota tx) dt \in U_1(\mathbb{C})$ .

A consequence:

Let  $h(x) = \int_0^1 f(-\iota tx) dt$ . Then

$$\int_{-1}^{1} f(\iota - tx) \, dt = \int_{0}^{1} f(-\iota tx) \, dt + \int_{-1}^{0} f(-\iota tx) \, dt = h(x) + \overline{h(x)} = 2\Re(h)$$

Since  $h \in U_1(\mathbb{C})$ , it follows that the integral from -1 to 1 is in **P**.

*Question* 186. If  $f \in \mathbf{P}_2^{\text{pos}}$ ,  $f = \sum f_i(x)y^i$  then all the polynomials  $\begin{vmatrix} f_i(x) & f_{i+1}(x) \\ f_{i-1}(x) & f_i(x) \end{vmatrix}$  are stable. In particular, all their coefficients are positive. Thus, the sequence of coefficients of f are *q-log-concave*. (This is not true if  $f \in P_2$ .) More generally, show that the d + 1 by d + 1 determinant of the matrix given below is stable.

*Question* 187. Although we can't graph a polynomial in  $U_2(\mathbb{C})$  since it has complex roots, we can look at the plots of the real and imaginary parts. Since all imaginary parts are negative, the graph of the imaginary parts lies in the lower half plane; the graph of the real parts has no such restriction. The graph of the real part looks like the graph of a polynomial in P<sub>2</sub>, while the graph of the imaginary part looks very different (Figure 1). Explain this behavior.

#### Stable polynomials

*Question* 188. Suppose that  $f = \sum f_i(x)y^i$  is in  $\mathbf{P}_2^{\text{pos}}(n)$ . Define the polynomial

$$p_{d}(x,y) = \sum_{k=0}^{n-2d} y^{k} \operatorname{det}(f_{k+i+j})_{0 \leqslant i,j < d}$$

Show that

- 1.  $p_d(x, \alpha)$  is a stable polynomial for all positive  $\alpha$ .
- 2. All coefficients (the determinants) are stable polynomials.

*Question* 189. If  $f(x, y) \in \mathbf{P}_2^{\text{pos}}$  then all minors of the matrix (21.10.2) are stable. *Question* 190. If  $f, g \in \mathbf{P}_2^{\text{pos}}$  then all entries of the Bezout matrix B(f, g) are stable polynomials in y.

$$B(f,g) = \frac{f(v,y)g(u,y) - f(u,y)g(v,y)}{u - v}$$

Question 191. If  $f = \sum f_i(x, y) z^i \in \mathbf{P}_3^{\text{pos}}$  and

$$M_{d} = \begin{pmatrix} f_{0}(x, y_{0}) & \dots & f_{0}(x, y_{d}) \\ \vdots & & \vdots \\ f_{d}(x, y_{0}) & \dots & f_{d}(x, y_{d}) \end{pmatrix}$$

then the coefficient of any monomial  $y_0^{e_0} \cdots y_d^{e_d}$  in  $|M_d|$  is stable. If d = 1 this is the Bezout matrix. In addition, the determinant of  $M_d$  is stable.

*Question* 192. A special case of the previous question. If  $\sum f_i(x)y^i \in \mathbf{P}_2^{\text{pos}}$  then the polynomial below has all positive coefficients. Moreover, all coefficients of  $x^iy^j$  are in  $\mathbf{P}^{\text{pos}}$ .

$$\frac{1}{(x-y)(x-z)(y-z)} \begin{vmatrix} f_0(x) & f_1(x) & f_2(x) \\ f_0(y) & f_1(y) & f_2(y) \\ f_0(z) & f_1(z) & f_2(z) \end{vmatrix}$$

*Question* 193. If  $g_1, \ldots, g_n \in \mathbf{P}^{pos}(r)$  and  $\mathfrak{m}_1, \cdots, \mathfrak{m}_n$  are positive definite matrices then

$$|g_1\mathfrak{m}_1 + \cdots + g_\mathfrak{m}\mathfrak{m}_n| \in \mathcal{H}$$

*Question* 194. Is the determinant of a stable matrix polynomial a stable polynomial?

*Question* 195. If  $f = \sum f_i(x)y^i \in \mathbf{P}_2^{\text{pos}}$  then is

$$\frac{f_1}{f_2} - \frac{f_0}{f_1} \colon \mathsf{Q}_{\mathrm{I}} \longrightarrow \mathsf{Q}_{\mathrm{I}}$$

*Question* 196. Suppose  $f = \prod (x + r_i) \in \mathbf{P}^{pos}$ . What polynomials in  $\mathbf{P}^{pos}$  interlace f in  $\mathcal{H}_1$ ?

*Question* 197. If  $f \xleftarrow{H} g$  then when are there positive definite D<sub>1</sub>, D<sub>2</sub>, skew-symmetric A, and positive  $\alpha$ ,  $\beta$  such that

$$|\mathbf{I} + \mathbf{x}\mathbf{D}_1 + \mathbf{y}\mathbf{D}_2 + \mathbf{A}| = \alpha \mathbf{f} + \beta \mathbf{g}\mathbf{y} + \cdots$$

*Question* 198. Suppose that  $f = \sum f_i(x)y^i$  is in  $\mathbf{P}_2^{\text{pos}}(n)$ . Define the polynomial

$$p_{d}(x,y) = \sum_{k=0}^{n-2d} y^{k} \operatorname{det}(f_{k+i+j})_{0 \leq i,j < d}$$

Show that

1.  $p_d(x, \alpha)$  is a stable polynomial for all positive  $\alpha$ .

2. All coefficients (the determinants) are stable polynomials.

*Question* 199. Define the infinite matrix

$$M = \begin{pmatrix} \cdot & \cdot & \cdot & \cdots \\ \binom{k}{k} & \cdot & \cdot & \cdots \\ \cdot & \binom{k+1}{k} & \cdot & \cdots \\ \cdot & \cdot & \binom{k+2}{k} & \cdots \\ \vdots & & & \ddots \end{pmatrix}$$

where the periods (.) mean zero. If

$$\exp(M) = \begin{pmatrix} \cdot & \cdot & \cdot & \cdots \\ a_{0,0} & \cdot & \cdot & \cdots \\ a_{0,1} & a_{1,1} & \cdot & \cdots \\ a_{0,2} & a_{1,2} & a_{2,2} & \cdots \\ \vdots & & & \ddots \end{pmatrix}$$

then show that the rows  $\sum a_{i,j} x^j$  are stable.

If k = 2 and  $f_n$  is the polynomial corresponding to the nth row then the recurrence

$$f_n = (x + n - 1)f_{n-1} - \frac{1}{2} {n \choose 2} f_{n-2}$$

shows that  $f_n \in P$ . Since all entries of the exponential are positive we have that  $f_n \in P^{\text{pos}} \subset \mathcal{H}_1$ .

*Question* 200. If  $f \in \mathcal{H}$  and  $f(x) \ge 0$  for all real vectors x then are there  $g \xleftarrow{P} h$  so that  $f = g^2 + h^2$ .

#### Transformations in the complex plane

*Question* 201. Show that the containment in Proposition 22.23 is an equality. That is, if  $T(x^n) = L_n(x)$  and  $|z| \leq 1$  then all roots of  $T(x - z)^n$  have non-negative real part.
#### QUESTIONS

*Question* 202. If  $T: x^n \mapsto L_n(x)$  then show that  $\Lambda(TP^{\Delta})$  is the closed right half plane. Equivalently, show that if Tf has a root with negative real part, then f has a root with absolute value greater than 1.

*Question* 203. Suppose  $f_{\beta}$  is defined as in (22.2.1). Is there a Sendov type result for the roots of  $f_{\beta}$ ? That is, if all the roots of f are in the unit disk, then how far away from a root of f are the roots of  $f_{\beta}$ ?

*Question* 204. What linear transformations have the property that they map polynomials with all roots on a horizontal line in the complex plane to polynomials with all roots on a horizontal line? The transformation  $x^n \mapsto H_n$  is such a transformation.

*Question* 205. If S is a region in  $\mathbb{C}$  and T is a linear transformation, then when is it the case that

1. 
$$\Lambda(\mathsf{T}\mathbf{P}^{\$}) = \Lambda(\mathsf{T}(\phi\$))$$
?  
2.  $\Lambda(\mathsf{T}\mathbf{P}^{\$}) \subset \operatorname{Hull}(\Lambda(\mathsf{T}(\phi\$)))$ ?

*Question* 206. If  $\Delta$  is the unit disk in the complex plane, then what are examples, other than dilations or powers of the derivative, of linear transformations that map  $\mathbf{P}^{\Delta}$  to itself?

*Question* 207. If  $T(x^n) = p_n(x)$  where  $p_n$  is the Legendre or Chebyshev polynomial, then is it the case that

$$\mathsf{T}(\mathbf{P}^{\Delta}) \subseteq \mathbf{P}^{\Delta}?$$

*Question* 208. What are examples, other than the derivative, of linear transformations that map  $\mathbf{P}^{\Delta}$  to itself?

*Question* 209. Let  $T(x^n) = L_n(x)$ . If  $\Delta$  is the unit disk and **R** the right half plane, then

 $\mathsf{T}(\mathbf{P}^{\Delta}) \subset \mathbf{P}^{\mathbf{R}}$ 

That is, if all roots of f have magnitude at most 1, then the roots of Tf have non-negative real part. Is it also the case that every point in the right half plane is the root of Tf, for some  $f \in \mathbf{P}^{\Delta}$ ?

# Notes

#### Polynomials

Interlacing has several synonyms in the literature. Gantmacher [65] uses the term *positive pair* for interlacing polynomials with all negative roots. In [63] a pair of polynomials f, g with all real roots satisfying f < g or  $f \gg g$  is called a *real pair*. If they satisfy  $f \gg g$  and all roots are negative, they are called a *positive pair*. Wagner [178] says that f *alternates left of* g if  $f \gg g$ . Bilodeau [14] says that q *separates* p if  $p \leq q$ . In [73] two polynomials with a common interlacing are called *complementary*. In [158] and [101] a sequence f < g < h is called a *Sturm sequence of polynomials*. [35] says that two polynomials interlace if the roots *se séperent* (separate themselves).

Apparently Markov was the first to prove that differentiation preserves interlacing. See [141].

Lemma 1.11 is in [124].

Quantitative sign interlacing occurs in several different places. It is implicit in [65, page 226]. The case  $f \gg g$  is found in [66]. Quantitative sign interlacing is used in [68] to prove that the characteristic polynomial of a graph is interlaced by the characteristic polynomial of a one point deletion. Exercise 7 on page 146 of [68] is an explicit statement of quantitative sign interlacing for f < g. Quantitative sign interlacing can also be interpreted as a property of rational functions. See [155, page 320].

[179] contains many of the sign-interlacing results of § 1.1.2

Corollary 1.56 is well known; it occurs for example as an exercise in [63, page 269].

Johnson [95] shows the convexity of the set of polynomials that interlace a given polynomial using sign interlacing.

The basis in Section 1.3 is sometimes called the "interpolation basis" [63], since they are, up to a multiple, the polynomials in Lagrange interpolation. Fuhrman [63] calls the basis  $f_i(x) = \frac{f(x)}{x - \alpha_i}$  the *spectral basis*. See [141, page 198] for a different proof of Proposition 1.35. [73] established

See [141, page 198] for a different proof of Proposition 1.35. [73] established Proposition 1.35 in the case of positive combinations for the special class of Hurwitz polynomials. These are polynomials  $f(x^2)$  where  $f \in \mathbf{P}^{\alpha lt}$  has no repeated roots.

The identity [32, page 24]  $\left| \begin{array}{c} p_n & p_{n+1} \\ p'_n & p'_{n+1} \end{array} \right| > 0$  where the  $p_i$  are orthogonal follows from Lemma 1.55 using the fact that  $p_{n+1} < p_n$ .

A consequence of Lemma 1.59 is that the associate polynomials of a family

of orthogonal polynomials interlace. See [32, page 86]. Lemma 1.59 is often used implicitly - see for example [119, Lemma 3.1]

[8] proved Lemma 1.55 in the more general case of differentiable functions. Corollary 4.31 is in [35].

[56] has many results about polynomials, with no references. Some of them are from [138]. This is translation of the book that first appeared in 1954. Here are a few of the items in the book:

problem 727 interlacing implies all linear combinations in P

problem 728 all linear combinations implies interlacing

problem 729 f, g interlace implies derivatives interlace.

problem 730  $\gamma f + (\lambda + x) f' \in \mathbf{P}$  if  $\gamma > 0$  or  $\gamma < -n$ 

problem 742 if  $f - a \in \mathbf{P}$  and  $f - b \in \mathbf{P}$ , then  $f - \gamma \in \mathbf{P}$  for  $a \leq \gamma \leq b$ [121] has a proof of an elementary interlacing property.

[131] has a proof of an elementary interlacing property.

Jahnke [93, page 153] observes that af + bf' and cf + df' interlace if ad - bc is not zero, where f is a Bessel function.

Lemma 1.75 is a version of Kharitonov's theorem [13] for polynomials in  $\mathbf{P}^{pos}$ .

See [87] or [136] for inequalities similar to Example 1.69.

Corollary 1.83 is in [117], along with many nice applications.

#### Matrices preserving interlacing

In [70] they use the term *interlacing negative zeros* for a set of mutually interlacing polynomials (with negative zeros).

The proof of Proposition 3.22 is based on the argument in Holtz[81].

#### Homogeneous

We could also derive Newton's inequalities from Corollary 4.31 using f and f'.

#### Analytic

Theorem 5.36 is in [14]. The class  $\widehat{\mathbf{P}}$  is also known as the Laguerre-Pólya class  $\mathcal{L} - \mathcal{P}$  [39]. The polynomials  $T(1 + x)^n$  are called Jensen polynomials. A polynomial in  $\widehat{\mathbf{P}^{pos}}$  is called type I.

In [45] they essentially prove Kurtz' result (Theorem 5.19), but with a coefficient of 4.07 instead of 4.

#### **Linear Transformations**

In [37] they discuss various questions concerning linear transformations, in particular what they call linear operators operating on entire functions. For instance, they ask for a classification of all linear transformations that take polynomials with zeros in a set S (which might be complex) to polynomials

with all zero in a set T. In [36]Corollary 2.4 they determine all linear transformations T for which f and Tf interlace, but just for transformations that are multipliers. The answer is not expressed in terms of derivatives, but rather that the multiplier sequence is a nonconstant arithmetic sequence all of whose terms have the same sign.

[124] proves a special case of Lemma 7.87.

[24] proved the identity of Example 6.46 by using properties of the diamond product.

#### **Examples of Linear Transformations**

Szegö[168] is the source for all the facts about orthogonal polynomials.

Iserles and Norsett [90] call the inverse of the falling factorial the Laguerre transformation.

In [51] they study the interlacing properties of the polynomials sequences determined by recurrences  $p_{n+1} = 2r_n x p_n - (1 - x^2)p'_n$  for interlacing sequences  $(r_n)$  of coefficients.

See [22] for an identity involving Hadamard products and Laguerre polynomials that is similar to (7.10.6).

Some results involving linear transformations of polynomials have been expressed in umbral notation, which, to my mind, is just confusing. For example, Rota [148] defines modified Hermite polynomials  $\tilde{H}_n(x)$  by

$$\sum_{n=0}^{\infty} \tilde{H}_n(x) \frac{y^n}{n!} = e^{-x^2 + xy}$$

If we define  $T(x^n) = \tilde{H}_n(x)$  and  $M(x) = \sqrt{2}x$  then we have an elegant identity  $T^2 = M^{-1}TM$ . However, this is written in [148] as

$$\tilde{H}_{n}(\mathbf{H}) = 2^{n/2} \tilde{H}_{n}(\frac{x}{\sqrt{2}})$$

[10] and [153] discuss asymptotic properties of the roots of eigenpolynomials of arbitrary differential operators of the form (7.4.1).

#### Affine Transformations

Johnson [95] proves Lemma 8.26 by a messy algebraic analysis. The idea for the q-Laguerre polynomials comes from [156], who considers various properties of more general polynomials. There is no discussion about the roots, however.

The affine derivative was called  $L_{(q,b)}$  in [77].

#### **Complex Transformations**

There is an analytic proof [91] that the linear transformation  $x^n \longrightarrow T_n$  maps  $\mathbf{P}^{(-1,1)}$  to itself.

## NOTES

Wagner [175] refers to a folklore theorem that is Corollary 22.48 with  $\alpha = \pi/3$ .

#### **Polynomials in two variables**

Lemma 9.26 is a standard result about asymptotes. See [132] Lemma 9.130 can be found in [134]

#### Polynomials in several variables

## Matrices

The calculation in Section 17.2 is from [85, page 187].

Smith [88, 160] extends results on interlacing of Schur complements to singular Hermitian matrices using the Moore-Penrose inverse.

#### Stable

Lemma 21.38 is in [186, page 33], but the proof is more involved. [118] showed that the Bell polynomials are q-log-convex.

NOTES

[\*]

749

Bibliography

# **Bibliography**

- [1] Martin Aigner and Günter Ziegler, Proofs from the book, Springer, 1998. 1421
- [2] Michael Aissen, I. J. Schoenberg, and A. M. Whitney, On the generating functions of totally positive sequences. I, J. Analyse Math. 2 (1952), 93–103. MR 0053174 (14,732d) ↑113
- [3] Waleed A. Al-Salam and Mourad E. H. Ismail, Polynomials orthogonal with respect to discrete convolution, J. Math. Anal. and Appl. 55 (1976), 125–139. ↑184
- Bruce Anderson, Polynomial root dragging, American Mathematical Monthly (1993), 864–866.
   <sup>1</sup>42
- [5] T. Ando, *Totally positive matrices*, Linear algebra and its applications 90 (1987), 165–219. ↑59, 534
- [6] George E. Andrews, Richard Askey, and Ranjan Roy, Special functions, Encyclopedia of Mathematics and its Applications, vol. 71, Cambridge University Press, Cambridge, 1999. MR 2000g:33001 122
- [7] Bernard A. Asner Jr., On the total nonnegativity of the Hurwitz matrix, SIAM J. Appl. Math. 18 (1970), 407–414. MR 0260768 (41 #5391) <sup>649</sup>
- [8] Louis C. Barrett and Gerald E. Bendixen, Separation and interlacing theorems, Quart. Appl. Math. 23 (1965), 69–78. MR 31 #2523 ↑746
- Bernard Beauzamy and Jérôme Dégot, Differential identities, Trans. Amer. Math. Soc. 347 (1995), no. 7, 2607–2619. MR 1277095 (96c:05009) ↑
- [10] Tanja Bergkvist and Hans Rullgøard, On polynomial eigenfunctions for a class of differential operators, Research reports in mathematics, 2001. ↑510, 747
- [11] Dennis S. Bernstein, *Matrix mathematics*, Princeton University Press, Princeton, NJ, 2005. Theory, facts, and formulas with application to linear systems theory. MR 2123424 (2005i:93001) ↑583
- [12] Rajendra Bhatia, Infinitely divisible matrices, Americal mathematical monthly (2006). ↑457
- [13] S.P. Bhattacharyya, H Chapellat, and L.H. Keel, Robust control: The parametric approach, Prentice Hall, 1995. <sup>↑</sup>746
- [14] G. G. Bilodeau, Generating polynomials all of whose roots are real, Mathematics Magazine 64 (1991), 263–270. ↑122, 124, 745, 746
- [15] Ralph Philip Boas Jr., Entire functions, Academic Press Inc., New York, 1954. MR 16,914f <sup>†</sup>125
- [16] Julius Borcea, Petter Brändén, and Thomas M. Liggett, Negative dependence and the geometry of polynomials. arXiv:0707.2340. ↑563
- [17] Julius Borcea, Petter Brändén, and Boris Shapiro, Applications of stable polynomials to mixed determinants: Johnson's conjectures, unimodality and symmetrized fischer products. math.SP/0607755. ↑
- [18] \_\_\_\_\_, Classification of hyperbolicity and stability preservers: the multivariate Weyl algebra case. arXiv:math.CA/0606360. <sup>↑</sup>398, 505, 506, 565, 568
- [19] Julius Borcea, Petter Br andèn, and Boris Shapiro, Polya-schur master theorems for circular domains and their boundaries. arXiv:math/0607416. ↑

- [20] Peter Borwein and Tamás Erdélyi, Polynomials and polynomial inequalities, Graduate Texts in Mathematics, vol. 161, Springer-Verlag, New York, 1995. MR 97e:41001 1616
- [21] Jean-Christophe Bourin, Matrix versions of some classical inequalities. arXiv:math.OA/0601543. ↑330
- [22] L. R. Bragg, Trigonometric integrals and Hadamard products, Amer. Math. Monthly 106 (1999), no. 1, 36–42. ↑747
- [23] P. Brändén, Polynomials with the half-plane property and matroid theory. arXiv:math.CO/0605678. ↑389, 565
- [24] Petter Brändén, On linear transformations preserving the Pólya Frequency property. arXiv:math.CO/0403364. <sup>↑</sup>747
- [25] Francesco Brenti, Unimodal, log-concave and Pólya frequency sequences in combinatorics, Memoirs, vol. 81, American Mathematical Society, 1989. ↑113
- [26] Daniel Bump, Kwok-Kwong Choi, Pär Kurlberg, and Jeffrey Vaaler, A local Riemann hypothesis. I, Math. Z. 233 (2000), no. 1, 1–19. MR 2001g:11073a 147, 186, 497, 666
- [27] David Cardon, Sums of exponential functions having only real zeros, Manuscripta Math. 113 (2004), 307–317. ↑668
- [28] David A. Cardon and Pace P. Nielsen, Convolution operators and entire functions with simple zeros, Number theory for the millennium, i (urbana, il, 2000), 2002, pp. 183–196. MR 1956225 (2003m:30012) ↑
- [29] David A. Cardon, Convolution operators and zeros of entire functions, Proc. of the Amer. Math. Soc. (2002), 1725–1734. ↑668
- [30] J. M. Carnicer, J. M. Peña, and A. Pinkus, On some zero-increasing operators, Acta Math. Hungar. 94 (2002), no. 3, 173–190. MR 1905723 (2003c:26020) <sup>142</sup>
- [31] F. W. Carroll, A polynomial in each variable separately is a polynomial, Amer. Math. Monthly 68 (1961), 42. ↑417
- [32] T. S. Chihara, Orthogonal polynomials, Gordon-Breach, 1978. <sup>↑</sup>745, 746
- [33] Young-Bin Choe, James G. Oxley, Alan D. Sokal, and David G. Wagner, *Homogeneous multi-variate polynomials with the half-plane property*, Adv. in Appl. Math. **32** (2004), no. 1-2, 88–187. Special issue on the Tutte polynomial. MR **2037144** (2005d:05043) ↑565, 568
- [34] Fritz Colonius, Uwe Helmke, Dieter Prtzel-Wolters, and Fabian Wirth (eds.), Advances in mathematical systems theory, Systems & Control: Foundations & Applications, Birkhäuser Boston Inc., Boston, MA, 2001. MR 1787335 (2001e:93003) ↑619
- [35] Florian Constantinescu, Relations entre les coefficients de deus polynômes dont les racines se séparent, Časopis pro pěstováni matematiky (1964), 1–3. ↑745, 746
- [36] T. Craven and G. Csordas, On the betweenness condition of rolle's theorem, Rocky Mountain. J. of Math. 15 (1985), 721–728. <sup>†</sup>747
- [37] Thomas Craven and George Csordas, Composition theorems, multiplier sequences and complex zero decreasing sequences. to appear. <sup>†</sup>746
- [38] \_\_\_\_\_, On the number of real roots of polynomials, Pacific Journal of Mathematics 102 (1982), no. 1, 15–28. ↑
- [39] \_\_\_\_\_, Complex zero decreasing polynomials, Methods Appl. Anal. (1995), 420–441. ↑746
- [40] George Csordas, Marios Charalambides, and Fabian Waleffe, A new property of a class of jacobi polynomials, Proc. Amer. Math. Soc. 133 (2005), 3551–3560. ↑
- [41] George Csordas and Dimitar K. Dimitrov, Conjectures and theorems in the theory of entire functions, Numerical Algorithms (2000), 1–14. ↑294, 295
- [42] George Csordas and Jack Williamson, The zeros of Jensen polynomials are simple, Proc. of the Amer. Math. Soc. 49 (1975), no. 1, 263–264. <sup>↑</sup>488

- [43] Jean Pierre Dedieu and R. J. Gregorac, Corrigendum. Obreshkoff's theorem revisited: What convex sets are contained in the set of hyperbolic polynomials?, Journal of Pure and Applied Algebra 93 (1994), 111–112. ↑
- [44] Jean Pierre Dedieu, Obreshkoff's theorem revisited: What convex sets are contained in the set of hyperbolic polynomials?, Journal of Pure and Applied Algebra 81 (1992), 269–278. ↑
- [45] Dimitar K. Dimitrov and Juan Manuel Pena, Almost strict total positivity, strong Turan inequalities and a class of Hurwitz polynomials, 2004. ↑746
- [46] Dimitar K. Dimitrov and A. Sri Ranga, Monotonicity of zeros of orthogonal Laurent polynomials, Methods Appl. Anal. 9 (2002), no. 1, 1–11. MR 1948463 (2003i:42034) ↑45
- [47] Dimitar K. Dimitrov, Counterrexamples to a problem of Pólya and to a problem of Karlin, East. J. Approx. (1998), 479–489. ↑294, 295
- [48] V. V. Dodonov and V. I. Man'ko, New relations for two-dimensional Hermite polynomials, J. Math. Phys. 35 (1994), no. 8, 4227–4288. ↑
- [49] Kathy Driver and Peter Duren, *Trajectories of the zeros of hypergeometric polynomials f(-n,b;2b;z)* for b < −1/2, Const. Approx. 17 (2001), 169–179. ↑</p>
- [50] \_\_\_\_\_, Zeros of ultraspherical polynomials and the Hilbert-Klein formulas, J. Compu. Appl. Math. 135 (2001), 293–301. ↑42
- [51] F. Dubeau and J. Savoie, More on the roots of Euler-Frobenius polynomials, Ann. Sci. Math. Québec 22 (1998), no. 1, 31–42. <sup>198</sup>, 747
- [52] Charles F. Dunkl and Yuan Hu, Orthogonal polynomials of several variables, Encyclopedia of Mathematics and Its Applications, vol. 81, Cambridge University Press, 2001. ↑
- [53] Charles F. Dunkl and Yuan Xu, Orthogonal polynomials of several variables, Cambridge University Press, 2001. ↑424, 428
- [54] Ömer Eğecioğlu, Timothy Redmond, and Charles Ryavec, From a polynomial Riemann hypothesis to alternating sign matrices, Electron. J. Combin. 8 (2001), no. 1, Research Paper 36, 51 pp. (electronic). MR 2002k:05251 ↑527
- [55] Ömer Eğecioğlu and Charles Ryavec, Polynomial families satisfying a Riemann hypothesis, Proceedings of the thirty-second southeastern international conference on combinatorics, graph theory and computing (baton rouge, la, 2001), 2001, pp. 177–191. MR 2002m:11019 ↑
- [56] D. K. Faddeev and I. S. Sominskii, *Problems in higher algebra*, Translated by J. L. Brenner, W. H. Freeman and Co., San Francisco, 1965. MR 31 #1258 ↑746
- [57] F.M. Fernández, Generating functions for Hermite polynomials of arbitrary order, Physics Letters A 237 (1998), 189–191. ↑188
- [58] Steve Fisk, Aspects of stable polynomials. arXiv:0803.0286. <sup>↑</sup>619
- [59] \_\_\_\_\_, An introduction to upper half plane polynomials. arXiv:0711.4043. ↑589
- [60] \_\_\_\_\_, Coloring theories, Contemporary Mathematics, American Mathematical Society, 1989. ↑
- [61] Sergey Fomin and Andrei Zelevinsky, Total positivity: tests and parameterizations. math.L.A/9912128. ↑59, 112
- [62] Peter E. Frenkel, *Pfaffians, hafnians, and products of linear functionals.* arXiv:math.CA/07040028. ↑591
- [63] Paul A. Fuhrmann, A polynomial approach to linear algebra, Springer, 1996. <sup>†745</sup>
- [64] William Fulton, Eigenvalues, invariant factors, highest weights, and Schubert calculus, Bull. Amer. Math. Soc. (N.S.) 37 (2000), no. 3, 209–249 (electronic). MR 2001g:15023 ↑325
- [65] F. R. Gantmacher, The theory of matrices, Vol. 2, Chelsea, 1984. ↑745
- [66] Jürgen Garloff and David G. Wagner, Hadamard products of stable polynomials are stable, Journal of mathematical analysis and applications 202 (1996), 797–809. <sup>+</sup>645, 733, 745

- [67] George Gasper and Mizan Rahman, *Basic hypergeometric series*, Encyclopedia of mathematics and its applications, Cambridge University press, 1990. <sup>1</sup>122
- [68] C. D. Godsil, Algebraic combinatorics, Chapman and Hall, 1993. <sup>155</sup>, 745
- [69] Javier Gomez-Calderon and David M. Wells, Why polynomials have roots, College Mathematics Journal 27 (1996), no. 2, 90–94. ↑270
- [70] T. N. T. Goodman and Qiyu Sun, Total positivity and refinable functions with general dilation, Appl. Comput. Harmon. Anal. 16 (2004), no. 2, 69–89. MR MR2038266 (2004k:42056) ↑92, 746
- [71] J. H. Grace, The zeros of a polynomial, Proc. cambridge Philos. Soc. 11 (1902), 352–357. ↑
- [72] I. S. Gradshteyn and J. M. Ryzhjk, Table of integrals, series, and products, Academic Press, 1994. 123, 190, 488
- [73] Jiří Gregor and Jaroslav Tišer, On convex combinations of Hurwitz polynomials, Appl. Math. and Comp. Sci 6 (1996), no. 1, 33–47. ↑745
- [74] \_\_\_\_\_, On Hadamard powers of polynomials, Math. Control Signals Systems (1998), 372–378. ↑722
- [75] Leonid Gurvits, *Combinatorial and algorithmic aspects of hyperbolic polynomials*. arXiv:math.CO/0404474. ↑
- [76] James Haglund, Ken Ono, and David G. Wagner, *Theorems and conjectures involving rook polynomials with only real zeros*, Topics in number theory (university park, pa, 1997), 1999, pp. 207–221. MR 1691320 (2000d:05097) ↑537, 731
- [77] W Hahn, Über orthogonalpolynome, die q-diffenenzengleichungen, Math. Nachr. 2 (1949), 4–34. ↑747
- [78] G. H. Hardy, J. E. Littlewood, and G. Pólya, *Inequalities*, Cambridge Mathematical Library, Cambridge University Press, Cambridge, 1988. Reprint of the 1952 edition. MR 944909 (89d:26016) ↑110
- [79] J. William Helton and Victor Vinnikov, Linear Matrix Inequality Representation of Sets. arXiv:math.OC/0306180. ↑282
- [80] V. E. Hoggatt Jr. and Marjorie Bicknell, Roots of Fibonacci polynomials, Fibonacci Quart. 11 (1973), no. 3, 271–274. MR 0323700 (48 #2056) ↑640
- [81] Olga Holtz, Hermite-Biehler, Routh-Hurwitz, and total positivity, Linear Algebra Appl. 372 (2003), 105–110. MR 1 999 142 ↑674, 746
- [82] \_\_\_\_\_, M-matrices satisfy Newton's inequalities, Proceeding of the AMS (2004August). ↑
- [83] Lars Hörmander, An introduction to complex analysis in several variables, Van Nostrand, 1966.

   <sup>†</sup>447
- [84] Alfred Horn, Eigenvalues of sums of Hermitian matrices, Pacific J. Math. 12 (1962), 225–241. MR 25 #3941 ↑
- [85] Roger A. Horn and Charles R. Johnson, Matrix analysis, Cambridge University Press, 1985. 1343, 347, 545, 556, 748
- [86] \_\_\_\_\_, Topics in matrix analysis, Cambridge University Press, 1991. ↑
- [87] Alan Horwitz, On the ratio vectors of polynomials, J. Math. Anal. Appl. 205 (1997), no. 2, 568– 576. MR 1428370 (97j:12001) ↑746
- [88] Shu-an Hu and Ronald L. Smith, The Schur complement interlacing theorem, Siam J. Matrix Anal. Appl. 16 (1995), no. 3, 1013–1023. ↑748
- [89] Ljubomir Iliev, Laguerre entire functions, Bulgarian Academy of Sciences, 1987. ↑374, 498, 615
- [90] A. Iserles and S. P. Norsett, Zeros of transformed polynomials, Math. Anal. (1990), no. 2, 483– 509. ↑160, 212, 747

- [91] A. Iserles and E. B. Saff, Zeros of expansions in orthogonal polynomials, Math. Proc. Camb. Phil. Soc. 105 (1989), 559–573. ↑747
- [92] Mourad Ismail and Martin Muldoon, A discrete approach to monotonicity of zeros of orthogonal polynomials, Transactions of the AMS 323 (1991), 65–78. ↑
- [93] Jahnke-Emde-Lösch, Tables of higher functions, sixth edition, McGraw-Hill, 1960. <sup>↑</sup>746
- [94] L. Jódar and E. Defez, A connection between Laguerre's and Hermite's matrix polynomials, Appl. Math. Lett. 11 (1998), no. 1, 13–17. ↑
- [95] Charles R. Johnson, Interlacing polynomials, Proceedings of the American Mathematical Society 100 (1987), 401–404. <sup>†</sup>40, 457, 745, 747
- [96] C.R. Johnson, A characteristic polynomial for matrix pairs, Linear and multilinear algebra 25 (1989), 289–290. ↑
- [97] Warren P. Johnson, *The curious history of Faà di Bruno's formula*, Amer. Math. Monthly **109** (2002), no. 3, 217–234. MR **1903577 (2003d:**01019) ↑39, 41
- [98] Victor Kac and Pokman Cheung, Quantum calculus, Universitext, Springer-Verlag, New York, 2002. MR 1865777 (2003i:39001) ↑219
- [99] Samuel Karlin and Gabor Szego, On certain determinants whose entries are orthogonal polynomials, J. Analyse Math. 8 (11960/61), 1–157. ↑730
- [100] Samuel Karlin, Total positivity, Stanford University Press, 1968. <sup>294</sup>, 299
- [101] Deok Kim and Kil Kwon, On a conjecture of Karlin and Szego, Proceedings of the American Mathematical Society 124 (1996), 227–231. ↑745
- [102] Alexander A. Klyachko, Random walks on symmetric spaces and inequalities for matrix spectra, Linear Algebra Appl. 319 (2000), no. 1-3, 37–59. MR 2002f:15023 ↑
- [103] Roelof Koekoek and Renè F. Swarttouw, The Askey-scheme of hypergeometric orthogonal polynomials and its q-analog, Technical Report 94-05, Technische Universiteit Delft, 1994. <sup>244</sup>
- [104] V. Kostov and B. Shapiro, On arrangements of roots for a real hyperbolic polynomial and its derivatives. to appear. ↑
- [105] Yu. Kozitsky, P. Oleszczuk, and L. Wolowski, Infinite Order Differential Operators in Spaces of Entire Functions, J. Math. Anal. Appl. 277 (2003), 423–437. arXiv:math.FA/0311213. ↑
- [106] Yuri Kozitsky and Lech B. Wolowski, Laguerre Entire Functions and Related Locally Convex Spaces. arXiv:math.CV/9812111. ↑454
- [107] Ilia Krasikov, On zeros of polynomials and allied functions satisfying second order differential equations. arXiv:math.CA/0211190. ↑137
- [108] C. Krattenthaler, Advanced determinant calculus, Sm. Lothar. Combin 42 (1999). ↑299, 582, 652
- [109] Greg Kuperberg, Special moments. arXiv:math.PR/0408360. 130
- [110] David C. Kurtz, A sufficient condition for all the roots of a polynomial to be real, Amer. Math. Monthly (1992), 259–263. <sup>↑</sup>128
- [111] Peter Lancaster and Panayiotis Psarrakos, The numerical range of self-adjoint quadratic matrix polynomials, SIAM J. Matrix Anal. Appl. 23 (2001/02), no. 3, 615–631 (electronic). MR 1896809 (2003e:15032) ↑559
- [112] Adrian S. Lewis, Pablo A. Parrilo, and Motakuri V. Ramana, The Lax conjecture is true. arXiv:math.OC/0304104. <sup>↑</sup>282
- [113] ChiKwong Li and Stephen Pierce, Linear preserver problems, Amer. Math. Monthly 108 (2001), no. 7, 591–605. ↑
- [114] Elliott H. Lieb and Robert Seiringer, Equivalent forms of the bessis-moussa-villani conjecture, Journal of Statistical Physics 115 (2004), no. 1-2, 185–190. ↑351
- [115] Elliott H. Lieb and Alan D. Sokal, A general Lee-Yang theorem for one-component and multicomponent ferromagnets, Comm. Math. Phys. 80 (1981), no. 2, 153–179. MR 623156 (83c:82008) <sup>↑</sup>453

- [116] John B. Little, On the zeros of two families of polynomials arising from certain rational integrals, Rocky Mountain J. Math. 35 (2005), no. 4, 1205–1215. MR 2178985 ↑45
- [117] Li Liu and Yi Wang, A unified approach to polynomial sequences with only real zeros. arXiv:math.CO/0509207. <sup>↑</sup>93, 94, 724, 746
- [118] \_\_\_\_\_, On the log-convexity of combinatorial sequences. arXiv:math.CO/0602672. ↑650, 651, 748
- [119] Franz Locher, Stability tests for linear difference forms, International series of numerical mathematics 112 (1993), 215–223. ↑746
- [120] C.C. MacDuffee, *Theory of equations*, Wiley, 1954. ↑
- [121] Morris Marden, The geometry of the zeros of a polynomial in a complex variable, American Mathematical Society, 1949. ↑661
- [122] A.S. Markus, Introduction to the spectral theory of polynomial operator pencils, Translations of Mathematical Monographs, vol. 71, Americam Mathematical Society, 1988. ↑547
- [123] Horacio J. Marquez and Panajotis Agathoklis, On the existence of robust strictly positive real rational functions, IEEE Trans. Circuits Systems I Fund. Theory Appl. 45 (1998), no. 9, 962– 967. MR 1647619 (99h:93039) ↑
- [124] Gisli Másson and Boris Shapiro, On polynomial eigenfunctions of a hypergeometric-type operator, Experiment. Math. 10 (2001), no. 4, 609–618. MR 2003e:33038 ↑745, 747
- [125] Anne de Médicis, Dennis W. Stanton, and Dennis E. White, The combinatorics of q-Charlier polynomials. arXiv:math.CA/9307208. ↑729
- [126] H. G. Meijer, Coherent pairs and zeros of Sobolev-type orthogonal polynomials, Indag. Mathem. 2 (1993), 163–176. ↑
- [127] Todor P. Mitev, New inequalities between elementary symmetric polynomials, JIPAM. J. Inequal. Pure Appl. Math. 4 (2003), no. 2, Article 48, 11 pp. (electronic). MR 1994261 (2004d:26015) ↑
- [128] M. E. Muldoon, Properties of zeros of orthogonal polynomials and related functions, J. Compu. Appl. Math. 48 (1993), 167–186. ↑42
- [129] editor (2003) N. J. A. Sloane, *The on-line encyclopedia of integer sequences*. http://www.research.att.com/njas/sequences/. ↑
- [130] Constantin P. Niculescu, A new look at Newton's inequalities, JIPAM. J. Inequal. Pure Appl. Math. 1 (2000), no. 2, Article 17, 14 pp. (electronic). MR MR1786404 (2001h:26020) ↑
- [131] Albert Nijenhuis, On permanants and the zeros of rook polynomials, J. Comb. Theory (A) 21 (1976), 240–244. ↑746
- [132] Jeffrey Nunemacher, Asymptotes, cubic curves, and the projective plane, Math. Magazine 72 (1999), no. 3, 183–192. ↑462, 748
- [133] P.S.S.N.V P. Rao, A characterization of merely subdefinite matrices and related results, Sankhya Ser. A 39 (1977), no. 4, 387–395. <sup>↑</sup>346
- [134] Beresford Parlett, The symmetric eigenvalue problem, Prentice-Hall, 1980. 1303, 748
- [135] Merrell L. Patrick, Extensions of inequalities of the Laguerre and Turán type, Pacific J. Math. 44 (1973), 675–682. MR 47 #3725 ↑
- [136] Gideon Peyser, On the roots of the derivative of a polynomial with real roots, Amer. Math. Monthly 74 (1967), 1102–1104. MR 0227332 (37 #2917) ↑149, 746
- [137] G. Pólya and J. Schur, Uber zwei arten von factorenfolgen in der theorie der algebraischen gleichungen, Journal fur die reine und angewandte mathematik band 144 (1914), 89–113. <sup>1</sup>135, 488
- [138] G. Pólya and G. Szego, Problems and theorems in analysis, Vol. 2, Springer-Verlag, 1972. <sup>1</sup>122, 124, 130, 171, 199, 746
- [139] \_\_\_\_\_, Problems and theorems in analysis, Vol. 1, Springer-Verlag, 1972. 130, 616

- [140] G. Pólya, Bemerkung uber die integraldarstellung der Riemannssche ζ-function, Acta Math. 48 (1926), 305–317. ↑
- [141] Q. I. Rahman and G. Schmeisser, Analytic theory of polynomials, Mathematical Society Monographs, Oxford Science Publications, 2002. ↑5, 20, 745
- [142] Timothy Redmond, A class of polynomial Riemann hypotheses. in prepartion. †147, 148, 726
- [143] Bruce Reznick, Sums of even powers of real linear forms, Memoirs, American mathematical society, 1992. ↑
- [144] \_\_\_\_\_, An inequality for products of polynomials, Proceedings of the American mathematical society 117 (1993), 1063–1073. ↑
- [145] J. Rivlin, The Chebychev polynomials, Wiley, 1974. ↑
- [146] Steve Roman, The umbral calculus, Academic Press, 1984. <sup>†</sup>144, 164, 188, 190, 490, 499
- [147] Shmuel Rosset, Normalized symmetric functions, Newton's inequalities and a new set of stronger inequalities, Amer. Math. Monthly 96 (1989), no. 9, 815–819. MR 1033349 (91e:05083) ↑117
- [148] Gian-Carlo Rota, Finite operator calculus, Academic Press, 1975. 164, 182, 183, 747
- [149] Lee Rubel, Entire and meromorphic functions, Springer-Verlag, 1996. <sup>125</sup>
- [150] David Ruelle, Grace-like polynomials, Foundations of computational mathematics (hong kong, 2000), 2002, pp. 405–421. MR 2021990 (2004k:30014) ↑581, 582
- [151] G. M. Shah, On the zeros of Van Vleck polynomials, Proc. Amer. Math. Soc. 19 (1968), 1421– 1426. MR 37 #6511 ↑17
- [152] A. Shapiro, S. Kaplan, and M. Teicher, Several Applications of Bezout Matrices. arXiv:math.AG/0601047. ↑308
- [153] B. Shapiro and G. Másson, On polynomial eigenfunctions of  $\frac{d^k}{dx^k}q(x)$ , IHES-preprint (2001).  $\uparrow$ 747
- [154] Boris Shapiro and Volkmar Welker, Combinatorics and topology of stratifications of the space of monic polynomials with real coefficients, Result. Math 33 (1998), 338–355. ↑
- [155] T. Sheil-Small, Complex polynomials, Cambridge Studies in Advanced Mathematics, vol. 75, Cambridge University Press, Cambridge, 2002. MR 1 962 935 ↑674, 745
- [156] D. P. Shukla, q-angelescu polynomials, Publications de l'Institut Math/'ematique 29 (1981), no. 43, 205–213. <sup>†</sup>747
- [157] Avram Sidi, Numerical quadrature and non-linear sequence transformations; unified rules for efficient computation of integrals with algebraic and logarithmic endpoint singularities, Math. Comp. 35 (1989), 851–874. ↑
- [158] Rodica Simion, A multiindexed Strum sequence of polynomials and unimodality of certain combinatorial sequences, J. Comb. Theory(A) 36 (1984), 15–22. ↑745
- [159] D. Singer, A bijective proof of Borchardt's identity, Electon. J. Combin. 11 (2004), no. R48. 1457
- [160] Ronald L. Smith, Some interlacing properties of the Schur complement of a Hermitian matrix, Linear algebra and its applications 177 (1992), 137–144. ↑535, 748
- [161] David E. Speyer, Horn's problem, Vinnikov curves, and the hive cone. preprint. <sup>257</sup>, 322, 324
- [162] Richard Stanley, Log-concave and unimodal sequences in algebra, combinatorics, and geometry, Annals New York Academy of Sciences 576 (1989), 500–535. ↑
- [163] \_\_\_\_\_, Enumerative combinatorics, Vol. II, Cambridge University Press, 1997. ↑
- [164] J. Michael Steele, The Cauchy-Schwarz master class, MAA Problem Books Series, Mathematical Association of America, Washington, DC, 2004. An introduction to the art of mathematical inequalities. MR 2062704 (2005a:26035) ↑110
- [165] A. Stoyanoff, Sur un theorem de M Marcel Riesz, Nouvelles Annales de Mathematique 1 (1926), 97–99. ↑234
- [166] P. K. Suetin, Orthogonal polynomials in two variables (analytical methods and special functions), Taylor and Francis, 1999. 1424, 426, 427

- [167] Otto Szász, On sequences of polynomials and the distribution of their zeros, Bull. Amer. Math. Soc. 49 (1943), 377–383. MR 0008274 (4,273c) ↑110
- [168] Gabor Szego, Orthogonal polynomials, American Mathematical Society, 1939. <sup>†</sup>44, 155, 186, 202, 234, 433, 477, 641, 736, 737, 747
- [169] \_\_\_\_\_, On an inequality of P. Turan concerning Legendre polynomials, Bull. Amer. Math. Soc (1948), 401–5. ↑124
- [170] Walter Van Assche and Els Coussement, Some classical multiple orthogonal polynomials, J. Comput. Appl. Math. 127 (2001), no. 1-2, 317–347. Numerical analysis 2000, Vol. V, Quadrature and orthogonal polynomials. MR 2001i:33012 ↑209, 210
- [171] V. A. Vassiliev, Topology of spaces of functions without compound singularities, Funct. Anal. Appl. 23 (1989), 277–286. ↑
- [172] \_\_\_\_\_, Complements of discriminants of smooth maps: topology and applications, Translations of Math. Mono., vol. 98, A.M.S., 1992. ↑
- [173] Gert Vegter, The apolar bilinear form in geometric modeling, Math. Comp. 69 (2000), no. 230, 691–720. MR 1654006 (2000i:65018) ↑
- [174] Victor Vinnikov, Selfadjoint determinantal representations of real plane curves, Math. Ann. 296 (1993), no. 3, 453–479. MR 94e:14038 ↑282
- [175] David G. Wagner, Enumeration of spanning subgraphs with degree constraints. arXiv:math.CO/0412059. ↑748
- [176] \_\_\_\_\_, Zeros of genus polynomials of graphs in some linear families. preprint. ↑
- [177] \_\_\_\_\_, Zeros of reliability polynomials and f-vectors of matroids. arXiv:math.CO/9802047. ↑
- [178] \_\_\_\_\_, Total positivity of Hadamard products, J. Math. Anal. Appl. 163 (1992), 459–483. <sup>↑</sup>167, 406, 745
- [179] \_\_\_\_\_, Multipartition series, Siam J. Discrete Math. 9 (1996), no. 4, 529–544. <sup>745</sup>
- [180] Peter Walker, Separation of zeros of translates of polynomials and entire functions, J. Math. Anal. Appl. 206 (1997), no. 1, 270–279. MR 97m:30004 ↑236
- [181] \_\_\_\_\_, Separation of zeros of translates of polynomials and entire functions, Journal of Mathematical Analysis and Applications 206 (1997), 270–279. ↑
- [182] Long Wang, Zhizhen Wang, and Wensheng Yu, Stability of polytopic polynomial matrices, 2002. arXiv:math/0202238. ↑
- [183] Steven Weller, W. Moran, Brett Ninness, and A.D. Pollington, Sampling zeros and the Euler-Frobenius polynomials, IEEE transactions on automatic control 46 (2001), no. 2, 340–343. <sup>↑</sup>198
- [184] Kohhei Yamaguchi, Spaces of polynomials with real roots of bounded multiplicity, J. Math. Kyoto Univ. 42 (2002), no. 1, 105–115. ↑
- [185] Wensheng Yu, Long Wang, and Jürgen Ackermann, Robust strictly positive real synthesis for polynomial families of arbitrary order, Sci. China Ser. F 47 (2004), no. 4, 475–489. MR 2126240 (2006a:93044) ↑
- [186] Jerzy Zabczyk, Mathematical control theory: an introduction, Systems & Control: Foundations & Applications, Birkhäuser Boston Inc., Boston, MA, 1992. MR 1193920 (93h:49001) ↑748
- [187] Stephen M. Zemyan, On the zeroes of the Nth partial sum of the exponential series, Amer. Math. Monthly 112 (2005), no. 10, 891–909. MR 2186832 ↑638, 689
- [188] A.S. Zhedanov, Gauss sums and orthogonal polynomials, Intl. J. of Modern Physics A 12 (1997), no. 1, 289–294. 118

# Index

Symbols	
$\langle \underline{\mathbf{x}} \rangle_{\mathrm{n}} \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots $	
$(\underline{\mathbf{x}})_{n}$	
D	
$\widehat{\mathbf{P}^{\text{pos}}}$	
<b>P</b> <sub>2</sub>	
$\overline{\mathbf{P}}_{d}$	
$\overline{\mathbf{P}}_{d}(\mathbf{n})$	
Pd	
interior	
$\mathbb{P}_d(\mathfrak{n})$	
P <sub>d</sub>	
$\mathbf{P}_{d}^{\text{pos}}$	
$\mathbf{P}_{2}^{\mathrm{pos}}$	
<b>P</b> <sup>2</sup>	
<b>P</b> <sup>I</sup>	
<b>P</b> <sup>pos</sup>	
<b>P</b> <sup>alt</sup>	
$\mathbf{P}^{\pm}$	
<b>Sub</b> <sub>d</sub>	
$SL_2(\mathbb{R})$	
TP	
TP <sub>2</sub>	
f <sup>H</sup>	
I	
<b>I</b>	
J <sub>i</sub> see Bessel function	
EXP	
f <sup>H</sup>	
$\int \dots \dots \dots 194$	
СР()534	
diag454	
diag <sub>1</sub> 454	
<b>Sub</b> <sub>2</sub> 257	
«2	
≫2	
defined for matrices534	

# A

additive Horn problem	325
adjugate	555
affine transformation	145
Appell polynomials	. 429
Appell sequence	166
arg sum	.616
arithmetic-geometric inequality	110

# В

Bôcher-Walsh theorem 681
Bell polynomial 41, 360
Bell polynomials184, 192, 357, 657,
659
Bernoulli polynomials 167
Bernstein polynomials
Bessel function 124, 127, 129, 378, 494,
500, 501, 752
Bezout identity74
Bezout matrix 311, 313, 748
Bezout polynomial647
bilinear forms in $\overline{\mathbf{P}}_{2d}$
binomial type164
Borchardt's identity
boundary polynomial problem325
_

cancellation
Cauchy's identity
Cauchy-Binet identity 114, 116
Cauchy-Schwartz inequality 663
characteristic polynomial 33, 534
Charlier polynomials 191, 192, 231
and stable polynomials652
diamond product
two variable438
Chebyshev polynomials 170, 204, 523
question
Christoffel-Darboux formula647
coefficients
alternating signs
leading 26, 27, 497, 513
Newton's inequalities 109
of interlacing polynomials116
sign conditions
common
interlacing
common interlacing 3. 5. 21. 22. 55.
81, 384
hyperbolic matrix polynomials
553
in <b>Hyp</b> <sub>1</sub> 553
Commutator 45, 732
commuting diagrams
composition
Charlier
falling factorial
Hermite
is ubiquitous
Laguerre 193
condition
positivity
condition number
cone
cone of interlacing
conjugate
convex
set in $\mathbb{R}^d$
in i-th coordinate
convolution
mutually interlacing 67
cross polytope 47

# 

# D

degree
has constant24
of x <sub>i</sub>
of a monomial
possible sequences
preserves
total
determinants
and interlacing squares 27, 28, 30
and low rank matrices387
and polynomials in $P_2 \dots 385$
and realizing $P_2 \dots 375$
and sign interlacing
and signs of coefficients 31
and the resultant
in $P_2$
involving f and f' $\dots 29, 30$
involving $f, f_x, f_y \dots 321$
of coefficients9, 117
of coefficients in $\mathbf{P}_2^{\text{pos}}$
not totally positive 303
of coefficients in $\mathbf{P}_2 \dots \dots 299$
of coefficients of $\mathbf{P}_2^{\text{pos}}$
of translates
diagonal polynomials454
diagonals
interlacing 144
diamond product
Charlier polynomials
exponential-question
falling factorial-question 731
general
Hermite
Laguerre-question
original signed 410
difference equation 191
difference equation
differential operators 180,290,340
514
direction of interlacing 403
division algorithm 73

domain

C	of a li	near	trans	form	ation	• • •	150

# Ε

Ehrhart polynomial
eigenpolynomial210, 515, 738
equivariant
Euclidean algorithm
Euler polynomials167, 659
Euler-Frobenius polynomials 200
Eulerian polynomials 199
question 730
even and odd parts 58
in U <sub>2</sub> 609
even part 85, 201, 205, 252, 456, 544,
689
not arbitrary 203
exponential function
exponential generating function193
exponential operator 51, 254
exponential polynomials184
exponential transformation 194
extension
of 2 interlacing polynomials . 294
of 3 polynomials 703
of two polynomials331
F
f <sup>neg</sup> 48
f <sup>pos</sup> 48
f(x,D)
factorization property
falling factorial 131, 161, 184, 185, 193,
230, 232, 442, 703
diamond product 413
$question \dots / 2/, / 30, / 34, / 44$
question / 2/, / 30, / 34, / 44 family
family of polynomials

Gamma function	125, 195
----------------	----------

Gauss-Lucas theorem 666, 681
Gaussian binomial coefficient 244
generating function 492, 494
generic polynomials in $P_2 \dots 304$
Grace polynomial394, 528, 589
Grace's theorem 401
graph
of P <sub>2</sub>
of P <sub>d</sub> 339
group

Н	
Hadamard product 176, 182, 194, 283	,
288, 396, 494, 633	<b>`</b>
for cones	3 ₄
and exponential 194	1
for $\mathcal{H}_1$	1
for ideals	•
general 353, 505	2
generalized 176	5
generating function 494	1
in $\mathbf{\tilde{P}}$	7
in $\mathbf{P}_2^{\text{pos}}$	9
in J 613	3
in P <sub>d</sub>	9
matrix polynomials 562, 564	4
proof	3
variant	3
generating function 494	4
Hankel matrix	7
Hartog's Theorem 452	2
Heine-Stieltjes polynomials 17	7
Hermite polynomial 523	3
Hermite polynomials189–191, 211	,
213, 278, 361, 470, 495, 501	,
504, 686, 690	
diamond product 412	2
equivariant690	0
generating function	4
identities	8
in $\mathbf{P}_{d}$	0
question	3
mapping in $\mathcal{H}_1$	5
modified	3
question	1
Rodrigues' formula	9

transformations 277, 280, 509
Hermite-Biehler 628, 680
Hermite-Biehler problem
Hermite-Poulin Theorem 292
Hermitian matrices
hive
hive theorem
homogeneous part 264, 374
homogenized polynomial
homogenizing
in d variables
multiply by y 628
Horn problem
additive
Horn's conjecture
Hurwitz matrix
Hurwitz polynomial
Hurwitz's theorem
for $\overline{\mathbf{P}}_{2}^{\text{pos}}$
question
for Z/2Z 202, 689
for Z/dZ 114, 202, 688
hyperbolic matrix polynomials546
common interlacing 553
interlacing549, 553
orthogonal sequences 558
root zones
sign interlacing560
Hypergeometric polynomials 208
Hypergeometric series
hypergeometric series
51 U

I
index set
induced transformation 277, 406
inequalities
arithmetic-geometric 110
for $P_2$
inequality
Muirhead 111
integral transformation
interlacing 2
A
coefficients of 116, 117
common3, 5, 21, 22, 55, 548
cone 420

doesn't preserve240
for ideals696
geometric meaning in $P_2 \dots 274$
hyperbolic matrix polynomials
549, 553
in <b>P</b> <sup>pos</sup> 54
in $\mathbf{P}^{pos}$ and $\mathcal{H}_1 \dots \dots \dots 642$
in $\overline{\mathbf{P}}_{d}$
in $\mathbf{P}_2$
in $\overline{\mathbf{P}}_{d}$
in P <sub>d</sub>
in <b>Sub</b> <sub>d</sub>
mutually 61, 240, 266
of exponentials
of squares
positive 52, 625, 662
preserves 3, 24
sign 6
strict 2
strict, in P <sub>2</sub> 274
transformations
interlacing square
interlacing family
interlacing square
interpolation12
basis 105, 233, 294
mutually interlacing
<i>,</i> 0

# J

Jacobi matrix	310, 660
Jacobi polynomials	205, 503
question	730, 731
Rodrigues' formula	429
Jacobi-Piñeiro polynomials	213

## Κ

Kakeya	622
Karlin's conjecture	298
Kronecker product	547

# L

Laguerre inequality	. 30, 136
Laguerre polynomials173,	193, 213,
471, 490, 500, 501, 51	17 <i>,</i> 659
q	246
generating function	494

identities 163, 164
mapping in $\mathcal{H}_1 \dots \dots \dots 635$
monic
question 733
Rodrigues' formula 193, 246, 429
Rodrigues' formula for $P_d$ 430
Laplace transform
Laurent orthogonal polynomials45,
213
Legendre polynomial
Legendre polynomials124, 205
generalized 432
question
Leibnitz 16, 383, 480
Linear combinations
$f(x) + g(y) \dots 387$
linear transformation
complex674
doesn't induce on $P_2 \dots 277$ , 406
doesn't preserve interlacing 25
has constant degree 24
preserves degree
preserves interlacing 3, 24
preserves roots3
preserves sign of leading coeffi-
cient
satisfies induction
satisfies induction on $\mathbf{P}_2^{\text{pos}} \dots 406$
singular value165
Liouville's theorem
locally interlacing family540
of matrices 540
Loewner-Whitney theorem61
log concave
in $\mathbf{P}_2^{\mathrm{pos}}$
strict
lower truncation operator254

### М

M-invariant587
Möbius transformation . 158, 201, 445
MacLaurin's inequality 110
majorization 110
matrix
negative subdefinite 346, 349, 413
preserving initial segments 698

#### Ν

Narayana polynomials ..... 45, 659 negative part of a polynomial ..... 48 negative subdefinite matrix . 346, 349, 413 Newton graph ...... 119 Newton quotient ..... 108, 130 Newton's inequalities ... 73, 107, 115, 124, 129, 139, 203, 283, 696 and stability.....649 and zero coefficients in  $\overline{\mathbf{P}}_2 \dots 388$ converse ...... 129, 130 cubi . 116

cubic.	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	116	,
for $\mathbf{P}^{\mathbb{A}}$																								249	,

for interlacing polynomials118	polynomials
hyperbolic matrix polynomials	mutual
565	of binor
interlacing version of344	Appell
matrix polynomials 566	Bell
norm	Charlie
of a matrix 555	Chebys
NX matrix	Euler
recurrence relations	Euler-F

0
odd part
open map 24, 25
operator
differential 180, 290, 340, 514
exponential 254
reversal 175
order ideal
Orthogonal matrices
orthogonal polynomials
analogs in $\mathbf{P}_2$
Appell 429
Charlier 191, 192
Hermite
higher analogs
hyperbolic matrix polynomials
558
Laurent
stable 646

# Р

<b>P</b> -positive semi-definite728
Pólya-Schur 125
partial fractions 13, 29, 222
partial sums 457
permanent 542, 588
question 737
Perron's theorem
Pincherlet derivative
Pochhammer symbol 123
polar derivative105, 142, 400
in d variables
polynomial
negative part
positive part48
stable 58, 203, 266, 699, 706

mutually interlacing 276, 728
of binomial type164
Appell 429
Bell
Charlier 231
Chebyshev 204, 730
Euler
Euler-Frobenius
Eulerian 199, 730
Hermite
homogeneous part 264
integral not in <b>P</b> 4
Jacobi
Jacobi-Piñeiro
Laguerre . 146, 173, 193, 500, 501,
659
Legendre
generalized 432
mutually interlacing 240, 531
Narayana
of Meixner type157, 184, 189, 191
q-Hermite 247, 734
Rogers-Szego121
totally positive coefficients 736
totally positive coefficients 301
positive coefficients
not stable 637
positive definite 534
ordering of matrices 462, 542
positive interlacing . 52, 374, 379, 448,
625, 662, 690, 728
positive part of a polynomial48
positive semi-definite matrix 386
positivity
of substitutions323
positivity condition 264, 316
power method
principle submatrices
principle submatrix . 31, 286, 364, 535
product
diamond 167, 234, 410, 504
purely imaginary roots46
0
Q

q-exponential ...... 174, 246

q-Hermite polynomials 247
question
q-log-concave657
q-log-convex657
quadratic form413
quadratic form 346, 350, 414, 433, 435,
449-451
quadratic forms346
quadratics in $P_2$
quadrilateral inequalities 268, 327
quantitative sign interlacing 12, 15,
420, 422, 751
· ·

# R

range
of a linear transformation 150
rank of a matrix
real upper half plane polynomial.571
rearrangement lemma
recurrence
Charlier polynomials 192
Chebyshev
Euler
Euler-Frobenius
for $x^i \mapsto (\underline{\alpha})_i x^i \dots 173$
Hermite
higher dimensional
Hypergeometric209
Laguerre
Mexier 157
monic Charlier
orthogonal polynomials 156
orthogonal polynomials inverse
157
rising factorial186
two dimensional 434
with polynomial coefficients.523
recurrence relations
and rational functions 528
from matrices
recurrences
from P <sub>3</sub> 526
resultant 419, 420
reversal constant
reverse
conjugated by 175, 190

in Hyp <sub>1</sub> 549
in $P_2$
in P <sub>d</sub>
of a polynomial 105, 159, 161,
175, 534, 540
rhombus inequalities 268, 303, 327,
376
for Hyp <sub>2</sub> 567
matrix polynomials
right half plane polynomial 571
rising factorial 131, 186, 193, 230, 231,
278, 291, 441, 442, 445, 495,
734
diamond product 413
guestion
Rodrigues' formula 429, 464
q-Laguerre
affine233
Euler
generalized Legendre 432
Hermite 189, 237, 429, 464
Jacobi
Laguerre 192, 193, 246, 429
Laguerre in $P_d$
orthogonal polynomials in $P_d$ 429
Rogers-Szego polynomials 121
Rolle's Theorem 4, 283, 495, 686
root zone
root zones
Ruelle polynomials
1 /

S
satisfies induction406
on $\mathbf{P}_{2}^{\text{pos}}$
Schur complement
Schur's Theorem
Schur-PólyaTheorem292
Schur-Szegö
theorem
Schur-Szegömap400
sign change
sign interlacing5, 6, 131
for A 223
hyperbolic matrix polynomials
560
-

ijuga by

signs

patterns of coefficients54
simultaneous negative pair
of matrices
of polynomials 415
sin and cos 124, 125, 174, 453, 502, 729
skew-symmetric 659
solution curves270, 271, 274, 284, 305,
308, 415, 437, 442, 447
square
interlacing19, 27, 481
stable
polynomial58
stable matrix polynomials661
Stirling's formula 127
strict interlacings2
strictly totally positive60
subdefinite matrix
negative 346, 349, 413
substitution
arising from determinants 257
x-substitution257, 314
Sylvester's identity 300
Sylvester-Gallai
symmetric derivative
symmetric polynomials

# Т

$T_{1/z} \dots 161, 280, 445, 446, 496, 497, 501$
$\Gamma(1+x)^n \dots 493$
Taylor series 278, 283, 298, 321, 342,
392, 409, 410, 464, 490, 500,
693
Taylor's theorem404
tensor product
Toeplitz matrix 297
totally non-negative
totally positive 60, 68, 73, 80, 114, 118,
302
coefficients
coefficients of <b>P</b> <sup>pos</sup>
function
function on $\mathbb{R}^2_+$
question
recurrence relations
strongly, of order 2 80
totally positive <sub>2</sub> 65, 89
~ .

.54	transformation
	exponential 194
115	induced 277, 406
115	integral 194
729	truncation
559	operator254
05,	Turan's inequality 124

# U

umbral notation	. 753
unimodular	58
upper half plane polynomial	. 571

# v

-	
Van Vleck polynomials	17
Vandermonde vector	311
very hyperbolic	179

### W

Whitney theorem	50
Wronskian	48

# .

Ζ	
Z-note-chap-20-1	.596
zero coefficients	
in P <sub>2</sub>	. 388

# 766