ON THE ALGEBRAIC HOLONOMY OF STABLE PRINCIPAL BUNDLES

INDRANIL BISWAS

ABSTRACT. Let E_G be a stable principal G-bundle over a compact connected Kähler manifold, where G is a connected reductive linear algebraic group defined over \mathbb{C} . Let $H \subset G$ be a complex reductive subgroup which is not necessarily connected, and let $E_H \subset E_G$ be a holomorphic reduction of structure group to H. We prove that E_H is preserved by the Einstein–Hermitian connection on E_G . Using this we show that if E_H is a minimal reductive reduction (which means that there is no complex reductive proper subgroup of H to which E_H admits a holomorphic reduction of structure group), then E_H is unique in the following sense: For any other minimal reduction of structure group $(H', E_{H'})$ of E_G to some reductive subgroup H', there is some element $g \in G$ such that $H' = g^{-1}Hg$ and $E_{H'} = E_Hg$. As an application, we show the following:

Let M be a simply connected, irreducible smooth complex projective variety of dimension n such that the Picard number of M is one. If the canonical line bundle K_M is ample, then the algebraic holonomy of the holomorphic tangent bundle $T^{1,0}M$ is $\operatorname{GL}(n,\mathbb{C})$. If K_M^{-1} is ample, the rank of the Picard group of M is one, the biholomorphic automorphism group of M is finite, and M admits a Kähler–Einstein metric, then the algebraic holonomy of $T^{1,0}M$ is $\operatorname{GL}(n,\mathbb{C})$.

These answer some questions posed in [BK].

1. INTRODUCTION

In [BK], Balaji and Kollár introduced the notion of algebraic holonomy of polystable vector bundles on normal projective varieties defined over \mathbb{C} . Given a polystable vector bundle E over M together with a smooth point $x \in M$, the algebraic holonomy of E is a canonically associated complex reductive subgroup of $\operatorname{Aut}(E_x)$, the group of all linear automorphisms of the fiber E_x , which is constructed using the restrictions of E to the general complete intersection curves of sufficiently large degrees that pass through x. Our aim here is to address some questions in [BK], which are recalled below, on the algebraic holonomy.

Let M be a simply connected, irreducible smooth complex projective variety equipped with a Kähler–Einstein metric. In [BK, p. 187, Question 9] it is asked whether the algebraic holonomy of the holomorphic tangent bundle $T^{1,0}M$ coincides with the complexification of the differential geometric holonomy of M.

Let M be a simply connected, irreducible smooth complex projective variety whose Néron–Severi group NS(M) is of rank one. We note that rank(NS(M)) = rank(Pic(M))because M is simply connected. In [BK, p. 210, Question 51] it is asked whether the

²⁰⁰⁰ Mathematics Subject Classification. 53C07, 32Q20, 14J60.

Key words and phrases. Algebraic holonomy, stability, Einstein-Hermitian connection.

condition that the canonical line bundle K_M is ample implies that the algebraic holonomy of $T^{1,0}M$ is the full $\operatorname{GL}(n,\mathbb{C})$. We note that from a theorem of Aubin and Yau it follows that the holomorphic tangent bundle $T^{1,0}M$ of M is polystable provided K_M is ample. If the anticanonical line bundle K_M^{-1} is ample, $\operatorname{Aut}(X) < \infty$, and the holomorphic tangent bundle $T^{1,0}M$ is polystable, the Question 48 of [BK, p. 209] asks whether the algebraic holonomy of $T^{1,0}M$ is $\operatorname{GL}(n,\mathbb{C})$.

We show that both Question 9 and Question 51 have an affirmative answer. We also show that Question 48 has an affirmative answer under the extra assumption that Madmits a Kähler–Einstein metric.

Let M be a compact connected Kähler manifold equipped with a Kähler form. Let G be a connected reductive linear algebraic group defined over \mathbb{C} . It is known that any stable principal G-bundle over M admits a unique Einstein-Hermitian connection.

We prove the following (see Theorem 2.3):

Theorem 1.1. Let E_G be a stable principal G-bundle over M. Let H be a complex reductive subgroup of G which is not necessarily connected, and let $E_H \subset E_G$ be a holomorphic reduction of structure group of E_G to H. Then the Einstein–Hermitian connection on E_G is induced by a connection on E_H .

Let H be a complex reductive subgroup of G which is not necessarily connected, and let $E_H \subset E_G$ be a holomorphic reduction of structure group to H of a stable G-bundle E_G defined over M. Assume that there is no complex reductive proper subgroup of Hto which E_H admits a holomorphic reduction of structure group. Such reductions will be called the minimal reductive ones. Theorem 1.1 says that E_H is preserved by the Einstein-Hermitian connection on E_G .

Fix a point $x_0 \in M$, and also choose a point in the fiber $z \in (E_G)_{x_0}$. Taking parallel translations, for the Einstein-Hermitian connection on E_G , of z along piecewise smooth paths in M based at x_0 we get a subset of E_G . The topological closure, in E_G , of this subset gives a smooth reduction of structure group of E_G to a compact subgroup $\overline{K}_E \subset G$. The corresponding smooth reduction of structure group $E_{\overline{K}_E}^{z_0}$ of E_G to the Zariski closure $\overline{K}_E^{\mathbb{C}} \circ f \overline{K}_E$: G is not all a laborated in W and the fully integrable of E_G to E_G to the Zariski closure

 $\overline{K}_E^{\mathbb{C}}$ of \overline{K}_E in G is actually holomorphic. We prove the following (see Theorem 3.1):

Theorem 1.2. There is a point $z_0 \in (E_G)_z$ such that the above minimal reductive reduction E_H coincides with $E_{\overline{K}_E^{\mathbb{C}}}^{z_0}$. In particular, H coincides with $\overline{K}_E^{\mathbb{C}}$ for such a base point z_0 .

Take any $g \in G$. If we replace the base point z_0 by z_0g , then the subgroup $\overline{K}_E \subset G$ gets replaced by $g^{-1}\overline{K}_E g$, and hence $\overline{K}_E^{\mathbb{C}}$ gets replaced by $g^{-1}\overline{K}_E^{\mathbb{C}}g$. Also,

$$E_{\overline{K}_E^{\mathbb{C}}}^{z_0g} = E_{\overline{K}_E^{\mathbb{C}}}^{z_0}g \subset E_G$$

Therefore, if $E_{\overline{K}_{E}^{\mathbb{C}}}^{z_{0}} = E_{\overline{K}_{E}^{\mathbb{C}}}^{z_{0}g}$, then $g \in \overline{K}_{E}^{\mathbb{C}}$. In particular, if z_{0} satisfies the condition in Theorem 1.2, then $z_{0}g$ also satisfies the condition if and only if $g \in H$.

The adjoint bundle of E_G is the fiber bundle $\operatorname{Ad}(E_G) \longrightarrow M$ associated to E_G for the adjoint action of G on itself (see [KN, p. 55, Proposition 5.4] for the construction of associated bundles). Hence the fibers of $\operatorname{Ad}(E_G)$ are groups isomorphic to G.

It follows from Theorem 1.2 and the above comments that if (H, E_H) and $(H', E_{H'})$ are two minimal reductive reductions of E_G , then there is an element $g \in G$ such that $H' = g^{-1}Hg$ and $E_{H'} = E_Hg$. This gives us the following (see Corollary 3.2):

Corollary 1.3. Let E_G be a stable principal G-bundle over M. Then there is a unique holomorphic sub-fiber bundle \mathcal{G}_{E_G} of the adjoint bundle $\operatorname{Ad}(E_G)$, with fibers being subgroups, that satisfies the following condition: For any minimal reductive reduction E_H of E_G , the adjoint bundle $\operatorname{Ad}(E_H)$, which is a sub-fiber bundle of $\operatorname{Ad}(E_G)$, coincides with \mathcal{G}_{E_G} .

The sub-fiber bundle $\mathcal{G}_{E_G} \subset \operatorname{Ad}(E_G)$ in Corollary 1.3 is the complexification of the holonomy of the Einstein-Hermitian connection on E_G . While the stability condition does not depend on the choice of a Kähler form in a given Kähler class, the Einstein-Hermitian connection on a stable bundle depends on the choice of the Kähler form. We note that the sub-fiber bundle \mathcal{G}_{E_G} does not depend either on the Kähler form or on the Kähler class as long as E_G remains stable. When M is a complex projective manifold, \mathcal{G}_{E_G} coincides with the algebraic holonomy of E_G . In particular, Theorem 1.2 gives an affirmative answer to Question 9 of [BK].

In Theorem 4.1 we prove the following:

Theorem 1.4. Let M be a simply connected, irreducible smooth complex projective variety of dimension n such that the canonical line bundle K_M is ample, and $\operatorname{rank}(NS(M)) = 1$. Then the algebraic holonomy of $T^{1,0}M$ is $\operatorname{GL}(n, \mathbb{C})$.

In Theorem 4.2 we prove the following:

Theorem 1.5. Let M be an irreducible smooth complex projective variety of dimension n satisfying the following three conditions:

- the anticanonical line bundle K_M^{-1} is ample, and M admits a Kähler-Einstein metric,
- $\operatorname{rank}(\operatorname{NS}(M)) = 1$, and
- the biholomorphic automorphism group of M is finite.

Then the algebraic holonomy of $T^{1,0}M$ is $\operatorname{GL}(n,\mathbb{C})$.

Acknowledgements. The author is very grateful to János Kollár and Dominic Joyce for helpful comments. The author is very grateful to the referees for their helpful comments.

2. EINSTEIN-HERMITIAN CONNECTION AND REDUCTION OF STRUCTURE GROUP

Let M be a compact connected Kähler manifold equipped with a Kähler form ω . The *degree* of a torsion-free coherent analytic sheaf V on M is defined to be

$$\operatorname{degree}(V) := (c_1(V) \cup \omega^{\dim_{\mathbb{C}} M - 1}) \cap [M] \in \mathbb{R}.$$

Let G be a linear algebraic group defined over \mathbb{C} . The Lie algebra of G will be denoted by \mathfrak{g} . For any $g_0 \in G$, let $\operatorname{Ad}(g_0) : G \longrightarrow G$ be the inner automorphism defined by $g \longmapsto g_0^{-1}gg_0$. The corresponding automorphism of \mathfrak{g} will also be denoted by $\operatorname{Ad}(g_0)$.

Let $E_G \longrightarrow M$ be a holomorphic principal G-bundle. The adjoint vector bundle of E_G will be denoted by $\operatorname{ad}(E_G)$. We recall that $\operatorname{ad}(E_G)$ is a quotient of $E_G \times \mathfrak{g}$, and two points (z, v) and (z', v') of $E_G \times \mathfrak{g}$ are identified in $\operatorname{ad}(E_G)$ if and only if there is some $g_0 \in G$ such that $z' = zg_0$ and $v' = \operatorname{Ad}(g_0)(v)$. Therefore, the fibers of $\operatorname{ad}(E_G)$ are Lie algebras isomorphic to \mathfrak{g} . If E is a vector bundle of rank r, and $E_{\operatorname{GL}(r,\mathbb{C})}$ is the corresponding principal $\operatorname{GL}(r,\mathbb{C})$ -bundle, then $\operatorname{ad}(E_{\operatorname{GL}(r,\mathbb{C})}) = \mathcal{E}nd(E) = E \otimes E^*$.

Let $\operatorname{Ad}(E_G)$ be the quotient of $E_G \times G$ where two points (z, g) and (z', g') are identified if and only if there is some $g_0 \in G$ such that $z' = zg_0$ and $g' = \operatorname{Ad}(g_0)(g)$. Therefore, the adjoint bundle $\operatorname{Ad}(E_G)$ is a fiber bundle over M, and the fibers are groups isomorphic to G. For any point $x \in M$, the fiber $\operatorname{Ad}(E_G)_x$ is the group of all automorphisms of the fiber $(E_G)_x$ that commute with the action of G on $(E_G)_x$. For the above principal $\operatorname{GL}(r, \mathbb{C})$ -bundle $E_{\operatorname{GL}(r,\mathbb{C})}$ associated to E, the fiber of $\operatorname{Ad}(E_{\operatorname{GL}(r,\mathbb{C})})$ over any point $x \in M$ is the group of automorphisms of the fiber E_x .

Henceforth, G will be a connected reductive group.

Since the degree has been defined, we have the notions of a stable G-bundle and a polystable G-bundle over M. See [RS], [AB], [Ra2] for definitions of stable and polystable principal G-bundles.

Fix a maximal torus $T \subset G$ and a Borel subgroup $B \subset G$ containing T. We also fix a maximal compact subgroup K of G. By a parabolic subgroup of G we will mean one containing B. Therefore, the Levi quotient L(Q) of any parabolic subgroup $Q \subset G$ is also a subgroup of Q. Let $Z_0(G)$ denote the connected component of the center of G that contains the identity element.

Take any polystable principal G-bundle E_G over M. From the definition of a polystable principal G-bundle (see [AB, p. 221, Definition 3.5], [RS, p. 23]) it follows that there is a Levi subgroup L(P) associated to some parabolic subgroup $P \subset G$ (the subgroup Pneed not be proper) along with a holomorphic reduction of structure group $E_{L(P)} \subset E_G$ to L(P), such that

- the principal L(P)-bundle $E_{L(P)}$ is stable, and
- the principal *P*-bundle $E_P := E_{L(P)}(P)$, obtained by extending the structure group of $E_{L(P)}$ using the inclusion L(P) in *P*, is an admissible reduction of E_G (the definition of an admissible reduction is recalled below).

5

Note that since $L(P) \subset P \subset G$, the *P*-bundle E_P is a reduction of structure group of E_G to *P*. The condition that E_P is an admissible reduction of structure group of E_G means that for each character χ of *P*, which is trivial on $Z_0(G)$, the associated line bundle $E_P(\chi)$ over *M* is of degree zero. The *G*-bundle E_G is stable if and only if P = G.

Since E_G and $E_{L(P)}$ are polystable, they admit unique Einstein–Hermitian connections [AB, p. 208, Theorem 0.1]. It should be clarified that the Einstein–Hermitian reduction of structure to a maximal compact subgroup depends on the choice of the maximal compact subgroup. Even for a fixed maximal compact subgroup, the Einstein–Hermitian reduction of structure group need not be unique. However, once the Kähler form on M is fixed, the Einstein–Hermitian connection on a polystable principal bundle over M is unique. Let $\nabla^{L(P)}$ be the Einstein–Hermitian connection on the stable L(P)–bundle $E_{L(P)}$. Let ∇^G be the connection on E_G induced by $\nabla^{L(P)}$. Note that a connection on L(P) induces a connection on any fiber bundle associated to $E_{L(P)}$. The principal G-bundle E_G is associated to $E_{L(P)}$ for the left translation action of L(P) on G, and ∇^G is the connection on it obtained from $\nabla^{L(P)}$.

Proposition 2.1. The induced connection ∇^G on E_G coincides with the unique Einstein-Hermitian connection on E_G .

Proof. Since the inclusion map $L(P) \hookrightarrow G$ need not take the connected component, containing the identity element, of the center of L(P) into $Z_0(G)$, the proposition does not follow immediately from [AB, p. 208, Theorem 0.1]. The connection $\nabla^{L(P)}$ on $E_{L(P)}$ is induced by a connection on a smooth reduction of structure group of $E_{L(P)}$ to a maximal compact subgroup of L(P) (this is a part of the definition of an Einstein–Hermitian connection). A maximal compact subgroup of L(P) is contained in a maximal compact subgroup of G. Consequently, the connection ∇^G on E_G is induced by a connection on a smooth reduction of structure group of E_G to a maximal compact subgroup of G. Therefore, to prove that ∇^G is the Einstein–Hermitian connection on E_G it suffices to show that ∇^G satisfies the Einstein–Hermitian equation.

Let $\mathfrak{z}(\mathfrak{l}(P))$ be the Lie algebra of the center of L(P). For any $\theta \in \mathfrak{z}(\mathfrak{l}(P))$, the holomorphic section of the adjoint bundle $\mathrm{ad}(E_{L(P)})$ given by θ will be denoted by $\hat{\theta}$; the vector bundle $\mathrm{ad}(E_{L(P)})$ is associated to $E_{L(P)}$ for the adjoint action of L(P) on its Lie algebra. Let Λ_{ω} be the adjoint of multiplication by ω of differential form on M. That the connection $\nabla^{L(P)}$ is Einstein–Hermitian means that there is an element $\theta \in \mathfrak{z}(\mathfrak{l}(P))$ such that the Einstein–Hermitian equation

(2.1)
$$\Lambda_{\omega} \mathcal{K}_{\nabla^{L(P)}} = \widehat{\theta}$$

holds.

If θ in (2.1) is in the Lie algebra $\mathfrak{z}(G)$ of $Z_0(G)$, then ∇^G is a Einstein-Hermitian connection on E_G . Therefore, in that case the proposition is proved. Assume that

(2.2)
$$\theta \notin \mathfrak{z}(G)$$
.

Fix a character χ of P which is trivial on $Z_0(G)$ but satisfies the following condition: the homomorphism of Lie algebras

$$(2.3) d\chi: \mathfrak{p} \longrightarrow \mathbb{C}$$

given by χ , where **p** is the Lie algebra of *P*, is nonzero on θ . (It is easy to check that the group of characters of *P* coincides with the group of characters of L(P).)

Consider the holomorphic line bundle $L_{\chi} := E_P(\chi)$ over M associated to E_P for χ . Let ∇^{χ} be the connection on L_{χ} induced by the connection on E_P given by $\nabla^{L(P)}$. Since $\nabla^{L(P)}$ is an Einstein–Hermitian connection, the connection ∇^{χ} is also Einstein–Hermitian. Indeed, if $\mathcal{K}(\nabla^{\chi})$ is the curvature of ∇^{χ} , then

$$\Lambda_{\omega} \mathcal{K}(\nabla^{\chi}) = \mathrm{d}\chi(\theta) \,,$$

where $d\chi$ is the homomorphism in (2.3) and θ is the element in (2.1). Therefore,

(2.4)
$$\operatorname{degree}(L_{\chi}) = \frac{\mathrm{d}\chi(\theta)\sqrt{-1}}{2\pi d} \int_{M} \omega^{d},$$

where $d = \dim_{\mathbb{C}} M$; see [Ko2, p. 103, Proposition 2.1].

The condition that $E_P \subset E_G$ is an admissible reduction of structure group says that

$$\operatorname{degree}(L_{\chi}) = 0.$$

This, in view of the assumption in (2.2) that $d\chi(\theta) \neq 0$, contradicts (2.4). Therefore, we conclude that $\theta \in \mathfrak{z}(G)$. This immediately implies that the connection ∇^G on E_G is Einstein-Hermitian. This completes the proof of the proposition.

Let E_G be a holomorphic principal G-bundle over M and ∇' a C^{∞} connection on E_G compatible with the holomorphic structure of E_G . Such a connection is called a complex connection; see [AB, p. 230, Definition 3.1(1)] for the precise definition of a complex connection. A connection $\widetilde{\nabla}$ is complex if and only if the (0, 2)-Hodge type component of the curvature of $\widetilde{\nabla}$ vanishes identically.

Definition 2.2. Let H be a closed complex subgroup of G and $E_H \subset E_G$ a C^{∞} reduction of structure group of E_G to H. We will say that E_H is *preserved* by the connection ∇' if ∇' induces a connection on E_H .

It follows immediately that E_H is preserved by ∇' if and only if there is a smooth connection ∇'' on E_H such that the connection ∇' on E_G coincides with the one given by ∇'' using extension of structure group. It is easy to see that E_H is preserved by ∇' if and only if for each point $z \in E_H$, the horizontal subspace in $T_z E_G$ for the connection ∇' is contained in $T_z E_H$. If E_H is preserved by ∇' , then E_H is a holomorphic reduction of structure group of E_G .

Theorem 2.3. Let E_G be a stable principal G-bundle over M and ∇^G the Einstein-Hermitian connection on E_G . Let H be a complex reductive subgroup of G which is not necessarily connected, and let $E_H \subset E_G$ be a holomorphic reduction of structure group of E_G to H. Then E_H is preserved by the connection ∇^G . *Proof.* Let $H_0 \subset H$ be the connected component containing the identity element. Set

$$X := E_H/H_0,$$

which is finite étale Galois cover of M with Galois group H/H_0 . Let

$$(2.5) p: X \longrightarrow M$$

be the projection. We note that p^*E_H has a canonical reduction of structure group to the subgroup $H_0 \subset H$. Set

$$F_G := p^* E_G$$

Let

$$F_{H_0} \subset F_G$$

be the reduction of structure group to H_0 obtained from the canonical reduction of structure group of p^*E_H to H_0 .

Equip X with the Kähler form $p^*\omega$. The Einstein-Hermitian connection on E_G pulls back to an Einstein-Hermitian connection on F_G for the Kähler form $p^*\omega$. Therefore, F_G is polystable with respect to $p^*\omega$. Let $\operatorname{ad}(F_G)$ be the adjoint bundle over X. We recall that $\operatorname{ad}(F_G)$ is associated to F_G for the adjoint action of G on its Lie algebra. The connection on the vector bundle $\operatorname{ad}(F_G)$ induced by the Einstein-Hermitian connection of F_G is clearly Einstein-Hermitian. Hence the adjoint vector bundle $\operatorname{ad}(F_G)$ is polystable.

Let Z(G) be the center of G. Set

$$H' := H_0 Z(G) / Z(G) .$$

Therefore, H' is a complex reductive subgroup of the complex semisimple group G' := G/Z(G). Let \mathfrak{h}_0 be the Lie algebra of H_0 . As before, \mathfrak{g} is the Lie algebra of G. Note that the adjoint action makes \mathfrak{g} (respectively, \mathfrak{h}_0) a G'-module (respectively, H'-module). We will also consider \mathfrak{g} as a H'-module using the inclusion of H' in G'.

Since \mathfrak{g} is a faithful G'-module, and H' is a reductive subgroup of G', there is a positive integer N and nonnegative integers $a_i, b_i, i \in [1, N]$, such that the H'-module \mathfrak{h}_0 is a direct summand of the H'-module

(2.6)
$$\bigoplus_{i=1}^{N} \mathfrak{g}^{\otimes a_i} \otimes (\mathfrak{g}^*)^{\otimes b_i} = \bigoplus_{i=1}^{N} \mathfrak{g}^{\otimes a_i} \otimes \mathfrak{g}^{\otimes b_i}$$

[De, p. 40, Proposition 3.1]; since H' is complex reductive, any exact sequence of H'-modules splits; also, $\mathfrak{g} = \mathfrak{g}^*$ as G is reductive. Therefore, the adjoint vector bundle $\mathrm{ad}(F_{H_0})$ is a direct summand of the vector bundle

(2.7)
$$\bigoplus_{i=1}^{N} \operatorname{ad}(F_G)^{\otimes a_i} \otimes \operatorname{ad}(F_G)^{\otimes b_i}$$

We note that the vector bundle in (2.7) is associated to F_{H_0} for the H_0 -module in (2.6).

Since the adjoint vector bundle $ad(F_G)$ is polystable of degree zero (recall that $ad(F_G) = ad(F_G)^*$), the vector bundle

$$\operatorname{ad}(F_G)^{\otimes a} \otimes \operatorname{ad}(F_G)^{\otimes b}$$

is polystable of degree zero for all $a, b \in \mathbb{N}$ [AB, p. 224, Theorem 3.9]. Therefore, the vector bundle in (2.7) is polystable of degree zero. Since $\operatorname{ad}(F_{H_0})$ is a direct summand of it of degree zero, we conclude that $\operatorname{ad}(F_{H_0})$ is also polystable. (See the proof of [AAB, Lemma 5] for a similar argument.)

The principal H_0 -bundle F_{H_0} over X is polystable because the vector bundle $\operatorname{ad}(F_{H_0})$ is polystable [AB, p. 224, Corollary 3.8]. Hence F_{H_0} admits a unique Einstein-Hermitian connection [AB, p. 208, Theorem 0.1]. Let ∇^{H_0} be the Einstein-Hermitian connection on F_{H_0} . So, there is an element $\nu \in \mathfrak{h}$ such that

(2.8)
$$\Lambda_{p^*\omega} \mathcal{K}(\nabla^{H_0}) = \widehat{\nu},$$

where $\mathcal{K}(\nabla^{H_0})$ is the curvature of ∇^{H_0} , and $\hat{\nu}$ is the holomorphic section of $\operatorname{ad}(p^*E_H) = \operatorname{ad}(F_{H_0})$ given by ν .

Since H/H_0 is a finite group, giving a connection on a principal H_0 -bundle is equivalent to giving a connection on the principal H-bundle obtained from it by extension of structure group. From the uniqueness of the Einstein-Hermitian connection on F_{H_0} it follows that the corresponding connection on p^*E_H is left invariant by the action of the Galois group H/H_0 . Consequently, the connection on p^*E_H given by ∇^{H_0} descends to a connection on E_H . Let ∇^H denote the connection on E_H obtained this way. It is clear that ∇^H is a complex connection.

Let ∇ be the complex connection on E_G induced by the above connection ∇^H on E_H . We will first show that ∇ is unitary, which means that ∇ is induced by connection on a smooth reduction of structure group of E_G to a maximal compact subgroup of G. Then we will show that ∇ satisfies the Einstein–Hermitian equation.

To prove that ∇ is unitary, fix a point $x_0 \in M$, and also fix a point $z_0 \in (E_G)_{x_0}$ in the fiber of E_G over x_0 . Taking parallel translations of z_0 , with respect to the connection ∇ , along piecewise smooth paths in M based at x_0 we get a subset S of E_G . Sending any $g \in G$ to the point $z_0g \in (E_G)_{x_0}$ we get an isomorphism $G \longrightarrow (E_G)_{x_0}$. Using this isomorphism, the intersection $(E_G)_{x_0} \cap S$ is a subgroup K_E of G. The condition that the connection ∇ is unitary is equivalent to the condition that K_E is contained in some compact subgroup of G.

Fix a point $x \in p^{-1}(x_0)$, and also fix a point $z \in (p^*E_G)_x$, where p is the covering map in (2.5). Consider parallel translations of z, with respect to the connection $p^*\nabla$ on $p^*E_G =: F_G$, along piecewise smooth paths in X based at x. As before, we get a subgroup $K_F \subset G$ from the resulting subset of F_G . It is easy to see that K_F is a finite index subgroup of the group K_E constructed above.

We note that the connection $p^*\nabla$ on F_G is induced by the unitary connection ∇^{H_0} on the reduction F_{H_0} of F_G . Therefore, the connection $p^*\nabla$ is unitary. Consequently, the subgroup $K_F \subset G$ is contained in a compact subgroup of G. Using this together with the observation that K_F is a finite index subgroup of the subgroup $K_E \subset G$ we conclude that K_E is also contained in a compact subgroup of G. Thus the connection ∇ is unitary.

We will now show that ∇ satisfies the Einstein–Hermitian equation.

Let $\mathcal{K}(\nabla)$ be the curvature of the connection ∇ . From (2.8) it follows immediately that $\Lambda_{p^*\omega}\mathcal{K}(\nabla^{H_0})$ is a holomorphic section of $\operatorname{ad}(p^*E_H)$. Consequently, $\Lambda_{\omega}\mathcal{K}(\nabla)$ is a holomorphic section of the adjoint vector bundle $\operatorname{ad}(E_G)$.

Since E_G is stable, it can be shown that all holomorphic sections of $\operatorname{ad}(E_G)$ are given by the Lie algebra $\mathfrak{z}(G)$ of $Z_0(G)$. To prove this consider the Einstein-Hermitian connection on $\operatorname{ad}(E_G)$ induced by the Einstein-Hermitian connection on E_G . We have $\operatorname{ad}(E_G)^* =$ $\operatorname{ad}(E_G)$ because G is reductive. Hence the constant in the Einstein-Hermitian equation for $\operatorname{ad}(E_G)$ vanishes (see [Ko2, p. 99, Proposition (1.4)(2)]). Therefore, the mean curvature for the Einstein-Hermitian connection on $\operatorname{ad}(E_G)$ vanishes (see [Ko2, p. 51, (1.7)] for the definition of mean curvature, and [Ko2, p. 99] for the expression of the Einstein-Hermitian equation in terms of mean curvature). Since the mean curvature of the Einstein-Hermitian connection on $\operatorname{ad}(E_G)$ vanishes, from [Ko2, p. 52, Theorem (1.9)] it follows that any holomorphic section of $\operatorname{ad}(E_G)$ is flat with respect to the Einstein-Hermitian connection on it. Now the fact that E_G is stable implies that any flat section of $\operatorname{ad}(E_G)$ for the Einstein-Hermitian connection on it is given by some element of $\mathfrak{z}(G)$; see the proof of [Ra1, p. 136, Proposition 3.2].

In other words, there is an element $\theta \in \mathfrak{z}(G)$ such that $\Lambda_{\omega}\mathcal{K}(\nabla)$ coincides with the section of $\mathrm{ad}(E_G)$ given by θ . Thus, we conclude that the connection ∇ is the unique Einstein-Hermitian connection on E_G . Since ∇ is induced by a connection on E_H , the proof of the theorem is complete. \Box

Proposition 2.1 and Theorem 2.3 together have the following:

Corollary 2.4. Let E_G be a polystable *G*-bundle over *M*. Take $E_{L(P)}$ as in Proposition 2.1. Let *H* be a complex reductive subgroup of L(P) (not necessarily connected), and let $E_H \subset E_{L(P)}$ be a holomorphic reduction of structure group of $E_{L(P)}$ to *H*. Then E_H is preserved by the Einstein–Hermitian connection on E_G .

3. PROPERTIES OF A MINIMAL REDUCTION

Let E_G be a stable principal G-bundle over M. Fix a point $x_0 \in M$, and also fix a point $z_0 \in (E_G)_{x_0}$ in the fiber over x_0 . Let ∇^G be the Einstein-Hermitian connection on E_G .

Taking parallel translations of z_0 , with respect to ∇^G , along piecewise smooth paths in M based at x_0 we get a subset S of E_G . Let \overline{S} be the topological closure of S in E_G . The map $G \longrightarrow (E_G)_{x_0}$ defined by $g \longmapsto z_0 g$ is an isomorphism. Using this isomorphism, the intersection $(E_G)_{x_0} \cap \overline{S}$ gives a compact subgroup $\overline{K}_E \subset G$. The subset $\overline{S} \subset E_G$ is a C^{∞} reduction of structure group of E_G to the subgroup \overline{K}_E . Let $\overline{K}_E^{\mathbb{C}}$ be the complex reductive subgroup of G obtained by taking the Zariski closure of \overline{K}_E in G. The subset

(3.1)
$$E_{\overline{K}_{E}^{\mathbb{C}}}^{z_{0}} := \mathcal{S}\overline{K}_{E}^{\mathbb{C}} \subset E_{G}$$

is a holomorphic reduction of structure group of E_G to $\overline{K}_E^{\mathbb{C}}$; see [Bi, Section 3] for the details. It is easy to see that the principal $\overline{K}_E^{\mathbb{C}}$ -bundle $E_{\overline{K}_E^{\mathbb{C}}}^{z_0}$ is the extension of structure group of the principal \overline{K}_E -bundle \overline{S} .

The subgroup $\overline{K}_E \subset G$ is called the (differential geometric) holonomy of the connection ∇^G with reference point z_0 (see [KN, p. 72]). The subset $\overline{\mathcal{S}}$ is called the (differential geometric) holonomy bundle through z_0 (see [KN, p. 85]).

We note that in [Bi], the point z_0 is taken to be in the subset of E_G given by an Einstein–Hermitian reduction of structure group (see [Bi, p. 71, (3.20)]). This was done only to make \overline{K}_E lie inside a fixed maximal compact subgroup of G. If we replace the base point z_0 by z_0g , where g is any point of G, then it is easy to see that the subset \mathcal{S} constructed above using z_0 gets replaced by $\mathcal{S}g$. Therefore, the subgroup $\overline{K}_E^{\mathbb{C}}$ in (3.1) gets replaced by $g^{-1}\overline{K}_E^{\mathbb{C}}g$, and the reduction $E_{\overline{K}_E^{\mathbb{C}}}^{z_0} \subset E_G$ in (3.1) gets replaced by $E_{\overline{K}_E^{\mathbb{C}}}^{z_0}g$.

We note that $(\overline{K}_E^{\mathbb{C}}, E_{\overline{K}_E^{\mathbb{C}}}^{z_0})$ in (3.1) is a minimal complex reduction of E_G in the following sense: There is no complex proper subgroup of $\overline{K}_E^{\mathbb{C}}$ to which $E_{\overline{K}_E^{\mathbb{C}}}^{z_0}$ admits a holomorphic reduction of structure group which is preserved by the connection ∇^G . Indeed, this follows immediately from the construction of $E_{\overline{K}_E^{\mathbb{C}}}^{z_0}$. Furthermore, if $E_{H'} \subset E_G$ is a holomorphic reduction of structure group of E_G , to a complex subgroup $H' \subset G$, satisfying the two conditions:

- $E_{H'}$ is preserved by ∇^G , and
- there is no complex proper subgroup of H' to which $E_{H'}$ admits a holomorphic reduction of structure group which is also preserved by ∇^G ,

then it follows immediately that there is a point $z_0 \in (E_G)_{x_0}$ such that $E_{H'} = E_{\overline{K_E^{\mathbb{C}}}}^{z_0}$. Indeed, z_0 can be taken to be any point of the fiber $(E_{H'})_{x_0}$.

Let H be a complex reductive subgroup of G which is not necessarily connected, and let $E_H \subset E_G$ be a holomorphic reduction of structure group to H of the stable G-bundle E_G . This reduction E_H will be called a *minimal reductive reduction* of E_G if there is no complex reductive proper subgroup of H to which E_H admits a holomorphic reduction of structure group.

In view of the above observations, using Theorem 2.3 we get the following:

Theorem 3.1. Let E_G be a stable principal G-bundle over a compact connected Kähler manifold M, where G is a connected reductive complex linear algebraic group. Let $E_H \subset$ E_G be a minimal reductive reduction of E_G . Fix a point $x_0 \in M$. Then there is a point z_0 in the fiber $(E_G)_{x_0}$ such that $E_{K_E}^{z_0}$ (defined in (3.1)) coincides with E_H . In particular, the subgroup H coincides with $\overline{K}_E^{\mathbb{C}}$ for such a base point z_0 .

Let $\operatorname{Ad}(E_G)$ be the adjoint bundle for E_G . So $\operatorname{Ad}(E_G)$ is the fiber bundle over M associated to E_G for the adjoint action of G on itself. The fibers of $\operatorname{Ad}(E_G)$ are groups

11

isomorphic to G. Let $\operatorname{Ad}(E_{\overline{K}_{E}^{C}}^{z_{0}})$ be the adjoint bundle of the principal $\overline{K}_{E}^{\mathbb{C}}$ -bundle $E_{\overline{K}_{E}^{C}}^{z_{0}}$ defined in (3.1). Since $E_{\overline{K}_{E}^{\mathbb{C}}}^{z_{0}}$ is a holomorphic reduction of structure group of E_{G} , the fiber bundle $\operatorname{Ad}(E_{\overline{K}_{E}^{z_{0}}})$ is a holomorphic sub–fiber bundle of $\operatorname{Ad}(E_{G})$ with the fibers of $\operatorname{Ad}(E_{\overline{K}_{E}^{z_{0}}})$ being subgroups of the fibers of $\operatorname{Ad}(E_{G})$.

We noted earlier that if z_0 is replaced by z_0g , where $g \in G$, then the subgroup $\overline{K}_E^{\mathbb{C}}$ gets replaced by $g^{-1}\overline{K}_E^{\mathbb{C}}g$, and $E_{\overline{K}_E^{\mathbb{C}}}^{z_0}$ gets replaced by $E_{\overline{K}_E^{\mathbb{C}}}^{z_0}g$. From this it follows immediately that the subbundle

$$\operatorname{Ad}(E^{z_0}_{\overline{K}^{\mathbb{C}}_{r}}) \subset \operatorname{Ad}(E_G)$$

is independent of the choice of the base point z_0 .

Therefore, from Theorem 3.1 we have the following:

Corollary 3.2. Let E_G be a stable principal G-bundle over a compact connected Kähler manifold M, where G is a connected reductive complex linear algebraic group. Then there is a unique holomorphic sub-fiber bundle \mathcal{G}_{E_G} of the adjoint bundle $\operatorname{Ad}(E_G)$, with fibers being subgroups, that satisfies the following condition: For any minimal reductive reduction E_H of E_G , the adjoint bundle $\operatorname{Ad}(E_H)$, which is a sub-fiber bundle of $\operatorname{Ad}(E_G)$, coincides with \mathcal{G}_{E_G} .

From Theorem 3.1 it follows that the sub-fiber bundle $\mathcal{G}_{E_G} \subset \operatorname{Ad}(E_G)$ in Corollary 3.2 is the complexification of the holonomy of the Einstein-Hermitian connection on E_G . We note that for defining stable bundles we need only the Kähler class $[\omega] \in H^2(M, \mathbb{R})$. In other words, the stability condition does not depend on the choice of the Kähler form in a given Kähler class. On the other hand, the Einstein-Hermitian connection on a stable bundle depends on the choice of the Kähler form. From Corollary 3.2 it follows that the sub-fiber bundle \mathcal{G}_{E_G} does not depend on the choice of the Kähler form in a given Kähler class. In fact, it does not even depend on the Kähler class as long as E_G remains stable.

Assume that M is a complex projective manifold. The algebraic holonomy $H_{x_0}(E_G)$ of E_G constructed in [BK] is an algebraic subgroup of $\operatorname{Ad}(E_G)_{x_0}$; recall that $\operatorname{Ad}(E_G)_{x_0}$ is the group of automorphisms of $(E_G)_{x_0}$ that commute with the action of G. Hence $H_{x_0}(E_G)$ gives a conjugacy class of algebraic subgroups of G. Take any subgroup $H_{x_0} \subset G$ in this conjugacy class. The principal G-bundle admits a minimal reductive reduction of structure group to this subgroup H_{x_0} [BK, p. 193, Theorem 20(3)]. Hence from Corollary 3.2 it follows that the algebraic holonomy of E_G coincides with the fiber $(\mathcal{G}_{E_G})_{x_0}$ when M is a complex projective manifold.

4. Algebraic holonomy and Kähler–Einstein metric

Let (M, g) be a simply connected compact Kähler manifold. The de Rham decomposition says that there is a biholomorphic isometry

(4.1)
$$(M,g) \xrightarrow{\sim} \prod_{i=1}^{\ell} (M_i,g_i),$$

where each (M_i, g_i) is a compact Kähler manifold whose holonomy group is irreducible (see [Ko1, § 4, pp. 327–328], [Jo, p. 49, Theorem 3.2.7]). Furthermore, for $i \in [1, \ell]$, the Kähler manifold (M_i, g_i) is either an irreducible Hermitian symmetric space or its holonomy is one of the following three:

(4.2) $U(m_i)$, $SU(m_i)$, and $Sp(m_i/2)$,

where $m_i = \dim_{\mathbb{C}} M_i$ [Jo, p. 58, § 3.4.2] (this also follows from the combination of [Ko1, p. 327, Corollary 4] and [Be, p. 294, § 10.66]).

Theorem 4.1. Let M be a simply connected, irreducible smooth complex projective variety, of complex dimension n, such that the canonical line bundle K_M is ample, and $\operatorname{rank}(\operatorname{NS}(M)) = 1$. Then the algebraic holonomy of the holomorphic tangent bundle $T^{1,0}M$ is $\operatorname{GL}(n, \mathbb{C})$.

Proof. Fix a Kähler–Einstein metric g on M, which exists by [Au], [Ya]. We know that the algebraic holonomy of M is a complexification of the holonomy of the Kähler–Einstein metric g (see Theorem 3.1). Therefore, it suffices to show that the holonomy of g is full U(n).

Consider the decomposition in (4.1). Since $\operatorname{rank}(\operatorname{NS}(M)) = 1$, we have $\ell = 1$. A simply connected compact Hermitian symmetric space is a Fano manifold. Hence the holonomy of g is one of the three groups in (4.2). Since the canonical line bundle K_M is ample, only $\operatorname{U}(n)$ in (4.2) can be the holonomy. This completes the proof of the theorem.

Theorem 4.2. Let M be an irreducible smooth complex projective variety of complex dimension n satisfying the following four conditions:

- the anticanonical line bundle K_M^{-1} is ample,
- $\operatorname{rank}(\operatorname{NS}(M)) = 1$,
- the biholomorphic automorphism group of M is finite, and
- M admits a Kähler–Einstein metric.

Then the algebraic holonomy of the holomorphic tangent bundle $T^{1,0}M$ is $GL(n, \mathbb{C})$.

Proof. Fix a Kähler–Einstein metric g on M. As in the proof of Theorem 4.1, we need to show that the holonomy of g is U(n).

Consider the decomposition in (4.1). As before, $\ell = 1$ because rank(NS(M)) = 1. Also, M is not a Hermitian symmetric space because the biholomorphic automorphism group of M is finite. Since K_M^{-1} is ample, except U(n) none of the other groups in (4.2) can be the holonomy of g. This completes the proof of the theorem.

References

[Au] T. Aubin: Équations du type Monge–Ampère sur les variétés kähleriennes compactes, Comp. Ren. Acad. Sci. Paris Sér. A 283 (1976), 119–121.

- [AB] B. Anchouche and I. Biswas: Einstein–Hermitian connections on polystable principal bundles over a compact Kähler manifold, *Amer. Jour. Math.* **123** (2001), 207–228.
- [AAB] B. Anchouche H. Azad and I. Biswas: Harder–Narasimhan reduction for principal bundles over a compact Kähler manifold, *Math. Ann.* **323** (2002), 693–712.
- [BK] V. Balaji and J. Kollár: Holonomy groups of stable vector bundles, Publ. Res. Inst. Math. Sci. 44 (2008), 183–211.
- [Be] A. Besse: *Einstein Manifolds*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3), Band 10, Springer-Verlag, Berlin, 1987.
- [Bi] I. Biswas: Stable bundles and extension of structure group, *Diff. Geom. Appl.* 23 (2005), 67–78.
- [De] P. Deligne: Hodge cycles on abelian varieties, (in: Hodge cycles, motives, and Shimura varieties, by P. Deligne, J. S. Milne, A. Ogus and K.-Y. Shih), Lecture Notes in Mathematics, 900, Springer-Verlag, Berlin-New York, 1982.
- [Jo] D. D. Joyce: *Compact Manifolds with Special Holonomy*, Oxford Mathematical Monographs. Oxford University Press, Oxford, 2000.
- [Ko1] S. Kobayashi: The first Chern class and holomorphic symmetric tensor fields, Jour. Math. Soc. Japan 32 (1980), 325–329.
- [Ko2] S. Kobayashi: Differential Geometry of Complex Vector Bundles, Publications of the Math. Society of Japan 15, Iwanami Shoten Publishers and Princeton University Press, 1987.
- [KN] S. Kobayashi and K. Nomizu: Foundations of differential geometry. Vol. I, Interscience Publishers, New York-London, 1963.
- [RS] A. Ramanathan and S. Subramanian: Einstein-Hermitian connections on principal bundles and stability, Jour. Reine Angew. Math. 390 (1988), 21–31.
- [Ra1] A. Ramanathan: Stable principal bundles on a compact Riemann surface, Math. Ann. 213 (1975), 129–152.
- [Ra2] A. Ramanathan: Moduli for principal bundles over algebraic curves: I, Proc. Ind. Acad. Sci. (Math. Sci.) 106 (1996), 301–328.
- [Ya] S.-T. Yau: On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation. I, *Comm. Pure Appl. Math.* **31** (1978), 339–411.

School of Mathematics, Tata Institute of Fundamental Research, Homi Bhabha Road, Bombay 400005, India

E-mail address: indranil@math.tifr.res.in