

Finite Fields Appl. 14(2008), no. 2, 470–481.

## On Value Sets of Polynomials over a Field

Zhi-Wei Sun

Department of Mathematics, Nanjing University  
Nanjing 210093, People's Republic of China

zwsun@nju.edu.cn

<http://math.nju.edu.cn/~zwsun>

### Abstract

Let  $F$  be any field. Let  $p(F)$  be the characteristic of  $F$  if  $F$  is not of characteristic zero, and let  $p(F) = +\infty$  otherwise. Let  $A_1, \dots, A_n$  be finite nonempty subsets of  $F$ , and let

$$f(x_1, \dots, x_n) = a_1x_1^k + \dots + a_nx_n^k + g(x_1, \dots, x_n) \in F[x_1, \dots, x_n]$$

with  $k \in \{1, 2, 3, \dots\}$ ,  $a_1, \dots, a_n \in F \setminus \{0\}$  and  $\deg g < k$ . We show that

$$|\{f(x_1, \dots, x_n) : x_1 \in A_1, \dots, x_n \in A_n\}| \geq \min \left\{ p(F), \sum_{i=1}^n \left\lfloor \frac{|A_i| - 1}{k} \right\rfloor + 1 \right\}.$$

When  $k \geq n$  and  $|A_i| \geq i$  for  $i = 1, \dots, n$ , we also have

$$\begin{aligned} & |\{f(x_1, \dots, x_n) : x_1 \in A_1, \dots, x_n \in A_n, \text{ and } x_i \neq x_j \text{ if } i \neq j\}| \\ & \geq \min \left\{ p(F), \sum_{i=1}^n \left\lfloor \frac{|A_i| - i}{k} \right\rfloor + 1 \right\}; \end{aligned}$$

consequently, if  $n \geq k$  then for any finite subset  $A$  of  $F$  we have

$$|\{f(x_1, \dots, x_n) : x_1, \dots, x_n \in A, \text{ and } x_i \neq x_j \text{ if } i \neq j\}| \geq \min\{p(F), |A| - n + 1\}.$$

In the case  $n > k$ , we propose a further conjecture which extends the Erdős-Heilbronn conjecture in a new direction.

## 1 Introduction

For a field  $F$  we use  $p(F)$  to denote the additive order of the multiplicative identity of  $F$ , which is either infinite or a prime. A field  $F$  is said to have characteristic zero if  $p(F) = +\infty$ , and have characteristic  $p$  if  $p(F)$  is a prime  $p$ .

By the Chevalley-Warning theorem (cf. pp. 50–51 of [10]), for any polynomial  $P(x_1, \dots, x_n)$  over a finite field  $F$ , if  $n > \deg P$  then the characteristic of  $F$  divides the number of solutions

---

2000 *Mathematics Subject Classification*. Primary 11T06; Secondary 05A05, 11B75, 11P99, 12E10.

Supported by the National Science Fund (Grant No. 10425103) for Distinguished Young Scholars in China.

to the equation  $P(x_1, \dots, x_n) = 0$  over  $F^n$ . However, this says nothing about the solvability of the equation over  $F^n$  unless there is an obvious solution.

Given a field  $F$ , we consider polynomials of the form

$$f(x_1, \dots, x_n) = a_1x_1^k + \dots + a_nx_n^k + g(x_1, \dots, x_n) \in F(x_1, \dots, x_n), \quad (1.1)$$

where

$$k \in \mathbb{Z}^+ = \{1, 2, 3, \dots\}, \quad a_1, \dots, a_n \in F^* = F \setminus \{0\} \text{ and } \deg g < k. \quad (1.2)$$

What can we say about the solvability of the equation  $f(x_1, \dots, x_n) = 0$  over  $F^n$ ?

Let  $F$  be the field  $F_p = \mathbb{Z}/p\mathbb{Z}$  with  $p$  a prime, and assume (1.1) and (1.2). In 1956 Carlitz [3] proved that the equation  $f(x_1, \dots, x_n) = 0$  has a solution in  $F_p^n$  when  $k \mid p-1$  and  $n \geq k$ . In 2006 Felszeghy [7] extended this result by showing that the equation is solvable if  $k < p$  and  $n \geq \lceil (p-1)/\lfloor (p-1)/k \rfloor \rceil$ . (For a real number  $\alpha$ ,  $\lceil \alpha \rceil$  denotes the least integer not smaller than  $\alpha$  while  $\lfloor \alpha \rfloor$  represents the largest integer not exceeding  $\alpha$ .) Note that the equation  $f(x_1, \dots, x_n) = 0$  over  $F_p^n$  is solvable if and only if the value set

$$\{f(x_1, \dots, x_n) : x_1, \dots, x_n \in F_p\}$$

contains 0. In 1959 Chowla, Mann and Straus (cf. pp. 60-61 of [10]) used Vosper's theorem (cf. pp. 52-57 of [10]) to deduce that if  $p > 3$ ,  $1 < k < (p-1)/2$  and  $k \mid p-1$ , then

$$|\{a_1x_1^k + \dots + a_nx_n^k : x_1, \dots, x_n \in F_p\}| \geq \min \left\{ p, (2n-1)\frac{p-1}{k} + 1 \right\}.$$

Let  $F_q$  be the finite field of  $q$  elements where  $q > 1$  is a prime power. In 1993 Wan, Shiue and Chen [14] showed that if  $P(x)$  is a polynomial over  $F_q$  and  $l \in \mathbb{N} = \{0, 1, 2, \dots\}$  is the least nonnegative integer with  $\sum_{x \in F_q} P(x)^l \neq 0$  then  $|\{P(x) : x \in F_q\}| \geq l+1$ ; in 2004 Das [4] extended this to multi-variable polynomials over  $F_q$ . By modifying the proof of Theorem 1.5 of [4] slightly, one gets the following assertion: If  $P(x_1, \dots, x_n) \in F_q[x_1, \dots, x_n]$ ,  $\emptyset \neq S \subseteq F_q^n$ , and  $l$  is the smallest nonnegative integer with  $\sum_{(x_1, \dots, x_n) \in S} P(x_1, \dots, x_n)^l \neq 0$ , then we have  $|\{P(x_1, \dots, x_n) : (x_1, \dots, x_n) \in S\}| \geq l+1$ . Here the lower bound depends heavily on values of  $P(x_1, \dots, x_n)$ .

In this paper we investigate two kinds of value sets of a polynomial in the form (1.1).

Here is our first theorem which includes Felszeghy's result as a special case.

**Theorem 1.1.** *Let  $f(x_1, \dots, x_n)$  be a polynomial over a field  $F$  given by (1.1) and (1.2). Then, for any finite nonempty subsets  $A_1, \dots, A_n$  of  $F$ , we have*

$$|\{f(x_1, \dots, x_n) : x_1 \in A_1, \dots, x_n \in A_n\}| \geq \min \left\{ p(F), \sum_{i=1}^n \left\lfloor \frac{|A_i| - 1}{k} \right\rfloor + 1 \right\}. \quad (1.3)$$

This result will be proved in Section 2 where we also give an example to show that the inequality (1.3) is sharp when  $F$  is an algebraically closed field.

**Corollary 1.1.** (Cauchy–Davenport Theorem) *Let  $A_1, \dots, A_n$  be finite nonempty subsets of a field  $F$ . Then*

$$|A_1 + \dots + A_n| \geq \min\{p(F), |A_1| + \dots + |A_n| - n + 1\},$$

where the sumset  $A_1 + \dots + A_n$  is given by

$$A_1 + \dots + A_n = \{x_1 + \dots + x_n : x_1 \in A_1, \dots, x_n \in A_n\}.$$

*Proof.* Simply apply Theorem 1.1 with  $f(x_1, \dots, x_n) = x_1 + \dots + x_n$ .

*Remark 1.1.* The original Cauchy–Davenport theorem (see p. 44 of [10] or p. 200 of [13]) is Corollary 1.1 in the case  $n = 2$  and  $F = \mathbb{Z}/p\mathbb{Z}$  with  $p$  a prime.

**Corollary 1.2.** *Let  $F$  be a field of characteristic zero, and assume (1.1) and (1.2). Then, for any finite nonempty subset  $A$  of  $F$ , we have*

$$|\{f(x_1, \dots, x_n) : x_1, \dots, x_n \in A\}| \geq n \left\lfloor \frac{|A| - 1}{k} \right\rfloor + 1.$$

*Proof.* Just apply Theorem 1.1 with  $A_1 = \dots = A_n = A$ .

**Corollary 1.3.** *Let  $F$  be a field with prime characteristic  $p$ , and let  $f(x_1, \dots, x_n)$  be given by (1.1) and (1.2). If  $A$  is a finite subset of  $F$  satisfying  $\lfloor (|A| - 1)/k \rfloor \geq (p - 1)/n$ , then*

$$|\{f(x_1, \dots, x_n) : x_1, \dots, x_n \in A\}| \geq p.$$

*Proof.* It suffices to apply Theorem 1.1 with  $A_1 = \dots = A_n = A$ .

*Remark 1.2.* In the case  $A = F = F_p$ , Corollary 1.3 yields Felszeghy’s result.

Now we state our second theorem.

**Theorem 1.2.** *Let  $f(x_1, \dots, x_n)$  be a polynomial over a field  $F$  given by (1.1) and (1.2) with  $n \leq k = \deg f$ . And let  $A_1, \dots, A_n$  be finite subsets of  $F$  with  $|A_i| \geq i$  for  $i = 1, \dots, n$ . Then, for the restricted value set*

$$V = \{f(x_1, \dots, x_n) : x_1 \in A_1, \dots, x_n \in A_n, \text{ and } x_i \neq x_j \text{ if } i \neq j\}, \quad (1.4)$$

*we have*

$$|V| \geq \min \left\{ p(F), \sum_{i=1}^n \left\lfloor \frac{|A_i| - i}{k} \right\rfloor + 1 \right\}. \quad (1.5)$$

**Corollary 1.4.** *Let  $A$  be a finite subset of a field  $F$ , and let  $f(x_1, \dots, x_n)$  be a polynomial given by (1.1) and (1.2). Write  $|A| = kq + r$  with  $q, r \in \mathbb{N}$  and  $r < k$ . Then we have*

$$\begin{aligned} & |\{f(x_1, \dots, x_n) : x_1, \dots, x_n \in A, \text{ and } x_i \neq x_j \text{ if } i \neq j\}| \\ & \geq \begin{cases} \min\{p(F), n(q - 1) + \min\{n, r\} + 1\} & \text{if } n \leq k, \\ \min\{p(F), |A| - n + 1\} & \text{if } n \geq k. \end{cases} \end{aligned} \quad (1.6)$$

In Section 3 we shall prove Theorem 1.2 and Corollary 1.4, and give an example to illustrate that the inequality (1.5) is essentially best possible when  $F$  is an algebraically closed field.

In 1964 Erdős and Heilbronn [6] conjectured that if  $A$  is a subset of  $\mathbb{Z}/p\mathbb{Z}$  with  $p$  a prime then

$$|\{x_1 + x_2 : x_1, x_2 \in A \text{ and } x_1 \neq x_2\}| \geq \min\{p, 2|A| - 3\}.$$

Thirty years later this deep conjecture was confirmed by Dias da Silva and Hamidoune [5] who used the representation theory of groups to show that if  $A$  is a finite subset of a field  $F$  then

$$\begin{aligned} & |\{x_1 + x_2 + \dots + x_n : x_1, \dots, x_n \in A, \text{ and } x_i \neq x_j \text{ if } i \neq j\}| \\ & \geq \min\{p(F), n(|A| - n) + 1\}. \end{aligned}$$

This suggests that Corollary 1.4 in the case  $n > k$  might be further improved. In Section 4 we will discuss our following conjecture which extends the Dias da Silva–Hamidoune result in a new way. (The reader may consult [2], [8], [9] and [12] for other generalizations of the Erdős–Heilbronn conjecture.)

**Conjecture 1.1.** *Let  $f(x_1, \dots, x_n)$  be a polynomial over a field  $F$  given by (1.1) and (1.2), and let  $A$  be any finite subset of  $F$ . Provided  $n > k$ , we have*

$$\begin{aligned} & |\{f(x_1, \dots, x_n) : x_1, \dots, x_n \in A, \text{ and } x_i \neq x_j \text{ if } i \neq j\}| \\ & \geq \min \left\{ p(F) - \delta, \frac{n(|A| - n)}{k} - k \left\{ \frac{n}{k} \right\} \left\{ \frac{|A| - n}{k} \right\} + 1 \right\}, \end{aligned} \quad (1.7)$$

where  $\{\alpha\}$  denotes the fractional part  $\alpha - \lfloor \alpha \rfloor$  of a real number  $\alpha$ , and

$$\delta = \begin{cases} 1 & \text{if } n = 2 \text{ and } a_1 = -a_2, \\ 0 & \text{otherwise.} \end{cases}$$

## 2 Proof of Theorem 1.1

We need a useful tool of algebraic nature.

**Combinatorial Nullstellensatz** (Alon [1]). *Let  $A_1, \dots, A_n$  be finite subsets of a field  $F$ , and let  $P(x_1, \dots, x_n) \in F[x_1, \dots, x_n]$ . Suppose that  $\deg P = k_1 + \dots + k_n$  where  $0 \leq k_i < |A_i|$  for  $i = 1, \dots, n$ . Then  $P(x_1, \dots, x_n) \neq 0$  for some  $x_1 \in A_1, \dots, x_n \in A_n$  if*

$$[x_1^{k_1} \dots x_n^{k_n}]P(x_1, \dots, x_n) \neq 0,$$

where  $[x_1^{k_1} \dots x_n^{k_n}]P(x_1, \dots, x_n)$  denotes the coefficient of  $x_1^{k_1} \dots x_n^{k_n}$  in  $P(x_1, \dots, x_n)$ .

*Proof of Theorem 1.1.* Let  $m$  be the largest nonnegative integer not exceeding  $n$  such that  $\sum_{0 < i \leq m} \lfloor (|A_i| - 1)/k \rfloor < p(F)$ . For each  $0 < i \leq m$  let  $A'_i$  be a subset of  $A_i$  with cardinality  $k \lfloor (|A_i| - 1)/k \rfloor + 1$ . In the case  $m < n$ ,  $p = p(F)$  is a prime and we let  $A'_{m+1}$  be a subset of  $A_{m+1}$  with

$$\begin{aligned} |A'_{m+1}| &= k \left( p - 1 - \sum_{0 < i \leq m} \left\lfloor \frac{|A_i| - 1}{k} \right\rfloor \right) + 1 \\ &< k \left\lfloor \frac{|A_{m+1}| - 1}{k} \right\rfloor + 1 \leq |A_{m+1}|. \end{aligned}$$

If  $m + 1 < j \leq n$  then we let  $A'_j \subseteq A_j$  be a singleton. Whether  $m = n$  or not, we have  $\sum_{i=1}^n (|A'_i| - 1) = k(N - 1)$ , where

$$N = \min \left\{ p(F), \sum_{i=1}^n \left\lfloor \frac{|A_i| - 1}{k} \right\rfloor + 1 \right\}.$$

Set

$$C = \{f(x_1, \dots, x_n) : x_1 \in A'_1, \dots, x_n \in A'_n\}.$$

Suppose that  $|C| \leq N - 1$ . Then

$$\begin{aligned} & [x_1^{|A'_1|-1} \cdots x_n^{|A'_n|-1}] f(x_1, \dots, x_n)^{N-1-|C|} \prod_{c \in C} (f(x_1, \dots, x_n) - c) \\ &= [x_1^{|A'_1|-1} \cdots x_n^{|A'_n|-1}] (a_1 x_1^k + \cdots + a_n x_n^k)^{N-1} \\ &= \frac{(N-1)!}{\prod_{i=1}^n ((|A'_i|-1)/k)!} a_1^{(|A'_1|-1)/k} \cdots a_n^{(|A'_n|-1)/k} \neq 0. \end{aligned}$$

By the Combinatorial Nullstellensatz, for some  $x_1 \in A'_1, \dots, x_n \in A'_n$  we have

$$f(x_1, \dots, x_n)^{N-1-|C|} \prod_{c \in C} (f(x_1, \dots, x_n) - c) \neq 0$$

which contradicts the fact  $f(x_1, \dots, x_n) \in C$ .

In view of the above,

$$|\{f(x_1, \dots, x_n) : x_1 \in A_1, \dots, x_n \in A_n\}| \geq |C| \geq N$$

and this concludes the proof.

**Example 2.1.** Let  $a_1, \dots, a_n \in F^* = F \setminus \{0\}$ , where  $F$  is an algebraically closed field. For each  $i = 1, \dots, n$  let

$$A_i = \{x \in F : f_k(x) \in \{a_i^{-1}, 2a_i^{-1}, \dots, q_i a_i^{-1}\}\} \setminus R_i,$$

where  $k$  and  $q_i < p(F)$  are positive integers,

$$f_k(x) = \begin{cases} x^k - x & \text{if } p(F) \text{ is a prime divisor of } k, \\ x^k & \text{otherwise,} \end{cases} \quad (2.1)$$

and  $R_i$  is a subset of  $\{x \in F : f_k(x) = q_i a_i^{-1}\}$  with  $|R_i| = r_i \leq k - 1$ . For each  $c \in F^*$ , the equation  $f_k(x) = c$  has exactly  $k$  distinct solutions in  $F$  since there is no  $x \in F$  satisfying  $f_k(x) - c = 0 = f'_k(x)$ , where  $f'_k$  is the formal derivative of  $f_k$ . So  $|A_i| = kq_i - r_i$  and hence

$$\left\lfloor \frac{|A_i| - 1}{k} \right\rfloor = \left\lfloor \frac{k(q_i - 1) + (k - 1 - r_i)}{k} \right\rfloor = q_i - 1.$$

For every  $i = 1, \dots, n$  we have

$$\{f_k(a) : a \in A_i\} = \{a_i^{-1}, 2a_i^{-1}, \dots, q_i a_i^{-1}\}.$$

Thus

$$\begin{aligned} & \{a_1 f_k(x_1) + \cdots + a_n f_k(x_n) : x_1 \in A_1, \dots, x_n \in A_n\} \\ &= \{a_1 (y_1 a_1^{-1}) + \cdots + a_n (y_n a_n^{-1}) : y_i \in \{1, \dots, q_i\} \text{ for } i = 1, \dots, n\} \\ &= \{(y_1 + \cdots + y_n)e : y_i \in \{1, \dots, q_i\} \text{ for } i = 1, \dots, n\} \\ &= \{ne, (n+1)e, \dots, (q_1 + \cdots + q_n)e\}, \end{aligned}$$

where  $e$  denotes the multiplicative identity of the field  $F$ . Observe that

$$q_1 + \cdots + q_n - n = \sum_{i=1}^n (q_i - 1) = \sum_{i=1}^n \left\lfloor \frac{|A_i| - 1}{k} \right\rfloor.$$

Therefore

$$\begin{aligned} & |\{a_1 f_k(x_1) + \cdots + a_n f_k(x_n) : x_1 \in A_1, \dots, x_n \in A_n\}| \\ &= \min \left\{ p(F), \sum_{i=1}^n \left\lfloor \frac{|A_i| - 1}{k} \right\rfloor + 1 \right\}. \end{aligned}$$

### 3 Proofs of Theorem 1.2 and Corollary 1.4

*Proof of Theorem 1.2.* Let  $q_i = \lfloor (|A_i| - i)/k \rfloor$  for  $i = 1, \dots, n$ . And let  $m$  be the largest nonnegative integer not exceeding  $n$  such that  $\sum_{0 < i \leq m} q_i < p(F)$ . For each  $0 < i \leq m$  let  $A'_i$  be a subset of  $A_i$  with cardinality  $kq_i + i$ . In the case  $m < n$ ,  $p = p(F)$  is a prime and we let  $A'_{m+1}$  be a subset of  $A_{m+1}$  with

$$\begin{aligned} |A'_{m+1}| &= k \left( p - 1 - \sum_{0 < i \leq m} q_i \right) + m + 1 \\ &< kq_{m+1} + m + 1 \leq |A_{m+1}|. \end{aligned}$$

If  $m + 1 < j \leq n$  then we let  $A'_j \subseteq A_j$  with  $|A'_j| = j$ . Whether  $m = n$  or not, we have  $\sum_{i=1}^n (|A'_i| - i) = k(N - 1)$ , where

$$N = \min\{p(F), q_1 + \cdots + q_n + 1\}.$$

Set

$$C = \{f(x_1, \dots, x_n) : x_1 \in A'_1, \dots, x_n \in A'_n, \text{ and } x_i \neq x_j \text{ if } i \neq j\}.$$

Suppose that  $|C| \leq N - 1$  and let  $P(x_1, \dots, x_n)$  denote the polynomial

$$f(x_1, \dots, x_n)^{N-1-|C|} \prod_{c \in C} (f(x_1, \dots, x_n) - c) \times \prod_{1 \leq i < j \leq n} (x_j - x_i).$$

Then

$$\deg P \leq k(N - 1) + \binom{n}{2} = \sum_{i=1}^n (|A'_i| - 1).$$

By linear algebra,

$$\prod_{1 \leq i < j \leq n} (x_j - x_i) = \det(x_i^{j-1})_{1 \leq i, j \leq n} = \sum_{\sigma \in S_n} \text{sign}(\sigma) \prod_{i=1}^n x_i^{\sigma(i)-1},$$

where  $S_n$  is the symmetric group of all permutations on  $\{1, \dots, n\}$ , and  $\text{sign}(\sigma)$  is 1 or  $-1$  according as  $\sigma$  is even or odd. Recall that

$$|A'_i| = \begin{cases} kq_i + i & \text{if } 1 \leq i \leq m, \\ k(p(F) - 1 - q_1 - \cdots - q_m) + i & \text{if } i = m + 1 \leq n, \\ i & \text{if } m + 1 < i \leq n. \end{cases}$$

For each  $i = 1, \dots, n$ , we clearly have  $|A'_i| - 1 \equiv i - 1 \pmod{k}$  and  $\sigma(i) - 1 < n \leq k$  for all  $\sigma \in S_n$ . Thus

$$\begin{aligned} & \left[ x_1^{|A'_1|-1} \cdots x_n^{|A'_n|-1} \right] P(x_1, \dots, x_n) \\ &= \left[ x_1^{|A'_1|-1} \cdots x_n^{|A'_n|-1} \right] (a_1 x_1^k + \cdots + a_n x_n^k)^{N-1} \sum_{\sigma \in S_n} \text{sign}(\sigma) \prod_{i=1}^n x_i^{\sigma(i)-1} \\ &= \left[ \prod_{i=1}^n x_i^{|A'_i|-1} \right] \sum_{j_1 + \cdots + j_n = N-1} \frac{(N-1)!}{j_1! \cdots j_n!} a_1^{j_1} \cdots a_n^{j_n} \prod_{i=1}^n x_i^{j_i k + i - 1} \\ &= \frac{(N-1)!}{q_1! \cdots q_m! q!} a_1^{q_1} \cdots a_m^{q_m} a \neq 0, \end{aligned}$$

where  $q = a = 1$  if  $m = n$ , and  $q = (p(F) - 1 - q_1 - \cdots - q_m)!$  and  $a = a_{m+1}^q$  if  $m < n$ . In light of the Combinatorial Nullstellensatz, there are  $x_1 \in A'_1, \dots, x_n \in A'_n$  such that  $P(x_1, \dots, x_n) \neq 0$ . Obviously this contradicts the fact  $f(x_1, \dots, x_n) \in C$ .

By the above,  $|V| \geq |C| \geq N$  and hence (1.5) holds.

**Example 3.1.** Let  $F$  be any algebraically closed field, and let  $a_1, \dots, a_n \in F^*$ . For each  $i = 1, \dots, n$  let

$$A_i = S_i \cup \{x \in F : f_k(x) \in \{j a_i^{-1} : 1 < j \leq q_i\}\},$$

where  $k \geq n$  and  $q_i < p(F)$  are positive integers,  $f_k(x)$  is given by (2.1), and  $S_i$  is a subset of  $\{x \in F : f_k(x) = a_i^{-1}\}$  with  $|S_i| \geq i$ . Observe that  $|A_i| = k(q_i - 1) + |S_i|$  and hence

$$\left\lfloor \frac{|A_i| - i}{k} \right\rfloor = q_i - 1.$$

For every  $i = 1, \dots, n$ , we have

$$\{f_k(a) : a \in A_i\} = \{a_i^{-1}, 2a_i^{-1}, \dots, q_i a_i^{-1}\}.$$

If  $y_i \in \{1, \dots, q_i\}$  for  $i = 1, \dots, n$ , we can find distinct  $x_1 \in A_1, \dots, x_n \in A_n$  such that  $f_k(x_i) = y_i a_i^{-1}$  for  $i = 1, \dots, n$ ; in fact, if  $x_1 \in A_1, \dots, x_{i-1} \in A_{i-1}$  are distinct with  $i \leq n$ , and  $f_k(x_j) = y_j a_j^{-1}$  for  $j = 1, \dots, i-1$ , then we can choose  $x_i \in A_i \setminus \{x_1, \dots, x_{i-1}\}$  satisfying  $f_k(x_i) = y_i a_i^{-1}$  because  $k \geq |S_i| > i - 1$ . Thus

$$\begin{aligned} & \{a_1 f_k(x_1) + \cdots + a_n f_k(x_n) : x_1 \in A_1, \dots, x_n \in A_n, \text{ and } x_i \neq x_j \text{ if } i \neq j\} \\ &= \{a_1 (y_1 a_1^{-1}) + \cdots + a_n (y_n a_n^{-1}) : y_i \in \{1, \dots, q_i\} \text{ for } i = 1, \dots, n\} \\ &= \{(y_1 + \cdots + y_n) e : y_i \in \{1, \dots, q_i\} \text{ for } i = 1, \dots, n\} \\ &= \{n e, (n+1)e, \dots, (q_1 + \cdots + q_n)e\}, \end{aligned}$$

where  $e$  is the multiplicative identity of  $F$ . Note that

$$q_1 + \cdots + q_n - n = \sum_{i=1}^n (q_i - 1) = \sum_{i=1}^n \left\lfloor \frac{|A_i| - i}{k} \right\rfloor.$$

So we have

$$\begin{aligned} & |\{a_1 f_k(x_1) + \cdots + a_n f_k(x_n) : x_1 \in A_1, \dots, x_n \in A_n, \text{ and } x_i \neq x_j \text{ if } i \neq j\}| \\ &= \min \left\{ p(F), \sum_{i=1}^n \left\lfloor \frac{|A_i| - i}{k} \right\rfloor + 1 \right\}. \end{aligned}$$

*Proof of Corollary 1.4.* The case  $|A| < n$  is trivial; below we assume  $|A| \geq n$ .

We first handle the case  $n \leq k$ . If  $1 \leq i \leq \min\{n, r\}$  then we let  $A_i$  be a subset of  $A$  with cardinality  $kq + i + \max\{r - n, 0\} \leq kq + r = |A|$ ; if  $r < j \leq n$  then we let  $A_j$  be a subset of  $A$  with cardinality  $k(q - 1) + j \leq k(q - 1) + n \leq kq$ . (Note that when  $r < n$  we have  $q \neq 0$  since  $|A| \geq n$ .) Obviously,

$$\sum_{i=1}^n \left\lfloor \frac{|A_i| - i}{k} \right\rfloor = \sum_{i=1}^{\min\{n, r\}} q + \sum_{r < j \leq n} (q - 1) = n(q - 1) + \min\{n, r\}.$$

Applying Theorem 1.2 we obtain that

$$\begin{aligned} & |\{f(x_1, \dots, x_n) : x_1, \dots, x_n \in A, \text{ and } x_i \neq x_j \text{ if } i \neq j\}| \\ & \geq |\{f(x_1, \dots, x_n) : x_1 \in A_1, \dots, x_n \in A_n, \text{ and } x_i \neq x_j \text{ if } i \neq j\}| \\ & \geq \min\{p(F), n(q - 1) + \min\{n, r\} + 1\}. \end{aligned}$$

In particular, if  $k = n$  then

$$\begin{aligned} & |\{f(x_1, \dots, x_n) : x_1, \dots, x_n \in A, \text{ and } x_i \neq x_j \text{ if } i \neq j\}| \\ & \geq \min\{p(F), n(q - 1) + r + 1\} = \min\{p(F), |A| - n + 1\}. \end{aligned}$$

Now suppose that  $n > k$ . Let  $c_{k+1}, \dots, c_n$  be  $n - k$  distinct elements of  $A$ . Then  $A' = A \setminus \{c_{k+1}, \dots, c_n\}$  has cardinality  $|A| - (n - k)$ . By what we have proved,

$$\begin{aligned} & |\{f(x_1, \dots, x_k, c_{k+1}, \dots, c_n) : x_1, \dots, x_k \in A', \text{ and } x_i \neq x_j \text{ if } i < j\}| \\ & \geq \min\{p(F), |A'| - k + 1\} = \min\{p(F), |A| - n + 1\}. \end{aligned}$$

So the desired inequality follows.

## 4. Discussion of Conjecture 1.1

Conjecture 1.1 in the case  $f(x_1, \dots, x_n) = x_1 + \dots + x_n$ , essentially gives the Dias da Silva–Hamidoune result mentioned in Section 1.

In the case  $f(x_1, \dots, x_n) = x_1^k + \dots + x_n^k$  with  $k > 1$ , we may explain the symmetry between  $n$  and  $|A| - n$  as follows: If  $|A| > n$  then

$$\begin{aligned} & |\{x_1^k + \dots + x_n^k : x_1, \dots, x_n \in A, \text{ and } x_i \neq x_j \text{ if } i \neq j\}| \\ & = \left| \left\{ \sum_{a \in A} a^k - y_1^k - \dots - y_{|A|-n}^k : y_1, \dots, y_{|A|-n} \in A, \text{ and } y_i \neq y_j \text{ if } i \neq j \right\} \right| \\ & = |\{y_1^k + \dots + y_{|A|-n}^k : y_1, \dots, y_{|A|-n} \in A, \text{ and } y_i \neq y_j \text{ if } i \neq j\}|. \end{aligned}$$

Conjecture 1.1 holds when  $n = 2$ . In fact, if we set  $A_1 = \{a_1x : x \in A\}$  and  $A_2 = \{a_2x : x \in A\}$ , then  $|A_1| + |A_2| - 3 = 2(|A| - 2) + 1$  and

$$\begin{aligned} & |\{a_1x_1 + a_2x_2 : x_1, x_2 \in A \text{ and } x_1 \neq x_2\}| \\ & = |\{y_1 + y_2 : y_1 \in A_1, y_2 \in A_2 \text{ and } a_1^{-1}y_1 \neq a_2^{-1}y_2\}| \\ & = |\{y_1 + y_2 : y_1 \in A_1, y_2 \in A_2 \text{ and } y_1 - a_1a_2^{-1}y_2 \neq 0\}| \\ & \geq \min\{p(F) - \delta, |A_1| + |A_2| - 3\} \quad (\text{by Corollary 3 of [11]}). \end{aligned}$$



The following example illustrates how the lower bound in (1.7) comes out.

**Example 4.1.** Let  $k, n \in \mathbb{Z}^+$  and  $q \in \mathbb{N}$ . Let

$$A = \{z \in \mathbb{C} : z^k \in \{1, \dots, q\}\} \cup R,$$

where  $\mathbb{C}$  is the field of complex numbers and  $R$  is a subset of  $\{z \in \mathbb{C} : z^k = q + 1\}$  with cardinality  $r < k$ . Suppose that  $n \geq k$  and  $|A| = kq + r \geq n = k\lfloor n/k \rfloor + s$  with  $s \in \{0, \dots, k-1\}$ . Clearly,

$$V = \{x_1^k + \dots + x_n^k : x_1, \dots, x_n \in A, \text{ and } x_i \neq x_j \text{ if } i \neq j\}$$

just consists of those sums  $\sum_{i=1}^n y_i$  with  $y_i \in \{1, \dots, q+1\}$ ,  $|\{i : y_i = q+1\}| \leq r$  and  $|\{i : y_i = j\}| \leq k$  for all  $j = 1, \dots, q$ . Thus the smallest element of  $V$  is

$$m_V = k \sum_{i=1}^{\lfloor n/k \rfloor} i + s \left( \left\lfloor \frac{n}{k} \right\rfloor + 1 \right) = \left( \frac{k}{2} \left\lfloor \frac{n}{k} \right\rfloor + s \right) \left( \left\lfloor \frac{n}{k} \right\rfloor + 1 \right),$$

while the largest element of  $V$  is

$$M_V = r(q+1) + k \sum_{i=0}^{\lfloor n/k \rfloor - 1} (q-i) + (s-r) \left( q - \left\lfloor \frac{n}{k} \right\rfloor \right) - d(r, s),$$

where

$$d(r, s) = \begin{cases} r - s & \text{if } r \geq s, \\ 0 & \text{if } r < s. \end{cases}$$

It follows that

$$\begin{aligned} & d(r, s) + M_V - m_V \\ &= r(q+1) + kq \left\lfloor \frac{n}{k} \right\rfloor - k \left\lfloor \frac{n}{k} \right\rfloor^2 + (s-r)q + (r-s) \left\lfloor \frac{n}{k} \right\rfloor - s \left( \left\lfloor \frac{n}{k} \right\rfloor + 1 \right) \\ &= \left\lfloor \frac{n}{k} \right\rfloor \left( kq + r - s - k \left\lfloor \frac{n}{k} \right\rfloor - s \right) + r + sq - s \\ &= \left\lfloor \frac{n}{k} \right\rfloor (|A| - s - n) + s(q-1) + r \\ &= \left\lfloor \frac{n}{k} \right\rfloor (|A| - n) + s \left( q - \left\lfloor \frac{n}{k} \right\rfloor - 1 \right) + r \\ &= \left\lfloor \frac{n}{k} \right\rfloor (|A| - n) + s \left\lfloor \frac{|A| - n}{k} \right\rfloor + \begin{cases} r - s & \text{if } r \geq s, \\ r & \text{if } r < s. \end{cases} \end{aligned}$$

Therefore

$$\begin{aligned} |V| &= |\{m_V, \dots, M_V\}| = M_V - m_V + 1 \\ &= 1 + \left\lfloor \frac{n}{k} \right\rfloor (|A| - n) + s \left\lfloor \frac{|A| - n}{k} \right\rfloor + \begin{cases} r & \text{if } r < s, \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Finally, we note that

$$\begin{aligned}
& \left\lfloor \frac{n}{k} \right\rfloor (|A| - n) + s \left\lfloor \frac{|A| - n}{k} \right\rfloor \\
&= \left( \frac{n}{k} - \left\{ \frac{n}{k} \right\} \right) (|A| - n) + k \left\{ \frac{n}{k} \right\} \left\lfloor \frac{|A| - n}{k} \right\rfloor \\
&= \frac{n(|A| - n)}{k} - k \left\{ \frac{n}{k} \right\} \left\{ \frac{|A| - n}{k} \right\}.
\end{aligned}$$

**Acknowledgments.** The author thanks the referees for their helpful comments.

## References

- [1] N. Alon, *Combinatorial Nullstellensatz*, *Combin. Probab. Comput.* **8**(1999), 7–29.
- [2] N. Alon, M. B. Nathanson and I. Z. Ruzsa, *The polynomial method and restricted sums of congruence classes*, *J. Number Theory* **56**(1996), 404–417.
- [3] L. Carlitz, *Solvability of certain equations in a finite field*, *Quart. J. Math.* **7**(1956), 3–4.
- [4] P. Das, *Value sets of polynomials and the Cauchy-Davenport theorem*, *Finite Fields Appl.* **10**(2004), 113–122.
- [5] J. A. Dias da Silva and Y. O. Hamidoune, *Cyclic spaces for Grassmann derivatives and additive theory*, *Bull. London Math. Soc.* **26**(1994), 140–146.
- [6] P. Erdős and H. Heilbronn, *On the addition of residue classes modulo  $p$* , *Acta Arith.* **9**(1964), 149–159.
- [7] B. Felszeghy, *On the solvability of some special equations over finite fields*, *Publ. Math. Debrecen* **68**(2006), 15–23.
- [8] Q. H. Hou and Z. W. Sun, *Restricted sums in a field*, *Acta Arith.* **102**(2002), 239–249.
- [9] G. Károlyi, *The Erdős-Heilbronn problem in abelian groups*, *Israel J. Math.* **139**(2004), 349–359.
- [10] M. B. Nathanson, *Additive Number Theory: Inverse Problems and the Geometry of Sumsets*, *Graduate Texts in Math.*, Vol. 165, Springer, New York, 1996.
- [11] H. Pan and Z. W. Sun, *A lower bound for  $|\{a + b : a \in A, b \in B, P(a, b) \neq 0\}|$* , *J. Combin. Theory Ser. A* **100**(2002), 387–393.
- [12] Z. W. Sun, *On Snevily’s conjecture and restricted sumsets*, *J. Combin. Theory Ser. A* **103**(2003), 288–301.
- [13] T. Tao and V. H. Vu, *Additive Combinatorics*, Cambridge Univ. Press, Cambridge, 2006.
- [14] D. Wan, P. J.-S. Shiue and C. S. Chen, *Value sets of polynomials over finite fields*, *Proc. Amer. Math. Soc.* **119**(1993), 711–717.