

# TWISTED HOCHSCHILD HOMOLOGY AND MACLANE HOMOLOGY

TEIMURAZ PIRASHVILI

ABSTRACT. We prove that  $H_i(A, \Phi(A)) = 0$ ,  $i > 0$ . Here  $A$  is a commutative algebra over the prime field  $\mathbb{F}_p$  of characteristic  $p > 0$  and  $\Phi(A)$  is  $A$  considered as a bimodule, where the left multiplication is the usual one, while the right multiplication is given via Frobenius endomorphism and  $H_\bullet$  denotes the Hochschild homology over  $\mathbb{F}_p$ . This result has implications in MacLane homology theory. Among other results, we prove that  $\text{HML}_\bullet(A, T) = 0$ , provided  $A$  is an algebra over a field  $K$  of characteristic  $p > 0$  and  $T$  is a strict homogeneous polynomial functor of degree  $d$  with  $1 < d < \text{Card}(K)$ .

## 1. INTRODUCTION

In this short note we study Hochschild and MacLane homology of commutative algebras over the prime field  $\mathbb{F}_p$  of characteristic  $p > 0$ . Let us recall that MacLane homology is isomorphic to the topological Hochschild homology [13] and to the stable  $K$ -theory as well [5].

Let  $A$  be a commutative algebra over the prime field  $\mathbb{F}_p$  of characteristic  $p > 0$  and let  $\Phi(A)$  be denote  $A$ - $A$ -bimodule, which is  $A$  as a left  $A$ -module, while the right multiplication is given via Frobenius endomorphism. We prove that the Hochschild homology vanishes  $H_i(A, \Phi(A)) = 0$ ,  $i > 0$ . The proof makes use a simple result on homotopy groups of simplicial rings, which says that if  $R_\bullet$  is a simplicial ring such that all rings involved in  $R_\bullet$  satisfy  $x^m = x$ ,  $m \geq 2$  identity then  $\pi_i(R_\bullet) = 0$  for all  $i > 0$ . These results has implications in MacLane homology theory. We extend the computation of Franjou-Lannes and Schwartz [4] of MacLane (co)homology of finite fields with coefficients in symmetric  $S^d$  and divided powers  $\Gamma^d$  to arbitrary commutative  $\mathbb{F}_p$ -algebras, provided that  $d > 1$ . As a consequence of our computations we show that  $\text{HML}_\bullet(A, T) = 0$ , provided  $T$  is a strict homogeneous polynomial functor of degree  $d > 1$  and  $A$  is an algebra over a field  $K$  of characteristic  $p > 0$  with  $\text{Card}(K) > d$ .

Many thanks to referee for his valuable comments and to Petter Andreas Bergh for his remarks.

## 2. WHEN IT IS TOO EASY TO COMPUTE HOMOTOPY GROUPS

It is well-known that the homotopy groups of a simplicial abelian group  $(A_\bullet, \partial_\bullet, s_\bullet)$  can be computed as the homology of the normalized chain complex  $(N_\bullet(A_\bullet), d)$ , where

$$N_n(A_\bullet) = \{x \in A_n \mid \partial_i(x) = 0, i > 0\}$$

---

\* Partially supported by the grant from MEC, MTM2006-15338-C02-02 (European FEDER support included).

and the boundary map  $N_n(A_\bullet) \rightarrow N_{n-1}(A_\bullet)$  is induced by  $\partial_0$ . Our first result shows that if  $A_\bullet$  has a simplicial ring structure and the rings involved in  $A_\bullet$  satisfy extra conditions then homotopy groups are zero in positive dimensions. This fact is an easy consequence of the following result which is probably well-known

**Lemma 1.** Let  $R_\bullet$  be a simplicial object in the category of not necessarily associative rings and let  $x, y \in N_n(R_\bullet)$  be two elements. Assume  $n > 0$  and  $x$  is a cycle. Then the cycle  $xy \in N_n(R_\bullet)$  is a coboundary.

*Proof.* Consider the element

$$z = s_0(xy) - s_1(x)s_0(y).$$

Then we have

$$\partial_0(z) = xy - (s_0\partial_0(x))y = xy.$$

Moreover,

$$\partial_1(z) = xy - xy = 0,$$

We also have

$$\partial_2(z) = (s_0\partial_1(x))(s_0\partial_1(y)) - x(s_0\partial_1(y)) = 0.$$

Similarly for all  $i > 2$  we have

$$\partial_i(z) = (s_0\partial_{i-1}(x))(s_0\partial_{i-1}(y)) - (s_1\partial_{i-1}(x))(s_0\partial_{i-1}(y)) = 0.$$

Hence  $z$  is an element of  $N_{n+1}(R_\bullet)$  with  $\partial(z) = xy$ .  $\square$

**Corollary 2.** Let  $R_\bullet$  be a simplicial ring. If the rings involved in  $R_\bullet$  satisfy  $x^m = x$  identity for  $m \geq 2$ , then

$$\pi_n(R_\bullet) = 0, \quad n > 0.$$

*Proof.* Take a cycle  $x \in N_n(R_\bullet)$ ,  $n > 0$ . Then the class of  $x = xx^{m-1}$  in  $\pi_n(R_\bullet)$  is zero.  $\square$

**Remark.** A more general fact is true. Let  $\mathbf{T}$  be a pointed algebraic theory [15] and let  $X_\bullet$  be a simplicial object in the category of  $\mathbf{T}$ -models [15]. Then  $\pi_1(X_\bullet)$  is a group object in the category of  $\mathbf{T}$ -models, while  $\pi_i(X_\bullet)$  are abelian group objects in the category of  $\mathbf{T}$ -models for all  $i > 1$ . Thus  $\pi_i(X_\bullet) = 0$ ,  $i \geq 1$  provided all group objects are trivial. This is what happens for the category of rings satisfying the identity  $x^m = x$ ,  $m \geq 2$ . Another interesting case is the category of Heyting algebras [3].

### 3. HOCHSCHILD HOMOLOGY WITH TWISTED COEFFICIENTS

In what follows the ground field is the prime field  $\mathbb{F}_p$  of characteristic  $p > 0$ . All algebras are taken over  $\mathbb{F}_p$  and they are assumed to be associative. For an algebra  $R$  and an  $R$ - $R$ -bimodule  $B$  we let  $\mathbf{H}_\bullet(R, B)$  and  $\mathbf{H}^\bullet(R, B)$  be the Hochschild homology and cohomology of  $R$  with coefficients in  $B$ . Let us recall that

$$\mathbf{H}_\bullet(R, B) = \mathrm{Tor}_\bullet^{R \otimes R^{op}}(R, B)$$

and

$$\mathbf{H}^\bullet(R, B) = \mathrm{Ext}_{R \otimes R^{op}}^\bullet(R, B).$$

Moreover, we let  $C_\bullet(R, B)$  be the standard simplicial vector space computing Hochschild homology

$$\pi_\bullet(C_\bullet(R, B)) \cong \mathbf{H}_\bullet(R, B).$$

Recall that  $C_n(R, B) = B \otimes R^{\otimes n}$ , while

$$\partial_0(b, r_1, \dots, r_n) = (br_1, \dots, r_n),$$

$$\partial_i(b, r_1, \dots, r_n) = (b, r_1, \dots, r_i r_{i+1}, \dots, r_n), \quad 0 < i < n$$

and

$$\partial_n(b, r_1, \dots, r_n) = (r_n b, r_1, \dots, r_{n-1}).$$

Here  $b \in B$  and  $r_1, \dots, r_n \in R$ .

Let  $n > 1$  be a natural number and let  $A$  be a commutative  $\mathbb{F}_p$ -algebra. The Frobenius homomorphism gives rise to the functors  $\Phi^n$  from the category of  $A$ -modules to the category of  $A$ - $A$ -bimodules, which are defined as follows. For an  $A$ -module  $M$  the bimodule  $\Phi^n(M)$  coincides with  $M$  as a left  $A$ -module, while the right  $A$ -module structure on  $\Phi^n(M)$  is given by

$$ma = a^{p^n} m, \quad a \in A, m \in M.$$

Having  $A$ - $A$ -bimodule  $\Phi^n(M)$  we can consider the Hochschild homology  $\mathbf{H}_\bullet(A, \Phi^n(M))$ . In this section we study these homologies. In order to state our results we need some notation. We let  $\psi^n(A)$  be the quotient ring  $A/(a - a^{p^n})$ ,  $n \geq 1$  which is considered as an  $A$ -module via the quotient map  $A \twoheadrightarrow \psi^n(A)$ . Thus  $\psi^n$  is the left adjoint of the inclusion of the category of commutative  $\mathbb{F}_p$ -algebras with identity  $x^m = x, m = p^n$  to the category of all commutative  $\mathbb{F}_p$ -algebras.

**Example 3.** Let  $n > 1$ . If  $K$  is a finite field with  $q = p^d$  element then  $\psi^n(K) = K$  if  $n = dt, t \in \mathbb{N}$  and  $\psi^n(K) = 0$  if  $n \neq dt, t \in \mathbb{N}$ .

**Lemma 4.** Let  $A$  is a commutative algebra over a field  $K$  of characteristic  $p > 0$  with  $\text{Card}(K) > p^n$ . Then  $\psi^n(A) = 0, n > 1$ .

*Proof.* By assumption there exists  $k \in K$  such that  $k^{p^n} - k$  is an invertible element of  $K$ . It follows then that the elements of the form  $a^{p^n} - a$  generates whole  $A$ .  $\square$

**Theorem 5.** Let  $A$  be a commutative  $\mathbb{F}_p$ -algebra and  $n > 1$ . Then

$$\mathbf{H}_i(A, \Phi^n(A)) = 0$$

for all  $i > 0$  and

$$\mathbf{H}_0(A, \Phi^n(A)) \cong \psi^n(A).$$

*Proof.* The proof consists of three steps.

**Step 1. The theorem holds if  $A = \mathbb{F}_p[x]$ .** In this case we have the following projective resolution of  $A$  over  $A \otimes A = \mathbb{F}_p[x, y]$ :

$$0 \rightarrow \mathbb{F}_p[x, y] \xrightarrow{\eta} \mathbb{F}_p[x, y] \xrightarrow{\epsilon} \mathbb{F}_p[x] \rightarrow 0.$$

Here  $\epsilon(x) = \epsilon(y) = x$  and  $\eta$  is induced by multiplication by  $(x - y)$ . Hence for any  $A$ - $A$ -bimodule  $B$ , we have  $H_i(A, B) = 0$  for  $i > 1$  and

$$H_0(A, B) \cong \text{Coker}(u) \text{ and } H_1(A, B) \cong \text{Ker}(u),$$

where  $u : B \rightarrow B$  is given by  $u(b) = xb - bx$ . If  $B = \Phi^n(\mathbb{F}_p[x])$ , then  $u : \mathbb{F}_p[x] \rightarrow \mathbb{F}_p[x]$  is the multiplication by  $(x^{p^n} - x)$  and we obtain  $H_1(A, \Phi^n(A)) = 0$  and  $H_0(A, \Phi^n(A)) = \psi^n(A)$

**Step 2. The theorem holds if  $A$  is a polynomial algebra.** Since Hochschild homology commutes with filtered colimits it suffices to consider the case when  $A = \mathbb{F}_p[x_1, \dots, x_d]$ . By the Künneth theorem for Hochschild homology (see [10, Theorem X.7.4]) we have  $H_\bullet(A, \Phi^n(A)) = H_\bullet(\mathbb{F}[x], \Phi^n(\mathbb{F}[x]))^{\otimes d}$  and the result follows.

**Step 3. The theorem holds for arbitrary  $A$ .** We use the same method as used in the proof of [9, Theorem 3.5.8]. First we choose a simplicial commutative algebra  $L_\bullet$  such that each  $L_n$  is a polynomial algebra,  $n \geq 0$  and  $\pi_i(L_\bullet) = 0$  for all  $i > 0$ ,  $\pi_0(L_\bullet) = A$ . Such a resolution exist thanks to [14]. Now consider the bisimplicial vector space  $C_\bullet(L_\bullet, \Phi^n(L_\bullet))$ . The  $s$ -th horizontal simplicial vector space is the simplicial vector space  $L_\bullet^{\otimes s+1}$ . By the Eilenberg-Zilber-Cartier and Künneth theorems it has zero homotopy groups in positive dimensions and  $\pi_0(L_\bullet^{\otimes s+1}) = A^{\otimes s+1}$ . On the other hand the  $t$ -th vertical simplicial vector space of  $C_\bullet(L_\bullet, \Phi^n(L_\bullet))$  is isomorphic to the Hochschild complex  $C_\bullet(L_t, \Phi^n(L_t))$  which has zero homology in positive dimensions by the previous step. Hence both spectral sequences corresponding to the bisimplicial vector space  $C_\bullet(L_\bullet, \Phi^n(L_\bullet))$  degenerate and we obtain the isomorphism

$$H_\bullet(A, \Phi^n(A)) \cong \pi_\bullet(\psi^n(L_\bullet)).$$

Now we can use Lemma 2 to finish the proof.  $\square$

**Corollary 6.** Let  $A$  be a commutative  $\mathbb{F}_p$ -algebra,  $M$  be an  $A$ -module and  $n > 1$ . Then there exist functorial isomorphisms

$$H_\bullet(A, \Phi^n(M)) \cong \text{Tor}_\bullet^A(\psi^n(A), M), \quad n \geq 0$$

and

$$H^\bullet(A, \Phi^n(M)) \cong \text{Ext}_A^\bullet(\psi^n(A), M), \quad n \geq 0.$$

In particular, if  $A$  is a commutative algebra over a field  $K$  of characteristic  $p > 0$  with  $\text{Card}(K) > p^n$ , then

$$H_\bullet(A, \Phi^n(M)) = 0 = H^\bullet(A, \Phi^n(M)).$$

*Proof.* Observe that  $C_\bullet(A, \Phi^n(A))$  is a complex of left  $A$ -modules. By Theorem 5 it is a free-resolution of  $\psi^n(A)$  in the category of  $A$ -modules. Hence it suffices to note that

$$C_\bullet(A, \Phi^n(M)) \cong M \otimes_A C_\bullet(A, \Phi^n(A)),$$

$$C^\bullet(A, \Phi^n(M)) \cong \text{Hom}_A(C_\bullet(A, \Phi^n(A)), M),$$

where  $C^*$  denotes the standard complex for Hochschild cohomology. The last assertion follows from Lemma 4.  $\square$

**Example 7.** It follows for instance that  $\mathbf{H}^i(A, \Phi^n(M)) = 0$ ,  $i > 0$ , provided  $M$  is an injective  $A$ -module and  $n > 1$ . In particular  $\mathbf{H}^i(A, \Phi^n(A)) = 0$  if  $A$  is a self-injective algebra. On the other hand if  $A = \mathbb{F}_p[x_1, \dots, x_d]$  then  $\mathbf{H}^i(A, \Phi^n(A)) = 0$ ,  $i \neq d$ ,  $n > 1$  and  $\mathbf{H}^d(A, \Phi^n(A)) = \psi^n(A)$ ,  $n > 1$ .

#### 4. APPLICATION TO MACLANE COHOMOLOGY

We recall the definition of MacLane (co)homology. For an associative ring  $R$  we let  $\mathbf{F}(R)$  be the category of finitely generated free left  $R$ -modules. Moreover, we let  $\mathcal{F}(R)$  be the category of all covariant functors from the category  $\mathbf{F}(R)$  to the category of all  $R$ -modules. The category  $\mathcal{F}(R)$  is an abelian category with enough projective and injective objects. By definition [8] the *MacLane cohomology* of  $R$  with coefficient in a functor  $T \in \mathcal{F}(R)$  is given by

$$\mathbf{HML}^\bullet(R, T) := \mathbf{Ext}_{\mathcal{F}(R)}^\bullet(I, T),$$

where  $I \in \mathcal{F}(R)$  is the inclusion of the category  $\mathbf{F}(R)$  into the category of all left  $R$ -modules. One defines MacLane homology in a dual manner (see [13, Proposition 3.1]). For an  $R$ - $R$ -bimodule  $B$ , one considers the functor  $B \otimes_R (-)$  as an object of the category  $\mathcal{F}(R)$ . For simplicity we write  $\mathbf{HML}_\bullet(R, B)$  instead of  $\mathbf{HML}_\bullet(R, B \otimes_R (-))$ . There is a binatural transformation

$$\mathbf{HML}_\bullet(R, B) \rightarrow \mathbf{H}_\bullet(R, B)$$

which is an isomorphism in dimensions 0 and 1.

In the rest of this section we consider MacLane (co)homology of commutative  $\mathbb{F}_p$ -algebras.

**Lemma 8.** For any commutative  $\mathbb{F}_p$ -algebra  $A$  one has an isomorphism

$$\mathbf{HML}_{2i}(A, \Phi^n(A)) = \psi^n(A), \quad i \geq 0, n > 1,$$

and

$$\mathbf{HML}_{2i+1}(A, \Phi^n(A)) = 0, \quad i \geq 0, n > 1.$$

*Proof.* According to [12, Proposition 4.1] there exists a functorial spectral sequence

$$E_{pq}^2 = \mathbf{H}_p(A, \mathbf{HML}_q(\mathbb{F}_p, B)) \implies \mathbf{HML}_{p+q}(A, B).$$

Here  $B$  is an  $A$ - $A$ -bimodule. By the well-known computation of Breen [2], Bökstedt [1] (see also [4]) we have

$$\mathbf{HML}_{2i}(\mathbb{F}_p, B) = B$$

and

$$\mathbf{HML}_{2i+1}(\mathbb{F}_p, B) = 0.$$

Now we put  $B = \psi^n(A)$  and use Theorem 5 to get  $E_{pq}^2 = 0$  for all  $p > 0$ . Hence the spectral sequence degenerates and the result follows.  $\square$

We now consider MacLane cohomology with coefficients in strict polynomial functors [6]. Let us recall that the strict homogeneous polynomial functors of degree  $d$  form an abelian category  $\mathcal{P}_d(A)$  and there exist an exact functor  $i : \mathcal{P}_d(A) \rightarrow \mathcal{F}(A)$  [7]. For an object  $T \in \mathcal{P}_d(A)$  we write  $\mathbf{HML}_\bullet(A, T)$  instead of  $\mathbf{HML}_\bullet(A, i(T))$ . Projective generators of the category  $\mathcal{P}_d$  are tensor products of the divided powers, while the injective cogenerators are symmetric powers. Let us recall that the  $d$ -th divided power functor  $\Gamma^d \in \mathcal{F}(A)$  and  $d$ -th symmetric functors  $S^n$  are defined by

$$\Gamma^d(M) = (M^{\otimes d})^{\Sigma_d}, \quad S^n(M) = (M^{\otimes d})_{\Sigma_d}.$$

Here tensor products are taken over  $A$ ,  $\Sigma_d$  is the symmetric group on  $d$ -letters, which acts on the  $d$ -th tensor power by permuting of factors,  $M \in \mathbf{F}(A)$  and  $X^G$  (resp.  $X_G$ ) denotes the module of invariants (resp. coinvariants) of a  $G$ -module  $X$ , where  $G$  is a group.

For a functor  $T \in \mathcal{F}(A)$  we let  $\tilde{T} \in \mathcal{F}(\mathbb{F}_p)$  be the functor defined by

$$\tilde{T}(V) = T(V \otimes A)$$

According to [13, Theorem 4.1] the groups  $\mathbf{HML}_i(\mathbb{F}_p, \tilde{T})$  have an  $A$ - $A$ -bimodule structure. The left action comes from the fact that  $T$  has values in the category of left  $A$ -modules, while the right action comes from the fact that  $T$  is defined on  $\mathbf{F}(A)$ . In particular it uses the action of  $T$  on the maps  $l_a : X \rightarrow X$ , where  $a \in A$ ,  $X \in \mathbf{F}(A)$  and  $l_a$  is the multiplication on  $a$ . Since  $T(l_a) = l_{a^d}$  if  $T$  is a strict homogeneous polynomial functor of degree  $d$  [6], the bimodule  $\mathbf{HML}_i(\mathbb{F}_p, \tilde{T})$  is of the form  $\Phi^n(M)$  provided  $d = p^n$ .

**Theorem 9.** Let  $d > 1$  be an integer and let  $A$  be a commutative  $\mathbb{F}_p$ -algebra. Then  $\mathbf{HML}_\bullet(A, \Gamma^d) = 0$  if  $d$  is not a power of  $p$ . If  $d = p^n$  and  $n > 0$ , then

$$\mathbf{HML}_i(A, \Gamma^d) = 0 \quad \text{if } i \neq 2p^{nt}, t \geq 0$$

and

$$\mathbf{HML}_i(A, \Gamma^d) = \psi^n(A) \quad \text{if } i = 2p^{nt}, t \geq 0.$$

In particular  $\mathbf{HML}_\bullet(A, \Gamma^d) = 0$  provided  $A$  is an algebra over a field  $K$  of characteristic  $p > 0$  with  $\text{Card}(K) > d$ .

*Proof.* According to [13, Theorem 4.1],[12] there exists a functorial spectral sequence:

$$E_{pq}^2 = \mathbf{H}_p(A, \mathbf{HML}_q(\mathbb{F}_p, \tilde{T})) \implies \mathbf{HML}_{p+q}(A, T).$$

For  $T = \Gamma_A^n$  one has  $\tilde{T} = \Gamma_{\mathbb{F}_p}^n \otimes A$ . Here we used the notation  $\Gamma_A^n$  in order to emphasize the dependence on the ring  $A$ . By the result of Franjou, Lannes and Schwartz [4]  $\mathbf{HML}_i(\mathbb{F}_p, \tilde{T})$  vanishes unless  $d = p^n$  and  $i = 2p^{nt}$ ,  $t \geq 0$ . Moreover in these exceptional cases  $\mathbf{HML}_i(\mathbb{F}_p, \tilde{T})$  equals to  $\Phi^n(A)$  (as an  $A$ - $A$ -module). Hence the spectral sequence together with Theorem 5 gives the result.  $\square$

**Corollary 10.** Let  $A$  be a commutative algebra over a field  $K$  of characteristic  $p > 0$  with  $\text{Card}(K) > d > 1$ . If  $T$  is a strong homogeneous polynomial functor of degree  $d$ . Then

$$\mathbf{HML}_\bullet(A, T) = 0 = \mathbf{HML}^\bullet(A, T).$$

*Proof.* We already proved that the result is true if  $T$  is a divided power. By the well-known vanishing result [11] the result is also true if  $T = T_1 \otimes T_2$  with  $T_1(0) = 0 = T_2(0)$ . Since any object of  $\mathcal{P}_d$  has a finite resolution which consists with finite direct sums of tensor products of divided powers [6] the result follows.  $\square$

## REFERENCES

- [1] M. Bökstedt. The topological Hochschild homology of  $\mathbb{Z}$  and  $\mathbb{Z}/p$ . Unpublished manuscript.
- [2] L. Breen. Extensions du groupe additif. Publ. IHES 48 (1978), 39–125.
- [3] L. Esakia. Heyting algebras. I. Duality theory. (Russian) “Metsniereba”, Tbilisi, 1985. 105 pp.
- [4] V. Franjou, J. Lannes and L. Schwartz. Autour de la cohomologie de MacLane des corps finis. Invent. Math. 115 (1994), no. 3, 513–538.
- [5] V. Franjou, E. Friedlander, T. Pirashvili and L. Schwartz. Rational representations, the Steenrod algebra and functor homology. Panoramas et Synthésés 16. Société Mathématique de France, Paris, 2003. xxii+132 pp.
- [6] E. Friedlander and A. Suslin. Cohomology of finite group schemes over a field. Invent. Math. 127 (1997), no. 2, 209–270
- [7] V. Franjou, E. Friedlander, A. Scorichenko and A. Suslin. General linear and functor cohomology over finite fields. Ann. of Math. (2) 150 (1999), no. 2, 663–728.
- [8] M. Jibladze and T. Pirashvili. Cohomology of algebraic theories. J. Algebra 137 (1991), no. 2, 253–296.
- [9] J.-L. Loday. Cyclic homology. Grundlehren der Mathematischen Wissenschaften, 301. Springer-Verlag, Berlin, 1998.
- [10] S. Mac Lane. Homology. Die Grundlehren der mathematischen Wissenschaften, Bd. 114, Springer, Berlin. 1963 x+422 pp.
- [11] T. Pirashvili. Higher additivizations. Trudy Tbiliss. Mat. Inst. Razmadze Akad. Nauk Gruzin. SSR 91 (1988), 44–54.
- [12] T. Pirashvili. Spectral sequennce for MacLane homology. J. of Algebra, 170(1994), 422-427.
- [13] T. Pirashvili and F. Waldhausen. MacLane homology and topological Hochschild homology. J. Pure Appl. Algebra 82 (1992), 81–98
- [14] D. Quillen. On the (co-) homology of commutative rings. Proc. Sympos. Pure Math., Vol. XVII, New York, 1968. 65–87.
- [15] S. Schwede. Stable homotopy of algebraic theories. Topology 40 (2001), no. 1, 1–41.