

A few remarks about operator theory, topology, and analysis on metric spaces

Stephen Semmes
Rice University

Abstract

Some basic facts about Fredholm indices are briefly reviewed, often used in connection with Toeplitz and pseudodifferential operators, and which may be relevant for operators associated to fractals.

Let \mathcal{H} be a complex infinite-dimensional separable Hilbert space, which is of course unique up to isomorphism. Let $\mathcal{B}(\mathcal{H})$ be the Banach algebra of bounded linear operators on \mathcal{H} , and let $\mathcal{C}(\mathcal{H})$ be the closed ideal in $\mathcal{B}(\mathcal{H})$ consisting of compact operators on \mathcal{H} . The quotient $\mathcal{A}(\mathcal{H}) = \mathcal{B}(\mathcal{H})/\mathcal{C}(\mathcal{H})$ is a C^* algebra known as the *Calkin algebra*.

A bounded linear operator T on \mathcal{H} is said to be *Fredholm* if the kernel of T is finite-dimensional and T maps \mathcal{H} onto a closed linear subspace of \mathcal{H} of finite codimension. Sometimes it is convenient to consider unbounded linear operators too, such as differential operators. It may also be helpful to include operators between different Hilbert spaces. One way to deal with unbounded linear operators is to treat them as bounded linear operators between different Hilbert spaces.

If $T \in \mathcal{B}(\mathcal{H})$ is Fredholm, then there is an $R \in \mathcal{B}(\mathcal{H})$ such that $RT - I$ and $TR - I$ are finite rank operators, where I is the identity operator on \mathcal{H} . Conversely, $T \in \mathcal{B}(\mathcal{H})$ is Fredholm if there is an $R \in \mathcal{B}(\mathcal{H})$ such that $RT - I$ and $TR - I$ are compact operators. Equivalently, $T \in \mathcal{B}(\mathcal{H})$ is Fredholm if and only if the corresponding element \hat{T} of $\mathcal{A}(\mathcal{H})$ is invertible.

The *index* of a Fredholm operator $T \in \mathcal{B}(\mathcal{H})$ is defined to be the difference between the dimension of the kernel of T and the codimension of $T(\mathcal{H})$ in \mathcal{H} . If $T, T' \in \mathcal{B}(\mathcal{H})$, T is Fredholm, and $T' - T$ is compact, then T' is a Fredholm operator with the same index as T . If $T, T' \in \mathcal{B}(\mathcal{H})$, T is Fredholm, and T' is sufficiently close to T in the operator norm topology, then T' is again Fredholm and has the same index as T . The composition of two Fredholm operators is Fredholm, and the index of the composition is equal to the sum of the indices.

Many questions in analysis are concerned with invertability of linear operators. One might start by showing that an operator is Fredholm, and then try to analyze the kernel and cokernel. These may be considered as nuisances to be minimized. By contrast, from the perspective of topology, nontrivial indices are an opportunity.

As in finite dimensions, invertible operators on \mathcal{H} form a connected open set in $\mathcal{B}(\mathcal{H})$. Fredholm operators form an open set that is not connected, with different components distinguished by the index.

The set of invertible operators on \mathcal{H} is in fact contractable [32]. The set of Fredholm operators is not, and can instead be used as a classifying space for topological K -theory [3].

Fredholm operators occur frequently in the study of integral equations, where they may be given as compact perturbations of multiples of the identity and hence have index 0 automatically. However, they may also have nontrivial kernels and cokernels, as in the context of elliptic boundary value problems.

A Toeplitz operator associated to a continuous complex-valued function on the unit circle is Fredholm when the function is nonvanishing. The index of the Toeplitz operator is equal to the winding number of the function around 0 in the complex plane.

There are very interesting ways in which to elaborate some Fredholm operators and get additional indices. One approach applies to operators whose commutators with multiplication operators by continuous functions are compact [2]. This permits the operator to be adjusted to act on spaces of sections of vector bundles. Another approach deals with homomorphisms from algebras of continuous functions on compact metric spaces into the Calkin algebra [12, 13, 14, 21]. In this case there are also operators corresponding to matrix-valued functions on the space.

Techniques like these are very important for increasing the information available from an operator or a collection of operators. A basic construction may lead to a whole family of indices.

It seems to me that there is a lot of room for development of operator theory on fractals. In particular, there are a lot of interesting fractals with topological dimension 1. Of course, many fundamental notions and results of analysis carry over to the broad setting of spaces of homogeneous type [16, 17]. Part of the point would be to take advantage of extra structure when it is present, even if it is not quite the same as the usual smooth case. There can still be significant interactions with the underlying geometry and topology.

Extensions of C^* -algebras as in [12, 13, 13, 21] are especially appealing for spaces with topological dimension 1, where the higher-dimensional cohomology groups are trivial. It is also nice to be able to view a 1-dimensional space as some sort of boundary of a 2-dimensional space. This includes fractals in the plane, and the spaces considered by Bourdon and Pajot [7, 8, 9, 10, 11], for instance. As in these examples, the 2-dimensional spaces may have singularities or other complications. At any rate, it would be interesting to have something like complex-analytic metric space involved.

References

- [1] L. Ambrosio and P. Tilli, *Topics on Analysis in Metric Spaces*, Oxford University Press, 2004.

- [2] M. Atiyah, *Global theory of elliptic operators*, in *Proceedings of the International Conference on Functional Analysis and Related Topics*, 21–30, University of Tokyo Press, 1970.
- [3] M. Atiyah, *K-Theory*, notes by D. Anderson, 2nd edition, Addison-Wesley, 1989.
- [4] W. Arveson, *A Short Course on Spectral Theory*, Springer-Verlag, 2002.
- [5] P. Auscher, T. Coulhon, and A. Grigoryan, editors, *Heat Kernels and Analysis on Manifolds, Graphs, and Metric Spaces*, American Mathematical Society, 2003.
- [6] R. Beals and P. Greiner, *Calculus on Heisenberg Manifolds*, Princeton University Press, 1988.
- [7] M. Bourdon, *Immeubles hyperboliques, dimension conforme et rigidité de Mostow*, *Geometric and Functional Analysis* **7** (1997), 245–268.
- [8] M. Bourdon and H. Pajot, *Poincaré inequalities and quasiconformal structure on the boundary of some hyperbolic buildings*, *Proceedings of the American Mathematical Society* **127** (1999), 2315–2324.
- [9] M. Bourdon and H. Pajot, *Rigidity of quasi-isometries for some hyperbolic buildings*, *Comentarii Mathematici Helvetici* **75** (2000), 701–736.
- [10] M. Bourdon and H. Pajot, *Quasi-conformal geometry and hyperbolic geometry*, in *Rigidity in Dynamics and Geometry*, 1–17, Springer-Verlag, 2002.
- [11] M. Bourdon and H. Pajot, *Cohomologie ℓ_p et espaces de Besov*, *Journal für die Reine und Angewandte Mathematik* **558** (2003), 85–108.
- [12] L. Brown, R. Douglas, and P. Fillmore, *Extensions of C^* -algebras, operators with compact self-commutators, and K -homology*, *Bulletin of the American Mathematical Society* **79** (1973), 973–978.
- [13] L. Brown, R. Douglas, and P. Fillmore, *Unitary equivalence modulo the compact operators and extensions of C^* -algebras*, in *Proceedings of a Conference on Operator Theory*, 58–128, *Lecture Notes in Mathematics* **345**, Springer-Verlag, 1973.
- [14] L. Brown, R. Douglas, and P. Fillmore, *Extensions of C^* -algebras and K -homology*, *Annals of Mathematics* (2) **105** (1977), 265–324.
- [15] J. Cheeger, *Differentiability of Lipschitz functions on metric measure spaces*, *Geometric and Functional Analysis* **9** (1999), 428–517.
- [16] R. Coifman and G. Weiss, *Analyse Harmonique Non-Commutative sur Certains Espaces Homogènes*, *Lecture Notes in Mathematics* **242**, Springer-Verlag, 1971.

- [17] R. Coifman and G. Weiss, *Extensions of Hardy spaces and their use in analysis*, Bulletin of the American Mathematical Society **83** (1977), 569–645.
- [18] A. Connes, *Noncommutative Geometry*, based in part on a translation from the French by S. Berberian, Academic Press, 1994.
- [19] M. Cotlar, *Convolution Operators and Factorization*, McGill University, 1972.
- [20] G. David and S. Semmes, *Fracture Fractals and Broken Dreams: Self-Similar Geometry through Metric and Measure*, Oxford University Press, 1997.
- [21] R. Douglas, *C^* -Algebra Extensions and K -Homology*, Princeton University Press, 1980.
- [22] R. Douglas, *Banach Algebra Techniques in Operator Theory*, 2nd edition, Springer-Verlag, 1998.
- [23] G. Folland, *Introduction to Partial Differential Equations*, 2nd edition, Princeton University Press, 1995.
- [24] G. Folland and E. Stein, *Hardy Spaces on Homogeneous Groups*, Princeton University Press, 1982.
- [25] P. Hajłasz and P. Koskela, *Sobolev Met Poincaré*, Memoirs of the American Mathematical Society **688**, 2000.
- [26] J. Heinonen, *Lectures on Analysis on Metric Spaces*, Springer-Verlag, 2001.
- [27] J. Heinonen, *Nonsmooth calculus*, Bulletin of the American Mathematical Society **44** (2007), 163–232.
- [28] W. Hurewicz and H. Wallman, *Dimension Theory*, Princeton University Press, 1941.
- [29] J. Kigami, *Analysis on Fractals*, Cambridge University Press, 2001.
- [30] S. Krantz, *A Panorama of Harmonic Analysis*, Mathematical Association of America, 1999.
- [31] S. Krantz and S.-Y. Li, *Boundedness and compactness of integral operators on spaces of homogeneous type and applications I, II*, Journal of Mathematical Analysis and Applications **258** (2001), 629–641, 642–657.
- [32] N. Kuiper, *The homotopy type of the unitary group of Hilbert space*, Topology **3** (1965), 19–30.
- [33] T. Laakso, *Ahlfors Q -regular spaces with arbitrary $Q > 1$ admitting weak Poincaré inequality*, Geometric and Functional Analysis **10** (2000), 111–123; erratum, **12** (2002), 650.

- [34] T. Laakso, *Plane with A_∞ -weighted metric not bi-Lipschitz embeddable to \mathbf{R}^N* , *Bulletin of the London Mathematical Society* **34** (2002), 667–676.
- [35] M. Lapidus, *Analysis on fractals, Laplacians on self-similar sets, noncommutative geometry and spectral dimensions*, *Topological Methods in Nonlinear Analysis* **4** (1994), 137–195.
- [36] M. Lapidus, *Towards a noncommutative fractal geometry? Laplacians and volume measures on fractals*, in *Harmonic Analysis and Nonlinear Differential Equations*, 211–252, American Mathematical Society, 1997.
- [37] N. Nikolski, *Treatise on the Shift Operator*, with an appendix by S. Hruščev and V. Peller, translated from the Russian by J. Peetre, Springer-Verlag, 1986.
- [38] N. Nikolski, *Operators, Functions, and Systems: An Easy Reading*, volume 1, *Hardy, Hankel, and Toeplitz*, volume 2, *Model Operators and Systems*, translated from the French by A. Hartmann and revised by the author, American Mathematical Society, 2002.
- [39] V. Peller, *Hankel Operators and their Applications*, Springer-Verlag, 2003.
- [40] D. Sarason, *Function Theory on the Unit Circle*, Virginia Polytechnic and State University, 1978.
- [41] S. Semmes, *Some Novel Types of Fractal Geometry*, Oxford University Press, 2001.
- [42] E. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton University Press, 1970.
- [43] E. Stein, *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, with the assistance of T. Murphy, Princeton University Press, 1993.
- [44] E. Stein and G. Weiss, *Introduction to Fourier Analysis on Euclidean Spaces*, Princeton University Press, 1971.
- [45] R. Strichartz, *Analysis on fractals*, *Notices of the American Mathematical Society* **46** (1999), 1199–1208.
- [46] R. Strichartz, *Differential Equations on Fractals: A Tutorial*, Princeton University Press, 2006.
- [47] N. Varopoulos, L. Saloff-Coste, and T. Coulhon, *Analysis and Geometry on Groups*, Cambridge University Press, 1992.
- [48] N. Weaver, *Lipschitz Algebras*, World Scientific, 1999.