

# ON FLEXIBLE QUADRATIC ALGEBRAS

S. PUMPLÜN

ABSTRACT. Let  $R$  be a ring. A construction method for flexible quadratic algebras with scalar involution over  $R$  is presented which unifies various classical constructions in the literature, in particular those to construct composition algebras.

## INTRODUCTION

Let  $R$  be a unital commutative associative ring. We present a construction method for  $R$ -algebras starting from an associative quadratic  $R$ -algebra with a scalar involution. Special cases of this construction yield flexible quadratic algebras over rings (an algebra  $A$  is *flexible* if  $x(yx) = (xy)x$  for all elements  $x, y \in A$ ). Composition algebras are certainly among the best-known classes of flexible quadratic algebras. Our method unifies the different approaches by Petersson [P2], Thakur [T] and [Pu] to construct composition algebras. In special cases it overlaps with the one described by Becker [B] which yields flexible quadratic division algebras containing subalgebras of half their rank.

## 1. PRELIMINARIES

Let  $A$  be an  $R$ -algebra. The term “ $R$ -algebra” always refers to a unital nonassociative algebra  $A$  which is faithfully projective as an  $R$ -module [K, p. 52].

$A$  is called *alternative* if its associator  $[x, y, z] = (xy)z - x(yz)$  is alternating. The *nucleus* of  $A$  is defined as  $N(A) = \{x \in A \mid [x, A, A] = [A, x, A] = [A, A, x] = 0\}$ . The nucleus is an associative subalgebra of  $A$  (it may be zero), and  $x(yz) = (xy)z$  whenever one of the elements  $x, y, z$  is in  $N(A)$ . An anti-automorphism  $\sigma : A \rightarrow A$  of period 2 is called an *involution* on  $A$ . If 2 is an invertible element in  $R$ , we have  $A = \text{Sym}(A, \sigma) \oplus \text{Skew}(A, \sigma)$  with  $\text{Skew}(A, \sigma) = \{x \in A \mid \sigma(x) = -x\}$  the set of skew symmetric elements

---

2000 *Mathematics Subject Classification*. Primary: 17A45; Secondary: 17A20, 17A75, 11E25.

*Key words and phrases*. Flexible quadratic algebras, composition algebras, cross products.

and  $\text{Sym}(A, \sigma) = \{x \in A \mid \sigma(x) = x\}$  the set of symmetric elements in  $A$  with respect to  $\sigma$ . An involution is called *scalar* if all *norms*  $\sigma(x)x$  are elements of  $R$ . For every scalar involution  $\sigma$ ,  $n_A(x) = \sigma(x)x$  (resp. the *trace*  $t_A(x) = \sigma(x) + x$ ) is a quadratic (resp. a linear) form on  $A$ .  $A$  is called *quadratic*, if there exists a quadratic form  $n: A \rightarrow R$  such that  $n(1_A) = 1$  and  $x^2 - n(1_A, x)x + n(x)1_A = 0$  for all  $x \in A$ , where  $n(x, y)$  denotes the induced symmetric bilinear form  $n(x, y) = n(x+y) - n(x) - n(y)$ . The form  $n$  is uniquely determined and called the *norm* of the quadratic algebra  $A$  [P2]. The existence of a scalar involution on an algebra  $A$  implies that  $A$  is quadratic [M1].

An  $R$ -algebra  $C$  is called a *composition algebra* if it carries a quadratic form  $n: C \rightarrow R$  whose induced symmetric bilinear form  $n(x, y)$  is *nondegenerate* (i.e., determines an  $R$ -module isomorphism  $C \xrightarrow{\sim} C^\vee = \text{Hom}_R(C, R)$ ) and which satisfies  $n(xy) = n(x)n(y)$  for all  $x, y \in C$  (it *permits composition*). Composition algebras are quadratic alternative; any nondegenerate quadratic form  $n$  on the composition algebra which permits composition is its norm as a quadratic algebra and thus is unique [P2]. It is called the *norm* of  $C$  and is also denoted by  $n_C$ . A quadratic alternative algebra is a composition algebra if and only if its norm is nondegenerate [M1, 4.6]. Composition algebras only exist in ranks 1, 2, 4 or 8. Those of rank 2 are exactly the quadratic étale  $R$ -algebras, those of rank 4 exactly the well-known quaternion algebras. The ones of rank 8 are called *octonion algebras*. A composition algebra  $C$  has a *canonical involution*  $\bar{\phantom{x}}$  given by  $\bar{x} = t_C(x)1_C - x$ , where  $t_C: C \rightarrow R$ ,  $t_C(x) = n(1_C, x)$ , is the *trace* of  $C$ . This involution is scalar.

## 2. FLEXIBLE QUADRATIC ALGEBRAS

Let  $N$  be a finitely generated projective  $R$ -module of constant rank. An alternating  $R$ -bilinear map  $\times: N \times N \rightarrow N$  is called a *cross product* on  $N$ . Let  $D$  be an associative  $R$ -algebra with a scalar involution  $\sigma = \bar{\phantom{x}}$  (e.g. a quadratic étale or a quaternion algebra over  $R$  with canonical involution). Let  $F$  be a locally free right  $D$ -module of constant finite rank  $s$  together with a sesquilinear form  $h: F \times F \rightarrow D$  (i.e.,  $h$  is a biadditive map such that  $h(ua, vb) = \bar{a}h(u, v)b$  for all  $a, b \in D$ ,  $u, v \in F$ ). Let  $\times$  be a cross product on  $F$ , where now  $F$  is viewed as an  $R$ -module, i.e. together with  $\times: F \times F \rightarrow F$ ,  $F$  is an anticommutative  $R$ -algebra. The  $R$ -module  $A = D \oplus F$  becomes a unital  $R$ -algebra denoted by  $A = (D, F, h, \times)$  via the multiplication

$$(\alpha, u)(\beta, v) = (\alpha\beta - h(v, u), v\alpha + u\bar{\beta} + v \times u),$$

for all  $\alpha, \beta \in D$ ,  $u, v \in F$ . The algebra  $D$  is a subalgebra of  $(D, F, h, \times)$ .

Define

$$\begin{aligned}\sigma_A : A &\rightarrow A, & (\alpha, u) &\rightarrow (\bar{\alpha}, -u), \\ n_A : A &\rightarrow D, & n_A((\alpha, u)) &= \sigma_A(\alpha, u)(\alpha, u), \\ t_A : A &\rightarrow R, & t_A((\alpha, u)) &= \sigma_A(\alpha, u) + (\alpha, u) = (t_D(\alpha), 0).\end{aligned}$$

Obviously,  $\sigma_A$  is an  $R$ -module isomorphism of order 2. We have

$$\begin{aligned}n_A((\alpha, u)) &= n_D(\alpha) + h(u, u), \\ n_A((\alpha, u)) &= (\alpha, u)\sigma_A(\alpha, u), \\ n_A((\sigma(\alpha, u))) &= n_A((\alpha, u)) \text{ and } t_A(\sigma(\alpha, u)) = t_A((\alpha, u)).\end{aligned}$$

The map  $n_A$  is quadratic with associated  $R$ -bilinear map  $n_A : A \times A \rightarrow D$  given by

$$\begin{aligned}n_A((\alpha, u), (\beta, v)) &= n_D(\alpha, \beta) + (h(u, v) + h(v, u)) = \\ &(\alpha, u)\sigma_A(\beta, v) + (\beta, v)\sigma_A(\alpha, u).\end{aligned}$$

Obviously,  $F = \ker(t_A) = \ker(t_D) \oplus F$  and  $u \times v = uv - \frac{1}{2}n_A(u, v)$ .

**Lemma 1.** (i)  $\sigma_A$  is a scalar involution if and only if  $h$  is a hermitian form (i.e.,  $h(u, v) = \overline{h(v, u)}$  for all  $u, v \in F$ ).

(ii) If  $h$  is a hermitian form, then  $t : A \times A \rightarrow R$ ,  $t(x, y) = t_A(xy)$  is a symmetric  $R$ -bilinear form.

Lemma 1 is a weak generalization of [B, Lemma 1].

**Corollary 1.** If  $h : F \times F \rightarrow D$  is a (perhaps degenerate) hermitian form, then  $A = (D, F, h, \times)$  is a quadratic  $R$ -algebra with scalar involution  $\sigma_A : A \rightarrow A$ ,  $\sigma_A((\alpha, u)) = (\bar{\alpha}, -u)$  and norm  $n_A : A \rightarrow R$ ,  $x \rightarrow x\sigma_A(x)$ , where  $n_A((\alpha, u)) = n_D(\alpha) + h(u, u)$ . Moreover, if  $n_A$  is isotropic (e.g., if  $h$  is isotropic) then  $A$  has zero divisors.

Note that even if  $D$  is an associative composition division algebra over a field, the fact that two quadratic algebras  $A = (D, F, h, \times)$  and  $A' = (D, F', h', \times')$  have isometric norms  $n_A \cong n_{A'}$  only implies that the non-degenerate hermitian forms  $(F, h)$  and  $(F', h')$  are isometric and yields no information on the cross products  $\times$  and  $\times'$ .

Quadratic  $R$ -algebras with a scalar involution were constructed using finite dimensional vector spaces which carry a bilinear form  $B$  by Osborn [O]. Finite-dimensional quadratic division algebras over a field of characteristic not 2 were shown to be even-dimensional by Petersson [P1]. Already earlier, Kuz'min [Ku] proved that any quadratic division algebra, over a field of

characteristic not 2 in which any two elements generate a subalgebra of dimension at most four, must have finite dimension and that, indeed, only the dimensions 1, 2, 4 or 8 are possible. Independently, Kunze and Scheinberg [KS, Theorem 2.4] enumerated all unital alternative algebras over fields of characteristic not 2 - including the infinite-dimensional ones - which have a scalar involution.

**Remark 1.** For  $A = (D, F, h, \times)$ ,  $2 \in R^\times$ , we obtain  $\text{Sym}(A, \sigma_A) = R$  and  $\text{Skew}(A, \sigma_A) = \text{Skew}(D, \sigma_D) \oplus F$ .

**Lemma 2.** *If  $h : F \times F \rightarrow D$  is a hermitian form, then  $(D, F, h, \times)$  is*  
 (i) *flexible if and only if  $h(u \times v, u) + \overline{h(u \times v, u)} = n_A(u \times v, u) = 0$  and  $(u \times v) \times u = u \times (v \times u)$  for all  $u, v \in F$ ;*  
 (ii) *alternative if and only if  $h(u, u \times v) = 0$  and  $u \times (u \times v) = -h(u, u)v + h(v, u)u$  for all  $u, v \in F$ , if and only if  $h(u \times v, v) = 0$  and  $(u \times v) \times v = h(v, v)u - h(u, v)v$  for all  $u, v \in F$ .*

If  $(D, F, h, \times)$  is flexible with a scalar involution, then it is a noncommutative Jordan algebra [M1, (3.3)]. Due to the nature of the multiplication in  $(D, F, h, \times)$  it suffices to work with elements of  $F$  to check flexibility. The identities in (ii) were already obtained in [T, (2.1)] in a more restricted setting (there,  $F$  has rank 3,  $D$  is quadratic étale, and  $h$  is a nondegenerate hermitian form). If  $\times$  is the zero-map, then  $(D, F, h, 0)$  is trivially flexible.

**Remark 2.** If  $D = R$ , then  $\sigma = id$  and  $h = B$  is a symmetric  $R$ -bilinear form on  $F$ .  $A = (R, F, B, \times)$  is the unital quadratic  $R$ -algebra described in [B, p. 26] (see also [O, Theorem 1]). It is well-known that any quadratic algebra  $A$  over a field of characteristic not 2 with a scalar involution can be of the type  $A = (R, F, B, \times)$  for a suitable  $F$ , a symmetric bilinear form  $B$  and a cross product  $\times$ .

(i) By Lemma 2 (i),  $A$  is flexible if and only if  $B(u, v \times u) = B(u, u \times v)$  and  $(u \times v) \times u = u \times (v \times u)$  for all  $u, v \in F$ , which is equivalent to

$$B(u, v \times u) = 0 \quad \text{and} \quad (u \times v) \times u = u \times (v \times u)$$

for all  $u, v \in F$ .

(ii) Let  $2 \in R^\times$ . If we apply [B, Lemma 2, Satz 1] to the residue class algebras, we see that  $A$  is flexible if and only if  $t_A$  is associative, if and only if  $B$  is associative; and that  $A$  is flexible and  $t_A$  and  $n_A$  are weakly nondegenerate if and only if  $B$  is associative and weakly nondegenerate. ( $q$  is *weakly nondegenerate* if  $\text{Rad}(q) = \{a \in A \mid q(a, A) = 0\} = 0$ .)

(iii) For a flexible quadratic  $R$ -algebra  $A = (R, F, B, \times)$  of rank greater or equal to 3 with a weakly nondegenerate associative bilinear form  $B$ , the residue class algebra  $A(P)$  is central-simple for each  $P \in \text{Spec}(R)$ . This follows from [B, Satz 1, Satz 2] applied to the residue class algebras. If  $A$  is a quadratic algebra of rank 3 over a domain  $R$  and  $B$  weakly nondegenerate and associative, then  $A$  is commutative. This follows from [B, p. 33].

**Theorem 1.** *Let  $2 \in R^\times$ .*

(i) *Let  $F$  be a finitely generated projective  $R$ -module of constant rank  $s$ . Let  $B : F \times F \rightarrow R$  be a nondegenerate symmetric bilinear form on  $F$  and  $\times : F \times F \rightarrow F$  a cross product, such that*

$$(1) \quad B(u \times v, u \times v) = B(u, u)B(v, v) - B(u, v)^2$$

$$(2) \quad B(u, u \times v) = 0$$

*for all  $u, v \in F$ . Then  $C = (R, F, B, \times)$  is a composition algebra over  $R$ . In particular  $s = 1, s = 3$ , or  $s = 7$ , and  $\det(F, N) = \bigwedge^s(F, N) = R$ .*

(ii) *Let  $C$  be a composition algebra over  $R$  with norm  $n$  of rank  $s + 1 \geq 1$ . Then  $C = R \oplus F$  with  $F = \text{Skew}(C, \bar{\phantom{x}})$ . The  $R$ -module  $F$  satisfies  $\det(F) = \bigwedge^s(F) = R$  with  $s = 1, 3$  or  $7$ . Define  $\times : F \times F \rightarrow F$  via  $u \times v = \frac{1}{2}[u, v] = \frac{1}{2}(uv - vu)$  and  $B : F \times F \rightarrow R$  via  $B(u, v) = -\frac{1}{2}(uv + vu) = \frac{1}{2}(u\bar{v} + v\bar{u}) = \frac{1}{2}n(u, v)$ . Then  $\times$  is a cross product and  $B$  is a nondegenerate symmetric bilinear form on  $F$  satisfying equations (1) and (2) in (i) for all  $u, v \in F$ .*

This is well-known if  $R$  is a field. Lemma 2 (ii) implies the first part of Theorem 1. Otherwise, the proof of [Mo, Theorem 10] can be used verbatim. Note that the intrinsic value of this theorem is impaired by the fact that producing a cross product on a finitely generated projective  $R$ -module  $F$  of rank 7 satisfying (1) and (2) is probably not any easier than producing an octonion algebra structure on  $R \oplus F$ .

If, in (ii),  $C$  is quadratic étale, then there exists a line bundle  $L$  over  $R$  and a nondegenerate quadratic form  $N_L : L \times L \rightarrow R$  such that  $C \cong \text{Cay}(R, L, N_L)$ . Since every bilinear map  $L \times L \rightarrow L$  is symmetric, this forces the cross product on  $L$  to be zero.

**Example 1.** Let  $A$  be a quadratic  $R$ -algebra of rank 2. Then  $A$  is commutative associative and the conjugation is a scalar involution (see [K, I.(1.3.6)] or check it locally). We get  $A = (R, L, B, \times)$  for a suitable  $L \in \text{Pic } R$ ,

$B : L \times L \rightarrow R$  a bilinear form.  $A$  is commutative, therefore  $\times$  is the zero-map, and  $A = \text{Cay}(R, L, N)$  with  $N : L \rightarrow R$  the - perhaps degenerate - quadratic form associated to  $B$ .

**Lemma 3.** *Let  $2 \in R^\times$ . Let  $A$  be a flexible quadratic  $R$ -algebra with a scalar involution  $\sigma$  which contains a composition subalgebra  $D$  of rank 2 and no other associative composition subalgebra of larger rank, or a composition subalgebra  $D$  of rank 4. Let  $F = D^\perp$ . Define  $-h$  to be the orthogonal projection of the multiplication  $vu$  for  $u, v \in F$  onto  $D$  and  $\times$  to be the orthogonal projection of the multiplication  $vu$  for  $u, v \in F$  onto  $F$ . Then:*

- (i)  $\times : F \times F \rightarrow F$  is a cross product.
- (ii)  $n_A(u, v) = t_D(h(u, v))$  and  $h(u, v) = \overline{h(v, u)}$ .

*Proof.* Let  $A$  be a quadratic  $R$ -algebra with a scalar involution  $\sigma$ , which contains a composition subalgebra  $D$  of rank 2 or 4 as subalgebra of maximal rank with nondegenerate norm. Since the norm  $n_A(x) = x\sigma(x)$  is nondegenerate when restricted to  $D$ , we can decompose  $A$  as  $A = D \perp D^\perp$  [K, (3.6.2), p. 17]. Since  $-h$  and  $\times$  are the orthogonal projections of the multiplication  $vu$  for  $u, v \in F$  onto  $D$  and onto  $F$ , we have  $vu = -h(u, v) + u \times v$  for  $u, v \in F$ .

(i) Since  $F \subset \text{Skew}(A, \sigma)$ , it follows that  $u^2 \in R$  for all  $u \in F$  and thus  $u \times u = u^2 + h(u, u) \in D \cap D^\perp = \{0\}$ . Hence  $\times$  is anticommutative and a cross product on  $F$ .

(ii) We have  $n_A(u) = -u^2 = h(u, u)$  for all  $u \in F$ . Since  $u \times v = -v \times u$ , it follows that  $h(u, v) + \overline{h(u, v)} = -vu - uv = v\sigma(u) + u\sigma(v) = n_A(u, v)$ , i.e.  $n_A(u, v) = t_D(h(u, v))$  and  $h(u, v) = \overline{h(v, u)}$ .  $\square$

**Example 2.** (a) Let  $2 \in R^\times$ . Let  $A = (D, F, h, \times)$  be an algebra with a scalar involution,  $D$  quadratic étale or a quaternion algebra. The algebra  $\hat{A} = (R, F, B, \times)$ ,  $B(u, v) = \frac{1}{2}(h(u, v) + \overline{h(u, v)}) = \frac{1}{2}n_A(u, v)$ , is flexible if and only if  $A$  is flexible. If  $A$  is alternative, the identity  $((u \times v) \times v) \times v = n_A(v)u \times v$  holds in  $\hat{A}$  for all  $u, v \in F$  (Lemma 2). In particular, if  $A = \text{Cay}(T, F, h, \times_\alpha)$  is an octonion algebra over a field  $R$  of characteristic not 2, then  $\hat{A} = (R, F, B, \times_\alpha)$  is the colour algebra defined in [E].

(b) If the cross product is trivial,  $A = (D, F, h, 0)$  with  $S$  a quadratic étale  $R$ -algebra or a quaternion algebra over  $R$ , is a unital noncommutative Jordan algebra. In particular,  $(R, F, -B, 0) = \text{JSpin}(F, B)$  is the well-known algebra of the Spin Factors [M2, p. 178].

(c) Let  $k$  be a field of characteristic not 2. Given a quadratic  $k$ -algebra  $A$

with scalar involution  $*$ , the map  $(a, b)^* = (a^*, -b)$  defines a scalar involution on the quadratic algebra  $C(A, \cdot_1, \cdot_2, \cdot_3) = A \oplus A$  with multiplication  $(a, b)(c, d) = (ac + d^* \cdot_1 b, d \cdot_2 a + b \cdot_2 c^* + b \cdot_3 d)$ , where the maps  $\cdot_i$  are certain bilinear products on  $A$  [B]. Using this construction all flexible quadratic division algebras of dimension  $2m$  over  $k$  which contain a unital subalgebra of dimension  $m$  can be obtained [B, p. 41]. The construction generalizes to quadratic algebras with scalar involutions over arbitrary rings where 2 is invertible. However, it will not produce all such algebras. If a flexible quadratic algebra over a field contains a composition subalgebra of dimension 2 or 4, the method coincides with ours whenever the algebra has twice its dimension, i.e. dimension 4, respectively 8. In that case  $d^* \cdot_1 b$  is given by the hermitian form  $h(d, b)$ ,  $\cdot_2$  is the usual algebra multiplication of  $A$ , and  $\cdot_3$  is our cross product  $\times$ .

Let  $S$  be a quadratic étale  $R$ -algebra. For the remainder of this section, we restrict our attention to the special case that  $F$  is a finitely generated projective right  $S$ -module of constant rank carrying a nondegenerate hermitian form  $h: F \times F \rightarrow S$ . For each invertible  $\mu \in S$ , the form  $\mu h: F \times F \rightarrow S$  is a nondegenerate  $\epsilon$ -hermitian form, with  $\epsilon = \bar{\mu}/\mu$ . In particular,  $\epsilon \neq 1$  if and only if  $\mu \notin R$ , and  $\epsilon = -1$  if and only if  $\mu \in \text{Skew}(S, \bar{\phantom{x}})$ .

Recall that conversely, the study of  $\epsilon$ -hermitian forms over  $S$  can be reduced to the study of hermitian ones by scaling in several important cases: Let  $\mu \in S^\times$  such that  $\frac{\bar{\mu}}{\mu} = \epsilon$ . Then  $\mu h$  is a hermitian form, for any  $\epsilon$ -hermitian form  $h$ . Such a  $\mu$  exists if  $H^1(\mathbb{Z}/2\mathbb{Z}, S^\times) = 0$  (Hilbert's Theorem 90), e.g. if  $\text{Pic } R = 0$  [K, p. 300].

**Proposition 1.** *Let  $A = (S, F, \mu h, \times)$  for some nondegenerate hermitian form  $h: F \times F \rightarrow S$ , and an invertible scalar  $\mu \in S \setminus R$ . If  $S$  is a domain (or if  $F$  is torsion-free and  $\bar{\mu} - \mu$  not a zero divisor in  $S$ ), then  $N(A) = S$ .*

*Proof.* A straightforward calculation shows that  $S \subset N(A)$ .

Now let  $(e, e') \in N(A)$  with  $e \in S$  and  $e' \in F$ . We have to show that this implies  $e' = 0$ . The equation

$$(e, e')((u, w)(u', w')) = ((e, e')(u, w))(u', w')$$

implies that

$$e' \bar{\mu} h(w, w') = w' \mu h(w, e')$$

for all  $w, w' \in F$  and thus (put  $w' = e'$ )

$$e'(\bar{\mu} - \mu)h(w, e') = 0$$

for all  $w \in F$ . If  $h(w, e') = 0$  for all  $w \in F$  then  $e' = 0$  since  $h$  is nondegenerate and thus  $(e, e') = (e, 0) \in S$ . Otherwise, there is one  $w \in F$  such that  $h(w, e') \neq 0$ . Then if  $S$  has no zero divisors, i.e., is a domain,  $F$  is torsion free and we have  $e' = 0$  since  $\bar{\mu} - \mu \neq 0$  here. The same holds if  $F$  is torsion free and  $\bar{\mu} - \mu$  not a zero divisor in  $S$ .  $\square$

### 3. CONSTRUCTIONS OF COMPOSITION ALGEBRAS

Petersson [P2] introduced the following *generalized Cayley-Dickson doubling process* for composition algebras over rings: Let  $D$  be a composition algebra of rank  $\leq 4$  over  $R$  with canonical involution  $\bar{\phantom{x}}$ . Let  $P$  be a locally free right  $D$ -module of rank one and norm one (cf. [P2, 2.3] for the definition; if  $D$  is a quaternion algebra, of norm one means that the reduced norm functor defined in [KOS] associates to  $P$  the free line bundle  $R$ ). Thus there exists a nondegenerate quadratic form  $N: P \rightarrow R$  satisfying  $N(w \cdot u) = N(w)N_D(u)$  for all elements  $w \in P$ ,  $u \in D$ , where  $\cdot$  denotes the right  $D$ -module structure of  $P$  (for  $D$  a quaternion algebra, see also [K, p. 161 ff]).  $N$  is uniquely determined up to an invertible factor in  $R^\times$  and determines a unique  $R$ -bilinear map  $P \times P \rightarrow D$ , written multiplicatively and satisfying  $(w \cdot u)(w \cdot v) = N(w)\bar{v}u$  for all  $u, v \in D$ ,  $w \in P$ . The  $R$ -module  $\text{Cay}(D, P, N) = D \oplus P$  becomes a composition algebra under the multiplication

$$(u, w)(u', w') = (uu' + ww', w' \cdot u + w \cdot \bar{u}')$$

for  $u, u' \in D$ ,  $w, w' \in P$ , with  $\cdot$  denoting the right  $D$ -module structure of  $P$ . Its norm is given by

$$n_{\text{Cay}(D, P, N)}((u, w)) = n_D(u) - N(w).$$

If  $F$  a locally free right  $D$ -module of rank 1 and  $h: F \times F \rightarrow D$  a nondegenerate hermitian form, then

$$(D, F, h, 0) = \text{Cay}(D, F, -N)$$

where  $N(u) = h(u, u)$  for all  $u \in F$ .

Let  $S$  be a quadratic étale  $R$ -algebra with canonical involution  $\bar{\phantom{x}}$ . Let  $(P, h)$  be a ternary nondegenerate  $\bar{\phantom{x}}$ -hermitian space such that  $\bigwedge^3(P, h) \cong \langle 1 \rangle$ . Choose an isomorphism  $\alpha: \bigwedge^3(P, h) \rightarrow \langle 1 \rangle$ . Define a cross product  $\times_\alpha: P \times P \rightarrow P$  as in [T, p. 5122]. Note that  $\times_\alpha$  can be derived solely from  $T, P, h$  and  $\alpha$ . The  $R$ -module  $\text{Cay}(S, P, h, \alpha) = S \oplus P$  becomes an octonion



algebra under the multiplication

$$(a, u)(b, v) = (ab - h(v, u), va + u\bar{b} + u \times_\alpha v)$$

for  $u, v \in P$  and  $a, b \in S$ , with norm

$$n((a, u)) = n_S(a) + h(u, u).$$

If the ternary hermitian space  $(P, h)$  is orthogonally decomposable this construction is independent of the choice of the isomorphism  $\alpha$ . Any octonion algebra which contains a quadratic étale subalgebra can be constructed like this (Pettersson-Racine [PR, 3.8] or Thakur [T]). We obtain

$$(S, P, h, \times_\alpha) = \text{Cay}(S, P, h, \alpha).$$

Let  $2 \in R^\times$ . There exists a general construction for quaternion algebras over  $R$  [Pu]: Let  $F$  be a finitely generated projective  $R$ -module of constant rank 3 such that  $\det F = \bigwedge^3 F \cong R$ , and  $N: F \rightarrow R$  a nondegenerate quadratic form, i.e., a selfdual isomorphism  $\varphi_N: F \xrightarrow{\sim} \check{F}$ ,  $x \rightarrow N(x, \_)$ . (Any selfdual isomorphism  $\varphi$  from  $F$  to its dual bundle  $\check{F}$  canonically induces a nondegenerate quadratic form on  $F$  via  $N_\varphi(x) := \frac{1}{2}\langle x, \varphi(x) \rangle$  where  $\langle \_, \_ \rangle: F \times \check{F} \rightarrow R$  is the canonical pairing, cf. [P2, 3.2].) Assume that  $N$  has trivial determinant. Choose an isomorphism  $\alpha: \det F \xrightarrow{\sim} R$ , then  $\alpha$  is unique up to an invertible factor in  $R$  and induces a bilinear map

$$F \times F \rightarrow \check{F}$$

$$(u, v) \rightarrow u \times_\alpha v := \alpha(u \wedge v \wedge \_)$$

as well as an isomorphism  $\beta: \det \check{F} \xrightarrow{\sim} R$  given by the condition

$$\alpha(u_1 \wedge u_2 \wedge u) \beta(\check{u}_1 \wedge \check{u}_2 \wedge \check{u}_3) = \det(\langle u_i, \check{u}_j \rangle)$$

for  $u_i$  in  $F$ ,  $\check{u}_j$  in  $\check{F}$ ,  $1 \leq i, j \leq 3$ , where again  $\langle \_, \_ \rangle: F \times \check{F} \rightarrow R$  is the canonical pairing. The  $R$ -module

$$\text{Quat}(F, N) = R \oplus F$$

becomes a quaternion algebra over  $R$  under the multiplication

$$(a, u)(b, v) = (ab - \frac{1}{2}N(u, v), av + bu + \frac{1}{2}\varphi_N^{-1}(u \times_\alpha v))$$

for all  $a, b$  in  $R$ ,  $u, v$  in  $F$ , with norm

$$N_{\text{Quat}(F, N)}((a, u)) = a^2 + N(u).$$

Every quaternion algebra over  $R$  is of the type  $\text{Quat}(F, N)$ . We obtain

$$\text{Quat}(F, N) = (R, F, B, \times_\alpha),$$

where  $B$  is the nondegenerate symmetric bilinear form on  $F$  associated with  $N$ .

We have thus unified the three constructions known for composition algebras.

#### REFERENCES

- [B] Becker, E., *Über eine Klasse flexibler quadratischer Divisionsalgebren*, J. für Reine u. Angew. Math. 256 (1972), 25-57.
- [E] Elduque, A., *Colour algebras and Cayley-Dickson algebras*, Proc. of the Royal Soc. of Edinburgh, 125A (1995), 1287-1303.
- [K] Knus, M. A., "Quadratic and Hermitian Forms", Springer Verlag, 1991.
- [KOS] Knus, M. A., Ojanguren, M., Sridharan, R., *Quadratic forms and Azumaya algebras*, J. Reine Angew. Math. 303/304 (1978), 231-248.
- [KS] Kunze, R. A., Scheinberg, S., *Alternative algebras having scalar involution*, Pac. J. Math. 124 (1986), 159-172.
- [Ku] Kuz'min, E. N., *A class of anticommutative algebras*, Algebra i Logika 6 (4) (1967), 31-50.
- [M1] McCrimmon, K., *Nonassociative algebras with scalar involution*, Pacific J. of Math. 116(1) (1985), 85-108.
- [M2] McCrimmon, K., "A Taste of Jordan Algebras", Universitext, Springer Verlag, New York 2004.
- [Mo] Morandi, P., *Lie algebras, composition algebras, and the existence of cross products on finite-dimensional vector spaces*, Expo Math. 17 (1999), 063-074.
- [O] Osborn, J. M., *Quadratic division algebras*, Trans. Amer. Math. Soc. 105 (1962), 202-221.
- [P1] Petersson, H.P., *Eine Bemerkung zu quadratischen Divisionsalgebren*, Arch. Math. 22 (1971), 59-61.
- [P2] Petersson, H. P., *Composition algebras over algebraic curves of genus 0*, Trans. Amer. Math. Soc. 337 (1993), 473-491.
- [PR] Petersson, H. P., Racine, M., *Reduced models of Albert algebras*, Math. Z. 223 (3) (1996), 367-385.
- [Pu] Pumplün, S., *Quaternion algebras over elliptic curves*, Comm. Algebra 26 (12) (1998), 4357-4373.
- [T] Thakur, M. L., *Cayley algebra bundles on  $\mathbb{A}_K^2$  revisited*, Comm. Algebra 23(13) (1995), 5119-5130.

SCHOOL OF MATHEMATICS, UNIVERSITY OF NOTTINGHAM, UNIVERSITY PARK, NOTTINGHAM NG7 2RD, UNITED KINGDOM, SUSANNE.PUMPLUEN@NOTTINGHAM.AC.UK