

# BATALIN-VILKOVISKY LIE ALGEBRA STRUCTURE ON THE LOOP HOMOLOGY OF COMPLEX STIEFEL MANIFOLDS

HIROTAKA TAMANOI

University of California, Santa Cruz

ABSTRACT. We determine the Batalin-Vilkovisky Lie algebra structure for the integral loop homology of special unitary groups and complex Stiefel manifolds. It is shown to coincide with the Poisson algebra structure associated to a certain odd symplectic form on a super vector space for which loop homology is the super algebra of functions. Over rationals, the loop homology of the above spaces splits into a tensor product of simple BV algebras, and it is shown to contain the super Lie algebra  $\pi sp$ .

## CONTENTS

1. Introduction .....	1
2. Intersection and Pontrjagin products in the homology of $SU(n+1)$ .....	4
3. Homology $S^1$ -action on the Hopf algebra $H_*(LSU(n+1))$ .....	7
4. BV Lie algebra structure on the loop homology $\mathbb{H}_*(LSU(n+1))$ .....	10
5. BV-algebra structure on the loop homology of complex Stiefel manifolds ...	14
References .....	16

## §1 Introduction

The homology of the free loop space  $LM$  of an oriented closed manifold  $M$  was shown to admit a structure of a Batalin-Vilkovisky (BV) algebra in which an associative algebra structure, a Lie algebra structure and a circle action are amalgamated into a single coherent algebraic structure [1]. Namely, the Lie bracket appears as the deviation of homology  $S^1$ -action from being a derivation with respect to the loop product which defines the associative algebra structure. The loop product structure in loop homology has been calculated for some spaces including spheres and complex projective spaces [3]. However, the nature of the Lie algebra structure on loop homology has not been well understood, except that one of the defining identities of BV algebras says that the Lie bracket acts as a derivation on the underlying associative algebra. The purpose of this paper is to describe the full BV algebra structure on the loop homology of complex Stiefel manifolds. For a brief description of a general BV algebra, see the beginning of section 4.

Since complex Stiefel manifolds are homogeneous spaces of special unitary groups, the description of BV Lie algebra structure on the loop homology of the Lie group  $SU(n+1)$  is fundamental for our purpose, and the results for Stiefel manifolds follow from it. In general, for a Lie group  $G$ , the associative algebra structure of its loop homology is given by a tensor product  $\mathbb{H}_*(LG) \cong \mathbb{H}_*(G) \otimes H_*(\Omega G)$ , where  $\mathbb{H}_*(G) \cong H^{-*}(G)$  is the intersection homology ring with degree shift so that the fundamental class has degree 0, and  $\Omega G$  is the group of based loops. Although the associative

---

1991 *Mathematics Subject Classification.* 55P35.

*Key words and phrases.* Batalin-Vilkovisky algebras, complex Stiefel manifold, free loop spaces, Hamiltonian derivation, Hopf algebra, loop homology, Poisson bracket, super Lie algebra, symplectic form,.

algebra structure is the obvious one, the BV Lie algebra structure on it is highly nontrivial because the circle action on the loop group  $LG$  mixes the homology of  $G$  and the homology of  $\Omega G$ . For the case of  $G = \mathrm{SU}(n+1)$ , we show that the BV Lie bracket coincides with the Poisson bracket associated to an odd symplectic form on a super (vector) space of dimension  $n|n$  whose super algebra of functions is the loop homology  $\mathbb{H}_*(\mathrm{LSU}(n+1))$ . Similar description is valid for the loop homology of all complex Stiefel manifolds.

To describe the result, the associative algebra structure of the integral loop homology is given by

$$(1-1) \quad \mathbb{H}_*(\mathrm{LSU}(n+1)) \cong \Lambda_{\mathbb{Z}}(\alpha_3, \alpha_5, \dots, \alpha_{2n+1}) \otimes \mathbb{Z}[e_2, e_4, \dots, e_{2n}],$$

where  $\mathbb{H}_*(\mathrm{SU}(n+1)) \cong \Lambda_{\mathbb{Z}}(\alpha_3, \alpha_5, \dots, \alpha_{2n+1})$  and  $H_*(\Omega\mathrm{SU}(n+1)) \cong \mathbb{Z}[e_2, e_4, \dots, e_{2n}]$  are also coalgebras whose generators have degree  $|\alpha_{2k+1}| = -2k - 1$ ,  $|e_{2k}| = 2k$  for  $1 \leq k \leq n$ . For sequences  $I = (2i_1 + 1, 2i_2 + 1, \dots, 2i_r + 1)$  with  $1 \leq i_1 < i_2 < \dots < i_r \leq n$ , and  $J = (j_1, j_2, \dots, j_n)$  with nonnegative integer entries, let  $\alpha_I = \alpha_{2i_1+1} \alpha_{2i_2+1} \dots \alpha_{2i_r+1}$  and  $e^J = e_2^{j_1} e_4^{j_2} \dots e_{2n}^{j_n}$ . We set  $|I| = r$ , the length of  $I$ .

We define a differential operator  $D_{2\ell}$  acting on the polynomial ring  $\mathbb{Z}[e_2, e_4, \dots, e_{2n}]$  by

$$(1-2) \quad D_{2\ell} = \sum_{k=\ell}^n e_{2k-2\ell} \frac{\partial}{\partial e_{2k}} = \frac{\partial}{\partial e_{2\ell}} + e_2 \frac{\partial}{\partial e_{2\ell+2}} + \dots + e_{2n-2\ell} \frac{\partial}{\partial e_{2n}}, \quad 1 \leq \ell \leq n,$$

where we set  $e_0 = 1$ . These  $n$  differential operators mutually commute. On the exterior algebra  $\Lambda_{\mathbb{Z}}(\alpha_3, \alpha_5, \dots, \alpha_{2n+1})$ , we have *odd* derivations  $\partial/\partial\alpha_{2\ell+1}$  for  $1 \leq \ell \leq n$  in the sense that when we move  $\partial/\partial\alpha_{2\ell+1}$  beyond  $\alpha_{2k+1}$ , we get a minus sign.

The circle action  $\Delta : S^1 \times LM \rightarrow LM$  induces the BV operator  $\Delta : \mathbb{H}_*(LM) \rightarrow \mathbb{H}_{*+1}(LM)$  given by  $\Delta(z) = \Delta_*([S^1] \otimes z)$  for  $z \in \mathbb{H}_*(LM)$ . Recall that the BV Lie bracket  $\{\cdot, \cdot\}$  and the BV operator  $\Delta$  are related by a formula

$$(1-3) \quad \Delta(a \cdot b) = \Delta(a) \cdot b + (-1)^{|a|} a \cdot \Delta(b) + (-1)^{|a|} \{a, b\},$$

for arbitrary two elements  $a, b$  in a BV algebra.

**Theorem A.** (1) The BV operator  $\Delta$  on the loop homology  $\mathbb{H}_*(\mathrm{LSU}(n+1))$  is given by

$$\Delta(\alpha_I e^J) = \sum_{\ell=1}^n \frac{\partial \alpha_I}{\partial \alpha_{2\ell+1}} \cdot D_{2\ell}(e^J).$$

(2) The BV Lie bracket in the loop homology  $\mathbb{H}_*(\mathrm{LSU}(n+1))$  is given by

$$(-1)^{|I|} \{\alpha_I e^J, \alpha_K e^L\} = \Delta(\alpha_I e^L) \cdot \alpha_K e^J + (-1)^{|I|} \alpha_I e^L \cdot \Delta(\alpha_K e^J).$$

It is interesting that the formula for the BV Lie bracket is given by a very clean derivation formula of  $\Delta$ , but with  $e$ -terms switched. To prove Theorem A, the homological  $S^1$ -action is first calculated in the Hopf algebra  $H_*(\mathrm{LSU}(n+1))$  with Pontrjagin product with respect to which  $\Delta$  is a derivation. Its coalgebra structure is used to determine  $\Delta$  on Pontrjagin ring generators. Then the result is translated to the loop homology.

To have a better understanding of the BV Lie bracket, we take a point of view that the algebra in (1-1) is a super algebra of functions on a super vector space in which  $e_2, e_4, \dots, e_{2n}$  are even coordinate functions and  $\alpha_3, \alpha_5, \dots, \alpha_{2n+1}$  are odd coordinate functions. See [4] for super algebras and super geometry. In the Hopf algebra  $H_*(\Omega\mathrm{SU}(n+1)) \otimes \mathbb{Q}$  over rationals, there exist primitive

elements  $h_{2\ell}$  such that  $h_{2\ell} = e_{2\ell} + (\text{decomposables})$  for  $1 \leq \ell \leq n$  so that  $H_*(\Omega\text{SU}(n+1)) \otimes \mathbb{Q} \cong \mathbb{Q}[h_2, h_4, \dots, h_{2n}]$ . We regard  $h_{2\ell}$ 's as new even coordinate functions, and as such a differentiation  $\partial/\partial h_{2\ell}$  makes sense for functions in  $e_{2k}$ 's. On the super vector space  $\mathbb{Q}^{n|n}$ , we consider an odd symplectic form

$$\omega = \sum_{\ell=1}^n d\alpha_{2\ell+1} \wedge dh_{2\ell}.$$

We consider the associated Poisson bracket on functions  $[\cdot, \cdot]_\omega : \mathbb{H}_*^{\mathbb{Q}} \otimes \mathbb{H}_*^{\mathbb{Q}} \rightarrow \mathbb{H}_*^{\mathbb{Q}}$ , where  $\mathbb{H}_*^{\mathbb{Q}} = \mathbb{H}_* \otimes \mathbb{Q}$ , given by  $[F, G]_\omega = X_F G$ , where  $X_F$  is the Hamiltonian vector field associated to  $F$  by  $\iota_X \omega = dF$ . Concretely, the Poisson bracket is given by

$$[\alpha_I e^J, \alpha_K e^L]_\omega = \sum_{\ell=1}^n \left( (-1)^{|I|} \frac{\partial(\alpha_I e^J)}{\partial \alpha_{2\ell+1}} \cdot \frac{\partial(\alpha_K e^L)}{\partial h_{2\ell}} + \frac{\partial(\alpha_I e^J)}{\partial h_{2\ell}} \cdot \frac{\partial(\alpha_K e^L)}{\partial \alpha_{2\ell+1}} \right).$$

The next theorem explains the nature of the BV Lie bracket.

**Theorem B.** *The Poisson bracket is well defined over  $\mathbb{H}_*$ , and the BV Lie bracket in the loop homology  $\mathbb{H}_*(\text{LSU}(n+1))$  coincides with the Poisson bracket associated to the symplectic form  $\omega$ . Namely,  $\{\alpha_I e^J, \alpha_K e^L\} = [\alpha_I e^J, \alpha_K e^L]_\omega$  for  $\alpha_I e^J, \alpha_K e^L \in \mathbb{H}_*$ .*

It is satisfying that symplectic even coordinate functions  $h_{2\ell}$ 's are given exactly by primitive elements in the function algebra  $\mathbb{H}_*^{\mathbb{Q}} = \mathbb{H}_* \otimes \mathbb{Q}$ .

Let  $\mathbb{H}(\ell) = \Lambda_{\mathbb{Z}}(\alpha) \otimes \mathbb{Z}[h]$  with  $|\alpha| = -2\ell - 1$  and  $|h| = 2\ell$  be a BV algebra associated to an odd symplectic form  $d\alpha \wedge dh$  and with a BV operator given by  $\Delta(h^k) = 0$  and  $\Delta(\alpha h^k) = kh^{k-1}$  for  $k \geq 0$ . Since the symplectic vector space  $(\mathbb{Q}^{n|n}, \omega)$  is a direct sum  $\bigoplus_{\ell=1}^n (\mathbb{Q}^{1|1}, d\alpha_{2\ell+1} \wedge dh_{2\ell})$ , over rationals, our BV algebra  $\mathbb{H}_*(\text{LSU}(n+1))$  admits a corresponding splitting. Namely,

**Corollary C.** *Over rationals, the loop homology of  $\text{SU}(n+1)$  splits into a following tensor product as BV algebras:*

$$\mathbb{H}_*(\text{LSU}(n+1)) \otimes \mathbb{Q} \cong \bigotimes_{\ell=1}^n \mathbb{H}(\ell) \otimes \mathbb{Q}.$$

The above corollary is related to a fact that over rationals, the Lie group  $\text{SU}(n+1)$  splits as a product of odd dimensional spheres, and the BV algebra  $\mathbb{H}(\ell)$  is precisely the loop homology of  $S^{2\ell+1}$  in view of Theorem 5-2.

In general, for a BV algebra  $\mathbb{H}_*$ , the Lie bracket is a derivation in each variable. For example,

$$\{a, b \cdot c\} = \{a, b\} \cdot c + (-1)^{|b|(|a|+1)} b \cdot \{a, c\},$$

for any three elements  $a, b, c \in \mathbb{H}_*$ . This means that there is a Lie algebra homomorphism from  $\mathbb{H}_*$  to the Lie algebra  $\text{Der}(\mathbb{H}_*)$  of all derivations on the associative algebra  $\mathbb{H}_*$ . In our case of  $\mathbb{H}_* = \mathbb{H}_*(\text{LSU}(n+1))$ , the Lie algebra  $\text{Der}(\mathbb{H}_*)$  is a free  $\mathbb{H}_*$ -module of rank  $n|n$ . Theorem B suggests that the description of the Lie algebra structure on the loop homology simplifies when tensored with  $\mathbb{Q}$  so that primitive elements are available. For a derivation  $X \in \text{Der}(\mathbb{H}_*^{\mathbb{Q}})$ , let  $\mathcal{L}_X$  be its Lie derivative given by  $\mathcal{L}_X = d \circ \iota_X + \iota_X \circ d$ .

**Corollary D.** (1) *The BV Lie algebra  $\mathbb{H}_*^{\mathbb{Q}}/\mathbb{Q}\cdot 1$  is isomorphic to the super Lie algebra of Hamiltonian derivations given by  $\{X \in \text{Der}(\mathbb{H}_*^{\mathbb{Q}}) \mid \mathcal{L}_X \omega = 0\}$ .*

(2) *Let  $Q \subset \mathbb{H}_*^{\mathbb{Q}}$  be the subspace spanned by quadratic elements in  $h_{2\ell}$ 's and  $\alpha_{2k+1}$ 's. Then  $Q$  is a super Lie subalgebra isomorphic to the super Lie algebra  $\pi\text{sp}(n|n)$ .*

Here the super Lie algebra  $\pi sp(n|n)$  consists of those linear maps on a vector space with basis  $\{h_2, \dots, h_{2n}, \alpha_3, \dots, \alpha_{2n+1}\}$  preserving the odd symplectic form  $\omega$  [5].

For the complex Stiefel manifold  $V_{n+1-k}(\mathbb{C}^{n+1}) = \text{SU}(n+1)/\text{SU}(k)$  of  $n+1-k$  orthonormal frames in  $\mathbb{C}^{n+1}$ , entirely analogous statements hold in which

$$\mathbb{H}_*(V_{n+1-k}(\mathbb{C}^{n+1})) \cong \Lambda_{\mathbb{Z}}(\alpha_{2k+1}, \dots, \alpha_{2n+1}) \otimes \mathbb{Z}[e_{2k}, \dots, e_{2n}],$$

with some obvious modifications on the operators  $D_{2\ell}$ 's.

In this article, all homology and cohomology groups have integer coefficients unless otherwise stated.

The organization of this paper is as follows. After reviewing the relationship of intersection product and Pontrjagin product in the homology of  $\text{SU}(n+1)$  in section 2, we compute the homological  $S^1$  action in the Pontrjagin ring in section 3. Here the Hopf algebra is effectively used to determine this action. Then in section 4, we describe the full BV algebra structure of the loop homology of  $\text{SU}(n+1)$  using the result of section 3. After deriving the formula for the BV Lie bracket, we describe its meaning in the context of the odd symplectic structure. In section 5, we describe BV algebra structure on the loop homology of complex Stiefel manifolds.

## §2 Pontrjagin and intersection products in $H_*(\text{SU}(n+1))$

In this section, we review the homology and cohomology groups of  $\text{SU}(n+1)$  as Hopf algebras, with a discussion on a relation between Pontrjagin product and intersection product in  $H_*(\text{SU}(n+1))$ . We start with the Pontrjagin structure. A basic reference for the topology of Lie groups is [6].

Let  $\Sigma\mathbb{C}P^n$  be the reduced suspension of  $\mathbb{C}P^n = \text{U}(n+1)/(\text{U}(1) \times \text{U}(n))$ . Let a map  $\beta_n : \Sigma\mathbb{C}P^n \rightarrow \text{SU}(n+1)$  be given by  $\beta_n(t, [A]) = A \cdot R_t(1, n) \cdot A^{-1} \cdot R_t(1, n)^{-1}$ , where  $A \in \text{U}(n+1)$  and  $R_t(1, n) = e^{\pi it} \oplus e^{-\pi it} I_n$ . It is easy to check that this definition is independent of the choice of the representative  $A$  of the equivalence class  $[A]$ . Let  $\bar{\beta}_n : \mathbb{C}P^n \rightarrow \Omega\text{SU}(n+1)$  be the adjoint of  $\beta_n$ . The Pontrjagin ring structures on the homology of  $\text{SU}(n+1)$  and of its based loop space  $\Omega\text{SU}(n+1)$  are given by the following well known theorem.

**Theorem 2-1.** (1) *Let  $x_{2k+1} = \beta_{n*}([\Sigma\mathbb{C}P^k])$  for  $1 \leq k \leq n$ . Then the Hopf algebra structure of the homology of  $\text{SU}(n+1)$  is given by*

$$H_*(\text{SU}(n+1)) \cong \Lambda_{\mathbb{Z}}(x_3, x_5, \dots, x_{2n+1}),$$

*and all generators are primitive.*

(2) *Let  $e_{2k} = \bar{\beta}_{n*}([\mathbb{C}P^k])$  for  $1 \leq k \leq n$ . Then the Pontrjagin ring structure and the coalgebra structure of the homology of  $\Omega\text{SU}(n+1)$  are given by*

$$H_*(\Omega\text{SU}(n+1)) \cong \mathbb{Z}[e_2, e_4, \dots, e_{2n}], \quad \phi_*(e_{2k}) = \sum_{i+j=k} e_{2i} \otimes e_{2j}.$$

We also regard homology elements  $x_{2k+1}$  and  $e_{2k}$  as maps  $x_{2k+1} : \Sigma\mathbb{C}P^k \rightarrow \text{SU}(n+1)$  and  $e_{2k} : \mathbb{C}P^k \rightarrow \Omega\text{SU}(n+1)$ , which are adjoint to each other, for  $1 \leq k \leq n$ .

Let  $U$  be the sequence  $U = (3, 5, \dots, 2n+1)$ . For any sequence  $I = (2i_1+1, 2i_2+1, \dots, 2i_r+1)$  with  $1 \leq i_1 < \dots < i_r \leq n$ , we let  $x_I = x_{2i_1+1} \cdot x_{2i_2+1} \cdot \dots \cdot x_{2i_r+1}$ . We orient  $\text{SU}(n+1)$  by the fundamental class  $x_U$ . Let  $y_{2k+1}$  be the cohomology element dual to  $x_{2k+1}$  with respect to the monomial basis  $\{x_I\}$ . It is straightforward to dualize the Hopf algebra structure of  $H_*(\text{SU}(n+1))$  to get the Hopf algebra structure of the cohomology.

**Theorem 2-2.** *As a Hopf algebra we have*

$$H^*(\mathrm{SU}(n+1)) \cong \Lambda_{\mathbb{Z}}(y_3, y_5, \dots, y_{2n+1}),$$

where all the exterior generators are primitive and the Kronecker pairing is given by

$$(2-1) \quad \langle y_I, x_J \rangle = (-1)^{\frac{|I|(|I|-1)}{2}} \delta_{I,J}.$$

The sign above is the result of the usual sign convention in which switching two odd degree objects gives a minus sign.

Next we discuss the intersection product in homology. For an oriented closed manifold  $M^d$ , the intersection product in homology can be defined as the Poincaré dual of the cup product in cohomology. Namely, let  $D : H_*(M) \xrightarrow{\cong} H^{d-*}(M)$  be the Poincaré duality isomorphism given by  $D(\alpha) \cap [M] = \alpha$  for  $\alpha \in H_*(M)$ . Then for  $\alpha, \beta \in H_*(M)$ , their intersection product  $\alpha \circ \beta$  is defined by  $D(\alpha \circ \beta) = D(\alpha) \cup D(\beta)$ . It is immediate that  $\beta \circ \alpha = (-1)^{(d-|\alpha|)(d-|\beta|)} \alpha \circ \beta$ .

**Remark.** If  $\alpha, \beta$  are homology classes represented by submanifolds  $K, N$  intersecting transversally, the orientation of  $\alpha \circ \beta = [K \cap N]$  is given as follows. At  $x \in K \cap N$ , choose a basis  $\vec{v} = \{v_1, \dots, v_k\}$  of  $T_x(K \cap N)$ . Extend it to an oriented basis of  $T_x K$  and of  $T_x N$  as  $\{v_1, \dots, v_k, u_1, \dots, u_\ell\}$  and as  $\{v_1, \dots, v_k, w_1, \dots, w_m\}$ , respectively. Then if  $\{\vec{v}, \vec{w}, \vec{u}\}$  is the oriented basis of  $T_x M$ , then  $\vec{v}$  gives the orientation of  $K \cap N$ . Otherwise, opposite orientation gives the orientation of  $K \cap N$ .

We come back to the homology of  $\mathrm{SU}(n+1)$ . For a sequence  $I$ , let  $\alpha_I \in H_*(\mathrm{SU}(n+1))$  be the Poincaré dual to  $y_I$ , that is  $D(\alpha_I) = y_I$ . Let  $I^c$  be the sequence complement to  $I$  in  $U$ . For two disjoint sequences  $I, J$  of strictly increasing integers, let  $I \cup J$  be the combined sequence of strictly increasing integers. Let  $\mathrm{sgn}(I, J) \in \{\pm 1\}$  be defined by

$$(2-2) \quad x_{I \cup J} = \mathrm{sgn}(I, J) x_I \cdot x_J$$

with respect to the Pontrjagin product. This is the same sign resulting from converting the juxtaposition  $IJ$  into a strictly increasing sequence  $I \cup J$ . Note that  $\mathrm{sgn}(I, J) = (-1)^{|I||J|} \mathrm{sgn}(J, I)$ .

**Lemma 2-3.** (1) *For any sequence  $I$ , the homology class  $\alpha_I$  which is the Poincaré dual of  $y_I$ , is identified as*

$$(2-3) \quad \alpha_I = (-1)^{\frac{|I|(|I|-1)}{2}} \mathrm{sgn}(I, I^c) x_{I^c}.$$

(2) *For two disjoint sequences  $I, J$ , the intersection product is given by*

$$\alpha_I \circ \alpha_J = \mathrm{sgn}(I, J) \alpha_{I \cup J}.$$

Thus, if  $I = (2i_1 + 1, 2i_2 + 1, \dots, 2i_r + 1)$  with  $1 \leq i_1 < \dots < i_r \leq n$ , we have

$$\alpha_I = \alpha_{2i_1+1} \circ \alpha_{2i_2+1} \circ \dots \circ \alpha_{2i_r+1}.$$

*Proof.* Since  $D(\alpha_I) = y_I$ , for any sequence  $J$ , we have  $\langle y_J, \alpha_I \rangle = \langle y_J, D(\alpha_I) \cap x_U \rangle = \langle y_J \cup y_I, x_U \rangle$ , where  $y_J \cup y_I = \mathrm{sgn}(J, I) y_{I \cup J}$  and the last pairing is nonzero only when  $J = I^c$ . Hence we have  $\langle y_J, \alpha_I \rangle = (-1)^{\frac{n(n-1)}{2}} \mathrm{sgn}(I^c, I) \delta_{J, I^c}$ , using (2-1). This means that the homology class  $\alpha_I$  is  $x_{I^c}$  up to a sign. Since  $\langle y_{I^c}, x_{I^c} \rangle = (-1)^{\frac{|I^c|(|I^c|-1)}{2}}$ , we see that  $\alpha_I = (-1)^{\frac{|I|(|I|-1)}{2}} \mathrm{sgn}(I, I^c) x_{I^c}$ .

For the second formula, we compute:

$$D(\alpha_I \circ \alpha_J) = D(\alpha_I) \cup D(\alpha_J) = y_I \cup y_J = \text{sgn}(I, J)y_{I \cup J} = \text{sgn}(I, J)D(\alpha_{I \cup J}).$$

This allows us to write  $\alpha_I$  as an intersection product of  $\alpha_{2i+1}$ 's.  $\square$

Since homology elements  $\alpha_I$ 's are dual to cohomology elements, they are more suitable for computing intersection products than  $x_I$ 's, and elements  $x_I$ 's are used for computing Pontrjagin products. In homology, the intersection product is denoted by a circle "o" and the Pontrjagin product is denoted by a dot ".".

Next we examine the relationship between the intersection product and the Pontrjagin product. Properties of  $\alpha_I$ 's are best described through Poincaré duality.

**Proposition 2-4.** *For any sequence  $I$ , we have*

$$y_{2i+1} \cup D(x_{2i+1} \cdot \alpha_I) = \begin{cases} y_I & \text{if } 2i+1 \in I, \\ 0 & \text{if } 2i+1 \notin I. \end{cases}$$

*In other words,*

$$x_{2i+1} \cdot \alpha_I = \begin{cases} (-1)^{r-1} \alpha_{I - \{2i+1\}} & \text{if } 2i+1 \in I, \\ 0 & \text{if } 2i+1 \notin I. \end{cases}$$

*Here, when  $2i+1$  appears in  $I$ , it appears as the  $r$ -th entry.*

*Proof.* Since  $\alpha_I = \pm x_{I^c}$ , if  $2i+1 \notin I$ , then  $x_{2i+1} \cdot \alpha_I = 0$ . Suppose  $2i+1 \in I$ . By (2-2) and (2-3), we have

$$x_{2i+1} \cdot \alpha_I = (-1)^{\frac{|I|(|I|-1)}{2}} \text{sgn}(I, I^c) \text{sgn}(\{2i+1\}, I^c) x_{I^c \cup \{2i+1\}}.$$

On the other hand, by (2-3) we have

$$\alpha_{I - \{2i+1\}} = (-1)^{\frac{(|I|-1)(|I|-2)}{2}} \text{sgn}(I - \{2i+1\}, I^c \cup \{2i+1\}) x_{I^c}.$$

Comparing the signs, we get  $x_{2i+1} \cdot \alpha_I = (-1)^{|I|-1} \text{sgn}(I - \{2i+1\}, \{2i+1\}) \alpha_{I - \{2i+1\}}$ . The last sign is  $(-1)^{|I|-r}$  if  $2i+1$  appears in  $I$  as the  $r$ -th entry. This proves the second part. For the first part, we have  $y_{2i+1} \cdot D(x_{2i+1} \cdot \alpha_I) = y_{2i+1} \cdot (-1)^{r-1} y_{I - \{2i+1\}} = y_I$ .  $\square$

This result means that the Pontrjagin product with  $x_{2i+1}$  coincides, in the intersection ring  $\mathbb{H}_*(\text{SU}(n+1)) = \Lambda_{\mathbb{Z}}(\alpha_3, \dots, \alpha_{2n+1})$ , with the odd derivation  $\partial/\partial\alpha_{2i+1}$ , where  $\alpha_{2k+1}$ 's are regarded as odd variables to get the sign right. Thus for any sequence  $I$ ,

$$(2-4) \quad x_{2i+1} \cdot \alpha_I = \frac{\partial \alpha_I}{\partial \alpha_{2i+1}}, \quad 1 \leq i \leq n.$$

The next corollary is an immediate consequence of this fact.

**Corollary 2-5.** *For any two sequences  $I, J$  not necessarily disjoint, in  $H_*(\text{SU}(n+1))$  we have*

$$x_{2i+1} \cdot (\alpha_I \circ \alpha_J) = (x_{2i+1} \cdot \alpha_I) \circ \alpha_J + (-1)^{|I|} \alpha_I \circ (x_{2i+1} \cdot \alpha_J).$$

### §3 Homology $S^1$ -action on the Hopf algebra $H_*(LSU(n+1))$

For a Lie group  $G$ , its free loop space is topologically a product  $\xi : G \times \Omega G \cong LG$  in which  $(g, \gamma)$  is mapped to  $g \cdot \gamma$  for an element  $g \in G$  and a loop  $\gamma$  based at the identity element, using the multiplication  $\mu : G \times G \rightarrow G$  in  $G$ . Although the map  $\xi$  commutes with diagonal maps, it does not commute with group multiplications on both sides unless  $G$  is abelian. Thus there may be a possibility that the Hopf algebra structure of  $H_*(LG)$  is different from the tensor product of Hopf algebras  $H_*(G)$  and  $H_*(\Omega G)$ . However, under mild assumptions on  $G$ , we show that indeed the Hopf algebra structure  $H_*(LG)$  is isomorphic to the tensor product of two Hopf algebras  $H_*(G)$  and  $H_*(\Omega G)$ .

To compare the group structures of both sides of the map  $\xi$ , we consider the conjugation map  $\text{Ad} : \Omega G \times G \rightarrow \Omega G$  given by  $\text{Ad}(\gamma, g) = g^{-1} \cdot \gamma \cdot g$ . Then the following diagram commutes:

$$(3-1) \quad \begin{array}{ccc} (G \times \Omega G) \times (G \times \Omega G) & \xrightarrow[\cong]{\xi \times \xi} & LG \times LG \\ \downarrow 1 \times 1 \times \phi \times 1 & & \downarrow \mu \\ G \times \Omega G \times G \times G \times \Omega G & & LG \\ \downarrow 1 \times \text{Ad} \times 1 \times 1 & & \cong \uparrow \xi \\ G \times \Omega G \times G \times \Omega G & \xrightarrow{(\mu \times \mu) \circ (1 \times T \times 1)} & G \times \Omega G \end{array}$$

where  $\phi$  is the diagonal map and  $T$  is the switching map. To study the homology of this diagram, we need to know the induced map  $\text{Ad}_*$  on homology, which turn out to be trivial in most cases.

**Lemma 3-1.** *Assume that the Pontrjagin ring  $H_*(G)$  is torsion free and generated by odd degree elements. Then positive degree elements of  $H_*(G)$  acts trivially on  $H_*(\Omega G)$  through  $\text{Ad}_*$ . Namely, for  $e \in H_*(\Omega G)$  and  $x \in H_*(G)$ ,*

$$\text{Ad}_*(e \otimes x) = \begin{cases} 0 & \text{if } |x| > 0, \\ e & \text{if } x = 1 \in H_0(G). \end{cases}$$

*Proof.* Since  $H_*(G) \cong \Lambda_{\mathbb{Z}}(x_a \mid a \in A)$ , where  $A$  is an index set, we have  $H_*(\Omega G) \cong \mathbb{Z}[e_a \mid a \in A]$  with  $|e_a| + 1 = |x_a|$ . Then for each  $a \in A$ , the element  $\text{Ad}_*(e \otimes x_a)$  has an odd degree, and hence must vanish since  $\mathbb{Z}[e_a]$  has only even degree elements. Since  $\text{Ad}_*$  defines a right action of the Pontrjagin ring  $H_*(G)$  on  $H_*(\Omega G)$ , we see that  $\text{Ad}_*$  action vanishes for all positive degree elements in  $H_*(G)$ .  $\square$

An immediate consequence is that the composition  $(\text{Ad} \times 1) \circ (1 \times \phi) : \Omega G \times G \rightarrow \Omega G \times G$  induces an identity map in homology. Thus, the commutative diagram (3-1) implies the following result.

**Proposition 3-2.** *Assume that  $H_*(G)$  is torsion free and generated by odd degree elements. Then  $H_*(LG) \cong H_*(G) \otimes H_*(\Omega G)$  as Hopf algebras.*

Next we examine the compatibility between the Hopf algebra structure of  $H_*(LG)$  and the homological  $S^1$  action  $\Delta : H_*(LG) \rightarrow H_{*+1}(LG)$  given by  $\Delta(z) = \Delta_*([S^1] \otimes z)$  for  $z \in H_*(LG)$ . Here we consider  $\Delta$  in the Pontrjagin ring  $H_*(LG)$ , not in the loop homology algebra  $\mathbb{H}_*(LG)$ , which is the topic of the next section.

**Proposition 3-3.** *Let  $G$  be a Lie group.*

(1) *The map  $\Delta$  is a derivation on the Pontrjagin ring  $H_*(LG)$ . Namely, for  $z, w \in H_*(LG)$ ,*

$$\Delta(z \cdot w) = \Delta(z) \cdot w + (-1)^{|z|} z \cdot \Delta(w).$$

(2) *The map  $\Delta$  commutes with the coalgebra map  $\phi : H_*(LG) \rightarrow H_*(LG) \otimes H_*(LG)$ . Namely, if  $\phi(z) = \sum_i z'_i \otimes z''_i$  for  $z \in H_*(LG)$ , then*

$$\phi(\Delta(z)) = \Delta(\phi(z)) = \sum_i \left\{ \Delta(z'_i) \otimes z''_i + (-1)^{|z'_i|} z'_i \otimes \Delta(z''_i) \right\}.$$

*Proof.* These are straightforward consequences of the following commutative diagrams.

$$\begin{array}{ccccc} S^1 \times (LG \times LG) & \xrightarrow{1 \times \mu} & S^1 \times LG & & \\ (1 \times T \times 1) \circ (\phi \times 1 \times 1) \downarrow & & \Delta \downarrow & & S^1 \times LG \xrightarrow{\Delta} LG \\ (S^1 \times LG) \times (S^1 \times LG) & & LG & & (1 \times T \times 1) \circ (\phi \times \phi) \downarrow \phi \downarrow \\ \Delta \times \Delta \downarrow & & \parallel & & (S^1 \times LG) \times (S^1 \times LG) \xrightarrow{\Delta \times \Delta} LG \times LG \\ LG \times LG & \xrightarrow{\mu} & LG & & \end{array}$$

For a loop  $\gamma \in LG$ , let  $\gamma_t$  denote the loop rotated by  $t \in S^1 = \mathbb{R}/\mathbb{Z}$ . Then the first diagram above says  $(\gamma \cdot \eta)_t = \gamma_t \cdot \eta_t$  for any two loops  $\gamma$  and  $\eta$ . The second diagram says  $(\gamma, \gamma)_t = (\gamma_t, \gamma_t)$ .  $\square$

If we can determine the effect of  $\Delta$  on the generators of the Pontrjagin ring  $H_*(LG)$ , then by derivation property of  $\Delta$ , we can determine  $\Delta$  on the entire ring  $H_*(LG)$ .

To describe  $\Delta$ , we need differential operators  $D_{2\ell}$  acting on the polynomial ring  $\mathbb{Z}[e_2, e_4, \dots, e_{2n}]$  introduced in (1-2). These differential operators have the following properties.

**Proposition 3-4.** *Let  $\phi$  be the coalgebra map in  $\mathbb{Z}[e_2, e_4, \dots, e_{2n}]$  given in Theorem 2-1.*

(1) *The differential operators  $D_2, D_4, \dots, D_{2n}$  are characterized by*

$$(3-2) \quad \phi(e^J) = 1 \otimes e^J + e_2 \otimes D_2 e^J + e_4 \otimes D_4 e^J + \dots + e_{2n} \otimes D_{2n} e^J + \sum_{I+K=J} a_{I,J} e^I \otimes e^K,$$

where in the last summation,  $e^I$  is a product of at least two elements.

(2) *The operators  $D_{2\ell}$ 's mutually commute, that is,  $[D_{2i}, D_{2j}] = 0$  for  $1 \leq i, j \leq n$ .*

*Proof.* The formula in (1) can be checked directly. The commutativity of  $D_{2\ell}$ 's can also be checked directly, but we can also use the fact that the coalgebra map  $\phi$  is cocommutative, and we compare terms of the form  $e_{2i} \otimes e_{2j} \otimes (D_{2j} D_{2i} e^J)$  in the both sides of  $(T \otimes 1)(1 \otimes \phi)\phi(e^J) = (1 \otimes \phi)\phi(e^J)$ .  $\square$

To determine the effect of  $\Delta$  on algebra generators of  $H_*(LSU(n+1))$ , we use the coalgebra structure.

**Theorem 3-5.** *In the Pontrjagin ring  $H_*(LSU(n+1)) \cong \Lambda(x_3, x_5, \dots, x_{2n+1}) \otimes \mathbb{Z}[e_2, e_4, \dots, e_{2n}]$ , the effect of  $\Delta$  on generators is given by*

$$\Delta(x_{2\ell+1}) = 0, \quad \Delta(e_{2\ell}) = \sum_{i=1}^{\ell} x_{2i+1} \cdot e_{2\ell-2i}, \quad 1 \leq \ell \leq n.$$



The general formula of the action  $\Delta$  is

$$(3-3) \quad \Delta(x_I e^J) = \sum_{\ell=1}^n (x_{2\ell+1} \cdot x_I) \cdot (D_{2\ell} e^J).$$

*Proof.* Since the map  $x_{2k+1} : \Sigma \mathbb{C}P^k \rightarrow \mathrm{SU}(n+1) \rightarrow \mathrm{LSU}(n+1)$  is a cycle consisting of constant loops, the  $S^1$ -action on it is trivial and we have  $\Delta(x_{2k+1}) = 0$  for all  $k$ . Since  $x_{2k+1}$  is the adjoint of the map  $e_{2k} : \mathbb{C}P^k \rightarrow \Omega \mathrm{SU}(n+1)$ ,  $x_{2k+1}$  can be thought of as the following composition:

$$x_{2k+1} : S^1 \times \mathbb{C}P^k \xrightarrow{1 \times e_{2k}} S^1 \times \Omega \mathrm{SU}(n+1) \rightarrow S^1 \times \mathrm{LSU}(n+1) \xrightarrow{\Delta} \mathrm{LSU}(n+1) \xrightarrow{p} \mathrm{SU}(n+1),$$

where  $p$  is the projection to the base of loops. This means  $x_{2k+1} = p_* (\Delta_* ([S^1] \otimes e_{2k})) = p_* (\Delta(e_{2k}))$ . Thus, in  $H_*(\mathrm{LSU}(n+1))$ , we have  $\Delta(e_{2k}) = x_{2k+1} + (\text{decomposables})$ , where each term in the decomposables has a nontrivial  $e$ -factor. We determine the decomposable part by induction on  $k$ . When  $k = 1$ , by dimensional reason, we must have  $\Delta(e_2) = x_3$ . Now assume that  $\Delta(e_{2k}) = \sum_{i=1}^k x_{2i+1} \cdot e_{2k-2i}$  for  $1 \leq k \leq \ell - 1$  for some  $\ell \geq 2$ . Since  $\Delta$  is a coalgebra map by Proposition 3-3, we have  $\phi(\Delta(e_{2\ell})) = \Delta(\phi(e_{2\ell}))$ . We compare both sides. We let  $\Delta(e_{2\ell}) = x_{2\ell+1} + Y$ , where  $Y$  is a decomposable element of homogeneous degree  $2\ell + 1$ . Since  $\phi(e_{2\ell}) = \sum_{k=0}^{\ell} e_{2k} \otimes e_{2\ell-2k}$  and  $x_{2\ell+1}$  is primitive, the above identity gives the following identity for  $Y$ :

$$\phi(Y) = Y \otimes 1 + 1 \otimes Y + \sum_{k=1}^{\ell-1} (\Delta(e_{2k}) \otimes e_{2\ell-2k} + e_{2k} \otimes \Delta(e_{2\ell-2k})).$$

Applying formula (3-2) to  $Y$ , and comparing it with the above identity, we see that  $Y$  must satisfy a system of PDEs given by  $D_{2k}(Y) = \Delta(e_{2\ell-2k})$  for  $1 \leq k \leq \ell - 1$ , and  $D_{2k}(Y) = 0$  for  $\ell \leq k \leq n$ . Using the explicit expression of operators  $D_{2k}$  given in (1-2), the above system can be rewritten in matrix form as

$$\begin{pmatrix} 1 & e_2 & \dots & e_{2\ell-2} & \dots & e_{2n-2} \\ & 1 & \dots & e_{2\ell-4} & \dots & e_{2n-4} \\ & & \ddots & \vdots & \ddots & \vdots \\ & & & 1 & \dots & e_{2n-2\ell} \\ & & & & \ddots & \vdots \\ & & & & & 1 \end{pmatrix} \begin{pmatrix} \partial_2 Y \\ \partial_4 Y \\ \vdots \\ \partial_{2\ell} Y \\ \vdots \\ \partial_{2n} Y \end{pmatrix} = \begin{pmatrix} \Delta(e_{2\ell-2}) \\ \Delta(e_{2\ell-4}) \\ \vdots \\ \Delta(e_2) \\ \vdots \\ 0 \end{pmatrix}$$

where the coefficient matrix has zeros below the diagonal. Since the coefficient matrix is invertible, we can uniquely determine partial derivatives  $\partial_{2k} Y = \partial Y / \partial e_{2k}$  for all  $1 \leq k \leq n$ . Since the constant term of  $Y$  is zero, if the solution to the system exists, it must be unique. On the other hand, it is easy to check that  $Y = \sum_{i=1}^{\ell-1} x_{2i+1} e_{2\ell-2i}$  satisfies the system using the induction hypothesis. Thus, we have  $\Delta(e_{2\ell}) = x_{2\ell+1} + Y = \sum_{i=1}^{\ell} x_{2i+1} e_{2\ell-2i} = \sum_{i=1}^{\ell} x_{2i+1} D_i(e_{2\ell})$ . This proves the first part.

The second part is a simple consequence of the derivation property of  $\Delta$  and differential operators  $D_{2k}$ , using the calculation of  $\Delta$  on algebra generators above.  $\square$

## §4 Batalin-Vilkovisky Lie algebra structure on the loop homology $\mathbb{H}_*(\mathbf{LSU}(n+1))$

In this section we describe the full BV algebra structure of the loop homology of  $\mathbf{SU}(n+1)$ . First we give a brief description of a general Batalin-Vilkovisky algebra.

**Definition.** A Batalin-Vilkovisky algebra  $A$  is a graded commutative associative algebra with a degree 1 operator  $\Delta$  with  $\Delta^2 = 0$  such that the bilinear product  $\{ , \}$  on  $A$  defined by

$$(4-1) \quad \Delta(ab) = \Delta(a) \cdot b + (-1)^{|a|} a \cdot \Delta(b) + (-1)^{|a|} \{a, b\}, \quad a, b \in A$$

is a derivation in each variable in the sense that for every  $a, b, c \in A$ ,

$$\begin{aligned} \{a, b \cdot c\} &= \{a, b\} \cdot c + (-1)^{|b|(|a|+1)} b \cdot \{a, c\}, \\ \{a \cdot b, c\} &= a \cdot \{b, c\} + (-1)^{|b|(|c|+1)} \{a, c\} \cdot b. \end{aligned}$$

It is a formal consequence that the bilinear product  $\{ , \}$  defined above satisfies identities for a Lie bracket of degree 1, meaning

$$\begin{aligned} \{a, b\} &= -(-1)^{(|a|+1)(|b|+1)} \{b, a\}, \\ \{a, \{b, c\}\} &= \{\{a, b\}, c\} + (-1)^{(|a|+1)(|b|+1)} \{b, \{a, c\}\}. \end{aligned}$$

We note that additively there is a bijective correspondence of elements between the Pontrjagin ring  $H_*(\mathbf{LSU}(n+1))$  and the loop homology  $\mathbb{H}_*(\mathbf{LSU}(n+1))$  preserving the homological  $S^1$ -action  $\Delta$ , although the ring structure and the grading are different between these two rings. Specifically, this correspondence is given by (2-3). Using this bijection, we can translate the homological  $S^1$ -action  $\Delta$  given in (3-3) in the Pontrjagin ring to the loop homology, and we get the following formula:

$$\Delta(\alpha_I e^J) = \sum_{\ell=1}^n \frac{\partial \alpha_I}{\partial \alpha_{2\ell+1}} \cdot (D_{2\ell} e^J),$$

for any sequences  $I, J$ , where  $\partial/\partial \alpha_{2\ell+1}$  is an odd derivation on odd variables  $\alpha_{2k+1}$ 's given in (2.4). Here are some special cases of the above formula.

$$\Delta(\alpha_I) = 0, \quad \Delta(e^J) = 0, \quad \Delta(\alpha_{2\ell+1} e^J) = D_{2\ell} e^J, \quad \Delta(\alpha_{2\ell+1} e_{2\ell}) = 1, \quad 1 \leq \ell \leq n.$$

We use the bold face  $\Delta$  for the BV operator in the loop homology to distinguish it from the  $\Delta$  operator in the Pontrjagin ring.

We can now compute the BV Lie bracket using the relation (4-1). For arbitrary two elements  $\alpha_I e^J, \alpha_K e^L \in \mathbb{H}_*(\mathbf{LSU}(n+1))$ , we have

$$\begin{aligned} \Delta(\alpha_I e^J \circ \alpha_K e^L) &= \sum_{i=1}^n \frac{\partial}{\partial \alpha_{2i+1}} (\alpha_I \alpha_K) \cdot D_{2i} (e^J e^L) \\ &= \sum_{i=1}^n \left( \frac{\partial \alpha_I}{\partial \alpha_{2i+1}} \cdot D_{2i} e^J \right) \cdot \alpha_K e^L + (-1)^{|I|} \alpha_I e^J \cdot \sum_{i=1}^n \left( \frac{\partial \alpha_K}{\partial \alpha_{2i+1}} \cdot D_{2i} e^L \right) \\ &\quad + \sum_{i=1}^n \left( \frac{\partial \alpha_I}{\partial \alpha_{2i+1}} \cdot D_{2i} e^L \right) \circ \alpha_K e^J + (-1)^{|I|} \alpha_I e^L \circ \sum_{i=1}^n \left( \frac{\partial \alpha_K}{\partial \alpha_{2i+1}} \cdot D_{2i} e^J \right) \\ &= \Delta(\alpha_I e^J) \circ \alpha_K e^L + (-1)^{|I|} \alpha_I e^J \circ \Delta(\alpha_K e^L) + \Delta(\alpha_I e^L) \circ \alpha_K e^J + (-1)^{|I|} \alpha_K e^L \circ \Delta(\alpha_I e^J). \end{aligned}$$

Comparing with the formula (4-1), we see that the last two terms give the BV Lie bracket. Collecting our results for the loop homology, we get the next Theorem.

**Theorem 4-1.** *In the loop homology  $\mathbb{H}_*(LSU(n+1)) \cong \Lambda_{\mathbb{Z}}(\alpha_3, \alpha_5, \dots, \alpha_{2n+1}) \otimes \mathbb{Z}[e_2, e_4, \dots, e_{2n}]$ , the BV operator  $\Delta$  and BV Lie bracket are given by the following formulae:*

$$\begin{aligned} \Delta(\alpha_I e^J) &= \sum_{\ell=1}^n \frac{\partial \alpha_I}{\partial \alpha_{2\ell+1}} \cdot (D_{2\ell} e^J), \\ (-1)^{|I|} \{\alpha_I e^J, \alpha_K e^L\} &= \Delta(\alpha_I e^L) \circ \alpha_K e^J + (-1)^{|I|} \alpha_I e^L \circ \Delta(\alpha_K e^J). \end{aligned}$$

It is rather surprising that the BV Lie bracket in our loop homology is given precisely by the  $\Delta$ -derivation formula with  $e$ -terms switched. We want to understand what is behind this rather intriguing formula.

So far, we have been examining the Batalin-Vilkovisky algebra structure in loop homology, but loop homology has more structures. For example, it admits so called string operations. To understand the BV bracket, a relevant structure in loop homology not contained in BV algebra structure is the *coalgebra* structure in  $H_*(\Omega SU(n+1))$ . Especially, its primitive elements play an essential role. In Theorem 2-1, we described its coalgebra structure. Primitive elements are spanned by  $s$ -classes  $s_{2\ell}(e_2, e_4, \dots, e_{2\ell})$  for  $1 \leq \ell \leq n$ , defined inductively by Newton's formula:

$$s_{2\ell} - e_2 s_{2\ell-2} + e_4 s_{2\ell-4} - \dots + (-1)^{\ell-1} e_{2\ell-2} s_2 + (-1)^\ell \ell e_{2\ell} = 0, \quad 1 \leq \ell \leq n.$$

Note that  $s_{2\ell} = (-1)^{\ell-1} \ell e_{2\ell} + (\text{decomposables})$ . We extend the ground ring from  $\mathbb{Z}$  to  $\mathbb{Q}$  and we define elements

$$h_{2\ell} = (-1)^{\ell-1} \frac{s_{2\ell}}{\ell} = e_{2\ell} + (\text{decomposables}) \in \mathbb{Q}[e_2, \dots, e_{2n}], \quad 1 \leq \ell \leq n.$$

Then the algebra  $\mathbb{Q}[e_2, e_4, \dots, e_{2n}] = \mathbb{Q}[h_2, h_4, \dots, h_{2n}]$  is primitively generated by  $h_{2\ell}$ 's and we may regard  $\{h_2, h_4, \dots, h_{2n}\}$  as new coordinate functions on  $n$ -dimensional vector space  $\mathbb{Q}^n$ . As such, differential operators  $\partial/\partial h_{2\ell}$  makes sense for functions in  $e_2, \dots, e_{2n}$ .

**Proposition 4-2.** *For each  $1 \leq \ell \leq n$ , the derivation  $\partial/\partial h_{2\ell}$  preserves the integral subring  $\mathbb{Z}[e_2, e_4, \dots, e_{2n}]$ , and is given by*

$$\frac{\partial}{\partial h_{2\ell}} = D_{2\ell} = \sum_{k=\ell}^n e_{2k-2\ell} \frac{\partial}{\partial e_{2k}}.$$

Furthermore, as a function in  $e_2, \dots, e_{2n}$ , all the partial derivatives of  $h_{2\ell}$  have integral coefficients:

$$\frac{\partial h_{2\ell}}{\partial e_{2k}} \in \mathbb{Z}[e_2, e_4, \dots, e_{2n}], \quad 1 \leq k, \ell \leq n.$$

*Proof.* For the first part, we need to show that the functions  $h_{2\ell}(e_2, \dots, e_{2\ell})$  satisfy the system  $D_{2k} h_{2\ell} = \delta_{k,\ell}$  for all  $1 \leq k, \ell \leq n$ . It is possible to verify this directly by induction, but it is rather long and tedious. Here we take the following approach, which gives a quick proof. Recall that the differential operators  $D_{2k}$  are defined in terms of coalgebra map  $\phi$  in  $\mathbb{Z}[e_2, \dots, e_{2n}]$  as in (3-2). Extending  $\phi$  to over  $\mathbb{Q}$ , we have

$$\phi(h_{2\ell}) = 1 \otimes h_{2\ell} + e_2 \otimes D_2(h_{2\ell}) + e_4 \otimes D_4(h_{2\ell}) + \dots + e_{2n} \otimes D_{2n}(h_{2\ell}) + \sum_{I,J} a_{I,J} e^I \otimes e^J.$$

On the other hand, the element  $h_{2\ell}$  is primitive. So we have  $\phi(h_{2\ell}) = 1 \otimes h_{2\ell} + h_{2\ell} \otimes 1 = 1 \otimes h_{2\ell} + e_{2\ell} \otimes 1 + (\text{decomposables}) \otimes 1$ . Comparing these two identities, we see that  $D_{2k}(h_{2\ell}) = \delta_{k,\ell}$  for all  $k, \ell$ .

For the second part, writing the system  $D_{2k}h_{2\ell} = \delta_{k,\ell}$  for a given  $\ell$  in a matrix form, we get

$$\begin{pmatrix} 1 & e_2 & \cdots & e_{2\ell-2} & \cdots & e_{2n-2} \\ & 1 & \cdots & e_{2\ell-4} & \cdots & e_{2n-4} \\ & & \ddots & \vdots & \ddots & \vdots \\ & & & 1 & \cdots & e_{2n-2\ell} \\ & & & & \ddots & \vdots \\ & & & & & 1 \end{pmatrix} \begin{pmatrix} \partial_2 h_{2\ell} \\ \partial_4 h_{2\ell} \\ \vdots \\ \partial_{2\ell} h_{2\ell} \\ \vdots \\ \partial_{2n} h_{2\ell} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ \vdots \\ 0 \end{pmatrix}$$

where in the right hand side, 1 is in the  $\ell$ th position. Since the coefficient matrix is invertible in the integral ring  $\mathbb{Z}[e_2, \dots, e_{2n}]$ , we see that all the partial derivatives of  $h_{2\ell}$  are in this integral ring.  $\square$

To understand BV Lie bracket in the loop homology, we take the point of view that  $\mathbb{H}_*^{\mathbb{Q}} = \mathbb{H}_*(\text{SU}(n+1)) \otimes \mathbb{Q}$  is the ring of functions on the super vector space  $\mathbb{Q}^{n|n}$  of dimension  $n|n$  in which  $h_2, h_4, \dots, h_{2n}$  are even coordinate functions, and  $\alpha_3, \alpha_5, \dots, \alpha_{2n+1}$  are odd coordinate functions. We can do usual differential geometry in this super setting and we can consider vector fields and differentials. In particular, we can consider an odd symplectic form

$$\omega = \sum_{\ell=1}^n d\alpha_{2\ell+1} \wedge dh_{2\ell} \in \bigwedge^2 \left( \bigoplus_{\ell=1}^n \mathbb{H}_*^{\mathbb{Q}} d\alpha_{2\ell+1} \oplus \bigoplus_{\ell=1}^n \mathbb{H}_*^{\mathbb{Q}} dh_{2\ell} \right).$$

The associated Poisson bracket  $[\ , \ ]_{\omega} : \mathbb{H}_*^{\mathbb{Q}} \otimes \mathbb{H}_*^{\mathbb{Q}} \rightarrow \mathbb{H}_*^{\mathbb{Q}}$ , is given by  $[F, G] = X_F G$  for  $F, G \in \mathbb{H}_*^{\mathbb{Q}}$ , where the vector field  $X = X_F$  is defined by the relation  $\iota_X \omega = dF$ . We follow [4] for a sign convention. For example, the exterior derivative is given by  $d = \sum_{\ell} d\alpha_{2\ell+1} \otimes (\partial/\partial\alpha_{2\ell+1}) + \sum_{\ell} dh_{2\ell} \otimes (\partial/\partial h_{2\ell})$ , and for  $F = \alpha_I h^J \in \mathbb{H}_*^{\mathbb{Q}}$  the associated vector field  $X_F$  is given by

$$X_F = \sum_{\ell=1}^n \left( (-1)^{|I|} \frac{\partial F}{\partial \alpha_{2\ell+1}} \cdot \frac{\partial}{\partial h_{2\ell}} + \frac{\partial F}{\partial h_{2\ell}} \cdot \frac{\partial}{\partial \alpha_{2\ell+1}} \right).$$

**Theorem 4-3.** (1) *The differentials  $dh_{2\ell}$  and derivations  $\partial/\partial h_{2\ell}$  for  $1 \leq \ell \leq n$  are defined over  $\mathbb{Z}$ . Consequently, the symplectic form and the Poisson bracket are defined over  $\mathbb{Z}$ :*

$$\omega \in \bigwedge^2 \left( \bigoplus_{\ell=1}^n \mathbb{H}_* d\alpha_{2\ell+1} \oplus \bigoplus_{\ell=1}^n \mathbb{H}_* dh_{2\ell} \right), \quad \text{and} \quad [\ , \ ]_{\omega} : \mathbb{H}_* \otimes \mathbb{H}_* \rightarrow \mathbb{H}_*.$$

(2) *The Poisson bracket  $[\ , \ ]_{\omega}$  defined on  $(\mathbb{H}_*, \omega)$  coincides with the Batalin-Vilkovisky Lie bracket. Namely,*

$$\{\alpha_I e^J, \alpha_K e^L\} = [\alpha_I e^J, \alpha_K e^L]_{\omega},$$

for  $\alpha_I e^J, \alpha_K e^L \in \mathbb{H}_*$ .

*Proof.* The first part is a consequence of Proposition 4-2. For the second part, using  $\partial/\partial h_{2\ell} = D_{2\ell}$ ,

we have

$$\begin{aligned}
[\alpha_I e^J, \alpha_K e^L]_\omega &= X_{\alpha_I e^J}(\alpha_K e^L) \\
&= \sum_{\ell=1}^n \left( (-1)^{|I|} \frac{\partial \alpha_I}{\partial \alpha_{2\ell+1}} \cdot \frac{\partial e^L}{\partial h_{2\ell}} \cdot \alpha_K e^J + \alpha_I e^L \cdot \frac{\partial \alpha_K}{\partial \alpha_{2\ell+1}} \frac{\partial e^J}{\partial h_{2\ell}} \right) \\
&= (-1)^{|I|} \left( \sum_{\ell=1}^n \frac{\partial \alpha_I}{\partial \alpha_{2\ell+1}} \cdot D_{2\ell} e^L \right) \cdot \alpha_K e^J + \alpha_I e^L \cdot \left( \sum_{\ell=1}^n \frac{\partial \alpha_K}{\partial \alpha_{2\ell+1}} \cdot D_{2\ell} e^J \right) \\
&= (-1)^{|I|} \mathbf{\Delta}(\alpha_I e^L) \circ \alpha_K e^J + \alpha_I e^L \circ \mathbf{\Delta}(\alpha_K e^J) \\
&= \{ \alpha_I e^J, \alpha_K e^L \}
\end{aligned}$$

This proves the second part.  $\square$

Let  $\text{Ham}(\mathbb{H}_*^{\mathbb{Q}}) = \{x \in \text{Der}(\mathbb{H}_*^{\mathbb{Q}}) \mid \mathcal{L}_X \omega = 0\}$  be the set of all Hamiltonian derivations, where  $\mathcal{L}_X$  is the Lie derivative by  $X$ . Since  $d[F, G] = \iota_{[X_F, X_G]} \omega$  for  $F, G \in \mathbb{H}_*^{\mathbb{Q}}$ , the correspondence  $\mathbb{H}_*^{\mathbb{Q}} \rightarrow \text{Ham}(\mathbb{H}_*^{\mathbb{Q}})$  given by  $F \mapsto X_F$  is a super Lie algebra homomorphism with kernel  $\mathbb{Q} \cdot 1$ . Since our symplectic manifold  $\mathbb{Q}^{n|n}$  is affine, the above correspondence is onto. This gives a description of  $\mathbb{H}_*^{\mathbb{Q}}$  modulo  $\mathbb{Q} \cdot 1$  as the super Lie algebra of Hamiltonian derivations.

Now consider a vector subspace  $Q$  of  $\mathbb{H}_*^{\mathbb{Q}}$  spanned by quadratic elements  $\alpha_{2\ell+1} h_{2k}$ ,  $\alpha_{2\ell+1} \alpha_{2j+1}$ , and  $h_{2\ell} h_{2k}$  with  $1 \leq \ell, k, j \leq n$ ,  $\ell \neq j$ . The corresponding Hamiltonian derivations are

$$\begin{aligned}
X_{\alpha_{2\ell+1} h_{2k}} &= \alpha_{2\ell+1} \frac{\partial}{\partial \alpha_{2k+1}} - h_{2k} \frac{\partial}{\partial h_{2\ell}}, \\
X_{\alpha_{2\ell+1} \alpha_{2j+1}} &= \alpha_{2k+1} \frac{\partial}{\partial h_{2\ell}} - \alpha_{2\ell+1} \frac{\partial}{\partial h_{2k}}, \\
X_{h_{2\ell} h_{2k}} &= h_{2\ell} \frac{\partial}{\partial \alpha_{2k+1}} + h_{2k} \frac{\partial}{\partial \alpha_{2\ell+1}}.
\end{aligned}$$

Let  $V = \bigoplus_{\ell=1}^n \mathbb{Q} \alpha_{2\ell+1} \oplus \bigoplus_{\ell=1}^n \mathbb{Q} h_{2\ell}$ . The above derivations induce linear maps  $f : V \rightarrow V$  annihilating the odd symplectic form  $\omega$  in the sense that

$$\sum_{\ell=1}^n \{d(f(\alpha_{2\ell+1})) \wedge dh_{2\ell} + d\alpha_{2\ell+1} \wedge d(f(h_{2\ell}))\} = 0,$$

and by counting the dimension, they form a basis for a super Lie algebra  $\pi sp(n|n)$ . This proves Corollary D in the introduction.

We consider the Lie algebra structure of the loop homology of  $\text{SU}(n+1)$  from another point of view. For  $0 \leq p \leq n$ , let  $L(p)$  be the subgroup of the (integral) loop homology  $\mathbb{H}_* = \mathbb{H}_*(\text{LSU}(n+1))$  spanned by elements  $\alpha_I e^J$  with  $|I| = p$  and arbitrary  $J$ . Then  $\mathbb{H}_*$  is a direct sum  $\mathbb{H}_* = \bigoplus_{p=1}^n L(p)$ . In view of Theorem 4-1, BV Lie bracket decreases the number of  $\alpha$ -factors by 1, and we have

$$\{L(p), L(q)\} \subset L(p+q-1), \quad 0 \leq p, q \leq n,$$

where  $L(p) = 0$  when  $p > n$ . These subgroups have the following properties:

- (1)  $L(1)$  is a Lie subalgebra.
- (2)  $L(0)$  is an abelian Lie subalgebra.
- (3)  $L(p)$  is a module over the Lie algebra  $L(1)$  for  $0 \leq p \leq n$ .
- (4)  $L(1)$  is the Lie algebra of all derivations on  $L(0) = \mathbb{Z}[e_2, e_4, \dots, e_{2n}]$ .

Only the last statement needs to be verified. By Theorem 4-1, the action of  $L(1)$  on  $L(0)$  is given by  $\{\alpha_{2\ell+1} e^J, e^L\} = -e^J \cdot D_{2\ell} e^L$ . Thus the effect of  $\alpha_{2\ell+1} e^J$  on  $L(0)$  is the same as the derivation  $-e^J D_{2\ell}$ . Since over  $\mathbb{Z}[e_2, \dots, e_{2n}]$ , operators  $D_2, \dots, D_{2n}$  span the same set of derivations as operators  $\partial/\partial e_2, \dots, \partial/\partial e_{2n}$  do,  $L(1)$  is the Lie algebra of all derivations on  $L(0)$ .

## §5 Batalin-Vilkovisky algebra structure on the loop homology of complex Stiefel manifolds

Let  $V_k(\mathbb{C}^n)$  be the complex Stiefel manifold of complex orthonormal  $k$ -frames in  $\mathbb{C}^n$  with respect to the standard Hermitian pairing in  $\mathbb{C}^n$ . We determine the Batalin-Vilkovisky algebra structure of its loop homology  $\mathbb{H}_*(LV_k(\mathbb{C}^n))$ . It is well known that the complex Stiefel manifolds are homogeneous spaces:

$$V_{n+1-k}(\mathbb{C}^{n+1}) = \mathrm{SU}(n+1)/\mathrm{SU}(k), \quad \text{for } 1 \leq k \leq n, \quad V_{n+1}(\mathbb{C}^{n+1}) = \mathrm{U}(n+1).$$

Note that  $V_n(\mathbb{C}^{n+1}) = \mathrm{SU}(n+1)$ , and  $V_1(\mathbb{C}^{n+1}) = S^{2n+1}$ . Since  $\mathrm{U}(n+1) = \mathrm{SU}(n+1) \times S^1$  as topological spaces, we have an isomorphism as BV algebras:

$$\mathbb{H}_*(L\mathrm{U}(n+1)) \cong \mathbb{H}_*(L\mathrm{SU}(n+1)) \otimes \mathbb{H}_*(LS^1)$$

Thus for the remainder of this section, we consider the BV algebra  $\mathbb{H}_*(LV_{n+1-k}(\mathbb{C}^{n+1}))$  with  $1 \leq k \leq n$ . We will see that its description is very similar to the one for  $\mathbb{H}_*(L\mathrm{SU}(n+1))$ .

First we review the homology and cohomology of the homogeneous space  $\mathrm{SU}(n+1)/\mathrm{SU}(k)$ . We define a set  $\mathcal{S}_k$  of sequences by

$$\mathcal{S}_k = \{(2i_1 + 1, \dots, 2i_r + 1) \mid k \leq i_1 < \dots < i_r \leq n\}.$$

Let  $U = U_k = (2k + 1, 2k + 3, \dots, 2n + 1) \in \mathcal{S}_k$ . Recall that for  $1 \leq i \leq n$  we have cycles  $x_{2i+1} : \Sigma\mathbb{C}P^i \rightarrow \mathrm{SU}(n+1)$  for  $\mathrm{SU}(n+1)$ . For  $I \in \mathcal{S}_k$ , we define a cycle  $x_I$  in  $\mathrm{SU}(n+1)/\mathrm{SU}(k)$  by

$$x_I = x_{2i_1+1} \times \dots \times x_{2i_r+1} : \prod_{t=1}^r \Sigma\mathbb{C}P^{i_t} \rightarrow \mathrm{SU}(n+1) \rightarrow \mathrm{SU}(n+1)/\mathrm{SU}(k).$$

In view of the cell decomposition of  $\mathrm{SU}(n+1)/\mathrm{SU}(k)$  using the product of top cells of  $x_{2i+1}$ 's, the above cycles span the entire homology group. Thus

$$H_*(\mathrm{SU}(n+1)/\mathrm{SU}(k)) \cong \bigoplus_{I \in \mathcal{S}_k} \mathbb{Z}x_I.$$

The element  $x_U$  defines the orientation class. Since  $\mathrm{SU}(n+1)/\mathrm{SU}(k)$  is not an  $H$ -space, there is no Pontrjagin ring structure in this homology group. However, there is a coalgebra structure given by  $\phi(x_I) = \sum_{J \cup K = I} \mathrm{sgn}(J, K)x_J \otimes x_K$ . Primitive elements are spanned by  $x_{2\ell+1}$  for  $k \leq \ell \leq n$ . Let  $y_{2\ell+1}$  be a cohomology element dual to the primitive  $x_{2\ell+1}$  with respect to the basis  $\{x_I\}$ . Then it is straightforward to see that

$$H^*(\mathrm{SU}(n+1)/\mathrm{SU}(k)) \cong \Lambda_{\mathbb{Z}}(y_{2k+1}, y_{2k+3}, \dots, y_{2n+1}).$$

Let  $D : H_*(\mathrm{SU}(n+1)/\mathrm{SU}(k)) \xrightarrow{\cong} H^{d-*}(\mathrm{SU}(n+1)/\mathrm{SU}(k))$  be the Poincaré duality map with respect to the orientation class  $x_U$  so that that  $D(x_I) \cap x_U = x_I$  for all  $I \in \mathcal{S}_k$ , where  $d$  is the dimension of the homogeneous space. As before, let  $\alpha_I$  be the homology class Poincaré dual to  $y_I$  for  $I \in \mathcal{S}_k$ . Then as for the case of  $\mathrm{SU}(n+1)$ , we have

$$(5-1) \quad \alpha_I = (-1)^{\frac{|I|(|I|-1)}{2}} \mathrm{sgn}(I, I^c)x_{I^c},$$

where the complement  $I^c$  is taken in  $U = U_k$ . The intersection ring is then given by

$$\mathbb{H}_*(\mathrm{SU}(n+1)/\mathrm{SU}(k)) \cong \Lambda_{\mathbb{Z}}(\alpha_{2k+1}, \alpha_{2k+3}, \dots, \alpha_{2n+1}),$$

where the generator  $\alpha_{2\ell+1} = (-1)^{\ell-k} x_{U-\{2\ell+1\}}$  has codimension  $2\ell+1$  in  $\mathrm{SU}(n+1)/\mathrm{SU}(k)$ , and has degree  $-2\ell-1$  in the intersection ring.

Next we describe the homology of the based loop space  $\Omega(\mathrm{SU}(n+1)/\mathrm{SU}(k))$ . Using the map  $e_{2\ell}$  for the  $\Omega\mathrm{SU}(n+1)$ , we let  $e_{2\ell} : \mathbb{C}P^\ell \xrightarrow{e_{2\ell}} \Omega\mathrm{SU}(n+1) \rightarrow \Omega(\mathrm{SU}(n+1)/\mathrm{SU}(k))$ , using the same symbol. Then  $x_{2\ell+1} : \Sigma\mathbb{C}P^\ell \rightarrow \mathrm{SU}(n+1)/\mathrm{SU}(k)$  is adjoint to  $e_{2\ell} : \mathbb{C}P^\ell \rightarrow \mathrm{SU}(n+1)/\mathrm{SU}(k)$ . By a standard argument, we have

$$H_*(\Omega(\mathrm{SU}(n+1)/\mathrm{SU}(k))) \cong \mathbb{Z}[e_{2k}, e_{2k+2}, \dots, e_{2n}],$$

and the map  $p_* : H_*(\Omega(\mathrm{SU}(n+1))) \rightarrow H_*(\Omega(\mathrm{SU}(n+1)/\mathrm{SU}(k)))$  is a quotienting map by an ideal  $(e_2, e_4, \dots, e_{2k-2})$ . Since the above map  $p_*$  is a Hopf algebra map, all the Hopf algebra structure of  $H_*(\Omega(\mathrm{SU}(n+1)/\mathrm{SU}(k)))$  is obtained by setting  $e_2 = \dots = e_{2k-2} = 0$  in the structure maps of  $H_*(\Omega(\mathrm{SU}(n+1)))$ .

Next, we consider the fibre bundle

$$\Omega(\mathrm{SU}(n+1)/\mathrm{SU}(k)) \rightarrow L(\mathrm{SU}(n+1)/\mathrm{SU}(k)) \rightarrow \mathrm{SU}(n+1)/\mathrm{SU}(k).$$

The associated homology and loop homology spectral sequences

$$\begin{aligned} E_{*,*}^2 &= H_*(\mathrm{SU}(n+1)/\mathrm{SU}(k)) \otimes H_*(\Omega(\mathrm{SU}(n+1)/\mathrm{SU}(k))) \Longrightarrow H_*(L(\mathrm{SU}(n+1)/\mathrm{SU}(k))), \\ E_{*,*}^2 &= \mathbb{H}_*(\mathrm{SU}(n+1)/\mathrm{SU}(k)) \otimes H_*(\Omega(\mathrm{SU}(n+1)/\mathrm{SU}(k))) \Longrightarrow \mathbb{H}_*(L(\mathrm{SU}(n+1)/\mathrm{SU}(k))), \end{aligned}$$

collapse by a standard argument, although the fibre bundle itself is not a product. Thus

$$H_*(L(\mathrm{SU}(n+1)/\mathrm{SU}(k))) \cong \bigoplus_{I,J} \mathbb{Z} x_I e^J,$$

where  $I \in \mathcal{S}_k$  and  $J$  runs over all sequences  $J = (j_k, \dots, j_n)$  with nonnegative integer entries. This homology is only a coalgebra and does not have a ring structure. We can think of the element  $x_I e^J$  as the image of the element of the same name in  $H_*(L\mathrm{SU}(n+1))$  under the projection map  $p_*$ . On the other hand, the loop homology algebra is given by

$$(5-2) \quad \mathbb{H}_*(L(\mathrm{SU}(n+1)/\mathrm{SU}(k))) \cong \Lambda_{\mathbb{Z}}(\alpha_{2k+1}, \dots, \alpha_{2n+1}) \otimes \mathbb{Z}[e_{2k}, \dots, e_{2n}],$$

but there is no coalgebra structure. Note that the map  $p : \mathrm{SU}(n+1) \rightarrow \mathrm{SU}(n+1)/\mathrm{SU}(k)$  does not induce a homomorphism in loop homology.

To determine the BV operator  $\Delta$  in loop homology, we first determine homological  $S^1$ -action in ordinary homology, and then translate it into loop homology using (5-1). Since  $S^1$  action commutes with the projection map  $p : L\mathrm{SU}(n+1) \rightarrow L(\mathrm{SU}(n+1)/\mathrm{SU}(k))$ , the homological  $S^1$  action in  $H_*(L(\mathrm{SU}(n+1)/\mathrm{SU}(k)))$  is obtained by projection from the formula of  $\Delta$  in  $H_*(\mathrm{SU}(n+1))$ . Thus for an element  $x_I e^J \in H_*(L(\mathrm{SU}(n+1)/\mathrm{SU}(k)))$ , we lift it to an element  $x_I e^J$  in  $H_*(\mathrm{SU}(n+1))$ , apply  $\Delta$ , then project down by  $p_*$ :

$$\Delta(x_I e^J) = p_*(\Delta(x_I e^J)) = \sum_{\ell=k}^n p_*(x_{2\ell+1} \cdot x_I) \cdot D_{2\ell}^{(k)} e^J,$$

where  $D_{2\ell}^{(k)}$  is the differential operator obtained from  $D_{2\ell}$  by setting  $e_2 = \dots = e_{2k-2} = 0$ . We can rewrite this formula in terms of elements  $\alpha_{2\ell+1}$ 's. As in (2-4) for  $\mathrm{SU}(n+1)$  case, we can show  $p_*(x_{2\ell+1} \cdot \alpha_I) = \partial\alpha_I/\partial\alpha_{2\ell+1}$ . Thus the BV structure of the loop homology of the complex Stiefel manifold  $V_{n+1-k}(\mathbb{C}^{n+1}) = \mathrm{SU}(n+1)/\mathrm{SU}(k)$  is formally entirely analogous to the BV structure of  $\mathbb{H}_*(L\mathrm{SU}(n+1))$ , and we obtain the following statement.

**Theorem 5-1.** *In the loop homology  $\mathbb{H}_*(L(\mathrm{SU}(n+1)/\mathrm{SU}(k)))$  of the complex Stiefel manifold given in (5-2), the BV operator  $\Delta$  is given by*

$$\Delta(\alpha_I e^J) = \sum_{\ell=k}^n \frac{\partial \alpha_I}{\partial \alpha_{2\ell+1}} \cdot D_{2\ell}^{(k)} e^J,$$

where differential operators  $D_{2\ell}^{(k)}$  on the polynomial ring  $\mathbb{Z}[e_{2k}, \dots, e_{2n}]$  is given by

$$D_{2\ell}^{(k)} = \begin{cases} \frac{\partial}{\partial e_{2\ell}} + \sum_{j=k}^{n-\ell} e_{2j} \frac{\partial}{\partial e_{2\ell+2j}}, & \text{if } \ell \leq n-k, \\ \frac{\partial}{\partial e_{2\ell}} & \text{if } \ell > n-k. \end{cases}$$

The BV Lie bracket is given by

$$\begin{aligned} \{\alpha_I e^J, \alpha_K e^L\} &= (-1)^{|I|} \Delta(\alpha_I e^L) \circ \alpha_K e^J + \alpha_K e^L \circ \Delta(\alpha_I e^J) \\ &= [\alpha_I e^J, \alpha_K e^L]_\omega, \end{aligned}$$

where  $\omega = \sum_{\ell=k}^n d\alpha_{2\ell+1} \wedge dh_{2\ell}^{(k)}$  is the odd symplectic form associated with primitive elements  $h_{2\ell}^{(k)}$  for  $k \leq \ell \leq n$  in  $\mathbb{Q}[e_{2k}, \dots, e_{2n}]$ , and  $[\cdot, \cdot]_\omega$  is the Poisson bracket associated to  $\omega$ .

The proof is entirely analogous to the  $\mathrm{SU}(n+1)$  case.

As a final result, we specialize to the case  $V_1(\mathbb{C}^{n+1}) = \mathrm{SU}(n+1)/\mathrm{SU}(n) = S^{2n+1}$ .

**Theorem 5-2.** *In the loop homology of an odd dimensional sphere*

$$\mathbb{H}_*(LS^{2n+1}) \cong \Lambda(\alpha) \otimes \mathbb{Z}[h],$$

where  $|\alpha| = -2n - 1$  and  $|h| = 2n$ , the BV operator is given by

$$\Delta(h^\ell) = 0, \quad \Delta(\alpha h^\ell) = \ell h^{\ell-1}, \quad \ell \geq 0.$$

As a BV Lie algebra,  $\mathbb{H}_{\mathrm{even}} = \mathbb{Z}[h]$  is an abelian Lie subalgebra, and  $\mathbb{H}_{\mathrm{odd}} = \alpha\mathbb{Z}[h]$  is the Lie algebra of derivations on the polynomial ring  $\mathbb{Z}[h]$ . Namely,

$$\{\alpha h^{k+1}, \alpha h^{\ell+1}\} = (k - \ell)\alpha h^{k+\ell+1}, \quad \{\alpha h^{k+1}, h^m\} = -m h^{k+m}, \quad k, \ell \geq -1, m \geq 0.$$

**Remark.** If we let  $L(k) = \alpha h^{k+1}$  for  $k \geq -1$ , then these elements satisfy

$$\{L(k), L(\ell)\} = (k - \ell)L(k + \ell), \quad k, \ell \geq -1.$$

This is the commutation relation of the positive half of the Virasoro algebra. Its generalization is the Lie algebra  $L(1)$  acting as the Lie algebra of derivations on the polynomial algebra  $\mathbb{Z}[e_2, e_4, \dots, e_{2n}]$  discussed at the end of section 4.

The BV algebra in Theorem 5-2 is a building block of the loop homology of  $\mathrm{SU}(n+1)$ , as we saw in Corollary C.

## REFERENCES

- [1] M. Chas and D. Sullivan, *String topology*, arXiv:math.GT/9911159.
- [2] R.L. Cohen, J.D.S. Jones, *A homotopy theoretic realization of string topology*, Math. Ann **324** (2002), 773–798.
- [3] R.L. Cohen, J.D.S. Jones, and J. Yan, *The loop homology algebra of spheres and projective spaces*, Progr. Math. **215** (2004), 77–92.
- [4] P. Deligne, et al., *Quantum Fields and Strings: A Course for Mathematicians*, Amer. Math. Society, RI, 1999.
- [5] V. Kac, *Lie superalgebras*, Adv. in Math. **26** (1977), 8–96.
- [6] M. Mimura and H. Toda, *Topology of Lie Groups I, II*, Translations of Math. Monographs, 91, Amer. Math. Society, RI, 1991.