

Curves and vector bundles on quartic threefolds

Enrique Arrondo and Carlo G. Madonna *

Abstract

In this paper we study arithmetically Cohen-Macaulay (ACM for short) vector bundles \mathcal{E} of rank $k \geq 3$ on hypersurfaces $X_r \subset \mathbf{P}^4$ of degree $r \geq 1$. We consider here mainly the case of degree $r = 4$, which is the first unknown case in literature. Under some natural conditions for the bundle \mathcal{E} we derive a list of possible Chern classes (c_1, c_2, c_3) which may arise in the cases of rank $k = 3$ and $k = 4$, when $r = 4$ and we give several examples.

1 Introduction

Let $X = X_r \subset \mathbf{P}^4$ be a smooth hypersurface of degree $r \geq 1$ and let H be the class of a hyperplane section. It is well known that

$$\mathrm{Pic}(X_r) \cong \mathbf{Z}[H] \cong \mathbf{Z} \quad (1.1)$$

and hence we may identify any line bundle \mathcal{L} on X_r with the sheaf $\mathcal{O}_{X_r}(nH) \cong \mathcal{O}_{X_r}(n)$ for some $n \in \mathbf{Z}$. We recall the following:

Definition 1.1. *Let \mathcal{E} be a rank k vector bundle on X_r , $r \geq 1$. We call \mathcal{E} arithmetically Cohen-Macaulay (ACM for short) if $h^i(X_r, \mathcal{E}(n)) = 0$ for all $i = 1, 2$ and $n \in \mathbf{Z}$, where $\mathcal{E}(n) := \mathcal{E} \otimes \mathcal{O}_{X_r}(n)$, under the identification of (1.1).*

In a previous work [19] the second author showed that a rank two ACM vector bundle on X_r splits as a direct sum of two line bundles with only few possible exceptions. Specifically, for each r , all the possible first Chern classes of such an indecomposable bundle (up to a twist with a line bundle) were given.

For low values of r , i.e. $r = 1, 2, 3, 4$, a complete classification of indecomposable rank two ACM vector bundles is known. We refer the interested reader to [15] [27] [4] [20] for more details on these cases. When $r = 5$ the possible Chern classes of such an indecomposable bundle were found in [21] while, when $r \geq 6$ and X_r is general, the results of [8] (for the case $r = 6$) and of [18] (for any $r \geq 6$) ensure the non existence of such bundles. Similar results were obtained for more general threefolds and in particular a complete classification of rank two ACM vector bundles on prime Fano threefolds of index 1 was given

*The first author was supported in part by the project BFM2003-03971/MATE funded by the Spanish MCYT. The second author was supported in part by the projects I3P-CSIC and MTM2007-67623 funded by the Spanish MEC.

in [12] and [5]. Also in some cases the corresponding moduli spaces are described (see [16] for a comprehensive account of the known results).

On the other hand, for the higher rank case very little is known even in the cases of low degree hypersurfaces. When the rank is bigger than or equal to 3, it is known that there are no indecomposable ACM vector bundles if $r = 1, 2$, while as far as we know, only the cubic case was considered in [4] under the following natural assumption (which we generalize to arbitrary r):

Definition 1.2. *We say that a rank $k \geq 2$ vector bundle on $X_r \subset \mathbf{P}^4$ satisfies condition \star if the following conditions hold:*

- (i) $h^0 \mathcal{E}(-1) = 0$ and $h^0 \mathcal{E} \geq k$;
- (ii) there exist $k - 1$ global sections of \mathcal{E} whose dependency locus has codimension equal to two;
- (iii) \mathcal{E} has not trivial summands.

In this paper we will consider mainly the cases of low rank, (precisely $k = 3, 4$) when $r = 4$, $X = X_4$ is general, and the above condition \star is satisfied. Notice, it is easy to produce examples of higher rank (> 3) ACM vector bundles by using extension classes. Specifically, starting with a pair $(\mathcal{E}', \mathcal{E}'')$ of rank k and k' ACM bundles on X_r , one may always consider extensions

$$0 \rightarrow \mathcal{E}' \rightarrow \mathcal{E} \rightarrow \mathcal{E}'' \rightarrow 0 \quad (1.2)$$

to get a rank $k' + k''$ ACM bundle on X_r . Of course, since line bundles on X_r are ACM, to get such non trivial extension classes it is necessary to have $k, k' \geq 2$. We will give several examples of rank four ACM bundles satisfying condition \star when $r = 4$ obtained in this way in Section 3.

Since such a construction was already used for the case $r = 3$ in [4] and for the case $r = 5$ in [22], one could state the following:

Question 1.3. *Let \mathcal{E} be a rank four ACM vector bundle satisfying condition \star on a general hypersurface $X_r \subset \mathbf{P}^4$. Is \mathcal{E} necessarily obtained as an extension class of a pair of rank two ACM bundles on X_r , as in (1.2)?*

We give here a negative answer to the above question when $r = 4$ and X_4 is general (see Proposition 4.4). We notice that Question 1.3 still remains open when $r = 3$. Moreover when $r = 6$ we have not examples at all of rank 4 ACM bundles, while when $r = 5$ we know actually only one example, given in [22], which satisfies condition \star .

Turning to the case of rank $k = 3$, we want to answer the following question:

Question 1.4. *Are there rank three ACM bundles satisfying condition \star on a general hypersurface $X_r \subset \mathbf{P}^4$ of degree $r \geq 4$?*

Since in [4] an affirmative answer (see Proposition 4.9) was already given for the case $r = 3$ we consider the case $r = 4$ here. We give an affirmative answer to this last question when $r = 4$. We notice that Question 1.4 remains open when $r \geq 5$. To answer the previous questions, we derive the possible first Chern

class of an ACM bundle of rank $k \geq 3$ satisfying condition \star on any smooth hypersurface of degree $r \geq 3$. Then we consider in detail the cases of low rank, i.e. $k = 3$ and $k = 4$, when $r = 4$, and we go on by a case by case analysis. In this direction our main result is the following:

Theorem 1.5. *Let \mathcal{E} be an ACM vector bundle of rank $k \in \{3, 4\}$ on a general quartic threefold $X_4 \subset \mathbf{P}^4$ which satisfies condition \star . Then the possible Chern classes $c_i = c_i(\mathcal{E})$ and the genus $g = g(C)$ of the curve C , dependency locus of $k - 1$ global section of \mathcal{E} , are given in the following table:*

k	c_1	$c_2 \in$	c_3	g
3	1	5	2	2
3	2	[8, 11]	$c_2 - 6$	$c_2 - 2$
3	3	[17, 18]	$2c_2 - 26$	$2c_2 - 12$
3	4	[27, 28]	$3c_2 - 66$	$3c_2 - 32$
4	1	6	4	3
4	2	[8, 12]	$c_2 - 4$	$c_2 - 1$
4	3	[16, 22]	$2c_2 - 24$	$2c_2 - 11$
4	4	[28, 32]	$3c_2 - 64$	$3c_2 - 31$
4	5	[44, 46]	$4c_2 - 132$	$4c_2 - 65$
4	6	64	84	203

Moreover there are examples of bundles with invariants $(k; c_1, c_2, c_3)$ for any of the following quadruples:

$$(4; 1, 6, 4), (4; 2, \alpha, \alpha - 4), (4; 3, \beta, 2\beta - 24), (4; 4, \gamma, 3\gamma - 64), \\ (4; 5, 46, 52), (4; 6, 64, 84), (3; 1, 5, 2),$$

where $\alpha \in \{10, 11, 12\}$, $\beta \in \{19, 20\}$, and $\gamma \in \{29, 30, 32\}$.

In the case of the rank four, almost all the examples are given by considering extension classes of rank two ACM bundles. It is not so for the case $(4; 1, 6, 4)$, which is constructed by means of the Hartshorne-Serre correspondence (see Proposition 2.1) starting from a suitable smooth projectively normal curve in X_4 and it is not given by any extension class of any two rank two ACM bundles on X_4 . The example for the rank three case is given in similar way, by showing the existence of the corresponding curve with prescribed invariants and then applying the above mentioned Hartshorne-Serre correspondence.

The paper is organized as follows. In Section 2 we recall some known facts and generalities that we need in the paper. In Section 3 (see Lemma 3.4), we find conditions such that the direct sum (or more generally any extension class) of ACM vector bundles satisfying condition \star is still an ACM vector bundle satisfying condition \star , and produce from this several examples of rank four bundles on X_4 (precisely all but one in the list of Theorem 1.5). In Section 4 we use the Hartshorne-Serre correspondence (see Proposition 2.1) between curves and vector bundles to give a characterization (see Proposition 4.1) of when the vector bundle obtained from a curve is ACM and satisfies condition \star ; using

this, we construct the example of rank three and the remaining one of rank four of ACM vector bundles satisfying condition \star on X_4 of the list of Theorem 1.5. Finally, in section 5 we give restrictions for the Chern classes of ACM vector bundles on X_r satisfying condition \star , which when particularized for $r = 4$ and rank $k = 3, 4$ yields the list given in Theorem 1.5 of all possible Chern classes in these ranks.

The preliminary draft of this paper was completed during the second author stay in the Departamento de Álgebra of University Complutense in Madrid, supported by a grant “Mensilità per l'estero” from the “Istituto Nazionale di Alta Matematica F. Severi” (10/2005-03/2006).

2 Preliminaries and basic facts

In this paper we work over the field of complex numbers \mathbf{C} . Let us consider a smooth hypersurface $X_r \subset \mathbf{P}^4$ of degree $r \geq 1$ in the complex 4-dimensional projective space \mathbf{P}^4 . Then, as recalled in the introduction $\text{Pic}(X_r) \cong \mathbf{Z}[H]$ where H is the class of a hyperplane section, and the canonical divisor of X_r is $K_{X_r} = (r - 5)H$.

If P denotes the class of a point and L the class of a line, then the intersection products on X_r are given by $H^3 = rP = r$, $HL = 1P = 1$, $H^2 = rL$. Let \mathcal{E} be a rank k vector bundle on X_r . We identify the first Chern class $c_1(\mathcal{E})$ of \mathcal{E} with the integer number c_1 which corresponds to $c_1(\mathcal{E})$ under the above isomorphism, i.e. $c_1(\mathcal{E}) = c_1H = c_1$. In a similar way we identify $c_2(\mathcal{E}) = \deg c_2(\mathcal{E}) = c_2$ and $c_3(\mathcal{E}) = c_3P = c_3$. Under this identification to any rank k vector bundle \mathcal{E} on X_r corresponds a quadruple $(k; c_1, c_2, c_3) \in \mathbf{N} \times \mathbf{Z}^3$. In this language, our main result gives all the possible quadruples of a rank three and four ACM vector bundle on a smooth quartic threefold which could arise under the condition \star of Definition 1.2.

For further computations we write down explicitly the Chern classes of the bundle $\mathcal{E}(n) = \mathcal{E} \otimes \mathcal{O}_{X_r}(n)$ in the following equations:

$$\begin{aligned} c_1(\mathcal{E}(n)) &= c_1 + kn \\ c_2(\mathcal{E}(n)) &= c_2 + r(k-1)nc_1 + r \binom{k}{2} n^2 = \\ &= c_2 + rn(k-1) \left(c_1 + \frac{1}{2}nk \right) \\ c_3(\mathcal{E}(n)) &= c_3 + (k-2)nc_2 + r \binom{k-1}{2} n^2 c_1 + r \binom{k}{3} n^3 = \\ &= c_3 + (k-2)n \left(c_2 + \frac{1}{2}(k-1)nrc_1 + \frac{1}{6}rn^2k(k-1) \right) \end{aligned}$$

In the sequel, to perform some computations, we will frequently use the following version of Riemann-Roch theorem for vector bundles:

Hirzebruch-Riemann-Roch Theorem. *If \mathcal{E} is a rank k vector bundle on a smooth hypersurface $X_r \subset \mathbf{P}^4$ of degree $r \geq 1$ with Chern classes $c_i(\mathcal{E}) = c_i \in \mathbf{Z}$*

for $i = 1, 2, 3$, then

$$\begin{aligned} \chi(\mathcal{E}) &= \frac{1}{6}rc_1^3 - \frac{1}{2}c_1c_2 + \frac{1}{2}c_3 + \frac{1}{4}rc_1^2(5-r) - \frac{1}{2}c_2(5-r) + \\ &+ \frac{1}{12}((r-5)^2 + (10-5r+r^2))c_1r + \frac{1}{24}rk(5-r)(10-5r+r^2) \end{aligned} \quad (2.1)$$

In particular

$$\begin{aligned} \chi(\mathcal{O}_{X_r}(a)) &= \frac{1}{6}a^3r + \frac{1}{4}a^2r(5-r) + \frac{1}{12}ar((r-5)^2 + \\ &+ (10-5r+r^2)) + \frac{r}{24}(5-r)(10-5r+r^2) \end{aligned} \quad (2.2)$$

Recall that a vector bundle \mathcal{E} is “decomposable” if there exist \mathcal{E}' and \mathcal{E}'' such that $\mathcal{E} \cong \mathcal{E}' \oplus \mathcal{E}''$. Otherwise \mathcal{E} is “undecomposable”. By the Serre’s duality, \mathcal{E} is ACM if and only if

$$h^1\mathcal{E}(n) = h^1\mathcal{E}^\vee(-n+r-5) = 0 \quad \forall n \in \mathbf{Z}$$

which is equivalent to the condition $h^1\mathcal{E}(n) = h^1\mathcal{E}^\vee(n) = 0 \quad \forall n \in \mathbf{Z}$.

From now on we will consider mainly vector bundles \mathcal{E} of rank $k \geq 3$ for which the condition \star (see Definition 1.2) is satisfied. If $h^0\mathcal{E}(-1) = 0$ and $h^0\mathcal{E} > 0$ we will say that \mathcal{E} is “normalized”. It is always possible to assume this condition since we may replace \mathcal{E} with $\mathcal{E}(-b)$, where $b = b(\mathcal{E})$ is defined as

$$b(\mathcal{E}) = b = \max\{n \in \mathbf{Z} : h^0\mathcal{E}(-n) = 0\}. \quad (2.3)$$

We recall also the notion of stability that we will use in the sequel. A rank k vector bundle \mathcal{E} on X_r is “stable” (resp. “semistable”) if for any subbundle $\mathcal{F} \subset \mathcal{E}$ of rank $m < k$ we have $c_1(\mathcal{F})/m < c_1(\mathcal{E})/k$ (resp. \leq). In particular if $b(\mathcal{E}) = b$ is defined as in (2.3) we have $kb(\mathcal{E}) - c_1(\mathcal{E}) < 0$ if \mathcal{E} is stable (resp. \leq). Indeed, any global section of $H^0\mathcal{E}(-b)$ gives rise to a morphism $\mathcal{O}_{X_r} \rightarrow \mathcal{E}(-b)$ and hence if \mathcal{E} is stable then $0 < c_1(\mathcal{E}(-b))/k = (c_1(\mathcal{E}) - kb)/k$ and hence $kb - c_1 < 0$. In particular if \mathcal{E} is normalized and stable then $c_1 > 0$. When $k = 2$ the condition $kb(\mathcal{E}) - c_1(\mathcal{E}) \leq 0$ is in fact equivalent to the semistability of \mathcal{E} . If $k \geq 3$ this is not an equivalence any more.

We can use condition (ii) in Definition 1.2 to translate properties of vector bundles satisfying \star to properties of curves in X_r . Indeed, for a vector bundle \mathcal{E} satisfying condition \star , the choice of $k-1$ global sections whose dependency locus is a curve $C \subset X_r$ yields the fundamental exact sequence:

$$0 \rightarrow \mathcal{O}_{X_r}^{k-1} \rightarrow \mathcal{E} \rightarrow \mathcal{J}_C(c_1) \rightarrow 0 \quad (2.4)$$

where \mathcal{J}_C is the ideal sheaf of $C \subset X_r$. When dualizing (2.4), we get the exact sequence

$$0 \rightarrow \mathcal{O}_{X_r}(-c_1) \rightarrow \mathcal{E}^\vee \rightarrow \omega_C^{k-1} \rightarrow \omega_C(5-r-c_1) \rightarrow 0 \quad (2.5)$$

where $\omega_C \cong \mathcal{E}xt^2(\mathcal{O}_C, \omega_{X_r})$ is the dualizing sheaf of C . This implies, in particular, that $\omega_C(5-r-c_1)$ is generated by $k-1$ global sections. Next we recall the

generalized Hartshorne-Serre correspondence (see [28] [3]), which states that is possible to reverse this process, in the sense that one recovers the vector bundle from the surjection in (2.5):

Proposition 2.1. *Let $C \subset X_r$ be a locally complete intersection curve and assume that, for some $c_1 \in \mathbf{Z}$, $\omega_C(5 - r - c_1)$ has $k - 1$ generating global sections. Then, there exists a rank k vector bundle \mathcal{E} on X_r fitting in the exact sequence (2.4) and the surjection of (2.5) corresponds to the choice of the $k - 1$ global sections of $\omega_C(5 - r - c_1)$. In particular $c_1(\mathcal{E}) = c_1$, $c_2(\mathcal{E}) = \deg C$ and C is the dependency locus of the $k - 1$ global sections of \mathcal{E} given by (2.4).*

This gives, any time we have an exact sequence like (2.4), the relation between the two first Chern classes of \mathcal{E} and the degree of C . Moreover, the third Chern class of \mathcal{E} is determined by the (arithmetic) genus of the curve C :

Corollary 2.2. *If $C \subset X_r$ is a locally complete intersection curve associated to a vector bundle on X_r of rank $k \geq 2$ as above, then*

$$g(C) = -\frac{5}{2}c_2 + \frac{1}{2}c_1c_2 + \frac{1}{2}c_3 + \frac{25}{12}r + \frac{1}{2}rc_2 - \frac{35}{24}r^2 + \frac{5}{12}r^3 - \frac{1}{24}r^4$$

where $c_i = c_i(\mathcal{E})$, $i = 1, 2, 3$. In particular when $r = 4$ we have

$$g(C) = 1 + \frac{1}{2}c_1c_2 - \frac{1}{2}c_2 + \frac{1}{2}c_3. \quad (2.6)$$

Proof. It follows from the equalities $g(C) = 1 - \chi(\mathcal{O}_C) = 1 - \chi(\mathcal{O}_{X_r}) + \chi(\mathcal{J}_C)$, $\chi(\mathcal{J}_C) = \chi(\mathcal{E}(-c_1)) - \chi(\mathcal{O}_{X_r}^{k-1}(-c_1))$ (by the exact sequence (2.4)) and the Riemann–Roch theorem. \square

Having in mind Hartshorne-Serre correspondence, we will frequently consider pairs

$$(\mathcal{E}, C) \quad (2.7)$$

given by a vector bundle \mathcal{E} of rank $k \geq 3$ which satisfies condition \star and a curve $C \subset X_r$ dependency locus of $k - 1$ global sections of \mathcal{E} . We will also say in this case that the pair (\mathcal{E}, C) is a *ACM pair* and satisfies condition \star (in Proposition 4.1 we will give a criterion for this property in terms of the curve C).

We recall the following generalization of Clifford’s theorem (observe that in our situation C is connected because the fact that \mathcal{E} is ACM implies from (2.4) the vanishing of $H^1\mathcal{J}_C$):

Theorem 2.3. *Let C be a connected locally complete intersection curve, and let \mathcal{L} be a line bundle on C such that $H^0\mathcal{L} \neq 0$ and $H^1\mathcal{L} \neq 0$. Then $2(h^0\mathcal{L} - 1) \leq \deg \mathcal{L}$.*

Proof. The same proof as in [11] Theorem A works in this case, since the irreducibility assumed there is not used to prove the inequality. In fact, we still have a nondegenerate bilinear map $H^0\mathcal{L} \otimes \text{Hom}(\mathcal{L}, \omega_C) \rightarrow H^0\omega_C$, so that the result follows, as in [11], by the so-called bilinear lemma, Serre’s duality and Riemann-Roch theorem. The connectedness of C is used to conclude that the dimension $h^0\omega_C = h^1\mathcal{O}_C$ is $g(C)$ because $h^0\mathcal{O}_C = 1$. \square

Let us finish this section by recalling the classification of indecomposable rank two ACM bundles on quartics X_4 obtained in [20] and on cubics X_3 obtained in [4].

Proposition 2.4. *Let \mathcal{E} be a normalized and indecomposable rank two ACM bundle on a smooth hypersurface $X_r \subset \mathbf{P}^4$ with Chern classes $c_i(\mathcal{E}) = c_i$, $i = 1, 2$.*

1. *If $r = 4$ then $(c_1, c_2) \in \{(-1, 1), (0, 2), (1, 3), (1, 4), (1, 5), (2, 8), (3, 14)\}$;*
2. *if $r = 3$ then $(c_1, c_2) = \{(0, 1), (1, 2), (2, 5)\}$.*

Moreover all the cases arise.

Remark 2.5. In particular in all the above cases condition \star is satisfied with the only exceptions of cases $(c_1, c_2) \in \{(-1, 1), (0, 2), (1, 5)\}$ when $r = 4$ and $(c_1, c_2) = (0, 1)$ when $r = 3$. Moreover, as shown in [20], it also holds that the vector bundles on X_4 with $(c_1, c_2) = (3, 14)$ and the general one with $(c_1, c_2) = (2, 8)$ are generated by their global sections.

3 Constructing bundles from extensions

We prove in this section that direct sums (and more generally extensions) of vector bundles satisfying condition \star are still ACM vector bundles satisfying condition \star . We will end by producing in this way examples of rank four ACM vector bundles satisfying condition \star on X_4 . Observe first that an ACM vector bundle \mathcal{E} satisfying condition \star cannot decompose as $\mathcal{E} \cong \mathcal{E}' \oplus \mathcal{O}_{X_r}(a)$ where \mathcal{E}' is a vector bundle of rank $k - 1$. Indeed, the possibility $a > 0$ is excluded by condition (i), the possibility $a = 0$ is excluded by condition (iii) and the possibility $a < 0$ is excluded by condition (ii), since the dependency locus of $k - 1$ sections of \mathcal{E} would be the dependency locus of $k - 1$ sections of \mathcal{E}' (hence of expected codimension one).

Of course it can be possible to have direct sums $\mathcal{E} \cong \mathcal{E}' \oplus \mathcal{E}''$ with $rk(\mathcal{E}') < k$ and $rk(\mathcal{E}'') \geq 2$. In fact, we are going to see that the second part in condition (i) implies that condition \star is preserved by direct sums (and more generally by extensions). To show this, we start with a standard fact:

Lemma 3.1. *Let \mathcal{E}' and \mathcal{E}'' be two vector bundles on X_r of rank k' and k'' respectively. Let $s'_1, \dots, s'_{k'}$ be linearly independent sections of \mathcal{E}' and let $s''_1, \dots, s''_{k''}$ be linearly independent sections of \mathcal{E}'' . Assume that the dependency locus of $s'_1, \dots, s'_{k'-1}$ is a curve C' and the dependency locus of $s''_2, \dots, s''_{k''}$ is a curve C'' . Then the dependency locus of the sections*

$$(s'_1, 0), \dots, (s'_{k'-1}, 0), (s'_{k'}, s''_1), (0, s''_2), \dots, (0, s''_{k''})$$

of $\mathcal{E}' \oplus \mathcal{E}''$ is $C' \cup C'' \cup V(s'_1 \wedge \dots \wedge s'_{k'}, s''_1 \wedge \dots \wedge s''_{k''})$, where $s'_1 \wedge \dots \wedge s'_{k'}$ is interpreted as a section of the line bundle $\wedge^{k'} \mathcal{E}'$ and $s''_1 \wedge \dots \wedge s''_{k''}$ as a section of $\wedge^{k''} \mathcal{E}''$.

Proof. It is enough to check the statement locally. We can thus restrict ourselves to an open subset of X_r on which the vector bundles trivialize. If on that open set the section s'_i is represented by the functions $f'_{i1}, \dots, f'_{ik'}$ and the section s''_j is represented by $f''_{j1}, \dots, f''_{jk''}$, then the dependency locus of $(s'_1, 0), \dots, (s'_{k'-1}, 0), (s'_{k'}, s''_1), (0, s''_2), \dots, (0, s''_{k''})$ is defined by the vanishing of the maximal minors of the $(k' + k'' - 1) \times (k' + k'')$ matrix

$$\begin{pmatrix} f'_{11} & \cdots & f'_{1k'} & 0 & \cdots & 0 \\ \vdots & & \vdots & \vdots & & \vdots \\ f'_{k'-1,1} & \cdots & f'_{k'-1,k'} & 0 & \cdots & 0 \\ f'_{k'1} & \cdots & f'_{k'k'} & f''_{11} & \cdots & f''_{1k''} \\ 0 & \cdots & 0 & f''_{21} & \cdots & f''_{2k''} \\ \vdots & & \vdots & \vdots & & \vdots \\ 0 & \cdots & 0 & f''_{k''1} & \cdots & f''_{k''k''} \end{pmatrix}$$

These $k' + k'' - 1$ minors take the form

$$\begin{vmatrix} f'_{11} & \cdots & f'_{1,i-1} & f'_{1,i+1} & \cdots & f'_{1k'} \\ \vdots & & \vdots & \vdots & & \vdots \\ f'_{k'-1,1} & \cdots & f'_{k'-1,i-1} & f'_{k'-1,i+1} & \cdots & f'_{k'-1,k'} \end{vmatrix} \begin{vmatrix} f''_{11} & \cdots & f''_{1k''} \\ \vdots & & \vdots \\ f''_{k''1} & \cdots & f''_{k''k''} \end{vmatrix}, \quad i = 1, \dots, k'$$

$$\begin{vmatrix} f'_{11} & \cdots & f'_{1k'} \\ \vdots & & \vdots \\ f'_{k'1} & \cdots & f'_{k'k'} \end{vmatrix} \begin{vmatrix} f''_{21} & \cdots & f''_{2,j-1} & f''_{2,j+1} & \cdots & f''_{2k''} \\ \vdots & & \vdots & \vdots & & \vdots \\ f''_{k''1} & \cdots & f''_{k'',j-1} & f''_{k'',j+1} & \cdots & f''_{k''k''} \end{vmatrix}, \quad j = 1, \dots, k''$$

Since the curve C' is locally defined by the minors

$$\begin{vmatrix} f'_{11} & \cdots & f'_{1,i-1} & f'_{1,i+1} & \cdots & f'_{1k'} \\ \vdots & & \vdots & \vdots & & \vdots \\ f'_{k'1} & \cdots & f'_{k',i-1} & f'_{k',i+1} & \cdots & f'_{k'k'} \end{vmatrix}, \quad i = 1, \dots, k',$$

the curve C'' is defined by the minors

$$\begin{vmatrix} f''_{11} & \cdots & f''_{1,j-1} & f''_{1,j+1} & \cdots & f''_{1k''} \\ \vdots & & \vdots & \vdots & & \vdots \\ f''_{k''1} & \cdots & f''_{k'',j-1} & f''_{k'',j+1} & \cdots & f''_{k''k''} \end{vmatrix}, \quad j = 1, \dots, k''$$

and the set

$$V(s'_1 \wedge \dots \wedge s'_{k'}, s''_1 \wedge \dots \wedge s''_{k''})$$

is defined by

$$\begin{vmatrix} f'_{11} & \cdots & f'_{1k'} \\ \vdots & & \vdots \\ f'_{k'1} & \cdots & f'_{k'k'} \end{vmatrix}$$

and

$$\begin{pmatrix} f''_{11} & \cdots & f''_{1k''} \\ \vdots & & \vdots \\ f''_{k''1} & \cdots & f''_{k''k''} \end{pmatrix},$$

the result follows at once. \square

As a corollary, we can prove the following (see also Lemma 4.1 of [4]):

Lemma 3.2. *Let \mathcal{E}' and \mathcal{E}'' be two vector bundles on X_r of rank k' and k'' respectively. Assume that \mathcal{E}' and \mathcal{E}'' satisfy condition \star and that there are sections $s'_1, \dots, s'_{k'}$ of \mathcal{E}' and $s''_1, \dots, s''_{k''}$ of \mathcal{E}'' such that the hypersurfaces $V(s'_1 \wedge \dots \wedge s'_{k'})$ and $V(s''_1 \wedge \dots \wedge s''_{k''})$ do not share a common component. Then a general \mathcal{E} fitting in an exact sequence*

$$0 \rightarrow \mathcal{E}' \rightarrow \mathcal{E} \rightarrow \mathcal{E}'' \rightarrow 0 \quad (3.1)$$

satisfies condition \star .

Proof. It is clear that \mathcal{E} satisfies conditions (i) and (iii) of Definition 1.2. Hence it is enough to check condition (ii). Since this condition is open, it suffices to prove it for $\mathcal{E} = \mathcal{E}' \oplus \mathcal{E}''$. But this follows immediately from Lemma 3.1, since our assumption implies that we can find sections $s'_1, \dots, s'_{k'}$ of \mathcal{E}' and $s''_1, \dots, s''_{k''}$ of \mathcal{E}'' such that $V(s'_1 \wedge \dots \wedge s'_{k'}, s''_1 \wedge \dots \wedge s''_{k''})$ is a curve. \square

Remark 3.3. We can use condition (ii) in Definition 1.2 to give a geometric interpretation (and a criterion) for the hypothesis in Lemma 3.2. For a vector bundle \mathcal{E} satisfying condition \star , the choice of $k - 1$ global sections s_1, \dots, s_{k-1} whose dependency locus is a curve $C \subset X_r$ yields the exact sequence (2.4). Since we are assuming $h^0 \mathcal{E} \geq k$, this means that there is at least a section s_k of \mathcal{E} independent of s_1, \dots, s_{k-1} . Any such s_k yields a hypersurface of degree c_1 containing C and defined by the global section $s_1 \wedge \dots \wedge s_k$ of $\mathcal{O}_{X_r}(c_1)$ (with the convention of Lemma 3.1). Notice if \mathcal{E}' and \mathcal{E}'' are ACM also \mathcal{E} is ACM.

Corollary 3.4. *Let (\mathcal{E}', C') and (\mathcal{E}'', C'') be two ACM pairs satisfying condition \star . Then a general \mathcal{E} appearing in an extension as (3.1) satisfies condition \star if at least one of the following conditions is satisfied:*

- (a) *there is a hypersurface of degree c'_1 containing C' and a hypersurface of degree c''_1 containing C'' such that these two hypersurfaces do not share any component;*
- (b) *at least one of \mathcal{E}' and \mathcal{E}'' is generated by its global sections.*

Proof. Part (a) is an immediate consequence of Lemma 3.2 and the previous remark. To prove part (b), assume for instance that \mathcal{E}' is generated by its global sections. We fix any hypersurface of degree k'' containing C'' and take a point in any of its components. Since \mathcal{E}' is generated by its global sections we can find k' sections of it whose dependency locus does not contain any of those points. This dependency locus is therefore a hypersurface of degree c'_1 containing C' and having no common components with the fixed hypersurface of degree c''_1 containing C'' . We thus conclude from (a). \square

We want now to apply Corollary 3.4 to obtain rank four ACM vector bundles on X_4 satisfying condition \star from vector bundles of rank two \mathcal{E}' and \mathcal{E}'' . We recall how to compute the invariants of any \mathcal{E} fitting in an extension (3.1). Let us denote by $c'_i \in \mathbf{Z}$ the Chern classes of \mathcal{E}' and by $c''_i \in \mathbf{Z}$ the Chern classes of \mathcal{E}'' . Then:

$$\begin{aligned} c_1(\mathcal{E}) &= c'_1 + c''_1 \\ c_2(\mathcal{E}) &= c'_2 + 4c'_1c''_1 + c''_2 \\ c_3(\mathcal{E}) &= c'_2c''_1 + c'_1c''_2 \end{aligned} \tag{3.2}$$

Checking Remark 2.5 for finding the list of rank two ACM bundles on X_4 satisfying condition \star we find the following list of examples.

Example 3.5. We take $\mathcal{E}', \mathcal{E}''$ to be rank two ACM vector bundles on X_4 with $c_1 = 3$, $c_2 = 14$. These are generated by their global sections, so that it follows from Corollary 3.4(b) that any general element in $Ext^1(\mathcal{E}'', \mathcal{E}')$ (which is a vector space of dimension at least 7) provides a rank four ACM vector bundle satisfying condition \star . In fact, in this case it is easier to observe that any extension provides a globally generated vector bundle, and hence it always satisfies condition \star . By the formulas (3.2), the invariants of such vector bundle are $(k; c_1, c_2, c_3) = (4; 6, 64, 84)$.

Example 3.6. We take now the rank two ACM vector bundles $\mathcal{E}', \mathcal{E}''$ in X_4 with $c_1(\mathcal{E}') = 3$, $c_2(\mathcal{E}') = 14$, and $c_1(\mathcal{E}'') = 2$. It follows again from Corollary 3.4(b) that a general element in $Ext^1(\mathcal{E}'', \mathcal{E}')$ (which is a vector space of dimension at least 7) provides a rank four ACM vector bundle \mathcal{E} satisfying condition \star . In this case, the invariants produced by the formulas (3.2) are $(k; c_1, c_2, c_3) = (4; 5, 46, 52)$.

Example 3.7. We repeat the same reasoning as in Example 3.6, but taking now \mathcal{E}'' a rank two ACM vector bundle with $c_1(\mathcal{E}'') = 1$, $c_2(\mathcal{E}'') = d$ (for $d = 3, 4$). As before, a general element in $Ext^1(\mathcal{E}'', \mathcal{E}')$ (which is a vector space of dimension at least $3d - 8$) yields an unstable rank four ACM vector bundle satisfying condition \star and with invariants $(k; c_1, c_2, c_3) = (4; 4, 26 + d, 3d + 14)$, $d = 3, 4$.

Example 3.8. Now we take $\mathcal{E}', \mathcal{E}''$ two normalized rank two ACM vector bundles that are globally generated on X_4 with $c_1 = 2$ and $c_2 = 8$. Again from Corollary 3.4(b) we get that a general element in $Ext^1(\mathcal{E}'', \mathcal{E}')$ (which is a vector space of dimension at least 4) provides a rank four ACM vector bundle satisfying condition \star . Using once more the equations (3.2) we find that this bundle has invariants $(k; c_1, c_2, c_3) = (4; 4, 32, 32)$.

Example 3.9. We replace now in Example 3.8 the vector bundle \mathcal{E}'' with a normalized rank two ACM vector bundle on X_4 with $c_1(\mathcal{E}'') = 1$ and $c_2(\mathcal{E}'') = d$ (with $d = 3, 4$). Hence we get that a general element in $Ext^1(\mathcal{E}'', \mathcal{E}')$ (which is a vector space of dimension at least $2d - 4$) provides an unstable rank four ACM vector bundle satisfying condition \star and with invariants $(k; c_1, c_2, c_3) = (4; 3, 16 + d, 8 + 2d)$, $d = 3, 4$.

Example 3.10. Take now \mathcal{E}' and \mathcal{E}'' to be two normalized rank two ACM bundles on X_4 with $c_1(\mathcal{E}') = 1$, $c_2(\mathcal{E}') = d'$ and $c_1(\mathcal{E}'') = 1$, $c_2(\mathcal{E}'') = d''$, (with $d', d'' = 3, 4$). The curves obtained as the zero loci of sections of \mathcal{E}' and \mathcal{E}'' are elliptic curves of degree d' and d'' , in any case contained in at least one hyperplane of \mathbf{P}^4 . A simple calculation shows that a hyperplane section of X_4 (which is a K3 surface) contains at most a pencil of elliptic curves of degree d , while the family of those curves in X_4 has dimension d (see [20]). Hence we can take pairs (\mathcal{E}', C') and (\mathcal{E}'', C'') such that the hyperplanes containing C' and C'' are different. Therefore we can apply Lemma 3.2 and take a general element in $\text{Ext}^1(\mathcal{E}', \mathcal{E}'')$ (which has dimension at least $d' + d'' - 6 \geq 0$) to produce an ACM vector bundle satisfying condition \star and with invariants $(k; c_1, c_2, c_3) = (4; 2, d' + d'' + 4, d' + d'')$, $d, d' = 3, 4$.

4 Constructing bundles from curves on the quartic threefold

In this section, we use first Hartshorne-Serre correspondence, to translate the property of being an ACM vector bundle satisfying condition \star to the property of curves in X_r to be associated to it. This will allow to produce several examples of ACM vector bundles of rank $k = 3, 4$ on X_4 satisfying condition \star from curves in X_4 .

We start by characterizing when a curve C determines an ACM vector bundle (observe that the condition $c_1 > 0$ will not be restrictive because of (5.1)). In the sequel we will denote by $\mathcal{O}_C(1)$ the restriction of the hyperplane class H of X_r to $C \subset X_r$.

Proposition 4.1. *Let $C \subset X_r$ be a locally complete intersection curve and let \mathcal{E} be the vector bundle of rank $k \geq 2$ obtained, as in Proposition 2.1, from $k - 1$ sections of $\omega_C(5 - r - c_1)$. Assume $c_1 > 0$. Then \mathcal{E} is an ACM bundle satisfying condition \star if and only if C is projectively normal and the following four conditions hold:*

- (a) *the $k - 1$ given sections form a basis of $H^0\omega_C(5 - r - c_1)$ (in particular $h^0\omega_C(5 - r - c_1) = k - 1$);*
- (b) *$h^0\mathcal{J}_C(c_1 - 1) = 0$ and $h^0\mathcal{J}_C(c_1) \geq 1$;*
- (c) *$h^0\omega_C(4 - r - c_1) = 0$, for which a sufficient condition is $2g(C) - 2 < (r + c_1 - 4) \deg C$;*
- (d) *the natural map $\alpha_n : H^0\omega_C(5 - r - c_1) \otimes H^0\mathcal{O}_C(n) \rightarrow H^0\omega_C(5 - r - c_1 + n)$ is surjective $\forall n \geq 1$.*

Proof. Since $h^i\mathcal{O}_{X_r}(n) = 0$ for all $n \in \mathbf{Z}$ and $i = 1, 2$ then the condition $h^1\mathcal{E}(n) = 0$ for all $n \in \mathbf{Z}$ is equivalent to the condition $h^1\mathcal{J}_C(c_1 + n) = 0$

for all $n \in \mathbf{Z}$, i.e. to the projective normality of C . On the other hand, splitting (2.5) into the following exact sequences

$$0 \rightarrow \mathcal{O}_{X_r}(-c_1) \rightarrow \mathcal{E}^\vee \rightarrow K \rightarrow 0 \quad (4.1)$$

and

$$0 \rightarrow K \rightarrow \mathcal{O}_{X_r}^{k-1} \rightarrow \omega_C(5-r-c_1) \rightarrow 0 \quad (4.2)$$

we get that $h^1 \mathcal{E}^\vee(n) = 0 \forall n \in \mathbf{Z}$ if and only if the map $r_n : H^0 \mathcal{O}_{X_r}^{k-1}(n) \rightarrow H^0 \omega_C(5-r-c_1+n)$ is surjective $\forall n \in \mathbf{Z}$. The surjectivity of r_n for all $n < 0$ is equivalent to $h^0 \omega_C(4-r-c_1) = 0$, for which a sufficient condition is $2g(C) - 2 < (r+c_1-4) \deg C$. For $n = 0$, the map r_0 is the one coming from the choice of $k-1$ sections of $\omega_C(5-r-c_1+n)$, so it is surjective if and only if we take a system of generators of $H^0 \omega_C(5-r-c_1+n)$. For $n > 0$, observe that r_n factors through

$$H^0 \mathcal{O}_{X_r}^{k-1} \otimes H^0 \mathcal{O}_{X_r}(n) \rightarrow H^0 \omega_C(5-r-c_1) \otimes H^0 \mathcal{O}_C(n) \rightarrow H^0 \omega_C(5-r-c_1+n).$$

If C is projectively normal and r_0 is surjective, the first map is surjective, and hence the surjectivity of r_n becomes equivalent to the surjectivity of the second map, which is precisely α_n .

Finally, observe that the map r_0 is not injective (i.e. condition (a) does not hold) is equivalent, by (4.1) and (4.2) and the fact that $c_1 > 0$, to the existence of a section of \mathcal{E}^\vee mapping to a nonzero section of $\mathcal{O}_{X_r}^{k-1}$. This is equivalent to say that \mathcal{E} has a trivial summand, which means that condition (iii) in Definition 1.2 does not hold. On the other hand, condition (b) is clearly equivalent, by (2.4), to $h^0 \mathcal{E}(-1) = 0$ and $h^0 \mathcal{E} \geq k$, i.e. condition (i) in Definition 1.2. \square

We give now some examples of the above construction applied to $X = X_4$ (for the rest of this section, unless otherwise specified, X will stand for X_4). We start by giving a method to construct curves contained in a hyperplane, which we will thus allow us to construct vector bundles with $c_1 = 1$.

Lemma 4.2. *Let \mathcal{C} be a family of degenerate curves in \mathbf{P}^4 such that some curve of \mathcal{C} is contained in a smooth degenerate (in \mathbf{P}^4) quartic surface and it is not the complete intersection of this surface and a hypersurface. Then the general quartic hypersurface $X \subset \mathbf{P}^4$ contains a curve of \mathcal{C} .*

Proof. Consider the set Σ of degenerate quartic surfaces of \mathbf{P}^4 containing a curve of \mathcal{C} . We define the natural map

$$\varphi : \Sigma \rightarrow (\mathbf{P}^4)^*$$

associating to each quartic surface the unique hyperplane containing it. Fix now a hyperplane H of \mathbf{P}^4 . It is a standard fact in the Noether-Lefschetz theory (see for example [6]), that the set of quartic surfaces in H containing a curve in \mathcal{C} has codimension one in the set of quartic surfaces of H . In other words, the set $\varphi^{-1}(H)$ has dimension 33, and hence Σ has dimension 37.

Consider now the set I of pairs (S, X) where $S \in \Sigma$ and X is a quartic hypersurface of \mathbf{P}^4 containing S . Since the fibers of the projection $I \rightarrow \Sigma$ are projective spaces of dimension 35, it follows that I has dimension 72.

We finally consider the second projection $p_2 : I \rightarrow \mathbf{P}^{69}$ (where we identify \mathbf{P}^{69} with the set of quartic hypersurfaces of \mathbf{P}^4). The lemma will be proved if we show that p_2 is surjective. Observe that, since any smooth degenerate quartic surface in \mathbf{P}^4 is the hyperplane section of some smooth quartic hypersurface of \mathbf{P}^4 , it follows from our hypotheses that there is a smooth quartic hypersurface X in the image of p_2 . Since a general hyperplane section of X has its Picard group generated by the hyperplane divisor (see for example [24]), such a hyperplane section cannot contain a curve of \mathcal{C} , and hence the set $p_2^{-1}(X)$ has dimension at most three. Hence the fiber of any element of the image of p_2 has necessarily dimension three, which shows that p_2 is surjective, as wanted. \square

In order to apply the previous lemma we show the following:

Lemma 4.3. *A smooth projectively normal space curve C of degree d and genus g with $d \geq g - 1$ is contained in a smooth quartic surface in \mathbf{P}^3 .*

Proof. The statement follows readily from the more general results of [23]. We give however a direct proof valid for our cases using the following simple standard argument. We first observe, from Castelnuovo-Mumford's criterion, that $\mathcal{J}_C(4)$ is globally generated. Indeed the projective normality of C provides the vanishing of $h^1\mathcal{J}_C(3)$, while the vanishings of $h^3\mathcal{J}_C(1)$ and $h^2\mathcal{J}_C(2)$ come from the equalities

$$h^3\mathcal{J}_C(3) = h^2\mathcal{O}_C(3) = 0,$$

$$h^2\mathcal{J}_C(2) = h^1\mathcal{O}_C(2) = h^0\omega_C(-2) = 0$$

(the latter coming from the assumption $d \geq g - 1$).

Hence the linear system $|H^0\mathcal{J}_C(4)|$ has no base-points outside C and therefore, by Bertini's theorem, a general element of it is smooth outside C . By a well known argument (see e.g. [25]), from the exact sequence

$$0 \rightarrow \mathcal{J}_C^2(4) \rightarrow \mathcal{J}_C(4) \rightarrow N_C^\vee(4) \rightarrow 0 \quad (4.3)$$

a surface in the linear system $|H^0\mathcal{J}_C(4)|$ provides a section of $N_C^\vee(4)$, and the singular points of the surface belonging to C are those in the zero locus of the section. The sequence (4.3) proves that the rank two vector bundle $N_C^\vee(4)$ is generated by the global sections coming from sections of $\mathcal{J}_C(4)$. Hence, a general such section of $N_C^\vee(4)$ will be nowhere vanishing, which implies that a general element of $|H^0\mathcal{J}_C(4)|$ is smooth also in the points of C . Such a general element gives thus the wanted smooth quartic surface containing C . \square

As a first application, we give a negative answer to Question 1.3.

Proposition 4.4. *There exists an ACM bundle on X satisfying condition \star with invariants $(4; 1, 6, 4)$. Moreover a general such bundle is not extension class of any rank two ACM bundles.*

Proof. Let C be a smooth projectively normal curve (hence non hyperelliptic) of degree $\deg C = 6$ and genus $g(C) = 3$ in \mathbf{P}^3 . By Lemma 4.3, C is contained in a smooth quartic surface and by Lemma 4.2 a general quartic hypersurface in \mathbf{P}^4 contains a degenerate curve C of degree 6 and genus 3. Since $h^0\omega_C = 3$, by Proposition 2.1 C defines a rank four vector bundle given by

$$0 \rightarrow \mathcal{O}_X^3 \rightarrow \mathcal{E} \rightarrow \mathcal{J}_C(1) \rightarrow 0 \quad (4.4)$$

with $c_1(\mathcal{E}) = 1$, $c_2(\mathcal{E}) = 6$, $c_3 = 4$ (by (2.6)) and $h^0\mathcal{E} = 4$. Proposition 4.1 applies once we show the map

$$\alpha_1 : H^0\omega_C \otimes H^0\mathcal{O}_C(1) \rightarrow H^0\omega_C(1)$$

is surjective, which follows by Castelnuovo's Lemma (see [1] pg.151 or [2] theorem (1.6)). Hence \mathcal{E} is ACM and satisfies condition \star .

Finally any such bundle is not an extension class of any two rank two ACM bundles on X . Suppose to the contrary that there exists a non trivial extension class

$$0 \rightarrow \mathcal{E}' \rightarrow \mathcal{E} \rightarrow \mathcal{E}'' \rightarrow 0$$

with \mathcal{E}' and \mathcal{E}'' ACM of rank two. Since \mathcal{E} satisfies condition \star , then both \mathcal{E}' and \mathcal{E}'' are normalized, i.e. $b(\mathcal{E}') = b(\mathcal{E}'') = 0$ and $h^0\mathcal{E}' \cdot h^0\mathcal{E}'' \geq 1$. Then a direct computation shows that equations (3.2) have not integral solutions for Chern classes of \mathcal{E}' and \mathcal{E}'' as in Proposition 2.4 and we are done. \square

Remark 4.5. Alternatively, to show the existence of space curves of degree 6 and genus 3 on general X one can start with a curve of degree 10 as in Example 3.10 and then taking the residual curve to it in a complete intersection $(2, 2, 4)$.

Similar to the previous case, we have also the following:

Proposition 4.6. *There exists an ACM bundle on X satisfying condition \star with invariants $(3; 1, 5, 2)$.*

Proof. Let C be a smooth curve of type $(2, 3)$ in a smooth quadric surface $C \subset Q \subset \mathbf{P}^3$. By Lemma 4.3, there exists a smooth quartic surface containing C . Hence, by Lemma 4.2, a general quartic hypersurface contains a curve C of degree 5 genus 2. Since $h^0\omega_C = 2$, by Proposition 2.1 we get a rank three vector bundle \mathcal{E} fitting in the exact sequence

$$0 \rightarrow \mathcal{O}_X^2 \rightarrow \mathcal{E} \rightarrow \mathcal{J}_C(1) \rightarrow 0$$

with $c_1 = 1$, $c_2 = 5$, $c_3 = 2$ and $h^0\mathcal{E} = 3$. Using Castelnuovo's Lemma it is easy to check that C satisfies the conditions of Proposition 4.1 and hence \mathcal{E} is ACM and satisfies condition \star . \square

Remark 4.7. A similar liaison argument as above shows the existence of rational space cubic curves in X . Specifically, let Q be the unique quadric surface containing a quintic $C \subset X$ of genus 2 as in Proposition 4.6. Then C is residual to a rational cubic curve $D \subset Q \cap X$.

5 Bounds of the Chern classes

In this section we will prove several restrictions for the Chern classes of ACM vector bundles, with special attention to the case $r = 4$. This will yield the list of possible Chern classes stated in Theorem 1.5 for the cases $k = 3, 4$. We start with bounds for c_1 and c_2 . The proof of the bound for c_1 is a straightforward extension of the method used in [19] for the rank two case.

Lemma 5.1. *Let (\mathcal{E}, C) be an ACM pair on X_r satisfying condition \star and having invariants $(k; c_1, c_2, c_3)$. Then*

$$1 \leq c_1 \leq \frac{k(r-1)}{2} \quad (5.1)$$

and

$$c_2 \leq \frac{r}{2}c_1^2 - \frac{r(r-2)}{2}c_1 + \frac{r(r-1)(r-2)}{6}k.$$

Proof. By the exact sequence (2.4), since $h^0\mathcal{E} \geq k$ it follows $h^0J_C(c_1) \geq 1$ and hence $c_1 \geq 1$. Let H be a general hyperplane section of X_r . Taking cohomology in the exact sequence

$$0 \rightarrow \mathcal{E}(-2) \rightarrow \mathcal{E}(-1) \rightarrow \mathcal{E}_H(-1) \rightarrow 0$$

(and its twists by any $\mathcal{O}_{X_r}(l)$), we get that \mathcal{E}_H is an ACM bundle on H . Since $h^0\mathcal{E}(-1)$ (because \mathcal{E} normalized) we also get $h^0\mathcal{E}_H(-1) = 0$. Therefore,

$$0 \leq h^2\mathcal{E}_H(-1) = \chi\mathcal{E}_H(-1) = -c_2 + \frac{r}{2}c_1^2 - \frac{r(r-2)}{2}c_1 + \frac{r(r-1)(r-2)}{6}k$$

and the bound for c_2 follows. Similarly, if Y is a general hyperplane section of H , we have an exact sequence

$$0 \rightarrow \mathcal{E}_H(-2) \rightarrow \mathcal{E}_H(-1) \rightarrow \mathcal{E}_Y(-1) \rightarrow 0$$

which implies $h^0\mathcal{E}_Y(-1) = 0$. Therefore $0 \geq \chi\mathcal{E}_Y(-1) = r(c_1 - k(r-1)/2)$ and the upper bound for c_1 follows immediately. \square

In the case of quartic hypersurfaces X_4 , which is the case we are interested in, we have stronger restrictions:

Proposition 5.2. *Let (\mathcal{E}, C) be an ACM pair satisfying condition \star on X_4 and having invariants $(k; c_1, c_2, c_3)$. Then*

$$c_3 = -\frac{4}{3}c_1^3 + 2c_1^2 - \frac{14}{3}c_1 + c_1c_2 - c_2 + 2k \quad (5.2)$$

and

$$g(C) = -\frac{2}{3}c_1^3 + c_1^2 - \frac{7}{3}c_1 + 1 + (c_1 - 1)c_2 + k. \quad (5.3)$$

Moreover

$$1 \leq c_1 \leq \frac{3k}{2}. \quad (5.4)$$

$$2c_1^2 - 2c_1 + k \leq c_2 \leq \min\{2c_1^2 - 4c_1 + 4k, 2c_1^2 + k\} \quad (5.5)$$

and if $c_1 > 1$ then $c_2 \geq 2c_1^2 - 4c_1 + 8$.

Proof. By hypothesis $h^0\mathcal{E}(-1) = 0$, and also $h^0\mathcal{E}^\vee = 0$ because \mathcal{E} has not trivial summands (see the proof of Proposition 4.1). Since $h^3\mathcal{E}(-1) = h^0\mathcal{E}^\vee$ (this is the point in which we use $r = 4$) it follows $\chi\mathcal{E}(-1) = 0$, which provides (5.2) by using Riemann-Roch. Now (5.3) comes from (2.6) performing the substitution of c_3 given by (5.2).

The bound (5.4) and first upper bound for c_2 in (5.5) are the ones of Lemma 5.1 for $r = 4$. For the other upper bound for c_2 , since $h^3\mathcal{E} = h^0\mathcal{E}^\vee(-1) = 0$ (again because $r = 4$) and $h^0\mathcal{E} \geq k$ by hypothesis, it follows that $\chi(\mathcal{E}) \geq k$. By Riemann-Roch, using the substitution (5.2) we get $\chi(\mathcal{E}) = -c_2 + 2c_1^2 + 2k$ which yields the wanted upper bound for c_2 .

For the lower bound of c_2 , we take a general linear projection $\pi : X \rightarrow \mathbf{P}^3$ from a point of \mathbf{P}^4 . It is immediate to observe that $\pi_*\mathcal{E}$ is an ACM vector bundle of rank $4k$, which should split completely by Horrocks theorem. Since $h^0\mathcal{E}(-1) = 0$, $\pi_*\mathcal{E}$ must contain a direct summand $\mathcal{O}_{X_4}^{h^0\mathcal{E}}$, and similarly a direct summand $\mathcal{O}_{X_4}(-2)^{h^3\mathcal{E}(-2)}$, since $h^3\mathcal{E}(-1) = 0$. Therefore, $h^0\mathcal{E} + h^3\mathcal{E}(-2) \leq 4k$, which gives the wanted inequality when using Riemann-Roch and the substitution (5.2).

Finally, let us prove the last lower bound for c_2 . Since $h^0\mathcal{J}_C(c_1 - 2) = 0$ (by Proposition 4.1(b)), it follows that

$$h^0\mathcal{O}_C(c_1 - 2) = h^0\mathcal{O}_{X_4}(c_1 - 2) = \frac{2}{3}c_1^3 - 3c_1^2 + \frac{19}{3}c_1 - 5.$$

Since by Riemann-Roch we have

$$\chi\mathcal{O}_C(c_1 - 2) = \frac{2}{3}c_1^3 - c_1^2 + \frac{7}{3}c_1 - k - c_2$$

we derive $h^0\omega_C(2 - c_1) = c_2 - 2c_1^2 + 4c_1 - 5 + k$. The proof concludes by applying the bilinear lemma (see [11]) to the nondegenerate bilinear map

$$H^0\omega_C(1 - c_1) \otimes H^0\mathcal{O}_C(1) \rightarrow H^0\omega_C(2 - c_1)$$

using that $h^0\omega_C(1 - c_1) = k - 1$ (by Proposition 4.1(a)) and that $h^0\mathcal{O}_C(1) \geq 5$ if $c_1 > 1$, because C is not contained in any hyperplane (by Proposition 4.1(b)). \square

We will study separately the cases $c_1 = 1$ and $2 \leq c_1 \leq 3k/2$. Proposition 5.2 gives immediate results for vector bundles of rank $k \geq 3$ with $c_1 = 1$:

Proposition 5.3. *Let (\mathcal{E}, C) be a pair on X_4 satisfying condition \star , where \mathcal{E} is ACM of rank $k \geq 3$ and $c_1 = 1$. Then $c_2 = k + 2$, $c_3 = 2k - 4$ and C is a curve of degree $k + 2$ and genus $g(C) = k - 1$.*

Proof. We immediately get from Proposition 5.2 $g(C) = k - 1$ and $\deg C = c_2 \leq k + 2$. Hence, C cannot be a plane curve and therefore $h^0 \mathcal{J}_C(1) = 1$ (it is at least 1 by Proposition 4.1). This implies $(h^3 \mathcal{E} = 0$, as proved in Proposition 5.2) $\chi(\mathcal{E}) = h^0 \mathcal{E} = k - 1 + h^0 \mathcal{J}_C(1) = k$. Since by Riemann-Roch $\chi(\mathcal{E}) = -c_2 + 2k + 2$ it follows $c_2 = k + 2$, which is the degree of C . Finally, from (5.2) we get $c_3 = 2k - 4$. \square

For the rank three case when $c_1 > 1$ we have the following result if $c_1 = 3, 4$:

Proposition 5.4. *Let (\mathcal{E}, C) be a pair satisfying condition \star , where \mathcal{E} is ACM of rank $k = 3$. If $c_1 \geq 3$ then $2c_1^2 - 4c_1 + 11 \leq c_2 \leq 2c_1^2 - 4c_1 + 12$.*

Proof. The upper bound follows from Proposition 5.2. For the lower bound, observe that $h^3 \mathcal{E}^\vee(1) = h^0 \mathcal{E}(-2) = 0$, and thus $h^0 \mathcal{E}^\vee(1) = \chi \mathcal{E}^\vee(1) = 2c_1^2 - 4c_1 - c_2 + 12$. Hence it will be enough to prove $h^0 \mathcal{E}^\vee(1) \leq 1$.

Assume for contradiction $h^0 \mathcal{E}^\vee(1) \geq 2$ and let s_1, s_2 be two independent global sections. Then $\wedge^2(\mathcal{E}^\vee(1)) \cong \mathcal{E}(2 - c_1)$ contains the nonzero section defined by $s_1 \wedge s_2$, which is absurd since \mathcal{E} is normalized and $2 - c_1 < 0$. \square

Remark 5.5. The above Proposition leaves only two possibilities for c_2 , depending on whether $h^0 \mathcal{E}^\vee(1)$ is 0 or 1. When $h^0 \mathcal{E}^\vee(1) = 0$ (i.e. $c_2 = 2c_1^2 - 4c_1 + 12$), it follows that both $\mathcal{E}(-1)$ and its dual have no global sections, so it follows (as in [26] Remark 1.2.6) that $\mathcal{E}(-1)$, and hence \mathcal{E} , is stable. Observe that if $c_1 = 3$ we get $c_2 = 18$, and if $c_1 = 4$ then $c_2 = 28$.

Propositions 5.2, 5.3 and 5.4 give immediately the following possibilities for the Chern classes of ACM vector bundles of rank three or four over X_4 :

Proposition 5.6. *Let (\mathcal{E}, C) be an ACM pair satisfying condition \star where \mathcal{E} is of rank $k = 3$. Then the possibilities for the Chern classes of \mathcal{E} and the genus $g = g(C)$ are:*

1. $c_1 = 1, c_2 = 5, c_3 = 2, g(C) = 2$;
2. $c_1 = 2, 8 \leq c_2 \leq 11, c_3 = c_2 - 6, g(C) = c_2 - 2$;
3. $c_1 = 3, 17 \leq c_2 \leq 18, c_3 = 2c_2 - 26, g(C) = 2c_2 - 12$;
4. $c_1 = 4, 27 \leq c_2 \leq 28, c_3 = 3c_2 - 66, g(C) = 3c_2 - 32$.

Proposition 5.7. *Let (\mathcal{E}, C) be an ACM pair satisfying condition \star where \mathcal{E} is of rank $k = 4$. Then the possibilities for the Chern classes of \mathcal{E} and the genus $g = g(C)$ are:*

1. $c_1 = 1, c_2 = 6, c_3 = 4, g(C) = 3$;
2. $c_1 = 2, 8 \leq c_2 \leq 12, c_3 = c_2 - 4, g(C) = c_2 - 1$;
3. $c_1 = 3, 16 \leq c_2 \leq 22, c_3 = 2c_2 - 24, g(C) = 2c_2 - 11$;
4. $c_1 = 4, 28 \leq c_2 \leq 32, c_3 = 3c_2 - 64, g(C) = 3c_2 - 31$;

5. $c_1 = 5$, $44 \leq c_2 \leq 46$, $c_3 = 4c_2 - 132$, $g(C) = 4c_2 - 65$;

6. $c_1 = 6$, $c_2 = 64$, $c_3 = 84$, $g(C) = 5c_2 - 117$.

We end this section with some remarks and comments. We notice the following result can be derived using similar arguments as in the above cases of rank three:

Corollary 5.8. *There are no rank three ACM vector bundles satisfying condition \star on a general quartic hypersurface in \mathbf{P}^n , $n \geq 5$. In other words, rank three ACM bundles satisfying condition \star on a general quartic threefold $X_4 \subset \mathbf{P}^4$ do not extend to a general quartic hypersurface $W \subset \mathbf{P}^5$ having X_4 as a hyperplane section.*

Since in [9] we showed that all rank two ACM bundles on a general quartic $W \subset \mathbf{P}^n$, $n \geq 5$, splits, then a natural question, which we hope to consider later, is the following:

Question 5.9. *Which is the minimum rank for a positive dimensional family of ACM bundles satisfying condition \star on a general quartic fourfold $W \subset \mathbf{P}^5$.*

References

- [1] E. Arbarello, M. Cornalba, P.A. Griffiths and J. Harris, *Geometry of Algebraic Curves*, Springer, 1985.
- [2] E. Arbarello and E. Sernesi, *Petri's approach to the study of the ideal associated to a special divisor*, *Inventiones Math.* **49**(1978), 99–119.
- [3] E. Arrondo, *A home-made Hartshorne-Serre correspondence*, *Rev. Mat. Complut.* **20** (2007), no. 2, 423–443.
- [4] E. Arrondo and L. Costa, *Vector bundles on Fano 3-folds without intermediate cohomology*, *Comm. Algebra* **28** (2000), no. 8, 3899–3911.
- [5] E. Arrondo and D. Faenzi, *Vector bundles with no intermediate cohomology on Fano threefolds of type V_{22}* , *Pacific J. Math.* **225** (2006), no. 2, 201–220.
- [6] J. Carlson, M. Green, P. Griffiths and J. Harris, *Infinitesimal variations of Hodge structures (I)*, *Comp. Math.* **50** (1983), 105–205.
- [7] L. Chiantini and C. Madonna, *ACM bundles on a general quintic threefold*, *Matematiche (Catania)* **55**(2000), no.2, 239–258.
- [8] L. Chiantini and C. Madonna, *A splitting criterion for rank 2 vector bundles on a general sextic threefold*, *Internat. J. Math.* **15**(2004), no.4, 341–359.

- [9] L. Chiantini and C. Madonna, *ACM bundles on general hypersurfaces in \mathbf{P}^5 of low degree*, Collect. Math. Vol. **56**(2005), no. 1, 85-96.
- [10] D. Eisenbud, *Commutative algebra with a view toward algebraic geometry*, Springer 1999.
- [11] D. Eisenbud, J. Koh, and M. Stillman, *Determinantal Equations for Curves of High Degree* Amer. J. Math. Vol. **110**(1988), no. 3, pp. 513-539.
- [12] D. Faenzi, *Bundles over the Fano threefold V_5* , Comm. Algebra **33**(2005), no. 9, 3061–3080.
- [13] W. Fulton, *Intersection theory*, Springer 1998.
- [14] J. Harris, M. Roth and J. Starr, *Curves of small degree on cubic threefolds*, Rocky Mountain J. Math. **35**(2005), no. 3, 761–817.
- [15] G. Horrocks, *Vector bundles on the punctured spectrum of a local ring*, Proc. London Math. Soc. **14** (1964), 689-713.
- [16] A. Iliev and D. Markushevich, *Quartic 3-folds: pfaffians, vector bundles, and half-canonical curves*, Mich. Math. J. **47** (2000), 385–394.
- [17] A. Iliev and L. Manivel, *Pfaffian lines and vector bundles on Fano threefolds of genus 8*, J. Alg. Geom., to appear.
- [18] N.M. Kumar, A.P. Rao and G.V. Ravindra, *Arithmetically Cohen-Macaulay bundles on three dimensional hypersurfaces*, Int. Math. Res. Not. IMRN 2007, no. 8, Art. ID rnm025, 11 pp.
- [19] C. Madonna, *A splitting criterion for rank 2 vector bundles on hypersurfaces in \mathbf{P}^4* , Rend. Sem. Mat. Univ. Pol. Torino **56** (1998), no.2, 43–54.
- [20] C. Madonna, *Rank-two vector bundles on general quartic hypersurfaces in \mathbf{P}^4* , Rev. Mat. Complut. **XIII** (2000), num.2, 287–301.
- [21] C.G. Madonna, *ACM vector bundles on prime Fano threefolds and complete intersection Calabi Yau threefolds*, Rev. Roumaine Math. Pures Appl. **47**(2002) no.2, 211-222.
- [22] C. Madonna, *Rank 4 vector bundles on the quintic threefold*, CEJM **3** (2005), no.3, 404–411.
- [23] S. Mori, *On degrees and genera of curves on smooth quartic surfaces in \mathbf{P}^3* , Nagoya Math. J. **96**(1984), 127–132.
- [24] B.G. Moı̄shezon, *Algebraic homology classes on algebraic varieties*, Izv. Akad. Nauk SSSR Ser. Mat. **31** (1967) 225–268.

- [25] P.E. Newstead, *A space curve whose normal bundle is stable*, J.London Math. Soc. **28** (1983), no.2, 428–434.
- [26] C. Okonek, M. Schneider and H. Splinder, *Vector bundles on complex projective spaces*, Birkhauser, 1980, Boston.
- [27] G. Ottaviani, *Some extension of Horrocks criterion to vector bundles on Grassmannians and quadrics*, Ann. Mat. Pura Appl. (4) **155** (1989), 317–341.
- [28] A. Vogeelar, *Constructing vector bundles from codimension-two subvarieties*, PhD thesis, Leiden, 1978.

Enrique Arrondo

Departamento de Álgebra, Facultad de Ciencias Matemáticas, Universidad Complutense de Madrid, 28040 Madrid (Spain)
email: Enrique_Arrondo@mat.ucm.es

Carlo G. Madonna

Departamento de Matemáticas, Instituto de Matemáticas y Física Fundamental, CSIC, C/ Serrano 121, 28006 Madrid (Spain)

current address:

Departamento de Álgebra, Facultad de Ciencias Matemáticas Universidad Complutense de Madrid, 28040 Madrid (Spain)

email: carlo.madonna@mat.csic.es