AN ASYMPTOTICS OF VARIANCE OF THE LATTICE POINTS COUNT

JIŘÍ JANÁČEK

ABSTRACT. The variance of the number of lattice points inside the dilated bounded set rD with random position in \mathbb{R}^d has asymptotics $\sim r^{d-1}$ if the rotational quadratic average of the modulus of the Fourier transform of the set is $O(\rho^{-d-1})$. The asymptotics follows from a Wiener's Tauberian theorem.

1. INTRODUCTION

The number of lattice points in a set with random position can be used for estimation of the volume of the set and its variance has been studied for a long time [6, 7]. If **T** is a *d*-periodical point lattice of spatial intensity α in the *d*-dimensional Euclidean space \mathbb{R}^d then the mean value of

$$\left(\mathbf{card}\left(\left(B_{d}\left(r\right)+x\right)\cap\mathbf{T}\right)-lpha\lambda^{d}\left(B_{d}\left(r\right)\right)\right)^{2},$$

where λ^d is Lebesgue measure, i.e. the variance of the lattice points count in the ball of radius r with uniform random position, is

$$C_{\mathbf{T}}H^{d-1}\left(\partial B_{d}\left(r\right)\right)\Phi\left(r\right).$$

Here $C_{\mathbf{T}}$ is a lattice constant, H^{d-1} is the surface measure and Φ defined by the above equality fulfills $\lim_{t\to\infty} \frac{1}{t} \int_0^t \Phi(t) dt = 1$ and is bounded, $\limsup_{t\to\infty} \Phi(t) \leq 2$ [4, 5]. Hence the variance of the lattice point count in the ball has asymptotics "in the mean" $C_{\mathbf{T}}H^{d-1}(\partial B_d(1)) r^{d-1}$ and is $O(r^{d-1}), r \to \infty$.

Let D be a compact body the isotropic covariogram $\overline{\gamma}_D$ (rotational average of $\gamma_D = I_D \star I_{-D}$) of which is a fractional integral of the Weyl type of a finite signed measure σ on \mathbb{R}^+ :

$$\overline{\gamma}_{D}\left(t\right) = \frac{1}{\Gamma\left(\frac{d+1}{2}+1\right)} \int_{t}^{\infty} \left(s-t\right)^{\frac{d+1}{2}} d\sigma\left(s\right)$$

for $t \in \mathbb{R}^+$. This means that the fractional derivative of $\overline{\gamma}_D$ of the order $\frac{d+1}{2}$ has bounded variation. Then the asymptotics "in the mean" of the form $C_{\mathbf{T}}H^{d-1}(\partial D) r^{d-1}$ of the variance of the lattice points count inside randomly rotated and shifted set rD follows easily from Theorem 2.8 in [3].

The variance of the lattice points count inside randomly rotated and shifted bounded set D dilated by r > 0 is $O(r^{d-1})$, $r \to \infty$, if the rotational quadratic average of the modulus of the Fourier transform of the set D, defined as $\int_{S_{d-1}} \left| \widehat{I_D}(\rho\xi) \right|^2 d\xi$ is $O(\rho^{-d-1})$, $\rho \to \infty$ [12]; the later property was proved for convex sets and sets with $C^{\frac{3}{2}}$ boundary [2]. The aim of this paper is to prove that

¹⁹⁹¹ Mathematics Subject Classification. 62D05, 11H06.

Key words and phrases. point lattice, Fourier transform, volume, variance,

JIŘÍ JANÁČEK

this assumption on the growth of the modulus of the Fourier transform also yields the asymptotics of the form $C_{\mathbf{T}}H^{d-1}(\partial D) r^{d-1}$ of the variance of the lattice points count inside randomly placed bounded full *d*-dimensional set rD with sufficiently regular boundary (locally finite union of sets of finite reach [9]). The above asymptotics can be thus used for compact bodies with piecewise \mathbf{C}^2 smooth boundary and *d*-dimensional convex sets in \mathbb{R}^d .

2. The Variance of the Lattice Points Count

Notation 2.1. Let **T** be a *d*-periodic lattice of points in the *d*-dimensional Euclidean space \mathbb{R}^d defined by the regular matrix $A \in \mathbb{R}^{d \times d}$ as $\mathbf{T}(A) = A\mathbb{Z}^d$, where \mathbb{Z}^d is set of all points in \mathbb{R}^d with integral co-ordinates. **T** has the fundamental region $F_{\mathbf{T}} = A [0, 1)^d$ of volume $\lambda^d (F_{\mathbf{T}}) = \det A$, where λ^d is the Lebesgue measure; hence the spatial intensity of **T** is $\alpha = (\det A)^{-1}$. The group dual to the group $\mathbf{T}(A)$ is $\mathbf{T}^* = A^{-1}\mathbb{Z}^d$.

Fourier transform of a function $f \in \mathbf{L}^1(\mathbb{R}^d)$ is

(2.1)
$$\widehat{f}(\xi) = \int_{\mathbb{R}^d} f(x) \exp\left(-2\pi i x \cdot \xi\right) dx.$$

If f is moreover spherically symmetric then $r^{d-1}f(r) \in \mathbf{L}^1(\mathbb{R}^+)$ and Fourier transform of f can be expressed as the Haenkel transform

(2.2)
$$\widehat{f}(\rho) = 2\pi\rho^{1-\frac{d}{2}} \int_0^\infty r^{\frac{d}{2}} J_{\frac{d}{2}-1}(2\pi\rho r) f(r) dr,$$

where $J_{\frac{d}{2}-1}$ is the Bessel function of the first kind.

 $\kappa_d = \pi^{\frac{d}{2}} \Gamma \left(\frac{d}{2} + 1\right)^{-1}$ is the volume of the unit ball $B_d(1)$ in \mathbb{R}^d , where Γ is the Euler gamma function.

 I_D is the characteristic function of the set D.

Proposition 2.2. Let **T** be a *d*-periodic lattice of points and let *D* be a bounded measurable set in \mathbb{R}^d . Then

(2.3)
$$\int_{F_{\mathbf{T}}} \left(\operatorname{\mathbf{card}} \left((D+x) \cap \mathbf{T} \right) \right) \alpha \, dx = \alpha \lambda^d \left(D \right),$$

and

(2.4)
$$\int_{F_{\mathbf{T}}} \left(\operatorname{\mathbf{card}} \left((D+x) \cap \mathbf{T} \right) - \alpha \lambda^{d} (D) \right)^{2} \alpha \, dx = \sum_{0 \neq \xi \in \mathbf{T}^{*}} \left| \widehat{I_{D}} (\xi) \right|^{2},$$

where α is the spatial intensity of **T**.

Proof. Equation (2.3) can be proved by standard arguments and equation (2.4) follows from the Parseval theorem, see Theorem 2.3 in [3]. \Box

Definition 2.3. Covariogram of a bounded measurable set D is the function

$$\gamma_D(x) = I_D \star I_{-D}(x) = \int_{\mathbb{R}^d} I_D(x) I_D(x-y) \, dy.$$

It follows from the properties of Fourier transform that $\widehat{\gamma_D} = \left|\widehat{I_D}\right|^2$ is a nonnegative function. The isotropic covariogram is $\overline{\gamma_D}(|x|) = \int_{\mathbf{SO}_d} \gamma_{MD}(x) \, dM$, where MD is

the set D rotated by $M \in \mathbf{SO}_d$ and the integration uses the invariant probabilistic measure on \mathbf{SO}_d , the group of rotations in \mathbb{R}^d ; an equivalent definition is $\overline{\gamma_D}(u) = \int_{S_d-1} \gamma_D(ux) dx$. The Haenkel transform of the isotropic covariogram is $\overline{\widehat{\gamma_D}}$.

Remark 2.4. As was already discussed in Remark 2.5 in [3], it follows from this definition that γ_D is bounded and, as $\widehat{\gamma_D} \ge 0$, the function $\widehat{\gamma_D}$ is integrable in \mathbb{R}^d (see [1] Theorem 9.). Further, $\rho^{d-1}\widehat{\gamma_D}(\rho) \ge 0$ is integrable in \mathbb{R}^+ by Fubini theorem. γ_D is then the inverse Fourier transform (2.1) of $\widehat{\gamma_D}$ ([1] Theorem 8.) and $\overline{\gamma_D}$ is the (inverse) Haenkel transform (2.2) of $\widehat{\gamma_D}(\rho)$. We have from (2.4) and (2.3) using the variance decomposition lemma [8]

$$\begin{split} \int_{\mathbf{SO}_d} \int_{F_{\mathbf{T}}} \left(\mathbf{card} \left((MD + x) \cap \mathbf{T} \right) - \alpha \lambda^d \left(D \right) \right)^2 \alpha \, dx \, dM = \\ &= \sum_{0 \neq \xi \in \mathbf{T}^*} \widehat{\gamma_D} \left(|\xi| \right). \end{split}$$

Lemma 2.5. Let $r^2 f(r) \ge 0$ be bounded measurable function on \mathbb{R}^+ . Then

$$\lim_{R \to \infty} \frac{1}{R} \int_0^R r^2 f(r) \, dr = \lim_{h \to 0+} \frac{\pi^{-\frac{3}{2}}}{h} \frac{\Gamma\left(\frac{d+1}{2}\right)}{\Gamma\left(\frac{d}{2}\right)} \int_0^\infty \left(1 - \Gamma\left(\frac{d}{2}\right) \frac{J_{\frac{d}{2}-1}\left(2\pi hr\right)}{(\pi hr)^{\frac{d}{2}-1}}\right) f(r) \, dr$$

whenever at least one of the two limits exists.

Proof. The case d = 1 was proved in [13] (Theorem 21) using Wiener's Tauberian theorem (see e.g. [11]): Let $\varphi \in \mathbf{L}^{\infty}(\mathbb{R})$, $K \in \mathbf{L}^{1}(\mathbb{R})$, \widehat{K} has no root and $K \star \varphi(t) \to a\widehat{K}(0)$ as $t \to \infty$. Then for each $g \in \mathbf{L}^{1}(\mathbb{R})$ $g \star \varphi(t) \to a\widehat{g}(0)$ as $t \to \infty$. What follows is an extension of this proof to higher dimensions.

By the substitution $r = \exp(t)$ and defining

$$R = \exp(\eta) = \frac{1}{h}, \quad r^2 f(r) = \varphi(t),$$

we obtain equivalent formulation of the theorem suitable for direct application of the Wiener's Tauberian theorem

$$\lim_{\eta \to \infty} \int_{-\infty}^{\infty} K_1(\eta - t) \varphi(t) dt = \lim_{\eta \to \infty} \int_{-\infty}^{\infty} K_2(\eta - t) \varphi(t) dt$$

whenever at least one of the two limits exists. Here

$$K_{1}(s) = I_{\{s|s>0\}} \exp(-s), \quad K_{2}(s) = \exp(s) L(\exp(-s)),$$
$$L(u) = \pi^{-\frac{3}{2}} \frac{\Gamma(\frac{d+1}{2})}{\Gamma(\frac{d}{2})} \left(1 - \Gamma(\frac{d}{2}) \frac{J_{\frac{d}{2}-1}(2\pi u)}{(\pi u)^{\frac{d}{2}-1}}\right).$$

We can easily see that $\widehat{K_1}(\tau) = \int_0^\infty \exp\left(-s - 2\pi i s \tau\right) ds = \frac{1}{1+2\pi i \tau}$ has no real root and $\widehat{K_1}(0) = 1$. If moreover $\widehat{K_2}(0) = \int_0^\infty u^{-2} L(u) du = 1$ and the left side limit exists, then also the right side limit exists and the equality follows from the Wiener's Tauberian theorem. To prove the statement in the oposite direction it remains to show that $\widehat{K_2}(\tau) = \int_0^\infty u^{-2+2\pi i \tau} L(u) du$ has no real root.

To complete the proof by establishing the validity of assumptions concerning the function \widehat{K}_2 , we will evaluate the integral

$$\int_0^\infty u^{-2+2\pi i\tau} \left(1 - \Gamma\left(\frac{d}{2}\right) \frac{J_{\frac{d}{2}-1}\left(2\pi u\right)}{(\pi u)^{\frac{d}{2}-1}} \right) du.$$

Using $\int t^{-\nu} J_{\nu+1}(t) dt = -t^{-\nu} J_{\nu}(t)$ we get

$$2\pi\Gamma\left(\frac{d}{2}\right)\int_{0}^{\infty}u^{-2+2\pi i\tau}\int_{0}^{u}(\pi s)^{1-\frac{d}{2}}J_{\frac{d}{2}}(2\pi s)\,ds\,du$$

and integration by parts gives

$$\frac{2\pi\Gamma\left(\frac{d}{2}\right)}{1-2\pi i\tau}\left(-\left[u^{-1+2\pi i\tau}\int_{0}^{u}\frac{J_{\frac{d}{2}}\left(2\pi s\right)}{(\pi s)^{\frac{d}{2}-1}}ds\right]_{0}^{\infty}+\pi^{1-2\pi i\tau}\int_{0}^{\infty}\frac{J_{\frac{d}{2}}\left(2\pi u\right)}{(\pi u)^{\frac{d}{2}-2\pi i\tau}}du\right).$$

The first term is zero, as the limits in 0 and infinity are zero by l'Hospital formula and asymptotic properties of the Bessel function. From the formula $\int_0^\infty t^a J_\nu(2t) dt = \frac{1}{2}\Gamma\left(\frac{\nu+a+1}{2}\right)\Gamma\left(\frac{\nu-a+1}{2}\right)^{-1}$, valid if $Re \ a < \frac{1}{2}$, $Re \ a + \nu > -1$, it follows that

$$\widehat{K_{2}}\left(\tau\right) = \frac{\pi^{-2\pi i\tau - \frac{1}{2}}}{1 - 2\pi i\tau} \frac{\Gamma\left(\frac{d+1}{2}\right)\Gamma\left(\frac{1}{2} + \pi i\tau\right)}{\Gamma\left(\frac{d+1}{2} - \pi i\tau\right)}$$

Now it is easy to see that $\widehat{K}_2(0) = 1$, and \widehat{K}_2 has no real root because gamma function has none and because all poles of gamma function are negative.

Remark 2.6. If *D* is a bounded full-dimensional locally finite union of sets of finite reach, then the derivative of the covariance from the right $\overline{\gamma_D}'^+(0) = -\frac{\kappa_{d-1}}{d\kappa_d}H^{d-1}(\partial D)$ [9].

Theorem 2.7. Let **T** be a d-periodic lattice of points, $r \in \mathbb{R}^+$, D a bounded measurable set such that finite $\overline{\gamma_D}'^+(0)$ exists and Φ a function on \mathbb{R}^+ defined by equation

$$\int_{\mathbf{SO}_d} \int_{F_{\mathbf{T}}} \left(\operatorname{card} \left((rMD + x) \cap \mathbf{T} \right) - \alpha \lambda^d (D) \right)^2 \alpha \, d\lambda^d (x) \, dM =$$
$$= \frac{-\overline{\gamma_D}'^+ (0)}{2\pi^2 \kappa_{d-1}} \left(\sum_{0 \neq \xi \in \mathbf{T}^*} |\xi|^{-d-1} \right) \Phi(r) \, r^{d-1}.$$

Further, let $\rho^{d+1}\widehat{\gamma_D}(\rho)$ be bounded measurable on \mathbb{R}^+ . Then Φ is bounded and

(2.5)
$$\lim_{t \to \infty} \frac{1}{t} \int_0^t \Phi(t) \, dt = 1.$$

If moreover $\lim_{\rho\to\infty} \rho^{d+1} \widehat{\gamma_D}(\rho)$ exists then

(2.6)
$$\lim_{t \to \infty} \Phi(t) = 1.$$

Proof. From Remark 2.4 we have

$$\int_{\mathbf{SO}_{d}} \int_{F_{\mathbf{T}}} \left(\operatorname{\mathbf{card}} \left(\left(rMD + x \right) \cap \mathbf{T} \right) - \alpha \lambda^{d} \left(D \right) \right)^{2} \alpha d\lambda^{d} \left(x \right) dM = \sum_{0 \neq \xi \in \mathbf{T}^{*}} \widehat{\gamma_{D}} \left(r \left| \xi \right| \right) r^{2d}.$$

We shall prove first that the auxiliary function Ψ defined by the equation

$$-\overline{\gamma_D}'^+(0)\Psi(t) = 2\pi^2\kappa_{d-1}t^{d+1}\widehat{\overline{\gamma_D}}(t)$$

has the property (2.5) or (2.6). It is easy to see that the function

$$\Phi(t) = \frac{\sum_{0 \neq \xi \in \mathbf{T}^{*}} |\xi|^{-d-1} \Psi(t|\xi|)}{\sum_{0 \neq \xi \in \mathbf{T}^{*}} |\xi|^{-d-1}}$$

is bounded and has then the same property (2.5) or (2.6) as the auxiliary function Ψ .

Let $\rho^{d+1}\widehat{\gamma_D}(\rho)$ be bounded measurable on \mathbb{R}^+ . From the Lemma 2.5, Proposition 2.2 and Remark 2.4 it follows that

$$\lim_{R \to \infty} \frac{1}{R} \int_0^R 2\pi^2 \kappa_{d-1} \rho^{d+1} \widehat{\overline{\gamma_D}}(\rho) \ d\rho =$$
$$= \lim_{h \to 0+} \frac{1}{h} \int_0^\infty \left(d\kappa_{d-1} - 2\pi \left(h\rho\right)^{1-\frac{d}{2}} J_{\frac{d}{2}-1}\left(2\pi h\rho\right) \right) \rho^{d-1} \widehat{\overline{\gamma_D}}(\rho) \ d\rho =$$
$$= \lim_{h \to 0+} \frac{1}{h} \left(\overline{\gamma_D}(0) - \overline{\gamma_D}(h) \right) = -\overline{\gamma_D}'^+(0)$$

and if the left side limit exists then

$$\lim_{r \to \infty} 2\pi^2 \kappa_{d-1} r^{d+1} \widehat{\overline{\gamma_D}}(r) = -\overline{\gamma_D}'^+(0) \,.$$

Corollary 2.8. From Theorem 2.7 and Remark 2.6 it follows that if D is a bounded full-dimensional locally finite union of sets of finite reach such that $\rho^{d+1}\widehat{\gamma_K}(\rho)$ is bounded (or has a limit in $+\infty$), then

(2.7)
$$\int_{\mathbf{SO}_d} \int_{F_{\mathbf{T}}} \left(\operatorname{card} \left((rMD + x) \cap \mathbf{T} \right) - \alpha \lambda^d (D) \right)^2 \alpha \, d\lambda^d (x) \, dM = \\ = C_{\mathbf{T}} H^{d-1} \left(\partial D \right) \Phi (r) \, r^{d-1},$$

where

$$C_{\mathbf{T}} = \frac{1}{2\pi^2 d\kappa_d} \sum_{0 \neq n \in \mathbb{Z}^{\mathbf{d}}} \left| A^{-1} n \right|^{-d-1}$$

is a lattice constant and Φ fulfills (2.5) (or has limit 1).

3. Discussion

The assumption of the Theorem 2.7 and Corollary 2.8, that $\widehat{\gamma_D}(\rho)$ is $O(\rho^{-d-1})$, $\rho \to \infty$, holds for convex sets and sets with $C^{\frac{3}{2}}$ boundary (the boundary of the set can be decomposed into finitely many neighbourhoods such that given any pair of points P, Q in the neighbourhood, $|(P-Q)n(Q)| \leq c |P-Q|^{\frac{3}{2}}$, where n(Q) is a unit normal to the set in Q) [2]. The asymptotics $\sim r^{d-1}$ of the variance of the lattice points count in the mean value (2.7) thus holds for sets with piecewise \mathbb{C}^2 smooth boundary and convex sets in \mathbb{R}^d .

As was already said in the introduction, Theorem 2.7 gives results similar to Theorem 2.8 in [3] for a compact body D with smooth isotropic covariogram $\overline{\gamma}_D$. Slightly different situation is studied in [3]: locally finite periodic measure is studied instead of the point lattice and the size of the body is fixed while the scale s of the lattice tends to zero and the counting measure is multiplied by factor s^d ; consequently, the asymptotics $\sim s^{-d-1}$ is obtained there. As in [3] we may generalize

JIŘÍ JANÁČEK

the results of the present paper to locally finite periodic measures. In [3], lattice constants $C_{\rm T}$ in Equation 2.7 were calculated for some important point lattices.

Acknowledgement. This study was supported by the Academy of Sciences of the Czech Republic, grant No. A100110502 and AV0Z 50110509.

References

- S. Bochner, K. Chandrasekharan: Fourier transforms, Princeton University Press, 1949, Zbl 0152.11902
- [2] L. Brandolini, S. Hofmann, A. Iosevich: Sharp rate of average decay of Fourier transform of a bounded set. Geom. Func. Anal. 13 (2003), 671-680, Zbl 1032.42017
- [3] J. Janáček: Variance of periodic measure of bounded set with random position. Comment. Math. Univ. Carolinae 47 (2006), 473-482, Zbl pre05119961
- [4] D. G. Kendall: On the number of lattice points inside a random oval. Quarterly J. Math. 19 (1948), 1-26, Zbl 0031.11201
- [5] D. G. Kendall, R. A. Rankin: On the number of points of a given lattice in a random hypersphere. Quarterly J. Math., 2nd Ser., 4 (1953), 178-189, Zbl 0052.14503
- [6] B. Matérn: Precision of area estimation: a numerical study. J. Microsc. 153 (1989), 269-283
- [7] G. Matheron: Les variables regionalisees et leur estimation. Masson et CIE, Paris, 1965
- [8] R. C. Rao: Linear statistical inference and its applications, 2nd ed., John Wiley & Sons, New York, 1973, Zbl 0256.62002
- [9] J. Rataj: On set covariance and three-point test sets. Czech Mat. J. 54 (2004), 205-214, Zbl 1049.52004
- [10] G. N. Watson: A treatise on the theory of Bessel functions. 2nd edition, Cambridge University Press, 1922, Zbl 0849.33001
- [11] W. Rudin: Functional Analysis. McGraw-Hill Book Company, 1973, Zbl 0253.46001
- [12] A. Varchenko: Number of lattice points in families of homothetic domains in \mathbb{R}^n . Func. Anal. Appl. 17 (1983), 79-83, Zbl 0522.10031
- [13] N. Wiener: The Fourier integral and certain of its applications. Dover Publications Inc., New York, 1933, Zbl 0656.42001

Institute of Physiology, Academy of Sciences of the Czech Republic, Vídeňská 1083, 142 20 Praha, Czech Republic