

# ON HOPF ALGEBRAS AND THEIR GENERALIZATIONS

GIZEM KARAALI

ABSTRACT. We survey Hopf algebras and their generalizations. In particular, we compare and contrast three well-studied generalizations (quasi-Hopf algebras, weak Hopf algebras, and Hopf algebroids), and two newer ones (Hopf monads and hopfish algebras). Each of these notions was originally introduced for a specific purpose within a particular context; our discussion favors applicability to the theory of dynamical quantum groups. Throughout the note, we provide several definitions and examples in order to make this exposition accessible to readers with differing backgrounds.

## 1. INTRODUCTION - GOALS AND MOTIVATION

The purpose of this note is to provide a historical and comparative study of the several notions of generalized Hopf algebras that have been introduced and studied in recent years. We start with a brief discussion of Hopf algebras. We then consider the three main contenders for the *correct* notion of a generalized Hopf algebra: quasi-Hopf algebras, weak Hopf algebras, and Hopf algebroids. We then discuss two newer notions that generalize Hopf algebras: Hopf monads and hopfish algebras. As these are newer concepts, our study of them is necessarily rather cursory.

In our pursuit we trace the steps of many researchers, both mathematically and philosophically. Hence, this is an expository note, with no claim of introducing new mathematics. Its sole purpose is to collect together several pieces of interesting mathematics and present them in one comprehensive historical narrative, so as to provide a broader perspective on the current status of research involving various generalizations of Hopf algebras.

---

*Date:* August 18, 2007.

*2000 Mathematics Subject Classification.* Primary 17B62, 17B20.

*Key words and phrases.* Hopf algebra, quantum group, quasi-Hopf algebra, weak Hopf algebra, Hopf algebroid, Hopf monad, hopfish algebra.

The technical details and basic examples provided are intended to make this note accessible to a beginner in the theory of Hopf algebras, while we hope that those with more experience will still enjoy reading the discussions.

**Acknowledgments.** The author would like to thank Marcelo Aguiar, Gabriella Böhm, Ignacio Lopez Franco, Ludmil Hadjiivanov, Mitja Mastnak, Susan Montgomery, Frederic Patras, Fernando Souza, Ivan Todorov, Alexis Virelizier, Alan Weinstein and Milen Yakimov for helpful comments and useful questions during various stages of the work that led to this paper.

## 2. HOPF ALGEBRAS

Naturally, we begin with the definition of a Hopf algebra:

**Definition 2.1.** A **Hopf algebra** over a commutative ring  $\mathbb{K}$  is a  $\mathbb{K}$ -module  $H$  such that:

- (1)  $H$  is an associative unital  $\mathbb{K}$ -algebra (with  $m : H \otimes H \rightarrow H$  as the *multiplication* and  $u : \mathbb{K} \rightarrow H$  as the *unit*) and a coassociative counital  $\mathbb{K}$ -coalgebra (with  $\Delta : H \rightarrow H \otimes H$  as the *comultiplication* and  $\epsilon : H \rightarrow \mathbb{K}$  as the *counit*);
- (2) The comultiplication  $\Delta$  and the counit  $\epsilon$  are both algebra homomorphisms;
- (3) The multiplication  $m$  and the unit  $u$  are both coalgebra homomorphisms;
- (4) There is a bijective  $\mathbb{K}$ -module map  $S : H \rightarrow H$ , called the *antipode*, such that for all elements  $h \in H$ :

$$S(h_{(1)})h_{(2)} = (u \circ \epsilon)(h) = h_{(1)}S(h_{(2)}). \quad (2.1)$$

We first note here that in the defining formula for the antipode, we are making use of the famous *Sweedler notation*<sup>1</sup>, which is quite well-established in the Hopf algebra literature. In short the Sweedler notation is a generalization of the Einstein notation (in that it intrinsically demands a summation). More specifically for any  $h \in H$ , we write:

$$\Delta(h) = h_{(1)} \otimes h_{(2)}.$$

---

<sup>1</sup>In the introduction to [79], Sweedler remarks that this notation was developed during years of joint work with Robert Heyneman. However, it is much more common to see references to Sweedler notation than to Heyneman-Sweedler notation. We follow the tradition here.

This presentation itself is purely symbolic; the terms  $h_{(1)}$  and  $h_{(2)}$  do not stand for particular elements of  $H$ . The comultiplication  $\Delta$  takes values in  $H \otimes H$ , and so we know that:

$$\Delta(h) = (h_{1,1} \otimes h_{1,2}) + (h_{2,1} \otimes h_{2,2}) + (h_{3,1} \otimes h_{3,2}) + \cdots + (h_{N,1} \otimes h_{N,2})$$

for some elements  $h_{i,j}$  of  $H$  and some integer  $N$ . The Sweedler notation is just a way to separate the  $h_{i,1}$  from the  $h_{j,2}$ . In other words, one can say that the notation  $h_{(1)}$  stands for the generic  $h_{i,1}$  and the notation  $h_{(2)}$  stands for the generic  $h_{j,2}$ . However the summation is inherent in the notation: Whenever one sees a term like  $S(h_{(1)})h_{(2)}$  (the left hand side of (2.1)), one has to realize that this stands for a sum of the form:

$$S(h_{1,1}) \otimes h_{1,2} + S(h_{2,1}) \otimes h_{2,2} + S(h_{3,1}) \otimes h_{3,2} + \cdots + S(h_{N,1}) \otimes h_{N,2}.$$

A second remark that should be made here involves commutative diagrams. In most texts introducing the notion of a Hopf algebra, one will come across commutative diagrams similar to the following:

$$\begin{array}{ccc} H \otimes H & \xrightarrow{S \otimes id} & H \otimes H \\ \Delta \uparrow & & \downarrow m \\ H & \xrightarrow{u \circ \epsilon} & H \end{array} \qquad \begin{array}{ccc} H \otimes H & \xrightarrow{id \otimes S} & H \otimes H \\ \Delta \uparrow & & \downarrow m \\ H & \xrightarrow{u \circ \epsilon} & H \end{array}$$

In fact these two diagrams are equivalent to the single-line equation (2.1), but for those comfortable with commutative diagrams, they provide a much more visual way to understand it. Incidentally an even more compact way to describe  $S$  would be as the convolution inverse of the identity homomorphism on  $H$ .

For the sake of completeness, we include here the diagrams representing the properties of the counit (the so-called *counit axiom*):

$$\begin{array}{ccccc} & & H & & \\ & \Delta \swarrow & | & \searrow \Delta & \\ H \otimes H & & id_H & & H \otimes H \\ & \epsilon \otimes 1 \searrow & \downarrow & \swarrow 1 \otimes \epsilon & \\ & & H & & \end{array}$$

(where we use  $H \cong H \otimes \mathbb{K} \cong \mathbb{K} \otimes H$ ), and the comultiplication (the coassociativity):

$$\begin{array}{ccc}
 H & \xrightarrow{\Delta} & H \otimes H \\
 \Delta \downarrow & & \downarrow \Delta \otimes 1 \\
 H \otimes H & \xrightarrow{1 \otimes \Delta} & H \otimes H \otimes H
 \end{array}$$

We also should point out that the second and third conditions of Definition 2.1 are equivalent to one another and, together with the first condition, define what is called a  $\mathbb{K}$ -bialgebra:

**Definition 2.2.** A **bialgebra** over a commutative ring  $\mathbb{K}$  is a  $\mathbb{K}$ -module  $B$  such that:

- (1)  $B$  is both an associative unital  $\mathbb{K}$ -algebra (with  $m : B \otimes B \rightarrow B$  as the *multiplication* and  $u : \mathbb{K} \rightarrow B$  as the *unit*) and a coassociative counital  $\mathbb{K}$ -coalgebra (with  $\Delta : B \rightarrow B \otimes B$  as the *comultiplication* and  $\epsilon : B \rightarrow \mathbb{K}$  as the *counit*);
- (2) The comultiplication  $\Delta$  and the counit  $\epsilon$  are both algebra homomorphisms.

Then we can define a Hopf algebra to be a bialgebra with an antipode, in other words, a bialgebra  $B$  with an antihomomorphism  $S : B \rightarrow B$  satisfying Equation (2.1).

Hopf algebras were first introduced in the early 1940s by Hopf in [44], where he was working on homology rings of certain compact manifolds. (For a modern exposition of his results using the language of Hopf algebras, we refer the reader to [60]). Later on, more examples were discovered and basic references started appearing, see for instance the early classics like Sweedler's [79] and Abe's [1]. For a more modern approach emphasizing actions of Hopf algebras, see Montgomery's [64].

For a simple example of Hopf algebras, consider the group algebra  $\mathbb{K}[G]$  of a group  $G$  over a field  $\mathbb{K}$ . Here we define the coalgebra structure by defining, on the elements  $g$  of  $G$ , the comultiplication and the counit by setting  $\Delta(g) = g \otimes g$  and  $\epsilon(g) = 1$ . (More generally, for an arbitrary coalgebra  $C$ , the elements  $g$  of  $C$  satisfying  $\Delta(g) = g \otimes g$  and  $\epsilon(g) = 1$  are called *group-like*). The antipode is defined on the group elements by  $S(g) = g^{-1}$ .

Dually, we can look at the algebra  $F_{\mathbb{K}}(G)$  of  $\mathbb{K}$ -valued functions on a group  $G$ .<sup>2</sup> There is a natural (commutative) multiplication  $m$  on this algebra given by:

$$m(f_1, f_2)(g) = f_1(g)f_2(g) \text{ for all } g \in G.$$

The unit just maps elements of the field  $\mathbb{K}$  to the associated constant functions. To define a comultiplication, one needs to use the natural embedding of  $F_{\mathbb{K}}(G) \otimes F_{\mathbb{K}}(G)$  into  $F_{\mathbb{K}}(G \times G)$ . Then the group multiplication  $m_G : G \times G \rightarrow G$  on  $G$  induces a comultiplication as follows:

$$\Delta(f)(g_1, g_2) = f(m_G(g_1, g_2)).$$

The counit is the map taking  $f \in F_{\mathbb{K}}(G)$  to  $f(e_G) \in \mathbb{K}$  where  $e_G$  is the identity element of the group  $G$ . Finally the antipode is the map  $S$  defined by  $S(f)(g) = f(g^{-1})$ .

Function algebras give us a large collection of examples of *commutative* Hopf algebras, (which are just Hopf algebras with commutative multiplications). In fact, one can show that any finitely generated commutative Hopf algebra over a field  $\mathbb{K}$  of characteristic zero is the function algebra of an affine algebraic group  $G$  over  $\mathbb{K}$ . We will come back to this in Section 8.

Another family of examples is given by universal enveloping algebras  $U(\mathfrak{g})$  of Lie algebras  $\mathfrak{g}$ . In this case, the coalgebra structure is defined on the elements  $x$  of  $\mathfrak{g}$  by:

$$\Delta(x) = x \otimes 1 + 1 \otimes x; \quad \epsilon(x) = 0.$$

(More generally, for an arbitrary coalgebra  $C$ , elements  $x$  of  $C$  satisfying  $\Delta(x) = x \otimes 1 + 1 \otimes x$  and  $\epsilon(x) = 0$  are called *primitive*). The antipode  $S$  is defined on the Lie algebra elements as  $S(x) = -x$ .

This last class of examples provides us with a large collection of *cocommutative* Hopf algebras. Cocommutativity is the property of the comultiplication  $\Delta$  described by the following commutative diagram:

$$\begin{array}{ccc} H \otimes H & \xleftarrow{T} & H \otimes H \\ \Delta \uparrow & & \uparrow \Delta \\ H & \xleftarrow{id} & H \end{array}$$

---

<sup>2</sup>We are intentionally being vague, in order to include several classes of algebras here; if, for instance,  $G$  is an (affine) algebraic group, we could be thinking of the coordinate ring of  $G$ ; if  $G$  is a Poisson group, we could be looking at the Poisson algebra  $C^\infty(G)$ .

which is exactly the diagram one obtains by switching the directions of the arrows and replacing  $m$  by  $\Delta$  in the diagram describing commutativity of a product  $m$ :

$$\begin{array}{ccc}
 H \otimes H & \xrightarrow{T} & H \otimes H \\
 m \downarrow & & \downarrow m \\
 H & \xrightarrow{id} & H
 \end{array}$$

In both of these diagrams  $T : H \otimes H \rightarrow H \otimes H$  is the usual twist map:  $T(a \otimes b) = b \otimes a$ .

After seeing the result above about commutative Hopf algebras, one may expect an analogous result regarding cocommutative Hopf algebras. In fact one can show that a cocommutative Hopf algebra generated by its primitive elements is indeed the universal enveloping algebra of a Lie algebra. This result will be mentioned in Section 8 as well.

Ever since their introduction, Hopf algebras have been studied by many mathematicians. In the early 1970s, Hochschild, while developing the theory of algebraic groups, translated much of representation theory into the language of Hopf algebras. (See for instance [42] for a classic written with this perspective; [33] is a more modern text using the same approach). In some sense, we can say that Hopf algebras provided mathematicians the ultimate framework to do representation theory.

The development of the theory of quantum groups brought a fresh revival of interest in Hopf algebras.<sup>3</sup> Since the early 1980s, many mathematicians have been working on structures which today we loosely call *quantum groups*. The earliest examples of quantum groups were particular deformations of universal enveloping algebras of simple Lie algebras and function algebras of simple algebraic groups. In mid 1980s, Drinfeld showed that the correct framework to use when studying quantum groups was that of Hopf algebras; see, for instance, his ICM address [23]. Thus the known world of Hopf algebras was significantly expanded to include all these new examples, and results from quantum groups began to add more spice and flavor to the classical theory.

Broadly speaking, a quantum group is a special type of noncommutative noncocommutative Hopf algebra. One obtains such Hopf algebras

---

<sup>3</sup>In the following paragraphs, we will provide only a brief sketch of the ideas of quantum group theory and refer the more interested reader to one of many textbooks and monographs in the subject (eg. [15, 19, 28, 43, 48, 50, 54, 58, 74]).

by deforming the multiplication or the comultiplication of a commutative or a cocommutative Hopf algebra. Since the end result of such a deformation is not commutative, it cannot properly be associated to a group and be its function algebra. Similarly, since it is not cocommutative, it is by no means the enveloping algebra of a Lie algebra. However, we can still view it as if it were the function algebra or the enveloping algebra of some phantom group or Lie algebra, and thus we have the term “*quantum group*”.<sup>4</sup>

For the sake of completeness we briefly describe the most well-known quantum group here: *Quantum  $sl_2(\mathbb{C})$* , denoted by  $U_h(sl_2(\mathbb{C}))$ . The notation is more illuminating than the name; this is a particular (Hopf algebra) deformation of the universal enveloping algebra  $U(sl_2(\mathbb{C}))$  of  $sl_2(\mathbb{C})$ .

As an algebra, it is generated by  $E, F, H$  subject to the following relations:

$$[H, E] = 2E, \quad [H, F] = -2F, \quad [E, F] = \frac{q^H - q^{-H}}{q - q^{-1}}$$

where  $h$  is viewed as a formal parameter,  $q = e^h = \sum_{n=0}^{\infty} \frac{h^n}{n!} \in \mathbb{C}[[h]]$ , and  $q^H, q^{-H}$  should be interpreted in a similar manner. The coalgebra structure is defined on the generators by:

$$\begin{aligned} \Delta(E) &= E \otimes q^H + 1 \otimes E; & \epsilon(E) &= 0; \\ \Delta(F) &= F \otimes 1 + q^{-H} \otimes F; & \epsilon(F) &= 0; \\ \Delta(H) &= H \otimes 1 + 1 \otimes H; & \epsilon(H) &= 0. \end{aligned}$$

Finally we define the antipode on the generators as:

$$S(E) = -Eq^{-H}; \quad S(F) = -q^H F; \quad S(H) = -H.$$

Extended linearly, these give us a Hopf algebra structure, the famous quantum  $sl_2$ .

Currently, there is no consensus on what the precise definition of a quantum group should be. Mainly, there are several well-known examples and an exponentially growing literature investigating their properties. For the purposes of this note, it will suffice to identify the term quantum group with quasitriangular or co-quasitriangular Hopf algebras, i.e. noncommutative noncocommutative Hopf algebras associated to solutions of a certain equation, the quantum Yang-Baxter Equation, which we will describe in the next section.

---

<sup>4</sup>This may remind some readers the philosophy of noncommutative geometry a la Connes. A similar approach to quantum groups would involve viewing them as symmetry objects of some quantum space; see [59].

## 3. LONG INTERLUDE - WHY GENERALIZE?

We believe that the previous section provides a sufficient overview of Hopf algebras, and prepares the reader for the discussion on the generalizations of this classical theory. For the reader who wishes to learn more about Hopf algebras, we recommend the references already mentioned above, as well as any textbook on quantum groups.<sup>5</sup>

Before we move on to the comparative study and technical details of the various generalizations, we would like to philosophize a bit about why we are interested in *any* generalization. To those who view mathematics as an intellectual pursuit merely interested in pure abstractions, the answer will be clear: *Why not?* However, for those who may need more motivation, we will provide one, which comes from the theory of quantum groups.

The study of quantum groups goes back to the well-known *quantum Yang-Baxter equation*, which in its simplest form is as follows:

$$R^{12}R^{13}R^{23} = R^{23}R^{13}R^{12}. \quad (3.1)$$

Here we can view  $R$  as a map  $R : G \times G \rightarrow G \times G$  for some factorizable Poisson-Lie group (i.e. a Lie group with a Poisson bracket on its function algebra). Then:

$$R^{12}(g, h, k) = (R(g, h), k),$$

and  $R^{13}$  and  $R^{23}$  can be defined likewise. In the realm of Hopf algebras, the *quantum Yang-Baxter equation* (henceforth referred to as the QYBE) is the same equation (3.1); however, this time its solutions, the so-called *quantum R-matrices*, are linear maps of  $\mathbb{K}$ -modules:  $R \in H \otimes H$ . For surveys on the quantum Yang-Baxter equation, displaying the several connections to the physics literature, see [45, 46].

The QYBE and its solutions give rise to quantum groups: in the geometric (Poisson) picture, they give a (Hopf algebra) deformation of the function algebra of the relevant group  $G$ ; in the Lie algebra picture, they give a (Hopf algebra) deformation of the associated universal enveloping algebra. In the case of quantum  $sl_2$  as presented above, the relevant quantum  $R$ -matrix is:

---

<sup>5</sup>Among the many books on quantum groups which also have a detailed exposition of Hopf algebras, we list here only a few: [19], [28], [48], [58], [74]. A quick introduction to Hopf algebras and their applications may be found in [40]. Other references about Hopf algebras which especially focus on combinatorial applications are [65, 2].



$$R = q^{\frac{1}{2}H \otimes H} \sum_{n \geq 0} q^{\frac{n(n-1)}{2}} \frac{[q - q^{-1}]^n}{[n]_q!} E^n \otimes F^n.$$

Here we are using the  $q$ -notation common in the literature:

$$[n]_q = \frac{q^n - q^{-n}}{q - q^{-1}} = q^{n-1} + q^{n-2} + \cdots + q^{-n+1} \text{ and } [n]_q! = \prod_{k=1}^n [k]_q$$

In 1984 [34], a modified version of the QYBE appeared in the mathematical physics literature. This new equation, later named the *quantum dynamical Yang-Baxter equation*, (henceforth labeled QDYBE), lives in  $V \otimes V \otimes V$  for a semisimple module  $V$  of an abelian Lie algebra  $\mathfrak{h}$ :

$$R^{12}(\lambda - h^{(3)})R^{13}(\lambda)R^{23}(\lambda - h^{(1)}) = R^{23}(\lambda)R^{13}(\lambda - h^{(2)})R^{12}(\lambda). \quad (3.2)$$

Here, the term  $h^{(i)}$  is to be substituted by  $\mu_i$  if  $\mu_i$  is the weight of the  $i$ th tensor component,  $i = 1, 2, 3$ . A solution  $R : \mathfrak{h}^* \rightarrow \text{End}_{\mathfrak{h}}(V \otimes V)$  to the QDYBE is called a *quantum dynamical  $R$ -matrix*. For an introduction to the dynamical Yang-Baxter equations along with some of their solutions, we refer the reader to [30]. A more geometric exposition can be found in [29].

Studying the QDYBE and its solutions, we get into the realm of *dynamical quantum groups*. The first examples of dynamical quantum groups that appeared in the literature are Felder's *elliptic quantum groups*, which were introduced in [30, 31]. A standard example that is studied in much detail is  $E_{\tau, \eta}(sl_2)$ , which is an algebra over  $\mathbb{C}$  generated by two kinds of generators: meromorphic functions of a single variable  $h$ , and matrix elements of a matrix  $L(\lambda, w) \in \text{End}(\mathbb{C}^2)$ . The two subscripts  $\tau, \eta$  are nonzero complex numbers with  $\text{Im}(\tau) > 0$ .

Without going into much detail, we note that the  $4 \times 4$  matrix solution to the QDYBE associated with  $E_{\tau, \eta}(sl_2)$  is in the following form:

$$\begin{aligned} R(\lambda, w, \tau, \eta) &= E_{1,1} \otimes E_{1,1} + E_{2,2} \otimes E_{2,2} \\ &+ \alpha(\lambda, w, \tau, \eta) E_{1,1} \otimes E_{2,2} + \beta(\lambda, w, \tau, \eta) E_{1,2} \otimes E_{2,1} \\ &+ \gamma(\lambda, w, \tau, \eta) E_{2,1} \otimes E_{1,2} + \delta(\lambda, w, \tau, \eta) E_{2,2} \otimes E_{1,1} \end{aligned}$$

Here the functions  $\alpha, \beta, \gamma, \delta$  are defined in terms of the theta function:

$$\theta(z, \tau) = -\sum_{j=-\infty}^{\infty} \exp(\pi i(j + \frac{1}{2})^2 \tau + 2\pi i(j + \frac{1}{2})(z + \frac{1}{2}))$$

by:

$$\alpha(\lambda, w, \tau, \eta) = \frac{\theta(w)\theta(\lambda + 2\eta)}{\theta(w - 2\eta)\theta(\lambda)} \quad , \quad \beta(\lambda, w, \tau, \eta) = \frac{\theta(-w - \lambda)\theta(2\eta)}{\theta(w - 2\eta)\theta(\lambda)}$$

$$\gamma(\lambda, w, \tau, \eta) = \frac{\theta(w - \lambda)\theta(2\eta)}{\theta(w - 2\eta)\theta(\lambda)} \quad , \quad \delta(\lambda, w, \tau, \eta) = \frac{\theta(w)\theta(\lambda - 2\eta)}{\theta(w - 2\eta)\theta(\lambda)}$$

For more on  $E_{\tau, \eta}(sl_2)$ , including an explicit presentation in terms of generators and relations, we refer the reader to [32].

The objects, like Felder's elliptic quantum groups, associated to solutions of the QDYBE turn out to be quite Hopf-like in many respects, but they are not all necessarily Hopf algebras. Or in other words, we can say that, just as in the case of quantum groups, some Hopf-like structures come into play when one studies the solutions of the QDYBE. And herein lies our motivation for the purpose of this particular note. A need for the *correct Hopf-like object* that will provide the framework for the theory of dynamical quantum groups leads us to the study of various generalizations of Hopf algebras.

#### 4. CANDIDATE 1 : QUASI-HOPF ALGEBRAS

Our first candidate in our search for the correct generalization of Hopf algebras is the quasi-Hopf algebra, first introduced by Drinfeld in [24, 25], and used to give a natural proof of Kohno's theorem relating the monodromy of the Knizhnik-Zamolodchikov equations to a representation of the braid group arising from a quantum group. In order to define quasi-Hopf algebras, we first need to introduce the notion of a *quasi-bialgebra*:

**Definition 4.1.** A **quasi-bialgebra**  $B$  over a commutative ring  $\mathbb{K}$  is a unital associative  $\mathbb{K}$ -algebra equipped with two algebra homomorphisms  $\epsilon : B \rightarrow \mathbb{K}$  (the *counit*) and  $\Delta : B \rightarrow B \otimes B$  (the *comultiplication*) together with an invertible element  $\Phi$  of  $B \otimes B \otimes B$ . Furthermore we require the following to hold:

(1)  $\Phi$  satisfies the *pentagon relation*:

$$(\Delta \otimes 1 \otimes 1)\Phi \cdot (1 \otimes 1 \otimes \Delta)\Phi = (\Phi \otimes 1) \cdot (1 \otimes \Delta \otimes 1)\Phi \cdot (1 \otimes \Phi)$$

(2)  $\Delta$  is *quasi-coassociative*:

$$(1 \otimes \Delta)\Delta(b) = \Phi^{-1}(\Delta \otimes 1)\Delta(b)\Phi \text{ for all } b \in B$$

(3)  $\Phi$  is compatible with the counit  $\epsilon$  in the following sense:

$$(1 \otimes \epsilon \otimes 1)\Phi = 1$$

(4)  $\epsilon$  satisfies the counit axiom.<sup>6</sup>

---

<sup>6</sup>We presented the commutative diagram version of the counit axiom earlier, in closed form it reads:

$$(\epsilon \otimes 1)\Delta = 1 = (1 \otimes \epsilon)\Delta.$$

From (1), (3) and (4), we get:

$$(\epsilon \otimes 1 \otimes 1)\Phi = (1 \otimes \epsilon \otimes 1)\Phi = (1 \otimes 1 \otimes \epsilon)\Phi = 1.$$

$\Phi$  of this definition is alternatively called an *associator* or a *co-associator*. We will call it the associator, merely to ease our typing efforts. However we should note that both versions have merit. The term “*co-associator*” makes sense because  $\Phi$  gives information on how far  $\Delta$  is from coassociativity. The term “*associator*” makes sense because using  $\Phi$ , one can define associativity isomorphisms in the representation category of  $B$  which makes it into a (non-strict) monoidal category. The pentagon relation (1) satisfied by  $\Phi$  in the above definition translates automatically to the pentagon relations of the associativity isomorphisms in the representation category of  $B$ .

Comparing Definitions 2.2 and 4.1, it is clear that this generalization has cost us only the coassociativity of the comultiplication. However this price is perfectly acceptable to many researchers, as quasibialgebras and their Hopf relatives fit very nicely into several theories. We will discuss these shortly.

Here is the definition of the Hopf version:<sup>7</sup>

**Definition 4.2.** A **quasi-Hopf algebra**  $H$  over a commutative ring  $\mathbb{K}$  is a quasi-bialgebra over  $\mathbb{K}$  equipped with an (algebra and coalgebra) antihomomorphism  $S : H \rightarrow H$  and two canonical elements  $\alpha, \beta \in H$  such that:

- (1)  $m \cdot (S \otimes \alpha)\Delta(b) = \epsilon(b)\alpha$  for all  $b \in B$ ;
- (2)  $m \cdot (1 \otimes \beta S)\Delta(b) = \epsilon(b)\beta$  for all  $b \in B$ ;
- (3)  $m(m \otimes 1) \cdot (S \otimes \alpha \otimes \beta S)\Phi = 1$ ;
- (4)  $m(m \otimes 1) \cdot (1 \otimes \beta S \otimes \alpha)\Phi^{-1} = 1$ .

We note that (1) and (2) are the modified versions of the antipode axiom (2.1). If  $\alpha = \beta = 1$ , and  $\Phi = 1 \otimes 1 \otimes 1$ ,  $H$  clearly becomes a Hopf algebra.

Drinfeld introduced quasi-Hopf algebras in his work relating the Knizhnik-Zamolodchikov equations to the theory of quantum groups.

---

<sup>7</sup>We note here that our definitions in this section are not quite as general as the original ones in [25]. Drinfeld’s original definitions involved two invertible elements  $l$  and  $r$ , or equivalently a left unit constraint  $l$  and a right unit constraint  $r$ , satisfying the so-called *Triangle Axiom*. However, Drinfeld showed, also in [25], that he could always reduce his quasi-Hopf algebras into quasi-bialgebras where  $r = l = 1$ . Therefore we will not be worried much about our more restrictive definitions.

However, these structures proved to be a lot more than mere tools in one proof. The large interest they invoked in the mathematicians in the field can easily be deduced, for example, from J. Stasheff's enthusiastic MathSciNet review *MR1091757* of Drinfeld's [25].

One general reason for this excitement can be explained, in category-theoretic terms, as follows: Quasi-Hopf algebras have the right kind of representation categories. In particular their representation categories are (not necessarily strict) rigid monoidal categories.<sup>8</sup>

Recall that a monoidal category is *strict* if its associativity isomorphism is canonical. It can be shown that any (nice enough) strict monoidal category is the representation category of a bialgebra, and any (nice enough) rigid strict monoidal category is the representation category of a Hopf algebra.<sup>9</sup> A well-known theorem of MacLane asserts that every non-strict monoidal category is in fact tensor equivalent to a strict one. So it is natural to look at algebraic objects whose representation categories are non-strict monoidal because these have in essence the same type of representation theory as that of bialgebras and Hopf algebras.

It turns out that quasi-Hopf algebras are just the right structures in this perspective! More specifically, quasi-bialgebras are precisely those algebras, equipped with counit and comultiplication, whose representation categories are monoidal. In fact in [48], quasi-bialgebras are defined to be those structures which have monoidal representation categories. (This is immediately followed by a proof of the equivalence of this definition to the definition of Drinfeld in [25]). Likewise quasi-Hopf algebras are structures with rigid monoidal representation categories.

When the first examples of dynamical quantum groups began to appear, people noticed that they were not Hopf algebras, but still looked very much like them. Therefore several researchers focused on quasi-Hopf algebras, as structures already under serious inspection for the

---

<sup>8</sup>Here we digress briefly and venture into category theory. In Section 8, we will give more details. The classic reference for the terms we use is [56]. [48] provides ample background and all the details within the context of Hopf algebras and quantum groups. Note that in [48], our monoidal categories are called *tensor categories*. These are exactly the “*categories with multiplication*” of [4, 55]. Also see the historical notes at the end of Chapter VII in [56].

<sup>9</sup>Obviously the phrase “*nice enough*” needs to be explained for this vague sentence to become a precise (and correct) mathematical statement. For instance, for a Tannaka-Krein type construction (cf. Section 8), we need to construct a fiber functor from our category into the category of vector spaces. We will not attempt this here.

reasons mentioned earlier, with the expectation that these could possibly provide the right Hopf-like theory to describe dynamical quantum groups. Indeed, it turns out that Felder’s elliptic quantum groups naturally fit into the framework of quasi-Hopf algebras, [26].

Besides Felder’s elliptic quantum groups, the fundamental example of a nontrivial quasi-Hopf algebra (i.e. one that is not a Hopf algebra) is the one constructed by Drinfeld. This construction involves the monodromy of the Knizhnik-Zamolodchikov equations and is beyond the scope of this note. We refer the interested reader to the original work of Drinfeld in [25] and the more pedagogical exposition in [28].

The representation theory of quasi-Hopf algebras is quite exciting. Category-theoretic arguments ensure that quasi-objects will generally be only a twist away from their non-quasi counterparts.<sup>10</sup> More precisely, MacLane’s theorem about the (tensor-)equivalence of non-strict monoidal categories to strict ones implies that the representation theory of a quasi-object is quite similar to that of a non-quasi one.

Tensor equivalence in the realm of monoidal categories translates into the language of equivalence of quasi-bialgebras and quasi-Hopf algebras in terms of twists in the algebraic realm. In many cases (when a certain cohomology vanishes), the twist will involve a trivial isomorphism on the level of the algebra. In other words, the quasi-structure will be “*twistable*” into a non-quasi one on the same underlying algebra. In other cases we will need to modify the underlying set. Once again the reader is referred to [28], for a discussion of the use of twists of quasi-bialgebras and quasi-Hopf algebras.

Overall, quasi-Hopf algebras are interesting structures, with many researchers still investigating their various applications. Their representation theory is also very appealing. Moreover as we mentioned earlier, they can be used to describe the elliptic quantum groups of Felder. Nonetheless, we continue our search and look for other alternatives. This is mainly due to the general fact that algebraists are not very much delighted by a non-associative or a non-coassociative structure. The relaxation of these properties makes description of actions and coactions a lot more cumbersome, and this is typically not very desirable. Another issue is the notion of duality for quasi-objects. Since the definition of a quasi-bialgebra is not self-dual (the underlying set is an associative algebra but the coalgebra structure is not coassociative), the natural object that should be the dual of a quasi-bialgebra is not a

---

<sup>10</sup>In this context, several authors prefer to use terms like “*gauge transformation*” or “*skrooching*” in place of the term “*twist*.”

quasi-bialgebra; and similarly the natural object which should be the dual of a quasi-Hopf algebra is not a quasi-Hopf algebra. One needs to define separately a dual quasi-bialgebra and hence a dual quasi-Hopf algebra; see for instance the preliminaries of [57]. As a consequence, we do not have a natural notion of a comodule or a Hopf module over a quasi-object and any other mathematical construction that relies on these cannot be generalized easily to the quasi-setup.<sup>11</sup>

## 5. CANDIDATE 2 : WEAK HOPF ALGEBRAS

Following the algebraists' intuition to avoid non-coassociativity and unnatural ways to define duals, we move on to our second candidate: Weak Hopf algebras. This time our objects are both algebras and coalgebras, associative and coassociative, respectively, but the relationship between the two types of structures is weakened. In particular we no longer force the coalgebra structure to respect the unit of the algebra structure. In other words, we drop the requirement that the comultiplication be unit preserving. We instead write:

$$\Delta(1) = 1_{(1)} \otimes 1_{(2)} \neq 1 \otimes 1.$$

This, in turn, forces the counit to be *weakly multiplicative*:

$$\epsilon(xy) = \epsilon(x1_{(1)})\epsilon(1_{(2)}y).$$

Here is our precise definition:<sup>12</sup>

**Definition 5.1.** A (finite) **weak Hopf algebra** over  $\mathbb{K}$  is a finite-dimensional<sup>13</sup>  $\mathbb{K}$ -vector space  $H$  with the structures of an associative algebra  $(H, m, 1)$ , and a coassociative coalgebra  $(H, \Delta, \epsilon)$  such that:

- (1) The comultiplication  $\Delta$  is a (not necessarily unit-preserving) homomorphism of algebras such that:

$$(\Delta \otimes id)\Delta(1) = (\Delta(1) \otimes 1)(1 \otimes \Delta(1)) = (1 \otimes \Delta(1))(\Delta(1) \otimes 1);$$

---

<sup>11</sup>However, also see [35, 73].

<sup>12</sup>We should point out that there is an alternative use in the literature for the term *weak Hopf algebra*. Some authors use the term to refer to a bialgebra  $B$  with a *weak antipode*  $S$  in  $\text{Hom}(B, B)$ , i.e.,  $id * S * id = id$  and  $S * id * S = S$ , where  $*$  is the convolution product; see for instance [51, 52]. [85] briefly discusses how the two concepts are related.

<sup>13</sup>We define only the finite-dimensional version here. In this context, it seems to be standard to restrict to finite dimensions because it is much easier to dualize the notion in that case. For a more detailed discussion of why finite-dimensionality is in general preferred, we refer the reader to [11].

(2) The counit  $\epsilon$  is a  $\mathbb{K}$ -linear map satisfying the identity:

$$\epsilon(fgh) = \epsilon(fg_{(1)})\epsilon(g_{(2)}h) = \epsilon(fg_{(2)})\epsilon(g_{(1)}h)$$

for all  $f, g, h \in H$ ;

(3) There is a linear map  $S : H \rightarrow H$ , called an **antipode**, such that for all  $h \in H$ ,

$$\begin{aligned} m(id \otimes S)\Delta(h) &= (\epsilon \otimes id)(\Delta(1)(h \otimes 1)); \\ m(S \otimes id)\Delta(h) &= (id \otimes \epsilon)((1 \otimes h)\Delta(1)); \\ m(m \otimes id)(S \otimes id \otimes S)(\Delta \otimes id)\Delta(h) &= S(h). \end{aligned}$$

We can easily see that the condition on the counit implies weak multiplicativity. We also note that the defining equations for the antipode given above would coincide with Equation (2.1) if  $\Delta$  were to preserve the unit. This would also imply that the counit would be a homomorphism of algebras, and we would thus end up with a Hopf algebra.

Weak Hopf algebras behave much better with respect to duality. In other words, given a weak Hopf algebra  $A$  over a field  $\mathbb{K}$ , the space  $Hom_{\mathbb{K}}(A, \mathbb{K})$  can be given a natural weak Hopf algebra structure, using the canonical pairing  $\langle \cdot, \cdot \rangle : Hom_{\mathbb{K}}(A, \mathbb{K}) \times A \rightarrow \mathbb{K}$ .

Weak Hopf algebras in the above sense were introduced in [11, 12, 68, 75]. Later on it was observed that the paragroups of [69] and the face algebras of [36, 37, 38] fit nicely into this description. Many of the original constructions of weak Hopf algebras were motivated by applications to operator algebras, but many saw from early on their relationship to “*dynamical deformations of quantum groups*.” We refer the reader to [67] for a more detailed exposition of weak Hopf algebras, and their relationship to various generalizations of the idea of quantum groups. This reference also contains discussion of various examples of weak Hopf algebras, one of which we present here.

Recall that group algebras were basic examples of Hopf algebras. We will look at a nice generalization of these to find our weak Hopf algebras. In particular we will need to generalize the notion of a *group* via the following:

**Definition 5.2.** A **groupoid over a set**  $X$  is a set  $G$  together with the following structure maps:

(1) A pair of maps  $s, t : G \rightarrow X$ , respectively called the *source* and the *target*.

- (2) A *product*  $m$ , i.e. a partial function  $m : G \times G \rightarrow G$  satisfying the following two properties:<sup>14</sup>
- (a)  $t(m(g, h)) = t(g)$ ,  $s(m(g, h)) = s(h)$  whenever  $m(g, h)$  is defined;
  - (b)  $m$  is associative:  $m(m(g, h), k) = m(g, m(h, k))$  whenever the relevant terms are defined.
- (3) An embedding  $\epsilon : X \rightarrow G$  called the *identity section* such that  $m(\epsilon(t(g)), g) = g = m(g, \epsilon(s(g)))$  for all  $g \in G$ .
- (4) An *inversion map*  $\iota : G \rightarrow G$  such that  $m(\iota(g), g) = \epsilon(s(g))$  and  $m(g, \iota(g)) = \epsilon(t(g))$  for all  $g \in G$ .

Note that a groupoid over a singleton  $X = \{x\}$  is a group.

We should point out here that there are other ways to define a groupoid. One way is to view it as a particular type of category  $(X, G)$  with  $X$  making up the set of objects, such that the morphisms (elements of  $G$ ) are all invertible. We choose not to consider this categorical description. However, simply by drawing some diagrams to represent the structure maps defined above, one can clearly see how the categorical notion may be derived easily.

Groupoids were first introduced in 1926, and since then found applications in differential topology and geometry, algebraic geometry and algebraic topology, and analysis. A very accessible introduction to groupoids can be found in [84]. For more rigorous accounts one may refer to the bibliography there.

Now we start with a finite groupoid  $G$  and consider the algebra  $\mathbb{K}G$ . Here we are considering the product  $m(g, h)$  of two elements of  $G$  to be defined as in the groupoid itself (and in cases when it is not defined, we set the relevant product equal to zero). We define the comultiplication  $\Delta$ , the counit  $\epsilon$  and the antipode  $S$  on  $G$  by:

$$\Delta(g) = g \otimes g; \quad \epsilon(g) = 1; \quad S(g) = \iota(g) \quad \text{for all } g \in G.$$

Note that when  $G$  is in fact a group,  $\mathbb{K}G$  is its group algebra and hence a Hopf algebra.

Incidentally the dual of the groupoid algebra is again a weak Hopf algebra and can be viewed as the function algebra on  $G$ . See [67] for more on this and other examples.

Due to the inherent symmetry in their construction, weak Hopf algebras have appealed to many researchers with strong algebraic preferences. Their theory has been studied in detail and many algebraically

---

<sup>14</sup>The domain of  $m$  is called the *set of composable pairs*.



natural constructions have been generalized to their context, see for instance [11, 13, 83]. Even the curious fact that the trivial representation of a weak Hopf algebra may or may not be indecomposable ([11, Prop.2.15]) has interesting implications; see [13].

Weak Hopf algebras have nice representation categories [13, 66]. In particular, representation categories of semisimple finite weak Hopf algebras are (multi-)fusion categories. A *fusion category* is a semisimple rigid monoidal category with finitely many simple objects and finite-dimensional homomorphism spaces, such that the endomorphism space of the unit object is one-dimensional. Relaxing the condition that the endomorphism space of the unit object be one-dimensional gives us the definition of a *multi-fusion category*. Fusion categories have been of much interest in the past few years. We refer the reader to [27] for a presentation of recent results about them, and [18] for another accessible exposition. The reader may also find [3] useful for background.

It can be shown that a multi-fusion category can be viewed as the representation category of a (non-unique) semisimple weak Hopf algebra [39, 76]. In fact it turns out that many natural category-theoretical constructions can be restated in the language of weak Hopf algebras. Thus in [27], we see many results on (multi-)fusion categories stated and proved in terms of weak Hopf algebras. The authors of [27] also show that a certain class of fusion categories can also be realized as the representation categories of finite-dimensional semisimple quasi-Hopf algebras. Hence we can see that the notion of a fusion category allows us to build a framework in which both quasi-Hopf algebras and weak Hopf algebras can be understood.

At this point, we move on to our third candidate, the Hopf algebroid, even though our second one is still quite promising. We provide the reader with only one reason, and that is generality. In [14], Böhm and Szlachányi showed that weak Hopf algebras with bijective antipodes are in fact Hopf algebroids. In [86], Xu constructed particular Hopf algebroids which fully encoded the information of certain quasi-Hopf algebras associated with solutions of the quantum dynamical Yang-Baxter equation. In other words, our third candidate will in fact provide us with a fine structure which incorporates most of the interesting weak Hopf algebras and all of the dynamical quantum groups realized as quasi-Hopf algebras, and then offers us some more.

## 6. CANDIDATE 3 : HOPF ALGEBROIDS

A natural question for an algebraist would be: What if we do not restrict ourselves to commutative rings  $\mathbb{K}$ ? Can one develop the theory of bialgebras in this more general setting? In 1977, Takeuchi [80] described and studied a new algebraic structure generalizing bialgebras to the noncommutative setting. His structures were originally called  $\times_A$ -bialgebras. Then in 1996, when developing a geometrically motivated generalization of the theory of quantum groups, Lu defined similar structures [53] and called them *bialgebroids*, to emphasize that the way these structures generalized bialgebras resembled the way Poisson groupoids generalized Poisson groups. A bit later, in 2001 [87], Xu gave a similar definition and also implied the equivalence of these three notions. [17] provides a complete algebraic proof of this result. [7, 14, 17] all include a historical account of these developments. In this note we will mainly be following [7] and [14] for our definitions.

In short, a bialgebroid should be the natural extension of the notion of a bialgebra to the world of groupoids. This then implies that a bialgebroid is no longer an algebra, but a bimodule over a non-commutative ring. Here we will describe two symmetrical notions, that of a *left bialgebroid* and a *right bialgebroid*, first developed in [47]. We begin first with a *left bialgebroid*  $\mathcal{B}_L$ , given by the following data:

- (1) Two associative unital rings: the *total ring*  $B$  and the *base ring*  $L$ .
- (2) Two ring homomorphisms: the *source*  $s_L : L \rightarrow B$  and the *target*  $t_L : L^{op} \rightarrow B$  such that the images of  $L$  in  $B$  commute, making  $B$  an  $L - L$  bimodule denoted by  ${}_L B_L$ .
- (3) Two maps  $\gamma_L : B \rightarrow B_L \otimes_L B$  and  $\pi_L : B \rightarrow L$ , which make  $({}_L B_L, \gamma_L, \pi_L)$  a comonoid in the category of  $L - L$  bimodules.

We also need the following to be satisfied:

$$\begin{aligned} b_{(1)}t_L(l) \otimes b_{(2)} &= b_{(1)} \otimes b_{(2)}s_L(l), \\ \gamma_L(1_B) = 1_B \otimes 1_B \quad \text{and} \quad \gamma_L(bb') &= \gamma_L(b)\gamma_L(b'), \\ \pi_L(1_B) &= 1_L, \\ \pi_L(bs_L \circ \pi_L(b')) &= \pi_L(bb') = \pi_L(bt_L \circ \pi_L(b')) \end{aligned}$$

Note that in the above, we have made use of the Sweedler notation for  $\gamma_L : B \rightarrow B_L \otimes_L B$ :

$$\gamma_L(b) = b_{(1)} \otimes b_{(2)}.$$

We also note that the source and target maps  $s_L$  and  $t_L$  may be used to define four commuting actions of  $L$  on  $B$ ; these in turn give us in an obvious way the new bimodules  ${}^L B^L$ ,  ${}^L B_L$ , and  ${}_L B^L$ .

Similarly we can define a *right bialgebroid*  $\mathcal{B}_R$  using the following data:

- (1) Two associative unital rings: the *total ring*  $B$  and the *base ring*  $R$ .
- (2) Two ring homomorphisms: the *source*  $s_R : R \rightarrow B$  and the *target*  $t_R : R^{op} \rightarrow B$  such that the images of  $R$  in  $B$  commute, making  $B$  an  $R - R$  bimodule denoted by  ${}^R B^R$ .
- (3) Two maps  $\gamma_R : B \rightarrow B^R \otimes {}^R B$  and  $\pi_R : B \rightarrow R$ , which make  $({}^R B^R, \gamma_R, \pi_R)$  a comonoid in the category of  $R - R$  bimodules.

As in the case of left bialgebroids, we need some conditions on  $s_R, t_R, \gamma_R, \pi_R$  analogous to the conditions on  $s_L, t_L, \gamma_L, \pi_L$ :

$$\begin{aligned} s_R(r)b^{(1)} \otimes b^{(2)} &= b^{(1)} \otimes t_R(r)b^{(2)}, \\ \gamma_R(1_B) &= 1_B \otimes 1_B \quad \text{and} \quad \gamma_R(bb') = \gamma_R(b)\gamma_R(b'), \\ \pi_R(1_B) &= 1_R, \\ \pi_R(s_R \circ \pi_R(b)b') &= \pi_R(bb') = \pi_R(t_R \circ \pi_R(b)b') \end{aligned}$$

Note that in the above, we have made use of a modified version of the Sweedler notation for  $\gamma_R : B \rightarrow B^R \otimes {}^R B$ :

$$\gamma_R(b) = b^{(1)} \otimes b^{(2)}.$$

Also we can define three other bimodule structures on  $B$  using the source and the target, and denote them by  ${}_R B_R$ ,  ${}^R B_R$ , and  ${}_R B^R$ .

These bimodule structures and the two notions of bialgebroids are related as expected. For instance if  $\mathcal{B}_L = (B, L, s_L, t_L, \gamma_L, \pi_L)$  is a left bialgebroid, then its co-opposite is again a left bialgebroid:  $(\mathcal{B}_L)_{cop} = (B, L^{op}, t_L, s_L, \gamma_L^{op}, \pi_L)$ , where  $\gamma_L^{op}$  is defined as  $T \circ \gamma_L$ . (Here, as before,  $T$  is the usual twist map, sending  $a \otimes b$  to  $b \otimes a$ ). The opposite  $(\mathcal{B}_L)^{op}$  defined by the data  $(B^{op}, L, t_L, s_L, \gamma_L, \pi_L)$  is a right bialgebroid.

We refer the reader to [17] for several concrete examples of bialgebroids. In this reference, Brzezinski and Militaru show a way to associate a bialgebroid to a braided commutative algebra in the category of Yetter-Drinfeld modules. Moreover they show that the smash product of a Hopf algebra with an algebra in the Yetter-Drinfeld category is a bialgebroid if and only if the algebra is braided commutative. Another interesting construction from [17] gives a generic method of obtaining bialgebroids from solutions of the quantum Yang-Baxter equation.

As mentioned above, Takeuchi's  $\times_A$ -bialgebras [80], Lu's bialgebroids [53] and Xu's bialgebroids [87] were all shown to be equivalent [17], and all these are compatible with the definition above which we took from [7, 14]. Currently, therefore, there is a universal consensus on what should be accepted as the correct structure generalizing bialgebras to the noncommutative base ring case. How, then, does one develop the theory of Hopf algebras in this same setting? In the following paragraphs, we briefly look at how several algebraists approached this problem. But first we would like to answer a different question: Why were algebraists interested in generalizing to the noncommutative base ring case in the first place?

In the mid 1990s, certain finite index depth 2 ring extensions from the theory of Von Neumann algebras were shown to be related to some Hopf algebras [78]. A search started to find connections with more and more general extensions. In 2003, some connections were discovered with the bialgebroids of Takeuchi [47]. Thus the case which would correspond to some bialgebroid with a generalized antipode in the noncommutative base ring case was naturally of some interest.

However, this problem proved to be somewhat complicated. The original antipode suggested in [53] was not universally accepted, and various other formulations followed. We refer the reader to [7] for a comparative study of these various antipodes.<sup>15</sup> The definition used in this note is from [14].

In particular, to define a Hopf algebroid, we need two associative unital rings  $H$  and  $L$ , and set  $R = L^{op}$ . We consider a left bialgebroid structure  $\mathcal{H}_L = (H, L, s_L, t_L, \gamma_L, \pi_L)$  and a right bialgebroid structure  $\mathcal{H}_R = (H, R, s_R, t_R, \gamma_R, \pi_R)$  associated to this pair of rings. We require that  $s_L(L) = t_R(R)$  and  $t_L(L) = s_R(R)$  as subrings of  $H$ , and:

$$\begin{aligned} (\gamma_L \otimes id_H) \circ \gamma_R &= (id_H \otimes \gamma_R) \circ \gamma_L, \\ (\gamma_R \otimes id_H) \circ \gamma_L &= (id_H \otimes \gamma_L) \circ \gamma_R. \end{aligned}$$

The last ingredient is the antipode; this will be a bijection  $S : H \rightarrow H$ , which will satisfy:<sup>16</sup>

$$\begin{aligned} S(t_L(l)ht_L(l')) &= s_L(l')S(h)s_L(l), \\ S(t_R(r')ht_R(r)) &= s_R(r)S(h)s_R(r') \end{aligned}$$

---

<sup>15</sup>A more category-theoretic discussion may be found in [21].

<sup>16</sup>We require that the antipode  $S$  be a bijection for the sake of simplicity. For a study of Hopf algebroids with antipodes which are not necessarily bijective, see [8].

for all  $l, l' \in L$ ,  $r, r' \in R$ , and  $h \in H$ . In other words, we require  $S$  to be a twisted isomorphism simultaneously of bimodules  ${}^L H_L \rightarrow {}_L H^L$  and of bimodules  ${}^R H_R \rightarrow {}_R H^R$ .

Our final constraint on the antipode  $S$  is as follows:

$$\begin{aligned} S(h_{(1)})h_{(2)} &= s_R \circ \pi_R(h), \\ h^{(1)}S(h^{(2)}) &= s_L \circ \pi_L(h) \end{aligned}$$

for any  $h \in H$ . Once again, the subscripts and the superscripts on  $h$  come from a generalized version of the Sweedler notation which we use to define the two maps  $\gamma_L$  and  $\gamma_R$ :

$$\begin{aligned} \gamma_L(h) &= h_{(1)} \otimes h_{(2)} \in H_L \otimes {}_L H \\ \gamma_R(h) &= h^{(1)} \otimes h^{(2)} \in H^R \otimes {}^R H. \end{aligned}$$

In this setup, then, we say that the triple  $\mathcal{H} = (\mathcal{H}_L, \mathcal{H}_R, S)$  is a *Hopf algebroid*.<sup>17</sup>

It is easy to see how this symmetric definition, in terms of two bialgebroids and a bijective map called an antipode, is analogous to the definition of a Hopf algebra from a bialgebra and a bijective map called an antipode. However, it may not be nearly as easy to see why the particular conditions above describe the “*right antipode*.” We accept the definition given above without further analysis; a sufficient discussion of this issue is provided already in the introduction to [7].

There are many examples of Hopf algebroids in recent literature. Here we will describe an interesting example from [14].

Let  $\mathbb{K}$  be a field of characteristic different from 2. Consider the group bialgebra  $\mathbb{K}\mathbb{Z}_2$  presented as a left bialgebroid with the relevant operations on the single generator  $t$  as follows:

$$t^2 = 1; \quad \Delta(t) = t \otimes t; \quad \epsilon(t) = 1.$$

The source and the target maps are the natural ones:  $s, t : \mathbb{K} \rightarrow \mathbb{K}\mathbb{Z}_2$  defined as  $s(\lambda) = \lambda 1 = t(\lambda)$ . Now if we define an antipode  $S : \mathbb{K}\mathbb{Z}_2 \rightarrow \mathbb{K}\mathbb{Z}_2$  by setting  $S(t) = -t$ , we obtain a Hopf algebroid. (Note that the given antipode is not a Hopf algebra antipode).

---

<sup>17</sup>We should note here that some of the above information in our definition is redundant. In fact one can start with a left bialgebroid  $\mathcal{H}_L = (H, L, s_L, t_L, \gamma_L, \pi_L)$  and an anti-isomorphism  $S$  of the total ring  $H$  satisfying certain conditions, and from here can reconstruct a right bialgebroid  $\mathcal{H}_R$  using the same total ring such that the triple  $(\mathcal{H}_L, \mathcal{H}_R, S)$  is a Hopf algebroid. See [7] for more details. An even more streamlined set of defining axioms is proposed in [10].

The notions of integrals and duals for a Hopf algebroid have already been studied [14, 8]. It turns out that the natural self-duality of weak Hopf algebra structures is not available in this theory. In order to define duals one needs a foray into the theory of integrals a la [14]. However the representation categories are still quite nice. In [8], it is shown that the category of (left-left) Hopf modules of a Hopf algebroid  $\mathcal{H} = (\mathcal{H}_L, \mathcal{H}_R, S)$  is equivalent to the category of (right) modules over the left base ring  $L$ .

Hopf algebroids can be used to describe the previous constructions discussed in this note. In other words, as we briefly mentioned at the end of the previous section, both quasi-Hopf objects and weak Hopf objects fit into the Hopf algebroid picture with some modifications. In particular, it can be shown that weak Hopf algebras with bijective antipodes are in fact Hopf algebroids [14]. Moreover, we already know that using the framework of fusion categories, we can describe quasi-Hopf objects in terms of weak Hopf algebras. We can see a connection between quasi-Hopf objects and Hopf algebroids even more directly, if we look at [86]. There, Xu constructs Hopf algebroids which fully encode the information of certain quasi-Hopf algebras associated with solutions of the quantum dynamical Yang-Baxter equation. Hence, it is clear that there are some very interesting connections between these three generalizations of the notion of Hopf algebras.

## 7. SHORT INTERLUDE - IS THIS ALL THAT THERE IS?

In Section 3, we motivated our interest in the search for *the correct generalization* of Hopf algebras by emphasizing the need for a Hopf-like object that can be used to develop sufficiently the theory of dynamical quantum groups. Indeed, several researchers already have investigated each of the three candidate structures we described so far with a view toward the theory of dynamical quantum groups.

We discussed, already, how these three fit into the theory of dynamical quantum groups and also have seen how they relate to one another. In particular, the theory of (multi-)fusion categories seems to provide a fresh point of view which can become the right framework for understanding all of these structures.

Nonetheless, the search for other generalizations of Hopf algebras still goes on. In the rest of this note, we will introduce two more recently developed structures, also generalizing Hopf algebras. These have come up in contexts which are not immediately connected to the theory of dynamical quantum groups. However they both are structures which

raise other new questions, and so they may be of interest to readers who are looking for general themes or for other new structures that may be useful for their own purposes.

## 8. A CATEGORY-THEORETICAL APPROACH: HOPF MONADS

As mentioned in Section 2, any finitely generated commutative Hopf algebra over a field  $\mathbb{K}$  with characteristic 0 is the function algebra of an affine group  $G$  over  $\mathbb{K}$ . Similarly a cocommutative Hopf algebra generated by its primitive elements is the universal enveloping algebra of a Lie algebra  $\mathfrak{g}$ . Readers may find material on these well-known results in classics like [42, 61] and in more modern texts like [28, 33].

Both these results are in fact particular instances of *Tannaka-Krein duality*. Traditionally, the origins of Tannaka-Krein theory are attributed to Grothendieck. We refer the reader to [41] for a comprehensive account relating the original works of Tannaka and Krein with more modern treatments. The first modern references in the subject that make extensive use of category theory are [71, 22]. Expositions of some Tannaka-Krein type theorems presented in the flavor closest to the perspective of this note may be found in [28]. For more details on the modern approach, with an emphasis on Hopf algebras and monoidal categories, the reader may find [58] useful.

In the language of categories, the philosophy underlying the various results that can be gathered under the heading of Tannaka-Krein theory can be stated as follows: it should be possible to view any *nice* category  $\mathcal{C}$  as the representation category of *some algebraic structure*. This vague assertion becomes accurate mathematics when one chooses appropriate descriptions for the italicized phrases.

For instance, one precise formulation of the statement of Section 2 about commutative Hopf algebras is as follows:

**Theorem 8.1.** [22, 28] *Let  $\mathcal{C}$  be a symmetric tensor category defined over an algebraically closed field  $\mathbb{K}$  of characteristic zero. Suppose that  $\mathcal{C}$  is abelian and has finitely many indecomposable objects. Suppose, in addition, that there is an exact faithful tensor functor  $\mathbf{F} : \mathcal{C} \rightarrow \text{Vec}_{\mathbb{K}}$ . Then the group  $G$  of tensor automorphisms of  $\mathbf{F}$  is a finite group and  $\mathcal{C} \simeq \text{Rep}(G)$ .*

In literature, one can find similar results for more general Hopf algebras [72, 82] and quasi-Hopf algebras [57]. In fact, the search for ever more general Hopf-like objects always includes category-theoretic

investigations which look at the representation categories of these objects and attempt to describe them as some nice types of categories.

Our fourth candidate for the correct generalization of Hopf algebras fits perfectly into the spirit of this categorical approach to Hopf algebras. This is the so-called *Hopf monad* structure.<sup>18</sup> Very briefly a **Hopf monad** is a Hopf-like object in a general category.

More precisely, we begin with a monoidal category  $\mathcal{C}$ , and assume that it is *rigid*; in other words, we assume that every object in  $\mathcal{C}$  has a left dual and a right dual. We consider first a *monad* in the sense of [56]. In short a **monad** on  $\mathcal{C}$  is an algebra in the monoidal category  $End(\mathcal{C})$ . More specifically a monad is an endofunctor  $\mathbf{T}$  of  $\mathcal{C}$  equipped with two functorial morphisms  $\mu : \mathbf{T}^2 \rightarrow \mathbf{T}$  and  $\eta : 1_{\mathcal{C}} \rightarrow \mathbf{T}$  which satisfy certain conditions analogous to those describing the product and the unit of an algebra.

The appropriate definition in this context for the notion generalizing bialgebras was introduced by Moerdijk in [63] and asserts that a **bimonad** is a monad  $\mathbf{T}$  which is also comonoidal. In the precise statement of the definition, there are functorial morphisms playing the analogous roles of the coproduct and the counit. The notion of a bimonad is not self-dual, but defining duals is not too complicated.

To make a bimonad  $\mathbf{T}$  into a Hopf monad, we only require the introduction of two new functorial morphisms, the so-called *left antipode* and the *right antipode*, which make use of the left and right duals of the category  $\mathcal{C}$  to encode the left and right duals in the category of  $\mathbf{T}$  modules. Thus a **Hopf monad** is a bimonad with two functorial morphisms, a left antipode and a right antipode.

Hopf monads in the above sense<sup>19</sup> were introduced and studied in detail in [16]. The authors' motivation there stems from the various topological invariants constructed via methods of what is now called quantum topology [49]. The original construction by Reshetikhin and Turaev [70] of invariants for 3-manifolds using quantum  $sl_2(\mathbb{C})$  and

---

<sup>18</sup>We note here that there is at least one other category-theoretic Hopf-like object studied in the recent years. We do not focus on the relevant constructions here; we simply refer the reader to [20, 21] where the ideas are developed in great detail. Some connections to the picture from Section 6 may be found in [14, 9].

<sup>19</sup>We need to remark on our specific terminology here. The term “*Hopf monad*” appeared earlier, in [63]. In particular, the “*Hopf monads*” of [63] are precisely the structures that we here called “*bimonads*.” We are following [16, 77] with this choice of terminology, because it seems quite natural linguistically to reserve the Hopf term for bimonads with antipode. However, for readers familiar with operad theory, the terminology choice of [63] may seem more reasonable; see also [62].



related (ribbon) Hopf algebras has been generalized in recent years to constructions in more and more general settings. The authors of [16] study Hopf monads as a generalization of Hopf algebras to categories with no braiding so as to provide the ultimate framework to understand all these newer invariants in a uniform manner.

Any finite-dimensional Hopf algebra  $H$  over  $\mathbb{K}$  can be used to construct a Hopf monad on the category of finite-dimensional  $\mathbb{K}$ -vector spaces. More generally, Hopf monads can be constructed using any rigid monoidal category with an underlying algebraic structure. Thus, the Hopf monad concept brings together all of the earlier structures we studied in this note: quasi-Hopf algebras, weak Hopf algebras and Hopf algebroids all can be used to construct Hopf monads in their relevant representation categories.<sup>20</sup>

## 9. A (POISSON-)GEOMETRIC APPROACH: HOPFISH ALGEBRAS

Finally we briefly describe a fifth Hopf-like object which generalizes the notion of a Hopf algebra. This object, playfully called a *hopfish algebra* by its creators, was introduced first in [81] and applied to the problem of describing modules of irrational rotation algebras in [5]. The ideas in its development fit in the context of (Poisson) geometry; the reader may find interesting background and motivational discussions in the recent survey [6] of algebraic structures in Poisson geometry.

Hopfish algebras live in a category  $\mathcal{A}$  whose objects are  $\mathbb{K}$ -unital algebras and whose morphisms are bimodules. Then as the authors of [81] show, the notions of comultiplication, counit, and the antipode can all be developed in this framework, as particular types of bimodules (i.e. morphisms in the new category).

More precisely, we start with the notion of a *sesquiunital sesquialgebra*, which is the relevant generalization of bialgebras in this context: A **sesquiunital sesquialgebra** over a commutative ring  $\mathbb{K}$  is a unital  $\mathbb{K}$ -algebra  $B$  equipped with a  $(B \otimes B, B)$ -bimodule  $\Delta$  (the **coproduct**), and a  $(\mathbb{K}, B)$ -bimodule (ie a right  $B$ -module)  $\epsilon$  (the **counit**), with the following properties:

- (1) (*coassociativity axiom*) The two  $(B \otimes B \otimes B, B)$  bimodules  $(B \otimes \Delta) \otimes_{B \otimes B} \Delta$  and  $(\Delta \otimes B) \otimes_{B \otimes B} \Delta$  are isomorphic; and
- (2) (*counit axiom*) The two  $(B, B)$  bimodules  $(\epsilon \otimes B) \otimes_{B \otimes B} \Delta$  and  $(B \otimes \epsilon) \otimes_{B \otimes B} \Delta$  are both isomorphic to  $B$ .

---

<sup>20</sup>For a clarification of the relationship between bimonads and bialgebroids, see [77]. The quasi- and weak Hopf algebra cases are explicitly discussed in [16].

To move on to the Hopf-like objects one needs an antipode appropriate for the context. In order to define the right structure that should correspond to the antipode, we begin with a *preantipode*. In particular a **preantipode** for a sesquiunital sesquialgebra  $B$  over  $\mathbb{K}$  is a left  $(B \otimes B)$ -module  $\mathbf{S}$  together with an isomorphism of its  $\mathbb{K}$ -dual with the right  $(B \otimes B)$ -module  $Hom_B(\epsilon, \Delta)$ . One can view  $\mathbf{S}$  as a  $(B, B^{op})$ -bimodule, and hence an  $\mathcal{A}$ -morphism in  $Hom(B, B^{op})$ . If a preantipode  $\mathbf{S}$  on  $B$  is a free left  $B$ -module of rank 1 when considered as a  $(B, B^{op})$ -bimodule, then it is called an **antipode** for  $B$ . A sesquiunital sesquialgebra  $H$  equipped with an antipode  $\mathbf{S}$  is called a **hopfish algebra**.

To see the standard examples, we need to begin with a well-known functor of Morita theory. This functor, called **modulation** in [81], goes from the category of  $\mathbb{K}$ -unital algebras with algebra homomorphisms as morphisms, to the category  $\mathcal{A}$ . Using this functor, one can see that the modulation of biunital bialgebras will be basic examples of sesquiunital sesquialgebras and the modulation of Hopf algebras will be basic examples of hopfish algebras. In [81] the authors also show that quasi-Hopf and weak Hopf algebras can be modulated to yield hopfish algebras.

For more details on hopfish algebras we refer the reader to [81, 5, 6]. Readers who prefer the more general setting of higher category theory may find that [20, 21] can perhaps complement the above references on hopfish algebras.

## 10. CONCLUSION - WHERE DO WE GO NEXT?

In this note, we focused on five structures that have been introduced in the recent years as generalizations of Hopf algebras. One thing is clear: The notion of Hopf algebras proved useful in so many diverse ways that mathematicians of many different persuasions decided that the search for *the correct generalization* was quite an important task. More significantly, a lot of new and interesting mathematics came up during these investigations.

Some readers may also have noticed along the way a few open problems that are looking for solutions. For instance, the question whether the Hopf algebroids from [53] are a proper subclass of the Hopf algebroids of Section 6 or not is still not settled. The example we gave at the end of Section 6 would not be a Hopf algebroid in the sense of [53]. So far, we do not have an example which would be a Hopf algebroid for [53] and would not be a Hopf algebroid in our sense, but it is not

certain that we will never find one. This is an interesting question, as the Hopf algebroids in [53] are quite natural structures themselves.

There are also many open questions about (multi-)fusion categories and hence about (weak) Hopf algebras. One of the most fascinating questions involves the existence (or non-existence) of finite dimensional semisimple Hopf algebras with non-group theoretical representation categories. For details on this and many other open questions about (multi-)fusion categories we refer the reader once again to [27].

The relationship between representation categories of all the various generalizations of Hopf algebras also needs to be studied further. Indeed, a lot of the pieces are out there, but there is still more work to be done.

In this note, we pointed out some such connections. We also provided some details on the motivations of the mathematicians who developed the relevant structures. Along the way, we gave our own reasons for being interested in generalizations of Hopf algebras. We hope that at this point, the readers have already decided if any of the five structures fits their own particular needs or mathematical inclinations. If not, they at least have an idea of what leads to follow to decide on this matter for themselves.

## REFERENCES

- [1] Abe, Eiichi; Hopf algebras, Cambridge Tracts in Mathematics, **74**, Cambridge University Press, Cambridge-New York, 1980.
- [2] Aguiar, Marcelo; Mahajan, Swapneel; Coxeter Groups and Hopf Algebras; Fields Institute Monographs, **23**, American Mathematical Society, Providence, RI, 2006.
- [3] Bakalov, Bojko; Kirillov, Alexander, Lectures on tensor categories and modular functors, AMS University Lecture Series, **21**, American Mathematical Society, Providence, RI, 2001.
- [4] Bénabou, Jean; “*Catégories avec multiplication*”, C. R. Acad. Sci. Paris **256** (1963), pp.1887–1890.
- [5] Blohmann, Christian; Tang, Xiang; Weinstein, Alan; “*Hopfish structures and modules over irrational rotation algebras*”, e-arXiv preprint, [arXiv:math.QA/0604405](https://arxiv.org/abs/math/0604405)
- [6] Blohmann, Christian; Weinstein, Alan; “*Group-like objects in Poisson Geometry and algebra*”, e-arXiv preprint, [arXiv:math.SG/0701499](https://arxiv.org/abs/math/0701499)
- [7] Böhm, Gabriella; “*An alternative notion of Hopf algebroid*”, Hopf Algebras in Noncommutative Geometry and Physics, Lecture Notes in Pure and Appl. Math. **239**, Dekker, New York, 2005, pp.31–53.
- [8] Böhm, Gabriella; “*Integral theory for Hopf algebroids*”, Algebr. Represent. Theory **8** (2005), no. 4, pp.563–599.

- [9] Böhm, Gabriella; “*Hopf algebroids*”, Handbook of Algebra, M. Hazewinkel (ed.), Elsevier, to appear.
- [10] Böhm, Gabriella; Brzeziński, Tomasz; “*Cleft extensions of Hopf algebroids*”, Appl. Categ. Structures **14** (2006), no. 5-6, pp.431–469.
- [11] Böhm, Gabriella; Nill, F.; Szlachányi, Kornél; “*Weak Hopf Algebras I. Integral Theory and  $C^*$  Structure*”, J. Algebra **221** (1999), pp.385–438.
- [12] Böhm, Gabriella; Szlachányi, Kornél; “*A Coassociative  $C^*$ -Quantum Group with Nonintegral Dimensions*”, Lett. Math. Phys. **38** (1996), no.4, pp.437–456.
- [13] Böhm, Gabriella; Szlachányi, Kornél; “*Weak Hopf Algebras II. Representation Theory, dimensions and the Markov trace*”, J. Algebra **233** (2000), pp.156–212.
- [14] Böhm, Gabriella; Szlachányi, Kornél; “*Hopf Algebroids with Bijective Antipodes: Axioms, Integrals and Duals*”, J. Algebra **274** (2004), no.2, pp.708–750.
- [15] Brown, Ken A.; Goodearl, Ken R.; Lectures on algebraic quantum groups; Advanced Courses in Mathematics. CRM Barcelona. Birkhäuser Verlag, Basel, 2002.
- [16] Bruguières, Alain; Virelizier, Alexis; “*Hopf Monads*”, e-arXiv preprint, [arXiv:math.QA/0604180](https://arxiv.org/abs/math.QA/0604180)
- [17] Brzeziński, Tomasz; Militaru, Gigel; “*Bialgebroids,  $\times_A$ -Bialgebras and Duality*”, J. Algebra **251** (2002), pp.279–294.
- [18] Calaque, Damien; Etingof, Pavel.; “*Lectures on tensor categories*”, e-arXiv preprint, [arXiv:math.QA/0401246](https://arxiv.org/abs/math.QA/0401246)
- [19] Chari, Vyjayanthi; Pressley, Andrew; A Guide to Quantum Groups, Cambridge University Press, 1995.
- [20] Day, Brian, McCrudden, Paddy, Street, Ross; “*Dualizations and antipodes*”, Appl. Categ. Structures **11** (2003), no. 3, pp.229–260.
- [21] Day, Brian, Street, Ross; “*Quantum categories, star autonomy, and quantum groupoids*”, Galois Theory, Hopf Algebras, and Semiabelian Categories, Fields Inst. Commun., **43**, Amer. Math. Soc., Providence, RI, 2004, pp.187–225.
- [22] Deligne, P.; Milne, R.; Tannakian Categories; Lecture Notes in Mathematics **900**; Springer-Verlag, 1982, pp.101–228.
- [23] Drinfeld, Vladimir; “*Quantum groups*”; Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Berkeley, Calif., 1986), Amer. Math. Soc., Providence, RI, 1987, pp.798–820.
- [24] Drinfeld, Vladimir; “*Quasi-Hopf algebras*” (Russian) Algebra i Analiz **1** (1989), no. 6, pp.114–148; translation in Leningrad Math. J. **1** (1990), no. 6, pp.1419–1457.
- [25] Drinfeld, Vladimir; “*Quasi-Hopf algebras and Knizhnik-Zamolodchikov equations*”; Problems of modern quantum field theory (Alushta, 1989), Res. Rep. Phys., Springer, Berlin, 1989, pp.1–13.
- [26] Enriquez, B.; Felder, Giovanni; “*Elliptic Quantum Groups  $E_{\tau,\eta}(sl_2)$  and Quasi-Hopf algebras*”, Comm. Math. Phys. **195** (1998), pp.651–689.
- [27] Etingof, Pavel; Nikshych, Dmitri; Ostrik, Viktor; “*On fusion categories*”, Ann. of Math. (2) **162** (2005), no. 2, pp.581–642.
- [28] Etingof, Pavel; Schiffman, Olivier; Lectures on Quantum Groups; International Press, 1998.

- [29] Etingof, Pavel; Varchenko, Alexander; “*Solutions of the quantum dynamical Yang-Baxter equation and dynamical quantum groups*”; Comm. Math. Phys. **196** (1998), no. 3, pp.591–640.
- [30] Felder, Giovanni; “*Conformal Field Theory and Integrable Systems associated to Elliptic Curves*”, Proceedings of the International Congress of Mathematicians (Zurich 1994), Birkhäuser, 1994, pp.1247–1255.
- [31] Felder, Giovanni; “*Elliptic Quantum Groups*”, Proceedings of the XIth International Congress of Mathematical Physics (Paris, 1994), International Press, 1995, pp.211–218.
- [32] Felder, Giovanni; Varchenko, Alexander; “*On Representations of the Elliptic Quantum Group  $E_{\tau,\eta}(sl_2)$* ”, Comm. Math. Phys. **181** (1996), pp.741–761.
- [33] Ferrer Santos, Walter; Rittatore, Alvaro; Actions and Invariants of Algebraic Groups, Pure and applied Mathematics **269**, Chapman & Hall (CRC Press), 2005.
- [34] Gervais, J. L.; Neveu, A.; “*Novel Triangle Relations and Absence of Tachyons in Liouville String Field Theory*”, Nucl Phys. B **238** (1984), pp.125–141.
- [35] Hausser, Frank, and Nill, Florian; “*Integral theory for quasi-Hopf algebras*”, e-arXiv preprint, [arXiv:math.QA/9904164](https://arxiv.org/abs/math/9904164)
- [36] Hayashi, Takahiro; “*Quantum group symmetry of partition functions of IRF models and its application to Jones’ index theory*”, Comm. Math. Phys. **157** (1993), no. 2, pp.331–345.
- [37] Hayashi, Takahiro; “*Face algebras and their Drinfeld doubles*”, Algebraic Groups and Their Generalizations: Quantum and Infinite-Dimensional Methods (University Park, PA, 1991), Proc. Sympos. Pure Math. **56**, Part 2, Amer. Math. Soc., Providence, RI, 1994, pp.49–61.
- [38] Hayashi, Takahiro; “*Face algebras. I. A generalization of quantum group theory*”, J. Math. Soc. Japan **50** (1998), no. 2, pp.293–315.
- [39] Hayashi, Takahiro; “*A canonical Tannaka duality for finite semisimple tensor categories*”, e-arXiv preprint, [arXiv:math.QA/99040731](https://arxiv.org/abs/math/99040731)
- [40] Hazewinkel, Michiel; “*Introductory recommendations for the study of Hopf algebras in mathematics and physics*; CWI Quarterly **4** (1991), no. 1, pp.3–26.
- [41] Hewitt, Edwin; Ross, Kenneth A.; Abstract Harmonic Analysis. Vol. II: Structure and analysis for compact groups, Analysis on locally compact Abelian groups; Die Grundlehren der mathematischen Wissenschaften **152**, Springer-Verlag, New York-Berlin 1970.
- [42] Hochschild, Gerhard P.; Basic Theory of Algebraic Groups and Lie Algebras, Graduate Texts in Mathematics **75**, Springer-Verlag, New York-Berlin, 1981.
- [43] Hong, Jin; Kang, Seok-Jin; Introduction to quantum groups and crystal bases; Graduate Studies in Mathematics **42**, American Mathematical Society, Providence, RI, 2002.
- [44] Hopf, Heinz; “*Über die Topologie der Gruppen-mannigfaltigkeiten und ihre Verallgemeinerungen*”, Ann. Math. **42** (1941), pp.22–52.
- [45] Jimbo, Michio; “*Introduction to the Yang-Baxter Equation*”, International J. Mod. Phys. A **4** (1989), pp.3759–3777.
- [46] Yang-Baxter Equation in Integrable Systems, Michio Jimbo ed., Advanced Series in Math. Phys., vol. 10, World Scientific, 1990.
- [47] Kadison, Lars; Szlachányi, Kornél; “*Bialgebroid actions on depth two extensions and duality*”, Adv. Math. **179** (2003), pp.75–121.

- [48] Kassel, Christian; Quantum groups; Graduate Texts in Mathematics **155**, Springer-Verlag, New York, 1995.
- [49] Kauffman, Louis; “*Introduction to quantum topology*”, Quantum Topology, Ser. Knots Everything, **3**, World Sci. Publ., River Edge, NJ, 1993, pp.1–77.
- [50] Lambe, Larry A.; Radford, David E.; Introduction to the quantum Yang-Baxter equation and quantum groups: an algebraic approach; Mathematics and its Applications **423**, Kluwer Academic Publishers, Dordrecht, 1997.
- [51] Li, Fang; “*Weak Hopf algebras and some new solutions of the quantum Yang-Baxter equation*”, J. Algebra **208** (1998), no. 1, pp.72–100.
- [52] Li, Fang, Duplij, Steven; “*Weak Hopf algebras and singular solutions of quantum Yang-Baxter equation*”, Comm. Math. Phys. **225** (2002), no. 1, pp.191–217.
- [53] Lu, J-H.; “*Hopf Algebroids and Quantum Groupoids*”, Int. J. Math. **7** (1996), no.1, pp.47–70.
- [54] Lusztig, George; Introduction to quantum groups; Progress in Mathematics **110**, Birkhäuser Boston Inc., Boston, MA, 1993.
- [55] MacLane, Saunders; “*Natural associativity and commutativity*”, Rice Univ. Studies **49** (1963) no. 4, pp.28–46.
- [56] MacLane, Saunders; Categories for the Working Mathematician; Second Edition; Springer-Verlag, New York, 1998.
- [57] Majid, Shahn; “*Tannaka-Krein Theorem for Quasi-Hopf algebras and Other Results*”, Contemp. Math. **134** (1992), pp.219–232..
- [58] Majid, Shahn; Foundations of quantum group theory; Cambridge University Press, Cambridge, 1995.
- [59] Manin, Yuri; Quantum Groups and Noncommutative Geometry; Université de Montréal, Centre de Recherches Mathématiques, Montreal, QC, 1988.
- [60] McCleary, John; A User’s Guide to Spectral Sequences; Second edition, Cambridge Studies in Advanced Mathematics, **58**, Cambridge University Press, Cambridge, 2001.
- [61] Milnor, John W.; Moore, John C.; “*On the Structure of Hopf Algebras*”; Ann. of Math. (2) **81** (1965), pp.211–264.
- [62] Moerdijk, Ieke; “*On the Connes-Kreimer construction of Hopf algebras*”, Homotopy Methods in Algebraic Topology (Boulder, CO, 1999), Contemp. Math., **271** (2001), pp.311–321.
- [63] Moerdijk, Ieke; “*Monads on Tensor categories*”, J. Pure Appl. Algebra **168** (2002), no 2-3, pp.189–208.
- [64] Montgomery, Susan; Hopf Algebras and their Actions on Rings; CBMS Lecture Notes **82**, American Math. Society, Providence, RI, 1993.
- [65] Nichols, Warren; Sweedler, Moss; “*Hopf algebras and combinatorics*”, Umbral Calculus and Hopf Algebras (Norman, Okla., 1978), Contemp. Math. **6**, Amer. Math. Soc., Providence, R.I., 1982, pp. 49–84.
- [66] Nikshych, D.; Turaev, V.; Vainerman, L.; “*Quantum Groupoids and Invariants of Knots and 3-manifolds*”, Proceedings of the Pacific Institute for the Mathematical Sciences Workshop ”Invariants of Three-Manifolds” (Calgary, AB, 1999). Topology Appl. **127** (2003), no. 1-2, pp.91–123.
- [67] Nikshych, D.; Vainerman, L.; “*Finite Quantum Groupoids and Their Applications*”, in: New Directions in Hopf Algebras, MSRI Publications Vol. **43**, 2002, pp.211–262.

- [68] Nill, F.; “*Axioms of Weak Bialgebras*”, e-arXiv preprint, arXiv:math.QA/9805104
- [69] Ocneanu, Adrian; “*Quantized groups, string algebras and Galois theory for algebras*”, Operator algebras and applications, Vol. 2, pp.119–172, London Math. Soc. Lecture Note Ser., **136**, Cambridge Univ. Press, Cambridge, 1988.
- [70] Reshetikhin, Nicolai; Turaev, Vladimir; “*Invariants of 3-manifolds via link polynomials and quantum groups*”, Invent. Math **103** (1991), no.3, pp.547–597.
- [71] Saavedra Rivano, Neantro; *Catégories Tannakiennes*; Lecture Notes in Mathematics, **265**, Springer-Verlag, Berlin-New York, 1972.
- [72] Schauenburg, Peter; *Tannaka Duality for Arbitrary Hopf Algebras*, Algebra Berichte **66**, Verlag Reinhard Fischer, Munich, 1992.
- [73] Schauenburg, Peter; “*Hopf modules and the double of a quasi-Hopf algebra*”, Trans. Amer. Math. Soc. **354** (2002), no. 8, pp.3349–3378
- [74] Shnider, Steven; Sternberg, Shlomo; *Quantum groups: From coalgebras to Drinfeld algebras: A guided tour*; Graduate Texts in Mathematical Physics, II. International Press, Cambridge, MA, 1993.
- [75] Szlachányi, Kornél; “*Weak Hopf algebras*”, Operator Algebras and Quantum Field Theory (Rome, 1996), Int. Press, Cambridge, MA, 1997, pp. 621–632.
- [76] Szlachányi, Kornél; “*Finite quantum groupoids and inclusions of finite type*”, Mathematical Physics in Mathematics and Physics (Siena, 2000), Fields Inst. Commun. **30**, AMS Providence, RI, 2001, pp.393–407.
- [77] Szlachányi, Kornél; “*The monoidal Eilenberg-Moore construction and bialgebroids*”, J. Pure Appl. Algebra **182** (2003), no. 2-3, pp.287–315.
- [78] Szymański, W.; “*Finite index subfactors and Hopf algebra cross products*”, Proc. Amer. Math. Soc. **120** 2 (1994), pp.519–528.
- [79] Sweedler, Moss E.; *Hopf algebras*; Mathematics Lecture Note Series, W. A. Benjamin, Inc., New York, 1969.
- [80] Takeuchi, Mitsuhiro; “*Groups of algebras over  $A \otimes \bar{A}$* ”; J. Math. Soc. Japan **29** (1977), no. 3, pp.459–492.
- [81] Tang, Xiang; Weinstein, Alan; Zhu, Chenchang; “*Hopf fish algebras*”, Pacific J. Math. **231** (2007), no. 1, pp.193–216.
- [82] Ulbrich, Karl-H.; “*Tannakian categories for non-commutative Hopf algebras*”, Israel J. Math **72** (1990), no 1-2, pp.252–256.
- [83] Vecseryné, P.; “*Larson-Sweedler theorem and the role of grouplike elements in weak Hopf algebras*”; J. Algebra **270** (2003), pp.471–520.
- [84] Weinstein, Alan; “*Groupoids: Unifying Internal and External Symmetry. A Tour through Some Examples*”, Notices Amer. Math. Soc. **43** (1996), no.7, pp.744–752.
- [85] Wisbauer, Robert; “*Weak corings*”, J. Algebra **245** (2001), no. 1, pp.123–160.
- [86] Xu, Ping; “*Quantum Groupoids associated to universal dynamical R-matrices*”, C. R. acad. Sci. Paris, t. 328 (1999), Série 1, pp.327–332.
- [87] Xu, Ping; “*Quantum groupoids*”, Comm. Math. Phys. **216** (2001), pp.539–581.

DEPARTMENT OF MATHEMATICS, POMONA COLLEGE, CLAREMONT CA 91711  
*E-mail address:* gizem.karaali@pomona.edu