

A NOTE ON THE q -GENOCCHI NUMBERS AND POLYNOMIALS

TAEKYUN KIM

ABSTRACT. In this paper we discuss the new concept of the q -extension of Genocchi numbers and give the some relations between q -Genocchi polynomials and q -Euler numbers.

§1. Introduction

A number given by the generating function

$$(1) \quad \frac{2t}{e^t + 1} = \sum_{n=0}^{\infty} G_n \frac{t^n}{n!}.$$

It satisfies $G_1 = 1, G_3 = G_4 = G_7 = \dots = 0$, and even coefficients are given by $G_{2m} = 2(1 - 2^{2n})B_{2n} = 2nE_{2n-1}(0)$, where B_n is a Bernoulli number and $E_n(x)$ is an Euler polynomial. The first few Genocchi numbers for $n = 2, 4, \dots$ are $-1, -3, 17, -155, 2073, \dots$. The Euler polynomials are well known as

$$(2) \quad \frac{2}{e^t + 1} e^{xt} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}, \text{ see [1, 3, 7, 8, 9].}$$

By (1) and (2) we easily see that

$$(3) \quad E_n(x) = \sum_{k=0}^n \binom{n}{k} \frac{G_{k+1}}{k+1} x^{n-k}, \text{ where } \binom{n}{k} = \frac{n(n-1)\cdots(n-k+1)}{k!}, \text{ cf. [4, 5, 6].}$$

For $m, n \geq 1$ and m odd, we have

$$(4) \quad (n^m - n)G_m = \sum_{k=1}^{m-1} \binom{m}{k} n^k G_k Z_{m-k}(n-1),$$

Key words and phrases. p -adic q -integrals, Genocchi numbers, q -Genocchi numbers, polynomials, sums of powers.

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\mathcal{T}\mathcal{E}\mathcal{X}$

where $Z_m(n) = 1^m - 2^m + 3^m - \dots + (-1)^{n-1}n^m$, see[3]. From (10) we derive

$$2t = \sum_{n=0}^{\infty} ((G+1)^n + G_n) \frac{t^n}{n!},$$

where we use the technique method notation by replacing G^m by $G_m (m \geq 0)$, symbolically. By (5), we see that

$$(6) \quad G_0 = 0, \quad (G+1)^n - G_n = \begin{cases} 2 & \text{if } n = 1 \\ 0 & \text{if } n > 1. \end{cases}$$

Let p be a fixed odd prime, and let \mathbb{C}_p denote the p -adic completion of the algebraic closure of $\mathbb{Q}_p (= p\text{-adic number field})$. For d a fixed positive integer with $(p, d) = 1$, let

$$\begin{aligned} X &= X_d = \varprojlim_{\mathbb{N}} \mathbb{Z}/dp^N \mathbb{Z}, \\ X_1 &= \mathbb{Z}_p, \\ X^* &= \bigcup_{\substack{0 < a < dp \\ (a, p) = 1}} (a + dp\mathbb{Z}_p), \\ a + dp^N \mathbb{Z}_p &= \{x \in X \mid x \equiv a \pmod{dp^N}\}, \end{aligned}$$

where $a \in \mathbb{Z}$ lies in $0 \leq a < dp^N$.

Ordinary q -calculus is now very well understood from many different point of views. Let us consider a complex number $q \in \mathbb{C}$ with $|q| < 1$ (or $q \in \mathbb{C}_p$ with $|1-q|_p < p^{-\frac{1}{p-1}}$) as an indeterminate. The q -basic numbers are defined by

$$[x]_q = \frac{q^x - 1}{q - 1} = 1 + q + q^2 + \dots + q^{x-1},$$

and

$$[x]_{-q} = \frac{-(-q)^x + 1}{q + 1} = 1 - q + q^2 + \dots + (-q)^{x-1}.$$

We say that f is a uniformly differentiable function at a point $a \in \mathbb{Z}_p$ and denote this property by $f \in UD(\mathbb{Z}_p)$, if the difference quotients

$$F_f(x, y) = \frac{f(x) - f(y)}{x - y}$$

have a limit $l = f'(a)$ as $(x, y) \rightarrow (a, a)$.

For $f \in UD(\mathbb{Z}_p)$, let us start with the expression

$$\frac{1}{[p^N]_q} \sum_{0 \leq j < p^N} q^j f(j) = \sum_{\substack{0 \leq j < p^N \\ 2}} f(j) \mu_q(j + p^N \mathbb{Z}_p),$$

representing a q -analogue of Riemann sums for f , cf. [5]. The integral of f on \mathbb{Z}_p will be defined as limit ($n \rightarrow \infty$) of those sums, when it exists. The p -adic q -integral of the function $f \in UD(\mathbb{Z}_p)$ is defined by

$$I_q(f) = \int_{\mathbb{Z}_p} f(x) d\mu_q(x) = \lim_{N \rightarrow \infty} \frac{1}{[p^N]_q} \sum_{0 \leq x < p^N} f(x) q^x, \quad (\text{see [5, 10, 11, 12]}).$$

In the previous paper [4] author constructed the q -extension of Euler polynomials in the fermionic sense of p -adic q -integral at $q = -1$ as follows:

$$(7) \quad E_{n,q}(x) = \int_{\mathbb{Z}_p} [t+x]_q^n d\mu_{-q}(t), \quad \text{where } \mu_{-q}(x + p^N \mathbb{Z}_p) = \lim_{q \rightarrow -q} \mu_q(x + p^N \mathbb{Z}_p).$$

From (7), we note that

$$(8) \quad E_{n,q}(x) = \frac{[2]_q}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} \frac{(-1)^l}{1+q^{l+1}} q^{lx}, \quad \text{see [4]}.$$

The q -extension of Genocchi numbers is defined as

$$(9) \quad g_q^*(t) = [2]_q t \sum_{n=0}^{\infty} (-1)^n q^n e^{[n]_q t} = \sum_{n=0}^{\infty} G_{n,q}^* \frac{t^n}{n!}, \quad \text{see [4]}.$$

The following formula is well known in [4, 7]:

$$(10) \quad E_{n,q}(x) = \sum_{k=0}^n \binom{n}{k} [x]_q^{n-k} q^{kx} \frac{G_{k+1,q}^*}{k+1}.$$

The modified q -Euler numbers are defined as

$$\xi_{0,q} = \frac{[2]_q}{2}, \quad (q\xi + 1)^k + \xi_{k,q} = \begin{cases} [2]_q & \text{if } k = 0 \\ 0 & \text{if } k \neq 0, \end{cases}$$

with the usual convention of replacing ξ^i by $\xi_{i,q}$, see [10]. Thus, we derive the generating function of $\xi_{n,q}$ as follows:

$$(11) \quad F_q(t) = [2]_q \sum_{k=0}^{\infty} (-1)^k e^{[k]_q t} = \sum_{n=0}^{\infty} \xi_{n,q} \frac{t^n}{n!}.$$

Now we also consider the q -Euler polynomials $\xi_{n,q}(x)$ as

$$(12) \quad F_q(t, x) = [2]_q \sum_{k=0}^{\infty} (-1)^k e^{[k+x]_q t} = \sum_{n=0}^{\infty} \xi_{n,q}(x) \frac{t^n}{n!}.$$

From (12) we note that

$$(13) \quad \xi_{n,q}(x) = \sum_{l=0}^n \binom{n}{l} \xi_{l,q} q^{lx} [x]_q^{n-l}, \text{ see [10].}$$

In the recent several authors studied the q -extension of Genocchi numbers and polynomials (see[1, 2, 5, 6, 7, 12]). In this paper we discuss the new concept of the q -extension of Genocchi numbers and give the some relations between q -Genocchi numbers and q -Euler numbers.

§2. q -extension of Genocchi numbers

In this section we assume that $q \in \mathbb{C}$ with $|q| < 1$. Now we consider the q -extension of Genocchi numbers as follows:

$$(14) \quad g_q(t) = [2]_q t \sum_{k=0}^{\infty} (-1)^k e^{[k]_q t} = \sum_{n=0}^{\infty} G_{n,q} \frac{t^n}{n!}.$$

In Eq.(14), it is easy to show that $\lim_{q \rightarrow 1} g_q(t) = \frac{2t}{e^t + 1} = \sum_{n=0}^{\infty} G_n \frac{t^n}{n!}$. From (14) we derive

$$\begin{aligned} g_q(t) &= [2]_q t \sum_{k=0}^{\infty} (-1)^k \sum_{m=0}^{\infty} [k]_q^m \frac{t^m}{m!} = [2]_q \sum_{k=0}^{\infty} (-1)^k \sum_{m=1}^{\infty} m [k]_q^{m-1} \frac{t^m}{m!} \\ &= [2]_q \sum_{k=0}^{\infty} (-1)^k \sum_{m=0}^{\infty} m [k]_q^{m-1} \frac{t^m}{m!}. \end{aligned} \quad (15)$$

By (15), we easily see that

$$(16) \quad g_q(t) = [2]_q \sum_{m=0}^{\infty} \left(m \left(\frac{1}{1-q} \right)^{m-1} \sum_{l=0}^{m-1} \binom{m-1}{l} (-1)^l \frac{1}{1+q^l} \right) \frac{t^m}{m!}.$$

From (14) and (16) we note that

$$(17) \quad \sum_{m=0}^{\infty} G_{m,q} \frac{t^m}{m!} = \sum_{m=0}^{\infty} \left(m [2]_q \left(\frac{1}{1-q} \right)^{m-1} \sum_{l=0}^{m-1} \binom{m-1}{l} \frac{(-1)^l}{1+q^l} \right) \frac{t^m}{m!}.$$

By comparing the coefficients on both sides in Eq.(17), we have the following theorem:

Theorem 1. For $m \geq 0$ we have

$$G_{m,q} = m[2]_q \left(\frac{1}{1-q} \right)^{m-1} \sum_{l=0}^{m-1} \binom{m-1}{l} \frac{(-1)^l}{1+q^l}.$$

From Theorem 1, we easily derive the following corollary:

Corollary 2. For $n \in \mathbb{N}$ we have

$$G_{0,q} = 0, \quad (qG + 1)^k + G_{k,q} = \begin{cases} \frac{[2]_q^2}{2} & \text{if } k = 1, \\ 0 & \text{if } k > 1, \end{cases}$$

with the usual convention of replacing G^i by $G_{i,q}$.

Remark. We note that Corollary 2 is the q -extension of Eq.(6). By (10) and Corollary 2, we obtain the following theorem:

Theorem 3. For $n \in \mathbb{N}$ we have

$$\xi_{n,q} = \frac{G_{n+1,q}}{n+1}.$$

From (12) we derive

$$\begin{aligned} F_q(x, t) &= [2]_q \sum_{n=0}^{\infty} (-1)^n e^{[n+x]_q t} = q^x t \frac{[2]_q}{q^x t} e^{[x]_q t} \sum_{n=0}^{\infty} (-1)^n e^{q^x [n]_q t} \\ &= e^{[x]_q t} \sum_{n=0}^{\infty} q^{nx} \frac{G_{n+1,q}}{n+1} \frac{t^n}{n!} = \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{n}{k} [x]_q^{n-k} q^{kx} \frac{G_{k+1,q}}{k+1} \right) \frac{t^n}{n!}. \end{aligned} \quad (18)$$

By (18), we easily see that

$$\xi_{n,q}(x) = \sum_{k=0}^n \binom{n}{k} [x]_q^{n-k} q^{kx} \frac{G_{k+1,q}}{k+1}.$$

This formula can be considered as the q -extension of Eq.(3). Let us consider the q -analogue of Genocchi polynomials as follows:

$$(19) \quad g_q(x, t) = [2]_q t \sum_{k=0}^{\infty} (-1)^k e^{[k+x]_q t} = \sum_{n=0}^{\infty} G_{n,q}(x) \frac{t^n}{n!}.$$

Thus, we note that $\lim_{q \rightarrow 1} g_q(x, t) = \frac{2t}{e^t + 1} e^{xt} = \sum_{n=0}^{\infty} G_n(x) \frac{t^n}{n!}$. From Eq.(19), we easily derive

$$(20) \quad G_{n,q}(x) = [2]_q n \left(\frac{1}{1-q} \right)^{n-1} \sum_{l=0}^{n-1} \frac{(-1)^l}{1+q^l} q^{lx} \binom{n-1}{l}.$$

By (19) we also see that

$$\begin{aligned} \sum_{n=0}^{\infty} G_{n,q}(x) \frac{t^n}{n!} &= [2]_q t \sum_{k=0}^{\infty} (-1)^k e^{[k+x]_q t} = [2]_q t \sum_{a=0}^{m-1} (-1)^a \sum_{k=0}^{\infty} (-1)^k e^{[k+\frac{a+x}{m}]_q t} [m]_q t \\ &= \frac{[2]_q}{[m]_q [2]_q^m} \sum_{a=0}^{m-1} (-1)^a \left([m]_q t [2]_q^m \sum_{k=0}^{\infty} (-1)^k e^{[m]_q t [k+\frac{a+x}{m}]_q t} \right) \\ &= \sum_{n=0}^{\infty} \left(\frac{[2]_q}{[m]_q [2]_q^m} \sum_{a=0}^{m-1} (-1)^a [m]_q^n G_{n,q^m} \left(\frac{x+a}{m} \right) \right) \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} \left(\frac{[2]_q}{[2]_q^m} [m]_q^{n-1} \sum_{a=0}^{m-1} (-1)^a G_{n,q^m} \left(\frac{x+a}{m} \right) \right) \frac{t^n}{n!}, \text{ where } m \in \mathbb{N} \text{ odd.} \end{aligned}$$

Therefore we obtain the following theorem:

Theorem 4. *Let $m(= \text{odd}) \in \mathbb{N}$. Then we have the distribution of the q -Genocchi polynomials as follows:*

$$G_{n,q}(x) = \frac{[2]_q}{[2]_q^m} [m]_q^{n-1} \sum_{a=0}^{m-1} (-1)^a G_{n,q^m} \left(\frac{x+a}{m} \right),$$

where n is positive integer.

Theorem 4 will be used to construct the p -adic q -Genocchi measures which will be treated in the next section. Let χ be a primitive Dirichlet character with a conductor $d(= \text{odd}) \in \mathbb{N}$. Then the generalized q -Genocchi numbers attached to χ are defined as

$$(21) \quad g_{\chi,q}(t) = [2]_q t \sum_{a=0}^{d-1} \chi(a) (-1)^a e^{[a]_q t} = \sum_{n=0}^{\infty} G_{n,\chi,q} \frac{t^n}{n!}.$$

From (21) we derive

$$(22) \quad G_{n,\chi,q} = \frac{[2]_q}{[2]_q^d} [d]_q^{n-1} \sum_{a=0}^{d-1} (-1)^a \chi(a) G_{n,q^d} \left(\frac{a}{d} \right).$$

§3. p -adic q -Genocchi measures

In this section we assume that $q \in \mathbb{C}_p$ with $|1 - q|_p < p^{-\frac{1}{p-1}}$ so that $q^x = \exp(x \log q)$. Let χ be a primitive Dirichlet's character with a conductor $d(= \text{odd}) \in \mathbb{N}$. For any positive integers N, k and $d(= \text{odd})$, let $\mu_k = \mu_{k,q;G}$ be defined as

$$(23) \quad \mu_k(a + dp^N \mathbb{Z}_p) = (-1)^a [dp^N]_q^{k-1} \frac{[2]_q}{[2]_{q^{dp^N}}} G_{k,q^{dp^N}} \left(\frac{a}{dp^N} \right).$$

By using Theorem 4 and (23), we show that

$$\sum_{i=0}^{p-1} \mu_k(a + idp^N + dp^{N+1} \mathbb{Z}_p) = \mu_k(a + dp^N \mathbb{Z}_p).$$

Therefore we obtain the following theorem:

Theorem 5. *Let d be odd positive integer. For any positive integers N, k , and let $\mu_k = \mu_{k,q;G}$ be defined as*

$$\mu_k(a + dp^N \mathbb{Z}_p) = (-1)^a [dp^N]_q^{k-1} \frac{[2]_q}{[2]_{q^{dp^N}}} G_{k,q^{dp^N}} \left(\frac{a}{dp^N} \right).$$

Then μ_k can be extended to a distribution on X .

From the definition of μ_k and (22) we note that

$$\int_X \chi(x) d\mu_k(x) = G_{k,\chi,q}.$$

By (14) and (16), it is not difficult to show that

$$(24) \quad G_{n,q}(x) = \sum_{k=0}^n \binom{n}{k} [x]_q^{n-k} q^{kx} G_{k,q}.$$

From (23) and (24) we derive

$$(25) \quad d\mu_k(a) = \lim_{N \rightarrow \infty} \mu_k(a + dp^N \mathbb{Z}_p) = k[a]_q^{k-1} d\mu_{-q}(a).$$

Therefore we obtain the following corollary:

Corollary 6. *Let k be a positive integer. Then we have*

$$G_{k,\chi,q} = \int_X \chi(x) d\mu_k(x) = k \int_X \chi(x) [x]_q^{k-1} d\mu_{-q}(x).$$

Moreover,

$$G_{k,q} = k \int_X [x]_q^{k-1} d\mu_{-q}(x).$$

Remark. In the recent paper(see [1]) Cenkci-Can-Kurt have studied q -Genocchi numbers and polynomials and p -adic q -Genocchi measures. Our q -Genocchi numbers and polynomials to treat in this paper are different their q -Genocchi numbers and polynomials.

REFERENCES

- [1] M. Cenkci, M. Can and V. Kurt, *q-extension of Genocchi Numbers*, J. Korean Math. Soc. **43** (2006), 183–198.
- [2] M. Cenkci, M. Can, *Some results on q-analogue of the Lerch zeta function*, Adv. Stud. Contemp. Math. **12** (2006), 213–223.
- [3] F. T. Howard, *Applications of a recurrence formula for the Bernoulli numbers*, J. Number Theory **52** (1995), 157–172.
- [4] T. Kim, *A note on q-Volkenborn integration*, Proc. Jangjeon Math. Soc. **8** (2005), 13–17.
- [5] T. Kim, *q-Volkenborn integration*, Russ. J. Math. Phys. **9** (2002), 288–299.
- [6] T. Kim, *A note on p-adic invariant integral in the rings of p-adic integers*, Advan. Stud. Contemp. Math. **13** (2006), 95–99.
- [7] T. Kim, L.-C. Jang, H. K. Pak, *A note on q-Euler and Genocchi numbers*, Proc. Japan Acad. Ser. A Math. Sci. **77** (2001), 139–141.
- [8] T. Kim, *A note on some formulas for the q-Euler numbers and polynomials*, Proc. Jangjeon Math. Soc. **9** (2006), 227–232.
- [9] T. Kim, *On the twisted q-Euler numbers and polynomials associated with basic q-l-functions*, J. Math. Anal. Appl. (in press).
- [10] T. Kim, *The modified q-Euler numbers and polynomials*, arXiv: math.NT/0702523 (2006).
- [11] T. Kim, *An invariant p-adic q-integral on \mathbb{Z}_p* , Appl. Math. Lett.(2007), doi:10.1016/j.aml.
- [12] H. M. Srivastava, T. Kim and Y. Simsek, *q-Bernoulli numbers and polynomials associated with multiple q-zeta functions and basic L-series*, Russ. J. Math. Phys. **12** (2005), 241–268.

Taekyun Kim

EECS, Kyungpook National University, Taegu 702-701, S. Korea

e-mail: tkim@knu.ac.kr; tkim64@hanmail.net