# Solvability of Backward Stochastic Differential Equations with Quadratic Growth

Revaz Tevzadze

Georgian-American University, Business School, 3, Alleyway II, Chavchavadze Ave. 17 a, Georgian Technical University, 77 Kostava str., 0175, Institute of Cybernetics, 5 Euli str., 0186, Tbilisi, Georgia (e-mail: reztev@yahoo.com)

We prove the existence of the unique solution of a general Backward Stochastic Differential Equation with quadratic growth driven by martingales. Some kind of comparison theorem is also proved.

**Key words and phrases**:Backward Stochastic Differential Equation, Contraction principle, BMO-martingale.

Mathematics Subject Classification (2000): 90A09, 60H30, 90C39.

## 1 Introduction

In this paper we show a general result of existence and uniqueness of Backward Stochastic Differential Equation (BSDE) with quadratic growth driven by continuous martingale. Backward stochastic differential equations have been introduced by Bismut [1] for the linear case as equations of the adjoint process in the stochastic maximum principle. A nonlinear BSDE (with Bellman generator) was first considered by Chitashvili [4]. He derived the semimartingale BSDE (or SBE), which can be considered as a stochastic version of the Bellman equation for a stochastic control problem, and proved the existence and uniqueness of a solution. The theory of BSDEs driven by the Brownian motion was developed by Pardoux and Peng [22] for more general generators. The results of Pardoux and Peng were generalized by Kobylansky [11], Lepeltier and San Martin [12] for generators with quadratic growth. In the work of Hu at all [8] BMO-martingales were used for BSDE with quadratic generators in Brownian setting and in [15], [16], [17], [18], [19], [21]

for BSDEs driven by martingales. By Chitashvili [4], Buckdahn [3], and El Karoui and Huang [7] the well posedness of BSDE with generators satisfying Lipschitz type conditions was established. Here we suggest new approach including an existence and uniqueness of the solution of general BSDE with quadratic growth. In the earlier papers [15], [16], [17], [18], [19] we studied, as well as Bobrovnytska and Schweizer [2], the particular cases of BSDE with quadratic nonlinearities related to the primal and dual problems of Mathematical Finance. In these works the solutions were represented as a value function of the corresponding optimization problems.

The paper is organized as follows. In Section 2 we give some basic definitions and facts used in what follows. In Section 3 we show the solvability of the system of BSDEs for sufficiently small initial condition and further prove the solvability of one dimensional BSDE for arbitrary bounded initial data. At the end of Section 4 we prove the comparison theorem, which generalizes the results of Mania and Schweizer [14], and apply this results to the uniqueness of the solution.

### 2 Some basic definitions and assumptions

Let  $(\Omega, \mathcal{F}, \mathbf{F} = (F_t)_{t \geq 0}, P)$  be filtered probability space satisfying the usual conditions. We assume that all local martingales with respect to  $\mathbf{F}$  are continuous. Here the time horizon  $T < \infty$  is a stopping time and  $\mathcal{F} = F_T$ . Let us consider Backward Stochastic Differential Equation (BSDE) of the form

$$dY_t = -f(t, Y_t, \sigma_t^* Z_t) dK_t - d\langle N \rangle_t g_t + Z_t^* dM_t + dN_t, \tag{2.1}$$

$$Y_T = \xi \tag{2.2}$$

We suppose that

- $(M_t, t \ge 0)$  is an  $R^n$ -valued continuous martingale with cross-variations matrix  $\langle M \rangle_t = (\langle M^i, M^j \rangle_t)_{1 \le i,j \le n}$ ,
- $(K_t, t \ge 0)$  is a continuous, adapted, increasing process, such that  $\langle M \rangle_t = \int_0^t \sigma_s \sigma_s^* dK_s$  for some predictable, non degenerate  $n \times n$  matrix  $\sigma$ ,
- $\xi$  is  $\mathcal{F}$ —measurable an  $\mathbb{R}^d$ -valued random variable,
- $f: \Omega \times R^+ \times R^d \times R^{n \times d} \to R^d$  is a stochastic process, such that for any  $(y, z) \in R^d \times R^{n \times d}$  the process  $f(\cdot, \cdot, y, z)$  is predictable,
- $g: \Omega \times \mathbb{R}^+ \to \mathbb{R}^{d \times d}$  is a predictable process.

The notation  $R^{n\times d}$  here denotes the space of  $n\times d$ -matrix C with Euclidian norm  $|C|=\sqrt{\operatorname{tr}(CC^*)}$ . For some stochastic process  $X_t$  and sopping times  $\tau$ ,  $\nu$ , such that  $\tau\geq\nu$  we denote  $X_{\nu,\tau}=X_{\tau}-X_{\nu}$ . For all unexplained notations concerning the martingale theory used below we refer [9], [5] and [13]. About BMO-martingales see [6] or [10].

A solution of the BSDE is a triple (Y, Z, N) of stochastic processes, such that (2.1), (2.2) is satisfied and

- Y is an adapted  $\mathbb{R}^d$ -valued continuous process,
- Z is an  $R^{n \times d}$ -valued predictable process,
- N is an  $\mathbb{R}^d$ -valued continuous martingale, orthogonal to the basic martingale M.

One says that  $(f, g, \xi)$  is a generator of BSDE (2.1),(2.2).

We introduce the following spaces

- $L^{\infty}(\mathbb{R}^d) = \{X : \Omega \to \mathbb{R}^d, \mathcal{F}_T \text{measurable}, ||X||_{\infty} = \underset{\Omega}{\text{ess sup}} |X(\omega)| < \infty\},$
- $S^{\infty}(R^d) = \{ \varphi : \Omega \times R^+ \to R^d, \text{ continuous, adapted, } ||\varphi||_{\infty} = \underset{[[0,T]]}{\operatorname{ess \, sup}} |\varphi(t,\omega)| < \infty \},$

•

$$H^{2}(R^{n\times d},\sigma) = \{\varphi: \Omega \times R^{+} \to R^{n\times d}, \text{ predictable,} \}$$
$$||\varphi||_{H}^{2} = \underset{[[0,T]]}{\text{ess sup}} E(\int_{t}^{T} |\sigma_{s}^{*}\varphi_{s}|^{2} dK_{s}|\mathcal{F}_{t}) \equiv \underset{[[0,T]]}{\text{ess sup}} E(\text{tr}\langle\varphi\cdot M\rangle_{tT}|\mathcal{F}_{t}) < \infty\}, (2.3)$$

• BMO(Q) = {N,  $R^d$  - valued Q - martingale  $||N||_Q^2 = \operatorname{ess\,sup} E^Q(tr\langle N \rangle_{tT} | \mathcal{F}_t) < \infty$ }

We also use the notation  $|r|_{2,\infty}$  for the norm  $||\int_0^T r_s^2 dK_s||_{\infty}$ . The norm of the triple is defined as

$$||(Y, Z, N)||^2 = ||Y||^2 + ||Z||_H^2 + ||N||_P^2.$$

Throughout the paper we use the condition

A) There exist a constant  $\theta$  and predictable processes

$$\alpha: \Omega \times R^+ \to R^d, \ \Gamma: \Omega \times R^+ \to Lin(R^{n \times d}, R^d), \ r: \Omega \times R^+ \to R,$$

such that the following conditions  $\int_0^T r_s dK_s$ ,  $\int_0^T r_s^2 dK_s \in L^{\infty}$ ,  $\Gamma(\sigma^{-1}) \in H_T^2$ ,  $|\alpha_t| \leq r_t$ ,  $|g_t| \leq \theta^2$  and

$$|f(t, y_1, z_1) - f(t, y_2, z_2) - \alpha_t(y_1 - y_2) - \Gamma_t(z_1 - z_2)|$$

$$\leq (r_t|y_1 - y_2| + \theta|z_1 - z_2|)(r_t(|y_1| + |y_2|) + \theta(|z_1| + |z_2|)).$$
(2.4)

are satisfied.

Sometimes we use the more restrictive conditions

B1) 
$$\int_0^T |f(t,0,0)| dK_t + |g_t| \le \theta^2$$
 for all  $t \in [0,T]$ ,

B2) 
$$|f_y(t, y, z)| \le r_t$$
,  $|f_z(t, y, z)| \le r_t + \theta |z|$  for all  $(t, y, z)$ ,

B3) 
$$|f_{yy}(t, y, z)| \le r_t^2$$
,  $|f_{yz}(t, y, z)| \le \theta r_t$ ,  $|f_{zz}(t, y, z)| \le \theta^2$  for all  $(t, y, z)$ .

**Remark 1.** Condition A) follow from conditions B1)-B3), since using notations  $\delta y = y_1 - y_2$ ,  $\delta z = z_1 - z_2$  for  $\alpha_t = f_y(t, 0, 0)$ ,  $\Gamma_t = f_z(t, 0, 0)$  by the mean value theorem we have

$$|f(t, y_1, z_1) - f(t, y_2, z_2) - \alpha_t \delta y - \Gamma_t(\delta z)|$$

$$= |f_y(t, \nu y_1 + (1 - \nu)y_2, \nu z_1 + (1 - \nu)z_2)\delta y - f_y(t, 0, 0)\delta y|$$

$$+ f_z(t, \nu y_1 + (1 - \nu)y_2, \nu z_1 + (1 - \nu)z_2)(\delta z) - f_z(t, 0, 0)(\delta z)|,$$

for some  $\nu \in [0,1]$ . Using again mean value theorem we obtain that

$$|f(t, y_{1}, z_{1}) - f(t, y_{2}, z_{2}) - \alpha_{t} \delta y - \Gamma_{t}(\delta z)|$$

$$\leq (|\nu y_{1} + (1 - \nu)y_{2}| \max_{y, z} |f_{yy}(t, y, z)| + |\nu z_{1} + (1 - \nu)z_{2}| \max_{y, z} |f_{yz}(t, y, z)|)|\delta y|$$

$$+(|\nu y_{1} + (1 - \nu)y_{2}| \max_{y, z} |f_{yz}(t, y, z)| + |\nu z_{1} + (1 - \nu)z_{2}| \max_{y, z} |f_{zz}(t, y, z)|)|\delta z|$$

$$\leq [r_{t}^{2}(|y_{1}| + |y_{2}|) + r_{t}\theta(|z_{1}| + |z_{2}|)]|\delta y| + [r_{t}\theta(|y_{1}| + |y_{2}|) + \theta^{2}(|z_{1}| + |z_{2}|)]|\delta z|$$

$$= (r_{t}|\delta y| + \theta|\delta z|)(r_{t}(|y_{1}| + |y_{2}|) + \theta(|z_{1}| + |z_{2}|).$$

**Remark 2.** If d = 1 the operator  $\Gamma_t$  is given by an n-dimensional vector  $\gamma_t$  such that  $\Gamma_t(z) = \gamma_t^* z$ . Thus inequality in A) can be rewritten as

$$|f(t, y_1, z_1) - f(t, y_2, z_2) - \alpha_t \delta y - \gamma_t^* \delta z|$$

$$\leq (r_t |\delta y| + \theta |\delta z|) (r_t (|y_1| + |y_2|) + \theta (|z_1| + |z_2|)).$$

The main statement of the paper is the following

**Theorem 1.** Let  $\xi \in L^{\infty}$ , d = 1 and conditions B1)-B3) are satisfied. Then there exists a unique triple (Y, Z, N), where  $Y \in S^{\infty}$ ,  $Z \in H^2$ ,  $N \in BMO$ , that satisfies equation (2.1), (2.2).

#### 3 Existence of the solution

First we prove the existence and uniqueness of the solution for a sufficiently small initial data.

**Proposition 1.** Let f and g satisfy condition A) with  $\alpha = 0$  and  $\gamma_t = 0$ . Then for  $\xi$  with the norm  $||\xi||_{\infty} < \frac{1}{32\beta}$ ,  $\beta = 8 \max(|r|_{2,\infty}^2, \theta^2)$  there exists a unique solution (Y, Z, N) of BSDE

$$dY_{t} = (f(t, 0, 0) - f(t, Y_{t}, \sigma_{t}^{*} Z_{t}))dK_{t} + d\langle N \rangle_{t} g_{t} + Z_{t}^{*} dM_{t} + dN_{t},$$

$$Y_{T} = \xi,$$
(3.1)

with the norm  $||(Y, Z, N)|| \le R$ , where R is a constant satisfying the inequality  $4||\xi||_{\infty}^2 + \beta^2 R^4 \le R^2$ , namely  $R = 2\sqrt{2}||\xi||_{\infty}$ .

Moreover if  $||\xi||_{\infty} + ||\int_0^{\infty} |f(s,0,0)| dK_s||_{\infty}$  is small enough then BSDE (2.1) admits a unique solution.

*Proof.* We define the mapping (Y, Z, N) = F(y, z, n), n is orthogonal to M,  $(y, z \cdot M + n) \in S_T^{\infty} \times BMO(P)$  by the relation

$$dY_{t} = (f(t, 0, 0) - f(t, y_{t}, \sigma_{t}^{*} z_{t}))dK_{t} + d\langle n \rangle_{t} g_{t} + Z_{t}^{*} dM_{t} + dN_{t},$$

$$Y_{T} = \xi.$$
(3.2)

Using the Ito formula for  $|Y_t|^2$  we obtain that

$$|Y_{t}|^{2} = |\xi|^{2} + 2 \int_{t}^{T} Y_{s}^{*}(f(s, y_{s}, \sigma_{s}^{*}z_{s}) - f(s, 0, 0)) dK_{t}$$

$$+2 \int_{t}^{T} Y_{s}^{*}d\langle n \rangle_{s}g_{s} - \int_{t}^{T} \operatorname{tr}Z_{s}^{*}d\langle M \rangle_{s}Z_{s} - \operatorname{tr}\langle N \rangle_{tT} - \int_{t}^{T} Y_{s}^{*}Z_{s}^{*}dM_{s} - \int_{t}^{T} Y_{s}^{*}dN_{s}.$$

If we take the conditional expectation and use (2.3) and the elementary inequality  $2ab \le \frac{1}{4}a^2 + 4b^2$  we get

$$|Y_{t}|^{2} + E\left(\int_{t}^{T} |\sigma_{s}^{*}Z_{s}|^{2} dK_{s} + tr\langle N \rangle_{tT} |\mathcal{F}_{t}\right) \leq ||\xi||^{2} + \frac{1}{4}||Y||_{\infty}^{2}$$

$$+4E^{2}\left(\int_{t}^{T} |f(s, y_{s}, \sigma_{s}^{*}z_{s}) - f(s, 0, 0)| dK_{s} + \int_{t}^{T} |g_{s}| d\text{tr}\langle n \rangle_{s} |\mathcal{F}_{t}\right). \tag{3.3}$$

Thus using condition A), identities

$$\operatorname{tr}\langle z \cdot M \rangle_t = \operatorname{tr} \int_0^t z_s^* d\langle M \rangle_s z_s = \int_0^t \operatorname{tr}(z_s^* \sigma_s \sigma_s^* z_s) dK_s = \int_0^t |\sigma_s^* z_s|^2 dK_s$$
 (3.4)

and explicit inequalities

$$\frac{1}{2}(||Y||_{\infty}^{2} + ||Z \cdot M + N||_{\text{BMO}}^{2}) \leq \max(||Y||_{\infty}^{2}, ||Z \cdot M + N||_{\text{BMO}}^{2}) 
\leq \underset{[[0,T]]}{\operatorname{ess sup}}[|Y_{t}|^{2} + E(\int_{t}^{T} |\sigma_{s}^{*}Z_{s}|^{2} dK_{s} + tr\langle N \rangle_{tT} |\mathcal{F}_{t})]$$

we obtain from (3.3)

$$\frac{1}{4}||Y||_{\infty}^{2} + \frac{1}{2}||Z \cdot M + N||_{\text{BMO}}^{2} \leq ||\xi||^{2}$$

$$+4 \operatorname{ess\,sup} E^{2} \left( \int_{t}^{T} |f(s, y_{s}, \sigma_{s}^{*} z_{s}) - f(s, 0, 0)| dK_{t} + \theta^{2} \operatorname{tr} \langle n \rangle_{tT} |\mathcal{F}_{t} \right)$$

$$\leq ||\xi||^{2} + 16 \operatorname{ess\,sup} E^{2} \left( \int_{t}^{T} r_{s}^{2} y_{s}^{2} dK_{s} + \theta^{2} \operatorname{tr} \langle z \cdot M + n \rangle_{tT} |\mathcal{F}_{t} \right)$$

$$\leq ||\xi||^{2} + 16 |r|_{2,\infty}^{4} ||y||_{\infty}^{4} + 16 \theta^{4} ||z \cdot M + n||_{\text{BMO}}^{4}.$$

Therefore

$$||Y||_{\infty}^{2} + ||Z \cdot M + N||_{\text{BMO}}^{2} \le 4||\xi||^{2} + 64|r|_{2,\infty}^{4}||y||_{\infty}^{4} + 64\theta^{4}||z \cdot M + n||_{\text{BMO}}^{4}$$

$$< 4||\xi||^{2} + \beta^{2}(||y||_{\infty}^{2} + ||z \cdot M + n||_{\text{PMO}}^{2})^{2}.$$

where  $\beta = 8 \max(|r|_{2,\infty}^2, \theta^2)$ . We can pick R such that

$$4||\xi||^2 + \beta^2 R^4 \le R^2$$

if and only if  $||\xi||_{\infty} \leq \frac{1}{4\beta}$ . For instance  $R = 2\sqrt{2}||\xi||_{\infty}$  satisfies this quadratic inequality. Therefore the ball

$$\mathcal{B}_R = \{ (Y, Z \cdot M + N) \in S^{\infty} \times BMO, \ N \perp M, \ ||Y||_{\infty}^2 + ||Z \cdot M + N||_{BMO}^2 \le R^2 \}$$

is such that  $F(\mathcal{B}_R) \subset \mathcal{B}_R$ .

Similarly for  $(y^j, z^j \cdot M + n^j) \in \mathcal{B}_R$ , j = 1, 2 using the notations  $\delta y = y^1 - y^2$ ,  $\delta z = z^1 - z^2$ ,  $\delta n = n^1 - n^2$  we can show that

$$||\delta Y||_{\infty}^{2} + ||\delta Z \cdot M + \delta N||_{\text{BMO}}^{2}$$

$$\leq 4 \operatorname{ess\,sup} E^{2} \Big( \int_{t}^{T} |f(s, y_{s}^{1}, \sigma_{s}^{*} z_{s}^{1}) - f(s, y_{s}^{2}, \sigma_{s}^{*} z_{s}^{2}) |dK_{s}$$

$$+ \int_{t}^{T} |g_{s}| d \operatorname{var} (\operatorname{tr} \langle \delta n, n^{1} + n^{2} \rangle)_{s} |\mathcal{F}_{t})$$

$$\leq 8 \operatorname{ess\,sup} E \Big( \int_{t}^{T} (r_{s}^{2} |\delta y_{s}|^{2} + \theta^{2} |\sigma_{s}^{*} \delta z_{s}|^{2} dK_{s}) |\mathcal{F}_{t})$$

$$\times E \Big( \int_{t}^{T} (r_{s} (|y_{s}^{1}| + |y_{s}^{2}|) + \theta (|\sigma_{s}^{*} z_{s}^{1}| + |\sigma_{s}^{*} z_{s}^{2}|))^{2} dK_{s}) |\mathcal{F}_{t})$$

$$+ \theta^{2} E (\operatorname{tr} \langle \delta n \rangle_{tT} |\mathcal{F}_{s}) E (\operatorname{tr} \langle n^{1} + n^{2} \rangle_{tT} |\mathcal{F}_{t})$$

Again using the equalities (3.4) we can pass to the norm. Thus

$$\begin{split} ||\delta Y||_{\infty}^{2} + ||\delta Z \cdot M + \delta N||_{\text{BMO}}^{2} \\ &\leq 8(|r|_{2,\infty}^{2}||\delta y||_{\infty}^{2} + \theta^{2}||\delta z \cdot M||_{\text{BMO}}^{2}) \\ \times (|r|_{2,\infty}^{2}(||y^{1}||_{\infty}^{2} + ||y^{2}||_{\infty}^{2}) + \theta^{2}(||z^{1} \cdot M||_{P}^{2} + ||z^{2} \cdot M||_{P}^{2}) \\ &+ 2\theta^{2}||\delta n||_{\text{BMO}}^{2}(||n^{1}||_{\text{BMO}}^{2} + ||n^{2}||_{\text{BMO}}^{2})^{2}). \end{split}$$

Since  $||z^1 \cdot M||, ||z^2 \cdot M|| \le R, ||n^1||, ||n^2|| \le R$  we get

$$||\delta Y||_{\infty}^{2} + ||\delta Z \cdot M + \delta N||_{\text{BMO}}^{2}$$

$$\leq 128\beta^{2}R^{2}(||\delta y||_{\infty}^{2} + ||\delta z \cdot M||_{\text{BMO}}^{2}) + 4\beta^{2}R^{2}||\delta n||_{\text{BMO}}^{2}$$

$$\leq 128\beta^{2}R^{2}(||\delta y||_{\infty}^{2} + ||\delta z \cdot M + \delta n||_{\text{BMO}}^{2}).$$
(3.5)

Now we can take  $R = 2\sqrt{2}||\xi||_{\infty} < \frac{1}{8\sqrt{2}\beta}$ . This means that  $||\xi||_{\infty} < \frac{1}{32\beta}$  and F is contraction on  $\mathcal{B}_R$ . By contraction principle the mapping F admits a unique fixed point, which is the solution of (3.1).

From now we suppose that d = 1.

**Lemma 1**. Let condition A) is satisfied. Then the generator  $(\bar{f}, \bar{g}, \bar{\xi})$ , where

$$\bar{f}(t, \bar{y}, \bar{z}) = e^{\int_0^t \alpha_s dK_s} (f(t, e^{-\int_0^t e^{\alpha_s dK_s}} \bar{y}, e^{-\int_0^t e^{\alpha_s dK_s}} \bar{z}) - f(t, 0, 0)) - \alpha_t \bar{y} - \gamma_t^* \bar{z},$$

$$\bar{g}_t = e^{-\int_0^t \alpha_s dK_s} g_t$$
 and  $\bar{\xi} = e^{\int_0^T \alpha_s dK_s} \xi_s$ 

satisfies condition A) with  $\alpha = 0$ ,  $\gamma = 0$ ,  $\bar{r}_t = r_t e^{||\int_0^\infty r_s dK_s||_\infty}$ , and  $\bar{\theta} = \theta e^{||\int_0^T r_s dK_s||_\infty}$ . Moreover, (Y, Z, N) is a solution of BSDE (3.1) if and only if

$$(\bar{Y}_t, \bar{Z}_t, \bar{N}_t) = (e^{\int_0^t \alpha_s dK_s} Y_t, e^{\int_0^t \alpha_s dK_s} Z_t, \int_0^t e^{\int_0^s \alpha_u dK_u} dN_s)$$

is a solution w.r.t. measure  $d\bar{P} = \mathcal{E}_T((\gamma \sigma^{-1}) \cdot M)dP$  of BSDE

$$d\bar{Y}_t = -\bar{f}(t, \bar{Y}_t, \sigma_t^* \bar{Z}_t) dK_t - d\langle \bar{N} \rangle_t \bar{g}_t + \bar{Z}_t^* d\bar{M}_t + d\bar{N}_t,$$

$$\bar{Y}_T = \bar{\xi},$$
(3.6)

where  $\bar{M}_t = M_t - \langle (\gamma \sigma^{-1}) \cdot M, M \rangle_t$ .

*Proof.* Condition A) for  $(\bar{f}, \bar{g}, \bar{\xi})$  is satisfied since by (2.4)

$$\begin{split} |\bar{f}(t,\bar{y}_{1},\bar{z}_{1}) - \bar{f}(t,\bar{y}_{2},\bar{z}_{2})| \\ &\leq e^{\int_{0}^{t} \alpha_{s} dK_{s}} (r_{t}|\delta \bar{y}| + \theta|\delta \bar{z}|) (r_{t}(|\bar{y}_{1}| + |\bar{y}_{2}|) + \theta(|\bar{z}_{1}| + |\bar{z}_{2}|)) \\ &\leq (\bar{r}_{t}|\delta \bar{y}| + \bar{\theta}|\delta \bar{z}|) (\bar{r}_{t}(|\bar{y}_{1}| + |\bar{y}_{2}|) + \bar{\theta}(|\bar{z}_{1}| + |\bar{z}_{2}|)). \end{split}$$

On the other hand using the Ito formula we have

$$d\bar{Y}_t = e^{\int_0^t \alpha_s dK_s} dY_t + \alpha_t e^{\int_0^t \alpha_s dK_s} Y_t dK_t$$

$$= e^{\int_0^t \alpha_s dK_s} (f(t, 0, 0) - f(t, Y_t, \sigma_t^* Z_t)) dK_t + e^{\int_0^t \alpha_s dK_s} d\langle N \rangle_t g_t$$

$$+ e^{\int_0^t \alpha_s dK_s} Z_t^* dM_t + e^{\int_0^t \alpha_s dK_s} dN_t + \alpha_t \bar{Y}_t dK_t$$

Taking into account that

$$\begin{split} e^{\int_0^t \alpha_s dK_s} (f(t,0,0) - f(t,Y_t,\sigma_t^* Z_t)) + \alpha_t \bar{Y}_t \\ &= -\bar{f}(t,\bar{Y}_t,\sigma_t^* \bar{Z}_t) - \gamma_t \sigma_t^* \bar{Z}_t, \\ e^{\int_0^t \alpha_s dK_s} d\langle N \rangle_t g_t &= d\langle \bar{N} \rangle_t e^{-\int_0^t \alpha_s dK_s} g_t = d\langle \bar{N} \rangle_t \bar{g}_t \end{split}$$

and

$$\bar{Z} \cdot M - \int_0^{\cdot} \gamma_t \sigma_t^* \bar{Z}_t dK_t = \bar{Z} \cdot M - \int_0^{\cdot} \gamma_t \sigma_t^{-1} \sigma_t \sigma_t^* \bar{Z}_t dK_t$$

$$= \bar{Z} \cdot M - \int_0^{\cdot} \gamma_t \sigma_t^{-1} d\langle M \rangle_t \bar{Z}_t = \bar{Z} \cdot M - \langle (\gamma \cdot \sigma^{-1}) \cdot M, \bar{Z} \cdot M \rangle = \bar{Z} \cdot \bar{M}$$

we obtain

$$d\bar{Y}_t = -\bar{f}(t, \bar{Y}_t, \sigma_t^* \bar{Z}_t) dK_t - d\langle \bar{N} \rangle_t \bar{g}_t + \bar{Z}_t d\bar{M}_t + d\bar{N}_t.$$

Here  $\bar{M}$  is a local martingale w.r.t.  $\bar{P}$  by Girsanov theorem.

Corollary 1. Let f and g satisfy condition A) and  $||\xi||_{\infty} \leq \frac{1}{32\beta} \exp(-2||\int_0^T r_s dK_s||_{\infty})$ . Then there exist the solution of (3.1) with the norm  $||Y||_{\infty}^2 + ||Z \cdot \bar{M} + N||_{\mathrm{BMO}(\bar{P})}^2 \leq \frac{1}{128\beta^2}$ . *Proof.* Obviously that

$$||Y||_{\infty}^{2} + ||Z \cdot \bar{M} + N||_{\mathrm{BMO}(\bar{P})}^{2} \leq \left(||\bar{Y}||_{\infty}^{2} + ||\bar{Z} \cdot \bar{M} + \bar{N}||_{\mathrm{BMO}(\bar{P})}^{2}\right) \exp(2||\int_{0}^{T} r_{s} dK_{s}||_{\infty})$$

$$\leq 8||\bar{\xi}||_{\infty}^{2} \exp(2||\int_{0}^{T} r_{s} dK_{s}||_{\infty}) \leq 8||\xi||_{\infty}^{2} \exp(4||\int_{0}^{T} r_{s} dK_{s}||_{\infty}).$$

From  $||\xi||_{\infty} \leq \frac{1}{32\beta} \exp(-2||\int_0^T r_s dK_s||_{\infty})$  follows that  $8||\xi||_{\infty} \exp(4||\int_0^T r_s dK_s||_{\infty}) \leq \frac{1}{128\beta^2}$ . Hence we get  $||Y||_{\infty}^2 + ||Z \cdot \bar{M} + N||_{\text{BMO}(\bar{\mathbb{P}})}^2 \leq \frac{1}{128\beta^2}$ .

Corollary 2. Let generator  $(f, g, \xi)$  satisfies conditions B1)-B3) and  $(\tilde{Y}_t, \tilde{Z}_t, \tilde{N}_t)$  be a solution of (3.1). Then BSDE

$$d\hat{Y}_{t} = (f(t, \tilde{Y}_{t}, \sigma_{t}^{*} \tilde{Z}_{t}) - f(t, \hat{Y}_{t} + \tilde{Y}_{t}, \sigma_{t}^{*} \hat{Z}_{t} + \sigma_{t}^{*} \tilde{Z}_{t}))dK_{t}$$

$$-d(\langle \hat{N} \rangle_{t} + 2\langle \tilde{N}, \hat{N} \rangle_{t})g_{t} + \hat{Z}_{t}^{*}dM_{t} + d\hat{N}_{t},$$

$$\hat{Y}_{T} = \hat{\xi}$$

$$(3.7)$$

satisfy condition A) with  $-\hat{f}(t,y,z) = f(t,\tilde{Y}_t,\sigma_t^*\tilde{Z}_t) - f(t,y+\tilde{Y}_t,z+\sigma_t^*\tilde{Z}_t)$ ,  $\alpha_t = f_y(t,\tilde{Y}_t,\sigma_t^*\tilde{Z}_t)$ ,  $\gamma_t = f_z(t,\tilde{Y}_t,\sigma_t^*\tilde{Z}_t)$  and the new probability measure  $\mathcal{E}_T(2g\cdot\tilde{N})dP$ . Moreover (3.7) admits a unique solution  $(\hat{Y}_t,\hat{Z}_t,\hat{N}_t)$  if  $||\hat{\xi}||_{\infty} \leq \frac{1}{32\beta}\exp(-2||\int_0^t r_s dK_s||_{\infty})$ .

*Proof.* Using a change of measure the equation (3.7) reduces to equation of type (3.1). By previous corollary we obtain the existence and uniqueness of the BSDE.

**Lemma 2.** Let conditions B1)-B3) be satisfied and random variables  $\tilde{\xi}$  and  $\hat{\xi}$  be such that  $\max(||\tilde{\xi}||_{\infty}, ||\hat{\xi}||_{\infty}) \leq \frac{1}{32\beta}e^{-2||\int_0^T r_s^2 dK_s||_{\infty}}$ . Then there exist solutions of BSDEs (3.7) and

$$d\tilde{Y}_t = (f(t,0,0) - f(t,\tilde{Y}_t,\sigma_t^*\tilde{Z}_t))dK_t - d\langle \tilde{N} \rangle_t g_t + \tilde{Z}_t^* dM_t + d\tilde{N}_t,$$

$$\tilde{Y}_T = \tilde{\xi}$$
(3.8)

and the triple  $(Y, Z, N) = (\tilde{Y} + \hat{Y}, \tilde{Z} + \hat{Z}, \tilde{N} + \hat{N})$  satisfies BSDE

$$dY_t = (f(t, 0, 0) - f(t, Y_t, \sigma_t^* Z_t))dK_t - d\langle N \rangle_t g_t + Z_t^* dM_t + dN_t,$$
$$Y_T = \tilde{\xi} + \hat{\xi}.$$

*Proof.* Similarly to the Remark from Section 1 we can show that for  $\hat{f}(t, y, z) = f(t, \tilde{Y}_t, \sigma_t^* \tilde{Z}_t) - f(t, y + \tilde{Y}_t, \sigma_t^* z + \sigma_t^* \tilde{Z}_t)$ ,  $\alpha_t = f_y(t, \tilde{Y}_t, \sigma_t^* \tilde{Z}_t)$ ,  $\gamma_t = f_z(t, \hat{Y}_t, \sigma_t^* \hat{Z}_t)$  the estimate

$$|\hat{f}(t, y_1, z_1) - \hat{f}(t, y_2, z_2) - \alpha_t \delta y - \gamma_t^* \delta z|$$

$$\leq (r_t |\delta y| + \theta |\delta z|) (r_t (|y_1| + |y_2|) + \theta (|z_1| + |z_2|)).$$

holds.

Now by Lemma 1 and Corollary 2 of Lemma 1 we obtain the solvability of both equations (3.8),(3.7).

**Proposition 2.** Let f and g satisfy condition B1)-B3) and  $\xi \in L^{\infty}$ . Then BSDE (2.1) admits a solution  $(Y, Z \cdot M + N) \in S^{\infty} \times BMO$ .

*Proof.* An arbitrary  $\xi \in L^{\infty}(R)$  can be represented as sum  $\xi = \sum_{i=1}^{m} \xi_i$  with  $||\xi_i||_{\infty} \le \frac{1}{32\beta} \exp(-2||\int_0^{\cdot} r_s dK_s||_{\infty})$ . Denote by  $(Y^j, Z^j, N^j), \ j=1,...,m$  the solution of

$$dY_{t}^{j} = (f(t, Y_{t}^{0} + \dots + Y_{t}^{j-1}, \sigma_{t}^{*}(Z_{t}^{0} + \dots + Z_{t}^{j-1}))$$

$$-f(t, Y_{t}^{0} + \dots + Y_{t}^{j}, \sigma_{t}^{*}(Z_{t}^{0} + \dots + Z_{t}^{j}))dK_{t}$$

$$-d(\langle N^{j} \rangle_{t} + 2\langle N^{j}, N^{0} + \dots + N^{j-1} \rangle_{t})g_{t} + Z_{t}^{j*}dM_{t} + dN_{t}^{j},$$

$$Y_{T}^{j} = \xi^{j}$$

$$Y^{0} = 0, \quad Z^{0} = 0 \quad N^{0} = 0.$$

$$(3.9)$$

By Corollary 1 we get

$$||Y^j||_{\infty}^2 + ||Z^j \cdot M^j + N^j||_{\mathrm{BMO}(\mathrm{P}^j)}^2 \le \frac{1}{128\beta^2},$$

where  $dP^j = \mathcal{E}_T(\int_0^{\cdot} f_z(s, Y_s^0 + ... Y_s^{j-1}, \sigma_s^*(Z_s^0 + ... + Z_s^{j-1}))\sigma_s^{-1}dM_s)dP$ , and  $M^j = M - \langle f_z(\cdot, Y^0 + ... + Y^{j-1}, \sigma^*(Z^0 + ... + Z^{j-1}))\sigma^{-1} \cdot M, M \rangle$ .

Using Lemma 2 we get the existence of a solution for BSDE

$$d\bar{Y}_t = (f(t,0,0) - f(t,\bar{Y}_t,\sigma_t^*Z_t))dK_t - d\langle N \rangle_t g_t + Z_t^*dM_t + dN_t,$$
$$\bar{Y}_T = \xi.$$

Since  $\int_0^T f(t,0,0)dK_t$  is bounded we can apply the above argument with f replaced by  $\bar{f}(t,y,z) = f(t,y-\int_0^t f(s,0,0)dK_s,z)$  to get the existence of solution

$$d\bar{Y}_{t} = (f(t,0,0) - f(t,\bar{Y}_{t} - \int_{0}^{t} f(s,0,0)dK_{s}, \sigma_{t}^{*}Z_{t}))dK_{t} - d\langle N \rangle_{t}g_{t} + Z_{t}^{*}dM_{t} + dN_{t},$$
$$\bar{Y}_{T} = \xi + \int_{0}^{T} f(s,0,0)dK_{s}.$$

Obviously  $Y_t = \bar{Y}_t - \int_0^t f(s, 0, 0) dK_s$  is a solution of BSDE (2.1),(2.2).

#### 4 A comparison theorem for BSDEs

Let us consider BSDE (2.1),(2.2) in the case d=1.

**Lemma 3.** Let  $\xi \in L_{\infty}$  and assume that there are positive constants C(f), C(g), increasing function  $\lambda : R^+ \to R^+$ , bounded on all bounded subsets and a predictable process  $k \in H^2(R, 1)$  such that

$$|f(t,y,z)| \le k_t^2 \lambda(|y|) + C(f)z^2,$$
 (4.1)

$$|g(t)| \le C(g). \tag{4.2}$$

Then the martingale part of any bounded solution of (2.1),(2.2) belongs to the space BMO(P).

*Proof.* Let Y be a solution of (2.1),(2.2) and there is a constant C>0 such that

$$|Y_t| \leq C$$
 a.s for all  $t$ .

Applying the Itô formula for  $\exp\{\beta Y_T\} - \exp\{\beta Y_\tau\}$  and using the boundary condition  $Y_T = \xi$  we have

$$\frac{\beta^2}{2} \int_{\tau}^{T} e^{\beta Y_s} Z_s^* d\langle M \rangle_s Z_s + \frac{\beta^2}{2} \int_{\tau}^{T} e^{\beta Y_s} d\langle N \rangle_s$$

$$-\beta \int_{\tau}^{T} e^{\beta Y_s} f(s, Y_s, Z_s) dK_s - \beta \int_{\tau}^{T} e^{\beta Y_s} g(s) d\langle N \rangle_s$$

$$+\beta \int_{\tau}^{T} e^{\beta Y_s} Z_s^* dM_s + \beta \int_{\tau}^{T} e^{\beta Y_s} dN_s = e^{\beta \xi} - e^{\beta Y_{\tau}} \le e^{\beta C}, \tag{4.3}$$

where  $\beta$  is a constant yet to be determined.

If  $Z \cdot M$  and N are square integrable martingales taking conditional expectations in (4.3) we obtain

$$\frac{\beta^{2}}{2}E\left(\int_{\tau}^{T}e^{\beta Y_{s}}Z_{s}^{*}d\langle M\rangle_{s}Z_{s}|F_{\tau}\right) + \frac{\beta^{2}}{2}E\left(\int_{\tau}^{T}e^{\beta Y_{s}}d\langle N\rangle_{s}|F_{\tau}\right)$$

$$\leq e^{\beta C} + \beta E\left(\int_{\tau}^{T}e^{\beta Y_{s}}|f(s,Y_{s},Z_{s})|dK_{s}|F_{\tau}\right) + \beta E\left(\int_{\tau}^{T}e^{\beta Y_{s}}|g(s)|d\langle N\rangle_{s}|F_{\tau}\right)$$

Now if we use the estimates (4.1),(4.2) we get

$$\frac{\beta^{2}}{2}E\left(\int_{\tau}^{T}e^{\beta Y_{s}}Z_{s}^{*}d\langle M\rangle_{s}Z_{s}|F_{\tau}\right) + \frac{\beta^{2}}{2}E\left(\int_{\tau}^{T}e^{\beta Y_{s}}d\langle N\rangle_{s}|F_{\tau}\right)$$

$$\leq e^{\beta C} + \beta\lambda(C)E\left(\int_{\tau}^{T}e^{\beta Y_{s}}k_{s}^{2}dK_{s}|F_{\tau}\right)$$

$$+\beta C(f)E\left(\int_{\tau}^{T}e^{\beta Y_{s}}|\sigma_{s}^{*}Z_{s}|^{2}dK_{s}|F_{\tau}\right) + \beta E\left(\int_{\tau}^{T}e^{\beta Y_{s}}|g(s)|d\langle N\rangle_{s}|F_{\tau}\right)$$

$$\leq e^{\beta C} + \beta\lambda(C)E\left(\int_{\tau}^{T}e^{\beta Y_{s}}k_{s}^{2}dK_{s}|F_{\tau}\right)$$

$$+\beta C(f)E\left(\int_{\tau}^{T}e^{\beta Y_{s}}|Z_{s}^{*}d\langle M\rangle_{s}Z_{s}|^{2}|F_{\tau}\right) + C(g)\beta E\left(\int_{\tau}^{T}e^{\beta Y_{s}}d\langle N\rangle_{s}|F_{\tau}\right).$$

Conditions (4.1) and (4.2) imply that

$$\left(\frac{\beta^{2}}{2} - \beta C(f)\right) E\left(\int_{\tau}^{T} e^{\beta Y_{s}} Z_{s}^{*} d\langle M \rangle_{s} Z_{s} | F_{\tau}\right) + \\
+ \left(\frac{\beta^{2}}{2} - \beta C(g)\right) E\left(\int_{\tau}^{T} e^{\beta Y_{s}} d\langle N \rangle_{s} | F_{\tau}\right) \leq \\
\leq e^{\beta C} + \beta \lambda(C) E\left(\int_{\tau}^{T} e^{\beta Y_{s}} k_{s}^{2} dK_{s} | F_{\tau}\right). \tag{4.4}$$

Taking  $\beta = 4\overline{C}$ , where  $\overline{C} = max(C(f), C(g))$ , from (4.4) we have

$$4\overline{C}^{2}[E(\int_{\tau}^{T} e^{\beta Y_{s}} Z_{s}^{*} d\langle M \rangle_{s} Z_{s} | F_{\tau}) + E(\int_{\tau}^{T} e^{\beta Y_{s}} d\langle N \rangle_{s} | F_{\tau})] \leq$$

$$< e^{4C\overline{C}} (4\overline{C}\lambda(C) ||k||_{H} + 1).$$

Since  $Y \geq -C$ , from the latter inequality we finally obtain the estimate

$$E(\langle Z \cdot M \rangle_{\tau T} | F_{\tau}) + E(\langle N \rangle_{\tau T} | F_{\tau}) \le$$

$$\le \frac{e^{8C\overline{C}} [4\overline{C}\lambda(C)||k||_{H} + 1]}{4\overline{C}^{2}}$$
(4.5)

for any stopping time  $\tau$ , hence  $Z \cdot M, N \in BMO$ .

For general  $Z \cdot M$  and N we stop at  $\tau_n$  and derive (4.5) with T replaced  $\tau_n$ . Letting  $n \to \infty$  then completes the proof.

Further we use some notations. Let  $(Y, Z), (\widetilde{Y}, \widetilde{Z})$  be two pairs of processes and  $(f, g, \xi), (\widetilde{f}, \widetilde{g}, \widetilde{\xi})$  two triples of generators. Then we denote:

$$\begin{split} \delta f &= f - \tilde{f}, \ \delta g = g - \tilde{g}, \ \delta \xi = \xi - \tilde{\xi}, \\ \partial_y f(t, Y_t, \widetilde{Y}_t, Z_t) &\equiv \partial f_y(t) = \frac{f(t, Y_t, Z_t) - f(t, \widetilde{Y}_t, Z_t)}{Y_t - \widetilde{Y}_t} \\ &\text{for all } \ j = 1, ..., n, \ \partial_j f(t, \widetilde{Y}_t, Z_t, \widetilde{Z}_t) \equiv \partial_j f(t) \\ &= \frac{f(t, \widetilde{Y}_t, Z_t^1, ..., Z_t^{j-1}, Z_t^j, \widetilde{Z}_t^{j+1}, ..., \widetilde{Z}_t^n) - f(t, \widetilde{Y}_t, Z_t^1, ..., Z_t^{j-1}, \widetilde{Z}_t^j, \widetilde{Z}_t^{j+1}, ..., \widetilde{Z}_t^n)}{Z_t^j - \widetilde{Z}_t^j}, \\ &\nabla f(t) = (\partial_1 f(t), ..., \partial_n f(t))^* \end{split}$$

Thus we have

$$f(t, Y_t, Z_t) - f(t, \widetilde{Y}_t, \widetilde{Z}_t) = \partial_y f(t) \delta Y_t + \nabla f(t)^* \delta Z_t.$$
(4.6)

**Theorem 2.** Let Y and  $\widetilde{Y}$  be the bounded solutions of SBE (2.1) with generators  $(f, g, \xi)$  and  $(\widetilde{f}, \widetilde{g}, \widetilde{\xi})$  respectively, satisfying the conditions of Lemma 3.

If  $\xi \geq \tilde{\xi}$  (a.s),  $f(t,y,z) \geq \tilde{f}(t,y,z)$  ( $\mu^{K}$ -a.e.),  $g(t) \geq \tilde{g}(t)$  ( $\mu^{\langle N \rangle}$ -a.e.) and f (or  $\tilde{f}$ ) satisfies the following Lipschitz condition:

L1) for any Y, Y, Z

$$\frac{f(t, Y_t, Z_t) - f(t, \widetilde{Y}_t, Z_t)}{Y_t - \widetilde{Y}_t} \in S^{\infty},$$

L2) for any  $Z,\widetilde{Z}\in H^2$  and any bounded process Y

$$(\sigma_t \sigma_t^*)^{-1} \nabla f(t, Y_t, Z_t, \widetilde{Z}_t) \in H^2(\mathbb{R}^n, \sigma),$$

then  $Y_t \geq \widetilde{Y}_t$  a.s. for all  $t \in [0, T]$ .

*Proof.* Taking the difference of the equations (2.1), (2.2) with generators  $(f, g, \xi)$  and  $(\widetilde{f}, \widetilde{g}, \widetilde{\xi})$  respectively, we have

$$Y_{t} - \widetilde{Y}_{t} = Y_{0} - \widetilde{Y}_{0}$$

$$- \int_{0}^{t} [f(s, Y_{s}, Z_{s}) - f(s, \widetilde{Y}_{s}, \widetilde{Z}_{s})] dK_{s}$$

$$- \int_{0}^{t} [f(s, \widetilde{Y}_{s}, \widetilde{Z}_{s}) - \widetilde{f}(s, \widetilde{Y}_{s}, \widetilde{Z}_{s})] dK_{s} - \int_{0}^{t} [g(s) - \widetilde{g}(s)] d\langle N \rangle_{s}$$

$$- \int_{0}^{t} \widetilde{g}(s) d(\langle N \rangle_{s} - \langle \widetilde{N} \rangle_{s}) + \int_{0}^{t} (Z_{s} - \widetilde{Z}_{s}) dM_{s} + N_{t} - \widetilde{N}_{t}.$$

$$(4.7)$$

Let us define the measure Q by  $dQ = \mathcal{E}_T(\Lambda)dP$ , where

$$\Lambda_t = \int_0^t \nabla f(s)^* (\sigma_s \sigma_s^*)^{-1} dM_s + \int_0^t \widetilde{g}(s) d(N_s + \widetilde{N}_s).$$

By Lemma 3  $Z, \widetilde{Z} \in H^2$  and  $N, \widetilde{N}$  are BMO- martingales. Therefore Condition L1), L2) and (4.2) imply that  $\Lambda \in BMO$  and hence Q is a probability measure equivalent to P. Denote by  $\Lambda$  the martingale part of  $\delta Y = Y - \widetilde{Y}$ , i.e.,

$$\bar{\Lambda} = (Z - \widetilde{Z}) \cdot M + N - \widetilde{N}.$$

Therefore, by Girsanov's Theorem and by (4.6) the process

$$\delta Y_t + \int_0^t (\partial_y f(s) \delta Y_s + \nabla f(s)^* \delta Z_s) dK_s$$

$$+ \int_0^t \delta f(s, \widetilde{Y}_s, \widetilde{Z}_s) dK_s + \int_0^t \delta g(s) d\langle N \rangle_s$$

$$= \delta Y_t + \int_0^t (\partial_y f(s) \delta Y_s + \delta f(s, \widetilde{Y}_s, \widetilde{Z}_s)) dK_s$$

$$+ \int_0^t \nabla f(s)^* (\sigma_s \sigma_s^*)^{-1} d\langle M \rangle_s \delta Z_s + \int_0^t \delta g(s) d\langle N \rangle_s$$

$$= -\int_0^t \widetilde{g}(s) d(\langle N \rangle_s - \langle \widetilde{N} \rangle_s) + \int_0^t (Z_s - \widetilde{Z}_s) dM_s + N_t - \widetilde{N}_t$$

$$= \overline{\Lambda}_t - \langle \Lambda, \overline{\Lambda} \rangle_t,$$

is a local martingale under Q. Moreover, since by Lemma 3  $\bar{N} \in BMO$ , Proposition 11 of [6] implies that

$$\bar{\Lambda}_t - \langle \Lambda, \bar{\Lambda} \rangle_t \in BMO(Q).$$

Thus, using the martingale property and the boundary conditions  $Y_T = \xi, \widetilde{Y}_T = \widetilde{\xi}$  we have

$$Y_t - \widetilde{Y}_t =$$

$$= E^{Q} \left( e^{\int_{t}^{T} \partial_{y} f_{s} dK_{s}} (\xi - \tilde{\xi}) \right.$$

$$+ \int_{t}^{T} e^{\int_{t}^{s} \partial_{y} f_{u} dK_{u}} \left( f(s, \widetilde{Y}_{s}, \widetilde{Z}_{s}) - \tilde{f}(s, \widetilde{Y}_{s}, \widetilde{Z}_{s}) \right) dK_{s} | F_{t} \right)$$

$$+ E^{Q} \left( \int_{t}^{T} e^{\int_{t}^{s} \partial_{y} f_{u} dK_{u}} (g(s) - \widetilde{g}(s)) d\langle N \rangle_{s} | F_{t} \right),$$

which implies that  $Y_t \geq \widetilde{Y}_t$  a.s. for all  $t \in [0, T]$ .

Corollary. Let condition A) be satisfied. Then if the solution of (2.1),(2.2) exists it is unique.

The proof of **Theorem 1** follows now from the last corollary and Proposition 2. **Remark.** Condition L1),L2) is satisfied if there is constant C > 0 such that

$$|f(t,y,z) - f(t,\tilde{y},\tilde{z})| \le C|y - \tilde{y}| + C|z - \tilde{z}|(|z| + |\tilde{z}|)$$

and  $tr(\sigma_t \sigma_t^*)^{-1} \leq C$  for all  $y, \tilde{y} \in R$ ,  $z, \tilde{z} \in R^n$   $t \in [0, T]$ . Conditions L1),L2) are also fulfilled if f(t, y, z) satisfies the global Lipschitz condition and  $M \in BMO$ .

#### Acknowledgements

I am thankful to M. Mania for useful discussions and remarks.

I also would like to thank the referee for valuable remarks and suggestions which lead to many improvements in the original version of the paper .

# References

- [1] Bismut J. M. (1973) Conjugate convex functions in optimal stochastic control, *J. Math. Anal. Appl.* **44**, 384–404.
- [2] Bobrovnytska O. and Schweizer M. (2004) Mean-Variance Hedging and Stochastic Control: Beyond the Brownian Setting, IEEE Transactions on Automatic Control 49, 396-408.
- [3] Buckdahn R., (1993) Backward stochastic differential equation driven by Martingale, Preprint.
- [4] Chitashvili R., (1983) Martingale ideology in the theory of controlled stochastic processes, *Lect. Notes in Math.*, Springer, Berlin, N. 1021, pp. 73-92.
- [5] Dellacherie C. and Meyer P. A., (1980) Probabilités et potentiel. Chapitres V a VIII. Théorie des martingales. Actualités Scientifiques et Industrielles Hermann, Paris.
- [6] Doleans-Dade K. and Meyer P. A., (1979) Inégalités de normes avec poinds, Séminaire de Probabilités XIII, Lect. Notes in Math., Springer, Berlin, N. 721, pp. 204-215.

- [7] El Karoui N., Huang S.J., (1997) A general result of existence and uniqueness of backward stochastic differential equations. Pitman Res. Notes Math. Ser.,364, Longman, Harlow, 27-36.
- [8] Hu Y., Imkeller P. and Müller M., (2005), Utility maximization in incomplete markets. Ann. Appl. Probab. 15, no. 3, 1691-1712.
- [9] Jacod J., (1979) Calcule Stochastique et problèmes des martingales. Lecture Notes in Math., Springer, Berlin, N.714.
- [10] Kazamaki N., (1994) Continuous exponential martingales and BMO, Lecture Notes in Math., Springer, Berlin, N. 1579.
- [11] Kobylanski M., (2000) Backward stochastic differential equation and partial differential equations with quadratic growth, The Annals of Probability, vol. 28, N2, 558-602.
- [12] Lepeltier J.P. and San Martin J., (1998) Existence for BSDE with superlinear-quadratic coefficient, Stoch. Stoch. Rep. 63, 227-240.
- [13] Liptser R.Sh. and Shiryaev A.N., (1986) Martingale theory, Nauka, Moscow.
- [14] Mania M. and Schweizer M., (2005) Dynamic exponential utility indifference valuation, Ann. Appl. Probab. 15, no. 3, 2113–2143.
- [15] Mania M. and Tevzadze R., (2000) A Semimartingale Bellman equation and the variance-optimal martingale measure, *Georgian Math. J.* **7** 4, 765–792.
- [16] Mania M. and R. Tevzadze R. (2003) A Unified Characterization of q-optimal and minimal entropy martingale measures by Semimartingale Backward Equations, Georgian Math. J. vol.10, No 2, 289-310.
- [17] Mania M. and Tevzadze R. (2003) Backward Stochastic PDE and Imperfect Hedging, International Journal of Theoretical and Applied Finance, vol.6, 7,663-692.
- [18] Mania M. Santacroce M. and Tevzadze R. (2003) A Semimartingale BSDE related to the minimal entropy martingale measure, Finance and Stochastics, vol.7, No 3, 385-402.
- [19] Mania M. and Tevzadze R. (2003) A Semimartingale Bellman equation and the variance-optimal martingale measure under general information flow, SIAM Journal on Control and Optimization, Vol. 42, N5, 1703-1726.
- [20] Mania M. and Tevzadze R. (2005) An exponential martingale equation, From Stochastic Calculus to Mathematical Finance, The Shiryaev Festschrift, Springer, 507-516.
- [21] Morlais M.A., Quadratic Backward Stochastic Differential Equations (BSDEs) Driven by a Continuous Martingale and Application to the Utility Maximization Problem, http://hal.ccsd.cnrs.fr/ccsd-00020254/

[22] Pardoux E. and Peng S.G. (1990) Adapted solution of a backward stochastic differential equation, Systems Control Lett. 14 ,55–61.