

Explicit generators for (conjectural) mixed motives (in Voevodsky's DM_{gm}^{eff}). The Kunneth decomposition of pure (numerical) motives

M.V. Bondarko

September 25, 2018

Introduction

The current version of this text is just an announcement of results. It includes the main ideas that are quite sufficient to prove the results announced; yet complete proofs will be written down later.

We recall that constructing the category of mixed motives MM is probably the most challenging problem of the modern algebraic geometry. Note that MM should be a full subcategory of Voevodsky's DM_{gm}^{eff} or $DM_{gm}^{eff}\mathbb{Q}$. Yet Voevodsky's motives that are currently assumed to be mixed do not generate DM_{gm}^{eff} ; they do not generate even motives of dimension ≤ 2 (to the knowledge of the author).

In this note we describe very explicitly a rich family of mixed motives that surely generates $DM_{gm}^{eff}\mathbb{Q}$ (as a triangulated category). They "should be" mixed since they have only one non-zero Betti cohomology group. Our method also allows to define a family of direct summands of the numerical motif of any smooth projective variety P . Modulo certain standard conjectures, this construction yields the Kunneth decomposition of the diagonal of P .

From the "motivic" point of view the main ideas of this paper are: repetitive hyperplane sections of a smooth affine variety give a resolution of its Voevodsky's motif by mixed motives; applying this fact to a Jouanolou's "replacement" of a smooth projective variety P one obtains the Kunneth decomposition of the diagonal of the numerical motif of P .

It seems that our ideas are related to the motivic ideas of M. Nori. Yet the results described in §2 seem to be completely new.

In this version we will assume that our base field k is fixed and canonically embedded in the field of complex numbers. We will denote by H_B the (rational) Betty cohomology of a variety or a motif. We will only consider motives and cohomology with rational coefficients.

We will use some definitions and notation of [7]. $DM_{gm}^{eff}\mathbb{Q}$ will denote the idempotent completion of DM_{gm}^{eff} with morphisms tensored by \mathbb{Q} . Similarly, $Chow$ ($Chow^{eff}$) will denote the category of rational (effective) Chow motives; Mot_{hom} will be the category of rational effective homological motives; Mot_{num} will be the category of effective rational numerical motives.

We will also mention the (conjectural) abelian category of (effective) mixed motives $MM \subset DM_{gm}^{eff}\mathbb{Q}$; so the (conjectural) mixed motivic cohomology functors will be covariant. $H_M^i(X) \in ObjMM$ will denote the i -th cohomology of $X \in ObjDM_{gm}^{eff}\mathbb{Q}$ with respect to the (conjectural!) mixed motivic structure. Often X will be the rational Voevodsky's motif of a (smooth) variety V (denoted by $M_{gm}(V)$).

1 A method for constructing mixed motives (in DM_{gm}^{eff})

The starting point of this work were certain results of [1] and [2].

Let U be a smooth Zarisky open subvariety of a smooth projective variety Y of dimension m and let H be its hyperplane section; let $V = U \cap H$ be non-empty.

We recall Theorem 6.1.1 of [1]. It states that the induced map $H^i(V) \rightarrow H^i(U)$ is bijective for $i < m - 1$ and is injective for $i = m - 1$. To the author's knowledge this means that (U, V) is a *good pair* in the sense of M. Nori.

Proposition 1.1. *Let U be affine. We define $M \in ObjDM_{gm}^{eff}\mathbb{Q}$ as a cone of $M_{gm}(V) \rightarrow M_{gm}(U)$ i.e. $M_{gm}(V) \rightarrow M_{gm}(U) \rightarrow M$ is a distinguished triangle. Then the only non-zero Betty cohomology group of M is H_B^m .*

Proof. For any i we have a long exact sequence

$$\dots \rightarrow H_B^{i-1}(U) \rightarrow H_B^{i-1}(V) \rightarrow H_B^i(M) \rightarrow H_B^i(U) \rightarrow H_B^i(V) \rightarrow \dots$$

It remains to recall that $H_B^j(X) = 0$ for $j > m$ and apply Theorem 6.1.1 of [1].

□

Remark 1.2. 1. It follows that $M[-m]$ a nice candidate for an (abelian) mixed motif. We will call motives of this type "our" motives. For example, the becomes mixed in Hanamura's construction (see §3 of [4]; note

that Hanamura's construction is purely conjectural!). Recall that Hanamura's motivic category is anti-equivalent to (Voevodsky's) $DM_{gm}\mathbb{Q}$; see the proof in §4 of [3]. If $M[-m]$ in MM should be an extension of $H_M^m(X)$ by $\text{Ker} : H_M^{m-1}(Y) \rightarrow H_M^{m-1}(X)$. Note that the corresponding fact holds in the category of (rational) mixed Hodge structures i.e. $H^i(M)_{Hodge}$ is an extension of $\text{Coker} : H_{Hodge}^{m-1}(X) \rightarrow H_{Hodge}^{m-1}(Y)$ by $H_{Hodge}^m(X)$.

2. One could easily construct a large variety of "1-extensions" of our motives. This leads to the following construction of mixed motives that is (at the moment) the most general among those known to the author.

We take a complex $X_i \in K^b(\text{SmCor} \otimes \mathbb{Q})$ with $\dim(X_i) = i$; X_i are affine. Let $Y_i \subset X_i$ be a compatible system of hyperplane sections (as in Proposition 1.1). Then the cone of $(Y_i) \rightarrow (X_i)$ (considered as a motif coming from $K^b(\text{SmCor} \otimes \mathbb{Q})$) cannot have any cohomology out of dimension 0. This statement follows from Proposition 1.1 by easy induction. In particular, in this way one could describe tensor products of "our" motives in $DM_{gm}^{eff}\mathbb{Q}$. (Recall that the tensor product of motives is compatible with the tensor product of varieties.)

It also seems that the "twisted dual" of M i.e a cone M' of the natural morphism $M_{gm}^c(U) \rightarrow M_{gm}^c(V)(1)[2]$ can be obtained from $K^b(\text{SmCor} \otimes \mathbb{Q})$ using this construction (see Theorem 4.3.7 of [7]). One could also easily check that M' has only one non-zero cohomology group using the Poincare duality and Proposition 1.1.

3. It is easily seen that "our" motives generate the whole $DM_{gm}^{eff}\mathbb{Q}$ as a triangulated category (after idempotent completion). Indeed, a repetitive application of Proposition 1.1 (to U , then to V , etc.) immediately yields that the triangulated category D generated by "our" motives contains all motives of smooth affine varieties. This method should correspond to the resolution of $M_{gm}(U)$ by mixed motives.

Next, the Mayer-Vietoris triangle (see §2 of [7]) yields that D contains motives of all smooth varieties. Lastly, recall that $DM_{gm}^{eff}\mathbb{Q}$ is the idempotent completion of a certain localization of $K^b(\text{SmCor} \otimes \mathbb{Q})$.

Moreover, it seems that any object of $DM_{gm}^{eff}\mathbb{Q}$ that comes from $K^b(\text{SmCor} \otimes \mathbb{Q})$ could be presented as a certain "complex" of "our" motives (i.e. it becomes equivalent in $DM_{gm}^{eff}\mathbb{Q}$ to such a complex constructed inside of $K^b(\text{SmCor} \otimes \mathbb{Q})$).

4. One could try to define the mixed motivic t -structure using "our" motives. Then it seems that the main problem is to prove that there are no morphisms of negative degrees between "our" motives (in $DM_{gm}^{eff}\mathbb{Q}$). Unfortunately, one probably cannot overcome this difficulty with assuming certain vanishing conjectures (as it was done in [4]). Still there is a hope to deduce everything from the (conjectural) conservativity of the Betti realization of

motives. Note that (by the conservativity of the weight complex functor, see Proposition 6.1.3 of [3], and by the existence of the weight spectral sequence for realizations of motives, see (14) of §7.3 *ibid.*) it suffices to check this conservativity on $K^b(\mathit{Chow}^{eff})$.

It seems that Proposition 1.1 should be connected with the ioga of relative motives (mainly of motives over $\mathbb{Q}(i)$).

2 A candidate for the Kunneth decomposition of pure motives

Unfortunately, it is not clear how to describe all $H_M^i(U)$ for all i using our method. Yet this seems to be easy if we restrict ourselves to numerical motives (of smooth projective varieties).

More precisely, we apply Theorem 2.1 of [2] for smooth projective $X = Y$ (in the notation of *loc.cit.*). Essentially, this is (more or less) equivalent to the composition of a repetitive application of Proposition 1.1 with Jouanolou's trick and Theorem 6.1.1 of [1]. In order to clarify the connection of the reasoning below with Proposition 1.1 we recall that for any smooth quasi-projective X there exists an affine line bundle (of some dimension $l \geq 0$) X'/X such that X' is affine over k (Jouanolou's trick). We will call X' a *Jouanolou's replacement* for X . Since X' is Zarisky locally isomorphic to $X \times \mathbb{A}^l$, we obtain that $M_{gm}(X') \cong M_{gm}(X)$ in $DM_{gm}^{eff}\mathbb{Q}$; cf. Proposition 3.5.1 of [7].

Now, Theorem 2.1 of [2] applied to smooth projective $X = Y$ (in the notation of *loc.cit.*) states exactly that the canonical filtration for the cohomology of X can be described by kernels of $H_B(X) \rightarrow H_B(X_p)$ for some morphism of smooth varieties $X_p \rightarrow X$ (here X_p are obtained by taking smooth hyperplane sections of any Jouanolou's replacement X' of X). Now one can look at the numerical motif of X . X_p is (probably) not projective; yet it corresponds to a complex of Chow motives (see §6 of [3] for the consideration of *weight complex* of motives). Now consider the cokernel of the maps induced by $X_p \rightarrow X$ on the (covariant) numerical motif of X . More precisely, by part 1 of Theorem 6.2.1 of [3] the weight complex $t(M_{gm}(X_p))$ is a complex of (effective) Chow motives concentrated in non-negative degrees. We consider the corresponding complex $T_p = t_{num\mathbb{Q}}(M_{gm}(X_p)) = T_0 \rightarrow T_1 \rightarrow \dots \in K^b(\mathit{Mot}_{num})$. Note that X_p could be taken to be a complement of one smooth projective varieties by another one; hence we may assume that $T_i = 0$ for $i > 1$. Since the category of rational (effective) numerical motives is abelian semi-simple (see Theorem 1 of [5]), the cokernel of $H^0(T_p) \rightarrow \mathit{Mot}_{num\mathbb{Q}}(X)$ gives a direct

summand of the motif of X . It seems to be a very reasonable candidate for $\sum_{p \leq i \leq 2n} H_M^i(X)[-i]$ (as a numerical motif with the grading compatible with the theory of mixed motives). Note that X_p are embedded into each other, so one can define each $H_M^i(X)$ separately (unconditionally). We obtain an explicit construction that (conjecturally) yields the Kunnet decomposition of the diagonal of the numerical motif of X . Indeed, the construction obviously gives the the Kunnet decomposition if the numerical equivalence is equivalent to the homological equivalence. Note also that (using easy finite dimension arguments similar to those of [5]) any direct summand of the numerical motif of X could be lifted to a direct summand of its homological motif (for example, one could consider the "minimal possible lift"). Yet it doesn't seem easy to prove that the homological motives obtained give the Kunnet decomposition on this level. Still even without this fact the decomposition constructed could help to describe the (conjectural) Tannakian category of pure motives.

It doesn't seem to be very difficult to prove (certain) functoriality of this construction; in particular, to verify that the construction is canonical. Possibly, to simplify the proof one should modify the construction a little. Probably this leads to the consideration of (a certain) coniveau filtration of the motif of X . Note that to this end we should consider some Jouanolou's replacement X' of X . Possibly it also makes sense to consider a certain "limit" with respect to all Jouanolou's replacements for X ; note the fiber product over X of two Jouanolou's replacements for X is also a replacement for it.

Lastly, we note that the construction could also be applied on the level of Chow motives. It does not necessarily yield a Chow motif since $Chow^{eff}$ is not abelian; yet it yields a factor of the functor represented by $Mot_{Chow}(X)$ in the abelian category $Chow_*^{eff}$ of additive contravariant functors $Chow^{eff} \rightarrow AbGr$ (we have a natural full embedding $Chow^{eff} \rightarrow Chow_*^{eff}$). It doesn't seem difficult to make this construction functorial (using the limit methods described above). Note that then our construction would automatically give canonical functors $F_{Hom}^i : Chow^{eff} \rightarrow Mot_{hom*}$ and $F_{num}^i : Chow^{eff} \rightarrow Mot_{num}$ (since Mot_{num} is abelian!). This approach could also yield (by applying Theorem 2.1 of [2]) that these functors could be constructed using the 'basic' construction (without passing to any limits); the corresponding "Chow fact" seems to be more difficult.

A curious observation: for any X_p all T_i (could be chosen to be) divisible by $\mathbb{Q}(i)$. Hence for the birational motives theory (where $\mathbb{Q}(1)$ is killed) it suffices to consider only the cokernel of $T_0 \rightarrow Mot_{num}(X)$; see [6].

3 Concluding remarks

Certainly, these arguments cannot prove all standard conjectures. Yet they are very explicit; this gives a hope to prove some parts of standard conjectures (or other interesting statements) unconditionally. Also, there should be a considerable impact on interrelations between conjectures. The author's knowledge of this field (of conjectures) is quite poor; so he will be deeply grateful for any ideas in this direction.

References

- [1] Intermediate Jacobians and Hodge Structures of Moduli Spaces, Arapura D., Sastry P. // Proc. Indian Acad. Sci. (Math Sci.), vol. 110, No. 1, 2000, 1–26; see <http://www.ias.ac.in/mathsci/vol110/feb2000/Pm1709.pdf>
- [2] The Leray spectral sequence is motivic, Arapura D.// Inv. Math., vol. 160 (2005), 567–589; see also <http://arxiv.org/abs/math.AG/0301140>.
- [3] Bondarko M.V., 'Weight enhancement' and truncations for Voevodsky motives; filtrations and motivic descent spectral sequence for differential graded realizations, electronic, <http://arxiv.org/abs/math.AG/0601713>.
- [4] Hanamura M. Mixed motives and algebraic cycles, III// Math. Res. Letters 6, 1999, 61–82.
- [5] Jannsen U., Motives, numerical equivalence, and semi-simplicity //Inv. Math., vol. 107 (1992), 447–452.
- [6] Kahn B., Sujatha R., Birational motives, I, preprint, available on the K-theory preprint archive, no 0596, <http://www.math.uiuc.edu/K-theory/0596/>
- [7] Voevodsky V. Triangulated category of motives, in: Voevodsky V., Suslin A., and Friedlander E., Cycles, transfers and motivic homology theories, Annals of Mathematical studies, vol. 143, Princeton University Press, 2000, 188–238, see also <http://www.math.uiuc.edu/K-theory/0074/>