

WALKS ON GROUPS, COUNTING REDUCIBLE MATRICES, POLYNOMIALS, AND SURFACE AND FREE GROUP AUTOMORPHISMS

IGOR RIVIN

ABSTRACT. We prove sharp limit theorems on random walks on graphs with values in finite groups. We then apply these results (together with some elementary algebraic geometry, number theory, and representation theory) to finite quotients of lattices in semisimple Lie groups (specifically $SL(n, \mathbb{Z})$ and $Sp(2n, \mathbb{Z})$) to show that a “random” element in one of these lattices has irreducible characteristic polynomials (over \mathbb{Z}). The term “random” can be defined in at least two ways (in terms of height and also in terms of word length in terms of a generating set) – we show the result using both definitions.

We use the above results to show that a random (in terms of word length) element of the mapping class group of a surface is pseudo-Anosov, and that a random free group automorphism is irreducible with irreducible powers (or *strongly irreducible*¹).

INTRODUCTION

This paper was inspired by the following question, first brought to the author’s attention by Ilya Kapovich:

Is it true that a random element of the mapping class group of a surface is pseudo-Anosov?

The definition of *random* in the question above is not explicitly given, but a reasonable way to define is to fix a generating set of the

Date: November 26, 2024.

Key words and phrases. irreducible, reducible, matrices, polynomials, surfaces, automorphisms.

I would like to thank Ilya Kapovich for asking the questions on the irreducibility of automorphisms of surfaces and free groups, and for suggesting that these questions may be fruitfully attacked by studying the action on homology. I would also like to thank Nick Katz for making him aware of Nick Chavdarov’s work, Sinai Robins for making him aware of Morris Newman’s classic book, and Akshay Venkatesh, Peter Sarnak, and Alex Lubotzky for enlightening conversations. The author would also like to thank Benson Farb, Ilya Kapovich, and Lee Mosher for comments on previous versions of this note.

mapping class group, and look at all the words of bounded length.¹ Kapovich had suggested that a reasonable way to attach this question was to study the action of the mapping class group \mathcal{M}_g on homology, which gives a symplectic representation of \mathcal{M}_g . Results of Casson and Bleiler (see [4]) then give a set of sufficient conditions for a surface automorphism to be pseudo-Anosov in terms of its image under that representation.

This paper, then, is the embodiment of the program as described above. Along the way, we show a number of results not explicitly related to low dimensional topology or geometric group theory.

Here is a summary of the results:

In my old preprint [19] I state, and sketch the proof of a general equidistribution theorem for products of elements of finite (and, more generally, compact) groups along long paths in finite graphs. In the current paper I give complete arguments. In the follow up-paper [18] much sharper results are given (with speed of convergence bounds). In particular, the convergence bounds are *uniform* for families of finite quotients of a group satisfying Lubotzky's property τ (or the stronger Kazhdan's property T).

We then show that the set of polynomials reducible over \mathbb{Z} with constant coefficient 1 is an algebraic subvariety of the set of polynomials, and using elementary theory of algebraic groups show that the set of matrices in $SL(n, R)$ (where R is the coefficient ring) with reducible characteristic polynomials is a finite union of Zarisky-closed sets, so in particular, when $R = \mathbb{Z}_p$, the proportion of matrices in $SL(n, R)$ with reducible characteristic polynomials decreases as $O(1/p)$, with constants independent of p (Section 5). Using these methods together with Weil's estimate for the number of points on a curve over \mathbb{F}_p we show that the probability that the characteristic polynomial of a matrix in $SL(n, \mathbb{Z}_p)$ has the full symmetric group S_n as Galois group goes is bounded (uniformly) away from 0. Along the way, we show that the probability that a polynomial with fixed constant term 1 has a prescribed splitting type T modulo p is the same as the probability that a *random* polynomial has splitting type T (for large p). (Section 10).

We quote results of Borel (proofs can be found in Nick Chavdarov's thesis [5]) which show that the probability that a matrix in $Sp(2n, \mathbb{Z}_p)$

¹Another way is to look at a combinatorial ball of radius N around identity; it turns out that our results, together with the results announced recently by Ursula Hammenstadt answer both questions.

has reducible characteristic polynomial is bounded away from 1 (uniformly in p .) (Section 6)

Combining these results with the results on walks on graphs, together with elementary local-to-global estimates, we show that for a given *undirected* graph G whose vertices are decorated with a symmetric generating set of Γ (where Γ is either $SL(n, \mathbb{Z})$ or $Sp(2n, \mathbb{Z})$), the probability that the product of generators along a random long walk of length N gives a matrix with reducible characteristic polynomial (or, in the case of $SL(n, \mathbb{Z})$, a polynomial with non-generic Galois group) goes to 0 as N goes to infinity.

From a number-theoretic standpoint, using the length of the representing word in generators is a less natural way to define the size of an integral matrix than a(ny) matrix norm. It turns out that with that definition of size the result follows by combining our results on finite matrix groups with the work of [7], and effective bounds can be obtained using the uniformity results of [16] (see Theorem 5.2).

Finally, we apply the above-mentioned results to automorphisms of surfaces (Section 8) and free group automorphisms (Section 9). For surface automorphisms we use the observations of Casson and Bleiler ([4]) to show that for an arbitrary generating set, most products of up to N generators are pseudo-Anosov, and if the generating set happens to be symmetric, then the fraction of the non-pseudo-Anosov products goes to zero exponentially fast with N . It can be argued that it is more natural to consider all elements in a combinatorial ball of radius N . Since our limit theorems for graphs are for undirected graphs, such a result will follow (for certain generating sets) once we know that the mapping class groups are bi-automatic.

Similarly, it is shown that most words in a generating set of the outer automorphism group of a free group F_n are irreducible with irreducible powers, or what Lee Mosher calls strongly irreducible.

1. GENERALITIES ON ALGEBRAIC GEOMETRY AND ALGEBRAIC GROUPS

First, recall the following:

Definition 1.1 ([10][p. 23]). *Let $V \subseteq k^n$, and $W \subseteq k^m$ be non-empty algebraic sets. We say that $\phi : V \rightarrow W$ is a regular map, if there exist $f_1, \dots, f_m \in k[X_1, \dots, X_n]$, (where X_1, \dots, X_n are indeterminates) such that*

$$\phi(x) = (f_1(x), \dots, f_m(x))$$

for all $x \in V$.

Example 1.2. Let $V = \mathrm{SL}(n, k) \subset k^{n^2}$, and let $W = k^n$. Then the map associating to each matrix the coefficients its characteristic polynomial is a regular map.

Consider the following setup: we have a parametrized set S in k^n (for k an algebraically closed field), that is:

$$\begin{aligned} x_1 &= f_1(s_1, \dots, s_m), \\ x_2 &= f_2(s_1, \dots, s_m), \\ &\vdots \\ x_n &= f_n(s_1, \dots, s_m), \end{aligned}$$

where f_1, \dots, f_n are polynomials in s_1, \dots, s_m . By the ‘‘Implicitization Theorem’’ [6][Chapter 3], the Zariski closure of S is an affine variety, whose dimension is bounded above by m (by considering the dimension of the tangent space, see [10][Theorem 1.4.11]).

1.1. Fibers of dominant morphisms. The following results are standard; the statement is taken from [10][pp. 116,118]. First, let $A[X]$ be the algebra of regular functions on X , and second, for $g \in A[X]$, let $X_g = \{x \in X \mid f(x) \neq 0\}$. Now

Theorem 1.3. *Let $\phi : X \rightarrow Y$ be a dominant morphism of irreducible affine varieties. In particular, $\phi^* : A[Y] \rightarrow A[X]$ is injective, and $d = \dim X - \dim Y \geq 0$. Then we have a factorization*

$$\phi|_{X_{\phi^*(g)}} : X_{\phi^*(g)} \xrightarrow{\bar{\phi}} Y_g \times k^d \xrightarrow{p_1} Y_g,$$

where $\bar{\phi}$ is a finite dominant morphism, and p_1 is the projection on the first factor.

Corollary 1.4. *Let $\alpha : X \rightarrow Y$ be a dominant morphism of irreducible varieties, and put $r = \dim X - \dim Y$. Then, there exists a non-empty open set $U \subseteq Y$, such that $U \subseteq \alpha(X)$, and such that $\dim \alpha^{-1}(y) = r$ for all $y \in U$.*

Remark 1.5. U can be taken to be of the form X_f , for some $f \in A[Y]$.

A dominant morphism α is a morphism such that the preimage of any open dense set in the image variety is dense. In particular, a surjective regular map is dominant.

Example 1.6. Consider $V \subset k^{n^2}$, where $V = \mathrm{SL}(n, k)$. The map χ , which associates to each matrix the coefficients of its characteristic polynomial is a dominant morphism onto k^{n-1} (by the companion matrix construction, it is surjective). By the results (and notation) above,

there is a $g \in A[V]$, such that the fibers of χ restricted to V_g are $n^2 - n$ -dimensional varieties.

Similarly, let $\text{Sp}(2n, k) = W \subset k^{4n^2}$, and let χ' be the map which associates to each matrix m the coefficients of x, \dots, x^n of the characteristic polynomial of m . By Theorem A.1, χ' is a dominant morphism, and by Fact 3.3 and the results of this section, there is an $h \in A[W]$, such that the fibers of χ' restricted to W_h are $2n^2$ -dimensional varieties.

2. COUNTING POINTS ON VARIETIES

Let S be a variety of dimension m over \mathbb{C} (or $\overline{\mathbb{Q}}$). Consider a reduction of S modulo p .

Theorem 2.1 (Lang-Weil, [15]). *The number of F_p points on S grows as $O(p^m)$. The implied constant is uniform (that is, it is a function of the dimension and codimension of the variety only).*

It should be noted that we are using this theorem for the upper bound only, and the upper bound is an easy result (see [10, Lemma 4.1.3]), unlike the full Lang-Weil result which gives both upper and lower bounds. Lang and Weil is deduced from: A. Weil's estimate on the number of \mathbb{F}_p points on a curve defined over \mathbb{F}_p :

Theorem 2.2 (A. Weil, [23]). *Let $f \in \mathbb{F}_p[X, Y]$ be an absolutely irreducible (that is, irreducible in $\overline{\mathbb{F}_p}[X, Y]$) polynomial of degree d . Then if*

$$C = \{(x, y) \in \mathbb{F}_p^2 \mid f(x, y) = 0\},$$

we have the estimate

$$\|C| - p| \leq 2g\sqrt{p} + d^2,$$

where g is the genus of the curve defined by f (which satisfies $g \leq (d - 1)(d - 2)$). This estimate is sharp.

Theorem 2.1, together with the results of Section 1.1 imply:

Theorem 2.3. *With the situation as described in Corollary 1.4, if the ground field is \mathbb{F}_p , if $S \subseteq Y$, then $|\alpha^{-1}(S)| \leq c_1|S|p^r + c_2(p^{\dim X - 1})$ where the constants c_1, c_2 depend only on the dimensions of X, Y .*

3. CLASSICAL GROUPS

In this paper we will be primarily concerned with the special linear group and the symplectic group over various domains. We will need the following facts:

Fact 3.1 (See [2, 10]). *The groups $GL(n, k)$, $SL(n, k)$ and $Sp(n, k)$ over an algebraically closed field k are irreducible as algebraic varieties. This is equivalent to saying that the groups in question are connected.*

Fact 3.2. *The dimension of $SL(n, k)$ equals $n^2 - 1$. The dimension of $Sp(2n, k)$ equals $2n^2 + n$. In both cases, dimension is meant in the sense of algebraic geometry.*

Let $\mathbb{F}_p = \mathbb{Z}/p\mathbb{Z}$.

The following goes back to Dickson:

Fact 3.3. *The order of $SL(n, \mathbb{F}_p)$ equals*

$$p^{(n^2-n)/2}(p^2 - 1)(p^3 - 1) \dots (p^n - 1) = p^{n^2-1} + O(p^{n^2-3}).$$

The order of $GL(n, \mathbb{F}_p)$ equals

$$p^{(n^2-n)/2}(p - 1)(p^2 - 1)(p^3 - 1) \dots (p^n - 1) = p^{n^2} + O(p^{n^2-1}).$$

The order of $Sp(2n, \mathbb{F}_p)$ equals

$$p^{n^2}(p^2 - 1)(p^4 - 1) \dots (p^{2n} - 1) = p^{2n^2+n} + O(p^{2n^2+n-2}).$$

4. APPLICATIONS TO POLYNOMIALS

Let $\mathcal{P}_d(k, \beta)$ be the set of all monic polynomials in one variable of degree d over a field k with constant coefficient β , and let $\mathcal{P}_d^m(k, \beta, \alpha) \subseteq \mathcal{P}_d(k, \beta)$ be the set of those polynomials which have a polynomial factor (over k) of degree m with constant term α . Let us identify $\mathcal{P}_d(k, \beta)$ with the affine space k^{d-1} . Then, we have the following:

Theorem 4.1. *The set $\mathcal{P}_d^m(k, \beta, \alpha)$ is contained in an affine hypersurface of $\mathcal{P}_d(k, \beta)$.*

Proof. Let

$$p(x) = x^d + \sum_{i=0}^{d-1} a_i x^i \in \mathcal{P}.$$

By assumption, $p(x) = q(x)r(x)$. Assume that the degree of $q(x) = m$, while the constant term of $q(x)$ equals α . Writing

$$q(x) = x^m + \sum_{j=1}^{m-1} b_j x^j,$$

and

$$r(x) = x^{d-m} + a_0/\alpha + \sum_{k=1}^{d-m-1} c_k x^k,$$

we see that $\mathcal{P}_d^m(k, \beta, \alpha)$ is a polynomially parametrized hypersurface of $\mathcal{P}_d(\beta, k)$. \square

Corollary 4.2. *The set of all reducible monic polynomials with fixed constant term β is the union*

$$\bigcup_{m=1}^d \mathcal{P}_d^m(k, \beta, \alpha)$$

of affine hypersurfaces of $\mathcal{P}_d(k, \beta) \simeq k^{d-1}$.

Theorem 4.3. *Let $P(d)(\mathbb{F}_p)$ be the set of polynomials of degree d with coefficients in absolute value by N and constant coefficient 1, and let $R_1(d)(\mathbb{F}_p)$ be the set of polynomials reducible over \mathbb{F}_p with some factor having constant term equal to 1 mod p . Then, $R_1(d)$ lies on an algebraic hypersurface \mathbb{C}^{d-1} (where the coordinates are the coefficients), and consequently*

$$\frac{R_1(d)}{P_1(d)} = O\left(\frac{1}{p}\right).$$

Proof. Immediate from Theorem 2.1. \square

Finally, one definition:

Definition 4.4. *We say that a polynomial $p(x) \in P_1(d)$ is reciprocal if $x^d p(1/x) = p(x)$ – in other words, the list of coefficients of p is the same read from left to right as from right to left. Reciprocal polynomials can also be defined as follows: A (monic) polynomial (of even degree $2n$) is reciprocal if it can be written as*

$$\prod_{j=1}^n (x - r_j)(x - 1/r_j) = \prod_{j=1}^n (x^2 - (r_j + r_j^{-1})x + 1).$$

5. APPLICATIONS TO MATRICES

5.1. The special linear group.

Lemma 5.1. *The probability that the characteristic polynomial of a matrix M in $SL(n, \mathbb{F}_p)$ has a factor with constant term 1 is of order $O(1/p)$.*

Proof. This follows immediately from Example 1.6 and Theorems 4.3 and 2.3 \square

Unfortunately, since the number of integral points on $SL(n, \mathbb{Z})$ of height (absolute value) bounded by B grows much slower than B^{n^2-1} the above results *do not* imply the following

Theorem 5.2. *The probability that a matrix in $\mathrm{SL}(n, \mathbb{Z})$ with coefficients bounded by B has reducible characteristic polynomial goes to 0 as B goes to infinity.*

Proof. For the “soft” result as above we use results of Duke Rudnick and Sarnak ([7]) or of Eskin and McMullen ([8]). These imply that the matrices in $\mathrm{SL}(n, \mathbb{Z})$ with coefficients bounded by B are asymptotically equidistributed among the cosets of any subgroup of finite index, in particular the cosets of the principal congruence subgroup $\Gamma_p(n, \mathbb{Z})$. Since reducibility modulo p depends only on the coset modulo Γ_p , Lemma 5.1 immediately replies the result. \square

Remark 5.3. To get an effective result, we need to know that the error terms in equidistribution modulo Γ_p are uniform (do not depend on p). Precisely such a result is shown in the preprint of Amos Nevo and Peter Sarnak [16].

6. RANDOM PRODUCTS OF MATRICES IN THE SYMPLECTIC AND SPECIAL LINEAR GROUPS

In the preceding section we defined the size of a matrix by (in essence) its L^1 norm (any other Banach norm will give the same results). However, it is sometimes more natural to measure size differently: In particular, if we have a generating set $\gamma_1, \dots, \gamma_l$ of our lattice Γ (which might be $\mathrm{SL}(n, \mathbb{Z})$ or $\mathrm{Sp}(2n, \mathbb{Z})$) we might want to measure the size of an element by the length of the (shortest) word in γ_i equal to that element – this is the combinatorial measure of size. The relationship between the size of elements and combinatorial length is not at all clear, so the results in this section are proved quite differently from the results in the preceding section.

We will be using Theorems 11.2 and 11.3.

Remark 6.1. We will be applying Theorem 11.2 to groups $\mathrm{SL}(n, \mathbb{Z}/p\mathbb{Z})$ and $\mathrm{Sp}(2n, \mathbb{Z}/p\mathbb{Z})$. Since those groups have no non-trivial one-dimensional representations, the assumption on ρ in the statement of the theorem is vacuous.

We will also need the following results of Nick Chavdarov and Armand Borel.

Theorem 6.2 (N. Chavdarov, A. Borel [5]). *Let $q > 4$, and let $R_q(n)$ be the set of $2n \times 2n$ symplectic matrices over the field F_q with reducible characteristic polynomials. Then*

$$\frac{|R_q(n)|}{|\mathrm{Sp}(2n, F_q)|} < 1 - \frac{1}{3n}.$$

Theorem 6.3 (N. Chavdarov, A. Borel [5]). *Let $q > 4$, and let $G_q(n)$ be the set of $n \times n$ matrices with determinant $\gamma \neq 0$ over the field F_q with reducible characteristic polynomials. Then*

$$\frac{|G_q(n)|}{|\mathrm{SL}(n, F_q)|} < 1 - \frac{1}{2n}.$$

Theorem 6.3 follows easily from the following result of A. Borel:

Theorem 6.4 (A. Borel). *Let F be a monic polynomial of degree N over $\mathbb{Z}/p\mathbb{Z}$ with nonzero constant term. Then, the number $\#F, p$ of matrices in $\mathrm{GL}(N, p)$ with characteristic polynomial equal to F satisfies*

$$(p - 3)^{N^2 - N} \leq \#(F, p) \leq (p + 3)^{N^2 - N}.$$

Theorem 6.4 will be used in Section 7.

We now have our results:

Theorem 6.5. *Let G and S_N , be as in the statement of Theorem 11.2, but with $\Gamma = \mathrm{Sp}(2n, \mathbb{Z})$, or $\Gamma = \mathrm{SL}(2, \mathbb{Z})$. Then the probability that a matrix in S_N has a reducible characteristic polynomial goes to 0 as N tends to infinity.*

Proof. Let Γ_l be the set of matrices in Γ reduced modulo l – it is known (see [17]) that Γ_l is $\mathrm{SL}(n, l)$ or $\mathrm{Sp}(2n, l)$ (depending on which Γ we took. Let p_1, \dots, p_k be distinct primes, let $K = p_1 \dots p_k$. We know that:

$$\Gamma_K = \Gamma_{p_1} \times \dots \times \Gamma_{p_k}.$$

(see [17] for the proof of the last equality). A generating set of $\mathrm{Sp}(2n, \mathbb{Z})$ projects via reduction modulo K to a generating set of Γ_K (see, again, Newman's book [17]), and also, via reduction mod p_i to generating sets of the $\mathrm{Sp}(2n, p_i)$. By Theorems 11.2, 11.3 and 6.2, the probability that the characteristic polynomial in a random product of $N \gg 1$ generators is reducible modulo all of the p_i is at most equal to $(1 - 3/n)^k$. Since this is an upper bound on the probability of being reducible modulo \mathbb{Z} , the result follows. \square

Remark 6.6. Using Lemma 5.1 instead of Theorem 6.3 for $\mathrm{SL}(n, \mathbb{Z})$ gives a sharper result, as well as a more elementary argument.

Remark 6.7. The proof of Theorem 5.2 translates *mutatis mutandis* into this setting – instead of the principal congruence subgroup Γ_p for a single prime p , we use it for $q = p_1 \times \dots \times p_k$.

An example of a graph G is a graph where every vertex is connected to every vertex (*including itself*), and the set of label is a *symmetric* generating set. In this case, we are just taking random products of generators or their inverses. Another is the graph (studied in [19])

where a vertex labelled by a generator a is connected to every vertex *except* the one labelled by a^{-1} , so that only reduced words in the generators are allowed, and so on.

7. STRONGER IRREDUCIBILITY

We might ask if something stronger than irreducibility of the characteristic polynomial can be shown. The answer is in the affirmative. Indeed, the methods of the preceding sections combined with the results of the Appendix give immediately:

Theorem 7.1. *The probability that a random word of length L in a generating set of $SL(N, \mathbb{Z})$ has characteristic polynomial with Galois group S_N goes to 1 as L goes to infinity.*

Aside from its intrinsic interest, Theorem 7.1 implies the following:

Theorem 7.2. *The probability that a random word w of length L in a generating set of $SL(n, \mathbb{Z})$ and all proper powers w^k have irreducible characteristic polynomials goes to 1 as L goes to infinity.*

Theorem 7.2 will follow easily from Theorem 7.1 together with the following Lemma²:

Lemma 7.3. *Let $M \in SL(n, \mathbb{Z})$ be such that the characteristic polynomial of M^k is reducible for some k . Then the Galois group of the characteristic polynomial of M is imprimitive, or the characteristic polynomial of M is cyclotomic.*

Remark 7.4. For the definition of *imprimitive* see, for example, [24, 11].

Proof. Assume that the characteristic polynomial $\chi(M)$ is irreducible (otherwise the conclusion of the Lemma obviously holds, since the Galois group of $\chi(M)$ is not even transitive). Let the roots of $\chi(M)$ (in the algebraic closure of \mathbb{Q}) be $\alpha_1, \dots, \alpha_n$. The roots of $\chi(M^k)$ are β_1, \dots, β_n , where $\beta_j = \alpha_j^k$. Suppose that $\chi(M^k)$ is reducible, and so there is a factor of $\chi(M^k)$ whose roots are β_1, \dots, β_l , for some $l < n$. Since $\text{Gal}(\chi(M))$ acts transitively on $\alpha_1, \dots, \alpha_n$, it must be true that for every $i \in \{1, \dots, n\}$, $\alpha_i^k = \beta_j$, for some $j \in \{1, \dots, l\}$. Let B_j be those i for which $\alpha_i^k = \beta_j$. This defines a partition of $\{1, \dots, n\}$ into blocks, which is stabilized by the Galois group of $\chi(M)$, and so G is an intransitive subgroup of S_n , unless $l = 1$. In that case, the characteristic polynomial of M^k equals $(x - \beta)^n$, and since $M^k \in SL(n, \mathbb{Z})$ it follows that $\beta = 1$, and all the eigenvalues of M are n -th roots of unity, so that $M^k = 1$. \square

²Compare with [5][Lemma 5.3]

8. THE MAPPING CLASS GROUP

Let S_g be a closed surface of genus $g \geq 1$, and let Γ_g be the mapping class group of S_g . The group Γ_g admits a homomorphism \mathfrak{s} onto $\mathrm{Sp}(2g, \mathbb{Z})$ (we associate to each element its action on homology; the symplectic structure comes from the intersection pairing). The following result can be found in [4]:

Theorem 8.1. *For $\gamma \in \Gamma_g$ to be pseudo-Anosov, it is sufficient that $g = \gamma$ satisfy all of the following conditions:*

- (1) *The characteristic polynomial of g is irreducible.*
- (2) *The characteristic polynomial of g is not cyclotomic.*
- (3) *The characteristic polynomial of g is not of the form $g = h(x^k)$, for some $k > 1$.*

The following is a corollary of our results on matrix groups:

Theorem 8.2. *Let g_1, \dots, g_k be a generating set of $\mathrm{Sp}(2n, \mathbb{Z})$. The probability that a random product of length N of g_1, \dots, g_k satisfies the conditions of Theorem 8.1 goes to 1 as N goes to infinity.*

Proof. We prove that the probability that the random word w_N not satisfy the conditions goes to 0. By Theorem 6.5, the probability that w_N has reducible characteristic polynomial goes to 0. In order for the characteristic polynomial to be of the form $g = h(x^k)$ it is necessary that the linear term (the trace) vanish. The set of traceless matrices is a proper subvariety of $\mathrm{Sp}(2g)$, so by Theorem 2.1, for a large p , the probability that a given matrix is traceless is $\ll 1/p$, and in particular is bounded away from 1. The proof of Theorem 6.5 now goes through verbatim to show that the set of traceless matrices is asymptotically negligible in $\mathrm{Sp}(2g, \mathbb{Z})$.

Finally, the set of cyclotomic polynomials of a given degree $2g$ is bounded by some function $h(g)$. \square

9. FREE GROUP AUTOMORPHISMS

An automorphism ϕ of a free group F_n is called *strongly irreducible*³ if no (positive) power of ϕ sends a free factor H of F_n to a conjugate. This concept was introduced by M. Bestvina and M. Handel [1], and many of the results of the theory of automorphisms of free groups are shown for such automorphisms (for a survey, and the relationship between strongly irreducible automorphisms and

³This terminology, with strong support from this author, has been introduced by L. Mosher and M. Handel for what was previously known as *irreducible with irreducible powers*

pseudo-Anosov automorphisms of surfaces, the reader is urged to read the very clear survey⁴[3]. By passing to the action of ϕ on the abelianization of F_n (equivalently, on $H_1(F_n, \mathbb{Z})$), Section 7⁵ shows the following:

Theorem 9.1. *Let f_1, \dots, f_k be a generating set of the automorphism group of F_n . Consider all words of length L in f_1, \dots, f_k . Then, for any n , the probability that such a word is irreducible tends to 1 as L tends to infinity and also the probability that such a word is strongly irreducible tends to 1 as L tends to infinity.*

10. GALOIS GROUPS OF GENERIC RESTRICTED POLYNOMIALS

Let $P_{N,d}(\mathbb{Z})$ be the set of monic polynomials of degree d with integral coefficients bounded by N in absolute value. It is a classical result of B. L. van der Waerden that the probability that the Galois group of $p \in P_{N,d}(\mathbb{Z})$ is the full symmetric group S_d tends to 1 as N tends to infinity. The argument is quite elegant: First, it is observed that a subgroup $H < S_d$ is the full symmetric group if and only if H intersects every conjugacy class of S_d . This means that H has an element with every possible cycle type. It is further noted that there is a cycle type (n_1, \dots, n_k) in the Galois group of p over $\mathbb{Z}/p\mathbb{Z}$ if and only if p factors over $\mathbb{Z}/p\mathbb{Z}$ into irreducible polynomials of degrees n_1, \dots, n_k . Using Dedekind's generating function for the number of irreducible polynomials over $\mathbb{Z}/p\mathbb{Z}$ of a given degree, it is shown that the probability of a fixed partition is bounded below by a constant (independent of the prime p), and the proof is finished by an application of a Chinese Remainder Theorem.

In this note, we ask the following simple-sounding question: Let $P_{N,d,a,k}(\mathbb{Z})$ be the set of all polynomials in $P_{N,d}(\mathbb{Z})$ where the coefficient of x^k equals a . Is it still true that the Galois group of a random such polynomial is the full symmetric group? The result would obviously follow if the probability that the Galois group of a random general polynomial is "generic" were to go to 1 sufficiently fast with N . In fact, the probability that an element of $P_{N,d}$ is *reducible* (which means that its Galois group is not transitive, hence not S_n) is of the order of $1/N$, so that approach does not work.

Mimicking the proof of van der Waerden's result does not appear to work (at least not easily): Dedekind's argument enumerates all

⁴ to appear, but available at

<http://www.math.cornell.edu/~vogtmann/papers/AutQuestions/Questions.html>

⁵We need to change $SL(n, \mathbb{Z})$ to $GL(n, \mathbb{Z})$ throughout

irreducible polynomials, and the result is not “graded” by specific coefficients. It is certainly possible that the argument can be pushed through, but this appears to be somewhat involved.

Given this sad state of affairs, we first use a simple trick and Dirichlet’s theorem on primes on arithmetic progressions to show first the following technical result:

Theorem 10.1. *The probability that a random element of $P_{N,d,a,k}(\mathbb{Z}/p\mathbb{Z})$ has a prescribed splitting type s approaches the probability that a random unrestricted polynomial of degree d has the splitting type s , as long as $p - 1$ is relatively prime to $(d - k)!$, and as p becomes large.*

which implies (by van der Waerden’s argument):

Theorem 10.2. *The probability that a random element of $P_{N,d,a,k}(\mathbb{Z})$ has S_d as the Galois group tends to 1 as N tends to infinity,*

It should be noted that the (multivariate) Large Sieve (as used by P. X. Gallagher in [9]) can be used to give an effective estimate on the probability in the statement of Theorems 10.2: that is: $p(N) \ll N^{-1/2} \log N$.

10.1. Proof of Theorem 10.2. We will need two ingredients other than van der Waerden’s original idea. The first is A. Weil’s estimate (Theorem 2.2), the second is the following classical result:

Theorem 10.3 ([14][Theorem VIII.9.1].) *Let k be a field, and $n \geq 2$ an integer. Let $a \in k$, $a \neq 0$. Assume that for all prime numbers p such that $p|n$ we have $a \notin k^p$, and if $4|n$, then $a \notin -4k^4$. Then $X^n - a$ is irreducible in $k[X]$.*

Theorem 10.3 goes essentially back to N. H. Abel’s foundational memoir.

We will need an additional observation:

Lemma 10.4. *Let $q = p^l$, and let $x_1, \dots, x_k \in \mathbb{F}_q$. Let $a, b \in \mathbb{F}_p$, with $(a, b) \neq (1, 0)$ and let $g(a, b)(x) = ax + b$ be a transformation of \mathbb{F}_q to itself. Then, it is not possible for $g(a, b)$ to permute x_1, \dots, x_k , if $k!$ is coprime to $p - 1$.*

Lemma 10.5. *Consider a polynomial f of degree d over \mathbb{F}_p , such that $d < p$, and such that the coefficient of x^{d-1} does not vanish. Then there is no pair $(a, b) \neq (1, 0)$, such that $f(ax + b) = a^d f(x)$, for all $x \in \mathbb{F}_p$.*

Proof. There are two distinct cases to analyze. The first is when $a = 1$. In that case, $f(x+b) = f(x)$ for all $x \in \mathbb{F}_p$, and since $p > d$, $f(x+b) = f(x)$, for all x in the algebraic closure of F_p . Let r be a root of f . Then, so are

$r + b, r + 2b, \dots, r + b(p - 1)$, which are all distinct as long as $b \neq 0$, but since p is greater than d that means that f is identically 0.

The second case is when $a \neq 1$. In that case, $x_0 = b/(1 - a)$ is fixed under the substitution $x \rightarrow ax + b$, so setting $z = x - x_0$, we see that $f(ax + b) = f(az)$, and so $f(ax + b) = a^d f(x)$ implies that $f(az) = a^d f(z)$, for all $z \in \mathbb{F}_p$. Since $p > \deg f$, the corresponding coefficients of the right and the left hand polynomials must be equal. Since the coefficient of x^{d-1} does not vanish, it follows that $a = 1$, which contradicts our assumption. \square

The argument now proceeds as follows. First, we note that if the polynomial $f(x)$ of degree d has a certain splitting type (hence Galois group) over \mathbb{F}_p then so does $f(ax + b)/a^d$, for any $a \neq 0, b \in \mathbb{F}_p$. The set of all linear substitutions forms a group \mathbb{A} , which acts freely on the set of polynomials of degree d , except for the (small) exceptional set of polynomials with a vanishing coefficient of x^{d-1} as long as $d < p$ (by Lemma 10.5), so the distribution of splitting types among the \mathbb{A} orbits is the same as among all of the polynomials of degree d . Now, consider polynomials with constant term 1. How many of them are there in the \mathbb{A} orbit of $f(x)$? It is easy to see that the number is equal to the number of solutions to

$$f(b) = a^d.$$

If the curve C_f given by $f(x) - y^d$ is absolutely irreducible, that number is $p + O(\sqrt{p})$, by Theorem 2.2. By Theorem 10.3, in order for C_f to not be absolutely irreducible, we must either have that $f(x) = g^q(x)$, for some $q|d$, or $f(x) = -4h^4(x)$, in case $4|d$. But the number of such polynomials is bounded by $O(p^{d/2})$, which is asymptotically negligible. So, we see that the distribution of splitting types amongst polynomials of degree d with constant term 1 is the same as for all polynomials, as long as $d < p$.

11. RANDOM WALKS ON GROUPS AND GRAPHS

We consider the following situation: G is a finite *undirected* (multi)graph with n vertices, and Γ is a finite group. Each vertex v_i of G is decorated with an element t_i of Γ ; we assume that the set $\{t_1, \dots, t_n\}$ generates Γ . We make the following assumptions:

Condition 11.1. (1) The graph G is *ergodic*: the adjacency matrix $A(G)$ of G is a Perron-Frobenius matrix (meaning that there is a unique eigenvalue of maximal modulus, and that some power $A^k(G)$ has all entries positive).

- (2) We assume further that for every nontrivial one-dimensional unitary representation ρ of Γ , there exists $1 \leq i, j \leq n$, such that $\rho(t_i) \neq \rho(t_j)$.

To each walk $w = (v_{i_1}, v_{i_2}, \dots, v_{i_k})$ on G we associate the element $\gamma_w = t_{i_k} \dots t_{i_2} t_{i_1}$. We now denote the set of all walks of length N by W_N .

Furthermore, we define a *probability distribution* P_N on Γ , as follows: the probability density $p_N : \Gamma \rightarrow \mathbb{R}_+$ is defined as $p_N(\gamma) = |\{w \in W_N \mid \gamma_w = \gamma\}|/|W_N|$.

A slightly more abstract way to think of this is as follows: P_N is the function on the group ring $\mathbb{R}[\Gamma]$, defined by:

$$P_N = \frac{1}{|W_N|} \sum_{w \in W_N} \gamma_w.$$

Our main result is the following:

Theorem 11.2. *The distributions P_N converge to the uniform distribution on Γ as N goes to infinity. Furthermore, there is a constant $c = c(G, \Gamma, T) < 1$, such that $p_N(\gamma) - 1/|\Gamma| < c^N$, for all $\gamma \in \Gamma$. Here, T refers to the assignment of generators of Γ to the vertices of G .*

We will actually define a slightly stronger result: we will pick $1 \leq i, j \leq n$, and consider the set of all walks $W_{N,i,j}$ of length N from the i th to the j th vertex of G . We define distributions $P_{N,i,j}$ in the obvious way. Then:

Theorem 11.3. *Theorem 11.2 holds with P_N replaced by $P_{N,i,j}$.*

The starting point for the proof of the theorems above is Fourier Transform on finite groups, which is discussed in Section 12. In particular, we will be using Theorem 12.2 and Corollary 12.3 to reduce the question of whether a probability distribution is close to uniform to the proving that the Fourier Transform is small at every *non-trivial* representation. The reader might well wonder how moving the problem to Fourier transform space helps us – the answer is that it turns out that we can reduce the estimation of the “fourier coefficients” to questions in linear algebra, through the construction in Section 13.

12. FOURIER TRANSFORM ON FINITE GROUPS

For a thorough introduction to the topic of this section the reader is referred to [20, 21]. Let Γ be a finite group, and let $f : \Gamma \rightarrow \mathbb{C}$ be a function on Γ . Furthermore, let $\widehat{\Gamma}$ be the *unitary dual* of Γ : the set of all irreducible complex unitary representations of Γ . To f we can

associate its *Fourier Transform* \hat{f} . This is a function which associates to each d -dimensional unitary representation ρ a $d \times d$ matrix $\hat{f}(\rho)$ as follows:

$$\hat{f}(\rho) = \sum_{\gamma \in \Gamma} f(\gamma) \rho(\gamma).$$

There is an inverse transformation, as well. Given a function g on $\widehat{\Gamma}$ which associates to each d -dimensional representation ρ a $d \times d$ matrix $g(\rho)$, we can write:

$$g^\sharp(\gamma) = \frac{1}{|\Gamma|} \sum_{\rho \in \widehat{\Gamma}} d_\rho \operatorname{tr}(g(\rho) \rho(\gamma^{-1})),$$

where d_ρ is the dimension of ρ . We mean “inverse” in the most direct way possible:

$$f^\sharp = f.$$

The following result is classical (see, eg, [21]):

Theorem 12.1.

$$\sum_{\rho \in \widehat{\Gamma}} d_\rho^2 = |\Gamma|,$$

and, together with the Fourier inversion formula, implies

Theorem 12.2. *Let g be a function on $\widehat{\Gamma}$, such that for every nontrivial $\rho \in \widehat{\Gamma}$,*

$$\|g(\rho)\|_{op} < \epsilon,$$

where $\|\bullet\|_{op}$ denotes the operator norm (see Section 14). Then, for any $\gamma_1, \gamma_2 \in \Gamma$,

$$|g^\sharp(\gamma_1) - g^\sharp(\gamma_2)| < 2\epsilon.$$

Proof. First, note that for the trivial representation ρ_0 , the quantity

$$d_{\rho_0} g(\rho_0) \rho_0(\gamma) = g(\rho_0),$$

so does not depend on γ . By the Fourier inversion formula, then,

$$\begin{aligned}
& |g^\sharp(\gamma_1) - g^\sharp(\gamma_2)| = \\
& \left| \frac{1}{|\Gamma|} \sum_{\substack{\rho \in \widehat{\Gamma} \\ \rho \neq \rho_0}} d_\rho \operatorname{tr}(g(\rho)(\rho(\gamma_1) - \rho(\gamma_2))) \right| \leq \\
& \sum_{i=1}^2 \left| \frac{1}{|\Gamma|} \sum_{\substack{\rho \in \widehat{\Gamma} \\ \rho \neq \rho_0}} d_\rho \operatorname{tr}(g(\rho)\rho(\gamma_i)) \right| \stackrel{\text{by Eq. (4)}}{\leq} \\
& \frac{2}{|\Gamma|} \sum_{\rho \in \widehat{\Gamma}} d_\rho^2 \|g(\rho)\|_{\text{op}} < 2\epsilon.
\end{aligned}$$

□

Corollary 12.3. *Under the assumption of Theorem 12.2, and assuming in addition that g is real valued, if*

$$\sum_{\gamma \in \Gamma} g(\gamma) = 1,$$

then

$$g(\gamma) - 1/|\Gamma| < 2\epsilon \quad \forall \gamma \in \Gamma.$$

Furthermore, if $\Omega \in \Gamma$,

$$(1) \quad \left| \sum_{\gamma \in \Omega} g(\gamma) - \frac{|\Omega|}{|\Gamma|} \right| < 2\epsilon |\Omega|.$$

Proof. Without loss of generality, suppose that $g(\gamma) > 1/|\Gamma|$. Then there is a γ_2 , such that $g(\gamma_2) < 1/|\Gamma|$. Thus,

$$g(\gamma) - 1/|\Gamma| < g(\gamma) - g(\gamma_2) < 2\epsilon.$$

The estimate (1) follows immediately by summing over Ω . □

13. FOURIER ESTIMATES VIA LINEAR ALGEBRA

In order to prove Theorem 11.3, we would like to use Theorem 12.2, and to show the equidistribution result, we would need to show that for every *nontrivial* irreducible representation ρ ,

$$(2) \quad \lim_{N \rightarrow \infty} \frac{1}{|W_{N,i,j}|} \operatorname{tr} \sum_{w \in W_{N,i,j}} \rho(\gamma_w) = 0.$$

To demonstrate Eq. (2), suppose that ρ is k -dimensional, so acts on a k -dimensional Hilbert space $H_\rho = H$. Let $Z = L^2(G)$ – the space of complex-valued functions from $V(G)$ to \mathbb{C} , let e_1, \dots, e_n be the standard basis of Z , and let P_i be the orthogonal projection on the i -th coordinate space. We introduce the matrix

$$U_\rho = \sum_{i=1}^n P_i \otimes \rho(t_i) = \begin{pmatrix} \rho(t_1) & 0 & \dots & 0 \\ 0 & \rho(t_2) & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & \rho(t_n) \end{pmatrix},$$

and also the matrix $A_\rho = A(G) \otimes I_H$, where I_H is the identity operator on H . Both U_ρ and A_ρ act on $Z \otimes H$. The following is immediate:

Lemma 13.1. *Consider the matrix $(U_\rho A_\rho)^l$, and think of it as an $n \times n$ matrix of $k \times k$ blocks. Then the ij -th block equals the sum over all paths w of length l beginning at v_i and ending of v_j of $\rho(\gamma_w)$.*

Now, let T_{ji} be the operator on Z which maps e_k to $\delta_{kj}e_i$.

Lemma 13.2.

$$\text{tr} \left[\left((T_{ji}^t P_j) \otimes I_H \right) (U_\rho \otimes A_\rho)^N (P_i \otimes I_H) \right] = \text{tr} \sum_{w \in W_{N,i,j}} \rho(\gamma_w)$$

Proof. The argument of trace on the left hand side simply extracts the ij -th $k \times k$ block from $(U_\rho \otimes A_\rho)^N$. \square

By submultiplicativity of operator norm, we see that

$$\| (T_{ji}^t P_j) \otimes I_H (U_\rho \otimes A_\rho)^N P_i \otimes I_H \|_{\text{op}} \leq \| (U_\rho \otimes A_\rho)^N \|_{\text{op}},$$

and so proving Theorem 11.3 reduces (thanks to Theorem 12.2) to showing

Theorem 13.3.

$$\lim_{N \rightarrow \infty} \frac{\| (U_\rho \otimes A_\rho)^N \|_{\text{op}}}{|W_{N,i,j}|} = 0,$$

for any non-trivial ρ .

Notation 13.4. We will denote the spectral radius of an operator A by $\mathcal{R}(A)$.

Since $|W_{N,i,j}| \asymp \mathcal{R}^N(A(G))$, and by Gelfand's Theorem (Theorem 14.3),

$$\lim_{N \rightarrow \infty} \|B^N\|^{1/N} = \mathcal{R}(B),$$

for any matrix B and any matrix norm $\|\bullet\|$, Theorem 13.3 is equivalent to the statement that the spectral radius of $U_\rho \otimes A_\rho$ is smaller than that of $A(G)$.

Theorem 13.3 is proved in Section 13.1.

13.1. Proof of Theorem 13.3.

Lemma 13.5. *Let A be a bounded hermitian operator $A : H \rightarrow H$, and $U : H \rightarrow H$ a unitary operator on the same Hilbert space H . Then the spectral radius of UA is smaller than the spectral radius of A , and the inequality is strict unless an eigenvector of A with maximal eigenvalue is also an eigenvector of U .*

Proof. The spectral radius of UA does not exceed the operator norm of UA , which is equal to the spectral radius of A . Suppose that the two are equal, so that there is a v , such that $\|UA v\| = \mathcal{R}(A)v$, and v is an eigenvector of UA . Since U is unitary, v must be an eigenvector of A , and since it is also an eigenvector of UA , it must also be an eigenvector of U . \square

In the case of interest to us, ρ is a k -dimensional irreducible representation of Γ , $U = \text{Diag}(\rho(t_1), \dots, \rho(t_n))$, while $A = A(G) \otimes I_k$. We assume that $A(G)$ is an irreducible matrix, so that there is a unique eigenvalue of modulus $\mathcal{R}(A(G))$, that eigenvalue λ_{\max} (the *Perron-Frobenius eigenvalue*) is positive, and it has a strictly positive eigenvector v_{\max} . We know that the spectral radius of A equals the spectral radius of $A(G)$, and the eigenspace of λ_{\max} is the set of vectors of the form $v_{\max} \otimes w$, where w is an arbitrary vector in \mathbb{C}^k . If $v_{\max} = (x_1, \dots, x_n)$, we can write $v_{\max} \otimes w = (x_1 w, \dots, x_n w)$, and so $U(v_{\max} \otimes w) = (x_1 \rho(t_1) w, \dots, x_n \rho(t_n) w)$. Since all of the x_i are nonzero, in order for the inequality in Lemma 13.5 to be nonstrict, we must have some w for which $\rho(t_i) w = c w$ (where the constant c does *not* depend on i .) Since the elements t_i generate Γ , the existence of such a w contradicts the irreducibility of ρ , unless ρ is one dimensional. This proves Theorem 13.3

14. SOME REMARKS ON MATRIX NORMS

In this note we use a number of matrix norms, and it is useful to summarize what they are, and some basic relationships and inequalities satisfied by them. For an extensive discussion the reader is referred to the classic [12]. All matrices are assumed square, and $n \times n$.

A basic tool in the inequalities below is the *singular value decomposition* of a matrix A .

Definition 14.1. *The singular values of A are the non-negative square roots of the eigenvalues of AA^* , where A^* is the conjugate transpose of A .*

Since AA^* is a positive semi-definite Hermitian matrix for any A , the singular values $\sigma_1 \stackrel{\text{def}}{=} \sigma_{\max} \geq \sigma_2 \geq \dots$ are non-negative real numbers. For a Hermitian A , the singular values are simply the absolute values of the eigenvalues of A .

The first matrix norm is the *Frobenius norm*, denoted by $\|\bullet\|$. This is defined as

$$\|A\| = \sqrt{\text{tr } AA^*} = \sqrt{\sum_i \sigma_i^2}.$$

This is also the sum of the square moduli of the elements of A .

The next matrix norm is the *operator norm*, $\|\bullet\|_{\text{op}}$, defined as

$$\|A\|_{\text{op}} = \max_{\|v\|=1} \|Av\| = \sigma_{\max}$$

Both the norms $\|\bullet\|$ and $\|\bullet\|_{\text{op}}$ are *submultiplicative* (submultiplicativity is part of the definition of matrix norm: saying that the norm $\|\bullet\|$ is submultiplicative means that $\|AB\| \leq \|A\| \|B\|$.)

From the singular value interpretation⁶ of the two matrix norms and the Cauchy-Schwartz inequality we see immediately that

$$(3) \quad \|A\| / \sqrt{n} \leq \|A\|_{\text{op}} \leq \|A\|$$

We will also need the following simple inequalities:

Lemma 14.2. *Let U be a unitary matrix:*

$$(4) \quad |\text{tr } AU| \leq \|A\| \sqrt{n} \leq n \|A\|_{\text{op}}.$$

Proof. Since U is unitary, $\|U\| = \|U^t\| = \sqrt{n}$. So, by the Cauchy-Schwartz inequality, $\text{tr } AU \leq \|A\| \|U\| = \sqrt{n} \|U\|$. The second inequality follows from the inequality (3). \square

The final (and deepest result) we will have the opportunity to use is:

Theorem 14.3 (Gelfand). *For any operator M , the spectral radius $\mathcal{R}(M)$ and any matrix norm $\|\bullet\|$,*

$$\mathcal{R}(M) = \lim_{k \rightarrow \infty} \|\|M^k\|\|^{1/k},$$

⁶A celebrated result of John von Neumann states that *any* unitarily invariant matrix norm is a symmetric gauge on the space of singular values - [22].

APPENDIX A. SYMPLECTIC MATRICES AND DAVID KIRBY'S THEOREM

Recall that the *symplectic quadratic form* in $2n$ dimensions is given by the matrix

$$J = \begin{pmatrix} 0 & \mathbf{I}_n \\ -\mathbf{I}_n & 0 \end{pmatrix},$$

where \mathbf{I}_n is the $n \times n$ identity matrix. A $2n \times 2n$ matrix M is called *symplectic* if it preserves the symplectic form J , that is

$$(5) \quad M^t J M = J.$$

The invariance relation Eq. (5) can be written slightly more explicitly if we write M in $n \times n$ block form, as

$$M = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$$

In that case, Eq. (5) is equivalent to the following system:

$$(6) \quad A^t C = C^t A (= (A^t C)^t),$$

$$(7) \quad B^t D = D^t B (= (B^t D)^t),$$

$$(8) \quad A^t D - C^t B = \mathbf{I}_n.$$

This simple observation is all that is needed to prove the following

Theorem A.1 (David Kirby, [13]). *Let \mathcal{R} be a commutative ring with 1, and let $p(x) \in \mathcal{R}[x]$ be a reciprocal polynomial. Then, there is a symplectic matrix M with coefficients in \mathcal{R} , such that*

$$\det(x\mathbf{I}_{2n} - M) = p(x).$$

Proof sketch. In the notation of Eq. (A), let $A = \mathbf{0}$, let $\det B = 1$, and let $C = -(B^t)^{-1}$. Then, in order for

$$M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} = \begin{pmatrix} \mathbf{0} & B \\ -(B^t)^{-1} & D \end{pmatrix}$$

to be symplectic, it is necessary and sufficient that $B^t D$ be symmetric (since Eq. (6) and Eq. (8) hold automatically).

Now, let

$$B_{ij} = \begin{cases} 0 & i + j < n \\ 1 & i + j = n \\ b_{i+j+1-n} & i + j > n, \end{cases}$$

and let $D = E + F$, where

$$E_{ij} = \begin{cases} 0 & |i - j| > 1 \text{ or } i = j > 1. \\ 1 & |i - j| = 1 \\ -1 & i = j = 1, \end{cases}$$

and

$$F_{ij} = \begin{cases} 0 & j < n \\ f_i & \text{otherwise} \end{cases}$$

for some $b_2, \dots, b_n, f_1, \dots, f_n \in \mathcal{R}$. It is not hard to check that this is always so if $n = 1$, and when $n > 1$, it is necessary and sufficient that

$$(9) \quad b_2 = f_1 + 1,$$

$$(10) \quad b_i = f_{i-1} + \sum_{j=2}^{i-1} b_j f_{i-j}.$$

Now, if $p(x) = (1 + x^{2n}) + a_1(x + x^{2n-1}) + \dots + a_n(x^n + x^{n+1})$, a computation shows that a matrix M as above with $f_i = a_{i-1} - a_i$ does the trick. We won't go through all the details, but the main idea in computing the determinant comes from the following:

$$\begin{pmatrix} x\mathbf{I} & -B \\ B^{-1} & xI - D \end{pmatrix} \begin{pmatrix} B & 0 \\ x\mathbf{I} & B^{-1} \end{pmatrix} = \begin{pmatrix} 0 & -\mathbf{I} \\ (x^2 + 1)\mathbf{I} - xD & xB^{-1} - DB^{-1} \end{pmatrix}$$

so by taking determinants:

$$\det(x\mathbf{I} - M) = \det((x^2 + 1)\mathbf{I} - xD).$$

□

REFERENCES

- [1] Mladen Bestvina and Michael Handel. Train tracks and automorphisms of free groups. *Ann. of Math. (2)*, 135(1):1–51, 1992.
- [2] Armand Borel. *Linear algebraic groups*, volume 126 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1991.
- [3] M. Bridson and K. Vogtmann. Automorphism groups of free, surface, and free abelian groups. Preprint, 2006, to appear in *Handbook of Questions around the Mapping Class Group*, edited by B. Farb.
- [4] Andrew J. Casson and Steven A. Bleiler. *Automorphisms of surfaces after Nielsen and Thurston*, volume 9 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 1988.
- [5] Nick Chavdarov. The generic irreducibility of the numerator of the zeta function in a family of curves with large monodromy. *Duke Math. J.*, 87(1):151–180, 1997.

- [6] David Cox, John Little, and Donal O'Shea. *Ideals, varieties, and algorithms*. Undergraduate Texts in Mathematics. Springer-Verlag, New York, second edition, 1997. An introduction to computational algebraic geometry and commutative algebra.
- [7] W. Duke, Z. Rudnick, and P. Sarnak. Density of integer points on affine homogeneous varieties. *Duke Math. J.*, 71(1):143–179, 1993.
- [8] Alex Eskin and Curt McMullen. Mixing, counting, and equidistribution in Lie groups. *Duke Math. J.*, 71(1):181–209, 1993.
- [9] P. X. Gallagher. The large sieve and probabilistic Galois theory. In *Analytic number theory (Proc. Sympos. Pure Math., Vol. XXIV, St. Louis Univ., St. Louis, Mo., 1972)*, pages 91–101. Amer. Math. Soc., Providence, R.I., 1973.
- [10] Meinolf Geck. *An introduction to algebraic geometry and algebraic groups*, volume 10 of *Oxford Graduate Texts in Mathematics*. Oxford University Press, Oxford, 2003.
- [11] Marshall Hall. *Group Theory*. MacMillan, 1959.
- [12] Roger A. Horn and Charles R. Johnson. *Matrix analysis*. Cambridge University Press, Cambridge, 1990. Corrected reprint of the 1985 original.
- [13] David Kirby. Integer matrices of finite order. *Rend. Mat. (6)*, 2:403–408, 1969.
- [14] Serge Lang. *Algebra*. Addison-Wesley Publishing Company Advanced Book Program, Reading, MA, second edition, 1984.
- [15] Serge Lang and André Weil. Number of points of varieties in finite fields. *Amer. J. Math.*, 76:819–827, 1954.
- [16] Amos Nevo and Peter Sarnak. Prime integral matrices. 2006. preprint.
- [17] Morris Newman. *Integral matrices*. Academic Press, New York, 1972. Pure and Applied Mathematics, Vol. 45.
- [18] . Igor Rivin. Equidistribution and irreducibility in lattices and groups of automorphisms. 2007. In preparation.
- [19] Igor Rivin. Growth on groups (and other stories). Technical Report math.CO/9911076, arxiv.org, 1999.
- [20] Jean-Pierre Serre. *Représentations linéaires des groupes finis*. Hermann, Paris, revised edition, 1978.
- [21] Barry Simon. *Representations of finite and compact groups*, volume 10 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 1996.
- [22] John von Neumann. Some matrix inequalities and metrization of matrix space. *Tomsk University Review*, 1:286–300. In collected works, Pergamon, Oxford, 1962, Volume IV, 205-218.
- [23] André Weil. *Sur les courbes algébriques et les variétés qui s'en déduisent*. Actualités Sci. Ind., no. 1041 = Publ. Inst. Math. Univ. Strasbourg 7 (1945). Hermann et Cie., Paris, 1948.
- [24] Helmut Wielandt. *Permutation Groups*. California Institute of Technology, Pasadena, CA, 1961.

DEPARTMENT OF MATHEMATICS, TEMPLE UNIVERSITY, PHILADELPHIA

E-mail address: rivin@math.temple.edu

Current address: Mathematics Department, Stanford University, Stanford, California

E-mail address: rivin@math.stanford.edu