

Quantum Hilbert matrices and orthogonal polynomials

Jørgen Ellegaard Andersen and Christian Berg

March 9, 2021

Abstract

Using the notion of quantum integers associated with a complex number $q \neq 0$, we define the quantum Hilbert matrix and various extensions. They are Hankel matrices corresponding to certain little q -Jacobi polynomials when $|q| < 1$, and for the special value $q = (1 - \sqrt{5})/(1 + \sqrt{5})$ they are closely related to Hankel matrices of reciprocal Fibonacci numbers called Filbert matrices. We find a formula for the entries of the inverse quantum Hilbert matrix.

2000 *Mathematics Subject Classification*: primary 33D45; secondary 11B39.

Keywords: Basic orthogonal polynomials, quantum integers, Fibonacci numbers.

1 Introduction

In [8] Hilbert introduced the *Hilbert matrices*

$$\mathcal{H}_n = (1/(l + j + 1)), \quad 0 \leq l, j \leq n, \quad n = 0, 1, \dots, \quad (1)$$

and found the following expression for their determinants

$$\det \mathcal{H}_n = \left(\prod_{k=1}^n (2k + 1) \binom{2k}{k}^2 \right)^{-1}, \quad (2)$$

showing that they are the reciprocal of integers. This fact is also a consequence of the observation that the inverse matrices \mathcal{H}_n^{-1} have integer entries and Choi [6] found the following integer expression for them

$$(\mathcal{H}_n^{-1})_{l,j} = (-1)^{l+j} (l + j + 1) \binom{n+l+1}{n-j} \binom{n+j+1}{n-l} \binom{l+j}{l} \binom{l+j}{j}. \quad (3)$$

This formula was generalized in [4, Theorem 4.1] to the one parameter extension $\mathcal{H}_n^{(\alpha)} = (\alpha/(l+j+\alpha))$ of the Hilbert matrices, where $\alpha > 0$,

$$(\mathcal{H}_n^{(\alpha)})_{l,j}^{-1} = (-1)^{l+j} \frac{l+j+\alpha}{\alpha} \binom{n+l+\alpha}{n-j} \binom{n+j+\alpha}{n-l} \binom{l+j+\alpha-1}{l} \binom{l+j+\alpha-1}{j}. \quad (4)$$

The idea of proof is to observe that the matrices are the Hankel matrices of a moment problem. After having determined the corresponding orthogonal polynomials and their kernel polynomials one uses the result that the matrix of coefficients of the kernel polynomial is the inverse of the Hankel matrix, see [4, Theorem 2.1].

In [13] Richardson noticed that the *Filbert matrices*

$$\mathcal{F}_n = (1/F_{l+j+1}), \quad 0 \leq l, j \leq n, \quad n = 0, 1, \dots, \quad (5)$$

where $F_n, n \geq 0$ is the sequence of Fibonacci numbers, have the property that all elements of the inverse matrices are integers. Richardson gave an explicit formula for the elements of the inverse matrices and proved it using computer algebra. It is the special case $\alpha = 1, \sinh \theta = \frac{1}{2}$ of (39). The formula shows a remarkable analogy with Choi's formula (3) in the sense that one shall replace the binomial coefficients $\binom{n}{k}$ by the analogous *Fibonomial coefficients*

$$\binom{n}{k}_{\mathbb{F}} = \prod_{j=1}^k \frac{F_{n-j+1}}{F_j}, \quad 0 \leq k \leq n, \quad (6)$$

with the usual convention that empty products are defined as 1. These coefficients are defined and studied in [11] and are integers. We recall that the sequence of Fibonacci numbers is $F_0 = 0, F_1 = 1, \dots$, with the recursion formula $F_{n+1} = F_n + F_{n-1}, n \geq 1$.

The Hilbert matrices are the Hankel matrices (s_{l+j}) corresponding to the moment sequence

$$s_n = 1/(n+1) = \int_0^1 x^n dx, \quad n \geq 0, \quad (7)$$

and the corresponding orthogonal polynomials are the Legendre polynomials for the interval $[0, 1]$, see [2, Section 7.7]. This was used in [4] and in the survey paper [3] to prove Choi's formula.

In [4] it was also proved that for $\alpha \in \mathbb{N}$ the sequence $(F_\alpha/F_{n+\alpha})_{n \geq 0}$ is the moment sequence of a signed measure of total mass one and the corresponding orthogonal polynomials were identified as special little q -Jacobi polynomials corresponding to the value $q = (1 - \sqrt{5})/(1 + \sqrt{5})$. From this Richardson's formula (39) for the elements of the inverse Filbert matrices was derived as the case $\alpha = 1$ of [4, (29)].

The results about Fibonacci numbers have been extended by Ismail in [10] to a one-parameter family of sequences $(F_n(\theta))_{n \geq 0}, \theta > 0$, determined by the recursion

$$F_{n+1}(\theta) = 2 \sinh \theta F_n(\theta) + F_{n-1}(\theta), \quad n \geq 1, \quad F_0(\theta) = 0, F_1(\theta) = 1. \quad (8)$$

When $\sinh \theta = \frac{1}{2}$ we have $F_n(\theta) = F_n$, and when $2 \sinh \theta$ is a positive integer then all $F_n(\theta)$ are integers.

The purpose of this paper is to give a common generalization of all these results by the use of *quantum integers* defined by

$$[n]_q = \frac{q^{n/2} - q^{-n/2}}{q^{1/2} - q^{-1/2}}, \quad n = 0, 1, \dots \quad (9)$$

where $q \in \mathbb{C} \setminus \{0\}$. To make the definition precise we consider $q \rightarrow [n]_q$ as a holomorphic function in the cut plane $\mathbb{C} \setminus]-\infty, 0]$ extended to $q \in]-\infty, 0[$ by

$$[n]_q = \lim_{\varepsilon \rightarrow 0^+} [n]_{q+i\varepsilon}.$$

Clearly $[n]_q = n$ for $n = 0, 1$ and $[n]_1 = n$ for all $n \in \mathbb{N}$. The quantum integer $[n]_q$ vanishes, when q is an n 'th root of unity different from 1.

Assume now $0 < |q| < 1$. We consider the complex measures in the complex plane depending on $\alpha \in \mathbb{N}$ and defined by

$$\mu^{(\alpha)}(q) = (1 - q^\alpha) \sum_{k=0}^{\infty} q^{k\alpha} \delta_{q^{k+1/2}}, \quad (10)$$

where δ_c is the Dirac measure concentrated at $c \in \mathbb{C}$. It is easy to see the following q -analogue of (7)

$$\frac{1}{[n+1]_q} = \int x^n d\mu^{(1)}(q)(x), \quad n \geq 0, \quad (11)$$

and in general

$$\frac{[\alpha]_q}{[n+\alpha]_q} = \int x^n d\mu^{(\alpha)}(q)(x), \quad n \geq 0. \quad (12)$$

The Hankel matrices corresponding to $\mu^{(1)}(q)$ are defined as

$$\mathcal{H}_n(q) = \left(\frac{1}{[l+j+1]_q} \right), \quad 0 \leq l, j \leq n, \quad n = 0, 1, \dots, \quad (13)$$

and are called the *quantum Hilbert matrices*, but we will also consider the *generalized quantum Hilbert matrices* ($\alpha \in \mathbb{N}$)

$$\mathcal{H}_n^{(\alpha)}(q) = \left(\frac{[\alpha]_q}{[l+j+\alpha]_q} \right), \quad 0 \leq l, j \leq n, \quad n = 0, 1, \dots \quad (14)$$

These matrices are well-defined for non-zero complex numbers q which are not roots of unity of order $\leq 2n + \alpha$. When $0 < q < 1$ the measure $\mu^{(1)}(q)$ is a probability measure on $[0, 1]$ which converges weakly to the Lebesgue measure on $[0, 1]$ for $q \rightarrow 1$. The quantum Hilbert matrices $\mathcal{H}_n(q)$ converge to the ordinary Hilbert matrices when $q \rightarrow 1$.

We prove in section 2 that the generalized quantum Hilbert matrices are regular and find a formula for the elements of the inverse matrix, see Theorem 2.1. This is a q -analogue of Choi's formula when $\alpha = 1$. The proof uses the method of [4] by identifying the orthogonal polynomials of $\mu^{(\alpha)}(q)$ as little q -Jacobi polynomials.

In section 3 we consider the special values $q = -e^{-2\theta}$, $\theta > 0$, which for $\sinh \theta = \frac{1}{2}$ gives $q = (1 - \sqrt{5})/(1 + \sqrt{5})$. We prove that $\mathcal{H}_n^{(\alpha)}(q)$ is unitarily related to the *generalized Hilbert matrix*

$$\mathcal{F}_n^{(\alpha)}(\theta) = (F_\alpha(\theta)/F_{l+j+\alpha}(\theta)), \quad 0 \leq l, j \leq n, \quad n = 0, 1, \dots \quad (15)$$

in the sense that

$$\mathcal{H}_n^{(\alpha)}(q) = U_n \mathcal{F}_n^{(\alpha)}(\theta) U_n, \quad (16)$$

where U_n is a unitary diagonal matrix with diagonal elements $i^l, l = 0, 1, \dots, n$. This makes it possible to deduce Richardson's formula for the elements of the inverse of \mathcal{F}_n and its generalizations $\mathcal{F}_n^{(\alpha)}(\theta)$ from the q -analogue of Choi's formula.

2 Quantum Hilbert matrices

The little q -Jacobi polynomials are given by

$$p_n(x; a, b; q) = {}_2\phi_1 \left(\begin{matrix} q^{-n}, abq^{n+1} \\ aq \end{matrix}; q, xq \right), \quad (17)$$

where a, b are complex parameters satisfying $|a|, |b| \leq 1$.

In [7, Section 7.3] one finds a discussion of the little q -Jacobi polynomials defined in (17), and it is proved that

$$\sum_{k=0}^{\infty} p_n(q^k; a, b; q) p_m(q^k; a, b; q) \frac{(bq; q)_k (aq)^k}{(q; q)_k} = \frac{\delta_{n,m}}{h_n(a, b; q)}, \quad (18)$$

where

$$h_n(a, b; q) = \frac{(abq; q)_n (1 - abq^{2n+1}) (aq; q)_n (aq; q)_\infty (aq)^{-n}}{(q; q)_n (1 - abq) (bq; q)_n (abq^2; q)_\infty}. \quad (19)$$

In [7] it is assumed that $0 < q, aq < 1$, but the derivation shows that it holds for $0 < |q| < 1, |a| \leq 1, |b| \leq 1$, in particular in the case of interest here: $a = q^{\alpha-1}, \alpha \in \mathbb{N}, b = 1$, in the case of which we get

$$\sum_{k=0}^{\infty} p_n(q^k; q^{\alpha-1}, 1; q) p_m(q^k; q^{\alpha-1}, 1; q) q^{\alpha k} = \delta_{n,m} \frac{q^{\alpha n} (q; q)_n^2}{(q^\alpha; q)_n^2 (1 - q^{2n+\alpha})}. \quad (20)$$

The Gaussian q -binomial coefficients

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = \frac{(q; q)_n}{(q; q)_k (q; q)_{n-k}}$$

are polynomials in q . Using the quantum integers (9) we get

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = \prod_{j=1}^k \frac{[n-j+1]_q q^{(n-j)/2}}{[j]_q q^{(j-1)/2}} = q^{k(n-k)/2} \prod_{j=1}^k \frac{[n-j+1]_q}{[j]_q},$$

and introducing the notation

$$[n]_q! = \prod_{k=1}^n [k]_q, \quad [0]_q! = 1,$$

we define the *quantum binomial coefficients* by

$$\binom{n}{k}_q = \prod_{j=1}^k \frac{[n-j+1]_q}{[j]_q} = \frac{[n]_q!}{[k]_q! [n-k]_q!}, \quad (21)$$

hence

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = q^{k(n-k)/2} \binom{n}{k}_q. \quad (22)$$

This formula shows that $\binom{n}{k}_q$ is a holomorphic function of q in the cut plane $\mathbb{C} \setminus]-\infty, 0]$ with a continuous extension to the upper part of the cut.

Defining

$$p_n^{(\alpha)}(q; x) := \binom{n+\alpha-1}{n}_q p_n(x/q^{1/2}; q^{\alpha-1}, 1; q), \quad (23)$$

some calculation leads to the simple expression

$$p_n^{(\alpha)}(q; x) = \sum_{j=0}^n \binom{n}{j}_q \binom{n+j+\alpha-1}{n}_q (-1)^j x^j, \quad (24)$$

which is a q -analogue of the polynomials $r_n^{(\alpha)}$ of [4].

The equation (20) can be written

$$\int p_n^{(\alpha)}(q; x) p_m^{(\alpha)}(q; x) d\mu^{(\alpha)}(q)(x) = \delta_{n,m} \frac{[\alpha]_q}{[2n+\alpha]_q} \quad (25)$$

showing that

$$P_n^{(\alpha)}(q; x) = ([2n+\alpha]_q / [\alpha]_q)^{1/2} p_n^{(\alpha)}(q; x),$$

are orthonormal polynomials. The corresponding kernel polynomials are defined by

$$K_n^{(\alpha)}(q; x, y) = \sum_{k=0}^n P_k^{(\alpha)}(q; x) P_k^{(\alpha)}(q; y) = \sum_{k=0}^n ([2k+\alpha]_q / [\alpha]_q) p_k^{(\alpha)}(q; x) p_k^{(\alpha)}(q; y). \quad (26)$$

While $P_n^{(\alpha)}(q; x)$ depends on the choice of a square root, the kernel polynomials $K_n^{(\alpha)}(q; x, y)$ are independent of this choice. Since the orthogonal polynomials exist with respect to $\mu^{(\alpha)}(q)$ and the moments are given by (12), it follows by [5, Theorem 3.1] that $\mathcal{H}_n^{(\alpha)}(q)$ is regular for each n . For $q \rightarrow 1$ and $\alpha = 1$ the expressions (24)–(26) tend to classical formulas for Legendre polynomials for $[0, 1]$.

Writing

$$K_n^{(\alpha)}(q; x, y) = \sum_{l,j=0}^n a_{l,j}^{(n)}(q; \alpha) x^l y^j,$$

it follows by (24) that the coefficients $a_{l,j}^{(n)}(q; \alpha)$ are given by

$$a_{l,j}^{(n)}(q; \alpha) = (-1)^{l+j} \sum_{k=\max(l,j)}^n \frac{[2k+\alpha]_q}{[\alpha]_q} \binom{k}{l}_q \binom{k}{j}_q \binom{k+l+\alpha-1}{k}_q \binom{k+j+\alpha-1}{k}_q. \quad (27)$$

Theorem 2.1 *The l, j 'th element of the inverse matrix of the generalized quantum Hilbert matrix $\mathcal{H}_n^{(\alpha)}(q)$ defined in (14) is given as*

$$(-1)^{l+j} \frac{[l+j+\alpha]_q}{[\alpha]_q} \binom{n+l+\alpha}{n-j}_q \binom{n+j+\alpha}{n-l}_q \binom{l+j+\alpha-1}{l}_q \binom{l+j+\alpha-1}{j}_q. \quad (28)$$

Furthermore,

$$\det \mathcal{H}_n^{(\alpha)}(q) = [\alpha]_q^n \left(\prod_{k=1}^n [2k + \alpha]_q \binom{2k+\alpha-1}{k}_q \right)^{-1}. \quad (29)$$

Proof. It is a general fact that the coefficients $a_{l,j}^{(n)}(q)$ of the kernel polynomial are the entries of the inverse of the Hankel matrix, cf. [4, Theorem 2.1].

Let $R(n; l, j)$ denote the number given in the right-hand side of (28), and define

$$C(k; l, j) = (-1)^{l+j} \frac{[2k+\alpha]_q}{[\alpha]_q} \binom{k}{l}_q \binom{k}{j}_q \binom{k+l+\alpha-1}{k}_q \binom{k+j+\alpha-1}{k}_q, \quad k \geq l, j.$$

We shall prove that

$$R(n; l, j) = \sum_{k=\max(l,j)}^n C(k; l, j) \quad (30)$$

by induction in n , and can assume $l \geq j$ without loss of generality. The equation (30) is easy for $n = k = l$ and is left to the reader. We shall establish the induction step

$$R(n+1; l, j) - R(n; l, j) = C(n+1; l, j). \quad (31)$$

The left-hand side of this expression can be written

$$(-1)^{l+j} \frac{[l+j+\alpha]_q}{[\alpha]_q} \binom{l+j+\alpha-1}{l}_q \binom{l+j+\alpha-1}{j}_q T,$$

where

$$\begin{aligned} T &= \binom{n+l+\alpha+1}{n+1-j}_q \binom{n+j+\alpha+1}{n+1-l}_q - \binom{n+l+\alpha}{n-j}_q \binom{n+j+\alpha}{n-l}_q \\ &= \frac{([n+l+\alpha]_q \cdots [l+j+\alpha+1]_q)([n+j+\alpha]_q \cdots [l+j+\alpha+1]_q)}{[n+1-j]_q! [n+1-l]_q!}. \end{aligned}$$

$$\{[n+l+\alpha+1]_q [n+j+\alpha+1]_q - [n+1-j]_q [n+1-l]_q\}.$$

The quantity in braces equals $[2n+2+\alpha]_q [l+j+\alpha]_q$, and now it is easy to complete the proof of (31).

From the general theory of orthogonal polynomials, cf. [1],[5],[9], it is known that the leading coefficient of the orthonormal polynomial $F_n^{(\alpha)}(q; x)$ is $\sqrt{D_{n-1}/D_n}$, where

$$D_n = \det \mathcal{H}_n^{(\alpha)}(q).$$

From (24) and (25) we then get

$$D_{n-1}/D_n = ([2n+\alpha]_q / [\alpha]_q) \binom{2n+\alpha-1}{n}_q^2,$$

hence

$$\frac{1}{D_n} = \prod_{k=1}^n \frac{D_{k-1}}{D_k} = \prod_{k=1}^n ([2k+\alpha]_q / [\alpha]_q) \binom{2k+\alpha-1}{k}_q^2,$$

and (29) follows. \square

Remark 2.2 By analytic continuation, the formulas (28) and (29) of Theorem 2.1 are valid for $q \neq 0$ which is not a root of unity of order $\leq 2n+\alpha$.

3 Filbert matrices

In this section we specialize to $q = -e^{-2\theta}$ for $\theta > 0$. It is easy to see that the unique solution $F_n(\theta), n \geq 0$ to (8) is given by

$$F_n(\theta) = \frac{e^{n\theta} - (-1)^n e^{-n\theta}}{e^\theta + e^{-\theta}}, \quad (32)$$

which is Ismail's definition in [10], hence

$$[n]_q = (-i)^{n-1} F_n(\theta). \quad (33)$$

For $\theta = \theta_0 > 0$ such that $\sinh \theta_0 = \frac{1}{2}$ we get $q = (1 - \sqrt{5}) / (1 + \sqrt{5})$ and $F_n = F_n(\theta_0)$. For information about Fibonacci numbers, see [11],[12]. Ismail [10] also considered the generalized Fibonomial coefficients

$$\binom{n}{k}_{\mathbb{F}(\theta)} = \prod_{j=1}^k \frac{F_{n-j+1}(\theta)}{F_j(\theta)}, \quad 0 \leq k \leq n, \quad (34)$$

with the usual convention that empty products are 1, and gave the following recursion

$$\binom{n}{k}_{\mathbb{F}(\theta)} = F_{k-1}(\theta) \binom{n-1}{k}_{\mathbb{F}(\theta)} + F_{n-k+1}(\theta) \binom{n-1}{k-1}_{\mathbb{F}(\theta)}, \quad n > k \geq 1, \quad (35)$$

which shows that they are integers when $2 \sinh \theta$ is an integer. Using (33) we next get that the quantum binomial coefficients can be expressed by the Fibonomial coefficients of (34) as

$$\binom{n}{k}_q = (-i)^{k(n-k)} \binom{n}{k}_{\mathbb{F}(\theta)}, \quad (36)$$

hence for $0 \leq l, j \leq n, \alpha \in \mathbb{N}$

$$\frac{[\alpha]_q}{[l+j+\alpha]_q} = i^{l+j} \frac{F_\alpha(\theta)}{F_{l+j+\alpha}(\theta)}. \quad (37)$$

Letting U_n denote the unitary diagonal $(n+1) \times (n+1)$ -matrix with l 'th diagonal element equal to $i^l, l = 0, 1, \dots, n$, then formula (37) implies

$$\mathcal{H}_n^{(\alpha)}(q) = U_n \mathcal{F}_n^{(\alpha)}(\theta) U_n, \quad \mathcal{F}_n^{(\alpha)}(\theta)^{-1} = U_n \mathcal{H}_n^{(\alpha)}(q)^{-1} U_n. \quad (38)$$

This leads to a new proof of Berg's and Ismail's generalizations of Richardson's formula, cf. [4], [10].

Theorem 3.1 *Let A be the matrix $(1/F_{l+j+\alpha}(\theta)), 0 \leq l, j \leq n$. Then A^{-1} has the entries*

$$(-1)^{n(l+j+\alpha) - \binom{l}{2} - \binom{j}{2}} F_{l+j+\alpha}(\theta) \binom{n+l+\alpha}{n-j}_{\mathbb{F}(\theta)} \binom{n+j+\alpha}{n-l}_{\mathbb{F}(\theta)} \binom{l+j+\alpha-1}{l}_{\mathbb{F}(\theta)} \binom{l+j+\alpha-1}{j}_{\mathbb{F}(\theta)}, \quad (39)$$

and

$$\det A = (-1)^{\alpha \binom{n+1}{2}} \left(F_\alpha(\theta) \prod_{k=1}^n F_{2k+\alpha}(\theta) \binom{2k+\alpha-1}{k}_{\mathbb{F}(\theta)}^2 \right)^{-1}. \quad (40)$$

Proof. We use

$$\det U_n = i^{\binom{n+1}{2}}, \quad \det \mathcal{H}_n^{(\alpha)}(q) = (-1)^{\binom{n+1}{2}} \det \mathcal{F}_n^{(\alpha)}(\theta)$$

and the formulas (33) and (36) to make the calculation, the only non-obvious thing being the sign in the two formulas. In the proof of (39) we get the following sign for the lj 'th element of A^{-1} :

$$\begin{aligned} & i^{l+j} (-1)^{l+j} (-i)^{l+j+(n-j)(l+j+\alpha)+(n-l)(l+j+\alpha)+l(j+\alpha-1)+j(l+\alpha-1)} \\ &= (-1)^{l+j} (-i)^{2n(l+j+\alpha)-l(l+1)-j(j+1)} = (-1)^{n(l+j+\alpha) - \binom{l}{2} - \binom{j}{2}}. \end{aligned}$$

In the proof of (40) we get the sign

$$\begin{aligned}
& (-1)^{\binom{n+1}{2}} (-i)^{(\alpha-1)n} \left(\prod_{k=1}^n (-i)^{2k+\alpha-1+2k(k+\alpha-1)} \right)^{-1} \\
&= (-1)^{\binom{n+1}{2}} \left(\prod_{k=1}^n (-i)^{2k(k+1)+2k(\alpha-1)} \right)^{-1} = (-1)^{\binom{n+1}{2}} \left(\prod_{k=1}^n (-1)^{k(\alpha-1)} \right)^{-1} \\
&= (-1)^{\alpha \binom{n+1}{2}}.
\end{aligned}$$

□

References

- [1] N. I. Akhiezer, *The classical moment problem*. Oliver and Boyd, Edinburgh, 1965.
- [2] G.E. Andrews, R. Askey and R. Roy, *Special functions*. Cambridge University Press, Cambridge 1999.
- [3] C. Berg, *Ortogonal polynomier og Hilbert matrixen*. (In Danish) *NORMAT* **54** no 3 (2006), 116–133.
- [4] C. Berg, *Fibonacci numbers and orthogonal polynomials*, *J. Comput. Appl. Math.* (To appear).
- [5] T. S. Chihara, *An introduction to orthogonal polynomials*. Gordon and Breach, New York-London-Paris, 1978.
- [6] Man-Duen Choi, *Tricks or Treats with the Hilbert Matrix*, *Amer. Math. Monthly* **90** (1983), 301–312.
- [7] G. Gasper and M. Rahman, *Basic hypergeometric series*. Cambridge University Press, Cambridge 1990, second edition 2004.
- [8] D. Hilbert, *Ein Beitrag zur Theorie des Legendreschen Polynoms*, *Acta Math.* **18** (1894), 155–159. (367–370 in “Gesammelte Abhandlungen II”, Berlin 1933.)
- [9] M. E. H. Ismail, *Classical and Quantum Orthogonal Polynomials in One Variable*. Cambridge University Press, Cambridge 2005.
- [10] M. E. H. Ismail, *One Parameter Generalizations of the Fibonacci and Lucas Numbers*. Preprint, August 2006.

- [11] D. E. Knuth, *The Art of Computer Programming*. Vol. 1, 2nd Ed., Addison-Wesley, 1973
- [12] T. Koshy, *Fibonacci and Lucas Numbers With Applications*. John Wiley, New York, 2001.
- [13] T. M. Richardson, *The Filbert matrix*, Fibonacci Quart. **39** no. 3 (2001), 268–275.

Jørgen Ellegaard Andersen, Department of Mathematics, University of Aarhus, Ny Munkegade, DK-8000 Aarhus C, Denmark. email: andersen@imf.au.dk

Christian Berg, Department of Mathematics, University of Copenhagen, Universitetsparken 5, DK 2100 Copenhagen Ø, Denmark. email: berg@math.ku.dk