# GOTZMANN IDEALS OF THE POLYNOMIAL RING

SATOSHI MURAI AND TAKAYUKI HIBI

ABSTRACT. Let  $A = K[x_1, \ldots, x_n]$  denote the polynomial ring in n variables over a field K. We will classify all the Gotzmann ideals of A with at most n generators. In addition, we will study Hilbert functions H for which all homogeneous ideals of A with the Hilbert function H have the same graded Betti numbers. These Hilbert functions will be called inflexible Hilbert functions. We introduce the notion of segmentwise critical Hilbert function and show that segmentwise critical Hilbert functions are inflexible.

#### 1. INTRODUCTION

Let K be an arbitrary field and let  $A = K[x_1, \ldots, x_n]$  denote the polynomial ring in n variables over K with each deg  $x_i = 1$ . For a homogeneous ideal I of A, the Hilbert function  $H(I, -) : \mathbb{Z}_{\geq 0} \to \mathbb{Z}_{\geq 0}$  of I is the numerical function defined by  $H(I, t) = \dim_K I_t$ , where  $I_t$  is the homogeneous component of degree t of I. Work with the lexicographic order  $<_{\text{lex}}$  on A induced by the ordering  $x_1 > x_2 > \cdots > x_n$ . Recall that a set V of monomials in A is said to be *lexsegment* if, for monomials u and v of A with  $u \in V$ , deg  $u = \deg v$  and  $u <_{\text{lex}} v$ , one has  $v \in V$ . Also, a monomial ideal I of A is called a *lexsegment ideal* if the set of monomials in I is lexsegment. Let I be a homogeneous ideal of A and  $I^{\text{lex}}$  the (unique) lexsegment ideal ([2] and [14]) with the same Hilbert function as I. A Gotzmann ideal introduced in [13] is a homogeneous ideal I for which the number of minimal generators of I is equal to that of  $I^{\text{lex}}$ . In this paper, we classify all the Gotzmann ideals of A generated by at most n homogeneous polynomials.

A numerical function  $H(-): \mathbb{Z}_{\geq 0} \to \mathbb{Z}_{\geq 0}$  is said to be *critical* [18] if it is equal to the Hilbert function of a lexsegment ideal of A which has at most n generators. Let  $1 \leq s \leq n$  and  $f_1, \ldots, f_s$  homogeneous polynomials with

$$f_i \in K[x_i, x_{i+1}, \dots, x_n]$$

for each  $1 \leq i \leq s$  and with deg  $f_s > 0$ . In [11] the ideal  $I_{(f_1,\ldots,f_s)}$  of A defined by

(1) 
$$I_{(f_1,\dots,f_s)} = (f_1x_1, f_1f_2x_2,\dots, f_1f_2\cdots f_{s-1}x_{s-1}, f_1f_2\cdots f_s)$$

was introduced. A homogeneous ideal I of A is called *canonical critical* if  $I = I_{(f_1,\ldots,f_s)}$  for some homogeneous polynomials  $f_1,\ldots,f_s$  with  $f_i \in K[x_i,x_{i+1},\ldots,x_n]$  for each  $1 \leq i \leq s$  and with deg  $f_s > 0$ , where  $1 \leq s \leq n$ . Our first result is

**Theorem 1.1.** Given a homogeneous ideal I of  $A = K[x_1, \ldots, x_n]$ , the following conditions are equivalent:

The first author is supported by JSPS Research Fellowships for Young Scientists.

- (i) I has a critical Hilbert function;
- (ii) there exists a linear transformation  $\varphi$  on A such that  $\varphi(I)$  is a canonical critical ideal;
- (iii) I is a Gotzmann ideal generated by at most n homogeneous polynomials.

It was shown in [13, Corollary 1.4] that a homogeneous ideal I is Gotzmann if and only if I and  $I^{\text{lex}}$  have the same graded Betti numbers. Thus Theorem 1.1 shows that if a homogeneous ideal has a critical Hilbert function H, then the graded Betti numbers are determined by H. In the second part of this paper, we generalize this fact. We introduce two classes of Hilbert functions H for which all ideals with the Hilbert function H have the same graded Betti numbers (Theorems 4.3 and 4.14).

We establish fundamental properties of critical functions and canonical critical ideals in Section 2, and give our proof of Theorem 1.1 in Section 3. Next, in Section 4, we introduce some Hilbert functions H such that all ideals with the Hilbert function H have the same graded Betti numbers. Finally, in the appendix, by using Theorem 1.1, we give a simple and purely algebraic proof of [11, Proposition 2].

## 2. Critical ideals

In this section, we establish fundamental properties of critical ideals. First, we recall basic facts on Gotzmann ideals and lexsegment ideals. Let K be a field and  $A = K[x_1, \ldots, x_n]$  the polynomial ring in n variables over a field K with each deg  $x_i = 1$ . Let  $d \ge 0$  be an integer and V a subspace of the K-vector space  $A_d$ . We write  $\text{Lex}(V) \subset A_d$  for the K-vector space spanned by the lexsegment set  $L \subset A_d$  of monomials with  $|L| = \dim_K V$ . A famous result of Macaulay (see e.g., [16, Corollary C4]) guarantees

(2) 
$$\dim_K(A_1 \cdot V) \ge \dim_K(A_1 \cdot \operatorname{Lex}(V)),$$

where  $A_1 \cdot V = \{ \ell g : \ell \in A_1 \text{ and } g \in V \}$ . A *K*-vector space *V* is called a *Gotzmann* space if  $\dim_K(A_1 \cdot V) = \dim_K(A_1 \cdot \operatorname{Lex}(V))$ . A homogeneous ideal *I* of *A* is said to be *Gotzmann* if  $I_k$  is Gotzmann for all  $k \geq 0$ . For any homogeneous ideal  $I \subset A$ , let  $I^{\operatorname{lex}} = \bigoplus_{k\geq 0} \operatorname{Lex}(I_k)$ . It follows from (2) that  $I^{\operatorname{lex}}$  is indeed a lexsegment ideal having the same Hilbert function as *I*. Clearly a homogeneous ideal *I* of *A* is Gotzmann if and only if the number of minimal generators of *I* is equal to that of  $I^{\operatorname{lex}}$ .

A monomial ideal I of A is said to be universal lexsegment [1] if, for any integer  $m \ge 0$ , the ideal of  $K[x_1, \ldots, x_{n+m}]$  having the same generators as I is a lexsegment ideal of  $K[x_1, \ldots, x_{n+m}]$ . It was shown in [18, Corollary 1.3] that a monomial ideal I of A is universal lexsegment if and only if I is lexsegment and  $|G(I)| \le n$ , where G(I) is the set of the minimal monomial generators of I. Moreover, it is known [18, Proposition 1.2] that a monomial ideal I of A is universal lexsegment if and only if, for some  $1 \le s \le n$  and for some nonnegative integers  $b_1, b_2, \ldots, b_s$ , one has

(3) 
$$I = (x_1^{b_1+1}, x_1^{b_1} x_2^{b_2+1}, \dots, x_1^{b_1} \cdots x_{s-1}^{b_{s-1}} x_s^{b_s+1}).$$

It follows from [18, Proposition 1.5] that the Hilbert function H(I, t) of the universal lexsegment ideal (3) is given by

(4) 
$$H(I,t) = \binom{t-a_1+n-1}{n-1} + \dots + \binom{t-a_s+n-s}{n-s},$$

where the sequence  $(a_1, a_2, \ldots, a_s)$  with

$$0 < a_1 \le a_2 \le \dots \le a_s$$

is defined by setting

$$a_i = \deg x_1^{b_1} \cdots x_{i-1}^{b_{i-1}} x_i^{b_i+1}, \quad 1 \le i \le s$$

or equivalently,

$$b_i = a_i - a_{i-1}, \qquad 1 \le i \le s$$

with  $a_0 = 1$ .

Since a lexsegment ideal with a given Hilbert function is uniquely determined, it follows that a numerical function  $H: \mathbb{Z}_{\geq 0} \to \mathbb{Z}_{\geq 0}$  is critical if and only if there exists a sequence  $(a_1, \ldots, a_s)$  of integers with  $0 < a_1 \leq a_2 \leq \cdots \leq a_s$ , where  $1 \leq s \leq n$ , such that H(t) is written in the form (4). A critical function of the form (4) will be called a critical function of type  $(a_1, a_2, \ldots, a_s)$ . For convenience, we say that a homogeneous ideal of A is a *critical ideal of type*  $(a_1, a_2, \ldots, a_s)$  if its Hilbert function is the critical function of type  $(a_1, a_2, \ldots, a_s)$ .

**Lemma 2.1.** Let  $1 < s \leq n$ . Fix homogeneous polynomials  $f_1, \ldots, f_{s-1}$  with each  $f_i \in K[x_i, \ldots, x_n]$ . Let  $g \in K[x_s, \ldots, x_n]$  be a homogeneous polynomial with deg g >0. Then

(5) 
$$f_1 f_2 \cdots f_{s-1} g \notin (f_1 x_1, f_1 f_2 x_2, \dots, f_1 f_2 \cdots f_{s-1} x_{s-1}).$$

*Proof.* One has (5) if and only if

(6) 
$$f_2 \cdots f_{s-1}g \notin (x_1, f_2 x_2, \dots, f_2 \cdots f_{s-1} x_{s-1}).$$

Since  $f_2 \cdots f_{s-1}g \in K[x_2, \ldots, x_n]$ , it follows that (6) is equivalent to saying

(7) 
$$f_2 \cdots f_{s-1}g \notin (f_2 x_2, \dots, f_2 \cdots f_{s-1} x_{s-1}).$$

Now, working with induction on 
$$s$$
, the desired result (7) is guaranteed.

**Lemma 2.2.** As a vector space over K the ideal (1) is the direct sum

(8) 
$$I_{(f_1,\dots,f_s)} = \left(\bigoplus_{j=1}^{s-1} (f_1\cdots f_j x_j) K[x_j,\dots,x_n]\right) \bigoplus (f_1\cdots f_s) K[x_s,\dots,x_n].$$

*Proof.* Let 
$$J = (f_1 x_1, f_1 f_2 x_2, \dots, f_1 f_2 \cdots f_{s-1} x_{s-1})$$
. Let  $g = f_{s-1} x_{s-1}$ . Then  
 $J = (f_1 x_1, \dots, f_1 f_2 \cdots f_{s-2} x_{s-2}, f_1 f_2 \cdots f_{s-2} g)$ .

Now, working with induction on s yields

$$J = \left(\bigoplus_{j=1}^{s-2} (f_1 \cdots f_j x_j) K[x_j, \dots, x_n]\right)_3 \bigoplus (f_1 f_2 \cdots f_{s-2} g) K[x_{s-1}, \dots, x_n].$$

In other words, one has

$$J = \bigoplus_{j=1}^{s-1} (f_1 \cdots f_j x_j) K[x_j, \dots, x_n].$$

On the other hand, by using Lemma 2.1, it follows that

$$(f_1 \cdots f_{s-1} f_s) K[x_s, \ldots, x_n] \bigcap J = \{0\}.$$

Clearly,  $f_1 \cdots f_{s-1} f_s x_j \in J$  for each j < s. Hence (8) follows.

**Corollary 2.3.** Let  $I_{(f_1,\ldots,f_s)}$  denote the ideal (1).

- (a)  $I_{(f_1,\ldots,f_s)}$  is a critical ideal of type  $(a_1,\ldots,a_s)$ , where  $a_i = \deg f_1 f_2 \cdots f_i x_i$ ,  $i = 1, \ldots, s 1$ , and  $a_s = \deg f_1 f_2 \cdots f_s$ .
- (b)  $I_{(f_1,\ldots,f_s)}$  is minimally generated by

(9) 
$$\{f_1x_1, \ldots, f_1f_2 \cdots f_{s-1}x_{s-1}, f_1f_2 \cdots f_s\}.$$

(c) 
$$I_{(f_1,\ldots,f_s)}$$
 is Gotzmann.

Proof. The direct sum decomposition (8) says that the Hilbert function of  $I_{(f_1,\ldots,f_s)}$  is of the form (4) and, in addition, that  $I_{(f_1,\ldots,f_s)}$  is minimally generated by (9). Thus (a) and (b) follow. Since the lexsegment ideal with the Hilbert function (4) is the universal lexsegment ideal (3), one has  $|G((I_{(f_1,\ldots,f_s)})^{\text{lex}})| = s$ . Thus  $I_{(f_1,\ldots,f_s)}$  is Gotzmann by (b).

If u is a monomial of A, then we write m(u) for the largest integer j for which  $x_j$  divides u. A monomial ideal I of A is called *stable* if, for each monomial u belonging to G(I) and for each  $1 \leq i < m(u)$ , one has  $(x_i u)/x_{m(u)} \in I$ .

**Lemma 2.4.** Let I be a stable ideal of A with  $G(I) = \{u_1, \ldots, u_s\}$ . Then the Hilbert function of I is

(10) 
$$H(I,t) = \sum_{j=1}^{s} \binom{t - \deg u_j + n - m(u_j)}{n - m(u_j)}.$$

*Proof.* By virtue of [8, Lemma 1.1] every monomial  $v \in I$  can be uniquely expressed of the form v = uw, where u and w are monomial of A with  $u \in G(I)$  and  $w \in K[x_{m(u)}, \ldots, x_n]$ . It then follows that

$$I = \bigoplus_{j=1}^{\circ} u_j K[x_{m(u_j)}, \dots, x_n].$$

Thus the desired formula (10) follows.

**Lemma 2.5.** A monomial ideal I of A which is both critical and stable is universal lexsegment.

*Proof.* Suppose that I is a critical ideal of type  $(a_1, \ldots, a_s)$ . Let  $a_0 = 1$  and  $b_i = a_i - a_{i-1} \ge 0$  for  $i = 1, 2, \ldots, s$ . What we must prove is that I coincides with the

lexsegment ideal (3). Since both I and the lexsegment ideal (3) have the Hilbert function (4), our claim follows from

$$\{x_1^{b_1+1}, x_1^{b_1}x_2^{b_2+1}, \dots, x_1^{b_1}\cdots x_{s-1}^{b_{s-1}}x_s^{b_s+1}\} \subset G(I).$$

Since  $H(I, a_1 - 1) = 0$  and  $H(I, a_1) > 0$  and since I is stable, it follows that  $x_1^{a_1} = x_1^{b_1+1} \in G(I)$ . Let  $1 \le p < s$  and

$$Q = \{x_1^{b_1+1}, x_1^{b_1} x_2^{b_2+1}, \dots, x_1^{b_1} \cdots x_{p-1}^{b_{p-1}} x_p^{b_p+1}\}.$$

Suppose that  $Q \subset G(J)$ . Our work is to show that  $x_1^{b_1} \cdots x_p^{b_p} x_{p+1}^{b_{p+1}+1}$  belongs to G(I).

We claim that each monomial  $u \in G(I) \setminus Q$  satisfies m(u) > p. Suppose that there is  $w \in G(I) \setminus Q$  with  $m(w) \leq p$ . Since I is stable, Lemma 2.4 says that

$$H(I,t) \ge \sum_{j=1}^{p} {\binom{t-a_j+n-j}{n-j}} + {\binom{t-\deg w+n-p}{n-p}}.$$

Since the Hilbert function H(I,t) is equal to (4), it follows that

(11) 
$$\sum_{j=p+1}^{s} \binom{t-a_j+n-j}{n-j} \ge \binom{t-\deg w+n-p}{n-p}$$

for all t. However, for  $t \gg 0$ , the right-hand side of (11) is a polynomial on t of degree n-p and the left-hand side of (11) is that of degree at most n-p-1. Hence the inequalities (11) cannot be valid for  $t \gg 0$ . This completes the proof of our claim that each monomial  $u \in G(I) \setminus Q$  satisfies m(u) > p.

Let J be the universal lexsegment ideal of A with G(J) = Q. Since H(I, t) =H(J,t) for  $t < a_{p+1}$  and  $H(I, a_{p+1}) > H(J, a_{p+1})$ , it follows that there is a monomial belonging to  $G(I) \setminus Q$  of degree  $a_{p+1}$  and that each monomial of A of degree  $a_{p+1}$ belonging to  $I \setminus J$  must belong to G(I).

Let  $u = x_1^{c_1} x_2^{c_2} \cdots x_n^{c_n}$  be a monomial belonging to  $G(I) \setminus Q$  of degree  $a_{p+1}$ . Then

(12) 
$$c_1 \ge b_1, \dots, c_{p-1} \ge b_{p-1}$$

To see why (12) is true, suppose that there is  $1 \leq i < p$  with  $c_i < b_i$ . Since I is stable, one has

$$v = x_1^{c_1} x_2^{c_2} \cdots x_i^{c_i} x_{i+1}^{a_{p+1} - \sum_{j=1}^i c_j} \in I.$$

Since  $u \notin J$ , it follows that the monomial  $x_1^{c_1} x_2^{c_2} \cdots x_i^{c_i}$  can be divided by none of the monomials  $x_1^{b_1} \cdots x_{j-1}^{b_{j-1}} x_j^{b_j+1}$  with  $1 \leq j \leq i$ . Since  $c_i < b_i$ , the monomial v can be divided by none of the monomials  $x_1^{b_1} \cdots x_{j-1}^{b_{j-1}} x_j^{b_j+1}$  with  $i < j \leq p$ . Hence  $v \notin J$ . Since deg  $v = a_{p+1}$ , one has  $v \in G(I)$ . Since  $v \in G(I) \setminus Q$ , the claim stated above says that m(v) > p, a contradiction. This completes the proof of (12).

On the other hand, if there is  $1 \leq i \leq p$  with  $c_i > b_i$ , then u is divided by  $x_1^{b_1} \cdots x_{i-1}^{b_{i-1}} x_i^{b_i+1}$ , a contradiction. Hence

$$c_1 = b_1, \dots, c_{p-1} = b_{p-1}, c_p \le b_p.$$

Since I is stable with deg  $w = a_{p+1}$ , the monomial

$$w = x_1^{b_1} \cdots x_{p-1}^{b_{p-1}} x_p^{b_p} x_{p+1}^{b_{p+1}+1}$$

must belong to I. Since  $w \notin J$ , it follows that  $w \in G(I)$ , as desired.

Let I be a homogeneous ideal of A. When K is infinite, given a monomial order  $\sigma$  on A, we write  $gin_{\sigma}(I)$  for the generic initial ideal ([7] and [12]) of I with respect to  $\sigma$ .

**Lemma 2.6.** Suppose that K is infinite. Let I be a critical ideal of A. Then, for an arbitrary monomial order  $\sigma$  on A induced by the ordering  $x_1 > \cdots > x_n$ , the generic initial ideal  $gin_{\sigma}(I)$  is stable. Thus in particular  $gin_{\sigma}(I)$  is universal lexsegment.

Proof. Since both I and  $\operatorname{gin}_{\sigma}(I)$  have the same Hilbert function, it follows that  $\operatorname{gin}_{\sigma}(I)$  is critical. Since  $\operatorname{gin}_{\sigma}(I)$  is a monomial ideal, it follows from [18, Corollary 1.7] that  $\operatorname{gin}_{\sigma}(I)$  is Gotzmann. Thus, by Gotzmann's persistence theorem (see Lemma 4.5),  $\operatorname{gin}_{\sigma}(I)$  is componentwise linear [13]. Then [5, Lemma 1.4] says that  $\operatorname{gin}_{\operatorname{crev}}(\operatorname{gin}_{\sigma}(I)) = \operatorname{gin}_{\sigma}(I)$  is stable. Here  $\operatorname{crev}$  is the reverse lexicographic order on A induced by the ordering  $x_1 > \cdots > x_n$ . Since  $\operatorname{gin}_{\sigma}(I)$  is both critical and stable, it follows from Lemma 2.5 that  $\operatorname{gin}_{\sigma}(I)$  is universal lexsegment.  $\Box$ 

Note that if char(K) = 0 then Lemma 2.6 is obvious since generic initial ideals are stable in characteristic 0.

**Lemma 2.7.** Suppose that a homogeneous ideal I of A is a critical ideal of type  $(a_1, \ldots, a_s)$ , where  $2 \leq s \leq n$ . Then there exists a homogeneous polynomial f of A with deg  $f = a_1 - 1$  together with a homogeneous ideal J of A such that

 $I = f \cdot J.$ 

*Proof.* To prove the statement, by considering an extension field, we may assume that the base field K is infinite. Then there is a linear transformation  $\varphi$  with  $\ln_{\leq_{\text{lex}}}(\varphi(I)) = \min_{\leq_{\text{lex}}}(I)$ . Considering  $\varphi(I)$  instead of I, one may assume that  $\ln_{\leq_{\text{lex}}}(I) = \min_{\leq_{\text{lex}}}(I)$ . Lemma 2.6 says that  $\ln_{\leq_{\text{lex}}}(I)$  is universal lexsegment. Hence

$$\operatorname{in}_{<_{\operatorname{lex}}}(I) = (x_1^{b_1+1}, x_1^{b_1} x_2^{b_2+1}, \dots, x_1^{b_1} \cdots x_{s-1}^{b_{s-1}} x_s^{b_s+1}).$$

where  $b_i = a_i - a_{i-1}, 1 \le i \le s$ , with  $a_0 = 1$ .

To simplify the notation, let  $u_i = x_1^{b_1} \cdots x_i^{b_i}$  for  $i = 1, \ldots, s$ . Thus  $\operatorname{in}_{<_{\operatorname{lex}}}(I) = (u_1 x_1, \ldots, u_s x_s)$ . Let  $\mathcal{G} = \{g_1, \ldots, g_s\}$  be a Gröbner basis of I, where  $g_i$  is a homogeneous polynomial of A with  $\operatorname{in}_{<_{\operatorname{lex}}}(g_i) = u_i x_i$  for each  $1 \leq i \leq s$ , and  $\mathcal{G}' = \{g_2, \ldots, g_s\}$ . We show that  $\mathcal{G}'$  is a Gröbner basis with respect to  $<_{\operatorname{lex}}$ . Let  $2 \leq i < j \leq s$  and divide the S-polynomial of  $g_i$  and  $g_j$  by  $\mathcal{G}$ , say,

$$(u_j/u_i)x_jg_i - x_ig_j = p_1g_1 + \dots + p_sg_s,$$

where  $p_1, \ldots, p_s$  are homogeneous polynomials of A with

$$\operatorname{in}_{\leq_{\operatorname{lex}}}(p_k g_k) \leq_{\operatorname{lex}} \operatorname{in}_{\leq_{\operatorname{lex}}}((u_j/u_i)x_j g_i - x_i g_j)$$

for each  $1 \le k \le s$ . Since

$$\inf_{\leq_{\text{lex}}}(p_1g_1) = \inf_{\leq_{\text{lex}}}(p_1)x_1^{b_1+1} <_{\text{lex}} \inf_{\substack{\leq_{\text{lex}}\\6}}(x_ig_j) = x_ix_ju_j = x_ix_j(x_1^{b_1}\cdots x_j^{b_j}),$$

it follows that  $p_1 = 0$ . In other words, a remainder of the *S*-polynomial of  $g_i$  and  $g_j$  with respect to  $\mathcal{G}'$  can be 0. Hence  $\mathcal{G}'$  is a Gröbner basis with respect to  $<_{\text{lex}}$ , as desired.

Now, we prove Lemma 2.7 by using induction on s. Let s = 2. Again, divide the S-polynomial of  $g_1$  and  $g_2$  by  $\mathcal{G}$ , say,

$$x_2^{b_2+1}g_1 - x_1g_2 = p_1g_1 + p_2g_2,$$

where  $p_1$  and  $p_2$  are homogeneous with deg  $p_1 = b_2 + 1$  and deg  $p_2 = 1$ , and where each of  $p_1$  and  $p_2$  satisfies

$$\operatorname{in}_{\leq_{\operatorname{lex}}}(p_k g_k) \leq_{\operatorname{lex}} \operatorname{in}_{\leq_{\operatorname{lex}}}(x_2^{b_2+1}g_1 - x_1g_2).$$

One has  $(x_2^{b_2+1} - p_1)g_1 = (x_1 + p_2)g_2$ . Since  $\ln_{<_{lex}}(x_2^{b_2+1} - p_1) = x_2^{b_2+1}$  and since  $\ln_{<_{lex}}(x_1 + p_2) = x_1$ , the polynomial  $x_1 + p_2$  must divide  $g_1$ . Let  $f = g_1/(x_1 + p_2)$ . Then deg  $f = a_1 - 1$  and  $I = f \cdot (x_1 + p_2, x_2^{b_2+1} - p_1)$ , as required.

Next, let s > 2 and write J for the ideal of A generated by  $\mathcal{G}'$ . Since

$$\operatorname{in}_{<_{\operatorname{lex}}}(J) = (u_2 x_2, \dots, u_s x_s) = x_1^{b_1} (x_2^{b_2+1}, x_2^{b_2} x_3^{b_3+1}, \dots, x_2^{b_2} \cdots x_{s-1}^{b_{s-1}} x_s^{b_s+1})$$

and since

$$(x_2^{b_2+1}, x_2^{b_2} x_3^{b_3+1}, \dots, x_2^{b_2} \cdots x_{s-1}^{b_{s-1}} x_s^{b_s+1})$$

is universal lexsegment by a permutation of the variables, the ideal J is a critical ideal of type  $(a_2, \ldots, a_s)$ . The induction hypothesis guarantees the existence of a homogeneous polynomial  $f_0$  of A with  $\deg(f_0) = a_2 - 1$  which divides each of  $g_2, \ldots, g_s$ . Since  $\inf_{<_{\text{lex}}}(f_0)$  divides  $\inf_{<_{\text{lex}}}(g_i) = u_i x_i$  for each  $1 < i \leq s$ , one has  $\inf_{<_{\text{lex}}}(f_0) = u_2$ . Let  $g'_i = g_i/f_0$  for  $i = 2, \ldots, s$ . Thus in particular  $\inf_{<_{\text{lex}}}(g'_2) = u_2 x_2/u_2 = x_2$ .

Now, divide the S-polynomial of  $g_1$  and  $g_2$  by  $\mathcal{G}$ , say,

$$x_2^{b_2+1}g_1 - x_1(f_0g_2') = q_1g_1 + q_2(f_0g_2') + \dots + q_s(f_0g_s'),$$

where  $q_1, \ldots, q_s$  are homogeneous polynomials of A with

$$\operatorname{in}_{\leq_{\operatorname{lex}}}(q_1g_1) \leq_{\operatorname{lex}} \operatorname{in}_{\leq_{\operatorname{lex}}}(x_2^{b_2+1}g_1 - x_1(f_0g_2'))$$

and with

$$\operatorname{in}_{\leq_{\operatorname{lex}}}(q_k(f_0g'_k)) \leq_{\operatorname{lex}} \operatorname{in}_{\leq_{\operatorname{lex}}}(x_2^{b_2+1}g_1 - x_1(f_0g'_2))$$

for each  $2 \leq k \leq s$ . Let

$$f_0h = q_2(f_0g'_2) + \dots + q_s(f_0g'_s).$$

Thus

$$(x_2^{b_2+1} - q_1)g_1 = f_0(x_1g_2' + h).$$

Since  $\operatorname{in}_{\leq_{\operatorname{lex}}}(x_2^{b_2+1}-q_1) = x_2^{b_2+1}$ ,  $\operatorname{in}_{\leq_{\operatorname{lex}}}(g_1) = x_1^{b_1+1}$  and  $\operatorname{in}_{\leq_{\operatorname{lex}}}(x_1g_2'+h) = x_1x_2$ , it follows that  $x_1g_2'+h$  can divide neither  $x_2^{b_2+1}-q_1$  nor  $g_1$ . Thus  $x_1g_2'+h$  is a product  $(x_1+h_1)(x_2+h_2)$ , where  $h_1$  and  $h_2$  are homogeneous polynomials of A with  $\deg h_1 = \deg h_2 = 1$ , such that  $x_1 + h_1$  divides  $g_1$  and  $x_2 + h_2$  divides  $x_2^{b_2+1}-q_1$ . Let  $f = g_1/(x_1+h_1)$ . Then  $\deg f = a_1 - 1$  and f divides both  $g_1$  and  $f_0$ .  $\Box$ 

# 3. Proof of Theorem 1.1

We are now in the position to give a proof of Theorem 1.1.

Proof of Theorem 1.1. First, Corollary 2.3 guarantees (ii)  $\Rightarrow$  (iii). Second, (iii)  $\Rightarrow$  (i) is clear by the definition of Gotzmann ideals and that of critical functions.

On the other hand, a proof of (i)  $\Rightarrow$  (ii) will be achieved by induction on s. Let  $I \subset A$  be a critical ideal of type  $(a_1, \ldots, a_s)$ . Let s = 1. Let  $f_1$  be a homogeneous polynomial of degree  $a_1$  belonging to I. Then the Hilbert function of the ideal  $(f_1)$  of A coincides with that of I. Thus  $I = (f_1)$ , as desired.

Let s > 1. Lemma 2.7 guarantees that  $I = f \cdot J$ , where f is a homogeneous polynomial of A with deg  $f = a_1 - 1$  and where J is a homogeneous ideal of A. The Hilbert function of J is  $H(J,t) = H(I,t+a_1-1)$ . Hence J is a critical ideal of type $(1, a_2 - a_1 + 1, \ldots, a_s - a_1 + 1)$ . Since  $H(J,1) \neq 0$ , there exists a linear transformation  $\varphi$  on A with  $x_1 \in \varphi(J)$ . Let J' be the ideal

$$J' = \varphi(J) \cap K[x_2, \dots, x_n]$$

of  $K[x_2,\ldots,x_n]$ . Then

$$\varphi(J) = x_1 K[x_1, \dots, x_n] \bigoplus J'.$$

Since the Hilbert function of J' is

$$H(J',t) = H(\varphi(J),t) - H(x_1K[x_1,\ldots,x_n],t),$$

the ideal J' of  $K[x_2, \ldots, x_n]$  is a critical ideal of type  $(a_2 - a_1 + 1, \ldots, a_s - a_1 + 1)$ . The induction hypothesis then guarantees the existence of a linear transformation  $\psi$  on  $K[x_2, \ldots, x_n]$  such that  $\psi(J')$  is a canonical critical ideal of  $K[x_2, \ldots, x_n]$ , say

$$\psi(J') = (f_2 x_2, \dots, f_2 \cdots f_{s-1} x_{s-1}, f_2 \cdots f_s)$$

where  $f_i \in K[x_i, x_{i+1}, \ldots, x_n]$  for each  $2 \leq i \leq s$  and where deg  $f_s > 0$ . Now, regarding  $\psi$  to be a linear transformation on A by setting  $\psi(x_1) = x_1$ , one has

$$\begin{aligned} (\psi \circ \varphi)(I) &= ((\psi \circ \varphi)(f)) \cdot ((\psi \circ \varphi)(J)) \\ &= ((\psi \circ \varphi)(f)) \cdot (\psi(x_1 A \bigoplus J')) \\ &= (\psi \circ \varphi)(f) \cdot (x_1 A \bigoplus \psi(J')). \end{aligned}$$

Let  $f_1 = (\psi \circ \varphi)(f)$ . Then it follows that

$$(\psi \circ \varphi)(I) = (f_1 x_1, f_1 f_2 x_2, \dots, f_1 f_2 \cdots f_{s-1} x_{s-1}, f_1 f_2 \cdots f_s)$$

as desired.

**Remark 3.1.** In Lemma 2.6 we assume that the base field K is infinite. However, this assumption is not required in Theorem 1.1 since Lemma 2.6 is only required to prove Lemma 2.7 and since we may assume that K is infinite to prove Lemma 2.7.

**Example 3.2.** For a permutation  $\pi$  on  $[n] = \{1, \ldots, n\}$  and for a monomial  $u = x_1^{a_1} \cdots x_n^{a_n}$  of A, we set  $\pi(u) = x_{\pi(1)}^{a_1} \cdots x_{\pi(n)}^{a_n}$ . A monomial ideal I of A with  $G(I) = \{u_1, \ldots, u_s\}$  is called *trivially Gotzmann* if there is a permutation  $\pi$  on [n] such that the monomial ideal  $(\pi(u_1), \ldots, \pi(u_s))$  is lexsegment.

Given an arbitrary integer m > n, there exists a Gotzmann monomial ideal I of A with |G(I)| = m which cannot be trivially Gotzmann. In fact, let  $q \ge 2$  be an integer and I the monomial ideal of A generated by  $x_1^{q-1}x_2, x_1^{q-2}x_2^2, \ldots, x_1x_2^{q-1}$  together with  $x_3, \ldots, x_n$ . Then I is a Gotzmann ideal with |G(I)| = n + q - 3 which cannot be trivially Gotzmann.

#### 4. INFLEXIBLE HILBERT FUNCTIONS

Throughout this section, we assume that  $A = K[x_1, \ldots, x_n]$  is the polynomial ring over an infinite field K with each deg  $x_i = 1$ . The graded Betti numbers of a homogeneous ideal I are the integers  $\beta_{i,j}(I) = \dim_K \operatorname{Tor}_i(I, K)_j$ . In other words, the graded Betti numbers appear in the minimal graded free resolution

$$\cdots \longrightarrow \bigoplus_{j} S(-j)^{\beta_{2,j}(I)} \longrightarrow \bigoplus_{j} S(-j)^{\beta_{1,j}(I)} \longrightarrow \bigoplus_{j} S(-j)^{\beta_{0,j}(I)} \longrightarrow I \longrightarrow 0$$

of I over A. By the famous result of Bigatti [2], Hulett [14] and Pardue [19], there exists an ideal having the largest graded Betti numbers among all ideals for a fixed Hilbert function. However, an ideal having the smallest graded Betti numbers among all ideals for a fixed Hilbert function need not exist. Recently, existence and non-existence of the smallest graded Betti numbers of ideals for a fixed Hilbert function are studied in several papers (see e.g., [4, 6, 21]). An extremal example of a Hilbert function having an ideal with the smallest graded Betti numbers is a Hilbert function H for which all ideals with the Hilbert function H have the same graded Betti numbers. Let  $H : \mathbb{Z}_{\geq 0} \to \mathbb{Z}_{\geq 0}$  be the Hilbert function of a homogeneous ideal of A. We say that H is an *inflexible Hilbert function of* A if all homogeneous ideals of A with the Hilbert function H have the same graded Betti numbers. It is known [13, Corollary 1.4] that H is an inflexible Hilbert function of A if and only if all ideals of A with the Hilbert function H are Gotzmann. Thus, in particular, Theorem 1.1 shows that critical functions are inflexible Hilbert functions. In this section, we introduce some more inflexible Hilbert functions.

First, we recall fundamental properties on Macaulay representations and the minimal growth of Hilbert functions. Given positive integers a and d, there exists the unique representation of a, called the *d*-th Macaulay representation of a, of the form

(13) 
$$a = \binom{a(d)+d}{d} + \dots + \binom{a(k)+k}{k},$$

where  $k \ge 1$  and where  $a(d) \ge \cdots \ge a(k) \ge 0$ . We recall the following easy fact (see [3, Lemma 4.2.7]).

**Lemma 4.1.** Let  $a_d \geq \cdots \geq a_1 \geq -1$  and  $b_d \geq \cdots \geq b_1 \geq -1$  be integers. Then

$$\binom{a_d+d}{d} + \dots + \binom{a_1+1}{1} > \binom{b_d+d}{d} + \dots + \binom{b_1+1}{1}$$

if and only if  $(a_d, \ldots, a_1) >_{\text{lex}} (b_d, \ldots, b_1)$ .

By using the d-th Macaulay representation (13) of a positive integer a, one defines

$$a^{\mathrm{MG}(d)} = \binom{a(d)+d+1}{d} + \dots + \binom{a(k)+k+1}{k}$$

and  $0^{MG(d)} = 0$ . This number is convenient to describe the minimal growth of Hilbert functions of homogeneous ideals. Let  $L \subset A_d$  be a K-vector space spanned by a lexsegment set of monomials of degree d. Let u be the minimal monomial in L with respect to  $\leq_{\text{lex}}$ . Then u can be written in the form

$$u = x_1^{a_0 - 1} x_2^{a_1 - a_0} \cdots x_k^{a_{k-1} - a_{k-2}} x_{k+1}^{a_k - a_{k-1}} x_n^{d - a_k},$$

where  $0 < a_0 \leq a_1 \leq \cdots \leq a_k \leq d$  and  $0 \leq k \leq n-2$ . Then it follows from [16, Lemma C.10] that

$$\dim_{K} L = \binom{d - a_0 + n - 1}{n - 1} + \dots + \binom{d - a_k + n - 1 - k}{n - 1 - k}$$

is the (n-1)-th Macaulay representation of  $\dim_K L$ . On the other hand, since  $A_1 \cdot L$  is also lexsegment and since the minimal element of  $A_1 \cdot L$  w.r.t.  $<_{\text{lex}}$  is  $x_n u$ , one has

(14) 
$$\dim_K(A_1 \cdot L) = (\dim_K L)^{\mathrm{MG}(n-1)}.$$

Hence by (2) any homogeneous ideal I of A satisfies

$$H(I, k+1) \ge H(I, k)^{\mathrm{MG}(n-1)} \quad \text{for all } k \ge 0.$$

Suppose that H is the Hilbert function of a homogeneous ideal of A. Peeva [20, Corollary 1.4] proved that, any numerical function H satisfying that, for all  $k \ge 1$ , if  $H(k)^{MG(n-1)} < H(k+1)$ , then  $H(k+1)^{MG(n-1)} = H(k+2)$ , is an inflexible Hilbert function. If n = 2, those functions essentially characterize inflexible Hilbert functions of A. Indeed, it is not hard to see that, in case of n = 2, H is an inflexible Hilbert function of A if and only if H satisfies that, for all  $k \ge 1$ , if  $H(k)^{MG(n-1)} < H(k+1)$  and  $H(k) \ne 1$ , then  $H(k+1)^{MG(n-1)} = H(k+2)$ . See [15] for further results on graded Betti numbers of homogeneous ideals of  $K[x_1, x_2]$  for a fixed Hilbert function. Peeva's result and Theorem 1.1 lead us to consider the following numerical functions.

**Definition 4.2.** A numerical function  $H : \mathbb{Z}_{\geq 0} \to \mathbb{Z}_{\geq 0}$  is said to be *segmentwise* critical if there exist integers  $0 = s_0 < s_1 < \cdots < s_\ell = \infty$  such that

- (i) for each  $j = 0, 1, ..., \ell 1$ , there exists a critical function  $H_j$  with  $H(t) = H_j(t)$  for all  $s_j \le t \le s_{j+1}$ ;
- (ii)  $H_j(t)$  for all  $s_j \le t \le s_{j+1}$ ; (ii)  $H(s_{j+1}-1)^{\mathrm{MG}(n-1)} = H(s_{j+1})$  for  $j = 0, 1, \dots, \ell - 2$ .

Let  $d \ge 0$  be an integer. We say that a subspace V of  $A_d$  is a *critical space* if there exists a critical ideal I of A such that  $V = I_d$ . Note that critical spaces are Gotzmann by Theorem 1.1. For any homogeneous ideal I of A, we write  $gin(I) = gin_{<_{rev}}(I)$ . The goal of this section is to prove the following result.

**Theorem 4.3.** If I has a segmentwise critical Hilbert function then  $I_k$  is a critical space for all  $k \ge 0$ . In particular, I is Gotzmann and gin(I) is lexsegment.

To prove Theorem 4.3, we first study critical spaces. For a homogeneous ideal Iof A, the ideal

$$I^{\text{sat}} = \bigcup_{k \ge 0} (I : (x_1, \dots, x_n)^k)$$

is called the saturation of I. The (Castelnuovo-Mumford) regularity of a homogeneous ideal I is the integer

$$\operatorname{reg}(I) = \max\{j : \beta_{i,i+j}(I) \neq 0 \text{ for some } i\}.$$

The following facts are known (see  $[12, \S2]$ ).

**Lemma 4.4.** Let I be a homogeneous ideal of A.

- (i)  $I_k^{\text{sat}} = I_k \text{ for all } k \ge \text{reg}(I);$ (ii)  $gin(I^{\text{sat}}) = \bigcup_{k\ge 0} (gin(I): x_n^k) = (gin(I))^{\text{sat}};$
- (iii)  $(Bayer-Stillman) \operatorname{reg}(I) = \operatorname{reg}(\operatorname{gin}(I)).$

Note that the second equation of (ii) follows from [7, Proposition 15.24].

A homogeneous ideal I is said to have a *d*-linear resolution if it is generated in degree d and reg(I) = d. The following result of Gotzmann is called Gotzmann's persistence theorem (see  $[3, \S4.3]$  and [12] for the proof).

**Lemma 4.5** (Gotzmann's persistence theorem [9]). Let  $V \subset A_d$  be a Gotzmann space and I the ideal generated by V. Then

- (i)  $A_1 \cdot V$  is Gotzmann;
- (ii) I has a d-linear resolution.

Recall that a homogeneous ideal  $I \subset A$  is saturated if  $I : (x_1, \ldots, x_n) = I$ . Note that a homogeneous ideal I of A is saturated if and only if the depth of A/I is positive (see [3, §1.2]). The next result immediately follows from [18, Corollary 1.4].

**Lemma 4.6.** A lexsegment ideal I of A is saturated if and only if  $|G(I)| \le n-1$ .

Let V be a subspace of  $A_d$  and I the ideal of A generated by V. Let gin(V) be the generic initial space of V, in other words,

$$gin(V) = gin(I)_d.$$

**Lemma 4.7.** Let V be a subspace of  $A_d$  and I the ideal of A generated by V. If I has a d-linear resolution and gin(V) = Lex(V), then

(i)  $I^{\text{sat}}$  is a critical ideal generated by at most n-1 homogeneous polynomials;

(ii) V is a critical space.

*Proof.* By the assumption and Lemma 4.4 (iii), it follows that gin(I) has a d-linear resolution. Thus gin(I) is generated by  $gin(I)_d = gin(V) = Lex(V)$ . Hence gin(I) is lexsegment. It is clear that the saturation of a lexsegment ideal is again a lexsegment ideal. Thus, by Lemma 4.6, the ideal  $gin(I)^{sat}$  is a universal lexisegment ideal with  $|G(gin(I)^{sat})| \leq n-1$ . Since  $gin(I^{sat}) = gin(I)^{sat}$  by Lemma 4.4 (ii), the Hilbert function of  $I^{\text{sat}}$  is equal to that of the universal lexsegment ideal  $gin(I)^{\text{sat}}$ . Hence  $I^{\text{sat}}$  is a critical ideal. Also since the number of generators of  $I^{\text{sat}}$  is smaller than or equal to that of  $gin(I^{sat}) = gin(I)^{sat}$ , the ideal  $I^{sat}$  is generated by at most n-1 homogeneous polynomials. Finally, since  $\operatorname{reg}(I) = d$ , we have  $V = I_d = (I^{\operatorname{sat}})_d$  by Lemma 4.4 (i), and therefore V is a critical space.

Lemma 4.7 enable us to characterize critical spaces in terms of generic initial spaces.

**Corollary 4.8.** A subspace V of  $A_d$  is a critical space if and only if V is Gotzmann and gin(V) = Lex(V).

*Proof.* If V is a critical space then V is Gotzmann and gin(V) = Lex(V) by Theorem 1.1 and Lemma 2.6. On the other hand, by Lemmas 4.5 and 4.7, if V is Gotzmann and gin(V) = Lex(V), then V is a critical space.

**Example 4.9.** Let  $A = K[x_1, x_2, x_3]$ ,  $I = (x_1^3, x_1^2 x_2, x_1 x_2^2, x_2^3, x_1^2 x_3, x_1 x_2 x_3, x_2^2 x_3)$ and  $V = I_3$ . It is easy to see that V is a Gotzmann space. However, V is not a critical space. Indeed,  $gin(V) = V \neq Lex(V)$  and  $I^{sat} = (x_1^2, x_1 x_2, x_2^2)$  does not have a critical Hilbert function. On the other hand, for any  $f \in A$ , the ideal  $J = (fx_1^2, fx_1 x_2, fx_1 x_3, f(x_2 + x_3) x_2, f(x_2 + x_3) x_3)$  is generated by a critical space. Indeed,  $J^{sat} = (fx_1, f(x_2 + x_3))$  is a canonical critical ideal and J is generated by the homogeneous component of  $J^{sat}$  of degree deg f + 2. Note also that these critical spaces are parametrized by some Hilbert schemes (see the appendix).

By Lemma 4.7, if  $V \subset A_d$  is a critical space, then the saturation of the ideal I generated by V is critical. We know the type of this critical ideal by using the Macaulay representation of  $\dim_K V$ .

**Lemma 4.10.** Let  $V \subset A_d$  be a critical space and I the ideal of A generated by V. Let  $\dim_K V = \sum_{j=1}^p {\binom{d-a_p+n-j}{n-j}}$  be the (n-1)-th Macaulay representation of  $\dim_K V$ . Then  $I^{\text{sat}}$  is a critical ideal of type  $(a_1, \ldots, a_p)$ .

*Proof.* Suppose that  $I^{\text{sat}}$  is a critical ideal of type  $(b_1, \ldots, b_q)$ . What we must prove is  $(a_1, \ldots, a_p) = (b_1, \ldots, b_q)$ .

By Theorem 1.1 and Corollary 2.3, the type of a critical ideal is equal to the sequence of degrees of its minimal generators. Thus, by Lemma 4.7, we have  $q \leq n-1$ . In particular,

$$H(I^{\text{sat}},k) = \sum_{j=1}^{q} \binom{k-b_j+n-j}{n-j}$$

is the (n-1)-th Macaulay representation of  $H(I^{\text{sat}}, k)$  for  $k \geq b_q$ . On the other hand, (14) and Gotzmann's persistence theorem imply

$$H(I,k) = \sum_{j=1}^{p} \binom{k-a_j+n-j}{n-j} \quad \text{for all } k \ge d.$$

Since the above equation is the (n-1)-th Macaulay representation of H(I,k) for all  $k \ge d$  and since  $H(I,k) = H(I^{\text{sat}},k)$  for  $k \gg 0$ , it follows from Lemma 4.1 that  $(a_1,\ldots,a_p) = (b_1,\ldots,b_q).$ 

Next, we study the relation between critical functions and Macaulay representations. Let H be the critical function of type  $(a_1, \ldots, a_p)$ . If  $p \leq n-1$  then (4) coincides with the (n-1)-th Macaulay representation. On the other hand, in case of p = n, the (n-1)-th Macaulay representation of H(k) is given by the following formula.

**Lemma 4.11.** Let  $a_0 = 0 < a_1 \le \dots \le a_n \le d$  be integers and let  $a = \sum_{j=1}^n {\binom{d-a_j+n-j}{n-j}}$ . Set  $s = \min\{k : a_k = a_{n-1}, k \le n-1\}$ . Then

$$a = \left\{ \sum_{j=1}^{s-1} \binom{d-a_j+n-j}{n-j} \right\} + \binom{t-(a_s-1)+n-s}{n-s}$$

is the (n-1)-th Macaulay representation of a.

*Proof.* The statement immediately follows from the next equation

$$\binom{t-a_s+n-s}{n-s} + \dots + \binom{t-a_s+1}{1} + \binom{t-a_n+0}{0} = \binom{t-(a_s-1)+n-s}{n-s}.$$

For 
$$\mathbf{a} = (a_1, \dots, a_p) \in \mathbb{Z}_{>0}^p$$
, where  $0 < a_1 \le \dots \le a_p$  and where  $1 \le p \le n$ , write  $\bar{\mathbf{a}} = \begin{cases} (a_1, \dots, a_p), & \text{if } p < n, \\ (a_1, \dots, a_{s-1}, a_s - 1), & \text{if } p = n, \end{cases}$ 

where  $s = \min\{k : a_k = a_{n-1}, k \le n-1\}$ . We also require the following fact, which immediately follows from Gotzmann's persistence theorem.

**Lemma 4.12.** Let I be a homogeneous ideal of A. If  $H(I,k)^{MG(n-1)} = H(I,k+1)$ , then  $I_k$  and  $I_{k+1}$  are Gotzmann spaces.

*Proof.* Clearly  $I_k$  is Gotzmann and I has no generators of degree k + 1. Hence  $I_{k+1} = A_1 \cdot I_k$  is also a Gotzmann space by Gotzmann's persistence theorem. 

Now we prove Theorem 4.3.

Proof of Theorem 4.3. For any integer  $k \geq 0$ , we write  $I_{\langle k \rangle}$  for the ideal generated by all polynomials in I of degree k. Let  $0 = s_0 < s_1 < \cdots < s_\ell = \infty$  be integers satisfying the conditions of Definition 4.2. Let

$$I[i] = (I_{\langle s_i \rangle})^{\operatorname{sat}} + I_{\langle s_i+1 \rangle} + \dots + I_{\langle s_{i+1} \rangle} \quad \text{for } i = 0, 1, \dots, \ell - 1.$$

We show that I[i] is a critical ideal with  $(I[i])_k = I_k$  for all  $s_i \leq k \leq s_{i+1}$ . Since  $I_0$  is a critical space, by using induction on i, it is enough to show that if  $I_{s_i}$  is a critical space, then I[i] is a critical ideal with  $(I[i])_k = I_k$  for all  $s_i \leq k \leq s_{i+1}$ .

First, we show  $(I[i])_k = I_k$  for all  $s_i \leq k \leq s_{i+1}$ . Since  $I_{s_i}$  is a critical space by the assumption,  $I_{s_i}$  is a Gotzmann space. Thus, by Lemma 4.5 (ii), we have  $\operatorname{reg}(I_{\langle s_i \rangle}) = s_i$ . Hence, by Lemma 4.4 (i), we have

(15) 
$$((I_{\langle s_i \rangle})^{\operatorname{sat}})_k = (I_{\langle s_i \rangle})_k \subset I_k \quad \text{for all } k \ge s_i.$$

Then it is clear that  $(I[i])_k = I_k$  for all  $s_i \leq k \leq s_{i+1}$ .

Next, we show that I[i] has a critical Hilbert function. Suppose that, for  $k = s_i, s_i + 1, \ldots, s_{i+1}, H(I, k)$  is equal to the critical function of type  $\mathbf{a} = (a_1, \ldots, a_p)$ . Thus

(16) 
$$H(I[i],k) = H(I,k) = \sum_{j=1}^{p} \binom{k-a_j+n-j}{n-j}$$
 for  $s_i \le k \le s_{i+1}$ .

We may assume  $a_p \leq s_{i+1}$  since  $\binom{k-a_p+n-p}{n-p} = 0$  if  $k < a_p$ . Set  $\bar{\mathbf{a}} = (b_1, \ldots, b_q)$ . We claim that

(17) 
$$H(I[i],k) = \sum_{j=1}^{q} \binom{k-b_j+n-j}{n-j} = \sum_{j=1}^{p} \binom{k-a_j+n-j}{n-j}$$
 for all  $k \ge s_{i+1}$ .

If  $i = \ell - 1$  then there is nothing to prove. Suppose  $i < \ell - 1$ . Since  $a_p \leq s_{i+1}$ , it follows from Lemma 4.11 and (16) that

$$\dim_K I_{s_{i+1}} = H(I, s_{i+1}) = \sum_{j=1}^q \binom{s_{i+1} - b_j + n - j}{n - j}$$

is the (n-1)-th Macaulay representation of  $\dim_K I_{s_{i+1}}$ . On the other hand, by Lemma 4.12 and condition (ii) of Definition 4.2, it follows that  $I_{s_{i+1}}$  is a Gotzmann space. Hence by Gotzmann's persistence theorem we have

$$H(I_{\langle s_{i+1} \rangle}, k) = \sum_{j=1}^{q} \binom{k - b_j + n - j}{n - j} \quad \text{for all } k \ge s_{i+1}$$

Then the desired equation (17) follows from Lemma 4.11, since (15) says that  $I[i]_k = (I_{\langle s_{i+1} \rangle})_k$  for  $k \geq s_{i+1}$ .

Now we show that I[i] has a critical Hilbert function.

Case 1: Suppose  $a_p \leq s_i$ . We show that I[i] has the critical Hilbert function of type  $(b_1, \ldots, b_q)$ . Since  $a_p \leq s_i$ , Lemma 4.11 together with (16) and (17) implies

(18) 
$$H(I[i],k) = \sum_{j=1}^{q} \binom{k-b_j+n-j}{n-j} \quad \text{for all } k \ge s_i$$

In particular,

$$\dim_{K} I_{s_{i}} = H(I[i], s_{i}) = \sum_{j=1}^{q} \binom{s_{i} - b_{j} + n - j}{n - j}$$

is the (n-1)-th Macaulay representation of  $\dim_K I_{s_i}$ . Thus, by Lemma 4.10, one has

$$H(I[i], k) = H((I_{\langle s_i \rangle})^{\text{sat}}, k) = \sum_{j=1}^{q} \binom{k - b_j + n - j}{n - j} \quad \text{for } k \le s_i.$$

Then the above equation together with (18) implies that the Hilbert function of I[i] is the critical function of type  $(b_1, \ldots, b_q)$ .

Case 2: Suppose  $a_p > s_i$ . We show that I[i] has the critical Hilbert function of type  $(a_1, \ldots, a_p)$ . Let  $\ell = \min\{k : a_k > s_i\} - 1$ . Then, by (16),

$$\dim_{K} I_{s_{i}} = H(I, s_{i}) = \sum_{j=1}^{\ell} \binom{s_{i} - a_{j} + n - j}{n - j}$$

is the (n-1)-th Macaulay representation of  $\dim_K I_{s_i}$ . Thus, by Lemma 4.10, we have

$$H(I[i],k) = H((I_{\langle s_i \rangle})^{\text{sat}},k) = \sum_{j=1}^{\ell} \binom{k-a_j+n-j}{n-j} \quad \text{for } k \le s_i.$$

Then the above equation together with (16) and (17) says that the Hilbert function of I[i] is the critical function of type  $(a_1, \ldots, a_p)$ . 

**Example 4.13.** Unfortunately, we cannot expect a simple canonical formula like Theorem 1.1 for homogeneous ideals having a segmentwise critical Hilbert function. If I has a segmentwise critical Hilbert function, then each  $I_k$  is a critical space by Theorem 4.3. However, there does not necessarily exist a linear transformation  $\varphi$ on S such that  $\varphi(I)_k$  is a canonical critical space for all k. Let

$$I = (x_1^3, x_1^2 x_2, x_1^2 x_3^2, x_1 (x_2^4 + x_3^4) x_3) \subset \mathbb{Q}[x_1, x_2, x_3].$$

Let  $J = (x_1^3, x_1^2 x_2, x_1^2 x_3^2)$  and  $J' = (x_1^2, x_1(x_2^4 + x_3^4)x_3)$ . Then J and J' are critical ideals and  $I_k = J_k$  for  $k \leq 5$  and  $I_k = J'_k$  for  $k \geq 5$ . Hence I has a segmentwise critical Hilbert function. However, for any linear transformation  $\varphi$  on S, if  $\varphi(I)_4$  is a canonical critical space, then  $\varphi(I)_6$  is not a canonical critical space.

Finally, we generalize Peeva's result [20, Corollary 1.4]. For the (n-1)-th Macaulay representation  $a = \sum_{j=1}^{s} {\binom{a(j)+n-j}{n-j}}$  of a positive integer a > 0, define

$$a^{+} = \begin{cases} a^{\mathrm{MG}(n-1)}, & \text{if } s \ge n-2, \\ a^{\mathrm{MG}(n-1)} + \binom{n-s-1}{n-s-1} + \dots + \binom{2}{2}, & \text{if } s < n-2. \end{cases}$$

**Theorem 4.14.** Let I be a homogeneous ideal satisfying, for all  $k \geq 1$ ,

if 
$$H(I,k)^{\mathrm{MG}(n-1)} < H(I,k+1)$$
 then  $H(I,k+2) \le H(I,k+1)^+$ .

Then I is Gotzmann.

*Proof.* We show that  $I_k$  is Gotzmann for all k by using induction on k. It is clear

that  $I_k$  is Gotzmann for  $k \gg 0$ . We show if  $I_{k+1}$  is Gotzmann then  $I_k$  is Gotzmann. Let  $H(I,k) = \sum_{j=1}^{s} {\binom{k-a_j+n-j}{n-j}}$  be the (n-1)-th Macaulay representation of H(I,k). If  $H(I,k)^{\mathrm{MG}(n-1)} = H(I,k+1)$  or  $H(I,k-1)^{\mathrm{MG}(n-1)} = H(I,k)$  then  $I_k$ is Gotzmann by Lemma 4.12. Thus we may assume  $H(I,k)^{MG(n-1)} < H(I,k+1)$ and  $H(I, k-1)^{MG(n-1)} < H(I, k)$ . Then, by the assumption, we have s < n-2 and  $H(I, k+1) \leq H(I, k)^+$ . Hence

$$H(I, k+1) = \sum_{j=1}^{s} \binom{k+1-a_j+n-j}{n-j} + \binom{n-s-1}{n-s-1} + \dots + \binom{n-\ell}{n-\ell}$$

for some  $\ell \leq n-2$ . Since  $I_{k+1}$  is Gotzmann by the induction hypothesis, Gotzmann's persistence theorem implies

$$H(I_{\langle k+1 \rangle}, t) = \sum_{j=1}^{s} \binom{t-a_j+n-j}{n-j} + \sum_{j=s+1}^{\ell} \binom{t-(k+1)+n-j}{n-j} \text{ for all } t \ge k+1.$$

Since  $n - \ell \geq 2$ , by Proposition A.1, which we will show later in the appendix, it follows that  $(I_{\langle k+1 \rangle})^{\text{sat}}$  is a critical ideal of type  $(a_1, \ldots, a_s, k+1, \ldots, k+1)$ . Then, since  $H(I,k) = \sum_{j=1}^{s} {\binom{k-a_j+n-j}{n-j}} = H((I_{\langle k+1 \rangle})^{\text{sat}}, k)$  and  $I_k \subset (I_{\langle k+1 \rangle})^{\text{sat}}$ , we have  $I_k = ((I_{\langle k+1 \rangle})^{\text{sat}})_k$ . Hence  $I_k$  is a critical space, and therefore  $I_k$  is Gotzmann as desired.  $\Box$ 

**Example 4.15.** By Macaulay's theorem, there is a one-to-one correspondence between Hilbert functions of homogeneous ideals of A and lexsegment ideals of A. Let  $I = (u_1, u_2, \ldots, u_t)$  be a lexsegment ideal. Set  $d_k = \deg u_k$  for  $k = 1, 2, \ldots, t$ . Suppose  $d_1 \leq d_2 \leq \cdots \leq d_t$ . If  $A = K[x_1, x_2]$ , then segmentwise critical Hilbert function corresponds to lexsegment ideal I satisfying  $d_2 < d_3 - 1 < d_4 - 2 < \cdots < d_t - t + 2$ and Hilbert functions satisfying the condition of Theorem 4.14 correspond to lexsegment ideals I satisfying  $d_1 = \cdots = d_{s_1} < d_{s_1+1} - 1 = \cdots = d_{s_2} - 1 < d_{s_2+1} - 2 = \cdots$ , where  $1 \leq s_1 < s_2 < \cdots$ . Note that if n = 2 then Theorem 4.14 is equivalent to Peeva's result since  $a^+ = a^{MG(n-1)}$ .

**Remark 4.16.** We can construct new inflexible Hilbert functions by using Theorems 4.3 and 4.14. Let d > 0 be an integer and H a segmentwise critical function such that  $H(k)^{\mathrm{MG}(n-1)} = H(k+1)$  for  $k \geq d$ . Let H' be a Hilbert function of a homogeneous ideal of A satisfying the condition of Theorem 4.14. If H(d) = H'(d) and H(d+1) = H'(d+1), then we define the new Hilbert function  $\tilde{H}$  by  $\tilde{H}(k) = H(k)$  for  $k \leq d$  and  $\tilde{H}(k) = H'(k)$  for  $k \geq d$ . This new function  $\tilde{H}$  is an inflexible Hilbert function.

Indeed, if a homogeneous ideal I of A has the Hilbert function H, then, by the same way as in the proof of Theorem 4.14, it follows that  $I_k$  is a Gotzmann space for  $k \ge d$ . On the other hand, by Gotzmann's persistence theorem, the ideal  $J = \sum_{k=0}^{d} I_{\langle k \rangle}$  has the Hilbert function H. Since H is a segmentwise critical function,  $I_k = J_k$  is a Gotzmann space for  $k \le d$ .

**Example 4.17.** Let d > 0 be an integer and let  $I = (x_1^d) + (x_1, \ldots, x_n)^{d+1} \subset A$ . Then all ideals of A having the same Hilbert function as I are Gotzmann. However, the Hilbert function of I is not segmentwise critical and does not satisfy the condition of Theorem 4.14.

### Appendix

Let I be a homogeneous ideal of  $A = K[x_1, \ldots, x_n]$  and write  $P_I(t)$  for the Hilbert polynomial of I. It follows from (14) that  $P_I(t)$  can be written uniquely in the form

(19) 
$$P_{I}(t) = \binom{t - a_{1} + n - 1}{n - 1} + \dots + \binom{t - a_{s} + n - s}{n - s},$$

where  $1 \le s \le n-1$  and where  $a_1, \ldots, a_s$  are integers with  $0 < a_1 \le \cdots \le a_s$ .

Gotzmann [11, Proposition 2] proved using the languages of algebraic geometry that if a homogeneous ideal I of A is saturated and if the Hilbert polynomial of Iof the form (19) satisfies  $n - s \ge 2$ , then there exists a linear transformation  $\varphi$  on A such that  $\varphi(I)$  is a canonical critical ideal. Gotzmann [10, 11] also show that the Hilbert scheme parametrizing ideals with Hilbert polynomial (19), which satisfies either (i)  $n - s \le 2$  or (ii) s = n - 1 and  $a_1 - a_n \le 1$ , is irreducible and simply connected, with a single Borel-fixed point. On the other hand, Mall [17] gave a combinatorial proof of the simple connectivity of these Hilbert schemes by showing that they have a single Borel-fixed point.

Theorem 1.1 together with Proposition A.1 stated below yields a simple and purely algebraic proof of [11, Proposition 2].

**Proposition A.1** (Gotzmann). Let I be a saturated ideal of A with the Hilbert polynomial (19) and suppose that one of the following conditions is satisfied:

(i)  $n - s \ge 2;$ 

(ii) 
$$s = n - 1$$
 and  $a_{n-1} - a_1 \le 1$ .

Then the Hilbert function of I is the critical function of type  $(a_1, a_2, \ldots, a_s)$ .

Note that a universal lexsegment ideal whose Hilbert function is a critical function of type  $(a_1, \ldots, a_s)$  with  $n - s \ge 2$  has at most n - 2 generators. In particular, by (3) such a universal lexsegment ideal has no generators which are divisible by  $x_{n-1}$ .

Proposition A.1 was proved in [10] but we include a proof for the sake of completeness. For the *d*-th Macaulay representation  $a = \sum_{j=k}^{d} {a(j)+j \choose j}$  of a positive integer a > 0, let

$$a_{\langle d \rangle} = \begin{pmatrix} a(d) + d - 1 \\ d \end{pmatrix} + \dots + \begin{pmatrix} a(k) + k - 1 \\ k \end{pmatrix}$$

and  $0_{\langle d \rangle} = 0$ . Then, by (2) and (14), any homogeneous ideal I of S satisfies

(20) 
$$H(I, k+1)_{\langle n-1 \rangle} \ge H(I, k), \quad k = 0, 1, 2, \dots$$

Proof of Proposition A.1. We define functions  $H : \mathbb{Z}_{\geq 0} \to \mathbb{Z}_{\geq 0}$  and  $H' : \mathbb{Z}_{\geq 0} \to \mathbb{Z}_{\geq 0}$ by setting

$$H(t) = \binom{t - a_1 + n - 1}{n - 1} + \dots + \binom{t - a_s + n - s}{n - s}$$

and

(21) 
$$H'(t) = \binom{t - a_1 + n - 2}{n - 2} + \dots + \binom{t - a_s + n - s - 1}{n - s - 1}.$$

Then a simple computation yields

(22) 
$$H'(t) = H(t) - H(t-1)$$
 for all  $t \ge 0$ .

By considering an extension field, we may assume that K is infinite. Then, since  $I : (x_1, \ldots, x_n) = I$ , it follows that the quotient algebra A/I possesses a homogeneous non-zero-divisor  $\theta$  with deg  $\theta = 1$ . Let  $B = K[y_1, \ldots, y_{n-1}]$  denote the polynomial ring in n-1 variables over K. Regarding  $(A/I)/\theta(A/I)$  as a quotient algebra of B by a homogeneous ideal J, a simple computation yields that

$$H(B/J,t) = H(A/I,t) - H(A/I,t-1)$$
 for all  $t \ge 0$ .

Then, since H(B,t) + H(A,t-1) = H(A,t), we have

(23) 
$$H(J,d) = H(I,t) - H(I,t-1)$$
 for all  $t \ge 0$ .

Since the Hilbert polynomial  $P_I(t)$  of I is (19), it follows that  $H(I,t) = P_I(t) = H(t)$  for all  $t \gg 0$ . Then we have H(J,t) = H'(t) for all  $t \gg 0$  by (22) and (23). We claim that if either (i) or (ii) is satisfied then

(24) 
$$H(J,t) \le H'(t) \quad \text{for all } t \ge 0.$$

Case 1: Suppose  $n - s \ge 2$ . Then (21) coincides with the (n - 2)-th Macaulay representation of H'(t). Thus  $H'(t+1)_{(n-2)} = H'(t)$  for all  $t \ge 0$ . Now, if  $H(J, t_0 + 1) \le H'(t_0 + 1)$  for some  $t_0$ , then it follows from (20) that

$$H(J, t_0) \le H(J, t_0 + 1)_{\langle n-2 \rangle} \le H'(t_0 + 1)_{\langle n-2 \rangle} = H'(t_0).$$

(The second inequality follows from Lemma 4.1.) Hence  $H(J, t_0) \leq H'(t_0)$ . Consequently, we have  $H(J, t) \leq H'(t)$  for all t as desired.

Case 2: Suppose that s = n-1,  $a_{n-1} - a_1 \le 1$  and  $a_p = a_{p+1} = \cdots = a_{n-1}$ , where either  $a_{p-1} < a_p$  or p = 1. Then it follows from Lemma 4.11 that

$$H'(t) = \sum_{k=1}^{p-1} {\binom{t-a_k+n-1-k}{n-1-k}} + {\binom{t-(a_p-1)+n-1-p}{n-1-p}}$$

is the (n-2)-th Macaulay representation of H'(t) for all  $t \ge a_p$ . Thus  $H'(t+1)_{\langle n-2 \rangle} = H'(t)$  for all  $t \ge a_p$ . Then, by the same way as in Case 1, we have  $H(J,t) \le H'(t)$  for all  $t \ge a_p - 1$ . On the other hand, since  $H(J, a_p - 1) \le H'(a_p - 1) < n - 1$ , it follows that H(J,t) = 0 for  $t < a_p - 1$ . Thus the desired inequality (24) follows.

Now (22), (23) and (24) imply

(25) 
$$H(I,t) = \sum_{k=1}^{t} H(J,t) \le \sum_{k=1}^{t} H'(t) = H(t) \text{ for all } t \ge 0.$$

Since  $H(I,t) = P_I(t) = H(t)$  for all  $t \gg 0$ , (25) implies H(J,t) = H'(t) for all  $t \ge 0$ . Hence, by (25), we have H(I,t) = H(t) for all  $t \ge 0$ . Thus the Hilbert function of I is the critical function of type  $(a_1, \ldots, a_p)$  as desired.  $\Box$ 

**Remark A.2.** If *I* has a critical Hilbert function of type  $(a_1, \ldots, a_p)$  with  $p \le n-1$ , then the Hilbert polynomial of *I* is given by the formula (19). However, this does not imply  $H(I,k) = P_I(k)$  for all  $k \ge 0$ . Indeed, if n = 3, p = 1 and  $a_1 = 4$  then  $P_I(t) = \binom{t-2}{2} = \frac{(t-2)(t-3)}{2}$ . Thus  $P_I(1) = 1$ , however,  $H(I,1) = \binom{-1}{2} = 0$ .

**Remark A.3.** Proposition A.1 is false if neither (i) nor (ii) is satisfied. Indeed, it follows from [17, Theorem 26] that if a sequence of integers  $0 < a_1 \leq \cdots \leq a_s$  with  $1 \leq s \leq n-1$  satisfies neither (i) nor (ii), then there exists a saturated non-lexsegment Borel-fixed ideal with the Hilbert polynomial (19). The Hilbert function of such an ideal is not critical by Lemma 2.6.

Acknowledgments: We would like to thank the referee for suggesting the study of segmentwise critical functions as well as for many helpful comments to improve the paper.

## References

- E. Babson, I. Novik and R. Thomas, Reverse lexicographic and lexicographic shifting, J. Algebraic Combin. 23 (2006), 107–123.
- [2] A. M. Bigatti, Upper bounds for the Betti numbers of a given Hilbert function, Comm. Algebra 21 (1993), 2317–2334.
- [3] W. Bruns and J. Herzog, "Cohen-Macaulay Rings," Revised Edition, Cambridge University Press, Cambridge, 1996.
- [4] H. Charalambous and E.G. Evans, Resolutions with a given Hilbert function, in "Commutative Algebra: Syzygies, Multiplicities, and Birational Algebra," (W.J. Heinzer, C. Huneke and J.D. Sally Eds.), Contemp. Math., vol 159, Amer. Math. Soc., Providence, 1994, pp. 19–26.
- [5] A. Conca, J. Herzog and T. Hibi, Rigid resolutions and big Betti numbers, Comment. Math. Helv. 79 (2004), 826–839.
- [6] C. Dodd, A. Marks, V. Meyerson and B. Richert, Minimal Betti numbers, Comm. Algebra 35 (2007), 759–772.
- [7] D. Eisenbud, "Commutative Algebra. With a View Toward Algebraic Geometry," Springer-Verlag, New York, 1995.
- [8] S. Eliahou and M. Kervaire, Minimal resolutions of some monomial ideals, J. Algebra 129 (1990), 1–25.
- [9] G. Gotzmann, Eine Bedingung f
  ür die Flachheit und das Hilbertpolynom eines graduierten Ringes, Math. Z. 158 (1978), 61–70.
- [10] G. Gotzmann, Einige einfach-zusammenhängende Hilbertschemata, Math. Z. 180 (1982), 291– 305.
- [11] G. Gotzmann, Some irreducible Hilbert schemes, Math. Z. 201 (1989), 13–17.
- [12] M. Green, Generic initial ideals, in "Six Lectures on Commutative Algebra" (J. Elias, J. M. Giral, R. M. Miró-Roig and S. Zarzuela, Eds.), Birhäuser, Basel, 1998, pp. 119–186.
- [13] J. Herzog and T. Hibi, Componentwise linear ideals, Nagoya Math. J. 153 (1999), 141–153.
- [14] H. A. Hulett, Maximum Betti numbers of homogeneous ideals with a given Hilbert function, Comm. Algebra 21 (1993), 2335–2350.
- [15] A. Iarrobino, Betti strata of height two ideals, J. Algebra 285 (2005), 835–855.
- [16] A. Iarrobino and S. L. Kleiman, The Gotzmann theorems and the Hilbert scheme, in A. Iarrobino and V Kanev, "Power Sums, Gorenstein Algebras, and Determinantal Loci," Springer-Verlag, Berlin, 1999, pp. 289–312.
- [17] D. Mall, Characterisations of lexicographic sets and simply-connected Hilbert schemes, in "Applied algebra, algebraic algorithms and error-correcting codes" (T. Mora and H. Mattson Eds.), Lecture Notes in Comput. Sci., vol. 1255, Springer, Berlin, 1977, pp. 221–236.
- [18] S. Murai and T. Hibi, The depth of an ideal with a given Hilbert function, arXiv:math.AC/0608188, Proc. Amer. Math. Soc., to appear.
- [19] K. Pardue, Deformation classes of graded modules and maximal Betti numbers, *Illinois J. Math.* 40 (1996), 564–585.
- [20] I. Peeva, Consecutive cancellations in Betti numbers, Proc. Amer. Math. Soc. 132 (2004), 3503–3507.
- [21] B. Richert, Smallest graded Betti numbers, J. Algebra 244 (2001), 236–259.

SATOSHI MURAI, DEPARTMENT OF PURE AND APPLIED MATHEMATICS, GRADUATE SCHOOL OF INFORMATION SCIENCE AND TECHNOLOGY, OSAKA UNIVERSITY, TOYONAKA, OSAKA 560-0043, JAPAN

*E-mail address*: s-murai@ist.osaka-u.ac.jp

TAKAYUKI HIBI, DEPARTMENT OF PURE AND APPLIED MATHEMATICS, GRADUATE SCHOOL OF INFORMATION SCIENCE AND TECHNOLOGY, OSAKA UNIVERSITY, TOYONAKA, OSAKA 560-0043, JAPAN

*E-mail address*: hibi@math.sci.osaka-u.ac.jp