Curvature semipositivity of relative pluricanonical systems

Hajime TSUJI*

March 28, 2007

Abstract

We prove semipositivity of the curvature of the Narashimhan-Simha metric on an arbitrary flat projective family of varieties with only canonical singularities. This also implies that the direct image of pluricanonical systems have natural continuous hermitian strucutres with semipositive curvature curvature currents in the sense of Nakano. This paper is a revised version of [T4] using [T5, T6]. MSC: 14J15,14J40, 32J18

Contents

1	Inti	roduction	2
2	Pre	reliminaries	
	2.1	Canonical singularities	3
	2.2	Singular hermitian metrics	4
	2.3	Lelong numbers	5
	2.4	Analytic Zariski decompositions (AZD)	7
3	Narashimhan-Simha metrics		8
	3.1	Narashimhan-Simha metrics	8
	3.2	Dependence of Narashimhan-Simha metrics on families	9
	3.3	Positivity of Narashimhan-Simha metrics on projective families $% \left(1\right) =\left(1\right) +\left($	10
4	Canonical AZD		13
	4.1	Construction of the canonical AZD	14
	4.2	Canonical AZD for families	15
	4.3	Birational invariance	17
	4.4	An application of the canonical AZD	17
5	Positivity of the direct images of pluricanonical systems		18
	5.1	Theorems of Maitani-Yamaguchi and Berndtsson	18
	5.2	Variation of hermitian adjoint bundles	19
	5.3	Proof of Theorem 1.2	22

1 Introduction

Let $f: X \longrightarrow S$ be a flat projective family of varieties over a complex manifold S. In [N-S], Narashimhan and Simha constructed a continuous hermitian metric h_m on $mK_{X/S}$ for every sufficiently large positive integer m, when f is smooth and $K_{X/S}$ is relatively ample. They studied the parameter dependence of the metrics ([N-S, p.122, Section 3]) and proved the existence of moduli space of smooth projective varieties with ample canonical bundle ([N-S, p.126, Theorem]).

This metric depends only on the complex structure of X, hence it is functorial under biholomorphisms. Although they argued only the case of relatively ample $K_{X/S}$, the same construction can be applied also to the case that a general fiber has nonnegative Kodaira dimension (cf. Section 3). We call the metric the Narashimhan-Simha metric. But in this case we see that the resulting metric is a singular hermitian metric. And the metric is functorial under birational morphism in an appropriate sense (cf. Section 4.3).

The semipositivity result of Y. Kawamata [Ka1, p.57, Theorem 1] proved that when X is smooth and S is a projective curve, then $f_*\mathcal{O}_X(mK_{X/S})$ is semipositive on S in the sense of algebraic geometry, i.e., any quotient sheaf of $f_*\mathcal{O}_X(mK_{X/S})$ has semipositive degree. We note that the semipositivity in [Ka1, p.66,Theorem 1] comes from a Finsler type metric but not from a hermitian metric on $f_*\mathcal{O}_X(mK_{X/S})$.

The purpose of this paper is to refine this semipositivity result in terms of the Narashimhan-Simha singular hemitian metric on the multi relative canonical sheaf $\mathcal{O}_X(mK_{X/S})$. In this case it is natural to consider flat projective families of varieties with only canonical singularities (cf. Definition 2.1). Namely we prove the following theorem.

Theorem 1.1 ([T4, Theorem 1.1]) Let $f: X \longrightarrow S$ be a flat projective family with connected fibers such that a general fiber has only canonical singularities over a complex manifold S. We set

 $S^{\circ} := \{ s \in S \mid X_s := f^{-1}(s) \text{ is reduced and has only canonical singularities} \}.$

Let h_m be the Narashimhan-Simha singular hermitian metric on $mK_{X/S}$ (cf. Section 3.1) on $X^\circ := f^{-1}(S^\circ)$

Then h_m has semipositive curvature in the sense of current on X° and the curvature $-dd^c \log h_m$ extends across $X - X^{\circ}$ as a closed positive current. In particular h_m extends to a singular hermitian metric on $mK_{X/S}$ across $X - X^{\circ}$.

The main difference between Theorem 1.1 and [Ka1, p.57, Theorem 1] is that in Theorem 1.1 the positivity is on the family X not on the base space S. This fact plays the key role in this paper. Also we note that it is clear that h_m has semipositive curvature in the vertical direction, i.e., the restriction of h_m to every fiber of f has semipositive curvature. Hence the main assertion of Theorem 1.1 is that the curvature of h_m is semipositive also in the horizontal direction. This theorem is new even when $K_{X/S}$ is relatively ample and $f: X \longrightarrow S$ is smooth.

We note that the statement of Theorem 1.1 is meaningful, only if $f_*\mathcal{O}_X(mK_{X/S})$ is not zero. Otherwise h_m is not defined (cf. Section 3.1).

As a direct consequence of Theorem 1.1, we have the following theorem.

Theorem 1.2 Let $f: X \longrightarrow S$ be a flat projective family with connected fibers such that a general fiber of f has only canonical singularities. We assume X is normal and S is smooth. We set

 $S^{\circ} := \{ s \in S \mid X_s := f^{-1}(s) \text{ is reduced and has only canonical singularities} \}.$

Then $K_{X/S}$ has a relative AZD h over S° such that Θ_h is semipositive on X (Theorem 4.6).

And $F_m := f_* \mathcal{O}_X(mK_{X/S})$ carries a hermitian metric h_{F_m} defined by

$$h_{F_m}(\sigma,\sigma')_s := (\sqrt{-1})^{n^2} \int_{X_s} \sigma \wedge \bar{\sigma}' \cdot h^{m-1} \qquad (\sigma,\sigma' \in F_{m,s}, s \in S^\circ)$$

with Nakano semipositive curvature in the sense of current over S° , where $n = \dim X - \dim S$. \square

Remark 1.3 According to the recent outstanding result on finite generation of canonical rings ([B-C-H-M]), one may use $h_m^{1/m}$ instead of h for the relative AZD for some positive integer m. By the definition $h_m^{1/m}$ (see Section 3.1) have only algebraic singularities (cf. Definition 2.6). In this case the resulting metric on F_m seems to be quite regular. This might be a great advantage for the future applications. \square

We note that there is another approach to obtain Theorem 1.2 using the dynamical construction of AZD's (cf [T6]).

This article consists of three parts. The first part (Section 1,2) consists of the introduction and preliminaries. The second part (Section 3,4) consists of the proof of the semipositivity theorem, Theorem 1.1 and the construction of the canonical AZD for the relative canonical bundle. The third part (Section 5) consists of the proof of Theorem 1.2.

The author would like to express scincere gratitude to Professor Mihai Paun who pointed out an error in the proof of Theorem 5.5 and the reference [B-P].

In this paper all the varieties are defined over \mathbb{C} and the Bergman kernels means the diagonal parts (we do not use the reproducing kernel properties in this paper).

2 Preliminaries

Here we shall collect basic notions which are used in this article.

2.1 Canonical singularities

The notion of canonical singularities were introduced by M. Reid to study canonical models of general type ([R]). Actually canonical singularities are caractertized as singularities which appear on the canonical models when the canonical rings are finitely generated.

Definition 2.1 Let X be a normal variety such that the canonical divisor K_X is **Q**-Caritier. If $\mu: Y \longrightarrow X$ is a resolution of X, then we can write

$$K_Y = \mu^* K_X + F$$

with $F = \sum_j e_j E_j$ for the exceptional divisors $\{E_j\}$. We call F the discrepancy and $e_j \in \mathbf{Q}$ the discrepancy coefficient for E_j .

X is said to have only terminal singularities (resp. canonical singularities, if $e_j > -1$ (resp. ≥ -1) for all j for a log resolution $\mu: Y \longrightarrow X$. One can also say that X is terminal (resp. canonical), or K_X is terminal (resp. canonical), when X has only terminal (resp. canonical) singularities. \square

Definition 2.2 Let X be a normal variety with only canonical singularities. Let x be a point on X. The **local index** of X at x is the least positive integer r such that rK_X is Cartier near x. The **global index** of X is the least positive integer r such that rK_X is Cartier on X. \square

The following theorem is fundamental in this article.

Theorem 2.3 ([Ka3, p.85,Main Theorem]) Let $\pi: X \longrightarrow S$ be a flat morphism from a germ of variety (X, x_0) to a germ of a smooth curve (S, s_0) whose spetial fiber $X_0 = \pi^{-1}(s_0)$ has only canonical singularities. Then X has only canonical singularities. In particular, the fibers $X_s = \pi^{-1}(s)$ have only canonical singularities. \square

2.2 Singular hermitian metrics

In this subsection L will denote a holomorphic line bundle on a complex manifold M.

Definition 2.4 A singular hermitian metric h on L is given by

$$h = e^{-\varphi} \cdot h_0,$$

where h_0 is a C^{∞} -hermitian metric on L and $\varphi \in L^1_{loc}(M)$ is an arbitrary function on M. We call φ a weight function of h. \square

The curvature current Θ_h of the singular hermitian line bundle (L, h) is defined by

$$\Theta_h := \Theta_{h_0} + \sqrt{-1}\partial\bar{\partial}\varphi,$$

where $\partial \bar{\partial}$ is taken in the sense of a current. The L^2 -sheaf $\mathcal{L}^2(L,h)$ of the singular hermitian line bundle (L,h) is defined by

$$\mathcal{L}^{2}(L,h) := \{ \sigma \in \Gamma(U, \mathcal{O}_{M}(L)) \mid h(\sigma, \sigma) \in L^{1}_{loc}(U) \},$$

where U runs over the open subsets of M. In this case there exists an ideal sheaf $\mathcal{I}(h)$ such that

$$\mathcal{L}^2(L,h) = \mathcal{O}_M(L) \otimes \mathcal{I}(h)$$

holds. We call $\mathcal{I}(h)$ the **multiplier ideal sheaf** of (L,h). If we write h as

$$h = e^{-\varphi} \cdot h_0,$$

where h_0 is a C^{∞} hermitian metric on L and $\varphi \in L^1_{loc}(M)$ is the weight function, we see that

$$\mathcal{I}(h) = \mathcal{L}^2(\mathcal{O}_M, e^{-\varphi})$$

holds. For $\varphi \in L^1_{loc}(M)$ we define the multiplier ideal sheaf of φ by

$$\mathcal{I}(\varphi) := \mathcal{L}^2(\mathcal{O}_M, e^{-\varphi}).$$

Example 2.5 Let $\sigma \in \Gamma(X, \mathcal{O}_X(L))$ be the global section. Then

$$h := \frac{1}{\mid \sigma \mid^2} = \frac{h_0}{h_0(\sigma, \sigma)}$$

is a singular hemitian metric on L, where h_0 is an arbitrary C^{∞} -hermitian metric on L (the right hand side is orbitally independent of h_0). The curvature Θ_h is given by

$$\Theta_h = 2\pi\sqrt{-1}(\sigma)$$

where (σ) denotes the current of integration over the divisor of σ . \square

Definition 2.6 If $\{\sigma_i\}$ are a finite number of global holomorphic sections of L, for every positive rational number α and a continuous function ϕ ,

$$h := e^{-\phi} \cdot \frac{1}{\sum_{i} |\sigma_{i}|^{2\alpha}}$$

defines a singular hermitian metric on αL . We call such a metric h a singular hermitian metric on αL with algebraic singularities. \square

Singular hermitian metrics with algebraic singularities are particularly easy to handle, because its multiplier ideal sheaf of the metric can be controlled by taking suitable successive blowing ups such that the total transform of the divisor $\sum_{i}(\sigma_{i})$ is a divisor with normal crossings.

Definition 2.7 L is said to be **pseudoeffective**, if there exists a singular hermitian metric h on L such that the curvature current Θ_h is a closed positive current. Also a singular hermitian line bundle (L,h) is said to be **pseudoeffective**, if the curvature current Θ_h is a closed positive current. \square

2.3 Lelong numbers

The Lelong number is a measure of singularities of closed positive currents. We note that a subvariety of codimension p can be identified with the current of integration over it. In this special case the Lelong number exactly corresponds to the multiplicity. Hence we can consider the Lelong number as a generalization of the notion of multiplicities.

Let $W \subset \mathbb{C}^n$ be a domain, and Θ a positive current of degree (q,q) on W. For a point $p \in W$ one defines

$$\nu(\Theta, p, r) = \frac{1}{r^{2(n-q)}} \int_{\|z-p\| < r} \Theta(z) \wedge (dd^c \|z\|^2)^{n-q}.$$

The **Lelong number** of Θ at p is defined as

$$\nu(\Theta, p) = \lim_{\substack{r \to 0 \\ r > 0}} \nu(\Theta, p, r).$$

If Θ is the curvature of $h = e^{-u}$, u plurisubharmonic, one has

$$\nu(\Theta, p) = \sup\{\gamma \ge 0; u \le \gamma \log(\|z - p\|^2) + O(1)\}. \tag{1}$$

The definition of a singular hermitian metric carries over to the situation of reduced complex spaces.

Definition 2.8 Let Z be a reduced complex space and L a holomorphic line bundle. A singular hermitian metric h on L is a singular hermitian metric h on $L|Z_{reg}$ with the following property: There exists a desingularization $\pi: \tilde{Z} \to Z$ such that h can be extended from Z_{reg} to a singular hermitian metric \tilde{h} on π^*L over \tilde{Z} . \square

The definition is independent of the choice of a desingularization under a further assumption. Suppose that $\Theta_{\tilde{h}} \geq -c \cdot \omega$ in the sense of currents, where c>0, and ω is a positive definite, real (1,1)-form on \tilde{Z} of class C^{∞} . Let $\pi_1:Z_1\to Z$ be a further desingularization. Then $\tilde{Z}\times_Z Z_1\to Z$ is dominated by a desingularization Z' with projections $p:Z'\to \tilde{Z}$ and $p_1:Z'\to Z_1$. Now $p^*\log\tilde{h}$ is of class L^1_{loc} on Z' with a similar lower estimate for the curvature. The push-forward $p_{1*}p^*\tilde{h}$ is a singular hermitian metric on Z_1 . In particular, the extension of h to a desingularization of Z is unique.

In [G-R] for plurisubharmonic functions on a normal complex space the Riemann extension theorems were proved, which will be essential for our application.

For a reduced complex space a plurisubharmonic function u is by definition an upper semi-continuous function $u:X\to [-\infty,\infty)$ whose restriction to any local, smoothly parameterized analytic curve is either identically $-\infty$ or subharmonic.

A locally bounded function $u: X \to [-\infty, \infty)$ from $L^1_{loc}(X)$ is called weakly plurisubharmonic, if its restriction to the regular part of X is plurisubharmonic.

Differential forms with compact support on a reduced complex space are by definition locally extendable to an ambient subspace, which is an open subset U of some \mathbb{C}^n . Hence the dual spaces of differential C^{∞} -forms on such U define currents on analytic subsets of U. The positivity of a real (1,1)-current is defined in a similar way as above.

For locally bounded functions of class L^1_{loc} the weak plurisubharmonicity is equivalent to the positivity of the current dd^cu . It was shown that these functions are exactly those, whose pull-back to the normalization of X are plurisubharmonic. We note

Definition 2.9 Let L be a holomorphic line bundle on a reduced complex space X. Then a singular hermitian metric h is called positive, if the functions, which define locally $-\log h$ are weakly plurisubharmonic. \square

This definition is compatible with Definition 2.4. Let L be a holomorphic line bundle on a complex space Z equipped with a positive, singular hermitian

metric h_r on $L|Z_{reg}$. If $\pi: \tilde{Z} \to Z$ is a desingularization, and \tilde{h} a positive, singular hermitian metric on π^*L , extending $h|Z_{reg}$, we see that $-\log h_r$ is locally bounded at the singularities of Z so that \tilde{h} induces a singular, positive metric on L over Z.

The Lelong number of the curvature current of a pseudoeffective singular hemitian line bundle (cf. Definition 2.7) can be observed as an obstruction for nefness by the following lemma.

Lemma 2.10 (cf. [T1, T2]) Let X be a smooth projective variety and let (L, h) be a singular hermitian line bundle such that Θ_h is semipositive in the sense of current and the Lelong number $\nu(\Theta_h)$ is identically 0 on X. Then L is nef.

Proof of Lemma 2.10. Let $x \in X$ be an arbitrary point on X. Let A be a positive line bundle on X and let h_A be a C^{∞} -hermitian metric on A such that $\omega := \Theta_{h_A}$ is a Kähler form. And let d_x denote the distance function from x with respect to a Kähler form ω on X. Let a be a positive integer such that

$$(2n+2)\log d_x + \mathrm{Ric}_\omega + a\Theta_{h_A}$$

is strictly positive for every $x \in X$, where n denotes the dimension of X. This implies that

$$d_x^{-(2n+2)} \cdot h_A^a \cdot (\omega^n) \cdot h_L^m$$

is a singular hermitian metric on $A - K_X + mL$ with strictly positive curvature. Then by Nadel's vanishing theorem [N, p.561] and the description of the Lelong number (1), we see that mL + aA is free at x for every m. Since m and x are arbitrary, this implies that L is nef. \square

Remark 2.11 It is easy to generalize Lemma 2.10, in the form :

If (L,h) is a pseudoeffective singular hermitian line bundle on a smooth projective variety X. Then for every $x \in X$ such that $\nu(\Theta_h, x) = 0$ and an irreducible curve C containing $x, L \cdot C \geq 0$ holds, i.e., L is nef modulo the positive Lelong number locus of Θ_h . \square

2.4 Analytic Zariski decompositions (AZD)

In this subsection we shall introduce the notion of analytic Zariski decompositions. By using analytic Zariski decompositions, we can handle big line bundles like nef and big line bundles.

Definition 2.12 Let M be a compact complex manifold and let L be a holomorphic line bundle on M. A singular hermitian metric h on L is said to be an analytic Zariski decomposition, if the followings hold.

- 1. Θ_h is a closed positive current,
- 2. for every $m \ge 0$, the natural inclusion

$$H^0(M, \mathcal{O}_M(mL) \otimes \mathcal{I}(h^m)) \to H^0(M, \mathcal{O}_M(mL))$$

is an isomorphim. \square

Remark 2.13 If an AZD exists on a line bundle L on a smooth projective variety M, L is pseudoeffective by the condition 1 above. \square

Theorem 2.14 ([T1, T2]) Let L be a big line bundle on a smooth projective variety M. Then L has an AZD. \square

As for the existence for general pseudoeffective line bundles, now we have the following theorem.

Theorem 2.15 ([D-P-S, Theorem 1.5]) Let X be a smooth projective variety and let L be a pseudoeffective line bundle on X. Then L has an AZD.

3 Narashimhan-Simha metrics

In this section, following [N-S], we shall define a singular hermitian metric on the canonical bundle of a smooth projective manifold with nonnegative Kodaira dimension and study the behavior of the metric under deformation.

3.1 Narashimhan-Simha metrics

Let X be a smooth projective variety with nonnegative Kodaira dimension of dimension n. Let m be a positive integer. For a section $\eta \in H^0(X, \mathcal{O}_X(mK_X))$ we define a nonnegative number $\| \eta \|_{\underline{1}}$ by

$$\|\eta\|_{\frac{1}{m}} = |\int_{V} (\eta \wedge \bar{\eta})^{\frac{1}{m}}|^{\frac{m}{2}}.$$

Then $\eta \mapsto \parallel \eta \parallel_{\frac{1}{m}}$ is a continuous pseudonorm on $H^0(X, \mathcal{O}_X(mK_X))$, i.e., it is a continuous and has the properties :

- 1. $\|\eta\|_{\frac{1}{m}} = 0 \Leftrightarrow \eta = 0$,
- 2. $\|\lambda\eta\|_{\frac{1}{2}} = |\lambda| \cdot \|\eta\|_{\frac{1}{2}}$ holds for all $\lambda \in \mathbb{C}$.

But it is not a norm on $H^0(X, \mathcal{O}_X(mK_X))$ except m=1. We define a continuous section K_m of $(K_X \otimes \bar{K}_X)^{\otimes m}$

$$K_m(x) := \sup\{|\eta(x)|^2 ; \|\eta\|_{\frac{1}{m}} = 1\} \quad (x \in X),$$

where $|\eta(x)|^2 = \eta(x) \otimes \bar{\eta}(x)$. We call K_m the *m*-th Narashimhan-Simha potential of X. The continuity of K_m follows from the expression

$$K_m(x) = \sup\{|\eta(x)|^2 / ||\eta||_{\frac{1}{m}}; \quad \eta = \sum_{i=0}^{N(m)} c_i \sigma_i^{(m)}, \sum_{i=0}^{N(m)} |c_i|^2 = 1\},$$

where $\{\sigma_0^{(m)}, \dots, \sigma_{N(m)}^{(m)}\}$ is a basis for $H^0(X, \mathcal{O}_X(mK_X))$. We define the singular hermitian metric h_m on $H^0(X, \mathcal{O}_X(mK_X))$ by

$$h_m := \frac{1}{K_m}.$$

We call h_m the Narashimhan-Simha metric on mK_X . This metric is introduced by Narashimhan and Seshadri for smooth canonically polarized variety to study the moduli space of canonically polarized variety ([N-S]). We note that the singularities of h_m is located exactly on the support of the base locus of $|mK_X|$. Hence in the case of canonically polarized variety, h_m is a nonsingular continuous metric on mK_X for every sufficiently large m. Since K_m is locally the supremum of the family of powers of absolute value of holomorphic functions we see that the curvature

$$\Theta_{h_m} := \sqrt{-1}\partial\bar{\partial}\log K_m$$

is a closed positive current.

For applications sometimes more convenient to consider the following regularization of Θ_{h_m} .

Let us define the L^2 -inner product on $H^0(X, \mathcal{O}_X(mK_X))$ by

$$(\eta,\tau)_m := (\sqrt{-1})^{n^2} \int_X \frac{\eta \cdot \bar{\tau}}{K_{-}^{\frac{m-1}{m}}} \qquad (\eta,\tau \in H^0(X,\mathcal{O}_X(mK_X))).$$

The inner product is well defined, because by the definition of K_m , the zero of K_m corresponds exactly the ideal of Bs $|mK_X|$. Let $\{\sigma_0^{(m)}, \cdots \sigma_{N(m)}^{(m)}\}$ be a complete orthonormalbasis of $H^0(X, \mathcal{O}_X(mK_X))$ with respect to the inner product $(,)_m$. We define the Bergman kernel \tilde{K}_m by

$$\tilde{K}_m = \sum_{i=0}^{N(m)} |\sigma_i^{(m)}|^2.$$

Then \tilde{K}_m is independent of the choice of the complete orthonormal basis $\{\sigma_0^{(m)}, \cdots \sigma_{N(m)}^{(m)}\}$. In fact

$$\tilde{K}_m(x) := \sup\{ |\sigma(x)|^2 | \sigma \in H^0(X, \mathcal{O}_X(mK_X)), ||\sigma|| = 1 \}$$

holds, where $\| \|$ denotes the L^2 -norm with respect to $(,)_m$ (see for example [Kr, p.46]).

Then the closed positive current

$$\omega_m := \sqrt{-1}\partial\bar{\partial}\log\tilde{K}_m$$

coincides (up to some normalized constant) the pullback of the Fubini-Study Kähler form on $\mathbb{P}^{N(m)}$ on $X\backslash \operatorname{Supp} \operatorname{Bs} | mK_X|$ by the rational map :

$$X \ni x - \dots \to [\sigma_0^{(m)}(x) : \dots : \sigma_{N(m)}^{(m)}(x)] \in \mathbb{P}^{N(m)}.$$

We call ω_m the Narashimhan-Shimha Kähler current.

3.2 Dependence of Narashimhan-Simha metrics on families

Let $f: X \longrightarrow S$ be a flat projective family of varieties with only canonical singularities. Let n denote the relative dimension of $f: X \longrightarrow S$. Let X_s denote the fiber $f^{-1}(s)(s \in S)$. Let m be a positive integer and let $K_{m,s}$

be the Narashimhan-Shimha potential of mK_{X_s} . Then we obtain the family $\{mK_{X_s}\}_{s\in S}$ of Narashimhan-Simha potentials over S and the family of Narashimhan-Simha metrics $\{h_{m,s}\}_{s\in S}$ on X_s .

In [N-S] Narashimhan-Simha proved the continuous dependence of $K_{m,s}$ with respect to $s \in S$, when $K_{X/S}$ is relatively ample and f is smooth. The key argument here is the invariance of the plurigenera $P_m(X_s) = h^0(X_s, \mathcal{O}_{X_s}(mK_{X_s}))$ with respect to $s \in S$ which is derived by the Kodaira vanishing theorem at that time

We note that by [Ka3, p.91,Theorem 6] and [S1, S2, T3], the plurigenera $P_m(X_s)$ is independent of S. Hence every global holomorphic section of mK_{X_s} extends holomorphically to nearby fibers.

Hence by the defiition of the Narashimhan-Simha potential, we have the following lemma.

Lemma 3.1 $\{K_{m,s}\}$ is continuous with on X. \square

By Lemma 3.1 we see that there exists a relative volume form K_m such that

$$K_m \mid_{X_s} = K_{m,s}$$

holds for all $s \in S$. We call K_m the Narashimhan-Simha potential of the family $f: X \longrightarrow S$ and also

$$h_m := \frac{1}{K_m}$$

the Narashimhan-Simha metric on $mK_{X/S}$ respectively.

3.3 Positivity of Narashimhan-Simha metrics on projective families

In this subsection we shall prove Theorem 1.1. Let $f: X \longrightarrow S$ be a flat projective family with connected fibers such that a general fiber has only canonical singularities over a complex manifold S. We set

 $S^{\circ} := \{s \in S \mid X_s := f^{-1}(s) \text{ is reduced and has only canonical singularities}\}.$

For $s \in S$, we denote X_s the fiber $f^{-1}(s)$. For the proof we may and do assume that S is the open unit disk Δ in \mathbb{C} . We note that $f^*\mathcal{O}_X(mK_{X/S})$ is not compatible with base change. Nevertheless this is enough to prove the statement of Theorem 1.1 on the extension of h_m across $X - X - \circ$ holds, by slicing.

Let us denote $f_*\mathcal{O}_X(mK_{X/S})$ by F_m .

Lemma 3.2 ([Ka1, p.63, Lemma 7 and p.64, Lemma 8]) Let $\eta \in H^0(S, \mathcal{O}_S(F_m))$ be a holomorphic section. Let $\|\eta(s)\|_{\frac{1}{m}}$ be the Narashimhan-Sihma pseudonorm of $\eta(s) \in H^0(X_s, \mathcal{O}_{X_s}(mK_{X_s}))$. Then

$$\log \|\eta(s)\|_{\frac{1}{m}} \ge \frac{1}{2\pi} \int_0^{2\pi} \log \|\eta(s + re^{\sqrt{-1}\theta})\|_{\frac{1}{m}} d\theta$$

holds for every $s \in S$ and r > 0 such that $\Delta(s,r) \subset S$, i.e. $\log \| \eta(s) \|_{\frac{1}{m}}$ is superharmonic on S, where $\Delta(s,r)$ denotes the open disk of radius r with centre s. \square

Remark 3.3 In [Ka1], Lemma 3.2 was considered only for smooth projective families. But by Theorem 2.3 the same proof works for flat projective families with only canonical singularities. We note that canonical singularities are rational ([E] or cf.[K-M]). \square

Let us fix an arbitrary $s \in S^{\circ}$. Let $x \in X_s$, r be a positive number such that $\Delta(s,r) \subset S$ and let $\tau: \Delta(s,r) \longrightarrow X$ be any holomorphic section such that $\tau(s) = x$. In general the such a section does not exist when x is a singular point on X_s . This case will be treated later.

Let $\eta \in H^0(S, \mathcal{O}_S(F_m))$ such that $\| \eta(s) \|_{\underline{1},s} = 1$ and

$$|\eta(s,x)|^2 = K_{m,s}(x)$$

holds, where $\eta(s)$ denotes the restriction $\eta\mid_{X_s}$ and $\eta(s,x)$ denotes the point value of $\eta(s)$ at x.

Let a be a positive integer such that $aK_{X/S}$ is Cartier. Let Ω be a local generator of the invertible sheaf $\mathcal{O}_X(aK_{X/S})$ on a neighbourhood U of x. Shrinking S, if necessary, we may assume that $\tau(\Delta(s,r)) \subset U$ holds. Let f be the function on $\Delta(s,r)$ defined by

$$f(z) = \left| \frac{\eta}{\Omega^{m/a}} \right|^2 (\tau(z)) \quad (z \in \Delta(s, r))$$

Then we see that

$$\log f(s) \le \frac{1}{2\pi} \int_0^{2\pi} \log f(s + re^{\sqrt{-1}\theta}) d\theta \tag{2}$$

holds by the subharmonicity of the logarithm of abolute value of holomorphic function. On the other hand we see that

$$\log \| \eta(s) \|_{\frac{1}{m}} \ge \frac{1}{2\pi} \int_0^{2\pi} \log \| \eta(s + re^{\sqrt{-1}\theta}) \|_{\frac{1}{m}} d\theta$$
 (3)

holds by Lemma 3.2.

By (1) and (2), we see that

$$\log \frac{f(s)}{\| \eta(s) \|_{\frac{1}{m}}} \le \frac{1}{2\pi} \int_0^{2\pi} \log \frac{f(s + re^{\sqrt{-1}\theta})}{\| \eta(s + re^{\sqrt{-1}\theta})) \|_{\frac{1}{m}}} d\theta$$

We note that

$$K_m(\tau(s + re^{\sqrt{-1}\theta})) \ge \frac{f(s + re^{\sqrt{-1}\theta})}{\|\eta(s + re^{\sqrt{-1}\theta})\|_{\frac{1}{a}}} |\Omega|^{\frac{m}{a}}$$

holds by the definition of K_m . Hence we see that

$$\log K_m(\tau(s)) \le \frac{1}{2\pi} \int_0^{2\pi} \log K_m(\tau(s + re^{\sqrt{-1}\theta})) d\theta$$

holds. Hence $\log K_m$ is plurisubharmonic on $\tau(\Delta(s,r))$.

Next we shall deal with the case that there exists no local section passing through x. In this case the above argument also holds replacing section by holomorphic disk $\tau: \Delta \longrightarrow X$ passing through x.

Since x an τ are arbitrary, the curvature $\Theta_{h_m} = \sqrt{-1}\partial\bar{\partial} \log K_m$ is semipositive everywhere on X° .

Since for every local holomorphic section σ of $f_*\mathcal{O}_X(mK_{X/S})$ the function

$$(\sqrt{-1})^{n^2} \int_{X_{\mathfrak{o}}} (\sigma \wedge \bar{\sigma})^{\frac{1}{m}}$$

is of algebraic growth along $S - S^{\circ}$. More precisely for $s_0 \in S - S^{\circ}$ as in [Ka1, p.59 and p. 66] there exist positive numbers C, α, β such that

$$(\sqrt{-1})^{n^2} \int_{X_s} (\sigma \wedge \bar{\sigma})^{\frac{1}{m}} \leq C \cdot |s - s_0|^{-\alpha} \cdot |\log(s - s_0)|^{\beta}$$

$$\tag{4}$$

holds. Hence h_m is also of algebraic growth by its definiton along the fiber on $S - S^{\circ}$. Hence the current $-dd^c \log h_m$ on X° extends to a closed positive current on X.

This completes the proof of Theorem 1.1. \Box

Remark 3.4 As one sees in the proof above, the semipositivity of the curvature Θ_{h_m} is the consequence of the Kawamata's semipositivity ([Ka1, p.63 Lamma 7] and the fact that the supremum of plurisubharmonic functions are again plurisubharmonic if we take the uppersemicontinuous envelope (cf. Theorem 4.4). \square

Let us consider the direction in which Θ_{h_m} is strictly positive. Let

$$p: \mathbb{P}(F_m^*) \longrightarrow S$$

be the projective bundle associated with F_m^* and let

$$L \longrightarrow \mathbb{P}(F_{m}^{*})$$

be the tautological bundle on L. Then

$$\|\eta\|_{\frac{1}{m}} = |\int_{X_s} (\eta \wedge \bar{\eta})^{\frac{1}{m}}|^{\frac{m}{2}} \quad (\eta \in H^0(X_s, \mathcal{O}_{X_s}(mK_{X_s})))$$

defines a continuous hermitian metric h_L on L. By [Ka1, p.63, Lemma 7 and p.64, Lemma 8] we see that the curvature current Θ_{h_L} is semipositive.

We note that on the fiber of p, Θ_{h_L} is semipositive and is the curvature of the tautological line bundle which is ample on the fiber. In fact let $\eta(t)(t \in \Delta)$ be a holomorphic family of sections on a fixed projective variety X. Then by the concavity of logarithm, for every $r \in [0,1)$

$$\begin{aligned} \log(|\int_{X} (\eta(0) \wedge \bar{\eta}(0))^{\frac{1}{m}}|) &= \log(\int_{X} (\frac{1}{2\pi} |\int_{0}^{2\pi} (\eta(re^{\sqrt{-1}\theta}) \wedge \bar{\eta}(re^{\sqrt{-1}\theta}))^{\frac{1}{m}}|) d\theta) \\ &\geq \frac{1}{2\pi} \int_{0}^{2\pi} \log(|\int_{X} \eta(re^{\sqrt{-1}\theta}) \wedge \bar{\eta}(re^{\sqrt{-1}\theta})^{\frac{1}{m}}|) d\theta \end{aligned}$$

hold and the similar inequality holds for every point on S (this is a special case of [Ka1, p. 63, Lemma 7 and p.64, Lemma 8]).

Hence $\Theta_{h_L}|_{p^{-1}(s)}$ is strictly positive on the generic point of the fiber $p^{-1}(s)$ for every $s \in S$. We note that Θ_{h_L} is strictly positive in the direction in which the coupling of the Kodaira-Spencer class of the logarithmic deformation in the direction and the m-th root of the m-canonical forms is nonzero as in the proof of [Ka1, p.63, Lemma 7] (this is not explicitly stated in the proof of [Ka1, p.63, Lemma 7], but the discussion in [Ka2] clarifies this standard fact). Hence to specify the strictly posive direction of Θ_{h_m} , we need the following infinitesimal Torelli theorem (see also Remark 3.4 above).

Theorem 3.5 ([Ka2]) Let X be a normal projective variety of dimension d having only canonical singularities. We assume that X is a good minimal algebraic variety, i.e., m_0K_X is Cartier and $\hat{\omega}_X^{m_0} = \mathcal{O}_X(m_0K_X)$ is generated by global sections for some positive integer m_0 . Let $\gamma: X \longrightarrow B$ be the morphism given by the base point free linear system $|m_0K_X|$. We take m_0 large enough so that B is normal and γ has connected fibers. Let $C = (\sigma)$ for some $\sigma \in H^0(X, \hat{\omega}_X^m)$ with $m = am_0$ and $a \in \mathbb{N}$.

Then for $0 < i < m, i \equiv 1$ and if 2i > m+1, then the kernel of the following homomorphism induced by the cup product

$$\lambda_X^i: \operatorname{Ext}_X^1(\Omega_X^1(\log C), \mathcal{O}_X) \to \operatorname{Hom}(H^0(X, \hat{\omega}_X^{m+1-i}), \operatorname{Ext}_X^1(\Omega_X^1(\log C), \hat{\omega}_X^{m+1-i}))$$

is contained in the image of the natural homomorphism

$$H^1(B, \gamma_* \mathcal{O}_X(T_{X/B})) \to Ext_X^1(\Omega_X^1(\log C), \mathcal{O}_X).$$

Hence by the proof of Theorem 1.1 and Theorem 3.5, we have the following refined semipositivity theorem.

Theorem 3.6 Let $f: X \longrightarrow S$ be a flat projective family of canonical models of general type with only canonical singularities. Let h_m be the Narashimhan-Simha singular hermitian metric on $mK_{X/S}$. Suppose that $mK_{X/S}$ is Cartier and $\mid mK_{X/S} \mid$ is relatively very ample. Then Θ_{h_m} is strictly positive on the generic point of the fiber in the effectively parametrized direction. \square

Remark 3.7 It is not difficult to generalized Theorem 3.6 for all $m \geq 2$ such that $f_*\mathcal{O}_X(mK_{X/S}) \neq 0$ by using the ring structure of $\bigoplus_{\ell=0}^{\infty} f_*\mathcal{O}_X(\ell K_{X/S})$. Similar stirict positivity theorem holds for the flat families of minimal projective varieties under the assumption that minimal model conjecture holds. \square

4 Canonical AZD

In this section, we shall prove that the canonical bundle of every smooth projective manifold of nonnegtive Kodaira dimension carries a canonical AZD which is functorial under any birational morphism. Namely we shall prove the following theorem.

Theorem 4.1 Let X be a smooth projective variety (not necessary of general type) of dimension n with nonnegative Kodaira dimension. Let K_m be the m-th Narashimhan-Simha potential for every positive integer m. We set

$$C_m := \int_X K_m^{\frac{1}{m}}.$$

We define a semipositive (n,n)-form K_{∞} on X by

$$K_{\infty}(x) := \sup_{m \ge 1} \frac{1}{C_m} K_m^{\frac{1}{m}}(x) \quad (x \in X).$$

Then K_{∞} exists as a bounded (n,n)-form on X $(n=\dim X)$ and

$$h := \frac{1}{K_{\infty}}$$

is an AZD of K_X . \square

As a corollary, we have the following:

Corollary 4.2 Let X be a smooth projective variety with nonnegative Kodaira dimension. Then there exists an AZD h on K_X which depends only on the complex structure of X. \square

Definition 4.3 Let X be a smooth projective variety with nonnegative Kodaira dimension. We call the AZD of K_X constructed as in Theorem 4.1 the canonical AZD of K_X . \square

4.1 Construction of the canonical AZD

The supremum of a family of plurisubharmonic functions uniformly bounded from above is known to be again plurisubharmonic, if we modify the supremum on a set of measure 0(i.e., if we take the uppersemicontinuous envelope) by the following theorem of P. Lelong.

Theorem 4.4 ([L, p.26, Theorem 5]) Let $\{\varphi_t\}_{t\in T}$ be a family of plurisubharmonic functions on a domain Ω which is uniformly bounded from above on every compact subset of Ω . Then $\psi = \sup_{t\in T} \varphi_t$ has a minimum uppersemicontinuous majorant ψ^* which is plurisubharmonic. We call ψ^* the uppersemicontinuous envelope of ψ . \square

Remark 4.5 In the above theorem the equality $\psi = \psi^*$ holds outside of a set of measure 0(cf./L, p.29). \square

Proof of Theorem 4.1. Since

$$\frac{1}{C_m} \int_{Y} K_m^{\frac{1}{m}} = 1$$

holds for every m such that $\mid mK_X \mid \neq \emptyset$. Since $h_m = 1/K_m$ has semipositive curvature in the sense of current, by the submeanvalue property of plurisubharmonic functions, we see that $\{C_m^{-1} \cdot K_m^{\frac{1}{m}}\}_{m=1}^{\infty}$ is a uniformly bounded family of semipositive (n,n)-forms on X. Hence the pointwise supremum

$$K_{\infty} := \sup_{m \ge 1} \frac{1}{C_m} K_m^{\frac{1}{m}}$$

exists as a semipositive (n, n)-form on X. By definition

$$K_{\infty} \ge \frac{1}{C_m} K_m^{\frac{1}{m}}$$

holds for every m. Hence

$$h :=$$
the lower envelope of $\frac{1}{K_{\infty}}$

is a well defined singular hermitian metric on K_X and

$$\mathcal{I}(h^m) \supseteq \mathcal{I}(h_m) \tag{5}$$

holds for every $m \ge 1$. We note that the definition of K_m ,

$$H^{0}(X, \mathcal{O}_{X}(mK_{X}) \otimes \mathcal{I}(h_{m})) = H^{0}(X, \mathcal{O}_{X}(mK_{X}))$$
(6)

holds for every $m \ge 1$. Then combining (5) and (6), we see that

$$H^0(X, \mathcal{O}_X(mK_X) \otimes \mathcal{I}(h^m)) = H^0(X, \mathcal{O}_X(mK_X))$$

holds for every $m \ge 1$.

On the other hand, by Theorem 4.4 , we see that the curvature current Θ_h of h is seimipositive in the sense of current.

Hence h is an AZD of K_X . \square

4.2 Canonical AZD for families

The construction of the canonical AZD in Theorem 4.1 can be applied to a flat projective family of varieties with only canonical singularities similarly as in Section 3.2.

Theorem 4.6 Let $f: X \longrightarrow S$ be a flat projective family of varieties with connected fibers such that a general fiber of f has only canonical singularities and nonnegative Kodaira dimension. We set

$$S^{\circ} := \{ s \in S \mid X_s := f^{-1}(s) \text{ is reduced with only canonical singularities} \}.$$

Then there exists a singular hermitian metric h on $K_{X/S}$ with semipostive curvature current such that for every $s \in S^{\circ}$, $h \mid_{X_s}$ is an AZD of K_{X_s} . \square

Proof of Theorem 4.6. The construction is essentially same as in the last subsection. Let $\{K_{m,s}\}(s \in S^{\circ})$ be the family of the m-th Narashimhan-Simha potentials over S° . We define the relative potential K_m on $f^{-1}(S^{\circ})$ by

$$K_m | X_s = K_{m,s} \quad (s \in S^\circ).$$

Suppose that there exists $\sigma_0 \in H^0(X_s, \mathcal{O}_{X_s}(m_0K_{X_s}))$ such that

$$|\int_{X_s} (\sigma_0 \wedge \bar{\sigma}_0)^{\frac{1}{m_0}}| = 1$$

Then for any point $x \in X_s$ and positive integer ℓ , we see that

$$K_{m_0\ell,s}(x) \ge |\sigma_0|^{2\ell} (x)$$

holds by the definition of $K_{m_0\ell,s}$. Hence by using the invariance of plurigenera ([S1, S2, T3, Ka3]), since we have assumed that every fiber on S° has nonnegative Kodaira dimension, there exists a positive continuous function c(s) on S° such that for every positive integer ℓ and $s \in S^{\circ}$

$$\int_{X_{-}} K_{m_0\ell,s}^{\frac{1}{m_0\ell}} \geqq c(s) \tag{7}$$

holds.

On the other hand, we obtain the upper estimate of

$$\int_{X_s} K_{m,s}^{\frac{1}{m}}$$

as follows. Let us fix a C^{∞} -volume form dV_s on X_s , such that dV_s^{-1} is a C^{∞} hermitian metric on the $\mathbb Q$ line bundle K_{X_s} . Then for any $\sigma \in H^0(X_s, \mathcal O_{X_s}(mK_{X_s}))$ with

$$|\int_{X_s} (\sigma \wedge \bar{\sigma})^{\frac{1}{m}}| = 1,$$

by the submean value property of plurisubharmonic functions, there exists a positive constant C'(s) independent of σ and m such that

$$\mid \frac{(\sigma \wedge \bar{\sigma})^{\frac{1}{m}}}{dV_s} \mid \leq C'(s)$$

holds on X_s . Hence there exists a positive continuous function C(s) on S° such that

$$\int_{X_s} K_{m,s}^{\frac{1}{m}} \le C(s) \tag{8}$$

holds for every $m \ge 1$.

Let $s_0 \in S^{\circ}$ be an arbitrary point. We set

$$C_m := \int_{X_{s_0}} K_{m,s_0}^{\frac{1}{m}}.$$

Let

$$K_{\infty} := \sup_{m \ge 1} \frac{1}{C_m} K_m^{\frac{1}{m}}$$

be the pointwise supremum. Then $K_{\infty}^{-1} \mid X_{s_0}$ is an AZD on $K_{X_{s_0}}$ as in Theorem 4.1. By the estimates (7) and (8), we see that K_{∞} is a nontrivial locally bounded realative volume form on $f^{-1}(S^{\circ})$. By the plurisubharmonic variation property of K_m (Theorem 1.1),

$$dd^c \log K_{\infty} \ge 0$$

holds on S° . Then again by the proof of Theorem 4.1, we see that $K_{\infty} \mid X_s$ is an AZD of K_{X_s} for every $s \in S^{\circ}$ by the above construction (especially by the estimates (7) and (8)).

Now we consider the extension of the closed positive current

$$dd^c \log K_{\infty}$$

across $X - f^{-1}(S^{\circ})$. But this is trivial, since each $K_m^{\frac{1}{m}}$ has only algebraic singularities along $X - f^{-1}(S^{\circ})$, in other words, for any local holomorphic section σ of $f_*\mathcal{O}_X(mK_{X/S})$,

$$(\sqrt{-1})^{n^2} \int_{X_-} (\sigma \wedge \bar{\sigma})^{\frac{1}{m}} \quad (n = \dim X - \dim S)$$

has algebraic growth glong $S-S^{\circ}$ as (4) in Section 3.2 by its definition (this has been already observed in [Ka1, pp.65-66]). Hence by definition, $dd^c \log K_{\infty}$ also extends across $X-f^{-1}(S^{\circ})$ as a closed positive current (one may also use Lemma 5.6 below). \square

By Theorem 1.1 and the recent result of finite generation canonical rings ([B-C-H-M]) and Theorem 3.5 we have the following theorem immediately (see Theorem 3.6).

Theorem 4.7 Let $f: X \longrightarrow S$ be a flat projective family of varieties of general type with only canonical singularities. Let h be the canonical AZD on $mK_{X/S}$. Then h has semipositive curvature Θ_h in the sense of current everywhere on X and Θ_h is strictly positive on the generic point of the fiber in the birationally effectively parametrized direction on X_s for every $s \in S_0$. \square

4.3 Birational invariance

By the definition of the canonical AZD in Theorem 4.1, one can see easily that it is invariant under birational morphism.

Let X, X' be smooth projective variety with nonnegative Kodaira dimension. Suppose that X and X' are birational. Then there exists a smooth projective variety Y which dominates birationally both X and X'. Let $f: Y \longrightarrow X$ and $f': Y \longrightarrow X'$ be the birational morphism.

For every positive integer m, let us denote the m-th Narashimhan-Simha potential of X and X' by K_m and K'_m respectively.

Then for every positive integer m, by the definition of the Narashimhan-Simha potential, we see that

$$f^*K_m = (f')^*K'_m$$

holds on Y. Hence the Narashimhan-Simha potential is invariant under birational morphisms and the canonical AZD for K_X is also invariant under birational morphisms.

4.4 An application of the canonical AZD

We shall show an easy application of the canonical AZD.

Theorem 4.8 Let $f: X \longrightarrow C$ be a semistable projective family over a projective curve C such that $K_{X/C} | F$ is ample for every fiber F.

Then $K_{X/C}$ is nef. \square

Proof of Theorem 4.8. Let $C_0 \subset C$ be the Zariski open subset such that f is smooth over C_0 . Let m be a positive integer such that $mK_{X/C}$ is relatively very ample. Let h_m be the Narashimhan-Simha metric on $mK_{X/C} \mid f^{-1}(C_0)$.

Let $F_0 = \sum D_i$ be a singular fiber. We note that in this case by the L^2 -extension theorem [O, Theorem], for every i, every element of $H^0(D_i, \mathcal{O}_{D_i}(mK_{D_i} + (m-1)\sum_{j\neq i}D_j \mid D_i)$ extends holomorphically to a section $H^0(f^{-1}(V), \mathcal{O}_X(mK_{X/C}))$, where V is a neighbourhood of $f(F_0)$. Hence we see that $h_m = K_m^{-1}$ extends to a singular hermitian metric on $mK_{X/C}$ on the whole X so that K_m extends continuously to X.

Then by the definition of h_m , for every sufficiently large m, $h_m|D_i$ is a smooth metric on the globally generated sheaf $\mathcal{O}_{D_i}(mK_{D_i}+(m-1)\sum_{j\neq i}D_j\mid D_i)$. By the construction of the canonical AZD h, we see that h is a singular hermitian metric on $K_{X/C}$ with the semipositive curvature current Θ_h satisfying $\nu(\Theta_h)\equiv 0$ on X by the relatively ampleness of $K_{X/C}$. This implies that $K_{X/C}$ is nef by Lemma 2.10. \square

5 Positivity of the direct images of pluricanonical systems

In this section we prove Theorem 1.2.

5.1 Theorems of Maitani-Yamaguchi and Berndtsson

In 2004, Maitani and Yamaguchi proved the following theorem.

Theorem 5.1 ([M-Y]) Let Ω be a pseudoconvex domain in $\mathbb{C}_z \times \mathbb{C}_w$ with C^1 boundary. Let $\Omega_t := \Omega \cap (\mathbb{C}_z \times \{t\})$ and Let K(z,t) be the Bergman kernel function of Ω_t .

Then $\log K(z,t)$ is a plurisubharmonic function on Ω . \square

Recently generalizing Theorem 5.1, B. Berndtsson proved the following higher dimensional and twisted version of Theorem 5.1.

Theorem 5.2 ([B1]) Let D be a pseudoconvex domain in $\mathbb{C}_z^n \times \mathbb{C}_t^k$. And let ϕ be a plurisubharmonic function on D. For $t \in \Delta$, we set $D_t := \Omega \cap (\mathbb{C}^n \times \{t\})$ and $\phi_t := \phi \mid D_t$. Let $K(z,t)(t \in \mathbb{C}_t^k)$ be the Bergman kernel of the Hilbert space

$$A^{2}(D_{t}, e^{-\phi_{t}}) := \{ f \in \mathcal{O}(\Omega_{t}) \mid \int_{D_{t}} e^{-\phi_{t}} \mid f \mid^{2} < +\infty \}.$$

Then $\log K(z,t)$ is a plurisubharmonic function on D. \square

As in mensioned in [B2], his proof also works for a pseudoconvex domain in a locally trivial family of manifolds which admits a Zariski dense Stein subdomain. Also he proved the following theorem.

Theorem 5.3 ([B2, Theorem 1.1]) Let us consider a domain $D = U \times \Omega$ and let ϕ be a plurisubharmonic function on D. For simplicity we assume that ϕ is smooth up to the boundary and strictly plurisubharmonic in D. Then for each $t \in U$, $\phi_t := \phi(\cdot, t)$ is plurisubharmonic on Ω . Let A_t^2 be the Bergman space of holomorphic functions on Ω with norm

$$|| f ||^2 = || f ||_t^2 := \int_{\Omega} e^{-\phi_t} |f|^2.$$

The spaces A_t^2 are all equal as vector spaces but have norms that vary with t. Then "infinite rank" vector bundle E over U with fiber $E_t = A_t^2$ is therefore trivial as a bundle but is equipped with a notrivial metric. Then $(E, || \cdot ||_t)$ is strictly positive in the sense of Nakano. \square

In Theorem 5.2 the assumption that D is a pseudoconvex domain in the product space is rather strong. And in Theorem 5.3, Berndtsson also assumed that D is a product.

5.2 Variation of hermitian adjoint bundles

Recently Berndtsson and I have independently generalized Theorems 5.2 and 5.3 to the case of projective families. For our purpose, we need the following version. Theorem 5.4 has been obtained independently by Berndtsson and Paun [B3, B-P] by a very different method.

Theorem 5.4 ([T5, T6, B-P]) Let $f: X \longrightarrow S$ be a projective family of projective varieties over a complex manifold S. Let S° be the maximal nonempty Zariski open subset such that f is smooth over S° .

Let (L,h) be a singular hermitian line bundle on X such that Θ_h is semi-positive on X.

Let $K_s := K(X_s, K_X + L \mid_{X_s}, h \mid_{X_s})$ be the Bergman kernel of $K_{X_s} + (L \mid X_s)$ with respect to $h \mid X_s$ for $s \in S^{\circ}$. Then the singular hermitian metric h_B of $K_{X/S} + L \mid f^{-1}(S^{\circ})$ defined by

$$h_B \mid X_s := K_s^{-1} (s \in S^{\circ})$$

has semipositive curvature on $f^{-1}(S^{\circ})$ and extends on X as a singular hermitian metric on $K_{X/S} + L$ with semipositive curvature current. \square

Theorem 5.5 ([T5, T6]) Let $f: X \longrightarrow S$ be a projective family of over a complex manifold S such that X is smooth. Let S° be a nonempty Zariski open subset such that f is smooth over S° . Let (L,h) be a hermitian line bundle on X such that Θ_h is semipositive on X. We define the hermitian metric h_E on $E:=f_*\mathcal{O}_X(K_{X/S}+L)\mid S^{\circ}$ by

$$h_E(\sigma,\tau) := (\sqrt{-1})^{n^2} \int_{X_s} h \cdot \sigma \wedge \bar{\tau} \qquad (\sigma,\tau \in H^0(X_s,\mathcal{O}_{X_s}(K_{X_s} + L\mid_{X_s}))),$$

where n denotes the relative dimension of $f: X \longrightarrow S$. Let S_0 be the maximal Zariski open subset of S° such that $E \mid S_0$ is locally free. Then $(E, h_E) \mid S_0$ is semipositive in the sense of Nakano. Moreover if Θ_h is strictly positive, then $(E, h_E) \mid S_0$ is strictly positive in the sense of Nakano. \square

Here we reproduce the proofs of Theorems 5.4 and 5.5 here for the completeness.

Proof of Theorems 5.4 and 5.5.

Let $f: X \longrightarrow S$ be a projective family. Since the statement is local we may assume that S is the unit open ball B with center O in \mathbb{C}^m . We may also and do assume that the family $f: X \longrightarrow B$ is a restriction of a projective family

$$\hat{f}: \hat{X} \longrightarrow B(O,2)$$

over the open ball B(O,2) of radius 2 with center O. Let

$$F: \hat{X} \times B(O,2) \longrightarrow B(O,2) \times B(O,2)$$

the fiber space defined by

$$F(x,t) = (f(x),t).$$

Let ε be a positive number less than 1. We set

$$T(\varepsilon) = \{(s,t) \in B(O,2) \times B(O,2) \mid t \in B, s \in B(t,\varepsilon)\}\$$

and

$$X(\varepsilon) := F^{-1}(T(\varepsilon)).$$

Let

$$f_{\varepsilon}: X(\varepsilon) \longrightarrow B$$

be the family defined by

$$f_{\varepsilon}(x,t) = t.$$

Since for $(x,t) \in X(\varepsilon)$, $x \in f^{-1}(B(t,\varepsilon))$ holds, we see that

$$X(\varepsilon)_t := f^{-1}(B(t,\varepsilon))$$

holds. Hence we may consider $X(\varepsilon)_t$ as a family

$$\pi_{\varepsilon,t}: X(\varepsilon,t) \longrightarrow B(t,\varepsilon).$$

We note that $T(\varepsilon)$ is a domain of holomorphy in \mathbb{C}^{2m} . Hence $X(\varepsilon)$ is a pseudo-convex domain in $X \times B(O,2)$. Since $X \times B(O,2)$ is a product manifold, the proof of Theorem 5.2 works without any essential change in this case (cf. [B1]). Hence if we define K_{ε} by

$$K_{\varepsilon} \mid X(\varepsilon)_t := K(X(\varepsilon)_t, K_X + L \mid X(\varepsilon)_t, h_L \mid X(\varepsilon)_t),$$

then

$$\sqrt{-1}\partial\bar{\partial}\log K_{\varepsilon} \ge 0 \tag{9}$$

holds on $X(\varepsilon)$. We note that

$$\lim_{\varepsilon \downarrow 0} \operatorname{vol}(B(t,\varepsilon)) \cdot K(X(\varepsilon)_t, K_X + L \mid X(\varepsilon)_t, h_L \mid X(\varepsilon)_t) = K(X_t, K_{X_t} + L \mid X_t, h \mid X_t) \wedge f^* d\mu(t)$$
(10)

holds for almost all $t \in B$, where $d\mu(t)$ denotes the standard volume form on \mathbb{C}^m . In fact, if we consider the family

$$\pi_{\varepsilon,t}:X(\varepsilon)\longrightarrow B(t,\varepsilon)$$

as a family over the unit open ball B in \mathbb{C}^m with center O by

$$t' \mapsto \varepsilon^{-1}(t'-t),$$

the limit as $\varepsilon \downarrow 0$ is nothing but the trivial family $X_t \times B$. And we note that for a L-valued canonical form σ on $f^{-1}(B(t,\varepsilon))$,

$$\int_{B(t,\varepsilon)} h_E(\sigma,\sigma)$$

is nothing but the L^2 -norm of the L-valued canonical form σ with respect h_L over $f^{-1}(B(t,\varepsilon))$ by Fubini's theorem, where we abbrebiate the standard Lebesgue measure on \mathbb{C}^m . In this way the L^2 -norm on $f^{-1}(B(t,\varepsilon))$ can be viewd the molification of the L^2 -norm on the fiber $f^{-1}(t)$. Hence as $\varepsilon \downarrow 0$, the molified norm converges to the norm on the fiber almost everywhere on B. Then the desired equality follows from the L^2 -extension theorm ([O-T, O]) and the extremal property of the Bergman kernels. In fact the L^2 -extension theorem implies

$$\lim_{\varepsilon \downarrow 0} \operatorname{vol}(B(t,\varepsilon)) \cdot K(X(\varepsilon)_t, K_X + L \mid X(\varepsilon)_t, h_L \mid X(\varepsilon)_t) \ge K(X_t, K_{X_t} + L \mid X_t, h \mid X_t) \wedge f^* d\mu(t)$$

for almost all $t \in B$ and the converse inequality follows from the convergence to the trivial family as above.

Combining (9) and (10), we see that h_B has semipositive curvature on $f^{-1}(S^{\circ})$.

To get an extension of h_B across $X - f^{-1}(S^{\circ})$, let us quote the following

Lemma 5.6 ([B-T, Corollary 7.3]) Let $\{u_j\}$ be a sequence of plurisubharmonic functions locally bounded above on the bounded open set Ω in \mathbb{C}^m . Suppose further

$$\limsup_{j\to\infty} u_j$$

is not identically $-\infty$ on any component of Ω . Then there exists a plurisubharmonic function u on Ω such that the set of points

$$\{x \in \Omega \mid u(x) \neq (\limsup_{j \to \infty} u_j)(x)\}$$

is pluripolar. \square

The extension of h_B in Theorem 5.4 follows from Lemma 5.6. In fact since $K(X(\varepsilon)_t, K_X + L \mid X(\varepsilon)_t, h_L \mid X(\varepsilon)_t)$ varies plurisubharmonic way in $t \in B$,

$$\{\operatorname{vol}(B(t,\varepsilon))\cdot K(X(\varepsilon)_t,K_X+L\mid X(\varepsilon)_t,h_L\mid X(\varepsilon)_t)\mid_{X_t}\}_{t\in B}$$

is uniformly bounded on B even when $t \in S - S^{\circ}$. Then letting ε tend to 0 and applying Lemma 5.6 we see that $-dd^{c} \log h_{B}$ extends across $X - f^{-1}(S^{\circ})$ as a closed positive current.

This completes the proof of Theorem 5.4.

The proof of Theorem 5.5, is quite similar. First we note that

$$f_{\varepsilon}: X(\varepsilon) \longrightarrow B$$

is everywhere smooth.

For the moment, we shall assume that E is locally free on B(O,2). Then there exists a global generator $\{\sigma_1, \cdots, \sigma_r\}$ of E on B(O,2), where $r = \operatorname{rank} E$. Then we see that every $t \in B$, the fiber of the vector bundle $(f_{\varepsilon})_* \mathcal{O}_{X(\varepsilon)}(K_{X(\varepsilon)/B} + L) \otimes \mathcal{I}(h_L)$ at t is canonically isomorphic to $\mathbb{C}^r \times \mathcal{O}(B(t,\varepsilon))$ in terms of the frame $\{\sigma_1, \cdots, \sigma_r\}$. And moreover the space $\mathcal{O}(B(t,\varepsilon))$ is canonically isomorphic to $\mathcal{O}(B(O,\varepsilon))$ by the parallel translation. In this case

$$E_{\varepsilon} := (f_{\varepsilon,(2)})_* \mathcal{O}_{X(\varepsilon)}(K_{X(\varepsilon)/B} \otimes p_1^* L \otimes \mathcal{I}(p_1^* h))$$

is a vector bundle of infinite rank on B, where

$$p_1:X(\varepsilon)\longrightarrow X$$

denotes the first projection

$$p_1(x,t) = x, \quad (x,t) \in X(\varepsilon)$$

and $(f_{\varepsilon,(2)})_*\mathcal{O}_{X(\varepsilon)}(K_{X(\varepsilon)/B}\otimes p_1^*L\otimes\mathcal{I}(p_1^*h))$ denotes the direct image of L^2 -holomorphic sections.

By the same proof as Theorem 5.3, we see that the curvature current of $h_{E,\varepsilon}$ is well defined everywhere on B and is semipositive in the sense of Nakano. Letting ε tend to 0, the curvature $\Theta_{h_{E_{\varepsilon}}}$ converges to the curvature $\Theta_{h_{E}}$ operating on $E_{t}\otimes \mathcal{O}_{B,t}$ in the obvious manner for every t such that f is smooth over t. Hence $\Theta_{h_{E}}$ is semipositive in the sense of Nakano.

This completes the proof of Theorems 5.4 and 5.5. \square

5.3 Proof of Theorem 1.2

Let $f: X \longrightarrow S$ be a flat projective family with connected fibers such that a general fiber of f has only canonical singularities. We assume X is normal and S is smooth. Let n denote the relative dimension of $f: X \longrightarrow S$. We set

 $S^{\circ} := \{ s \in S \mid X_s := f^{-1}(s) \text{ is reduced and has only canonical singularities} \}.$

Then $K_{X/S}$ has a canonial AZD h over S such that Θ_h is semipositive on X (Theorem 4.6). And over S° , F_m is locally free by [Ka3, S2, T3].

Then the hermitian inner product h_{F_m} defined by

$$h_{F_m}(\sigma, \sigma')_s := (\sqrt{-1})^{n^2} \int_{X_s} \sigma \wedge \bar{\sigma}' \cdot h^{m-1}$$

is continuous on S° , because $h^m(\sigma, \sigma')$ is bounded on X_s by the construction of h (cf. Proof of Theorem 4.6). And the curvature current of h_{F_m} is semipositive in the sense of Nakano on S° by Theorem 5.5 (here X need not be smooth but Theorem 5.5 is applicable by resolution and the continuity of metrics ([Ka3, S2, T3])). This completes the proof of Theorem 1.2. \square

Remark 5.7 In Theorem 1.2, we have only considered the relative canonical AZD h constructed in Theorem 4.6. But of course the smilar statement holds for the L^2 -inner product:

$$h'_{F_m}(\sigma,\sigma')_s:=(\sqrt{-1})^{n^2}\int_{X_s}\sigma\wedge\bar{\sigma}'\cdot h_m^{\frac{m-1}{m}},$$

where h_m is as in Theorem 1.1. This may be useful, since h_m has only algebraic singularities. One may also consider many other variants. See also Remark 1.3. \square

References

[B-T] Bedford, E. and Taylor, B.A.: A new capacity of plurisubharmonic functions, Acta Math. 149 (1982), 1-40.

- [B1] Berndtsson, B.: Subharmonicity properties of the Bergman kernel and some other functions associated to pseudoconvex domains, math.CV/0505469 (2005).
- [B2] Berndtsson, B.: Curvature of vector bundles and subharmonicity of vector bundles, math.CV/050570 (2005).
- [B3] Berndtsson, B.: Curvature of vector bundles associated to holomorphic fibrations, math.CV/0511225 (2005).
- [B-P] Berndtsson, B. and Paun, M.: Bergman kernels and the pseudoeffectivity of relative canonical bundles, math.AG/0703344 (2007).
- [B-C-H-M] Birkar, C, Cascini, P., Hacon, C., McKernan, J.: Existence of miniaml models for varieties of log general type, math. AG/0610203 (2006).
- [D-P-S] Demailly, J.P.-Peternell, T.-Schneider, M.: Pseudo-effective line bundles on compact Kähler manifolds, math. AG/0006025 (2000).
- [E] Elkik, R.: Singularités rationelles et déformations, Invent. math. 47 (1978), 139-147.
- [Fu1] Fujita, T.: On the structure of polarized varieties with Δ -genera zero. J. Fac. Sci. Univ. Tokyo Sect. IA Math. **22** (1975), 103–115.
- [Fu2] Fujita, T.: Approximating Zariski deecomposition of big line bundle , Kodai Math. J. 17 (1994), 1-4.
- [G-R] Grauert, H., Remmert, R.: Plurisubharmonische Funktionen in komplexen Räumen. Math. Z. 65 175–194 (1956).
- [Ka1] Kawamata, Y.: Kodaira dimension of Algebraic giber spaces over curves, Invent. Math. 66 (1982), pp. 57-71.
- [Ka2] Kawamata, Y., Minimal models and the Kodaira dimension, Jour. für Reine und Angewande Mathematik 363 (1985), 1-46.
- [Ka3] Kawamata, Y.: Deformation of canonical singularities, Jour. of A.M.S. 12 (1999), 85-92.
- [K-M] Kollár, J.-Mori, S.: Birational geometry of algebraic varieties, Cambridge Tracts in Math., Cambridge University Press (1998).
- [Kr] Krantz, S.: Function theory of several complex variables, John Wiley and Sons (1982).
- [L] Lelong, P.: Fonctions Plurisousharmoniques et Formes Differentielles Positives, Gordon and Breach (1968).
- [M-Y] Maitani, and Yamaguchi, S.: Variation of Bergman metrics on Riemann surfaces, Math. Ann. **330** (2004) 477-489.
- [Mo] Mori, S.: Flip conjecture and the existence of minimal model for 3-folds, J. of A.M.S. 1 (1988), 117-253.

- [N-T] Nadel, A.M. and Tsuji, H.: Compactification of complete Kähler manifolds of negative Ricci curvature, Jour. Diff. Geom. 28 (1988), 503-512.
- [N] Nadel, A.M.: Multiplier ideal sheaves and existence of Kähler-Einstein metrics of positive scalar curvature, Ann. of Math. 132 (1990),549-596.
- [N-S] Narashimhan, M.S. and Simha, R.R.: Manifolds with ample canonical class, Invent. math. 5 (1968), 120-128.
- [O] Ohsawa, T.: On the extension of L^2 holomorphic functions V, effects of generalization, Nagoya Math. J. 161(2001) 1-21.
- [O-T] Ohsawa, T. and Takegoshi, K.: L^2 -extension of holomorphic functions, Math. Z. 195 (1987),197-204.
- [R] Ried, M.: Canonical 3-folds, In Géométric Algébrique, Angers 1979, Sijthoff and Nordhoff International Publishers, Alphen and Rijn (1980), 273-310.
- [S1] Siu, Y.-T.: Invariance of plurigenera, Invent. Math. 134 (1998), 661-673.
- [S2] Siu, Y.-T.: Extension of twisted pluricanonical sections with plurisubharmonic weight and invariance of semipositively twisted plurigenera for manifolds not necessarily of general type, Collected papers Dedicated to Professor Hans Grauert (2002), pp. 223-277.
- [T1] Tsuji H.: Analytic Zariski decomposition, Proc. of Japan Acad. 61(1992), 161-163.
- [T2] Tsuji, H.: Existence and Applications of Analytic Zariski Decompositions, Trends in Math. Analysis and Geometry in Several Complex Variables, (1999), 253-272.
- [T3] Tsuji, H.: Deformation invariance of plurigenera, Nagoya Math. J. 166 (2002), 117-134.
- [T4] Tsuji, H.: Refined semipositivity and Moduli of canonical models, preprint, (2005).
- [T5] Tsuji, H.: Variation of Bergman kernels of adjoint line bundles, math.CV/0511342 (2005).
- [T6] Tsuji, H.: Dynamical construction of Kähler-Einstein metrics, math.AG/0606023 (2006).

Author's address
Hajime Tsuji
Department of Mathematics
Sophia University
7-1 Kioicho, Chiyoda-ku 102-8554
Japan
e-mail address: tsuji@mm.sophia.ac.jp