

# ON EXTENSIONS OF GROUP WITH INFINITE CONJUGACY CLASSES, I

JEAN-PHILIPPE PRÉAUX<sup>1 2</sup>

ABSTRACT. We give a characterization of the group property of being with infinite conjugacy classes (or *icc*, *i.e.*  $\neq 1$  and of which all conjugacy classes beside 1 are infinite) for extensions of abelian, centerless, *icc*, or word hyperbolic groups.

## INTRODUCTION

A group is said to be with *infinite conjugacy classes* (or *icc*) if it is non trivial, and if all its conjugacy classes beside  $\{1\}$  are infinite. This property is motivated by the theory of Von Neumann algebra, since for any group  $\Gamma$ , a necessary and sufficient condition for its Von Neumann algebra  $W_\lambda^*(\Gamma)$  to be a type  $II - 1$  factor is that  $\Gamma$  be *icc* (cf. [ROIV]).

The property of being *icc* has been characterized in several classes of groups : 3-manifolds and  $PD(3)$  groups in [HP], groups acting on Bass-Serre trees in [Co], wreath products, semi-direct products and finite extensions in [P1, P2, P3]. We will focus here on groups defined by some specific extensions of a group, namely extensions of abelian groups, of centerless groups, of *icc* groups, or of word hyperbolic groups. We give a necessary and sufficient condition for those groups to satisfy the *icc* property.

We first consider the case of extensions of abelian groups. In a group  $G$ ,  $FC(G)$  denotes the unions of finite conjugacy classes in  $G$  ; it turns out that  $FC(G)$  is a characteristic subgroup of  $G$ . Obviously,  $G$  is *icc* if and only if  $FC(G) = 1$ .

**Theorem 1.** *Let  $G \neq 1$  be a group defined by an extension of an abelian group :*

$$1 \longrightarrow K \text{ abelian} \longrightarrow G \longrightarrow Q \longrightarrow 1$$

*and  $\theta : Q \longrightarrow \text{Aut}(K)$  the associated homomorphism. Then  $G$  is *icc* if and only if both :*

- (i)  $K \setminus 1$  contains only infinite  $\theta(Q)$ -orbits,
- (ii) the induced homomorphism  $\theta : FC(Q) \longrightarrow \text{Aut}(K)$  is injective.

On the opposite, we consider the case of extensions of centerless groups. For any extension  $G$  of a group  $K$  we consider the homomorphism  $\pi : G \longrightarrow \text{Aut}(K)$  defined by, for all  $g \in G$ ,  $k \in K$ ,  $\pi(g)(k) = k^g$ .

**Theorem 2.** *Let  $G \neq 1$  be a group defined by an extension of a centerless group :*

$$1 \longrightarrow K \text{ centerless} \longrightarrow G \longrightarrow Q \longrightarrow 1$$

*and  $\theta : Q \longrightarrow \text{Out}(K)$  the associated homomorphism. Then  $G$  is *icc* if and only if both :*

- (i)  $K$  does not contain a non trivial normal subgroup  $N$  either finite or  $\mathbb{Z}^n$  which is preserved by  $\pi(G)$  and contains only finite  $\pi(G)$ -orbits,
- (ii) the induced homomorphism  $\theta : FC(Q) \longrightarrow \text{Out}(K)$  is injective.

<sup>1</sup>Centre de Recherche de l'Armée de l'air, Ecole de l'air, F-13661 Salon de Provence air

<sup>2</sup>Centre de Mathématiques et d'Informatique, Université de Provence, 39 rue F.Joliot-Curie, F-13453 marseille cedex 13

*E-mail :* preaux@cmi.univ-mrs.fr

*Mathematical subject classification :* 20E45 , 20E22

Since in condition (i)  $N$  necessarily lies in  $FC(K)$ , and since an icc group is centerless, it follows immediatly :

**Theorem 3.** *Let  $G$  be a group defined by an extension of an icc group :*

$$1 \longrightarrow K_{icc} \longrightarrow G \longrightarrow Q \longrightarrow 1$$

and  $\theta : Q \longrightarrow Out(K)$  the associated homomorphism. Then  $G$  is icc if and only if the induced homomorphism  $\theta : FC(Q) \longrightarrow Out(K)$  is injective.

Together with a few observations on word hyperbolic groups, one obtains as another corollary the following characterization of icc property for extensions of word hyperbolic groups.

**Theorem 4.** *Let  $G$  be defined by an extension of a non trivial word hyperbolic group :*

$$1 \longrightarrow K_{hyperbolic} \longrightarrow G \longrightarrow Q \longrightarrow 1$$

and  $\theta : Q \longrightarrow Out(K)$  the associated homomorphism. Then  $G$  is icc if and only if both :

- (i)  $K$  is icc,
- (ii) the induced homomorphism  $\theta : FC(Q) \longrightarrow Out(K)$  is injective.

## 1. PROOF OF THE RESULTS.

Let  $G$  be a group,  $H$  a non empty subset of  $G$ , and  $u, g \in G$ . We use the notations  $u^g = g^{-1}ug$  and  $u^H = \{u^g | g \in H\}$ , so that  $g^G$  denotes the conjugacy class of  $g$  in  $G$  ;  $Z_G(g)$  stands for the centralizer of  $g$  in  $G$  and  $Z(G)$  for the center of  $G$ . It follows immediatly that  $g^G$  is finite if and only if  $Z_G(g)$  has a finite index in  $G$ . Let  $\rho : G \longrightarrow Q$  denote the projection given by the extension.

*Proof of theorem 1.* Note that that the conjugacy class in  $G$  of  $k \in K$ , coincide with its  $\theta(Q)$ -orbit. Obviously  $G$  icc implies that both conditions (i) and (ii) are satisfied. We prove the converse by showing that if  $G$  is not icc then either condition (i) or condition (ii) is not satisfied. So suppose that  $G$  is not icc ; since  $G \neq 1$  there exists  $u \neq 1$  in  $G$  such that its conjugacy class  $u^G$  in  $G$  is finite. If  $u \in K$  ; then condition (i) is not satisfied, so suppose in the following that  $u \in G \setminus K$ . Then  $\rho(u)$  lies in  $FC(Q)$ , for  $\rho(u)^Q = \rho(u^G)$ , and  $\rho(u) \neq 1$ . Let  $k \in K$ , then  $[u, k] \in K$  has a finite conjugacy class in  $G$ , for  $Z_G(u)$  and  $Z_G(kuk^{-1})$  both have a finite index in  $G$  and their intersection lies in  $Z_G([u, k])$ . So that either, for all  $k \in K$ ,  $[u, k] = 1$ , which implies that  $\pi(u) = \theta(\rho(u)) = Id_K$  and then  $\theta : FC(Q) \longrightarrow Aut(K)$  is non injective, or as above, condition (i) is not satisfied.  $\square$

*Proof of theorem 2.* Suppose that  $G$  is not icc ; since  $G \neq 1$  there exists  $u \neq 1$  in  $G$  such that  $u^G$  is finite. If  $u$  lies in  $K$ , let  $N'$  be the subgroup of  $K$  generated by  $u^G$  ;  $N'$  is a non trivial subgroup of  $FC(K)$  preserved by  $\pi(G)$  and with only finite  $\pi(G)$ -orbits. Moreover  $N'$  is a finitely generated  $FC$ -group and hence a finite central extension of a f.g. abelian group (cf. [Ne]) ; it follows that the set  $Tor(N')$  of torsion elements of  $N'$  is a characteristic finite subgroup of  $N'$  and  $N'/Tor(N')$  is  $\mathbb{Z}^n$ . Hence condition (i) is not satisfied by picking  $N = Tor(N')$ , if  $Tor(N') \neq 1$ , and  $N = N'$  otherwise. If  $u$  lies in  $G \setminus K$ , then  $\rho(u)$  is a non trivial element of  $FC(Q)$ . For any  $k \in Q$ ,  $[u, k]$  lies in  $K$  and has a finite conjugacy class in  $G$ , so that, as in the proof of theorem 1, we can suppose that  $\pi(u)$  is the identity, and hence  $\theta(\rho(u)) = 1$ , for otherwise condition (i) is not satisfied. It follows that  $\theta : FC(Q) \longrightarrow Out(K)$  is non injective, so that condition (ii) is not satisfied. We have proved that if both condition (i) and (ii) are satisfied then  $G$  is icc ; we now prove the converse. If  $K$  contains a non trivial subgroup preserved by  $\pi(G)$  and with only finite  $\pi(G)$ -orbits then each element of  $N$  has a finite conjugacy class in  $G$  and then  $G$  is not icc. If  $\theta : FC(Q) \longrightarrow Out(K)$  is non injective, let  $q \neq 1$  in  $FC(Q)$  be such that  $\theta(q) = 1$ . Then there exists a lift  $\bar{q}$  of  $q$  in  $G$  such that  $\pi(\bar{q})$  is the identity of  $K$ , so that

$K \subset Z_G(\bar{q})$ . For any  $q_1 \in Z_Q(q)$ , if  $\bar{q}_1$  denotes a lift of  $q_1$  in  $G$ , then  $\bar{q}_1 \bar{q} \bar{q}_1^{-1} = \bar{q}k$  for some  $k \in K$ , which must belong to  $Z(K)$  since  $\pi(\bar{q}_1 \bar{q} \bar{q}_1^{-1}) = \pi(k)$  is the identity on  $K$ . Since  $Z(K) = 1$ ,  $Z_G(\bar{q})$  contains  $\rho^{-1}(Z_Q(q))$ , and hence has a finite index in  $G$ , so that  $G$  is not icc.  $\square$

*Proof of theorem 4.* If  $K$  is a non trivial elementary group then  $K$  is not icc, and neither  $G$ , for  $\text{Aut}(K)$  is finite. So that the assumption in theorem 4 is satisfied and we suppose in the following that  $K$  is non elementary. A non elementary word hyperbolic has a finite center (cf. [Gr]). It follows that each element of  $Z(K)$  has a finite conjugacy class both in  $K$  and in  $G$ . Then, if  $Z(K) \neq 1$  neither  $K$  nor  $G$  are icc so that the assumption of theorem 4 is satisfied. So suppose in the following that  $K$  is centerless, so that theorem 2 applies. In a non elementary word hyperbolic group  $K$ ,  $FC(K)$  turns out to be a finite characteristic subgroup of  $K$ . Hence theorem 2 shows that  $G$  is icc if and only both  $\theta : FC(Q) \rightarrow \text{Out}(K)$  is injective and  $FC(K) = 1$ .  $\square$

## REFERENCES

- [Co] Y.DE CORNULIER, *Infinite conjugacy classes in groups acting on trees*, preprint (2005).
- [Gr] M.GROMOV, *Hyperbolic groups* in "Essays in group theory", MSRI Publications, Springer (1987), 75–263.
- [HP] P.DE LA HARPE et J.-P.PRÉAUX, *Groupes fondamentaux des variétés de dimension 3 et algèbres d'opérateurs*, preprint arXiv:math.GR/0509449 v1 (2005).
- [ROIV] F.J.MURRAY et J.VON NEUMANN, *On rings of operators*, IV, *Annals of Math.* **44** (1943), 716–808.
- [Ne] W.NEUMANN, *Groups with only finite conjugacy classes*, *Proc. of the L.M.S.* **1** (1951), 178–187.
- [P1] J.-P.PRÉAUX, *Wreath product of groups with infinite conjugacy classes*, preprint (2006), arXiv:math.GR/0612685.
- [P2] J.-P.PRÉAUX, *Finite extension of group with infinite conjugacy classes*, preprint (2007), arXiv:math.GR/0703314.
- [P3] J.-P.PRÉAUX, *Split extensions of group with infinite conjugacy classes*, preprint (2007).