

Stably Just Infinite Rings

Jason Bell¹

*Department of Mathematics
Simon Fraser University
Burnaby, BC V5A 1S6. Canada*

John Farina^{*}

*Department of Mathematics
University of California, San Diego
La Jolla, CA 92093*

Cayley Pendergrass-Rice²

*Department of Mathematics
Albion College
Albion, MI 49224*

Abstract

We study just infinite algebras which remain so upon extension of scalars by arbitrary field extensions. Such rings are called *stably just infinite*. We show that just infinite rings over algebraically closed fields are stably just infinite provided that the ring is either right noetherian (4.2) or countably generated over a large field (6.4). We give examples to show that, over countable fields, a just infinite algebra which is either affine or non-noetherian need not remain just infinite under extension of scalars. We also give a concrete classification of PI stably just infinite rings (5.5) and give two characterizations of non-PI stably just infinite rings in terms of Martindale's *extended center* (3.4, 3.5).

Key words: just infinite, stably just infinite, scalar extension, extended center, extended centroid, central closure.

1991 MSC: 16N60, 16P40

^{*} Corresponding author.

Email addresses: jpb@math.sfu.ca (Jason Bell), jfarina@math.ucsd.edu (John Farina), cpendergrass@albion.edu (Cayley Pendergrass-Rice).

¹ The first author thanks NSERC for its generous support.

² The third author received partial support from a grant from the Hewlett Mellon

1 Introduction

Simple rings are so-called because their two-sided ideal structure is as simple as possible. A natural way of generalizing the notion of a simple ring is to allow the presence of two-sided ideals, but to insist that the nonzero ones be very “large.” Specifically, we consider algebras for which all nonzero ideals have finite codimension. We begin, as always, with a definition.

Definition 1.1 A k -algebra A is called *just infinite dimensional*, or *just infinite* for short, if $\dim_k(A) = \infty$ and each of its nonzero two-sided ideals has finite codimension.

The nomenclature which has been adopted is to call such rings *projectively simple* in case the ring is \mathbb{N} -graded. There are many examples of just infinite algebras. Aside from any infinite dimensional simple ring, the most immediate example is a polynomial ring in one variable $k[x]$. Reichstein, Rogalski, and Zhang [15] give non-PI examples constructed using abelian varieties and Bartholdi [1] constructs interesting just infinite rings using groups acting on infinite trees.

Much is known for just infinite algebras over uncountable fields; for example, Farkas and Small [6] show they are either primitive, have nonzero Jacobson radical, or satisfy a polynomial identity. Over countable fields much less is known and many of the basic questions that are easily answered over uncountable fields are still open. For this reason we are interested in when we can extend scalars in a just infinite algebra and remain just infinite. We thus give the following definition.

Definition 1.2 A just infinite k -algebra A is called *stably just infinite* if $A \otimes_k K$ is just infinite over K , for every field extension K/k .

Just infinite algebras, even affine ones, may behave rather poorly upon extension of scalars. For example, $\mathbb{C}[x]$ is an affine just infinite \mathbb{R} -algebra, but $\mathbb{C}[x] \otimes_{\mathbb{R}} \mathbb{C}$ contains $\mathbb{C} \otimes_{\mathbb{R}} \mathbb{C}$, which is not a domain. Thus $\mathbb{C}[x] \otimes_{\mathbb{R}} \mathbb{C}$ is not even prime and hence $\mathbb{C}[x]$ fails to be just infinite over \mathbb{R} [5]. This example shows that in case the ground field is not algebraically closed, things can go horribly awry even upon extending scalars by a finite dimensional field extension! Bergman proved the tensor product of any two prime k -algebras is again prime when k is algebraically closed [3, Proposition 17.2]. It is therefore reasonable to restrict our attention to the case that the ground field is algebraically closed. In this way we can, at the very least, avoid the unpleasantness of having a just infinite algebra A for which extension of scalars results in a ring which fails to be prime.

Fund for Faculty Development at Albion College.

We summarize our main results: We show that right noetherian just infinite algebras are well-behaved under extension of scalars.

Theorem 1.3 *Let k be an algebraically closed field and let A be a non-PI right noetherian just infinite k -algebra. Then A is stably just infinite.*

In addition, we show that the hypothesis that A is a right noetherian ring cannot be weakened when the ground field is countable—even if the algebra is finitely generated.

Theorem 1.4 *Let k be a countable algebraically closed field. Then there exists an affine non-simple right Goldie just infinite k -algebra which is not stably just infinite.*

It is necessary to take k to be a countable algebraically closed field in this theorem. In fact, we show that if A is a countably generated just infinite k -algebra over an uncountable algebraically closed field, then A is stably just infinite.

Theorem 1.5 *Let k be an uncountable algebraically closed field and let A be a countably generated just infinite k -algebra. Then A is stably just infinite.*

We remark that this gives an analog of a result of Reichstein et al. [15, Lemma 1.8] for ungraded rings. To complete our study of stably just infinite algebras, we look at the PI case and show that a PI just infinite k -algebra A is stably just infinite if and only if the center of the quotient division algebra of A is a finitely generated field extension of k of transcendence degree 1.

This paper is organized as follows. In §2 and §3, we give some useful facts about just infinite algebras and extended centers, which we later use in our consideration of stably just infinite algebras. In §4, we prove Theorem 1.3. In §5, we present our characterization of PI stably just infinite algebras. In §6, we turn our attention to just infinite algebras over uncountable fields and prove Theorem 1.5. Finally, in §7, we construct interesting just infinite algebras that are not stably just infinite to obtain Theorem 1.4.

All rings under consideration will be associative algebras, with 1, over a field k , and homomorphisms are assumed to be unital.

2 Just Infinite Algebras

Reichstein et al. [15] show that a non-PI projectively simple algebra has a unique maximal ideal, namely the augmentation ideal. A similar result holds for just infinite algebras with nonzero Jacobson radical.

Proposition 2.1 *Let A be a just infinite k -algebra with nonzero Jacobson radical. Then A has only finitely many prime ideals.*

Proof. We denote the Jacobson radical of the algebra A by $J(A)$. Since A is just infinite and $J(A) \neq (0)$, $A/J(A)$ is finite dimensional. It follows that there are only finitely many prime ideals in A that properly contain $J(A)$ [18, Theorem 2.3.9 (ii)]. But every nonzero prime ideal in A is necessarily maximal and thus primitive and so it contains $J(A)$. The result follows. \square

The hypothesis that $J(A) \neq (0)$ is necessary, as $\mathbb{C}[x]$ demonstrates. Proposition 2.1 motivates the following

Question 2.2 If A is a finitely generated non-PI just infinite k -algebra, does A have only finitely many prime ideals?

It is known that a PI just infinite algebra is a finitely generated module over its center and the center is itself just infinite [5]. In stark contrast to the PI case, a just infinite algebra that doesn't satisfy a polynomial identity necessarily has a small center (assuming the ring isn't simple). In fact, we next show that the center of any non-simple non-PI just infinite algebra is a finite dimensional field extension of the ground field. This result is essentially a consequence of two lemmas of Farkas and Small [6]. For a fixed positive integer n , we denote by $I_n(A)$ the two-sided ideal of the algebra A generated by all specializations in A of all polynomial identities of $n \times n$ matrices.

Proposition 2.3 *Let A be a non-PI just infinite k -algebra that is not simple. Then $Z(A)$ is a finite dimensional field extension of k .*

Proof. Since A is just infinite, it is prime [5], and so $Z(A)$ is a domain. Choose $0 \neq z \in Z(A)$ and set $J = \bigcap_{d \in \mathbb{N}} Az^d$. Since A is just infinite, the two-sided ideal Az has finite codimension in A . A result of Farkas and Small [6, Lemma 1.1] gives $I_n(A) \subseteq J$ for some n . Since A doesn't satisfy a polynomial identity, $I_n(A)$ is a nonzero ideal of A , and so J is nonzero as well. It follows that z is a unit of A [6, Lemma 1.2] and thus is invertible in $Z(A)$. Thus $Z(A)$ is a field extension of k . Next, since A is not simple, there is a nonzero proper ideal I of A . Since $Z(A)$ is a field, $I \cap Z(A) = (0)$ and so the natural projection map $A \rightarrow A/I$ induces an injection from $Z(A)$ to A/I , and the latter is finite dimensional. \square

The fact that the center of a (non-simple) non-PI just infinite algebra is a finite dimensional field extension of k shows that non-PI just infinite algebras are a reasonable generalization of central simple algebras. In particular, showing that a just infinite algebra is stably just infinite can be regarded as an analog of the fact that a central simple algebra remains simple under extension of scalars.

3 Central Closure

To prove our results we make use of the so-called *extended center*. This idea was largely inspired by some of the methods of Resco [16], to which we owe a great debt.

Let A be a prime ring and consider the set of all right A -module homomorphisms $f: I \rightarrow A$, where I ranges over all nonzero two-sided ideals of A . Martindale [13] shows how one can endow this set of maps with a useful algebra structure. For the reader's convenience we include the full definition of Martindale's ring of quotients, but we refer to other sources [8, 10, 18] for the proof that this construction actually yields a ring with the stated properties.

Definition 3.1 Let A be a prime k -algebra. The *(right) Martindale ring of quotients of A* , denoted $Q_r(A)$, consists of equivalence classes of pairs (I, f) where $I \triangleleft A$, $I \neq (0)$, and $f \in \text{Hom}_A(I_A, A_A)$. Here two pairs (I, f) , (J, g) are defined to be equivalent if $f = g$ on the intersection $I \cap J$. Addition and multiplication are given by

$$\begin{aligned} (I, f) + (J, g) &= (I \cap J, f + g), \\ (I, f) \cdot (J, g) &= (JI, f \circ g). \end{aligned}$$

Definition 3.2 Let A be a prime k -algebra. The *extended center of A* , written $C(A)$, is defined to be $Z(Q_r(A))$.

Remark 3.3 $C(A)$ is also called the *extended centroid* in the literature. Hopefully our terminology won't cause confusion, as the centroid of a unital ring coincides with its center. $C(A)$ is a field extension of k and $C(A) \cap A = Z(A)$. The *central closure of A* , denoted $AC(A)$, is the $C(A)$ -linear subspace of $Q_r(A)$ generated by A . Lastly, A is called *centrally closed* if $C(A) = Z(A)$, (equivalently, if $AC(A) = A$). Note that the central closure of any ring is itself centrally closed.

There is an entirely internal characterization of $C(A)$ which bears mentioning. The extended center of A consists precisely of those pairs (I, f) where $f: I \rightarrow A$ is an (A, A) -bimodule homomorphism:

$$C(A) = \{ (I, f) \mid (0) \neq I \triangleleft A, f \in \text{Hom}_A({}_A I_A, {}_A A_A) \}.$$

Centrally closed prime algebras behave particularly well with respect to extension of scalars, and as a result we can show that a non-PI just infinite algebra is stably just infinite if and only if it is centrally closed.

Theorem 3.4 *Let k be algebraically closed, and let A be a non-PI just infinite algebra with $Z(A) = k$. Then A is stably just infinite iff A is centrally closed.*

Proof. When A is simple (and hence centrally closed) the result follows from the well-known fact that central simple rings remain simple upon extension of scalars. Thus we may assume that A is not simple. Suppose that A is centrally closed. Choose a nonzero ideal $I \triangleleft A \otimes_k K$ and set $J = I \cap A$, so $J \otimes_k K \subseteq I$. Since A is centrally closed, J is a nonzero ideal of A [4, Lemma 3.4]. Since

$$(A \otimes_k K)/(J \otimes_k K) \cong A/J \otimes_k K$$

and the latter is finite dimensional over K , we see that I has finite codimension in $A \otimes_k K$.

For the other direction, suppose that A is stably just infinite but not centrally closed. Choose $(\varphi, I) \in C(A) \setminus k$. We may assume that I is generated by a single element a , and so φ is completely determined by $b := \varphi(a)$. Since $(\varphi, I) \notin k$, we may further assume that $b \neq \lambda a$ for any $\lambda \in k$. Set $B := A \otimes_k k(t)$ where t is an indeterminate. We identify B with the localization of $A[t]$ obtained by inverting all nonzero polynomials in t with coefficients in k . Consider the ideal $(a+tb)$ of B generated by $a+tb$. We will show that $A \cap (a+tb) = (0)$, so suppose there is some nonzero element $c \in A \cap (a+tb)$. We may write $c = \sum_i r_i(a+tb)s_i$, with $r_i, s_i \in B$. If $q(t) \in k[t]$ is a common right denominator for the r_i and s_i , we see that

$$cq(t) = \sum_i f_i(t)(a+tb)g_i(t), \quad \text{for some } f_i, g_i \in A[t].$$

Now, writing $f_i(t) = \sum_j p_{ij}t^j$ and $g_i(t) = \sum_l q_{il}t^l$ where $p_{ij}, q_{il} \in A$, we see that

$$\begin{aligned} cq(t) &= \sum_{i,j,l} p_{ij}aq_{il}t^{j+l} + p_{ij}bq_{il}t^{j+l+1} \\ &= \sum_{i,j,l} p_{ij}aq_{il}t^{j+l} + t \sum_{i,j,l} p_{ij}\varphi(a)q_{il}t^{j+l}. \end{aligned}$$

Set $u(t) := \sum_{i,j,l} p_{ij}aq_{il}t^{j+l} \in A[t]$, so

$$cq(t) = u(t) + t\varphi(u(t)),$$

where here φ denotes the (A, A) -bimodule endomorphism of $A[t]$ defined by $\varphi(a) = b$ and $\varphi(t) = t$. Note that for all $\lambda \in k$,

$$cq(\lambda) = u(\lambda) + \lambda\varphi(u(\lambda)) \in A,$$

and so for all $\lambda \in k$ we have

$$c(q(\lambda) - q(0)) = u(\lambda) + \lambda\varphi(u(\lambda)) - u(0). \quad (\dagger)$$

Let V denote the k -span of $\{u(\lambda) \mid \lambda \in k\}$, $W = V + kc$, and note that W is

a nonzero subspace of A with $\dim_k(W) < \infty$. Moreover, by equation (\dagger) ,

$$\varphi(W) \subseteq W,$$

and so φ is algebraic over k by the Cayley-Hamilton theorem. Since k is algebraically closed, we see that $\varphi \in k$. Thus $A \cap (a + tb) = (0)$ and hence A embeds in $B/(a + tb)$. However, A is stably just infinite so $B/(a + tb)$ is finite dimensional over $k(t)$, and thus A is PI. This contradiction completes the proof. \square

The careful reader might note that we only used the non-PI hypothesis in one direction of the preceding proof. But note that if A is PI and centrally closed, then the center of $Q(A)$ is reduced to scalars. Thus $Q(A)$ is finite dimensional over k by Kaplansky's theorem and hence A cannot be just infinite.

If we remove the condition that $Z(A) = k$ in Theorem 3.4, then it seems that to decide whether or not a non-PI just infinite algebra is stably just infinite requires consideration of arbitrary field extensions K/k . In fact, the next proposition shows that one only need consider the extended center.

Proposition 3.5 *Let A be a non-PI just infinite k -algebra, and let $C(A)$ denote the extended center of A . Then A is stably just infinite iff $A \otimes_k C(A)$ is just infinite over $C(A)$.*

Proof. One direction is trivial. For the other, suppose that $A \otimes_k C(A)$ is just infinite over $C(A)$. Note that we have a surjective k -algebra homomorphism from $A \otimes_k C(A)$ onto the central closure of A (given by multiplication). Since A is just infinite and not PI, $\dim_{C(A)} AC(A) = \infty$, and thus the above map is an isomorphism: $A \otimes_k C(A) \cong AC(A)$. We see that $AC(A)$ is just infinite over $C(A)$. We claim then that $AC(A)$ is stably just infinite over $C(A)$. If A is simple then $C(A) = Z(A)$, so $AC(A) = A$ is central simple over $C(A)$ and thus A is stably just infinite. If A is not simple, then, as in the first part of the proof of Theorem 3.4, it is easy to see that $AC(A)$ is stably just infinite over $C(A)$.

Now, let K/k be any field extension, and let L denote a compositum (over k) of K and $C(A)$. We then have that

$$A \otimes_k L \cong (A \otimes_k C(A)) \otimes_{C(A)} L$$

is just infinite over L , and Lemma 5.2 shows that $A \otimes_k K$ is just infinite over K , completing the proof. \square

Theorem 3.4 and Proposition 3.5 give entirely internal characterizations of non-PI stably just infinite algebras. The difficulty is that in practice it is often hard to compute the extended center.

4 Noetherian Just Infinite Algebras

In this section we prove that a non-PI right noetherian just infinite algebra over an algebraically closed field is stably just infinite. To obtain this result, we use central closure.

Proposition 4.1 *Let A be a right noetherian non-PI just infinite algebra with center k . Then $C(A)$ is an algebraic extension of k .*

Proof. Since A is right Goldie, the right maximal ring of quotients of A coincides with its right Goldie quotient ring, which we denote by $Q(A)$. Since A is prime, $C(A) = Z(Q(A))$ [11, Prop 14.17]. Choose a nonzero element $t \in C(A)$ and let B be the subalgebra of $Q(A)$ generated by A and t . Let $p = p(t)$ be any nonzero polynomial in t with coefficients in k and let $\pi : B \rightarrow B/pB$ denote the natural projection map. There are nonzero elements $a, b \in A$ with b regular such that $p = ab^{-1}$ and so $pb = a$ in B . Hence $pB \cap A \neq (0)$. Moreover,

$$B/pB \cong \left(\frac{A}{pB \cap A} \right) \{\bar{t}\}, \quad \text{where } \bar{t} = t + pB.$$

Thus every element of B/pB can be written as a polynomial in \bar{t} of degree $< \deg(p)$ with coefficients in $A/(pB \cap A)$. Since A is just infinite, this shows that $\dim_k(B/pB) < \infty$. Set $n = \dim_k(B/pB)$. By a result of Farkas and Small [6, Lemma 1.1], $I_n(B) \subseteq \bigcap_{d \geq 1} p^d B$, and since A , and hence also B , is not PI, $I_n(B) \neq (0)$.

Choose any nonzero regular element

$$y \in \bigcap_{d \geq 1} p^d B$$

(such a regular element exists because B is prime right Goldie). Since p is central, we can then find regular elements $a_1, a_2, \dots \in B$ with

$$y = a_d p^d, \quad \text{for all } d \in \mathbb{N},$$

and since p is regular in B , we have $a_d = a_{d+1}p$ for all d . In particular, $a_d B \subseteq a_{d+1} B$. Since B is right noetherian, the chain of right ideals

$$a_1 B \subseteq a_2 B \subseteq \dots$$

must terminate, so $a_i B = a_{i+1} B$ for some i . In particular, $a_i B = a_{i+1} B = a_{i+1} p B$, and since a_{i+1} is regular, the second equality shows that p has a right inverse in B . Since B is right noetherian, p is a unit in B .

Let $\mathfrak{m} \in \text{mspec}(B)$ be any nonzero maximal ideal of B . We have a k -algebra isomorphism

$$\frac{B}{\mathfrak{m}} \cong \left(\frac{A}{\mathfrak{m} \cap A} \right) \{\bar{t}\}, \text{ where } \bar{t} = t + \mathfrak{m}.$$

Since A is just infinite, $A/(\mathfrak{m} \cap A)$ is finite dimensional over k , and since \bar{t} is central, B/\mathfrak{m} is a simple PI ring, and thus a finite module over its center, Z , by Kaplansky's theorem. Since B/\mathfrak{m} is clearly an affine k -algebra, the Artin-Tate lemma then shows that Z is affine as well. But Z is a field, and thus $\dim_k(Z) < \infty$. This shows that $\dim_k(B/\mathfrak{m}) < \infty$.

Finally, since $B/\mathfrak{m} \cong (A/(\mathfrak{m} \cap A))\{\bar{t}\}$ is finite dimensional, we see that \bar{t} is algebraic over k , and we claim that this implies that t is algebraic over k as well. To see this, suppose that t were transcendental. Since any nonzero polynomial $p(t) \in k\{t\}$ is a unit in B , B contains the field $k(t)$, which has transcendence degree 1. Since $\mathfrak{m} \neq B$, $\mathfrak{m} \cap k(t) = (0)$, and so B/\mathfrak{m} contains an isomorphic copy of $k(t)$, namely $k(\bar{t})$. Since \bar{t} is algebraic over k , this is a contradiction. \square

Since being centrally closed and being stably just infinite are equivalent properties for non-PI just infinite algebras, we obtain as a corollary Theorem 1.3.

Corollary 4.2 *Let k be an algebraically closed field, and let A be a right noetherian non-PI just infinite algebra with $Z(A) = k$. Then A is stably just infinite.*

We can obtain further stability results by combining the above ideas with the Nullstellensatz.

Proposition 4.3 *Let k be an algebraically closed field and let A be a primitive non-PI just infinite k -algebra. If A satisfies the Nullstellensatz, then A is stably just infinite.*

Proof. By a result of Martindale [12], the extended center $C(A)$ embeds in $\text{End}_A(M)$ for any faithful simple A -module M . Since A satisfies the Nullstellensatz, this implies that $C(A) = k$, so A is centrally closed and the result follows from Theorem 3.4. \square

5 PI Just Infinite Algebras

In this section, we characterize PI stably just infinite rings over algebraically closed fields. We begin by considering the commutative case.

Lemma 5.1 *Let k be an algebraically closed field and let A be a commutative noetherian domain such that $Q(A)$ is a finitely generated field extension of k*

with $\text{trdeg}(Q(A)) \leq 1$, then A is just infinite.

Proof. We denote the (classical) Krull dimension of a ring A by $\text{Kdim}(A)$. Since A is a domain with $\text{Kdim}(A) \leq 1$, $\text{Kdim}(A/P) = 0$ for every nonzero $P \in \text{spec}(A)$. Thus A/P is a field extension of k . If A is not just infinite, then, since A is noetherian, we may choose an ideal I maximal with respect to $\dim_k(A/I) = \infty$. It is easy to see that I is then prime and it suffices to show that $A/P = k$ for every nonzero prime ideal P of A . Fix a nonzero $P \in \text{spec}(A)$ and suppose there is some element $x + P \in A/P$ which is transcendental over k . Since k is algebraically closed and A is a domain, every nonzero element of P is transcendental over k . Fix a nonzero $t \in P$. Then x and t are algebraically independent over k since x is regular mod P , and so $\text{trdeg}(Q(A)) \geq 2$, a contradiction. Thus we see that A/P is an algebraic extension of k . Since k is algebraically closed field, we see that $A/P \cong k$. \square

To characterize commutative stably just infinite rings, we first require another lemma.

Lemma 5.2 *Let A be a k -algebra and let K/k be a field extension. If $A \otimes_k K$ is just infinite over K , then A is just infinite over k .*

Proof. Choose a nonzero ideal $I \triangleleft A$. Then $I \otimes_k K$ is a nonzero two-sided ideal of $A \otimes_k K$ with $(A \otimes_k K)/(I \otimes_k K) \cong (A/I) \otimes_k K$. Taking dimensions then yields

$$\begin{aligned} \dim_k(A/I) &= \dim_K((A/I) \otimes_k K) \\ &= \dim_K((A \otimes_k K)/(I \otimes_k K)) < \infty. \end{aligned} \quad \square$$

Proposition 5.3 *Let k be an algebraically closed field and let A be a commutative just infinite k -algebra. Then A is stably just infinite iff $Q(A)$ is a finitely generated field extension of k of transcendence degree at most 1.*

Proof. Suppose first that A is stably just infinite. Set $F = Q(A)$, the quotient field of A . If A is stably just infinite, then $A \otimes_k K$ is noetherian for every field extension K/k since a commutative just infinite ring is necessarily commutative. Since $F \otimes_k K$ is a localization of $A \otimes_k K$, it is also noetherian for every field extension K/k . By a result of Vámos [20], F is a finitely generated field extension of k , and it suffices to show that $\text{trdeg}_k(F) \leq 1$. Note that $\text{Kdim}(A \otimes_k F) \leq 1$ since $A \otimes_k F$ is a just infinite F -algebra. Since $F \otimes_k F$ is a localization of $A \otimes_k F$, $F \otimes_k F$ is a just infinite F -algebra as well, and so $\text{Kdim}(F \otimes_k F) \leq 1$, and thus $\text{trdeg}_k(F) \leq 1$ [14, 6.4.8 & 6.6.17].

Next suppose that F is a finitely generated field extension of k with $\text{trdeg}_k(F) \leq 1$. We claim that A is stably just infinite. Let K/k be a field extension, and let \overline{K} denote the algebraic closure of K . Since F is finitely generated, there is a finitely generated purely transcendental extension E/k with $E \subseteq F$ and

$[F : E] < \infty$. Then $\text{trdeg}_k(E) \leq 1$ and so the field of fractions of $E \otimes_k \overline{K}$ is a purely transcendental extension of \overline{K} of transcendence degree at most one. But $Q(F \otimes_k \overline{K})$ is a finite extension of $E \otimes_k \overline{K}$ and thus also has transcendence degree at most one. Since $Q(A \otimes_k \overline{K}) = Q(F \otimes_k \overline{K})$, $A \otimes_k \overline{K}$ has Krull dimension at most 1 over \overline{K} . Applying Lemma 5.1, we see that $A \otimes_k \overline{K}$ is just infinite over \overline{K} . Lemma 5.2 then implies that $A \otimes_k K$ is just infinite over K , completing the proof. \square

Corollary 5.4 *Affine commutative just infinite algebras over algebraically closed fields are stably just infinite.*

Suppose now that A is a PI just infinite algebra over an algebraically closed field k such that $Q(Z(A))$ is a finitely generated field extension of k with $\text{trdeg}_k(Q(Z(A))) \leq 1$. Then A is a finitely generated module over $Z(A)$, which is itself just infinite [5, Corollary 2]. By Proposition 5.3, $Z(A)$ is stably just infinite.

If K/k is a field extension, then $Z(A) \otimes_k K$ is a (noetherian) just infinite K -algebra. Since $A \otimes_k K$ is a finitely generated $Z(A) \otimes_k K$ -module, $A \otimes_k K$ is also noetherian. Choose a nonzero two-sided ideal $I \triangleleft A \otimes_k K$. As $A \otimes_k K$ is prime noetherian, I contains a nonzero regular element x . Since $A \otimes_k K$ is a finite $Z(A) \otimes_k K$ -module, x is algebraic over $Z(A) \otimes_k K$. If

$$x^t + z_{t-1}x^{t-1} + \dots + z_0 = 0, \quad z_j \in Z(A) \otimes_k K$$

has t minimal, then $z_0 \neq 0$. It follows that $z_0 \in I \cap (Z(A) \otimes_k K)$, and so $I \cap (Z(A) \otimes_k K) \neq (0)$. Finally, $(A \otimes_k K)/I$ is a finitely generated module over $(Z \otimes_k K)/(I \cap (Z \otimes_k K))$, hence $A \otimes_k K$ is just infinite over K . To summarize, we have the following

Theorem 5.5 *Let k be an algebraically closed field, and let A be a PI just infinite k -algebra. Then A is stably just infinite iff $Q(Z(A))$ is a finitely generated field extension of k with $\text{trdeg}_k(Q(Z(A))) \leq 1$.*

Corollary 5.6 *Affine PI just infinite algebras over algebraically closed fields are stably just infinite.*

6 Algebras over Uncountable Fields

In this section we show that countably generated just infinite algebras over uncountable algebraically closed fields are stably just infinite, proving Theorem 1.5. We first consider the PI case.

Proposition 6.1 *Let k be an uncountable algebraically closed field and let*

A be a countably generated PI just infinite k -algebra. Then A is stably just infinite.

Proof. We begin by showing we can reduce to the commutative case. Since A is just infinite and PI, we have that A is a finite module over its center and that its center is just infinite [5]. Moreover, since an algebra is countably generated iff it is countable dimensional, we see that $Z(A)$ is countably generated as well. By Theorem 5.5, we may then assume that A is commutative.

Note that $\text{Kdim}(A) \leq 1$ since A is just infinite. In fact, $\text{Kdim}(A) = 1$ because there are no proper field extensions of k which are countably generated as algebras. By Proposition 5.3 it is enough to show that $Q(A)$ is a finitely generated field extension of k of transcendence degree at most 1. A is noetherian, so $A \otimes_k Q(A)$ is noetherian [2, Theorem 1.2]. Thus $Q(A) \otimes_k Q(A)$ is noetherian since it is a localization of $A \otimes_k Q(A)$. It follows from a result of Vámos [20] that $Q(A)$ is a finitely generated field extension of k .

Let R be any affine commutative k -algebra with $Q(R) = Q(A)$. Since A is countably generated, we may choose elements $a_1 b_1^{-1}, a_2 b_2^{-1}, \dots$, with $a_i, b_i \in R$, which generate A as a k -algebra. Set $B = RS^{-1}$, where S is the multiplicatively closed subset of R generated by the set $\{b_i \mid i \in \mathbb{N}\}$. Note that B satisfies the Nullstellensatz since it is countably generated and k is uncountable. Since k is algebraically closed, $B/M \cong k$ for some maximal ideal M of B . Set $P = M \cap R$. Then P is a maximal ideal of R . Note also that R_P , the localization of R at P , contains B and thus also contains A . Hence $\text{Kdim}(A) = \text{trdeg}_k(A)$ [7, Theorem 1.5]. \square

Next we dispose of the non-PI case. The techniques are rather different, and we need a preliminary result which may be of independent interest.

Proposition 6.2 *Let A be a countably generated non-PI just infinite k -algebra. Then $C(A)$ is a countable dimensional field extension of k .*

Proof. If A is simple, then $Z(A)$ is a field. But A is countable dimensional over k , so $Z(A)$ is countable dimensional over k as well, and in a simple ring, $C(A) = Z(A)$.

Now suppose that A is not simple. Since A doesn't satisfy any polynomial identity, each $I_n(A)$ is nonzero. Moreover, A is just infinite, and so every nonzero ideal $I \triangleleft A$ contains some $I_n(A)$. Suppose there is an uncountable collection of elements

$$\{(I_\alpha, f_\alpha) \mid \alpha \in \mathcal{I}\} \subseteq C(A)$$

which is k -linearly independent. Note that if $(I, f) \in C(A)$, then

$$(I, f) = (I_n(A), f|_{I_n(A)}),$$

so every bimodule map is determined by its action on some I_n . Thus for each α , we may assume that $(I_\alpha, f_\alpha) = (I_n(A), f_\alpha)$ for some n . Since there are only countably many $I_n(A)$, there is some $d \in \mathbb{N}$ for which the uncountable set $\{(I_d(A), f_\alpha) \mid \alpha \in \mathcal{I}\}$ is k -linearly independent. Next, choose any nonzero element $a \in I_d(A)$ and note that by replacing $I_d(A)$ with AaA , we may assume that there is an uncountable k -linearly independent set of the form $\{(AaA, f_\alpha) \mid \alpha \in \mathcal{I}\}$. Being an (A, A) -bimodule map, each f_α is completely determined by $f_\alpha(a)$. However, since A is a countably generated k -algebra, $\dim_k(A)$ is countable, and it follows that the set

$$\{f_\alpha(a) \mid \alpha \in \mathcal{I}\}$$

is k -linearly dependent. This proves the theorem. \square

Remark 6.3 Though Proposition 6.2 requires A to be non-PI and just infinite, the conclusion still holds for any countably generated prime algebra with the property that there is a countable set of nonzero elements such that every nonzero ideal contains at least one of these elements; this is often referred to as having a *countable separating set* in the literature. Note that, for example, any primitive homomorphic image of an enveloping algebra of a finite dimensional Lie algebra has a countable separating set, as does any countably generated prime non-PI right Goldie ring of GKdim 2. We also remark that the preceding proof, with the obvious modifications, yields a generalization of [19, Theorem 1].

Corollary 6.4 *Let k be an uncountable algebraically closed field and let A be a countably generated non-PI just infinite k -algebra. Then A is stably just infinite.*

Proof. If A is simple then $C(A) = k$ and the result follows from Theorem 3.4. Otherwise A is not simple, and so $C(A)$ is a countable dimensional field extension of k by Proposition 6.2. Since k is uncountable, $C(A)$ must be algebraic over k . But k is algebraically closed, so A is centrally closed, and the result follows from Theorem 3.4. \square

Combining Proposition 6.1 and Corollary 6.4, we obtain Theorem 1.5.

7 An Example

In this short section we prove Theorem 1.4. We construct an interesting example showing that the right noetherian hypothesis in Theorem 1.3 and the uncountable hypothesis in Theorem 1.5 are necessary.

Let k be a countable algebraically closed field. There exists a simple Ore

domain R , affine over k , with $\text{trdeg}_k(Z(R)) = \infty$ [9]. Thus R is affine, simple, and infinite dimensional, and so R cannot satisfy a polynomial identity. The ring R is obtained as a skew laurent extension of an Ore domain with respect to an automorphism of infinite order [9]. In particular, R is not a division ring. Fix a nonzero element $x \in R$ which is not a unit, and set $A = k + xR$. A is affine [17, Proposition 2], and since xR is simple as a (non-unital) ring, A is just infinite. As R is not PI, neither is A , and so we see that $Z(A)$ is a finite dimensional field extension of k by Proposition 2.3. But note that $Z(A) \cap xR = (0)$ because xR is a proper right ideal of R . Hence $Z(A) = k$. Finally, it is easy to see that A is right Ore, and so the extended center of A coincides with the center of its right Goldie quotient ring. Since $Q(A) = R$, we see that $C(A) = Z(R)$. Thus A is not centrally closed, and so A is not stably just infinite by Theorem 3.4.

Acknowledgments

We thank Lance Small for many helpful conversations.

References

- [1] L. Bartholdi. Branch rings, thinned rings, tree enveloping rings. *available at: arxiv.org/pdf/math.RA/0410226*, 2004.
- [2] J. P. Bell. Noetherian algebras over algebraically closed fields. *J. Algebra*, 310(1):148 – 155, 2006.
- [3] G. M. Bergman. Zero-divisors in tensor products. In *Noncommutative ring theory (Internat. Conf., Kent State Univ., Kent, Ohio, 1975)*, pages 32–82. Lecture Notes in Math., Vol. 545. Springer, Berlin, 1976.
- [4] T. S. Erickson, W. S. Martindale, 3rd, and J. M. Osborn. Prime nonassociative algebras. *Pacific J. Math.*, 60(1):49–63, 1975.
- [5] J. Farina and C. Pendergrass. A few properties of just infinite rings. *accepted, Comm. Algebra*, 2006.
- [6] D. R. Farkas and L. W. Small. Algebras which are nearly finite dimensional and their identities. *Israel J. Math.*, 127:245–251, 2002.
- [7] R. Gilmer, B. Nashier, and W. Nichols. The prime spectra of subalgebras of affine algebras and their localizations. *J. Pure Appl. Algebra*, 57(1):47–65, 1989.
- [8] M. Hazewinkel, editor. *Handbook of algebra. Vol. 1*. North-Holland Publishing Co., Amsterdam, 1996.
- [9] R. S. Irving. Finitely generated simple Ore domains with big centres. *Bull. London Math. Soc.*, 12(3):197–201, 1980.

- [10] T. Y. Lam. *Lectures on modules and rings*, volume 189 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1999.
- [11] T. Y. Lam. *A first course in noncommutative rings*, volume 131 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 2001.
- [12] W. S. Martindale, III. Lie isomorphisms of prime rings. *Trans. Amer. Math. Soc.*, 142:437–455, 1969.
- [13] W. S. Martindale, III. Prime rings satisfying a generalized polynomial identity. *J. Algebra*, 12:576–584, 1969.
- [14] J. C. McConnell and J. C. Robson. *Noncommutative Noetherian rings*, volume 30 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, revised edition, 2001. With the cooperation of L. W. Small.
- [15] Z. Reichstein, D. Rogalski, and J. J. Zhang. Projectively simple rings. *Adv. Math.*, 203(2):365–407, 2006.
- [16] R. Resco. A reduction theorem for the primitivity of tensor products. *Math. Z.*, 170(1):65–76, 1980.
- [17] R. Resco. Affine domains of finite Gelfand-Kirillov dimension which are right, but not left, Noetherian. *Bull. London Math. Soc.*, 16(6):590–594, 1984.
- [18] L. H. Rowen. *Ring theory. Vol. I*, volume 127 of *Pure and Applied Mathematics*. Academic Press Inc., Boston, MA, 1988.
- [19] L. H. Rowen and L. Small. Primitive ideals of algebras. *Comm. Algebra*, 25(12):3853–3857, 1997.
- [20] P. Vámos. On the minimal prime ideal of a tensor product of two fields. *Math. Proc. Cambridge Philos. Soc.*, 84(1):25–35, 1978.