

On J. Borwein's Concept of Sequentially Reflexive Banach Spaces

by P. Ørno

A Banach space X is reflexive if the Mackey topology $\tau(X^*, X)$ on X^* agrees with the norm topology on X^* . Borwein [B] calls a Banach space X *sequentially reflexive* provided that every $\tau(X^*, X)$ convergent *sequence* in X^* is norm convergent. The main result in [B] is that X is sequentially reflexive if every separable subspace of X has separable dual, and Borwein asks for a characterization of sequentially reflexive spaces. Here we answer that question by proving

Theorem. *A Banach space X is sequentially reflexive if and only if ℓ_1 is not isomorphic to a subspace of X .*

Proof: Assume first that ℓ_1 is not isomorphic to a subspace of X and let $\{x_n^*\}_{n=1}^\infty$ be a weak*-null sequence in X^* for which the sequence $\{\langle x_n^*, x_n \rangle\}_{n=1}^\infty$ converges to zero for every weakly null sequence $\{x_n\}_{n=1}^\infty$ in X ; by the easy Lemma 2.1 in [B] it is enough to check that such a sequence $\{x_n^*\}_{n=1}^\infty$ must converge in norm to zero. If not, by passing to a subsequence we can select a sequence $\{x_n\}_{n=1}^\infty$ in the unit ball of X with $\{\langle x_n^*, x_n \rangle\}_{n=1}^\infty$ bounded away from zero. By passing to a further subsequence, we can assume by Rosenthal's theorem [R], [D, chpt. XI] on Banach spaces which do not contain isomorphs of ℓ_1 that $\{x_n\}_{n=1}^\infty$ is weakly Cauchy. Since $\{x_n^*\}_{n=1}^\infty$ converges weak* to zero, by passing to further subsequences and replacing $\{x_n\}_{n=1}^\infty$ with a subsequence of differences $\frac{x_{2n} - x_{2n-1}}{2}$, we can assume moreover that $\{x_n\}_{n=1}^\infty$ is weakly null. This contradiction completes the proof of the first direction.

To go the other way, suppose that Y is a subspace of X which is isomorphic to ℓ_1 and let $\{e_n\}_{n=1}^\infty$ be the image of the unit vector basis under some isomorphism from ℓ_1 onto Y . Define a bounded linear operator from Y into $L_\infty[0, 1]$ by mapping e_n to the n -th Rademacher function r_n . By the injective property of $L_\infty[0, 1]$, this operator extends to a bounded linear operator T from X into $L_\infty[0, 1]$. Let r_n^* be the n -th Rademacher function in $L_1[0, 1]$ considered as a subspace of $L_\infty[0, 1]^*$. Thus the sequence $\{r_n^*\}_{n=1}^\infty$, being equivalent to an orthonormal sequence in a Hilbert space, converges weakly to zero. Since $L_\infty[0, 1]$ has the Dunford-Pettis property (cf. [D, p. 113]), $\{r_n^*\}_{n=1}^\infty$ converges in the Mackey topology to zero, *a fortiori* $\{T^*r_n^*\}_{n=1}^\infty$ converges $\tau(X^*, X)$ to zero. But $\langle T^*r_n^*, e_n \rangle = \langle r_n^*, r_n \rangle = 1$, so $\{T^*r_n^*\}_{n=1}^\infty$ does not converge to zero in norm.

References

- [B] J. Borwein, *Asplund spaces are "sequentially reflexive"*, (preprint).
- [D] J. Diestel, *Sequences and series in Banach spaces*, Springer-Verlag Graduate Texts in Mathematics 92, (1984).
- [R] H. P. Rosenthal, *A characterization of Banach spaces containing ℓ_1* , **Proc. Nat. Acad. Sci.** **71**, (1974), 2411–2413.