

**FACTOR = QUOTIENT,  
UNCOUNTABLE BOOLEAN ALGEBRAS,  
NUMBER OF ENDOMORPHISM AND WIDTH**

SAHARON SHELAH

June 22, 1991

ABSTRACT. We prove that assuming suitable cardinal arithmetic, if  $B$  is a Boolean algebra every homomorphic image of which is isomorphic to a factor, then  $B$  has locally small density. We also prove that for an (infinite) Boolean algebra  $B$ , the number of subalgebras is not smaller than the number of endomorphisms, and other related inequalities. Lastly we deal with the obtainment of the supremum of the cardinalities of sets of pairwise incomparable elements of a Boolean algebra.

We show in the first section:

**0.1 Conclusion.** *It is consistent, that for every  $Q = F$  Boolean algebra  $B^*$ , for some  $n < \omega$ ,  $\{x : B^*|x \text{ has a density } \leq \aleph_n\}$  is dense (so  $B^*$  has no independent subset of power  $\aleph_n$ ).*

Where:

**0.2 Definition.** *A B.A. is  $Q = F$  (quotient equal factor) if: every homomorphic image of  $B$  is isomorphic to some factor of  $B$  i.e  $B|a$  for some  $a \in B$ .*

The “consistent” is really a derivation of the conclusion from a mild hypothesis on cardinal arithmetic (1.2). The background of this paper is a problem of Bonnet whether every  $Q = F$  Boolean algebra is superatomic.

Noting that : “ $B|x$  has density  $\leq \aleph_n$ ” is a weakening of “ $x$  is an atom of  $B'$ ”, we see that **0.1** is relevant.

The existence of non trivial example is proved in R. Bonnet, S. Shelah [2].

M. Bekkali, R. Bonnet and M. Rubin [1] characterized all interval Boolean algebras with this property.

In the second section we give a more abstract version. In a paper in preparation, Bekkali and the author use theorem **2.1** to show that every  $Q = F$  tree Boolean algebras are superatomic.

In the third section we deal with the number of endomorphism (e.g.  $aut(B)^{\aleph_0} \leq end(B)$ ) and in the fourth with the width of a Boolean algebra.

We thank D. Monk for detecting an inaccuracy in a previous version.

**Notation**

$B$  denote a Boolean algebra.

---

Partially supported by Basic Research Fund of the Israeli Academy of Sciences. Publ. No. 397

## FACTOR = QUOTIENT, UNCOUNTABLE BOOLEAN ALGEBRAS

$B^+$  is the set of non zero members of  $B$ .

$B|x$  where  $x \in B^+$  is  $B|\{y : y \leq x\}$ .

$\text{comp} B$  is the completion of  $B$ , so it is an extension of  $B$ .

Remember : if  $B_1$  is a subalgebra of  $B_2$ ,  $B$  is complete,  $h$  a homomorphism from  $B_1$  to  $B$ , then  $h$  can be extended to a homomorphism from  $B_2$  to  $B$ .

$\text{id}_A$  is the identity function on  $A$ .

$\gamma\alpha = \{\eta : \eta \text{ is a sequence of lengths } \gamma \text{ of ordinals } < \alpha\}$ .

$\gamma > \alpha = \bigcup_{\beta < \alpha} \beta\alpha$ .

§1 Maybe every “quotient equal factor”  $BA$  has locally small density .

**1.1 Hypothesis.**

- (1)  $B^*$  is a  $Q = F$  Boolean algebra.
- (2) for each  $n < \omega$ ,  $B^*$  has a factor  $B_n$  s.t.:  $0 < x \in B_n \Rightarrow \text{density}(B_n|x) \geq \aleph_n$ .

**1.2 Hypothesis.** For  $\alpha \geq \omega$  we have  $2^{|\alpha|} > \aleph_\alpha$ .

1.3 Desired Conclusion. Contradiction.

We shall use **1.1** all the times, but **1.2** only in **1.17**.

**1.4 Definition.**  $K_\lambda^*$  is the class of Boolean algebra such that:

$$(\forall x \in B^+)[\text{density}(B|x) = \lambda].$$

**1.5 Claim.** If  $B$  is atomless,  $x \in B^+$  then for some  $y$ ,  $0 < y \leq x$ , and infinite cardinal  $\lambda$  we have  $B|y \in K_{\text{density}(B|x)}^*$ .

**1.6 Claim.** If  $B \in K_\lambda^*$ ,  $B \subseteq B' \subseteq \text{comp}(B)$  then,

$$B' \in K_\lambda^*.$$

**1.7 Claim.** If  $B \in K_\lambda^*$ ,  $\aleph_0 \leq \mu < \lambda$ ,  $\mu$  regular then some subalgebra  $B'$  of  $B$  is in  $K_\mu^*$ . If in addition  $B$  is a  $Q = F$  algebra, then some homomorphic image  $B''$  of  $B$  is in  $K_\mu^*$ .

*Proof.* Choose by induction on  $i < \mu$ ,  $B_i \subseteq B$ ,  $|B_i| \leq \mu$ ,  $[i < j \Rightarrow B_i \subseteq B_j]$  such that:

if  $x \in B_i^+$  then there is  $y(x, i) \in B_{(i+1)}$  satisfying:

- (1)  $0 < y(x, i) \leq x$ ,
- (2) for no  $z \in B_i$ ,  $0 < z \leq y(x, i)$ .

(possible as for each  $i$  and  $x \in B_i^+$   $\text{density}(B_i|x) = \lambda$ ). Let now  $B' = \bigcup_{i < \mu} B_i$ ; it is a subalgebra of  $B$ . Now  $x \in (B')^+ \Rightarrow \text{density}(B'|x) = \mu$  as on the one hand  $|B'| \leq \sum_{i < \mu} |B_i| \leq \mu \times \mu = \mu$  implies  $\text{density}(B'|x) \leq \mu$  for every  $x \in B'$  and on the other hand if  $x \in (B')^+$ ,  $A \subseteq B'|x$ ,  $|A| < \mu$  then for some  $i < \mu$ ,  $A \subseteq B_i$ , hence  $y(x, i) \in (B')^+$  witness  $A$  is not dense in  $B'|x$ . Now, if  $B$  is a  $Q = F$  Boolean Algebra, then  $\text{id}_{B'}$  can be extended to a homomorphism  $h'$  from  $B$  into  $\text{comp}(B')$ , so  $h'(B)$  is as required.

FACTOR = QUOTIENT, UNCOUNTABLE BOOLEAN ALGEBRAS

**1.8 Conclusion.**  $\{\lambda : \lambda \text{ regular and } B^* \text{ has a factor in } K_\lambda^*\}$  is an initial segment of  $\{\aleph_\alpha : \aleph_\alpha \text{ regular}\}$ .

**1.9 Definition.**

- (1)  $\alpha(*)$  is minimal such that for no  $\lambda > \aleph_{\alpha(*)}$  does  $B^*$  has a factor in  $K_\lambda^*$ . Let  $\kappa^* = \kappa(*) =: |\alpha(*)|$ .
- (2) Let for  $\alpha < \alpha^*$ ,  $b_\alpha \in B^*$  be such that  $B^*|b_\alpha \in K_{\aleph_{\alpha+1}}^*$ .
- (3) Let  $J_\alpha$  be the ideal  $\{b \in B^* : B^*|b \in K_{\aleph_{\alpha+1}}^*\}$ .

**1.10 Definition.**  $B^{[\omega > \lambda]}$  is the Boolean algebra generated freely by  $\{x_\eta : \eta \in \omega > \lambda\}$  except  $x_\eta \leq x_{\eta|m}$  ( $m \leq \lg(\eta), \eta \in \omega > \lambda$ ).

**1.11 Claim.**

- (1) For  $\alpha < \beta < \alpha^*$ ,  $J_\alpha \cap J_\beta = \{0\}$  and  $J_\alpha$  is an ideal.
- (2) For no  $B' \subseteq B^*$  and proper ideal  $I$  of  $B'$ ,  $[x \in B' \setminus I \Rightarrow \text{density}(B'|x/I) > \aleph_{\alpha(*)}]$ .

*Proof.*

(1) Trivial.

(2) By **1.5** for some  $\beta > \alpha(*)$  and proper ideal  $J$  of  $B'$  (of the form  $\{x \in B' : x - b \in I\}$ ) we have  $B'/I \in K_{\aleph_\beta}^*$  so there is a homomorphism  $h$  from  $B^*$  into  $\text{comp}(B'/I)$  extending  $x \mapsto x/I$  ( $x \in B'$ ). So  $B^*$  has a factor isomorphic to  $\text{Rang } h$ , but this Boolean algebra is in  $K_{\aleph_\beta}^*$ ,  $\aleph_\beta > \aleph_{\alpha(*)}$ . So by **1.7** we get contradiction to the choice of  $\alpha(*)$ .

**1.12 Definition.** 1)  $I^* =: \{x \in B^* : \bigcup_{(\alpha < \alpha(*)} J_\alpha \text{ is dense below } x\}$ .

2) For  $A \subseteq \alpha(*) : I_A^* =: \{x \in B^* : \bigcup \{J_\alpha : \alpha \in A\} \text{ is dense below } x\}$ .

**1.13 Claim.**

- (1)  $I^*, I_A^*$  are ideals of  $B$ .
- (2)  $A \subseteq B \Rightarrow I_A^* \subseteq I_B^* \subseteq I^*$ .

**1.14 Claim.** For every  $A \subseteq \alpha(*)$ , there are  $c_A, h_A$  such that:

- (1)  $c_A \in B^*$  and  $I_A^*$  is dense below  $c_A$ ,
- (2)  $h_A$  is a homomorphism from  $B^*$  onto  $B^*|c_A$ ,
- (3)  $h_A|I_A^*$  is one to one,
- (4) If  $B|x \cap I_A^* = \{0\}$  then  $h_A(x) = 0$ ,
- (5)  $B^*|c_A$  is a subalgebra of the completion of the subalgebra  $\{h_A(x) : x \in I_A^*\}$ .

*Proof.* Let  $ba(A)$  be

$$I_A^* \cup \{1 - x : x \in I_A^*\},$$

this is a subalgebra of  $B^*$ .

Let  $h_1$  be a homomorphism from  $B^*$  to  $\text{comp}(ba(A))$  extending  $id_{ba(A)}$  and  $h_1(x) = 0$  if  $B^*|x \cap I_A^* = \{0\}$ .

Now  $h_1(B^*)$  is a quotient of  $B^*$  hence there is an isomorphism  $h_2$  from  $h_1(B^*)$  onto some  $B^*|c_A$ .

FACTOR = QUOTIENT, UNCOUNTABLE BOOLEAN ALGEBRAS

Let

$$h_A = h_2 \circ h_1$$

so (1), (2), (3), (4), (5) holds.

**1.15 Claim.** Let  $A \subseteq \alpha(*)$

- (1) If  $x \in \bigcup_{\alpha \in A} J_\alpha$  then  $h_A(x) \in \bigcup_{\alpha \in A} J_\alpha$ .
- (2) If  $x \in I_A^* \setminus \bigcup_{\alpha \in A} J_\alpha$  then  $h_A(x) \notin \bigcup_{\alpha \in A} J_\alpha$ .
- (3)  $\bigcup_{\alpha \in A} h_A(J_\alpha)$  is dense and downward closed in  $B^*|c_A$ .
- (4)  $c_A \in I_A^*$ .
- (5) If  $x \in J_\alpha$  and  $\alpha \in A$  then  $h_A(x) \in J_\alpha$ .

*Proof.*

- (1) By (5) of **1.14**.
- (2) is easy too.
- (3) is easy too.
- (4) is easy too.
- (5) is easy too.

**1.16 Claim.** We can find  $x_\eta \in B^*$  for  $\eta \in \omega^{>\lambda}$  where  $\lambda = 2^{\kappa(*)}$  such that:

- (1)  $m < lg(\eta) \Rightarrow 0 < x_\eta \leq x_{(\eta|m)}$ .
- (2) If  $\bigwedge_{e=1}^k \nu_e \not\leq \eta$ ,  $\eta \in \omega^{>\lambda}$ ,  $\bigwedge_e \nu_e \in \omega^\lambda$  then

$$B^* \vDash x_\eta \not\leq \bigcup_{e=1}^k x_{\nu_e}.$$

*Proof.* Now we can choose  $\langle A_\eta^0 : \eta \in \omega^{>\lambda} \rangle$  which is a family of subsets of  $\kappa(*)$  such that any non trivial Boolean combination of them has cardinality  $\kappa(*)$ .

Let for  $\eta \in \omega^{>\lambda}$   $A_\eta = \text{def} \bigcap_{e \leq lg(\eta)} A_{\eta|e}$ . Let

$$(a) \quad x_{\langle \rangle} = c_{A_{\langle \rangle}} = h_{A_{\langle \rangle}}(1_{B^*}).$$

$$(b) \quad x_{\langle i \rangle} = h_{A_{\langle \rangle}}(c_{A_{\langle i \rangle}}) = h_{A_{\langle \rangle}} h_{A_{\langle i \rangle}}(1_{B^*})$$

and generally,

$$x_{\langle i_0, i_1, \dots, i_{n-1} \rangle} = h_{A_{\langle \rangle}} h_{A_{\langle i_0 \rangle}} h_{A_{\langle i_0, i_1 \rangle}} \dots h_{A_{\langle i_0, i_1, \dots, i_{n-1} \rangle}}(1_{B^*}).$$

We prove (a) by induction of  $lg \eta$ .

The reader may check

- 1.17 Final Contradiction.**  $\{x_\eta : \eta \in \omega^{>\lambda}\}$  from **Claim 1.16** contradict by **1.11(2)** and the choice of  $\aleph_{\alpha(*)}$ , because  $\lambda = 2^{\kappa(*)} = 2^{|\alpha(*)|} > \aleph_{\alpha(*)}$ .  
 [of course  $\aleph_{\alpha(*)} \leq |B^*|$ ]  
 Actually, we have prove more.

**1.18 Remark.** (1) So we have in **1.17** prove that if set theory is as in **Hypothesis 1.2**, then there is no Boolean algebra as in **1.1**, hence proving **Conclusion 0.1**

(2) Note: if **1.2**, any  $Q = F$  Boolean algebra has no factor  $\mathcal{P}(\omega)$ .

§2  $Q = F$  Boolean algebras: a general theorem .

**2.1 Theorem.** Suppose:

- (1)  $B^*$  is a  $(Q = F)$  Boolean algebra.
- (2)  $N$  is a family of (non zero) members of  $B^*$  (the “nice” elements).
- (3)  $\kappa$  a cardinal ( $\geq \aleph_0$ ),  $\langle K_\alpha : \alpha < \kappa \rangle$  a sequence.
- (4)  $K_\alpha$  is family of Boolean algebras closed under isomorphism and for  $\alpha \neq \beta$  we have  $K_\alpha \cap K_\beta = \emptyset$ .
- (5) for every  $\alpha$  some factor of  $B^*$  is in  $K_\alpha$ .
- (6) if  $x \in (B^*)^+$ ,  $(B^*|x) \in K_\alpha$  then for some  $y \leq x$ ,  $B^*|y \in K_\alpha$ ,  $y \in N$ .
- (7) if  $x_1, x_2 \in N$ ,  $B^*|x_1 \in K_{\alpha_1}$ ,  $B^*|x_2 \in K_{\alpha_2}$ ,  $\alpha_1 \neq \alpha_2$  then  $x_1 \cap x_2 = 0$ .
- (8) if  $x \in B^*$ ,  $N' \subseteq N$  then
  - ( $\alpha$ ) for some  $y_1, \dots, y_n \in N'$ , for every  $z \in N'$  we have  $B^* \models x \cap z \subseteq y_1 \cup \dots \cup y_n$
  - or
  - ( $\beta$ ) for some  $y \in N' : y \leq x$ .
 Actually we use (8) only for  $N'$  of the form  $\{y \in N : (\exists \alpha)[\alpha \in A \ \& \ B^*|y \in K_\alpha]\}$ .
- (9) if  $x < y \in B^*$ ,  $B^*|x \in K_\alpha$  and  $B^*|y \in K_\beta$ , then  $\alpha = \beta$ .

Then the Boolean algebra  $B[\omega^{>}(2^\kappa)]$  can be embedded into  $B^*$ , remember  $\omega^{>}(2^\lambda)$  is

the tree  $\{\eta : \eta \text{ a finite sequence of ordinals } < 2^\kappa\}$  and Def 1.10.

*Proof of theorem 2.1.*

Let for  $\alpha < \kappa$ ,  $Y_\alpha =: \{y \in N : B^*|y \in K_\alpha\}$ .

For  $A \subseteq \kappa$  let us define

$I_A =$  the ideal generated by  $Y_A = \bigcup_{\alpha \in A} Y_\alpha = \{y \in N : B^*|y \in K_\alpha \text{ for some } \alpha \in A\}$  and

$J_A =: \{z \in B_\alpha : \text{for every } y \in Y_A \text{ we have } z \cap y = 0\}$ .

Clearly  $J_A$  is an ideal.

Now for each  $A \subseteq \kappa$   $B^*/J_A$  is a quotient of  $B^*$ . Hence by condition (1) there are  $y_A^* \in B^*$  and an isomorphism  $h_A : B^*/J_A \rightarrow B^*|y_A^*$  onto.

Let  $g_A : B^* \rightarrow B^*/J_A$  be canonical, so  $h_A \circ g_A(1_{B^*}) = y_A^*$ . Let  $f_A = h_A \circ g_A$ .

Define for  $y \in B^*$  the following:  $\text{cont}(y) =: \{\alpha : (\exists y' \leq y)[B^*|y' \in K_\alpha]\}$ .

(i.e. the content of  $y$ ). We next prove

(\*)<sub>1</sub>  $\text{cont}(y_A^*) \supseteq A$ .

*Proof.* By condition (5) for each  $\alpha \in A$ , there is  $x_\alpha \in B^*$ , such that:  $B^*|x_\alpha \in K_\alpha$ . By condition (6) wlog  $x_\alpha \in N$ , hence  $x_\alpha \in I_A$ , hence  $g_A(B^*|x_\alpha)$ , is one to one, hence  $B^*|x_\alpha \cong B^*|f_A(x_\alpha)$ , hence by condition (4)  $B^*|f_A(x_\alpha) \in K_\alpha$ ; now  $\text{Rang} f_A = B^*|y_A^*$ , so  $\alpha \in \text{cont}(y_A^*)$ . So we have prove (\*)<sub>1</sub>.

(\*)<sub>2</sub>  $\text{cont}(y_A^*) \subseteq A$

*Proof.* Suppose  $\alpha \in \text{cont}(y_A^*)$ , so there is  $z \leq y_A^*$ , such that  $B^*|z \in K_\alpha$ . As  $f_A$  is a homomorphism from  $B^*$  onto  $B^*|y_A^*$ , there is  $x \in B^*$  such that  $f_A(x) = z$ .

FACTOR = QUOTIENT, UNCOUNTABLE BOOLEAN ALGEBRAS

Now the kernel of  $f_A$  is  $J_A$ , and  $B^*|0 \notin K_\alpha$  so  $x \notin J_A$ ; and clearly  $(B^*|x)/J_A$  is in  $K_\alpha$ .

Hence by  $(*)_3$  below  $\alpha \in A$ .

$(*)_3$  if  $x \in B^* \setminus J_A$  and  $(B^*/J_A)|f_A(x) \in K_\alpha$  then  $\alpha \in A$

*Proof.* We apply condition (8) to  $x$  and  $N' = Y_A$ . So one of the following two cases occurs:

**Case  $\alpha$ :** There are  $n < \omega$ ,  $y_1, \dots, y_n \in N'$  such that:

$(\forall z \in N') x \cap z \subseteq y_1 \cup \dots \cup y_n$ .

So  $x - (y_1 \cup \dots \cup y_n) \in J_A$  (by definition of  $J_A$ ).

Let  $x_1 = x \cap (y_1 \cup \dots \cup y_n)$  hence  $B^*|f_A(x) = (B^*|x)/J_A \cong (B^*|x_1)/J_A \cong B^*|x_1$ , (last isomorphism as  $\wedge_e y_e \in Y_A$  hence  $y_1 \cup \dots \cup y_n \in I_A$  hence  $f_A|(B^*|x_1)$  is one to one). So  $B^*|x_1 \in K_\alpha$ , hence by condition (6) for some  $x_2 \leq x_1$ , we have  $x_2 \in N \& B^*|x_2 \in K_\alpha$ .

Let  $B^*|y_e \in K_{\alpha_e}$  where  $\alpha_e \in A$  (by definition of  $Y_A$ ). Clearly  $x_2 \leq x_1 \leq y_1 \cup \dots \cup y_n$ , so for some  $e$ ,  $y_e \cap x_2 \neq 0$  hence  $\alpha = \alpha_e \in A$  (by condition (7)) so we get the trivial desired conclusion.

**Case  $\beta$ :** There is  $y \in N'$ ,  $y \leq x$ .

As  $y \in N' = Y_A = \cup_{\beta \in A} Y_\beta$  for some  $\beta \in A$  we have  $y \in K_\beta$ , also  $y \in N' \subseteq N$  so  $J_A \cap (B^*|y) = \{0\}$  so  $(B^*/J_A)|g_A(y) \in K_\beta$ . Remembering  $(B^*/J_A)|g_A(x) \in K_\alpha$ , as  $B^*/J_A \cong B^*|y_A^*$  we get by condition (9) that  $\alpha = \beta$ . So  $(*)_3$  hence  $(*)_2$  is proved.

Next we prove

$(*)_4$  if  $B \subseteq A \subseteq \kappa$ , and  $\text{cont}(y) = B$ , then  $\text{cont}[f_A(y)] = B$ .

*Proof.*

**inclusion  $\supseteq$**

First let  $\alpha \in B$ , then for some  $x \leq y$ ,  $B^*|x \in K_\alpha$ , and by condition (6)  $wlog$   $x \in N$ , hence  $x \in I_A$  (as  $\alpha \in B \subseteq A$ ) hence  $f_A|(B^*|x)$  is one to one and onto  $B^*|f_A(x)$  so  $f_A(x) \leq f_A(y)$ ,  $B^*|f_A(x) \in K_\alpha$ , so  $\alpha \in \text{cont}(f_A(y))$ .

**inclusion  $\subseteq$**

Second let us assume  $\alpha \in \text{cont}[f_A(y)]$ . So (as  $f_A$  is onto  $B^*|y_A^*$ , and if  $f_A(x) \leq f_A(y)$  then  $f_A(x \cap y) = f_A(y)$ ,  $x \cap y \leq y$ ) there is  $x \leq y$  such that  $B^*|f_A(x) \in K_\alpha$ . Now apply condition (8) to  $x$  and  $Y_A$ . So case  $(\alpha)$  or case  $(\beta)$  below holds.

**Case  $\alpha$ :** There are  $n < \omega$ ,  $y_1, \dots, y_n \in Y_A$  such that for every  $z \in Y_A$  we have  $x \cap z \subseteq y_1 \cup \dots \cup y_n$ .

Hence  $x - (y_1 \cup \dots \cup y_n) \in J_A$  let  $x_1 = x \cap (y_1 \cup \dots \cup y_n)$  so  $f_A(x) = f_A(x_1)$  so  $B^*|f_A(x_1) \in K_\alpha$  and of course  $x_1 \leq y_1 \cup \dots \cup y_n$ , (and  $x_1 \leq y$ ) so  $f_A|(B^*|x_1)$  is one to one.

Now  $f_A$  is one to one on  $B^*|x_1$  hence  $B^*|x_1 \cong B^*|f_A(x_1) \in K_\alpha$ . Now  $x_1 \leq x \leq y$ , so  $x_1$  witness  $\alpha \in \text{cont}(y)$ , which is  $B$ .

**Case  $\beta$ :** There is  $t \leq x$ ,  $t \in Y_A$ .

Now  $t \leq x \leq y$ ,  $B^*|t \in K_\beta$  for some  $\beta \in A$  as  $t \in Y_A$ . Now  $f_A$  is one to one on  $B^*|f_A(t) \in K_\beta$ .

Also  $f_A(t) \leq f_A(x)$  hence by assumption (9) we have  $\beta = \alpha$ . Also  $t \leq x \leq y$  so  $t$  witness  $\beta \in \text{cont}(y)$ , so  $\alpha = \beta \in \text{cont}(y) = B$  as required.

So we have proved  $(*)_4$

## FACTOR = QUOTIENT, UNCOUNTABLE BOOLEAN ALGEBRAS

*end of proof of theorem 2.1:* Let  $\lambda = 2^\kappa$ .

Let  $\langle \mathcal{U}_\eta : \eta \in {}^\omega \lambda \rangle$  be a family of subsets of  $\kappa$ , any finite Boolean combination of them has power  $\kappa$  (or just  $\neq \emptyset$ ).

Let  $\mathcal{U}_\eta^* = \bigcap_{e \leq l g \eta} \mathcal{U}_{\eta|e}$ . Now define for every  $\eta \in {}^\omega \kappa$  and  $e \leq l g(\eta)$  an element  $y_\eta^e$  of  $B^*$ :

$$y_\eta^e =^{def} f_{\mathcal{U}_{\eta|e}^*} f_{\mathcal{U}_{\eta|(e+1)}^*} \cdots f_{\mathcal{U}_{\eta|(n-1)}^*} f_{\mathcal{U}_\eta^*}(1_{B^*}) \text{ and } y_\eta^\infty =^{def} y_\eta^\circ$$

Now: (a) prove for each  $\eta \in {}^n \kappa$  by downward induction on  $e \in \{0, 1, \dots, n\}$  that  $cont(y_\eta^e) = \mathcal{U}_\eta^*$ ; for  $e = n$  this is  $(*)_1 + (*)_2$  as  $y_\eta^n = y_{\mathcal{U}_\eta^*}^*$ ;

for  $e < n$  (assuming for  $e + 1$ ) this is by  $(*)_4$ .

Next note: (b) if  $\nu = \eta \hat{ } \alpha >$  then  $y_\nu^\infty \leq y_\eta^\infty$

[prove by downward induction for  $e \in \{0, 1, \dots, l g \eta\}$  we have:  $y_\nu^{e+1} \leq y_\eta^e$ ; remember  $f_{\mathcal{U}}$  is order preserving].

Lastly note (c) if  $\eta \in {}^\omega \lambda$ ,  $n < \omega$ , and  $\nu_e \in {}^\omega \lambda$  is not initial segment of  $\eta$  for  $e = 1, \dots, n$  then  $y_\eta^\circ - \bigcup_{e=1}^n y_{\nu_e}^\circ \neq \emptyset$ ; this follows by (a) and the definition of  $cont(y_\eta^\circ)$ .

Now by (a), (b), (c) there is an embedding  $g$  from the subalgebra of  $B^*$  which  $\{y_\eta^\circ : \eta \in {}^\omega \lambda\}$  generates mapping  $y_\eta^\circ$  to  $x_\eta$ .

### §3 The Number of Subalgebras .

**3.1 Definition.** For a BA  $A$

- (1)  $Sub(A)$  is the set of subalgebras of  $A$ .
- (2)  $Id(A)$  is the set of ideals of  $A$ .
- (3)  $End(A)$  is the set of endomorphisms of  $A$ .
- (4)  $Pend(A)$  is the set of partial endomorphisms ( i.e. homomorphisms from a subalgebra of  $A$  into  $A$ ).
- (5)  $Psub(A)$  is the family of subsets of  $A$  closed under union ,intersection and substruction but 1 may be not in it though 0 is [so not necessarily closed under complementation].
- (6) We let  $sub(A)$ ,  $id(A)$ ,  $end(A)$ ,  $aut(A)$ ,  $pend(A)$ ,  $psub(A)$  be the cardinality of  $Sub(A)$ ,  $Id(A)$ ,  $End(A)$ ,  $Aut(A)$ ,  $Pend(A)$  and  $Psub(A)$  respectively

In D. Monk [4] list of open problems appear:

**PROBLEM 63.** Is there a BA  $A$  such that  $aut(A) > sub(A)$  ?

See [4] page 125 for background.

**3.2 Theorem.** For a BAA we have:  $aut(A)$  is not bigger than  $sub(A)$ .

**3.3 Conclusion.**

- (1) For a BAA we have  $end(A)$  is not bigger than  $sub(A)$ .
- (2) For a BAA we have  $pend(A)$  is exactly  $sub(A)$ .
- (3) For a BAA and  $a$  in  $A$ ,  $0 < a < 1$  we have  $sub(A) = Max\{sub(A|a), sub(A|-a)\}$ .

We shall prove it in 3.5

**Remark.** Of course -  $A$  is infinite- we many times forget to say so.

*3.4 Proof of the theorem.* Let  $\mu$  be  $sub(A)$ .

**3.4A Observation:**  $Psub(A)$  has cardinality  $Sub(A)$  [why? for the less trivial inequality,  $\leq$ , for every  $X$  in  $Psub(A)$  which is not a subset of  $\{0, 1\}$  choose a member  $a \neq 0, 1$  in  $X$  and let  $Y_a[X]$  be the subalgebra generated by  $\{x : x \in$

FACTOR = QUOTIENT, UNCOUNTABLE BOOLEAN ALGEBRAS

$X, x \leq a$  let  $Z_a[X]$  be the subalgebra generated by  $\{x : x \in X, x \cap a = 0\}$ ; now  $X$  can be reconstructed from  $\langle a, Y, Z \rangle$  as  $\{x \in A : x \cup a \in Y \text{ and } x - a \in Z\}$ . So  $|Psub(A)| \leq |A| \times |Sub(A)|^2 + 4 = |Sub(A)| + \aleph_0 = |Sub(A)|$ . (remember that  $|A| \geq \aleph_0$ , and that  $|A| \leq |Sub(A)|$ , as the number of non-atoms of  $A$  is  $\leq |\{Z_a[A] : a \in A\}|$ ).

For any automorphism  $f$  of  $A$  we shall choose a finite sequence of members of  $Psub(A)$  ( in particular of ideals of  $A$  ), and this mapping is one to one, thus we shall finish.

Let  $J =_{def} \{x : x \in A, \text{ and for every } y \in A \text{ below } x, \text{ we have } f(y) = y\}$ ; let  $I^*$  be the ideal of  $A$  generated by  $J$ , the set of elements  $x$  for which  $f(x) \cap x = 0$ .

**Observe** :  $x \in I$  implies  $f(x) \in I$  [why? as  $y = f(x), x \in I$  implies  $f(x) \cap x = 0$  hence  $f(y) \cap y = f(f(x)) \cap f(x) = f(f(x) \cap x) = f(0) = 0$ ].

**Observe**:  $J \cup I$  is a dense subset of  $A$  [if  $x \in B^+$  and there is no  $y \in J, 0 < y \leq x$  then wlog  $x \neq f(x)$ . If  $x \not\leq f(x)$  then  $z =: x - f(x)$  satisfies  $0 < z \leq x, f(z) \cap z \leq f(x) \cap z = f(x) \cap (x - f(x)) = 0$ ; so  $z \in I$ . If  $x \leq f(x)$  then  $x < f(x)$ ; as  $f$  is an automorphism of  $B^*$ , for some  $z$  in  $B^*$  we have  $f(z) = x$ , so  $x < f(x)$  means  $f(z) < f(f(z))$  hence  $z < f(z)$ , and  $z' = x - z = f(z) - z$  is in  $I$ , is  $> 0$  but  $< x$ , as required.]

Next let  $\{x_i : i < \alpha\}$  be a maximal sequence of distinct members of  $I$  satisfying : for any  $i, j < \alpha$  we have  $x_i \cap f(x_j) = 0$ , let  $I_1$  be the ideal generated by  $\{x_i : i < \alpha\}$  and let  $I_2$  be the ideal generated by  $\{f(x_i) : i < \alpha\}$ .

Clearly  $I_1 \cap I_2 = \{0\}$  let  $I_0 = \{y : y \in I^*, \text{ and for every } x \in I_1, y \cap x = 0 \text{ and } f(y) \in I_1\}$  and let  $I_3$  be the ideal of  $A$  generated by  $\{f(x) : x \in I_2\}$

**Observe** : each  $I_t$  ( $t = 0, 1, 2, 3$ ) is an ideal of  $A$ , contained in  $I^*$  [why? for  $t = 1$  by choice each  $x_i$  is in  $I$ ,  $t = 0$  by it's definition,  $I_2, I_3$  by their definition and an observation above, i.e.,  $x \in I \Rightarrow f(x) \in I$ ]

**Observe**: for  $t = 0, 1, 2$  we have:  $I_t \cap I_{t+1} = \{0\}$  [why ? for  $t = 1$ , by the choice of  $\{x_i : i < \alpha\}$ , for  $t = 0$  by definition of  $I_0$ , lastly for  $t = 2$  applies its definition +  $f$  being an automorphism.]

**Observe**:  $I_0 \cup I_1 \cup I_2$  is a dense subset of  $I^*$  [why? assume  $x$  in  $I^*$  but below it there is no non zero member of this union, so we can replace it by any non zero element below it; as  $x \in I^*$ , there is below it a non zero element  $y$  with  $y \cap f(y) = 0$  so wlog  $x \cap f(x) = 0$ ; why have we not choose  $x_\alpha = x$ ? there are two case:

**Case 1**: For some  $j < \alpha$ ,  $x \cap f(x_j)$  is not zero  
so there is a non zero element below  $x$  in  $I_2$ .

**Case 2**: for some  $j < \alpha$ ,  $f(x) \cap x_j$  is not zero  
so there is a non zero  $y$  in  $I_1$  below  $f(x)$  hence (as  $f$  is an automorphism) there is  $x'$  below  $x$  such that  $f(x') = y$  so wlog  $f(x)$  is in  $I_1$ , but then by its definition, either below  $x$  there is a member of  $I_1$  or  $x$  is in  $I_0$  so we have finished proving the observation.]

**Observe**: for  $t = 0, 1, 2$ ,  $x \in I_t \Rightarrow f(x) \in I_{t+1}$  [check].

Now we define  $C_t = C_t^f$ , a member of  $Psub(A)$  for  $t = 0, 1, 2$  as follows:  $C_t$  is the set  $\{x \cup f(x) : x \in I_t\}$ . The closure under the relevant operations follows as  $I_t$  is closed under them and  $f$  is an isomorphism and for  $x, y$  in  $I_t$ ,  $x \cap f(y) = 0$ , this is needed for subtraction.

Also for every automorphism  $f$  of  $A$  we assign the sequence  $\langle J, I, I_0, I_1, I_2, I_3, C_0, C_1, C_2 \rangle$  (some redundancy). Suppose for  $f_1, f_2 \in Aut(A)$  we get the same tuple; it is enough to show that their restriction to  $J$  and to  $I$  are equal -as the union is



## FACTOR = QUOTIENT, UNCOUNTABLE BOOLEAN ALGEBRAS

dense. Concerning  $J$  this is trivial - they are the identity on it so we discuss  $I$ , by an observation above it is enough to check it for  $I_t, t = 0, 1, 2$  but for each  $t$ , from  $C_t$  and  $I_t, I_{t+1}$  we can [or see below] reconstruct  $f|I_t$ .

So we have finished the proof.

**3.4B Remark.** We can phrase this argument as a claim: let  $I, J$ , be ideals of  $A$  with intersection  $\{0\}$ ; for every  $f$ , a one to one homomorphism from  $I$  to  $J$  let  $X_f$  be the set  $\{x \cup f(x) : x \in I\}$  then “ $f$  mapped to  $X_f$ ” is a one to one mapping from  $HOM(I, J)$  to  $Psub(A)$  (the former include  $\{g|I : g \in Aut(A), g \text{ maps } I \text{ onto } J\}$ , for which we use this). For subalgebra relative to  $\cup, \cap$  only  $f$  needs not be one to one.

*3.5 Proof of the conclusion from the theorem.*

- (1) For a given Boolean Algebra  $B$  assume  $\mu =: sub(A)$  is  $< end(A)$ . For any endomorphism  $f$  of  $A$  we attach the pair  $(Kernel(f), Range(f))$ . The number of possible such pairs is at most  $id(A) \times sub(A)$ , which is at most  $\mu$  (we are dealing with infinite  $BAs$  and  $id(A) \leq psub(A) \leq sub(A)$ ) so as we assume  $\mu < end(A)$ , there are distinct  $f_i$ , endomorphisms of  $A$  for  $i < \mu^+$ , an ideal  $I$  and a subalgebra  $R$  of  $B$  such that for every  $i$  we have  $Kernel(f_i) = I$  and  $Range(f_i) = R$ .

We now define a homomorphism  $g_i$  from  $B/I$  to  $R$  by :  $g_i(x/I) = f_i(x)$ . Easily the definition does not depend on the representative, so  $g_i$  is as required and it is one to one and onto. So  $\{g_i \circ (g_0)^{-1} : i < \mu^+\}$  is a set of  $\mu^+$  distinct automorphisms of  $R$ .

So

$$(*)_1 \quad \mu < aut(R)$$

but, by the theorem

$$(*)_2 \quad aut(R) \leq sub(R)$$

obviously

$$(*)_3 \quad sub(R) \leq sub(A)$$

remember

$$(*)_4 \quad \mu = sub(A)$$

together a contradiction

- (2) As  $R \mapsto (\text{identity map on } R)$  is one to one from  $sub(A)$  into  $Pend(A)$ , obviously  $sub(A) \leq pend(A)$ , so we are left with proving the other inequality. Same proof, only the domain is a subalgebra too and it has an ideal. So for every such partial endomorphism  $h$  of  $A$  we attach two subalgebras  $D_h = Domain(h)$  and  $R_h = Range(h)$  an ideal  $I_h$  of  $D_h$   $\{x : x \in D_h \text{ and}$

FACTOR = QUOTIENT, UNCOUNTABLE BOOLEAN ALGEBRAS

$f(x) = 0\}$ . They are all in  $Psub(A)$ , so their number is at most  $sub(A)$  and if we fixed them the amount of freedom we have left is : an automorphism of  $R$  (and  $aut(R) \leq sub(R) \leq sub(A)$ ).

- (3) Let  $B$  be a subalgebra of  $A$ . We shall attached to it several ideals and subalgebras of  $B|a, B|(-a)$  such that  $B$  can be reconstructed from them; this clearly suffice. Let  $C$  be the subalgebra  $\langle B, a \rangle, C_o = \{x \in C : x \leq a\}, C_1 = \{x \in C : x \leq 1 - a\}$ . The number of possible  $C$  is clearly the number of pairs  $\langle C_o, C_1 \rangle$  which is clearly  $sub(A|a) + sub(A|-a)$ ; fix  $C$ . Let  $I =: \{x : x \in B, x \leq a\}$ , it is an ideal of  $C|a$ ; so the number of such  $I$  is at most

$$id(C|a) \leq pend(C|A) \leq sub(C|a) \leq sub(A|a).$$

So we can fix it . Similarly we can fix  $J = \{y : y \in B, y \leq (1 - a)\}$ , now  $I$  and  $J$  are subsets of  $B$ , now check : the amount of freedom we have left is an isomorphism  $g$  from  $(C|a)/I$  onto  $(C|-a)/J$  such that  $B = \{$  the subalgebra of  $C$  generated by  $I \cup J \cup \{x \cup y : x \in (C|a), y \in (C|-a)$  and  $g(x/I) = (y/J)\}$ .

So we can finish easily.

We originally want to prove  $Aut(A)^{\aleph_0} \leq sub A$  and even

$|\{f \in End(A) : f \text{ is onto}\}|^{\aleph_0} \leq sub A$ . But we get more: intermediate invariants with reasonable connections.

**3.6 Definition:** For a Boolean algebra  $A$

- (0) A partial function  $f$  from  $A$  to  $A$  is everywhere onto if:  
 $x \in Dom(f) \& y \in Rang(f) \& y \leq f(x) \Rightarrow (\exists z)[z \leq x \& f(z) = y]$ .
- (1)  $End_0(A) = End(A)$ .  
 $End_1(A) = \{f \in End(A) : Rang(f) \text{ include a dense ideal}\}$ .  
 $End_2(A)$  is the set of endomorphism  $f$  of  $A$  which are onto.  
 $End_3(A) = \{f : \text{for some dense ideal } I, f \text{ is an onto endomorphism of } I\}$ .  
 $End_4(A) = \{f : \text{for some ideal } I \text{ of } A, f \text{ is an onto endomorphism of } I\}$ .  
 $End_5(A) = \{f : \text{for some dense ideals } I, J, f \text{ is an homorphism from } I \text{ onto } J\}$ .  
 $End_6(A) = \{f : \text{for some ideals } I, J \text{ of } A, f \text{ is an endomorphism from } I \text{ onto } J\}$ .
- Note**  $I = A$  is allowed. All kinds of endomorphism, commute with  $\cap, \cup$ , preserve 0 but not necessarily  $-$ .
- (2) For  $l = 1 \dots, 6$  we let  $Aut_l(A) = \{f \in End_l(A) : f(x) = 0 \Leftrightarrow x = 0 \text{ for } x \in Dom f\}$ .
- (3) For function  $f, g$  whose domain is  $\subseteq A$  let:  $f \sim g$  if  $Ker f = Ker g$  and  $\{x : f(x) = g(x) \text{ or both are defined not } \}$  include a dense ideal of  $B/Ker f$ .
- (4) Let  $end_e(A) = |End_e(A)|, aut_e(A) = |Aut_e(A)|$ .  
 Let  $end_e^{\sim}(A) = |\{f / \sim : f \in End_e(A)\}|$  and  $aut_e^{\sim}(A) = |\{f / \sim : f \in Aut_e(A)\}|$ .
- (5) We allow to replace  $A$  by an ideal  $I$  with the natural changes.
- (6) We define  $Endv_e(A), endv_e(A)$  similarly replacing “onto” by “everywhere onto” and define  $Endl_e(A), endl_e(A)$  similarly omitting “onto”. We define  $Endu_e(A)$  as the set of  $f \in End_e(A)$  such that for every  $x \in Dom f, f(x) \neq 0$  and ideal  $I$  of  $A|x$  which is dense, we have  $f(x) = sup_A\{f(y) : y \in I\}$ . We defined naturally  $Autv_e(A), Autl_e(A), Autu_e(A)$  etc.

FACTOR = QUOTIENT, UNCOUNTABLE BOOLEAN ALGEBRAS

**Note:** In  $Endv_1(A)$  we mean: for some dense ideal  $I$  of  $A, y \leq f(x) \in I \Rightarrow (\exists z \leq x)(f(z) = y)$  and  $Endl_1(A) = End(A)$ .

**3.7 Claim.**

- (1)  $End_6(A) \supseteq End_5(A) \cup End_4(A) \supseteq End_5(A) \cap End_4(A) \supseteq End_3(A) \supseteq End_2(A)$  and  $End_2(A) \subseteq End_1(A) \subseteq End_0(A) = EndA$ .
- (2)  $end_6(A) \geq \max\{end_5(A), end_4(A)\} \geq \min\{end_5(A), end_4(A)\} \geq end_3(A) \geq end_2(A)$ , and  $end_2(A) \leq end_1(A) \leq end(A)$ .
- (3) In (1) we can replace  $End$  by  $Aut$  or  $Endv$  or  $aEndl$  or  $Autv$  or  $Autl$ , or  $Endu$  or  $Autu$  and in (2) replace by  $aut$  etc. We can in (2) replace  $end_l$  (or  $aut_l$ ) by  $end_l^\sim$  (or  $aut_l^\sim$ ) etc.
- (4)  $Aut_l(A) \subseteq End_l(A)$  hence  $aut_l(A) \leq end_l(A)$ ; and  $Endv_e(A) \subseteq End_e(A) \subseteq Endl_e(A)$  hence  $endv_e(A) \leq end_e(A) \leq endl_e(A)$ ;  $Endv_e(A) = End_e(A)$  if  $e = 0$ . Also  $Aut(A) = Aut_2^\sim(A)$  hence  $aut(A) = aut_2^\sim(A)$ .
- (5) If  $f, g, \in Endv_6(A)$ ,  $f \sim g$  then  $f, g$ , are compatible functions, i.e.  $x \in Domf \cap Domg \Rightarrow f(x) = g(x)$ .
- (6)  $sub(A) \geq aut_6(A)$  and  $aut(A) \leq aut_3^\sim(A)$ .
- (7)  $Aut_e(A) = Autv_e(A)$ .
- (8)  $end_e(A) \leq id(A) + end_e^\sim(A)$  etc.

*Proof.* E.g.

- (5) Suppose  $x \in Domf \cap Domg$ ,  $f(x) \neq g(x)$ , so  $wlog f(x) \not\leq g(x)$  so let  $z =: f(x) - g(x) \neq 0$ . As  $z \leq f(x)$  for some  $t \in Domf$  we have  $0 < f(t) \leq z$ , and  $wlog f(t) = g(t)$ , and we get contradiction.
- (6) As in proof of 3.2 or see proof of 3.12 (noting  $|\{I : I \text{ a dense ideal of } A\}| \leq sub(A)$  by 3.4A).

**3.8 Definition.** For a Boolean Algebra  $A$

- (1)  $Idc(A) = \{I : I \text{ an ideal of } A, \text{ and } I^c = I\}$  where  $I^c = \{x \in A : I \text{ is dense below } x\}$ .
- (2)  $idc(A) = |Idc(A)|$ .
- (3)  $Did(A) = \{I : I \text{ a dense ideal of } A\}$ .
- (4) For ideals  $I, J$   $I + J$  is the minimal ideal  $I$  of  $A$  which include  $I \cup J$  i.e.  $\{x \cup y : x \in I \text{ and } y \in J\}$ . Similarly  $\sum_{\zeta < \xi_0} I_\zeta$ .
- (5) If we replace  $A$  by an ideal  $I$  (in 3.8 (1),(2),(3), 3.1 (2)) means we restrict ourselves to subideals of it.

**3.9 Claim.** For a Boolean algebra  $A$ :

- (1)  $id(A) \geq idc(A) = |comp(A)|$ .
- (2)  $|A| \leq idc(A) = idc(A)^{\aleph_0}$  (when  $A$  is infinite, of course).
- (3) If  $f \in Endu_5(A)$ , then  $f$  has a unique extension to an endomorphism  $f^+$  of  $comp(A)$  where  $f^+(x) = sup_A\{f(y) : y \leq x, y \in Domf\}$ . If  $f$  is everywhere onto it is the unique extension of  $f$  in  $End(compA)$ .
- (4) For  $g \in End(A)$ ,  $(\exists f \in End_5(A))f^+ \supseteq g$  iff  $g \in Endu(A)$ .
- (5) For  $f, g \in End_5(A)$  we have  $f \sim g \Leftrightarrow f^+ = g^+$ .
- (6) For  $f \in Endu_6(A)$ , letting  $a = sup_{comp(A)}\{x : B|x \subseteq Domf\} \in comp(A)$  and  $b = sup_{comp(A)}\{x : A|x \subseteq Rangf\} \in comp(A)$ , we define  $f^+ \in HOM(comp(A|a), comp(A|b))$  extending  $f$  by  $f(x) = sup\{f(y) : y \leq x, y \in Domf\}$ , also  $f^+$  is onto.

**3.10 Claim.**

- (1)  $id(A) \leq endl_3(A)$ .
- (2)  $idc(A) \leq endl_3^\sim(A)$ .
- (3)  $endl_e(A) = id(A) + endl_e^\sim(A)$  for  $e = 3, 4, 5, 6$ .
- (4) If  $f \in End_5(A)$  then  $f^+ \in End_1(compA)$ .
- (5)  $aut_e(A) \leq id(A) + aut_m^\sim(A)$  when  $e, m \in \{3, 5\}$  or  $e, m \in \{4, 6\}$ .

*Proof.*

- (1) For  $I \in Id(A)$  let  $J_I = \{x \in B : (B|x) \cap I = \{0\}\}$ , so  $J_I$  is an ideal,  $J_I \cap I = \{0\}$ ,  $J_I \cup I$  is dense. Let  $F_I$  be the following map:  $Dom f_I = I + J_I$ ,  $f_I|I = id_I$ ,  $f_I|J_I = 0_{J_I}$ . Now  $I \mapsto f_I$  is a one to one mapping from  $Id(A)$  to  $Endl_3(A)$ .
- (2) The mapping above works.
- (3) Note that  $id(A) \leq endl_3(A)$  by part(1), and  $endl_3^\sim(A) \leq endl_3(A)$  trivially.  $A$  is infinite hence all those cardinals are infinite so  $\chi =: id(A) + endl_3^\sim(A) \leq endl_3(A)$ . The inverse inequality is easy too.
- (4), (5) Left to the reader.

**3.11 Claim.**

- (1) For  $x \in A$ :  
 $(\exists f \in Aut(A))[x \neq f(x)]$  iff  
 $(\exists f \in Aut_6(A))(x \in Dom f \& x \neq f(x))$  iff  $(\exists y, z)[0 < y \leq x \& 0 < z \leq 1 - x \& A|y \cong A|z]$ .
- (2) If  $\langle I_\zeta : \zeta < \alpha \rangle$  is a sequence of ideals of  $A$ ,  $[\zeta \neq \xi \Rightarrow I_\zeta \cap I_\xi = \{0\}]$  and  $I = \sum_{i < \alpha} I_i$  then:  
 $id(I) \geq \pi_{\zeta < \alpha} id(I_\zeta)$ .  
 $idc(I) \geq \pi_{\zeta < \alpha} idc(I_\zeta)$ .  
 $aut(I) \geq \pi_{\zeta < \alpha} aut(I_\zeta)$ .  
 $end(I) \geq \pi_{\zeta < \alpha} end(I_\zeta)$ .  
 Similarly for  $end_l, aut_l, end_l^\sim, aut_l^\sim$  etc.

There are many more easy relations, but for our aim the main point is

**3.12 Main Lemma.** For an infinite Boolean Algebra  $A$ :

- (1)  $aut_e^\sim(A)$  for  $e = 3, 5$  are equal or both finite (and we can restrict ourselves to automorphisms of order 2).
- (2) If for some  $e \in \{3, 4, 5, 6\}$  we have  $aut_e^\sim(A) > idc(A)$  then  $aut_e^\sim(A)$  for  $e = 3, 4, 5, 6$  are all equal.
- (3)  $aut_3^\sim(A) = aut_3^\sim(A)^{\aleph_0}$  or  $aut_3(A)$  is finite.
- (4)  $autv_e^\sim(A) = autv_3(A) + idc(A)$  for  $e = 4, 6$ .

*Proof.* Let  $J =: \{x \in A : \text{for every } f \in Aut(A), f|(A|x) = id_{A|x}\}$ .

The function  $F_1, \dots, F_5$  satisfying  $y \leq x \Rightarrow F_e(y) \leq F_e(x)$  are functions from  $A$  to  $ord$  defined below; and we let:

$K =: \{x \in A :$

- (i) for some  $y, x \cap y = 0$  and  $A|x \cong A|y$ .

FACTOR = QUOTIENT, UNCOUNTABLE BOOLEAN ALGEBRAS

(ii) for  $e = 1, \dots, 5$  and  $0 < y < x$  we have  $F_e(y) = F_e(y)$ .

Where:

$F_1(x)$  = the cardinality of  $A|x$ .

$F_2(x) = \text{idc}(A|x)$ .

$F_3(x) = \text{aut}(A|x)$ .

$F_4(x) = \text{aut}_3(A|x)$ .

$F_5(x) = \text{aut}_3^\sim(A|x)$ .

Now

(\*)<sub>1</sub>  $J$  is an ideal of  $B$ .

(\*)<sub>2</sub>  $K$  is downward closed.

(\*)<sub>3</sub>  $K \cup J$  is dense.

Choose  $\{x_i : i < \alpha\}$  maximal such that:

(a)  $x_i \in K, x_i > 0$ ,

(b) if  $i \neq j, 0 < y' \leq x_i, 0 < y'' \leq x_j$  then  $A|y' \not\cong A|y''$ .

Let  $K_i = \{y : \text{for some } y' \leq x_i, A|y \cong A|y'\}$ , and  $K_i^+$  : the ideal  $K_i$  generate.

Now

(\*)<sub>4</sub>  $\bigcup_{i < \alpha} K_i$  is dense in  $K$  [hence  $J \cup \bigcup_{i < \alpha} K_i$  is dense in  $A$ ].

(\*)<sub>5</sub> For  $i \neq j$  we have  $K_i \cap K_j = \{0\}, K_i \cap J = \{0\}$ .

Clearly,

(\*)<sub>6</sub> for  $f \in \text{Aut}_6(A)$  we have:

(a)  $f$  is the identity on  $J \cap \text{Dom} f$ .

(b) for  $x \in \text{Dom} f$  we have  $x \in K_i \Leftrightarrow f(x) \in K_i$ .

So

(\*)<sub>7</sub>  $\text{aut}_e^\sim(A) = \pi_{i < \alpha} \text{aut}_e^\sim(K_i^+)$  for  $e = 3, 5$  and  $\text{aut}_e^\sim(A) = \text{idc}(J) \times \pi_{i < \alpha} \text{aut}_e^\sim(K_i^+)$  ■

for  $e = 4, 6$

We shall prove:

(\*)<sub>8</sub> For each  $i$ , one of the following occurs:

(a)  $\text{aut}_e^\sim(K_i^+)$  for  $e = 3, 4, 5, 6$  are all finite  $> 1$ ,

(b) For some infinite  $\kappa$  we have  $\text{aut}_e^\sim(K_i^+) = \text{aut}_e^\sim(K_i^+)^{\kappa} \geq \text{idc}(K_i)$  for  $e = 3, 4, 5, 6$

(really we can use  $F_6(i) = \text{sup}\{\kappa^+ : B|x \text{ has } \kappa \text{ pairwise disjoint non zero members}\}$  and any such  $\kappa$  is OK for (b)).

**Case 1.**  $x_i$  is an atom.

This is clear: let  $\lambda_i = |K_i|$ , if it is infinite,  $\text{aut}_e^\sim(K_i^+) = 2^{\lambda_i}$  for  $e = 3, 4, 5, 6$  so case (b) in (\*)<sub>8</sub> occurs.

If  $\lambda_i$  is finite,  $1 < \text{aut}_e^\sim(K_i^+) < \aleph_0$  (we can compute exactly), so case (a) in (\*)<sub>8</sub>. In fact in all cases we can use just automorphism of order 2.

So we can assume

**Case 2** . not Case 1, so  $B|x_i$  is atomless, hence  $\text{idc}(B|x_i) \geq 2^{\aleph_0}$

Let  $\langle J_{i,\zeta}, J'_{i,\zeta} : \zeta < \zeta_i \rangle$  be a sequence such that:

( $\alpha$ )  $J_{i,\zeta}, J'_{i,\zeta}$  are ideals  $\subseteq K_i^+$  and  $\neq \{0\}$ ,

( $\beta$ )  $\{0\} \neq J_{i,\zeta} \subseteq A|x_i$ ,

( $\gamma$ )  $J_{i,\zeta} \cong J'_{i,\zeta}$ , an  $h_{i,\zeta}$  an isomorphism from  $J_{i,\zeta}$  onto  $J'_{i,\zeta}$ ,

( $\delta$ )  $\wedge_{\zeta < \xi < \zeta_i} J'_{i,\zeta} \cap J'_{i,\xi} = \{0\}$ ,

FACTOR = QUOTIENT, UNCOUNTABLE BOOLEAN ALGEBRAS

- ( $\epsilon$ ) if  $y \in K_\zeta^+$  is disjoint to all members  $\cup_{\xi < \zeta} J'_{i,\xi}$  of then for some  $y' \leq y$  and  $z \in J_{i,\zeta}, B|z' \cong B|y'$  hence
- ( $\zeta$ ) if  $\zeta < \xi < \zeta_i$  then  $J_{i,\xi} \cap J_{i,\xi}$  is a dense subset of  $J_{i,\xi}$

Now

- (\*)<sub>9</sub>  $\cup_{\zeta < \zeta_i} J'_{i,\zeta}$  is a dense subset of  $K_i^+$ .
  - (\*)<sub>10</sub>  $\zeta_i \geq 2$  (as there is  $y, x_i \cap y = 0$  and  $A|y \cong A|x_i$  as  $x_i \in K$ ).
  - (\*)<sub>11</sub>  $idc(K_i^+) = \pi_{\zeta < \zeta_i} idc(J'_{i,\zeta})$ .
- By the definition of  $K$  (see choice of  $F_2$ ), we have  $0 < y \leq x_i \Rightarrow idc(A|y) = (A|x)$  hence  $idc(J'_{i,\zeta}) = idc(A|x_i)$  so
- (\*)<sub>12</sub>  $idc(K_i^+) = [idc(A|x_i)]^{|\zeta_i|}$  hence  $[idc(K_i^+)]^{|\zeta_i|} = idc(K_i^+)$ .

Easily

- (\*)<sub>13</sub>  $aut_6^{\sim}(K_i^+) \geq aut_3^{\sim}(K_i^+) \geq idc(A|x_i)$ .
- (and even by automorphism of order 2).

Last inequality: for each  $z \in Idc(A|x_i)$  there is  $z', z' \cap x_i = 0$ , and  $A|z \cong A|z'$  and let  $g$  be such isomorphism, let  $I_z$  be the ideal of  $A$  generated by  $\{x \in K_i^+ : x \leq z \text{ or } x \leq z' \text{ or } x \cap z = x \cap z' = 0\}$   $g_z \in Aut(I_z) \subseteq Aut_3(K_3^+)$ ,  $g_z(y) =: (y - z \cup z') \cup g(y \cap z) \cup g^{-1}(y \cap z')$ .

**Case 2A.**  $\zeta_i < \aleph_0$  (and not Case 1)

Let for  $n < \omega$   $x_{i,n} \leq x_i, x_{i,n} \neq 0$ , and  $[n \neq m \Rightarrow x_{i,n} \cap x_{i,m} = 0]$ , and  $I_{i,\omega} = \{y \in K_i^+ : \wedge_n y \cap x_{i,n} = 0\}$  and  $I_{i,n} = \{z : z \leq x_{i,n}\}$  : So  $aut_3^{\sim}(K_i^+) \geq aut_3^{\sim}(\sum_{\alpha \leq \omega} I_{i,\alpha}) \geq \pi_{n < \omega} aut_3^{\sim}(B|x_{i,n}) = aut_3^{\sim}(B|x_i)^{\aleph_0}$ .

Similarly to the proof of Case 2B below (but easier) we can show that Case (b) of (\*)<sub>8</sub> occurs. From now on we assume

**Case 2B.** Not Case 2A, so  $\zeta_i \geq \aleph_0$

For every  $f \in Aut_6(K_i^+)$  let, for  $\zeta, \xi < \zeta_i$ :

$$L_{\zeta,\xi}^f = \{x \in J'_{i,\zeta} : f(x) \in J'_{i,\xi}\}.$$

$$M_{\zeta,\xi}^f = \{f(x) : x \in J'_{i,\zeta}, f(x) \in J'_{i,\xi}\}.$$

So the number of possible  $\bar{L}^f = \langle sup L_{\zeta,\xi}^f : \zeta \neq \xi < \zeta_i \rangle$  and  $\bar{M}^f = \langle sup(M_{\zeta,\xi}^f) : \zeta \neq \xi < \zeta_i \rangle$  is  $\leq [idc(A|x_i)]^{|\zeta_i|}$  and for fixed  $\bar{L}, \bar{M}$  the number of  $f \in aut_6^{\sim}(K_i^+)$  for which  $\bar{L}^f = \bar{L}, \bar{M}^f = \bar{M}$  is  $\leq \pi_{\zeta,\xi} |\{f/ \sim : f \text{ an isomorphism from a dense subset of } L_{\zeta,\xi} \text{ onto a dense subset of } M_{\zeta,\xi}\}| \leq \pi_{\zeta,\xi} aut_3^{\sim}(L_{\zeta,\xi}) \leq \pi_{\zeta,\xi} aut_3^{\sim}(A|x_i) = [aut_3^{\sim}(A|x_i)]^{|\zeta_i|}$ .

(In the last equality we use  $F_5$  in the definition of  $K$ : for the last  $\leq$ , note we can replace  $L_{\zeta,\xi}$  by isomorphc ideal  $L_{\zeta,\xi}^*$  of  $A|x_i$ , and letting  $L_{\zeta,\xi}^- = \{y : y \in A|x_i, (\forall z \in L_{\zeta,\xi}^*) [y \cap z = 0]\}$  we can extend every  $f \in Aut_3(L^*(\zeta, \xi))$  to  $f' \in Aut_3(A|x)$  by letting  $f'(x \cup y) = f(x) \cup y$  for  $x \in Dom(f), y \in L_{\zeta,\xi}^-$ )

So

$$(*)_{14} \quad aut_6^{\sim}(K_i^+) \leq [aut_3^{\sim}(A|x_i) + idc(A|x_i)]^{|\zeta_i|}.$$

Now for each  $\xi$  such that  $2\xi + 1 < \zeta_i$  we can choose  $y_{i,2\xi} \in J_{i,2\xi}, y_{i,2\xi+1} \in J_{i,2\xi+1}$  such that  $A|y_{i,2\xi} \cong A|y_{i,2\xi+1}$  and  $\langle g_{i,\xi,\alpha} : \alpha < aut_3^{\sim}(A|x_i) \rangle$  such that  $g_{i,\xi,\alpha}$  is an isomorphism from a dense subset of  $A|y_{i,\xi,\alpha}$  onto a dense subset of  $A|y_{i,2\xi+1}$ ,  $\langle g_{i,\xi,\alpha}^+ : \alpha < aut_3^{\sim}(A|x_i) \rangle$  pairwise distinct. Let  $\zeta_i^*$  be minimal such that  $2\zeta_i^* \leq \zeta_i$ .

Now for every sequence  $\bar{\alpha} = \langle \alpha_\xi : \xi < \zeta_i^* \rangle$ ,  $\alpha_\xi < aut_3^{\sim}(A|x_i)$ , we define  $g_{i,\bar{\alpha}} \in Aut_3(K_i^+)$  of order two (see condition ( $\gamma$ )):

$$g_{i,\bar{\alpha}}|(y_{i,2\xi} \cup y_{i,2\xi+1}) = h_{2\xi+1} \circ g_{i,\xi,\alpha} \circ h_{2\xi}^{-1} \cup h_{2\xi} \circ g_{i,\xi,\alpha}^{-1} \circ h_{2\xi+1}^{-1}$$

## FACTOR = QUOTIENT, UNCOUNTABLE BOOLEAN ALGEBRAS

and if  $y \in K_i^+ \setminus \{0\}$  and  $\bigwedge_{\xi < \zeta_i^*} (y \cap y_{i,\xi} = 0)$  then  $g_{i,\bar{\alpha}}(y) = 0$ .

Lastly if  $y$  is the disjoint union of  $y_0, \dots, y_n$  and each  $g_{\bar{\alpha}}(y_e)$  was defined then we define  $g_{\bar{\alpha}}(y_0 \cup \dots \cup y_n) = g_{\bar{\alpha}}(y_0) \cup \dots \cup g_{\bar{\alpha}}(y_n)$ . The reader may check that  $g_{\bar{\alpha}} \in \text{Aut}_3(K_i^+)$ .

The mapping  $\bar{\alpha} \mapsto g_{i,\bar{\alpha}}$  show:

$$(*)_{15} \text{aut}_3^{\sim}(K_i^+) \geq \text{aut}_3^{\sim}(A|x_i)^{|\zeta_i|}.$$

Easily, choosing  $y_i \in J_{i,1}$  (possible as  $\zeta_i \geq 2$ )

$$(*)_{16} \text{aut}_3^{\sim}(K_i^+) \geq \text{aut}_3^{\sim}(A|(x_i \cup y_i)) \geq \text{idc}(A|y) = \text{idc}(A|x_i) \geq \aleph_0.$$

Together by  $(*)_{13}, (* )_{14}, (* )_{15}, (* )_{16}$  for  $e = 3, 6$ :

$$\text{aut}_e^{\sim}(K_i^+) = \text{aut}_3^{\sim}(A|x_i)^{|\zeta_i|}.$$

By 3.7(1) this holds for  $e = 4, 5$ , hence  $(*)_8$  has been proved.

Now by  $(*)_7 + (* )_8$  the four parts of 3.12 follows.

**3.13 Conclusion :**  $\text{aut}(A)^{\aleph_0} \leq \text{sub}(A)$ , also  $\text{aut}_e(A)^{\aleph_0} \leq \text{sub}(A)$  for  $e = 3, 4, 5, 6$ .

Note: even if  $\text{aut}(A)$  is finite,  $A$  infinite, still  $2^{\aleph_0} \leq \text{sub}(A)$  (for  $A$  infinite).

*Proof.* By 3.7(3) + 3.7(6)  $\text{aut}(A) = \text{aut}_0^{\sim}(A) \leq \text{id}(A) + \text{aut}_3^{\sim}(A) \leq \text{id}(A) + \text{Pend}(A) \leq \text{sub}(A)$  but by S. Shelah [6]  $\text{id}(A)^{\aleph_0} = \text{id}(A)$  and by 3.12(3)  $\text{aut}_3^{\sim}(A)^{\aleph_0} = \text{aut}_3^{\sim}(A)$ , together we can finish the first inequality, the second is similar using 3.10(5).

**3.14 Claim :** For an (infinite) Boolean algebra  $A$  we have:

- (1)  $\text{end}_e(A) = \text{id}(A) + \text{aut}_3(A)$  for  $e = 3, 4, 5, 6$ .
- (2)  $\text{end}_6(A)^{\aleph_0} \leq \text{sub}(A)$ .
- (3)  $\text{end}_e(A)^{\aleph_0} \leq \text{sub}(A)$  for  $e = 2, 3, 4, 5, 6$ .

*Proof.*

- (1) Clearly  $\text{end}_3(A) \geq \text{aut}_3(A)$  and  $\text{end}_3(A) \geq \text{id}(A)$  (as the mapping in the proof of 3.10(1) exemplify).

On the other hand we can attach to every  $f \in \text{End}_6(A)$  three ideals  $I_1(f) = \text{Dom } f$ ,  $I_2(f) = \text{Rang}(f)$  and  $I_3(f) = \text{Ker}(f)$ . Now the number of triples  $\bar{I}$  of ideal of  $A$  has cardinality  $\text{id}(A)$  and for each such  $\bar{I}$ :

$\{f \in \text{End}_6(A) : I_e(f) = I_e \text{ for } e = 1, 2, 3\}$  has cardinality  $|\text{Aut}(I_2)|$  which is  $\leq \text{aut}_3(A)$ . By 3.7(2) we can finish.

- (2) Remember also  $\text{id}(A)^{\aleph_0} = \text{id}(A)$  by [6] and part (1) and 3.13.
- (3) By part (2) and 3.7.

### §4 The width of the Boolean algebra .

**4.1 Definition.** For a Boolean algebra  $B$  let: (1)  $A \subseteq B$  is an antichain if  $x \in A \& y \in A \& x \neq y \Rightarrow x \not\leq y$  (i.e.  $A$  is a set of pairwise incomparable elements).

(2) Width of  $B$ ,  $w(B)$  is  $\sup\{|A| : A \subseteq B \text{ antichain}\}$ ,  $w^+(B) = \cup\{|A|^+ : A \subseteq B \text{ antichain}\}$

E. C. Milner and M. Poizat [3], answering a question of E. K. van Dowen, D. Monk and M. Rubin [7] proved  $\text{cf}(w^+(B)) \neq \aleph_0$ .

In S. Shelah [5] we claim: if  $\lambda > \text{cf } \lambda > \aleph_0$ , for some generic extension of the universe preserving cardinalities and cofinalities, for some  $B$ ,  $w^+(B) = \lambda$ . We retract this and replace it by the theorem 4.2 below.

FACTOR = QUOTIENT, UNCOUNTABLE BOOLEAN ALGEBRAS

For weakly inaccessible we still have the consistency. Moreover, if  $\lambda$  is a limit uncountable regular cardinal,  $S \subseteq \lambda$  stationary not reflecting and  $\diamond_S$  it then we have such an example for  $\lambda$ .

**4.2 Theorem.** *For an infinite Boolean algebra  $B$ ,  $w^+(B)$  is an uncountable regular cardinal.*

*Proof.* As  $B$  is infinite it has an antichain  $A$ ,  $|A| = \aleph_0$ , [if  $B$  has finitely many atoms clear, if not it has a subalgebra which is atomless, without loss of generality countable and check]. So  $\lambda =: w^+(B) > \aleph_0$ . Assume  $\kappa =: cf\lambda < \lambda$ ; let  $\lambda = \sum_{i < \kappa} \lambda_i$ ,  $cf\lambda + \sum_{j < i} \lambda_j < \lambda_i < \lambda$  and let  $A_i \subseteq B$  be an antichain of cardinality  $\lambda_i^+$  (exist by the choice of  $\lambda$ ). Let  $A = \bigcup_{i < \kappa} A_i$ , so  $|A| = \lambda$ . Choose such  $\langle A_i : i < \kappa \rangle$  such that, if possible

(\*)  $i < j < \kappa$ ,  $x \in A_i$ ,  $y \in A_j \Rightarrow y \not\leq x$ .

For  $x \in B$  let  $A[>, x] = \{y \in A : y > x\}$ ,  $A[<, x] = \{y \in A : y < x\}$ ,  
 $A[>, \mu] = \{x \in A : |A[>, x]| < \mu\}$ ,  $A[<, \mu] = \{x \in A : |A[<, x]| < \mu\}$ .

**Case 1.** For some  $\mu < \lambda$ ,  $A[>, \mu]$  has cardinality  $\lambda$ .

By Hajnal free subset theorem, there is a set  $E \subseteq A[>, \mu]$  of cardinality  $\lambda$  such that:

$x \neq y \& x \in E \& y \in E \Rightarrow x \notin A[>, y] \& y \notin A[>, x]$ . So  $E$  witness  $w^+(B) > \lambda$ .

**Case 2.** For some  $\mu < \lambda$ ,  $A[<, \mu]$  has cardinality  $\lambda$ .

Same proof.

**Case 3.** For every  $i < \kappa$  there is  $x \in A$  such that  $\lambda_i < |A[>, x]| < \lambda$ .

Let for  $i < \kappa$ ,  $x_i \in A$  be such that  $\lambda_i < |A[>, x_i]| < \lambda$ . Let  $u \subseteq \kappa$  be such that:  $\kappa = supu$  and for  $i \in u$ ,  $\lambda_i > \sum_{j \in u \cap i} |A[>, x_j]|$  (choose the members of  $u$  inductively). By renaming without loss of generality  $u = \kappa$ . Clearly  $A[>, x_i] \setminus \bigcup_{j < i} A[>, x_j]$  has cardinality  $> \lambda_i$ .

As  $\lambda_i > \kappa$  (by its choice) and  $A = \bigcup_{j < \kappa} A_j$ , clearly for each  $i$  there is  $\alpha(i) < \kappa$  such that  $(A[>, x_i] \setminus \bigcup_{j < i} A[>, x_j]) \cap A_{\alpha(i)}$  has cardinality  $> \lambda_i$ ; necessarily  $\alpha(i) \geq i$ .

For some unbounded  $u \subseteq \kappa$  we have  $[i \in u \& j \in u \& i < j \Rightarrow \alpha(i) < j]$ ; without loss of generality  $u = \kappa$ ,  $\alpha_i = i$ . Let  $A_i^*$  be a subset of  $(A[>, x_i] \setminus \bigcup_{j < i} A[>, x_j]) \cap A_{\alpha(i)}$  of cardinality  $\lambda_i^+$ . Now  $\langle A_i^* : i < \kappa \rangle$  satisfies:  $A_i^* \subseteq B$  is an antichain of cardinality  $\lambda_i^+$  and

(\*)'  $i < j$ ,  $x \in A_i^*$ ,  $y \in A_j^* \Rightarrow x \not\leq y$  (otherwise  $x_i \leq x \leq y \notin A[>, x_i]$ , contradiction).

So  $\langle A'_i =: \{1_B - x : x \in A_i^*\} : i < \kappa \rangle$  satisfies  $A'_i \subseteq B$  is an antichain of  $B$  of cardinality  $\lambda_i^+$  and also (\*) above (check). So by the choice of  $\langle A_i : i < \kappa \rangle$ , it satisfies (\*). By (\*) + (\*)',  $A^* = \bigcup_{i < \kappa} A_i^*$  is an antichain of  $B$  of cardinality  $\lambda$ , so  $w^+(B) > \lambda$ .

**Case 4.** For every  $i < \kappa$  there is  $x \in A$  such that  $\lambda_i < |A[<, x]| < \lambda$ .

Similar to Case 3.

**Case 5.** None of the previous cases.

By “not Case 3” for some  $i(*) < \kappa$ , for no  $x \in A$  is  $\lambda_{i(*)} < |A[>, x]| < \lambda$ . By not Case 2,  $A[<, \lambda_{i(*)}^+]$  has cardinality  $< \lambda$ . By not Case 1  $A[>, \lambda_{i(*)}^+]$  has cardinality  $< \lambda$ .

Choose  $x^* \in A \setminus A[<, \lambda_{i(*)}^+] \setminus A[>, \lambda_{i(*)}^+]$  so  $A[>, x^*]$  has cardinality  $\geq \lambda_{i(*)}^+ > \lambda_{i(*)}$ , hence by the choice of  $i(*)$  we have  $A[>, x^*]$  has cardinality  $\lambda$ .



## FACTOR = QUOTIENT, UNCOUNTABLE BOOLEAN ALGEBRAS

As  $\lambda_{i(*)} > \kappa$ , for some  $j(*)$ ,  $A[<, x^*] \cap A_{j(*)}$  has cardinality  $> \lambda_{i(*)}$ , so choose distinct  $y_i \in A[<, x^*] \cap A_{j(*)}$  for  $i < \kappa$ . Now  $y_i < x^*$  (as  $y_i \in A[<, x^*]$ ), and  $[i \neq j \Rightarrow y_i \not\leq y_j]$  (as they are distinct and in  $A_{j(*)}$ ).

Let  $A'_i = A_i \cap A[>, x^*]$ , so  $A'_i \subseteq A_i$  hence is an antichain of  $B$ , and

$$\cup_i A'_i = (\cup A_i) \cap A[>, x^*] = A \cap A[>, x^*] = A[>, x^*]$$

So each  $A'_i$  is an antichain, its member are  $> x^*$  and  $|\cup_i A'_i|$  is  $\lambda$  as  $|A[>, x^*]|$  is.

Now

$$A' = \cup_{i < \kappa} \{y_i \cup (x - x^*) : x \in A'_i\}$$

is an antichain of  $B$  of cardinality  $\lambda$ , so  $w^+(B) > \lambda$ , as required.

### REFERENCES

1. M. Bekkali, R. Bonnet and M. Rubin, *Spaces for which every closed subspace is homeomorphic to a closed subspace*, to appear Order.
2. R. Bonnet and S. Shelah, *On HCO spaces. An uncountable compact  $T_2$  space different from  $\aleph_1 + 1$  which is homeomorphic to every of its uncountable closed subspaces*, to appear Israel J. Math.
3. E. C. Milner and M. Pouzet, *On the width of ordered sets and Boolean Algebras*, Algebra Universatis **23** (1986), 242–253.
4. J. D. Monk, *Cardinal Functions on Boolean Algebra, Lectures in Mathematics*, ETH Zurich, Birkhäuser Verlag, Basel, Boston, Berlin, 1990.
5. S. Shelah, *Construction of many complicated uncountable structures and Boolean Algebras*, Israel J. Math. **23** (1983), 100–146, [Sh 136].
6. S. Shelah, *Remarks on the number of ideals of Boolean algebras and open sets of a topology*, Springer-Verlag Lecture Notes **1182** (1982), 151–187, [Sh 233].
7. E. K. van Dowen, D. Monk and M. Rubin, *Some questions about Boolean Algebras*, Algebra Universatis **11** (1980), 220–243.

THE HEBREW UNIVERSITY OF JERUSALEM, ISRAEL, RUTGERS UNIVERSITY, NEW BRUNSWICK, ■  
 NJ USA, MSRI, BERKELEY, CALIF, U.S.A.