

APPEARED IN BULLETIN OF THE
AMERICAN MATHEMATICAL SOCIETY
Volume 26, Number 2, April 1992, Pages 258-263

THE A_ℓ AND C_ℓ BAILEY TRANSFORM AND LEMMA

STEPHEN C. MILNE AND GLENN M. LILLY

ABSTRACT. We announce a higher-dimensional generalization of the Bailey Transform, Bailey Lemma, and iterative “Bailey chain” concept in the setting of basic hypergeometric series very well-poised on unitary A_ℓ or symplectic C_ℓ groups. The classical case, corresponding to A_1 or equivalently $U(2)$, contains an immense amount of the theory and application of one-variable basic hypergeometric series, including elegant proofs of the Rogers-Ramanujan-Schur identities. In particular, our program extends much of the classical work of Rogers, Bailey, Slater, Andrews, and Bressoud.

1. INTRODUCTION

The purpose of this paper is to announce a higher-dimensional generalization of the Bailey Transform [2] and Bailey Lemma [2] in the setting of basic hypergeometric series very well-poised on unitary [19] or symplectic [14] groups. Both types of series are directly related [14, 18] to the corresponding Macdonald identities. The series in [19] were strongly motivated by certain applications of mathematical physics and the unitary groups $U(n)$ in [10, 11, 15, 16]. The unitary series use the notation A_ℓ , or equivalently $U(\ell + 1)$; the symplectic case, C_ℓ . The classical Bailey Transform, Lemma, and very well-poised basic hypergeometric series correspond to the case A_1 , or equivalently $U(2)$.

The classical Bailey Transform and Bailey Lemma contain an immense amount of the theory and application of one-variable basic hypergeometric series [2, 12, 25]. They were ultimately inspired by Rogers’ [24] second proof of the Rogers-Ramanujan-Schur identities [23]. The Bailey Transform was first formulated by Bailey [8], utilized by Slater in [25], and then recast by Andrews [4] as a fundamental matrix inversion result. This last version of the Bailey Transform has immediate applications to connection coefficient theory and “dual” pairs of identities [4], and q -Lagrange inversion and quadratic transformations [13].

The most important application of the Bailey Transform is the Bailey Lemma. This result was mentioned by Bailey [8; §4], and he described how the proof would work. However, he never wrote the result down explicitly and thus missed the full power of *iterating* it. Andrews first established the Bailey Lemma explicitly in [5] and realized its numerous possible applications in terms of the iterative “Bailey chain” concept. This iteration mechanism enabled him to derive many

1991 *Mathematics Subject Classification*. Primary: 33D70, 05A19.

S. C. Milne was partially supported by NSF grants DMS 86-04232, DMS 89-04455, and DMS 90-96254

G. M. Lilly was fully supported by NSA supplements to the above NSF grants and by NSA grant MDA 904-88-H-2010

Received by the editors April 28, 1991

©1992 American Mathematical Society
0273-0979/92 \$1.00 + \$.25 per page

q -series identities by “reducing” them to more elementary ones. For example, the Rogers-Ramanujan-Schur identities can be reduced to the q -binomial theorem. Furthermore, general multiple series Rogers-Ramanujan-Schur identities are a direct consequence of iterating suitable special cases of Bailey’s Lemma. In addition, Andrews notes that Watson’s q -analog of Whipple’s transformation is an immediate consequence of the second iteration of one of the simplest cases of Bailey’s Lemma. Continued iteration of this same case yields Andrews’ [3] infinite family of extensions of Watson’s q -Whipple transformation. Even Whipple’s original work [26, 27] fits into the $q = 1$ case of this analysis. Paule [22] independently discovered important special cases of Bailey’s Lemma and how they could be iterated. Essentially all the depth of the Rogers-Ramanujan-Schur identities and their iterations is embedded in Bailey’s Lemma.

The process of iterating Bailey’s Lemma has led to a wide range of applications in additive number theory, combinatorics, special functions, and mathematical physics. For example, see [2, 5, 6, 7, 9].

The Bailey Transform is a consequence of the terminating ${}_4\phi_3$ summation theorem. The Bailey Lemma is derived in [1] directly from the ${}_6\phi_5$ summation and the matrix inversion formulation [4, 13] of the Bailey Transform. We employ a similar method in the A_ℓ and C_ℓ cases by starting with a suitable, higher-dimensional, terminating ${}_6\phi_5$ summation theorem extracted from [19] and [14], respectively. The A_ℓ proofs appear in [20, 21], and the C_ℓ case is established in [17]. Many other consequences of the A_ℓ and C_ℓ generalizations of Bailey’s Transform and Lemma will appear in future papers. These include A_ℓ and C_ℓ q -Pfaff-Saalschütz summation theorems, q -Whipple transformations, connection coefficient results, and applications of iterating the A_ℓ or C_ℓ Bailey Lemma.

2. RESULTS

Throughout this article, let $\mathbf{i}, \mathbf{j}, \mathbf{N}$, and $\mathbf{y}$ be vectors of length ℓ with nonnegative integer components. Let q be a complex number such that $|q| < 1$. Define

$$(2.1a) \quad (\alpha)_\infty \equiv (\alpha; q)_\infty := \prod_{k \geq 0} (1 - \alpha q^k)$$

and, thus,

$$(2.1b) \quad (\alpha)_n \equiv (\alpha; q)_n := (\alpha)_\infty / (\alpha q^n)_\infty.$$

Define the Bailey transform matrices, M and M^* , as follows.

Definition (M and M^* for A_ℓ). Let $a, x_1, \dots, x_\ell$ be indeterminate. Suppose that none of the denominators in (2.2a–b) vanishes. Then let

$$(2.2a) \quad M(\mathbf{i}; \mathbf{j}; A_\ell) := \prod_{r,s=1}^{\ell} \left(q \frac{x_r}{x_s} q^{j_r - j_s} \right)_{i_r - j_r}^{-1} \prod_{k=1}^{\ell} \left(a q \frac{x_k}{x_\ell} \right)_{i_k + (j_1 + \dots + j_\ell)}^{-1};$$

and

(2.2b)

$$\begin{aligned} M^*(\mathbf{i}; \mathbf{j}; A_\ell) &:= \prod_{k=1}^{\ell} \left[1 - a \frac{x_k}{x_\ell} q^{i_k + (i_1 + \dots + i_\ell)} \right] \prod_{k=1}^{\ell} \left(a q \frac{x_k}{x_\ell} \right)_{j_k + (i_1 + \dots + i_\ell) - 1} \\ &\times \prod_{r,s=1}^{\ell} \left(q \frac{x_r}{x_s} q^{j_r - j_s} \right)_{i_r - j_r}^{-1} (-1)^{(i_1 + \dots + i_\ell) - (j_1 + \dots + j_\ell)} q^{\binom{(i_1 + \dots + i_\ell) - (j_1 + \dots + j_\ell)}{2}}. \end{aligned}$$

Definition (M and M^* for C_ℓ). Let $x_1, \dots, x_\ell$ be indeterminate. Suppose that none of the denominators in (2.3a–b) vanishes. Then let

$$(2.3a) \quad M(\mathbf{i}; \mathbf{j}; C_\ell) := \prod_{r,s=1}^{\ell} \left[\left(q \frac{x_r}{x_s} q^{j_r - j_s} \right)_{i_r - j_r}^{-1} (q x_r x_s q^{j_r + j_s})_{i_r - j_r}^{-1} \right];$$

and

(2.3b)

$$\begin{aligned} M^*(\mathbf{i}; \mathbf{j}; C_\ell) &:= \prod_{r,s=1}^{\ell} \left[\left(q \frac{x_r}{x_s} q^{j_r - j_s} \right)_{i_r - j_r}^{-1} (x_r x_s q^{j_r + j_s})_{i_r - j_r}^{-1} \right] \prod_{1 \leq r < s \leq \ell} \left[\frac{1 - x_r x_s q^{j_r + j_s}}{1 - x_r x_s q^{i_r + i_s}} \right] \\ &\times (-1)^{(i_1 + \dots + i_\ell) - (j_1 + \dots + j_\ell)} q^{\binom{(i_1 + \dots + i_\ell) - (j_1 + \dots + j_\ell)}{2}}. \end{aligned}$$

As in the classical case [1], we have the following theorem.

Theorem (Bailey Transform for A_ℓ and C_ℓ). Let $G = A_\ell$ or C_ℓ . Let M and M^* be defined as in (2.2) and (2.3), with rows and columns ordered lexicographically. Then M and M^* are inverse, infinite, lower-triangular matrices. That is,

$$(2.4) \quad \prod_{k=1}^{\ell} \delta(i_k, j_k) = \sum_{\substack{j_k \leq i_k \leq i_k \\ k=1,2,\dots,\ell}} M(\mathbf{i}; \mathbf{y}; G) M^*(\mathbf{y}; \mathbf{j}; G),$$

where $\delta(r, s) = 1$ if $r = s$, and 0 otherwise.

Equations (2.2) and (2.3) motivate the definition of the A_ℓ and C_ℓ Bailey pair.

Definition (G -Bailey Pair). Let $G = A_\ell$ or C_ℓ . Let $N_k \geq 0$ be integers for $k = 1, 2, \dots, \ell$. Let $A = \{A_{(\mathbf{y}; G)}\}$ and $B = \{B_{(\mathbf{y}; G)}\}$ be sequences. Let M and M^* be as above. Then we say that A and B form a G -Bailey Pair if

$$(2.5) \quad B_{(\mathbf{N}; G)} = \sum_{\substack{0 \leq y_k \leq N_k \\ k=1,2,\dots,\ell}} M(\mathbf{N}; \mathbf{y}; G) A_{(\mathbf{y}; G)}.$$

As a consequence of the Bailey transform, (2.4), and the definition of the G -Bailey pair, (2.5), we have the following result.

Corollary (Bailey Pair Inversion). *A and B satisfy equation (2.5) if and only if*

$$(2.6) \quad A_{(\mathbf{N}; G)} = \sum_{\substack{0 \leq y_k \leq N_k \\ k=1, 2, \dots, \ell}} M^*(\mathbf{N}; \mathbf{y}; G) B_{(\mathbf{y}; G)}.$$

Define the sequences $A' = \{A'_{(\mathbf{y}; A_\ell)}\}$ and $B' = \{B'_{(\mathbf{y}; A_\ell)}\}$ by

$$(2.7a) \quad \begin{aligned} A'_{(\mathbf{N}; A_\ell)} &:= \prod_{k=1}^{\ell} \left(\frac{aq}{\rho} \frac{x_k}{x_\ell} \right)^{-1}_{N_k} \prod_{k=1}^{\ell} \left(\sigma \frac{x_k}{x_\ell} \right)_{N_k} \\ &\times \frac{(\rho)_{N_1+\dots+N_\ell}}{(aq/\sigma)_{N_1+\dots+N_\ell}} (aq/\rho\sigma)^{N_1+\dots+N_\ell} A_{(\mathbf{N}; A_\ell)} \end{aligned}$$

and

$$(2.7b) \quad \begin{aligned} B'_{(\mathbf{N}; A_\ell)} &:= \sum_{\substack{0 \leq y_k \leq N_k \\ k=1, 2, \dots, \ell}} \left\{ \prod_{k=1}^{\ell} \left[\left(\sigma \frac{x_k}{x_\ell} \right)_{y_k} \left(\frac{aq}{\rho} \frac{x_k}{x_\ell} \right)^{-1}_{N_k} \right] \prod_{r,s=1}^{\ell} \left(q \frac{x_r}{x_s} q^{y_r-y_s} \right)^{-1}_{N_r-y_r} \right. \\ &\times \frac{(aq/\rho\sigma)_{(N_1+\dots+N_\ell)-(y_1+\dots+y_\ell)}}{(aq/\sigma)_{N_1+\dots+N_\ell}} \frac{(\rho)_{y_1+\dots+y_\ell}}{(aq/\rho\sigma)^{y_1+\dots+y_\ell}} B_{(\mathbf{y}; A_\ell)} \left. \right\} \end{aligned}$$

Define the sequences $A' = \{A'_{(\mathbf{y}; C_\ell)}\}$ and $B' = \{B'_{(\mathbf{y}; C_\ell)}\}$ by

$$(2.8a) \quad A'_{(\mathbf{N}; C_\ell)} := \prod_{k=1}^{\ell} \left[\frac{(\alpha x_k)_{N_k}}{(\beta x_k)_{N_k}} \frac{(qx_k \beta^{-1})_{N_k}}{(qx_k \alpha^{-1})_{N_k}} \right] \left(\frac{\beta}{\alpha} \right)^{N_1+\dots+N_\ell} A_{(\mathbf{N}; C_\ell)}$$

and

$$(2.8b) \quad \begin{aligned} B'_{(\mathbf{N}; C_\ell)} &:= \sum_{\substack{0 \leq y_k \leq N_k \\ k=1, 2, \dots, \ell}} \left\{ cr \prod_{k=1}^{\ell} \left[\frac{(\alpha x_k)_{y_k}}{(\beta x_k)_{N_k}} \frac{(qx_k \beta^{-1})_{y_k}}{(qx_k \alpha^{-1})_{N_k}} \right] \prod_{r,s=1}^{\ell} \left(q \frac{x_r}{x_s} q^{y_r-y_s} \right)^{-1}_{N_r-y_r} \right. \\ &\times \prod_{1 \leq r < s \leq \ell} \left[(qx_r x_s q^{y_r+y_s})^{-1}_{N_s-y_s} (qx_r x_s q^{N_s-y_s})^{-1}_{N_r-y_r} \right] \\ &\times \left(\frac{\beta}{\alpha} \right)_{(N_1+\dots+N_\ell)-(y_1+\dots+y_\ell)} \left(\frac{\beta}{\alpha} \right)^{y_1+\dots+y_\ell} B_{(\mathbf{y}; C_\ell)} \left. \right\} \end{aligned}$$

These definitions lead to our generalization of Bailey's lemma.

Theorem (The G -generalization of Bailey's Lemma). *Let $G = A_\ell$ or C_ℓ . Suppose $A = \{A_{(\mathbf{N}; G)}\}$ and $B = \{B_{(\mathbf{N}; G)}\}$ form a G -Bailey Pair. If $A' = \{A'_{(\mathbf{N}; G)}\}$ and $B' = \{B'_{(\mathbf{N}; G)}\}$ are as above, then A' and B' also form a G -Bailey Pair.*

3. SKETCHES OF PROOFS

Proof of (2.4). In each case, A_ℓ and C_ℓ , we begin with a terminating ${}_4\phi_3$ summation theorem. In the C_ℓ case, it is first necessary to specialize Gustafson's C_ℓ ${}_6\psi_6$ summation theorem, see [14], terminate it from below and then from above, and further specialize the resulting terminating ${}_6\phi_5$ to yield a terminating ${}_4\phi_3$. In both the A_ℓ and C_ℓ cases, the ${}_4\phi_3$ is modified by multiplying both the sum and product sides by some additional factors. Finally, that result is transformed term-by-term to yield the sum side of (2.4). $\square$

Proof of (2.6). Equation (2.6) follows directly from the definition, (2.5), and the termwise nature of the calculations in the proof of (2.4). $\square$

Proof of Bailey's Lemma. The definitions in (2.7) and (2.8) are substituted into (2.5). After an interchange of summation, the inner sum is seen to be a special case of the appropriate ${}_6\phi_5$. The ${}_6\phi_5$ is then summed, and the desired result follows. $\square$

Detailed proofs of the C_ℓ case will appear in [17], as will a discussion of the C_ℓ Bailey chain and a connection coefficient result associated with the C_ℓ Bailey Transform.

REFERENCES

1. A. K. Agarwal, G. Andrews, and D. Bressoud, *The Bailey lattice*, J. Indian Math. Soc. **51** (1987), 57–73.
2. G. E. Andrews, *q-Series: Their development and application in analysis, number theory, combinatorics, physics and computer algebra*, CBMS Regional Con. Ser. in Math., no. 66, Conf. Board Math. Sci., Washington, DC, 1986.
3. ———, *Problems and prospects for basic hypergeometric functions*, Theory and Applications of Special Functions (R. Askey, ed.), Academic Press, New York, 1975, pp. 191–224.
4. ———, *Connection coefficient problems and partitions*, Proc. Sympos. Pure Math., (D. Ray-Chaudhuri, ed.), vol. 34, Amer. Math. Soc., Providence, RI, 1979, pp. 1–24.
5. ———, *Multiple series Rogers-Ramanujan type identities*, Pacific J. Math. **114** (1984), 267–283.
6. G. E. Andrews, R. J. Baxter, and P. J. Forrester, *Eight-vertex SOS model and generalized Rogers-Ramanujan-type identities*, J. Statist. Phys. **35** (1984), 193–266.
7. G. E. Andrews, F. J. Dyson, and D. Hickerson, *Partitions and indefinite quadratic forms*, Invent. Math. **91** (1988), 391–407.
8. W. N. Bailey, *Identities of the Rogers-Ramanujan type*, Proc. London Math. Soc. (2) **50** (1949), 1–10.
9. R. J. Baxter, *Exactly solved models in statistical mechanics*, Academic Press, London and New York, 1982.
10. L. C. Biedenharn and J. D. Louck, *Angular momentum in quantum physics: Theory and applications*, Encyclopedia of Mathematics and Its Applications, (G.-C. Rota, ed.), vol. 8, Addison-Wesley, Reading, MA, 1981.
11. ———, *The Racah-Wigner algebra in quantum theory*, Encyclopedia of Mathematics and Its Applications, (G.-C. Rota, ed.), vol. 9, Addison-Wesley, Reading, MA, 1981.
12. G. Gasper and M. Rahman, *Basic hypergeometric series*, Encyclopedia of Mathematics and Its Applications, (G.-C. Rota, ed.), vol. 35, Cambridge University Press, Cambridge, 1990.
13. I. Gessel and D. Stanton, *Applications of q-Lagrange inversion to basic hypergeometric series*, Trans. Amer. Math. Soc. **277** (1983), 173–201.
14. R. A. Gustafson, *The Macdonald identities for affine root systems of classical type and hypergeometric series very well-poised on semi-simple Lie algebras*, Ramanujan International Symposium on Analysis (December 26th to 28th, 1987, Pune, India) (N. K. Thakare, ed.), 1989, pp. 187–224.

15. W. J. Holman, III, *Summation Theorems for hypergeometric series in $U(n)$* , SIAM J. Math. Anal. **11** (1980), 523–532.
16. W. J. Holman III, L. C. Biedenharn, and J. D. Louck, *On hypergeometric series well-poised in $SU(n)$* , SIAM J. Math. Anal. **7** (1976), 529–541.
17. G. M. Lilly and S. C. Milne, *The C_ℓ Bailey transform and Bailey lemma*, preprint.
18. S. C. Milne, *An elementary proof of the Macdonald identities for $A_\ell^{(1)}$* , Adv. in Math. **57** (1985), 34–70.
19. ———, *Basic hypergeometric series very well-poised in $U(n)$* , J. Math. Anal. Appl. **122** (1987), 223–256.
20. ———, *Balanced ${}_3\phi_2$ summation theorems for $U(n)$ basic hypergeometric series*, (in preparation).
21. ———, *A $U(n)$ generalization of Bailey’s lemma*, in preparation.
22. P. Paule, *Zwei neue Transformationen als elementare Anwendungen der q -Vandermonde Formel*, Ph.D. thesis, 1982, University of Vienna.
23. L. J. Rogers, *Second memoir on the expansion of certain infinite products*, Proc. London Math. Soc. **25** (1894), 318–343.
24. ———, *On two theorems of combinatory analysis and some allied identities*, Proc. London Math. Soc. (2) **16** (1917), 315–336.
25. L. J. Slater, *Generalized hypergeometric functions*, Cambridge University Press, London and New York, 1966.
26. F. J. W. Whipple, *On well-poised series, generalized hypergeometric series having parameters in pairs, each pair with the same sum*, Proc. London Math. Soc. (2) **24** (1924), 247–263.
27. ———, *Well-poised series and other generalized hypergeometric series*, Proc. London Math. Soc. (2) **25** (1926), 525–544.

DEPARTMENT OF MATHEMATICS, THE OHIO STATE UNIVERSITY, COLUMBUS, OHIO 43210
E-mail address: milne@function.mps.ohio-state.edu