

REFLECTION AND WEAKLY COLLECTIONWISE HAUSDORFF SPACES

Tim LaBerge

Avner Landver

October 20, 1992

ABSTRACT. We show that $\square(\theta)$ implies that there is a first countable $<\theta$ -collectionwise Hausdorff space that is not weakly θ -collectionwise Hausdorff. We also show that in the model obtained by Levy collapsing a weakly compact (supercompact) cardinal to ω_2 , first countable \aleph_1 -collectionwise Hausdorff spaces are weakly \aleph_2 -collectionwise Hausdorff (weakly collectionwise Hausdorff). In the last section we show that assuming E_θ^ω , a certain θ -family of integer valued functions exists and that in the model obtained by Levy collapsing a supercompact cardinal to ω_2 , these families do not exist.

§1 Introduction. Reflection is a central theme in modern set-theoretic topology. As Alan Dow points out in [Do], we often prove theorems when some type of reflection principle holds, and we build counter-examples when reflection fails. This paper contains both types of results, on questions related to the failure of collectionwise Hausdorff.

We say that a subset A of a topological space X can be *separated* if there is a collection $\{U_x : x \in A\}$ of disjoint open sets with $x \in U_x$ for every $x \in A$. A space X is *$<\theta$ -collectionwise Hausdorff* ($<\theta$ -cwH) if every closed discrete subset of size $<\theta$ can be separated. A space X is *$\leq\theta$ -collectionwise Hausdorff* ($\leq\theta$ -cwH) if every closed discrete subset of size $\leq\theta$ can be separated. X is *collectionwise Hausdorff* (cwH) if it is $\leq\theta$ -cwH for every cardinal θ .

In particular, we are concerned with Fleissner's questions:

Question 1. *Is there a ZFC example of a first countable $<\aleph_2$ -cwH space that is not $\leq\aleph_2$ -cwH?*

Question 2. *Is there a ZFC example of a first countable $<\aleph_2$ -cwH space that is not cwH?*

Fleissner asks for ZFC examples because he showed [F] that $E_{\omega_2}^\omega$ (a non-reflecting stationary subset of $\{\alpha < \omega_2 : cf(\alpha) = \omega\}$) can be used to construct an example of a first countable $<\aleph_2$ -cwH space that is not $\leq\aleph_2$ -cwH. Since the failure of $E_{\omega_2}^\omega$ is equiconsistent with the existence of a Mahlo cardinal ([De] and [HS]), large cardinals are required for a negative answer to Question 1. In fact, we'll see that

1990 *Mathematics Subject Classification.* Primary 54D15; Secondary 54A35 04A72.

Key words and phrases. reflection, weakly collectionwise Hausdorff, Levy forcing, Mitchell forcing, fans.

by a result of Todorćević [To2], at least a weakly compact cardinal will be needed to get a negative answer to Question 1. Shelah has shown [S] that in the model obtained by Levy collapsing a weakly compact cardinal to ω_2 , first countable $< \aleph_2$ -cwH spaces that are locally of size at most \aleph_1 are $\leq \aleph_2$ -cwH, and that in the model obtained by collapsing a supercompact cardinal, such spaces are cwH. For a more complete history of the general problem of reflecting the failure of cwH, see [W] and [FS]. We would like to thank Bill Fleissner for his valuable input and Gary Gruenhage for letting us use the handwritten notes [GT].

§2 Squares of Fans and First Countable Collectionwise Hausdorff Spaces. ■

Recently, Gruenhage and Tamano [GT] have discovered a connection between Fleissner's questions and the problem of determining the tightness of the squares of certain fans. Before we discuss this connection, we make some definitions.

The fan $F_{\theta, \omega}$ is the quotient space obtained by identifying the non-isolated points of θ -many copies of the convergent ω -sequence. To be precise, $F_{\theta, \omega} = \{*\} \cup (\theta \times \omega)$, topologized so that points of $\theta \times \omega$ are isolated and so that an open base at $*$ is the family of all

$$B_f = \{*\} \cup \{(\alpha, n) : n \geq f(\alpha)\} \quad (f \in {}^\theta \omega).$$

When working with the square of $F_{\theta, \omega}$, one can always replace two functions $f_1, f_2 \in {}^\theta \omega$ with $g = \max(f_1, f_2)$, so we use the family of all

$$V_g = B_g \times B_g \quad (g \in {}^\theta \omega)$$

as an open base for $(*, *)$ in $F_{\theta, \omega} \times F_{\theta, \omega}$.

We say that a set $S \subseteq F_{\theta, \omega} \times F_{\theta, \omega}$ is θ -good if $(*, *) \in \overline{S}$, but for all $T \in [S]^{< \theta}$, $(*, *) \notin \overline{T}$. Thus, "there is a θ -good set in $F_{\theta, \omega} \times F_{\theta, \omega}$ " means that the tightness of $F_{\theta, \omega} \times F_{\theta, \omega}$ is θ and that this tightness is actually attained.

Finally, we say that a set $H \in [\omega \times \omega]^{< \omega}$ is *closed downward* (c.d.w.) if whenever $(n, m) \in H$, then $n' \leq n$ and $m' \leq m$ implies $(n', m') \in H$.

The Gruenhage-Tamano result is the equivalence (1) \iff (2) in Theorem 1. We have added the combinatorial equivalence (3) and the superficially stronger topological characterization (4).

Theorem 1. *Let θ be an uncountable cardinal. TFAE*

- (1) *There exists a θ -good subset of $F_{\theta, \omega} \times F_{\theta, \omega}$.*
- (2) *There exists a space X that is first countable, $< \theta$ -cwH and not $\leq \theta$ -cwH.*
- (3) *There exists $\mathcal{H} = \{H_{\alpha\beta} : \alpha < \beta < \theta\} \subseteq [\omega \times \omega]^{< \omega}$ with each $H_{\alpha\beta}$ c.d.w. such that*
 - (a) *For every $A \in [\theta]^{< \theta}$ there is a function $f : \theta \rightarrow \omega$ such that for all $\alpha < \beta$ in A , $(f(\alpha), f(\beta)) \notin H_{\alpha\beta}$.*
 - (b) *For every $f : \theta \rightarrow \omega$ there are $\alpha < \beta < \theta$ such that $(f(\alpha), f(\beta)) \in H_{\alpha\beta}$.*
- (4) *There exists a space X that is first countable, zero-dimensional, $< \theta$ -cwH and not $\leq \theta$ -cwH.*

In order to put Theorem 1 in perspective, we need the definition of a difficult-to-deny combinatorial principle. For θ an uncountable regular cardinal, $\square(\theta)$ is the assertion that there is a family $\{C_\alpha : \alpha < \theta\}$ satisfying the following conditions:

- (i) $C_\alpha \subset \alpha$ is a club subset of α .
- (ii) If α is a limit point of C_β , then $C_\alpha = C_\beta \cap \alpha$.
- (iii) There is no club $C \subset \theta$ such that for every limit point α of C , $C_\alpha = C \cap \alpha$.

$\square(\theta)$ is true for every regular θ which is not weakly compact in L (see [To1]).

Todorcevic showed (see [To2] and [Be]) that $\square(\theta)$ can be used to construct a θ -good set. Also, combining Theorem 1 with Fleissner's construction of a first countable, $< \theta$ -cwH, not $\leq \theta$ -cwH space from E_θ^ω , we see that E_θ^ω can be used to construct a θ -good set.

Because the proof of Theorem 1 is essentially the same as the proof of Theorem 2, we only prove Theorem 2. Before the statement of the theorem we define a weakening of collectionwise Hausdorff introduced by Tall [Ta]. We say that a subset A of a space X is *weakly separated* if it has a subset of size $|A|$ that is separated. X is *weakly θ -cwH* if every closed discrete subset of size θ is weakly separated. X is *weakly cwH* if it is weakly θ -cwH for every cardinal θ .

Theorem 2. *Let $\kappa \leq \theta$ be uncountable cardinals. TFAE*

- (1) *There exists a space X that is first countable, $< \kappa$ -cwH and not weakly θ -cwH.*
- (2) *There exists $\mathcal{H} = \{H_{\alpha\beta} : \alpha < \beta < \theta\} \subset [\omega \times \omega]^{<\omega}$, with each $H_{\alpha\beta}$ c.d.w., such that*
 - (a) *For every $A \in [\theta]^{<\kappa}$ there is a function $f : \theta \rightarrow \omega$ such that for all $\alpha < \beta$ in A , $(f(\alpha), f(\beta)) \notin H_{\alpha\beta}$.*
 - (b) *For every $B \in [\theta]^\theta$ and every $f : \theta \rightarrow \omega$ there are $\alpha < \beta$ in B such that $(f(\alpha), f(\beta)) \in H_{\alpha\beta}$.*
- (3) *There exists a space X that is first countable, zero-dimensional, $< \kappa$ -cwH and not weakly θ -cwH.*

Proof. (1) \Rightarrow (2): Let $\theta \subset X$ be a subset which is not weakly separated. For every $\alpha \in \theta$ let $\{U_n(\alpha) : n \in \omega\}$ be a decreasing neighbourhood base at α . Let

$$S = \{(\langle \alpha, n \rangle, \langle \beta, m \rangle) : U_n(\alpha) \cap U_m(\beta) \neq \emptyset\}.$$

S is a subset of $F_{\theta,\omega} \times F_{\theta,\omega}$. For every $B \subset \theta$, let $S_B = \{(\langle \alpha, n \rangle, \langle \beta, m \rangle) \in S : \alpha, \beta \in B\}$. We claim that S satisfies the following two conditions

- (i) For every $A \in [\theta]^{<\kappa}$, $(*, *) \notin \overline{S}_A$.
- (ii) For every $B \in [\theta]^\theta$, $(*, *) \in \overline{S}_B$.

To show (i), let $A \in [\theta]^{<\kappa}$. Let $f : A \rightarrow \omega$ be such that $\{U_{f(\alpha)}(\alpha) : \alpha \in A\}$ is a separation of A in X . Let V_f be the open neighbourhood of $(*, *)$ in $F_{\theta,\omega} \times F_{\theta,\omega}$ that is determined by f . It is easy to check that $V_f \cap S_A = \emptyset$. For (ii), let $B \in [\theta]^\theta$. Let $f : \theta \rightarrow \omega$, and let V_f be as above. Now, $f \upharpoonright B$ is not a code for a separation of B in X , therefore there are $\alpha < \beta$ in B such that $U_{f(\alpha)}(\alpha) \cap U_{f(\beta)}(\beta) \neq \emptyset$. But then $(\langle \alpha, f(\alpha) \rangle, \langle \beta, f(\beta) \rangle) \in S_B \cap V_f$.

Using the fact that $(*, *)$ is not in the closure of any countable subset of S , it is not hard to verify the following facts.

Fact 1. *For every $\alpha < \theta$ there is $h(\alpha) \in \omega$ such that for every $n > h(\alpha)$ and for every $\beta < \theta$ the set $\{m \in \omega : (\langle \alpha, n \rangle, \langle \beta, m \rangle) \in S\}$ is finite.*

Fact 2. *For every $\beta < \theta$ there is $g(\beta) \in \omega$ such that for every $m > g(\beta)$ and for every $\alpha < \theta$ the set $\{n \in \omega : (\langle \alpha, n \rangle, \langle \beta, m \rangle) \in S\}$ is finite.*

Let $T = S \cap (B_h \times B_g)$. Clearly, T satisfies (i) and (ii). Now, for every $\alpha < \beta < \theta$ define

$$H'_{\alpha\beta} = \{(n, m) \in \omega \times \omega : (\langle \alpha, n \rangle, \langle \beta, m \rangle) \in T\}.$$

Let $H_{\alpha\beta}$ be the downward closure of $H'_{\alpha\beta}$. Let us show that the $H_{\alpha\beta}$'s are finite. It is enough to show that the $H'_{\alpha\beta}$'s are finite. Assume that this is false and let $\alpha < \beta < \theta$ be such that $H'_{\alpha\beta}$ is infinite. It follows that for every $k \in \omega$ there are $n, m > k$ such that $(n, m) \in H'_{\alpha\beta}$. This implies that $(*, *)$ is in the closure of $\{(\langle \alpha, n \rangle, \langle \beta, m \rangle) : (n, m) \in H'_{\alpha\beta}\}$, therefore $(*, *)$ is in the closure of a countable subset of T , a contradiction.

Finally, it is not hard to see that $\{H_{\alpha\beta} : \alpha < \beta < \theta\}$ satisfies (a) and (b) of (2) precisely because T satisfies (i) and (ii).

(2) \Rightarrow (3): As in [GT], first let $I = \{(\langle \alpha, n \rangle, \langle \beta, m \rangle) : \alpha < \beta < \theta \wedge (n, m) \in H_{\alpha\beta}\}$. Then let $X = I \cup \theta$. Points in I are isolated, and for every $\gamma \in \theta$ a base at γ is given by

$$U_k(\gamma) = \{\gamma\} \cup \{(\langle \alpha, n \rangle, \langle \beta, m \rangle) \in I : (\alpha = \gamma \wedge n \geq k) \vee (\beta = \gamma \wedge m \geq k)\} \quad (k \in \omega).$$

Clearly, X is first countable. To see that X is $< \kappa$ -cwH let $A \in [\theta]^{< \kappa}$. Let $f : \theta \rightarrow \omega$ be given by (2a). Then check that $\{U_{f(\alpha)}(\alpha) : \alpha \in A\}$ is a separation of A . To see that X is not weakly θ -cwH let $B \in [\theta]^\theta$ and let $f : \theta \rightarrow \omega$. By (2b), there are $\alpha < \beta$ in B such that $(f(\alpha), f(\beta)) \in H_{\alpha\beta}$. So,

$$(\langle \alpha, f(\alpha) \rangle, \langle \beta, f(\beta) \rangle) \in U_{f(\alpha)}(\alpha) \cap U_{f(\beta)}(\beta),$$

therefore f does not code a separation of B .

Finally, the finiteness of the $H_{\alpha\beta}$'s implies that each $U_k(\alpha)$ is clopen and therefore X is zero-dimensional.

(3) \Rightarrow (1): Trivial. \square

§3 Independence Results.

In this section, we first show that $\square(\theta)$ implies that there is a first countable, $< \theta$ -cwH space that is not weakly θ -cwH. We then demonstrate the consistency of “first countable $< \aleph_2$ -cwH spaces are weakly \aleph_2 -cwH (weakly cwH)”, assuming the consistency of “there is a weakly compact (supercompact) cardinal”.

Since $\square(\theta)$ is true unless θ is weakly compact in L , this gives that “first countable $< \aleph_2$ -cwH spaces are weakly \aleph_2 -cwH” is equiconsistent with “there is a weakly compact cardinal” and that the consistency of “first countable $< \aleph_2$ -cwH spaces are weakly cwH” implies the existence of an inner model with many measurable cardinals [KM].

Theorem 3. *Let θ be a regular uncountable cardinal and suppose that $\square(\theta)$ holds. Then there is a zero-dimensional, first countable, $< \theta$ -cwH space that is not weakly θ -cwH.*

Proof. In [Be], Todorcevic constructs, from the assumption of $\square(\theta)$, a function ρ_2 that maps pairs $\alpha < \beta < \theta$ into ω . In §4, we will study integer-valued families of functions, so to keep our notation consistent, we define for each $\beta < \theta$ a function $h_\beta : \beta \rightarrow \omega$ by

$$h_\beta(\alpha) = \rho_2(\alpha, \beta) \quad (\alpha < \beta).$$

By the properties of ρ_2 cited in [Be], the family $\{h_\beta : \beta < \theta\}$ satisfies

- (1) For all $\alpha < \beta < \theta$, there is an $n_{\alpha\beta} \in \omega$ such that for all $\xi < \alpha$, $h_\alpha(\xi) < h_\beta(\xi) + n_{\alpha\beta}$. (This is monotonicity [DW].)
- (2) For all $B \in [\theta]^\theta$ and all $n \in \omega$, there exist $\alpha < \beta$ in B such that $h_\beta(\alpha) > n$.

For every $\alpha < \beta < \theta$, define $H_{\alpha\beta} = \{(n, m) : n + m \leq h_\beta(\alpha)\}$. Let us show that $\mathcal{H} = \{H_{\alpha\beta} : \alpha < \beta < \theta\}$ satisfies (2) of Theorem 2 with $\kappa = \theta$.

For part (a) let $A \in [\theta]^{<\kappa}$. Let $\gamma < \theta$ be a bound for A and define $f : \theta \rightarrow \omega$ by

$$f(\alpha) = \begin{cases} h_\gamma(\alpha) + n_{\alpha\gamma} & \text{if } \alpha < \gamma, \\ 0 & \text{otherwise.} \end{cases}$$

Now let $\alpha < \beta$ in A , then $f(\alpha) + f(\beta) \geq h_\gamma(\alpha) + n_{\beta\gamma} > h_\beta(\alpha)$. Therefore $(f(\alpha), f(\beta)) \notin H_{\alpha\beta}$.

For part (b) let $B \in [\theta]^\theta$ and $f : \theta \rightarrow \omega$. Then there exist $B' \in [B]^\theta$ and $n \in \omega$ such that for every $\alpha \in B'$, $f(\alpha) = n$. But then there are $\alpha < \beta$ in B' such that $h_\beta(\alpha) > 2n$. Hence $f(\alpha) + f(\beta) = 2n < h_\beta(\alpha)$ and so $(f(\alpha), f(\beta)) \in H_{\alpha\beta}$. \square

Remark. Since the h_β 's also satisfy

(3) Whenever $A, B \in [\theta]^\theta$ and $n \in \omega$, there are $\alpha < \beta$ with $\alpha \in A$ and $\beta \in B$ such that $h_\beta(\alpha) > n$,

it follows that the space constructed above is badly non-normal.

The next two theorems will use the method of forcing + reflection. A good reference for this technique is [DTW1] and [DTW2]. We first prove a preservation lemma, whose proof is motivated by Lemma 3.13 in [Be].

Main Lemma. *Assume that \mathbb{P} is an ω_1 -closed partial order and that $\theta \geq \omega_2$ is a regular cardinal. Assume that $\mathcal{H} = \{H_{\alpha\beta} : \alpha < \beta < \theta\} \subset [\omega \times \omega]^{<\omega}$ are c.d.w. and the following condition from Theorem 2 holds:*

(b) *For every $B \in [\theta]^\theta$ and every $f : \theta \rightarrow \omega$ there are $\alpha < \beta$ in B such that $(f(\alpha), f(\beta)) \in H_{\alpha\beta}$.*

Assume that G is a \mathbb{P} -generic filter over V . Then in $V[G]$, for every $f : \theta \rightarrow \omega$ there are $\alpha < \beta < \theta$ such that $(f(\alpha), f(\beta)) \in H_{\alpha\beta}$.

Proof. Assume that the lemma is false. Then there is $p_0 \in \mathbb{P}$ and a \mathbb{P} -name \dot{g} such that

$$p_0 \Vdash \text{“}\dot{g} : \theta \rightarrow \omega \wedge (\forall \alpha < \beta < \theta) (\dot{g}(\alpha), \dot{g}(\beta)) \notin H_{\alpha\beta}\text{”}.$$

Let M be an elementary substructure of some $H(\lambda)$ (λ large enough) with $\theta, \mathcal{H}, \mathbb{P}, p_0, \dot{g}$, and $\Vdash_{\mathbb{P}}$ in M and $M \cap \theta = \beta < \theta$.

If possible, choose $p_1 \leq p_0$ and $\alpha_0 < \beta$ such that $p_1 \in M$ and $p_1 \Vdash \text{“}(\dot{g}(\alpha_0), 0) \in H_{\alpha_0\beta}\text{”}$. Note that this would imply that $p_1 \Vdash \text{“}\dot{g}(\beta) > 0\text{”}$.

Similarly construct $p_0 \geq p_1 \geq \dots \geq p_{n+1} \dots$ such that $p_{n+1} \in M$ and $\alpha_n < \beta$ with

$$p_{n+1} \Vdash \text{“}(\dot{g}(\alpha_n), n) \in H_{\alpha_n\beta}\text{”}.$$

In particular this would imply that $p_{n+1} \Vdash \text{“}\dot{g}(\beta) > n\text{”}$. This process must stop at a finite stage since otherwise one could choose q with $q \leq p_n$ for every $n \in \omega$ and get that for every $n \in \omega$, $q \Vdash \text{“}\dot{g}(\beta) > n\text{”}$, which is impossible. \square

So, there is $\bar{n} \in \omega$ such that for every $q \leq p_{\bar{n}}$ and for every $\alpha < \beta$, if $q \in M$, then

$$q \not\Vdash \text{“}(\dot{g}(\alpha), \bar{n}) \in H_{\alpha\beta}\text{”}.$$

Claim 1. For every $\alpha < \beta$, $p_{\bar{n}} \Vdash “(\dot{g}(\alpha), \bar{n}) \notin H_{\alpha\beta}”$.

Proof. Assume not. Then there is $q \leq p_{\bar{n}}$ and $\alpha < \beta$ such that $q \Vdash “(\dot{g}(\alpha), \bar{n}) \in H_{\alpha\beta}”$. Fix such $\alpha < \beta$ and the corresponding $H_{\alpha\beta}$. Then $H(\lambda) \models (\exists q \leq p_{\bar{n}}) q \Vdash “(\dot{g}(\alpha), \bar{n}) \in H_{\alpha\beta}”$. But $p_{\bar{n}}, \Vdash_{\mathbb{P}}, \dot{g}, \alpha$ and $H_{\alpha\beta}$ are all in M , therefore

$$M \models \exists q \leq p_{\bar{n}} (q \Vdash “(\dot{g}(\alpha), \bar{n}) \in H_{\alpha\beta}”),$$

which is a contradiction. \square

Let $B = \{\beta \in \theta : \forall \alpha < \beta (p_{\bar{n}} \Vdash “(\dot{g}(\alpha), \bar{n}) \notin H_{\alpha\beta}”)\}$. By elementarity of M , $|B| = \theta$.

Now, for every $\alpha \in \theta$ let $q_\alpha \leq p_{\bar{n}}$ and $m_\alpha \in \omega$ be such that $q_\alpha \Vdash “\dot{g}(\alpha) = m_\alpha”$. In V , define a function $f : \theta \rightarrow \omega$ by:

$$f(\alpha) = m_\alpha + \bar{n} \quad (\alpha \in \theta).$$

The following claim will give us the desired contradiction.

Claim 2. For every $\beta \in B$ and every $\alpha < \beta$, $(f(\alpha), f(\beta)) \notin H_{\alpha\beta}$.

Proof. Fix $\beta \in B$ and $\alpha < \beta$. By definition, $f(\alpha) \geq m_\alpha$ and $f(\beta) \geq \bar{n}$, therefore it is enough to show that $(m_\alpha, \bar{n}) \notin H_{\alpha\beta}$. But $p_{\bar{n}} \Vdash “(\dot{g}(\alpha), \bar{n}) \notin H_{\alpha\beta}”$, and $q_\alpha \leq p_{\bar{n}}$, and $q_\alpha \Vdash “\dot{g}(\alpha) = m_\alpha”$. Therefore $q_\alpha \Vdash “(m_\alpha, \bar{n}) \notin H_{\alpha\beta}”$, and this implies that indeed $(m_\alpha, \bar{n}) \notin H_{\alpha\beta}$. \square

Let κ be strongly inaccessible and let $\mathbb{P} = \mathbb{P}_\kappa$ be the Levy collapse of κ to ω_2 with countable conditions (for a definition and proofs of the facts below, see [K]). Let G be a \mathbb{P} -generic filter. We will use the following facts about \mathbb{P} : \mathbb{P} is an ω_1 -closed partial order. \mathbb{P} is κ -c.c. and $\kappa = \omega_2$ in $V[G]$. For every $\lambda < \kappa$, \mathbb{P} can be factored as $\mathbb{P} = \mathbb{P}_\lambda \times \mathbb{P}^\lambda$, $\mathbb{P} \cap V_\lambda = \mathbb{P}_\lambda$ and in $V[G_\lambda]$, \mathbb{P}^λ is forcing equivalent to \mathbb{P} (in particular it is ω_1 -closed). For a definition and discussion of weakly compact cardinals see [K] and [KM]. The following type of proof was invented by Baumgartner [Ba], see also [DTW2].

Theorem 4. Assume that κ is a weakly compact cardinal. Let \mathbb{P} be the Levy collapse of κ to ω_2 with countable conditions and let G be a \mathbb{P} -generic filter over V . Then in $V[G]$, every first countable $< \aleph_2$ -cwH space is weakly \aleph_2 -cwH.

Proof. By way of contradiction, suppose that in $V[G]$ there is a first countable $< \aleph_2$ -cwH space that is not weakly \aleph_2 -cwH. In $V[G]$, $\omega_2 = \kappa$, therefore there is $\mathcal{H} = \{H_{\alpha\beta} : \alpha < \beta < \kappa\}$ that satisfies (2) of Theorem 2 with $\kappa = \theta$. This fact must be forced and the forcing statement is Π_1^1 over (V_κ, \in) (with few extra parameters). Using the facts that κ is weakly compact (Π_1^1 -reflection) and that \mathbb{P} is κ -c.c., we find an inaccessible $\lambda < \kappa$ such that in $V[G_\lambda]$, $\{H_{\alpha\beta} : \alpha < \beta < \lambda\}$ satisfies (2) of Theorem 2 (with λ playing the role of both κ and θ).

Now, $V[G_\lambda] \models \mathbb{P}^\lambda$ is ω_1 -closed. Therefore, by the Main Lemma, for every function $f : \lambda \rightarrow \omega$ in $V[G]$, there are $\alpha < \beta < \lambda$ such that $(f(\alpha), f(\beta)) \in H_{\alpha\beta}$. But $\lambda < \kappa$, therefore this contradicts the fact that (a) of part (2) of Theorem 2 holds for \mathcal{H} in $V[G]$. \square

For a definition and discussion of supercompact cardinals see [KM], [J], and [DTW1].

Theorem 5. *Assume that κ is a supercompact cardinal and that the GCH holds above κ . Let \mathbb{P} be the Levy collapse of κ to ω_2 with countable conditions and let G be a \mathbb{P} -generic filter over V . Then in $V[G]$ every first countable $<\aleph_2$ -cwH space is weakly cwH.*

Proof. We have to show that in $V[G]$, if $\theta \geq \omega_2$ is a cardinal and X is a first countable, $<\aleph_2$ -cwH space, then X is weakly θ -cwH. It is a consequence of a result of Watson [W] that if θ is a singular strong limit cardinal, and X is first countable and weakly $<\theta$ -cwH, then X is weakly θ -cwH. Therefore, we may assume that θ is regular. Since $\kappa = \omega_2$ in $V[G]$, we have $\theta \geq \kappa$. The plan is to show that (2) of Theorem 2, with $\kappa = \omega_2$, fails in $V[G]$.

Let $j : V \rightarrow M$ be an elementary embedding such that

- (i) $j(\alpha) = \alpha$, for all $\alpha < \kappa$.
- (ii) $j(\kappa) > \theta$.
- (iii) ${}^\theta M \cap V \subset M$.

Now, $\mathbb{P} \subset V_\kappa$ and $j \upharpoonright V_\kappa$ is the identity map, therefore $\mathbb{P} \subset M$ and by (iii), $\mathbb{P} \in M$. So, $M[G]$ makes sense. In M (and also in V), $j(\mathbb{P})$ is the Levy collapse of $j(\kappa)$ to ω_2 . In the usual way $j(\mathbb{P})$ can be factored, so $j(\mathbb{P}) = j(\mathbb{P})_\kappa \times j(\mathbb{P})^\kappa$. But $j(\mathbb{P})_\kappa = \mathbb{P}$, so $j(\mathbb{P}) = \mathbb{P} \times j(\mathbb{P})^\kappa$. Moreover,

$$M[G] \models j(\mathbb{P})^\kappa \text{ is an } \omega_1\text{-closed partial order.}$$

Let I be a $j(\mathbb{P})^\kappa$ -generic filter over $V[G]$. I is also $j(\mathbb{P})^\kappa$ -generic over $M[G]$, and there is a filter $K \subset j(\mathbb{P})$ such that $M[G][I] = M[K]$ and $V[G][I] = V[K]$. Using the fact that for every $p \in \mathbb{P}$, $j(p) = p$, we can define, in $V[K]$,

$$j^* : V[G] \rightarrow M[K]$$

that extends j and show that it is an elementary embedding (see [KM] §25).

Let $\mathcal{H} \in V[G]$ with $\mathcal{H} = \{H_{\alpha\beta} : \alpha < \beta < \theta\} \subset [\omega \times \omega]^{<\omega}$ c.d.w. and assume that the following condition from Theorem 2 holds in $V[G]$:

- (b) For every $B \in [\theta]^\theta$ and every $f : \theta \rightarrow \omega$ there are $\alpha < \beta$ in B such that $(f(\alpha), f(\beta)) \in H_{\alpha\beta}$.

We will show that (a) from Theorem 2 (with $\kappa = \omega_2$) is false for \mathcal{H} in $V[G]$.

\mathcal{H} has a name of size θ in V that is a subset of M , therefore by (iii) this name is in M . Hence, $\mathcal{H} \in M[G]$ and it is clear that condition (b) holds for \mathcal{H} in $M[G]$ as well. Notice that formally \mathcal{H} is a function given by $\mathcal{H} = \{(\langle \alpha, \beta \rangle, H_{\alpha\beta}) : \alpha < \beta < \theta\}$. Let $\mathcal{H}' = j^*(\mathcal{H}) \in M[K]$. Then $\mathcal{H}' = \{(\langle \gamma, \delta \rangle, H'_{\gamma\delta}) : \gamma < \delta < j(\theta)\}$, where each $H'_{\gamma\delta}$ is a c.d.w. subset of $[\omega \times \omega]^{<\omega}$. Notice that for every $\alpha < \beta < \theta$, $H_{\alpha\beta} = j(H_{\alpha\beta}) = H'_{j(\alpha)j(\beta)}$. Let $A = j''\theta$, then $j''\mathcal{H} = j''\mathcal{H}' = \{(\langle j(\alpha), j(\beta) \rangle, H_{\alpha\beta}) : \alpha < \beta < \theta\} = \{(\langle \gamma, \delta \rangle, H'_{\gamma\delta}) : (\gamma < \delta) \wedge (\gamma, \delta \in A)\} \subset \mathcal{H}'$. The map $j \upharpoonright \mathcal{H}$ is a subset of M of size θ , but it is not in V , therefore we can not conclude that it is in M . Fortunately, ${}^\theta M[G] \cap V[G] \subset M[G]$ (see [J] p. 463), therefore $j \upharpoonright \mathcal{H} \in M[G] \subset M[K]$.

All that implies that the following two statements are equivalent in $M[K]$:

- (*) for every $f : j(\theta) \rightarrow \omega$ there are $\gamma < \delta$ in A such that $(f(\gamma), f(\delta)) \in H'_{\gamma\delta}$.
- (**) for every $f : \theta \rightarrow \omega$ there are $\alpha < \beta < \theta$ such that $(f(\alpha), f(\beta)) \in H_{\alpha\beta}$.

As was mentioned before, (b) holds for \mathcal{H} in $M[G]$ and $M[G] \models \text{“}j(\mathbb{P})^\kappa \text{ is an } \omega_1\text{-closed partial order”}$. Therefore, by the main lemma, (**) (and hence (*)) holds in $M[K]$.

We conclude that in $M[K]$, there is $A \in [j(\theta)]^{<j(\kappa)}$ such that for every $f : j(\theta) \rightarrow \omega$ there are $\gamma < \delta$ in A with $(f(\gamma), f(\delta)) \in j^*(\mathcal{H})_{\gamma\delta} (= H'_{\gamma\delta})$.

Therefore, by elementarity of j^* , in $V[G]$ there is an $A \in [\theta]^{\leq \aleph_1}$ such that for every $f : \theta \rightarrow \omega$ there are $\alpha < \beta$ in A with $(f(\alpha), f(\beta)) \in H_{\alpha\beta}$. \square

We can also prove Theorems 4 and 5 replacing the Levy collapse by the Mitchell collapse ([Mi] and [DJW]), and thus obtain the conclusions of these theorems in models where CH fails.

§4 Integer-Valued Functions.

As was mentioned before, Todorcevic proved that $\square(\theta)$ implies that there is a θ -good subset of $F_{\theta,\omega} \times F_{\theta,\omega}$ [To2]. In [DW], it is proved that all one needs is a non-extendible monotone θ -family of functions (which exists under $\square(\theta)$ but not in the model of Theorem 5).

Let g, h be functions with range a subset of ω . We say that g *weakly bounds* h if there is $n \in \omega$ such that for every $x \in \text{dom}(g) \cap \text{dom}(h)$

$$g(x) + n > h(x) \quad (\text{in short, } g + n > h).$$

We say that $\mathcal{H} = \{h_\beta : \beta < \theta\}$ is a θ -family of functions if the domain of h_β is β . The θ -family \mathcal{H} is *weakly bounded* if there is a g that weakly bounds each element of \mathcal{H} . We say that \mathcal{H} is an *initially weakly bounded (i.w.b.) θ -family* if for every $A \in [\theta]^{<\theta}$, $\{h_\beta : \beta \in A\}$ is weakly bounded. The family is *non-extendible* if $\{h_\beta : \beta < \theta\}$ is not weakly bounded.

We can now repeat the construction of a θ -good subset of $F_{\theta,\omega} \times F_{\theta,\omega}$ from a non-extendible i.w.b. θ -family $\{h_\beta : \beta < \theta\}$, using i.w.b. in place of monotonicity. To see this, define for every $\alpha < \beta < \theta$ the finite set $H_{\alpha\beta} = \{(n, m) \in \omega \times \omega : n + m \leq h_\beta(\alpha)\}$ and check that $\mathcal{H} = \{H_{\alpha\beta} : \alpha < \beta < \theta\}$ satisfies (3) of Theorem 1.

Question 3. *Does the existence of a θ -good subset of $F_{\theta,\omega} \times F_{\theta,\omega}$ imply the existence of a non-extendible i.w.b. θ -family of functions?*

Fleissner [F] used the combinatorial principle E_θ^ω to construct a locally countable, locally compact Moore space that is $<\theta$ -cwH but not $\leq\theta$ -cwH. In particular, by [GT] (see (1) \iff (2) of Theorem 1), E_θ^ω implies the existence of a θ -good subset of $F_{\theta,\omega} \times F_{\theta,\omega}$.

Theorem 6. *Let θ be an uncountable regular cardinal, and let $E \subset \{\alpha \in \theta : \text{cof}(\alpha) = \omega\}$ be a non-reflecting stationary set. Then there is a θ -family $\{h_\beta : \beta < \theta\}$ such that for every $A \subset \theta$, $\{h_\beta : \beta \in A\}$ is weakly bounded if and only if $A \cap E$ is non-stationary. (In particular $\{h_\beta : \beta < \theta\}$ is a non-extendible i.w.b. θ -family.)*

Proof. For every $\alpha \in E$, fix $\{a_\alpha(n) : n \in \omega\}$ an increasing ω -sequence unbounded in α . For every $\alpha < \beta < \theta$ define

$$h_\beta(\alpha) = \begin{cases} \min\{n \in \omega : a_\alpha(n) \neq a_\beta(n)\} & \text{if } (\alpha \in E) \wedge (\beta \in E), \\ 0 & \text{otherwise.} \end{cases}$$

It is enough to consider $A \subset E$. Let A be stationary and assume, by way of contradiction, that $g : \theta \rightarrow \omega$ is a weak bound for $\{h_\beta : \beta \in A\}$. There exists a stationary $B \subset A$ and $n \in \omega$ such that for every $\beta \in B$, $g + n > h_\beta$. Let $f = g + n$, then for every $\beta \in B$, $f > h_\beta$.

Now, there exists a stationary $S \subset B$, and $m \in \omega$ such that for every $\alpha, \beta \in S$, $f(\alpha) = m$ and $\langle a_\alpha(0), \dots, a_\alpha(m) \rangle = \langle a_\beta(0), \dots, a_\beta(m) \rangle$. Let $\alpha < \beta$ in S . Let $k = \min\{n \in \omega : a_\alpha(n) \neq a_\beta(n)\}$. Then $k > m$, therefore $f(\alpha) = m < k = h_\beta(\alpha)$, contradiction.

Next we prove, by induction on $\gamma < \theta$, that there exists $g_\gamma : \gamma \rightarrow \omega$ that weakly bounds $\{h_\beta : \beta < \gamma\}$.

The successor case is easy so let us assume that γ is a limit. let $C \subset \gamma$ be a club with $C \cap E \cap \gamma = \emptyset$. For every $\beta \in E \cap \gamma$ let

$$\begin{aligned}\gamma^+(\beta) &= \min(C \setminus \beta) \\ \gamma^-(\beta) &= \max(C \cap \beta).\end{aligned}$$

Notice that for every $\beta \in E \cap \gamma$ we have $\gamma^-(\beta) < \beta < \gamma^+(\beta) < \gamma$. Now define

$$g_\gamma(\beta) = \begin{cases} g_{\gamma^+(\beta)}(\beta) & \text{if } \beta \in E, \\ 1 & \text{otherwise.} \end{cases}$$

Let us show that g_γ weakly bounds $\{h_\beta : \beta < \gamma\}$. If $\beta \notin E$, then h_β is identically zero, so it suffices to consider $\beta \in E \cap \gamma$. Let $n = k + m + 1$, where k is the least integer such that $a_\beta(k) > \gamma^-(\beta)$ and m is satisfies $g_{\gamma^+(\beta)} + m > h_\beta$. Let us show that $g_\gamma + n > h_\beta$. Let $\alpha \in E \cap \beta$.

Case 1: $\gamma^-(\alpha) = \gamma^-(\beta)$. In this case, $\gamma^+(\alpha) = \gamma^+(\beta)$, therefore $g_\gamma(\alpha) = g_{\gamma^+(\alpha)}(\alpha) = g_{\gamma^+(\beta)}(\alpha)$. But $g_{\gamma^+(\beta)}(\alpha) + m > h_\beta(\alpha)$, therefore $g_\gamma(\alpha) + n > h_\beta(\alpha)$.

Case 2: $\alpha < \gamma^-(\beta)$. Then $k + 1 > h_\beta(\alpha)$ because $a_\beta(k) > \gamma^-(\beta)$, therefore $g_\gamma(\alpha) + n > h_\beta(\alpha)$.

Finally, let $A \subset E$ be non-stationary. Let us use $\{g_\gamma : \gamma < \theta\}$ to produce $g : \theta \rightarrow \omega$ that weakly bounds $\{h_\beta : \beta \in A\}$. Let $C \subset \theta$ be a club such that $C \cap A = \emptyset$. For every $\beta \in E$ let

$$\begin{aligned}\delta^+(\beta) &= \min(C \setminus (\beta + 1)) \\ \delta^-(\beta) &= \sup(C \cap \beta), \\ g(\beta) &= \begin{cases} g_{\delta^+(\beta)}(\beta) & \text{if } \beta \in E, \\ 1 & \text{otherwise.} \end{cases}\end{aligned}$$

Let $\beta \in A$. As before, let $n = k + m + 1$, where k is the least such that $a_\beta(k) > \delta^-(\beta)$ (notice, $\beta \in A$ implies that $\delta^-(\beta) < \beta$), and m is such that $g_{\delta^+(\beta)} + m > h_\beta$. Let us show that $g + n > h_\beta$. Let $\alpha \in E \cap \beta$.

Case 1: $\delta^-(\alpha) = \delta^-(\beta)$. So, $\delta^+(\alpha) = \delta^+(\beta)$, therefore $g(\alpha) = g_{\delta^+(\alpha)}(\alpha) = g_{\delta^+(\beta)}(\alpha)$. But $g_{\delta^+(\beta)}(\alpha) + m > h_\beta(\alpha)$, therefore $g(\alpha) + n > h_\beta(\alpha)$.

Case 2: $\delta^-(\alpha) < \delta^-(\beta)$. In this case $\delta^-(\beta) \geq \alpha$. But $a_\beta(k) > \delta^-(\beta)$, therefore $a_\beta(k) > \alpha$ and hence $a_\beta(k) \neq a_\alpha(k)$. So, $k + 1 > h_\beta(\alpha)$, therefore $g(\alpha) + n > h_\beta(\alpha)$.

□

We remark here that $\neg E_{\omega_2}^\omega$ is equiconsistent with a Mahlo cardinal ([D] and [HS]), while $\neg \square(\omega_2)$ is equiconsistent with a weakly compact cardinal ([Ma], [To1] and [Be]), (and therefore $\square(\omega_2)$ does not imply $E_{\omega_2}^\omega$). And so consistency-wise, a result from $\square(\omega_2)$ is better (more difficult to deny) than one from $E_{\omega_2}^\omega$. On the other hand, $E_{\omega_2}^\omega$ does not imply $\square(\omega_2)$ (PFA is consistent with $E_{\omega_2}^\omega$ but implies $\neg \square(\omega_2)$ [Be]).

Theorem 7. *Let κ be a supercompact cardinal, let \mathbb{P} be the Levy collapse of κ to ω_2 with countable conditions and let G be a \mathbb{P} -generic filter over V . Then in $V[G]$ the following holds: For every regular $\theta \geq \omega_2$, and every θ -family $\{h_\beta : \beta < \theta\}$, if for all $A \in [\theta]^{\aleph_1}$, $\{h_\beta : \beta \in A\}$ is weakly bounded, then for every stationary set $S \subset \theta$, there exists a stationary $T \subset S$ such that $\{h_\beta : \beta \in T\}$ is weakly bounded.*

The proof of Theorem 7 is very much like the proof of Theorem 5. The following lemma is the analog of the Main Lemma, the rest of the proof of the theorem is left to the reader.

Lemma. *Assume that \mathbb{P} is an ω_1 -closed partial order, and that $\theta \geq \omega_2$ is a regular cardinal. Assume that $\mathcal{H} = \{h_\beta : \beta < \theta\}$ is a θ -family, and $S \subset \theta$ is stationary such that for every stationary $T \subset S$ the family $\{h_\beta : \beta \in T\}$ is not weakly bounded. Assume that G is a \mathbb{P} -generic filter over V . Then in $V[G]$, $\{h_\beta : \beta \in S\}$ is not weakly bounded.*

Proof. We proceed as in our Main Lemma and assume the lemma is false. Let $p_0 \in \mathbb{P}$ and \dot{g} , a \mathbb{P} -name, be such that $p_0 \Vdash \dot{g} : \theta \rightarrow \omega$ weakly bounds $\{h_\beta : \beta \in S\}$.

Let M be an elementary substructure of $H(\lambda)$ for some large enough regular λ , with $S, \theta, \mathcal{H}, \mathbb{P}, p_0, \dot{g}$, and $\Vdash_{\mathbb{P}}$ all members of M and $M \cap \theta = \beta \in S$.

Construct $p_0 \geq p_1 \geq \dots \geq p_{n+1} \dots$ such that $p_{n+1} \in M$ and $\alpha_n < \beta$ with

$$p_{n+1} \Vdash \dot{g}(\alpha_n) + n \leq h_\beta(\alpha_n).$$

This process must stop at a finite stage since otherwise one could choose q with $q \leq p_n$ for every $n \in \omega$, and get that for every $n \in \omega$ there is $\alpha_n < \beta$ such that $q \Vdash \dot{g}(\alpha_n) + n \leq h_\beta(\alpha_n)$, which is impossible.

So, there is $\bar{n} \in \omega$ such that for every $q \leq p_{\bar{n}}$ and for every $\alpha < \beta$, if $q \in M$, then $q \not\Vdash \dot{g}(\alpha) + \bar{n} \leq h_\beta(\alpha)$. As in the Main Lemma, it follows that for every $\alpha < \beta$,

$$p_{\bar{n}} \Vdash \dot{g}(\alpha) + \bar{n} > h_\beta(\alpha).$$

Let $T = \{\beta \in S : (\forall \alpha < \beta) p_{\bar{n}} \Vdash \dot{g}(\alpha) + \bar{n} > h_\beta(\alpha)\}$. By elementarity of M , T is stationary.

Now, for every $\alpha \in \theta$, let $q_\alpha \leq p_{\bar{n}}$ and $m_\alpha \in \omega$ be such that $q_\alpha \Vdash \dot{g}(\alpha) = m_\alpha$. In V , define $f : \theta \rightarrow \omega$ by $f(\alpha) = m_\alpha + \bar{n}$. Finally, continue as in the Main Lemma to get a contradiction by showing that for every $\beta \in T$ and every $\alpha < \beta$, $h_\beta(\alpha) < f(\alpha)$. In particular, f weakly bounds $\{h_\beta : \beta \in T\}$. \square

By Theorem 6, in the model of Theorem 7, E_θ^ω fails for every regular $\theta \geq \omega_2$, (this is a result of Shelah). As was the case for Theorems 4 and 5, Theorem 7 can also be proved for the Mitchell collapse instead of the Levy collapse. We finish with a question.

Question 4. *Is it consistent (relative to a large cardinal) that every i.w.b. ω_2 -family extends?*

References

- [Ba] J. Baumgartner, *A new class of order types*, Annals Math. Logic **9** (1976), no. 3, 187–222.
- [Be] M. Bekkali, *Topics in Set Theory*, Springer-Verlag, 1991.
- [De] K.J. Devlin, *Aspects of Constructibility*, Lecture Notes in Mathematics, 354, Springer-Verlag, 1973.

- [DJW] A. Dow, I. Juhasz, and W. Weiss, *Integer-valued functions and increasing unions of first countable spaces*, Israel Jour. Math. **67** (1989), no. 1, 181–192.
- [Do] A. Dow, *Set-theory in topology*, Preprint (1992).
- [DTW1] A. Dow, F. Tall, and W. Weiss, *New proofs of the consistency of the normal Moore space conjecture I*, Top. Appl. **37** (1990), 33–51.
- [DTW2] A. Dow, F. Tall, and W. Weiss, *New proofs of the consistency of the normal Moore space conjecture II*, Top. Appl. **37** (1990), 115–129.
- [DW] A. Dow and S. Watson, *A subcategory of TOP*, Preprint.
- [GT] G. Gruenhage and Tamano, *handwritten notes*.
- [F] W. G. Fleissner, *On λ -collectionwise Hausdorff spaces*, Top. Proc. **2** (1977), 445–456.
- [FS] W. G. Fleissner and S. Shelah, *Collectionwise Hausdorff: Incompactness at Singulars*, Top. Appl. **31** (1989), 101–107.
- [HS] L. Harrington and S. Shelah, *Some Exact Equiconsistency Results in Set Theory*, Notre Dame J. Formal Logic **26** (1985), no. 2, 178–188.
- [J] T. Jech, *Set Theory*, Academic Press, 1978.
- [KM] A. Kanamori and M. Magidor, *The development of large cardinals axioms in set theory*, Higher Set Theory, Lecture Notes in Mathematics, 699, Springer-Verlag, 1978, pp. 99–275.
- [K] K. Kunen, *Set Theory*, North-Holland, 1980.
- [Ma] M. Magidor, *Reflecting Stationary Sets*, J. Symbolic Logic **47** (1982), no. 4, 755–771.
- [Mi] W. Mitchell, *Aronszajn Trees and the Independence of the Transfer Property*, Annals Math. Logic **5** (1972), 21–46.
- [S] S. Shelah, *Remarks on λ -collectionwise hausdorff spaces*, Top. Proc. **2** (1977), 583–592.
- [Ta] F.D. Tall, *Weakly collectionwise Hausdorff spaces*, Top. Proc. **1** (1976), 295–304.
- [To1] S. Todorcevic, *Partitioning pairs of countable ordinals*, Acta Math. **159** (1987), 261–294.
- [To2] S. Todorcevic, *My new fan*, handwritten notes (1989).
- [W] S. Watson, *Comments on Separation*, Top. Proc. **14** (1989), no. 2, 315–372.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF KANSAS, LAWRENCE, KS 66045
E-mail address: LaBerge@kuhub.cc.ukans.edu Landver@kuhub.cc.ukans.edu