On necessary multiplier conditions for Laguerre expansions II

George $Gasper^1$ and $Walter Trebels^2$

Dedicated to Dick Askey and Frank Olver

(March 5, 1992 version)

Abstract. The necessary multiplier conditions for Laguerre expansions derived in Gasper and Trebels [3] are supplemented and modified. This allows us to place Markett's Cohen type inequality [6] (up to the log-case) in the general framework of necessary conditions.

Key words. Laguerre polynomials, necessary multiplier conditions, Cohen type inequalities, fractional differences, weighted Lebesgue spaces

AMS(MOS) subject classifications. 33C65, 42A45, 42C10

1 Introduction

The purpose of this sequel to [3] is to obtain a better insight into the structure of Laguerre multipliers on L^p spaces from the point of view of necessary conditions. We recall that in [3] there occurs the annoying phenomenon that, e.g., the optimal necessary conditions in the case p=1 do not give the "right" unboundedness behavior of the Cesàro means. By slightly modifying these conditions we can not only remedy this defect but can also derive Markett's Cohen type inequality [6] (up to the log-case) as an immediate consequence.

For the convenience of the reader we briefly repeat the notation; we consider the Lebesgue spaces

$$L_{w(\gamma)}^p = \{ f : \| f \|_{L_{w(\gamma)}^p} = \left(\int_0^\infty |f(x)e^{-x/2}|^p x^{\gamma} \, dx \right)^{1/p} < \infty \} , \quad 1 \le p < \infty,$$

denote the classical Laguerre polynomials by $L_n^{\alpha}(x)$, $\alpha > -1$, $n \in \mathbb{N}_0$ (see Szegö [8, p. 100]), and set

$$R_n^{\alpha}(x) = L_n^{\alpha}(x)/L_n^{\alpha}(0), \quad L_n^{\alpha}(0) = A_n^{\alpha} = \binom{n+\alpha}{n} = \frac{\Gamma(n+\alpha+1)}{\Gamma(n+1)\Gamma(\alpha+1)}.$$

¹Department of Mathematics, Northwestern University, Evanston, IL 60208, USA. The work of this author was supported in part by the National Science Foundation under grant DMS-9103177.

²Fachbereich Mathematik, TH Darmstadt, D-6100 Darmstadt, Germany.

Associate to f its formal Laguerre series

$$f(x) \sim (\Gamma(\alpha+1))^{-1} \sum_{k=0}^{\infty} \hat{f}_{\alpha}(k) L_k^{\alpha}(x),$$

where the Fourier Laguerre coefficients of f are defined by

$$\hat{f}_{\alpha}(n) = \int_0^\infty f(x) R_n^{\alpha}(x) x^{\alpha} e^{-x} dx \tag{1}$$

(if the integrals exist). A sequence $m = \{m_k\}$ is called a (bounded) multiplier on $L^p_{w(\gamma)}$, notation $m \in M^p_{w(\gamma)}$, if

$$\| \sum_{k=0}^{\infty} m_k \hat{f}_{\alpha}(k) L_k^{\alpha} \|_{L_{w(\gamma)}^p} \leq C \| f \|_{L_{w(\gamma)}^p}$$

for all polynomials f; the smallest constant C for which this holds is called the multiplier norm $\parallel m \parallel_{M^p_{w(\gamma)}}$. The necessary conditions will be given in certain "smoothness" properties of the multiplier sequence in question. To this end we introduce a fractional difference operator of order δ by

$$\Delta^{\delta} m_k = \sum_{j=0}^{\infty} A_j^{-\delta - 1} m_{k+j}$$

(whenever the sum converges), the first order difference operator Δ_2 with increment 2 by

$$\Delta_2 m_k = m_k - m_{k+2},$$

and the notation

$$\Delta_2 \Delta^{\delta} m_k = \Delta^{\delta+1} m_k + \Delta^{\delta+1} m_{k+1}.$$

Generic positive constants that are independent of the functions (and sequences) will be denoted by C. Within the setting of the $L^p_{w(\gamma)}$ -spaces our main results now read (with 1/p + 1/q = 1):

Theorem 1.1 Let α , a > -1 and $\alpha + a > -1$. If $f \in L^p_{w(\gamma)}$, $1 \le p < 2$, then

$$\left(\sum_{k=0}^{\infty} |(k+1)^{(\gamma+1)/p-1/2} \Delta_2 \Delta^a \hat{f}_{\alpha}(k)|^q\right)^{1/q} \le C \|f\|_{L^p_{w(\gamma)}},\tag{2}$$

provided

$$\frac{\gamma+1}{p} \leq \frac{\alpha+a}{p} + 1 \qquad \qquad \textit{if} \ \ \alpha+a \leq 1/2,$$

$$\frac{\gamma+1}{p} \le \frac{\alpha+a}{2} + 1 + \frac{1}{2}\left(\frac{1}{p} - \frac{1}{2}\right)$$
 if $\alpha + a > 1/2$.

As in [3] (see there the proof of Lemma 2.3) we immediately obtain

Theorem 1.2 Let $m = \{m_k\} \in M^p_{w(\gamma)}, 1 \le p < 2$, and let α and a be as in Theorem 1.1. Then

$$\sup_{n} \left(\sum_{k=n}^{2n} |(k+1)^{(2\gamma+1)/p - (2\alpha+1)/2} \Delta_2 \Delta^a m_k|^q \frac{1}{k+1} \right)^{1/q} \le C \parallel m \parallel_{M_{w(\gamma)}^p}, \tag{3}$$

provided that in the case $\alpha + a \leq 1/2$ the condition

$$\frac{\alpha + a}{p} + 1 \ge \frac{\gamma + 1}{p} > \begin{cases} (\alpha + 1)/2 + 1/3p & \text{if } 1 \le p < 4/3\\ (\alpha + 1)/2 + 1/4 & \text{if } 4/3 \le p < 2 \end{cases}$$

holds, and in the case $\alpha + a > 1/2$ the condition

$$\frac{\alpha + a}{2} + 1 + \frac{1}{2} \left(\frac{1}{p} - \frac{1}{2} \right) \ge \frac{\gamma + 1}{p} > \begin{cases} (\alpha + 1)/2 + 1/3p & \text{if } 1 \le p < 4/3 \\ (\alpha + 1)/2 + 1/4 & \text{if } 4/3 \le p < 2. \end{cases}$$

In view of the results in [6], [3] and for an easy comparison we want to emphasize the cases $\gamma = \alpha$ and $\gamma = \alpha p/2$. Therefore, we state

Corollary 1 a) Let $m \in M_{w(\alpha)}^p$, $1 \le p < 2$, and let $\alpha > -1$ be such that $\max\{1/(3p), 1/4\} < (\alpha + 1)(1/p - 1/2)$. Then, with $\lambda := (2\alpha + 1)(1/p - 1/2)$,

$$\sup_{n} \left(\sum_{k=n}^{2n} |(k+1)^{\lambda} \Delta_2 \Delta^{\lambda-1} m_k|^q \frac{1}{k+1} \right)^{1/q} \le C \parallel m \parallel_{M_{w(\alpha)}^p}.$$

b) Let $m \in M^p_{w(\alpha p/2)}$, $1 \le p < 4/3$, and $(\alpha - 1)(1/p - 1/2) \ge -1/2$. Then

$$\sup_{n} \left(\sum_{k=n}^{2n} |(k+1)^{1/p-1/2} \Delta_2 m_k|^q \frac{1}{k+1} \right)^{1/q} \le C \parallel m \parallel_{M_{w(\alpha p/2)}}.$$

Remarks. 1) For polynomial $f(x) = \sum_{k=0}^{n} c_k L_k^{\alpha}(x)$ Theorem 1.1 yields, by taking only the (k=n)-term on the left hand side of (2),

$$|c_n|(n+1)^{(\gamma+1)/p-1/2} \le C \parallel f \parallel_{L^p_{w(\gamma)}}, \quad 1 \le p < 2$$

(under the restrictions on γ of Theorem 1.1). In particular, if we choose $\gamma = \alpha$, this comprises formula (1.13) in Markett [6] for his basic case $\beta = \alpha$. For $\gamma = \alpha p/2$, it even extends formula (1.14) in [6] to negative α 's as described in Corollary 1.3, b). The case 2 can be done by an application of a Nikolskii inequality, see [6].

2) Analogously, Cohen type inequalities follow from Theorem 1.2; in particular, Corollary 1.3 yields

Corollary 2 Let $m = \{m_k\}_{k=0}^n$ be a finite sequence, $1 \le p < 2$, and $\alpha > -1$.

a) If $m \in M_{w(\alpha)}^p$ then

$$(n+1)^{(2\alpha+2)(1/p-1/2)-1/2} |m_n| \le C \parallel m \parallel_{M^p_{w(\alpha)}}, \quad 1 \le p < \frac{4\alpha+4}{2\alpha+3},$$

provided $\max\{1/3p, 1/4\} < (\alpha + 1)(1/p - 1/2)$.

b) If $m \in M_{w(\alpha p/2)}^p$ and $(\alpha - 1)(1/p - 1/2) \ge -1/2$, then

$$(n+1)^{2/p-3/2}|m_n| \le C \parallel m \parallel_{M^p_{w(\alpha p/2)}}, \quad 1 \le p < 4/3.$$

With the exception of the crucial log-case, i.e. $p_0 = (4\alpha + 4)/(2\alpha + 3)$ or $p_0 = 4/3$, resp., Corollary 1.4 contains Markett's Theorem 1 in [6] and extends it to negative α 's. In particular we obtain for the Cesàro means of order $\delta \geq 0$, represented by its multiplier sequence $m_{k,n}^{\delta} = A_{n-k}^{\delta}/A_n^{\delta}$, the "right" unboundedness behavior (see [4])

$$\| \{ m_{k,n}^{\delta} \} \|_{M_{w(\alpha)}^p} \ge C(n+1)^{(2\alpha+2)(1/p-1/2)-1/2-\delta}, \quad 1 \le p < \frac{4\alpha+4}{2\alpha+3+2\delta}.$$

3) There arises the question, in how far the type of necessary conditions in [3] are comparable with the present ones. Let $\lambda > 1$. Since $\Delta_2 m_k = \Delta m_k + \Delta m_{k+1}$ we obviously have

$$\sup_{n} \left(\sum_{k=n}^{2n} |(k+1)^{(2\gamma+1)/p - (2\alpha+1)/2} \Delta_2 \Delta^{\lambda-1} m_k|^q \frac{1}{k+1} \right)^{1/q} \tag{4}$$

$$\leq C \sup_{n} \left(\sum_{k=n}^{2n} |(k+1)^{(2\gamma+1)/p-(2\alpha+1)/2} \Delta^{\lambda} m_{k}|^{q} \frac{1}{k+1} \right)^{1/q}.$$

In general, a converse cannot hold as can be seen by the following example: choose $\gamma = \alpha$, $\lambda = (2\alpha + 1)(1/p - 1/2)$ and $m_k = (-1)^k k^{-\varepsilon}$, $0 < \varepsilon < 1$. Then

$$\sup_{n} \left(\sum_{k=n}^{2n} |(k+1)\Delta m_{k}|^{q} \frac{1}{k+1} \right)^{1/q} = \infty$$

and hence by the embedding properties of the wbv-spaces, see [2], the right hand side of (4) cannot be finite for all $\lambda > 1$. But since $\Delta_2 \Delta^{\lambda-1} m_k = \Delta^{\lambda-1} \Delta_2 m_k \sim (k+1)^{-\varepsilon-\lambda}$, the left hand side of (4) is finite for all $\lambda > 1$.

Theorem 1.1 will be proved in Section 2 by interpolating between (L^1, l^{∞}) and (L^2, l^2) estimates. The $a \neq 0$ case is an easy consequence of the case a = 0 when

one uses the basic formula (see formula (3) in [3] and Remark 3 preceding Section 3 there)

$$\Delta^{a} R_{k}^{\alpha}(x) = \frac{\Gamma(\alpha+1)}{\Gamma(\alpha+a+1)} x^{a} R_{k}^{\alpha+a}(x), \quad x > 0, \ a > -1 - \min\{\alpha, \alpha/2 - 1/4\}, \quad (5)$$

where in the case $a > -(2\alpha + 1)/4$ the series for the fractional difference converges absolutely. In Section 3, a necessary (L^1, l^1) -estimate is derived and it is compared with a corresponding sufficient (l^1, L^1) -estimate.

2 Proof of Theorem 1.1

Let us first handle the (L^2, l^2) -estimate. Since

$$\Delta_2 \Delta^a \hat{f}_{\alpha}(k) = \Delta^{1+a} \hat{f}_{\alpha}(k) + \Delta^{1+a} \hat{f}_{\alpha}(k+1)$$

it follows from the Parseval formula preceding Corollary 2.5 in [3] that

$$\left(\sum_{k=0}^{\infty} |\sqrt{A_k^{\alpha+1+a}} \Delta_2 \Delta^a \hat{f}_{\alpha}(k)|^2\right)^{1/2} \le C \left(\int_0^{\infty} |f(t)e^{-t/2}t^{(\alpha+1+a)/2}|^2 dt\right)^{1/2}.$$
 (6)

Concerning the (L^1, l^{∞}) -estimate we first restrict ourselves to the case a = 0. Define $\mu \in \mathbf{R}$ by

$$2\left(\frac{1}{p} - \frac{1}{2}\right)\mu = \frac{\gamma}{p} - \frac{\alpha + 1}{2};$$

with the notation $\mathcal{L}_k^{\alpha}(t) = (A_k^{\alpha}/\Gamma(\alpha+1))^{1/2} R_k^{\alpha}(t) e^{-t/2} t^{\alpha/2}$ it follows that

$$\begin{split} |\Delta_2 \hat{f}_\alpha(k)| &= C |\int_0^\infty f(t) \{ \mathcal{L}_k^\alpha(t) / \sqrt{A_k^\alpha} - \mathcal{L}_{k+2}^\alpha(t) / \sqrt{A_{k+2}^\alpha} \} e^{-t/2} t^{\alpha/2} \, dt | \\ &\leq C (k+1)^{-1-\alpha/2} \int_0^\infty |f(t)| |t^{-\mu-1/2} \mathcal{L}_k^\alpha(t)| e^{-t/2} t^{(\alpha+1)/2+\mu} \, dt \\ &+ C (k+1)^{-\alpha/2} \int_0^\infty |f(t)| |t^{-\mu-1/2} \{ \mathcal{L}_k^\alpha(t) - \mathcal{L}_{k+2}^\alpha(t) \} |e^{-t/2} t^{(\alpha+1)/2+\mu} \, dt = I + II. \end{split}$$

We distinguish the two cases $\alpha \leq 1/2$ and $\alpha > 1/2$:

First consider the case $\alpha \leq 1/2$. By the asymptotic estimates for $\mathcal{L}_k^{\alpha}(t) - \mathcal{L}_{k+2}^{\alpha}(t)$ in Askey and Wainger [1, p.699], see formula (2.12) in [6], it follows for $\gamma \leq \alpha + p - 1$ that

$$\| t^{-\mu-1/2} \{ \mathcal{L}_k^{\alpha}(t) - \mathcal{L}_{k+2}^{\alpha}(t) \} \|_{\infty} \le C(k+1)^{-1-\mu}$$

so that

$$II \le C(k+1)^{-1-\mu-\alpha/2} \int_0^\infty |f(t)| e^{-t/2} t^{(\alpha+1)/2+\mu} dt, \quad \gamma \le \alpha + p - 1.$$
 (7)

By Lemma 1, 4th case, in [5]

$$\|t^{-\mu-1/2}\mathcal{L}_{k}^{\alpha}(t)\|_{\infty} < C(k+1)^{-\mu-5/6}$$

so that trivially

$$I \le C(k+1)^{-1-\mu-\alpha/2} \int_0^\infty |f(t)| e^{-t/2} t^{(\alpha+1)/2+\mu} dt, \quad \frac{\gamma+1}{p} \le \frac{\alpha+1}{2} - \frac{1}{3p} + \frac{2}{3}.$$

By Lemma 1, 5th case, in [5]

$$\| t^{-\mu-1/2} \mathcal{L}_k^{\alpha}(t) \|_{\infty} \le C(k+1)^{\mu+1/2}$$

so that

$$I \le C(k+1)^{\mu-1/2-\alpha/2} \int_0^\infty |f(t)| e^{-t/2} t^{(\alpha+1)/2+\mu} dt,$$

$$\leq C(k+1)^{-1-\mu-\alpha/2} \int_0^\infty |f(t)| e^{-t/2} t^{(\alpha+1)/2+\mu} dt, \quad \frac{\gamma+1}{p} > \frac{\alpha+1}{2} - \frac{1}{3p} + \frac{2}{3},$$

provided that $\mu - (\alpha + 1)/2 \le -1 - \mu - \alpha/2$ which is equivalent to $\mu \le -1/4$ or $\gamma \le 3p/4 - 1/2 + \alpha p/2$. But this is no further restriction since for $\alpha \le 1/2$ there holds $\alpha + p - 1 \le 3p/4 - 1/2 + \alpha p/2$. Summarizing, for $-1 < \alpha \le 1/2$, $\gamma \le \alpha + p - 1$ and $\mu = (\gamma/p - (\alpha + 1)/2)/2(1/p - 1/2)$ we have that

$$\sup_{k} |(k+1)^{1+\mu+\alpha/2} \Delta_2 \hat{f}_{\alpha}(k)| \le C \int_0^\infty |f(t)| e^{-t/2} t^{(\alpha+1)/2+\mu} dt.$$
 (8)

Now consider the case $\alpha > 1/2$. Then, by formula (2.12) in [6], (7) is obviously true when $(\gamma + 1)/p \le \alpha/2 + 1 + (1/p - 1/2)/2$. Again, the application of Lemma 1 in [5] requires $\gamma \le \alpha + p - 1$, which for $\alpha > 1/2$ is less restrictive than $(\gamma + 1)/p \le \alpha/2 + 1 + (1/p - 1/2)/2$. Its 4th case now leads to

$$I \le C(k+1)^{-11/6-\mu-\alpha/2} \int_0^\infty |f(t)| e^{-t/2} t^{(\alpha+1)/2+\mu} dt, \quad \frac{\gamma+1}{p} \le \frac{\alpha+1}{2} - \frac{1}{3p} + \frac{2}{3},$$

and its 5th case to

$$I \le C(k+1)^{\mu-1/2-\alpha/2} \int_0^\infty |f(t)| e^{-t/2} t^{(\alpha+1)/2+\mu} dt, \quad (\gamma+1)/p > \frac{\alpha+1}{2} - \frac{1}{3p} + \frac{2}{3}.$$

But $\mu - 1/2 - \alpha/2 \le -\mu - 1 - \alpha/2$ if $(\gamma + 1)/p \le \alpha/2 + 1 + (1/p - 1/2)/2$; so that, summarizing, (8) also holds under this restriction for $\alpha > 1/2$.

Now an application of the Stein and Weiss interpolation theorem (see [7]) with $Tf = \{Tf(k)\}$ and $Tf(k) = \sqrt{A_k^{\alpha+1}} \Delta_2 \hat{f}_{\alpha}(k)$ gives the assertion of Theorem 1.1 in the case a = 0.

If $a \neq 0$ then by (1), the definition of $\Delta_2 \Delta^a$, and by (5)

$$\Delta_2 \Delta^a \hat{f}_{\alpha}(k) = C\{\Delta \hat{f}_{\alpha+a}(k) + \Delta \hat{f}_{\alpha+a}(k+1)\} = C\Delta_2 \hat{f}_{\alpha+a}(k),$$

since already the condition $\gamma < \alpha + a + 1$ (which implies no new restriction) gives absolute convergence of the infinite sum and integral involved (see the formula following (9) in [3]) and Fubini's Theorem can be applied. Hence all the previous estimates remain valid when α is replaced by $\alpha + a$.

3 A variant for integrable functions

Theorem 1.1 gives a necessary condition for a sequence $\{f_k\}$ to generate with respect to L_k^{α} an $L_{w(\gamma)}^1$ -function. But this condition is hardly comparable with the following sufficient one which is a slight modification of Lemma 2.2 in [3].

Theorem 3.1 Let $\alpha > -1$ and $\delta > 2\gamma - \alpha + 1/2 \ge 0$. If $\{f_k\}$ is a bounded sequence with $\lim_{k\to\infty} f_k = 0$ and

$$\sum_{k=0}^{\infty} (k+1)^{\delta+\alpha-\gamma} |\Delta^{\delta+1} f_k| \le K_{\{f_k\}},$$

then there exists a function $f \in L^1_{w(\gamma)}$ with $\hat{f}_{\alpha}(k) = f_k$ for all $k \in \mathbb{N}_0$ and

$$|| f ||_{L^1_{w(\gamma)}} \le C K_{\{f_k\}}$$

for some constant C independent of the sequence $\{f_k\}$.

The proof follows along the lines of Lemma 2.2 in [3] since the norm of the Cesàro kernel

$$\chi_n^{\alpha,\delta}(x) = (A_n^{\delta} \Gamma(\alpha+1))^{-1} \sum_{k=0}^n A_{n-k}^{\delta} L_k^{\alpha}(x) = (A_n^{\delta} \Gamma(\alpha+1))^{-1} L_n^{\alpha+\delta+1}(x)$$

can be estimated with the aid of Lemma 1 in [5] by

$$\|\chi_k^{\alpha,\delta}\|_{L^1_{w(\gamma)}} \le C(k+1)^{\alpha-\gamma}, \quad \delta > 2\gamma - \alpha + 1/2$$

The variant of Theorem 1.1 in the case p = 1 is

Theorem 3.2 If $\alpha > -1$ and $\gamma > \max\{-1/3, \alpha/2 - 1/6\}$, then

$$\sum_{k=0}^{\infty} (k+1)^{\gamma-2/3} |\Delta^{2\gamma-\alpha+1/3} \hat{f}_{\alpha}(k)| \le C \| f \|_{L^{1}_{w(\gamma)}}.$$

A comparison of the sufficient condition and the necessary one nicely shows where the $L^1_{w(\gamma)}$ -functions live; in particular we see that the "smoothness" gap (the difference of the orders of the difference operators) is just greater than 7/6. It is clear that Theorem 3.2 can be modified by using the Δ_2 -operator. Theorem 3.2 does not follow from the p=1 case of Lemma 2.1 in [3] since that estimate would lead to the divergent sum $\sum_{k=0}^{\infty} (k+1)^{-1} \parallel f \parallel_{L^1_{w(\gamma)}}$.

Proof

By formula (5) we have

$$\Delta^{2\gamma - \alpha + 1/3} \hat{f}_{\alpha}(k) = C \int_{0}^{\infty} f(t) R_{k}^{2\gamma + 1/3}(t) t^{2\gamma + 1/3} e^{-t} dt$$
$$= C(k+1)^{-\gamma - 1/6} \int_{0}^{\infty} f(t) \mathcal{L}_{k}^{2\gamma + 1/3}(t) t^{\gamma + 1/6} e^{-t/2} dt$$

and hence

$$\sum_{k=0}^{\infty} (k+1)^{\gamma-2/3} |\Delta^{2\gamma-\alpha+1/3} \hat{f}_{\alpha}(k)| \le C \int_{0}^{\infty} |f(t)| \sum_{k=0}^{\infty} (k+1)^{-5/6} |t^{1/6} \mathcal{L}_{k}^{2\gamma+1/3}(t)| t^{\gamma} e^{-t/2} dt$$

if the right hand side converges. To show this we discuss for $j \in \mathbf{Z}$

$$\sup_{2^{j} \le t \le 2^{j+1}} \sum_{k=0}^{\infty} (k+1)^{-5/6} |t^{1/6} \mathcal{L}_{k}^{2\gamma+1/3}(t)|$$

and prove that this quantity is uniformly bounded in j, whence the assertion.

First consider those $j \geq 0$ for which there exists a nonnegative integer n such that $0 \leq k \leq 2^n$ implies $3\nu/2 := 3(2k+2\gamma+4/3) \leq 2^j$ but such that this inequality fails to hold for $k \geq 2^{n+1}$; the latter assumption in particular implies that essentially $\nu/2 \geq 2^{j+1}$ for $k \geq 2^{n+4}$. Since $\|t^{1/6}\mathcal{L}_k^{2\gamma+1/3}(t)\|_{\infty} \leq C(k+1)^{-1/6}$ by Lemma 1 in [5], we obviously have

$$\sum_{k=0}^{\infty} (k+1)^{-5/6} |t^{1/6} \mathcal{L}_k^{2\gamma+1/3}(t)| \le \left(\sum_{k=0}^{2^n} + \sum_{k=2^{n+4}}^{\infty}\right) \dots + O(1).$$
 (9)

For $k = 0, ..., 2^n$ we can now apply the fourth case of formula (2.5) in [5] to obtain $|t^{1/6}\mathcal{L}_k^{2\gamma+1/3}(t)| \leq Ce^{-\mu 2^j}$ for some positive constant μ and the first sum on the right

hand side of (9) is bounded uniformly in j. In consequence of the choice of n the second case of formula (2.5) in [5] can be used for $k \ge 2^{n+4}$, giving

$$\sum_{k=2^{n+4}}^{\infty} (k+1)^{-5/6} |t^{1/6} \mathcal{L}_k^{2\gamma+1/3}(t)| \le Ct^{-1/12} \sum_{k=2^{n+4}}^{\infty} (k+1)^{-13/12} = O(1)$$

since $2^j \le t \le 2^{j+1}$ and j and n are comparable.

Now consider the remaining j's: We have to split up the sum $\sum_{k=0}^{\infty} \dots$ into two parts, one where k is such that $2^{j}\nu \geq 1$ (this contribution has just been seen to be uniformly bounded in j), the other where k is such that $2^{j}\nu \leq 1$. To deal with the last case choose again n to be the greatest integer such that $2^{n+2} + 4\gamma + 8/3 \leq 2^{-j}$; this time, n and -j are comparable and we obtain by the first case of (2.5) in [5]

$$\sum_{k=0}^{2^{n}} (k+1)^{-5/6} |t^{1/6} \mathcal{L}_{k}^{2\gamma+1/3}(t)| \le Ct^{\gamma+1/3} \sum_{k=0}^{2^{n}} (k+1)^{\gamma-2/3} = O(1)$$

if $2^{j} \le t \le 2^{j+1}$, $\gamma > -1/3$, which completes the proof.

References

- [1] R. Askey and S. Wainger, Mean convergence of expansions in Laguerre and Hermite series, Amer. J. Math., 87 (1965), pp. 695 708.
- [2] G. GASPER AND W. TREBELS, A characterization of localized Bessel potential spaces and applications to Jacobi and Hankel multipliers, Studia Math., 65 (1979), pp. 243 278.
- [3] G. Gasper and W. Trebels, Necessary multiplier conditions for Laguerre expansions, Canad. J. Math., 43 (1991), pp. 1228 1242.
- [4] E. GÖRLICH AND C. MARKETT, A convolution structure for Laguerre series, Indag. Math., 44 (1982), pp. 161 171.
- [5] C. Markett, Mean Cesàro summability of Laguerre expansions and norm estimates with shifted parameter, Anal. Math., 8 (1982), pp. 19 37.
- [6] C. MARKETT, Cohen type inequalities for Jacobi, Laguerre and Hermite expansions, SIAM J. Math. Anal., 14 (1983), pp. 819 833.
- [7] E.M. Stein and G. Weiss, Interpolation of operators with change of measures, Trans. Amer. Math. Soc., 87 (1958), pp. 159 172.
- [8] G. SZEGÖ, Orthogonal Polynomials, 4th ed., Amer. Math. Soc. Colloq. Publ. 23, Providence, R.I., 1975.