

# CONTRACTIVE PROJECTIONS IN NONATOMIC FUNCTION SPACES

BEATA RANDRIANANTOANINA

ABSTRACT. We prove that there is no 1-complemented subspaces of finite codimension in separable rearrangement-invariant function spaces.

We study contractive projections onto finite-codimensional subspaces of real nonatomic function spaces. In general such projections are not common. It is well known that only in Hilbert space there exists a contractive projection onto every subspace of fixed finite codimension (cf. [1]).

The study of contractive projections (and more general projections with minimal norm) is important in approximation theory (cf. the survey of Cheney and Price [3]).

It is known that there is no 1-complemented subspaces of finite-codimension in  $C[0, 1]$  (Wulbert [13]) and in  $L_1(\mu)$  if the measure  $\mu$  is nonatomic ([6, Corollary IV.1.15], [4]). De Figueiredo and Karlovitz [5] (1970) proved that if  $\mu$  is nonatomic then there is no 1-complemented hyperplanes in  $L_p(\Omega, \mu)$ ,  $1 < p < \infty$ ,  $p \neq 2$ , (cf. also [2]).

In this paper we prove that in rearrangement-invariant nonatomic function spaces not isometric to  $L_2$  there is no 1-complemented subspaces of any finite codimension.

We use the terminology and notation as in [9].

Our method of proof is surprisingly simple – it is based on the following observation:

**Proposition 1.** (cf. [7, 11]) *In a real Banach space  $X$  if  $P$  is a projection then  $\|I - P\| = 1$  (where  $I$  denotes identity operator) if and only if  $x^*(Px) \geq 0$  for all  $x \in X$  and  $x^* \in X^*$  norming for  $x$ .*

In [7, Theorem 4.3] (cf. [8, 10]) Kalton and the author proved the nonexistence of 1-complemented hyperplanes in a wide class of nonatomic function spaces. However the original theorem uses special technical frazeology so we state it below in the language of projections:

---

1991 *Mathematics Subject Classification.* 46B,46E.

**Theorem 2.** *Suppose  $X$  is a real order-continuous Köthe function space on  $(\Omega, \mu)$  and  $\mu$  is nonatomic. Then the hyperplane  $H$  in  $X$  is 1-complemented if and only if there exists a nonnegative measurable function  $w$  with  $\text{supp } w = B = \text{supp } f$ , where  $f \in H^\perp \subset X^*$ , so that for any  $x \in X$  with  $\text{supp } x \subset B$*

$$\|x\| = \left( \int |x|^2 w \, d\mu \right)^{1/2}.$$

Hence there is no 1-complemented hyperplanes in  $X$  unless  $L_2$  is isometric to a band in  $X$ . In particular there is no 1-complemented hyperplanes in separable r.i. spaces on  $[0, 1]$  ([7, Theorem 4.4]).

As a part of proof of Theorem 2 we proved the following fact which we state separately for the future use.

**Proposition 3.** ([10, Proposition 2.8]) *Let  $X$  be an order-continuous Köthe function space on  $(\Omega, \mu)$ , where  $\mu$  is nonatomic. Suppose that the set*

$$\Lambda = \left\{ \frac{x^*}{x} : x \in X, x^* \in X^* \text{ norming for } x \right\}$$

*is one-dimensional i.e.  $\Lambda \subset \{aw : a \in \mathbb{R}\}$  for some  $w \in L_0(\Omega, \mu)$ . Then  $X$  is isometric to  $L_2(wd\mu)$ .*

Now we are ready to prove our main result.

**Theorem 4.** *Suppose  $\mu$  is nonatomic and  $X$  is a separable r.i. space on  $([0, 1], \mu)$  not isometric to  $L_2$ . Then there is no 1-complemented subspaces of any finite codimension in  $X$ .*

For the proof we need the following measure-theoretic lemma.

**Lemma 5.** *Suppose  $\mu$  is nonatomic and suppose  $f_1, \dots, f_n, g_1, \dots, g_n \in L_1(\mu)$  are such that  $g_1, \dots, g_n$  are linearly independent and*

$$\sum_{i=1}^n \left( \int h f_i \, d\mu \right) \left( \int h g_i \, d\mu \right) \geq 0$$

*whenever  $|h| = 1$  a.e. Then  $\{f_j\}_{j=1}^n \subset \text{span } \{g_j\}_{j=1}^n$ .*

*Proof.* Consider the subset  $\Gamma$  of  $\mathbb{R}^{2n}$  of all  $2n$ -tuples  $(a_1, b_1, \dots, a_n, b_n)$  such that for some  $h \in L_\infty(\mu)$  with  $|h| = 1$  a.e. we have  $a_j = \int h f_j d\mu$ ,  $b_j = \int h g_j d\mu$ ,  $j = 1, \dots, n$ . Let  $H = \text{span } \Gamma \subset \mathbb{R}^{2n}$ .

We immediately see that  $\Gamma = -\Gamma$  and that by Liapunoff's theorem ([12])  $\Gamma$  is convex. Hence for every  $(s_1, t_1, \dots, s_n, t_n) \in H$  we have  $\sum_{j=1}^n s_j t_j \geq 0$ .

On the other hand consider  $V = \{v_k = e_{2k-1} - e_{2k} : k = 1, \dots, n\} \subset \mathbb{R}^{2n}$  where  $e_j$  denotes the natural basis of  $\mathbb{R}^{2n}$ . Clearly  $V \cap H = \{0\}$  so  $\dim H \leq n$ . Therefore there exist  $c_1, \dots, c_n \in \mathbb{R}$  such that  $s_i = \sum_{j=1}^n c_j t_j$  for  $i = 1, \dots, n$ , for every  $(s_1, t_1, \dots, s_n, t_n) \in H$ . In particular  $\int h f_i d\mu = \int h \left( \sum_{j=1}^n c_j g_j \right) d\mu$  for all  $h$  with  $|h| = 1$  a.e. and the lemma follows.  $\square$

*Proof of Theorem 4.* Suppose that  $F$  is a closed linear subspace of codimension  $n$  in  $X$  and let  $u_1, \dots, u_n$  be linearly independent functions in  $X$  such that  $\text{span}\{F, u_1, \dots, u_n\} = X$ . Denote  $P : X \rightarrow F$  a contractive projection onto  $F$  and consider  $Q = I - P$ . Then  $Q = \sum_{j=1}^n f_j \otimes u_j$  for some linearly independent  $f_1, \dots, f_n \in X^*$ .

By Proposition 1 for any  $x \in X$  and  $x^* \in X^*$  norming for  $x$   $x^*(Qx) \geq 0$ . Next for any  $h$  with  $|h| = 1$  a.e.  $hx^*$  is norming for  $hx$  if  $x^*$  is norming for  $x$ . Hence  $hx^*(Q(hx)) \geq 0$  i.e.

$$\sum_{j=1}^n \left( \int f_j h x d\mu \right) \left( \int u_j h x^* d\mu \right) \geq 0.$$

By Lemma 5  $f_1 x \in \text{span}\{u_j x^* : j = 1, \dots, n\}$ . So if  $B = \text{supp } f_1$  ( $\mu(B) > 0$ ) then  $\frac{x}{x^*} \Big|_B \in \text{span}\left\{ \frac{u_j}{f_1} : j = 1, \dots, n \right\}$ . By re-arrangement invariance of  $X$  for every measure preserving map  $\sigma : [0, 1] \rightarrow [0, 1]$   $x^* \circ \sigma$  is norming for  $x \circ \sigma$  and so

$$\frac{x \circ \sigma}{x^* \circ \sigma} \Big|_B \in \text{span}\left\{ \frac{u_j}{f_1} : j = 1, \dots, n \right\}$$

i.e. the set  $\left\{ \left[ \left( \frac{x}{x^*} \right) \circ \sigma \right] \Big|_B \mid \sigma : [0, 1] \rightarrow [0, 1] \text{ measure preserving} \right\}$  is finite dimensional which is impossible unless  $\frac{x}{x^*}$  is a constant. But then by Proposition 3 we conclude that  $X$  is isometric to  $L_2[0, 1]$  contrary to our assumption.  $\square$

*Remark .* Notice that in the proof of Theorem 4 we use re-arrangement invariance of  $X$  only in the final step to conclude that if the set  $\left\{ \frac{x^*}{x} : x \in X, x^* \in X^* \text{ norming for } x \right\}$  is

finite-dimensional then it is one-dimensional. Similar conclusion is true also in spaces of the form  $X(Y)$ ,  $X_1(X_2(\dots(X_m)\dots))$ , where  $X, Y, X_j$  are r.i. and Theorem 4 holds as stated also for those spaces.

**Acknowledgments** . I wish to express my gratitude to Professor Nigel Kalton for his interest in this work and many valuable discussions.

#### REFERENCES

- [1] D. AMIR. *Characterizations of inner product spaces*. Birkhäuser Verlag, Basel, Boston Stuttgart, 1986.
- [2] B. BEAUZAMY AND B. MAUREY. *Points minimaux et ensembles optimaux dans les espaces de Banach*. J. Functional Anal. **24** (1977) 107–139.
- [3] E. W. CHENEY AND K. H. PRICE. *Minimal projections*. In: A. Talbot, editor, *Approximation Theory, Proceedings of a symposium held at Lancaster, July 1969*, pages 261 – 289. Academic Press, 1970.
- [4] R. G. DOUGLAS. *Contractive projections on an  $L_1$  space*. Pacific J. Math. **15** (1965) 443–462.
- [5] D.G. DE FIGUEIREDO AND L. A. KARLOVITZ. *On the extention of contractions of normed spaces*. In: *Nonlinear Functional Analysis Proceedings of symposia in Pure Mathematics, Volume 18, Part 1*, pages 95–104, Providence, 1970. Amer. Math. Soc.
- [6] P. HARMAND, D. WERNER, AND W. WERNER. *M-ideals in Banach Spaces and Banach Algebras*. Lecture Notes in Math. 1547. Springer, Berlin-Heidelberg-New York, 1993.
- [7] N. J. KALTON AND B. RANDRIANANTOANINA. *Surjective isometries of rearrangement-invariant spaces*. to appear in Quart. J. Math. Oxford.
- [8] N. J. KALTON AND B. RANDRIANANTOANINA. *Isometries of rearrangement-invariant spaces*. C. R. Acad. Sc. Paris **316** (1993) 351–355.
- [9] J. LINDENSTRAUS AND L. TZAFRIRI. *Classical Banach spaces, Vol. 2, Function spaces*. Springer–Verlag, Berlin–Heidelberg–New York, 1979.
- [10] B. RANDRIANANTOANINA. *Isometries of function spaces, Ph.D. Dissertation*. University of Missouri–Columbia, 1993.
- [11] H. P. ROSENTHAL. *Contractively complemented subspaces of Banach spaces with reverse monotone (transfinite) bases*. Longhorn Notes, The University of Texas Functional Analysis Seminar (1984-5) 1–14.
- [12] W. RUDIN. *Functional Analysis*. McGraw–Hill, New York, second edition, 1991.
- [13] D. E. WULBERT. *Projections of norm 1 on  $C(X)$* . Notices Amer. Math. Soc. **15** (1968) 362.

DEPARTMENT OF MATHEMATICS, BOWLING GREEN STATE UNIVERSITY, BOWLING GREEN, OH 43403

*E-mail address:* `brandri@andy.bgsu.edu`