Q.H.I. spaces

V.Ferenczi

July 5, 2021

Abstract

A Banach space X is said to be Q.H.I. if every infinite dimensional quotient space of X is H.I.: that is, a space is Q.H.I. if the H.I. property is not only stable passing to subspaces, but also passing to quotients and to the dual. We show that Gowers-Maurey's space is Q.H.I.; then we provide an example of a reflexive H.I. space \mathcal{X} whose dual is not H.I., from which it follows that \mathcal{X} is not Q.H.I.:

1 Introduction

In the following, by *space (resp. subspace)*, we shall always mean infinite dimensional Banach space (resp. closed subspace). We recall the central definition of [GM] A Banach space is said to be *hereditarily indecomposable (or H.I.)* if it has no decomposable subspace. In others words, a space X is H.I. if for any $\epsilon > 0$, any subspaces Y and Z of X, there exist vectors $y \in Y, z \in Z$, such that ||y|| = ||z|| = 1 and $||y - z|| \le \epsilon$.

We now give some notation that is useful for the construction of Gowers-Maurey's space and similar spaces. Let c_{00} be the space of sequences of scalars all but finitely many of which are zero. Let e_1, e_2, \ldots be its unit vector basis. If $E \subset \mathbf{N}$, then we shall also use the letter E for the projection from c_{00} to c_{00} defined by $E(\sum_{i=1}^{\infty} a_i e_i) = \sum_{i \in E} a_i e_i$. If $E, F \subset \mathbf{N}$, then we write E < F to mean that $\sup E < \inf F$. An *interval* of integers is a subset of \mathbf{N} of the form $\{a, a + 1, \ldots, b\}$ for some $a, b \in \mathbf{N}$. For N in \mathbf{N}, E_N denotes the interval $\{1, \ldots, N\}$. The *range* of a vector x in c_{00} , written ran(x), is the smallest interval E such that Ex = x. We shall write x < y to mean ran(x) < ran(y); notice that this is only defined on c_{00} . If $x_1 < \cdots < x_n$ we shall say that x_1, \ldots, x_n are successive.

Let \mathcal{X} be the class of Banach sequence spaces such that $(e_i)_{i=1}^{\infty}$ is a normalized bimonotone basis. Notice that for $p \geq 1$, l_p is in \mathcal{X} . We denote by $B(l_p)$ the unit ball of $l_p \cap c_{00}$.

By a *block basis* in a space $X \in \mathcal{X}$ we mean a sequence x_1, x_2, \ldots of successive non-zero vectors in X (such a sequence must be a basic sequence)

and by a *block subspace* of a space $X \in \mathcal{X}$ we mean a subspace generated by a block basis.

Let f be the function $\log_2(x+1)$. If $X \in \mathcal{X}$, and all successive vectors x_1, \ldots, x_n in X satisfy the inequality $f(n)^{-1} \sum_{i=1}^n ||x_i|| \le ||\sum_{i=1}^n x_i||$, then we say that X satisfies an *f*-lower estimate. We denote by $\mathcal{X}(f)$ the set of Banach spaces in \mathcal{X} satisfying an *f*-lower estimate.

A function $h: [1, +\infty) \to [1, +\infty)$ belongs to the Schlumprecht class \mathcal{F} of functions if it satisfies the following five conditions:

(i) h(1) = 1 and h(x) < x for every x > 1;

(ii) h is strictly increasing and tends to infinity;

(iii) $\lim_{x\to\infty} x^{-q}h(x) = 0$ for every q > 0;

(iv) the function x/h(x) is concave and non-decreasing;

(v) $h(xy) \le h(x)h(y)$ for every $x, y \ge 1$.

We notice that f and \sqrt{f} belong to \mathcal{F} .

Given X in \mathcal{X} , given g in \mathcal{F} , a functional x^* in X^* is an (M,g) - formif $||x^*||^* \leq 1$ and $x^* = \sum_{j=1}^M x_j^*$ for some sequence $x_1^* < \cdots < x_M^*$ of successive functionals such that $||x_j^*||^* \leq g(M)^{-1}$ for each j. Notice that if $X \in \mathcal{X}(f)$, then every (M, f)-form is in the unit ball of X^* .

Let X be a Banach space. Let y be a subspace of X. We shall denote by Id_Y the identity map from Y to X. An operator from Y to X is said to be *finitely singular* if its restriction to some finite codimensional subspace is an isomorphism into. It is said to be *infinitely singular* if it is not finitely singular. An operator S from Y to X is said to be *strictly singular* if the restriction of S to a subspace is never an isomorphism into. This is equivalent to saying that for any $\epsilon > 0$, any Z, there exists z in Z such that $||S(z)|| \le \epsilon ||z||$.

By Proposition 2.c.4 of [LT], if S is strictly singular, then for every Z and every $\epsilon > 0$, there exists $Z' \subset Z$ such that $||S_{/Z'}|| \leq \epsilon$. If S is strictly singular and T is an operator, then TS and ST are strictly singular whenever they are defined. We denote by $\mathcal{S}(Y, X)$ the space of strictly singular operators from Y to X.

Let T be an operator from Y to X. By Proposition 2.c.10 of [LT], if T is of the form Id + S, where S is strictly singular, then it is finitely singular, that is, an isomorphism on some finite codimensional subspace. It is said to be an Id + S-isomorphism if it is of the form Id + S and an isomorphism on the whole subspace Y. If T is an Id + S-isomorphism, then so is $T^{-1} = Id - (T - Id)T^{-1}$. If T and U are Id + S-isomorphisms, then so is TU when it is defined. The subspaces Y and Z of X are said to be Id + S-isomorphic if there exists an Id + S isomorphism from Y onto Z.

We finally recall a classical notion: two Banach spaces X and X' are totally incomparable if no subspace of X is isomorphic to a subspace of X'.

2 Definition: Q.H.I. spaces

2.1 Definition

A Banach space X is quotient hereditarily indecomposable (or Q.H.I.) if every infinite dimensional quotient space of X is H.I..

Remark Any Q.H.I. space is H.I. as a quotient space of itself.

Proposition 1 Let X be a Banach space. Assume that for every infinite codimensional and infinite dimensional subspace Y, X/Y is H.I.. Then X is Q.H.I..

<u>Proof</u> It is enough to prove that X is H.I.. Assume X is not H.I.. Then X contains a direct sum $Y \oplus Z$. Let W be an infinite dimensional and infinite codimensional subspace of Y (for example the space generated by the even vectors of a basic sequence in Y). Then X/W contains a space isomorphic to the sum $Y/W \oplus Z$, so X/W is not H.I..

Proposition 2 If X is reflexive and Q.H.I., then so is X^* .

<u>Proof</u> A space X is Q.H.I. if and only if any subspace of a quotient of X is indecomposable. For a reflexive space, this property passes to the dual.

2.2 Notation

We shall often refer to the space defined by Gowers and Maurey in [GM], and to the techniques developped in their article. When we shall refer to a lemma of [GM], we shall write it with the letters GM.

By space (resp. subspace), we shall always mean infinite dimensional space (resp. subspace). Let X be a Banach space, Y be a subspace of X.

An operator on X is said to be *finitely singular* if its restriction to some finite codimensional subspace is an isomorphism into. It is said to be *infinitely singular* if it is not finitely singular. It is said to be *strictly singular* if all its restrictions are infinitely singular.

An operator from Y to X is said to be an Id + S-isomorphism if it is an isomorphism of the form Id + S, where S is strictly singular.

The subspace Y is said to be *quasi-maximal* if Y and any subspace W of X have Id + S-isomorphic subspaces.

We know from [F] that if X is H.I., then every subspace of X is quasimaximal; as an easy consequence, if X has a quasi-maximal H.I. subspace then X is H.I..

Two Banach spaces X and X' are totally incomparable if no subspace of X is isomorphic to a subspace of X'.

For $p \ge 1$, $B(l_p)$ denotes the unit ball of $l_p \cap c_{00}$. An interval of integers E is a subset of **N** of the form $\{m, m+1, \ldots, n\}$. The letter E will also

denote the projection defined on c_{00} by $E(\sum_{i=1}^{+\infty} \lambda_i e_i) = \sum_{i \in E} \lambda_i e_i$. The range of a vector x in c_{00} , denoted ran(x) is the smallest interval E such that Ex = x. We say that E and F are successive if max $E < \min F$. Two vectors in c_{00} are said to be successive if their ranges are successive.

Let \mathcal{X} be the set of Banach spaces with a bimonotone basis. Let f be the function $x \to \log_2(x+1)$. A space X in \mathcal{X} is said to satisfy an f-lower estimate if any successive vectors $x_1 < \cdots < x_n$ in X satisfy the inequality $\|\sum_{i=1}^n x_i\| \ge f(n)^{-1} \sum_{i=1}^n \|x_i\|$. Let \mathcal{X}_f be the set of Banach spaces satisfying an f-lower estimate. An (M, f)-form is a vector of X^* of the form $f(M)^{-1} \sum_{i=1}^M x_i^*$, where the x_i^* 's are successive vectors in the unit ball of X^* . If $X \in \mathcal{X}_f$, then every (M, f)-form is in the unit ball of X^* .

3 Successive vectors in a quotient space

In this section, we want to define a notion of successive vectors and of l_1^{n+} -vectors in a quotient space of a reflexive space in \mathcal{X}_f . As a quotient space needs not have a basis, we cannot use the usual definition. Furthermore, our new definition does not necessarily coincide with the usual one when the quotient space has a basis. The first lemmas are rather technical; they allow us to show Lemma 7, which is, with the definition of successive vectors and of l_1^{n+} -averages in a quotient space, the main tool used afterwards.

3.1 Notation

Let $0 < \Delta < 1$.

Definition Let W be a Banach space. Let (w_n) and (w'_n) be two sequences in W. They are said to be Δ -equivalent if for all n,

$$||w_n - w'_n|| \le \Delta^n \inf(||w_n||, ||w'_n||).$$

Definitions Let W be a Banach space with a basis. A sequence (w_n) in W is said to be *almost successive* if it is Δ -equivalent to a sequence of successive vectors in W. Let V be a subset of W. A sequence (w_n) in W is said to be *almost in* V if it is Δ -equivalent to a sequence of vectors in V.

Lemma 1 Let W be a Banach space in \mathcal{X} . Let (w_i) be a successive sequence in W and (w'_i) be Δ -equivalent to (w_i) . Then

$$\left\|\sum_{i=1}^{n} w_{i}'\right\| \le (1+2\Delta) \left\|\sum_{i=1}^{n} w_{i}\right\|.$$

<u>Proof</u> By definition,

$$\left\|\sum_{i=1}^{n} w_{i}'\right\| \leq \left\|\sum_{i=1}^{n} w_{i}\right\| + \sum_{i=1}^{n} \Delta^{i} \|w_{i}\|,$$

so, as the basis in W is bimonotone,

$$\|\sum_{i=1}^{n} w_i'\| \le (1+2\Delta) \|\sum_{i=1}^{n} w_i\|.$$

Lemma 2 Let W be a space in \mathcal{X} . Let (w_n) be a sequence in W such that $w_n/||w_n|| \stackrel{w}{\to} 0$. Then (w_n) has an almost successive subsequence.

<u>Proof</u> We may assume (w_n) is a norm 1 sequence. Assume we have already selected $w_{n_1}, \ldots, w_{n_{k-1}}$ satisfying the conclusion; let t_1, \ldots, t_{k-1} be the associated successive sequence. Let E be an interval containing the first vector of the basis and the range of $t_1 + \cdots + t_{k-1}$. There exists n_k such that $||Ew_{n_k}|| \leq \Delta^k/4$. Let $t'_k = w_{n_k} - Ew_{n_k}$. There exists an interval F such that Ft'_k is equal to t'_k up to $\Delta^k/4$. If we let $t_k = Ft'_k$, we have that $t_k > t_{k-1}$, and $||t_k - w_{n_k}|| \leq \Delta^k/2$, from which it follows that

$$||t_k - w_{n_k}|| \le \Delta^k \inf(||t_k||, ||w_{n_k}||),$$

so the sequence w_{n_1}, \ldots, w_{n_k} and the successive sequence t_1, \ldots, t_k are Δ -equivalent.

¿From now on, X stands for a reflexive Banach space in \mathcal{X}_f , Y for an infinite dimensional and infinite codimensional subspace of X, and we assume that $\Delta = 10^{-4}$.

3.2 Successive vectors in X/Y

Definitions Given z in X, we denote by \hat{z} its class in X/Y.

A couple denotes a sequence $((\widetilde{z_n}, z_n^*))$ in $X \times X^*$.

For an element (\tilde{z}, z^*) in $X \times X^*$, the range is the smallest interval containing $ran(\tilde{z})$ and $ran(z^*)$.

A couple is said to be *successive* if the sequence of ranges is successive.

Let $(z_n)_{n \in \mathbb{N}}$ be a sequence in X/Y. A couple $(\widetilde{z_n}, z_n^*)$ is said to be associated to (z_n) if $(\widehat{\overline{z_n}})$ and (z_n) are Δ -equivalent and (z_n^*) is almost in $B(Y^{\perp})$.

Let $\lambda > 1$. A couple associated to (z_n) is said to be λ -associated to (z_n) if for all n, $\|\widetilde{z_n}\| \leq \lambda \|z_n\|$ and $z_n^*(\widetilde{z_n}) \geq (1/3) \|z_n\|$.

Definition Let $\lambda > 1$. Let $(z_n)_{n \in \mathbb{N}}$ be a sequence in X/Y. We say that (z_n) is λ -successive if it is λ -associated to some successive couple $((\widetilde{z_n}, z_n^*))$. We shall then call $(\widetilde{z_n})$ a λ -lifting, (z_n^*) a λ -functional.

Remark If (z_n) is λ -successive, then for any $\lambda' \geq \lambda$, any subsequence (z'_n) of (z_n) , z'_n is λ' -successive.

Lemma 3 Let (z_n) be a 400-successive sequence in X/Y. Then for all n,

$$(4f(n))^{-1}\sum_{i=1}^{n} ||z_i|| \le ||\sum_{i=1}^{n} z_i||.$$

<u>Proof</u> Let (z_n) be a 400-successive sequence. Let (z_n^*) in X^* be the successive functional associated to (z_n) , let (y_n^*) in $B(Y^{\perp})$ be the almost successive functional Δ -equivalent to (z_n^*) . By Lemma 1, we have that

$$\|\sum_{i=1}^{n} y_i^*\| \le (1+2\Delta) \|\sum_{i=1}^{n} z_i^*\|,$$

 \mathbf{SO}

$$\|\sum_{i=1}^{n} y_i^*\| \le (1+2\Delta)(1+\Delta)f(n) \le (1+4\Delta)f(n),$$

It follows that

$$\begin{split} \|\sum_{i=1}^{n} z_{i}\| &\geq ((1+4\Delta)f(n))^{-1} \left(\sum_{i=1}^{n} y_{i}^{*}\right) \left(\sum_{i=1}^{n} z_{i}\right), \\ \|\sum_{i=1}^{n} z_{i}\| &\geq ((1+4\Delta)f(n))^{-1} \left((\sum_{i=1}^{n} z_{i}^{*})(\sum_{i=1}^{n} \widetilde{z}_{i}) - (\sum_{i=1}^{n} \|z_{i}^{*} - y_{i}^{*}\|)(\sum_{i=1}^{n} \|\widetilde{z}_{i}\|) \right), \\ \|\sum_{i=1}^{n} z_{i}\| &\geq ((1+4\Delta)f(n))^{-1}(1/3 - 800\Delta) \sum_{i=1}^{n} \|z_{i}\| \geq (4f(n))^{-1} \sum_{i=1}^{n} \|z_{i}\|. \end{split}$$

Lemma 4 Let (z_n) be a sequence in X/Y such that $z_n/||z_n|| \xrightarrow{w} 0$. Then (z_n) has a 3-successive subsequence.

<u>Proof</u> We may assume that (z_n) is a norm 1 sequence. Let $(\widetilde{z_n}')$ be a lifting for (z_n) such that $\|\widetilde{z_n}'\| \leq 1 + \Delta^n$. The sequence $(\widetilde{z_n}')$ is bounded, so, passing to a subsequence, we may assume that $(\widetilde{z_n}')$ converges weakly. Let y be the weak limit of $(\widetilde{z_n}')$. The vector y has norm 1, and belongs to Y, because for every y^* in Y^{\perp} , $y^*(\widetilde{z_n}') = y^*(z_n)$ tends to 0.

Let $\widetilde{y_n} = \widetilde{z_n}' - y$. We have

$$\|\widetilde{y_n}\| \le 2 + \Delta^n.$$

We have that $\widetilde{y_n} \stackrel{w}{\to} 0$ so passing to a further subsequence, we may assume by Lemma 1 that $(\widetilde{y_n})$ is almost successive; let $(\widetilde{z_n})$ be a successive sequence equivalent to $(\widetilde{y_n})$. We have that

$$\|\widetilde{z_n}\| \le (1+\Delta^n) \|\widetilde{y_n}\| \le 3 \|z_n\|.$$

Now let (y'_n) be a dual sequence in $B(Y^{\perp})$ such that for all $n, y'_n(z_n) = 1$. Passing to a subsequence, we may assume that $y'_n \stackrel{w}{\to} y^*$ and that for all $n, |y^*(z_n)| \leq 1/6$. Let $y_n^* = 1/2(y'_n - y^*)$. We have that $y_n^* \in B(Y^{\perp})$ and

$$y_n^*(z_n) = 1/2(y_n'^*(z_n) - y^*(z_n)) \ge 5/12$$

for n greater than some N. Furthermore, $y_n^* \xrightarrow{w} 0$. Passing to a new subsequence, by Lemma 3, we may assume that (y_n^*) is almost successive. Let (z_n^*) be the equivalent successive sequence. Then

$$z_n^*(\widetilde{z_n}) \ge y_n^*(z_n) - ||z_n^* - y_n^*|| ||\widetilde{z_n}||,$$

so that

$$z_n^*(\widetilde{z_n}) \ge 5/12 - 3.\Delta^n \ge 1/3.$$

Definition Let $\lambda > 1$. Let (z_n) be a sequence in X/Y. For any finite subset E of N, z_E denotes the sum $\sum_{i \in E} z_i$. For $j \ge 0$, $n \ge 0$, let $E_n(j)$ be the interval $[2^{j}n + 1, 2^{j}n + 2^{j}]$. We say that (z_{n}) is λ -supersuccessive if for every j, the sequence $(z_{E_n(j)})_{n\geq 1}$ is λ -successive.

Lemma 5 Let (z_n) be a norm 1 sequence in X/Y such that $z_n \xrightarrow{w} 0$. Then (z_n) has a 3-supersuccessive subsequence.

Proof It is enough to prove by induction that there exists an inclusion decreasing sequence of subsequences $(z_n^1)_{n \in \mathbb{N}}, \ldots, (z_n^k)_{n \in \mathbb{N}}$ such that for all k, all $j \leq k$, the sequence $(z_{E_n(j)}^k)_{n\geq 1}$ is 3-successive. Indeed, the result then follows by taking a diagonal subsequence.

Assume the induction hypothesis is true for k-1. The sequence $(z_{E_n(k)}^{k-1})$ converges weakly to 0 and by Lemma 3, it is bounded below, so by Lemma 4, it admits a 3 - successive subsequence $(z_{E_{n_i}(k)}^{k-1})_{i \in \mathbb{N}}$. Let $E = \bigcup_i E_{n_i}$. We let $(z_n^k)_{n \in \mathbb{N}}$ be the sequence $(z_n^{k-1})_{n \in E}$. It is a subsequence of $(z_n^{k-1})_{n \in \mathbb{N}}$. Furthermore, it is easy to check that the sequence $(z_{E_n(j)}^k)_{n \geq 1}$ is still successive for $j \leq k - 1$, and by construction, this is also true for j = k.

3.3 l_1^{n+} vectors in X/Y

Definitions: l_1^{n+} -vectors Let $\epsilon_0 = 1/40$, let C > 0.

An l_1^{n+} -vector with constant C in X is a vector x of the form $\sum_{i=1}^n x_i$ such that the sequence (x_i) is successive and for all i, $||x_i|| \le C ||x||/n$.

An l_1^{n+} -average in X is a norm 1 l_1^{n+} -vector in X.

An l_1^{n+} -vector in X/Y is a vector x of the form $\sum_{i=1}^n x_i$ such that the sequence (x_i) is 3-successive and for all i, $||x_i|| \le (1 + \epsilon_0) ||x||/n$.

Let x be a l_1^{n+} -vector in X/Y of the form $\sum_{i=1}^n x_i$. An l_1^{n+} -lifting for x

is a lifting $\tilde{x} = \sum_{i=1}^{n} \tilde{x}_i$, where $(\tilde{x}_i)_{1 \le i \le n}$ is a 3-lifting for $(x_i)_{1 \le i \le n}$. An l_1^{n+} -average in X/Y is an l_1^{n+} -vector in X/Y the l_1^{n+} -lifting of which has norm 1.

Lemma 6 Let $n \in \mathbf{N}$. Let $x = \sum_{i=1}^{n} x_i$ be a l_1^{n+} -average in X/Y, let \tilde{x} be an l_1^{n+} -lifting for x. Then \tilde{x} is a l_1^{n+} -average with constant 4 and $\|\tilde{x}\| \le 4\|x\|.$

Proof We know that the vectors (\tilde{x}_i) are successive in X. Now

$$\|x - \hat{\tilde{x}}\| \le \sum_{i=1}^{n} \|x_i - \hat{\tilde{x}}_i\| \le \sum_{i=1}^{n} \Delta^i \|x_i\| \le 2\Delta(1 + \epsilon_0)/n \|x\|$$

It follows that

$$\|x\| \le (1 - 2\Delta(1 + \epsilon_0)/n)^{-1} \|\hat{\tilde{x}}\| \le (1 - 2\Delta(1 + \epsilon_0)/n)^{-1} \|\tilde{x}\|.$$

So

$$\|\tilde{x}_i\| \le 3\|x_i\| \le 3(1+\epsilon_0)\|x\|/n \le 4\|\tilde{x}\|/n,$$

so \tilde{x} is a l_1^{n+} with constant 4. Furthermore,

$$\|\tilde{x}\| \le \sum_{i=1}^{n} \|\tilde{x}_i\| \le 3 \sum_{i=1}^{n} \|x_i\| \le 3(1+\epsilon_0) \|x\| \le 4 \|x\|.$$

Lemma 7 Let $Z \subset X/Y$. Then Z contains a 4-successive basic sequence (Z_n) of $l_1^{2^n+}$ -averages, such that for all n, the 4-lifting $\widetilde{Z_n}$ is a $l_1^{2^n+}$ -lifting. Such a sequence will be called an average basic sequence in Z.

<u>Proof</u> As Z is reflexive, there exists a basic sequence (z_n) of unit vectors such that $z_n \xrightarrow{w} 0$. By Lemma 5, we may assume that (z_n) is 3-supersuccessive.

We now prove the result by induction. Assume we have chosen the first n-1 terms Z_1, \ldots, Z_{n-1} of the sequence. Let $(\widetilde{Z}_j, Z_j^*)_{1 \leq j \leq n-1}$ be the associated 4-successive couple. Let T satisfy $4f(2^{nT}) < (1 + \epsilon_0)^T$ and let N = nT. Let Z'_1, \ldots, Z'_{2^N} be the sequence $Z_{2^N r+1}, \ldots, Z_{2^N r+2^N}$, for some r such that, with this notation,

$$\min_{j \le n, p \le 2^{N-j}} ran((\widetilde{Z'_{E_p(j)}}, {Z'_{E_p(j)}}^*)) > ran((\widetilde{Z_{n-1}}, Z_{n-1}^*)).$$

For any $j \leq N$, the sequence (Z'_{E_nj}) is 3-successive. Now assume no decomposition of any $Z'_{E_p(j)}$ as $\sum_{k=1}^{2^n} Z'_{E_k(j-1)}$ is an $l_1^{2^n+}$ decomposition. It follows by induction that

$$||Z'_{E_p(j)}|| \le (2^n/(1+\epsilon_0))^j,$$

so that

$$||Z'_{E_1(T)}|| \le 2^N / (1 + \epsilon_0)^T.$$

But (z_n) is 3-successive so by Lemma 3,

$$||Z'_{E_1(T)}|| = ||Z'_1 + \dots + Z'_{2^N}|| \ge 2^N/4f(2^N),$$

a contradiction by choice of N.

It follows that one of the $Z'_{E_p(j)}$ is a $l_1^{2^n+}$ -vector; we choose Z_n to be the associated $l_1^{2^n+}$ -average. By choice of r, the couple $(\widetilde{Z_n}, Z_n^*)$ satisfies $(\widetilde{Z_n}, Z_n^*) > (\widetilde{Z_{n-1}}, Z_{n-1}^*)$.

4 Rapidly Increasing Sequences

4.1 R.I.S.-vectors in a space of the class \mathcal{X}_f

We now define R.I.S.-vectors in a Banach space X in \mathcal{X}_f . In fact, the properties of R.I.S. are not interesting in all spaces in \mathcal{X}_f , but only on those of Gowers-Maurey's type; we give a sense to this expression in the following paragraph, defining GM-type spaces, and then state several lemmas true in GM-type spaces.

Definitions Let J be a set of integers $\{j_n, n \in \mathbf{N}\}$, such that $f(j_1) > 256$ and for all n, $\log \log \log j_{n+1} \ge 4j_n^2$, let $K = \{j_1, j_3, j_5, \ldots\}$, let $L = \{j_2, j_4, j_6, \ldots\}$.

Let $L' \subset L$. An L'-sequence is a successive sequence $x_1^* < \cdots < x_k^*$ with $k \in K$, such that for all i, x_i^* is an (M_i, f) -form where M_i is an element in L' greater than j_{2k} . An L'-sum is a vector of the form $1/\sqrt{f(k)} \sum_{i=1}^k x_i^*$, where x_1^*, \ldots, x_k^* is a L'-sequence.

A space X is of GM-type if it belongs to \mathcal{X}_f , and if there exists a set \mathcal{S} of L-sums in $B(X^*)$ such that every vector in X has either the supremum norm, or is normed by an (M, f)-form, or by an element in \mathcal{S} .

Notice that Gowers-Maurey's space is of GM-type (the set S being the set of special sums), but that a GM-type space needs not be H.I.. Notice also that a space of GM-type satisfies a f-lower estimate, so by Lemma GM3, every block subspace contains l_1^{n+} -vectors with arbitrary constants and lengths.

Definition We recall that a *R.I.S.* of length *N* with constant ϵ in *X* is a successive sequence $(x_i)_{i=1}^N$ of $l_1^{n_i}$ -averages with constant *C* in *X* such that (n_i) satisfies $n_1 \ge 4(1+\epsilon)M_f(N/\epsilon')/\epsilon'$ and $\epsilon'/2f(n_k)^{1/2} \ge |ran(x_{k-1})|$ for $k = 2, \ldots, N$, where $\epsilon' = \min\{\epsilon, 1\}$ and $M_f(x) = f^{-1}(36x^2)$.

A R.I.S.-vector is a non-zero multiple of the sum of a R.I.S..

We now show some lemmas very similar to those of [GM]; we have to state them because we shall use different constants, and because they can be applied to any GM-type space, which will be useful in the last part of the article.

Lemma 8 Let X have GM-type. Let $N \in L$, let $n \in [\log N, \exp N]$, let $(x_i)_{i=1}^N$ be a R.I.S. of length M with constant ϵ in X. Then $\|\sum_{i=1}^n x_i\| \leq (1 + \epsilon + \epsilon')nf(n)^{-1}$.

<u>Proof</u> As X has GM-type, all the hypothesis of Lemma GM7 are satisfied. The conclusion then follows from Lemma GM9.

Lemma 9 Let X have GM-type. Let $N \in L$. Let $M = N^{1/40}$. Let x_1, \ldots, x_N be a R.I.S. in X with constant 4. Then $\sum_{i=1}^N x_i$ is a l_{1+}^M -vector

with constant 6.

 $\frac{\text{Proof}}{\sum_{i=(j-1)m+1}^{jm} x_i} \text{ Let } m = N/M, \text{ let } x = \sum_{i=1}^N x_i, \text{ and for } 1 \le j \le M \text{ let } y_j = \sum_{i=(j-1)m+1}^{jm} x_i. \text{ By Lemma 8, } \|y_j\| \le 5mf(m)^{-1}, \text{ while } \|\sum_{j=1}^m y_j\| = \|x\| \ge Nf(N)^{-1}, \text{ so } x \text{ is an } l_1^{M+} \text{-vector with constant } 5f(N)/f(m) \le 6.$

Lemma 10 Every GM-type space is reflexive.

<u>Proof</u> We follow the proof that Gowers-Maurey's space is reflexive. Let X have GM-type. We show that the canonical basis e_1, e_2, \ldots of X is boundedly complete and shrinking (we refer to [LT] for the definition of these notions). It follows from the fact that X belongs to \mathcal{X}_f that the basis is boundedly complete. Now assume it is not shrinking. Then we can find $\epsilon > 0$, a norm-1 functional x^* in X^* , and a sequence of successive vectors x_1, x_2, \ldots such that $x^*(x_n) \ge \epsilon$ for every n. It follows that $\sum_{n \in A} x_n$ is a $l_{1+}^{|A|}$ -vector with constant ϵ^{-1} for any $A \subset \mathbf{N}$. Given $N \in L$, we may construct a R.I.S. y_1, \ldots, y_N with constant ϵ^{-1} with such l_1^{n+} -vectors, and we have $x^*(y_1 + \cdots + y_N) \ge \epsilon N$. For N large enough, this contradicts Lemma 8.

Definition Let X have GM-type. Let x_1^*, \ldots, x_k^* be an L-sequence of lenght k, where each x_i^* is an (M_i, f) -form. A sequence of successive vectors $x_1 < \cdots < x_k$ in X is said to be a R.I.S. associated to x_1^*, \ldots, x_k^* if for every i, x_i is a normalized R.I.S. of lenght M_i and constant 4, $M_1 = j_{2k}$ and $1/2f((M_i)^{1/40})^{1/2} \ge |ran(x_{i-1})|$.

Remark Because of the increasing condition, and by Lemma 9, a R.I.S. associated to an L-sequence of length k is a R.I.S. with constant 6.

Lemma 11 Let X have GM-type. Let x be a norm 1 R.I.S.-vector in X of length $N_1 \in L$ and constant 4 and let x^* be an $(N_2, f) - f$ orm in X^* with $N_2 \in L$, and assume $N_1 \neq N_2$. Let $k \in K$ be such that $N_1 \geq j_{2k}$, $N_2 \geq j_{2k}$. Then for every interval E, $|x^*(Ex)| \leq 1/k^2$.

<u>Proof</u> The proof relies on Lemmas GM4 and GM5, which we may apply since X satisfies an f-lower estimate. First, by Lemma 9, x is a $l_1^{N'_1+}$ -average with constant 6, where $N'_1 = N_1^{1/40}$.

Now if $N_2 < N_1$, then $N_2 < N'_1$ by the lacunarity of *L*. By Lemma GM4, $|x^*(Ex)| = |(Ex^*)(x)| \le 18/f(N_2) \le 18/f(j_{2k}) \le k^{-2}$.

If $N_1 < N_2$, then $M_f(N_1) < N_2$ by the lacunarity of L, so we may apply Lemma GM5 to x', the sum of the R.I.S. whose normalized sum is x. It follows that $|x^*(Ex')| \leq 5$, while $||x'|| \geq N_1/f(N_1) \geq 5k^2$.

Lemma 12 Let X have GM-type. Let $k \in K$. Let $x_1 < \ldots < x_k$ in X be a R.I.S. associated to some L-sequence. Let $x = \sum_{i=1}^k x_i$.

Assume that for every L-sum in S, every interval E, $|z^*(Ex)| \leq 1/4$. Then

$$\|x\| \le 7k/f(k).$$

<u>Proof</u> By Lemma GM9 of [GM], for any $K_0 \subset K$, there exists a function g_{K_0} belonging to the "Schlumprecht" class \mathcal{F} of functions, such that $g_{K_0}(x) = \sqrt{f(x)}$ when $x \in K_0$, and $g_{K_0}(x) = f(x)$ when $x \in [\log N, \exp N]$, for any $N \in J \setminus K_0$.

Now take $K_0 = K \setminus \{k\}$, and let g be the function associated to K_0 . By the hypothesis, for any interval E,

$$||Ex|| \le 1 \lor \sup\{|x^*(Ex)| : M \ge 2, x^*(M,g) - form\}$$

As x_1, \ldots, x_k is a R.I.S. with constant 6, it is an easy consequence of Lemma GM7 that $||x|| \leq \frac{7k}{f(k)}$.

4.2 R.I.S.-vectors in a quotient space

We finally define R.I.S. in a quotient space X/Y where X has GM-type and link them with the R.I.S. in X.

Definition Let X have GM-type. Let $Z \subset X/Y$. Let (Z_n) be an average basic sequence in Z.

A *R.I.S.* of length M in Z is a subsequence $(Z_{n_i})_{i=1}^N$ of (Z_n) such that $2^{n_1} \ge 16M_f(N)$ and $1/2f(2^{n_i})^{1/2} \ge ran(\widetilde{Z_{n_{i-1}}})$ for $i = 2, \ldots, N$.

Lemma 13 Let X have GM-type. Let $Z \subset X/Y$. Let $M \in L$. Let $z = \sum_{i=1}^{M} Z_{n_i}$ be a R.I.S.-vector of length M in Z. Let $\tilde{z} = \sum_{i=1}^{M} \widetilde{Z_{n_i}}$ be the sum of the sequence of the associated $l_1^{n_i+}$ -liftings. Then \tilde{z} is a R.I.S.-vector in X with constant 4. Furthermore $\|\tilde{z}\| \leq 80\|z\|$.

<u>**Proof**</u> The first part is a direct consequence of Lemma 6 and of the definitions. For the second, notice that

$$\|\widetilde{z}\| = \|\widetilde{Z_{n_1}} + \dots + \widetilde{Z_{n_M}}\| \le 5M/f(M)$$

by Lemma 8, while

$$||z|| = ||Z_{n_1} + \dots + Z_{n_M}|| \ge (\sum_{i=1}^M ||Z_{n_i}||)/4f(M)$$

by Lemma 3. As $||Z_{n_i}|| \ge (1/4) ||\widetilde{Z_{n_i}}|| \ge 1/4$ by Lemma 6, it follows that $||\tilde{z}|| \le 80 ||z||$.

5 Gowers-Maurey's space is Q.H.I.

We refer to [GM] for the definition of Gowers-Maurey's space, which we shall denote by X. Gowers and Maurey have proved that X is H.I. Now let Y be any infinite dimensional and infinite codimensional subspace of X. We shall prove that X/Y is H.I.. We first show a Lemma similar to Lemma GM2.

Lemma 14 Let x_1^*, \ldots, x_k^* be a special sequence in X. Let $x_1 < \ldots < x_k$ be a R.I.S associated to x_1^*, \ldots, x_k^* . Let $x = \sum_{i=1}^k x_i$. Assume that for every interval E, $|(\sum_{i=1}^k x_i^*)(Ex)| \le 2$, then

$$\|x\| \le 7k/f(k).$$

<u>Proof</u> We already know that X has GM-type. By Lemma 12, it is enough to prove that for any special function z^* , every interval $E, |z^*(Ex)| \leq$ 1/4. We follow the proof of Lemma GM2. Let z^* be such a function of the form $f(k)^{-1/2} \sum_{i=1}^{k} z_i^*$. Let t be maximal such that $z_t^* = x_t^*$, or 0 if no such t exists. Assume $i \neq j$ or one of i, j is greater than t + 1. Then since σ is an injection, we can find $l \neq l'$ in L such that z_i^* is an (l, f)-form and x_i is a norm 1 R.I.S.-vector of length l'. It follows then from Lemma 11 that $|z_i^*(Ex_j)| \leq k^{-2}$. Now choose an interval F such that

$$\left| \left(\sum_{i=1}^{t} z_i^* \right) (Ex) \right| = \left| \left(\sum_{i=1}^{k} x_i^* \right) (Fx) \right| \le 2.$$

It follows that

$$\left| \left(\sum_{i=1}^{k} z_{i}^{*} \right) (Ex) \right| \le 2 + |z_{t+1}^{*}(x_{t+1})| + k^{2} \cdot k^{-2} \le 4,$$

and that $|z^*(Ex)| \le 4f(k)^{-1/2} < 1/4$.

Conclusion Let Z and Z' be two subspaces of X/Y. We want to prove that their sum is not direct. Lemma 7 allows us to consider an average basic sequence (Z_n) (resp. (Z'_n)) in Z (resp. Z'). Let $\delta > 0$, let $k \in K$ be such that $2.10^4/\sqrt{f(k)} \leq \delta$.

Let $M_1 = j_{2k}$. We may build with vectors of the average basic sequence (Z_n) a R.I.S. vector in Z of length M_1 , of the form $z_1 = \sum_{j=1}^{M_1} z_1^j$. Let $\tilde{z_1} = \sum_{j=1}^{M_1} z_1^j$. By Lemma 13, $\tilde{z_1}$ is the sum of a R.I.S. with constant 4, and we may assume it is of norm 1.

Let (z_1^{j*}) be the successive functional in X^* associated to $(z_1^j), (y_1^{j*})$ be the almost successive functional in $B(Y^{\perp})$ Δ -equivalent to (z_1^{j*}) . Let $z_1^{\prime *} = f(M_1)^{-1} (\sum_{j=1}^{M_1} z_1^{j*}).$ We have that $(1 + \Delta)^{-1} z_1^{\prime *}$ is a (M_1, f) -form, and

$$z_1^{\prime *}(\widetilde{z_1}) = f(M_1)^{-1} \sum_{j=1}^{M_1} z_1^{j *}(\widetilde{z_1^j}),$$

 \mathbf{SO}

$$z_1^{\prime *}(\widetilde{z_1}) \ge (3f(M_1))^{-1} \sum_{j=1}^{M_1} ||z_1^j||,$$

and by Lemma 13 and Lemma 2,

$$z_1^{\prime *}(\tilde{z_1}) \ge 1/240 \sum_{j=1}^{M_1} \|\tilde{z_1^j}\| \ge 1/960.$$

It follows that we may find a (M_1, f) -form z_1^* in **Q** such that $z_1^*(\tilde{z}_1) = 10^{-3}$ up to 1/k. Furthermore there is an element y_1^* in $B(Y^{\perp})$, such that

$$||y_1^* - z_1^*|| \le 1/f(M_1) \sum_{j=1}^{M_1} \Delta^j \le \Delta .10^{-3}/2 \le \Delta \min\{||y_1^*||, ||z_1^*||\}.$$

Let $M_2 = \sigma(z_1^*)$. As \mathbf{Q} is dense, we had infinitely many choices for z_1^* , so we may assume we chose z_1^* such that $1/2f((M_2)^{1/40})^{1/2} \ge |ran(\tilde{z_1})|$. As in above, we may find a R.I.S. vector z_2 in Z', whose lifting $\tilde{z_2}$ is of norm 1, and a (M_2, f) -form z_2^* in \mathbf{Q} such that $|z_2^*(z_2) - 10^{-3}| \le 1/k$, and such that the couple $(\tilde{z_2}, z_2^*)$ is successive to $(\tilde{z_1}, z_1^*)$. Going on in the same way, we build sequences $(z_i), (\tilde{z_i}), (z_i^*)$ and (y_i^*) such that for all i, z_i is in Z if i is odd, in Z' if i is even, z_i^* is a (M_i, f) -form, $||y_i^* - z_i^*|| \le \Delta^i \min\{||y_i^*||, ||z_i^*||\}$ $1/2f((M_i)^{1/40})^{1/2} \ge |ran(\tilde{z_{i-1}})|$, and $(\tilde{z_i}, z_i^*)$ is successive to $(\tilde{z_{i-1}}, z_{i-1}^*)$.

So by construction, z_1^*, \ldots, z_k^* is a special sequence, (z_i^*) and (y_i^*) are Δ -equivalent (so that the sequence (z_i^*) is almost in $B(Y^{\perp})$), and z_1, \ldots, z_k is a R.I.S. associated to z_1^*, \ldots, z_k^* .

It follows from Lemma 1 that $\|\sum_{i=1}^{k} y_i^*\| \leq (1+2\Delta)\sqrt{f(k)}$ so

$$\begin{split} \|\sum_{i=1}^{k} z_{i}\| &\geq \left((1+2\Delta)\sqrt{f(k)}\right)^{-1} \left(\sum_{i=1}^{k} y_{i}^{*}\right) \left(\sum_{i=1}^{k} z_{i}\right), \\ \|\sum_{i=1}^{k} z_{i}\| &\geq \left((1+2\Delta)\sqrt{f(k)}\right)^{-1} \left(\sum_{i=1}^{k} z_{i}^{*}(\tilde{z}_{i}) - 2\Delta\|\sum_{i=1}^{k} \tilde{z}_{i}\|\right), \\ \|\sum_{i=1}^{k} z_{i}\| &\geq \left((1+2\Delta)\sqrt{f(k)}\right)^{-1} (10^{-3}k - 1 - 2\Delta k) \geq 4.10^{-4}k/\sqrt{f(k)}. \end{split}$$

On the other hand, for all interval E, $|(\sum_{i=1}^{k} z_i^*)(\sum_{i=1}^{k} (-1)^i \tilde{z}_i)| \leq 2$, so by Lemma 14,

$$\left\|\sum_{i=1}^{k} (-1)^{i} \widetilde{z}_{i}\right\| \leq 7k/f(k).$$

It follows that

$$\left\|\sum_{i=1}^{k} (-1)^{i} z_{i}\right\| \leq \left\|\sum_{i=1}^{k} (-1)^{i} \widetilde{z}_{i}\right\| + \sum_{i=1}^{k} \left\|z_{i} - \widehat{\widetilde{z}}_{i}\right\| \leq \frac{7k}{f(k)} + 2\Delta \leq \frac{8k}{f(k)}$$

If z denotes the sum of the odd vectors, z' the sum of the even vectors, we have that $z \in Z, z' \in Z'$, and

$$||z - z'|| \le 2.10^4 f(k)^{-1/2} ||z + z'|| \le \delta ||z + z'||.$$

As δ is arbitrary, it follows that the sum of Z and Z' is not direct, and finally, that X/Y is H.I..

Remark As X is reflexive, it follows from Proposition 2 that X^* is H.I..

6 There exists a H.I. space \mathcal{X} such that \mathcal{X}^* is not H.I.

In this section, we build a H.I., not Q.H.I. space \mathcal{X} as a certain sum of two GM-type H.I. spaces X_1 and X_2 . The space \mathcal{X}^* will contain a direct sum of two subspaces. By a simple generalization explained in the Appendix, it is even possible to build a H.I. space \mathcal{X} such that \mathcal{X}^* contains a direct sum of n subspaces.

6.1 Proposition 3

For i = 1, 2, let X_i be a H.I. Banach space, let Z_i be a subspace of X_i . Assume that Z_1 and Z_2 are isometric, and that X_1/Z_1 and X_2/Z_2 are infinite dimensional and totally incomparable. By abuse of notation, we identify both Z_1 and Z_2 with a same space Z. Let \mathcal{X} be the quotient space $(X_1 \times X_2)/\{(z, -z), z \in Z\}$. Then \mathcal{X} is H.I. and \mathcal{X}^* is not H.I.

<u>Proof</u> For x_i in X_i , i = 1, 2, we denote by \hat{x}_i the class of x_i in X_i/Z_i , by $(\widehat{x_1, x_2})$ the class of (x_1, x_2) in \mathcal{X} . By definition,

$$\|(\widehat{x_1, x_2})\| = \inf_{z \in Z} (\|x_1 + z\| + \|x_2 - z\|).$$

It follows that the space $\mathcal{X}_1 = \{(\widehat{x_1, 0}), x_1 \in X_1\}$ is isometric to X_1 , the space $\mathcal{X}_2 = \{(\widehat{0, x_2}), x_2 \in X_2\}$ is isometric to X_2 , and the space $\mathcal{Z} = \{(\widehat{z, 0}), z \in Z\} = \{(\widehat{0, z}), z \in Z\}$ is isometric to Z. As an easy consequence, we have the relation

$$\mathcal{X}/\mathcal{Z} = \mathcal{X}_1/\mathcal{Z} \oplus \mathcal{X}_2/\mathcal{Z} \simeq X_1/Z_1 \oplus X_2/Z_2,$$

 \mathbf{SO}

$$\mathcal{Z}^{\perp} \simeq (X_1/Z_1)^* \oplus (X_2/Z_2)^*,$$

and this proves that \mathcal{X}^* is not H.I.. It remains to show that \mathcal{X} is H.I.. For i = 1, 2, we define a linear operator $\phi_i : \mathcal{X} \to X_i/Z_i$ by $\phi_i((\widehat{x_1, x_2})) = \widehat{x_i}$. It is easy to check that ϕ_i is well defined. Now let \mathcal{W} be a subspace of \mathcal{X} . There exists an i such that $\phi_{i/\mathcal{W}}$ is infinitely singular: indeed, if $\phi_{1/\mathcal{W}}$ and $\phi_{2/\mathcal{W}}$ are both finitely singular, then there exists a subspace \mathcal{V} of \mathcal{W} on which ϕ_1 and ϕ_2 are isomorphisms into, so that X_1/Z_1 and X_2/Z_2 have isomorphic subspaces, a contradiction.

Now assume for example that $\phi_{1/\mathcal{W}}$ is infinitely singular. As a consequence, there exists a norm 1 sequence $(w_n)_{n \in \mathbf{N}}$ such that $\phi_1(w_n) \xrightarrow{+\infty} 0$. By definition of ϕ_1 , this means that $d(w_n, \mathcal{X}_2) \xrightarrow{+\infty} 0$. It follows easily that \mathcal{W} and \mathcal{X}_2 have Id + S-isomorphic subspaces. As \mathcal{X}_2 is isometric to X_2 , it is H.I.; it follows that \mathcal{Z} is quasi-maximal in \mathcal{X}_2 , so \mathcal{W} and \mathcal{Z} have also Id + S-isomorphic subspaces.

We have now proved that for every subspace \mathcal{W} of \mathcal{X} , \mathcal{W} and \mathcal{Z} have Id + S-isomorphic subspaces. This means that \mathcal{Z} is quasi-maximal in \mathcal{X} . As \mathcal{Z} is H.I., this implies that \mathcal{X} is H.I..

6.2 Proposition 4

For i = 1, 2, there exist X_i reflexive Q.H.I. Banach space, Z_i subspace of X_i , such that Z_1 and Z_2 are isometric, and such that X_1/Z_1 and X_2/Z_2 are totally incomparable.

6.3 Definition of X_1 and X_2

We shall define two spaces X_i of the form $(c_{00}, \|.\|_i)$ for i = 1, 2, following a Gowers-Maurey's method, in which we force X_1 and X_2 to have isometric subspaces Z_1 and Z_2 . The Banach spaces involved in Proposition 4 are the completion of the spaces X_1 and X_2 .

Let $(e_n)_{n \in \mathbb{N}}$ be the canonical unit basis of c_{00} .

Let \mathbf{Q} be the set of sequences with finite range, rational coordinates and maximum at most one in modulus. We recall that J is a set of integers $\{j_n, n \in \mathbf{N}\}$, such that $f(j_1) > 256$ and for all n, log log log $j_{n+1} \ge 4j_n^2$, that $K = \{j_1, j_3, j_5, \ldots\}$, and $L = \{j_2, j_4, j_6, \ldots\}$. Furthermore, we let $L_1 =$ $\{j_2, j_6, j_{10}, \ldots\}$, $L_2 = \{j_4, j_8, \ldots\}$. For i = 1, 2, let σ_i be an injection from the collection of finite sequences of successive elements of \mathbf{Q} to L_i . For i =1, 2, we choose X_i to be of the form $(c_{00}, \|.\|_i)$, and $Z_i = span\{e_{2n+1}, n \in$ $\mathbf{N}\}$. We may identify Z_1 and Z_2 with the same algebraic space Z, and Z_1^{\perp} and Z_2^{\perp} with the same algebraic space $Z^{\perp} = span\{e_{2n}^*, n \in \mathbf{N}\}$. We now need some definitions.

Definitions A *dual couple* is a couple (G, H) of balanced bounded convex subsets of c_{00} .

Let (G, H) be a dual couple.

A *N*-Schlumprecht sum in *G* is a sum of the form $1/f(N) \sum_{i=1}^{N} x_i^*$, where the x_i^* 's are successive in *G*. A Schlumprecht sum in *G* is a *N*-Schlumprecht sum in *G* for some *N*. The set of Schlumprecht sums in *G* is denoted by $\Sigma(G)$. In the same way, we define Schlumprecht sums in *H*.

A special sequence in G is a sequence of successive vectors $x_1^* < \ldots < x_k^*$, with $k \in K$, such that for $i = 1, \ldots, k$, x_i^* is an M_i -Schlumprecht sum in G with $M_i \geq j_{2k}$, and $M_i = \sigma_1(x_1^*, \ldots, x_{i-1}^*)$ for $i = 2, \ldots, k$.

A special sum in G is a sum of the form $1/\sqrt{f(k)} \sum_{i=1}^{k} x_i^*$, where $x_1^* < \ldots < x_k^*$ is a special sequence in G. The set of special sums in G is denoted by S(G).

We similarly define special sequences and special sums in H replacing σ_1 by σ_2 in the above definition.

An associated dual couple is a dual couple (G, H) such that there exist two multivalued functions $a : G \to H$ and $b : H \to G$ satisfying the four following properties.

- (a) for all $x^* \in G$, all $y^* \in a(x^*)$, $y^* x^*$ is in Z^{\perp} ;
- (b) for all $x^* \in G$, all $y^* \in a(x^*)$, $ran(y^*) \subset ran(x^*)$;
- (c) for all $x^* \in G \cap Z^{\perp}$, $a(x^*) = \{0\}$;

(d) for all N-Schlumprecht sum x^* in G, $a(x^*)$ contains an N-Schlumprecht sum in H,

and the four similar properties for b.

Definitions Let (G, H) be an associated dual couple.

A shadow sequence in G is a sequence of successive vectors $x_1^* < \ldots < x_k^*$ such that there exists a special sequence $y_1^* < \ldots < y_k^*$ in H such that for all i, x_i^* is an M_i -Schlumprecht sum in G belonging to $b(y_i^*)$, where M_i is the integer associated to y_i^* in the definition of the special sequence. A shadow sequence in G can be thought of as the "shadow" of a special sequence in H.

A shadow sum in G is a sum of the form $1/\sqrt{f(k)}\sum_{i=1}^{k} x_i^*$, where $x_1^* < \ldots < x_k^*$ is a shadow sequence in G. The set of shadow sums in G is denoted by s(G, H).

We similarly define shadow sequences and shadow sums in H, and denote the set of shadow sums by s(H, G).

To define the norms, we shall now build by induction an associated dual couple (C, D) where C (resp. D) is meant to be almost the dual unit ball of X_1 (resp. X_2). We shall build C as $\bigcup_{n \in \mathbb{N}} C_n$, building the increasing sequence C_n by induction. We shall also build a by induction, defining a function a_n from C_n to D_n at each step n; but to simplify the notation, we shall denote all the terms of the sequence by a (and we shall do symmetrically the same for D and b). In this situation, Property (a) ensures that the subspaces Z_1 and Z_2 are isometric. Properties (b) and (d) allow us to give convenient properties to the images by a of the special sequences, that is the shadow sequences. Property (c) introduces an asymmetry, and as a consequence, the quotient spaces X_1/Z_1 and X_2/Z_2 will be totally incomparable.

Construction At the first step, we define (C_0, D_0) to be $(B(l_1), B(l_1))$, a and b by $a(\sum_{i \in \mathbf{N}} \lambda_i e_i^*) = b(\sum_{i \in \mathbf{N}} \lambda_i e_i^*) = \sum_{i \text{ odd }} \lambda_i e_i^*$. It is easy to check that (C_0, D_0) is an associated dual couple.

Now assume we are given an associated dual couple (C_{n-1}, D_{n-1}) , with functions $a: C_{n-1} \to D_{n-1}$ and $b: D_{n-1} \to C_{n-1}$.

We define C'_{n-1} to be $\Sigma(C_{n-1}) \cup S(C_{n-1}) \cup S(C_{n-1}, D_{n-1})$, and C_n to be the set of elements of the form $E(\sum_{i=1}^{M} \lambda_i x_i^*)$, where E is an interval projection, $\sum_{i=1}^{M} |\lambda_i| = 1$, and for all i, x_i^* is in C'_{n-1} . We define D_n in a similar way.

We now extend a to C_n . Let $x^* \in C_n$. If $x^* \in Z^{\perp}$, then we let $a(x^*) = \{0\}$. We now define a construction if x^* is not in Z^{\perp} .

The set $a(x^*)$ may be already defined or not (it is when x^* is in C_{n-1}); if not we may assume $a(x^*) = \emptyset$. Then we add new values to the set $a(x^*)$ in each of the following cases (notice that at least one of the possibilities happens, so that a is well defined on the whole of C_n , but that the possibilities are not exclusive).

- If x^* is a Schlumprecht sum of the form $f(N)^{-1} \sum_{i=1}^{N} x_i^*$ with $x_i^* \in C_{n-1}$ then we add to $a(x^*)$ the set $f(N)^{-1} \sum_{i=1}^{N} a(x_i^*)$. - If x^* is a special sum of the form $f(k)^{-1/2} \sum_{i=1}^{k} x_i^*$ where x_i^* is an

- If x^* is a special sum of the form $f(k)^{-1/2} \sum_{i=1}^{k} x_i^*$ where x_i^* is an (M_i, f) -form in C_{n-1} then we add to the set $a(x^*)$ the set of all sums of the form $f(k)^{-1/2} \sum_{i=1}^{k} y_i^*$, where y_i^* is an (M_i, f) -form in $a(x_i^*)$. - If x^* is a shadow sum of the form $f(k)^{-1/2} \sum_{i=1}^{k} x_i^*$ with $x_i^* \in b(y_i^*)$

- If x^* is a shadow sum of the form $f(k)^{-1/2} \sum_{i=1}^k x_i^*$ with $x_i^* \in b(y_i^*)$ and y_1^*, \ldots, y_k^* is a special sum in D_{n-1} , then we add to the set $a(x^*)$ the singleton $\{f(k)^{-1/2} \sum_{i=1}^k y_i^*\}$.

- If x^* is the projection of a convex combination of elements of the three previous forms, that is, $x^* = ran(x^*)(\sum_i \lambda_i x_i^*)$, then we add to the set $a(x^*)$ the set $ran(x^*)(\sum_i \lambda_i a(x_i^*))$, $a(x_i^*)$ being defined as above whether x_i^* is a Schlumprecht sum, a special sum, or a shadow sum in C_{n-1} . Notice that we only use this construction when x^* is not in Z^{\perp} .

It is then easy to check that a (resp. b) takes its values in D_n (resp. in C_n) and that it still satisfies the four properties (a)-(d), so (C_n, D_n) is a dual couple.

We finally define C as $\bigcup_{n \in \mathbb{N}} C_n$ and D as $\bigcup_{n \in \mathbb{N}} D_n$; the multifunction a (resp. b) is defined on C (resp. D), so (C, D) is a dual couple.

We define X_1 by its norm $\|.\|_1 = \sup_{x^* \in C} \langle x^*, . \rangle, X_2$ by its norm $\|.\|_2 = \sup_{y^* \in D} \langle y^*, . \rangle$.

6.4 Z_1 and Z_2 are isometric subspaces

Let z be an element of Z. Then

$$\|z\|_1 = \sup_{x^* \in C} () \leq \sup_{x^* \in C, y^* \in a(x^*)} (+ < y^*, z>).$$

Now by definition of $a, x^* - y^* \in Z^{\perp}$, so $\langle x^* - y^*, z \rangle = 0$; and as y^* is in $D, (\langle y^*, z \rangle) \leq ||z||_2$. It follows that $||z||_1 \leq ||z||_2$, and by symmetry, $||z||_1 = ||z||_2$.

6.5 X_1 and X_2 are reflexive and Q.H.I.

By symmetry, it is enough to show that for example X_1 is reflexive and Q.H.I..

Remarks With the definition following Lemma 7, a special sequence in X_1 (resp. X_2) is an L_1 (resp. L_2)-sequence, a shadow sequence in X_1 (resp. X_2) is an L_2 (resp. L_1)-sequence. Notice also that the space X_1 (resp. X_2) has GM-type, the set S_1 (resp. S_2) being the set of special and shadow sums, and so it is reflexive.

Lemma 15 Let X have GM-type. Let x_1^*, \ldots, x_k^* be a L_1 -sequence in X^* . Let $x_1 < \ldots < x_k$ in X be a R.I.S. associated to x_1^*, \ldots, x_k^* . Let $x = x_1 + \ldots + x_k$. Then for every L_2 -sum z^* of length k in X^* , every interval E, $|z^*(Ex)| \leq 1/4$.

Proof

Let z^* be an L_2 -sum, E be an interval. Then $z^* = 1/\sqrt{f(k)} \sum_{i=1}^k z_i^*$, where z_i^* is a (l_i, f) -form, and l_i is in L_2 . For every j, x_j is a norm 1 R.I.S. of length in L_1 . As L_1 and L_2 are disjoint, and the lengths are greater than j_{2k} by definition of an L-sequence and of an associated R.I.S., it follows from Lemma 11 that $|z_i^*(Ex_j)| \leq 1/k^2$. Finally,

$$|z^*(Ex)| \le 1/\sqrt{f(k)} \le 1/4.$$

Remark The result is also true for an L_1 -sum and a R.I.S. associated to a L_2 -sequence.

Lemma 16 Let x_1^*, \ldots, x_k^* be a special sequence in X_1^* . Let $x_1 < \ldots < x_k$ be associated to x_1^*, \ldots, x_k^* . Let $x = \sum_{i=1}^k x_i$. Assume that for every interval E, $|(\sum_{i=1}^k x_i^*)(Ex)| \le 2$, then $||x|| \le 7k/f(k)$.

<u>Proof</u> We already know that X_1 has GM-type, so from Lemma 15, for every shadow sum z^* , every interval E, $|z^*(Ex)| \leq 1/4$. We may now apply the same proof as in Lemma 14, replacing σ by σ_1 . Lemma 12 then gives the conclusion.

Proposition X_1 is Q.H.I..

<u>Proof</u> As we said above, X_1 is of GM-type, like Gowers-Maurey's space. Furthermore, we have proved Lemma 16, which is similar to Lemma 14 in the case of Gowers-Maurey's space. It follows that our final proof that Gowers-Maurey's space is Q.H.I. is still valid in the case of X_1 .

6.6 X_1/Z_1 and X_2/Z_2 are totally incomparable.

Lemma 17 Let y_1^*, \ldots, y_k^* be a special sequence in Z_2^{\perp} . Let $x_1 < \ldots < x_k$ in X_1 be associated to y_1^*, \ldots, y_k^* . Let $x = \sum_{i=1}^k x_i$. Then

$$\|x\| \le 7k/f(k)$$

<u>Proof</u> The space X_1 is of GM-type, so, because of Lemma 12, it is enough to show that for every interval E, every special or shadow sum z^* in X_1^* , $|z^*(Ex)| \leq 1/4$. By Lemma 15, this is true for every special sum. Now let E be an interval and let z^* be a shadow sum in X_1^* .

For every *i*, let M_i be such that y_i^* is an (M_i, f) -form. There exists a special sequence t_1^*, \ldots, t_k^* in X_2^* such that $z^* = 1/\sqrt{f(k)} \sum_{i=1}^k z_i^*$ with for every $i, z_i^* \in b(t_i^*)$; let N_i be such that t_i^* is an (N_i, f) -form; by construction z_i^* is also an (N_i, f) -form. Let $I = \sup\{i/M_i = N_i\}$, or 0 if no such I exists. For i < I, because σ_2 is an injection, we have that $t_i^* = y_i^*$. It follows that t_i^* is in Z_2^{\perp} , so $b(t_i^*) = \{0\}$, and $z_i^* = 0$. For i > I, z_i^* is an (N_i, f) -form, and N_i is an element of L_2 greater than j_{2k} and different from all M_j , $j = 1, \ldots, k$. It follows by Lemma 11 that $|z_i^*(Ex_j)| \leq 1/k^2$. Finally,

$$|z^*(Ex)| \le 1/\sqrt{f(k)}(0+|z_I^*(x_I)|+k^2.k^{-2}) \le 2/\sqrt{f(k)} \le 1/4.$$

Total incomparability We now assume that there exists an isomorphism α between a subspace W_1 of X_1/Z_1 and a subspace W_2 of X_2/Z_2 and we intend to find a contradiction.

First we notice that X_2/Z_2 has a basis (namely the basis $(e'_{2n})_{n \in \mathbb{N}}$ dual to the basis $(e^*_{2n})_{n \in \mathbb{N}}$). Let (w_n) be an average basic sequence in W_1 . Up to a perturbation on α , we may assume that the sequence $(\alpha(w_n))$ is a sequence of unit vectors, successive with respect to (e'_{2n}) .

Now let $k \in K$. In $span\{(w_n)\}$, we may find a R.I.S vector $x_1 = \sum_{i=1}^{M_1} x_1^i$ in X_1 of length $M_1 = j_{2k}$, such that $\widetilde{x_1} = \sum_{i=1}^{M_1} \widetilde{x_1^i}$ is of norm 1. For $i = 1, \ldots, M_1$, let $y_1^{i*} \in Z_2^{\perp}$ be a functional that norms $\alpha(x_1^i)$ and such that $ran(y_1^{i*}) \subset ran(\alpha(x_1^i))$, and let $y_1'^*$ be the (M_1, f) -form $f(M_1)^{-1} \sum_{i=1}^{M_1} y_1^{i*}$. We have that $y_1'^*$ is in Z_2^{\perp} and

$$y_1^{\prime *}(\alpha(x_1)) = f(M_1)^{-1} \sum_{i=1}^{M_1} \|\alpha(x_1^i)\|,$$

so by Lemma 7 and Lemma 9,

$$y_1^{\prime *}(\alpha(x_1)) \ge (4\|\alpha^{-1}\|f(M_1))^{-1} \sum_{i=1}^{M_1} \|\widetilde{x_1^i}\| \ge (20\|\alpha^{-1}\|)^{-1}.$$

We may then choose a (M_1, f) -form y_1^* in $B(Z_2^{\perp})$ such that y_1^* is in \mathbf{Q} , such that $ran(y_1^*) \subset ran(\alpha(x_1))$, and such that $y_1^*(\alpha(x_1)) \geq (25 \|\alpha^{-1}\|)^{-1}$.

We then define $M_2 = \sigma_2(y_1^*)$, and we may assume we chose y_1^* such that $1/2f((M_2)^{1/40})^{1/2} \ge |ran(\widetilde{x_1})|$. We now let x_2 be a R.I.S. of length M_2 in $span\{(w_n)\}$, such that $ran(\widetilde{x_2}) > ran(\widetilde{x_1})$. Following in the same way, we obtain sequences x_i, y_i^* for $i = 1, \ldots, k$, such that the sequence y_1^*, \ldots, y_k^* is a special sequence in Z_2^{\perp} and $\widetilde{x_1}, \ldots, \widetilde{x_k}$ is associated to y_1^*, \ldots, y_k^* .

It follows that

$$\|\alpha(\sum_{i=1}^{k} x_i)\| \ge f(k)^{-1/2} \sum_{i=1}^{k} y_i^*(\alpha(x_i)) \ge (25\|\alpha^{-1}\|)^{-1} k f(k)^{-1/2},$$

while as (\tilde{x}_i) is associated to (y_i^*) , by Lemma 17,

$$\left\|\sum_{i=1}^{k} \widetilde{x}_i\right\| \le 7kf(k)^{-1},$$

and

$$\left\|\sum_{i=1}^{k} x_{i}\right\| \le 7kf(k)^{-1} + 2\Delta \le 8kf(k)^{-1},$$

It follows that $\|\alpha\| \ge (200\|\alpha^{-1}\|)^{-1}\sqrt{f(k)}$, and this for any k, contradicting the boundedness of the operator α .

7 Appendix

We give a sketch of the proof of the existence of a H.I. space \mathcal{X} such that \mathcal{X}^* contains a direct sum of n subspaces.

7.1 Proposition

For i = 1, ..., n, let X_i be a H.I. Banach space, let Z_i be a subspace of X_i . Assume that the spaces Z_i are all isometric to a same space Z, and that for any $i \neq j$, X_i/Z_i and X_j/Z_j are infinite dimensional and totally incomparable. Let $Z_{[1,n]} = \{(z_1, ..., z_n) \in Z_1 \times \cdots \times Z_n / \sum_{i=1}^n z_i = 0\}$. Let \mathcal{X} be the quotient space $(X_1 \times \cdots \times X_n)/Z_{[1,n]}$. Then \mathcal{X} is H.I. and \mathcal{X}^* contains a direct sum of n subspaces.

<u>Proof</u>

We use the same notation as in the case n = 2, in particular we let $\mathcal{Z} = \{(\widehat{z, \ldots, 0}), z \in Z\}$, and we show that

$$\mathcal{Z}^{\perp} \simeq \bigoplus_{i=1}^{n} (X_i/Z_i)^*.$$

Now we consider \mathcal{W} a subspace of \mathcal{X} . There exists at most one value i_W of i such that $\phi_{i/\mathcal{W}}$ is finitely singular, otherwise two quotient spaces X_i/Z_i and X_j/Z_j would have isomorphic subspaces.

It follows easily that \mathcal{W} and \mathcal{X}_{i_W} have Id + S-isomorphic subspaces, and finally that \mathcal{X} is H.I..

7.2 Proposition

For i = 1, ..., n, there exist X_i Q.H.I. reflexive Banach space, Z_i subspace of X_i , such that all Z_i are isometric, and such that for any $i \neq j$, X_i/Z_i and X_j/Z_j are totally incomparable.

We make a construction similar to the case n = 2, using a partition of L in n subsets L_1, \ldots, L_n . We build n balanced bounded convex subsets C_1, \ldots, C_n of c_{00} , and multifunctions $a_{ij} : C_i \to C_j$ for $i \neq j$, such that for all $i \neq j$, (C_i, C_j) is an associated dual couple. The difference is that we have n - 1 kinds of shadow sequences in each C_i (those coming special sequences in C_j for all $j \neq i$). Defining a_{ij} for the new kinds of shadow sums is not difficult. The proof of the proposition then follows exactly the case n = 2.

7.3 Remark

We recall a definition from [F]: a Banach space is said to be HD_n if the maximum number of subspaces in a direct sum is finite and equal to n. Here, for i = 1, ..., n, X_i is reflexive Q.H.I., so by Proposition 2, X_i^* is H.I., so it follows from Corollary 1 of [F] that $X_1^* \oplus ... \oplus X_n^*$ is HD_n . Now \mathcal{X}^* is a subspace of $X_1^* \oplus ... \oplus X_n^*$, so it is HD_m for some $m \leq n$, and it contains the direct sum of n subspaces \mathcal{Z}^{\perp} , so m = n; but \mathcal{X}^* is not decomposable, otherwise \mathcal{X} would be decomposable. So \mathcal{X}^* is an example of a non decomposable HD_n space.

References

- [F] V.FERENCZI, Hereditarily finitely decomposable spaces, preprint.
- [GM] W.T. GOWERS and B. MAUREY, The unconditional basic sequence problem, J.Amer.Math.Soc. 6 (1993), 851-874.
- [LT] J. LINDENSTRAUSS and L.TZAFRIRI, Classical Banach spaces II, Springer-Verlag, New York (1977).