EFFECTIVE FORMULAS FOR INVARIANT FUNCTIONS – CASE OF ELEMENTARY REINHARDT DOMAINS

Peter Pflug & Włodzimierz Zwonek

ABSTRACT. In the paper we find effective formulas for the invariant functions, appearing in the theory of several complex variables, of the elementary Reinhardt domains. This gives us the first example of a large family of domains for which the functions are calculated explicitly.

0. Introduction. Holomorphically invariant functions and pseudometrics have proved to be very useful in the theory of several complex variables. Nevertheless, the problem of finding effective formulas for the objects has turned out to be very difficult. So far there have been very few examples of domains for which the formulas for these functions are known explicitly.

Among many different invariant functions and pseudometrics let us mention the Lempert and Green functions, the Kobayashi and Carathéodory pseudodistances as well as their infinitesimal versions i.e. the Kobayashi–Royden, Carathéodory and Azukawa pseudometrics.

Due to Lempert's theorem (see [L 1,2]) all holomorphically invariant functions and pseudometrics coincide in the class of convex domains, therefore, these are the non-convex domains, which may deliver us a great deal of different invariant functions, not the convex ones. But even in convex case it is difficult to find explicit formulas for the objects involved. Among few results in this direction let us mention here the special case of convex (see [BFKKMP], [JP 2]) and non-convex (see [PZ]) ellipsoids. The other class of non-convex domains for which some of the functions were calculated is a class of elementary Reinhardt domains (see [JP 1,2]). In our paper we extend the results obtained in those domains for all invariant functions and pseudometrics mentioned earlier. The formulas obtained enable us to understand better the mutual relations between the invariant objects and give surprising solutions to some problems.

1. Definitions, notations and main results. By E we will always denote the unit disc in \mathbb{C} . We put $m(\lambda_1, \lambda_2) := \frac{|\lambda_1 - \lambda_2|}{|1 - \lambda_1 \lambda_2|}$ for $\lambda_1, \lambda_2 \in E$ and $\gamma_E(\lambda; \alpha) := \frac{|\alpha|}{1 - |\lambda|^2}$, $\lambda \in E, \alpha \in \mathbb{C}$.

Typeset by \mathcal{AMS} -T_EX

The research started while the second author was visiting Universität Oldenburg, whose stay was enabled by Volkswagen Stiftung Az. I/71 062. The second author was also supported by KBN Grant No 2 PO3A 060 08

Let D be a domain in \mathbb{C}^n . Following [L 1], [Ko 1], [Kl 1,2], [C], [A], [R] and [JP2] for $(w, z) \in D$ and $(w; X) \in D \times \mathbb{C}^n$ we define the following functions:

$$\tilde{k}_D^*(w,z) := \inf\{m(\lambda_1,\lambda_2) : \exists \varphi \in \mathcal{O}(E,D), \ \varphi(\lambda_1) = w, \ \varphi(\lambda_2) = z\} \\ k_D^*(w,z) := \tanh k_D(w,z),$$

where k_D is the largest pseudodistance smaller or equal than $\tilde{k}_D := \tanh^{-1} \tilde{k}_D^*$, $g_D(w, z) := \sup\{u(z):$

 $\log u \in \mathrm{PSH}(D, [-\infty, 0)), \exists M, R > 0 : u(\zeta) \leq M ||\zeta - w|| \text{ for } \zeta \in D, \ ||\zeta - w|| < R \},$ $c_D^*(w, z) := \sup\{m(\varphi(w), \varphi(z)) : \ \varphi \in \mathcal{O}(D, E)\};$

and also their infinitesimal versions:

$$\kappa_D(w;X) := \inf\{\gamma_E(\lambda;\alpha) : \exists \varphi \in \mathcal{O}(E,D), \ \varphi(\lambda) = z, \alpha\varphi'(\lambda) = X\},\$$
$$A_D(w;X) := \limsup_{\lambda \neq 0} \frac{g_D(w,w+\lambda X)}{|\lambda|},\$$
$$\gamma_D(w;X) := \sup\{\gamma_E(\varphi(w);\varphi'(w)X) : \ \varphi \in \mathcal{O}(D,E)\}.$$

The function k_D^* (respectively, g_D , k_D^* , c_D^*) is called the Lempert function (respectively, the Green function, the Kobayashi and Carathéodory pseudodistance). The function κ_D (respectively, A_D and γ_D) is called the Kobayashi–Royden (respectively, Azukawa and Carathéodory–Reiffen) pseudometric.

Note that the functions k_D^* , k_D^* and c_D^* are always symmetric, whereas g_D need not have the property. For the basic properties of the functions defined we refer the interested reader to [JP 2]. Let us mention here only some basic relations between the objects involved:

$$\begin{split} \dot{k}_D^* \geq k_D^* \geq c_D^*, \quad \dot{k}_D^* \geq g_D \geq c_D^*, \\ \kappa_D \geq A_D \geq \gamma_D. \end{split}$$

A mapping $\varphi \in \mathcal{O}(E, D)$ is called a \tilde{k}_D -geodesic for (w, z), $w \neq z$ if $\varphi(\lambda_1) = w$, $\varphi(\lambda_2) = z$ and $m(\lambda_1, \lambda_2) = \tilde{k}_D^*(w, z)$ for suitable $\lambda_1, \lambda_2 \in E$.

The class of domains we are intersted in is defined below.

For $\alpha = (\alpha_1, \ldots, \alpha_n) \in \mathbb{R}^n_+$, n > 1, $(\mathbb{R}_+ := (0, \infty))$ define

$$D_{\alpha} := \{ z \in \mathbb{C}^n : |z_1|^{\alpha_1} \cdot \ldots \cdot |z_n|^{\alpha_n} < 1 \}.$$

We say that α is of a rational type if there are t > 0, $\beta = (\beta_1, \ldots, \beta_n) \in \mathbb{N}^n_*$ such that $\alpha = t\beta$. We say that α is of irrational type if α is not of a rational type. Remark that in case when α is of rational type we may without loss of generality assume that all α_i 's are relatively prime natural numbers. We define also

$$\hat{D}_{\alpha} := \{ z \in D_{\alpha} : z_1 \cdot \ldots \cdot z_n \neq 0 \},\$$

If $\alpha \in \mathbb{N}^n_*$, then we denote

$$z^{\alpha} := z_1^{\alpha_1} \cdot \ldots \cdot z_n^{\alpha_n}, \quad F^{\alpha}(z) := z^{\alpha},$$
$$F^{\alpha}_{(r)}(z)X := \sum_{\beta_1 + \ldots + \beta_n = r} \frac{1}{\beta_1! \cdot \ldots \cdot \beta_n!} \frac{\partial^{\beta_1 + \ldots + \beta_n} F^{\alpha}(z)}{\partial z_1^{\beta_1} \ldots \partial z_n^{\beta_n}} X^{\beta}, \quad z, X \in \mathbb{C}^n$$

Note that the domain D_{α} is always unbounded, Reinhardt, complete, and pseudoconvex but not convex.

As mentioned in Introduction some of the invariant functions for domains D_{α} are explicitly known. We gather the results known so far in the following theorem **Theorem 1** (see [JP 2]). If $\alpha \in \mathbb{N}^n_*$, where α_j 's are relatively prime, then :

$$c_{D_{\alpha}}^{*}(w,z) = m(w^{\alpha}, z^{\alpha}),$$

$$g_{D_{\alpha}}(w,z) = m(w^{\alpha}, z^{\alpha})^{\frac{1}{r}},$$

$$\gamma_{D_{\alpha}}(w;X) = \gamma_{E}(w^{\alpha}, (F^{\alpha})'(w)X),$$

$$A_{D_{\alpha}}(w;X) = \left(\gamma_{E}(w^{\alpha}, F_{(r)}^{\alpha}(w)X)\right)^{\frac{1}{r}}, \qquad (w,z) \in D_{\alpha} \times D_{\alpha}, (w;X) \in D_{\alpha} \times \mathbb{C}^{n}$$

where r is the order of vanishing of the function $F^{\alpha}(\cdot) - F^{\alpha}(w)$ at w. If α is of irrational type, then

$$\begin{aligned} c^*_{D_\alpha}(w;X) &= 0, \\ \gamma_{D_\alpha}(w;X) &= 0, \qquad (w,z) \in D \times D, (w;X) \in D \times \mathbb{C}^n. \end{aligned}$$

In our paper we extend the results of Theorem 1 to other invariant functions and pseudometrics and we find the remaining formulas for the Green function (and Azukawa pseudometric) in the irrational case. The results are presented in two theorems. One of them concerns with rational, while the other one with irrational α . In both theorems in case of the Lempert function the formulas may seem to be incomplete (not all the cases are covered); nevertheless, because of the symmetry of both functions one easily obtains the formulas in remaining cases.

Theorem 2. Assume that $\alpha \in \mathbb{N}^n_*$ with α_j 's relatively prime. Let $(w, z) \in D_\alpha \times D_\alpha$, $(w; X) \in D_\alpha \times \mathbb{C}^n$. Denote $\mathcal{J} := \{j \in \{1, \ldots, n\} : w_j = 0\} = \{j_1, \ldots, j_k\}$. Then we have

$$\tilde{k}_{D_{\alpha}}^{*}(w,z) = \begin{cases} \min\{m(\lambda_{1},\lambda_{2}):\lambda_{1},\lambda_{2} \in E, \lambda_{1}^{\min\{\alpha_{j}\}} = w^{\alpha}, \lambda_{2}^{\min\{\alpha_{j}\}} = z^{\alpha}\}, & \text{if } w, z \in \tilde{D}_{\alpha} \\ |z^{\alpha}|^{\frac{1}{\alpha_{j_{1}}+\ldots+\alpha_{j_{k}}}}, & \text{if } \mathcal{J} \neq \emptyset \end{cases};$$

$$k_{D_{\alpha}}^{*}(w,z) = \min\{m((w^{\alpha})^{\frac{1}{\min\{\alpha_{j}\}}}, (z^{\alpha})^{\frac{1}{\min\{\alpha_{j}\}}})\}.$$

where the minimum is taken over all possible roots;

in the infinitesimal case we have

$$\kappa_{D_{\alpha}}(w;X) = \begin{cases} \gamma_{E}((w^{\alpha})^{\frac{1}{\min\{\alpha_{k}\}}}, (w^{\alpha})^{\frac{1}{\min\{\alpha_{k}\}}} \frac{1}{\min\{\alpha_{k}\}} \sum_{j=1}^{n} \frac{\alpha_{j}X_{j}}{w_{j}}), & \text{if } \mathcal{J} = \emptyset, \\ (|w_{1}|^{\alpha_{1}} \cdots |X_{j_{1}}|^{\alpha_{j_{1}}} \cdots |X_{j_{k}}|^{\alpha_{j_{k}}} \cdots |w_{n}|^{\alpha_{n}})^{\frac{1}{\alpha_{j_{1}}+\cdots+\alpha_{j_{k}}}}, & \text{if } \mathcal{J} \neq \emptyset \end{cases}$$

Observe that if $\min\{\alpha_j\} = 1$, then $\tilde{k}^*_{D_\alpha}(w, z) = g_{D_\alpha}(w, z)$ for $w, z \in \tilde{D}_\alpha$; otherwise, if $w^\alpha \neq z^\alpha$, then the Green function is strictly less than the Lempert function.

In the irrational case unlike in the rational one, these are not only the Lempert function, Kobayashi pseudodistance and Kobayashi–Royden pseudometric, which have not been calculated so far but also the Green function and the Azukawa pseudometric. **Theorem 3.** Assume that α is of irrational type. Let $(w, z) \in D_{\alpha} \times D_{\alpha}$, $(w; X) \in D_{\alpha} \times \mathbb{C}^{n}$. Denote $\mathcal{J} := \{j \in \{1, \ldots, n\} : w_{j} = 0\} = \{j_{1}, \ldots, j_{k}\}$. Then we have

$$\tilde{k}_{D_{\alpha}}^{*}(w,z) = \begin{cases} m((|w_{1}|^{\alpha_{1}} \cdot \ldots \cdot |w_{n}|^{\alpha_{n}})^{\frac{1}{\min\{\alpha_{j}\}}}, (|z_{1}|^{\alpha_{1}} \cdot \ldots \cdot |z_{n}|^{\alpha_{n}})^{\frac{1}{\min\{\alpha_{j}\}}}), & \text{if } w, z \in \tilde{D}_{\alpha} \\ (|z_{1}|^{\alpha_{1}} \cdot \ldots \cdot |z_{n}|^{\alpha_{n}})^{\frac{1}{\alpha_{j_{1}}+\ldots+\alpha_{j_{k}}}}, & \text{if } \mathcal{J} \neq \emptyset \end{cases};$$

$$k_{D_{\alpha}}^{*}(w,z) = m\left(\left(\prod_{j=1}^{n} |w_{j}|^{\alpha_{j}}\right)^{\frac{1}{\min\{\alpha_{j}\}}}, \left(\prod_{j=1}^{n} |z_{j}|^{\alpha_{j}}\right)^{\frac{1}{\min\{\alpha_{j}\}}}\right),$$
$$g_{D_{\alpha}}(w,z) = \begin{cases} 0, & \text{if } \mathcal{J} = \emptyset, \\ (|z_{1}|^{\alpha_{1}} \cdot \ldots \cdot |z_{n}|^{\alpha_{n}})^{\frac{1}{\alpha_{j_{1}} + \ldots + \alpha_{j_{k}}}}, & \text{if } \mathcal{J} \neq \emptyset \end{cases};$$

in the infinitesimal case we have:

$$\kappa_{D_{\alpha}}(w;X) = \left\{ \begin{array}{l} \gamma_{E}(\left(\prod_{j=1}^{n} |w_{j}|^{\alpha_{j}}\right)^{\frac{1}{\min\{\alpha_{k}\}}}, \left(\prod_{j=1}^{n} |w_{j}|^{\alpha_{j}}\right)^{\frac{1}{\min\{\alpha_{k}\}}} \frac{1}{\min\{\alpha_{k}\}} \sum_{j=1}^{n} \frac{\alpha_{j}X_{j}}{w_{j}}) \quad , \text{if } \mathcal{J} = \emptyset, \\ (|w_{1}|^{\alpha_{1}} \cdot \ldots \cdot |X_{j_{1}}|^{\alpha_{j_{1}}} \cdot \ldots \cdot |X_{j_{k}}|^{\alpha_{j_{k}}} \cdot \ldots \cdot |w_{n}|^{\alpha_{n}})^{\frac{1}{\alpha_{j_{1}}+\ldots+\alpha_{j_{k}}}} \quad , \text{if } \mathcal{J} \neq \emptyset \end{array} \right\}$$

$$A_{D_{\alpha}}(w;X) =$$

$$\begin{cases} 0 , & \text{if } \mathcal{J} = \emptyset, \\ (|w_1|^{\alpha_1} \cdot \ldots \cdot |X_{j_1}|^{\alpha_{j_1}} \cdot \ldots \cdot |X_{j_k}|^{\alpha_{j_k}} \cdot \ldots \cdot |w_n|^{\alpha_n})^{\frac{1}{\alpha_{j_1} + \ldots + \alpha_{j_k}}} , & \text{if } \mathcal{J} \neq \emptyset \end{cases}.$$

Observe that for an arbitrary balanced pseudoconvex domain D we always have that $\tilde{k}_D^*(0,z) = h_D(z), z \in D$, where h_D denotes the Minkowski function for D. In the above formula we have that $k_{D_\alpha}^*(0,z) < h_{D_\alpha}(z), 0 \neq z \in D_\alpha$. It would be interesting to find the general form of $k_D^*(0,\cdot)$ in the case when D is an arbitrary balanced pseudoconvex domain.

2. Auxiliary results.

For
$$z \in \mathbb{C}^n$$
 put

$$T_z := \{ (e^{i\theta_1} z_1, \dots, e^{i\theta_n} z_n) : \theta_j \in \mathbb{R} \}.$$

Note that T_z is a group with the multiplication defined as follows:

$$(e^{i\theta_1}z_1,\ldots,e^{i\theta_n}z_n)\circ(e^{i\tilde{\theta}_1}z_1,\ldots,e^{i\tilde{\theta}_n}z_n):=(e^{i(\theta_1+\tilde{\theta}_1)}z_1,\ldots,e^{i(\theta_n+\tilde{\theta}_n)}z_n).$$

Define $T_{z,\alpha}$ as the subgroup of T_z generated by the set

$$\{(e^{i\frac{\alpha_{j_1}}{\alpha_1}2k_1\pi}z_1,\ldots,e^{i\frac{\alpha_{j_n}}{\alpha_n}2k_n\pi}z_n): j_1,\ldots,j_n \in \{1,\ldots,n\}, k_1,\ldots,k_n \in \mathbb{Z}\}$$

Note that if α is of a rational type, then $T_{z,\alpha}$ is finite; more precisely, if we assume that $\alpha \in \mathbb{N}^n_*$ and α_j 's are relatively prime, then

$$T_{z,\alpha} = \{ (\varepsilon_1 z_1, \ldots, \varepsilon_n z_n), \text{ where } \varepsilon_i^{\alpha_j} = 1 \}.$$

However, if α is of irrational type, then a well-known theorem of Kronecker (see [HW], theorem 439) gives that

(1)
$$\overline{T_{z,\alpha}} = T_z.$$

For $\mu \in E_*$ we define

$$\Phi_{\mu}: \mathbb{C}^{n-1} \ni (\lambda_1, \dots, \lambda_{n-1}) \to (e^{\alpha_n \lambda_1}, \dots, e^{\alpha_n \lambda_{n-1}}, \mu e^{-\alpha_1 \lambda_1} \cdot \dots \cdot e^{-\alpha_{n-1} \lambda_{n-1}}) \in D_{\alpha}$$

 Put

$$V_{\mu} := \Phi_{\mu}(\mathbb{C}^{n-1}), \ \mu \in E_*, \quad V_0 := \{ z \in \mathbb{C}^n : z_1 \cdot \ldots \cdot z_n = 0 \}.$$

Note that

$$\bigcup_{\mu \in E} V_{\mu} = D_{\alpha}.$$

Remark 4. Let $\mu \in E_*$. Assume that $w, z \in V_{\mu}$, and $X \in \mathbb{C}^n$ fulfis the equality $\sum_{j=1}^n \frac{\alpha_j X_j}{w_j} = 0$. Then

$$\tilde{k}_{D_{\alpha}}^{*}(w,z) = 0,$$

$$\kappa_{D_{\alpha}}(w;X) = 0.$$

In fact, $w = \Phi_{\mu}(\lambda)$, $z = \Phi_{\mu}(\gamma)$ for some $\lambda, \gamma \in \mathbb{C}^{n-1}$, so

$$\tilde{k}_{D_{\alpha}}^{*}(w,z) = \tilde{k}_{D_{\alpha}}^{*}(\Phi_{\mu}(\lambda), \Phi_{\mu}(\gamma)) \leq \tilde{k}_{\mathbb{C}^{n-1}}^{*}(\lambda,\gamma) = 0.$$

To see the second equality note that assuming $\Phi_{\mu}(\lambda) = w$ we have

$$\Phi'_{\mu}(\lambda)(Y) = \left[\alpha_n w_1 Y_1, \dots, \alpha_n w_{n-1} Y_{n-1}, -\sum_{j=1}^{n-1} \alpha_j w_n Y_j\right], \quad Y \in \mathbb{C}^{n-1}.$$

One may easily verify that

$$\Phi'_{\mu}(\lambda)(\mathbb{C}^{n-1}) = \left\{ X \in \mathbb{C}^n : \sum_{j=1}^n \frac{\alpha_j X_j}{w_j} = 0 \right\}.$$

Note that

$$0 = \kappa_{\mathbb{C}^{n-1}}(\lambda; Y) \ge \kappa_{D_{\alpha}}(\Phi_{\mu}(\lambda), \Phi'_{\mu}(\lambda)Y), \quad Y \in \mathbb{C}^{n-1}$$

which finishes the proof.

In the proof of Lemma 5 below we shall replace E in the definition of the Lempert function with $H := \{x + iy : 1 > x > -1\}.$

Lemma 5. Fix $w, z \in D_{\alpha}$. Take any $\tilde{z} \in T_{z,\alpha}$. Then for any $\varphi \in \mathcal{O}(E, D_{\alpha})$ such that $\varphi(\lambda_1) = w$, $\varphi(\lambda_2) = z$, $\lambda_1 \neq \lambda_2$ there is $\tilde{\varphi} \in \mathcal{O}(E, D_{\alpha})$ such that $\tilde{\varphi}(\lambda_1) = w$ and $\tilde{\varphi}(\lambda_2) = \tilde{z}$.

Consequently,

$$\tilde{k}^*_{D_{\alpha}}(w,z) = \tilde{k}^*_{D_{\alpha}}(w,\tilde{z}) \text{ for any } \tilde{z} \in T_{z,\alpha}.$$

Proof. For the proof of the lemma it is enough to take any mapping

$$\varphi \in \mathcal{O}(H, D_{\alpha}), \quad \varphi(0) = w, \quad \varphi(it) = z, \ t > 0.$$

Define the mapping $(k_n \in \mathbb{Z} \text{ is fixed})$

$$\tilde{\varphi}: H \ni \lambda \to (\varphi_1(\lambda), \dots, \varphi_{n-2}(\lambda), e^{-2k_n \pi \frac{\lambda}{t}} \varphi_{n-1}(\lambda), e^{\frac{\alpha_{n-1} 2k_n \pi \lambda}{\alpha_n t}} \varphi_n(\lambda)) \in D_{\alpha}.$$

We have

$$\tilde{\varphi}(0) = w, \quad \tilde{\varphi}(it) = (z_1, \dots, z_{n-1}, e^{i\frac{\alpha_{n-1}}{\alpha_n}2k_n\pi}z_n)$$

Note that we may replace α_{n-1} above with any other α_j and z_n with $e^{i\frac{\alpha_j}{\alpha_n}2k_n\pi}z_n$, and also we may continue the procedure as above with the next components z_j to be varied, which would finish the proof. \Box

Remark 6. From the proof of Lemma 5 we have also the following property:

Fix $\alpha \in \mathbb{N}_{*}^{n}$, α_{j} 's relatively prime and $0 < \delta_{1} \leq m(\lambda_{1}, \lambda_{2}) \leq \delta_{2} < 1$. Take any $\psi \in \mathcal{O}(E, \mathbb{C}^{n}), \ \psi(E) \subset \subset (\mathbb{C}_{*})^{n}$ and choose $z \in \mathbb{C}^{n}$ such that $z_{j}^{\alpha_{j}} = \psi_{j}^{\alpha_{j}}(\lambda_{2})$, for $j = 1, \ldots, n$. Then there is a mapping $\tilde{\psi} \in \mathcal{O}(E, \mathbb{C}^{n})$ such that $\tilde{\psi}(E) \subset \subset (\mathbb{C}_{*})^{n}$, $\psi(\lambda_{1}) = \tilde{\psi}(\lambda_{1}), \ \tilde{\psi}(\lambda_{2}) = z$ and

$$\begin{split} \psi_1^{\alpha_1}(\lambda) \cdot \ldots \cdot \psi_n^{\alpha_n}(\lambda) &= \tilde{\psi}_1^{\alpha_1}(\lambda) \cdot \ldots \cdot \tilde{\psi}_n^{\alpha_n}(\lambda), \quad \lambda \in E \\ m ||\psi_j||_E &\le ||\tilde{\psi}_j||_E \le M ||\psi_j||_E, \quad j = 1, \dots, n \end{split}$$

where m, M > 0 depend only on δ_1 and α .

Lemma 7. Fix $L_1^1, L_1^2 \subset \mathbb{C}$, $L_2 \subset \mathbb{C}_*$ and $\alpha \in (\mathbb{R}_+)^n$. Assume that there is $\delta > 0$ such that for any $\lambda_1 \in L_1^1$, $\lambda_2 \in L_1^2$ we have $m(\lambda_1, \lambda_2) \geq \delta$.

Then there is $L_2 \subset K \subset \mathbb{C}_*$ such that for any $z_1, z_2 \in L_2$ and for any $\lambda_1 \in L_1^1$, $\lambda_2 \in L_1^2$ there is $\psi \in \mathcal{O}(E, \mathbb{C}_*)$ with $\psi(\lambda_j) = z_j$, j = 1, 2, and $\psi(E) \subset K$.

Moreover, there is $\tilde{K} \subset \mathbb{C}_*$ such that for any numbers $z_1, \ldots, z_n \in L_2$, $w_1, \ldots, w_k \in L_2$, k < n with

$$|z_1|^{\alpha_1}\cdot\ldots\cdot|z_n|^{\alpha_n}=1$$

there are functions

$$\psi_j \in \mathcal{O}(E, \mathbb{C}_*), \quad \psi_j(E) \subset K, \ j = 1, \dots, n, \quad \psi_1^{\alpha_1}(\lambda) \cdot \dots \cdot \psi_n^{\alpha_n}(\lambda) = e^{i\theta}, \ \lambda \in E,$$
$$\psi_j(\lambda_1) = z_j, \ j = 1, \dots, n, \quad \psi_j(\lambda_2) = w_j, \ j = 1, \dots, k.$$

Proof. For the proof of the first part of the lemma it is sufficient to prove it for $L_1^1 = \{\lambda_1\}, L_1^2 = \{\lambda_2\}$ with $m(\lambda_1, \lambda_2) = \delta$. This is so because the general case one obtains from that special one by composing the functions with automorphisms of E and dilatation $R\lambda$, where $0 \le R < 1$ and as we see the images of new functions are contained in that of the starting one.

Define

$$L := \exp^{-1}(L_2) \cap (\mathbb{R} \times [0, 2\pi)) \subset \subset \mathbb{C}.$$

Now put

$$K := \{ \exp(h(\lambda)) : \lambda \in E \}$$

and h is of the type $h(\lambda) = a\lambda + b$, $a, b \in \mathbb{C}$, $h(\lambda_1) = \tilde{z}_1$, $h(\lambda_2) = \tilde{z}_2$, $\tilde{z}_1, \tilde{z}_2 \in L$.

Observe that $K \subset \mathbb{C}_*$. The mappings we are looking for are of the form $\exp \circ h$, where h is one of the functions appearing in the definition of K.

For the proof of the second part of the lemma we put w_j for j = k + 1, ..., n - 1as any number from L_2 and we take mappings $\psi_1, ..., \psi_{n-1}$ as in the first part of the lemma. Define

$$\psi_n(\lambda) := \frac{e^{i\theta}}{(\psi_1^{\alpha_1}(\lambda) \cdot \ldots \cdot \psi_{n-1}^{\alpha_{n-1}}(\lambda))^{1/\alpha_n}},$$

where the branches of powers are chosen arbitrarily and $\tilde{\theta} \in \mathbb{R}$ is chosen so that $\psi_n(\lambda_1) = z_n$. \Box

Lemma 8. Let $L_1^1, L_1^2, L_2, \delta$ be as in Lemma 7. Fix $\alpha \in \mathbb{N}_*^n$, where α_j 's are relatively prime. Then there is $K \subset \mathbb{C}_*$ such that for any mappings $\psi_j \in \mathcal{O}(E, \mathbb{C}_*)$, $j = 1, \ldots, n$ with

$$\psi_1^{\alpha_1} \cdot \ldots \cdot \psi_n^{\alpha_n} = 1, \quad \lambda \in E$$

and $\psi_j(\lambda_1), \psi_j(\lambda_2) \in L_2$, where $\lambda_1 \in L_1^1, \lambda_2 \in L_1^2$ there are functions $\tilde{\psi}_j \in \mathcal{O}(E, \mathbb{C}_*)$ such that

$$\psi_1^{\alpha_1} \cdot \ldots \cdot \psi_n^{\alpha_n} = 1, \quad \lambda \in E,$$

$$\tilde{\psi}_j(\lambda_1) = \psi_j(\lambda_1), \quad \tilde{\psi}_j(\lambda_2) = \psi_j(\lambda_2), \quad \tilde{\psi}_j(E) \subset K, \quad j = 1, \dots, n.$$

Proof. For the proof put $z_j := \psi_j(\lambda_1), w_j := \psi_j(\lambda_2), j = 1, ..., n$. From Lemma 7 there are $\tilde{\psi}_j, j = 1, ..., n - 1$ as desired. Put

$$\tilde{\psi}_n(\lambda) := \frac{1}{(\tilde{\psi}_1^{\alpha_1}(\lambda) \cdot \ldots \cdot \tilde{\psi}_{n-1}^{\alpha_{n-1}}(\lambda))^{1/\alpha_n}}$$

We choose the branch of the power $\frac{1}{\alpha_n}$ so that $\tilde{\psi}_n(\lambda_1) = z_n$, note also that $\tilde{\psi}_n^{\alpha_n}(\lambda_2) = w_n^{\alpha_n}$, from Remark 6 we may change $\tilde{\psi} := (\tilde{\psi}_1, \ldots, \tilde{\psi}_n)$ so that all the desired properties are preserved and, additionally, $\tilde{\psi}_n(\lambda_2) = w_n$. \Box

Below we present a lemma, which is a weaker infinitesimal version of Lemma 7. Lemma 9. Let $w \in \mathbb{C}_*$, $X \in \mathbb{C}$ and $\lambda_1 \in E$. Then there is a mapping $\psi \in \mathcal{O}(E, \mathbb{C}_*)$ such that

$$\psi(\lambda_1) = w, \quad \psi'(\lambda_1) = X.$$

Moreover, for given numbers $w_1, \ldots, w_n \in \mathbb{C}_*$, $X_1, \ldots, X_k \in \mathbb{C}$ (k < n) and $\alpha \in (\mathbb{R}_+)^n$, where $|w_1|^{\alpha_1} \cdot \ldots \cdot |w_n|^{\alpha_n} = 1$ there are mappings $\psi_j \in \mathcal{O}(E, \mathbb{C}_*)$, $j = 1, \ldots, n$ such that

$$\psi_j(\lambda_1) = w_j, \ j = 1, \dots, n, \quad \psi'_j(\lambda_1) = X_j, \ j = 1, \dots, k, \ and$$
$$\psi_1^{\alpha_1}(\lambda) \cdot \dots \cdot \psi_n^{\alpha_n}(\lambda) = e^{i\theta}, \quad \lambda \in E.$$

Proof. The first part goes as in the proof of Lemma 7 (note that we do not need to specify more, since we do not demand so much about the mapping ψ as in Lemma 7). The mapping we are looking for is of the form $\exp(a\lambda + b)$.

For the second part of the lemma put X_j as any number from \mathbb{C} $(j = k + 1, \ldots, n-1)$. Take ψ_j as given in the first part of the lemma (for $j = 1, \ldots, n-1$) with w replaced with w_j and X replaced with X_j . Put

$$\psi_n(\lambda) := \frac{e^{i\bar{\theta}}}{(\psi_1^{\alpha_1}(\lambda) \cdot \ldots \cdot \psi_{n-1}^{\alpha_{n-1}}(\lambda))^{1/\alpha_n}},$$

where the branches of powers are chosen arbitrarily and $\tilde{\theta} \in \mathbb{R}$ is chosen so that $\psi_n(\lambda_1) = w_n$. \Box

Now we are able to give formulas for the Lempert function and the Kobayashi–Royden metric for special points.

Lemma 10. Fix $w \in V_0$. Let $z \in D_\alpha$ and $X \in \mathbb{C}^n$. Then

$$\tilde{k}_{D_{\alpha}}^{*}(w,z) = \left(|z_{1}|^{\alpha_{1}} \cdot \ldots \cdot |z_{n}|^{\alpha_{n}}\right)^{\frac{1}{\alpha_{j_{1}}+\ldots+\alpha_{j_{k}}}},$$
$$\kappa_{D_{\alpha}}(w;X) = \left(|w_{1}|^{\alpha_{1}} \cdot \ldots \cdot |X_{j_{1}}|^{\alpha_{j_{1}}} \cdot \ldots \cdot |X_{j_{k}}|^{\alpha_{j_{k}}} \cdot \ldots \cdot |w_{n}|^{\alpha_{n}}\right)^{\frac{1}{\alpha_{j_{1}}+\ldots+\alpha_{j_{k}}}}$$

where $\mathcal{J} := \{j \in \{1, \ldots, n\} : w_j = 0\} = \{j_1, \ldots, j_k\}.$

Proof. Without loss of generality we may assume that $w_1 = \ldots = w_k = 0$, $w_{k+1}, \ldots, w_n \neq 0$, $n \geq k \geq 1$. We prove both equalities simultaneously.

First we consider the case

$$z \in \tilde{D}_{\alpha}$$
 (respectively, $X_j \neq 0$ for any $j = 1, \ldots, k$).

Take any $\varphi \in \mathcal{O}(\bar{E}, D_{\alpha})$ such that

$$\varphi(0) = w, \quad \varphi(t) = z \text{ (respectively, } \varphi(0) = w, \quad t\varphi'(0) = X), \text{ for some } t > 0.$$

We have that

$$\varphi(\lambda) = (\lambda \psi_1(\lambda), \dots, \lambda \psi_k(\lambda), \psi_{k+1}(\lambda), \dots, \psi_n(\lambda)), \quad \psi_j \in \mathcal{O}(\bar{E}, \mathbb{C}), \ j = 1, \dots, n.$$

Put

$$u(\lambda) := \prod_{j=1}^{n} |\psi_j(\lambda)|^{\alpha_j}.$$

We know that $\log u \in SH(\overline{E})$ and $u \leq 1$ on ∂E , so the maximum principle for subharmonic functions implies that $u \leq 1$ on E. In particular, $u(t) \leq 1$ (respectively, $u(0) \leq 1$), so

$$\frac{\prod_{j=1}^{n} |z_j|^{\alpha_j}}{t^{\alpha_1 + \dots + \alpha_k}} \le 1, \text{ (respectively, } \frac{\prod_{j=1}^{k} |X_j|^{\alpha_j} \prod_{j=k+1}^{n} |w_j|^{\alpha_j}}{t^{\alpha_1 + \dots + \alpha_k}} \le 1 \text{)},$$

which gives us the inequality

$$t \ge \left(\prod_{j=1}^n |z_j|^{\alpha_j}\right)^{\frac{1}{\alpha_1 + \ldots + \alpha_k}}, \text{ (respectively, } t \ge \left(\prod_{j=1}^k |X_j|^{\alpha_j} \prod_{j=k+1}^n |w_j|^{\alpha_j}\right)^{\frac{1}{\alpha_1 + \ldots + \alpha_k}}).$$

Therefore,

$$\tilde{k}_{D_{\alpha}}^{*}(w,z) \geq \left(\prod_{j=1}^{n} |z_{j}|^{\alpha_{j}}\right)^{\frac{1}{\alpha_{1}+\ldots+\alpha_{k}}},$$
(respectively, $\kappa_{D_{\alpha}}(w;X) \geq \left(\prod_{j=1}^{k} |X_{j}|^{\alpha_{j}} \prod_{j=k+1}^{n} |w_{j}|^{\alpha_{j}}\right)^{\frac{1}{\alpha_{1}+\ldots+\alpha_{k}}}$).

To get above the equality put

$$t := \left(\prod_{j=1}^{n} |z_j|^{\alpha_j}\right)^{\frac{1}{\alpha_1 + \ldots + \alpha_k}},$$
(respectively, $t := \left(\prod_{j=1}^{k} |X_j|^{\alpha_j} \prod_{j=k+1}^{n} |w_j|^{\alpha_j}\right)^{\frac{1}{\alpha_1 + \ldots + \alpha_k}}$)

and let us consider the following mapping:

$$\varphi(\lambda) := (\lambda \psi_1(\lambda), \dots, \lambda \psi_k(\lambda), \psi_{k+1}(\lambda), \dots, \psi_n(\lambda)), \quad \lambda \in E,$$

where $\psi_j \in \mathcal{O}(E, \mathbb{C}_*), j = 1, \dots, n, \prod_{j=1}^n \psi_j(\lambda)^{\alpha_j} = e^{i\theta}$ on E and

$$\begin{split} \psi_j(t) &= z_j/t, \ j = 1, \dots, k, \quad \psi_j(t) = z_j, \ j = k+1, \dots, n; \\ \psi_j(0) &= w_j, \ j = k+1, \dots, n, \ \text{(see Lemma 7)}, \\ \text{(respectively, } \psi_j(0) &= \frac{X_j}{t}, \ j = 1, \dots, k, \ \psi_j(0) = w_j, \ j = k+1, \dots, n, \\ \psi'_j(0) &= \frac{X_j}{t}, \ j = k+1, \dots, n - \text{see Lemma 9} \ \text{)}. \end{split}$$

Then $\varphi \in \mathcal{O}(E, D_{\alpha}), \, \varphi(0) = w, \, \varphi(t) = z$ (respectively, $t\varphi'(0) = X$), which finishes that case.

We are remained with the case $z \in V_0$ (respectively, $X_j = 0$ for some $1 \le j \le k$). If there is j such that $w_j = z_j = 0$ (respectively, $w_j = X_j = 0$), then the mapping

$$\mathbb{C}^{n-1} \ni (z_1, \dots, \check{z}_j, \dots, z_n) \to (z_1, \dots, 0, \dots, z_n) \in D_{\alpha}$$

gives us the following

$$0 = \tilde{k}_{\mathbb{C}^{n-1}}^*((w_1, \dots, \check{w}_j, \dots, w_n), (z_1, \dots, \check{z}_j, \dots, z_n)) \ge \tilde{k}_{D_{\alpha}}^*(w, z),$$

(respectively,
$$0 = \kappa_{\mathbb{C}^{n-1}}((w_1, \dots, \check{w}_j, \dots, w_n); (X_1, \dots, \check{X}_j, \dots, X_n)) \ge \kappa_{D_{\alpha}}(w; X)).$$

Therefore, we are remained only with the Lempert function and then we may assume that for all j we have $|w_j| + |z_j| > 0$.

Define for fixed $\beta > 0$ the mapping $\varphi := (\varphi_1, \ldots, \varphi_n)$ as follows

$$\begin{array}{l} \text{if } w_j = 0, \, \text{then } \varphi_j(\lambda) := \frac{\lambda - \beta}{1 - \beta \lambda} \psi_j(\lambda), \\ \text{if } z_j = 0, \, \text{then } \varphi_j(\lambda) := \frac{\lambda + \beta}{1 + \beta \lambda} \psi_j(\lambda), \\ \text{if } w_j z_j \neq 0, \, \text{then } \varphi_j(\lambda) := \psi_j(\lambda), \end{array}$$

where $\psi_j \in \mathcal{O}(E, \mathbb{C}_*)$, $\prod_{j=1}^n \psi_j(\lambda)^{\alpha_j} = e^{i\theta}$ on E and $\varphi(\beta) = w$, $\varphi(-\beta) = z$ (the values of $\psi_j(\beta)$ and $\psi_j(-\beta)$ are prescribed if only $w_j z_j \neq 0$; for those j for which $w_j z_j = 0$ only one from the values $\psi_j(\beta)$ and $\psi_j(-\beta)$ is prescribed, more precisely take j_1 such that $z_{j_1} = 0$, then we define $\psi_{j_1}(-\beta)$ so that $|\psi_1(-\beta)|^{\alpha_1} \cdot \ldots |\psi_n(-\beta)|^{\alpha_n} = 1$; note also that there is j_2 such that $w_{j_2} = 0$, so $\psi_{j_2}(\beta)$ has no fixed value it is the reason why we are allowed to use Lemma 7). Note also that $\varphi \in \mathcal{O}(E, D_\alpha)$. As $\beta > 0$ may be chosen arbitrarily small this completes the proof of the lemma. \Box

In the next step we will prove a formula for the Lempert function in the special case of the domain $D_{(1,...,1)}$. Following (to some extent) the ideas from [JPZ] and [PZ] we shall propagate the formulas to the general case using a technic, which could be called a transport of geodesics. Roughly speaking, the idea relies on transporting the formulas from simpler domains to more complex ones with the help of 'good' mappings. In [JPZ] and [PZ] it was the Euclidean ball that was a model domain. In our paper it is a domain $D_{(1,...,1)}$.

Lemma 11. If $w, z \in V_0$, then

$$\tilde{k}^*_{D_{(1,\dots,1)}}(w,z) = 0.$$

Assume that $w \in D_{(1,...,1)}$. Then the following equality holds:

$$\tilde{k}^*_{D_{(1,\dots,1)}}(w,z) = m(w_1\dots w_n, z_1\dots z_n)^{1/k}$$

where

$$k := \max\{\#\{j : z_j = 0\}, 1\}$$

Proof. The first part of the lemma is a consequence of Lemma 10. Moreover, also the case $z \in V_0$ is a consequence of Lemma 10.

Consider now the case $w, z \in \tilde{D}_{(1,...,1)}$. We may assume that $w_1 \cdot \ldots \cdot w_n \neq z_1 \cdot \ldots \cdot z_n$ (the other case is covered by Remark 4).

Let us consider the following mapping (see Lemma 7):

$$\varphi(\lambda) := \left(\psi_1(\lambda), \dots, \psi_{n-1}(\lambda), e^{-i\theta}\lambda\psi_n(\lambda)\right),$$

where

$$\lambda_1 := w_1 \cdot \ldots \cdot w_n, \quad \lambda_2 := z_1 \cdot \ldots \cdot z_n,$$

$$\psi_j \in \mathcal{O}(E, \mathbb{C}_*), \ j = 1, \ldots, n, \quad \psi_1(\lambda) \cdot \ldots \cdot \psi_n(\lambda) = e^{i\theta}, \quad \lambda \in E,$$

$$\psi_j(\lambda_1) = w_j, \quad \psi_j(\lambda_2) = z_j, \quad j = 1, \ldots, n-1,$$

(using Lemma 7 we may assume even that $\psi_j(E) \subset K \subset \mathbb{C}_*, \ j = 1, \ldots, n$ – compare Remark 12).

Note that

$$\varphi \in \mathcal{O}(E, D_{(1,\dots,1)}), \quad \varphi(\lambda_1) = w, \quad \varphi(\lambda_2) = z.$$

Therefore, combining these pieces of information with the formula of the Green function for $D_{(1,...,1)}$ (see Theorem 1) we have:

$$m(w_1 \cdot \ldots \cdot w_n, z_1 \cdot \ldots \cdot z_n) \ge \tilde{k}^*_{D(1,\ldots,1)}(w,z) \ge g_{D_{(1,\ldots,1)}}(w,z) = m(w_1 \cdot \ldots \cdot w_n, z_1 \cdot \ldots \cdot z_n).$$

This completes the proof. \Box

Remark 12. From the proof of Lemma 11 we get that for any $w, z \in \tilde{D}_{(1,...,1)}$ with $w_1 \cdot \ldots \cdot w_n \neq z_1 \cdot \ldots \cdot z_n$ there is a $\tilde{k}_{D_{(1,...,1)}}$ -geodesic for (w, z), which is of the form

$$(\psi_1(\lambda),\ldots,\psi_{n-1}(\lambda),e^{i\theta}\frac{\lambda-\beta}{1-\bar{\beta}\lambda}\psi_n(\lambda))$$

with $\psi_1(\lambda) \cdot \ldots \cdot \psi_n(\lambda) = 1$ and $\psi_j(E) \subset \mathbb{C}_*$.

The domains D_{α} although very regular have not got one property, which is crucial in the theory of the holomorphically invariant functions; namely, they are not taut. Therefore, we have no certainty that they admit $\tilde{k}_{D_{\alpha}}$ -geodesics. However, as Lemma 13 will show, at least in the rational case and for points, which are 'seperated' by the Lempert function, it holds. The existence of the geodesics will play a great role in the proof of the formula for the Lempert function in the rational case.

Lemma 13. Assume that $\alpha \in \mathbb{N}^n_*$ and α_j 's are relatively prime. Let $w, z \in \tilde{D}_\alpha$, $w^{\alpha} \neq z^{\alpha}$. Then there is a bounded $\tilde{k}_{D_{\alpha}}$ -geodesic $\varphi \in \mathcal{O}(E, D_{\alpha})$ for (w, z).

Proof. We know that (see Theorem 1)

$$t := \tilde{k}_{D_{\alpha}}^*(w, z) \ge g_{D_{\alpha}}(w, z) = m(w^{\alpha}, z^{\alpha}) > 0;$$

consequently, there are mappings $\varphi^{(k)} = (\varphi_1^{(k)}, \dots, \varphi_n^{(k)}), \ k = 1, 2, \dots$ such that

$$\varphi^{(k)} \in \mathcal{O}(\bar{E}, D_{\alpha}), \ \varphi^{(k)}(0) = w, \ \varphi^{(k)}(t_k) = z, \text{ where } t_k \ge t_{k+1} \to t > 0.$$

We have

$$\varphi_j^{(k)} = B_j^{(k)} \psi_j^{(k)}, \quad j = 1, \dots, n,$$

where $B_j^{(k)}$ is a Blaschke product and $\psi_j^{(k)} \in \mathcal{O}(E, \mathbb{C}_*)$.

Put $\psi^{(k)} := (\psi_j^{(k)})_{j=1}^n$. There are two possibilities (due to maximum principle of subharmonic functions – remember about the pseudoconvexity of the domain D_{α}):

(2)
$$\psi^{(k)}(E) \subset D_{\alpha},$$

(3)
$$\psi^{(k)}(E) \subset \partial D_{\alpha}.$$

Below we shall prove that without loss of generality we may reduce our attention only to the case, which is some kind of generalization of (3).

Take any k such that (2) is fulfilled. First, notice that the mapping $\tilde{\psi}^{(k)} := ((\psi_1^{(k)})^{\frac{\alpha_1}{\alpha_1 \cdots \alpha_n}}, \dots, (\psi_n^{(k)})^{\frac{\alpha_n}{\alpha_1 \cdots \alpha_n}})$ is a mapping from $\mathcal{O}(E, D_{(1,\dots,1)})$. From Remark 12 there is a $\tilde{k}_{D_{(1,\dots,1)}}$ -geodesic for $(\tilde{\psi}^{(k)}(0), \tilde{\psi}^{(k)}(t_k))$ of the form $\mu^{(k)} := (\hat{\psi}_1^{(k)}, \dots, \hat{\psi}_{n-1}^{(k)}, e^{i\theta_k} \frac{\lambda - \beta_k}{1 - \beta_k \lambda} \hat{\psi}_n^{(k)})$, where $\hat{\psi}_1^{(k)} \cdots \hat{\psi}_n^{(k)} = 1$ on E such that $\mu^{(k)}(0) = \tilde{\psi}^{(k)}(0)$ and $\mu^{(k)}(R_k t_k) = \tilde{\psi}^{(k)}(t_k), \beta_k \in E, R_k \leq 1$.

Coming back to the domain D_{α} we see that instead of considering $\varphi^{(k)}$ with the property (2) we may consider the mapping (note that $\frac{\alpha_1 \cdots \alpha_n}{\alpha_i} \in \mathbb{N}$)

$$\tilde{\varphi}^{(k)}(\lambda) := \left(B_j^{(k)}(\lambda)(\mu_j^{(k)})^{\frac{\alpha_1 \cdots \alpha_n}{\alpha_j}}(R_k\lambda)\right)_{j=1}^n$$

because $\tilde{\varphi}^{(k)} \in \mathcal{O}(E, D_{\alpha}), \, \tilde{\varphi}^{(k)}(0) = w \text{ and } \tilde{\varphi}^{(k)}(t_k) = z.$

Therefore we may assume that (irrespective of which case we start (2) or (3))

$$\varphi_j^{(k)} = B_j^{(k)} \psi_j^{(k)}, \quad j = 1, \dots, n,$$

where $(\psi_1^{(k)})^{\alpha_1} \cdot \ldots \cdot (\psi_n^{(k)})^{\alpha_n} = 1$ and $|B_j^{(k)}| \leq 1$ (although we have no longer that $B_j^{(k)}$'s are the Blaschke products).

Choosing, if necessary, a subsequence we may assume that for all $j = 1, \ldots, n$ $\{B_j^{(k)}\}_{k=1}^{\infty}$ converges locally uniformly on E. Keeping in mind that $\varphi^{(k)}(0) = w$ and $\varphi^{(k)}(t_k) = z$ we have in view of Lemma 8 that there is $K \subset \mathbb{C}$ such that we may assume that $\psi_j^{(k)}(E) \subset K$ for any j, k (we may apply Lemma 8 because $L_2 := \{\psi_j^{(k)}(t_k), \psi_j^{(k)}(0)\}_{j,k} \subset \mathbb{C}_*$, which follows from convergence and boundedness of $\{B_j^{(k)}\}_{k=1}^{\infty}$, the fact that $w_j z_j \neq 0, j = 1, \ldots, n$ and the equality $(\psi_1^{(k)})^{\alpha_1} \cdot \ldots \cdot (\psi_n^{(k)})^{\alpha_n} = 1)$, and then choosing, if necessary, a subsequence we get that the sequence $\varphi^{(k)}$ is convergent to a mapping $\varphi \in \mathcal{O}(E, \bar{D}_{(\alpha)})$ with $\varphi(E) \subset \subset (\mathbb{C}_*)^n$ such that $\varphi(0) = w, \varphi(t) = z$. The maximum principle for subharmonic functions implies, however, that $\varphi(E) \subset D_{\alpha}$. This completes the proof of the lemma. \Box

3. The rational case – Proof of Theorem 2. In the present section we provide the proof of Theorem 2. Since the theorem consists of a number of formulas we prove them below one by one. We start with the Lempert function, which is basic in the calculation of other functions.

We begin with a formula for the Möbius function, which seems to be very probable; nevertheless, we were not able to find some references in the literature. Its proof is elementary but it needs tedious calculations, so we skip the proof.

Lemma 14. Fix $0 < s \leq 1$. Then for any $\lambda_1 \in (0,1)$, $\lambda_2 \in E$ we have

$$m(\lambda_1^s, \lambda_2^s) \ge m(\lambda_1, \lambda_2),$$

where $\lambda_1^s \in (0,1)$ and the power λ_2^s is chosen so that the left-hand side of the formula is smallest possible.

Proof of formula for $\tilde{k}_{D_{\alpha}}^{*}$ in rational case. The case $w_{1} \cdot \ldots \cdot w_{n} = 0$ is a consequence of Lemma 10. The case $w, z \in \tilde{D}_{\alpha}, w^{\alpha} = z^{\alpha}$ follows from Remark 4. We are

remained with the case $w, z \in D_{\alpha}, w^{\alpha} \neq z^{\alpha}$. Due to Lemma 13 there is a bounded $\tilde{k}_{D_{\alpha}}$ -geodesic $\varphi \in \mathcal{O}(E, D_{\alpha})$ for $(w, z) = (\varphi(\lambda_1), \varphi(\lambda_2))$. Proceeding as in the proof of Lemma 13 we may assume that

$$\varphi_j = B_j \psi_j, \quad j = 1, \dots, n,$$

where B_j is the Blaschke product (up to a constant $|c_j| = 1$), $\psi_j(E) \subset K \subset \mathbb{C}_*$ and $\psi_1^{\alpha_1} \cdot \ldots \cdot \psi_n^{\alpha_n} = 1$. In fact, let us consider the decomposition of φ_j as above with the Blaschke product B_j . Put

$$\tilde{\psi} := ((\psi_j)^{\frac{\alpha_j}{\alpha_1 \cdots \alpha_n}})_{j=1}^n.$$

Consider two cases. If $\psi_1^{\alpha_1} \cdot \ldots \cdot \psi_n^{\alpha_n}$ is not constant on E, then $\tilde{\psi} \in \mathcal{O}(E, D_{(1,\ldots,1)})$ and it is a $\tilde{k}_{D_{(1,\ldots,1)}}$ -geodesic for $(\tilde{\psi}(\lambda_1), \tilde{\psi}(\lambda_2))$, otherwise, there would be $\hat{\psi} \in \mathcal{O}(E, D_{(1,\ldots,1)})$ such that $\hat{\psi}(\lambda_1) = \tilde{\psi}(\lambda_1), \hat{\psi}(\lambda_2) = \tilde{\psi}(\lambda_2)$ and $\hat{\psi}(E) \subset \subset D_{(1,\ldots,1)}$, taking then $\hat{\varphi}(\lambda) := (B_j(\lambda)\hat{\psi}_j^{\frac{\alpha_1\cdots\alpha_n}{\alpha_j}}(\lambda))_{j=1}^n$ we get a mapping such that $\hat{\varphi}(\lambda_1) = \varphi(\lambda_1), \hat{\varphi}(\lambda_2) = \varphi(\lambda_2)$ and $\hat{\varphi}(E) \subset \subset D_{\alpha}$ – contradiction. Due to Remark 12 we know that there is a $\tilde{k}_{D_{(1,\ldots,1)}}$ -geodesic for $(\tilde{\psi}(\lambda_1), \tilde{\psi}(\lambda_2)) = (\mu(\lambda_1), \mu(\lambda_2))$, where $\hat{\psi}_1 \cdot \ldots \cdot \hat{\psi}_n = 1$ and $\hat{\psi}_j(E)$'s are relatively compact in \mathbb{C}_* . Taking now $(B_j(\lambda)(\mu_j(\lambda))^{\frac{\alpha_1\cdots\alpha_n}{\alpha_j}})_{j=1}^n$ instead of φ we get the desired property.

In case $\psi_1^{\alpha_1} \cdot \ldots \cdot \psi_n^{\alpha_n} = e^{i\theta}$, we may assume that $\psi_j(E) \subset K \subset \mathbb{C}_*$ for some K because of Lemma 8 (and then without loss of generality we may assume that $e^{i\theta} = 1$).

Therefore, $\varphi(E)$ is contained in some polydisk. Consequently, $\varphi(E)$ is contained in some smooth bounded pseudoconvex complete Reinhardt domain $G \subset D_{\alpha}$, which arises from the domain D_{α} by 'cutting the ends' and 'smoothing the corners'. Therefore, φ is a \tilde{k}_G -geodesic for (w, z). Using the results of [E], [Pa] we have that there are mappings $h_j \in H^{\infty}(E, \mathbb{C}), j = 1, \ldots, n$ and $\rho : \partial E \to (0, \infty)$ such that

$$\frac{1}{\lambda}h_j^*(\lambda)\varphi_j^*(\lambda) = \rho(\lambda)\alpha_j |(\varphi^*(\lambda))^{\alpha}|, \quad j = 1, \dots, n, \text{ for almost all } \lambda \in \partial E$$

(we easily exclude the case $(\varphi^*(\lambda))^{\alpha} = 0$ for λ from some subset of ∂E with nonzero Lebesgue measure). Using the result of Gentili (see [Ge]) we get that for some $b_j \in \mathbb{C}_*, j = 1, \ldots, n, \beta \in E$,

$$\varphi_j(\lambda)h_j(\lambda) = b_j(1 - \bar{\beta}\lambda)(\lambda - \beta), \quad j = 1, \dots, n, \ \lambda \in E,$$

where $b_j/\alpha_j = b_k/\alpha_k$, j, k = 1, ..., n. Consequently, we may take

$$B_j(\lambda) = c_j \left(\frac{\lambda - \beta}{1 - \bar{\beta}\lambda}\right)^{r_j}, \quad |c_j| = 1,$$

where $r_j \in \{0, 1\}$ and not all r_j 's are equal to 0. Without loss of generality we may assume that $\beta = 0$ (we change then only λ_1 and λ_2).

Now we are coming back to the domain D_{α} . We may assume that $r_1 = \ldots = r_k = 1$ and $r_{k+1} = \ldots = r_n = 0$ $(1 \le k \le n)$. We want to have for some $\lambda_1, \lambda_2 \in E$ that (without loss of generality we may assume that $c_j = 1$ – if necessary we change

w and z with the help of rotations of suitable components, so the Lempert function does not change)

$$\lambda_1 \psi_j(\lambda_1) = w_j, \ j = 1, \dots, k, \quad \psi_j(\lambda_1) = w_j, \ j = k+1, \dots, n, \lambda_2 \psi_j(\lambda_2) = z_j, \ j = 1, \dots, k, \quad \psi_j(\lambda_2) = z_j, \ j = k+1, \dots, n.$$

Taking the α_i -th power and multiplying the equalities we get that

$$\lambda_1^{\alpha_1 + \dots + \alpha_k} = w^{\alpha}, \quad \lambda_2^{\alpha_1 + \dots + \alpha_k} = z^{\alpha}.$$

The formulas above describe all possibilities, which may deliver us the candidates for the realization of the Lempert function. Now for all possible pairs of numbers λ_1, λ_2 given as above we find mappings, which map λ_1 and λ_2 in w and z. Note that there are mappings $\psi_j \in \mathcal{O}(E, \mathbb{C}_*), j = 2, \ldots, n$ such that (see Lemma 7)

$$\psi_j(\lambda_1) = \frac{w_j}{(w^{\alpha})^{\frac{1}{\alpha_1 + \dots + \alpha_k}}} = \frac{w_j}{\lambda_1}, \quad j = 2, \dots, k,$$

$$\psi_j(\lambda_2) = \frac{z_j}{(z^{\alpha})^{\frac{1}{\alpha_1 + \dots + \alpha_k}}} = \frac{z_j}{\lambda_2}, \quad j = 2, \dots, k,$$

$$\psi_j(\lambda_1) = w_j, \quad j = k + 1, \dots, n,$$

$$\psi_j(\lambda_2) = z_j, \quad j = k + 1, \dots, n.$$

Define also

$$\psi_1(\lambda) := \frac{1}{\left(\psi_2^{\alpha_2}(\lambda) \cdot \ldots \cdot \psi_n^{\alpha_n}(\lambda)\right)^{\frac{1}{\alpha_1}}}, \quad \lambda \in E.$$

Put

$$\varphi(\lambda) := (\lambda \psi_1(\lambda), \dots, \lambda \psi_k(\lambda), \psi_{k+1}(\lambda), \dots, \psi_n(\lambda)).$$

The $\frac{1}{\alpha_1}$ -st root in definition of ψ_1 is chosen so that $\varphi_1(\lambda_1) = w_1$, and we know that $\varphi_1^{\alpha_1}(\lambda_2) = z_1^{\alpha_1}$. One may also easily verify that $\varphi(\lambda_1) = w$ and $\varphi_j(\lambda_2) = z_j$, $j = 2, \ldots, n$, which, however, in view of Lemma 5 shows that there is also a mapping $\tilde{\varphi} \in \mathcal{O}(E, D_\alpha)$ such that $\tilde{\varphi}(\lambda_1) = w$, $\tilde{\varphi}(\lambda_2) = z$. Therefore we have proved that

$$\tilde{k}_{D_{\alpha}}^{*}(w,z) = \min\{m(\lambda_{1},\lambda_{2}):\lambda_{1},\lambda_{2}\in E,\lambda_{1}^{\alpha_{j_{1}}+\ldots+\alpha_{j_{k}}}=w^{\alpha},\lambda_{2}^{\alpha_{j_{1}}+\ldots+\alpha_{j_{k}}}=z^{\alpha}\},$$

where the minimum is taken over all possible subsets $\{j_1, \ldots, j_k\} \subset \{1, \ldots, n\}$. And now Lemma 14 finishes the proof (remark that without loss of generality we may assume that $w_j > 0, j = 1, \ldots, n$). \Box

Proof of the formula for $k_{D_{\alpha}}^{*}$ in rational case. Note that \tanh^{-1} of the desired formula is equal to \tanh^{-1} of the Lempert function off the axis, satisfies the triangle inequality and is continuous. The definition of the Kobayashi pseudodistance and its continuity (see [JP 2]) finish the proof. \Box

To finish the proof we are remained only with the problem of computing the Kobayashi–Royden pseudometric $\kappa_{D_{\alpha}}$. We get that formula from that of the Kobayashi pseudodistance. But to see that we have to define an operator, which connects these both functions.

Following M. Jarnicki and P. Pflug (see [JP 2]) for a domain $D \subset \mathbb{C}^n$ we define the following function

$$\mathfrak{D}k_D(w;X) := \limsup_{\lambda \neq 0} \frac{k_{D_\alpha}^*(w, w + \lambda X)}{|\lambda|}, \quad w \in D, X \in \mathbb{C}^n.$$

The function defined above differs from that in [JP 2], nevertheless, since our version is not larger than that from [JP 2] the inequality below, which is crucial for our considerations, remains true

(4)
$$\mathfrak{D}k_D(w;X) \le \kappa_D(w;X), \quad w \in D, X \in \mathbb{C}^n$$

Lemma 15. Let $\alpha \in \mathbb{N}^n_*$, where α_j 's are relatively prime. Then

$$\mathfrak{D}k_{D_{\alpha}}(w;X) = \gamma_E\left(\left(\prod_{j=1}^n |w_j|^{\alpha_j}\right)^{\frac{1}{\min\{\alpha_k\}}}, \left(\prod_{j=1}^n |w_j|^{\alpha_j}\right)^{\frac{1}{\min\{\alpha_k\}}} \frac{1}{\min\{\alpha_k\}} \sum_{j=1}^n \frac{\alpha_j X_j}{w_j}\right)$$

 $w \in \tilde{D}_{\alpha}, X \in \mathbb{C}^n.$

Proof. Without loss of generality we may assume that $w_j > 0, j = 1, ..., n$ and $\alpha_n = \min{\{\alpha_k\}}$. Using the formula for $k_{D_{\alpha}}^*$ we get

(5)
$$\mathfrak{D}k_{D_{\alpha}}(w;X) = \limsup_{\lambda \neq 0} \frac{\left|\prod_{j=1}^{n} (w_j + \lambda X_j)^{\alpha_j/\alpha_n} - \prod_{j=1}^{n} w_j^{\alpha_j/\alpha_n}\right|}{\left|1 - \prod_{j=1}^{n} (w_j + \lambda X_j)^{\alpha_j/\alpha_n} \prod w_j^{\alpha_j/\alpha_n}\right| |\lambda|}.$$

Applying the Taylor formula we get for λ close to 0

$$(w_j + \lambda X_j)^{\alpha_j/\alpha_n} = w_j^{\alpha_j/\alpha_n} + \frac{\alpha_j}{\alpha_n} w_j^{\alpha_j/\alpha_n} \frac{\lambda X_j}{w_j} + \varepsilon_j(\lambda), \quad j = 1, \dots, n,$$

where $\frac{\varepsilon_j(\lambda)}{\lambda} \to 0$ as $\lambda \to 0$. Substituting the last equalities in (5) we get that

$$\mathfrak{D}k_{D_{\alpha}}(w;X) = \limsup_{\lambda \neq 0} \frac{\left(\prod_{j=1}^{n} |w_{j}^{\alpha_{j}}|^{1/\alpha_{n}}\right) |\lambda| \left|\sum_{j=1}^{n} \frac{\alpha_{j}X_{j}}{\alpha_{n}w_{j}}\right|}{\left(1 - \prod_{j=1}^{n} |w_{j}|^{2\alpha_{j}/\alpha_{n}}\right) |\lambda|},$$

which equals the desired value. \Box

Proof of the formula for $\kappa_{D_{\alpha}}$ in rational case. If $\mathcal{J} \neq \emptyset$, then, in view of Lemma 10 we are done. The case $\sum_{j=1}^{n} \frac{\alpha_j X_j}{w_j} = 0$ follows from Remark 4.

Take $w \in \tilde{D}_{\alpha}$. Without loss of generality we may assume that $w_j \in \mathbb{R}_+$, $j = 1, \ldots, n$ and $\alpha_n = \min\{\alpha_j\}$. Below, for $X \in \mathbb{C}^n$, $\sum_{j=1}^n \frac{\alpha_j X_j}{w_j} \neq 0$ we shall construct a mapping $\varphi \in \mathcal{O}(E, D_{\alpha})$ such that

$$\varphi(\lambda_1) = w, \quad t\varphi'(\lambda_1) = X,$$

where $\lambda_1 := (w_1^{\alpha_1} \cdot \ldots \cdot w_n^{\alpha_n})^{1/\alpha_n} > 0, t := (w_1^{\alpha_1} \cdot \ldots \cdot w_n^{\alpha_n})^{1/\alpha_n} \sum_{j=1}^n \frac{\alpha_j X_j}{\alpha_n w_j}$. Note that the existence of such a φ would finish the proof because of Lemma 15

Note that the existence of such a φ would limit the proof because of Lemma 1: and (4).

Define the mapping

$$\varphi(\lambda) := (\psi_1(\lambda), \dots, \psi_{n-1}(\lambda), \frac{\lambda}{(\psi_1^{\alpha_1}(\lambda) \cdot \dots \cdot \psi_{n-1}^{\alpha_{n-1}}(\lambda))^{1/\alpha_n}})$$

where (see Lemma 9)

$$\psi_j(\lambda_1) = w_j, \ j = 1, \dots, n-1, \quad t\psi'_j(\lambda_1) = X_j, \ j = 1, \dots, n-1.$$

We choose the $\frac{1}{\alpha_n}$ -th power so that $\varphi_n(\lambda_1) = w_n$, after some elementary transformation we get that

$$t\varphi_n'(\lambda_1) = X_n,$$

which finishes the proof. \Box

4. The irrational case – Proof of Theorem **3.** As in rational case we start with the proof of the formula of the Lempert function. First, we make use of the special properties of the domains of irrational type to get:

Lemma 16. Let α be of irrational type. Then for any $w, z \in D_{\alpha}$

$$\tilde{k}_{D_{\alpha}}^{*}(w,z) = \tilde{k}_{D_{\alpha}}^{*}(\tilde{w},\tilde{z}), \quad \tilde{w} \in T_{w}, \ \tilde{z} \in T_{z}.$$

Proof. Certainly it is enough to prove that

$$\tilde{k}_{D_{\alpha}}^{*}(w,z) = \tilde{k}_{D_{\alpha}}^{*}(w,\tilde{z}), \text{ whenever } \tilde{z} \in T_{z}$$

Assume that

(6)
$$\tilde{k}_{D_{\alpha}}^{*}(w,\tilde{z}_{1}) < \tilde{k}_{D_{\alpha}}^{*}(w,\tilde{z}_{2}) =: \varepsilon$$

for some $\tilde{z}_1, \tilde{z}_2 \in T_z$. Then in view of Lemma 5

(7)
$$\tilde{k}_{D_{\alpha}}^{*}(w,\tilde{z}) = \varepsilon$$

for all $\tilde{z} \in T_{\tilde{z}_2,\alpha}$. Because of (1) we have that $\tilde{z}_1 \in T_z = T_{\tilde{z}_2} = T_{\tilde{z}_2,\alpha}$. The last statement contradicts, in connection with (6) and (7), however, the upper-semicontinuity of the Lempert function. \Box

As an immediate corollary of Lemma 16 we get

Corollary 17. Let α be of irrational type, then for any $z \in D_{\alpha}$

$$\tilde{k}_{D_{\alpha}}^{*}(z,\tilde{z}) = 0 \text{ for any } \tilde{z} \in T_{z}.$$

Proof of the formula for $\tilde{k}_{D_{\alpha}}^{*}$ in irrational case. The case $\mathcal{J} \neq \emptyset$ is covered by Lemma 10. Consider now the remaining case. In view of Lemma 16 we have that

$$\hat{k}_{D_{\alpha}}^{*}(w,z) = \hat{k}_{D_{\alpha}}^{*}((|w_{1}|,\ldots,|w_{n}|),(|z_{1}|,\ldots,|z_{n}|)).$$

16

Let us choose a sequence $\{\alpha^{(k)}\}_{k=1}^{\infty} \subset (\mathbb{Q}_+)^n$ such that

$$\alpha^{(k)} \to \alpha.$$

First notice that in view of Theorem 2 we know that if $x, y \in (\mathbb{R}_+)^n \cap D_{\alpha^{(k)}}$, then

(8)
$$\tilde{k}_{D_{\alpha^{(k)}}}^*(x,y) = m((x_1^{\alpha_1^{(k)}} \cdot \ldots \cdot x_n^{\alpha_n^{(k)}})^{\frac{1}{\min\{\alpha_j^{(k)}\}}}, (y_1^{\alpha_1^{(k)}} \cdot \ldots \cdot y_n^{\alpha_n^{(k)}})^{\frac{1}{\min\{\alpha_j^{(k)}\}}}).$$

We may assume that $\min\{\alpha_j\} = \alpha_n$ and $\min\{\alpha_j^{(k)}\} = \alpha_n^{(k)}$. First we prove that

$$\tilde{k}_{D_{\alpha}}^{*}(w,z) \geq m((|w_{1}|^{\alpha_{1}} \cdot \ldots \cdot |w_{n}|^{\alpha_{n}})^{1/\alpha_{n}}, (|z_{1}|^{\alpha_{1}} \cdot \ldots \cdot |z_{n}|^{\alpha_{n}})^{1/\alpha_{n}}).$$

Suppose it does not hold, so there is a mapping $\varphi \in \mathcal{O}(\overline{E}, D_{\alpha})$ such that $\varphi(\lambda_1) = (|w_1|, \ldots, |w_n|), \varphi(\lambda_2) = (|z_1|, \ldots, |z_n|)$ and

$$m(\lambda_1, \lambda_2) < m((|w_1|^{\alpha_1} \cdot \ldots \cdot |w_n|^{\alpha_n})^{1/\alpha_n}, (|z_1|^{\alpha_1} \cdot \ldots \cdot |z_n|^{\alpha_n})^{1/\alpha_n}).$$

Then we may choose k so large that $\varphi(E) \subset D_{\alpha^{(k)}}$ and $m(\lambda_1, \lambda_2) < m((|w_1|^{\alpha_1^{(k)}} \cdot \ldots \cdot |w_n|^{\alpha_n^{(k)}})^{1/\alpha_n^{(k)}}, (|z_1|^{\alpha_1^{(k)}} \cdot \ldots \cdot |z_n|^{\alpha_n^{(k)}})^{1/\alpha_n^{(k)}})$, which, however, contradicts (8).

To get the equality consider the mapping $\varphi(\lambda) := (\psi_1(\lambda), \dots, \psi_{n-1}(\lambda), \lambda \psi_n(\lambda))$, where (see Lemma 7)

$$\psi_j \in \mathcal{O}(E, \mathbb{C}_*), \quad j = 1, \dots, n-1,$$

$$\lambda_1 := (|w_1|^{\alpha_1} \cdot \dots \cdot |w_n|^{\alpha_n})^{\frac{1}{\alpha_n}} > 0, \lambda_2 := (|z_1|^{\alpha_1} \cdot \dots \cdot |z_n|^{\alpha_n})^{\frac{1}{\alpha_n}} > 0;$$

$$\psi_j(\lambda_1) = |w_j|, \quad \psi_j(\lambda_2) = |z_j|, \ j = 1, \dots, n-1,$$

Define also

$$\psi_n(\lambda) := \frac{1}{\left(\psi_1^{\alpha_1}(\lambda) \cdot \ldots \cdot \psi_{n-1}^{\alpha_{n-1}}(\lambda)\right)^{\frac{1}{\alpha_n}}}, \quad \lambda \in E.$$

The $\frac{1}{\alpha_n}$ -th root is chosen so that $\varphi_n(\lambda_1) = |w_n|$. One may also easily check from the form of ψ_j 's in the proof of Lemma 7 that then $\varphi_n(\lambda_2) > 0$, so $\varphi_n(\lambda_2) = |z_n|$. This completes the proof. \Box

Identically as in the rational case we have:

Proof of the formula for $k_{D_{\alpha}}^*$ in irrational case. Note that \tanh^{-1} of the desired formula satisfies the triangle inequality and coincides with the \tanh^{-1} of Lempert function off the axis. The continuity of the Kobayashi pseudodistance (see [JP 2]) as well as the definition of the Kobayashi pseudodistance finish the proof. \Box

Having the formula for the Lempert function we get

Proof of the formula for $g_{D_{\alpha}}$ in the irrational case. **Case I.** $\mathcal{J} = \emptyset$. Corollary 16 implies that

 $g_{D_{\alpha}}(w,z) = 0$ for any $z \in T_w$.

Maximum principle for plurisubharmonic functions (applied to $g_{D_\alpha}(w,\cdot))$ implies that

 $g_{D_{\alpha}}(w,z) = 0$ for any z with $|z_j| \le |w_j|$,

which, however, means that $g_{D_{\alpha}}(w, \cdot)$ vanishes on a set with non-empty interior (remember that $w_1 \cdot \ldots \cdot w_n \neq 0$) but $g_{D_{\alpha}}(w, \cdot)$ is logarithmically plurisubharmonic, so it must vanish on D_{α} .

Case II. $\mathcal{J} \neq \emptyset$.

This case is a simple consequence of Lemma 10, the inequality $g \leq \tilde{k}^*$, definition of the Green function and the fact that the function $(|z_1|^{\alpha_1} \cdot \ldots \cdot |z_n|^{\alpha_n})^{\frac{1}{\alpha_{j_1} + \ldots + \alpha_{j_k}}}$ is logarithmically plurisubharmonic on D_{α} . \Box

Proof of the formula for $A_{D_{\alpha}}$ in irrational case. The result follows from the formula for the Green function and definition of the Azukawa pseudometric. \Box

And now similarly as in the rational case we finish up the proof by showing the formula for $\kappa_{D_{\alpha}}$.

Lemma 18. Let α be of irrational type. Then

$$\mathfrak{D}k_{D_{\alpha}}(w;X) = \gamma_E\left(\left(\prod_{j=1}^n |w_j|^{\alpha_j}\right)^{\frac{1}{\min\{\alpha_k\}}}, \left(\prod_{j=1}^n |w_j|^{\alpha_j}\right)^{\frac{1}{\min\{\alpha_k\}}} \frac{1}{\min\{\alpha_k\}} \sum_{j=1}^n \frac{\alpha_j X_j}{w_j}\right)$$

,

for $w \in \tilde{D}_{\alpha}, X \in \mathbb{C}^n$.

Proof. Without loss of generality we may assume that $\alpha_n = \min{\{\alpha_k\}}$. The formula for the Kobayashi pseudodistance gives us

(9)
$$\mathfrak{D}k_{D_{\alpha}}(w;X) = \limsup_{\lambda \neq 0} \frac{\left|\prod_{j=1}^{n} |w_{j} + \lambda X_{j}|^{\alpha_{j}/\alpha_{n}} - \prod_{j=1}^{n} |w_{j}|^{\alpha_{j}/\alpha_{n}}\right|}{\left|1 - \prod_{j=1}^{n} |w_{j} + \lambda X_{j}|^{\alpha_{j}/\alpha_{n}} \prod |w_{j}|^{\alpha_{j}/\alpha_{n}}\right| |\lambda|}$$

Note that $\alpha_j/\alpha_n \ge 1$. Therefore, applying the Taylor formula we get, for λ close to 0,

$$|w_j + \lambda X_j|^{\alpha_j/\alpha_n} = |w_j|^{\alpha_j/\alpha_n} + \frac{\alpha_j}{\alpha_n} |w_j|^{\alpha_j/\alpha_n} \left| \operatorname{Re}\left(\frac{\lambda X_j}{w_j}\right) \right| + \varepsilon_j(\lambda), \quad j = 1, \dots, n,$$

where $\frac{\varepsilon_j}{\lambda} \to 0$ as $\lambda \to 0$. Substituting the last equalities in (9) we get that

$$\mathfrak{D}k_{D_{\alpha}}(w;X) = \limsup_{\lambda \neq 0} \frac{\prod_{j=1}^{n} (|w_j|^{\alpha_j})^{1/\alpha_n} \operatorname{Re}\left(\lambda\left(\sum_{j=1}^{n} \frac{\alpha_j X_j}{\alpha_n w_j}\right)\right)}{\left(1 - \prod_{j=1}^{n} |w_j|^{2\alpha_j/\alpha_n}\right) |\lambda|}$$

which equals the desired value. \Box

Proof of the formula for $\kappa_{D_{\alpha}}$ in irrational case. If $\mathcal{J} \neq \emptyset$, then, in view of Lemma 10, we are done. Also the case $\sum_{j=1}^{n} \frac{\alpha_j X_j}{w_j} = 0$ follows from Remark 4. Below we deal with the remaining cases.

Take $w \in D_{\alpha}$. Without loss of generality we may assume that $w_j \in \mathbb{R}_+$, $j = 1, \ldots, n$ and $\alpha_n = \min\{\alpha_j\}$. Below, for $X \in \mathbb{C}^n$ with $\sum_{j=1}^n \frac{\alpha_j X_j}{w_j} \neq 0$ we shall construct a mapping $\varphi \in \mathcal{O}(E, D_{\alpha})$ such that

$$\varphi(\lambda_1) = w, \quad t\varphi'(\lambda_1) = X,$$

where $\lambda_1 := (w_1^{\alpha_1} \cdot \ldots \cdot w_n^{\alpha_n})^{1/\alpha_n} > 0, t := (w_1^{\alpha_1} \cdot \ldots \cdot w_n^{\alpha_n})^{1/\alpha_n} \sum_{j=1}^n \frac{\alpha_j X_j}{\alpha_n w_j}$. Note that the existence of such a φ would finish the proof because of Lemma 18

Note that the existence of such a φ would finish the proof because of Lemma 18 and (4).

Define the mapping

$$\varphi(\lambda) := \left(\psi_1(\lambda), \dots, \psi_{n-1}(\lambda), \frac{\lambda}{(\psi_1^{\alpha_1}(\lambda) \cdot \dots \cdot \psi_{n-1}^{\alpha_{n-1}}(\lambda))^{1/\alpha_n}}\right),$$

where (see Lemma 9)

$$\psi_j(\lambda_1) = w_j, \ j = 1, \dots, n-1, \quad t\psi'_j(\lambda_1) = X_j, \ j = 1, \dots, n-1.$$

We choose the $1/\alpha_n$ -th power so that $\varphi_n(\lambda_1) = w_n$. After some elementary transformation we get that

$$t\varphi_n'(\lambda_1) = X_n,$$

which finishes the proof. \Box

5. Some applications. Having proven the formulas for the invariant functions for the elementary Reinhardt domains we may formulate some conclusions, which follow from them. They show how irregularly the functions can behave although the domains considered are very regular.

For a given domain $D \subset \mathbb{C}^n$ we define a relation \mathcal{R} on D as follows: $w\mathcal{R}z$ for $w, z \in D$ if $k_D^*(w, z) = 0$. In [Ko2], S. Kobayashi asked a question whether the quotient D/\mathcal{R} has always a complex structure. The answer is 'no', however the examples showing this are artificial (see [Ko 1], page 130 also [HD] and [Gi]). ¿From Theorem 3 we know that if α is of irrational type, then D_{α}/\mathcal{R} is equal to [0, 1). This gives the first very simple example of a very regular domain, for which the answer to the above question is 'no'.

One may consider some generalizations of the Carathéodory pseudodistance, called the k-th Möbius function, denoted by m^k (for k = 1, 2, ...) (for definitions see [JP 2]). S. Nivoche (see [N]) has proved that if a domain is strictly hyperconvex, then the functions m^k tend to g. One may easily verify that if α is of irrational type, then all the m^k 's vanish on $D_{\alpha} \times D_{\alpha}$. Therefore we see that no such convergence takes place in domains D_{α} (α of irrational type), so one may not expect a similar result as in [N] in the class of Reinhardt complete pseudoconvex domains.

In general, the Lempert function seems to be very distant from the Green function. The definition of the Kobayashi pseudodistance makes an impression that the Kobayashi pseudodistance should be larger or equal than the Green function. Nevertheless, if $\alpha \in \mathbb{N}_*^n$ is such that all α_j 's are relatively prime and $\min\{\alpha_j\} = 1$, then we have the following inequalities (see Theorem 2):

$$c_{D_{\alpha}}^* \equiv k_{D_{\alpha}}^* \le g_{D_{\alpha}} \le k_{D_{\alpha}}^*, \quad k_{D_{\alpha}}^* \ne g_{D_{\alpha}}.$$

In the papers [Pa] and [L1] a notion of stationary maps was introduced and studied. In the class of strongly convex domains these mappings are exactly the \tilde{k} -geodesics. In case of strongly pseudoconvex domains godesics must be necessarily stationary maps. One knows that in general the inverse implication does not hold (see [Pa] and [PZ]). From the proof of Theorem 2 we may construct also other examples disproving the implication. One may find even domains, which are strongly pseudoconvex (one produces them by cutting the 'ends' and then smoothing the corners).

6. Open problems. It would be interesting to find formulas of all invariant functions discussed above for domains of the following more general type:

$$D_{\alpha^1} \cap \ldots \cap D_{\alpha^k} \cap ((R_1 E) \times \ldots \times (R_n E)),$$

where $\alpha^j \in (\mathbb{R}_+)^n$, $j = 1, \ldots, k$.

References

K. Azukawa, Two intrinsic pseudo-metrics with pseudoconvex indicatrices and star-
like domains, J. Math. Soc. Japan 38 (1986), 627–647.
B.E.Blank, D.Fan, D.Klein, S.G.Krantz, D.Ma, MY.Pang, The Kobayashi metric $% \mathcal{A}_{\mathcal{A}}$
of a complex ellipsoid in \mathbb{C}^2 , Experimental Math. 1 (1992), 47–55.
C. Carathéodory, Über eine spezielle Metrik die in der Theorie der analytischen
Funktionen auftritt, Atti Pontifica Acad. Sc., Nuovi Lincei 80 (1927), 135–141.
A. Edigarian, On extremal mappings in complex ellipsoids, Annales Pol. Math.
LXII.1 (1995), 83–96.
${\rm G.} \qquad {\rm Gentili}, \qquad {\it Regular} \qquad {\it complex} \qquad {\it geodesics} \qquad {\it in} \qquad {\it the} \qquad {\it domain}$
$D_n = \{(z_1, \ldots, z_n) \in \mathbf{C}^n : z_1 + \ldots + z_n < 1\}$, Springer Lecture Notes in Math.
1277 (1987), 35-45.
B. Gilligan, On the Kobayashi pseudometric reduction of homogeneous spaces, Can.
Math. Bull. 31 (1988), 45–51.
G. H. Hardy & E. M. Wright, An Introduction to the Theory of Numbers, Oxford
Science Publ., 1978.
V. Z. Hristov & T. Davidov, Examples of typical Carathéodory and Kobayashi pseu-
dodistances, Comptes rend. de l'Acad. Bulg. Sc. 39 (1986), 23–25.
M. Jarnicki & P. Pflug, Some remarks on the product property, Proc. Symp. Pure Math. 52 (Dant 2) (1001) 262-272
Math. 52 (Fart 2) (1991), 205-212. M. Jamieli & P. P. Hug, Invariant Distances and Matrices in Complex Analysis Welton
de Gruvter, 1993.
M. Jarnicki, P. Pflug & R. Zeinstra, Geodesics for convex complex ellipsoids, Ann.
d. Sc. Norm. Sup. di Pisa XX Fasc 4 (1993), 535–543.
M. Klimek, Extremal plurisubharmonic functions and invariant pseudodistances,
Bull. Soc. Math. France 113 (1985), 231-240.
M. Klimek, <i>Pluripotential Theory</i> , Oxford University Press, 1991.
S. Kobayashi, Hyperbolic Manifolds and Holomorphic Mappings, Pure and Appl.
Math. 2, M. Dekker, 1970.
S. Kobayashi, Intrinsic distances, measures and geometric function theory, Bull.
Amer. Math. Soc. 82 (1976), 457-416.
${\tt L.Lempert}, \ La \ m\acute{e}trique \ de \ Kobayashi \ et \ la \ représentation \ des \ domaines \ sur \ la \ boule,$
Bull. Soc. Math. France 109 (1981), 427-479.

- [L 2] L. Lempert, Intrinsic distances and holomorphic retracts, Complex Analysis and Applications '81 (1984), Sophia, 341-364.
- S. Nivoche, Pluricomplex Green function, capacitative notions and approximation problems in Cⁿ,, Ind. Univ. Math. J. 44 (1995), 489–510.
- [Pa] M.-Y.Pang, Smoothness of the Kobayashi metric of non-convex domains, International Journal of Mathematics 4 (1993), 953-987.

- [PZ] P. Pflug & W. Zwonek, The Kobayashi Metric for Non-Convex Complex Ellipsoids, Complex Variables 29 (1996), 59-71.
- [R] H.-J. Reiffen, Die Carathéodory Distanz und ihre zugehörige Differentialmetrik, Math. Annalen 161 (1965), 315–324.

CARL VON OSSIETZKY UNIVERSITÄT OLDENBURG, FACHBEREICH MATHEMATIK, POSTFACH 2503, D-26111 OLDENBURG, GERMANY

 $E\text{-}mail\ address:\ \texttt{pflug@mathematik.uni-oldenburg.de}$

Uniwersytet Jagielloński, Instytut Matematyki, Reymonta 4, 30-059 Kraków, Poland

E-mail address: zwonek@im.uj.ed.pl