ON PERTURBATIVE *PSU(N)* INVARIANTS OF RATIONAL HOMOLOGY 3-SPHERES

THANG T. Q. LE

ABSTRACT. We construct power series invariants of rational homology 3-spheres from quantum PSU(n)-invariants. The power series can be regarded as perturbative invariants corresponding to the contribution of the trivial connection in the hypothetical Witten's integral. This generalizes a result of Ohtsuki (the n = 2 case) which led him to the definition of finite type invariants of 3-manifolds. The proof utilizes some symmetry properties of quantum invariants (of links) derived from the theory of affine Lie algebras and the theory of the Kontsevich integral.

0. INTRODUCTION

In this paper we construct power series invariants of rational homology 3-spheres from the quantum PSU(n)-invariants. This generalizes a result of Ohtsuki (the n = 2 case) which led him to the definition of finite type invariants of 3-manifolds.

For a fixed compact Lie group and an integer r, Witten defined an invariant of 3manifolds using Feynman path integral which is not mathematically rigorous. Reshetikhin and Turaev [28], and later Turaev and Wenzl [34] defined (mathematically) quantum invariants of 3-manifolds associated with simple Lie algebras (see [33] and references therein). The quantum invariants can be thought of as a mathematical realization of Witten's path integral.

One could use formal perturbation theory to approximate the Witten integral. The coefficients of the approximations are known as perturbative invariants, which also need a mathematically rigorous definition. One might want to get perturbative invariants directly from (Reshetikhin-Turaev) quantum invariants. The difficulty here is that quantum invariant can be defined only when the "quantum" parameter q is roots of unity, while in order to get perturbative invariants, one need to make the formal expansion $q = \exp(h)$, with h an *indeterminate*. This substitution does not make sense if q takes only values roots of unity.

Ohtsuki [24] showed that one can extract power series invariants (of homology 3spheres) from quantum SU(2)-invariants which can be considered as the perturbative invariants. Ohtsuki's result led him to the important definition of finite type invariant of 3-manifolds, a counterpart of Vassiliev invariants for homology 3-spheres. The Ohtsuki series can be considered, on the physics level, as the contribution of the trivial connection in the perturbative expansion of the Witten integral (see [29]). Further investigation

This is NSF grant DMS-9626404. work partially supported by This and related preprints can obtained at http://www.math. be [0]buffalo.edu/~letu/.

of the Ohtsuki's series was carried out by Ohtsuki [24, 26], Lawrence and Rozansky [12, 13, 29, 30], Lin and Wang [20], Kricker and Spence [11], and others. A result of H. Murakami [23] says that the first term of the Ohtsuki series is 6 times the Casson-Walker invariant.

In [18] it was conjectured that similar power series invariants exist for Lie group SU(n), n > 2. It is the main goal of this paper to prove this conjecture. We show that one can extract power series invariants of rational homology 3-spheres from quantum PSU(n)-invariants. Here the quantum PSU(n)-invariant is a refined version of the SU(n)-invariant which was introduced by Kirby and Melvin for the n = 2 case (see [9]), and by Kohno and Takata for the n > 2 case (see [8]). It will be clear from the proof that for rational homology 3-spheres, one should use the PSU(n) version instead of the SU(n) one. It is because of some integrality and symmetry properties of quantum link invariants related to the PSU(n) case. In a sense, the SU(n) version is based on the weight lattice, and the PSU(n) version is based on the root lattice; and quantum invariants of links on the root lattice are nicer: they have integrality property and more symmetry.

A few words about the proof. The proof of the main result differs from that of Ohtsuki for the case n = 2, since Ohtsuki [24] used some identities which are either specific to the n = 2 case or hard to generalize to the n > 2 case (see also [20, 29]). Some of our arguments are generalizations of those in [24, 20, 29] which treat the n = 2 case. We will make use of some symmetry properties of quantum invariants of links which are derived from affine Lie algebra theory and the Kontsevich integral theory. Along the way we also investigate the dependence of quantum invariants of links on the colors of the link components. The proof also suggests how to generalize the result to other Lie algebras, and we will try to formulate everything in terms of Lie algebra theory.

In §1 we introduce the notation and formulate the main result. In section 2 we give the proof of the main theorem using some results proved in sections 3 and 4. In section 3 we investigate the dependence of quantum invariants of links on the colors (which are modules of Lie algebras). In section 4 we prove some properties of the quadratic Gauss sums based on the root lattice. In Appendix we show that our definition of quantum PSU(n)-invariant coincides with that of Kohno and Takata, and calculate the value of a multi-variable quadratic Gauss sum.

Acknowledgment. Much of this work was carried out while the author was visiting the Mathematical Sciences Research Institute in Berkeley in 1996-1997. Research at MSRI is supported in part by NSF grant DMS-9022140. The author thanks J. Murakami, T. Takata and H. Wenzl for helpful discussions.

1. Preliminaries. Main result

1.1. Notations. We will use the following notations for objects related to the Lie algebra sl_n . See, for example, [2, 5], for the terminologies used here.

- $\alpha_1, \ldots, \alpha_{n-1}$: the standard root basis.
- Φ : the root system.
- Φ_+ : the set of positive roots. There are n(n-1)/2 positive roots.
- $\lambda_1, \ldots, \lambda_{n-1}$: the standard fundamental weights.

A: the integral weight lattice, $\Lambda = \{\sum_{i=1}^{n-1} k_i \lambda_i, k_i \in \mathbb{Z}\}$. All the weights λ_i and roots α_j are elements of Λ . In $\Lambda \otimes \mathbb{R}$ there is the standard scalar product, for which $(\alpha_i | \alpha_j) = A_{ij}$. Here A is the Cartan matrix. One has $(\alpha_i | \lambda_j) = \delta_{ij}$.

 ρ : half-sum of positive roots. One has that $\rho = \lambda_1 + \cdots + \lambda_{n-1}$, and $(\rho|\rho) = n(n^2 - 1)/12$.

 \mathbb{Z}_+ : the set of non-negative integers.

 Λ_+ : the set of dominant weights, $\Lambda_+ = \{k_1\lambda_1 + \dots + k_{n-1}\lambda_{n-1} \mid k_1, \dots, k_{n-1} \in \mathbb{Z}_+\}$

 Λ_{++} : the set of strongly dominant weights, $\Lambda_{++} = \rho + \Lambda_{+}$.

 Λ^{root} : The root lattice, $\Lambda^{\text{root}} = \{\sum_{i=1}^{n-1} k_i \alpha_i, k_i \in \mathbb{Z}\}$. The root lattice is a subgroup of the weight lattice Λ of index n.

 θ : the longest root, $\theta = \alpha_1 + \cdots + \alpha_{n-1}$.

 \mathcal{R} : the ring of finite dimensional sl_n -modules.

 V_{μ} , with $\mu \in \Lambda_{++}$: the irreducible sl_n -module with highest weight $\lambda - \rho$. Note that this is slightly different from the usual notation. The shift by ρ is more convenient for us.

The Weyl group \mathcal{W} acts on $\Lambda \otimes \mathbb{R}$, and a fundamental domain is the fundamental chamber

$$C = \{ z \in \Lambda \times \mathbb{R} \mid (z, \alpha_i) \ge 0, \quad i = 1, \dots, n-1 \} = \{ \sum_{i=1}^{n-1} a_i \lambda_i \mid a_i \ge 0, a_i \in \mathbb{R} \}.$$

So, Λ_+ is the intersection of Λ and C. The Weyl group is generated by reflections along the boundary facets of C. For an element $w \in \mathcal{W}$, let $\operatorname{sn}(w)$ be the sign of w. Similarly, for a non-zero number b, let $\operatorname{sn}(b)$ be the sign of b.

Throughout the paper, the number n (of sl_n) is fixed. We will use r to denote the "level" of the theory (it's just a positive integer). Let

$$\bar{r} = \frac{r-1-n(n-1)}{2}$$

Note that if r is odd, which we always assume, then \bar{r} is an integer.

We will use q, x as indeterminates, with q = 1 + x. We use ζ to denote the *r*-th root of unity, $\zeta = \exp(2\pi i/r)$. When *r* is an odd prime, we will identify $\mathbb{Z}[\zeta]$ with $\mathbb{Z}[q]/(q^{r-1} + \cdots + q + 1)$.

For any positive integer ℓ let

$$\mathbb{Z}_{(\ell)} = \mathbb{Z}\left[\frac{1}{(\ell-1)!\,n!}\right].$$

For a rational homology 3-sphere M (i.e. a closed oriented 3-manifold whose homology group $H_1(M, \mathbb{Z})$ is finite), let $\mathbb{Q}[[x]]_{(M)}$ be the set of all power series $\sum_{\ell} d_{\ell} x^{\ell}$ such that d_{ℓ} is a number in $\mathbb{Z}[\frac{1}{(2\ell+n(n-1))!|H_1(M,\mathbb{Z})|}]$.

1.2. Quantum invariants of oriented framed links and 3-manifolds. For a framed oriented link L with m ordered components in S^3 let $J_L(\mu_1, \ldots, \mu_m; q)$ be the sl_n -quantum invariant of the link (see [27, 33]), where the components of L are colored by $V_{\mu_1}, \ldots, V_{\mu_m}$. Here q is an indeterminate and the μ_j 's are in Λ_{++} , so that V_{μ_j} are finite-dimensional

irreducible sl_n -modules of highest weight $\mu_j - \rho$. Actually, J_L can be regarded as a linear homomorphism from $\mathcal{R}^{\otimes m}$ to $\mathbb{Z}[q^{\pm 1/2n}]$.

We extend the definition of $J_L(\mu_1, \ldots, \mu_m; q)$ to the case when the μ_j 's are in Λ as follows. If one of the μ_j is on the boundary of a chamber w(C), where w is in the Weyl group, put $J_L(\mu_1, \ldots, \mu_m; q) = 0$. Otherwise, there exist unique w_1, \ldots, w_m in the Weyl group \mathcal{W} such that the elements $w_i(\mu_j)$ are in Λ_{++} . Then put

$$J_L(\mu_1, \dots, \mu_m; q) = \operatorname{sn}(w_1) \dots \operatorname{sn}(w_m) J_L(w_1(\mu_1), \dots, w_m(\mu_m); q).$$

It's more convenient to use the following normalization of J_L :

$$Q_L(\mu_1,\ldots,\mu_m;q)=J_L(\mu_1,\ldots,\mu_m;q)J_U(\mu_1;q)\ldots J_U(\mu_m;q),$$

where U is the unknot with framing 0. The invariant Q_L enjoys the symmetry described in the following proposition, which follows immediately from the definition.

Proposition 1.1. a) $Q_L(\mu_1, \ldots, \mu_{n-1}; q)$ is invariant under the action of the Weyl group, *i.e.* for every $w_1, \ldots, w_m \in \mathcal{W}$, one has

$$Q_L(\mu_1, \ldots, \mu_m; q) = Q_L(w_1(\mu_1), \ldots, w_m(\mu_m); q).$$

b) If one of the μ_j , j = 1..., m, is on the boundary of the fundamental chamber C, then $Q_L(\mu_1, \ldots, \mu_m; q) = 0$.

For a positive integer r let

$$F'_L(r) = \sum_{j=1}^m \sum_{\mu_j \in \Lambda'_r} Q_L(\mu_1, \dots, \mu_m; \zeta),$$

where $\zeta^a = \exp(2a\pi i/r)$ for every rational number a, and

$$\Lambda'_r = \{ \mu \in \Lambda_+ \mid (\mu | \theta) \le r \} = \{ \sum_{i=1}^{n-1} k_i \lambda_i \in \Lambda_+ \mid \sum k_i \le r \}.$$

Suppose the closed oriented 3-manifold M^3 is obtained from S^3 by surgery along a framed unoriented link L. Providing L with arbitrary orientation, the quantum SU(n)-invariant of M at the r-th root of unity is defined by (see, for example, [7, 34])

$$\tau_r^{SU(n)}(M) = \frac{F'_L(r)}{F'_{U_+}(r)^{\sigma_+} F'_{U_-}(r)^{\sigma_-}}.$$

Here σ_+, σ_- are the numbers of positive and negative eigenvalues of the linking matrix of L and U_{\pm} are the trivial knots with framing ± 1 .

1.3. Quantum invariants of links at roots of unity. Let C_r be the simplex which is the convex hull of the points $0, r\lambda_1, \ldots, r\lambda_{n-1}$. In other words,

$$C_r = \{ z \in C \mid (z, \theta) \le r \} = \{ \sum_{i=1}^{n-1} a_i \lambda_i \in C \mid \sum_{i=1}^{n-1} a_i \le r \},\$$

(we call it the fundamental alcove of level r). Then $\Lambda'_r = C_r \cap \Lambda_+$. The affine Weyl group at level r, by definition, is the group $\mathcal{W}_{(r)}$ generated by all the reflections along boundary facets of the simplex C_r . It is known that $\mathcal{W}_{(r)}$ is the semi-direct product of \mathcal{W} and the translation group $r\Lambda^{\text{root}}$ (see [5], chapter 6).

Lemma 1.2. Each of the lattices Λ^{root} and $\rho + \Lambda^{root}$ is invariant under the action of $\mathcal{W}_{(r)}$, for every positive integer r.

Proof. The root lattice Λ^{root} is certainly invariant under the action of the Weyl group and the translation group $r\Lambda^{\text{root}}$. Hence Λ^{root} is invariant under the action of $\mathcal{W}_{(r)}$.

If α_i is a basis root, then the reflection along the hyperplane perpendicular to α_i maps ρ to $\rho - \alpha_i$, which belongs to $\rho + \Lambda^{\text{root}}$. These reflections generate the Weyl group, hence $\rho + \Lambda^{\text{root}}$ is invariant under the action of the Weyl group. It is obviously invariant under the translation group $r\Lambda^{\text{root}}$.

When specialized to the r-th root of unity, Q_L has more symmetry. The proof of the following important proposition is given in §3.

Proposition 1.3. Let $\zeta = \exp(2\pi i/r)$, where r is a positive integer.

a) $Q_L(\mu_1, \ldots, \mu_m; \zeta)$ is invariant under the action of the affine Weyl group $\mathcal{W}_{(r)}$, i.e. for every w_1, \ldots, w_m in $\mathcal{W}_{(r)}$, one has

$$Q_L(\mu_1,\ldots,\mu_m;\zeta) = Q_L(w_1(\mu_1),\ldots,w_m(\mu_m);\zeta).$$

b) If one of the μ_j , j = 1..., m, is on the boundary of the fundamental alcove C_r , then $Q_L(\mu_1, \ldots, \mu_m; \zeta) = 0$.

1.4. On the definition of PSU(n)-quantum invariants. Quantum PSU(n)-invariants were introduced by Kirby-Melvin [9] in the n = 2 case, and by Kohno-Takata [8] in the n > 2 case. We will give here another definition which is more natural from our point of view. We will prove that our definition coincides with that of Kohno and Takata in Appendix.

Let Λ_r^{root} be the set of all elements of the root lattice in the half-open parallelepiped spanned by $r\alpha_1, \ldots, r\alpha_{n-1}$, i.e.

$$\Lambda_r^{\text{root}} = \{ k_1 \alpha_1 + \dots + k_{n-1} \alpha_{n-1} \in \Lambda^{\text{root}} \mid 0 \le k_i < r \}.$$

Define $F_L(r)$ using the same formula of $F'_L(r)$, only replacing Λ'_r by $\rho + \Lambda^{\text{root}}_r$:

$$F_L(r) = \sum_{j=1}^m \sum_{\mu_j \in (\rho + \Lambda_r^{\text{root}})} Q_L(\mu_1, \dots, \mu_m; \zeta).$$

Suppose that r and n are coprime. We will show later that $F_{U_{\pm}}(r) \neq 0$. Suppose that M is obtained by surgery along the framed link L. Define the quantum PSU(n)-invariant $\tau_r^{PSU(n)}(M)$ by

$$\tau_r^{PSU(n)}(M) = \frac{F_L(r)}{F_{U_+}(r)^{\sigma_+} F_{U_-}(r)^{\sigma_-}}.$$
(1.1)

Remark. a) In the set $\rho + \Lambda_r^{\text{root}}$ there are weights which are not highest weights of any finite-dimensional sl_n -modules.

b) In the definition of F_L , the μ_j 's run the set $\rho + \Lambda_r^{\text{root}}$ since we use V_{μ} for the sl_n module with highest weight $\mu - \rho$. Also if n is *odd*, then ρ is in the root lattice, hence $\rho + \Lambda^{\text{root}} = \Lambda^{\text{root}}$; and using the symmetry of Q_L we can replace $\rho + \Lambda_r^{\text{root}}$ by Λ_r^{root} .

c) If in the definition of F_L , we let the μ_j 's run the set of all integral weights in the same parallelepiped, then from the above formula we get exactly the SU(n)-invariant. So both SU(n) and PSU(n)-invariants can be defined by one formula, only for the SU(n) invariant, we use the weight lattice, while for the PSU(n) invariant, we use the root lattice. It is known that $\Lambda/\Lambda^{\text{root}}$ is the cyclic group of order n.

c) Kohno and Takata [8] showed that the PSU(n) version is finer than the SU(n) version.

The following important integrality property was established by H. Murakami [23] for n = 2, and by Takata-Yokota [32] and Masbaum-Wenzl [22] for n > 2; this result can also be obtained by the method of this paper.

Theorem 1.4. Suppose r is a prime not dividing $n!|H_1(M,\mathbb{Z})|$. Then the PSU(n)quantum invariant $\tau_r^{PSU(n)}(M)$ is a number in $\mathbb{Z}[\zeta]$. Here $\zeta = \exp(2\pi i/r)$.

1.5. Existence of perturbative expansion. Unlike the link case, quantum invariants of 3-manifolds can be defined only at roots of unity. In perturbative theory, we want to expand $q = e^h$, with h an indeterminate. However, in the 3-manifold case, q must be specified at the r-th root of unity, and it seems difficult to make meaning the substitution $q = e^h$, with h an indeterminate. One way around is the following. Instead of h, we use x = q - 1. Suppose r is a prime not dividing $n!|H_1(M,\mathbb{Z})|$.

Let $f(q) \in \mathbb{Z}[q]$ be a representative of $\tau_r^{PSU(n)}(M) \in \mathbb{Z}[\zeta] = \mathbb{Z}[q]/(q^{r-1} + \cdots + q + 1)$. Using the substitution q = 1 + x, we get

$$f(q)|_{q=x+1} = c_{r,0} + c_{r,1}x + \dots + c_{r,\ell}x^{\ell} + \dots$$

The integer coefficients $c_{r,\ell}$ are, in general, dependent on the representative f(q). However, it's easy to prove the following (see [24]).

Lemma 1.5. For $\ell \leq r-2$, the classes $c_{r,\ell} \pmod{r}$ do not depend on the representative f(q) in $\mathbb{Z}[q]$.

Hence for each prime r, the classes $c_{r,\ell} \pmod{r}$, $\ell = 0, 1, \ldots, r-2$ are invariants of M. Let us fix ℓ and let r vary (but r must be an odd prime). For each r we have a residue class $c_{r,\ell}(M) \pmod{r}$. We want to show that these classes (when r varies but ℓ fixed) can be unified in the following sense: they are derived from the same rational number not depending on r. More precisely, (see Theorem 1.7 below) we will show that there is a rational number $c_{\ell}(M) \in \mathbb{Z}[\frac{1}{(2\ell+n(n-1))!|H_1(M,\mathbb{Z})|}]$ such that

$$c_{\ell} \equiv c_{r,\ell} \pmod{r} \tag{1.2}$$

for sufficiently large prime r (in fact, $r > 2\ell + n(n-1)$ is enough). The following is easy to prove (see [24]).

Lemma 1.6. For each ℓ , there exists at most one rational number $c_{\ell} \in \mathbb{Z}[\frac{1}{(2\ell+n(n-1))! |H_1(M,\mathbb{Z})|}]$ such that (1.2) holds for all sufficiently large prime r.

It follows that if such c_{ℓ} exist, then it is an invariant of M. The existence of such c_{ℓ} is a quite non-trivial fact. If all the c_{ℓ} 's do exist, then the series $\sum c_{\ell} x^{\ell}$ should be considered the perturbative expansion of $\tau_r^{PSU(n)}(M)$.

Another way to look at the existence of c_{ℓ} is the following. For $f(q) \in \mathbb{Z}[q]$ (or in $\mathbb{Z}_{(r)}[q]$) we will define $\mathfrak{p}_r(f(q))$ as follows. First we substitute q = x + 1 in f, then cancel any powers of x with degree greater than $\bar{r} = (r - n(n-1) - 1)/2$, finally reduce all the coefficients modulo r. The result is $\mathfrak{p}_r(f)$. So \mathfrak{p}_r is an algebra homomorphism

$$\mathfrak{p}_r\colon \mathbb{Z}_{(r)}[q]\to \mathbb{Z}[x]/(r,x^{\bar{r}+1}).$$

It is easy to see that \mathfrak{p}_r descends to an algebra homomorphism, also denoted by \mathfrak{p}_r , on

$$\mathbb{Z}_{(r)}[q]/(q^{r-1}+\cdots+q+1) = \mathbb{Z}_{(r)}[\zeta]$$

Hence there is defined $\mathfrak{p}_r(\tau_r^{PSU(n)}(M))$. Certainly $\mathfrak{p}_r(x^d) = 0$ if $d > \bar{r}$.

Recall that $\mathbb{Q}[[x]]_{(M)}$ is the set of all formal power series $\sum d_{\ell}x^{\ell}$ such that $d_{\ell} \in \mathbb{Z}[\frac{1}{(2\ell+n(n-1))!|H_1(M,\mathbb{Z})|}]$. So, when r is a prime greater than both $|H_1(M,\mathbb{Z})|$ and $2\ell + n(n-1)$, we can reduce d_{ℓ} modulo r.

For a series $g = \sum d_{\ell} x^{\ell} \in \mathbb{Q}[[x]]_{(M)}$, where $|H_1(M,\mathbb{Z})| < r$, we can also define $\mathfrak{p}_r(g)$ in a similar way, as follows. First cancel powers of x with degree greater than \bar{r} , then reduce all the coefficients modulo r. The result is $\mathfrak{p}_r(g)$. A similar operator was also considered by Ohtsuki and Rozansky in connection with the n = 2 case.

Then the equality $c_{\ell} \equiv c_{r,\ell} \pmod{r}$, for every ℓ and $r > 2\ell + n(n-1)$, means that

$$\mathfrak{p}_r(\tau_r^{PSU(n)}(M)) = \mathfrak{p}_r(\sum_{\ell} c_\ell x^\ell).$$

The following is the main result of this paper.

Theorem 1.7. Suppose that M is a rational homology 3-sphere. There exists a power series

$$\tau^{PSU(n)}(M) = \sum_{\ell=0}^{\infty} c_{\ell}(M) x^{\ell} \in \mathbb{Q}[[x]]_{(M)}$$

such that for any prime number r greater than $\max\{n(n-1), |H_1(M,\mathbb{Z})|\}$, one has

$$\mathfrak{p}_r(\tau^{PSU(n)}(M)) = \mathfrak{p}_r(\tau_r^{PSU(n)}(M)).$$

Here $\tau_r^{PSU(n)}(M)$ is the PSU(n)-quantum invariant of M at the r-th root of unity.

Remark. a) In the case n = 2, the existence of c_{ℓ} was proved by Ohtsuki [24]. Our normalization of $\tau^{PSU(n)}$ differs from that of Ohtsuki by the factor $\left(\frac{|H_1(M,\mathbb{Z})|}{r}\right)$, the Legendre symbol.

b) The above equality means that $\tau^{PSU(n)}(M)$ is the Fermat limit of $\tau_r^{PSU(n)}(M)$, in the terminology of [20].

The theorem shows that we can recover part of the quantum invariant $\tau_r^{PSU(n)}(M)$ from the perturbative invariant $\tau^{PSU(n)}(M)$. We have the following conjecture.

Conjecture 1.8. The perturbative PSU(n)-invariant $\tau^{PSU(n)}(M)$ dominates the quantum invariants $\tau_r^{SU(n)}(M)$, for every positive integer r, not necessarily prime.

Another conjecture is the integrality of $c_{\ell}(M)$, which is a generalization of conjectures made and investigated by Lawrence, Lin, Rozansky and Wang (see [12, 13, 30, 20]).

Conjecture 1.9. The perturbative invariants $c_{\ell}(M)$ are in $\mathbb{Z}[\frac{1}{n! |H_1(M\mathbb{Z})|}]$.

One can show that $c_1(M) = (\frac{|H_1(M,\mathbb{Z})|}{r})|H_1(M,\mathbb{Z})|^{-n(n-1)/2}n(n^2-1)\lambda_C(M)$, where λ_C is the Casson-Walker invariant in Lescop normalization [19].

2. Proof of the main theorem

We will give the proof of the main theorem based on some technical results which will be proved later.

2.1. Integrality of the exponents. It is known that for fixed μ_1, \ldots, μ_m , the quantum invariant $J_L(\mu_1, \ldots, \mu_m; q)$ is a polynomial in $q^{1/2n}, q^{-1/2n}$ with *integer* coefficients (see, for example, [22] for a proof), i.e. $J_L(\mu_1, \ldots, \mu_m; q) \in \mathbb{Z}[q^{1/2n}, q^{-1/2n}]$. When the linking matrix of L is 0, we have the following stronger result, whose proof will presented in §3.

Proposition 2.1. Suppose that the linking matrix of L is 0. Then $Q_L(\mu_1, \ldots, \mu_m; q)$ is a polynomial in q, q^{-1} with integer coefficients. In other words, $Q_L(\mu_1, \ldots, \mu_m; q)$ does not contain any non-integer fractional power of q.

Remark. The proposition does not hold true if the linking matrix of L is not 0.

Corollary 2.2. Suppose that the linking matrix of L' is diagonal and that μ_1, \ldots, μ_m are in $(\rho + \Lambda^{root})$. Then $Q_{L'}(\mu_1, \ldots, \mu_m; q)$ is in $\mathbb{Z}[q, q^{-1}]$.

Proof. Let L be the same link as L', only with 0 framing on each component. By the previous proposition, $Q_L(\mu_1, \ldots, \mu_m)$ is $\mathbb{Z}[q, q^{-1}]$. From the general theory of quantum invariants, it is known that increasing the framing by 1 on the *j*-th component results in a factor $q^{(|\mu_j|^2 - |\rho|^2)/2}$ in Q_L (see, for example, [33]). Here $|\mu|^2 = (\mu|\mu)$. We have:

$$Q_{L'}(\mu_1, \dots, \mu_m; q) = \prod_{j=1}^m q^{b_j \frac{|\mu_j|^2 - |\rho|^2}{2}} Q_L(\mu_1, \dots, \mu_m; q).$$
(2.1)

where b_i 's are the framing of components of L'. It remains to notice that when μ is in $\rho + \Lambda^{\text{root}}$, $(|\mu|^2 - |\rho|^2)/2$ is an integer.

The corollary shows an advantage of using the root lattice and links with diagonal linking matrix (which correspond to rational homology 3-spheres). If μ_j are in the bigger lattice Λ , then, in general, $Q_{L'}(\mu_1, \ldots, \mu_m; q)$ contains fractional powers of q.

2.2. Polynomial functions on the weight lattice. Let us introduce a new indeterminate x = q - 1. Then $q^{-1} = 1 - x + x^2 - x^3 + \ldots$, and Q_L becomes a formal power series in x:

$$Q_L(\mu_1, \dots, \mu_m; q) \Big|_{q=x+1} = \sum_{\ell=0}^{\infty} f_\ell(\mu_1, \dots, \mu_m) x^{\ell}.$$
 (2.2)

This kind of expansion, actually, a version of it for the n = 2 case, was first considered by Melvin and Morton [21].

Each monomial $\alpha_{i_1}\alpha_{i_2}\ldots\alpha_{i_s}$, where α_i 's are the basis roots, defines a function on Λ by

$$\alpha_{i_1}\alpha_{i_2}\ldots\alpha_{i_s}(\mu) = (\alpha_{i_1}|\mu)(\alpha_{i_2}|\mu)\ldots(\alpha_{i_s}|\mu)$$

for $\mu \in \Lambda$. Using linearity, every polynomial in the α_i 's can be regarded as a function on Λ ; we call them the *polynomial functions* on Λ . The degree of the polynomial function is the degree of the corresponding polynomial. A function of many variables, $f(\mu_1, \ldots, \mu_m)$, with $\mu_1, \ldots, \mu_m \in \Lambda$, is a *polynomial function* if it is a linear combination of functions of the form $g_1(\mu_1) \ldots g_m(\mu_m)$, where each g_j is a polynomial function.

Let Δ be the following polynomial function on Λ of degree n(n-1)/2:

$$\Delta = \frac{\prod_{\alpha \in \Phi_+} \alpha}{\prod_{\alpha \in \Phi_+} (\rho | \alpha)}.$$

The denominator of Δ is $\prod_{\alpha \in \Phi_+} (\rho | \alpha) = 1^{n-1} 2^{n-2} \dots (n-1)^1$.

Lemma 2.3. a) If $\mu \in \Lambda_{++}$, then the dimension of V_{μ} is given by

$$\dim(V_{\mu}) = \Delta(\mu)$$

b) One has that $\Delta(\mu) = 0$ if μ is on the boundary of the fundamental chamber C. For every $w \in \mathcal{W}$,

$$\Delta(w(\mu)) = \operatorname{sn}(w)\Delta(\mu)$$

Part a) is the famous dimension formula of Weyl. Part b) follows from elementary Lie algebra theory (see [2]).

The proof of the following proposition will be given in $\S3$.

Proposition 2.4. Suppose that the link L has m components and 0 linking matrix. a) All the functions f_i in (2.2) are polynomial functions divisible by $\Delta^2(\mu_1) \dots \Delta^2(\mu_m)$:

$$f_\ell(\mu_1,\ldots,\mu_m) = \Delta^2(\mu_1)\ldots\Delta^2(\mu_m)f'_\ell(\mu_1,\ldots,\mu_m).$$

Moreover, f_{ℓ} are polynomial functions taking integer values when $\mu_1, \ldots, \mu_m \in \Lambda$.

b) Suppose the degree of μ_j in f'_{ℓ} is s_j . Then $\sum_{j=1}^m s_j \leq 3\ell/2$ and $\max_j s_j \leq 2\ell - \sum_{j=1}^m s_j$. For $\ell = 0$ we have $f'_0 = 1$.

2.3. An expression of Q_L in terms of exponential functions. Let q^{β} , for $\beta \in \Lambda^{\text{root}}$, be the formal power series in x obtained by formally expanding $q^{\beta} = (1 + x)^{\beta}$, the coefficients being polynomial in β :

$$q^{\beta} = \sum_{\ell=0}^{\infty} \binom{\beta}{\ell} x^{\ell}.$$

Here $\begin{pmatrix} \beta \\ \ell \end{pmatrix}$ is the usual binome polynomial: $\begin{pmatrix} \beta \\ \ell \end{pmatrix} = \frac{\beta(\beta-1)\dots(\beta-\ell+1)}{\ell!}$.

Then $q^{\beta}(\mu)$, for $\beta \in \Lambda^{\text{root}}$ and $\mu \in \Lambda$, is a formal power series in x with integer coefficients:

$$q^{\beta}(\mu) = \sum_{\ell=0}^{\infty} {\beta \choose \ell} (\mu) x^{\ell} \in \mathbb{Z}[[x]],$$

since $(\beta|\mu)$ is an integer for every $\mu \in \Lambda$ and $\beta \in \Lambda^{\text{root}}$.

For each $i = 1, \ldots, n - 1$, the element

$$\eta_i = 1 - q^{-\alpha_i}$$

can be regarded as a function from Λ to $\mathbb{Z}[[x]]$. Moreover, the degree 0 term of $\eta_i(\mu)$ is 0, i.e. $\eta_i(\mu)$ is divisible by x.

For
$$\mathbf{a} = (a_1, \dots, a_{n-1}) \in (\mathbb{Z}_+)^{n-1}$$
, let $|\mathbf{a}| = a_1 + \dots + a_{n-1}$, and $\boldsymbol{\eta}^{\mathbf{a}} = \prod_{i=1}^{n-1} \eta_i^{a_i}$. Then for $\mu \in \Lambda, \, \boldsymbol{\eta}^{\mathbf{a}}(\mu) \in x^{|\mathbf{a}|}\mathbb{Z}[[x]].$

Proposition 2.5. Suppose that L is a framed link with m components and 0 linking matrix. As formal power series in x we have

$$Q_L(\mu_1,\ldots,\mu_m;x+1) = \sum c_{\mathbf{a}_1,\ldots,\mathbf{a}_m;\ell} \boldsymbol{\eta}^{\mathbf{a}_1}(\mu_1)\ldots\boldsymbol{\eta}^{\mathbf{a}_m}(\mu_m)x^{\ell-\sum_j|\mathbf{a}_j|}, \qquad (2.3)$$

where the sum is over the set of $\mathbf{a}_j \in (\mathbb{Z}_+)^{n-1}$ and $\ell \in \mathbb{Z}_+$ such that

$$|\mathbf{a}_j| \ge n(n-1)$$
 for every $j = 1, \dots, m$, (2.4)

$$\sum_{j=1}^{m} (|\mathbf{a}_j| - n(n-1)) \le 3\ell/2, \quad and \tag{2.5}$$

$$\max_{j}(|\mathbf{a}_{j}| - n(n-1)) \le 2\ell - \sum_{j}(|\mathbf{a}_{j}| - n(n-1)).$$
(2.6)

Moreover, the coefficient $c_{\mathbf{a}_1,\ldots,\mathbf{a}_m;\ell}$ is in $\mathbb{Z}_{(2\ell+(m+1)n(n-1)-\sum |\mathbf{a}_j|+1)}$.

Remark. Since $\eta^{\mathbf{a}}$ is a power series divisible by $x^{|\mathbf{a}|}$, the right hand side of (2.3)contains only non-negative powers of x. The range of the sum is a bit complicated, the reader should just keep in mind that Q_L has the above form with appropriate restrictions on the indices and on denominators of the coefficients.

Proof. The proof consists of a few changes of variables. It is well known that if a polynomial $p(t_1, \ldots, t_s)$ takes integer values whenever t_i 's are integers, then $p(t_1, \ldots, t_s)$ is a \mathbb{Z} -linear combination of terms of the form $\binom{t_1}{l_1} \ldots \binom{t_s}{l_s}$, where l_1, \ldots, l_s are non-negative integers.

Recall that $(\alpha_i | \lambda_j) = \delta_{ij}$. For $\mathbf{a} = (a_1, \dots, a_{n-1}) \in (\mathbb{Z}_+)^{n-1}$, let

$$\begin{pmatrix} \boldsymbol{\alpha} \\ \mathbf{a} \end{pmatrix} = \prod_{i=1}^{n-1} \begin{pmatrix} \alpha_i \\ a_i \end{pmatrix}$$

By Proposition 2.4, $f_{\ell}(\mu_1, \ldots, \mu_m)$ is a polynomial function on μ_1, \ldots, μ_m which takes integer values whenever $\mu_j \in \Lambda$. Hence

$$f_{\ell}(\mu_1,\ldots,\mu_m) = \sum^{(\ell)} c'_{\mathbf{a}_1,\ldots,\mathbf{a}_m;\ell} \begin{pmatrix} \boldsymbol{\alpha} \\ \mathbf{a}_1 \end{pmatrix} (\mu_1) \ldots \begin{pmatrix} \boldsymbol{\alpha} \\ \mathbf{a}_m \end{pmatrix} (\mu_m), \qquad (2.7)$$

10

with $c'_{\mathbf{a}_1,\ldots,\mathbf{a}_m;\ell}$ in \mathbb{Z} . Here $\sum^{(\ell)}$ means the sum is over the set of all $\mathbf{a}_j \in (\mathbb{Z}_+)^{n-1}$ and $\ell \in \mathbb{Z}_+$ satisfying (2.4), (2.5) and (2.6) These restrictions on $|\mathbf{a}_j|$ and ℓ follows from the restriction on the degree of f'_{ℓ} (see Proposition 2.4, here $|\mathbf{a}_j| = s_j + n(n-1)$).

Recall that $\eta_i = 1 - (1 + x)^{-\alpha_i}$. We can easily express α_i in terms of x and η_i :

$$\alpha_i = \frac{\ln(1 - \eta_i)}{\ln(1 + x)},$$

and hence

$$x\alpha_{i} = \frac{x\ln(1-\eta_{i})}{\ln(1+x)}.$$
(2.8)

Note that the right hand side can be expressed as a formal power series in x and η_i ; moreoever, the coefficient of a term of total degree d (in x and η_i) is in $\mathbb{Z}_{(d+1)}$.

Using (2.7) in formula (2.2), we get

$$Q_{L}(\mu_{1},...,\mu_{m};q)\Big|_{q=x+1} = \sum_{\ell=0}^{\infty} \sum^{(\ell)} c'_{\mathbf{a}_{1},...,\mathbf{a}_{m};\ell} x^{\ell} \prod_{j=1}^{m} \binom{\boldsymbol{\alpha}}{\mathbf{a}_{j}} (\mu_{j})$$
$$= \sum_{\ell=0}^{\infty} \sum^{(\ell)} c'_{\mathbf{a}_{1},...,\mathbf{a}_{m};\ell} x^{\ell-\sum_{j} |\mathbf{a}_{j}|} \prod_{j=1}^{m} \binom{\boldsymbol{\alpha}}{\mathbf{a}_{j}} (\mu_{j}) x^{|\mathbf{a}_{j}|}$$

Using the change of variable (2.8) in the above formula, with some simple calculation, we get (2.3). The denominators of the coefficients come from the denominators of the expressions $\begin{pmatrix} \alpha \\ \mathbf{a}_j \end{pmatrix}$ and from the denominators of the right hand side of (2.8). It is easy to check that $c_{\mathbf{a}_1,\ldots,\mathbf{a}_m;\ell}$ is in $\mathbb{Z}_{(2\ell+(m+1)n(n-1)-\sum |\mathbf{a}_j|+1)}$.

2.4. Linear operator Γ_b . We now introduce the linear operator Γ_b which is a kind of Laplace transform. For $\mu \in \Lambda$ let

$$\psi(\mu;q) = \prod_{\alpha \in \Phi_+} (1 - q^{-(\mu|\alpha)}) = \prod_{\alpha \in \Phi_+} (1 - q^{-\alpha})(\mu).$$
(2.9)

This function plays important role in Lie theory, it appeared in Weyl's character formula.

Let \mathcal{S} be the free \mathbb{Z} -module generated by q^{β} , with $\beta \in \Lambda^{\text{root}}$. For each integer b not divisible by the odd prime r let Γ_b be the linear operator

$$\Gamma_b \colon \mathcal{S} \to \mathbb{Z}[\frac{1}{n!b}][[x]], \quad \text{defined by}$$

 $\Gamma_b(q^\beta) = (1+x)^{(-|\beta|^2/2b)} y_b.$

Here y_b is an element in $\mathbb{Z}[\frac{1}{n!}][[x]]$, not depending on β , and is given by

$$y_b = \frac{1}{n!} \left(\frac{|b|}{r}\right)^{n-1} (1+x)^{\frac{\operatorname{sn}(b)-b}{2}|\rho|^2} \psi\left(-\operatorname{sn}(b)\rho; 1+x\right),$$

(sn(b) is the sign of b).

Note that $\eta^{\mathbf{a}}$ is in \mathcal{S} , hence there is defined $\Gamma_b(\eta^{\mathbf{a}})$. A property of Γ_b is stated in the following proposition, whose proof will be presented in §4.6.

Proposition 2.6. The power series $\Gamma_b(\boldsymbol{\eta}^{\mathbf{a}})$ is divisible by $x^{\lfloor (n(n-1)+|\mathbf{a}|+1)/2 \rfloor}$. Here $\lfloor z \rfloor$ is the integer part of z.

2.5. Formula of perturbative invariants $\tau^{PSU(n)}$. We continue to assume that L has 0 linking matrix with m components. Suppose that L' is the same as L, except that the framing on components are *non-zero* integers b_1, \ldots, b_m . Suppose for L we have the expansion (2.3) of Q_L .

Let $\tau(L')$ be obtained from Q_L (i.e. the right hand side of (2.3)) by replacing $\eta^{\mathbf{a}_j}(\mu_j)$ by $\Gamma_{b_j}(\eta^{\mathbf{a}_j})$:

$$\tau(L') = \sum c_{\mathbf{a}_1,\dots,\mathbf{a}_m;\ell} \Gamma_{b_1}(\boldsymbol{\eta}^{\mathbf{a}_1}) \dots \Gamma_{b_m}(\boldsymbol{\eta}^{\mathbf{a}_m}) x^{\ell - \sum_j |\mathbf{a}_j|}.$$
 (2.10)

The range of the sum is the same as in formula (2.3), i.e. the sum is over the set of all $\mathbf{a}_j \in \mathbb{Z}_+^{n-1}$ and $\ell \in \mathbb{Z}_+$ satisfying (2.4), (2.5) and (2.6).

By Proposition 2.6, the term $\Gamma_{b_1}(\boldsymbol{\eta}^{\mathbf{a}_1}) \dots \Gamma_{b_m}(\boldsymbol{\eta}^{\mathbf{a}_m}) x^{\ell - \sum_j |\mathbf{a}_j|}$ is divisible by the power of x with exponent

$$\ell - \sum_{j} |\mathbf{a}_{j}| + \frac{mn(n-1)}{2} + \sum_{j} \lfloor \frac{|\mathbf{a}_{j}|}{2} \rfloor,$$

which is greater than or equal to $\ell - \sum_j |\mathbf{a}_j|/2 + mn(n-1)/2$. The latter, by (2.6), is greater than or equal to $\frac{1}{2} \max_j (|\mathbf{a}_j| - n(n-1))$. For each fixed number a, there are only a finite number of $\mathbf{a}_j \in (\mathbb{Z}_+)^{n-1}$ such that $\frac{1}{2} \max_j (|\mathbf{a}_j| - n(n-1)) < a$. Hence the right hand side of (2.10) is really a formal power series in x.

The denominators of coefficients of $\tau(L')$ come from the factor $n!b_j$ and the denominators of $c_{\mathbf{a}_1,\ldots,\mathbf{a}_m;\ell}$. Hence, using the property of the denominators of $c_{\mathbf{a}_1,\ldots,\mathbf{a}_m;\ell}$ in Proposition 2.5 one can easily shows that in $\tau(L')$ the coefficient of x^{ℓ} is in $\mathbb{Z}[\frac{1}{(2\ell+n(n-1))!b_1\ldots b_m}]$. In other words, $\tau(L')$ is in $\mathbb{Q}[[x]]_M$, where M is the 3-manifold obtained by surgery along L'.

We will show that $\tau(L')$ is an invariant of M, which is a rational homology 3-sphere. Later in §4.5 we will show that the constant term of the power series $\tau(L')$ is

$$\left(\frac{|H_1(M,\mathbb{Z})|}{r}\right)|H_1(M,\mathbb{Z})|^{-n(n-1)/2}$$

Hence there always exists the inverse $\tau(L')^{-1} \in \mathbb{Q}[[x]]_{(M)}$.

Let M be an arbitrary rational homology 3-sphere. Obtsuki showed (see [24], see also [23]) that there are lens spaces $M(d_1), \ldots, M(d_s)$ such that $M \# M(d_1) \# \ldots \# M(d_s)$ can be obtained from S^3 by surgery on a link L' with diagonal linking matrix. Here M(d) is the lens space obtained by surgery along U_d , the unknot with framing d. Moreover each d_i is less than or equal to $|H_1(M;\mathbb{Z})|$. Define

$$\tau^{PSU(n)}(M) = \tau(L') \, \tau(U_{d_1})^{-1} \dots \tau(U_{d_s})^{-1}$$

We have not proved that $\tau^{PSU(n)}(M)$ is an invariant of M yet. So, for the time being, $\tau^{PSU(n)}(M)$ means any formal power series obtained by the above procedure, which a priori depends on the choice of d_1, \ldots, d_s and L'.

The main theorem can be formulated in a more precise form as follows.

Theorem 2.7. Suppose M is rational homology 3-sphere, and r a prime number greater than $\max\{n(n-1), |H(M,\mathbb{Z})|\}$. Then

$$\mathfrak{p}_r\left(\tau_r^{PSU(n)}(M)\right) = \mathfrak{p}_r\left(\tau^{PSU(n)}(M)\right)$$

Now the uniqueness of Lemma 1.6, with this theorem, shows that $\tau^{PSU(n)}(M)$ is really an invariant of the 3-manifold M, i.e. it does not depend on the choice of d_1, \ldots, d_s and L' in the construction of $\tau^{PSU(n)}(M)$.

Since the series $\Gamma_b(\boldsymbol{\eta}^{\mathbf{a}})$ has coefficients in $\mathbb{Z}[\frac{1}{n!b}]$, we see that if all the coefficients $c_{\mathbf{a}_1,\dots,\mathbf{a}_m;\ell}$ are in $\mathbb{Z}[\frac{1}{n!}]$, then the power series $\tau^{PSU(n)}(M)$ has coefficients in $\mathbb{Z}[\frac{1}{n!H_1(M,\mathbb{Z})}]$. Hence Conjecture 1.9 follows from the following conjecture.

Conjecture 2.8. All the coefficients $c_{\mathbf{a}_1,\ldots,\mathbf{a}_m;\ell}$ are in $\mathbb{Z}[\frac{1}{n!}]$.

This conjecture, for the special case n = 2 and in a slightly different form, had been formulated by Rozansky. He also proved that the conjecture holds true in the special case n = 2 and the link L has one component (a knot), using explicit formula of the *R*-matrix (see [30]).

2.6. **Proof of the main theorem.** Case 1: M is obtained by surgery along a framed link L' with diagonal linking matrix and $\tau^{PSU(n)}(M) = \tau(L')$. Let L be the same link as L' except that the framing of each component is 0. Suppose that the framing on components of L' are b_1, \ldots, b_m . We will fix an odd prime number r greater than $\max\{n(n-1), |H_1(M,\mathbb{Z})|\}$. Let us consider the expression (2.3) of $Q_L(\mu_1, \ldots, \mu_m; x+1)$, and let $Q_L^{(1)}$ be the part of Q_L in (2.3), where the indices satisfy

$$2\ell - \sum_{j} (|\mathbf{a}_{j}| - n(n-1)) \le r - n(n-1).$$
(2.11)

In other words,

$$Q_L^{(1)}(\mu_1, \dots, \mu_m; x+1) = \sum_{j=1}^{n} c_{\mathbf{a}_1, \dots, \mathbf{a}_m; \ell} \, \boldsymbol{\eta}^{\mathbf{a}_1}(\mu_1) \dots \, \boldsymbol{\eta}^{\mathbf{a}_m}(\mu_m) x^{\ell - \sum_j |\mathbf{a}_j|},$$
(2.12)

where $\sum^{(1)}$ means the sum over the set of all $\mathbf{a}_j \in \mathbb{Z}^{n-1}_+$ and $\ell \in \mathbb{Z}_+$ satisfying (2.4), (2.5), (2.6) and (2.11). The reason we choose the new restriction (2.11) is that, by Proposition 2.5, all the coefficients $c_{\mathbf{a}_1,\ldots,\mathbf{a}_m;\ell}$ in (2.12) are in $\mathbb{Z}_{(r)}$. Note that $\sum^{(1)}$ is a finite sum.

Lemma 2.9. There is the following splitting for Q_L (with q = x + 1):

$$Q_L(\mu_1,\ldots,\mu_m;q) = Q_L^{(1)}(\mu_1,\ldots,\mu_m;q) + Q_L^{(2)}(\mu_1,\ldots,\mu_m;q) + Q_L^{(3)}(\mu_1,\ldots,\mu_m;q)$$

in which :
a)
$$Q_L^{(1)}$$
 is defined by (2.12),
b) $Q_L^{(2)}$ is a polynomial in $(q-1) = x$ of the form
 $Q_L^{(2)}(\mu_1, \dots, \mu_m; q) = \sum_{m=1}^{(2)} \tilde{c}_{\mathbf{a}_1, \dots, \mathbf{a}_m; \ell} \begin{pmatrix} \boldsymbol{\alpha} \\ \mathbf{a}_1 \end{pmatrix} (\mu_1) \dots \begin{pmatrix} \boldsymbol{\alpha} \\ \mathbf{a}_m \end{pmatrix} (\mu_m) x^{\ell},$
(2.13)

where $\Sigma^{(2)}$ is the sum over the set of all $\mathbf{a}_j \in \mathbb{Z}^{n-1}_+$ and $\ell \in \mathbb{Z}_+$ satisfying (2.4), (2.5), (2.6), and, in addition,

$$2\ell - \sum_{j} (|\mathbf{a}_{j}| - n(n-1)) > r - n(n-1), \qquad (2.14)$$

$$\ell < N. \tag{2.15}$$

c) $Q_L^{(3)}(\mu_1, \dots, \mu_m; q)$ is a polynomial in q, q^{-1} which is divisible by $(q-1)^N$ in $\mathbb{Z}_{(r)}[q, q^{-1}]$. Here $N = m(r-n-1)(n-1)/2 + \bar{r} + 1$.

Note that (2.11) and (2.14) are complementary.

Proof. With the substitution q = x + 1 and

$$\eta_i(\mu) = 1 - (1+x)^{-(\alpha_i|\mu)}, \qquad (2.16)$$

 $Q_L^{(1)}$ becomes a formal power series in x. The right hand side of (2.16) is a power series with *integer* coefficients, hence the denominators of the power series $Q_L^{(1)}$ comes from the coefficients $c_{\mathbf{a}_1,\ldots,\mathbf{a}_m;\ell}$ which are in $\mathbb{Z}_{(r)}$. Hence $Q_L^{(1)}$ is a power series in x with coefficients in $\mathbb{Z}_{(r)}$. On the other hand, Q_L is a power series in x with integer coefficients. Hence, for fixed $\mu_1,\ldots,\mu_m, Q_L - Q_L^{(1)}$ is a power series in x with coefficients in $\mathbb{Z}_{(r)}$.

It follows that

$$Q_L - Q_L^{(1)} = \sum \tilde{c}_{\mathbf{a}_1,\dots,\mathbf{a}_m;\ell} \begin{pmatrix} \boldsymbol{\alpha} \\ \mathbf{a}_1 \end{pmatrix} (\mu_1)\dots, \begin{pmatrix} \boldsymbol{\alpha} \\ \mathbf{a}_m \end{pmatrix} (\mu_m) x^{\ell}, \qquad (2.17)$$

where the sum is over the set of all $\mathbf{a}_j \in \mathbb{Z}_+^{n-1}$ and $\ell \in \mathbb{Z}_+$ satisfying (2.4), (2.5), (2.6) and (2.14), and the coefficients $\tilde{c}_{\mathbf{a}_1,\ldots,\mathbf{a}_m;\ell}$ are in $\mathbb{Z}_{(r)}$.

Let us define $Q_L^{(2)}$ by using the right hand side of the (2.17), imposing additional restriction (2.15) on the indices \mathbf{a}_j , ℓ . It is easy to see that there are only a finite number of terms in $Q_L^{(2)}$. Finally, let $Q_L^{(3)} = Q_L - Q_L^{(1)} - Q_L^{(2)}$. A term of the right hand side of (2.17), for which (2.15) is not satisfied, must be divisible by the power of x with exponent $\ell + 1$ which is greater than or equal to N. Hence $Q_L^{(3)}$ is divisible by x^N .

 $\ell + 1$ which is greater than or equal to N. Hence $Q_L^{(3)}$ is divisible by x^N . Let us return to the indeterminate q = x+1. The advantage of using this indeterminate is that all $Q_L, Q_L^{(1)}, Q_L^{(2)}, Q_L^{(3)}$ are *polynomials* in q, q^{-1} (i.e. they don't have infinite order), which can be seen as follows.

First of all, Q_L is a polynomial in q and q^{-1} with coefficients in \mathbb{Z} , by Proposition 2.1. For fixed μ_1, \ldots, μ_m , replacing x = q - 1 in the expression of $Q_L^{(2)}$, we see that $Q_L^{(2)}$ is a polynomial in q with coefficients in $\mathbb{Z}_{(r)}$.

Note that, for i = 1, ..., n-1, the number $(\alpha_i | \mu)$ is always an integer for every $\mu \in \Lambda$. Hence $\eta_i(\mu) = 1 - q^{-(\alpha_i|\mu)}$ is a polynomial in q, q^{-1} with integer coefficients, and η_i is divisible by (q-1) in $\mathbb{Z}[q, q^{-1}]$. Now formula (2.12) shows that $Q_L^{(1)}$ is a polynomial in q, q^{-1} with coefficients in $\mathbb{Z}_{(r)}$.

It follows that the remaining $Q_L^{(3)}$ is also a polynomial in q, q^{-1} with coefficients in $\mathbb{Z}_{(r)}$. Moreover, $Q_L^{(3)}$ is divisible by $(q-1)^N$ in $\mathbb{Z}_{(r)}[q, q^{-1}]$:

$$Q_L^{(3)}(\mu_1,\ldots,\mu_m;q) = (q-1)^N \sum_{s=-d}^d g(\mu_1,\ldots,\mu_m)q^s,$$

for some positive integer d. Here $g(\mu_1, \ldots, \mu_m)$, possibly not a polynomial function, takes values in $\mathbb{Z}_{(r)}$ when $\mu_1, \ldots, \mu_m \in \Lambda$. This completes the proof of the proposition.

Recall that L' differs from L only by the framing on components. By formula (2.1):

$$L'(\mu_1,\ldots,\mu_m;q) = \prod_{j=1}^m q^{b_j \frac{|\mu_j|^2 - |\rho|^2}{2}} Q_L(\mu_1,\ldots,\mu_m;q),$$

and hence

$$F_{L'}(r) = \sum_{\mu_j \in (\rho + \Lambda_r^{\text{root}})} \prod_{j=1}^m q^{b_j \frac{|\mu_j|^2 - |\rho|^2}{2}} Q_L(\mu_1, \dots, \mu_m; \zeta).$$
(2.18)

Let $F_{L'}^{(1)}(r), F_{L'}^{(2)}(r), F_{L'}^{(3)}(r)$ be the value of the right hand side of (2.18), if we replace Q_L by, respectively, $Q_L^{(1)}, Q_L^{(2)}, Q_L^{(3)}$. Then we have

$$F_{L'} = F_{L'}^{(1)} + F_{L'}^{(2)} + F_{L'}^{(3)}, \quad \text{and}$$
$$\tau_r^{PSU(n)}(M) = \frac{F_{L'}(r)}{F_{U_+}(r)^{\sigma_+} F_{U_-}(r)^{\sigma_-}}$$
$$= \tau^{(1)} + \tau^{(2)} + \tau^{(3)},$$

where

$$\tau^{(j)} = \frac{F_{L'}^{(j)}(r)}{F_{U_+}(r)^{\sigma_+} F_{U_-}(r)^{\sigma_-}}$$

The following 3 lemmas prove the main theorem (Theorem 2.7) in case 1.

Lemma 2.10. The number $\tau^{(3)}$ is divisible by $(\zeta - 1)^{\overline{r}+1}$ in $\mathbb{Z}_{(r)}[\zeta]$. Hence $\mathfrak{p}_r(\tau^{(3)}) = 0$.

Lemma 2.11. The number $\tau^{(2)}$ is divisible by $(\zeta - 1)^{\bar{r}+1}$ in $\mathbb{Z}_{(r)}[\zeta]$. Hence $\mathfrak{p}_r(\tau^{(2)}) = 0$.

Lemma 2.12. One has that $\mathfrak{p}_r(\tau^{(1)}) = \mathfrak{p}_r(\tau(L')).$

The expression $F_{U_{\pm}}(r)$ appears in the denominators of the formulas of $\tau^{(j)}$. We will need the following proposition whose proof will be given in §4.

Proposition 2.13. Each of $F_{U_+}(r)$ and $F_{U_-}(r)$ is proportional to $(\zeta - 1)^{(n-1)(r-n-1)/2}$ by a proportional factor which is a unit in $\mathbb{Z}_{(1)}[\zeta] = \mathbb{Z}[\frac{1}{n!}][\zeta]$.

Proof. [Proof of Lemma 2.10] Since each b_j is non-zero, $\sigma_+ + \sigma_- = m$. By Proposition 2.13, $F_{U_+}(r)^{\sigma_+}F_{U_-}(r)^{\sigma_-}$ is proportional to $(\zeta - 1)^{m(n-1)(r-n-1)/2}$ by a unit in $\mathbb{Z}_{(r)}[\zeta]$. But $F_{L'}^{(3)}(r)$ is divisible by $(\zeta - 1)^N$ in $\mathbb{Z}_{(r)}[\zeta]$ by Lemma 2.9. Hence

$$\tau^{(3)} = \frac{F_{L'}^{(3)}(r)}{F_{U_+}(r)^{\sigma_+}F_{U_-}(r)^{\sigma_-}}$$

is divisible by the power of $\zeta - 1$ with exponent $N - m(n-1)(r-n-1)/2 = \bar{r} + 1$. \Box

To prove Lemma 2.11 we will need the following lemma whose proof will be given in $\S4$.

Lemma 2.14. Let $\mathbf{a} \in (\mathbb{Z}_+)^{n-1}$ and $b \in \mathbb{Z}_+$. Then the number

$$\sum_{\mu \in \rho + \Lambda_r^{root}} \zeta^{b\frac{|\mu|^2 - |\rho|^2}{2}} \binom{\boldsymbol{\alpha}}{\mathbf{a}}(\mu)$$

is proportional to $(\zeta - 1)^{(n-1)(r-1)/2 - \lfloor |\mathbf{a}|/2 \rfloor}$ by a factor in $\mathbb{Z}_{(r)}[\zeta]$.

Proof. [Proof of Lemma 2.11] Using the expression (2.13) we have that

$$F_{L'}^{(2)}(r) = \sum_{\mu_j \in \rho + \Lambda_r^{\text{root}}} \prod_{j=1}^m \zeta^{b_j \frac{|\mu_j|^2 - |\rho|^2}{2}} \sum^{(2)} \tilde{c}_{\mathbf{a}_1, \dots, \mathbf{a}_m; \ell} \binom{\boldsymbol{\alpha}}{\mathbf{a}_1} (\mu_1) \dots \binom{\boldsymbol{\alpha}}{\mathbf{a}_m} (\mu_m) x^\ell$$
$$= \sum^{(2)} \tilde{c}_{\mathbf{a}_1, \dots, \mathbf{a}_m; \ell} x^\ell \prod_{j=1}^m \sum_{\mu_j \in \rho + \Lambda_r^{\text{root}}} \zeta^{b_j \frac{|\mu_j|^2 - |\rho|^2}{2}} \binom{\boldsymbol{\alpha}}{\mathbf{a}_j} (\mu_j)$$

Applying Lemma 2.14, we see that $F_{L'}^{(2)}(r)$ is divisible, in $\mathbb{Z}_{(r)}[\zeta]$, by the power of $\zeta - 1$ with the exponent

$$\ell + m \frac{(n-1)(r-1)}{2} - \sum_{j} \lfloor \frac{|\mathbf{a}_j|}{2} \rfloor,$$

which, by (2.14), is greater than or equal to N. As in the proof of Lemma 2.10, it follows that $\tau^{(2)}$ is divisible by $(\zeta - 1)^{\bar{r}+1}$.

In order to prove Lemma 2.12, we will need the following proposition, whose proof will be presented in §4.

Proposition 2.15. Let

$$g_b(\mathbf{a}) = \frac{\sum_{\mu \in (\rho + \Lambda_r^{root})} \zeta^{b^{\frac{|\mu|^2 - |\rho|^2}{2}}} \boldsymbol{\eta}^{\mathbf{a}}(\mu)}{F_{U_{\mathrm{sn}(b)}}(r)}$$

Then $\mathfrak{p}_r(g_b(\mathbf{a})) = \mathfrak{p}_r(\Gamma_b(\boldsymbol{\eta}^{\mathbf{a}})).$

Proof. [Proof of Lemma 2.12]. Recall that

$$Q_L^{(1)} = \sum_{\mu} {}^{(1)} c_{\mathbf{a}_1,\dots,\mathbf{a}_m;\ell} \, \boldsymbol{\eta}^{\mathbf{a}_1}(\mu_1) \dots \, \boldsymbol{\eta}^{\mathbf{a}_m}(\mu_m) \, (q-1)^{\ell - \sum_j |\mathbf{a}_j|}.$$

Here $c_{\mathbf{a}_1,\ldots,\mathbf{a}_m;\ell} \in \mathbb{Z}_{(r)}$. Hence

$$F_{L'}^{(1)}(r) = \sum_{j=1}^{n} c_{\mathbf{a}_1,\dots,\mathbf{a}_m;\ell} (\zeta - 1)^{\ell - \sum_j |\mathbf{a}_j|} \prod_{j=1}^m \sum_{\mu_j \in (\rho + \Lambda_r^{\text{root}})} \zeta^{b_j \frac{|\mu_j|^2 - |\rho|^2}{2}} \boldsymbol{\eta}^{\mathbf{a}_j}(\mu_j),$$

and

$$\tau^{(1)} = \frac{F_{L'}^{(1)}(r)}{F_{U_+}(r)^{\sigma_+}F_{U_-}(r)^{\sigma_-}}$$
$$= \sum_{i=1}^{(1)} c_{\mathbf{a}_1,\dots,\mathbf{a}_m;\ell} (\zeta - 1)^{\ell - \sum_j |\mathbf{a}_j|} \prod_{j=1}^m g_{b_j}(\mathbf{a}_j).$$

By Proposition 2.15, $\mathfrak{p}_r(g_b(\mathbf{a})) = \mathfrak{p}_r(\Gamma_b(\boldsymbol{\eta}^{\mathbf{a}}))$. It follows that

$$\mathfrak{p}_r(\tau^{(1)}) = \mathfrak{p}_r\left(\sum_{a_1,\dots,a_m;\ell} x^{\ell-\sum_j |\mathbf{a}_j|} \prod_{j=1}^m \Gamma_{b_j}(\boldsymbol{\eta}^{\mathbf{a}_j})\right).$$

The argument of \mathfrak{p}_r on the right hand side of this formula, by definition, is a part of $\tau(L')$ (see formula (2.10) for the definition of $\tau(L')$). This part differs from $\tau(L')$ only by elements of degree $> \bar{r}$, which are annihilated by \mathfrak{p}_r , hence

$$\mathfrak{p}_r(\tau^{(1)}) = \mathfrak{p}_r(\tau(L')).$$

This completes the proof of Lemma 2.12, and hence that of Main theorem in case 1. \Box

Case 2: M is an arbitrary rational homology 3-sphere, $M' = M \# M(d_1) \# \dots \# M(d_s)$ is obtained by surgery along a link L' with diagonal linking matrix, and

$$\tau^{PSU(n)}(M) = \tau(L')\tau(U_{d_1})^{-1}\dots\tau(U_{d_s})^{-1}.$$

The multiplicative property of quantum invariants says that

$$\tau_r^{PSU(n)}(M') = \tau_r^{PSU(n)}(M) \,\tau_r^{PSU(n)}(M(d_1)) \dots \tau_r^{PSU(n)}(M(d_s)).$$
(2.19)

Applying the result of case 1 to L' and U_{d_1}, \ldots, U_{d_s} , and using (2.19) we see that $\mathfrak{p}_r(\tau_r^{PSU(n)}(M)) = \mathfrak{p}_r(\tau_r^{PSU(n)}(M))$. This completes the proof of the main theorem.

3. QUANTUM INVARIANTS OF LINKS AND THE KONTSEVICH INTEGRAL

In this section we always assume that L is a framed oriented link of m components, and that r is a positive integer. We will investigate the dependence of the quantum invariant $J_L(\mu_1, \ldots, \mu_m; q)$ on the μ_j 's and prove Propositions 1.3, 2.1, and 2.4.

3.1. Some general facts about quantum invariants. Recall that $J_L(\mu_1, \ldots, \mu_m; q)$, or $J_L(V_{\mu_1}, \ldots, V_{\mu_m}; q)$ is the quantum invariant of L, when the components of L are colored by the sl_n -modules $V_{\mu_1}, \ldots, V_{\mu_m}$. Here V_{μ} is the irreducible module of highest weight $\mu - \rho$, for $\mu \in \Lambda_{++}$.

Recall that for $\mu \in \Lambda$,

$$\psi(\mu;q) = \prod_{\alpha \in \Phi_+} (1 - q^{-(\mu|\alpha)}) = \prod_{\alpha \in \Phi_+} (1 - q^{-\alpha})(\mu).$$

From the Weyl denominator formula (see, for example, [5]), we have the following.

Lemma 3.1. One has that
$$\psi(\mu;q) = \sum_{w \in \mathcal{W}} \operatorname{sn}(w) q^{(\mu|w(\rho)-\rho)}$$

The value of the quantum invariant J_L of the unknot U is (see, for example [4]):

$$J_U(\mu;q) = q^{(\mu-\rho|\rho)} \frac{\psi(\mu;q)}{\psi(\rho;q)}.$$
(3.1)

Note that $(\mu|\alpha) \in \mathbb{Z}$ for $\mu \in \Lambda$ and $\alpha \in \Lambda^{\text{root}}$. Moreover, it's known that $2\rho \in \Lambda^{\text{root}}$, and hence $(\mu|\rho) \in \frac{1}{2}\mathbb{Z}$. It follows that $J_U(\mu;q)$ is either in $q^{1/2}\mathbb{Z}[q,q^{-1}]$ or in $\mathbb{Z}[q,q^{-1}]$.

Let $L^{(2)}$ be the framed link obtained from L by replacing the first component by two its parallel push-offs (using the frame). Suppose that the color of these 2 push-offs are modules V and V'. Then we have the following formula (see [33]):

$$J_{L^{(2)}}(V,V',\ldots;q) = J_L(V \otimes V',\ldots;q)$$

Hence for Q_L we have a similar result:

$$Q_{L^{(2)}}(V, V', \dots; q) = Q_L(V \otimes V', \dots; q).$$
 (3.2)

In particular, when L = U, noting that $U^{(2)}$ is the trivial link with 2 components, we have that

$$J_U(V \otimes V'; q) = J_{U^{(2)}}(V, V'; q)$$
$$= J_U(V; q) J_U(V'; q)$$

This means

$$J_U(\cdot;q): \mathcal{R} \to \mathbb{Z}[q^{\pm 1/2n}]$$

is a *ring* homomorphism. Recall that \mathcal{R} is the ring of finite-dimensional sl_n -modules. Let I_r be the ideal in \mathcal{R} generated by all the V_{μ} , with $(\mu|\theta) = r$, (here $\mu \in \Lambda_{++}$). Recall that $\theta = \alpha_1 + \cdots + \alpha_{n-1}$ is the longest root. The quotient ring \mathcal{R}/I_r is known as the fusion algebra (see [1, 7] and reference therein).

Lemma 3.2. If μ is in I_r , then $J_U(\mu; \zeta) = 0$.

Proof. Since $J_U(\cdot; \zeta) : \mathcal{R} \to \mathbb{C}$ is a ring homomorphism, it suffices to show that $J_U(\mu; \zeta) = 0$ if $(\mu|\theta) = r$. We have that

$$1 - \zeta^{-(\mu|\theta)} = 1 - \zeta^{-r} = 0.$$

Since θ is one of the positive roots, it follows from the definition of ψ that $\psi(\mu; \zeta) = 0$, and hence $J_U(\mu; \zeta) = 0$.

Proposition 3.3. Suppose that one of μ_1, \ldots, μ_m is in I_r . Then $J_L(\mu_1, \ldots, \mu_m; \zeta) = 0$.

Proof. A proof for the case n = 2 is given in [9], Lemma 3.29. Suppose $\mu_1 \in I_r$. The argument of Lemma 3.29 of [9] shows that $J_L(\mu_1, \ldots, \mu_m; \zeta)$ is proportional to $J_U(\mu_1; \zeta)$ which is 0. Hence $J_L(\mu_1, \ldots, \mu_m; \zeta) = 0$.

Remark. This proposition also follows from the theory of the Kontsevich integral, more precisely, from Proposition 3.6 and formula (3.8) below.

3.2. **Proof of Proposition 1.3.** We will fix μ_2, \ldots, μ_m and will write $J_L(\mu_1; q)$ instead of $J_L(\mu_1, \mu_2, \ldots, \mu_m; q)$.

Proposition 3.3 shows that $J_L(\mu_1; \zeta) = J_L(\mu'_1; \zeta)$ if $\mu_1 \equiv \mu'_1 \pmod{I_r}$. From the theory of fusion algebra it is known that for every $w \in \mathcal{W}_{(r)}$,

$$V_{\mu} \equiv \operatorname{sn}(w)V_{w(\mu)} \pmod{I_r}.$$

Hence

$$J_L(\mu;\zeta) = \operatorname{sn}(w)J_L(w(\mu);\zeta).$$

18

Put L = U we get

$$J_U(\mu;\zeta) = \operatorname{sn}(w)J_U(w(\mu);\zeta).$$

Take the product of the last two identities, we get

$$Q_L(\mu;\zeta) = Q_L(w(\mu);\zeta),$$

which proves part a) of Proposition 1.3. Part b) follows from a) and Proposition 3.3.

3.3. **Proof of proposition 2.1.** We have to show that $Q_L(\mu_1, \ldots, \mu_m; q) \in \mathbb{Z}[q^{\pm 1}]$ if the linking matrix of L is 0. It's enough to consider the case when $\mu_j \in \Lambda_{++}$, since Q_L is invariant under the action of the Weyl group and equal to 0 if one of the μ_j is on the boundary of the fundamental chamber C.

The invariant Q_L satisfies the doubling formula (3.2), and the representation ring \mathcal{R} is generated (as algebra over \mathbb{Z}) by $V_{\lambda_1+\rho}, \ldots, V_{\lambda_{n-1}+\rho}$. Hence it suffices to prove that $Q_L(\mu_1, \ldots, \mu_m; q) \in \mathbb{Z}[q^{\pm 1}]$ for the case when each μ_j is in $\{\lambda_1 + \rho, \ldots, \lambda_{n-1} + \rho\}$. This fact follows immediately from the following lemma.

Lemma 3.4. Suppose that the linking matrix of L is 0 and $\mu_j = \lambda_{k_j} + \rho$, with $j = 1, \ldots, m$. Then $J_L(\mu_1, \ldots, \mu_m; q)$ is in $q^{(k_{j_1} + \cdots + k_{j_m})\frac{n-1}{2}}\mathbb{Z}[q, q^{-1}]$.

Remark. Without the assumption that the linking matrix is 0, the lemma does not hold true.

Proof. [Proof of Lemma 3.4] For any framed oriented link K let us defined $J_K^{Hom}(q)$ as follows. First let K' be any link obtained from K by changing the framings so that $K' \cdot K' = 0$, i.e. the sum of all the entries of the linking matrix of K' is 0. Then let $J_K^{Hom}(q) = J_{K'}(\mu_1, \ldots, \mu_m; q)$ with all the μ_j equal $\lambda_1 + \rho$ (i.e. when each V_{μ_j} is the fundamental representation of sl_n). This is a non-framed version of quantum invariants. For the link L with 0 linking matrix we choose L' = L. It is known that J^{Hom} is a version of the Homfly polynomial and can be calculated by the skein theory as follows.

$$J_{L_1 \sqcup L_2}^{Hom}(q) = J_{L_1}^{Hom}(q) J_{L_2}^{Hom}(q),$$
$$q^{n/2} J_{L_+}^{Hom} - q^{-n/2} J_{L_-}^{Hom} = (q^{1/2} - q^{-1/2}) J_{L_0}^{Hom},$$

where (L_0, L_-, L_+) is the standard skein triple. By using the skein relation and induction on the number of crossing points of link diagram, one easily proves

Lemma 3.5. If L has m components, then $J_L^{Hom}(q) \in q^{m(n-1)/2}\mathbb{Z}[q^{\pm 1}]$.

This shows that Lemma 3.4 holds true when all the μ_j 's are equal to $\lambda_1 + \rho$.

The case when all the μ_j 's are in the set $\{\lambda_1 + \rho, \dots, \lambda_{n-1} + \rho\}$ can be reduced to the case when all μ_j are equal to $\lambda_1 + \rho$ by using cabling as follows. (See similar arguments in Appendix 1 of [22]).

Let B_k be the braid group on k strands. In what follows we will consider linear combinations of elements of B_k with coefficients in $\mathbb{Z}(q^{1/2})$, the ring of rational functions in $q^{1/2}$ with integer coefficients. The anti-symmetrizer $g^{(k)}$ is defined, for $k = 1, \ldots, n-1$, by induction as follows (see [36]): $g^{(1)} = 1 \in B_1$,

$$g^{(k)} = \frac{1 - q^{-1}}{1 - q^{-k}} (g^{(k-1)} \otimes 1) - q^{(n-1)/2} \frac{1 - q^{1-k}}{1 - q^{-k}} (g^{(k-1)} \otimes 1) \sigma_{k-1} (g^{(k-1)} \otimes 1),$$
(3.3)

where $\sigma_1, \ldots, \sigma_{k-1}$ are the standard generators of the braid group B_k , and $z \otimes 1$ is obtained from z by adding a vertical strand to the right of the braid z.

Now suppose $\mu_j = \lambda_{k_j} + \rho$, where $1 \le k_j \le n - 1$. Then (see [36, 22])

$$J_L(\mu_1, \dots, \mu_m; q) = J_{\mathcal{L}}(\rho + \lambda_1, \dots, \rho + \lambda_1; q), \qquad (3.4)$$

where \mathcal{L} is a linear combination of links with coefficients in $\mathbb{Z}(q^{1/2})$ obtained as follows. First we replace the *j*-th component of *L* by k_j its push-offs, then we cut these push-offs at one place and glue in the anti-symmetrizer element $g^{(k_j)}$. One gets a linear combinations of links. Modify the links in this linear combination by changing all the framings to 0. The result is \mathcal{L} .

Note that since L has 0 linking matrix, the right hand side of (3.4) is equal to $J_{\mathcal{L}}^{Hom}(q)$. Claim: $g^{(k)} = \sum_{z} f_{z}(q)z$,

where the sum is over a finite subset of the braid group B_k , and $f_z(q)$ is a rational function in $q^{(1-\operatorname{sn}(z))(n-1)/4}\mathbb{Z}(q)$. Here $\operatorname{sn}(z)$ is the sign of the permutation corresponding to the braid z. Note that $1 - \operatorname{sn}(z)$ is always even.

The claim can be proved easily by induction on k, using (3.3) and noting that $\operatorname{sn}(zz') = \operatorname{sn}(z) \operatorname{sn}(z')$.

For a braid $z \in B_k$ let c(z) be the number of cycle in the permutation corresponding to z. The closure of z is a link of c(z) components.

It is easy to see that, for $z \in B_k$,

$$q^{(1-\operatorname{sn}(z))(n-1)/4}\mathbb{Z}(q) = q^{(k-c(z))(n-1)/2}\mathbb{Z}(q)$$
(3.5)

Hence the right hand side of (3.4), by Lemma 3.5 and equality (3.5), must belong to $q^{(k_{j_1}+\dots+k_{j_m})(n-1)/2}\mathbb{Z}(q)$. On the other hand, $J_L(\mu_1,\dots,\mu_m;q)$ is a polynomial in $q^{1/2n}, q^{-1/2n}$. It follows that $J_L(\mu_1,\dots,\mu_m;q)$ is in $q^{(k_{j_1}+\dots+k_{j_m})(n-1)/2}\mathbb{Z}[q,q^{-1}]$.

3.4. The Kontsevich integral and weight systems. The Kontsevich integral is a very powerful invariant of links which was found by Kontsevich [10]. We will use here the version for *framed* oriented link introduced in [16, 17], (with exactly the same normalization). In [17] the invariant was denoted by $\hat{Z}_f(L)$, but here for simplicity we will use the notation Z(L). This invariant Z(L) takes values in the (completed) graded vector space $\mathcal{A}(\sqcup^m S^1)$ of Chinese character diagrams on m circles, i.e. the support of the Chinese character diagrams is m circles. Here m is the numbers of components of L. See, for example, [17], for the definition Chinese character diagrams. In brief, a *Chinese character diagram* consists of a 1-dimensional compact oriented manifold X and a graph every vertex of which is either univalent or trivalent. The components of X are supposed be ordered; and a cyclic order at every trivalent vertex of the graph is fixed. All the univalent vertex of the graph must be on the manifold X. Usually, the components of X is drawn using solid line, while the components of the graph – dashed lines. So we sometime refer to components of X as solid components, and components of the graph –

dashed. A univalent (resp. trivalent) vertex of the graph is also called an external (resp. external) vertex of the Chinese character diagram. The degree of a Chinese character diagram is half the number of (internal and external) vertices. The space of Chinese character diagrams on X is the vector space generated by Chinese character whose solid part is X, subject to the antisymmetry and Jacobi (also known as IXH) relations.

On the Lie algebra sl_n there is the standard invariant bilinear form defined by (y|z) = tr(yz), where tr is the trace in the fundamental representation.

Suppose D is a Chinese character diagram on m solid circles. Using the above defined bilinear form, we can define the weight $W_D(V_1, \ldots, V_m)$, which is a number, of the Chinese character diagram D when the m components of D are colored by sl_n -modules V_1, \ldots, V_m (see, for example, [10, 17]). In fact, W_D can be regarded as a multi-linear map :

 $W_D: \mathcal{R}^{\otimes m} \to \mathbb{Q}.$

For $\mu_1, \ldots, \mu_m \in \Lambda_{++}$, let $W_D(\mu_1, \ldots, \mu_m)$ stand for $W_D(V_{\mu_1}, \ldots, V_{\mu_m})$.

If Y is a linear combination of chord diagrams on m solid circles, then we can define W_Y using linearity. The relation between quantum invariants of framed oriented links and the Kontsevich integral is expressed in the following proposition.

Proposition 3.6. For $\mu_1, \ldots, \mu_m \in \Lambda_{++}$, one has that

$$J_L(\mu_1, \dots, \mu_m; q) \Big|_{q=e^h} = \sum_{\ell=0}^{\infty} W_{Z_\ell(L)}(\mu_1, \dots, \mu_m) h^{\ell}.$$
 (3.6)

Here Z_{ℓ} is the degree ℓ part of the Kontsevich integral Z.

This is a simple corollary of Drinfeld's theory of quasi-Hopf algebra and was observed by Le-Murakami [17] and Kassel [6]. This shows that all quantum invariants are special values of the Kontsevich integral.

The following is an easy exercise in the theory of the Kontsevich integral and we omit the proof (see similar results in [14], Lemma 5.6, and [15], Proposition 5.3).

Lemma 3.7. Suppose that L has 0 linking matrix. Then the degree ℓ part Z_{ℓ} of the Kontsevich integral can be represented in the form

$$Z_{\ell}(L) = \sum_{D} a_{D}D, \quad \text{with } a_{D} \in \mathbb{Q}$$

where the sum is over the set of Chinese character diagrams D which has at most $3\ell/2$ external vertices. Moreover, if s_j is the number of external vertices of D on the *j*-th component, then $\max_j s_j \leq \ell - \sum_j s_j$.

3.5. Dependence of W_D on μ_1, \ldots, μ_m . Suppose that D is a Chinese character diagram on m circles. At first we will fix μ_2, \ldots, μ_m and investigate the dependence of W_D on μ_1 , we will write $W_D(\mu_1)$ instead of $W_D(\mu_1, \ldots, \mu_m)$.

Break the first component of D at an arbitrary point which is not an external vertex. The result is a Chinese character diagram D' whose support (i.e. the solid part) is the union of an interval and m - 1 circle. It is known that D', modulo the anti-symmetry and Jacobi relations, depends only on D, but not on the point where we break the first component (see [17], section 1).

Assign $V_{\mu_2}, \ldots, V_{\mu_m}$ to the circle components of D', and take of the weight of D'. The result is not a number, but an element z(D') which lies in the center of the universal enveloping algebra $U(sl_n)$ of sl_n (see, for example, [6], Proposition XX.8.2). Since D' is determined by D, we will write z(D) for z(D'). By the definition of the weight, for every $\mu_1 \in \Lambda_{++}$, we have that

$$W_D(\mu_1) = tr_{\mu_1} z(D), \tag{3.7}$$

where tr_{μ_1} is the trace taken in the representation V_{μ_1} .

Since z(D) is a central element, and V_{μ_1} is an irreducible sl_n -module, z(D) acts on V_{μ_1} as a scalar times the identity. The scalar, denoted by $\xi_D(\mu_1)$, is a function on μ_1 , which is known as *character* in Lie theory. Harish-Chandra theory (see, for example, [2]) says that $\xi_D(\mu_1)$ is a polynomial function on μ_1 of degree not exceeding the degree of z(D)in $U(sl_n)$. Moreover the polynomial function $\xi_D(\mu_1)$ is invariant under the action of the Weyl group.

Formula (3.7) now shows that

$$W_D(\mu_1) = \dim(V_{\mu_1})\,\xi_D(\mu_1),\tag{3.8}$$

where the dimension of V_{μ_1} , by the Weyl formula, is equal to $\Delta(\mu_1)$ (see Lemma 2.3).

From the definition of the weight, it follows that if D has k external vertices on the first component, then z(D) has degree $\leq k$, and hence $\xi_D(\mu_1)$ is a polynomial function of degree at most k.

We we return to the general case, when the other variables μ_2, \ldots, μ_m are not fixed. Since all the variables μ_1, \ldots, μ_m are independent, we get the following

Proposition 3.8. For every $\mu_1, \ldots, \mu_m \in \Lambda_{++}$, one has that

$$W_D(\mu_1, \dots, \mu_m) = \Delta(\mu_1) \dots \Delta(\mu_m) p_D(\mu_1, \dots, \mu_m), \qquad (3.9)$$

where $p_D(\mu_1, \ldots, \mu_m)$ is a polynomial function invariant under the action of the Weyl group. Moreover, the total degree of p_D is less than or equal to the number of external vertices of D, and the degree of μ_j in p_D is less than or equal to the number of external vertices on the *j*-th component.

3.6. **Proof of Proposition 2.4.** From Proposition 3.8 and Lemma 3.7 we see that, for every $\mu_1, \ldots, \mu_m \in \Lambda_{++}$,

$$W_{Z_{\ell}(L)}(\mu_1,\ldots,\mu_m) = \Delta(\mu_1)\ldots\Delta(\mu_m)p_{\ell}(\mu_1,\ldots,\mu_m),$$

where $p_{\ell}(\mu_1, \ldots, \mu_m)$ is a polynomial function of degree $\leq 3\ell/2$, and if s_j is the degree of μ_j in p_{ℓ} , then $\max_j s_j \leq 2\ell - \sum_j s_j$.

Applying Proposition 3.6, we get

$$J_L(\mu_1, \dots, \mu_m; q) \Big|_{q=e^h} = \Delta(\mu_1) \dots \Delta(\mu_m) \sum_{\ell=0}^{\infty} p_\ell(\mu_1, \dots, \mu_m) h^{\ell}.$$
 (3.10)

This is true for $\mu_1, \ldots, \mu_m \in \Lambda_{++}$. We will show that (3.10) holds true for $\mu_1, \ldots, \mu_m \in \Lambda$.

Suppose one of the μ_j , say μ_1 , is on the boundary of the fundamental chamber C. Then the left hand side is 0 by definition, while $\Delta(\mu_1) = 0$ by Lemma 2.3. Hence (3.10) also holds true in this case.

If we replace μ_1 by $w(\mu_1)$, where w is an element of the Weyl group, then the left hand side of (3.10) must be multiplied by $\operatorname{sn}(w)$. On the right hand side, all the functions p_ℓ 's are invariant under the action of the Weyl group, while $\Delta(w(\mu_1)) = \operatorname{sn}(w)\Delta(\mu_1)$ (see Lemma 2.3). Hence if (3.10) is true for μ_1 , then it holds true for $w(\mu_1)$. Similarly for other μ_j . It follows that (3.10) holds true for every $\mu_1, \ldots, \mu_m \in \Lambda$.

Now substituting $h = \ln(1+x)$, we get

$$Q_L(\mu_1, \dots, \mu_m; q)\Big|_{q=x+1} = J_L(\mu_1, \dots, \mu_m; q)\Big|_{q=x+1} J_{U^{(m)}}(\mu_1, \dots, \mu_m; q)\Big|_{q=x+1}$$
$$= \Delta^2(\mu_1) \dots \Delta^2(\mu_m) \sum_{\ell=0}^{\infty} f'_{\ell}(\mu_1, \dots, \mu_m) x^{\ell},$$
(3.11)

for some polynomial functions $f'_{\ell}(\mu_1, \ldots, \mu_m)$.

Note that h, as a formal power series in x, has the first non-trivial term x. Hence each f'_{ℓ} is a polynomial function of total degree $\leq 3\ell/2$, and if s_j is the degree of μ_j in f'_{ℓ} , then $s_j \leq 2\ell - \sum_j s_j$.

That $f_{\ell}(\mu_1, \ldots, \mu_m) \in \mathbb{Z}$ when $\mu_1, \ldots, \mu_m \in \Lambda$ follows from the fact that $Q_L(\mu_1, \ldots, \mu_m)$ is a Laurent polynomial in q with *integer* coefficients. This completes the proof of Proposition 2.4.

4. Sums related to the root lattice

We will always assume that r is an odd prime, and $\zeta = e^{2\pi i/r}$. Recall that ρ is the half-sum of positive roots, and $|\rho|^2 = n(n^2 - 1)/12$ is always in $\frac{1}{2}\mathbb{Z}$. If β is in the root lattice, then $|\beta|^2$ is an even integer.

4.1. Gauss sum on the root lattice. For an integer b not divisible by r let

$$\mathcal{G}(b) = \sum_{\mu \in (\rho + \Lambda_r^{\text{root}})} \zeta^{b^{\frac{|\mu|^2 - |\rho|^2}{2}}}$$

Note that for every $\mu \in (\rho + \Lambda_r^{\text{root}})$, one has that $\frac{|\mu|^2 - |\rho|^2}{2} \in \mathbb{Z}$, hence $\mathcal{G}(b) \in \mathbb{Z}[\zeta]$. The fact that ζ is an *r*-th of unity has the following consequence.

The fact that ζ is an *i*-th of unity has the following consequence

Proposition 4.1. For every β in the root lattice Λ^{root} , one has that

 μ

$$\sum_{\substack{\in (\rho + \Lambda_r^{root})}} \zeta^{b\frac{|\mu + \beta|^2 - |\rho|^2}{2}} = \mathcal{G}(b).$$
(4.1)

Proof. For $i = 1, \ldots, n - 1$, we have

$$|\mu + r\alpha_i|^2 - |\rho|^2 = 2r(\mu|\alpha_i) + r^2|\alpha_i|^2.$$

Since $|\alpha_i|^2 = 2$ is an even number and $(\mu | \alpha_i) \in \mathbb{Z}$ for every $\mu \in \Lambda$, the right hand side is divisible by 2r. Hence

$$\zeta^{b\frac{|\mu+r\alpha_i|^2-|\rho|^2}{2}} = \zeta^{b\frac{|\mu|^2-|\rho|^2}{2}}.$$
(4.2)

Let us denote the left hand side of (4.1) by $\mathcal{G}(b;\beta)$. It's enough to show that $\mathcal{G}(b;\beta+\alpha_i) =$ $\mathcal{G}(b;\beta)$ for every $\beta \in \Lambda^{\text{root}}$ and $i = 1, \ldots, n-1$. We have

$$\sum_{\mu \in (\rho + \Lambda_r^{\text{root}})} \zeta^{b\frac{|\mu + \beta|^2 - |\rho|^2}{2}} = \sum_{\mu \in (\beta + \rho + \Lambda_r^{\text{root}})} \zeta^{b\frac{|\mu|^2 - |\rho|^2}{2}}$$

By definition, $\beta + \rho + \Lambda_r^{\text{root}}$ and $\beta + \alpha_i + \rho + \Lambda_r^{\text{root}}$ have the same number of elements, and μ belongs to the first set if and only if either μ or $\mu + r\alpha_i$ belongs to the second set (see the definition of Λ_r^{root} in §1.4) Hence, equality (4.2) shows that

$$\sum_{\mu \in (\beta + \rho + \Lambda_r^{\text{root}})} \zeta^{b\frac{|\mu|^2 - |\rho|^2}{2}} = \sum_{\mu \in (\beta + \alpha_i + \rho + \Lambda_r^{\text{root}})} \zeta^{b\frac{|\mu|^2 - |\rho|^2}{2}},$$

$$\beta) = \mathcal{G}(b; \beta + \alpha_i).$$

or $\mathcal{G}(b; \beta)$

The exact value of $\mathcal{G}(b)$ is given in the next proposition. Let us first explain a notation. For a rational number a/b, with $a, b \in \mathbb{Z}$ and b not divisible by r, let $(a/b)^{\vee}$ be the natural image of a/b in \mathbb{Z}/rZ .

Proposition 4.2. Suppose that b is not divisible by the prime r and n < r. Then $\mathcal{G}(b)$ has absolute value $r^{(n-1)/2}$, and its phase is given by

$$\frac{\mathcal{G}(b)}{r^{(n-1)/2}} = \left(\frac{n}{r}\right) \left[\left(\frac{b}{r}\right) e^{\pi i (1-r)/4}\right]^{n-1} \zeta^{\left(-\frac{b|\rho|^2}{2}\right)^{\vee}}.$$
(4.3)

Here $\left(\frac{n}{r}\right)$ and $\left(\frac{b}{r}\right)$ are the Legendre symbols. In particular,

$$\frac{\mathcal{G}(b)}{\mathcal{G}(\operatorname{sn}(b))} = \left(\frac{|b|}{r}\right)^{n-1} \zeta^{(\frac{\operatorname{sn}(b)-b}{2}|\rho|^2)^{\vee}}.$$
(4.4)

Here sn(b) is the sign of b.

Proof. Recall that $|\rho|^2 = n(n^2 - 1)/12$. Note that r + 1 is an even number, and hence $|\rho|^2(r+1)^2/2$ is an integer. Then we have

$$(-\frac{b|\rho|^2}{2})^{\vee} \equiv -b|\rho|^2(r+1)^2/2 \pmod{r}.$$
(4.5)

Direct calculation shows that

$$\frac{|\mu+\rho|^2 - |\rho|^2}{2} - \frac{|\mu+(r+1)\rho|^2 - (r+1)^2|\rho|^2}{2} = -r(\mu|\rho)$$

For $\mu \in \Lambda^{\text{root}}$, $(\mu|\rho)$ is in \mathbb{Z} , and hence the right hand side is divisible by r. It follows that

$$\zeta^{b\frac{|\mu+\rho|^2 - |\rho|^2}{2}} = \zeta^{b\frac{|\mu+(r+1)\rho|^2 - (r+1)^2|\rho|^2}{2}}$$
$$= \zeta^{b\frac{|\mu+(r+1)\rho|^2}{2}} \zeta^{\left(-\frac{b|\rho|^2}{2}\right)^{\vee}}, \qquad (4.6)$$

where the second identity follows from (4.5). Now we have

$$\begin{aligned} \mathcal{G}(b) &= \sum_{\mu \in (\rho + \Lambda_r^{\text{root}})} \zeta^{b^{\frac{|\mu|^2 - |\rho|^2}{2}}} \\ &= \sum_{\mu \in \Lambda^{\text{root}}} \zeta^{b^{\frac{|\mu + \rho|^2 - |\rho|^2}{2}}} \\ &= \zeta^{(-\frac{b|\rho|^2}{2})^{\vee}} \sum_{\mu \in \Lambda^{\text{root}}} \zeta^{b^{\frac{|\mu + (r+1)\rho|^2}{2}}} \\ &= \zeta^{(-\frac{b|\rho|^2}{2})^{\vee}} \sum_{\mu \in \Lambda^{\text{root}}} \zeta^{b^{\frac{|\mu|^2}{2}}} \end{aligned}$$
(4.7)

where the third identity follows from (4.6) and the fourth from Proposition 4.1, noting that $(r+1)\rho$ is in the root lattice (since r+1 is even).

Recall that $(\alpha_i | \alpha_j) = A_{ij}$, where A is the Cartan matrix. If $\mu = k_1 \alpha_1 + \cdots + k_{n-1} \alpha_{n-1}$, then $|\mu|^2 = \mathbf{k}^t A \mathbf{k}$, where \mathbf{k} is the column vector transpose to $\mathbf{k}^t = (k_1, \ldots, k_{n-1})$. Hence

$$\sum_{u \in \Lambda_r^{\text{root}}} \zeta^{b\frac{|\mu|^2}{2}} = \sum_{\mathbf{k} \in (\mathbb{Z}/r\mathbb{Z})^{n-1}} \zeta^{b\mathbf{k}^t A \mathbf{k}/2}$$

The right hand side is a multi-variable Gauss sum, and its value is:

$$\sum_{\mathbf{k}\in(\mathbb{Z}/r\mathbb{Z})^{n-1}}\zeta^{b\,\mathbf{k}^t A\mathbf{k}/2} = \left(\frac{n}{r}\right)\,\left[\left(\frac{b}{r}\right)\,e^{\pi i(1-r)/4}\,\sqrt{r}\right]^{n-1}.\tag{4.8}$$

(A proof of this formula is given in Appendix). This formula, together with (4.7), proves the proposition. $\hfill \Box$

Corollary 4.3. The number $\mathcal{G}(b)$ is proportional to $\zeta^{(n-1)(r-1)/2}$ by a unit in $\mathbb{Z}[\zeta]$.

Proof. Note that ζ is invertible in $\mathbb{Z}[\zeta]$, since $\zeta^{-1} = \zeta^{r-1}$. It is well-known that $e^{(\pi i(1-r)/4)}\sqrt{r}$ is proportional to $\zeta^{(r-1)/2}$ by a unit in $\mathbb{Z}[\zeta]$ (see, for example, [3, 23]). The corollary now follows immediately from formula (4.3).

4.2. Completing the square.

Proposition 4.4. For every β in the root lattice Λ^{root} , we have

$$\sum_{\mu \in (\rho + \Lambda_r^{root})} \zeta^{b\frac{|\mu|^2 - |\rho|^2}{2}} \zeta^{(\mu|\beta)} = \zeta^{(-|\beta|^2/2b)^{\vee}} \mathcal{G}(b).$$
(4.9)

Proof. The proof uses the trick of completing the square. Suppose b^* is a number such that $bb^* \equiv 1 \pmod{r}$. Noting that $(\mu|\beta) \equiv b(\mu|b^*\beta) \pmod{r}$, we have

$$\frac{b}{2}(|\mu|^2 - |\rho|^2) + (\mu|\beta) \equiv \frac{b}{2}(|\mu + b^*\beta|^2 - |\rho|^2) - (\frac{|\beta|^2}{2b})^{\vee} \pmod{r}.$$
(4.10)

It follows that the left hand side of (4.9) is equal to

$$\zeta^{(-|\beta|^2/2b)^{\vee}} \sum_{\mu \in (\rho + \Lambda_r^{\text{root}})} \zeta^{b\frac{|\mu + b^*\beta|^2 - |\rho|^2}{2}}$$

which is equal to the right hand side of (4.9) by Proposition 4.1.

4.3. The value of F_{U_b} . Recall that U_b is the unknot with framing b. We have

$$J_{U_b}(\mu;q) = q^{b\frac{|\mu|^2 - |\rho|^2}{2}} J_U(\mu;q).$$

Using the value of the unknot (3.1) and the definition of Q_L we get

$$Q_{U_b}(\mu;q) = q^{b\frac{|\mu|^2 - |\rho|^2}{2}} q^{2(\mu - \rho|\rho)} \frac{\psi(\mu;q)^2}{\psi(\rho;q)^2}.$$

Hence

$$F_{U_b}(r) = \sum_{\mu \in (\rho + \Lambda_r^{\text{root}})} Q_L(\mu; \zeta)$$

= $\frac{\zeta^{-2|\rho|^2}}{\psi(\rho; \zeta)^2} \sum_{\mu \in (\rho + \Lambda_r^{\text{root}})} \zeta^{b\frac{|\mu|^2 - |\rho|^2}{2}} [\zeta^{(\mu|\rho)}\psi(\mu; \zeta)]^2.$ (4.11)

Proposition 4.5. The value of $F_{U_b}(r)$ is given by

$$F_{U_b}(r) = n! \mathcal{G}(b) \frac{\psi(b^*\rho;\zeta)}{\psi(\rho;\zeta)\psi(-\rho;\zeta)}.$$
(4.12)

In particular, when $b = \pm 1$, we have

$$F_{U_{\pm}} = \frac{n! \mathcal{G}(\pm 1)}{\psi(\mp \rho; \zeta)} \tag{4.13}$$

Proof. The Weyl denominator formula (see Lemma 3.1) shows that

$$\zeta^{(\mu|\rho)}\psi(\mu;\zeta) = \sum_{w\in\mathcal{W}}\operatorname{sn}(w)\zeta^{(\mu|w(\rho))}.$$
(4.14)

Hence

$$[\zeta^{(\mu|\rho)}\psi(\mu;\zeta)]^2 = \sum_{w,w'\in\mathcal{W}} \operatorname{sn}(ww')\zeta^{(\mu|w(\rho)+w'(\rho))}.$$
(4.15)

Although ρ may not be in the root lattice, the sum $w(\rho) + w'(\rho)$ is always in Λ^{root} , since each of $w(\rho), w'(\rho)$ is in $\rho + \Lambda^{\text{root}}$ (see Lemma 1.2) and $2\rho \in \Lambda^{\text{root}}$.

Using (4.15) in (4.11) and applying Proposition 4.4, we get

$$F_{U_b}(r) = \frac{\zeta^{-2|\rho|^2} \mathcal{G}(b)}{\psi(\rho;\zeta)^2} \sum_{w,w' \in \mathcal{W}} \operatorname{sn}(ww') \zeta^{(-|(w(\rho)+w'(\rho)|^2/2b)^{\vee}}.$$
(4.16)

Note that

$$|w(\rho) + w'(\rho)|^{2} = |w(\rho)|^{2} + |w'(\rho)|^{2} + 2(w(\rho)|w'(\rho))$$
$$= 2|\rho|^{2} + 2(\rho|w^{-1}w'(\rho)).$$

26

It's easy to check that, modulo r,

$$\left(\frac{2|\rho|^2 + 2(\rho|w^{-1}w'(\rho))}{2b}\right)^{\vee} \equiv 2b^*|\rho|^2 + (b^*\rho|w^{-1}w'(\rho) - \rho),$$

where b^* is any integer such that $bb^* \equiv 1 \pmod{r}$. Using this in the right hand side of (4.16), we get

$$F_{U_b}(r) = \frac{\zeta^{(-2b^*|\rho|^2 - 2|\rho|^2)}\mathcal{G}(b)}{\psi(\rho;\zeta)^2} \sum_{w,w' \in \mathcal{W}} \operatorname{sn}(ww')\zeta^{(-b^*\rho|w^{-1}w'(\rho))}.$$

Note that $\operatorname{sn}(ww') = \operatorname{sn}(w^{-1}w')$. Hence

$$\sum_{w,w'\in\mathcal{W}}\operatorname{sn}(ww')\zeta^{(-b^*\rho|w^{-1}w'(\rho))} = |\mathcal{W}|\sum_{w\in\mathcal{W}}\zeta^{(-b^*\rho|w(\rho))}.$$

The right hand side, again by Lemma 3.1 and $|\mathcal{W}| = n!$, is equal to $n! \psi(-b^*\rho; \zeta)$. This shows that

$$F_{U_b}(r) = n! \mathcal{G}(b) \,\zeta^{(-2|\rho|^2 - 2b^*|\rho|^2)} \,\frac{\psi(-b^*\rho;\zeta)}{\psi(\rho;\zeta)^2}.$$
(4.17)

For every number k,

$$\psi(k\rho;\zeta) = \prod_{\alpha\in\Phi_+} (1-\zeta^{-(k\rho|\alpha)})$$
$$= \prod_{\alpha\in\Phi_+} \zeta^{-(k\rho|\alpha)} \prod_{\alpha\in\Phi_+} (\zeta^{(k\rho|\alpha)}-1)$$
$$= \zeta^{-2k|\rho|^2} (-1)^{n(n-1)/2} \psi(-k\rho;\zeta).$$

Using this formula, with $k = b^*$ and k = -1, in formula (4.17), we get (4.12).

4.4. **Proof of Proposition 2.13.** If k is an integer between 1 and r-1, then $\zeta^k - 1$ is proportional to $\zeta - 1$ by a unit in $\mathbb{Z}[\zeta]$. In fact, both

$$\frac{\zeta^k - 1}{\zeta - 1} \quad \text{and} \quad \frac{\zeta - 1}{\zeta^k - 1} = \frac{\zeta^{kk^*} - 1}{\zeta^k - 1}$$

are in $\mathbb{Z}[\zeta]$.

For every positive root α , the number $(\alpha | \rho)$ is an integer between 1 and n - 1. This fact follows from the explicit formulas of α and ρ (see [2]). There are n(n-1)/2 positive roots, and hence if r > n, the number

$$\psi(\mp\rho;\zeta) = \prod_{\alpha\in\Phi_+} (1-\zeta^{\pm(\rho|\alpha)})$$

is proportional to $(\zeta - 1)^{n(n-1)/2}$ in $\mathbb{Z}[\zeta]$.

Formula (4.13) and Corollary 4.3 shows that $F_{U_{\pm}}(r)$ is proportional to

$$\zeta^{(r-1)(n-1)/2 - n(n-1)/2}$$

by a proportional factor which is a unit in $\mathbb{Z}[\frac{1}{n!}][\zeta]$. This proves Proposition 2.13.

4.5. Simple lens spaces and the constant term. With the knowledge of $F_{U_b}(r)$, we can calculate the value $\tau_r^{PSU(n)}$ and $\tau^{PSU(n)}$ of lens spaces M(b).

Proposition 4.6. Let M(b) be the lens space obtained by surgery on the unknot with framing b, where b is a non-zero integer with absolute value less than the prime r. Then

$$\tau_r^{PSU(n)}(M(b)) = \left(\frac{|b|}{r}\right)^{n-1} \zeta^{(\frac{\operatorname{sn}(b)-b}{2}|\rho|^2)\vee} \frac{\psi(b^*\rho;\zeta)}{\psi(\operatorname{sn}(b)\rho;\zeta)}.$$
$$\tau^{PSU(n)}(M(b)) = \left(\frac{|b|}{r}\right)^{n-1} (1+x)^{(\frac{\operatorname{sn}(b)-b}{2}|\rho|^2)} \frac{\psi(\rho/b;1+x)}{\psi(\operatorname{sn}(b)\rho;1+x)}.$$
(4.18)

Proof. The first identity follows from $\tau_r^{PSU(n)}(M(b)) = F_{U_b}(r)/F_{U_{sn(b)}}(r)$, and the explicit formula (4.12) of F_{U_b} . Actually the value of $\tau_r^{PSU(n)}(M(b))$ had been calculated by Takata, see [31] for the values of $\tau_r^{PSU(n)}$ of Seifert fibered space.

The value of $\tau^{PSU(n)}(\dot{M}(b))$ can also be calculated directly from the explicit formula of $Q_{U_b}(\mu;q)$. However, it's simpler to notice that the right hand side of (4.18), which we denote by g for a moment, satisfies

$$\mathfrak{p}_r(g) = \mathfrak{p}_r\left(\tau_r^{PSU(n)}(M(b))\right).$$

Hence by the uniqueness (see Lemma 1.6), we can conclude that $g = \tau^{PSU(n)}(M(b))$.

From formula (4.18) one can calculate the constant term of the power series $\tau(U_b)$; the result is $\left(\frac{|b|}{r}\right)^{n-1} |b|^{-n(n-1)/2}$. Suppose L' is a link with diagonal linking matrix, with b_1, \ldots, b_m on the diagonal. From the definition it follows that the constant term of the power series $\tau(L')$ is the same as that of $\tau(U_{b_1}) \times \cdots \times \tau(U_{b_m})$. Hence the constant term of $\tau(L')$ is is $\left(\frac{|b_1\dots b_m|}{r}\right)^{n-1} |b_1\dots b_m|^{-n(n-1)/2}$. It follows that for every rational homology 3-sphere M, the constant term of the power series $\tau^{PSU(n)}(M)$ is $\left(\frac{|H_1(M,\mathbb{Z})|}{r}\right)^{n-1} |H_1(M,\mathbb{Z})|^{-n(n-1)/2}$.

4.6. **Proofs of Proposition 2.6 and 2.15.** The following follows directly from the definitions of \mathfrak{p}_r .

Lemma 4.7. For every $a, b \in \mathbb{Z}$ and every prime r not dividing b, one has that

$$\mathfrak{p}_r\left((1+x)^{a/b}\right) = \mathfrak{p}_r\left((1+x)^{(a/b)^\vee}\right) = \mathfrak{p}_r\left(\zeta^{(a/b)^\vee}\right)$$

Let $\beta \in \Lambda^{\text{root}}$ be an element of the root lattice. Then Proposition 4.4 and formula (4.13) show that

$$\frac{\sum_{\substack{\mu \in (\rho + \Lambda_r^{\text{root}})}} \zeta^{b\frac{|\mu|^2 - |\rho|^2}{2}} \zeta^{(\mu|\beta)}}{F_{U_{\text{sn}(b)}}} = \frac{1}{n!} \frac{\mathcal{G}(b)}{\mathcal{G}(\text{sn}(b))} \psi \left(-\operatorname{sn}(b)\rho;\zeta\right) \, \zeta^{\left(-\frac{|\beta|^2}{2b}\right)^{\vee}} \\ = \left(\frac{|b|}{r}\right)^{n-1} \frac{\zeta^{\left(\frac{\text{sn}(b) - b}{2}|\rho|^2\right)^{\vee}}}{n!} \, \psi \left(-\operatorname{sn}(b)\rho;\zeta\right) \, \zeta^{\left(-\frac{|\beta|^2}{2b}\right)^{\vee}} \tag{4.19}$$

The second equality follows from (4.4).

The right hand side, if the checks are droped and ζ replaced by x + 1, is, by definition, the same as $\Gamma_b(q^\beta)$. Hence Lemma 4.7 and (4.19) show that:

$$\mathfrak{p}_r\left(\frac{\sum\limits_{\mu\in(\rho+\Lambda_r^{\mathrm{root}})}\zeta^{b\frac{|\mu|^2-|\rho|^2}{2}}\zeta^{(\mu|\beta)}}{F_{U_{\mathrm{sn}(b)}}}\right) = \mathfrak{p}_r(\Gamma_b(q^\beta)). \tag{4.20}$$

This proves Proposition 2.15, since the set of all q^{β} spans \mathcal{S} .

Now we prove Proposition 2.6. Recall that

$$\Gamma_b(q^\beta) = q^{\frac{-|\beta|^2}{2b}} y_b$$

where

$$y_b = \left(\frac{|b|}{r}\right)^{n-1} \frac{(1+x)^{(\operatorname{sn}(b)-b)|\rho|^2/2}}{n!} \psi(-\operatorname{sn}(b)\rho; 1+x)$$

does not depend on β . Since $1 - (1 + x)^{\operatorname{sn}(b)(\alpha|\rho)}$ is divisible by x, and since there are n(n-1)/2 positive roots, $\psi(-\operatorname{sn}(b)\rho; 1+x)$ is divisible by $x^{n(n-1)/2}$ in $\mathbb{Z}[\frac{1}{n!b}][x]$. Since y_b is divisible by $\psi(-\operatorname{sn}(b)\rho; 1+x)$, it is divisible by $x^{n(n-1)/2}$ in $\mathbb{Z}[\frac{1}{n!b}][x]$.

Let $\mathbf{a} = (a_1, ..., a_{n-1})$. Then

$$\boldsymbol{\eta}^{\mathbf{a}} = \prod_{i=1}^{n-1} (1 - q^{-\alpha_i})^{a_i}$$
$$= \prod_{i=1}^{n-1} \sum_{s_i=0}^{a_i} {a_i \choose s_i} (-1)^{a_i - s_i} q^{(s_i - a_i)\alpha_i}$$
$$= \sum_{0 \le \mathbf{s} \le \mathbf{a}} (-1)^{|\mathbf{a} - \mathbf{s}|} {\mathbf{a} \choose \mathbf{s}} q^{\sum_i (s_i - a_i)\alpha_i}.$$

Here $\mathbf{s} = (s_1, \ldots, s_{n-1}), \mathbf{s} \leq \mathbf{a}$ means $s_i \leq a_i$ for every i, and $\binom{\mathbf{a}}{\mathbf{s}}$ means $\prod_i \binom{a_i}{s_i}$. It follows that

$$\frac{\Gamma_b(\boldsymbol{\eta}^{\mathbf{a}})}{y_b} = (-1)^{|\mathbf{a}|} \sum_{0 \le \mathbf{s} \le \mathbf{a}} (-1)^{|\mathbf{s}|} \binom{\mathbf{a}}{\mathbf{s}} (1+x)^{-|\sum (s_i - a_i)\alpha_i|^2/2b}.$$
 (4.21)

So we need only to show that the right hand side of (4.21) is divisible by $x^{\lfloor (|\mathbf{a}|+1)/2 \rfloor}$, or that the coefficient of x^d in the right hand side of (4.21) is equal to 0 whenever $2d < |\mathbf{a}|$. The coefficient of x^d is

$$v_d = (-1)^{|\mathbf{a}|} \sum_{0 \le \mathbf{s} \le \mathbf{a}} (-1)^{|\mathbf{s}|} \binom{\mathbf{a}}{\mathbf{s}} \binom{-|\sum(s_i - a_i)\alpha_i|^2/2b}{d}.$$

The expression

$$\binom{-|\sum(s_i - a_i)\alpha_i|^2/2b}{d}$$

is a polynomial in s_1, \ldots, s_{n-1} of total degree 2d. Hence when $2d < |\mathbf{a}|, v_d$ is equal to 0 by Lemma 4.9 below. This proves Proposition 2.6.

4.7. A technical result. The following result is well-known, see [35], Lemma 5.19.

Lemma 4.8. Suppose that a is a positive integer. Then

$$\sum_{s=0}^{a} (-1)^s \binom{a}{s} s^d = 0$$

for $d = 1, 2, \ldots, a - 1$.

From this lemma one can prove the following.

Lemma 4.9. Suppose $p(s_1, \ldots, s_{n-1})$ is a polynomial of total degree less than $|\mathbf{a}|$, where $\mathbf{a} \in (\mathbb{Z}_+)^{n-1}$. Then

$$\sum_{0 \le \mathbf{s} \le \mathbf{a}} (-1)^{|\mathbf{s}|} \begin{pmatrix} \mathbf{a} \\ \mathbf{s} \end{pmatrix} p(s_1, \dots, s_{n-1}) = 0.$$

Proof. It suffices to consider the case when $p(s_1, \ldots, s_{n-1}) = s_1^{d_1} \ldots s_{n-1}^{d_{n-1}}$. We have

$$\sum_{0 \le \mathbf{s} \le \mathbf{a}} (-1)^{|\mathbf{s}|} \binom{\mathbf{a}}{\mathbf{s}} p(s_1, \dots, s_{n-1}) = \prod_{i=1}^{n-1} \sum_{s_i=0}^{a_i} (-1)^{s_i} \binom{a_i}{s_i} s_i^{d_i}$$

Since

$$d_1 + \dots + d_{n-1} < a_1 + \dots + a_{n-1},$$

there must be an index *i* for which $d_i < a_i$. Apply the previous lemma we get the result.

4.8. **Proof of Proposition 2.14.** Proposition 2.14, for the case n = 2, is similar to Lemma 2.2 of [29]. We begin with the following lemma.

Lemma 4.10. Suppose r is an odd prime. For any $d \in \mathbb{Z}_+$, the sum $\sum_{k=0}^{r-1} \binom{k}{d}$ is divisible by $(\zeta - 1)^{\frac{r-1}{2} - \lfloor \frac{d}{2} \rfloor}$ in $\mathbb{Z}_{(r)}[\zeta]$, i.e the quotient $\frac{\sum_{k=0}^{r-1} \binom{k}{d}}{(\zeta - 1)^{(r-1)/2 - \lfloor \frac{d}{2} \rfloor}}$ is in $\mathbb{Z}_{(r)}[\zeta]$.

Proof. If $d \ge r - 1$, then $(r - 1)/2 - \lfloor d/2 \rfloor \le 0$. The above quotient is in $\mathbb{Z}[\zeta]$, and we are done.

Suppose that $d \leq r-2$. Note that $\binom{k}{d}$ is a polynomial in k of degree d. It is well-known that

$$\sum_{k=0}^{r-1} k^d = \frac{B_{d+1}(r) - B_{d+1}(0)}{d+1},$$
(4.22)

where $B_{d+1}(z)$ is the Bernoulli polynomial (see, for example [3]). It is also known that for $d \leq r-2$, the polynomial B_{d+1} has coefficients in $\mathbb{Z}_{(r)}$. Hence the right hand side of (4.22) is divisible by r, which is divisible by $(\zeta - 1)^{(r-1)}$, and hence by $(\zeta - 1)^{(r-1)/2}$, in $\mathbb{Z}_{(r)}[\zeta]$. This proves the lemma. **Corollary 4.11.** Suppose $p(k_1, \ldots, k_{n-1})$ is a polynomial of degree d which takes values in $\mathbb{Z}_{(r)}$ whenever $k_i \in \mathbb{Z}$. Then

$$\sum_{i=1}^{n-1} \sum_{k_i=0}^{r-1} p(k_1, \dots, k_{n-1})$$

is divisible by $(\zeta - 1)^{(n-1)\frac{r-1}{2} - \lfloor d/2 \rfloor}$ in $\mathbb{Z}_{(r)}[\zeta]$.

Proof. The polynomial $p(k_1, \ldots, k_{n-1})$ is a $\mathbb{Z}_{(r)}$ -linear combination of terms of the form

$$\binom{k_1}{d_1} \cdots \binom{k_{n-1}}{d_{n-1}}$$

Applying Lemma 4.10, we get the result.

[Proof of Proposition 2.14]. We have to show that the number

$$u = \sum_{\mu \in (\rho + \Lambda_r^{\text{root}})} \zeta^{b \frac{|\mu|^2 - |\rho|^2}{2}} \begin{pmatrix} \boldsymbol{\alpha} \\ \mathbf{a} \end{pmatrix} (\mu)$$

is divisible by

$$(\zeta - 1)^{(n-1)\frac{r-1}{2} - \lfloor \frac{|\mathbf{a}|}{2} \rfloor}$$

in $\mathbb{Z}_{(r)}[\zeta]$. Note that

$$u = \sum_{\mu \in \Lambda_r^{\text{root}}} \zeta^{b^{\frac{|\mu+\rho|^2 - |\rho|^2}{2}}} \begin{pmatrix} \boldsymbol{\alpha} \\ \mathbf{a} \end{pmatrix} (\mu+\rho).$$

Writing $\zeta = 1 + (\zeta - 1)$ and using the expansion $\zeta^c = \sum_l {c \choose l} (\zeta - 1)^l$ in the above formula, we can express u as a polynomial in $\zeta - 1$:

$$u = \sum_{\ell} u_{\ell} (\zeta - 1)^{\ell}.$$

The coefficient of $(\zeta - 1)^{\ell}$ is

$$u_{\ell} = \sum_{\mu \in \Lambda_r^{\text{root}}} \binom{b(|\mu + \rho|^2 - |\rho|^2)/2}{\ell} \binom{\boldsymbol{\alpha}}{\mathbf{a}} (\mu + \rho).$$

Suppose $\mu = k_1 \alpha_1 + \dots + k_{n-1} \alpha_{n-1}$, then $\sum_{\mu \in \Lambda_r^{\text{root}}}$ becomes $\sum_{i=1}^{n-1} \sum_{k_i=0}^{r-1}$, and u_ℓ can be written as

$$u_{\ell} = \sum_{i=1}^{n-1} \sum_{k_i=0}^{r-1} p(k_1, \dots, k_{n-1}),$$

where $p(k_1, \ldots, k_{n-1})$ is a polynomial taking values in $\mathbb{Z}_{(r)}$ whenever $k_i \in \mathbb{Z}$. The degree of p is $2\ell + |\mathbf{a}|$. Hence by Corollary 4.11, u_ℓ is divisible by $(\zeta - 1)^{(n-1)\frac{r-1}{2} - \lfloor \frac{|\mathbf{a}|}{2} \rfloor - \ell}$ in $\mathbb{Z}_{(r)}[\zeta]$.

Now remember that u_{ℓ} is the coefficient of $(\zeta - 1)^{\ell}$ in u. Hence taking $(\zeta - 1)^{\ell}$ into account, we see that each term $u_{\ell}(\zeta - 1)^{\ell}$, and hence u, is divisible by $(\zeta - 1)^{(n-1)\frac{r-1}{2} - \lfloor \frac{|\mathbf{a}|}{2} \rfloor}$ in $\mathbb{Z}_{(r)}[\zeta]$. This completes the proof of Proposition 2.14.

31

APPENDIX A.

A.1. On the definition of PSU(n)-quantum invariant. Here we prove that $\tau_r^{PSU(n)}(M)$ is coincident with the one introduced by Kohno and Takata [8] (see also [36]). We first recall Kohno and Takata's definition of quantum PSU(n) invariant, which we will denote by $\tilde{\tau}_r^{PSU(n)}(M)$.

Recall that Λ'_r is the intersection of the simplex C_r and the weight lattice Λ . Let us consider the subset Λ''_r of Λ'_r which consists of all weight $\mu = \rho + k_1\lambda_1 + \cdots + k_{n-1}\lambda_{n-1}$ such that

$$k_1 + 2k_2 + \dots + (n-1)k_{n-1} \quad \text{is divisible by } n \tag{(*)}$$

Now we define $F''_L(r)$ by the same formula of $F'_L(r)$, only replacing Λ'_r by Λ''_r :

$$F_L''(r) = \sum_{j=1}^m \sum_{\mu_j \in \Lambda_r''} Q_L(\mu_1, \dots, \mu_m; \zeta).$$

Then, by definition (see [8])

$$\tilde{\tau}_r^{PSU(n)}(M) = \frac{F_L''(r)}{F_{U_+}''(r)^{\sigma_+} F_{U_-}''(r)^{\sigma_-}}.$$
(A.1)

We have the following simple observation.

Lemma A.1. The tuple (k_1, \ldots, k_{n-1}) satisfies (*) if and only if $k_1\lambda_1 + \cdots + k_{n-1}\lambda_{n-1}$ is in the root lattice. In other words,

$$\Lambda_r'' = C_r \cap (\rho + \Lambda^{root})$$

Proof. If $\beta = l_1\alpha_1 + \ldots + l_{n-1}\alpha_{n-1}$, then $\beta = k_1\lambda_1 + \cdots + k_{n-1}\lambda_{n-1}$, where $\mathbf{k} = A\mathbf{l}$. Here $\mathbf{k} = (k_1, \ldots, k_{n-1})$ and $\mathbf{l} = (l_1, \ldots, l_{n-1})$, and A is the Cartan matrix. The inverse A^{-1} of A is the symmetric matrix whose entries are given by $(A^{-1})_{ij} = (n-i)j/n$ for $n-1 \ge i \ge j \ge 1$.

Now it is easy to show that **k** satisfies (*) if and only if $\mathbf{l} = A^{-1}\mathbf{k}$ has *integer* entries.

The translation group $r\Lambda^{\text{root}}$ acts on $\Lambda \otimes \mathbb{R}$; and a fundamental domain is the parallelepiped

$$P_r = \{ t_1 \alpha_1 + \dots + t_{n-1} \alpha_{n-1} \mid 1 \le t_i < r+1, t_i \in \mathbb{R} \}.$$

The affine group $\mathcal{W}_{(r)}$ is the semi-direct product of the Weyl group \mathcal{W} and the translation group $r\Lambda^{\text{root}}$. Hence $r\Lambda^{\text{root}}$ has index n! in $\mathcal{W}_{(r)}$. The group $\mathcal{W}_{(r)}$ acts on $\Lambda \otimes \mathbb{R}$, and a fundamental domain in C_r .

In a sense, P_r is n! times C_r . More precisely, each point in the *interior* of C_r has exactly n! points in P_r in its $\mathcal{W}_{(r)}$ orbit (see [5], Lemma 6.6). For a point on the boundary of C_r , the cardinality of its $\mathcal{W}_{(r)}$ orbits in P_r may be different from n!. However, if one of the μ_j is on the boundary of C_r , then $Q_L(\mu_1, \ldots, \mu_m; \zeta) = 0$, by Proposition 1.3. By that same proposition, $Q_L(\mu_1, \ldots, \mu_m, \zeta)$ is invariant under the action of $\mathcal{W}_{(r)}$.

Notice that $\Lambda_r'' = C_r \cap (\rho + \Lambda^{\text{root}})$, while $\rho + \Lambda_r^{\text{root}} = P_r \cap (\rho + \Lambda^{\text{root}})$. Hence $\sum_{\mu_i \in (\rho + \Lambda^{\text{root}})} Q_L(\mu_1, \dots, \mu_m; \zeta) = (n!)^m \sum_{\mu_j \in \Lambda''_r} Q_L(\mu_1, \dots, \mu_m; \zeta).$

 $F_L(r) = (n!)^m F_L''(r)$. When the linking matrix of L has non-zero determinant, one Or has that $\sigma_+ + \sigma_- = m$. Hence

$$\frac{F_L(r)}{F_{U_+}(r)^{\sigma_+} F_{U_+}(r)^{\sigma_+}} = \frac{F_L''(r)}{F_{U_+}''(r)^{\sigma_+} F_{U_+}''(r)^{\sigma_+}}$$

This shows that our definition agrees with that of Kohno and Takata.

A.2. The value of Gauss sums. For the Legendre symbols we have $\left(\frac{b}{r}\right)\left(\frac{b'}{r}\right) = \left(\frac{bb'}{r}\right)$. Hence we always have $\left(\frac{b}{r}\right) = \left(\frac{b^*}{r}\right)$, where b^* is any number such that $bb^* \equiv 1 \pmod{r}$. Let us consider the quadratic Gauss sum

$$\gamma = \sum_{k \in \mathbb{Z}/r\mathbb{Z}} \zeta^{k^2},$$

where $\zeta = e^{2\pi i/r}$. The value of γ is well-known (see, for example, [3]):

$$\gamma = \begin{cases} \sqrt{r} & \text{if } r \equiv 1 \pmod{4}, \\ i\sqrt{r} & \text{if } r \equiv 3 \pmod{4}. \end{cases}$$

Noting that $\left(\frac{2}{r}\right) = (-1)^{(n^2-1)/2}$ (see [3]) we can reformulate the value of γ as follows.

Lemma A.2. For an odd prime r, the value of the quadratic Gauss sum is given by

$$\gamma = \left(\frac{2}{r}\right) e^{\pi i (1-r)/4} \sqrt{r}.$$

The following is well-known (see [3], Proposition 6.3.1).

Lemma A.3. Suppose that b is an integer not divisible by r, then

$$\sum_{k\in\mathbb{Z}/r\mathbb{Z}}\zeta^{bk^2} = \left(\frac{b}{r}\right)\gamma.$$

Now let us consider the following multi-variable Gauss sum

$$\gamma_{A,b} = \sum_{\mathbf{k} \in (\mathbb{Z}/r\mathbb{Z})^{n-1}} \zeta^{b\mathbf{k}^t A \mathbf{k}/2}$$

where A is the Cartan matrix of sl_n . Recall that $A_{ii} = 2$, $A_{i,i+1} = A_{i+1,i} = -1$, and other entries are 0. Since the entries on the diagonal of A are even, $\mathbf{k}^t A \mathbf{k}/2$ is always an integer. Hence $\gamma_{A,b} \in \mathbb{Z}[\zeta]$.

Let D be the diagonal $(n-1) \times (n-1)$ -matrix with entries $D_{jj} = \frac{j+1}{2j}$. Let P be the upper-triangle $(n-1) \times (n-1)$ -matrix with entries $P_{ii} = 1$ for $i = 1, \ldots, n-1$, $P_{i,i+1} = -\frac{i}{i+1}$ for i = 1, ..., n-2, and other entries equal to 0. Then it easy to check that

$$\frac{1}{2}A = P^t D P \tag{A.2}$$

i.e., the bilinear form corresponding to A can be diagonalized (over \mathbb{Q}) using P.

Recall that "check" is the natural reduction modulo r map from $\mathbb{Z}_{(r)}$ to $Z/r\mathbb{Z}$. Since all the entries of P, D, P^{-1} are in $\mathbb{Z}_{(r)}$ (recall that r > n), there are defined D^{\vee} and $P^{\vee}, (P^{-1})^{\vee}$. It follows that the symmetric bilinear form $(A/2)^{\vee}$ can be diagonalized over $\mathbb{Z}/r\mathbb{Z}$ by P^{\vee} . The resulting diagonal matrix is D^{\vee} . So we have

$$\gamma_{A,b} = \sum_{\mathbf{k} \in (\mathbb{Z}/r\mathbb{Z})^{n-1}} \zeta^{b \, \mathbf{k}^t D^{\vee} \mathbf{k}}$$

The matrix D^{\vee} is diagonal, with $D_{jj}^{\vee} = (j+1)2^*j^*$. Hence

$$\gamma_{A,b} = \prod_{j=1}^{n-1} \sum_{k \in \mathbb{Z}/r\mathbb{Z}} \zeta^{b(j+1)2^*j^*k^2}.$$

Using Lemmas A.2 and A.3 and the fact that $\left(\frac{j}{r}\right) \left(\frac{j^*}{r}\right) = 1$ we get

$$\gamma_{A,b} = \left(\frac{n}{r}\right) \left[\left(\frac{b}{r}\right) e^{\pi i (1-r)/4} \sqrt{r}\right]^{n-1}$$

This proves formula (4.8).

References

- F. Goodman, H. Wenzl, Littlewood-Richardson coefficients for Hecke algebras at roots of unity, Adv. Math. 82 (1990), no. 2, 244–265.
- [2] J. E. Humphreys, Introduction to Lie algebras and representation theory, Graduate text in Mathenatics, 9, Springer-Verlag, New York.
- [3] K. Ireland, M. Rosen, A classical introduction to modern number theory, Graduate Text in Mathematics No 84, Springer-Verlag, 2nd edition, 1990.
- [4] V. Jones M. Rosso, On the invariants of torus knots derived from quantum groups, J. knot theory ramifications 2 (1993), 97–112.
- [5] V. Kac, Infinite-dimensional Lie algebras, 3rd edition, Cambridge University Press, 1993.
- [6] C. Kassel, Quantum Groups, Graduate Text in Mathematics 155, Springer-Verlag, New York 1995.
- [7] T. Kohno, T. Takata, Symmetry of Witten's 3-manifold invariant for $sl_n(\mathbb{C})$, J. Knot Theory Ramifications, **2** (1993),149–169.
- [8] T. Kohno, T. Takata, Level-rank duality for Witten's 3-manifold invariant Adv. Studies Pure Math., 24, (1996): Progress in Algebraic Combinatorics, 243–264.
- [9] R. Kirby , P. Melvin, The 3-manifold invariants of Witten and Reshetikhin-Turaev for sl(2, ℂ), Invent. Math. 105 (1991), 473–545.
- [10] M. Kontsevich, Vassiliev's knot invariants, Advances in Soviet Mathematics, 16 (1993), 137-150.
- [11] A. Kricker, B. Spence, Ohtsuki's Invariants are of Finite Type, preprint q-alg/9608007.

- [12] R. Lawrence, Asymptotic expansions of Witten-Reshetikhin-Turaev invariants for some simple 3-manifolds, J. Math. Phys. 36 (1995), pp. 6106–6129.
- [13] R. Lawrence and L. Rozansky, Witten-Reshetikhin-Turaev invariants of Seifert manifolds, preprint, University of Michigan, 1997.
- [14] T. T. Q. Le, An invariant of integral homology 3-spheres which is universal for all finite types invariants, in "Soliton, Geometry, and Topology: On the Crossroad", AMS Translations series 2, v. 179 Eds. V. Buchstaber and S. Novikov, pp 75–100; available at http://www.math.buffalo.edu/~letu/.
- [15] T. T. Q. Le, On denominators of the Kontsevich integral and the universal perturbative invariant of 3-manifolds, Invent. Math., to appear; available at http://www.math.buffalo.edu/~letu/.
- [16] T. T. Q. Le, J. Murakami, Representations of the category of tangles by Kontsevich's iterated integral, Commun. Math. Phys., 168 (1995), 535–562.
- [17] T. T. Q. Le, J. Murakami, The universal Vassiliev-Kontsevich invariant for framed oriented links, Compositio Math., 102 (1996), pp. 41–64.
- [18] T. T. Q. Le, J. Murakami and T. Ohtsuki, On a universal perturbative invariant of 3-manifolds, Topology, 37 (1998), 539–574.
- [19] C. Lescop, Global surgery formula for the Casson-Walker invariant, Annals of Mathematics Studies, 140. Princeton University Press, Princeton, NJ, 1996.
- [20] X. S. Lin, Z. Wang, On Ohtsuki's invariants of integral homology 3-spheres, preprint 1996.
- [21] P. Melvin, H. Morton, The coloured Jones function, Commun. Math. Phys., 169 (1995), 501–520.
- [22] G. Masbaum and H. Wenzl, Integral modular categories and integrality of quantum invariants at roots of unity of prime order, preprint 1997.
- [23] H. Murakami, Quantum SO(3)-invariants dominate the SU(2)-invariant of Casson and Walker, Math. Proc. Camb. Phil. Soc. 117 (1995), pp. 237–249.
- [24] T. Ohtsuki, A polynomial invariant of rational homology 3-spheres, Invent. Math., 123 (1996), no. 2, 241–257.
- [25] T. Ohtsuki, Finite type invariants of integral homology 3-spheres, Jour. of Knot Theory and its Ramifications 5 (1996), 101–115.
- [26] T. Ohtsuki, The perturbative SO(3) invariant of rational homology 3-spheres recovers from the universal perturbative invariant, preprint Tokyo Institute of Technology, 1996.
- [27] N. Yu. Reshetikhin, V. Turaev, Ribbon graphs and their invariants derived from quantum groups, Commun. Math. Phys., 127 (1990), 1–26.
- [28] N. Yu. Reshetikhin, V. Turaev, Invariants of 3-manifolds via link polynomials and quantum groups, Invent. Math., 103 (1991), 547–598.

- [29] L. Rozansky, Witten's Invariants of rational Homology 3-spheres at prime values of K and trivial connection contribution, Comm. Math. Phys. 180 (1996), no. 2, 297–324.
- [30] L. Rozansky, Higher order terms in the Melvin-Morton expansion of the colored Jones polynomial, Comm. Math. Phys. 183 (1997), no. 2, 291–306.
- [31] T. Takata, On quantum PSU(n) invariants for Seifert manifolds, J. Knot Theory Ramifications,
 6 (1997), no. 3, 417–426.
- [32] T. Takata and Y. Yokota, The PSU(N) invariants of 3-manifolds are polynomials, preprint Kyushu University, 1996.
- [33] V. G. Turaev, Quantum invariants of knots and 3-manifolds, de Gruyter Studies in Mathematics 18, Walter de Gruyter, Berlin New York 1994.
- [34] V. G. Turaev, H. Wenzl, Quantum invariants of 3-manifolds associated with classical simple Lie algebras, Internat. J. Math. 4 (1993), no. 2, 323–358.
- [35] L. Washington, Introduction to cyclotomic fields, GTM v. 83, Springer-Verlag 1982.
- [36] Y. Yokota, Skeins and quantum SU(N) invariants of 3-manifolds, Math. Ann., to appear.

DEPT. OF MATHEMATICS, SUNY AT BUFFALO, BUFFALO, NY 14214, USA *E-mail address*: letu@newton.math.buffalo.edu