# Cartan Spinor Bundles on Manifolds. \*

Thomas Friedrich, Berlin

July 26, 2021

## 1 Introduction.

Spinor fields and Dirac operators on Riemannian manifolds  $M^n$  can be introduced by means of a reduction of the frame bundle from the structure group O(n) to one of the groups Spin(n), Pin(n) or  $Spin^{\mathbb{C}}(n)$ . In any case the existence of a corresponding reduction imposes topological restrictions on the manifold  $M^n$ . The aim of the present paper is the construction of spinor bundles of Cartan type over certain non-orientable manifolds. The bundles under consideration do not split into subbundles invariant under the action of the Clifford algebra and therefore they are not induced from  $Pin^{\mathbb{C}}$  structure of the manifold. Moreover, we study the case of the real projective space  $\mathbb{RP}^n$  and its spinor bundle in more detail. These bundles over  $\mathbb{RP}^n$  admit a suitable metric connection  $\nabla^S$  and the corresponding Dirac operator. In particular it turns out that there are Killing spinors in the constructed twisted spinor bundle over  $\mathbb{RP}^n$  for all dimensions.

Let us fix some notations. In case  $n \equiv 0 \mod 2$  we denote by  $\gamma$ :  $\operatorname{Cliff}^{\mathbb{C}}(n) \xrightarrow{\sim} \operatorname{End}(\Delta_n)$  the isomorphism between the Clifford algebra and the algebra of endomorphisms of the space  $\Delta_n$  of all Dirac spinor. In this way we obtain the so called *Dirac* representation  $\gamma$  of the Clifford algebra. In case  $n \equiv 1 \mod 2$  we have the *Pauli* representation  $\gamma$ :  $\operatorname{Cliff}^{\mathbb{C}}(n) \to \operatorname{End}(\Delta_n)$ . Denote by  $\alpha$ :  $\operatorname{Cliff}^{\mathbb{C}}(n) \to \operatorname{Cliff}^{\mathbb{C}}(n)$  the canonical involution. Then

$$\gamma \oplus (\gamma \circ \alpha) : \operatorname{Cliff}^{\mathbb{C}}(n) \xrightarrow{\sim} \operatorname{End}(\Delta_n) \oplus \operatorname{End}(\Delta_n)$$

is an isomorphism and the corresponding representation of the Clifford algebra in  $\Delta_n \oplus \Delta_n$  is the so called *Cartan representation* (see [Fr2], [Tr]).

### 2 The construction of spinor bundles of Cartan type.

We consider odd-dimensional Riemannian manifolds  $M^{2k+1}$  only. A spinor bundle of Cartan type is a  $2^{k+1}$ -dimensional complex Clifford bundle S realizing at any point the Cartan representation.  $\gamma \oplus (\gamma \circ \alpha) : \operatorname{Cliff}^{\mathbb{C}}(\mathbb{R}^{2k+1}) \xrightarrow{\sim} \operatorname{End}(\Delta_{2k+1}) \oplus \operatorname{End}(\Delta_{2k+1})$  of the Clifford algebra (see [Tr]). Any spinor bundle of Cartan type over an orientable

<sup>\*</sup>Supported by the SFB 288 of the DFG.

manifold splits into two Clifford subbundles. Indeed, the volume form  $dM^{2k+1} = e_1 \cdot \ldots \cdot e_{2k+1} \in \text{Cliff}(T(M^{2k+1}))$  commutes with the Clifford multiplication. Since

$$\alpha(e_1 \cdot \ldots \cdot e_{2k+1}) = -e_1 \cdot \ldots \cdot e_{2k+1}$$

 $dM^{2k+1}$  acts on S with two different eigenvalues. Therefore  $dM^{2k+1}$  defines a splitting of the bundle S into two Clifford subbundles.

Consider a simply-connected, odd-dimensional Riemannian spin manifold  $M^{2k+1}$ and denote by S its spinor bundle. Any isometry  $\gamma: M^{2k+1} \to M^{2k+1}$  admits two lifts  $\hat{\gamma}_{\pm}: S \to S$  into the spinor bundle S. We once again describe the construction of  $\hat{\gamma}_{\pm}$ . The differential  $d\gamma$  (resp.  $(-d\gamma) - M$  is odd-dimensional !) maps the frame bundle into itself if  $\gamma$  preserves the orientation (resp. does not preserve the orientation). Moreover,  $d\gamma$  (or  $(-d\gamma)$ ) lifts into the spin structure of  $M^{2k+1}$  and defines two lifts  $\hat{\gamma}_{\pm}$ . The lifts  $\hat{\gamma}_{\pm}$  commute (anticommute) with the Clifford multiplication, i.e.

$$\hat{\gamma}_{\pm}(X \cdot \varphi) = \deg(\gamma) d\gamma(X) \cdot \hat{\gamma}_{\pm}(\varphi)$$

(see [Fr1]).

Let  $\Gamma$  be a discrete subgroup of isometries and suppose that  $\Gamma$  acts freely on  $M^{2k+1}$ . We consider the spinor bundle of Cartan type

$$S^* = S \oplus S$$

over  $M^{2k+1}$ . The Clifford multiplication in  $S^*$  is given by the formula

$$X \cdot \begin{pmatrix} \varphi \\ \psi \end{pmatrix} = \begin{pmatrix} X \cdot \varphi \\ -X \cdot \psi \end{pmatrix} \qquad \qquad X \in T(M^{2k+1})$$

where  $X \cdot \varphi$  denotes the Clifford multiplication in the spinor bundle S. Any isometry  $\gamma \in \Gamma$  admits the following 4 lifts into the bundle  $S^*$ :

$$\begin{pmatrix} \hat{\gamma}_{\pm} & 0\\ 0 & \hat{\gamma}_{\pm} \end{pmatrix}$$
 in case  $\gamma$  preserves the orientation.  
$$\begin{pmatrix} 0 & \hat{\gamma}_{\pm}\\ \hat{\gamma}_{\pm} & 0 \end{pmatrix}$$
 in case  $\gamma$  reverses the orientation.

Any of these lifts commutes with the Clifford multiplication in the bundle  $S^*$ .

We introduce now a family of automorphisms  $\Pi(\Theta_1, \Theta_2, \varepsilon_1, \varepsilon_2, \gamma)$  of the bundle  $S^*$  depending on four parameters  $0 \leq \Theta_1 \leq 2\pi$ ,  $0 \leq \Theta_2 \leq 2\pi$  and  $\varepsilon_1, \varepsilon_2 \in \{\pm 1\}$ . If  $\gamma$  preserves the orientation of the manifold we define

$$\Pi(\Theta_1, \Theta_2, \varepsilon_1, \varepsilon_2, \gamma) = \begin{pmatrix} e^{i\Theta_1}\hat{\gamma}_{\varepsilon_1} & 0\\ & & \\ 0 & e^{i\Theta_2}\hat{\gamma}_{\varepsilon_2} \end{pmatrix},$$

otherwise let us introduce the automorphism

$$\Pi(\Theta_1, \Theta_2, \varepsilon_1, \varepsilon_2, \gamma) = \begin{pmatrix} 0 & e^{i\Theta_2} \hat{\gamma}_{\varepsilon_2} \\ & & \\ e^{i\Theta_1} \hat{\gamma}_{\varepsilon_1} & 0 \end{pmatrix}.$$

Suppose that  $\varepsilon : \Gamma \to \operatorname{Aut}(S^*)$  is a homomorphism such that for any  $\gamma \in \Gamma$  the automorphism  $\varepsilon(\gamma)$  coincides with one of the lifts  $\Pi(\Theta_1, \Theta_2, \varepsilon_1, \varepsilon_2, \gamma)$  of  $\gamma$  into  $S^*$ . Then

$$\bar{S} = S^* / \varepsilon(\Gamma)$$

is a Clifford bundle of Cartan type over the Riemannian manifold  $\overline{M}^{2k+1} = M^{2k+1}/\Gamma$ . The bundle  $\overline{S}$  splits into two parts invariant under the action of the Clifford bundle Cliff $(T(M^{2k+1}))$  if and only if  $\overline{M}^{2k+1}$  is orientable. For a non-orientable manifold  $\overline{M}^{2k+1}$  the non-splitting bundle  $\overline{S}$  is not induced from a  $Pin^{\mathbb{C}}(2k+1)$  structure of  $\overline{M}^{2k+1}$ .

#### Example: $\Gamma = \mathbb{Z}_p$ .

Suppose  $\Gamma$  is generated by one involution  $\gamma: M^{2k+1} \to M^{2k+1}$  with the lifts  $\hat{\gamma}_{\pm}$ . If  $\gamma$  preserves the orientation and  $\hat{\gamma}_{\pm}^2 = \text{Id}$  the manifold  $\bar{M}^{2k+1} = M^{2k+1}/\mathbb{Z}_2$  admits a spin structure. In case  $\hat{\gamma}_{\pm}^2 = -\text{Id}$   $\bar{M}^{2k+1}$  is orientable, but does not admit a spin structure. Nevertheless we can construct a spinor bundle using the homomorphism

$$\varepsilon(\gamma) = \left(\begin{array}{cc} i\hat{\gamma}_+ & 0\\ 0 & i\hat{\gamma}_+ \end{array}\right)$$

If  $\gamma$  does not preserve the orientation we define the homomorphism  $\varepsilon : \mathbb{Z}_2 \to \operatorname{Aut}(S^*)$  by the formula

$$\varepsilon(\gamma) = \begin{pmatrix} 0 & \hat{\gamma}_+ \\ \hat{\gamma}_+ & 0 \end{pmatrix} \quad \text{if } \hat{\gamma}_+^2 = \text{Id}$$
$$\varepsilon(\gamma) = \begin{pmatrix} 0 & i\hat{\gamma}_+ \\ i\hat{\gamma}_+ & 0 \end{pmatrix} \quad \text{if } \hat{\gamma}_+^2 = -\text{Id}.$$

A similar construction for an arbitrary cyclic group  $\mathbb{Z}_p$  yields the result:

Let  $\bar{M}^{2k+1}$  be a Riemannian manifold with fundamental group  $\pi_1(\bar{M}^{2k+1}) = \mathbb{Z}_p$ and suppose that the universal covering  $M^{2k+1}$  admits a spin structure. Then over  $\bar{M}^{2k+1}$  there exists a spinor bundle  $\bar{S}$  of Cartan type. The bundle  $\bar{S}$  splits into two subbundles invariant under the action of the Clifford bundle  $\operatorname{Cliff}(T(\bar{M}^{2k+1}))$  if and only if  $\bar{M}^{2k+1}$  is orientable.

Let  $D: \Gamma(S) \to \Gamma(S)$  be the Dirac operator acting on sections of the spinor bundle S over  $M^{2k+1}$ . The Dirac operator  $\mathfrak{D}$  of the spinor bundle  $S^*$  of Cartan type is given by

$$\mathfrak{D} = \left( \begin{array}{cc} D & 0 \\ 0 & -D \end{array} \right)$$

Therefore the eigenspaces  $E_{\lambda}(\mathfrak{D})$  consists of pairs  $\begin{pmatrix} \varphi \\ \psi \end{pmatrix}$  of eigenspinors of D, i.e.

$$E_{\lambda}(\mathfrak{D}) = E_{\lambda}(D) \oplus E_{-\lambda}(D).$$

The group  $\Gamma$  acts on  $E_{\lambda}(\mathfrak{D})$  by the matrices  $\varepsilon(\gamma)$  and the eigenspace of the Dirac operator  $\overline{\mathfrak{D}}$  on  $\overline{M}^{2k+1}$  coincides with the subspace of  $\varepsilon(\Gamma)$ -invariant pairs in  $E_{\lambda}(\mathfrak{D})$ . In particular, in case  $\Gamma = \mathbb{Z}_2$  and  $\overline{M}^{2k+1}$  is non-orientable we have only one transformation

$$\varepsilon(\gamma) = \begin{pmatrix} 0 & \hat{\gamma}_+ \\ \hat{\gamma}_+ & 0 \end{pmatrix} \quad \text{or} \quad \varepsilon(\gamma) = \begin{pmatrix} 0 & i\hat{\gamma}_+ \\ i\hat{\gamma}_+ & 0 \end{pmatrix}.$$

The map  $E_{\lambda}(D) \ni \varphi \longmapsto \begin{pmatrix} \varphi \\ \hat{\gamma}_{+}(\varphi) \end{pmatrix} \in E_{\lambda}(\bar{\mathfrak{D}}) \text{ (resp. } \varphi \longmapsto \begin{pmatrix} \varphi \\ i\hat{\gamma}_{+}(\varphi) \end{pmatrix} \text{) defines an isomorphism between the corresponding eigenspaces. Therefore the spectrum of the operator <math>(\bar{\mathfrak{D}}, \bar{S}, \bar{M}^{2k+1})$  coincides with the spectrum of the operator  $(D, S; M^{2k+1})$ .

We thus constructed non-splitting spinor bundles over non-orientable manifolds  $\overline{M}^{2k+1}$  via coverings. We mention that other situations give similarly bundles with the described properties. For example, any immersion of  $\overline{M}^{2k+1}$  into an oriented spin manifold  $N^{2k+2}$  also defines a spinor bundle of Cartan type over  $\overline{M}^{2k+1}$ . Indeed, let  $f: \overline{M}^{2k+1} \to N^{2k+2}$  be the immersion and consider the Dirac spinor bundle S of  $N^{2k+2}$ . Then the induced bundle  $\overline{S} = f^*(S)$  inherits the structure of a spinor bundle of Cartan type over  $\overline{M}^{2k+1}$ .

### 3 The Spinor Bundle over $\mathbb{RP}^n$ .

We start with the n-dimensional sphere

$$S^{n} = \{ x \in \mathbb{R}^{n+1} : |x| = 1 \}$$

and we denote by  $S^*$  the  $2^{\left[\frac{n+1}{2}\right]}$ -dimensional bundle

$$S^* = S^n \times \Delta_{n+1}.$$

The tangent bundle of the sphere

$$T(S^n) = \{(x,t) \in S^n \times \mathbb{R}^{n+1} : \langle x,t \rangle = 0\}$$

acts by Clifford multiplication

$$\mu((x,t)\otimes(x,\varphi)):=(x,t\cdot\varphi)$$

on  $S^*$ . Therefore  $S^*$  is a Clifford bundle over  $S^n$ . Let  $g: S^n \to S^n, g(x) = -x$  be the antipodal map. Its differential  $dg: T(S^n) \to T(S^n)$  acts on a tangent vector  $(x,t) \in T(S^n)$  by

$$dg(x,t) = (-x,-t).$$

Let us define a lift  $\hat{g}$  of the antipodal map into the bundle  $S^*$  by the formula

$$\hat{g}(x,\varphi) = (-x, x \cdot \varphi).$$

Then  $\hat{g}$  is again an involution,  $(\hat{g})^2 = \text{Id.}$  Moreover,  $\hat{g}$  respects the Clifford multiplication, i.e. the diagram

$$T(S^{n}) \otimes S^{*} \xrightarrow{\mu} S^{*}$$

$$\downarrow^{(dg) \otimes \hat{g}} \qquad \qquad \downarrow^{\hat{g}}$$

$$T(S^{n}) \otimes S^{*} \xrightarrow{\mu} S^{*}$$

commutes. Indeed, given  $(x,t) \otimes (x,\varphi) \in T(S^n) \otimes S^*$  we have

$$\mu \circ (dg \otimes \hat{g})((x,t) \otimes (x,\varphi)) = \mu((-x,-t) \otimes (-x,x \cdot \varphi)) = (-x,-t \cdot x \cdot \varphi).$$

An the other hand,

$$\hat{g}\mu((x,t)\otimes(x,\varphi)) = \hat{g}((x,t\cdot\varphi)) = (-x,x\cdot t\cdot\varphi).$$

Since x and t are orthogonal vectors in  $\mathbb{R}^{n+1}$  we have in the Clifford algebra the relation

$$x \cdot t + t \cdot x = 0$$

i.e. the mentioned diagram commutes.

Over the real projective space  $\mathbb{RP}^n = S^n/g$  we define the bundle

$$S = S^* / \hat{g}.$$

The tangent bundle  $T(\mathbb{RP}^n)$  can be identified with  $T(\mathbb{RP}^n) = T(S^n)/dg$  and the above discussed property of the lift  $\hat{g} : S^* \to S^*$  implies that we obtain a well-defined Clifford multiplication

$$\mu: T(\mathbb{RP}^n) \otimes S \to S.$$

At any point of  $\mathbb{RP}^n$  the bundle S realizes the Dirac (in case  $n \equiv 0 \mod 2$ ) or the Cartan (in case  $n \equiv 1 \mod 2$ ) representation of the *n*-dimensional Clifford algebra.

Let us consider the even-dimensional case, n = 2k. The Dirac representation of the Clifford algebra decomposes into two irreducible representations with respect to the action of the even part  $\operatorname{Cliff}_{o}^{\mathbb{C}}(n)$  of the Clifford algebra. Therefore the bundle S decomposes at any fixed point as a Clifford module too. However, globally the bundle S over  $\mathbb{RP}^n$  does not split as a  $\operatorname{Cliff}_o(T(\mathbb{RP}^n))$ -bundle. Indeed suppose that there exists a decomposition  $S = S_1 \oplus S_2$  invariant under the action of the Clifford bundle. Then we obtain a corresponding decomposition  $S^* = S_1^* \oplus S_2^*$  of the bundle  $S^*$  over  $S^n$ . The spin module  $\Delta_{n+1}$  decomposes in a unique way as a  $\operatorname{Cliff}_o(T_x S^n)$ module

$$\Delta_{n+1} = \Delta_{n+1}^+(x) \oplus \Delta_{n+1}^-(x).$$

We describe this decomposition in an explicit way. Let  $e_1, \ldots, e_n$  an orthonormal basis of the tangent space  $T_x(S^n)$  and consider the element

$$f = i^{n/2} e_1 \cdot \ldots \cdot e_n : \Delta_{n+1} \to \Delta_{n+1}.$$

Then  $f^2 = \text{Id}$  and  $\Delta_{n+1}$  decomposes into the eigenspaces of  $f, \Delta_{n+1} = \Delta_{n+1}^+(x) \oplus \Delta_{n+1}^-$  with

$$\Delta_{n+1}^{\pm}(x) = \{ \varphi \in \Delta_{n+1} : \quad f(\varphi) = \pm \varphi \}.$$

Now the volume form  $e_1 \cdot \ldots \cdot e_n \cdot x$  of  $\mathbb{R}^{n+1}$  acts on  $\Delta_{n+1}$  by multiplication

$$e_1 \cdot \ldots \cdot e_n \cdot x_{|\Delta_{n+1}} = \alpha_{n+1}.$$

Then  $f = i^{n/2} e_1 \cdot \ldots \cdot e_n = -i^{n/2} \alpha_{n+1} x$  and the subspaces  $\Delta_{n+1}^{\pm}(x)$  are given by

$$\Delta_{n+1}^{\pm}(x) = \{\varphi \in \Delta_{n+1} : \ \pm x\varphi = i^{n/2} \alpha_{n+1}\varphi\}.$$

The bundles  $S_1^*$  and  $S_2^*$  coincide therefore with the subbundles  $\Delta_{n+1}^{\pm}(x)$ . Since  $S_i^*(i=1,2)$  are induced bundles by the covering  $S^n \to \mathbb{RP}^n = S^n/g$  they are invariant under the lift  $\hat{g} : S^* \to S^*$ . However, the lift  $\hat{g}$  defined by the Clifford multiplication does not preserve the decomposition  $\Delta_{n+1} = \Delta_{n+1}^+(x) \oplus \Delta_{n+1}^-(x)$ , a contradiction.

We discuss now the odd-dimensional case, n = 2k + 1. Again we start with an orthonormal basis  $e_1, \ldots, e_n \in T_x(S^n)$  and we introduce the automorphism

$$f = e_1 \cdot \ldots \cdot e_n : \Delta_{n+1} \to \Delta_{n+1}.$$

Then we have

$$f^2 = (-1)^{k+1}$$
 ,  $f \cdot x = -x \cdot f$ 

and

$$f \cdot t = t \cdot f$$
 for all tangent vectors  $t \in T_x(S^n)$ .

We decompose  $\Delta_{n+1}$  into the eigenspaces of f

$$\Delta_{n+1} = \Delta_{n+1}^+(x) \oplus \Delta_{n+1}^-(x) \quad , \quad \Delta_{n+1}^\pm(x) = \{\varphi \in \Delta_{n+1} : e_1 \cdot \ldots \cdot e_n \varphi = \pm i^{k+1} \varphi\}.$$

Since g commutes with all tangent vectors  $t \in T_x(S^n)$  the subspaces  $\Delta_{n+1}^{\pm}(x)$  are invariant under the action of the Clifford algebra  $\operatorname{Cliff}(T_x(S^n))$ . The lift  $\hat{g}$  preserves

now (n = 2k + 1) the decomposition  $\Delta_{n+1} = \Delta_{n+1}^+(x) \oplus \Delta_{n+1}^-(x)$ , i.e. the Clifford bundle S over  $\mathbb{RP}^n$  decomposes into two Clifford subbundles.

Vector fields V on the projective space  $\mathbb{RP}^n$  can be identified with maps  $V:S\to\mathbb{R}^{n+1}$  such that

$$V(-x) = -V(x) \quad , \quad \langle V(x), x \rangle = 0 \quad x \in S^n.$$

The bundle S over  $\mathbb{RP}^n$  arises from  $S^* = S^n \times \Delta_{n+1}$  and the identification  $\hat{g}(x, \varphi) = (-x, x \cdot \varphi)$ . Therefore a section  $\Phi \in \Gamma(S; \mathbb{RP}^n)$  is a map  $\Phi : S^n \to \Delta_{n+1}$  with the property

$$\Phi(-x) = x \cdot \Phi(x) . \tag{(*)}$$

The formula

$$(\nabla_V^S \Phi)(x) = d\Phi(V)(x) + \frac{1}{2}V(x) \cdot x \cdot \Phi(x)$$

defines a covariant derivative

$$\nabla^S : \Gamma(T(\mathbb{RP}^n)) \times \Gamma(S) \to \Gamma(S).$$

Indeed, a simple calculation shows that  $\nabla_V^S \Phi$  has the invariance property (\*) in case  $\Phi$  and V satisfy the corresponding transformation rules. Moreover, the covariant derivative  $\nabla^S$  is compatible with the Clifford multiplication, i.e.

$$\nabla_V^S(W \cdot \Phi) = (\nabla_V W) \cdot \Phi + W \cdot (\nabla_V^S \Phi),$$

and  $\nabla^S$  preserves the hermitian metric of the bundle S. To sum up, we see that the triple  $(S, \nabla^S, \langle, \rangle)$  is a Dirac bundle over  $\mathbb{RP}^n$  (see [LM] for the general definition of a Dirac bundle). A direct calculation yields the formula

$$R^{S}(V,W)\Phi = \frac{1}{2}\left(W \cdot V + \langle V,W \rangle\right) \cdot \Phi$$

for the curvature tensor  $R^S$  of the connection  $\nabla^S$ .

**Remark:** Let us compare the above defined covariant derivative  $\nabla^S$  with the usual covariant derivative  $\nabla$  in the usual spin bundle of a hypersurface in  $\mathbb{R}^{n+1}$ . The formula for  $\nabla$  is

$$\nabla_V \Phi = d\Phi(V) + \frac{1}{2} \Pi(V) \cdot \vec{N} \cdot \Phi$$

i.e. over the sphere  $S^n$  the introduced Dirac bundle coincides with the usual spinor bundle of the sphere.

The Dirac operator  $D = \sum_{i=1}^{n} e_i \nabla_{e_i}^S$  acting on sections  $\Gamma(S; \mathbb{RP}^n)$  is related to the Laplace operator  $\Delta = -\sum_{i=1}^{n} \nabla_{e_i}^S \nabla_{e_i}^S - \sum_{i=1}^{n} \operatorname{div}(e_i) \nabla_{e_i}^S$  by the well-known formula

$$D^2 = \Delta + R$$

where the endomorphism R (in case of a general Dirac bundle) is given by

$$R = \frac{1}{2} \sum_{j,k} e_j e_k R^S(e_j, e_k).$$

In our situation we simply obtain

$$D^2 = \Delta + \frac{1}{4}n(n-1) = \Delta + \frac{\tau}{4}$$

where  $\tau$  is the scalar curvature of  $\mathbb{RP}^n$ . The bundle S over  $\mathbb{RP}^n$  admits Killing spinors, i.e. spinor fields  $\Phi$  satisfying the differential equation

$$\nabla_V^S \Phi = \lambda V \cdot \Phi$$

for some  $\lambda \in \mathbb{R}^1$ . Indeed, suppose that  $\Phi$  is a Killing spinor in  $\Gamma(S; \mathbb{RP}^n)$ . Then we obtain

$$R^{S}(V,W)\Phi = 2\lambda^{2}(WV + \langle V,W\rangle) \cdot \Phi$$

using the differential equation. On the other hand, the formula for the curvature tensor implies

$$R^{S}(V,W)\Phi = \frac{1}{2}(WV + \langle V,W\rangle) \cdot \Phi,$$

i.e.  $\lambda = \pm \frac{1}{2}$ . Let us consider the spinor field

$$\Phi(x) = (1-x)\Phi_0$$

where  $\Phi_0 \in \Delta_{n+1}$  is constant. Then

$$\Phi(-x) = (1+x)\Phi_0 = x(1-x)\Phi_0 = x\Phi(x)$$

and therefore  $\Phi \in \Gamma(S; \mathbb{RP}^n)$  is a section in the spinor bundle over  $\mathbb{RP}^n$ . We calculate the covariant derivative  $\nabla_V^S \Phi$ :

$$(\nabla_V^S \Phi)(x) = -V(x) \cdot \Phi_0 + \frac{1}{2} V(x) \cdot x \cdot (1-x) \Phi_0 = = -V(x) \cdot \Phi_0 + \frac{1}{2} V(x) \cdot (1+x) \Phi_0 = = \frac{1}{2} V(x) \cdot (-1+x) \cdot \Phi_0 = -\frac{1}{2} V(x) \cdot \Phi(x)$$

,

i.e.  $\Phi(x)$  is a Killing spinor with Killing number  $\lambda = -\frac{1}{2}$ .

**Remark:** Killing spinors with Killing number  $\lambda = +\frac{1}{2}$  on  $\mathbb{RP}^n$  appear if we use the identification  $\hat{g}_-: S^* \to S^*$  defined by the formula

$$\hat{g}(x,\varphi) = (-x, -x \cdot \varphi).$$

 $\hat{g}_-$  defines in a similar way a Clifford bundle  $S_- = S^*/\hat{g}_-$  whose sections are maps  $\Phi: S^n \to \Delta_{n+1}$  with the property

$$\Phi(-x) = -x\Phi(x)$$

The spinor fields

$$\Phi(x) = (1+x)\Phi_0$$

are Killing spinors in the bundle  $S_{-}$  over  $\mathbb{RP}^{n}$  with Killing number  $\lambda = +\frac{1}{2}$ .

Killing spinors on compact Riemannian spin manifolds with positive scalar curvature correspond to eigenspinors of the Dirac operator related to the smallest eigenvalue. In case  $\mathbb{RP}^{4k+3}$  the first eigenvalue of the Dirac operator on usual spinors has been calculated  $\left(\lambda_1 = \pm \frac{1}{2}\sqrt{\frac{n}{n-1}\tau}\right)$  and there are Killing spinors (see [F1]). More general, we can calculate the spectrum of the Dirac operator acting on sections  $\Phi \in \Gamma(S; \mathbb{RP}^n)$ using the computation of the eigenvalues of D on the sphere. Let  $\Phi: S^n \to \Delta_{n+1}$ be a section in the bundle  $S^*$ . We decompose  $\Phi$  into  $\Phi = \Phi_+ + \Phi_-$  with

$$\Phi_{+}(x) = \frac{\Phi(x) - x \cdot \Phi(-x)}{2} \quad , \quad \Phi_{-}(x) = \frac{\Phi(x) + x \cdot \Phi(-x)}{2}.$$

The sections  $\Gamma(S; \mathbb{RP}^n)$  are described by the condition  $\Phi_- \equiv 0$ . Now we apply the calculation of the spectrum of the Dirac operator on spheres as well as the realization of eigenspinors by polynomials (see [Su], [Tr]). Imposing the additional condition  $\Phi_- \equiv 0$  we obtain the spectrum of the Dirac operator acting on the sections  $\Gamma(S; \mathbb{RP}^n)$  for arbitrary n.

### References

- [F1] Th. Friedrich. Der erste Eigenwert des Dirac Operators einer kompakten Riemannschen Mannigfaltigkeit nichtnegativer Skalarkrümmung, Math. Nachr. 97 (1980), 117-146.
- [F2] Th. Friedrich. Dirac-Operatoren in der Riemannschen Geometrie, Vieweg-Verlag Braunschweig/ Wiesbaden 1997.
- [FT] Th. Friedrich and A. Trautman. Clifford structures and spinor bundles, Preprint SFB 288 No. 251, Berlin 1996.
- [LM] H.B. Lawson and M.L. Michelsohn. Spin Geometry, Princeton 1989.
- [Su] S. Sulanke. Berechnung des Spektrums des Quadrates des Dirac-Operators auf Sphären und Untersuchungen zum ersten Eigenwert von D auf 5dimensionalen Räumen konstanter positiver Schnittkrümmung, Dissertation, Humboldt-Universität zu Berlin 1980.
- [Tr] A. Trautman. The Dirac operator on hypersurfaces, Acta Phys. Polon. B 26 (1995), 1283–1310.

Thomas Friedrich Humboldt-Universität zu Berlin Institut für Reine Mathematik Sitz: Ziegelstraße 13a D-10099 Berlin e-mail: friedric@mathematik.hu-berlin.de