# Drinfeldians

Valeriy N. Tolstoy

Institute of Nuclear Physics Moscow State University 119899 Moscow&Russia<sup>1</sup>

#### Abstract

We construct two-parameter deformation of an universal enveloping algebra  $U(q[u])$  of a polynomial loop algebra  $q[u]$ , where q is a finite-dimensional complex simple Lie algebra (or superalgebra). This new quantum Hopf algebra called the Drinfeldian  $D_{q\eta}(g)$  can be considered as a quantization of  $U(g[u])$  in the direction of a classical r-matrix which is a sum of the simple rational and trigonometric r-matrices. The Drinfeldian  $D_{q\eta}(g)$  contains  $U_q(g)$  as a Hopf subalgebra, moreover  $U_q(g[u])$  and  $Y_\eta(g)$  are its limit quantum algebras when the  $D_{q\eta}(g)$  deformation parameters  $\eta$  goes to 0 and q goes to 1, respectively. These results are easy generalized to a supercase, i.e. when  $g$  is a finite-dimensional contragredient simple superalgebra.

#### 1 Introduction

As it is well known, an universal enveloping algebra  $U(g[u])$  of a polynomial loop (current) Lie algebra  $g[u]$ , where g is a finite-dimensional complex simple Lie algebra, admits two type deformations: a trigonometric deformation  $U_q(g[u])$  and a rational deformationor Yangian  $Y_{\eta}(g)$  [[1, 2\]](#page-11-0). (In the case  $g = sl_n$  there also exists an elliptic quantum deformation of  $U(sl_n[u])$ . The algebras  $U_q(g[u])$ , and  $Y_q(g)$  are quantizations of  $U(g[u])$  in the direction of the simplest trigonometric and rational solutions of the classical Yang-Baxter equation over  $g$ , respectively. These deformations are one-parameter ones. It turns out that  $U(q[u])$  also admits two-parameter deformation which is called the Drinfeldian or the rational-trigonometric quantum algebra.

<sup>1</sup> *e-mail: tolstoy@anna19.npi.msu.su*

The Drinfeldian  $D_{q\eta}(g)$  is a quantization of  $U(g[u])$  in the direction of a classical r-matrix which is a sum of the simplest rational and trigonometric r-matrices. The Drinfeldian  $D_{q\eta}(g)$  contains  $U_q(g)$  as a Hopf subalgebra, and  $U_q(g[u])$  and  $Y_\eta(g)$  are its limit quantum algebras when the deformation parameters of  $D_{qn}(g)$   $\eta$  goes to 0 and  $q$  goes to 1, respectively. These results are easy generalized to a supercase, i.e. when g is a finite-dimensional contragredient simple superalgebra.

# 2 Quantum algebra  $U_q(g[u])$

Let g be a finite-dimensional complex simple Lie algebra of a rank  $r$  with a standard Cartan matrix  $A = (a_{ij})_{i,j=1}^r$ , with a system of simple roots  $\Pi := {\alpha_1, \ldots, \alpha_r}$  and a maximal positive root  $\theta$ , and with a Chevalley basis  $\{h_{\alpha_i}, e'_{\pm \alpha_i}, (i = 1, 2, \ldots, r)\}\.$  Let  $g[u]$  be a polynomial loop algebra (or a Lie algebra of polynomial currents over g), and g[u] be a "central extension" of g[u]:  $g[u] \simeq g[u] \oplus \mathbb{C}c$ , where c is a central element.<sup>2</sup> The Lie algebra  $g[u]$  (and its universal enveloping algebra  $U(g[u])$ ) is generated by the Chevalley basis of g and the affine element  $e'_{(\delta-\theta)} := ue_{-\theta}$  and  $h_{\delta} := c$  with the following defining relations:

$$
[h_{\delta}, \text{everything}] = 0 , \qquad [h_{\alpha_i}, h_{\alpha_j}] = 0 , \qquad (2.1)
$$

$$
[h_{\alpha_i}, e'_{\pm \alpha_j}] = \pm (\alpha_i, \alpha_j) e'_{\pm \alpha_j} , \qquad [e'_{\alpha_i}, e'_{-\alpha_j}] = \delta_{ij} h_{\alpha_i} , \qquad (2.2)
$$

$$
(\operatorname{ad} e'_{\pm \alpha_i})^{1 - a_{ij}} e'_{\pm \alpha_j} = 0 \qquad \text{for } i \neq j , \qquad (2.3)
$$

$$
[h_{\alpha_i}, e'_{\delta - \theta}] = -(\alpha_i, \theta) e'_{\delta - \theta}, \qquad [e'_{-\alpha_i}, e'_{\delta - \theta}] = 0 , \qquad (2.4)
$$

$$
(\text{ad } e'_{\alpha_i})^{1-a_{i0}} e'_{\delta-\theta} = 0, \qquad (a_{i0} = -2(\alpha_i, \theta)/(\alpha_i, \alpha_i)), \qquad (2.5)
$$

$$
[[e'_{\alpha_i}, e'_{\delta-\theta}], e'_{\delta-\theta}] = 0 \qquad \text{for } g \neq sl_2 \text{ and } (\alpha_i, \theta) \neq 0 , \qquad (2.6)
$$

$$
[[[e'_{\alpha}, e'_{\delta - \alpha}], e'_{\delta - \alpha}], e'_{\delta - \alpha}] = 0 \qquad \text{for } g = sl_2 .
$$
 (2.7)

Here "ad" is the adjoint action of  $\widetilde{g[u]}$  in  $\widetilde{g[u]}$ , i.e.  $(\text{ad } x)y = [x, y]$  for any  $x, y \in \widetilde{g[u]}$ . The relations (2.6) relate to the case  $g \neq sl_2$ , and the relation (2.7) belongs to the case  $g = sl_2$  (in this case  $\theta = \alpha_{r=1}$  and we set  $\alpha := \alpha_{r=1}$ ).

*Remark*. The defining relations for  $g[u]$  can be obtained from defining relations of the corresponding non-twisted affine Lie algebra  $\hat{q}$  by removing relations with a negative affine root vector  $e'_{-\delta+\theta}$ .

Let  $U_q(g)$  be a standard q-deformation of the universal enveloping algebra  $U(g)$ with Chevalley generators  $k_{\alpha_i}^{\pm 1}$ ,  $e_{\pm \alpha_i}$   $(i = 1, 2, \ldots, r)$  and with the defining relations

<sup>&</sup>lt;sup>2</sup>More correctly, the element  $c$  is a central element of a central extension of the total loop algebra  $g[u, u^{-1}].$ 

<span id="page-2-0"></span>
$$
[k_{\alpha_i}, k_{\alpha_j}] = 0 , \qquad k_{\alpha_i} e_{\pm \alpha_j} k_{\alpha_i}^{-1} = q^{\pm (\alpha_i, \alpha_j)} e_{\pm \alpha_j} , \qquad (2.8)
$$

$$
[e_{\alpha_i}, e_{-\alpha_i}] = \frac{k_{\alpha_i} - k_{\alpha_i}^{-1}}{q - q^{-1}}, \qquad (\text{ad}_q e_{\pm \alpha_i})^{1 - a_{ij}} e_{\pm \alpha_j} = 0 \quad \text{for } i \neq j ,
$$
 (2.9)

where  $(\text{ad}_q e_\beta) e_\gamma$  is the q-commutator:

$$
(\text{ad}_q e_\beta) e_\gamma := [e_\beta, e_\gamma]_q := e_\beta e_\gamma - q^{(\beta, \gamma)} e_\gamma e_\beta . \tag{2.10}
$$

A Hopf structure of  $U_q(g)$  is given the following formulas for a comultiplication  $\Delta_q$ , an antipode  $S_q$ , and a co-unite  $\varepsilon_q$ :

$$
\Delta_q(k_{\alpha_i}^{\pm 1}) = k_{\alpha_i}^{\pm 1} \otimes k_{\alpha_i}^{\pm 1} , \qquad \Delta_q(e_{\alpha_i}) = e_{\alpha_i} \otimes 1 + k_{\alpha_i}^{-1} \otimes e_{\alpha_i} ,
$$
  

$$
\Delta_q(e_{-\alpha_i}) = e_{-\alpha_i} \otimes k_{\alpha_i} + 1 \otimes e_{-\alpha_i} ,
$$
 (2.11)

$$
S_q(k_{\alpha_i}^{\pm 1}) = k_{\alpha_i}^{\mp 1} , \qquad S_q(e_{\alpha_i}) = -k_{\alpha_i}e_{\alpha_i} , \qquad S_q(e_{-\alpha_i}) = -e_{-\alpha_i}k_{\alpha_i}^{-1} , \qquad (2.12)
$$

$$
\varepsilon_q(k_{\alpha_i}^{\pm 1}) = 1 , \qquad \varepsilon_q(e_{\pm \alpha_i}) = 0 . \qquad (2.13)
$$

**Definition 2.1** *The quantum algebra*  $U_q(\widetilde{g[u]})$  *(or a q-deformation of*  $U(\widetilde{g[u]})$ *) is generated (as an associative algebra) by the algebra*  $U_q(g)$  *and the elements*  $e_{\delta-\theta}$ ,  $k_{\delta}^{\pm1}$ *with the relations:*

$$
[k_{\delta}^{\pm 1}, \text{everything}] = 0 , \qquad k_{\alpha_i} e_{\delta - \theta} k_{\alpha_i}^{-1} = q^{-(\alpha_i, \theta)} e_{\delta - \theta} , \qquad (2.14)
$$

$$
[e_{-\alpha_i}, e_{\delta-\theta}] = 0 , \qquad (ad_q e_{\alpha_i})^{n_{i0}} e_{\delta-\theta} = 0 , \qquad (2.15)
$$

*where*  $n_{i0} = 1 + 2(\alpha_i, \theta)/(\alpha_i, \alpha_i)$ , and

$$
[[e_{\alpha_i}, e_{\delta - \theta}]_q, e_{\delta - \theta}]_q = 0 \qquad \text{for } g \neq sl_2 \text{ and } (\alpha_i, \theta) \neq 0 , \qquad (2.16)
$$

$$
[[[e_{\alpha}, e_{\delta - \alpha}]_q, e_{\delta - \alpha}]_q, e_{\delta - \alpha}]_q = 0 \quad \text{for } g = sl_2 .
$$
 (2.17)

*The Hopf structure of*  $U_q(g[u])$  *is defined by the formulas*  $\Delta_q(x) = \Delta_q(x)$ *,*  $S_q(x) =$  $S_q(x)$  ( $x \in U_q(g)$ ), and  $\Delta_q(k_\delta^{\pm})$  $\lambda_{\delta}^{\pm}$  =  $k_{\delta}^{\pm 1}$ ,  $S_q(k_{\delta}^{\pm 1}) = k_{\delta}^{\mp 1}$ . The comultiplication, *the antipode and the co-unite of the element*  $e_{\delta-\alpha}$  are given by

$$
\Delta_q(e_{\delta-\theta}) = e_{\delta-\theta} \otimes 1 + k_{\delta-\theta}^{-1} \otimes e_{\delta-\theta} , \qquad (2.18)
$$

$$
S_q(e_{\delta-\theta}) = -k_{\delta-\theta}e_{\delta-\theta} , \qquad \varepsilon(e_{\delta-\theta}) = 0 . \qquad (2.19)
$$

*Here we put*

$$
k_{\delta-\theta} = k_{\delta} k_{\alpha_1}^{-n_1} k_{\alpha_2}^{-n_2} \cdots k_{\alpha_r}^{-n_r}
$$
 (2.20)

 $if \theta = n_1 \alpha_1 + n_2 \alpha_2 + \cdots + n_r \alpha_r.$ 

<span id="page-3-0"></span>*Remark*. The defining relations for  $U_q(\widetilde{g(u)})$  can be obtained from defining relations of the corresponding quantum non-twisted affine algebra  $U_q(\hat{g})$  by removing relations with the negative affine root vector  $e_{-\delta+\theta}$ .

It is easy to check the following result.

**Proposition 2.1** *There is a one-parameter group of Hopf algebra automorphisms*  $\mathcal{T}_a$ *of*  $U_q(g[u])$ *,*  $a \in \mathbb{C}$ *, given by* 

$$
\mathcal{T}_a(k_\delta^{\pm 1}) = k_\delta^{\pm 1} , \qquad \mathcal{T}_a(k_{\alpha_i}^{\pm 1}) = k_{\alpha_i}^{\pm 1} ,
$$
  

$$
\mathcal{T}_a(e_{\pm \alpha_i}) = e_{\pm \alpha_i} , \qquad \mathcal{T}_a(e_{\delta - \theta}) = a e_{\delta - \theta} .
$$
 (2.21)

# 3 Drinfeldian  $D_{qn}(g)$

Here we keep the notations of the previous Section and begin with the following important definition.

**Definition 3.1** *The Drinfeldian*  $D_{q\eta}(g)$  *is generated* (*as an associative algebra over*  $\mathcal{C}[[\eta]]$ ) by the algebra  $U_q(g)$  and the elements  $\xi_{\delta-\theta}$ ,  $k_{\delta}^{\pm 1}$  with the relations:

$$
[k_{\delta}^{\pm 1}, \text{everything}] = 0 \qquad k_{\alpha_i} \xi_{\delta - \theta} k_{\alpha_i}^{-1} = q^{-(\alpha_i, \theta)} \xi_{\delta - \theta} , \qquad (3.1)
$$

$$
[e_{-\alpha_i}, \xi_{\delta-\theta}] = \tau [e_{-\alpha_i}, \tilde{e}_{-\theta}], \qquad (\text{ad}_q e_{\alpha_i})^{n_{i0}} \xi_{\delta-\theta} = \tau (\text{ad}_q e_{\alpha_i})^{n_{i0}} \tilde{e}_{-\theta} \qquad (3.2)
$$

 $for n_{i0} = 1 + 2(\alpha_i, \theta)/(\alpha_i, \alpha_i)$ , and  $[[e_{\alpha_i},\xi_{\delta-\theta}]_q,\xi_{\delta-\theta}]_q\!=\!-\tau^2[[e_{\alpha_i},\tilde{e}_{-\theta}]_q,\tilde{e}_{-\theta}]_q$  $+ \tau \left[[e_{\alpha_i}, \tilde{e}_{-\theta}]_q, \xi_{\delta-\theta}]_q + \tau \left[[e_{\alpha_i}, \xi_{\delta-\theta}]_q, \tilde{e}_{-\theta} \right]$  $(3.3)$ 

 $for g \neq sl_2 \text{ and } (\alpha_i, \theta) \neq 0,$ 

$$
\begin{split}\n[[[e_{\alpha}, \xi_{\delta-\alpha}]_q, \xi_{\delta-\alpha}]_q, \xi_{\delta-\alpha}]_q &= \tau^3 [[[e_{\alpha}, \tilde{e}_{-\alpha}]_q, \tilde{e}_{-\alpha}]_q, \tilde{e}_{-\alpha}]_q \\
&\quad - \tau^2 [[[e_{\alpha}, \tilde{e}_{-\alpha}]_q, \tilde{e}_{-\alpha}]_q, \xi_{\delta-\alpha}]_q - \tau^2 [[[e_{\alpha}, \tilde{e}_{-\alpha}]_q, \xi_{\delta-\alpha}]_q, \tilde{e}_{-\alpha}]_q \\
&\quad - \tau^2 [[[e_{\alpha}, \xi_{\delta-\alpha}]_q, \tilde{e}_{-\alpha}]_q, \tilde{e}_{-\alpha}]_q + \tau [[[e_{\alpha}, \tilde{e}_{-\alpha}]_q, \xi_{\delta-\alpha}]_q, \xi_{\delta-\alpha}]_q \\
&\quad + \tau [[[e_{\alpha}, \xi_{\delta-\alpha}]_q, \tilde{e}_{-\alpha}]_q, \xi_{\delta-\alpha}]_q + \tau [[[e_{\alpha}, \xi_{\delta-\alpha}]_q, \xi_{\delta-\alpha}]_q, \tilde{e}_{-\alpha}]_q\n\end{split} \tag{3.4}
$$

*for*  $g = sl_2$ . The Hopf structure of  $D_{q\eta}(g)$  is defined by the formulas  $\Delta_{q\eta}(x)$  =  $\Delta_q(x)$ ,  $S_{q\eta}(x) = S_q(x)$  ( $x \in U_q(g)$ ) and  $\Delta_q(k_{\delta}^{\pm})$  $\delta(\bar{k}_{\delta}^{\pm 1}) = k_{\delta}^{\pm 1} \otimes k_{\delta}^{\pm 1}, \ S_q(k_{\delta}^{\pm 1}) = k_{\delta}^{\pm 1}.$  The *comultiplication and the antipode of*  $\xi_{\delta-\alpha}$  are given by

$$
\Delta_{q\eta}(\xi_{\delta-\theta}) = \xi_{\delta-\theta} \otimes 1 + k_{\delta-\theta}^{-1} \otimes \xi_{\delta-\theta}
$$
  
+
$$
a\left(\Delta_q(\tilde{e}_{-\theta}) - \tilde{e}_{-\theta} \otimes 1 - k_{\delta-\theta}^{-1} \otimes \tilde{e}_{-\theta}\right),
$$
(3.5)

$$
S_{q\eta}(\xi_{\delta-\theta}) = -k_{\delta-\theta}\xi_{\delta-\theta} + a\left(S_q(\tilde{e}_{-\theta}) + k_{\delta-\theta}\tilde{e}_{-\theta}\right) \tag{3.6}
$$

*Here*  $\tau := \eta/(q - q^{-1})$ ,  $(\text{ad}_q e_\beta)e_\gamma = [e_\beta, e_\gamma]_q$ , and the vector  $\tilde{e}_{-\theta}$  is any  $U_q(g)$  element *of the weight*  $-\theta$ *, such that*  $g \ni \lim_{q \to 1} \tilde{e}_{-\theta} \neq 0$ *.* 

<span id="page-4-0"></span>Theright-hand sides of the relations ([3.2\)](#page-3-0)-([3.6\)](#page-3-0) are nonsingular at  $q = 1$ .

**Theorem 3.1** *(i) The Drinfeldian*  $D_{q\eta}(g)$  *is a two-parameter quantization of*  $U(g[u])$ *in the direction of a classical r-matrix which is a sum of the simplest rational and trigonometric r-matrices.*

(*ii*) The Hopf algebra  $D_{q=1,\eta}(g)$  is isomorphic to the Yangian  $Y'_{\eta}(g)$  (with the addi*tional central element*  $c = h_{\delta}$ *). Moreover,*  $D_{qn=0}(g) = U_q(g[u])$ *.* 

*Remark*. Since the defining relations for  $D_{q\eta}(g)$  and  $U(\widetilde{g[u]})$  in terms of the Chevalley basis differ only in the right-hand sides of the relations([3.2\)](#page-3-0)-([3.4\)](#page-3-0), therefore the Dynkin diagram of  $g[u]$  can be also used for classification of the Drinfeldian  $D_{q\eta}(g)$ and the Yangian  $Y_n(g)$ .

An analog of Proposition [2.1](#page-3-0) is the following result.

**Proposition 3.1** *There is a one-parameter group of Hopf algebra automorphisms*  $\mathcal{T}_a$ *of*  $D_{qn}(g)$ *,*  $a \in \mathbb{C}$ *, given by* 

$$
\mathcal{T}_a(k_\delta^{\pm 1}) = k_\delta^{\pm 1} , \qquad \mathcal{T}_a(k_{\alpha_i}^{\pm 1}) = k_{\alpha_i}^{\pm 1} , \qquad \mathcal{T}_a(e_{\pm \alpha_i}) = e_{\pm \alpha_i} ,
$$
  

$$
\mathcal{T}_a(\xi_{\delta - \theta}) = \left(1 - (q - q^{-1})a\right) \xi_{\delta - \theta} + \eta \, a \, \tilde{e}_{-\theta} . \tag{3.7}
$$

In the next section we give an explicit description of the right-hand of the relations ([3.2\)](#page-3-0)-([3.6\)](#page-3-0) for the Drinfeldians  $D_{q\eta}(g)$  and the Yangians  $Y_{\eta}(g)$  of the Lie algebras g of rank 2.

## 4 Drinfeldians and Yangians over Lie algebras of rank 2

Explicit description of the Drinfeldians  $D_{q\eta}(g)$  and the Yangians  $Y_{\eta}(g)$  for the cases  $g = sl_2$  $g = sl_2$  $g = sl_2$  and  $sl_3$  were given in [[5\]](#page-11-0)-[\[7](#page-11-0)]. Here we consider the cases  $g = C_2(\simeq B_2)$  and  $G_2$ . *1. The Drinfeldian*  $D_{q\eta}(C_2)$  *and the Yangian*  $Y_{\eta}(C_2)$ . In the case of the Lie algebra  $g = C_2$  there are two positive simple roots  $\alpha$  and  $\beta$ , and the maximal positive root is  $\theta = \alpha + 2\beta$ .

As we already noted the Drinfeldian  $D_{q\eta}(g)$  of the Yangian  $Y_{\eta}(g)$  can be characterized the Dynkin diagram of the corresponding non-twisted Kac-Moody affine Lie algebra  $g^{(1)}$ . In the case  $g = C_2$  the Dynkin diagram of the corresponding affine Lie algebra  $C_2^{(1)}$  $2^{(1)}$  is presented by the picture [\[3\]](#page-11-0)

$$
\overset{\delta-\theta}{\circlearrowright} \overset{\beta}{\circlearrowright} \overset{\alpha}{\bullet}
$$

Fig.1. Dynkin diagram of the Lie algebra  $C_2^{(1)}$ 

The quantum Hopf algebra  $U_q(C_2)$  is generated by the elements  $k_\alpha^{\pm 1}$ ,  $k_\beta^{\pm 1}$ ,  $e_{\pm\alpha}$ ,  $e_{\pm\beta}$ withthe defining relations  $(2.8)-(2.13)$  $(2.8)-(2.13)$  $(2.8)-(2.13)$  $(2.8)-(2.13)$  (see also details in [\[4](#page-11-0)]). In the relations  $(3.1)$  $(3.1)$ -([3.6\)](#page-3-0) we set

$$
\tilde{e}_{-\theta} = k_{\delta - \alpha - 2\beta}^{-1} e_{-\alpha - 2\beta} , \qquad (4.1)
$$

where

$$
e_{-a-2\beta} := [e_{-\alpha-\beta}, e_{-\beta}]_q , \qquad e_{-a-\beta} := [e_{-\alpha}, e_{-\beta}]_q . \qquad (4.2)
$$

Using explicit relations for the Cartan-Weyl basis of  $U_q(C_2)$  ( $\simeq U_q(B_2)$ )) (see [\[4\]](#page-11-0)) it is not difficult to calculate the right-hand sides of the relations [\(3.2](#page-3-0))-[\(3.6](#page-3-0)). We obtain the result which is formulated as a definition of the Drinfeldian  $D_{q\eta}(C_2)$ .

**Definition 4.1** *The Drinfeldian*  $D_{q\eta}(C_2)$  *associated to*  $C_2(\simeq B_2)$  *is the Hopf algebra generated by the quantum algebra*  $U_q(C_2)$  *and the elements*  $k_{\delta}^{\pm 1}$ ,  $\xi_{\delta-\alpha-2\beta}$  *with the defining relations:*

$$
[k_{\delta}^{\pm 1}, \text{everything}] = 0 \tag{4.3}
$$

$$
k_{\alpha} \xi_{\delta - \alpha - 2\beta} k_{\alpha}^{-1} = \xi_{\delta - \alpha - 2\beta} \tag{4.4}
$$

$$
k_{\beta}\xi_{\delta-\alpha-2\beta}k_{\beta}^{-1} = q^{-(\alpha,\beta)}\xi_{\delta-\alpha-2\beta} , \qquad (4.5)
$$

$$
[e_{-\alpha}, \xi_{\delta-\alpha-2\beta}] = -\eta q^{-\frac{1}{2}(\alpha,\beta)} \left[\frac{(\alpha,\beta)}{2}\right] k_{\delta-\alpha-2\beta}^{-1} e_{-\alpha-\beta}^2 \tag{4.6}
$$

$$
[e_{-\beta}, \xi_{\delta-\alpha-2\beta}] = 0 \tag{4.7}
$$

$$
[e_{\alpha}, \xi_{\delta-\alpha-2\beta}]_q = \eta q^{\frac{3}{2}(\alpha,\beta)} [(\alpha,\beta)] \Big[ \frac{(\alpha,\beta)}{2} \Big] k_{\delta-\alpha-2\beta}^{-1} k_{\alpha+2\beta} e_{-\beta}^2 , \qquad (4.8)
$$

$$
[e_{\beta}, [e_{\beta}, [e_{\beta}, \xi_{\delta - \alpha - 2\beta}]_q]_q]_q = 0 , \qquad (4.9)
$$

$$
[[e_{\alpha}, \xi_{\delta-\alpha-2\beta}]_q, \xi_{\delta-\alpha-2\beta}]_q = 0.
$$
\n
$$
(4.10)
$$

*The Hopf structure of the Drinfeldian*  $D_{q\eta}(C_2)$  *is defined by the formulas*  $\Delta_{q\eta}(x)$  =  $\Delta_q(x)$ *,*  $S_{q\eta}(x) = S_q(x)$  ( $x \in U_q(C_2)$ ) and

$$
\Delta_{q\eta}(\xi_{\delta-\alpha-2\beta}) = \xi_{\delta-\alpha-2\beta} \otimes 1 + k_{\delta-\alpha-2\beta}^{-1} \otimes \xi_{\delta-\alpha-2\beta}
$$
  
+
$$
\eta(k_{\delta-\alpha-2\beta}^{-1} \otimes k_{\delta-\alpha-2\beta}^{-1}) \Big( e_{-\alpha-2\beta} \otimes \frac{k_{\alpha+2\beta}-k_{\delta-\alpha-2\beta}}{q-q^{-1}} - [(\alpha,\beta)]e_{-\alpha-\beta} \otimes k_{\alpha+\beta}e_{-\beta}
$$
  
+
$$
(q-q^{-1})q^{\frac{3}{2}(\alpha,\beta)} [(\alpha,\beta)] \Big[ \frac{(\alpha,\beta)}{2} \Big] e_{-\alpha} \otimes k_{\alpha}e_{-\beta}^{2} \Big), \qquad (4.11)
$$

$$
S_{q\eta}(\xi_{\delta-\alpha-2\beta}) = -k_{\delta-\alpha-2\beta}\xi_{\delta-\alpha-2\beta}
$$
  
+
$$
\eta \left(e_{-\alpha-2\beta} \frac{k_{\alpha+2\beta}k_{d-\alpha-2\beta}^{-1}}{q-q^{-1}} - \left[ (\alpha, \beta) \right] e_{-\alpha-\beta} e_{-\beta}
$$
  
-
$$
\left(q - q^{-1}\right) q^{\frac{1}{2}(\alpha,\beta)} \left[ (\alpha, \beta) \right] \left[ \frac{(\alpha,\beta)}{2} \right] e_{-\alpha} e_{-\beta}^{2} k_{\alpha+2\beta} k_{\delta-\alpha-2\beta} , \qquad (4.12)
$$

*Here we use the standard notation*  $[a] := (q^a - q^{-a})/(q - q^{-1})$ *.* 

At the limit  $q = 1$  we obtain the Yangian  $Y_{\eta}(C_2)' = D_{q=1,\eta}(C_2)$  with the central element  $h_{\delta}$ . We formulate this result as the following proposition.

**Proposition 4.1** *The Yangian*  $Y_n(C_2)$  *(as an associative algebra Hopf algebra over*  $\mathbb{E}[\eta]]$  *is generated by the algebra*  $U(C_2)$  *and the elements*  $h_\delta$ ,  $\xi_{\delta-\alpha-2\beta}$  *with the defining relations:*

$$
[h_{\delta}, \text{everything}] = 0 \tag{4.13}
$$

$$
[h_{\alpha}, \xi_{\delta - \alpha - 2\beta}] = \xi_{\delta - \alpha - 2\beta} \tag{4.14}
$$

$$
[h_{\beta}, \xi_{\delta - \alpha - 2\beta}] = (\alpha, \beta)\xi_{\delta - \alpha - 2\beta} , \qquad (4.15)
$$

$$
[e_{-\alpha}, \xi_{\delta - \alpha - 2\beta}] = -\eta \frac{1}{2} (\alpha, \beta) e_{-\alpha - \beta}^{2} , \qquad (4.16)
$$

$$
[e_{-\beta}, \xi_{\delta - \alpha - 2\beta}] = 0 \tag{4.17}
$$

$$
[e_{\alpha}, \xi_{\delta - \alpha - 2\beta}] = \eta \frac{1}{2} (\alpha, \beta)^2 e_{-\beta}^2 , \qquad (4.18)
$$

$$
[e_{\beta}, [e_{\beta}, [e_{\beta}, \xi_{\delta - \alpha - 2\beta}]]] = 0 , \qquad (4.19)
$$

$$
[[e_{\alpha}, \xi_{\delta-\alpha-2\beta}], \xi_{\delta-\alpha-2\beta}] = 0.
$$
\n(4.20)

*The nontrivial coproduct*  $\Delta_{\eta}$  *and antipode*  $S_{\eta}$  *for the element*  $\xi_{\delta-\alpha-2\beta}$  *is given by the formulas*

$$
\Delta_{\eta}(\xi_{\delta-\alpha-2\beta}) = \xi_{\delta-\alpha-2\beta} \otimes 1 + 1 \otimes \xi_{\delta-\alpha-2\beta}
$$
  
+ 
$$
\eta \left( e_{-\alpha-2\beta} \otimes \left( h_{\alpha+2\beta} - \frac{h_{\delta}}{2} \right) - (\alpha, \beta) e_{-\alpha-\beta} \otimes e_{-\beta} \right) , \qquad (4.21)
$$

$$
S_{\eta}(\xi_{\delta-\alpha-2\beta})=-\xi_{\delta-\alpha-2\beta}+\eta\Big(e_{-\alpha-2\beta}\Big(h_{\alpha+2\beta}-\frac{h_{\delta}}{2}\Big)-(\alpha,\beta)e_{-\alpha-\beta}e_{-\beta}\Big). \qquad(4.22)
$$

*1. The Drinfeldian*  $D_{q\eta}(G_2)$  *and the Yangian*  $Y_{\eta}(G_2)$ . In the case of the Lie algebra  $g = G_2$  there are two positive simple roots  $\alpha$  and  $\beta$ , and the maximal positive root is  $\theta = 2\alpha + 3\beta$ . The Dynkin diagram of the corresponding affine Lie algebra  $G_2^{(1)}$  $_2^{(1)}$  is presented by the picture [\[3\]](#page-11-0)

$$
\begin{array}{cc}\n\delta - \theta & \alpha & \beta \\
\circ & \circ \quad \bullet\n\end{array}
$$

Fig.2. Dynkin diagram of the Lie algebra  $G_2^{(1)}$ 

The quantum Hopf algebra  $U_q(G_2)$  is generated by the elements  $k_\alpha^{\pm 1}$ ,  $k_\beta^{\pm 1}$ ,  $e_{\pm\alpha}$ ,  $e_{\pm\beta}$ withthe defining relations  $(2.8)-(2.13)$  $(2.8)-(2.13)$  $(2.8)-(2.13)$  $(2.8)-(2.13)$  (see also details in [\[4](#page-11-0)]). In the relations  $(3.1)$  $(3.1)$ -([3.6\)](#page-3-0) we set

$$
\tilde{e}_{-\theta} = k_{d-2\alpha-3\beta}^{-1} e_{-2\alpha-3\beta} , \qquad (4.23)
$$

where

$$
e_{-2a-3\beta} := [e_{-\alpha-2\beta}, e_{-\alpha-\beta}]_q , \qquad e_{-a-\beta} := [e_{-\beta}, e_{-\alpha}]_q ,
$$
  

$$
e_{-a-2\beta} := [e_{-\beta}, e_{-\alpha-\beta}]_q , \qquad e_{-a-3\beta} := [e_{-\beta}, e_{-\alpha-2\beta}]_q .
$$
 (4.24)

Using explicit relations for the Cartan-Weyl basis of  $U_q(G_2)$  (see [\[4\]](#page-11-0)) we can calculate the right-hand sides of the relations([3.2\)](#page-3-0)-([3.6\)](#page-3-0). We obtain the result which is formulated as a definition of the Drinfeldian  $D_{\mathfrak{a}}(G_2)$ .

**Definition 4.2** *The Drinfeldian*  $D_{q\eta}(G_2)$  *associated to*  $G_2(\simeq B_2)$  *is the Hopf algebra generated by the quantum algebra*  $\hat{U}_q(G_2)$  *and the elements*  $k_{\delta}^{\pm 1}$ ,  $\xi_{\delta-2\alpha-3\beta}$  *with the defining relations:*

$$
[k_{\delta}^{\pm 1}, \text{everything}] = 0 \tag{4.25}
$$

$$
k_{\alpha}\xi_{\delta-2\alpha-3\beta}k_{\alpha}^{-1} = q^{-(\alpha,\beta)}\xi_{\delta-2\alpha-3\beta} , \qquad (4.26)
$$

$$
k_{\beta}\xi_{\delta-2\alpha-3\beta}k_{\beta}^{-1} = \xi_{\delta-2\alpha-3\beta} \tag{4.27}
$$

$$
[e_{-\alpha}, \xi_{\delta - 2\alpha - 3\beta}] = -\eta (q - q^{-1}) q^{-(\alpha, \beta)} bc k_{\delta - 2\alpha - 3\beta}^{-1} e_{-\alpha - \beta}^{3} , \qquad (4.28)
$$

$$
[e_{-\beta}, \xi_{\delta - 2\alpha - 3\beta}] = -\eta q^{-\frac{1}{3}(\alpha,\beta)} ab^{-1} c k_{\delta - 2\alpha - 3\beta}^{-1} e_{-\alpha - 2\beta}^2 ,
$$
 (4.29)

$$
[e_{\beta}, \xi_{\delta - 2\alpha - 3\beta}]_q = -\eta q^{\frac{1}{3}(\alpha,\beta)} ab k_{\delta - 2\alpha - 3\beta}^{-1} k_{\beta} e_{-\alpha - \beta}^2 , \qquad (4.30)
$$

$$
[e_{\alpha}, [e_{\alpha}, \xi_{\delta - 2\alpha - 3\beta}]_q]_q = \eta(q - q^{-1}) q^{-(\alpha, \beta)} abcd k_{\delta - 2\alpha - 3\beta}^{-1} k_{\alpha}^{-2} e_{-\beta}^3 , \qquad (4.31)
$$

$$
[[e_{\alpha}, \xi_{\delta-\alpha-2\beta}]_q, \xi_{\delta-\alpha-2\beta}]_q = \eta^2 q^{-(\alpha,\beta)} ab^{-1} c^2 dk_{\delta-2\alpha-3\beta}^{-2} k_{\alpha}^{-1} e_{-\alpha-2\beta}^3.
$$
 (4.32)

*The Hopf structure of the Drinfeldian*  $D_{q\eta}(G_2)$  *is defined by the formulas*  $\Delta_{q\eta}(x)$  =  $\Delta_q(x)$ *,*  $S_{q\eta}(x) = S_q(x)$   $(x \in U_q(G_2))$  and

$$
\Delta_{q\eta}(\xi_{\delta-2\alpha-3\beta}) = \xi_{\delta-2\alpha-3\beta} \otimes 1 + k_{\delta-2\alpha-3\beta}^{-1} \otimes \xi_{\delta-2\alpha-3\beta}
$$
  
\n
$$
+ \eta (k_{\delta-2\alpha-3\beta}^{-1} \otimes k_{\delta-2\alpha-3\beta}^{-1}) \Big( e_{-2\alpha-3\beta} \otimes \frac{k_{2\alpha+3\beta}-k_{\delta-2\alpha-3\beta}}{q-q^{-1}} - q^{-\frac{1}{3}(\alpha,\beta)} a e_{-\alpha-2\beta} \otimes k_{\alpha+2\beta} e_{-\alpha-\beta} + q^{\frac{4}{3}(\alpha,\beta)} (1 - (q - q^{-1})b) e_{-\alpha-3\beta} \otimes k_{\alpha+3\beta} e_{-\alpha})
$$
  
\n
$$
+ (q - q^{-1}) a \Big( q^{(\alpha,\beta)} a e_{-\beta} e_{-\alpha-2\beta} \otimes k_{\alpha+2\beta} e_{-\alpha} + q^{\frac{1}{3}(\alpha,\beta)} b e_{-\beta} \otimes k_{\beta} e_{-\alpha-\beta}^2 \Big)
$$
  
\n
$$
- (q - q^{-1})^2 q^{\frac{7}{3}(\alpha,\beta)} a^2 b e_{-\beta}^2 \otimes k_{\beta}^2 e_{-\alpha-\beta} e_{-\alpha}
$$
  
\n
$$
+ (q - q^{-1})^3 q^{4(\alpha,\beta)} a^2 b c e_{-\beta}^3 \otimes k_{\beta}^3 e_{-\alpha}^2 \Big), \qquad (4.33)
$$

$$
S_{q\eta}(\xi_{\delta-2\alpha-3\beta}) = -k_{\delta-2\alpha-3\beta}\xi_{\delta-2\alpha-3\beta}
$$
  
+
$$
\eta \left(e_{-2\alpha-3\beta} \frac{k_{2\alpha+3\beta}k_{\delta-2\alpha-3\beta}^{-1}}{q-q^{-1}} + q^{\frac{2}{3}(\alpha,\beta)}a e_{-\alpha-3\beta}e_{-\alpha} + q^{-\frac{1}{3}(\alpha,\beta)}a e_{-\alpha-2\beta}e_{-\alpha-\beta} + (q-q^{-1})ab\left(q^{\frac{2}{3}(\alpha,\beta)}e_{-\alpha-3\beta}e_{-\alpha} - q^{-\frac{1}{3}(\alpha,\beta)}e_{-\beta}e_{-\alpha-\beta}^2\right) + (q-q^{-1})^2a^2b\left(q^{\frac{1}{3}(\alpha,\beta)}e_{-\beta}e_{-\alpha-2\beta}e_{-\alpha} - q^{-\frac{1}{3}(\alpha,\beta)}e_{-\beta}^2e_{-\alpha-\beta}e_{-\alpha}\right) - (q-q^{-1})^3a^2bc\,e_{-\beta}^3e_{-\alpha}^2\right)k_{2\alpha+3\beta}^{-1}k_{\delta-2\alpha-3\beta} . \tag{4.34}
$$

Here we use the notations

$$
a := \frac{q^{(\alpha,\beta)} - q^{-(\alpha,\beta)}}{q - q^{-1}}, \qquad d := \frac{q^{2(\alpha,\beta)} - q^{-2(\alpha,\beta)}}{q - q^{-1}},
$$
  

$$
b := \frac{q^{\frac{2}{3}(\alpha,\beta)} - q^{-\frac{2}{3}(\alpha,\beta)}}{q - q^{-1}}, \qquad c := \frac{q^{\frac{1}{3}(\alpha,\beta)} - q^{-\frac{1}{3}(\alpha,\beta)}}{q - q^{-1}}, \qquad (4.35)
$$

At the limit  $q = 1$  we obtain the Yangian  $Y_{\eta}(G_2)' = D_{q=1,\eta}(G_2)$  with the central element  $h_{\delta}$ . We formulate this result as the following proposition.

**Proposition 4.2** *The Yangian*  $Y_{\eta}(G_2)$  *(as an associative algebra Hopf algebra over*  $\mathbb{E}[\![\hbar,\eta]\!]$  *is generated by the algebra*  $U(C_2)$  *and the elements*  $h_\delta$ ,  $\xi_{\delta-\alpha-2\beta}$  *with the defining relations:*

$$
[h_{\delta}, \text{everything}] = 0 \tag{4.36}
$$

$$
[h_{\alpha}, \xi_{\delta - 2\alpha - 3\beta}] = (\alpha, \beta)\xi_{\delta - 2\alpha - 3\beta} , \qquad (4.37)
$$

$$
[h_{\beta}, \xi_{\delta - 2\alpha - 3\beta}] = \xi_{\delta - 2\alpha - 3\beta} \tag{4.38}
$$

$$
[e_{-\alpha}, \xi_{\delta - 2\alpha - 3\beta}] = 0 , \qquad (4.39)
$$

$$
[e_{-\beta}, \xi_{\delta - 2\alpha - 3\beta}] = -\eta \frac{1}{2} (\alpha, \beta) e_{-\alpha - 2\beta}^2 ,
$$
 (4.40)

$$
[e_{\beta}, \xi_{\delta - 2\alpha - 3\beta}] = \eta \frac{2}{3} (\alpha, \beta)^2 e_{-\alpha - \beta}^2 , \qquad (4.41)
$$

$$
[e_{\alpha}, [e_{\alpha}, \xi_{\delta - \alpha - 2\beta}]] = 0 , \qquad (4.42)
$$

$$
[[e_{\alpha}, \xi_{\delta - 2\alpha - 3\beta}], \xi_{\delta - 2\alpha - 3\beta}] = \eta^2 \frac{1}{3} (\alpha, \beta)^3 e_{-\alpha - 2\beta}^3 , \qquad (4.43)
$$

*The nontrivial coproduct*  $\Delta_{\eta}$  *and antipode*  $S_{\eta}$  *for the element*  $\xi_{\delta-2\alpha-3\beta}$  *is given by the formulas*

$$
\Delta_{\eta}(\xi_{\delta-2\alpha-3\beta}) = \xi_{\delta-2\alpha-3\beta} \otimes 1 + 1 \otimes \xi_{\delta-2\alpha-3\beta} \tag{4.44}
$$

$$
+ \eta \left( e_{-\alpha-2\beta} \otimes \left( h_{2\alpha+3\beta} - \frac{h_{\delta}}{2} \right) + (\alpha, \beta)(e_{-\alpha-3\beta} \otimes e_{-\alpha} - e_{-\alpha-2\beta} \otimes e_{-\alpha-\beta}) \right)
$$
  

$$
S_{\eta}(\xi_{\delta-2\alpha-3\beta}) = -\xi_{\delta-2\alpha-3\beta}
$$
  

$$
+ \eta \left( e_{-2\alpha-3\beta} \left( h_{2\alpha+3\beta} - \frac{h_{\delta}}{2} \right) + (\alpha, \beta)(e_{-\alpha-3\beta}e_{-\alpha} - e_{-\alpha-2\beta}e_{-\alpha-\beta}) \right).
$$
  
(4.45)

#### 5 Super Drinfeldian and super Yangian

It is obvious that results of the Sections 3 are easy extended to the supercase. In the supercase, i.e. when  $g$  is a simple finite-dimensional contragredient Lie superalgebra all the commutators and the q-commutators of the relations [\(3.2](#page-3-0))-[\(3.4](#page-3-0)) are replaced by the supercommutators and the q-supercommutators. For example, the q-commutator ([2.10\)](#page-2-0) is replace by the q-supercommutator

$$
[e_{\beta}, e_{\gamma}]_q = e_{\beta} e_{\gamma} - (-1)^{\vartheta(\beta)\vartheta(\gamma)} q^{(\beta, \gamma)} e_{\gamma} e_{\beta} , \qquad (5.1)
$$

where  $\vartheta(\cdot)$  is a standard parity function  $(\vartheta(\gamma) = 0$  for any even root  $\gamma$ , and  $\vartheta(\gamma) = 1$ for any odd root  $\gamma$ ). Moreover we have to add some additional Serre relations if they exist (see[[4, 8](#page-11-0)], for example).

Let us construct the super Drinfeldian and the super Yangian of the superalgebra osp(1|2) ( $\simeq$  B(0,1)) as an example in an explicit form. The Dynkin diagram of the affine superalgebra  $\widehat{osp}(1|2)$  ( $\simeq B(0,1)^{(1)}$ ) is represented by the picture

$$
\overset{\delta-2\alpha}{\smile} \overset{\alpha}{\Longrightarrow}
$$

Fig.3. Dynkin diagram of the Lie superalgebra  $\widehat{osp}(1|2)$  ( $\simeq B(0,1)^{(1)}$ ).

where  $\alpha$  is the odd root  $(\vartheta(\alpha) = 1)$  and  $\delta - 2\alpha$  is even one  $(\vartheta(\delta - 2\alpha) = 0)$ .

The quantum Hopf algebra  $U_q(osp(1|2))$  is generated by the elements  $k^{\pm 1}_\alpha$ , and  $e_{\pm\alpha}$  with the defining relations

$$
k_{\alpha}e_{\pm\alpha}k_{\alpha}^{-1} = q^{\pm(\alpha,\alpha)}e_{\pm\alpha} , \qquad [e_{\alpha}, e_{-\alpha}] = \frac{k_{\alpha} - k_{\alpha}^{-1}}{q - q^{-1}} , \qquad (5.2)
$$

where the brackets  $[\cdot, \cdot]$  is the fermion commutator. The Hopf structure of  $U_q(osp(1|2))$ isgiven by the formulas  $(2.11)-(2.13)$  $(2.11)-(2.13)$  (see details in [[4](#page-11-0)]).

Inthe relations  $(3.1)-(3.6)$  $(3.1)-(3.6)$  $(3.1)-(3.6)$  $(3.1)-(3.6)$  we set

$$
\tilde{e}_{-\theta} = e_{-\alpha}^2 \tag{5.3}
$$

Aftercalculations of the right-hand sides of the relations  $(3.2)-(3.6)$  $(3.2)-(3.6)$  $(3.2)-(3.6)$  $(3.2)-(3.6)$  $(3.2)-(3.6)$  we obtain the result which is formulated as a definition of the Drinfeldian  $D_{q\eta}(osp(1|2)).$ 

**Definition 5.1** *The Drinfeldian*  $D_{q\eta}(osp(1|2))$  *associated to osp*(1|2) *is the Hopf algebra generated by the quantum algebra*  $U_q(osp(1|2))$  *and the elements and*  $k_0^{\pm 1}$ ,  $\xi_{\delta-2\alpha}$ *with the defining relations:*

$$
[k_{\delta}^{\pm}, \text{everything}] = 0 \tag{5.4}
$$

$$
k_{\alpha}\xi_{\delta-2\alpha}k_{\alpha}^{-1} = q_{\alpha}^{-2}\xi_{\delta-2\alpha} , \qquad (5.5)
$$

$$
[e_{-\alpha}, \xi_{\delta - 2\alpha}] = 0 \tag{5.6}
$$

$$
[e_{\alpha}, [e_{\alpha}, [e_{\alpha}, [e_{\alpha}, [e_{\alpha}, \xi_{\delta - 2\alpha}]_q]_q]_q]_q]_q = 0 ,
$$
\n(5.7)

$$
\begin{split}\n\left[ [e_{\alpha}, \xi_{\delta-2\alpha}]_q, \xi_{\delta-2\alpha} \right]_q &= \eta^2 [(\alpha, \alpha)] \left( [\alpha, \alpha] \right) e_{-\alpha}^4 e_{\alpha} + \left[ \frac{(\alpha, \alpha)}{2} \right] \left[ h_{\alpha} + \frac{7}{2} (\alpha, \alpha) \right] e_{-\alpha}^3 \\
-\eta (q - q^{-1}) \left( [(\alpha, \alpha)]^2 e_{-\alpha}^2 \{ e_{\alpha}, \xi_{\delta-2\alpha} \} - \left[ \frac{(\alpha, \alpha)}{2} \right] \left[ 2(\alpha, \alpha) \right] \left[ h_{\alpha} + \frac{5}{2} (\alpha, \alpha) \right] e_{-\alpha} \xi_{\delta-2\alpha} \right) .\n\end{split} \tag{5.8}
$$

*The Hopf structure of the Drinfeldian*  $D_{q\eta}(osp(1|2))$  *is defined by the formulas*  $\Delta_{q\eta}(x)$  =  $Δ<sub>q</sub>(x)$ *,*  $S<sub>qη</sub>(x) = S<sub>q</sub>(x)$  (x ∈  $U<sub>q</sub>(osp(1|2))$ *)* and

$$
\Delta_{q\eta}(\xi_{\delta-2\alpha}) = \xi_{\delta-2\alpha} \otimes 1 + k_{\delta-2\alpha}^{-1} \otimes \xi_{\delta-2\alpha}
$$
  
+
$$
\eta \left( e_{-\alpha}^2 \otimes \frac{k_{\alpha}^2 - 1}{q - q^{-1}} - \frac{k_{\delta-2\alpha}^{-1} - 1}{q - q^{-1}} \otimes e_{-\alpha}^2 - q_{\alpha}^{\frac{1}{2}} \left[ \frac{(\alpha, \alpha)}{2} \right] e_{-\alpha} \otimes k_{\alpha} e_{-\alpha} \right) , \qquad (5.9)
$$

$$
S_{q\eta}(\xi_{\delta-2\alpha}) = -k_{\delta-2\alpha}\xi_{\delta-2\alpha} + \eta q_{\alpha} \frac{q_{\alpha}^3 k_{\delta} - 1}{q - q^{-1}} e_{-\alpha}^2 k_{\alpha}^{-2} , \qquad (5.10)
$$

$$
\varepsilon_{q\eta}(\xi_{\delta-2\alpha}) = 0 , \qquad \varepsilon_{q\eta}(k_{\delta}) = \varepsilon_{q\eta}(1) = 1 . \tag{5.11}
$$

*Here*  $q_{\alpha} := q^{(\alpha,\alpha)}$ ,  $\{x, y\} := xy + yx$ , and  $[a] := (q^a - q^{-a})/(q - q^{-1})$ .

At the limit  $q = 1$  we obtain the super Yangian  $Y'_{\eta}(osp(1|2)) = D_{q=1,\eta}(osp(1|2))$ with the defining relations:

$$
[h_{\delta}, \text{everything}] = 0 \;, \qquad [e_{\alpha}, e_{-\alpha}] = h_{\alpha} \;, \tag{5.12}
$$

$$
[h_{\alpha}, e_{\pm \alpha}] = \pm (\alpha, \alpha) e_{\pm \alpha} , \qquad [h_{\alpha}, \xi_{\delta - 2\alpha}] = -2(\alpha, \alpha) \xi_{\delta - 2\alpha} , \qquad (5.13)
$$

$$
[e_{-\alpha}, \xi_{\delta - 2\alpha}] = 0 , \t [e_{\alpha}, [e_{\alpha}, [e_{\alpha}, [e_{\alpha}, \xi_{\delta - 2\alpha}]]]] = 0 , \t (5.14)
$$

$$
[[e_{\alpha}, \xi_{\delta-2\alpha}], \xi_{\delta-2\alpha}] = \eta^2 \frac{(\alpha, \alpha)^2}{2} \left( 2e_{-\alpha}^4 e_{\alpha} + \left( h_{\alpha} + \frac{7}{2}(\alpha, \alpha) \right) e_{-\alpha}^3 \right)
$$
(5.15)

with the non-trivial comultiplication  $\Delta_{\eta}$  and the antipode  $S_{\eta}$  for the affine root vector  $\xi_{\delta-2\alpha}$  given by

$$
\Delta_{\eta}(\xi_{\delta-2\alpha}) = \xi_{\delta-2\alpha} \otimes 1 + 1 \otimes \xi_{\delta-2\alpha} \n+ \eta \left( e_{-\alpha}^2 \otimes h_{\alpha} + \frac{1}{2} h_{\delta-2\alpha} \otimes e_{-\alpha}^2 - \frac{(\alpha, \alpha)}{2} e_{-\alpha} \otimes e_{-\alpha} \right) ,
$$
\n(5.16)

$$
S_{\eta}(\xi_{\delta-2\alpha}) = -\xi_{\delta-2\alpha} + \frac{\eta}{2} (h_{\delta} + 3(\alpha, \alpha)) e_{-\alpha}^{2}.
$$
 (5.17)

The Proposition [3.1](#page-4-0) is also valid for the super Drinfeldians and the super Yangians.

### <span id="page-11-0"></span>Acknowledgments

The author is thankful to Arnold Sommerfeld Institute for Mathematical Physics, Technical University of Clausthal, and the Organizing Committee of the International Workshop "Lie Theory and Its Applications in Physics II", H.-D. Doebner, V.K. Dobrev and J. Hilgert, for the support of his visit on this Workshop. This work was supported by the Russian Foundation for Fundamental Research, grant No. 96-01- 01421.

### References

- [1] V.G. Drinfeld, Hopf algebras and quantum Yang-Baxter equation, *Soviet Math. Dokl.,* 283 (1985) 1060-1064.
- [2] V.G. Drinfeld, Quantum groups, *Proc. ICM-86 (Berkeley USA) Amer. Math. Soc. Providence, RI* 1 (1987) 798-820.
- [3] V.G. Kac, Infinite demensional Lie algebras, Cambridge University Press, 1985.
- [4] S.M. Khoroshkin and V.N. Tolstoy, Universal R-matrix for quantized (super) algebras, *Commun. Math. Phys.* 141 (1991) 599-614.
- [5] V.N. Tolstoy, q-Deformed Yangian Y (sl2), *In; "Lie Theory and Its Applications in Physics", (Proc. of the Inter. Workshop; Clausthal, Germany, 1995; eds. H.-D. Doebner, V.K. Dobrev and J. Hilgert), World Scientific Publ., Singapore* (1996), 179-186
- [6] V.N. Tolstoy and S.M. Khoroshkin, Rational-trigonometric quantum groups, *In: "Quantum Group Symposium at Group 21" (Proc. of the XXI Inter. Colloquium on Group Theoretical Methods in Physics; Goslar, Germany, 1996; eds. H.-D. Doebner and V.K. Dobrev) Heron Press, Sofia; ISBN 954-590-027-5* (1997), 227- 234.
- [7] V.N. Tolstoy, Connection between Yangians and Quantum Affine Algebras, *In: New Symmetry in the Theories of Fundamental Interactions (Proc. of the IX Max Born Symposium; Karpacz, Poland, 1996; eds. J. Lukierski, M. Mozrzymas) Polish Scientific Publishers.- Warsaw* (1997), 99-117.
- [8] H. Yamane, A Serre type theorem for affine Lie superalgebras and their quantized universal enveloping superalgebras, *Proc. Japan. Acad.,* 70 (1994) 31.