

VANISHING THEOREMS AND UNIVERSAL COVERINGS OF PROJECTIVE VARIETIES

FEDOR BOGOMOLOV

Courant Institute for Mathematical Sciences

ABSTRACT. This article contains a new argument which proves vanishing of the first cohomology for negative vector bundles over a complex projective variety if the rank of the bundle is smaller than the dimension of the base. Similar argument is applied to the construction of holomorphic functions on the universal covering of the complex projective variety .

1. INTRODUCTION

There are many flavors of vanishing theorems for negative or semi-negative line bundles (see [2] for a survey of the existing results). In contrast the vanishing of the cohomology of negative vector bundles is not so well understood. In this paper I propose a new approach leading to a simple proof of the vanishing of the first cohomology of negative vector bundles whose rank is smaller than the dimension of the base. This approach also provides some insight into the problem of quasi convexity of the universal coverings of projective varieties.

For the results on Stein spaces used throughout the paper the reader may wish to consult A.L.Onishchik's survey [5]. Similarly to [5] the present article emphasizes mainly the holomorphic and cohomological features of Stein spaces and keeps the references to differential geometry and plurisubharmonic functions to a minimum. It is clear that most of the results here can be strengthened by applying differential geometric tools or by using some of the more elaborate properties of the vector bundles in question.

The main idea of this article stems from Lemma 2.3 in my joint work [1] with L.Katzarkov on the fundamental groups of projective and symplectic manifolds.

I am grateful to L.Katzarkov and T.Pantev for useful comments.

2. A VANISHING THEOREM FOR NEGATIVE VECTOR BUNDLES

Let X be a normal irreducible complex projective variety.

For a vector bundle $E \rightarrow X$ denote by $t(E) := \text{Spec}(S^\bullet E^\vee) \rightarrow X$ its total space scheme. Similarly let $\pi : \mathbb{P}(E) := \mathbf{Proj}(S^\bullet E^\vee) \rightarrow X$ denote the projectivization of E and let $\mathcal{O}_{\mathbb{P}(E)}(-1) \subset \pi^*E$ denote the tautological line bundle.

Partially supported by NSF grant DMS-9801591.

Recall (see e.g. [3]) that a vector bundle F is called *negative* if the regular functions on $t(F)$ separate points and tangent directions outside of the zero section. In other words F is negative if the natural affinization map

$$a : t(F) \rightarrow \text{Spec}(\mathbb{C}[t(F)]) = \text{Spec}(H^0(X, S^\bullet F^\vee))$$

contracts precisely the zero section.

Our main result is the following theorem

Theorem 1. *If F is a negative vector bundle on X with $\text{rk } F < \dim X$, then $H^1(X, F) = 0$.*

Proof. Put $d := \dim X$ and $r := \text{rk } F$. Assume that $H^1(X, F) \neq 0$. Then there is a cohomology class $s \in H^1(X, F)$, $s \neq 0$. Let \tilde{F} be the corresponding extension of F by \mathcal{O}_X . For a nonzero constant section $m \in \mathcal{O}_X$ the preimage of m in \tilde{F} is an affine bundle over X modeled on F . We will denote it by F_s since it does not depend on the choice of m . Consider the closure $\overline{t(F)}$ of $t(F)$ and $\overline{F_s}$ of F_s respectively inside the projective fibration $\mathbb{P}(\tilde{F} \oplus \mathcal{O}_X)$. Note that both divisors at infinity $\overline{t(F)} \setminus t(F)$ and $\overline{F_s} \setminus F_s$ are isomorphic to $\mathbb{P}(F)$ with the same positive normal bundle $N \cong \mathcal{O}_{\mathbb{P}(F)}(1)$.

By Grauert's criterion for ampleness [3] the line bundles $\mathcal{O}_{F_s}(\mathbb{P}(F))$ and $\mathcal{O}_{t(F)}(\mathbb{P}(F))$ are both ample. Thus the linear system $\mathcal{O}_{F_s}(\mathbb{P}(F))$ defines a projection $r : F_s \rightarrow \text{Spec}(\mathbb{C}[F_s]) =: F_s^r$ which is a proper morphism. Since by construction r is birational there are at most finitely many points $q_i \in F_s^r$ which have as preimages some positive dimensional compact subvarieties $X_i \subset F_s$.

Since each X_i 's is proper the natural map $p|_{X_i} : X_i \rightarrow X$, where p is the affine projection $p : F_s \rightarrow X$, will be finite on its image. In the next lemma we prove that $\dim_{\mathbb{C}} X_i < \dim_{\mathbb{C}} X = d$ for any q_i , which leads to a contradiction. Indeed $F_s^r = F_s / (\prod_i X_i)$ as a topological space (in the classical topology) and so for the reduced singular homology of F_s with say \mathbb{Q} -coefficients we have $\tilde{H}_i(F_s^r, \mathbb{Q}) = H_i((F_s, \prod_i X_i), \mathbb{Q})$. Now the long exact homology sequence of the pair $(F_s, \prod_i X_i)$ together with the fact that $\prod_i X_i$ is compact of complex dimension strictly less than d gives that $H_{2d}(F_s, \mathbb{Q}) \cong \tilde{H}_{2d}(F_s^r, \mathbb{Q}) = H_{2d}(F_s^r, \mathbb{Q})$. On the other hand since F_s^r is an affine variety we know that F_s^r has the homotopy type of a simplicial complex of real dimension which does not exceed $\dim_{\mathbb{C}} F_s^r = \dim_{\mathbb{C}} t(F) = \dim_{\mathbb{C}} X + \text{rk } F = d + r$. This yields $H_{2d}(F_s^r, \mathbb{Q}) = 0$ due to the hypothesis $r < d$. This however is impossible since F_s is an affine bundle over X and hence homotopically equivalent to X which in turn implies that $H_{2d}(F_s, \mathbb{Q})$ contains a non-zero homology class - namely the fundamental class of X . This gives us the desired contradiction and finishes the proof of the theorem modulo the fact that $\dim_{\mathbb{C}} X_i < \dim_{\mathbb{C}} X$ for all i .

We will derive this fact from the following general lemma.

Lemma 1. *Let X and F be as in the statement of Theorem 1 and let $y : Y \rightarrow X$ be a finite morphism. Then for any $0 \neq s \in H^1(X, F)$ we have $y^*s \neq 0$.*

Proof. Assume first that X is a smooth compact curve. By taking the Galois closure of $y : Y \rightarrow X$ and normalizing one gets a smooth finite Galois covering $y_G : X_G \rightarrow X$ with a finite Galois group G which factors a $y_G = y \circ y'$ with $y' : X_G \rightarrow Y$. By assumption $y_G^*s = 0$. Thus $y_G^*\tilde{F} \cong F \oplus \mathcal{O}_X$. Moreover since the property of being negative is preserved by finite base change Proposition 4.3, [3] we

have that y_G^*F is a negative vector bundle on X_G . Consequently $H^0(X_G, y_G^*F) = 0$ and so $H^0(X_G, y_G^*\tilde{F})$ injects into $H^0(X_G, \mathcal{O}_{X_G}) \cong \mathbb{C}$. Therefore

$$H^0(X_G, y_G^*\tilde{F}) = H^0(X_G, \mathcal{O}_{X_G}) \cong \mathbb{C}$$

and so G acts trivially on the one dimensional space $H^0(X_G, y_G^*\tilde{F})$. This implies that $(y_G^*y_G^*\tilde{F})^G = (\tilde{F} \otimes y_{G^*}\mathcal{O}_{X_G})^G = \tilde{F}$ has a nowhere vanishing section. Hence the exact sequence $0 \rightarrow F \rightarrow \tilde{F} \rightarrow \mathcal{O}_X \rightarrow 0$ splits and so $s = 0$ - a contradiction.

To finish the proof of the lemma it remains just to observe that for any X and any vector bundle F we can find a sufficiently ample divisor $D \subset X$ so that the cohomology group $H^1(X, F)$ injects into $H^1(D, F|_D)$. Since negativity is preserved by restrictions to subvarieties we can then find a smooth curve C (cut out by finitely many general hyperplane sections) for which the restriction of s on C is nonzero.

The lemma is proven. \square

We can now finish the proof of the theorem. Indeed since X_i are contracted by the affinization map r we have that $p^*s|_{X_i} = 0$ for all i . Moreover since by construction $p|_{X_i} : X_i \rightarrow X$ is finite onto its image Lemma 1 implies that $p|_{X_i}$ cannot be surjective. Thus $\dim X_i < \dim X$ which concludes the proof of the theorem. \square

Remark 1. It is pretty easy to construct many negative bundles of rank $\geq \dim X$ with nontrivial first cohomology. One possible construction goes as follows. Let L be a very ample line bundle on X which gives an embedding $X \subset \mathbb{P}^n$. There is a surjective map $h : \mathcal{O}_X^{\oplus n+1} \rightarrow L$ which defines a rank n subbundle $\ker h = F \subset \mathcal{O}_X^{\oplus n+1}$. The vector bundle $F \otimes L^{-1}$ is a negative bundle with $H^1(X, F \otimes L^{-1}) \neq 0$. Namely there is a nontrivial element $s \in H^1(X, F \otimes L^{-1}) \neq 0$ which corresponds to the extension $0 \rightarrow F \rightarrow \mathcal{O}_X^{\oplus n+1} \rightarrow L \rightarrow 0$. To see that the latter is non-split one only needs to observe that from the Euler sequence we have an isomorphism $F \otimes L^{-1} \cong \Omega_{\mathbb{P}^n|X}^1$ and so the extension class s is just the first Chern class of L .

Remark 2. In the process of proving Theorem 1 we showed that for a negative vector bundle F of arbitrary rank and any element $s \in H^1(X, F)$ the union of all subvarieties $Y \subset X$ for which $s|_Y = 0$ is actually an algebraic subvariety of X .

This condition is somewhat reminiscent of the Lang conjecture and it seems likely there is some deeper relation between them. Because of this analogy we will formulate it in a separate lemma

Lemma 2. *Let F be a negative bundle on a projective variety X and let $s \in H^1(X, F)$. Then there exists a finite set of projective schemes $\{X_i\}_{i \in I}$, $\dim X_i < \dim X$ and affine morphisms $f_i : X_i \rightarrow X$ with the following property. Let Y be an irreducible projective variety and let $f : Y \rightarrow X$ be a map with $f^*s = 0$ s.t. $f(Y)$ is not a point. Then there exists a $i \in I$ f factors as $f = f_i \circ g$ where $g : Y \rightarrow X_i$.*

Proof. Indeed in the notations of the proof of Theorem 1 we have that $f^*\tilde{F}$ splits into a direct sum $f^*F \oplus \mathcal{O}_Y$ and therefore f^*F_s is trivial as an affine bundle, i.e. $F_s \times_X Y$ contains Y as a closed subvariety. Consider the projection $p_1 : F_s \times_X Y \rightarrow F_s$. As we saw in the proof of Theorem 1 the affinization morphism of F_s contracts only a finite set of proper subvarieties $X_i \in F_s$. Hence $p_1(Y) = X_i$ for some i and therefore $f = f_i \circ p_1$ which proves the lemma. \square

Remark 3. As the proof shows the test variety Y in the above lemma can be any connected variety without non constant holomorphic functions.

Remark 4. If $E \rightarrow X$ is a positive vector bundle and $0 \neq s \in H^1(X, E)$, then we can always find a variety Y with a surjective finite map $f : Y \rightarrow X$ for which $f^*s = 0$. Indeed in this case the divisor $\mathbb{P}(E) \subset \overline{E}_s$ can be contracted and so by cutting down $\overline{E}_s/\mathbb{P}(E)$ by a sequence of hyperplane sections we will eventually get a subvariety $Y \subset E_s$ which has dimension equal to $\dim X$. Since the natural projection $p : E_s \rightarrow X$ is an affine morphism it follows that $p|_Y : Y \rightarrow X$ will be a finite map and so we can take $f = p|_Y$.

3. INFINITE COVERS OF PROJECTIVE VARIETIES

Now we can apply the above results to the study of infinite covering spaces of complex projective varieties.

Theorem 2. *Let F be a negative bundle over a complex projective variety X and let $0 \neq s \in H^1(X, F)$ be a nontrivial cocycle. Let Y be a complex space with a map $f : Y \rightarrow X$ which is locally finite and locally compact. Assume that $f^*s = 0$. Then there is a proper algebraic subset $Z \in X$ such that the holomorphic functions on Y separate points on $Y \setminus f^{-1}(Z)$.*

Proof. Consider the variety F_s as in the proof of Theorem 1. Holomorphic functions (even regular functions) separate points of F_s modulo a finite subset of algebraic subvarieties X_i which project finite to one onto proper subvarieties in X . Denote by Z the union of the images of all X_i in X . Consider the manifold $F_{f^*s} = F_s \times_X Y$ which is an affine fibration $p_Y : F_{f^*s} \rightarrow Y$ over Y .

The map f induces a locally finite and locally compact map $f' : F_{f^*s} \rightarrow F_s$. Thus the holomorphic functions on F_s locally separate points on F_{f^*s} outside of $f^{-1}(Z)$. Since $f^*s = 0$ the corresponding extension of F splits over Y . Thus there is a section $r : Y \rightarrow F_{f^*s}$ of f' and so the restriction of the holomorphic functions F_{f^*s} to $r(Y)$ separate points outside of $r(f^{-1}Z)$. This finishes the proof of the lemma. \square

There are some immediate corollaries of this theorem.

Corollary 1. *Let X, F and s be as in Theorem 2. Let $f : \tilde{X} \rightarrow X$ be the universal cover of X and assume that $f^*s = 0$. Then the holomorphic functions on \tilde{X} separate points outside a preimage of an algebraic subset $Z \in X$.*

Proof. Clear. \square

Corollary 2. *Let X, F and s be as in Theorem 2. Assume furthermore that F_s is an affine variety. Let $f : Y \rightarrow X$ be any infinite unramified covering s.t. $f^*s = 0$. Then Y is Stein.*

Proof. Since any non-ramified covering of a Stein space is Stein [5] the assumption that F_s is affine yields that $F_s \times_X Y$ is Stein. On the other in the proof of Theorem 2 we saw that $Y \subset F_s \times_X Y$ is a closed analytic subset and so Y is Stein. \square

Remark 5. The condition that F_s is an affine variety can be easily fulfilled in examples. For instance if X, F and s are as in Remark 1 the variety F_s is affine since by construction it is a closed subvariety in the affine variety

$$\mathbb{P}^n \times \mathbb{P}^{n\vee} \setminus \{(x, h) \in \mathbb{P}^n \times \mathbb{P}^{n\vee} | x \in h\}.$$

Corollary 2 suggests that the result of Theorem 2 may also be applicable to orbicoverings of X .

Let us first describe precisely the notion of orbicovering in the case of a complex variety.

Let X be a complex variety and $S \subset X$ be a proper analytic subset. Consider for any point $q \in S$ the local fundamental group $\pi_q = \pi_1(U(q) \setminus S)$ where $U(q)$ is a small ball in X centered at q . Let $L \subset \pi_1(X \setminus S)$ be a subgroup with the property that $L \cap \pi_q$ is of finite index in π_q for all $q \in S$. Then the nonramified covering of $X \setminus S$ corresponding to L can be naturally completed into a normal complex variety Y_L with a locally finite and locally compact surjective map $f_L : Y_L \rightarrow X$. The map $f_L : Y_L \rightarrow X$ is called an *orbicovering* of X with a ramification set S . Now we have the following

Corollary 3. *Let X , F and s be as in Theorem 2. Assume furthermore that F_s is an affine variety. Let $f : Y \rightarrow X$ be any orbicovering s.t. $f^*s = 0$. Then Y is Stein.*

Proof. Since every orbicovering of a Stein space is also Stein (see Theorem 4.6 of [5]) the proof is exactly the same as the proof of Corollary 2. \square

Remark 6. The prototype of Corollary 3 is Lemma 2.3 in our joint work [1] with L.Katzarkov.

In view of the previous results I would like to formulate a conjecture which, I believe, should be the correct substitute of the Shafarevich holomorphic convexity conjecture. First we need the following definition.

Definition. Let X be a smooth projective variety and let $f : Y \rightarrow X$ be a non-ramified covering of X . We will say that the covering satisfies the property (N) (after M.Nori, T.Napier, R.Narasimhan) if there exists a proper algebraic subvariety $Z \subset X$ such that there exists normal complex space M satisfying

- (a) The holomorphic functions on M separate points.
- (b) There exists a proper map with connected fibers $g : Y \setminus f^{-1}(Z) \rightarrow M$
- (c) Holomorphic functions on Y separate the fibers of g after restriction to $Y \setminus f^{-1}(Z)$

The conjecture now reads

Conjecture 1. *Let X be a smooth compact projective variety. Then the universal cover of X satisfies the property (N).*

Remark 7. We have shown above that if there exists a negative vector bundle $F \rightarrow X$ with a cohomology class $s \in H^1(X, F)$ which becomes trivial on \tilde{X} , then the conjecture is true and moreover $\dim M = \dim X$. Thus the identification on \tilde{X} of the two "infinitesimally" close bundles $F \oplus \mathcal{O}$ and \tilde{F} implies the property (N) for the universal covering.

It is worth to point out that in many of the cases for which the Shafarevich conjecture is known the proof relies on the comparison of two vector bundles on X which become equal when they are pulled back to \tilde{X} [3]. For example the theorem of M.Gromov uses the fact that some positive line bundle $E \rightarrow X$ becomes trivial when pulled back to the universal cover. The theorem of L.Katzarkov [4] establishes

the holomorphic convexity of \tilde{X} for projective surface X under the assumption of the existence of an almost faithful linear representation of $\pi_1(X)$. In this case all the bundles on X corresponding to the representations of the fundamental group of the same dimension are becoming equal on \tilde{X} .

Remark 8. In the case of surfaces the space \tilde{X} can be obtained as a union of two rather simple Stein manifolds with Stein intersection. This implies that the structure of the space of the moduli space of vector bundles on \tilde{X} in this case is somewhat similar to the structure of the moduli space of vector bundles on a curve. Namely any bundle of rank greater than 2 has a complete flag of subbundles, thus reducing the K-group $K_0(\tilde{X})$ to $\text{Pic}(X) \times \mathbb{Z}$. In particular one expects that many different bundles on X coincide after lifting to \tilde{X} .

REFERENCES

- [1] **F.Bogomolov, L.Katzarkov**, *Symplectic four-manifolds and projective surfaces.*, Topology and its applications **20** (1997), 1-31.
- [2] **H.Esnault, E.Viehweg**, *Lectures on vanishing theorems*, DMV 20, Birkhauser, Boston, 1992.
- [3] **R.Hartshorne**, *Ample vector bundles*, Publ. Math. I.H.E.S. **29** (1966), 63-94.
- [4] **L.Katzarkov**, *Shafarevich maps*, Algebraic geometry–Santa Cruz 1995, Proc. Sympos. Pure Math., 62, Part 2, Amer. Math. Soc., Providence, RI, 1997, p. 173–216.
- [5] **A.L.Onishchik**, *Methods in the theory of Sheaves and Stein spaces*, Several Complex Variables IV, Encyclopedia of Mathematical Sciences vol 10, Springer Verlag, 1986, pp. 2–61.

COURANT INSTITUTE FOR MATHEMATICAL SCIENCES, NEW YORK UNIVERSITY
E-mail address: bogomol@cims.nyu.edu