

Braided n -categories and Σ -structures

Lawrence Breen*

Abstract

We associate to any braided 2-groupoid with vanishing intermediate homotopy group a principal bundle (or torsor) endowed with a so-called Σ -structure, and show that this is the natural generalization to the 2-category context of the familiar quadratic invariant describing a braided groupoid. The corresponding structures for higher braided n -groupoids are also examined. This leads to the concept of Γ_3 -torsor pairs, which are in the same relation to cubic forms as torsors endowed with a Σ -structure are to quadratic ones. The discussion also covers the corresponding properties of braided 2- and n -stacks in groupoids.

1 Introduction

This text is one of a series in which we explore the commutativity conditions which may be imposed on tensor laws in monoidal n -categories. We showed in [6] that the structure of a monoidal groupoid \mathcal{C} could be analyzed in terms of a simpler geometric object, the associated commutator biextension, which measures the obstruction to the full commutativity of the tensor law of \mathcal{C} . This commutator biextension, which mimicks the well-known commutator map associated to a central extension of abelian groups, is endowed with a very strong anti-symmetry property, which we termed its alternating structure [4]. Such alternating biextensions are classified up to isomorphism by the groups $\text{Ext}^1(L\Lambda^2 B, A)$, where B and A are a pair of abelian groups, and $L\Lambda^2 B$ is the (non-additive) derived functor of the second exterior power functor $\Lambda_{\mathbf{Z}}^2$. In [6], we explored the manner in which monoidal n -groupoids may be described for $n \geq 2$ in terms of higher analogs of alternating biextensions, related to the higher derived exterior powers $L\Lambda_{\mathbf{Z}}^i B$. We always supposed there, as we will here, that the only nonvanishing homotopy groups of the n -groupoids considered were the extremal ones, the group $\pi_0(\mathcal{C}) = B$ of isomorphism classes of objects and the group $\pi_n(\mathcal{C}) = A$ of self-arrows of the identity $(n-1)$ -arrow, since without this hypothesis the associated geometric objects are no longer standard ones. Such n -groupoids, together with their monoidal structure, are entirely determined up to equivalence by the associated k -invariant of their geometric realization, an element which lives in one of the cohomology groups $H^{r+n}(K(B, r), A)$. The integer r is determined by the level of commutativity of the given tensor multiplication law on \mathcal{C} .

*UMR CNRS 7539, Institut Galilée, Université Paris 13, F-93430 Villetaneuse

In the present text, we consider monoidal n -groupoids \mathcal{C} endowed with a braiding structure, in other words those classified up to equivalence by an element of the cohomology groups $H^{n+2}(K(B, 2), A)$. We address the problem of determining the obstruction to the full commutativity of such a monoidal structure. In this situation, which is more restrictive than that examined in [6] where no braiding axiom was postulated in \mathcal{C} , the role previously assigned to the alternating commutator map associated to a central extension is taken up by the map $\tau : \Gamma_2^{\mathbf{Z}}B \longrightarrow A$ introduced by A. Joyal and R. Street in [18], whose source is the second divided power of the \mathbf{Z} -module B . The latter associates to an object X of \mathcal{C} whose class in B is x the automorphism $\tau(x)$ of $X \otimes X$ determined by the braiding, viewed as an element of A . In the first situation examined here, that of a braided 2-groupoid \mathcal{C} , this map τ is replaced by the (trivial) A -principal bundle (or torsor) on B whose fiber above an element $x \in B$ is the A -torsor of 2-arrows from the identity 1-arrow of $X \otimes X$ to the braiding 1-arrow $\tau(x)$. The additional structure which this A -torsor possesses asserts that it satisfies the theorem of the cube, and that it is symmetric (*i.e.* invariant in a strong sense under pull-back by the inverse map of B). Such A -torsors were said in [3] to be endowed with a Σ -structure. They are much more classical objects than the alternating biextensions encountered above, since they correspond in an algebro-geometric context, when A is the multiplicative group G_m , to symmetric polarizations of the group scheme B .

In the higher categorical situations, one is led to geometric objects associated to the higher divided power functors $\Gamma_i^{\mathbf{Z}}$. Since braided 3-categories do not yield any suprisingly new multilinear structures, we mainly consider those structures associated to braided 4-categories. To these 4-categories correspond somewhat exotic geometric structures, which may be thought of as higher analogs of the torsors with Σ -structures defined by braided 2-categories. While Σ -structure A -torsors on B were classified by the group $\text{Ext}^1(L\Gamma_2 B, A)$ associated to the derived functor $L\Gamma_2 B$ of Γ_2 , these new structures, which we have termed Γ_3 - A -torsor pairs on B , are classified by the group $\text{Ext}^1(L\Gamma_3 B, A)$. These Γ_3 -torsor pairs therefore stand in the same relation to cubic forms as the Σ -structure torsors do to quadratic ones. We unravel their definition here, but do not give a complete proof of the fact that the structures in question are indeed classified by this Ext group. Instead, we limit ourselves to the very strong plausibility argument given by proposition 5.2.

While we will speak freely in this text of n -categories for rather large integers n , ($n \leq 5$) will not make explicit the requisite definitions of such n -categorical structures. Even when this is not explicitly stated, our n -categories will always be assumed to be n -groupoids, in other words categories in which, for each $i < n$, i -arrows are invertible up to an $i + 1$ -arrow, and n -arrows are strictly invertible. The progress recently made by various authors in achieving a workable definition of an n -category, which is reflected in other texts [2], [23] of these Proceedings, justifies in our opinion this choice of the n -category terminology for what are in fact specific homotopy types determined by 2-stage Postnikov fibrations. A reader unwilling to endorse this point of view should consider this text as providing a very explicit description of a natural filtration on certain cohomology groups $H^{n+2}(K(B, 2), A)$. Nor will we make explicit the corresponding definition of an n -stack, and simply refer to [5] for an illustration of what we have in mind in the case $n = 2$. An n -stacks in groupoids with trivial non extremal homotopy sheaves corresponds to a simplicial 2-stage Postnikov simplicial

sheaf, described up to equivalence by the hyper-cohomology groups $\mathbf{H}^{r+n}(K(B, r), A)$. Here the extremal homotopy sheaves $\pi_i(\mathcal{C})$ associated to \mathcal{C} for are defined in the standard manner by sheafification of the naive homotopy presheaves associated to \mathcal{C} . The drawback with such a definition is that much of the geometry underlying the idea of an n -category, or an n -stack, viewed as a family of objects together with an assortment of arrows recedes into the background. We refer to [22],[17] for an up to date discussion of the theory of n -stacks. In the present context, the theory which we develop for n -groupoids extends to n -stacks in groupoids (which we will henceforth simply refer to as n -stacks). The biextensions, Σ -structures, and other geometric structures which we encounter in the n -groupoid case still exist, but no longer possess global sections which trivialize the underlying torsors. For the rest the situation is unchanged, with the proviso that additional non trivial terms in the filtration of the cohomology must be considered, which correspond to non vanishing higher Ext groups.

2 Braided categories and stacks

We begin by reviewing the classification of braided groupoids. Such a category may be defined directly, as in [18], as a groupoid endowed with a monoidal structure for which there exist inverses for objects and endowed with braiding morphisms¹

$$\tau_{X,Y} : YX \longrightarrow XY \tag{2.1}$$

compatible with the monoidal structure. It may also be viewed as a 3-groupoid whose set of objects and of 1-arrows are both reduced to a single element. The latter description makes it clear how to classify such categories. Let \mathcal{C} be a braided groupoid. The group $\pi_0(\mathcal{C})$ of isomorphism classes of objects of such a braided groupoid \mathcal{C} and the group $\pi_1(\mathcal{C})$ of automorphisms of the identity object I of \mathcal{C} are both abelian, and are respectively denoted B and A . The monoidal structure on \mathcal{C} determines for every object $X \in \mathcal{C}$, by left multiplication by X , an identification

$$(X \otimes -) : B = \text{Aut}_{\mathcal{C}}(I) \longrightarrow \text{Aut}_{\mathcal{C}}(X) \tag{2.2}$$

The nerve \mathcal{X} of \mathcal{C} is a homotopy commutative H -space, which admits a double delooping to a space $\mathcal{Y} = B^2\mathcal{X}$ whose only non-trivial homotopy groups are

$$\pi_i(\mathcal{Y}) = \begin{cases} B & i = 2 \\ A & i = 3 \end{cases}$$

Such a space is entirely described, up to equivalence, by its unique k -invariant, an element $k(\mathcal{Y})$ in the cohomology group $H^4(K(B, 2), A)$. The entire H -space structure of \mathcal{X} , in other words the braided category structure of \mathcal{C} , can be recovered from \mathcal{Y} by the double delooping $\mathcal{X} \sim \Omega^2\mathcal{Y}$, so that this invariant yields a complete classification for equivalence classes of braided categories with associated groups B and A .

¹We henceforth will denote by XY , rather than by the more customary $X \otimes Y$, the product of two objects X and Y in a monoidal category \mathcal{C} . On the other hand we will use the additive notation for the composition law in B .

Let us now examine the integral homology of $K(B, 2)$ in low degrees. This computation, which is functorial in B and goes back to [13] II, yields the following result

$$H_i(K(B, 2)) = \begin{cases} B & i = 2 \\ 0 & i = 3 \\ \Gamma_2(B) & i = 4 \end{cases} \quad (2.3)$$

A convenient method for studying the integral homology of $K(B, 2)$ in all degrees, along the lines of the computations for $K(B, 1)$ in [7], consists in observing that the homology of $K(B, 2)$ (with the algebra structure determined by the Pontrjagin product) is endowed with a natural divided power structure on even degree elements. The natural map $B \simeq H_2(K(B, 2))$ determined by the fundamental class therefore extends to a functorial algebra homomorphism

$$\Gamma_*(B) \longrightarrow H_*(K(B, 2)) \quad (2.4)$$

When the abelian group B is torsion-free, the entire homology of $K(B, 2)$ is easy to compute. The following result, which is implicit in [8] (exposé 11, theorems 4 and 6), is explicitly stated in [14] (proposition 3.4).

Lemma 2.1 *Let B be a torsion-free abelian group. The canonical map (2.4) is an isomorphism.*

Proof: For such a B , the homology $H_*(K(B, 1))$ of the group B is isomorphic to the exterior algebra $\Lambda^*(B)$ of the \mathbf{Z} -module B . A bar-construction (or Eilenberg-Moore spectral sequence) argument implies directly that

$$H_*(K(B, 2)) \simeq \text{Tor}^{\Lambda^*B}(\mathbf{Z}, \mathbf{Z})$$

It is an immediate consequence of the existence of the Koszul complex that the latter Tor term is isomorphic, as required, to Γ_*B .

The methods of [7] now allow us to compute the homology of $K(B, 2)$. Let $B_* \longrightarrow B$ be a torsion-free resolution of B . This determines a simplicial resolution $K(B_*, 0)$ of the constant simplicial group $K(B, 0)$, as well as a torsion-free simplicial resolution $K(B_*, 2)$ of the simplicial abelian group $K(B, 2)$. By the previous lemma, the E_1 term of the homology spectral sequence associated as in [7] §1 to $K(B_*, 2)$ can be identified with the simplicial abelian group $\Gamma_*(B_*, 0)$, obtained by applying the functor Γ_* componentwise to $K(B_*, 0)$. Since the homology of $K(B_*, 2)$ coincides with that of $K(B, 2)$, the following proposition is obtained.

Proposition 2.2 *There exists a functorial spectral sequence*

$$E_{p,q}^2 = L_p\Gamma_q(B, 0) \implies H_{p+2q}(K(B, 2))$$

which degenerates at E^2 .

The degeneracy of the spectral sequence was essentially proved in [7] (with a different description of the initial term). The latter is here the p th left (non-additive) derived functor of Γ_q , as defined in [12], applied to B placed in degree 0, which will henceforth be simply denoted $L_p\Gamma_q B$ rather than the morse standard $L_p\Gamma_q(B, 0)$. In low degrees, this proposition immediately yields the computations (2.3), as well as the isomorphism

$$H_5(K(B, 2)) \simeq L_1\Gamma_2 B \tag{2.5}$$

We refer to [3] §9 for a discussion of the functor $L_1\Gamma_2 B$, and its identification with the functor $R(B)$ which by [13] theorem 22.1 describes the group $H_5(K(B, 2))$.

We now consider the cohomology of $K(B, 2)$. It follows from the universal coefficient theorem and the computation (2.3) that

$$H^4(K(B, 2), A) \simeq \text{Hom}(\Gamma_2(B), A) \tag{2.6}$$

so that braided categories \mathcal{C} with associated groups B and A are classified up to equivalence by quadratic maps $\tau_{\mathcal{C}} : B \rightarrow A$. As observed by A. Joyal and R. Street ([18] theorem 3.3), this identification, which follows from the explicit description of cycles for $H_4(K(B, n))$ (as in [13] theorem 26.1), associates to a category \mathcal{C} with a braiding 2.1 the map

$$\tau = \tau_{\mathcal{C}} : B \rightarrow A \tag{2.7}$$

which sends an element $x \in B$ to the automorphism τ_{X_x, X_x} of $X_x X_x$, for some representative object $X_x \in \mathcal{C}$ of x . The fact that $\tau_{\mathcal{C}}$ is independent of this choice of representative, and that it is quadratic, follows directly from (2.6). It is, however, of some interest to verify directly the quadraticity of this map in geometric terms. To do this conveniently, we begin by assuming that the multiplication law in our braided category \mathcal{C} is strictly associative, a situation which can always be achieved by replacing \mathcal{C} by an equivalent strictly associative monoidal category \mathcal{D} , to which the braiding may be transported. In this new category, the commutative hexagons which determine the braided structure reduce to triangles such as the triangles

$$\begin{array}{ccc}
 & X_x X_y X_z & \\
 & \swarrow \quad \searrow & \\
 X_y X_x X_z & \xrightarrow{\quad} & X_y X_z X_x
 \end{array}
 \qquad
 \begin{array}{ccc}
 & X_x X_y X_z & \\
 & \swarrow \quad \searrow & \\
 X_x X_z X_y & \xrightarrow{\quad} & X_z X_x X_y
 \end{array}$$

which respectively shuffle X_x from the left through $X_y X_z$ and X_z from the right through $X_x X_y$. We denote the first of these ‘hexagon’ diagrams by $H_{[x|_2 y, z]}$ and the second by $H_{[x, y|_2 z]}$ and observe that these notations are consistent with Eilenberg-Mac Lane’s for cells in $K(B, 2)$. Let us choose, for each pair of elements $x, y \in B$, an arrow

$$X_x X_y \xrightarrow{c_{x, y}} X_{x+y} \tag{2.8}$$

and henceforth denote, for any pair of elements $x, y \in B$, by $\tau_{x,y}$ the braiding map τ_{X_x, X_y} (2.1). We now consider the commutative diagram

$$\begin{array}{ccccccc}
 X_{x+y} X_{x+y} & \longrightarrow & X_{x+y} X_x X_y & \longrightarrow & X_x X_y X_x X_y & \xrightarrow{\tau_{x,y}} & X_x X_x X_y X_y & (2.9) \\
 \downarrow \tau(x+y) & & \downarrow & \searrow & & & \downarrow \tau(x) \\
 & & & & X_x X_{x+y} & \longrightarrow & X_x X_x X_y X_y \\
 & & & \swarrow & & & \downarrow \tau(y) \\
 X_{x+y} X_{x+y} & \longrightarrow & X_x X_y X_{x+y} & \longrightarrow & X_x X_y X_x X_y & \xleftarrow{\tau_{y,x}} & X_x X_x X_y X_y
 \end{array}$$

Here the middle triangle is $H_{[x+y|_2 x, y]}$, and the top and bottom right-hand ones are respectively $H_{[x, y|_2 x]}$ and $H_{[x, y|_2 y]}$, the commutative squares being the obvious ones which follow from the functoriality of the maps τ . The commutativity of the outer square, when interpreted via arrows (2.8) as a square with the same object X_{2x+2y} at all vertices, asserts that the identity

$$\tau(x+y) - \tau(x) - \tau(y) = g(x, y) + g(y, x)$$

is satisfied in B , where $g(x, y) \in B$ is defined by the commutative diagram

$$\begin{array}{ccc}
 X_y X_x & \xrightarrow{\tau_{x,y}} & X_x X_y & (2.10) \\
 c_{y,x} \downarrow & & \downarrow c_{x,y} \\
 X_{x+y} & \xrightarrow{g(x,y)} & X_{x+y}
 \end{array}$$

Asserting that the map τ (2.7) is of degree two is therefore equivalent to showing that that the map $\phi(x, y) : B \times B \longrightarrow B$ defined by

$$\phi(x, y) = g(x, y) + g(y, x) \quad (2.11)$$

is bilinear. In order to prove this, consider first of all the well-known ‘‘Yang-Baxter’’ hexagon² associated to any three objects X, Y, Z in the braided category \mathcal{D} . This can be built up as follows from a pair of triangles $H_{[x|_2 y, z]}$ and $H_{[x|_2 z, y]}$

$$\begin{array}{ccccc}
 & & XZY & \longrightarrow & ZXY & & (2.12) \\
 & & \swarrow & & \searrow & & \\
 XYZ & & & & & & ZYX \\
 & & \searrow & & \swarrow & & \\
 & & YXZ & \longrightarrow & YZX & &
 \end{array}$$

²This hexagon (which is actually a hexagon when the associativity in the monoidal category is no longer assumed to be strict) is also referred to in the physics literature as the ‘‘ABC = CBA’’ identity.

(the other method for constructing the identical Yang-Baxter hexagon is by the scheme

$$\begin{array}{ccc}
 & XZY \longrightarrow ZXY & \\
 & \nearrow \quad \dashrightarrow & \\
 XYZ & & ZYX \\
 & \searrow \quad \dashrightarrow & \\
 & YXZ \longrightarrow YZX &
 \end{array} \tag{2.13}$$

the upper and lower commutative triangles now being $H_{[x,y|_2z]}$ and $H_{[y,x|_2z]}$). In both instances, the commutativity of the central square follows from the functoriality of the braiding map (2.1)). Returning to the study of the map (2.11), its linearity in the first variable may be obtained by considering the following commutative diagram, built up out from the Yang-Baxter hexagon associated to the three objects X_x, X_y, X_z , to which have been glued a pair of triangles $H_{[x,z|_2y]}$, $H_{[y|_2z,x]}$ and a square whose commutativity follows, as always, from the functoriality of the braiding map:

$$\begin{array}{ccccc}
 & & X_x X_z X_y & \xrightarrow{\tau_{z,x}} & X_z X_x X_y & \\
 & & \nearrow \tau_{z,y} & & \searrow \tau_{y,x} & \\
 & X_x X_y X_z & & & X_z X_y X_x & \\
 \nearrow \tau_{y,z} & & \searrow \tau_{y,x} & & \nearrow \tau_{z,y} & \searrow \tau_{x,y} \\
 X_x X_z X_y & \xrightarrow{\tau_{X_y, X_x X_z}} & X_y X_x X_z & \xrightarrow{\tau_{z,x}} & X_y X_z X_x & \xrightarrow{\tau_{X_z X_x, X_y}} & X_z X_x X_y \\
 & & \searrow \tau_{X_x X_z, X_y} & & \nearrow \tau_{z,x} & \\
 & & & & X_x X_z X_y &
 \end{array} \tag{2.14}$$

The arrows (2.8) allows us to transform each of the eight vertices of the outer diagram into the object X_{x+y+z} . The left hand (*resp.* right-hand) composite arrow then becomes our map $\phi(z, y)$ (*resp.* $\phi(x, y)$), and the lower composite one is $\phi(xz, y)$. The two final arrows $\tau_{z,x}$ encountered, when going around this diagram are both transformed into the maps $g_{x,z}$ (2.10), with opposite orientations. Since the group A of automorphisms of an object X in a braided category is abelian, these cancel out, and the required identity

$$\phi(xz, y) = \phi(x, y) + \phi(z, y)$$

is satisfied. The linearity of ϕ in the second variable y follows automatically, since the map ϕ is symmetric by construction.

In order to verify the quadraticity of the map τ (2.7), it is still necessary to ascertain that τ satisfies the identity

$$\tau(x) = \tau(-x) \tag{2.15}$$

Consider, for any $x \in B$, the commutative square

$$\begin{array}{ccc}
 X_x X_x X_{-x} X_{-x} & \xrightarrow{\tau(-x)} & X_x X_x X_{-x} X_{-x} \\
 \tau(x) \downarrow & \searrow & \downarrow \tau_{x,-x} \\
 X_x X_x X_{-x} X_{-x} & \xrightarrow{\tau_{x,-x}} & X_x X_{-x} X_x X_{-x}
 \end{array} \tag{2.16}$$

built from an upper triangle $H_{[x,-x]_2 |_{x,-x}}$ and a lower triangle $H_{[x]_2 |_{x,-x}}$. The common diagonal, when viewed after contraction to the identity object I of the middle term $X_x X_{-x}$ at its source and one or the other of the expressions $X_x X_{-x}$ at its target, is essentially the identity map from $X_x X_x$ to itself. The triangles may therefore be viewed as having a common diagonal edge along which they may be attached as shown. By the process previously applied to the diagram (2.14), the outer square in (2.16) transforms to one in which all four vertices equal to $I = X_0$. The identity

$$\tau(x) + \phi(x, -x) = \tau(-x) + \phi(x, -x)$$

may be read off from its outer edges, so that the identity (2.15) is proved.

Remark 2.3 *i)* In a category in which the multiplication law is not strictly associative, it is possible to carry out the above argument directly, without passing to a strictly associative equivalent category \mathcal{D} . A direct verification that $H_4(K(B, 2)) \simeq \Gamma_2(B)$ makes it clear that the only significant pentagonal cell to be added in that case to diagram (2.9) is an associativity pentagon $P_{x,y,x,y}$, inserted at the common vertex $X_x X_y X_x X_y$ appearing in both the top and bottom position. A pentagon $P_{x,-x,x,-x}$ must similarly be added at the vertex $X_x X_{-x} X_x X_{-x}$ to diagram (2.16)

ii) It also follows from (2.3), and the universal coefficient theorem, that

$$H^3(K(B, 2), A) \simeq \text{Ext}^1(B, A) \tag{2.17}$$

$$H^2(K(B, 2), A) \simeq \text{Hom}(B, A) \tag{2.18}$$

and these relations have geometric interpretations analogous to that given above for (2.6). The first implies, as mentioned in [18] remark 3.3, that the equivalences $F : \mathcal{C} \longrightarrow \mathcal{D}$ between \mathcal{C} and another braided category \mathcal{D} equivalent to it, when viewed up to natural equivalence, form a principal homogenous space under the group $\text{Ext}^1(B, A)$ of auto-equivalences of \mathcal{C} . The second relation asserts that the set of natural transformations between the identity equivalence $1_{\mathcal{C}} : \mathcal{C} \longrightarrow \mathcal{C}$ and some other fixed equivalence $u : \mathcal{C} \longrightarrow \mathcal{C}$ is either empty, or a principal homogeneous space under the group $\text{Hom}(B, A)$ of natural transformations between $1_{\mathcal{C}}$ and itself. We will not make explicit the geometrical realization of these assertions, but observe that this yields, together with the interpretation given above of (2.6), a complete description up to 2-equivalence of the (symmetric monoidal) 2-category of braided categories with associated abelian groups B and A , analogous to the one given in [18] for the underlying 1-category.

The previous discussion may also be carried out in the more general context of a braided stack \mathcal{C} . B and A are now abelian groups in the topos T considered, with the proviso that a

section x of B is only locally represented by an object X_x in \mathcal{C} . The isomorphisms (2.17) are essentially unchanged in this new situation, despite the fact that $H^i(K(B, 2), A)$ now denotes the A -valued hypercohomology groups $\mathbf{H}^i(K(B, 2), A)$ of the simplicial object $K(B, 2)$ of T . The higher Ext^i groups no longer vanish for $i > 1$ in categories of abelian sheaves, as they did in the category of abelian groups, so that the isomorphism (2.6) must now be replaced by the exact sequence

$$0 \longrightarrow \text{Ext}^2(B, A) \longrightarrow H^4(K(B, 2), A) \longrightarrow \text{Hom}(\Gamma_2(B), A) \longrightarrow \text{Ext}^3(B, A) \quad (2.19)$$

which is a low degree consequence of the universal coefficient spectral sequence

$$\text{Ext}^p(H_q(K(B, 2)), A) \implies H^{p+q}(K(B, 2), A)$$

and of computation (2.3). The map $H^4(K(B, 2), A) \longrightarrow \text{Hom}(\Gamma_2(B), A)$ is essentially the one studied above. While it is *a priori* only defined locally, the fact that the explicit definition of the map (2.7) was independent of the choice of representing object X_x for a section $x \in B$ ensures that these local versions of $\tau_{\mathcal{C}}$ are compatible. They therefore glue in the stack context to a genuine, globally defined, sheaf map $\tau_{\mathcal{C}} : B \longrightarrow A$. While the map

$$H^4(K(B, 2), A) \longrightarrow \text{Hom}(\Gamma_2(B), A) \quad (2.20)$$

is no longer injective, its kernel $\text{Ext}^2(B, A)$ once again has a natural interpretation as the group of equivalence classes of strictly symmetric monoidal categories with associated groups B and A [11]. The quadratic maps from B to A which live in the image of the map (2.20) thus still have, in the stack situation, a natural interpretation as the group of obstructions to the existence, on a braided monoidal category \mathcal{C} with associated groups B and A , of a strictly symmetric structure. This may be directly understood by observing that, the vanishing of the map (2.7) implies that $\tau_{X,X}$ (2.1) is the identity map for all objects X in \mathcal{C} . It then follows from diagram (2.9) that for any pair of objects X and Y in \mathcal{C} the composite map

$$YX \xrightarrow{\tau_{x,y}} XY \xrightarrow{\tau_{y,x}} YX$$

is also the identity map, so that \mathcal{C} is indeed, as asserted, a strictly symmetric monoidal category.

3 Braided 2-categories and Σ -structures

Let us now pass from the study of categories to that of 2-categories. Braiding structures on such categories were introduced by Kapranov and Voevodsky [19] (see also [1], [10] definition 12, [9]). We restrict ourselves here to 2-groupoids (*resp.* 2-stacks) whose intermediate homotopy group $\pi_1(\mathcal{C})$ is trivial, in other words those for which, for any given pair of arrows $f, g : X \longrightarrow Y$ between a pair of objects X, Y , there exists (*resp.* there exists locally) a 2-arrow $u : f \implies g$ connecting f to g . The first of the remaining homotopy groups of such a braided 2-groupoid \mathcal{C} is the abelian group $\pi_0(\mathcal{C}) = B$ of isomorphism classes of objects of \mathcal{C} . For a given object X of \mathcal{C} (which we may choose to be the identity object I of \mathcal{C}), the

previous hypothesis ensures that the symmetric monoidal category $\mathcal{A} = \text{Aut}_{\mathcal{C}}(X)$ is equivalent to the category with a unique object and an abelian group A of arrows. To make things specific, we may think of A as the group of 2-arrows

$$u : 1_I \Longrightarrow 1_I$$

from the identity 1-arrow $1_I : I \longrightarrow I$ of the identity object I to itself. In the stack case, both B and A are abelian sheaves, and \mathcal{A} is equivalent as a symmetric monoidal stack to the stack $\text{Tors}(A)$ of A -torsors. As discussed in [5] §8, the unique extra invariant necessary for a full description of \mathcal{C} as a braided 2-category or stack is “Postnikov invariant” of \mathcal{C} , an element k in the cohomology (or hypercohomology group $H^4(K(B, 2), \mathcal{A})$). Under our vanishing hypothesis for $\pi_1(\mathcal{C})$, this reduces to an element of the more traditional cohomology (or hypercohomology) group $H^5(K(B, 2), A)$. A direct analysis of this group *via* the universal coefficient theorem is easy to achieve since the values of the requisite homology groups of B are known to us by (2.3) and (2.5). Let us consider instead, along the lines of the discussion carried out for monoidal 2-categories in [6], the filtration on the cohomology group $H^5(K(B, 2), A)$ induced by the filtration on homology determined by proposition 2.2. In the so-called punctual case in which 2-categories (rather than to 2-stacks) are considered, the only term in the filtration which contributes something non trivial is the one involving Γ_2 , so that one obtains an isomorphism

$$H^5(K(B, 2), A) \simeq \text{Ext}^1(L\Gamma_2 B, A) \tag{3.1}$$

Both left and right hand term can be subjected to a further analysis via universal coefficient theorem. To the standard decomposition

$$0 \longrightarrow \text{Ext}^1(H_4(K(B, 2), A) \longrightarrow H^5(K(B, 2), A) \longrightarrow \text{Hom}(H_5(K(B, 2)), A) \longrightarrow 0$$

of the left-hand term of (3.1) corresponds the decomposition

$$0 \longrightarrow \text{Ext}^1(\Gamma_2(B), A) \longrightarrow \text{Ext}^1(L\Gamma_2 B, A) \longrightarrow \text{Hom}(L_1\Gamma_2(B), A) \longrightarrow 0$$

of the right-hand one, and this correspondence is consistent with the computation (2.3), (2.5) of the low degree homology of $K(B, 2)$.

The term $\text{Ext}^1(L\Gamma_2 B, A)$ which appeared in (3.1) may be interpreted geometrically as the group of isomorphism classes of A -torsors on B endowed with a so-called Σ -structure. For the reader’s convenience we rapidly review this concept, and refer to [3] for additional information. It is a strengthening of the more familiar notion of a cube structure on such an A -torsor P on B . The latter is determined by a trivialization s of the “second difference” $\Theta(P)$ of the given torsor P , which is the A -torsor on B^3 defined by

$$\Theta(P) = m_{123}^* P \wedge \left(\bigwedge_{ij} m_{ij}^* P \right)^{-1} \wedge \left(\bigwedge_i p_i^* P \right) \tag{3.2}$$

where \wedge is the contracted product of A -torsors, and the maps m_{123}, m_{ij}, p_i from B^3 to B are respectively the iterated sum, the partial (i, j) th sum and the projection on the i th factor. In addition, the trivialization s is required to satisfy a cocycle condition, for which we refer to

[3] §2. An alternate method for describing a cube structure on the A -torsor P is to introduce its “first difference” of the A -torsor P . This is the A -torsor $\Lambda(P)$ on B^2 defined by

$$\Lambda(P) = m^*(P) \wedge p_1^*(P)^{-1} \wedge p_2^*(P)^{-1} \quad (3.3)$$

(where m is the group law and p_i are the projections from $B \times B$ to B). The A -torsor P is endowed with a cube structure if and only if $\Lambda(P)$ is not simply an A -torsor above B^2 but actually a biextension³ of $B \times B$ by A .

A Σ -structure on an A -torsor P endowed with a cube structure is determined by the additional requirement that the torsor P be symmetric, in other words that there exists an isomorphism of A -torsors on B

$$\lambda : P \longrightarrow i^*P \quad (3.4)$$

(where $i : B \longrightarrow B$ is the inverse map for the group law of B) compatible in an appropriate sense with the cube structure of P (see [3] §5).

Example 3.1 In an algebro-geometric context, when A is the multiplicative group G_m and B a commutative group variety, an A -torsor on B is nothing else than a line bundle on B . In particular, when the group B is an abelian variety, we are assured of the existence of a trivialization s of the induced line bundle $\Theta(L)$ by the classical theorem of the cube [20]. In that situation, the cocycle condition mentioned above is satisfied, so that any such line bundle is automatically endowed with a cube structure. Furthermore, a symmetrization isomorphism λ (3.4) is then always compatible with the cube structure, whenever λ respects the rigidification, in other words sends the point $s(e, e, e)$ in the fibre P_e of P above the unit element $e \in B$ to itself.

Before explaining the correspondence (3.1) between braided 2-groupoids and Σ -structures in more geometric terms, let us examine in purely cohomological terms the corresponding situation in the 2-stack case. The identification (3.1) is now replaced by an exact sequence

$$\rightarrow \mathrm{Hom}(\Gamma_2(B), A) \rightarrow \mathrm{Ext}^3(B, A) \rightarrow H^5(K(B, 2), A) \xrightarrow{\tau} \mathrm{Ext}^1(L\Gamma_2 B, A) \rightarrow \mathrm{Ext}^4(B, A) \quad (3.5)$$

which prolongs the exact sequence (2.19). The term $\mathrm{Ext}^1(L\Gamma_2 B, A)$ once more describes A -torsors endowed with Σ -structures above B , but in the richer, sheaf-theoretic context in which a torsor does not, as in the punctual case, always possess a global section. The exactness of this sequence shows that the Σ -structure $\tau(\mathcal{C})$ associated by the middle map τ to a braided 2-stack \mathcal{C} is trivial if and only if \mathcal{C} is Picard strict, in the terminology of [5] definition 8.5.

This Picard strict definition is the strongest possible commutativity which can be imposed on a 2-stack, and it is of little interest in the punctual case since the group $\mathrm{Ext}^3(B, A)$ which classifies such structures for abstract abelian groups B and A vanishes. Let us briefly review from [5], for the reader’s convenience, the relations between this and other possible commutativity conditions on 2-categories. This Picard strict commutativity condition

³ We refer to [16] for a definition of this concept.

is strictly stronger than the condition on monoidal 2-categories which we called “strongly involutory”, which produces the 2-categories which R. Day and R. Street refer to as “symmetric Gray monoids” [10], and S. Crans as “symmetric monoidal 2-categories” [9]. The latter terminology for such 2-categories, which are classified under the vanishing hypothesis for $\pi_1(\mathcal{C})$ by elements in the group $H^7(K(B, 4), A)$, seems the most appropriate one since the cohomology group in question lies in the stable range. This condition is in turn more restrictive than the condition which we refer to as “weakly involutory”, and which applies to those categories which Day and Street call “sylleptic Gray monoids” and Crans refers to as “sylleptic monoidal 2-categories”. The latter are classified, when $\pi_1(\mathcal{C})$ vanishes, by elements in $H^6(K(B, 3), A)$. Finally, the commutativity conditions defining the original braided monoidal 2-categories of [19] are even weaker, since the latter are classified by elements of the group $H^5(K(B, 2), A)$. In each of these instances the universal coefficient theorem yields, in the stack case, an edge-homomorphism map

$$\mathrm{Ext}^3(B, A) \longrightarrow H^{i+3}(K(B, i), A)$$

and these maps are compatible with the suspension maps

$$H^{i+3}(K(B, i), A) \xrightarrow{S} H^{i+2}(K(B, i-1), A)$$

as i varies.

If we now reason in geometric rather than homological terms, the Σ -structure associated to a braided 2-stack \mathcal{C} may be understood by introducing the A -torsor $\mathcal{S}_{\mathcal{C}}$ on B , whose sections above an section $x \in B$ are the 2-arrows

$$\begin{array}{ccc} & 1_{X_x X_x} & \\ & \curvearrowright & \\ X_x X_x & & X_x X_x \\ & \curvearrowleft & \\ & \tau(x) & \end{array} \quad (3.6)$$

The A -torsor structure on $\mathcal{S}_{\mathcal{C}}$ is defined by composing such a locally defined 2-arrow with a 2-arrow from the identity map $1_{X_x X_x}$ to itself. In order to verify that $\mathcal{S}_{\mathcal{C}}$ is endowed with a Σ -structure, we must first verify that it is endowed with a cube structure, in other words that its “first difference”

$$\Lambda(\mathcal{S}_{\mathcal{C}}) = \mathrm{Isom}(p_1^* \mathcal{S}_{\mathcal{C}} \wedge p_2^* \mathcal{S}_{\mathcal{C}}, m^* \mathcal{S}_{\mathcal{C}})$$

is a biextension of $B \times B$ by A . By definition, a local section of $\Lambda(\mathcal{S}_{\mathcal{C}})$ is a rule which assigns to every (x, y) in some open set of $B \times B$, a triple of 2-arrows (S_x, S_y, S_{x+y}) (3.6), in a manner consistent with the indicated action of A . Reinterpreting diagram (2.9) in a 2-categorical context, we observe that the commutative triangles $H_{[x+y]_2 x, y}$, $H_{[x, y]_2 x}$ and $H_{[x, y]_2 y}$ have now been replaced by the corresponding hexagon 2-arrows determined by the braiding structure (see [19]). The other more trivial commutative cells in diagram (2.9) are also replaced by corresponding 2-arrows, so that diagram (2.9) determines, by composition

of all these 2-arrows, a composite 2-arrow⁴

$$\begin{array}{ccc}
 X_{x+y} X_{x+y} \longrightarrow X_x X_y X_x X_y \xrightarrow{\tau_{x,y}} X_x X_x X_y X_y & & (3.7) \\
 \downarrow \tau(x+y) & \swarrow & \downarrow \tau(x) \\
 & & X_x X_x X_y X_y \\
 & & \downarrow \tau(y) \\
 X_{x+y} X_{x+y} \longrightarrow X_x X_y X_x X_y \xleftarrow{\tau_{y,x}} X_x X_x X_y X_y & &
 \end{array}$$

Composing it with the given 2-arrows (S_x, S_y, S_{x+y}) according to the scheme indicated in the diagram

$$\begin{array}{ccc}
 X_{x+y} X_{x+y} \longrightarrow X_x X_y X_x X_y \xrightarrow{\tau_{x,y}} X_x X_x X_y X_y & & (3.8) \\
 \downarrow \tau(x+y) & \swarrow & \downarrow \tau(x) \Big| \leftarrow \Big) 1 \\
 & & X_x X_x X_y X_y \\
 & & \downarrow \tau(y) \Big| \leftarrow \Big) 1 \\
 X_{x+y} X_{x+y} \longrightarrow X_x X_y X_x X_y \xleftarrow{\tau_{y,x}} X_x X_x X_y X_y & &
 \end{array}$$

this yields a corresponding family of 2-arrows

$$\begin{array}{ccc}
 & 1_{X_y X_x} & \\
 X_y X_x & \xrightarrow{\quad} & X_y X_x \\
 & \Downarrow S_{x,y} & \\
 & \tau_{y,x} \tau_{x,y} &
 \end{array}
 \quad (3.9)$$

Let us now reconsider, in the 2-categorical context, the ‘‘Yang-Baxter’’ diagram (2.14), where all cells have now been replaced by 2-arrows determined by the braiding structure of \mathcal{C} . It may be viewed as a rule which assigns to every pair of 2-arrows (3.9)

$$\begin{array}{ccc}
 X_x X_y & \xrightarrow{1_{X_x X_y}} & X_x X_y & X_z X_y & \xrightarrow{1_{X_z X_y}} & X_z X_y \\
 \Downarrow S_{y,x} & & \Downarrow S_{y,z} & & & \\
 \tau_{x,y} \tau_{y,x} & & \tau_{z,y} \tau_{y,z} & & &
 \end{array}
 \quad (3.10)$$

a corresponding 2-arrow

$$\begin{array}{ccc}
 & 1_{X_{xz} X_y} & \\
 X_{xz} X_y & \xrightarrow{\quad} & X_{xz} X_y \\
 & \Downarrow S_{y,xz} & \\
 & \tau_{xz,y} \tau_{y,xz} &
 \end{array}
 \quad (3.11)$$

⁴This 2-arrow is roughly half of the diagram of 2-arrows denoted $((\bullet \otimes \bullet) \otimes (\bullet \otimes \bullet))$ in [19], when specialized from diagrams with vertices of the general form $XYZW$ and appropriate permutations thereof to those for which $X = Z$ and $Y = W$.

Remark 3.2 It is now easy to understand in what sense $\mathcal{S}_{\mathcal{C}}$ is an obstruction to commutativity of the group law of \mathcal{C} : when $\mathcal{S}_{\mathcal{C}}$ is trivial, there exists a global family of 2-arrows S_x . The axioms for a strict Picard 2-category, or 2-stack, as given in [5] definition 8.5, may now be read off as follows from the previous Σ -structure conditions. The requirement that S_x be additive with respect to x , in the sense briefly described in [5] page 150, corresponds in our context to the assertion that the family of 2-arrows $S_{x,y}$ (3.9) coincide with that determined by the braiding structure on \mathcal{C} . The extra requirement that the 2-arrow displayed in [5] definition 8.5 coincide with $S_{x,x}$ is equivalent to the assertion above that the composite 2-arrow associated by (3.14) to a given 2-arrow S_{-x} is equal to S_x . The geometric content of the previous discussion may thus be summarized in very compact form as the assertion that a braided 2-stack endowed with a family of 2-arrows S_x satisfying the two condition just mentioned, is a strict Picard 2-stack, *i.e.* automatically satisfies the intermediate ‘Picard’ commutativity axioms for a symmetric monoidal 2-stack. In particular, this asserts in the category context that a braided 2-category endowed with a family of 2-arrows S_x satisfying these conditions (and for which $\pi_1(\mathcal{C}) = 0$) is equivalent to the trivial symmetric monoidal 2-category determined by a pair of abelian groups B and A .

4 Comparison with the non-commutative obstructions

It was shown in [6] §8 how to associate to any monoidal 2-category (or 2-stack) \mathcal{C} determined as above by a pair of abelian groups B and A a so-called weak (2,2)-extension \mathcal{E} of $B \times B$ by A , which under an additional condition was found to be alternating in a requisite sense. By the discussion at the very end of [6], a trivialization of \mathcal{E} compatible with this entire structure makes \mathcal{C} into a strict Picard stack (and therefore in the punctual case trivializes \mathcal{C} compatibly with its monoidal structure). We now compare this (2,2)-extension obstruction \mathcal{E} (which we will denote by $\mathcal{E}_{\mathcal{C}}$ when we wish to emphasize its dependence on the given 2-category \mathcal{C}) with the corresponding Σ -structure obstruction $\mathcal{S}_{\mathcal{C}}$ arising under the stronger assumption that \mathcal{C} is braided.

Before passing to this 2-category situation, it is convenient to examine the corresponding assertion for monoidal and braided 1-categories. As recalled in §1, the obstruction to commutativity in a braided category or stack \mathcal{C} with associated invariant groups B and A is determined by the quadratic map $\tau_{\mathcal{C}}$ (2.7) from B to A . On the other hand, by [6] proposition 3.1, there corresponds to any monoidal category or stack \mathcal{C} with associated invariants B and A an A -torsor $E_{\mathcal{C}} = E$ above $B \times B$, defined above $(x, y) \in B \times B$ by

$$E_{x,y} = \text{Isom}_{\mathcal{C}}(X_y X_x, X_x X_y) \quad (4.1)$$

and which is endowed with the structure on a weak biextension⁶. Furthermore, when B is commutative, the associated commutator map, which examines whether the partial group laws on E are commutative, is a homomorphism $\phi_{\mathcal{C}}$ from $\Lambda^3 B$ to A , whose vanishing ensures that E is a (standard) biextension. The biextension E is then automatically an alternating

⁶A weak biextension is simply a (standard) biextension in which the pair of partial composition laws are no longer required to be commutative, see [16] exposé VII 2.10.1.

one, a concept for which we refer to [6] §2, [4]. By [6] corollaries 5.3 and 6.1, E then measures the obstruction to the existence on \mathcal{C} of a strict Picard structure. When the category \mathcal{C} is braided, the relation between the quadratic map $\tau_{\mathcal{C}}$ and the alternating biextension $E_{\mathcal{C}}$ goes as follows

Proposition 4.1 *Let \mathcal{C} be a braided category with associated abelian groups B and A . The weak biextension $E_{\mathcal{C}}$ (4.1) associated to \mathcal{C} is a genuine biextension, and in fact a trivial one. The quadratic map which by [4] proposition 1.5 determines the alternating structure on this trivial biextension E is, up to sign, the map $\tau_{\mathcal{C}}$ (2.7).*

Proof: The braiding maps $\tau_{x,y}$ (2.1) determine a section s of the A -torsor E (4.1) above $B \times B$. The hexagon axioms assert that this section is bimultiplicative, and therefore trivializes E as a biextension. This shows *a posteriori* that the weak biextension $E_{\mathcal{C}}$ is in fact a genuine one. An alternating structure on $E_{\mathcal{C}}$ is determined by a trivialization of ΔE which differs from the trivialization Δs by a quadratic map τ . The equation

$$\tau(x) \circ \tau_{x,x} = 1_{X_x X_x}$$

in $\text{Isom}_{\mathcal{C}}(X_x X_x, X_x X_x)$ proves that the the quadratic map τ associated to the canonical alternating structure on E determined as in [6] §5 by the identity arrow $1_{X_x X_x}$ on $X_x X_x$ is the opposite of the braiding automorphism $\tau_{x,x}$ of $1_{X_x X_x}$.

Remark 4.2 This proposition translates in cohomological terms to the assertion that there exists, for any pair of abelian groups (or sheaves of abelian groups) B and A , a diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Ext}^2(B, A) & \longrightarrow & H^4(K(B, 2), A) & \longrightarrow & \text{Hom}(\Gamma_2(B), A) & (4.2) \\ & & \parallel & & \downarrow s & & \downarrow \delta & \\ & & \text{Ext}^2(B, A) & \longrightarrow & G & \longrightarrow & \text{Ext}^1(L\Lambda^2 B, A) \end{array}$$

with exact lines, and for which the left-hand square commutes, and the right-hand one anti-commutes. Here G is the kernel of the composite map

$$H^3(K(B, 1), A) \longrightarrow \text{Hom}(H_3(B), A) \longrightarrow \text{Hom}(\Lambda^3 B, A) \quad (4.3)$$

determined by the universal coefficient theorem and the iterated Pontrjagin product map $\Lambda^3 B \longrightarrow H_3(B)$. The vertical maps S and δ are respectively the suspension map and the boundary map induced on Ext^1 by the distinguished triangle

$$L\Gamma_2 B \longrightarrow B \otimes B \longrightarrow L\Lambda^2 B \xrightarrow{+1} L\Gamma_2 B[1] \quad (4.4)$$

Finally the lower horizontal exact sequence is described in detail, in geometric terms, in [6]. Its constituents are low degree terms in the spectral sequence

$$E_2^{p,q} = \text{Ext}^p(L\Lambda^q B, A) \implies H^{p+q}(K(B, 1), A)$$

associated in cohomology to the homological spectral sequence (1.10) of [7].

Passing now from categories to 2-categories, the situation is governed, when the intermediate homotopy group $\pi_1(\mathcal{C})$ of the monoidal 2-category \mathcal{C} is trivial, by the diagram

$$\begin{array}{ccccc}
\longrightarrow & \text{Ext}^3(B, A) & \longrightarrow & H^5(K(B, 2), A) & \longrightarrow & \text{Ext}^1(L\Gamma_2(B), A) & (4.5) \\
& \parallel & & \downarrow s & & \downarrow \delta & \\
\longrightarrow & \text{Ext}^3(B, A) & \longrightarrow & K & \longrightarrow & \text{Ext}^2(L\Lambda^2 B, A) &
\end{array}$$

Let us briefly review the structure of the group $H^4(K(B, 1), A)$ which classifies such 2-categories \mathcal{C} [6] §8. This group is endowed with a 4-term filtration, so that its subgroup K displayed above can no longer be expressed (as the corresponding group G in diagram (4.2)), as the kernel of some edge-homomorphism analogous to the map (4.3). Instead, K may be described as follows: to any element of $H^4(K(B, 1), A)$, representing a given 2-category \mathcal{C} , is associated a so-called weak (2,2)-extension $\mathcal{E}_{\mathcal{C}} = \mathcal{E}$, whose fiber above a section (x, y) of $B \times B$, is defined by the formula

$$\mathcal{E}_{x,y} = \text{Isom}_{\mathcal{C}}(X_y X_x, X_x X_y)$$

together with appropriate pair of partial tensor laws determined by the hexagon 2-arrows in \mathcal{C} . For the element in question to live in the subgroup K of $H^4(K(B, 1), A)$, it is necessary that a certain quadrilinear map $\psi_{\mathcal{C}} : \Lambda^4 B \longrightarrow A$, and an induced alternating triextension $\Phi_{\mathcal{C}} \in \text{Ext}^1(L\Lambda^3 B, A)$ both be trivial. In that case \mathcal{E} is endowed with a genuine alternating (2,2)-extension structure, so that it determines as required by diagram (4.5) an element in $\text{Ext}^2(L\Lambda^2 B, A)$. As explained in [6] §8, the trivialization of this (2,2)-alternating biextension ensures that the monoidal 2-stack \mathcal{C} is strict Picard (and so is equivalent to the trivial monoidal category determined by the pair of groups B and A in the 2-category case). Since such strict Picard stacks are classified by elements of the group $\text{Ext}^3(B, A)$, the present discussion gives a complete geometric interpretation of the lower line of diagram (4.5). The upper line in the diagram was introduced in (3.5) above, where it was described in similar geometrical terms.

The comparison between the A -torsor with Σ -structure $\mathcal{S}_{\mathcal{C}}$ (3.6) associated to a braided 2-stack \mathcal{C} , and the alternating (2,2)-biextension $\mathcal{E}_{\mathcal{C}}$ associated to the underlying monoidal 2-stack may now be carried out along the same lines as in the corresponding discussion in the 1-category case. The given braiding morphisms $\tau_{x,y}$ on \mathcal{C} determine a global section of the A -gerbe $\mathcal{E}_{\mathcal{C}}$. The axioms satisfied by the hexagon 2-arrows in \mathcal{C} imply that this global section is compatible with the partial group laws on $\mathcal{E}_{\mathcal{C}}$, and therefore trivializes $\mathcal{E}_{\mathcal{C}}$ as a (2,2)-extension. In particular, $\mathcal{E}_{\mathcal{C}}$ is represented by a class which lives in the subgroup K of $H^4(K(B, 1), A)$, since this is where genuine (as opposed to weak) (2,2)-biextensions live. The exactness of the sequence

$$\cdots \rightarrow \text{Ext}^1(L\Gamma_2(B), A) \rightarrow \text{Ext}^2(L\Lambda^2 B, A) \rightarrow \text{Ext}^2(B \overset{L}{\otimes} B, A) \rightarrow \text{Ext}^2(L\Gamma_2(B), A) \rightarrow \cdots \quad (4.6)$$

associated to the distinguished triangle (4.4) makes it clear that the alternating structure on $\mathcal{E}_{\mathcal{C}}$ is determined by a trivialization of its restriction $\Delta\mathcal{E}_{\mathcal{C}}$ above the diagonal in B . Such a restriction to the diagonal of a (2,2)-extension \mathcal{E} is an A -gerbe on B canonically endowed with

a structure which we might call a 2- Σ structure, and which is classified up to equivalence by the group $\text{Ext}^2(L\Gamma_2 B, A)$. The complete definition of such a 2- Σ structure may be obtained along the same lines as that of alternating (2,2)-extensions in [6], by examining the dévissage of this Ext^2 group associated to the distinguished triangle

$$LSym^2 B \longrightarrow L\Gamma^2 B \longrightarrow B \otimes^L \mathbf{Z}/2 \xrightarrow{+1} LSym^2 B [1]$$

We do not work this out in detail here, since such a discussion was already carried out in a related context in [6] §8. We simply recall from [6] that the sought-after trivialization of $\Delta\mathcal{E}_{\mathcal{C}}$, which determines the alternating structure on $\mathcal{E}_{\mathcal{C}}$, is given by the identity map

$$1_{X_x X_x} : X_x X_x \longrightarrow X_x X_x$$

viewed as an object in $(\Delta\mathcal{E}_{\mathcal{C}})_x = \text{Isom}(X_x X_x, X_x X_x)$ for a varying section $x \in B$. The A -torsor with Σ -structure which describes this alternating structure on the torsor $\mathcal{E}_{\mathcal{C}}$ (trivialized by the given braiding arrows $\tau_{x,y}$ in \mathcal{C}) is, up to a sign depending on our conventions, the A -torsor $\mathcal{S}_{\mathcal{C}}$ of (locally defined) 2-arrows S_x (3.6) which compare the induced trivialization $\tau(x)$ of $\Delta\mathcal{E}_{\mathcal{C}}$ with the canonical trivialization $1_{X_x X_x}$. It follows from this discussion that the right-hand square in diagram (4.5) commutes up to sign, where the vertical map δ is defined either cohomologically as in (4.6), or geometrically as above.

5 Higher invariants: the higher Σ -structure torsors

We now briefly examine the invariants associated to higher homotopy types. We think of the homotopy types described by the groups $H^6(K(B, 2), A)$ as being represented by appropriately defined braided 3-categories \mathcal{C} (or rather, 3-groupoids since we assume that every n -arrow in \mathcal{C} is invertible up to an $n + 1$ -arrow) with the usual additional assumptions: the group $\pi_0(\mathcal{C})$ of isomorphism classes of objects is isomorphic to B , the group $\pi_3(\mathcal{C})$ of self 3-arrows of any identity 2-arrow is isomorphic to A , and the intermediate homotopy groups $\pi_i(\mathcal{C})$ are trivial. We will not spell out the braiding axioms here, but merely observe that one method for obtaining them would be to think of \mathcal{C} as a 5-category with a single object and a single 1-arrow. The multilinear objects attached, as in the lower dimensional case, to an appropriate filtration of $H^6(K(B, 2), A)$ are now the groups $\text{Hom}(\Gamma_3 B, A)$, $\text{Ext}^2(L\Gamma_2 B, A)$ and $\text{Ext}^4(B, A)$. We have seen that the group $\text{Ext}^2(L\Gamma_2 B, A)$ describes classes of what might be termed 2- Σ structures, the higher (1-categorical) analog of the Σ -structure A -torsors on B . We do not spell out here the definition of such structures, nor of the strict Picard 3-categorical structures classified by $\text{Ext}^4(B, A)$ since this is routine. Let us merely observe that 2- Σ -structures are to (ordinary) Σ -structures as their anti-symmetric analog, the alternating (2,2)-extensions of [6] are to (ordinary) alternating biextensions. The only essentially new feature which occurs when braided 3-categories are considered is thus the appearance of the degree three component $\Gamma_3 B$ of the divided power algebra $\Gamma_* B$ of B , together with the projection

$$H^6(K(B, 2), A) \longrightarrow \text{Hom}(\Gamma_3 B, A) \tag{5.1}$$

The latter map is readily understood from a topological point of view, since it is induced by the isomorphism $\Gamma_3 B \simeq H_6(K(B, 2))$ [13]⁷ theorem 21.1 determined by the divided power algebra structure of $H_*(K(B, 2))$, and by the universal coefficient theorem. In categorical terms, it can be understood by observing that in a braided 3-category, the two possible methods of construction of a Yang-Baxter 2-arrow described by diagrams (2.12, 2.13) no longer yield the same result, as in a braided 2-category. Instead, they can only be compared by a 3-arrow, which is part of the braiding axioms for 3-categories, and which we denote, for representative objects X_x, X_y, X_z in \mathcal{C} of the isomorphism classes $x, y, z \in B$, by $(YB)_{x,y,z}$. Such a 3-arrow corresponds to the degree 6 cell in Eilenberg-Mac Lane's chain complex for $K(B, 2)$ traditionally denoted by $[x|_2 y|_2 z]$. It corresponds, in Getzler-Jones' cell decomposition [15] §5.4 of the quotiented configuration space $\overset{\circ}{F}_3(2)$ of pairs of points in \mathbf{R}^3 , to the cell $[1||2||3]$. It now follows from the explicit computation of the elements of $H_6(K(B, 2))$ that the map (5.1) associates to the given braided 3-category \mathcal{C} the homomorphism

$$\Upsilon_{\mathcal{C}} : \Gamma_3 B \longrightarrow A \quad (5.2)$$

which sends an element $\gamma_3(x)$ to the 3-arrow $(YB)_{x,x,x}$. The latter specialized Yang-Baxter 3-arrow is, as required an element of the group A of self 3-arrows in \mathcal{C} from an appropriate trivial 2-arrow to itself.

The group $\Gamma_3 B$ differs from the previously encountered group $\Gamma_2 B$, since it is no longer generated by indecomposable elements of the form $\gamma_3(x)$, additional generators of the form $\gamma_2(x)y$ with $x, y \in B$ being required. The map (5.2) can therefore no longer be entirely described by the values of the elements $(YB)_{x,x,x} \in A$ for varying $x \in B$. Instead, one is also led to consider the composite map

$$\Gamma_2 B \otimes B \longrightarrow \Gamma_3 B \xrightarrow{\Upsilon_{\mathcal{C}}} A$$

determined by the algebra structure in $\Gamma_* B$. The latter may be viewed as a map

$$B \times B \longrightarrow A$$

which is quadratic in the first variable x and linear in the second variable y . It is constructed by assembling the three Yang-Baxter 3-arrows $(YB)_{x,x,y}, (YB)_{x,y,x}, (YB)_{y,x,x}$ into a self 3-arrow, in a manner which we will not make explicit here.

We now pass to the groups $H^7(K(B, 2), A)$, whose elements describes the braided 4-categories with trivial intermediate homotopy groups. There are now three corresponding multilinear invariants, which live in the groups

$$\text{Ext}^1(L\Gamma_3 B, A) \quad \text{Ext}^3(L\Gamma_2 B, A) \quad \text{Ext}^5(B, A)$$

While the last two represent some significant structure in the 4-stack case, they both vanish when 4-categories are considered, so we will restrict ourselves to a discussion of the geometric

⁷In [13], the standard degree i component $\Gamma_i B$ of the divided power algebra $\Gamma_* B$ is somewhat confusingly denoted $\Gamma_{2i} B$.

objects representing the remaining group $\text{Ext}^1(L\Gamma_3 B, A)$. This is in any case the most interesting of the three, since it involves the degree three functor $\Gamma_3 B$. This functor lives in the exact sequence

$$\Gamma_2 B \otimes B \longrightarrow \Gamma_3 B \longrightarrow B/3B \longrightarrow 0 \quad (5.3)$$

reminiscent of the short exact sequence

$$0 \longrightarrow \text{Sym}^2 B \longrightarrow \Gamma_2 B \longrightarrow B/2B \longrightarrow 0$$

by which one analyzes a Σ -structures in terms of the associated symmetric biextension [3] §8. In the present situation, the first arrow is the multiplication map considered above, and the second the map which sends $\gamma_3(x)$ to the class of $x \bmod (3B)$.

Since there exist two sorts of generators for $\Gamma_3 B$, the geometric structure classified by the group $\text{Ext}^1(L\Gamma_3 B, A)$ involves a pair of objects (E, L) where E an A -torsor on $B \times B$ and L is an A -torsor on B . When such a pair (E, L) arises from a braided 4-category \mathcal{D} with trivial intermediate homotopy groups, the corresponding A -torsor $L_{\mathcal{D}}$ is defined by

$$L_{\mathcal{D}} = \text{Isom}_{\mathcal{D}}(1, \Delta_{123}^*(YB))$$

Its fiber above $x \in B$ is the torsor of 4-arrows from the identity 3-arrow to $(YB)_{x,x,x}$. The fibre above $x, y \in B^2$ of the corresponding A -torsor $E_{\mathcal{D}}$ similarly consists of the torsor of 4-arrows in \mathcal{D} from the identity 3-arrow to the 3-arrow assembled from $(YB)_{x,x,y}$, $(YB)_{x,y,x}$ and $(YB)_{y,x,x}$ to which we previously alluded (without making it explicit) in the 3-category case.

Returning to the general situation, we now describe the additional structure to be imposed on the pairs (E, L) . The map from $\text{Ext}^1(\Gamma_3 B, A)$ to $\text{Ext}^1(L\Gamma_2 B \overset{L}{\otimes} B, A)$ induced the multiplication in $\Gamma_* B$ is the first projection from (E, L) to E , so that the first element of structure to be imposed on the pair (E, L) is that E be quadratic in the first variable and linear in the second. This means that for each $y \in B$, $E_{(-,y)}$, viewed as a torsor above B , is endowed with a Σ -structure, and that $E_{(x,-)}$, viewed for each $x \in B$ as an A -torsor on B , is a commutative extension of B by A . We also require the compatibility of these two structures, in other words that the partial multiplication map

$$E_{x,y_1} \wedge E_{x,y_2} \longrightarrow E_{x,y_1+y_2} \quad (5.4)$$

determined by the second variable group law respect the Σ -structures.

The next set of conditions on the pair (E, L) arises from the relations

$$\gamma_3(x + x') - \gamma_3(x) - \gamma_3(x') = \gamma_2(x) x' + \gamma_2(x') x \quad (5.5)$$

$$3\gamma_3(x) = \gamma_2(x) x \quad (5.6)$$

satisfied by the divided powers. The first of these relations translates into the requirement that there exists an isomorphism of A -torsors

$$\Lambda(L) \xrightarrow{\alpha} E \wedge s^* E \quad (5.7)$$

where s is the map which permutes the factors in B^2 , and $\Lambda(L)$ is the “first difference” A -torsor of L , defined above B^2 formula (3.3). Similarly, the relation (5.6) translates to the requirement that there exists an isomorphism of A -torsors

$$\Delta E \xrightarrow{\beta} L^3 \quad (5.8)$$

between the restriction of E to the diagonal and the contracted third power $L \wedge L \wedge L$ of the torsor L . It is obvious that such a requirement is satisfied by the pair $(E_{\mathcal{D}}, L_{\mathcal{D}})$ since sections of the restriction of $E_{\mathcal{D}}$ to the diagonal involve three copies of $(YB)_{x,x,x}$.

Consider the pullback

$$(\Lambda \times 1)^*(E \wedge s^*E) \xrightarrow{(\Lambda \times 1)^*\alpha} (\Lambda \times 1)^*(\Lambda(L))$$

by $(\Lambda \times 1)$ of the map α (5.7). The target of this map is the “first difference of the first difference” of L , which is canonically isomorphic to the “second difference” $\Theta(L)$ of L (3.2). By linearity of E in the first variable, the source of this map simplifies so that it reduces to an isomorphism

$$(\Lambda \times 1)^*E \longrightarrow \Theta(L) \quad (5.9)$$

Since E possesses a Σ -structure with respect to its first variable, and is linear with respect to the second one, the source of this map is canonically endowed with a triextension structure above $B \times B \times B$, and in fact one which is symmetric with respect to the interchange of the first two variables. In particular the A -torsor L is endowed *via* this isomorphism with a hypercube structure, since its second difference is multilinear⁸.

We now list the conditions which the maps α and β must satisfy. Observe that the target $\Theta(L)$ of the map (5.9) is invariant up to canonical isomorphism under any permutation of the three factors in B^3 . The canonical isomorphism $\Theta(L) \longrightarrow \sigma_{123}^* \Theta(L)$ induced by the cyclic permutation σ_{123} of the factors of B^3 transports to an isomorphism

$$\psi : (\Lambda \times 1)^*E \simeq \sigma_{123}^*((\Lambda \times 1)^*E)$$

It is convenient to consider this in the fiber above a general element $(x, y, z) \in B^3$. It can then be displayed as an isomorphism⁹

$$\psi_{x,y,z} : E_{x+y,z} E_{x,z}^{-1} E_{y,z}^{-1} \simeq E_{y+z,x} E_{y,x}^{-1} E_{z,x}^{-1} \quad (5.10)$$

Taking into account the maps $E_{z,x} E_{z,y} \simeq E_{z,x+y}$ and $E_{x,y} E_{x,z} \simeq E_{x,y+z}$ which expresses the linearity of E in the second variable, one therefore obtains an isomorphism ϕ between the requisite pullbacks of E , whose fiber $\phi_{x,y,z}$ above (x, y, z) is an isomorphism of A -torsors

$$E_{x+y,z} E_{z,x+y} E_{y,x} E_{x,y} \longrightarrow E_{y+z,x} E_{x,y+z} E_{z,y} E_{y,z} \quad (5.11)$$

⁸so that its third difference is trivial.

⁹The contracted product symbols \wedge will henceforth mostly be omitted.

We impose on the map α (5.7) the requirement that the following diagram commutes.

$$\begin{array}{ccc}
(E_{x+y,z} E_{z,x+y}) (E_{y,x} E_{x,y}) & \xrightarrow{\phi_{x,y,z}} & (E_{y+z,x} E_{x,y+z}) (E_{z,y} E_{y,z}) \\
\alpha_{x+y,z} \alpha_{x,y} \downarrow & & \downarrow \alpha_{x,y+z} \alpha_{y,z} \\
\Lambda(L)_{x+y,z} \Lambda(L)_{x,y} & \xrightarrow{\quad\quad\quad} & \Lambda(L)_{x,y+z} \Lambda(L)_{x,y}
\end{array} \tag{5.12}$$

the lower horizontal map being the (obvious) canonical isomorphism. A second constraint on the map (5.7) is much simpler to state. It is the requirement that the following commutative square, in which the horizontal maps are the canonical ones, also commutes:

$$\begin{array}{ccc}
\Lambda(L) & \xrightarrow{\cong} & s^* \Lambda(L) \\
\alpha \downarrow & & s^* \alpha \downarrow \\
E \wedge s^* E & \longrightarrow & s^* E \wedge E
\end{array} \tag{5.13}$$

A final axiom involves both the arrows α (5.7) and β (5.8). Observe first of all that the map ψ (5.10) specializes to

$$\psi_{y,y,x} : E_{2y,x} E_{y,x}^{-2} \simeq E_{x+y,y} E_{y,y}^{-1} E_{x,y}^{-1}$$

On the other hand, the given Σ -structure on the first variable of E determines an isomorphism [3] (6.3.2)

$$E_{2y,x} \simeq E_{y,x}^4$$

so that the map in question determines an isomorphism

$$E_{x+y,y} \simeq E_{x,y} E_{y,y} E_{y,x}^2$$

Exchanging the variables x and y this yields

$$E_{x+y,x} \simeq E_{y,x} E_{x,x} E_{x,y}^2$$

These two isomorphisms, and the linearity of E in the second variable therefore determine a composite map

$$E_{x+y,x+y} \simeq E_{x+y,x} E_{x+y,y} \simeq E_{x,x} E_{y,y} E_{x,y}^3 E_{y,x}^3$$

which may also be viewed as a map

$$\Lambda \Delta E \xrightarrow{\gamma} E^3 \wedge s^* E^3 \tag{5.14}$$

The final condition which we impose on the pair (E, L) is the commutativity of the following diagram in which the top horizontal arrow is the canonical one

$$\begin{array}{ccc}
\Lambda(L^3) & \xrightarrow{\cong} & \Lambda(L)^3 \\
\Lambda(\beta) \downarrow & & \downarrow \alpha^3 \\
\Lambda \Delta E & \xrightarrow{\gamma} & E^3 \wedge s^* E^3
\end{array} \tag{5.15}$$

We summarize the previous discussion as follows

Definition 5.1 *Let A and B be two abelian groups (or sheaves of abelian groups). A Γ_3 - A -torsor pair above B consists of an A -torsor L on B and an A -torsor E on $B \times B$ satisfying the following conditions:*

i) E is endowed with a Σ -structure with respect to the first variable and with a commutative extension structure with respect to the second. These conditions are compatible, in the manner previously described (5.4).

ii) There exists a pair of isomorphisms of A -torsors (5.7), (5.8)

$$\Lambda(L) \xrightarrow{\alpha} E \wedge s^*E \qquad \Delta E \xrightarrow{\beta} L^3$$

iii) The diagrams (5.12), (5.13), and (5.15) commute.

In order to obtain a complete proof that the category of such quadruples (E, L, α, β) is described by the truncated complex $t_{\leq 0} \mathrm{RHom}(L\Gamma_3(B), A[1])$, whose homology groups are the groups $\mathrm{Ext}^i(L\Gamma_3(B), A)$ for $i = 0, 1$, one would have to construct a simple realization of the object $L\Gamma_3(B)$, as was done in [3] §8 for the object $L\Gamma_2(B)$. We do not carry this out here, but observe the following fact, which provides, in view of the exactness of the sequence (5.3), very strong evidence for this assertion. For this reason, and for lack of a better name, we will call these quadruples “ Γ_3 - A -torsor pairs on B ”, or simply “ Γ_3 -torsor pairs”

Proposition 5.2 *Let (E, L, α, β) be a Γ_3 torsor pair for which there exists a global section of E , compatible with both the Σ -structure of E in the first variable and with the linearity in the second one. Then L is endowed with a group law, for which it is a commutative extension of B by A . Furthermore the pullback of the extension*

$$0 \longrightarrow A \longrightarrow L \longrightarrow B \longrightarrow 0$$

by the map $B \xrightarrow{3} B$ is canonically split.

Proof: The given trivialization of E induces by (5.7) a trivialization of the torsor $\Lambda(L)$, and this in turn determines a composition law on L which is compatible with the projection from L to B . The commutativity of diagram (5.12) implies that the induced trivialization of the source and target of the lower horizontal map in this diagram are compatible, a statement which is equivalent to the associativity of the corresponding composition law on L . Similarly, the commutativity of diagram (5.13) implies that the trivializations of the source and target of the upper horizontal map in the latter diagram are compatible, and this in turn is equivalent to the commutativity of the composition law on L . This shows that L is a commutative extension of B by A (since the existence of an inverse map on L , the existence of a unit element and the compatibility of the composition law in L with the inclusion of A in L follow automatically). Finally, the commutativity of diagram (5.15) implies that the trivialization of the A -torsor L^3 determined by the map β (5.8) is compatible with the group law on L^3 induced by the group law of L , and therefore splits the pushout L^3 of L by the map $A \xrightarrow{3} A$ as a group extension. This is equivalent to the assertion regarding the pullback of L by the map $B \xrightarrow{3} B$.

References

- [1] J. Baez and M. Neuchl, *Higher-dimensional algebra I: Braided Monoidal 2-Categories*, Adv. in Math. **121** (1996) 196-244.
- [2] J. Baez and J. Dolan, *Categorification*, these Proceedings.
- [3] L. Breen, *Fonctions thêta et théorème du cube*, Lecture notes in math. **980**, Springer Verlag (1983).
- [4] L. Breen, *Biextension alternées*, Compositio math. **63** (1987) 99-122.
- [5] L. Breen, *Classification of 2-gerbes and 2-stacks*, Astérisque **225**, Société mathématique de France (1994).
- [6] L. Breen, *Monoidal categories and multiextensions*, math.CT/9809104.
- [7] L. Breen, *On the functorial homology of abelian groups*, to appear in the J. of Pure and Applied Algebra.
- [8] H. Cartan, Séminaire H. Cartan E.N.S. 7e année: 1954/55, *Algèbres d'Eilenberg-MacLane et Homotopie*, Secrétariat Mathématique, Paris (1956).
- [9] S. Crans, *Generalized centers of braided and sylleptic monoidal 2-categories*, Advances in Math. **136** (1998), 183-223.
- [10] B. Day and R. Street, *Monoidal bicategories and Hopf algebroids*, Advances in Math. **129** (1997), 99-157.
- [11] P. Deligne, *La formule de dualité globale* exposé XVIII of *Théorie des Topos et Cohomologie Etale des Schémas* vol. 3 (SGA 4) Lecture notes in math. **305**, Springer Verlag (1973).
- [12] A. Dold and D. Puppe, *Homologie nicht-additiven funktoren. Anwendungen* Ann. Inst. Fourier **11** (1961), 201-312.
- [13] S. Eilenberg and S. Mac Lane, *On the groups $H(\Pi, n)$ I, II*, Ann. of Math. **58** (1953) 53-106, **60** (1954) 49-139.
- [14] G. Decker *The integral homology algebra of an Eilenberg-Mac Lane Space*, Ph. D. thesis, University of Chicago (1974).
- [15] E. Getzler and J.D.S Jones, *Operads, homotopy algebra, and iterated integrals for double loop spaces*, preprint
- [16] A. Grothendieck, *Séminaire de Géométrie Algébrique du Bois-Marie, 1967-69 (SGA7) I* Lecture notes in math. **288**, Springer-Verlag (1972).
- [17] A. Hischowitz and C. Simpson, *Descente pour les n -champs*, preprint (1998), math.AG/9807049.

- [18] A. Joyal and R. Street, *Braided tensor categories*, Advances in Math. **102** (1993) 20-78.
- [19] M. Kapranov and V. Voevodsky, *2-categories and Zamolodchikov equations*, Proc. Symp. Pure Math. **56** (2) (1994) 177-259.
- [20] D. Mumford, *Geometric Invariant Theory*, Ergebnisse, N.S. **34**, Springer-Verlag 1965.
- [21] M. Neuchl, Ph.D. thesis, Munich (1997).
- [22] C. Simpson, *A closed model structure for n -categories, internal Hom, n -stacks and generalized Seifert-Van Kampen*, preprint (1997), available as alg-geom/9704006.
- [23] R. Street, *The role of Michael Batanin's monoidal globular categories*, these Proceedings.