THE GRAFTING MAP OF TEICHMÜLLER SPACE

Kevin P. Scannell Dept. of Mathematics and Computer Science St. Louis University St. Louis, MO 63103

> Michael Wolf* Dept. of Mathematics Rice University Houston, TX 77251

> > October 11, 1998

§1. Introduction.

1.1 Statement and Context.

One of the underlying principles in the study of Kleinian groups is that aspects of the complex projective geometry of quotients of $\hat{\mathbf{C}}$ by the groups reflect properties of the threedimensional hyperbolic geometry of the quotients of \mathbf{H}^3 by the groups. Yet, even though it has been over thirty-five years since Lipman Bers wrote down a holomorphic embedding of the Teichmüller space of Riemann surfaces in terms of the projective geometry of a Teichmüller space of quasi-Fuchsian manifolds, no corresponding parametrization in terms of the three-dimensional hyperbolic geometry has been presented. One of the goals of this paper is to give such a parametrization. This parametrization is straightforward and has been expected for some time ([Ta97], [Mc98]): to each member of a Bers slice of the space QF of quasi-Fuchsian 3-manifolds, we associate the bending measured lamination of the convex hull facing the fixed "conformal" end.

The geometric relationship between a boundary component of a convex hull and the projective surface at infinity for its end is given by a process known as grafting, an operation on projective structures on surfaces that traces its roots back at least to Klein [Kl33;§50, p. 230], with a modern history developed by many authors ([Ma69], [He75], [Fa83], [ST83], [Go87], [GKM95], [Ta97], [Mc98]). The main technical tool in our proof that bending measures give coordinates for Bers slices, and the second major goal of this paper, is the completion of the proof of the "Grafting Conjecture". This conjecture states that for a fixed measured lamination λ , the self-map of Teichmüller space induced by grafting a surface along λ is a homeomorphism of Teichmüller space; our contribution to this argument is a proof of the injectivity of the grafting map. While the principal application

^{*}Partially supported by NSF Grants DMS 9626565, DMS 9707770 (SCREMS)

of this result that we give is to geometric coordinates on the Bers slice of QF, one expects that the grafting homeomorphism might lead to other systems of geometric coordinates for other families of Kleinian groups (see §5.2); thus we feel that this result is of interest in its own right.

A difficulty in proving injectivity results for maps of Teichmüller space is that Teichmüller space is a quotient space with no canonical sections; our approach is to choose a section in the space of metrics over Teichmüller space defined via harmonic maps. Indeed, our principal tool in proving the grafting conjecture is a study of the the differential equation governing the infinitesimal form of the energy density of such maps; this study is complicated somewhat by the grafted metrics having a mild singularity, and the infinitesimal form having a more serious singularity. Nevertheless, this equation is amenable to nearly a complete solution, and it is estimates based on this solution which are the technical linchpins of our argument.

We now state our results and methods more precisely. Throughout, S will denote a fixed differentiable surface which is closed, orientable, and of genus $g \ge 2$. Let T_g be the corresponding Teichmüller space of marked conformal structures on S, and let P_g denote the deformation space of (complex) projective structures on S (see §2 for definitions).

There are two well-known parametrization of P_g , each reflecting a different aspect of the general theory. The first uses the Schwarzian derivative of the developing map to obtain a quadratic differential on S, holomorphic with respect to the complex structure underlying the given projective structure. This identifies P_g with the total space of the bundle $Q_g \to T_g$ of holomorphic quadratic differentials over Teichmüller space. This identification is representative of the complex analytic side of the theory; see for instance [Ea81], [Gu81], [He75], [Kr69], [Kr71], [KM81], [MV94], [Sh87], [ST95].

The second parametrization is due to Thurston and is more geometric in nature. To describe it, fix a hyperbolic metric $\sigma \in T_g$, a simple closed geodesic $\gamma \subset S$ of length ℓ , and a positive real number s. Let

$$\tilde{A}_s = \{ (r, \theta) \in \mathbb{C}^* : |\theta - \pi/2| \le s \}$$

and

$$A_s = \tilde{A}_s / \langle z \mapsto e^\ell z \rangle.$$

Of course, if $s \ge 2\pi$, we must interpret the projective structure on A_s as being defined by a developing map which is no longer an embedding; in any case, we call A_s a (projective) *s-annulus*. A new projective structure on S is defined by cutting the original hyperbolic surface (S, σ) open along the simple closed curve γ and gluing in A_s . This is the grafting operation; it provided the first examples [Ma68] of projective structures for which the developing map is not a covering of its image. Grafting extends by continuity from pairs (γ, s) to general measured laminations, defining a map $\Theta : \mathcal{ML} \times T_g \to P_g$. Thurston has shown (in unpublished work) that Θ is a homeomorphism (see [KT92], [La92]).

A natural problem is to understand how these geometric and complex analytic aspects interact. For instance, a measured lamination $\lambda \in \mathcal{ML}$ defines a slice $\Theta(\{\lambda\} \times T_g) \subset P_g$; following this inclusion with the projection $P_g \to T_g$ defines a self-map of Teichmüller space $\operatorname{Gr}_{\lambda}: T_g \to T_g$. Our main result can be stated concisely as follows:

Theorem A. Gr_{λ} is a homeomorphism.

This result was obtained in special cases by McMullen [Mc98] (one-dimensional Teichmüller spaces), and Tanigawa [Ta97] (for integral points of \mathcal{ML} , using a result of Faltings [Fa83]). Our result will hold for all elements of \mathcal{ML} and all Teichmüller spaces of finitely punctured Riemann surfaces of finite genus. (For the sake of expositional ease, we write the proof for Teichmüller spaces of closed Riemann surfaces, but the extension to Teichmüller spaces of finitely punctured surfaces is mostly a matter of additional notation: see the remark at the end of §4.)

Theorem A allows one to understand various complex analytic constructions in the theory of Teichmüller spaces and Kleinian groups in terms of measured geodesic laminations and the grafting construction. As an example, we obtain the following corollary in §5.1:

Corollary. Let B_Y be a Bers slice with fixed conformal structure Y, and define a map $\beta : B_Y \to \mathcal{ML}$ which assigns the bending lamination on the component of the convex hull boundary facing Y. Then β is a homeomorphism onto its image.

The space of projective structures is intimately related with the space of locally convex pleated maps of \tilde{S} into \mathbb{H}^3 (as detailed for instance in [EM87]). The dual notions are explored in [Sc96], where it is shown that P_g classifies causally trivial de Sitter structures on $S \times \mathbb{R}$; here the grafting operation corresponds to a "stretching" of the causal horizon. We give an application of Theorem A to this situation in §5.3.

Finally, in [Mc98] McMullen observes that Theorem A follows from the conjectural rigidity of hyperbolic cone 3-manifolds (see [HK98]); hence our result can be viewed as further positive evidence for the validity of this conjecture.

1.2 Outline of the Argument. It has been shown that $\operatorname{Gr}_{\lambda}$ is real analytic [Mc98] and proper [Ta97], therefore it suffices, as T_g is a cell, to prove local injectivity. For simplicity, assume the measured lamination λ is given by some simple closed curve γ on S and a nonzero transverse measure $s \in \mathbf{R}_+$. Fix a hyperbolic metric $\sigma_0 \in T_g$ and a small deformation σ_t of σ_0 with the property that the grafted surfaces are conformal; i.e. $\operatorname{Gr}_{\lambda}(\sigma_0) = \operatorname{Gr}_{\lambda}(\sigma_t)$ in T_g . Each grafted surface may be equipped with a $C^{1,1}$ metric $\operatorname{gr}(\sigma_t)$ which is flat on the inserted cylinder and hyperbolic elsewhere (see §2.2). Thus, it is a consequence of the singular harmonic maps theory of [GS92] that for every t there is a unique harmonic map $w_t : (S, \operatorname{gr}(\sigma_t)) \to (S, \operatorname{gr}(\sigma_0))$ homotopic to the identity; indeed we may adjust the metrics by an isotopy and assume that w_t is the identity map (it is straightforward that w_t is a homeomorphism). After this normalization, the two boundary curves of the inserted cylinder may move around S as t varies; the variation vector fields V^{ℓ} and V^r so-defined (one for each of the two boundary components γ^{ℓ} and γ^r of the inserted cylinder) are key pieces of information in the proof.

With this setup, the proof proceeds by first recognizing that the conformal factor $\mathcal{H}_t = \frac{gr(\sigma_0)}{gr(\sigma_t)}$ has a second role as the holomorphic energy density of the harmonic map w_t ; in that second role, it satisfies the Bochner equation (given as equation (3.1.3) below). It turns out to be easier to analyze the linearized equation for \mathcal{H} at t = 0 (equation (3.1.5)).

Our strategy for solving (3.1.5) is straightforward. We think of (3.1.5) as representing two different equations; the first on the open inserted flat cylinder S_0 (where K = 0) and

the second on the cut-open hyperbolic surface S_{-1} (where K = -1). A general solution to the first equation can be found easily. To study the solution to the second equation and the global solution to (3.1.5) (note that the complete equation (3.1.5) contains a term given as a measure $2\dot{K}$ supported on γ^{ℓ} and γ^{r}), we begin with an observation: the normal derivative of $\dot{\mathcal{H}}$ across the boundary curves γ^{ℓ} and γ^{r} appears as the inhomogeneous term in an ordinary differential equation (see §2.4) which can be solved for the variational fields V^{ℓ} and V^{r} . These vector fields V^{ℓ} and V^{r} in turn determine the normal derivatives of the global solution $\dot{\mathcal{H}}$ as computed from the hyperbolic side S_{-1} . Finally, integrating $\dot{\mathcal{H}}\Delta\dot{\mathcal{H}}$ by parts on S_{-1} and using our knowledge of the boundary terms forces $\dot{\mathcal{H}}$ to vanish identically. This then implies that the original metrics, σ_{t} and σ_{0} , are infinitesimally conformal, proving the desired local injectivity.

It is helpful in understanding the overall argument to note that we ignore the fact that the length of the inserted cylinder (whose length is always denoted s) is constant in t until the very end of the proof. This is discussed in some detail in §3.3.

The case where λ is a general measured lamination and not just a simple closed curve (or a system of disjoint simple closed curves) follows from approximating the general lamination by simple closed curves, approximating the conformal deformation $Gr_{\lambda}(\sigma_t)$ by quasi-conformal deformations $Gr_{s_m\gamma_m}(\sigma_t)$, and then extending the previous argument for the simple closed curves and conformal deformations to find identities involving only quantities that are continuous on the space \mathcal{ML} .

Acknowledgments. The authors wish to acknowledge several pleasant and very useful conversations with John Polking on regularity issues, with Robert Hardt on properties of harmonic maps to singular spaces, and with Jim Anderson on his and Dick Canary's work on limits of Kleinian groups.

\S **2.** Notation and Background.

2.1 Teichmüller Space, Bers embedding. Let S denote a smooth surface of genus $g \geq 2$, and let $\mathcal{M}_{-1} = \mathcal{M}_{-1}(S)$ denote the space of metrics $\rho |dw|^2$ on S with Gaussian curvature identically -1. The group Diff_o of diffeomorphisms of S homotopic to the identity acts on \mathcal{M}_{-1} by pullback: if $\phi \in \text{Diff}_o$, then $\phi \cdot \rho = \phi^* \rho$. We define the Teichmüller space of genus g, T_g , to be the quotient space $T_g = \mathcal{M}_{-1}/\text{Diff}_o$, i.e., equivalence classes of metrics in \mathcal{M}_{-1} under the action of Diff_o. A metric (S, ρ) represents a conformal class of metrics on S, hence a Teichmüller equivalence class of Riemann surfaces. Let $QD(\sigma)$ denote the 3g-3 dimensional complex vector space of holomorphic quadratic differentials on (S, ρ) .

There are a number of continuous and real-analytic parametrizations of the Teichmüller space T_g and one complex analytic parametrization given by Lipman Bers [Be64]. The Bers embedding, as it is usually known (see [Na88] for a comprehensive account), is defined as follows. Fix a point Y in T_g . Then, for any (variable) point $X \in T_g$, consider the quasi-Fuchsian manifold Q(X, Y) with conformal boundaries X and Y and fundamental group $\Gamma(X, Y)$. There is a simultaneous uniformization homeomorphism $F : \hat{\mathbf{C}} \to \hat{\mathbf{C}}$ of the sphere $\hat{\mathbf{C}}$ which does the following: 1) it equivariantly and conformally maps the unit disk Δ to the universal cover of Y, 2) it equivariantly and quasi-conformally maps the complement Δ^* of the unit disk to the universal cover of X, and 3) it conjugates $\Gamma(Y, Y)$ to $\Gamma(X, Y)$. As $F \mid_{\Delta}$

is conformal, we may take its Schwarzian derivative, say $\mathcal{S}(F \mid_{\Delta}) = \Psi_X$. The holomorphic function Ψ_X descends to a holomorphic quadratic differential on the Riemann surface Y: the correspondence $X \in T_g \mapsto \Psi_X \in QD(Y)$ is the Bers embedding $B_Y : T_g \to QD(Y)$. As the name suggests, it is an embedding [Be64] of the 3g – 3-dimensional Teichmüller space T_g into the 3g – 3-dimensional complex vector space QD(Y); the point Y maps to the origin, and it follows from results of Nehari [Ne49] that the image is contained in a ball of radius 6 and contains a ball of radius $\frac{3}{2}$.

Within the space QF of quasi-Fuchsian manifolds, the family $\{Q(X,Y)|X \in T_g\}$ is known as the Bers slice of QF based at Y.

2.2 Grafting, Thurston Metric. Recall that a *(complex) projective structure* on S is a maximal atlas of charts from S into $\mathbb{C}P^1$ such that all transition maps are restrictions of elements of $PSL(2,\mathbb{C})$ (i.e. a $(PSL(2,\mathbb{C}),\mathbb{C}P^1)$ -structure in the sense of Thurston). Such a structure yields in the usual way a holonomy representation $hol : \pi_1(S) \to PSL(2,\mathbb{C})$ and an equivariant developing map $dev : \tilde{S} \to \mathbb{C}P^1$. We will write P_g for the moduli space of projective structures on S (as defined and topologized, for instance, in [Go88]).

Let S denote the set of isotopy classes of essential simple closed curves on S. There is a well-defined intersection pairing $i : S \times S \to \mathbb{Z}$ given by the minimum number of intersection points among pairs of representative curves in the isotopy classes. This in turn defines an embedding of $\mathbb{R}_+ \times S$ into \mathbb{R}^S by sending a weighted simple closed curve (s, γ) to the S-tuple $(s \cdot i(\gamma, \alpha))_{\alpha \in S}$. The space of measured laminations \mathcal{ML} is defined to be the closure of $\mathbb{R}_+ \times S$ in \mathbb{R}^S . For simplicity, a measured lamination coming from a pair (s, γ) will be denoted $s\gamma$.

In the presence of a hyperbolic structure on S, it is typical to define measured laminations in terms of geodesic laminations equipped with a measure on transverse arcs (see [Th82] or [Bo88] for more details). We can also use a hyperbolic structure on S to define a notion of the length $L(\lambda)$ of a measured lamination λ : one defines $L(s\gamma)$ to be the product of s and the hyperbolic length of γ on S, and then extends $L : \mathbb{R}_+ \times S \to \mathbb{R}_+$ to all of \mathcal{ML} by continuity (see e.g. [Ke85]).

In §1, grafting was defined in terms of a map $\Theta : \mathcal{ML} \times T_g \to P_g$; for laminations in the subset $\mathbb{R}_+ \times S$ of weighted simple closed curves the projective structure $\Theta(s\gamma, \sigma)$ was defined by gluing together the Fuchsian projective structure associated to σ and a projective s-annulus along γ . The proof that Θ extends continuously to all of $\mathcal{ML} \times T_g$ can be found in [KT92].

In order to understand the surjectivity of Θ , let us briefly recall the *canonical stratification* associated to a projective structure (originally due to Thurston – see also [KP94], [Ap88], [Sc96], [KT92]). First note that, via the developing map dev, \tilde{S} inherits a notion of open round ball from $\mathbb{C}P^1$. Furthermore, also using dev, we can pull back the usual metric on $\mathbb{C}P^1$ to an (incomplete) metric on \tilde{S} – the metric completion depends only on the projective structure and is called the *Möbius completion* of \tilde{S} [KP94]. The closure of an open round ball in the Möbius completion is conformally equivalent to compactified hyperbolic space $\mathbb{H}^2 \cup S^1_{\infty}$, so the usual notion of "hyperbolic convex hull" transfers. Thus,

given an open round ball, we write C(U) for the intersection of U and the convex hull of $\overline{U} \setminus U$ in \overline{U} . The key observation is the following:

Lemma 2.2.1. [KP94] For every $p \in \tilde{S}$, there is a unique open round ball U_p such that $p \in C(U_p)$.

The sets U_p given by the lemma are called *maximal balls*, and define a stratification of \tilde{S} into the sets $C(U_p)$ (this descends in turn to a stratification of S). It is easy to verify that in the case of a projective structure obtained by grafting along $\lambda \in \mathcal{ML}$, this stratification is the basically the same as the one given by the leaves and complementary regions of λ .

We also obtain a canonical Riemannian metric defined to be the restriction to $C(U_p)$ of the hyperbolic metric on the open round ball U_p [KP94]. We call this metric the grafted metric or the Thurston metric; if the projective structure is obtained by grafting the hyperbolic surface σ along the measured lamination λ (i.e. $\Theta(\lambda, \sigma)$) we write this metric as $gr_{\lambda}(\sigma)$. Chasing through the definitions in the case of grafting along a weighted simple closed curve $s\gamma$, one sees that $gr_{\lambda}(\sigma)$ coincides with σ on $S \setminus \gamma$ and is flat on the inserted annulus.

2.3 Harmonic maps from surfaces. Let $(M, \sigma |dz|^2)$ and $(N, \rho(w)|dw|^2)$ denote M and N equipped with smooth Riemannian structures; here z refers to a local conformal coordinate on the surface M, and w refers to a local conformal coordinate on the surface N. For a Lipschitz map $w : (M, \sigma |dz|^2) \to (N, \rho(w)|dw|^2)$, we define the energy $E(w; \sigma, \rho)$ of the map w to be

$$\begin{split} E(w;\sigma,\rho) &= \int_M \frac{1}{2} \|dw\|^2 dv(\sigma) \\ &= \int_M \frac{1}{\sigma(z)} \{ \|w_* \partial_z\|_\rho^2 + \|w_* \partial_{\bar{z}}\|_\rho^2 \} \sigma(z) dz d\bar{z} \end{split}$$

Evidently, while the total energy depends upon the metric structure of the target surface (N, ρ) , it only depends upon the conformal structure of the source $(M, \sigma(z))$.

A critical point of this functional is called a harmonic map. We will be interested in the situation where M = N = S, a fixed surface of finite analytic type, with a fixed homotopy class $w_0 : S \to S$ of maps into the target S, where (S, ρ) is possibly singular, but non-positively curved in the sense of Alexandrov. In that case, (see [GS92;Lemma 1.1, Theorem 2.3]) there is a unique (if $w_*(\pi_1 M)$ is non-abelian) harmonic map $w(\sigma) : (S, \sigma) \to$ (S, ρ) in the homotopy class of w_0 ; in the next section, we will specialize to a case where we will find additional smoothness for w.

For harmonic maps $w : (\mathcal{R}, \sigma) \to (N, \rho)$ from a Riemann surface \mathcal{R} to a smooth target, one can characterize the harmonicity of w in terms of conformal objects on \mathcal{R} . The pullback metric $w^*\rho$ decomposes by type as

$$\begin{split} w^*\rho &= \langle w_*\partial_z, w_*\partial_z \rangle_\rho \, dz^2 + (\|w_*\partial_z\|_\rho^2 + \|w_*\partial_{\bar{z}}\|_\rho^2)\sigma dz d\bar{z} + \langle w_*\partial_{\bar{z}}, w_*\partial_{\bar{z}} \rangle_\rho \, d\bar{z}^2 \\ &= \varphi dz^2 + \sigma e(w) dz d\bar{z} + \bar{\varphi} d\bar{z}^2 \end{split}$$

where $e = \frac{1}{2} ||dw||^2$ is the energy density of the map w. It is easy to show (see [Sa78]) that if w is harmonic then $\Phi = \varphi dz^2$ is a holomorphic quadratic differential on \mathcal{R} . In particular, Schoen [Sc84] has emphasized that even for harmonic maps to singular metric spaces (S, ρ) , it is a consequence of Weyl's lemma that the differential $\Phi = \varphi dz^2 = \langle w_* \partial_z, w_* \partial_z \rangle_{\rho} dz^2$ is holomorphic.

The expression $\mathcal{H} = ||w_*\partial_z||_{\rho}^2$ plays a special role in harmonic maps between surfaces (see, for instance [Wo91a]). First, we can rewrite the pullback metric $w^*\rho$ entirely in terms of $\Phi = \varphi dz^2$ and \mathcal{H} as follows:

$$w^*\rho = \varphi dz^2 + (\mathcal{H} + \frac{|\Phi|^2}{\sigma \mathcal{H}})dz d\bar{z} + \bar{\varphi} d\bar{z}^2.$$

Moreover, the function $\mathcal{H} = \mathcal{H}(z)$ satisfies the Bochner equation (this is basically a Liouville equation for prescribed curvature, using the harmonic map gauge)

(2.3.1)
$$\Delta_{\sigma} \log \mathcal{H}(z) = -2K_{\rho}(w(z))\{\mathcal{H}(z) - \frac{|\Phi(z)|^2}{\sigma(z)\mathcal{H}(z)}\} + 2K_{\sigma}(z).$$

Here K_{ρ} and K_{σ} refer to the Gauss curvatures of (S, σ) and (S, ρ) , respectively, and we are stating the equation only in the context of smooth maps; we will later extend the meaning of this equation to the singular context which is our principal interest in this paper.

2.3.2 Smoothness of Harmonic Maps Families. We will be interested in harmonic maps between surfaces equipped with the grafted (Thurston) metrics; in particular, we will carefully study one-parameter families of such maps. This study relies on the background result that these maps are reasonably smooth, and that the family of maps is reasonably smooth in the family parameter, for a smooth family of grafted metrics. In this section, we establish these basic smoothness results: the proofs are completely straightforward generalizations of those found in the literature (see [Jo97], [EL81], [Sa78]), but as the precise versions we need do not seem to be present already in print, we include them here for the sake of completeness.

First let us record the regularity of the Thurston metrics; a proof can be found in [KP94].

Lemma 2.3.1. For $\lambda \in \mathcal{ML}$, the grafted metrics $gr_{\lambda}(\sigma)$ are of class $C^{1,1}$.

Next, we consider the regularity of an individual harmonic map $w : (S, \operatorname{gr}(\sigma_0)) \to (S, \operatorname{gr}(\sigma_1)).$

Lemma 2.3.2. There exists a harmonic map $w : (S, gr(\sigma_0)) \to (S, gr(\sigma_1))$ homotopic to the identity; this map is of class $C^{2,\alpha}$.

Proof: As S is compact, and $gr(\sigma_1)$ is an NPC space (see [GS92]), it is straightforward that there is an energy minimizer w in the given homotopy class. Then we are able to make considerable use of the literature: Theorem 2.3 of [GS92] then ensures that $w \in H^1(S, S)$

is locally Lipschitz. The rest of the proof is straightforward bootstrapping applied to the harmonic map equation (see, e.g. [Jo97], proof of Theorem 3.2.4). \Box

Finally, we come to the smoothness of the families of the maps. We begin by recording the fact that $gr_{\lambda}(\sigma_t)$ varies analytically in t, for an analytic family of hyperbolic metrics σ_t .

Lemma 2.3.3. [Mc98] Let $\{\sigma_t\}$ be an analytic family (in t) of hyperbolic metrics. Then the family $\{gr_{\lambda}(\sigma_t)\}$ of grafted metrics is also analytic in t.

We omit the proof. Consider such an analytic family $\{\operatorname{gr}_{\lambda}(\sigma_t)\}\$ and the family $\{w_t\}$ of harmonic maps $w_t : (S, \operatorname{gr}(\sigma_t)) \to (S, \operatorname{gr}(\sigma_0))\$ which we know to exist and be of class $C^{2,\alpha}$.

Lemma 2.3.4. The family $w_t : (S, gr(\sigma_t)) \to (S, gr(\sigma_0))$ of harmonic maps is analytic in t, for small values of t. Any individual map $w_t : (S, gr(\sigma_t)) \to (S, gr(\sigma_0))$ is a homeomorphism.

Proof: We mimic an allied proof in [EL81]: see also [Sa78]. Given such a family, the first variation at t = 0 of the tension $\tau_t = \tau(w_t)$ can be computed to be

(2.3.2)
$$\frac{d\tau}{dt} = \Delta \dot{w} + K_0 \dot{w} - \operatorname{Tr}\left(\frac{\partial \Gamma(t)}{\partial t}\right).$$

 $\Gamma(t)$ refers to the Christoffel symbols of the family $\operatorname{gr}(\sigma_t)$ and where we have simplified the formula considerably by applying it at t = 0, where $w_0 : (S, \operatorname{gr}(\sigma_0)) \to (S, \operatorname{gr}(\sigma_0))$ is the identity map.

We aim to apply the analytic implicit function theorem (see [Be77]): the formal setting is that we regard the tension τ as a functional

$$\tau: C^{2,\alpha}(S,S) \times (-\epsilon,\epsilon) \longrightarrow C^{0,\alpha}(T(S))$$

where $C^{0,\alpha}(T(S))$ denotes $C^{0,\alpha}$ sections of the tangent bundle to S, the map associates to a map $w \in C^{2,\alpha}(S,S)$ and a metric $\operatorname{gr}(\sigma_t)$ the tension field $\tau(w,\operatorname{gr}(\sigma_t))$ of the map $w: (M,\operatorname{gr}(\sigma_t)) \to (M,\operatorname{gr}(\sigma_0))$. This functional is evidently analytic in t, so our attention turns to formula (2.3.2): we assert that

$$\left\|\frac{d\tau}{dt}\right\| > 0$$

where the norm is that taken on functionals between $C^{2,\alpha}(S,S)$ and $C^{0,\alpha}(T(S))$. It is enough to prove that $(\Delta + K_0)$ is invertible on $C^{0,\alpha}(T(S))$; i.e. that given $f \in C^{0,\alpha}(T(S))$, there is a $u \in C^{2,\alpha}(S,S)$ so that $\Delta u + K_0 u = f$. As $K_0 \leq 0$, this result follows from standard estimates: e.g. [GT83, Theorems 8.3, 8.8] give estimates on $||u||_{W^{2,2}}$ in terms of $||f||_{L^2}$, and since dim_{**R**} S = 2, this yields a C^{α} estimate on u, with higher regularity following from bootstrapping as in Lemma 2.3.2.

That an individual map $w_t : (S, \operatorname{gr}(\sigma_t)) \to (S, \operatorname{gr}(\sigma_0))$ is a homeomorphism follows from the map w_t being a perturbation of the identity. \Box

Remark: Here we restrict to families of grafted metrics where the grafting locus λ remains fixed in t. If we were to vary the grafting locus $\lambda = \lambda_t$ in \mathcal{ML} , we would need to deal with issues arising from \mathcal{ML} having but a piecewise linear structure and not a differentiable structure.

2.4 Variation of geodesics.

This section contains a brief discussion of the equations governing the variation fields of a geodesic in family of conformally related Riemannian metrics. We begin by setting some notation. Consider a smooth family of Riemannian metrics g_t on S and a family of g_t -geodesics $\gamma_t : [0, 1] \to S$. We adopt Fermi coordinates along the curve γ_0 so that

$$g_0 = F(x_2)^2 dx_1^2 + dx_2^2$$

The geodesic equation for γ_t in these coordinates is given by

(2.4.1)
$$\gamma_{t,11}^k + \Gamma_{t,ij}^k (\gamma_t(x_1)) \gamma_{t,1}^i \gamma_{t,1}^j = 0$$

where $\Gamma_{t,ij}^k$ are the g_t -Christoffel symbols. We differentiate (2.4.1) in time t to obtain the following equation for the vector field $\dot{\gamma}^k \partial_k = \frac{d}{dt} \Big|_{t=0} \gamma_t^k \partial_k$

$$\dot{\gamma}_{11}^{k} + \left(\frac{d}{dt}\Big|_{t=0} \Gamma_{t,ij}^{k}(\gamma_{0}(x_{1}))\right) \gamma_{0,1}^{i} \gamma_{0,1}^{j} + D_{m} \Gamma_{0,ij}^{k}(\gamma_{0}(x_{1})) \dot{\gamma}^{m} \gamma_{0,1}^{i} \gamma_{0,1}^{j} \\ + \Gamma_{0,ij}^{k}(\gamma_{0}(x_{1})) \dot{\gamma}_{1}^{i} \gamma_{0,1}^{j} + \Gamma_{0,ij}^{k}(\gamma_{0}(x_{1})) \gamma_{0,1}^{i} \dot{\gamma}_{1}^{j} = 0.$$

In the Fermi coordinates chosen, we have that $\Gamma_{0,ij}^k = 0$ and for a constant speed geodesic, we have $\gamma_{0,1}^i = \ell \delta_1^i$ where ℓ is the length of the geodesic and δ_1^i is the Kronecker delta. Thus the previous equation simplifies to

(2.4.2)
$$\dot{\gamma}_{11}^k + \left(\frac{d}{dt}\Big|_{t=0}\Gamma_{t,11}^k\right)\ell^2 + \left(D_m\Gamma_{0,11}^k\right)\dot{\gamma}^m\ell^2 = 0.$$

We are principally interested in the normal component of the variation field $\frac{d}{dt}\gamma_t$, so we set k = 2 and compute

$$\Gamma_{t,11}^2 = \frac{1}{2} g_t^{2\alpha} \left(2\partial_1 g_{t,1\alpha} - \partial_\alpha g_{t,11} \right).$$

Moreover, we will be interested only in the situation where $g_t = \frac{1}{\mathcal{H}_t}g_0$ is a family of conformal metrics (see §3.1) and where g_0 , being written in Fermi coordinates, is diagonal; this also forces g_t to be diagonal which simplifies the above description to

$$\Gamma_{t,11}^2 = -\frac{1}{2} \mathcal{H}_t \left(\partial_2 ((F(x_2)^2)/\mathcal{H}_t) \right).$$

It is then straightforward to compute from this equation and from $\mathcal{H}_0 \equiv 1$ that

$$\begin{array}{c} (2.4.3) \\ D_m \Gamma_{11}^2 = K \delta_m^2 \\ 9 \end{array}$$

and

(2.4.4)

$$\frac{d}{dt}\Big|_{t=0} \Gamma^{2}_{t,11} = -\frac{1}{2}\dot{\mathcal{H}} \left(\partial_{2} ((F(x_{2})^{2})/\mathcal{H}_{0}) - \frac{1}{2} \left\{ \left[-\partial_{2} ((F(x_{2}))^{2})\dot{\mathcal{H}}/\mathcal{H}_{0}^{2} + 2(F(x_{2}))^{2}(\partial_{2}\mathcal{H}_{0})\dot{\mathcal{H}}/\mathcal{H}_{0}^{3} \right] \right\} + \frac{1}{2} (F(x_{2}))^{2} \frac{\partial_{2}\dot{\mathcal{H}}}{\mathcal{H}_{0}^{2}} = \frac{1}{2} \partial_{2}\dot{\mathcal{H}}$$

where the first and second terms vanish because $\partial_2 F(x_2) = 0$, and the third term vanishes because $\partial_2 \mathcal{H}_0 = \partial_2(1) = 0$.

We conclude from (2.4.2), (2.4.3) and (2.4.4) that the variational field $V = \frac{d}{dt} \Big|_{t=0} \gamma_t^2$ satisfies

(2.4.5)
$$V_{11} + K_0 V \ell^2 = -\frac{1}{2} \ell^2 \partial_2 \dot{\mathcal{H}}.$$

Remark: The reader should recognize how, in the case of the Thurston metric defined above where K_0 is a discontinuous function, the equation (2.4.5) is really a pair of equations for a single variational field V. That is, on the flat cylinder $K_0 \equiv 0$ while on the hyperbolic portion of the surface $K_0 \equiv -1$; in our solution for $\dot{\mathcal{H}}$ below, this is reflected in a jump in the normal derivative of $\dot{\mathcal{H}}$ across the two geodesics bounding the grafted cylinder.

$\S3$. The Case of Simple Closed Curves.

In §3, we prove the main theorem in the model case when the measured lamination is a weighted simple closed curve. We begin by describing the problem in terms of harmonic maps and deriving our basic equation of study (3.1.5). The proof effectively becomes a computation, which we undertake in §3.2. As noted earlier, our setup applies quite generally to families of grafted metrics in which the length of the inserted annulus is allowed to vary. We only use the information that this length is constant in t at the very end of the proof – this is the content of section §3.3.

We begin with a precise statement of our objective.

Theorem 3.1. (Model Case). Let S be a closed differentiable surface of genus g > 1, let γ be an essential simple closed curve on S and let $s \in \mathbf{R}_+$ be a positive real number. Then the grafting map $Gr_{s\gamma}: T_g \to T_g$ is a homeomorphism.

Proof of Theorem 3.1.

As discussed in $\S1.2$, we need only show

Lemma 3.2. The grafting map $Gr_{s\gamma}$ is locally injective.

Proof of Lemma 3.2. We suppose, in order to obtain a contradiction, that there is a hyperbolic surface (S, σ) and an approximating sequence $\langle (S, \sigma_n) \rangle \to (S, \sigma)$ of hyperbolic

surfaces so that $\operatorname{Gr}_{s\gamma}(\sigma_n) = \operatorname{Gr}_{s\gamma}(\sigma)$ for all n. By passing to a subsequence, and using the differentiability of $\operatorname{Gr}_{s\gamma} : T_g \to T_g$, we find a tangent vector $[\dot{\mu}] \in T_{[\sigma]}T_g$ so that $d\operatorname{Gr}_{s\gamma}(\sigma)[\dot{\mu}] = [0] \in T_{[\operatorname{Gr}_{s\gamma}(\sigma)]}T_g$.

With this in mind, it is psychologically convenient to solve instead a formally easier problem: we imagine a differentiable family $\langle (S, \sigma_t) \rangle$ of hyperbolic surfaces converging to $(S, \sigma) = (S, \sigma_0)$ with the property that the tangent vector to the family is given by

$$\frac{d}{dt}\Big|_{t=0} \langle (S,\sigma_t) \rangle = [\dot{\mu}] \in T_{\sigma_0} T_g$$

and that $\operatorname{Gr}_{s\gamma}(\sigma_t) = \operatorname{Gr}_{s\gamma}(\sigma_0)$ in T_q . We then seek a contradiction to this situation.

Our method is to use harmonic maps to "fix the gauge" in comparing the surfaces $\operatorname{Gr}_{s\gamma}(\sigma_t)$. In particular, we imagine $\operatorname{Gr}_{s\gamma}(\sigma_t)$ as being realized by a metric $\operatorname{gr}(\sigma_t)$ on the underlying differentiable surface S. Of course, we have a choice for these representative metrics, as the group Diff_o of diffeomorphisms isotopic to the identity acts on metrics on S, with the orbit of $\operatorname{gr}(\sigma_t)$ consisting of isometric metrics. However, by the results in §2, and because $\operatorname{gr}(\sigma_0)$ is non-positively curved, there is a unique harmonic map $w: (S, \operatorname{gr}(\sigma_t)) \to (S, \operatorname{gr}(\sigma_0))$ homotopic to the identity for any of our choices of $\operatorname{gr}(\sigma_t)$. In particular, we can choose this representative metric $\operatorname{gr}(\sigma_t)$ on S so that the identity map

$$\operatorname{id}: (S, \operatorname{gr}(\sigma_t)) \longrightarrow (S, \operatorname{gr}(\sigma_0))$$

is harmonic for all $t \ge 0$. Since conformal maps are always harmonic and our harmonic map is unique, we may conclude that the identity map above is both harmonic and conformal. Let

(3.1.1)
$$\mathcal{H}_t = \|\operatorname{id}_{z_t}\|^2 = \frac{\operatorname{gr}(\sigma_0)}{\operatorname{gr}(\sigma_t)} |\operatorname{id}_{z_t}|^2$$

denote the holomorphic energy density of the harmonic conformal map, where here we have snuck in the local coordinate convention that the metric $\operatorname{gr}(\sigma_t)$ admits an expansion in the local conformal coordinates z_t (themselves smooth in t) as $\operatorname{gr}(\sigma_t) = \operatorname{gr}(\sigma_t)|dz_t|^2$. Then because the identity map is conformal, we conclude that

(3.1.2)
$$\operatorname{gr}(\sigma_t)|dz_t|^2 = \frac{1}{\mathcal{H}_t}\operatorname{gr}(\sigma_0)|dz_0|^2.$$

Furthermore, because the identity map is harmonic, we apply the Bochner equation (2.3.1) to conclude that

(3.1.3)
$$\Delta_{\operatorname{gr}(\sigma_t)} \log \mathcal{H}_t = -2K_0(\operatorname{id}(z))\mathcal{H}_t + 2K_t(z).$$

We then use (3.1.2) to rewrite (3.1.3) as

$$\mathcal{H}_t \Delta_{\operatorname{gr}(\sigma_0)} \log \mathcal{H}_t = -2K_0(z)\mathcal{H}_t + 2K_t(z).$$
11

We then divide by \mathcal{H}_t to obtain the equation

(3.1.4)
$$\Delta_{\operatorname{gr}(\sigma_0)} \log \mathcal{H}_t = -2K_0 + \frac{2K_t}{\mathcal{H}_t}$$

which is the precursor to our basic equation of study. To obtain the basic equation of study, we differentiate equation (3.1.4) in time to obtain an equation for $\dot{\mathcal{H}} = \frac{d}{dt}\Big|_{t=0} \mathcal{H}_t$

$$\Delta_{\mathrm{gr}(\sigma_0)}\dot{\mathcal{H}}/\mathcal{H}_0 = 2\dot{K}/\mathcal{H}_0 - 2K_0\dot{\mathcal{H}}/\mathcal{H}_0^2$$

where \dot{K} denotes the measure $\frac{d}{dt}\Big|_{t=0} K_t$ (see extended discussion below). Since $\mathcal{H}_0 \equiv 1$ by construction, we summarize our equation as

(3.1.5)
$$\left(\Delta_{\operatorname{gr}(\sigma_0)} + 2K_0\right)\dot{\mathcal{H}} = 2\dot{K}.$$

This equation requires some discussion. The term 2K refers to a measure supported on the pair of images of the geodesic γ which bound the grafted cylinder. We can imagine \dot{K} being constructed as follows. Since we have a well-defined family of metrics $\operatorname{gr}(\sigma_t)$ each with a pair $\{\gamma_0^\ell, \gamma_0^r\}$ (left and right) boundaries of the grafted cylinder, we see that there is then a pair of well-defined families of curves $\{\gamma_t^\ell, \gamma_t^r\}$ of left and right boundaries of the grafted cylinders. These curves a priori may vary smoothly on the surface S, and form the frontier of the support of the function K_t which is identically -1 on the hyperbolic portion of $\operatorname{gr}(\sigma_t)$. Of course, on any open set \mathcal{O} which avoids the family γ_t for some interval of time $t \in [0, \epsilon)$, we have that $\frac{d}{dt}K_t = 0$ on \mathcal{O} . Thus, \dot{K} is supported only on the pair $\{\gamma_0^\ell, \gamma_0^r\}$. It is straightforward, but not required for our work here, to compute \dot{K} in terms of the variational vector fields $\{\frac{d}{dt} \mid_{t=0} \gamma_t^\ell, \frac{d}{dt} \mid_{t=0} \gamma_t^r\}$ along the curves $\{\gamma_0^\ell, \gamma_0^r\}$. In fact, our solution $\frac{d}{dt}\mathcal{H}_t$ to equation (3.1.5) is analytic away from $\{\gamma_0^\ell, \gamma_0^r\}$

In fact, our solution $\frac{d}{dt}\mathcal{H}_t$ to equation (3.1.5) is analytic away from $\{\gamma_0^\ell, \gamma_0^r\}$ and Lipschitz in a neighborhood of those curves. To see that, begin with equation (3.1.1) and observe that, by definition, the harmonic map id : $(S, \operatorname{gr}(\sigma_t)) \to (S, \operatorname{gr}(\sigma_0))$ is conformal hence (locally) complex analytic (in the local coordinates). Thus, the expression $\frac{d}{dt} \operatorname{id}_{z_t}$ is real analytic for an analytic path of metrics $\{\sigma_t\}$. This fact implies that when we take $\frac{d}{dt}\Big|_{t=0}$ $\mathcal{H}_t = \frac{d}{dt}\Big|_{t=0} \frac{\operatorname{gr}(\sigma_0)}{\operatorname{gr}(\sigma_t)}\Big|\operatorname{id}_{z_t}\Big|^2$, the only possibly non-analytic contribution comes from the term $\frac{d}{dt}\operatorname{gr}(\sigma_t)(z_t) = \frac{\partial}{\partial z}\operatorname{gr}(\sigma_0) \cdot \frac{d}{dt}z_t$. Finally, from Lemma 2.3.1, we see that $\frac{\partial}{\partial z}\operatorname{gr}(\sigma_0)$ is Lipschitz continuous in a neighborhood of $\{\gamma_0^\ell, \gamma_0^r\}$, and therefore, so is $\dot{\mathcal{H}}$. Thus, much depends on understanding the "jump" in derivatives of a solution $\dot{\mathcal{H}}$ across the curves $\{\gamma_0^\ell, \gamma_0^r\}$.

3.2 Computation of \mathcal{H} .

The goal of this section is a proof of

Lemma 3.3. Any function $\frac{d}{dt}\Big|_{t=0} \mathcal{H}_t$ which solves (3.1.5) and infinitesimally solves (3.1.2) must vanish identically on S. Thus id : $(S, gr(\sigma_t)) \to (S, gr(\sigma_0))$ is an isometry, up to order $\mathcal{O}(t^2)$.

Recall the basic plan from §1.2: We consider equation (3.1.5) as really a pair of equations, the first saying that $\dot{\mathcal{H}}$ is harmonic on the Euclidean cylinder, and the second saying that $\dot{\mathcal{H}}$ satisfies the linearized Liouville equation on the hyperbolic portion of $\operatorname{gr}(\sigma_0)$. We then solve for the general expression for a harmonic function on the cylinder, and this determines both boundary values for $\dot{\mathcal{H}}$ and normal derivatives $\partial_n \dot{\mathcal{H}}$ on the cylinder. These derivatives $\partial_n \dot{\mathcal{H}}$ on the cylinder can be used to find general solutions V^{ℓ} and V^r to (2.4.5) with $K \equiv 0$ which is compatible with our general solution to (3.1.5) on the cylinder, and then the version of (2.4.5) with $K \equiv -1$ gives a general expression for $\partial_n \dot{\mathcal{H}}$ as viewed from the hyperbolic side of $\operatorname{gr}(\sigma_0)$.

We then take this general solution to (3.1.5) and integrate $\dot{\mathcal{H}}\Delta_{\mathrm{gr}(\sigma_0)}\dot{\mathcal{H}}$ by parts to find that equation (3.1.5) forces $\dot{\mathcal{H}}$ to vanish identically. This will prove Lemma 3.3.

We now carry out this outline. To begin, write equation (3.1.5) as

$$(3.1.5)_0 \qquad \qquad \Delta_E \dot{\mathcal{H}} = 0$$

and

$$(3.1.5)_{-1} \qquad (\Delta_h - 2)\dot{\mathcal{H}} = 0$$

where Δ_E and Δ_h denote the Laplace-Beltrami operators on the Euclidean $(K \equiv 0)$ and hyperbolic $(K \equiv -1)$ open submanifolds of $(S, \operatorname{gr}(\sigma_0))$, respectively. Let us solve $(3.1.5)_0$ by writing the Euclidean grafted cylinder as $\{(x, y) \mid -\frac{s}{2} \leq x \leq \frac{s}{2}, 0 \leq y \leq \ell\}$ and then writing

$$\mathcal{H} = \Sigma a_n(x) \exp(2\pi i n y/\ell).$$

Then equation $(3.1.5)_0$ implies that

$$\Sigma\left(a_n''(x) - \frac{4\pi^2 n^2}{\ell^2}a_n(x)\right)\exp(2\pi i ny/\ell) = 0.$$

We conclude that

$$a_n(x) = \begin{cases} c_n \cosh \frac{2\pi n}{\ell} x + d_n \sinh \frac{2\pi n}{\ell} x & n \neq 0\\ c_0 x + d_0 & n = 0 \end{cases}$$

so that, for z = x + iy,

(3.2.1)
$$\dot{\mathcal{H}}(z) = c_0 x + d_0 + \Sigma' \left(c_n \cosh \frac{2\pi n}{\ell} x + d_n \sinh \frac{2\pi n}{\ell} x \right) \exp(2\pi i n y/\ell).$$

Since $\dot{\mathcal{H}}(z)$ is real, we find that

(3.2.2)
$$\bar{c}_n = c_{-n}$$

 $\bar{d}_n = -d_{-n}.$

When we specialize (3.2.1) to the boundaries $x = \pm \frac{s}{2}$ of the Euclidean cylinder, we obtain the expansions

$$(3.2.3) \qquad \begin{aligned} \dot{\mathcal{H}}\Big|_{x=-\frac{s}{2}} &= -c_0 \frac{s}{2} + d_0 + \Sigma' \left(c_n \cosh \frac{\pi ns}{\ell} - d_n \sinh \frac{\pi ns}{\ell} \right) \exp(2\pi i ny/\ell) \\ \dot{\mathcal{H}}\Big|_{x=\frac{s}{2}} &= c_0 \frac{s}{2} + d_0 + \Sigma' \left(c_n \cosh \frac{\pi ns}{\ell} + d_n \sinh \frac{\pi ns}{\ell} \right) \exp(2\pi i ny/\ell). \end{aligned}$$

As remarked in the last paragraph of §3.1, $\dot{\mathcal{H}}$ is real analytic away from the boundary of the inserted cylinder and Lipschitz across the boundary. In particular, the *x*-derivative of our general solution (from the cylinder side) exists; we compute it by differentiating (3.2.1):

$$\frac{\partial \dot{\mathcal{H}}}{\partial x} = \dot{\mathcal{H}}_x = c_0 + \Sigma' (\frac{2\pi n}{\ell} c_n \sinh \frac{2\pi n}{\ell} x + \frac{2\pi n}{\ell} d_n \cosh \frac{2\pi n}{\ell} x) \exp(2\pi i n y/\ell)$$

which we specialize to $x = \pm \frac{s}{2}$ to obtain

(3.2.4)
$$\begin{aligned} \dot{\mathcal{H}}_x\Big|_{x=-\frac{s}{2}} &= c_0 + \Sigma' \frac{2\pi n}{\ell} \left(-c_n \sinh \frac{\pi ns}{\ell} + d_n \cosh \frac{\pi ns}{\ell} \right) \exp(2\pi i ny/\ell) \\ \dot{\mathcal{H}}_x\Big|_{x=\frac{s}{2}} &= c_0 + \Sigma' \frac{2\pi n}{\ell} \left(c_n \sinh \frac{\pi ns}{\ell} + d_n \cosh \frac{\pi ns}{\ell} \right) \exp(2\pi i ny/\ell). \end{aligned}$$

Corresponding to the pair $(3.1.5)_0$ and $(3.1.5)_{-1}$ of versions of (3.1.5) there is a pair of versions of equation (2.4.5); we intend to rewrite those equations in terms of the xy-coordinates, which requires some interpretation beforehand. First we decide that the x_2 -direction in §2.4 will be interpreted as the $\frac{\partial}{\partial x}$ direction and the x_1 -direction will be interpreted as the $\frac{\partial}{\partial x}$ direction and the x_1 -direction will be interpreted as the $\frac{\partial}{\partial x}$ direction and the x_1 -direction will be forcing $\{\frac{\partial}{\partial x_1}, \frac{\partial}{\partial x_2}\}$ to be a frame with the opposite orientation than $\{\frac{\partial}{\partial x}, \frac{\partial}{\partial y}\}$. (All this choice of direction amounts to in (2.4.5) is a choice of direction normal to γ^{ℓ} and γ^r in which V is measured positively.) Next we observe that the arclength parameter x_1 was defined on the domain [0, 1], while the coordinate y, which we are presently using to parametrize the geodesics γ^{ℓ} and γ^r , varies over the domain $[0, \ell]$; we conclude that $V_{yy} = \ell^{-2}V_{11}$. Thus we translate equation (2.4.5) to:

$$(2.4.5)_0 V_{yy} = -\frac{1}{2} (\partial_x \dot{\mathcal{H}})_0$$

$$(2.4.5)_{-1} V_{yy} - V = -\frac{1}{2} (\partial_x \dot{\mathcal{H}})_{-1}$$

where here we have written the pair of derivatives of $\dot{\mathcal{H}}$ as $(\partial_x \dot{\mathcal{H}})_0$ and $(\partial_x \dot{\mathcal{H}})_{-1}$ depending on which side of γ^{ℓ} or γ^r we are considering. (In this notation, formula (3.2.4) refers to $(\partial_x \dot{\mathcal{H}})_0$.) Once again, we are using the regularity of the solution $\dot{\mathcal{H}}$ for the existence of the *x*-derivatives $(\partial_x \dot{\mathcal{H}})_0$ and $(\partial_x \dot{\mathcal{H}})_{-1}$.

At this point, we need to consider that there are two boundary components γ^{ℓ} and γ^{r} , and hence two variation vector fields with normal components V_{-} defined along $\gamma^{\ell} = \{x =$ $-\frac{s}{2}$ and V_+ defined along $\gamma^r = \{x = +\frac{s}{2}\}$. If we then set

(3.2.5)
$$V_{-} = \Sigma \lambda_n \exp(2\pi i n y/\ell)$$
 and $V_{+} = \Sigma \rho_n \exp(2\pi i n y/\ell)$

(where λ_n and ρ_n suggest "left" and "right", respectively), we substitute (3.2.4) and (3.2.5) into $(2.4.5)_0$ to obtain

$$\Sigma \frac{-4\pi^2 n^2}{\ell^2} \lambda_n \exp(2\pi i n y/\ell) = \frac{-c_0}{2} - \frac{1}{2} \Sigma' \frac{2\pi n}{\ell} \left(-c_n \sinh \frac{\pi n s}{\ell} + d_n \cosh \frac{\pi n s}{\ell} \right) \exp(2\pi i n y/\ell).$$

We solve for λ_n in the above to obtain, for $n \neq 0$,

(3.2.6)_-
$$\lambda_n = \frac{\ell}{4\pi n} \left(-c_n \sinh \frac{\pi ns}{\ell} + d_n \cosh \frac{\pi ns}{\ell} \right)$$

Similarly, we find, for $n \neq 0$,

$$(3.2.6)_+ \qquad \qquad \rho_n = \frac{\ell}{4\pi n} \left(c_n \sinh \frac{\pi ns}{\ell} + d_n \cosh \frac{\pi ns}{\ell} \right)$$

Along the way, we also find that, setting n = 0,

$$(3.2.7) c_0 = 0.$$

The formulae $(3.2.6)_{-}$ and $(3.2.6)_{+}$ when substituted into (3.2.5) yield the expansions

$$V_{-} = \lambda_{0} + \Sigma' \frac{\ell}{4\pi n} \left(-c_{n} \sinh \frac{\pi ns}{\ell} + d_{n} \cosh \frac{\pi ns}{\ell} \right) \exp(2\pi i ny/\ell)$$

and $V_{+} = \rho_{0} + \Sigma' \frac{\ell}{4\pi n} \left(c_{n} \sinh \frac{\pi ns}{\ell} + d_{n} \cosh \frac{\pi ns}{\ell} \right) \exp(2\pi i ny/\ell).$

We now use the crucial observation that V_{-} (and V_{+} , of course) solves both equations $(2.4.5)_0$ and $(2.4.5)_{-1}$. Thus, from our knowledge of V_- and V_+ , we can apply $(2.4.5)_{-1}$ to this expansion and obtain $(\partial_x \dot{\mathcal{H}})_{-1}$. That is, $(2.4.5)_{-1}$ is equivalent to

$$(\partial_x \dot{\mathcal{H}})_{-1} \Big|_{x=-\frac{s}{2}} = -2\{(V_-)_{yy} - (V_-)\}$$

$$(3.2.8)_- = 2\lambda_0 - 2\Sigma' \frac{\ell}{4\pi n} \left(-c_n \sinh \frac{\pi ns}{\ell} + d_n \cosh \frac{\pi ns}{\ell}\right) \left(-\frac{4\pi^2 n^2}{\ell^2} - 1\right) \exp(2\pi i ny/\ell)$$

$$= 2\lambda_0 + \Sigma' \frac{1}{2\pi n\ell} \left(4\pi^2 n^2 + \ell^2\right) \left(-c_n \sinh \frac{\pi ns}{\ell} + d_n \cosh \frac{\pi ns}{\ell}\right) \exp(2\pi i ny/\ell).$$

Similarly

$$(3.2.8)_{+} \\ (\partial_x \dot{\mathcal{H}})_{-1} \Big|_{x=+\frac{s}{2}} = 2\rho_0 + \Sigma' \frac{1}{2\pi n\ell} \left(4\pi^2 n^2 + \ell^2 \right) \left(c_n \sinh \frac{\pi ns}{\ell} + d_n \cosh \frac{\pi ns}{\ell} \right) \exp(2\pi i ny/\ell)$$

We pause and observe that we have obtained in equations (3.2.3) and (3.2.8) the Dirichlet and Neumann conditions, respectively, for the linearized Liouville/Bochner equation $(3.1.5)_{-1}$. This permits us to focus for the rest of the computation on the compact hyperbolic surface $(S_{-1}, \operatorname{gr}(\sigma_0)|_{S_{-1}})$ where S_{-1} denotes the closure of the $\{K_0 = -1\}$ subdomain of S.

It is a reflex in this situation to integrate by parts:

$$(3.2.9) \qquad 0 = \iint_{S_{-1}} \dot{\mathcal{H}} \Delta_h \dot{\mathcal{H}} - 2\dot{\mathcal{H}}^2 dA_{\mathrm{gr}(\sigma_0)}$$
$$= \iint_{S_{-1}} - \|\nabla \dot{\mathcal{H}}\|^2 - 2\dot{\mathcal{H}}^2 dA_{\mathrm{gr}(\sigma_0)} + \int_{\partial S_{-1}} \dot{\mathcal{H}} (\partial_n \dot{\mathcal{H}}) ds_{\mathrm{gr}(\sigma_0)}$$

where $\partial_n \dot{\mathcal{H}}$ denotes the outward normal. The boundary term we compute with (3.2.3) and (3.2.8), using that the normal derivatives $\partial_n \dot{\mathcal{H}} \Big|_{x=-\frac{s}{2}} = \partial_x \dot{\mathcal{H}}$ and $\partial_n \dot{\mathcal{H}} \Big|_{x=+\frac{s}{2}} = -\partial_x \dot{\mathcal{H}}$ in our coordinates. Thus

$$\int_{\partial S_{-1}} \dot{\mathcal{H}}(\partial_n \dot{\mathcal{H}}) ds_{\mathrm{gr}(\sigma_0)} = \int_{\{x=-\frac{s}{2}\}} \dot{\mathcal{H}}(\partial_x \dot{\mathcal{H}}) dy + \int_{\{x=\frac{s}{2}\}} \dot{\mathcal{H}}(-\partial_x \dot{\mathcal{H}}) dy$$

$$= 2\ell d_0 \lambda_0 + \int_{\{x=-\frac{s}{2}\}}^{y=\ell} \Sigma' \frac{1}{2\pi n\ell} \left(4\pi^2 n^2 + \ell^2\right) \left(-c_n \sinh \frac{\pi ns}{\ell} + d_n \cosh \frac{\pi ns}{\ell}\right) \times \left(c_{-n} \cosh \frac{\pi (-n)s}{\ell} - d_{-n} \sinh \frac{\pi (-n)s}{\ell}\right) + \left\{\operatorname{terms involving non-trivial powers of } \exp(2\pi i ny/\ell)\right\} dy + \left(2\ell d_0 \rho_0 + \int_{\{x=+\frac{s}{2}\}}^{y=\ell} \Sigma'(-1) \frac{1}{2\pi n\ell} \left(4\pi^2 n^2 + \ell^2\right) \left(c_n \sinh \frac{\pi ns}{\ell} + d_n \cosh \frac{\pi ns}{\ell}\right) \times \left(c_{-n} \cosh \frac{\pi (-n)s}{\ell} + d_{-n} \sinh \frac{\pi (-n)s}{\ell}\right) + \left(c_{-n} \cosh \frac{\pi (-n)s}{\ell} + d_{-n} \sinh \frac{\pi (-n)s}{\ell}\right) + \left(c_{-n} \cosh \frac{\pi (-n)s}{\ell} + d_{-n} \sinh \frac{\pi (-n)s}{\ell}\right) + \left(c_{-n} \cosh \frac{\pi (-n)s}{\ell} + d_{-n} \sinh \frac{\pi (-n)s}{\ell}\right) + \left(c_{-n} \cosh \frac{\pi (-n)s}{\ell} + d_{-n} \sinh \frac{\pi (-n)s}{\ell}\right) + \left(c_{-n} \cosh \frac{\pi (-n)s}{\ell} + d_{-n} \sinh \frac{\pi (-n)s}{\ell}\right) + \left(c_{-n} \cosh \frac{\pi (-n)s}{\ell} + d_{-n} \sinh \frac{\pi (-n)s}{\ell}\right) + \left(c_{-n} \cosh \frac{\pi (-n)s}{\ell} + d_{-n} \sinh \frac{\pi (-n)s}{\ell}\right) + \left(c_{-n} \cosh \frac{\pi (-n)s}{\ell} + d_{-n} \sinh \frac{\pi (-n)s}{\ell}\right) + \left(c_{-n} \cosh \frac{\pi (-n)s}{\ell} + d_{-n} \sinh \frac{\pi (-n)s}{\ell}\right) + \left(c_{-n} \cosh \frac{\pi (-n)s}{\ell}\right) +$$

{terms involving non-trivial powers of $\exp(2\pi i n y/\ell)$ }dy

after substituting (3.2.3) and (3.2.8). After integration and applying (3.2.2), the above

expression simplifies to (3, 2, 10)

$$\begin{aligned} &\int_{\partial S_{-1}} \dot{\mathcal{H}}(\partial_n \dot{\mathcal{H}}) ds_{\mathrm{gr}(\sigma_0)} = 2\ell d_0 (\lambda_0 - \rho_0) + \\ \ell \Sigma' \frac{1}{2\pi n\ell} \left(4\pi^2 n^2 + \ell^2 \right) \left(-c_n \sinh \frac{\pi ns}{\ell} + d_n \cosh \frac{\pi ns}{\ell} \right) \left(\bar{c}_n \cosh \frac{\pi ns}{\ell} - \bar{d}_n \sinh \frac{\pi ns}{\ell} \right) \\ &- \ell \Sigma' \frac{1}{2\pi n\ell} \left(4\pi^2 n^2 + \ell^2 \right) \left(c_n \sinh \frac{\pi ns}{\ell} + d_n \cosh \frac{\pi ns}{\ell} \right) \left(\bar{c}_n \cosh \frac{\pi ns}{\ell} + \bar{d}_n \sinh \frac{\pi ns}{\ell} \right) \\ &= 2\ell d_0 (\lambda_0 - \rho_0) + \\ \frac{1}{2\pi} \Sigma' \frac{1}{n} \left(4\pi^2 n^2 + \ell^2 \right) \left(-(|c_n|^2 + |d_n|^2) \sinh \frac{\pi ns}{\ell} \cosh \frac{\pi ns}{\ell} + c_n \bar{d}_n \sinh^2 \frac{\pi ns}{\ell} + \bar{c}_n d_n \cosh^2 \frac{\pi ns}{\ell} \right) \\ &- \frac{1}{2\pi} \Sigma' \frac{1}{n} \left(4\pi^2 n^2 + \ell^2 \right) \left((|c_n|^2 + |d_n|^2) \sinh \frac{\pi ns}{\ell} \cosh \frac{\pi ns}{\ell} + c_n \bar{d}_n \sinh^2 \frac{\pi ns}{\ell} + \bar{c}_n d_n \cosh^2 \frac{\pi ns}{\ell} \right) \\ &= 2\ell d_0 (\lambda_0 - \rho_0) - \frac{1}{\pi} \Sigma' \frac{1}{n} \left(4\pi^2 n^2 + \ell^2 \right) \left(|c_n|^2 + |d_n|^2) \sinh \frac{\pi ns}{\ell} \cosh \frac{\pi ns}{\ell} \cosh \frac{\pi ns}{\ell} \cosh \frac{\pi ns}{\ell} . \end{aligned}$$

Thus this final integral is the negative of the sum of positive terms summed with a mystery term $2\ell d_0(\lambda_0 - \rho_0)$. In Lemma 3.4, we will use the "slice condition" that our family of grafted metrics have Euclidean cylinders of unvarying length s to conclude that this term $2\ell d_0(\lambda_0 - \rho_0)$ is non-positive, which will force the integral in (3.2.10) to be non-positive.

Lemma 3.4. $\lambda_0 - \rho_0 = -\frac{sd_0}{2}$

We postpone the proof of Lemma 3.4 until $\S3.3$, preferring to assume it for now to finish the proof of Lemma 3.3. Now, assuming Lemma 3.4 and applying (3.2.10) to (3.2.9), we find that

$$\iint_{S_{-1}} (3.2.11)^* \int_{S_{-1}} -\|\nabla \dot{\mathcal{H}}\|^2 - 2\dot{\mathcal{H}}^2 dA_{\mathrm{gr}(\sigma_0)} - \frac{1}{\pi} \Sigma' \frac{1}{n} \left(4\pi^2 n^2 + \ell^2\right) \left(|c_n|^2 + |d_n|^2\right) \sinh \frac{\pi ns}{\ell} \cosh \frac{\pi ns}{\ell} - \ell s d_0^2 = 0$$

with all terms being nonpositive: this forces $c_n = d_n = 0$ for all n and $\dot{\mathcal{H}}$ to vanish identically in S_{-1} . Because $\dot{\mathcal{H}}$ is continuous across the boundary of the cylinder, it vanishes on all of S. Thus $\operatorname{gr}(\sigma_t) = \operatorname{gr}(\sigma_0)(1 + \mathcal{O}(t^2))$, so that id : $(S, \operatorname{gr}(\sigma_t)) \to (S, \operatorname{gr}(\sigma_0))$ is an isometry, up to order $\mathcal{O}(t^2)$. This concludes the proof of Lemma 3.3. \Box

From $(3.2.11)^*$ we also obtain, as a corollary of the proof,

Lemma 3.5. For a family of conformal grafted metrics $gr(\sigma_t)$, we have $V_+ \equiv V_- \equiv 0$.

Proof: We see that $c_n = d_n = 0$, so formulae (3.2.6) show that $\lambda_n = \rho_n = 0$ for $n \neq 0$. But then also equation $(2.4.5)_{-1}$ forces $\lambda_0 = \rho_0 = 0$. \Box

It remains to conclude the proof of Lemma 3.2. We already have from Lemma 3.3 that $\operatorname{gr}(\sigma_t) = \operatorname{gr}(\sigma_0)(1 + \mathcal{O}(t^2))$; thus id : $(S, \operatorname{gr}(\sigma_t)) \to (S, \operatorname{gr}(\sigma_0))$ is an isometry, up to order $\mathcal{O}(t^2)$.

Now, if (S,g) is a metric space which can be written as the graft of a hyperbolic surface with geodesic boundary γ to a flat cylinder of height s, then there is a welldefined inverse operation $\operatorname{UnGr}_{s\gamma}$ to the grafting operation $\operatorname{Gr}_{s\gamma}$: the inverse operation excises the flat cylinder and reidentifies the hyperbolic surface with boundary along γ by projecting the flat cylinder along its longitudinal geodesics. Of course, we have the property that $\operatorname{UnGr}_{s\gamma} \circ \operatorname{Gr}_{s\gamma}$ is the identity isometry. We apply this operation $\operatorname{UnGr}_{s\gamma}$ to $(S, \operatorname{gr}(\sigma_t))$: because id : $(S, \operatorname{gr}(\sigma_t)) \to (S, \operatorname{gr}(\sigma_0))$ is an isometry to order $\mathcal{O}(t^2)$, and Lemma 3.5 guarantees that the geodesic γ is not moving (to $\mathcal{O}(t^2)$), we find that the hyperbolic portions $(S_{-1}, \operatorname{gr}(\sigma_0))$ and $(S_{-1}, \operatorname{gr}(\sigma_t))$ agree to $\mathcal{O}(t^2)$. Moreover, the flat portions $(S_0, \operatorname{gr}(\sigma_0))$ and $(S_0, \operatorname{gr}(\sigma_t))$ also agree to $\mathcal{O}(t^2)$, and so again citing Lemma 3.5, we conclude that the longitudinal geodesics across the flat cylinders are also unchanged, to $\mathcal{O}(t^2)$. Thus $\sigma_0 = \operatorname{UnGr}_{s\gamma}\{\operatorname{gr}(\sigma_t) + \mathcal{O}(t^2)\} = \operatorname{UnGr}_{s\gamma}\{\operatorname{gr}(\sigma_t)\} + \mathcal{O}(t^2) = \sigma_t + \mathcal{O}(t^2),$ proving the lemma. \Box

3.3 Slice Condition. We have yet to use the hypothesis that the grafted cylinder has constant length s in the family $\operatorname{gr}(\sigma_t)$. Certainly it is necessary to use this hypothesis to prove Lemma 3.2, as Teichmüller space is 6g-6 (real) dimensional and the space $T_g \times \mathbf{R}_+$ of grafted hyperbolic metrics (up to Diff_o equivalence) is 6g-5 (real) dimensional. Thus we might expect that the map $T_g \times \mathbf{R}_+ \to T_g$ which records the conformal equivalence class of an equivalence class of grafted metrics would pullback points to one-dimensional families of grafted metrics. The content to Lemma 3.2 is that such families would meet level sets $T_g \times \{s_0\} \subset T_g \times \mathbf{R}_+$ in points; thus we must somehow use the fact that we are restricted to such a level set in the proof of Lemma 3.2.

Let us extend the notation of §3.1 somewhat and allow the Euclidean portion of the grafted metric $gr(\sigma_t)$ to be a Euclidean cylinder of length s = s(t), where we permit the length s(t) to vary in t. In the notation of §3.2, we claim that

Lemma 3.6.
$$\lambda_0 - \rho_0 = -\frac{sd_0}{2} - \frac{ds(t)}{dt}$$

From this lemma, we see the

Proof of Lemma 3.4. The Euclidean length s(t) is constant, hence $\frac{ds(t)}{dt} = 0$, and result follows.

Proof of Lemma 3.6. The plan is to compute the *t*-derivative of the area $A(\text{gr}(\sigma_t))$ of the family $\text{gr}(\sigma_t)$ of grafted metrics two ways. In the first method, we use that $\text{gr}(\sigma_t)$ is a metric which is composed of a portion which is hyperbolic with geodesic boundary and a portion which is composed of a Euclidean cylinder, and so the area is compatible via Gauss-Bonnet and elementary geometry. In the second method, we use the analytical formulae (3.1.2) and (3.1.5).

First Method. If we remove the cylindrical portion of the grafted metric and glue the resulting hyperbolic surface-with-geodesic-boundary together across its pair of geodesic boundary components, we obtain a closed hyperbolic surface of area $-2\pi\chi$, where χ is the Euler characteristic. Thus, using that the cylinder has length $\ell_{\gamma}(\operatorname{gr}(\sigma_t)) \cdot s(t)$, we find that the area $A(\operatorname{gr}(\sigma_t))$ of the grafted metric satisfies

$$A(\operatorname{gr}(\sigma_t)) = -2\pi\chi + \ell_{\gamma}(\operatorname{gr}(\sigma_t)) \cdot s(t)$$

so that

(3.3.1)
$$\frac{d}{dt}\Big|_{t=0} A(\operatorname{gr}(\sigma_t)) = \left(\frac{d}{dt}\Big|_{t=0} \ell_{\gamma}(\operatorname{gr}(\sigma_t))\right) s(0) + \ell_{\gamma}(\operatorname{gr}(\sigma_t)) \cdot \frac{d}{dt}\Big|_{t=0} s(t).$$

To find the derivative $\frac{d}{dt}\Big|_{t=0} \ell_{\gamma}(\operatorname{gr}(\sigma_t))$, we first observe that since the length $\ell_{\gamma}(\operatorname{gr}(\sigma_t))$ is that of a geodesic γ_t which varies smoothly in a family containing the *geodesic* γ_0 , then we must have that

(3.3.2)
$$\frac{d}{dt}\Big|_{t=0} \ell_{\gamma}(\operatorname{gr}(\sigma_t)) = \frac{d}{dt}\Big|_{t=0} \int_{\gamma_0} ds_{\operatorname{gr}(\sigma_t)}.$$

Yet the term $ds_{gr(\sigma_t)}$ is computable from (3.1.2) as

$$ds_{\operatorname{gr}(\sigma_t)} = \sqrt{\frac{1}{\mathcal{H}_t}} ds_{\operatorname{gr}(\sigma_0)}.$$

We combine this equation with (3.3.2) and differentiate to find

$$\begin{aligned} \frac{d}{dt} \Big|_{t=0} \ell_{\gamma}(\operatorname{gr}(\sigma_{t})) &= -\frac{1}{2} \int_{\gamma} \dot{\mathcal{H}} \mathcal{H}_{0}^{-3/2} ds_{\operatorname{gr}(\sigma_{0})} \\ &= -\frac{1}{2} \int \dot{\mathcal{H}} ds_{\operatorname{gr}(\sigma_{0})} \end{aligned}$$

as $\mathcal{H}_0 \equiv 1$.

We next apply our formulae (3.2.3) for $\dot{\mathcal{H}}$ and (3.2.7) for c_0 to this last equation to find that

(3.3.3)
$$\frac{d}{dt}\Big|_{t=0} \ell_{\gamma}(\operatorname{gr}(\sigma_{t})) = -\frac{1}{2} \int_{0}^{\ell} d_{0} + \text{ terms involving powers of } \exp(2\pi i n y/\ell) dy$$
$$= -\frac{1}{2} d_{0}\ell.$$

Combining (3.3.1) and (3.3.3) yields

(3.3.4)
$$\frac{d}{dt}A(gr(\sigma_t)) = -\frac{1}{2}d_0\ell s(0) + \ell \frac{d}{dt} \Big|_{t=0} s(t)$$

Second Method. Formula (3.1.2) suggests another method, as the area $A(gr(\sigma_t))$ may be expressed as

$$A(\operatorname{gr}(\sigma_t)) = \iint_{S} dA(\operatorname{gr}(\sigma_t))$$
$$= \iint_{S} \frac{dA(\operatorname{gr}(\sigma_0))}{\mathcal{H}_t}.$$

Thus

(3.3.5)
$$\frac{d}{dt}\Big|_{t=0} A(\operatorname{gr}(\sigma_t)) = -\iint_{S} \dot{\mathcal{H}} \mathcal{H}_0^{-2} dA(\operatorname{gr}(\sigma_0))$$
$$= -\iint_{S_{-1}} \dot{\mathcal{H}} dA(\operatorname{gr}(\sigma_0)) - \iint_{S_0} \dot{\mathcal{H}} dA(\operatorname{gr}(\sigma_0))$$

The two terms in formula (3.3.5) require separate treatments. To evaluate the first term, begin with equation $(3.1.5)_{-1}$ and integrate to find

$$0 = \iint_{S_{-1}} (\Delta_h - 2) \dot{\mathcal{H}} dA_{\mathrm{gr}(\sigma_0)} = \int_{\partial S_{-1}} \partial_n \dot{\mathcal{H}} ds_{\mathrm{gr}(\sigma_0)} - 2 \iint_{S_{-1}} \dot{\mathcal{H}} dA_{\mathrm{gr}(\sigma_0)}.$$

We rearrange to find

$$-\iint_{S_{-1}} \dot{\mathcal{H}} = -\frac{1}{2} \int_{\partial S_{-1}} \partial_n \dot{\mathcal{H}} ds_{\mathrm{gr}(\sigma_0)}$$
$$= -\frac{1}{2} \int_{\{x=-\frac{s}{2}\}} \dot{\mathcal{H}}_x dy + \frac{1}{2} \int_{\{x=\frac{s}{2}\}} \dot{\mathcal{H}}_x dy$$

(by ∂_n referring to the outward normal)

$$= -\frac{1}{2} \int_0^{\ell} 2\lambda_0 + \{\text{terms involving non-trivial powers of } \exp(2\pi i n y/\ell)\} dy \\ + \frac{1}{2} \int_0^{\ell} 2\rho_0 + \{\text{terms involving non-trivial powers of } \exp(2\pi i n y/\ell)\} dy$$

(by $(3.2.8)_{-}$ and $(3.2.8)_{+}$)

(3.3.6)
$$= -\ell(\lambda_0 - \rho_0).$$

To find the second term in (3.3.5), we simply use formula (3.2.1), again using (3.2.7) to set $c_0 = 0$. We find (3.3.7)

$$-\iint_{S_0} \dot{\mathcal{H}} dA_{\mathrm{gr}(\sigma_0)} = -\int_{-\frac{s}{2}}^{\frac{s}{2}} \int_0^\ell d_0 + \{\text{terms involving non-trivial powers of } \exp(2\pi i n y/\ell)\} \\ = -d_0 \ell s(0).$$

We combine (3.3.5), (3.3.6) and (3.3.7) to find

(3.3.8)
$$\frac{d}{dt}A(\operatorname{gr}(\sigma_t)) = -\ell(\lambda_0 - \rho_0) - d_0\ell s(0).$$

Summary. Formulae (3.3.4) and (3.3.8) combine to yield

(3.3.9)
$$-\frac{1}{2}d_0\ell s(0) + \ell \frac{d}{dt}\Big|_{t=0} s(t) = -\ell(\lambda_0 - \rho_0) - d_0\ell s(0)$$

from which the statement of the lemma follows immediately. \Box

§4 The General Case.

In this section we will prove the main theorem in the case of grafting on a measured lamination which is not necessarily a weighted simple closed curve.

Theorem. For any $\lambda \in \mathcal{ML}$, $Gr_{\lambda} : T_g \to T_g$ is a homeomorphism.

As in the model case, we need only prove the local injectivity; to that end, we suppose the theorem is false and get a variation σ_t of σ_0 such that $Gr_\lambda\sigma_t = Gr_\lambda\sigma_0$ in T_g . The harmonic map setup is the same as the model case; we isotope the grafted metrics and assume that the identity map $(S, gr_\lambda(\sigma_t)) \to (S, gr_\lambda(\sigma_0))$ is harmonic and conformal for each t. The functions \mathcal{H}_t and $\dot{\mathcal{H}}$ are defined in the usual way.

The first step in the proof, of course, is to approximate $\lambda \in \mathcal{ML}$ by a sequence of weighted simple closed curves $s_m \gamma_m \to \lambda$ and attempt to use our computations from the model case. The main difficulty is that the family of grafted metrics $Gr_{s_m\gamma_m}(\sigma_t)$ can no longer be assumed conformal and we must generalize some results from §2.4 and §3 to allow for this possibility.

Our plan is to carry out the derivation of §3 for a single non-conformal deformation; thus we will suppress the subscripts m in the notation until section §4.5. Let us observe that there were four basic steps in §3: 1) The derivation of the equation (3.1.5) for $\dot{\mathcal{H}}$, 2) the computation of $(\partial_x \dot{\mathcal{H}})_{-1}$ using the geodesic variational vector field equation (2.4.5), 3) the relating of the difference $\lambda_0 - \rho_0$ of the constant terms in the variational vector field to global quantities in grafting and $\dot{\mathcal{H}}$, and 4) the derivation of the identity for the norm of $\dot{\mathcal{H}}$ in terms of ℓ , s and $\frac{ds}{dt}$, as embodied in formulae (3.2.9), (3.2.10) and the formula (3.2.11)*. We carry out these steps in the next four sections §§4.1-4.4, culminating in a formula like that in formula (3.2.11)*. We will then interpret this generalized formula (3.2.11)* in terms of quantities which are continuous on \mathcal{ML} .

4.1 The Infinitesimal Bochner Equation. In this section, we show that the basic global equations are unchanged.

Lemma 4.1.1. \mathcal{H} satisfies equation (3.1.5).

Thus our main equation of study is unchanged despite allowing the family of harmonic maps to stray from conformality.

Proof: To see this, we begin with the Bochner equation (2.3.1)

$$\Delta_t \log \mathcal{H}_t = -2K_0 \mathcal{H}_t + 2K_0 \frac{t^2 |\Phi|^2}{\rho_t \mathcal{H}_t} + 2K_t$$

and differentiate once with respect to t at t = 0. We see immediately that

$$\frac{d}{dt}2K_0\frac{t^2|\Phi|^2}{\rho_t\mathcal{H}_t} = 0$$

so the derivative of the right hand side becomes $-2K_0\dot{\mathcal{H}} + 2\dot{K}$. Because $\log \mathcal{H}_0 \equiv 0$, all terms of $\frac{d}{dt}\Delta_t$ vanish except those for which the derivative passes through. Thus

$$\frac{d}{dt}\Delta_t \log \mathcal{H}_t = \Delta_0 \dot{\mathcal{H}},$$

yielding (3.1.5).

4.2. The geodesic variational vector field. A straightforward computation confirms the formula

(4.2.1)
$$\operatorname{gr}(\sigma_t) = -\Phi(t)dz^2 + \frac{\operatorname{gr}(\sigma_0)}{\mathcal{H}_t} - \overline{\Phi(t)}d\overline{z}^2 + \mathcal{O}(t^2).$$

From this formula, we can compute the expressions we need in order to apply formula (2.4.1) to the present case. In particular, when we differentiate (2.4.1) in time, we observe that formula (2.4.2) (and those formulae following it) continues to be valid; we are left to evaluate $\frac{d}{dt}\Big|_{t=0} \Gamma^2_{t,11}$, where we adopt Fermi coordinates for $\operatorname{gr}(\sigma_0)$. In terms of those Fermi coordinates $\{x^1, x^2\}$ for $\operatorname{gr}(\sigma_0)$ along the curve γ (here representing one of $\{\gamma^{\ell}, \gamma^r\}$), the formula (4.2.1) becomes

(4.2.2)
$$\operatorname{gr}(\sigma_t) = \frac{\operatorname{gr}(\sigma_0)}{\mathcal{H}_t} + t\phi_{ij}dx_i dx_j + \mathcal{O}(t^2)$$

where the tensor $\phi_{ij} dx_i dx_j$ may be represented as

(4.2.3)
$$\phi_{ij} = \begin{pmatrix} -2\operatorname{Re}\phi & 2\operatorname{Im}\phi\\ 2\operatorname{Im}\phi & 2\operatorname{Re}\phi \end{pmatrix} + \mathcal{O}(x_2^2)\mathcal{O}(t) + \mathcal{O}(t^2).$$

Here we have written $\Phi(t) = \{t\phi + \mathcal{O}(t^2)\} \left(dx_1 + i\frac{dx_2}{F(x_2)}\right)^2$, noting that $dx_1 + i\frac{dx_2}{F(x_2)}$ is a conformal coordinate up to order $\mathcal{O}(x_2^2)$. We continue to consider both $\Phi(t)$ and ϕ as small quantities, since we regard $gr(\sigma_t)$ as nearly conformal to $gr(\sigma_0)$ to first order in t.

Thus, since $F(x_2) = 1 + \mathcal{O}(x_2^2)$, we may compute along the curve γ that,

$$\frac{d}{dt}\Big|_{t=0} \Gamma^2_{t,11} = \frac{1}{2}\partial_2 \dot{\mathcal{H}} + \frac{1}{2} \left[2\frac{\partial}{\partial x_1} (2\operatorname{Im}\phi) + \frac{\partial}{\partial x_2} (2\operatorname{Re}\phi) \right] + \mathcal{O}(x_2^2)$$
$$= \frac{1}{2}\partial_2 \dot{\mathcal{H}} + \frac{\partial}{\partial x_1} \operatorname{Im}\phi$$

where we evaluate along the curve $\gamma = \{x_2 = 0\}$ and we use the Cauchy-Riemann equations for ϕ to simplify the second term. Next, jump to section 3.2 where we use these equations to solve for $(\partial_x \dot{\mathcal{H}})_{-1}$ in terms of $(\partial_x \dot{\mathcal{H}})_0$: we adapt our work there to the present (more general) case by expanding the harmonic function Im ϕ in the expansion (compare (3.2.1))

(4.2.4)
$$\operatorname{Im} \phi = u_0 x + v_0 + \Sigma' \left(u_n \cosh \frac{2\pi n}{\ell} x + v_n \sinh \frac{2\pi n}{\ell} x \right) \exp(2\pi i n y/\ell).$$

Let W solve the amended variational equation

$$(2.4.5)_0^{\Phi} \qquad \qquad W_{yy} = -\frac{1}{2} (\partial_x \dot{\mathcal{H}})_0 - (\partial_y \operatorname{Im} \phi)$$

in analogy to equation $(2.4.5)_0$.

We find that if W is of the forms $W_{-} = \Sigma \lambda_n^* \exp(2\pi i n y/\ell)$ and $W_{+} = \Sigma \rho_n^* \exp(2\pi i n y/\ell)$, then for $n \neq 0$

(4.2.5)_-
$$\lambda_n^* = \lambda_n + \frac{\ell}{2\pi i n} \left(u_n \cosh \frac{\pi n s}{\ell} - v_n \sinh \frac{\pi n s}{\ell} \right).$$

$$(4.2.5)_+ \qquad \qquad \rho_n^* = \rho_n + \frac{\ell}{2\pi i n} \left(u_n \cosh \frac{\pi n s}{\ell} + v_n \sinh \frac{\pi n s}{\ell} \right)$$

again finding that

$$c_0 = 0.$$

Next, in analogy to (3.2.8), we find that

(4.2.6)
$$(\partial_x \dot{\mathcal{H}})_{-1} = (\partial_x \dot{\mathcal{H}})_0 - 2W$$

and so has the expansions

$$(4.2.7)_{-} \qquad \left. (\partial_x \dot{\mathcal{H}})_{-1} \right|_{x=-\frac{s}{2}} = 2\lambda_0 + \Sigma' (4\pi^2 n^2 + \ell^2) \left\{ \frac{1}{2\pi n\ell} \left(-c_n \sinh \frac{\pi ns}{\ell} + d_n \cosh \frac{\pi ns}{\ell} \right) + \frac{1}{\pi i n\ell} \left(u_n \cosh \frac{\pi ns}{\ell} - v_n \sinh \frac{\pi ns}{\ell} \right) \right\}$$

and

$$(4.2.7)_{+} \qquad \left. \begin{array}{c} (\partial_{x}\dot{\mathcal{H}})_{-1} \Big|_{x=\frac{s}{2}} = 2\rho_{0} + \Sigma'(4\pi^{2}n^{2} + \ell^{2}) \left\{ \frac{1}{2\pi n\ell} \left(c_{n} \sinh \frac{\pi ns}{\ell} + d_{n} \cosh \frac{\pi ns}{\ell} \right) \right. \\ \left. + \frac{1}{\pi i n\ell} \left(u_{n} \cosh \frac{\pi ns}{\ell} + v_{n} \sinh \frac{\pi ns}{\ell} \right) \right\}$$

We will use these formulas in 4.4 when we combine all of our modifications to 3 to get a new version of formula $(3.2.11)^*$.

4.3. The slice condition. In this section, we generalize §3.3 to the (general) case of a non-conformal deformation. Here the metric $gr(\sigma_t)$ is involved in the computation of the first variation of arclength (3.3.3) and in the computation of the first variation of area (3.3.5). In the latter, we note from (4.2.1) that

$$dA(\operatorname{gr}(\sigma_t)) = \frac{dA(\operatorname{gr}(\sigma_0))}{\mathcal{H}_t(1 - |\nu(t)|^2)}$$

where $\nu(t) = \mathcal{O}(t)$ denotes the Beltrami differential. Thus

(4.3.1)
$$dA(\operatorname{gr}(\sigma_t)) = \frac{dA(\operatorname{gr}(\sigma_0))}{\mathcal{H}_t} + \mathcal{O}(t)^2.$$

In the computation of the first variation of arclength, we have, from (4.2.2) and (4.2.3) that

(4.3.2)
$$\frac{d}{dt}\Big|_{t=0} \ell_{\gamma}(\operatorname{gr}(\sigma_{t})) = \frac{d}{dt}\Big|_{t=0} \int_{\gamma_{0}} ds_{\operatorname{gr}(\sigma_{t})} \\ = \frac{d}{dt} \int_{\gamma} \sqrt{\mathcal{H}_{t}^{-1} + 2\operatorname{Re} t\phi + \mathcal{O}(t^{2})} \, ds_{\operatorname{gr}(\sigma_{0})} \\ = -\frac{1}{2} \int_{\gamma} (\dot{\mathcal{H}} - 2\operatorname{Re} \phi) ds_{\operatorname{gr}(\sigma_{0})} \\ = \frac{1}{2} \int_{\gamma} \dot{\mathcal{H}} ds_{\operatorname{gr}(\sigma_{0})} + \ell \mathcal{O}(\|\dot{\Phi}(t)\|).$$

The last expression may require some explanation. For any given infinitesimal holomorphic quadratic differential $\dot{\Phi}(t)$, on the fixed compact surface $\operatorname{gr}(\sigma_0)$, the Harnack inequality bounds the supremum of $|\operatorname{Re} \phi|$ over the curve γ in terms of the integral norm of $\dot{\Phi}(t)$ and the length of the curve. As there is but a compact set of such unit norm quadratic differentials $\dot{\Phi}(t)$, the result holds in general, even in a precompact family $\{\gamma\}$ of grafting loci.

Finally we collect terms, as in the summary (3.3.9), but with the addition of the considerations from (4.3.1) and (4.3.2), to find

Lemma 4.3.1.
$$\ell(\lambda_0 - \rho_0) = -\frac{1}{2}d_0\ell s(0) - \ell \frac{ds}{dt} + \ell s(0)\mathcal{O}(\|\dot{\Phi}\|).$$

4.4. The extended identity. Finally, we combine the results of our previous sections into an identity analogous to that of $(3.2.11)^*$. We extend the computation of (3.2.9) and

(3.2.10) to obtain

$$\begin{split} &\int_{\partial S_{-1}} \dot{\mathcal{H}}(\partial_n \dot{\mathcal{H}}) ds_{\mathrm{gr}(\sigma_0)} = 2\ell d_0 (\lambda_0 - \rho_0) \\ &\quad -\frac{1}{\pi} \Sigma' \frac{1}{n} \left(4\pi^2 n^2 + \ell^2 \right) \left(|c_n|^2 + |d_n|^2 \right) \sinh \frac{\pi ns}{\ell} \cosh \frac{\pi ns}{\ell} \\ &\quad + \ell \Sigma' \left(4\pi^2 n^2 + \ell^2 \right) \frac{1}{\pi i n \ell} \Biggl\{ \left(u_n \cosh \frac{\pi ns}{\ell} - v_n \sinh \frac{\pi ns}{\ell} \right) \left(\bar{c}_n \cosh \frac{\pi ns}{\ell} - \bar{d}_n \sinh \frac{\pi ns}{\ell} \right) \Biggr\} \\ &\quad - \left(u_n \cosh \frac{\pi ns}{\ell} + v_n \sinh \frac{\pi ns}{\ell} \right) \left(\bar{c}_n \cosh \frac{\pi ns}{\ell} + \bar{d}_n \sinh \frac{\pi ns}{\ell} \right) \Biggr\} \\ &= 2\ell d_0 (\lambda_0 - \rho_0) - \frac{1}{\pi} \Sigma' \frac{1}{n} \left(4\pi^2 n^2 + \ell^2 \right) \left(|c_n|^2 + |d_n|^2 \right) \sinh \frac{\pi ns}{\ell} \cosh \frac{\pi ns}{\ell} \\ &\quad - \Sigma' \left(4\pi^2 n^2 + \ell^2 \right) \frac{2}{\pi i n} (v_n \bar{c}_n + u_n \bar{d}_n) \sinh \frac{\pi ns}{\ell} \cosh \frac{\pi ns}{\ell} \\ &= 2\ell d_0 (\lambda_0 - \rho_0) - \frac{1}{\pi} \Sigma' \frac{1}{n} \left(4\pi^2 n^2 + \ell^2 \right) \left(|c_n|^2 + |d_n|^2 \right) \sinh \frac{\pi ns}{\ell} \cosh \frac{\pi ns}{\ell} \\ &\quad - \sum_{n>0} \left(4\pi^2 n^2 + \ell^2 \right) \frac{2}{\pi n} \left(\mathrm{Im} \left[v_n \bar{c}_n + u_n \bar{d}_n \right] \right) \sinh \frac{\pi ns}{\ell} \cosh \frac{\pi ns}{\ell} . \end{split}$$

When we combine this with Lemma 4.3.1, we find the analogue to $(3.2.11)^*$: $(4.4.1)^*$

$$\iint_{S_{-1}} -\|\nabla \dot{\mathcal{H}}\|^2 - 2\dot{\mathcal{H}}^2 dA_{\mathrm{gr}(\sigma_0)} - \frac{1}{\pi} \Sigma' \frac{1}{n} \left(4\pi^2 n^2 + \ell^2\right) \left(|c_n|^2 + |d_n|^2\right) \sinh \frac{\pi ns}{\ell} \cosh \frac{\pi ns}{\ell} \\ -\sum_{n>0} \left(4\pi^2 n^2 + \ell^2\right) \frac{2}{\pi n} \left(\mathrm{Im} \left[v_n \bar{c}_n + u_n \bar{d}_n\right]\right) \sinh \frac{\pi ns}{\ell} \cosh \frac{\pi ns}{\ell} \\ -\ell s d_0^2 - 2\ell s d_0 \frac{\dot{s}}{s} + \ell s \mathcal{O}(\|\dot{\Phi}\|) = 0.$$

4.5. An identity on \mathcal{ML} and the conclusion of the proof. We define the length $L = L(\gamma)$ for the grafted curve γ to be

$$L(\gamma) = \ell_{\gamma} \cdot s;$$

if we regard the transverse measure on γ as being given by $s \cdot i(0, \gamma)$, then we see that $L(\gamma)$ is simply the length of $[\gamma]$ as an element of the measured lamination space \mathcal{ML} . It is well-known then that the function $L: \mathcal{S} \times \mathbf{R}_+ \subset \mathcal{ML} \to \mathbf{R}$ extends to a continuous function $L: \mathcal{ML} \to \mathbf{R}$. Our main observation is that $(4.4.1)^*$ extends to an identity in terms of $\dot{\mathcal{H}}, \dot{\Phi}, \frac{d}{dt}(\log s)$ and quantities that are continuous on \mathcal{ML} . To see this we rewrite

$$(4.4.1)^{*} \text{ as} (4.5.1)^{*} \iint_{S_{-1}} -\|\nabla \dot{\mathcal{H}}\|^{2} - 2\dot{\mathcal{H}}^{2} dA_{\text{gr}(\sigma_{0})} - \frac{1}{\pi} \Sigma' \frac{1}{n} \left(4\pi^{2}n^{2} + \ell^{2}\right) \left(|c_{n}|^{2} + |d_{n}|^{2}\right) \sinh \frac{\pi nL}{\ell^{2}} \cosh \frac{\pi nL}{\ell^{2}} - \sum_{n>0} \left(4\pi^{2}n^{2} + \ell^{2}\right) \frac{2}{\pi n} \left(\text{Im} \left[v_{n}\bar{c}_{n} + u_{n}\bar{d}_{n}\right]\right) \sinh \frac{\pi nL}{\ell^{2}} \cosh \frac{\pi nL}{\ell^{2}} - Ld_{0}^{2} - 2Ld_{0}\frac{d}{dt} \log s + L\mathcal{O}(\|\dot{\Phi}(t)\|) = 0.$$

It is now straightforward to complete the proof of the main technical result.

Proof of Theorem A. We have already established the result in the case where λ is a lamination supported on a finite set $\{\gamma_1, \ldots, \gamma_k\}$ of simple closed curves. For the general case consider a sequence $\langle s_m \gamma_m \rangle$ of measured laminations supported on simple closed curves of lengths $\ell_n = \ell(\gamma_n)$ which approximates λ . In our notation, s_m denotes the multiple of the transverse (intersection) measure for the geodesic γ_m , so we may express the transverse measure for λ as $\lim_{m\to\infty} s_m i(\cdot, \gamma_m)$ (see the discussion in §2.2).

In analogy with the opening of the proof of Lemma 3.2, we suppose (in order to obtain a contradiction) that there is a family of surfaces σ_t so that

(4.5.2)
$$\operatorname{Gr}_{\lambda}(\sigma_t) = \operatorname{Gr}_{\lambda}(\sigma_0)$$

at least to an order $\mathcal{O}(t^2)$ in T_g . We then consider the family $\operatorname{Gr}_{s_m\gamma_m}(\sigma_t)$: it is of course no longer necessary that $\operatorname{Gr}_{s_m\gamma_m}(\sigma_t)$ should equal $\operatorname{Gr}_{s_m\gamma_m}(\sigma_0)$ to $\mathcal{O}(t^2)$, but the condition (4.5.2) should be asymptotically true in m, by construction. This implies that the Hopf differential (which we will denote $\Phi_m(t)$) and which measures the quasiconformality between $\operatorname{Gr}_{s_m\gamma_m}(\sigma_t)$ and $\operatorname{Gr}_{s_m\gamma_m}(\sigma_0)$) should have first variation $\frac{d}{dt}\Big|_{t=0} \Phi_m(t)$ in t which tends to zero as $m \to \infty$. We write

$$\lim_{n \to \infty} \frac{d}{dt} \Big|_{t=0} \Phi_m(t) = 0$$

which implies that, in the formula $(4.5.1)^*$, we can take $|u_n|, |v_n| \to 0$, along with the final term $L\mathcal{O}(||\Phi||)$, since L is bounded. With the exception of the term involving $\frac{d}{dt}\Big|_{t=0} \log s(t)$, all of the term involve quantities which are continuous on \mathcal{ML} : this follows immediately, once we observe that $\ell^2 \sinh \frac{\pi n L}{\ell^2}$ is bounded for, say, $\ell > 1$. Of the exceptional term, we have the lemma

Lemma 4.5.1. $\frac{d}{dt}\Big|_{t=0}\log(s(t)) = 0.$

Proof: We simply need to interpret the "slice condition" for λ correctly. In particular, suppose we allow the transverse measure of λ to vary as our grafted metrics vary, i.e. let $\lambda_t = \rho_t \lambda$ with ρ_t differentiable and $\rho_0 = 0$. The slice condition is given by

$$\dot{\rho} = \frac{d}{dt} \Big|_{\substack{t=0\\26}} \rho_t = 0.$$

Now we can approximate λ_t by scaling the weighted simple closed curves $\{s_m \gamma_m\}$ in exactly the same way:

$$\lambda_t = \lim_{m \to \infty} \rho_t s_m \gamma_m.$$

Having done so, we have $s(t) = \rho_t s_m$ in the calculations above, and $0 = \dot{\rho} = \frac{\dot{s}}{s_m}$. \Box

The rest of the proof is straightforward. As $m \to \infty$, we have $\dot{\Phi} \to 0$ and the expression collapses to the previous version $(3.2.11)^*$. In particular, we see from $(4.5.1)^*$ that $\dot{\mathcal{H}}_m = \mathcal{O}(\|\dot{\Phi}_m(t)\|)$, where we have added subscripts to $\dot{\mathcal{H}}$ and $\Phi(t)$ to emphasize the dependence of these quantities upon the approximating sequence. Thus, the argument at the end of §3.2 extends to show that $\sigma_{t,m}$ agrees with $\sigma_{0,m}$ to $o_m(1) + \mathcal{O}(t^2)$ (where $o_m(1)$ goes to zero as m tends to infinity), and hence that σ_t agrees with σ_0 to $\mathcal{O}(t^2)$, as required. This proves the local injectivity, as desired. \Box

Remark: We have unnecessarily restricted ourselves to unpunctured surfaces, primarily for notational simplicity and expositional cleanliness. The proofs all extend to the punctured case once we make three observations: i) all of the measured laminations under consideration avoid a neighborhood of the punctures, ii) there is a unique harmonic map of finite energy between surfaces of bounded non-positive curvature and some negative curvature [Al64] ([Wo91b]), and (iii) the holomorphic energy function \mathcal{H} for such a map is bounded in C^1 across the punctures, so no new non-vanishing boundary terms would arise in a modified formula (3.2.9), or therefore in any of the subsequent starred formulae (3.2.11)*, (4.4.1)*, and (4.5.1)*.

§5 Applications.

5.1. Geometric Coordinates on the Bers Slice. Consider the Bers slice

$$B_Y = T_g \times \{Y\} \subset T_g \times T_g \cong QF.$$

There is a natural map $\beta : B_Y \to \mathcal{ML}$ which assigns to a quasi-Fuchsian group the bending lamination λ on the component of the convex hull boundary facing the fixed structure Y(continuity of β is proved in [KS95]). The hyperbolic structure on this component of the convex hull boundary defines a point μ in Teichmüller space, and the relevant observation is that $Y = \operatorname{Gr}_{\lambda}(\mu)$. Theorem A shows that the metric μ is determined by Y and λ ; therefore since the Thurston homeomorphism $\Theta : \mathcal{ML} \times T_g \to P_g$ (described in the the introduction §1) is one-one, the map β is also. This is a simple way of assigning "bending coordinates" to B_Y .

Corollary 5.1. Let B_Y be a Bers slice with fixed conformal structure Y. Then the map assigning the bending lamination on the component of the convex hull boundary facing Y is a homeomorphism onto its image.

5.2 Generalized Bers Slices. Deformation Spaces of Books of I-bundles. Any geometrically finite, freely indecomposable Kleinian group G has a space $\mathcal{T}(G)$ of quasiconformal deformations parametrized by the product $\prod T(S_i)$ of the Teichmüller spaces $T(S_i)$ of its boundary components $\{S_1, \ldots, S_K\}$ at infinity [Ma74]. As in the example of the quasi-Fuchsian groups above, one can define slices $S(t_1, \ldots, t_{K-1})$ of these deformation spaces by simply fixing, say, the conformal structures at infinity of the first K-1boundary components, and letting the last conformal structure vary over its Teichmüller space $T(S_K)$.

Let us focus our attention on a class of geometrically finite three-manifolds homeomorphic to the interior of a book of I-bundles. The deformation spaces of these three-manifolds are studied in detail in [AC96] and are important because of the discovery by Anderson and Canary that the closures of those deformation spaces exhibit previously unexpected phenomena. The simplest of these manifolds has the following description. Begin with a solid torus with three disjoint parallel annuli on the boundary; here we choose the annuli so that their central curve is homotopic within the solid torus to the core curve of the solid torus. Attach, along those annuli, thickenings $\{H_1, H_2, H_3\}$ of one-holed surfaces $\{M_1, M_2, M_3\}$ of genera h_1, h_2 , and h_3 (respectively). The new three-manifold N has boundary surfaces $\{S_1, S_2, S_3\}$ of genera $h_1 + h_2, h_2 + h_3$ and $h_3 + h_1$; indeed, these bounding surfaces S_1, S_2 and S_3 are obtained by gluing M_i to M_{i+1} (with cyclic indexing) along the single boundary $\gamma = \partial M_i = \partial M_{i+1}$. Because all the thickenings of the surfaces are glued along neighborhoods of curves which retract to the core curve, we see that all of the curves γ_i are homotopic to each other and to the core curve γ of the central solid torus.

Now consider the space $\mathcal{T}(G)$ of quasi-conformal deformations of the Kleinian group G obtained as the holonomy of the hyperbolization of this three-manifold N. We consider the slice $\mathcal{S}(t_1, t_2) \subset \mathcal{T}(G) = T(S_1) \times T(S_2) \times T(S_3)$ defined by the coordinate description

$$\mathcal{S}(t_1, t_2) = \{t_1\} \times \{t_2\} \times T(S_3)$$

We then consider a map $\beta_{AC} : \mathcal{S}(t_1, t_2) \to \mathcal{ML} \times \mathcal{ML}$, analogous to the map $\beta : B_Y \to \mathcal{ML}$ above, which assigns to an element $(t_1, t_2, t_3) \in \mathcal{S}(t_1, t_2)$ the pair of bending measures of the boundary components $(C_1 \text{ and } C_2)$ of the convex hulls facing the conformal structures at infinity represented by t_1 and t_2 . Our application of Theorem A is the following

Corollary 5.2. The map $\beta_{AC} : \mathcal{S}(t_1, t_2) \to \mathcal{ML} \times \mathcal{ML}$ is injective.

Proof: Suppose $\beta_{AC}(t_1, t_2, t_3) = \beta_{AC}(t_1, t_2, t'_3)$. Then by Theorem A, not only do the bending measures on the convex hull boundary components C_i and C'_i $(i = 1, 2, \text{ facing the ends } t_i \text{ and } t'_i$, respectively), but so do the hyperbolic structures. Lift to the quasi-Fuchsian covers \mathcal{Q}_i and \mathcal{Q}'_i of (t_1, t_2, t_3) and (t_1, t_2, t'_3) corresponding to the surfaces C_i and C'_i and observe that these are identical by Corollary 5.1. Thus the holonomy representations of $\pi_1(C_i)$ are conjugate to those of $\pi_1(C'_i)$. But as there is a common element γ in $\pi_1(C_1) \subset \pi_1(N)$ and $\pi_1(C_2) \subset \pi_1(N)$, we see that the pair of representations of $\pi_1(S_1) *_{[\gamma]} \pi_1(M_3) = \pi_1(N(t_1, t_2, t_3))$ (in the obvious notation) are conjugate. Thus $[(t_1, t_2, t_3)] = [(t_1, t_2, t'_3)] \in \mathcal{T}(G)$, as desired. \Box

5.3. 2+1 de Sitter Spacetimes. We finish with an application to the structure of (2+1)-dimensional de Sitter spacetimes, following [Sc96]. Recall that 3-dimensional de Sitter space is defined to be the set of unit spacelike vectors in Minkowski space:

$$\mathbb{S}_1^3 = \{ v \in \mathbb{R}_1^4 | < v, v > = +1 \}.$$
28

This is the model space for Lorentzian 3-manifolds of constant positive curvature. Projectivizing \mathbb{R}_1^4 to $\mathbb{R}P^3$, we get the Klein model of hyperbolic space from the unit timelike vectors, the sphere at infinity S_{∞}^2 from the light cone, and a projective model of \mathbb{S}_1^3 as the remainder of $\mathbb{R}P^n$. Taking polar duals with respect to the sphere at infinity gives a correspondence between points in the projectivized de Sitter space and planes in hyperbolic space (and thus with round circles on S_{∞}^2).

Now imagine a projective structure on a closed hyperbolic surface S close to a Fuchsian structure (the construction works for any projective structure but is easiest to describe for the quasi-Fuchsian case). Using the polarity mentioned above, the set of all closed round balls contained within $dev(\tilde{S})$ defines a certain open subset \mathcal{U} of \mathbb{S}_1^3 . The holonomy $hol(\pi_1(S))$ acts discontinuously on \mathcal{U} and the quotient is a de Sitter spacetime homeomorphic to $S \times \mathbb{R}$. Any example arising from a projective structure on S in this way is called a *standard de Sitter spacetime*. Standard de Sitter spacetimes are well-behaved from the point of view of causality – in particular, we can choose the product structure so that each slice $S \times \{t\}$ is spacelike and every timelike or lightlike curve crosses $S \times \{t\}$ exactly once (we say $S \times \mathbb{R}$ is a *domain of dependence*).

The main result of [Sc96] is that every de Sitter spacetime $S \times \mathbb{R}$ which is a domain of dependence embeds in a standard de Sitter spacetime. Now suppose we have an example coming from a projective structure with Thurston coordinates $(\lambda, \sigma) \in \mathcal{ML} \times T_g$. A domain of dependence has a well-defined *causal horizon*; it follows easily that the causal horizon corresponds to the space of *maximal* open round balls, which is in turn isometric to the \mathbb{R} -tree dual to λ [Sc96].

We are now able to refine our classification of de Sitter spacetimes, by providing coordinates in terms of naturally-arising data in the future (the future causal horizon) and the past (the conformal structure on S at past infinity). More precisely, we have the following reworking of Theorem A:

Corollary 5.3. Let $\lambda \in \mathcal{ML}$ be a measured lamination with dual \mathbb{R} -tree λ . Let $\mathbb{S}_1^3(S; \lambda)$ be the family of standard de Sitter spacetimes with future causal horizon $\tilde{\lambda}$, and define a map $c_{\infty} : \mathbb{S}_1^3(S; \tilde{\lambda}) \to T_g$ which assigns the conformal structure on S at past infinity. Then c_{∞} is one-one.

Proof: By definition, any two standard de Sitter spacetimes \mathcal{M}_i (i = 1, 2) in $\mathbb{S}^3_1(S; \tilde{\lambda})$ come from projective structures on S (say with Thurston coordinates (λ, σ_i)). By examining the construction above, we have

$$c_{\infty}(\mathcal{M}_i) = Gr_{\lambda}(\sigma_i).$$

Because Gr_{λ} is one-one (Theorem A), c_{∞} is also. \Box

References

- [Al64] S.I. Al'ber, On n-dimensional Problems in the Calculus of Variations, Sov. Math. Dokl. 5 (1964), 700–704.
- [AC96] J. W. Anderson and R. D. Canary, Algebraic limits of Kleinian groups which rearrange the pages of a book, Invent. Math. 126 (1996), no. 2, 205–214.

- [Ap88] B. N. Apanasov, The geometry of Nielsen's hull for a Kleinian group in space and quasiconformal mappings, Ann. Global Anal. Geom. 6 (1988), no. 3, 207–230.
- [Be77] M. Berger, Nonlinearity and Functional Analysis, Academic Press, New York, 1977.
- [Be64] L. Bers, On moduli of Riemann surfaces, ETH Lecture Notes, Zürich (1964).
- [Bo88] F. Bonahon, The geometry of Teichmüller space via geodesic currents, Invent. Math. 92 (1988), 139–162.
- [Ea81] C. J. Earle, On variation of projective structures, in Riemann Surfaces and Related Topics: Proceedings of the 1978 Stony Brook Conference, Ann. of Math. Stud., vol. 97 (1981), Princeton Univ. Press, Princeton, 87–99.
- [EL81] J. Eells and L. Lemaire, Deformations of metrics and associated harmonic maps, Proc. Indian Acad. Sci. 90 (1981), no. 1, 33–45.
- [EM87] D. B. A. Epstein and A. Marden, Convex hulls in hyperbolic space, a theorem of Sullivan, and measured pleated surfaces, in Analytical and Geometric Aspects of Hyperbolic Space, London Math. Soc. Lecture Note Ser., vol. 111 (1987), Cambridge Univ. Press, Cambridge, 113–254.
- [Fa83] G. Faltings, Real projective structures on Riemann surfaces, Compositio Math. 48 (1983), no. 2, 223–269.
- [GKM95] D. M. Gallo, M. E. Kapovich, and A. Marden, On monodromy of Schwarzian differential equations on Riemann surfaces, Preprint (1995).
- [GT83] D. Gilbarg and N. S. Trudinger, Elliptic partial differential equations of second order, Grund. Math. Wiss., vol. 224, Springer-Verlag, New York-Berlin-Heidelberg, 1983.
- [Go87] W. M. Goldman, Projective structures with Fuchsian holonomy, J. Differential Geom. 25 (1987), 297–326.
- [Go88] _____, Geometric structures on manifolds and varieties of representations, in Geometry of Group Representations, Contemp. Math., vol. 74 (1988), Amer. Math. Soc., Providence, RI, 169-197.
- [GS92] M. L. Gromov and R. M. Schoen, Harmonic maps into singular spaces and p-adic superrigidity for lattices in groups of rank one, Inst. Hautes Études Sci. Publ. Math. **76** (1992), 165–246.
- [Gu81] R. C. Gunning, Affine and projective structures on Riemann surfaces, in Riemann Surfaces and Related Topics: Proceedings of the 1978 Stony Brook Conference, Ann. of Math. Stud., vol. 97 (1981), Princeton Univ. Press, Princeton, 225–244.
- [He75] D. A. Hejhal, Monodromy groups and linearly polymorphic functions, Acta Math. 135 (1975), 1–55.
- [HK98] C. D. Hodgson and S. P. Kerckhoff, Rigidity of hyperbolic cone-manifolds and hyperbolic Dehn surgery, J. Differential Geom. 48 (1998), no. 1, 1–59.
- [Jo97] J. Jost, Compact Riemann Surfaces, Springer-Verlag, New York-Berlin-Heidelberg, 1997.
- [KT92] Y. Kamishima and S. P. Tan, Deformation spaces on geometric structures, in Aspects of lowdimensional manifolds, Adv. Stud. Pure Math., vol. 20 (1992), Kinokuniya, Tokyo, 263–299.
- [KS95] L. Keen and C. M. Series, Continuity of convex hull boundaries, Pacific J. Math. 168 (1995), no. 1, 183–206.
- [Ke85] S. P. Kerckhoff, Earthquakes are analytic, Comm. Math. Helv. 60 (1985), no. 1, 17–30.
- [Kl33] F. Klein, Vorlesungen Über die Hypergeometrische Funktion, Springer-Verlag, Berlin, 1933.
- [Kr69] I. Kra, Deformations of Fuchsian groups, Duke Math. J. **36** (1969), 537–546.
- [Kr71] _____, Deformations of Fuchsian groups, II, Duke Math. J. **38** (1971), 499–508.
- [KM81] I. Kra and B. Maskit, *Remarks on projective structures*, in Riemann Surfaces and Related Topics: Proceedings of the 1978 Stony Brook Conference, Ann. of Math. Stud., vol. 97 (1981), Princeton Univ. Press, Princeton, 343–359.
- [KP94] R. S. Kulkarni and U. Pinkall, A canonical metric for Möbius structures and its applications, Math. Z. 216 (1994), 89–129.
- [La92] F. Labourie, Surfaces convexes dans l'espace hyperbolique et CP¹-structures, J. London Math. Soc. 45 (1992), 549–565.
- [Ma74] A. Marden, The geometry of finitely generated Kleinian groups, Ann. of Math. **99** (1974), 383–462.

- [Ma69] B. Maskit, On a class of Kleinian groups, Ann. Acad. Sci. Fenn. Ser. A I Math. 442 (1969), 1–8.
- [MV94] K. Matsuzaki and J. A. Velling, Notes on projective structures and Kleinian groups, Osaka J. Math. 31 (1994), no. 1, 165–175.
- [Mc98] C. T. McMullen, Complex earthquakes and Teichmüller theory, J. Amer. Math. Soc. 11 (1998), no. 2, 283–320.
- [Na88] S. Nag, The complex analytic theory of Teichmüller spaces, Wiley, New York, 1988.
- [Ne49] Z. Nehari, The Schwarzian derivative and schlicht functions, Bull. Amer. Math. Soc. 55 (1949), 545–551.
- [Sa78] J. H. Sampson, Some properties and applications of harmonic mappings, Ann. Sci. École Norm. Sup. 11 (1978), no. 2, 211–228.
- [Sc84] R. M. Schoen, Analytic aspects of the harmonic map problem, in Seminar on nonlinear partial differential equations, Math. Sci. Res. Inst. Publ., vol. 2 (1984), Springer-Verlag, New York-Berlin-Heidelberg, 321–358.
- [Sc96] K. P. Scannell, Flat conformal structures and the classification of de Sitter manifolds, To appear, Comm. Anal. Geom. (1998).
- [Sh87] H. Shiga, Projective structures on Riemann surfaces and Kleinian groups, J. Math. Kyoto Univ. 27 (1987), no. 3, 433–438.
- [ST95] H. Shiga and H. Tanigawa, *Projective structures with discrete holonomy representations*, Trans. Amer. Math. Soc., to appear (1995).
- [ST83] D. P. Sullivan and W. P. Thurston, Manifolds with canonical coordinates: some examples, Enseign. Math. 29 (1983), 15–25.
- [Ta97] H. Tanigawa, Grafting, harmonic maps, and projective structures on surfaces, J. Differential Geom. 47 (1997), no. 3, 399–419.
- [Th82] W. P. Thurston, *The Geometry and Topology of Three-Manifolds*, Princeton Univ., Princeton, 1982.
- [Wo91a] M. Wolf, High energy degeneration of harmonic maps between surfaces and rays in Teichmüller space, Topology 30 (1991), no. 4, 517–540.
- [Wo91b] _____, Infinite energy harmonic maps and degeneration of hyperbolic surfaces in moduli space, J. Differential Geom. **33** (1991), 487-539.