

# BLOW-UPS OF CANONICAL SINGULARITIES

YU. G. PROKHOROV

## INTRODUCTION

In this paper we apply Shokurov's inductive method to study terminal and canonical singularities. As an easy consequence of the Log Minimal Model Program we show that for any three-dimensional log terminal singularity  $(X \ni P)$  there exist some special, so called, plt blow-ups  $Y \rightarrow X$  (Proposition 2.9). We discuss properties of them and construct some examples. Roughly speaking a plt blow-up  $Y \rightarrow X$  is a blow-up with an irreducible exceptional divisor  $S$  such that  $S$  is normal and the pair  $(S, \text{Diff}_S(0))$  is log terminal, where  $\text{Diff}_S(0)$  is the different [Sh, §3]. These blow-ups are very useful for inductive approach to the study of singularities and, more general, extremal contractions [P2]. We also obtain necessary conditions for a log surface to be an exceptional divisor of a plt blow-up of a terminal singularity (Corollary 3.3, Proposition 3.7). Unfortunately a plt blow-up of a terminal singularity is never unique (see 4.8). However there are canonical singularities for which it is unique. These singularities are called weakly exceptional (see Definition 4.1) and have the most interesting and complicated structure (cf. [MP], [MP1]). We obtain the criterion for a singularity to be weakly exceptional, in terms of the exceptional divisor of some plt blow-up (Theorem 4.3) and construct few examples.

**Acknowledgements.** I am grateful to V. V. Shokurov for useful discussions concerning complements and exceptionality. I wish to thank also D. Markushevich for many helpful conversations during my stay at the University of Lille 1 in the spring of 1998. This work was partially supported by a grant 96-01-00820 from the Russian Foundation of Basic Research and a grant PECO-CEI from the Ministry of Higher Education of France.

## 1. PRELIMINARY RESULTS

All varieties are algebraic and are assumed to be defined over  $\mathbb{C}$ , the complex number field. We follow the terminology and notation of the log Minimal Model Program [KMM], [Sh], [Ut] (see also [Ko] for an introduction to singularities of pairs). We will use the following abbreviations:

lc	log canonical [KMM]
klt	Kawamata log terminal [Sh], [Ut], [Ko]
plt	purely log terminal [Sh], [Ut], [Ko]
dlt	divisorial log terminal [Sh], [Ut]

$LCS(X, D)$  the locus of lc singularities of  $(X, D)$  [Sh, 3.14], [K1]

A *contraction* (or *extraction*, if we start with  $X$  instead of  $Y$ ) is a projective morphism of normal varieties  $f: Y \rightarrow X$  such that  $f_*\mathcal{O}_Y = \mathcal{O}_X$ . A *blow-up* is a birational extraction. Usually we write  $f: (Y, S) \rightarrow X$  to specify that  $S$  is the (reduced) exceptional divisor of  $f$ . Two blow-ups  $f: Y \rightarrow X$  and  $f': Y' \rightarrow X$  are said to be *birationally equivalent* if the map  $(f^{-1} \circ f'): Y' \dashrightarrow Y$  is an isomorphism in codimension one. We will work in the category of algebraic varieties, however many results can be easily modified to the category of analytic spaces.

**Theorem 1.1** ([Sh2, 3.1], [Ut, 17.10], cf. [Sh, 5.19]). *Let  $X$  be a normal variety of dimension  $\leq 3$  and let  $D$  be a boundary on it such that  $(X, D)$  is lc. Then there exists a blow-up  $f: Y \rightarrow X$  such that*

- (i) *every exceptional divisor  $E_i$  of  $f$  has the discrepancy  $a(E_i, D) = -1$  (and therefore one can write  $f^*(K_X + D) = K_Y + D_Y + \sum E_i$ , where  $D_Y$  is the proper transform of  $D$ );*
- (ii)  *$(Y, D_Y + \sum E_i)$  is dlt;*
- (iii)  *$Y$  is  $\mathbb{Q}$ -factorial.*

We call such a blow-up *minimal log terminal blow-up* of  $(X, D)$ .

**Definition 1.2** ([Sh, §3], [Ut, Ch. 16]). Let  $X$  be a normal variety and let  $S+B$  be a boundary on  $X$ , where  $S = \lfloor S+B \rfloor \neq \emptyset$  and  $B = \{S+B\}$ . Assume that  $K_X + S + B$  is lc in codimension two. Then the *different* of  $B$  on  $S$  is defined by

$$K_S + \text{Diff}_S(B) \equiv (K_X + S + B)|_S$$

**Theorem 1.3** ([Sh, 3.9], [Ut, 16.6]). *Let  $(X, S)$  be a plt pair, where  $S$  is reduced. Then  $\text{Diff}_S(0) = \sum (1 - 1/m_i)R_i$ , where  $R_i$  are irreducible components of  $\text{Sing}(X)$  contained on  $S$  and having codimension one in  $S$ . The numbers  $m_i \in \mathbb{N}$  can be characterized by one of the following properties:*

- (i)  *$m_i$  is the index of  $K_X + S$  at the general point of  $R_i$ ;*
- (ii)  *$m_i$  is the index of  $S$  at the general point of  $R_i$ .*

We say that coefficients of a  $\mathbb{Q}$ -divisor  $D$  are *standard* if they have the form  $(1 - 1/m_i)$ , where  $m_i \in \mathbb{N} \cup \{\infty\}$ .

**Theorem 1.4** (Inversion of Adjunction [Sh, 3.3], [Ut, 17.6]). *Let  $X$  be a normal variety and let  $D$  be a boundary on it. Write  $D = S + B$ ,*

where  $S = \lfloor D \rfloor$  and  $B = \{D\}$ . Assume that  $K_X + S + B$  is  $\mathbb{Q}$ -Cartier. Then  $(X, S + B)$  is plt near  $S$  iff  $S$  is normal and  $(S, \text{Diff}_S(B))$  is klt.

**Definition 1.5** ([Sh, 5.1]). Let  $X$  be a normal variety and let  $D = S + B$  be a boundary, where  $S = \lfloor D \rfloor$  is the integer part and  $B = \{D\}$  is the fractional part. Then one says that  $K_X + D$  is *n-complementary*, if there exists a  $\mathbb{Q}$ -divisor  $D^+$ , such that the following conditions are verified:

- (i)  $n(D^+ + K_X) \sim 0$  in particular  $nD^+$  has integer coefficients;
- (ii)  $nD^+ \geq nS + \lfloor (n+1)B \rfloor \geq 0$ ;
- (iii)  $K_X + D^+$  is lc.

The log canonical divisor  $K_X + D^+$  is called an *n-complement* of  $K_X + D$ .

Note that in general it is not true that  $D^+ \geq D$ . This however is true if the coefficients of  $D$  are standard [Sh1, 2.7].

**Proposition 1.6** ([Sh, 5.4], [Sh1, 4.4]). *Let  $f: Y \rightarrow X$  be a blow-up and let  $D$  be a boundary on  $Y$ . Then*

- (i)  $K_Y + D$  is *n-complementary*  $\implies K_X + f(D)$  is *n-complementary*.
- (ii) Assume additionally that  $K_Y + D$  is *f-nef* and  $f(D)$  has only standard coefficients. Then

$$K_X + f(D) \text{ is } n\text{-complementary} \implies K_Y + D \text{ is } n\text{-complementary}.$$

Combining [Ut, 19.6] with 1.6 we get the following.

**Proposition 1.7** ([P1, 2.1], cf. [Ut, 19.6]). *Let  $(X, S)$  be a purely log terminal pair with reduced  $S \neq 0$  and let  $f: X \rightarrow Z$  be a projective morphism such that  $-(K_X + S)$  is *f-nef* and *f-big*. Given an *n-complement*  $K_S + \text{Diff}_S(0)^+$  of  $K_S + \text{Diff}_S(0)$ , then in a neighborhood of any fiber of  $f$  meeting  $S$  there exists an *n-complement*  $K_X + S + B$  of  $K_X + S$  such that  $\text{Diff}_S(0)^+ = \text{Diff}_S(B)$ .*

*n-complements* with  $n \in \{1, 2, 3, 4, 6\}$  are called *regular*.

**Proposition 1.8** ([Sh1]). *Let  $(S, \Theta)$  be a projective log surface such that  $(S, \Theta)$  is lc but is not klt and  $-(K_S + \Theta)$  is nef and big (i. e.  $(S, \Theta)$  is a weak log Del Pezzo surface, see [Sh1]). Then  $K_S + \Theta$  has a regular non-klt complement  $K_S + \Theta^+$ . Moreover, we can take  $\Theta^+$  so that  $a(E, \Theta) = -1$  implies  $a(E, \Theta^+) = -1$  for every (not necessary exceptional divisor  $E$ ).*

*Proof.* Let  $f: \tilde{S} \rightarrow S$  be a log resolution of  $(S, \Theta)$ . Consider the crepant pull back  $K_{\tilde{S}} + E + D = f^*(K_S + \Theta)$ , where  $E$  is a reduced divisor,  $D$  is an effective  $\mathbb{Q}$ -divisor and  $\lfloor D \rfloor \leq 0$ . By [Sh, 5.2]  $K_E + \text{Diff}_E(D)$  is

1, 2, 3, 4 or 6-complementary and by [Ut, 19.6] this complement can be extended on  $\tilde{S}$ . Finally, we can push down it on  $S$  by 1.6.  $\square$

The following proposition can be proved in the same style.

**Proposition 1.9** ([Sh1], [P]). *Let  $f: S \rightarrow Z$  be a contraction from a surface  $S$  to a curve  $Z$  and let  $\Theta$  be a boundary on  $S$ . Assume that  $(S, \Theta)$  is lc and  $-(K_S + \Theta)$  is  $f$ -ample. Then near a fiber  $f^{-1}(o)$ ,  $o \in Z$  the log divisor  $K_S + \Theta$  has a regular non-klt complement  $K_S + \Theta^+$ . Moreover, if  $(S, \Theta)$  is not klt, then we can take  $\Theta^+$  so that  $a(E, \Theta) = -1$  implies  $a(E, \Theta^+) = -1$  for every (not necessary exceptional divisor  $E$ ).*

## 2. DEFINITIONS, EXAMPLES AND PROPERTIES OF PLT BLOW-UPS

**Definition 2.1.** Let  $X$  be a normal algebraic variety and let  $f: Y \rightarrow X$  be a blow-up such that the exceptional locus of  $f$  contains only one irreducible divisor, say  $S$ . Then  $f: (Y, S) \rightarrow X$  is called a *plt blow-up* of  $X$  if  $(Y, S)$  is plt and  $-(K_Y + S)$  is  $f$ -ample.

This definition can be easily modified for blow-ups of analytic spaces in a usual way (recall that we are working in the algebraic category).

*Remark 2.2.* (i) Note that if  $X$  is  $\mathbb{Q}$ -factorial, then  $\rho(Y/X) = 1$  and  $Y$  is also  $\mathbb{Q}$ -factorial. Indeed, let  $D \neq S$  be any effective Weil divisor on  $Y$  intersecting  $S$ . Then we have  $f^*(f(D)) \sim_{\mathbb{Q}} D + \alpha S$ , where  $\alpha > 0$ . Thus  $D \sim_{\mathbb{Q}} -\alpha S$  over  $X$ . Since  $\text{Weil}(Y)/\text{Weil}(X)$  is generated by  $S$ , the variety  $Y$  is  $\mathbb{Q}$ -factorial.

(ii) If  $f: (Y, S) \rightarrow X$  is a blow-up of a variety  $X$  with only klt singularities such that  $\rho(Y/X) = 1$  and  $(Y, S)$  is plt, then  $-(K_Y + S)$  is  $f$ -ample automatically, so  $f$  is a plt blow-up.

(iii) Any plt blow-up  $f: (Y, S) \rightarrow X$  is uniquely defined up to isomorphism over  $X$  by the discrete valuation corresponding to  $S$  (cf. [Ut, Proof of 6.2]).

*Remark 2.3.* By Inversion of Adjunction  $S$  is normal and  $(S, \text{Diff}_S(0))$  is klt. In dimension two  $S$  is an irreducible non-singular curve and since  $-(K_S + \text{Diff}_S(0))$  is ample,  $S \simeq \mathbb{P}^1$ . In dimension three two cases are possible:  $f(S)$  is a point and then  $(S, \text{Diff}_S(0))$  is a log Del Pezzo surface, or  $\dim(f(S)) = 1$  and then  $S \rightarrow f(S)$  is generically  $\mathbb{P}^1$ -fibration.

**Example 2.4.** We describe below the class of all  $(S, \text{Diff}_S(0))$ 's for two-dimensional Du Val singularities (see e. g. [P]). In this case  $S \simeq \mathbb{P}^1$  and  $\text{Diff}_S(0) = \sum_{i=1}^r (1 - 1/m_i) P_i$ , where  $P_1, \dots, P_r \in \mathbb{P}^1$  are distinct points,  $r \geq 0$  and  $m_i \in \mathbb{N}$ . Since  $-(K_S + \text{Diff}_S(0))$  is ample,  $\sum (1 -$

$1/m_i) < 2$ . It is easy to show that for  $(m_1, \dots, m_r)$  one has one of the following possibilities.

- $r = 0, 1$  or  $2$ . In this case  $(X \ni P)$  is a cyclic quotient singularity (type  $A_n$ ) and  $f$  is toric;
- $(m_1, \dots, m_r) = (2, 2, m)$ ,  $m$  is arbitrary (type  $D_n$ );
- $(m_1, \dots, m_r) = (2, 3, 3)$  (type  $E_6$ );
- $(m_1, \dots, m_r) = (2, 3, 4)$  (type  $E_7$ );
- $(m_1, \dots, m_r) = (2, 3, 5)$  (type  $E_8$ ).

In cases  $D_n$  and  $E_n$  a plt blow-up is unique (see 4.7). Note that in two-dimensional case all (one-dimensional) log Fano varieties with standard boundaries can be realized as exceptional divisors of some plt blow-up. Similar description can be obtained for two-dimensional lt singularities.

We are interested in the three-dimensional case.

**Problem 2.5.** Describe the class of all log Del Pezzo surfaces (resp. generically  $\mathbb{P}^1$ -fibrations) which can be exceptional divisors of some plt blow-up of a terminal (resp. canonical,  $\epsilon$ -lt) singularity.

**Example 2.6.** Let  $(X \ni P)$  be a cyclic quotient singularity  $\mathbb{C}^3/\mathbb{Z}_m$ , where  $\mathbb{Z}_m$  acts on  $\mathbb{C}^3$  with weights  $(a_1, a_2, a_3)$ ,  $(a_1, a_2, a_3) = 1$ . Then any weighted blow-up  $f: Y \rightarrow X$  is a plt blow-up. The exceptional divisor  $S$  is a weighted projective space  $\mathbb{P}(a_1, a_2, a_3)$ ,  $\text{Diff}_S(0)$  is supported in the triangle  $\{x_1x_2x_3 = 0\}$  and has coefficients  $1 - \frac{1}{(a_i, a_j)}$ .

**Example 2.7.** Let  $\widetilde{\mathbb{C}^3} \rightarrow \mathbb{C}^3$  be the blow-up of  $x$ -axis and let  $\widetilde{\mathbb{C}^3}/\mathbb{Z}_m \rightarrow \mathbb{C}^3/\mathbb{Z}_m$  be a quotient by a cyclic group acting on  $\mathbb{C}^3$  free in codimension 1. Then  $\widetilde{\mathbb{C}^3}/\mathbb{Z}_m \rightarrow \mathbb{C}^3/\mathbb{Z}_m$  is a plt blow-up. The exceptional divisor  $S$  is a generically  $\mathbb{P}^1$ -bundle with two singular points on  $f^{-1}(0)$ .

Combining 1.7 with 1.6 (i) we get the following assertion which shows that the existence of complements can be established inductively.

**Corollary 2.8.** *Let  $f: (Y, S) \rightarrow X$  be a plt blow-up of a klt singularity. Assume that  $(S, \text{Diff}_S(0))$  is  $n$ -complementary. Then  $K_X$  is  $n$ -complementary.*

The following proposition shows that plt blow-ups are easy to construct.

**Proposition 2.9** (cf. [Sh2, 3.1], [Ut, 17.10], [Sh, 5.19]). *Let  $X$  be a normal  $\mathbb{Q}$ -factorial variety of dimension  $\leq 3$  and let  $D$  be a boundary on  $X$  such that  $K_X + D$  is lc, but is not plt. Assume that  $D \neq 0$  and  $X$  has at*

worse klt singularities. Then there exists a plt blow-up  $f: (Y, S) \rightarrow X$  such that

- (i)  $K_Y + S + D_Y = f^*(K_X + D)$  is lc;
- (ii)  $K_Y + S + (1 - \varepsilon)D_Y$  is plt and anti-ample over  $X$  for any  $\varepsilon > 0$ ;
- (iii)  $Y$  is  $\mathbb{Q}$ -factorial and  $\rho(Y/X) = 1$ .

This blow-up  $f: (Y, S) \rightarrow X$  is called a *plt blow-up* of  $(X, D)$ . Note that the proposition holds in arbitrary dimension modulo the Log Minimal Model Program.

*Proof.* By Theorem 1.1 there exists a blow-up  $h: W \rightarrow X$  with the reduced exceptional divisor  $E = \sum E_i$  such that  $W$  is  $\mathbb{Q}$ -factorial and  $K_W + D_W + E = h^*(K_X + D)$  is dlt. Note that  $\rho(W/X)$  is the number of components of  $E$ , because  $X$  is  $\mathbb{Q}$ -factorial. Moreover, since  $K_X + D$  is not plt, we have  $E \neq 0$ .

Now we apply the  $(K_W + E)$ -Minimal Model Program over  $X$ . The divisor  $K_W + E \equiv h^*(K_X + D) - D_W$  cannot be nef over  $X$ . Therefore on the last step we get a divisorial extremal contraction  $f: Y \rightarrow X$  with  $\rho(Y/X) = 1$ , with a unique exceptional divisor  $S$  and negative with respect to  $K_Y + S$ . Because  $K_W + E + D_W$  is numerically trivial, the divisor  $f^*(K_X + D) = K_Y + S + D_Y$  is lc on  $Y$ . Since  $K_Y + S$  is plt, so is  $K_Y + E + (1 - \varepsilon)D$  by [Ut, 2.17.5]. This completes the proof.  $\square$

**Proposition 2.10.** *Let  $f: (Y, S) \rightarrow X$  be a plt blow-up of a variety  $X$  with only klt singularities. Then there exists a boundary  $D$  on  $X$  such that  $(S, D)$  is lc,  $a(S, D) = -1$  and  $S$  is the only divisor over  $X$  with discrepancy  $a(S, D) \leq -1$ .*

*Proof.* For sufficiently big and divisible  $n$  the divisor  $-n(K_Y + S)$  is very ample over  $X$ . Take a general member  $L \in |-n(K_Y + S) + f^*(H)|$ , where the divisor  $H$  is ample enough on  $X$ . Put  $B := (1/n)L$ ,  $D := f(B)$ . Then  $(Y, S + B)$  is plt and since  $f^*(K_X + D) = K_Y + S + B$ , the pair  $(X, D)$  is lc (see [Ko, 3.10]).  $\square$

**Corollary 2.11.** *Notation as in Proposition 2.10. Then*

- (i)  $f(S)$  is the minimal center of  $LCS(X, D)$ ;
- (ii)  $f(S)$  is normal;
- (iii) if  $\text{codim}(f(S)) = 2$ , then there exists a boundary  $D_{f(S)}$  on  $f(S)$  such that  $(f(S), D_{f(S)})$  is klt.

*Proof.* Since  $S$  is the only divisor over  $X$  with discrepancy  $a(S, f(B)) = -1$ ,  $f(S)$  is the minimal center of lc singularities of  $(X, D)$ . It is normal by [K1], (iii) follows by [K2].  $\square$

The following is converse to Proposition 2.10.

**Proposition 2.12.** *Let  $X$  be a normal  $\mathbb{Q}$ -factorial variety of dimension  $\leq 3$  and let  $W \subset X$  be an irreducible subvariety. Assume there exists a boundary  $D$  such that  $(X, D)$  is lc and the minimal center of  $LCS(X, D)$  is  $W$ . Then there exists a plt blow up with center along  $W$ .*

*Proof.* Apply Proposition 2.9 to  $(X, D)$ . □

**Corollary 2.13.** *Let  $X$  be a normal three-dimensional variety and let  $f: (Y \supset S) \rightarrow X$  be a plt blow-up with center along a curve  $W = f(S)$ . Then for any point  $P \in W$  there exists a plt blow-up with center at  $P$ .*

### 3. THE TERMINAL CASE

**Proposition 3.1.** *Let  $(X \ni P)$  be a three-dimensional klt singularity and let  $f: (Y, S) \rightarrow X$  be a plt blow-up. Assume either*

- (i)  $\dim f(S) = 1$ , or
- (ii) *there exists a boundary  $D$  on  $X$  such that  $\lfloor D \rfloor \neq \emptyset$  near  $P$  and  $(X, D)$  is plt.*

*Then  $K_S + \text{Diff}_S(0)$  has a regular non-klt complement (near the fiber  $f^{-1}(P)$  in the case  $\dim f(S) = 1$ ).*

*Proof.* The assertion (i) follows from Proposition 1.9. To prove (ii) we assume first that  $X$  is  $\mathbb{Q}$ -factorial. Write  $f^*(K_X + D) = K_Y + D_Y + aS$ , where  $D_Y$  is the proper transform of  $D$  on  $Y$  and  $a < 1$ . By Connectedness Lemma [Sh, 5.7], [Ut, 17.4] the set of lc singularities  $LCS(Y, D_Y + S)$  is connected. It is clear that  $LCS(Y, D_Y + S)$  contains  $S$  and  $\lfloor D_Y \rfloor$ . Whence  $K_Y + D_Y + S$  is not plt (see [Sh, 3.6]). Since  $K_Y + S$  is plt,  $K_Y + bD_Y + S$  is lc but is not plt for some  $0 < b \leq 1$ . Therefore  $K_S + \text{Diff}_S(bD_Y)$  is lc but not klt (see Theorem 1.4 and [Ut, 17.7]). On the other hand  $-(K_S + \text{Diff}_S(bD_Y))$  is ample. The assertion follows by Proposition 1.8. If  $X$  is not  $\mathbb{Q}$ -factorial, then we consider a  $\mathbb{Q}$ -factorialization  $h: Y' \rightarrow Y$  (see [Ut, 6.11.1]) and let  $f': (Y', S') \rightarrow X$  be the composition map. The same arguments show that  $K_{S'} + \text{Diff}_{S'}(0)$  has a regular non-klt complement  $K_{S'} + \text{Diff}_{S'}(0)^+$  (we have to remark that in this case  $-(K_{S'} + \text{Diff}_{S'}(0))$  is nef and big because  $h$  is a small morphism). Finally, to conclude the proof we can push  $K_{S'} + \text{Diff}_{S'}(0)^+$  down on  $S$  by (i) of Proposition 1.6. □

**Corollary 3.2.** *In notation of Proposition 3.1  $K_Y + S$  has a regular non-plt complement.*

The following is a necessary condition for Problem 2.5 in the terminal case.

**Corollary 3.3.** *Let  $(X \ni P)$  be a three-dimensional terminal singularity and let  $f: (Y, S) \rightarrow X$  be a plt blow-up. Then  $K_S + \text{Diff}_S(0)$  has a regular non-plt complement.*

*Proof.* Consider  $f: (Y \supset S) \rightarrow (X \ni P)$  as a morphism of analytic spaces. By [R, 6.4.B] the general element  $D \in |-K_X|$  has only a Du Val singularity at  $P$ . Inversion of Adjunction gives us that  $(X, D)$  is plt, so to complete the proof we can apply the analytic analog of Proposition 3.1.  $\square$

Note that 3.3 has no generalizations for canonical singularities (see [MP], [MP1] and Example 4.17).

Using the same arguments as in the proof of Proposition 3.1 one can show the following

**Proposition 3.4.** *Let  $(X \ni P)$  be a three-dimensional klt singularity, let  $f: (Y, S) \rightarrow X$  be a plt blow-up, and let  $K_X + D$  be an  $n$ -complement near  $P$ . Then one of the following holds*

- (i)  $a(S, D) = -1$  and  $K_Y + S + D_Y := f^*(K_X + D)$  is an  $n$ -complement of  $K_Y + S$ ;
- (ii)  $a(S, D) > -1$  and then there exists a regular complement of  $K_Y + S$  which is not plt.

**Corollary 3.5.** *Let  $(X \ni P)$  be a three-dimensional klt singularity, let  $f: (Y, S) \rightarrow X$  be a plt blow-up. Assume that  $K_X$  has a regular complement  $K_X + D$ . Then  $K_Y + S$  also has a regular complement  $K_Y + S + B$  such that  $K_Y + S + B = f^*(K_X + D)$  or  $K_Y + S + B$  is not plt.*

**Example 3.6.** Let  $(X \ni P)$  be the terminal  $cE_8$ -singularity given by the equation  $x_1^2 + x_2^3 + x_3^5 + x_4^5 = 0$  and let  $f: (Y, S) \rightarrow X$  be the weighted blow-up with weights  $(15, 10, 6, 6)$ . Then  $S$  is isomorphic to the cone over a rational normal curve of degree 5 and  $\text{Diff}_S(0) = (1/2)H_1 + (2/3)H_2$ , where  $H_1, H_2$  are hyperplane sections in a general position. Since  $(S, (1/2)H_1 + (2/3)H_2)$  is klt,  $f: (Y, S) \rightarrow X$  is a plt blow-up. It is easy to check that  $-K_S = (7/5)H$ . Since  $\text{Diff}_S(0)^+ \geq \text{Diff}_S(0)$ ,  $K_S + \text{Diff}_S(0)$  is not 1 or 2-complementary. On the other hand there exists a 3-complement  $K_S + (2/3)H_1 + (2/3)H_2 + 1/3L$ , where  $L$  is a line on  $S$ . Note however that this complement is klt. The only regular non-klt  $n$ -complement is  $K_S + (1/2)H_1 + (2/3)H_2 + L_1 + (1/6)L_2$ , where  $L_1, L_2$  are lines on  $S$ . Here  $n = 6$ .

Unfortunately the proof of [R, 6.4.B] uses the classification of terminal singularities and has no generalizations in higher dimensions. To



avoid this difficulty one can apply the method proposed by Shokurov in [Sh1, 7.8]. This method uses the classification of surface complements and computations of discrepancies. Remark also that the necessary condition of 3.3 is not sufficient. Below we give another necessary condition for Problem 2.5.

**Proposition 3.7.** *Let  $f: (Y, S) \rightarrow X$  be a plt blow-up of a terminal singularity of index  $r$  and let  $L$  be a curve on  $S$  such that  $L \cap \text{Sing}(S) = L \cap \text{Sing}(\text{Supp}(\text{Diff}_S(0))) = \emptyset$ . Write  $\text{Diff}_S(0) = \sum(1 - 1/m_i)\Delta_i$ , where  $m_i \in \mathbb{N}$  (see 1.3) and denote  $m := \text{lcm}\{m_i\}$ . Then*

$$-m(K_S + \text{Diff}_S(0)) \cdot L \geq 1 + 1/r. \quad (1)$$

Note that  $-m(K_S + \text{Diff}_S(0)) \cdot L$  can be computed on  $S$ , without using any information about whole  $Y$ .

*Proof.* We can write

$$K_Y + S = f^*K_X + (1 + a)S, \quad \text{where } a = a(S, 0, X).$$

Then along  $L$  divisors  $m(K_Y + S)$  and  $mS$  are Cartier by the following lemma (cf. 1.3).

**Lemma 3.8** ([P2]). *Let  $(Y, S)$  be a plt pair with a reduced boundary  $S \neq 0$ . Assume that near some point  $Q$  the surface  $S$  is non-singular and  $\text{Diff}_S(0) = (1 - 1/m)\Delta_1$ , where  $\Delta_1$  is a non-singular curve. Moreover, assume that  $Y$  is nonsingular outside  $S$ . Then near this point there is an analytic isomorphism  $(Y, S) \simeq (\mathbb{C}^3, \{x_1 = 0\})/\mathbb{Z}_m(0, 1, a)$ , where  $(a, m) = 1$ .*

*Proof.* Let  $\varphi: Y' \rightarrow Y$  be a cyclic cover such that  $S' := \varphi^{-1}(S)$  is a Cartier divisor (see [R, 3.6]). Then  $(Y', S')$  is plt by [Sh, §2], [Ut, 16.3]. Therefore the singularity  $Y'$  at  $Q' := \varphi^{-1}(Q)$  is terminal. Consider its canonical cyclic cover  $\psi: Y'' \rightarrow Y'$  and let  $S'' := \psi^{-1}(S')$ . The finite morphism of normal surfaces  $S'' \rightarrow S$  is étale outside  $\text{Supp}(\text{Sing}(Y)) = \text{Supp}(\text{Diff}_S(0))$ . Since the fundamental group  $\pi_1(S \setminus \text{Supp}(\text{Diff}_S(0)))$  is isomorphic to  $\mathbb{Z}$  (because  $(S, \text{Supp}(\text{Diff}_S(0))) \simeq (\mathbb{C}^2, \mathbb{C}^1)$ ), we have that  $\psi: Y'' \rightarrow Y'$  is a cyclic cover. Moreover,  $S'' \rightarrow S$  has a smooth ramification divisor. Hence  $S''$  is non-singular and so is  $Y''$ . The rest is obvious.  $\square$

Thus we have

$$-m(K_S + \text{Diff}_S(0)) \cdot L = -m(1 + a)S \cdot L,$$

where  $-m(K_S + \text{Diff}_S(0)) \cdot L = -m(K_Y + S) \cdot L$  and  $-mS \cdot L$  are positive integers. Since  $(X \ni P)$  is terminal of index  $r$ , we have  $a \geq 1/r$ . This gives us the desired inequality (1).  $\square$

More delicate method was proposed by Shokurov in [Sh1, 7.8]. It uses computations of all discrepancies of  $(X \ni P)$ .

**Example 3.9.** Assume that  $S$  is non-singular and  $\text{Diff}_S(0) = \emptyset$  (so  $S$  is a usual Del Pezzo surface). Then any curve on  $S$  satisfies conditions of 3.7. By (1)  $-K_S \cdot L > 1$ , so we have only two cases:  $S = \mathbb{P}^2$  or  $S = \mathbb{P}^1 \times \mathbb{P}^1$ . Note that in these cases  $Y$  is non-singular and  $Y \rightarrow X$  is Mori's extremal contraction. Similar, but more complicated, description can be done in the case when  $S$  has only Du Val singularities. In particular, then we have  $K_S^2 \neq 1$  and 7.

Corollary 3.3 and Proposition 3.7 show that log Del Pezzo surfaces which are exceptional divisors of plt blow-ups of a terminal singularities form a very restricted class. Regular complements of surfaces were classified in [Sh1, §6]. They are divided into types  $\mathbb{A}_m^n, \mathbb{E}1_m^n, \mathbb{D}_m^n, \mathbb{E}2_m^n, \mathbb{E}3_m^n, \mathbb{E}4_m^n$  and  $\mathbb{E}6_m^n$ . It is a very interesting problem to establish a correspondence between this classification and classification of terminal singularities [R]. For example, the complements  $(S, \text{Diff}_S(0)^+)$  from 3.6 are of types  $\mathbb{E}3_0^0$  and  $\mathbb{E}6_0^1$ .

#### 4. ON UNIQUENESS OF PLT BLOW-UPS

In this section we discuss the uniqueness property of plt blow-ups.

**Definition 4.1.** Let  $(X \ni P)$  be a klt singularity. It said to be *weakly exceptional* if there exists only one plt blow-up (up to birational equivalence). A lc pair  $(X, D)$  is said to be *exceptional* if there exists at most one divisor  $E$  over  $X$  with discrepancy  $a(E, D) = -1$ . A normal klt singularity  $(X \ni P)$  is said to be *exceptional* if  $(X, D)$  is exceptional for any boundary  $D$  whenever  $K_X + D$  is log canonical (see [Sh1, 1.5]).

**Proposition 4.2** ([MP1]). *Let  $(X \ni P)$  be an exceptional  $\mathbb{Q}$ -factorial singularity. Then the divisor  $S$  from Definition 4.1 with discrepancy  $a(S, D) = -1$  is birationally unique. Therefore if a singularity is exceptional, then it is also weakly exceptional.*

Below we give criterions for a klt singularity to be (weakly) exceptional in terms of plt blow-ups.

**Theorem 4.3.** *Let  $(X \ni P)$  be a  $\mathbb{Q}$ -factorial klt singularity of dimension  $\leq 3$  and let  $f: (Y, S) \rightarrow X$  be a plt blow-up of  $P$ . Then the following are equivalent*

- (i)  $(X \ni P)$  is not weakly exceptional;
- (ii) there is an effective  $\mathbb{Q}$ -divisor  $\Theta \geq \text{Diff}_S(0)$  such that  $-(K_S + \Theta)$  is ample and  $(S, \Theta)$  is not lc;

(iii) *there is an effective  $\mathbb{Q}$ -divisor  $\Theta \geq \text{Diff}_S(0)$  such that  $-(K_S + \Theta)$  is ample and  $(S, \Theta)$  is not klt<sup>1</sup>.*

*Proof.* Note that  $\rho(Y/X) = 1$  and  $Y$  is  $\mathbb{Q}$ -factorial (because so is  $X$ ). First we show the implication (i)  $\implies$  (ii). Assume that  $(X \ni P)$  is not weakly exceptional. By Proposition 2.10 there exists a boundary  $D$  such that  $(X, D)$  is lc and  $a(S, D) > -1$ ,  $a(E, D) = -1$  for some  $E \neq S$ . Let us write  $K_Y + \alpha S + D_Y = f^*(K_X + D)$ , where  $D_Y$  is the proper transform of  $D$  and  $\alpha < 1$ . This divisor is lc but is not plt. Then  $-(K_Y + S + D_Y)$  is ample over  $X$  and  $(Y, S + D_Y)$  is not lc. Take  $\Theta = \text{Diff}_S(D_Y)$ .

The implication (ii)  $\implies$  (iii) is clear. To show (iii)  $\implies$  (ii) we replace  $\Theta$  on  $\mathbb{Q}$ -Cartier divisor  $\Theta + \epsilon(\Theta - \text{Diff}_S(0))$  for some small  $\epsilon \geq 1$ .

Finally, we prove (ii)  $\implies$  (i). By 2.13 we may assume that  $f(S)$  is a point. For sufficiently large and divisible  $n$  the integer Cartier divisor  $-n(K_S + \Theta)$  is a very ample. Take  $H \in |-n(K_S + \Theta)|$  and denote  $\Xi := \Theta + \frac{1}{n}H$ . Then  $K_S + \Xi$  is numerically trivial and is not lc. Moreover, we may assume that  $n(K_S + \Xi) \sim 0$ . From the exact sequence

$$\begin{aligned} 0 \longrightarrow \mathcal{O}_Y(K_Y - (n+1)(K_Y + S)) \longrightarrow \mathcal{O}_Y(-n(K_Y + S)) \\ \longrightarrow \mathcal{O}_S(-n(K_S + \text{Diff}_S(0))) \longrightarrow 0 \end{aligned}$$

and the Kawamata-Viehweg vanishing  $H^1(Y, \mathcal{O}_Y(K_Y - (n+1)(K_Y + S))) = 0$  we get that the map

$$H^0(Y, \mathcal{O}_Y(-n(K_Y + S))) \longrightarrow H^0(S, \mathcal{O}_S(-n(K_S + \text{Diff}_S(0))))$$

is surjective. Thus there exists an integer effective divisor  $L \in |- (K_Y + S)|$  such that  $L|_S = n(\Xi - \text{Diff}_S(0))$ . Consider  $\mathbb{Q}$ -Cartier divisor  $B := \frac{1}{n}L$ . Then  $n(K_Y + S + B) \sim 0$  and  $(K_Y + S + B)|_S = K_S + \Xi$ . By [Ut, 17.7]  $K_Y + S + B$  is not lc. We can take  $B' = \alpha B$ ,  $\alpha < 1$  such that  $K_Y + S + B'$  is lc but is not plt. It is easy to see that

$$-(K_Y + S + B') \equiv (1 - \alpha)B \equiv -(1 - \alpha)(K_Y + S) \quad (2)$$

is  $f$ -ample. Further, we take a minimal log terminal blow-up (see 1.1)  $g: W \rightarrow Y$ . We have  $K_W + S_W + E + B'_W = g^*(K_Y + S + B')$ , where  $S_W$  and  $B'_W$  are proper transforms of  $S$  and  $B'$ , respectively, and  $E \neq \emptyset$  is the reduced exceptional divisor. One can see by (2)

$$K_W + S_W + E + B'_W \equiv -(1 - \alpha)(g^*(B)). \quad (3)$$

Apply the Log Minimal Model Program to  $(W, S_W + E + B'_W)$  over  $X$ . On each step by (3)  $K_W + S_W + E + B'_W$  cannot be nef over  $X$ .

<sup>1</sup>The author would like to thank Dr. O. Fujino, who pointed out some mistakes in this theorem

Let  $V \rightarrow X$  be the last step (i. e.  $V \rightarrow X$  contracts an irreducible exceptional divisor, say  $R$ ). By construction  $K_V + R + B'_V$  is plt. Since  $X$  is  $\mathbb{Q}$ -factorial,  $\rho(V/X) = 1$  and  $(V, R) \rightarrow X$  is a plt blow-up. It is sufficient to show that  $R$  is the proper transform of some of components of  $E$  (otherwise  $(X \ni P)$  is not weakly exceptional). Indeed, in the opposite case  $(V, R) \rightarrow X$  is birationally equivalent to  $(Y, S) \rightarrow X$ . Hence they are isomorphic because  $\rho(Y/X) = \rho(V/X) = 1$ . But then  $K_Y + S + B'$  is plt. This contradiction concludes the proof.  $\square$

**Corollary 4.4.** *Notation as in Theorem 4.3. If a three-dimensional singularity  $(X \ni P)$  is not weakly exceptional, then  $K_S + \text{Diff}_S(0)$  has a regular non-klt complement  $K_S + \text{Diff}_S(0)^+$ . Moreover, one can choose  $\text{Diff}_S(0)^+$  in such a way that  $a(E, \text{Diff}_S(0)^+) = -1$ ,  $a(E, \Theta) \leq -1$  for some (not necessary exceptional divisor  $E$ ).*

*Proof.* Apply Proposition 1.8 or 1.9.  $\square$

**Corollary 4.5.** *Notation as in 4.3 and 4.4. If a three-dimensional singularity  $(X \ni P)$  is not weakly exceptional, then  $K_X$  has a regular complement. Moreover,  $K_Y + S$  has a regular non-plt complement.*

**Corollary 4.6.** *Let  $(X \ni P)$  be a klt singularity and let  $D$  be a boundary on  $X$  such that  $\lfloor D \rfloor \neq \emptyset$  near  $P$ . If  $(X, D)$  is plt and  $\dim(X) \leq 3$ , then  $(X \ni P)$  is not weakly exceptional.*

*Proof.* Let  $f: (Y, S) \rightarrow X$  be a plt blow-up. Write  $K_Y + D_Y + \alpha S = f^*(K_X + D)$ , where  $\alpha < 1$  and  $D_Y$  is the proper transform of  $D$ . Then  $K_Y + D_Y + S$  is anti-ample over  $X$  and it is not plt. Therefore in Theorem 4.3 we can take  $\Theta = \text{Diff}_S(D_Y)$ .  $\square$

**Example 4.7.** Let  $(X \ni P)$  be a two-dimensional Du Val singularity. By 4.6 and by [K, §9] it is not weakly exceptional only if it is of type  $A_n$ . On the other hand singularities of type  $A_n$  are not weakly exceptional (cf. 2.6). In cases  $D_n$ ,  $E_6$ ,  $E_7$  and  $E_8$  the unique plt blow-up is the blow-up of the "central" vertex of the dual graph.

Similar to Proposition 3.1 one can prove

**Corollary 4.8.** *Let  $(X \ni P)$  be a three-dimensional terminal singularity. Then it is not weakly exceptional.*

Now we prove a criterion of the exceptionality. In contrast to Theorem 4.3 it does not depend on the Minimal Model Program.

**Theorem 4.9.** *Let  $(X \ni P)$  be a klt singularity and let  $f: (Y, S) \rightarrow X$  be a plt blow-up of  $P$ . Then the following are equivalent*

- (i)  $(X \ni P)$  is non-exceptional;
- (ii) there is an effective  $\mathbb{Q}$ -divisor  $\Theta \geq \text{Diff}_S(0)$  such that  $n(K_S + \Theta) \sim 0$  for some  $n \in \mathbb{N}$  and  $K_S + \Theta$  is not klt;
- (iii) there is an effective  $\mathbb{Q}$ -divisor  $\Xi \geq \text{Diff}_S(0)$  such that  $-(K_S + \Xi)$  is nef (in particular  $\mathbb{Q}$ -Cartier) and  $(S, \Xi)$  is not klt;
- (iv) (in dimension 3 only) there is a regular (i. e. 1, 2, 3, 4 or 6) complement of  $K_S + \text{Diff}_S(0)$  which is not klt;
- (v) (in dimension 3 only) there is a regular complement  $K_X + D$  of  $K_X$  which is not exceptional and such that  $a(S, D) = -1$  (and then the set of divisors with discrepancy  $-1$  must be infinite).

*Proof.* Note that  $S$  and  $K_Y$  are  $\mathbb{Q}$ -Cartier, because  $(X \ni P)$  is klt. First we prove (ii)  $\Rightarrow$  (i). Assume that  $(X \ni P)$  is exceptional. Let  $\Theta$  be a boundary such as in (ii). Then  $\Theta \geq \text{Diff}_S(0)$  and  $K_S + \Theta \equiv 0$ . Take  $n \in \mathbb{N}$  such that  $n(\Theta - \text{Diff}_S(0))$  and  $n(K_Y + S)$  are integer Cartier divisors. It is clear that  $n(\Theta - \text{Diff}_S(0)) \in |-n(K_S + \text{Diff}_S(0))|$ . We have the exact sequence

$$\begin{aligned} 0 \longrightarrow \mathcal{O}_Y(K_Y - (n+1)(K_Y + S)) \longrightarrow \mathcal{O}_Y(-n(K_Y + S)) \\ \longrightarrow \mathcal{O}_S(-n(K_S + \text{Diff}_S(0))) \longrightarrow 0 \end{aligned}$$

By the Kawamata-Viehweg vanishing  $H^1(Y, \mathcal{O}_Y(K_Y - (n+1)(K_Y + S))) = 0$ , because  $-(K_Y + S)$  is ample. So there exists an integer effective Weil divisor  $L$  on  $Y$  such that  $(K_Y + S + \frac{1}{n}L)_S = K_S + \Theta$ . Denote  $B := \frac{1}{n}L$ . If  $K_S + \Theta$  is not klt, then by Inversion of Adjunction  $K_Y + S + B$  is not plt. Put  $D := f(B)$ . Thus we have either  $K_X + D$  is lc (and then there is an exceptional divisor  $E \neq S$  over  $X$  with  $a(E, D) = a(S, D) = -1$ ) or  $K_X + \alpha D$  is lc but is not klt for some  $\alpha < 1$ . Since  $a(S, D) = -1$ , there is an exceptional divisor  $E \neq S$  over  $X$  with  $a(E, \alpha D) = -1$ . Both cases gives us a contradiction.

The implications (ii)  $\Rightarrow$  (iii) and (iv)  $\Rightarrow$  (ii) are obvious. As for (iii)  $\Rightarrow$  (ii) we can take  $\Theta = \Xi + (1/n)F$ , where  $F \in |-n(K_S + \Xi)|$ , the linear system  $|-n(K_S + \Xi)|$  is not empty for  $n \gg 0$  by Base Point Free Theorem [KMM, 3-1].

Let us show (i)  $\Rightarrow$  (ii). By the assumption  $(X \ni P)$  is not exceptional. Therefore there exists a boundary  $D$  such that  $(X, D)$  is lc and  $a(E, D) = -1$  for some  $E \neq S$ . Let us write  $K_Y + \alpha S + B = f^*(K_X + D)$ , where  $\alpha \leq 1$ . Then  $-(K_Y + S + B)$  is nef over  $X$  and  $(Y, S + B)$  is not plt. Take  $\Theta = \text{Diff}_S(B)$ .

To see (v)  $\Rightarrow$  (iv) we can take the regular complement  $f^*(K_X + D)|_S$  on  $K_S + \text{Diff}_S(0)$  which is not klt by Inversion of Adjunction 1.4. The converse implication follows from Propositions 1.7 and 1.6 (i).

Finally, we prove (ii)  $\Rightarrow$  (iv). We may assume that  $f(S)$  is a point (otherwise there exists a regular non-klt complement by Proposition 1.9). Thus  $S$  is projective. Denote  $\text{Diff}_S(0)$  by  $\Delta = \sum \delta_i \Delta_i$ . Assume that there exists a boundary  $\Theta \geq \Delta$  on  $S$  such that  $K_S + \Theta$  is numerically trivial and is not klt. The implication (ii)  $\Rightarrow$  (iv) is an immediate consequence of Corollary 4.11 below which is a generalization of 1.8.  $\square$

**Theorem 4.10** ([Sh1, 2.3]). *Let  $S$  be a projective surface and let  $\Theta$  be a boundary on it with standard coefficients such that  $-(K_S + \Theta)$  is nef. Assume that  $(S, \Theta)$  is lc but is not klt and assume that the Mori cone  $\text{NE}(S)$  is polyhedral and has contractible faces. Then  $K_S + \Theta$  has a regular complement.*

**Corollary 4.11.** *Let  $S$  be a projective surface and let  $\Theta \geq \Delta$  be a boundaries on it. Assume that*

- (i)  $\Delta$  has only standard coefficients;
- (ii)  $(S, \Delta)$  is klt;
- (iii)  $-(K_S + \Delta)$  is ample;
- (iv)  $-(K_S + \Theta)$  is nef;
- (v)  $(S, \Delta)$  is lc but is not klt.

*Then  $K_S + \Theta$  has a regular complement which is not klt.*

*Proof.* Note that  $\text{NE}(S)$  is polyhedral and has contractible faces by Cone Theorem [KMM, 4-2-1]. First we consider the case when  $[\Theta] \neq 0$ . Let  $\Delta_1$  be a component of  $[\Theta]$ . Replace  $\delta_1$  with 1:  $\Delta' := \Delta_1 + \sum_{i \neq 1} \delta_i \Delta_i$ , so  $\Delta \leq \Delta' \leq \Theta$ . If  $-(K_S + \Delta')$  is nef, then we can apply 4.10 (because  $[\Delta'] \neq 0$  and  $\Delta'$  has only standard coefficients). Further, we assume that  $-(K_S + \Delta')$  is not nef. Then there exists a  $(K_S + \Delta)$ -extremal ray  $R$  such that  $(K_S + \Delta') \cdot R \geq 0$ . If it is birational, then we contract it. Since  $K_S + \Theta$  is trivial on  $R$ , it preserves the lc property of  $K_S + \Theta$  and  $K_S + \Delta'$ . Since  $\Delta'$  has only standard coefficient, we can pull-back complements of  $K_S + \Delta'$  (but here we are looking for regular complements of  $K_S + \Delta'$ , see Lemma 1.6). Note also that  $(\Delta' - \Delta) \cdot R > 0$ . Therefore the curve  $\Delta_1$  is not contracted. Continue the process we get a non-birational extremal ray  $R$  on  $S$  such that  $(K_S + \Delta') \cdot R > 0$ . But on another hand,  $(K_S + \Delta') \cdot R \leq (K_S + \Theta) \cdot R = 0$ , a contradiction.

Let now  $[\Theta] = 0$ , then at some point, say  $P$ , the divisor  $K_S + \Theta$  is lc, but is not plt. In this point  $K_S + \Delta$  is klt. Similar to Proposition 2.9 we can construct a blow-up  $f: \tilde{S} \rightarrow S$  with an irreducible exceptional divisor  $E$  over  $P$  such that  $a(E, \Theta) = -1$ , the crepant pull back  $K_{\tilde{S}} + \tilde{\Theta} + E = f^*(K_S + \Theta)$  is lc and  $K_{\tilde{S}} + \tilde{\Theta} + E$  is plt for any  $\tilde{\Theta} =$

$\tilde{\Delta} + \epsilon(\tilde{\Theta} - \tilde{\Delta})$ ,  $\epsilon < 1$ . Recall that  $\tilde{\Theta}$  is the proper transform of  $\Theta$ . Note also that  $\rho(\tilde{S}/S) = 1$ .

$$K_{\tilde{S}} + \tilde{\Delta} + \alpha E = f^*(K_S + \Delta),$$

where  $\alpha < 1$ . Assume that there exists a curve  $C$  such that  $(K_{\tilde{S}} + \tilde{\Delta} + E) \cdot C > 0$ . Then  $(\tilde{\Theta} - \tilde{\Delta}) \cdot C < 0$ . Therefore  $C$  is a component of  $(\tilde{\Theta} - \tilde{\Delta})$  and  $C^2 < 0$ . Further,  $C \cdot E > 0$ . Hence  $C \neq E$  and we can chose  $\tilde{\Theta}' < \tilde{\Theta}$  so that  $(K_{\tilde{S}} + \tilde{\Theta}') \cdot C < 0$ . Therefore the  $C$  is an  $(K_{\tilde{S}} + \tilde{\Theta}')$ -extremal curve and its contraction preserves the lc property of  $K_{\tilde{S}} + \tilde{\Theta} + E$ . Again we can pull-back complements of  $K_{\tilde{S}} + \tilde{\Delta} + E$  (see Lemma 1.6). Continue the process we get the situation when  $-(K_{\tilde{S}} + \tilde{\Delta} + E)$  is nef. All the steps preserve the nef and big property of  $-(K_{\tilde{S}} + \tilde{\Delta} + \alpha E)$ . Now we can apply Shokurov's Theorem 4.10. It is sufficient to prove that the cone  $\text{NE}(\tilde{S})$  is polyhedral and its faces are contractible. This an easy consequence of the following easy lemma which is a corollary of Cone Theorem [KMM, 4-2-1].

**Lemma 4.12** ([Sh1, 2.5]). *Let  $S$  be a projective surface and let  $D$  be a boundary on  $S$  such that  $(S, D)$  is klt and  $-(K_S + D)$  is nef and big (in particular,  $(S, D)$  is a weakly log Del Pezzo). Then the Mori cone  $\text{NE}(S)$  is polyhedral and its faces are contractible.*

*Proof.* By Base Point Free Theorem [KMM, 3-1-1] the linear system  $| -n(K_S + D) |$  for  $n \gg 0$  gives a birational contraction  $S \rightarrow S'$ . Let  $\sum E_i$  be the (reduced) exceptional divisor. For some small positive  $\epsilon_i$ 's the  $\mathbb{Q}$ -divisor  $-\sum \epsilon_i E_i$ , is ample over  $S'$ . The assertion follows by Cone Theorem [KMM, 4-2-1] applied to  $(S, D + \sum \epsilon_i E_i)$ .  $\square$

Further, if  $\alpha \geq 0$ , then we apply Lemma to  $(\tilde{S}, \tilde{\Delta} + \alpha E)$ . If  $\alpha < 0$ , then by the monotonicity  $-(K_{\tilde{S}} + \tilde{\Delta})$  is nef and big. Again apply Lemma 4.12. This concludes the proof of Corollary 4.11.  $\square$

**Corollary 4.13** ([Sh1]). *Let  $(X \ni P)$  be a three-dimensional klt non-exceptional singularity. Then it is 1, 2, 3, 4 or 6 complementary.*

**Corollary 4.14.** *Let  $(X \ni P)$  be a three-dimensional klt singularity. Then it is non-exceptional if and only if there exists a regular complement  $K_X + D$  such that the set  $\{E\}$  of divisors with discrepancy  $a(E, D) = -1$  is infinite.*

**Corollary 4.15** ([MP]). *Let  $(X \ni P)$  be a klt singularity and let  $D$  be a boundary on  $X$  such that  $[D] \neq \emptyset$  near  $P$ . If  $(X, D)$  is lc, then  $(X \ni P)$  is not exceptional,*

V. V. Shokurov pointed out to the author that one can replace the condition  $[D] \neq 0$  in Corollaries 4.6 and 4.15 on  $\max\{d_i\} \geq 41/42$  (see [Ko1]).

**Example 4.16** ([Sh, 5.2.3]). Let  $(X \ni P)$  be a two-dimensional Du Val singularity. By 4.15 and by [K, §9] it is exceptional if only if it is of type  $E_6$ ,  $E_7$  or  $E_8$ .

Theorems 4.3 and 4.9 allow us to construct many examples of three-dimensional exceptional and weakly exceptional singularities singularities (cf. [MP], [MP1]).

**Example 4.17.** Let  $X \subset \mathbb{C}^4$  be the hypersurface singularity  $x_1^3 + x_2^4 + x_3^4 + x_4^4 = 0$ . It is canonical by [R1]. Consider the weighted blow-up  $(Y, S) \rightarrow X$  with weights  $(4, 3, 3, 3)$ . The exceptional divisor  $S$  is defined into  $\mathbb{P}(4, 3, 3, 3)$  by the equation  $x_1^3 + x_2^4 + x_3^4 + x_4^4 = 0$ . It is easy to see that the projection  $S \rightarrow \mathbb{P}(3, 3, 3)$  is an isomorphism, so  $S \simeq \mathbb{P}^2$ . The variety  $Y$  is singular along the curve  $x_1 = x_2^4 + x_3^4 + x_4^4 = 0$  and along this curve  $(Y \supset S)$  is isomorphic to  $(\mathbb{C}^3 \supset \mathbb{C}^2)/\mathbb{Z}_3$ . Therefore  $(Y, S)$  is plt and  $(Y, S) \rightarrow X$  is plt blow-up. Moreover,  $\text{Diff}_S(0) = (2/3)C$ , where  $C$  is a non-singular curve of degree 4 on  $S = \mathbb{P}^2$ . Among regular complements there are only the following

- 3-complement  $K_{\mathbb{P}^2} + (2/3)C + (1/3)L$ , where  $L$  is a line,
- 6-complement  $K_{\mathbb{P}^2} + (2/3)C + (1/6)Q$ , where  $Q$  is a conic, and
- 6-complement  $K_{\mathbb{P}^2} + (2/3)C + (1/6)L_1 + (1/6)L_2$ , where  $L_1, L_2$  are lines.

All these complements are klt by the simple lemma below. Therefore the singularity is exceptional by Theorem 4.9.

**Lemma 4.18.** *Let  $D = \sum d_i D_i$  be a boundary on  $\mathbb{C}^2$ , where  $D_i$  are non-singular curves. If  $\sum d_i \leq 1$  (resp.  $< 1$ ), then  $(\mathbb{C}^2, D)$  is canonical (resp. terminal).*

**Example 4.19.** Let  $X \subset \mathbb{C}^4$  be the canonical hypersurface singularity  $x_1^3 + x_2^3 + x_3^3 + x_4^n = 0$ , where  $(n, 3) = 1$ ,  $n \geq 7$ . The hyperplane section  $x_4 = 0$  is lc. Therefore by Corollary 4.15  $(X \ni 0)$  is not exceptional. Consider the weighted blow-up  $(Y, S) \rightarrow X$  with weights  $(n, n, n, 3)$ . As in Example 4.17 we have  $S \simeq \mathbb{P}^2$ ,  $(Y, S)$  is plt and  $\text{Diff}_S(0) = (1 - 1/n)C$ , where  $C$  is a smooth cubic curve. By Theorem 4.3 and Proposition 1.8 the singularity is weakly exceptional.



## REFERENCES

- [K] Kawamata Y. *The crepant blowing-up of 3-dimensional canonical singularities and its application to the degeneration of surfaces*, Ann. Math. **127** (1988), 93–163
- [K1] Kawamata Y. *On Fujita’s freeness conjecture for 3-folds and 4-folds*, e-print alg-geom/9712014
- [K2] Kawamata Y. *Subadjunction of log canonical divisors for a subvariety of codimension 2*, in ”Birational Algebraic Geometry”, Proc. of Conf. in Memory of Wei-Liang Chow (1996), Contemporary Math. AMS **207**, 79–88
- [KMM] Kawamata Y., Matsuda K., Matsuki K. *Introduction to the minimal model program*, Adv. Stud. in Pure Math. **10** (1987), 283–360
- [Ko] Kollár J. *Singularities of pairs*, Proc. Symp. Pure Math. **62** (1997), 221–287
- [Kol1] Kollár J. *Log surfaces of general type; some conjectures*, Contemporary Math. **162** (1994), 261–275
- [MP] Markushevich D., Prokhorov Yu. G. *Klein’s group defines an exceptional singularity of dimension 3*, e-print math.AG/9805004, to appear in *J. Math. Sci.* (1999)
- [MP1] Markushevich D., Prokhorov Yu. G. *Exceptional quotient singularities*, e-print math.AG/9806029, to appear in *Amer. J. Math.*
- [P] Prokhorov Yu. G. *Lectures on complements on surfaces*, Lecture Note, Moscow State University 1997 (Russian)
- [P1] Prokhorov Yu. G. *Mori conic bundles with a reduced log terminal boundary*, e-print math.AG/9802120, to appear in *J. Math. Sci.* (1999)
- [P2] Prokhorov Yu. G. *Complements on conic bundles, I*, Proc. of I. R. Shafarevich seminar, 1996–1998, Steklov Inst. Math., Moscow, (1998), 161–175, (Russian; English transl. to appear in *J. Math. Sci.*)
- [R] Reid M. *Young persons guide to canonical singularities*, in ” Algebraic Geometry, Bowdoin, 1985,” Proc. Symp. Pure Math. **46** (1987) 345–414
- [R1] Reid M. *Canonical threefolds’ Géométrie Algébrique Angers*, A. Beauville ed., (1980) Sijthoff and Noordhoff., 273–310
- [Sh] Shokurov V. V. *3-fold log flips*, Izv. AN SSSR Ser. mat. **56** (1992), 105–201 & **57** (1993), 141–175; English transl. Russian Acad. Sci. Izv. Math. **40** (1993), 93–202 & **43** (1994), 527–558
- [Sh1] Shokurov V. V. *Complements on surfaces*, e-print alg-geom/9711024
- [Sh2] Shokurov V. V. *3-fold log models*, *J. Math. Sci.* **81** (1996), 2666–2699
- [Ut] Kollár J. et al. *Flips and abundance for algebraic threefolds*, A summer seminar at the University of Utah, Salt Lake City, 1991. Astérisque. **211** (1992)

*E-mail address:* prokhoro@mech.math.msu.su, prokhoro@math.pvt.msu.su,

DEPARTMENT OF MATHEMATICS (ALGEBRA SECTION), MOSCOW STATE UNIVERSITY, 117234 MOSCOW, RUSSIA