

MONOMIAL BASES OF QUANTIZED ENVELOPING ALGEBRAS

VYJAYANTHI CHARI AND NANHUA XI

ABSTRACT. We construct a monomial basis of the positive part \mathbf{U}^+ of the quantized enveloping algebra associated to a finite-dimensional simple Lie algebra. As an application we give a simple proof of the existence and uniqueness of the canonical basis of \mathbf{U}^+ .

0. Introduction

In [L1], Lusztig showed that the positive part \mathbf{U}^+ of the quantized enveloping algebra associated to a finite-dimensional simple Lie algebra \mathfrak{g} , had a remarkable basis called the canonical basis. The main idea in proving its existence and uniqueness was the following. Corresponding to every reduced expression \mathbf{i} of the longest element w_0 of the Weyl group of \mathfrak{g} one constructs a Poincaré–Birkhoff–Witt basis $B_{\mathbf{i}}$ of \mathbf{U}^+ . Lusztig proved in [L1] that the $\mathbf{Z}[q^{-1}]$ -lattice \mathcal{L} spanned by $B_{\mathbf{i}}$ is independent of the choice of \mathbf{i} and that the image of $B_{\mathbf{i}}$ in the \mathbf{Z} -module $\mathcal{L}/q^{-1}\mathcal{L}$ is a basis B independent of \mathbf{i} . Let $\overline{\mathcal{L}}$ be the image of \mathcal{L} under the bar map (a certain \mathbf{Q} -algebra involution) of \mathbf{U}^+ . The canonical basis \mathbf{B} is the preimage of B in $\mathcal{L} \cap \overline{\mathcal{L}}$.

In [K1], Kashiwara introduced the notion of crystal bases for the quantized algebras of classical type and later generalized it to quantized algebras associated to an arbitrary symmetrizable Kac–Moody algebra. The crystal basis is a base ‘at $q = 0$ ’ of \mathbf{U}^+ with certain properties. Later, he proved that the crystal bases could be ‘melted’ to give a basis of \mathbf{U}^+ itself, which is called the global crystal base. The main tool used here was a certain bilinear form on the algebra, and the global crystal basis can be characterized as a bar-invariant quasi-orthonormal basis with respect to this form. In [X2], Xi proved that the bases $B_{\mathbf{i}}$ are quasi-orthonormal with respect to this form.

The quantized enveloping algebra also admits another symmetric bilinear form introduced by Drinfeld, and Lusztig proved in [L2] that the bases $B_{\mathbf{i}}$ are quasi-orthonormal and that the canonical basis can be characterized as the bar-invariant, quasi-orthonormal basis of \mathbf{U}^+ with respect to the Drinfeld form. It is now proved [GL] that in fact the global crystal basis and the canonical basis are the same.

In this paper we construct a basis of \mathbf{U}^+ whose terms are monomials in the Chevalley generators and hence is bar-invariant but not quasi-orthonormal. We are then able to give a very simple proof of the existence and uniqueness of the canonical basis. Our construction of the monomial basis depends on picking a specific reduced expression for w_0 . We conjecture that in fact there exists a monomial basis corresponding to every reduced expression. In view of [BCP] we expect also that similar results should be true for the quantized affine algebras.

1991 *Mathematics Subject Classification.* 17B.

Key words and phrases. quantized enveloping algebra, monomial basis.

1. PRELIMINARIES


Throughout this paper \mathfrak{g} will denote a finite-dimensional complex simple Lie algebra, $(a_{ij})_{i,j \in I}$, $I = \{1, \dots, n\}$, will denote its Cartan matrix and $\mathbf{N}[I]$ will denote the set of linear combinations $\nu = \sum_i \nu_i i$, $\nu_i \in \mathbf{Z}, \nu_i \geq 0$. Let W be the Weyl group of \mathfrak{g} . It is well-known that W is a Coxeter group generated by simple reflections s_i for $i \in I$. Let $l(w)$ denote the length of a reduced expression of w and let w_0 be the unique element of maximal length in W . Let R^+ denote a set of positive roots of \mathfrak{g} and let $\alpha_1, \dots, \alpha_n$ be the set of simple roots. Corresponding to any reduced expression of w_0 , say $w_0 = s_{k_1} s_{k_2} \dots s_{k_r}$, $r = |R^+|$, we have a total order on R^+ ,

$$\beta_1 < \beta_2 < \dots < \beta_r$$


where $\beta_i = s_{k_1} s_{k_2} \dots s_{k_{i-1}} \alpha_{k_i}$.

For the rest of the paper we shall be working with two specific reduced expressions of w_0 . Since the numbering of the nodes is important for our purposes, we include below, for the readers convenience, the Dynkin diagrams of the various finite-dimensional complex simple Lie algebras. We assume that the node α_1 is short if \mathfrak{g} is of type F_4 or G_2 .


Type A_n ($n \geq 1$).




Type B_n ($n \geq 2$).




Type C_n ($n \geq 3$).




Type D_n ($n \geq 4$).




Type F_4




Type G_2




Type E_6



Type E_7



Type E_8



Let $\gamma_n \in W$ be defined as follows:

$$\gamma_n = \begin{cases} s_1 s_2 \cdots s_n & \text{if } \mathfrak{g} \text{ is of type } A_n, \\ s_n s_{n-1} \cdots s_2 s_1 s_2 s_3 \cdots s_{n-1} s_n & \text{if } \mathfrak{g} \text{ is of type } B_n \text{ or } C_n, \\ s_n s_{n-1} \cdots s_3 s_1 s_2 s_3 \cdots s_{n-1} s_n & \text{if } \mathfrak{g} \text{ is of type } D_n. \end{cases}$$

It is easy to see that

$$\mathbf{j} = \gamma_n \gamma_{n-1} \cdots \gamma_1$$

is a reduced expression of w_0 .

For the exceptional algebras, we take \mathbf{j} as below:

$$\begin{array}{ll} G_2 & s_2 s_1 s_2 s_1 s_2 s_1, \\ F_4 & s_4 s_3 s_2 s_3 s_4 s_1 s_2 s_3 s_2 s_1 s_4 s_3 s_2 s_3 s_4 s_1 s_2 s_3 s_2 s_1 s_2 s_3 s_2 s_3, \end{array}$$

$$\begin{aligned}
E_6 & s_1 s_3 s_4 s_2 s_5 s_4 s_3 s_1 s_6 s_5 s_4 s_2 s_3 s_4 s_5 s_6 u_1, \\
E_7 & s_7 s_6 s_5 s_4 s_2 s_3 s_4 s_5 s_6 s_7 s_1 s_3 s_4 s_5 \\
& \times s_2 s_4 s_3 s_1 s_6 s_5 s_4 s_2 s_3 s_4 s_5 s_6 s_7 u_2, \\
E_8 & s_8 s_7 s_6 s_5 s_4 s_2 s_3 s_4 s_5 s_6 s_7 s_8 s_1 s_3 s_4 s_2 s_5 s_4 s_3 s_1 s_6 s_5 s_7 s_6 s_4 s_3 s_2 s_5 s_4 s_5 s_3 s_2 s_4 \\
& \times s_6 s_5 s_7 s_6 s_1 s_3 s_4 s_2 s_5 s_4 s_3 s_1 s_8 s_7 s_6 s_5 s_4 s_2 s_3 s_4 s_5 s_6 s_7 s_8 u_3,
\end{aligned}$$

where u_1 (resp. u_2, u_3) is the reduced expression of the longest element of D_5 (resp. E_6, E_7) obtained by dropping the node 6 (resp. 7,8) which has been chosen previously.

If \mathfrak{g} is of type A_n, D_n or E_6 , let τ be the non-trivial diagram automorphism of order 2 and let \mathbf{i} be the reduced expression for w_0 obtained by applying τ to \mathbf{j} ; for \mathfrak{g} of other types, we take $\mathbf{j} = \mathbf{i}$. In what follows we assume that the roots $\beta_i, i = 1, \dots, r$ are defined with respect to the reduced expression \mathbf{i} .

Let q be an indeterminate, let $\mathbf{Q}(q)$ be the field of rational functions in q with rational coefficients, and let $\mathbf{Z}[q, q^{-1}]$ be the ring of Laurent polynomials with integer coefficients. For $r, m, d \in \mathbf{N}, m \geq r$, define

$$[m]_d = \frac{q^{dm} - q^{-dm}}{q^d - q^{-d}}, \quad [m]_d! = [m]_d [m-1]_d \dots [2]_d [1]_d, \quad \begin{bmatrix} m \\ r \end{bmatrix}_d = \frac{[m]_d!}{[r]_d! [m-r]_d!}.$$

Then $\begin{bmatrix} m \\ r \end{bmatrix}_d \in \mathbf{Z}[q, q^{-1}]$ for all $m \geq r \geq 0$. Choose $d_i \in \{1, 2, 3\}$ such that $(d_i a_{ij})$ is symmetric and such that $\sum_i d_i$ is minimal.

Proposition 1.1. *There is a Hopf algebra \mathbf{U} over $\mathbf{Q}(q)$ which is generated as an algebra by elements $E_{\alpha_i}, F_{\alpha_i}, K_i^{\pm 1}$ ($i \in I$), with the following defining relations:*

$$\begin{aligned}
K_i K_i^{-1} &= K_i^{-1} K_i = 1, \quad K_i K_j = K_j K_i, \\
K_i E_{\alpha_j} K_i^{-1} &= q^{d_i a_{ij}} E_{\alpha_j}, \\
K_i F_{\alpha_j} K_i^{-1} &= q^{-d_i a_{ij}} F_{\alpha_j}, \\
[E_{\alpha_i}, F_{\alpha_j}] &= \delta_{ij} \frac{K_i - K_i^{-1}}{q^{d_i} - q^{-d_i}}, \\
\sum_{r=0}^{1-a_{ij}} (-1)^r \begin{bmatrix} 1-a_{ij} \\ r \end{bmatrix}_{d_i} (E_{\alpha_i})^r E_{\alpha_j} (E_{\alpha_i})^{1-a_{ij}-r} &= 0 \quad \text{if } i \neq j, \\
\sum_{r=0}^{1-a_{ij}} (-1)^r \begin{bmatrix} 1-a_{ij} \\ r \end{bmatrix}_{d_i} (F_{\alpha_i})^r F_{\alpha_j} (F_{\alpha_i})^{1-a_{ij}-r} &= 0 \quad \text{if } i \neq j.
\end{aligned}$$

The comultiplication of \mathbf{U} is given on generators by

$$\Delta(E_{\alpha_i}) = E_{\alpha_i} \otimes 1 + K_i \otimes E_{\alpha_i}, \quad \Delta(F_{\alpha_i}) = F_{\alpha_i} \otimes K_i^{-1} + 1 \otimes F_{\alpha_i}, \quad \Delta(K_i) = K_i \otimes K_i,$$

for $i \in I$. \square

Let \mathbf{U}^+ be the $\mathbf{Q}(q)$ -subalgebra of \mathbf{U} generated by the E_{α_i} for $i \in I$.

Definition 1.1. An element $x \in \mathbf{U}^+$ is called bar-invariant if it is fixed by the \mathbf{Q} -algebra homomorphism $\bar{\cdot} : \mathbf{U}^+ \rightarrow \mathbf{U}^+$ defined by extending:

$$\overline{E_{\alpha_i}} = E_{\alpha_i}, \quad \overline{q} = q^{-1}.$$

For $\nu \in \mathbf{N}[I]$, let \mathbf{U}_ν^+ be the subspace of \mathbf{U}^+ spanned by the monomials $E_{\alpha_{s_1}} E_{\alpha_{s_2}} \cdots E_{\alpha_{s_t}}$ such that for any $i \in I$, the number of occurrences of i in the sequence s_1, \dots, s_t is equal to ν_i . An element $x \in \mathbf{U}^+$ is said to have homogeneity ν if $x \in \mathbf{U}_\nu^+$ and we denote its homogeneity by $|x|$.

It is convenient to use the following notation:

$$E_{\alpha_i}^{(r)} = \frac{E_{\alpha_i}^r}{[r]_{d_i}!}.$$

The elements $F_{\alpha_i}^{(r)}$ are defined similarly.

Set $\mathcal{A} = Z[q, q^{-1}]$ and let ${}_{\mathcal{A}}\mathbf{U}^+$ be the \mathcal{A} -subalgebra of \mathbf{U}^+ generated by $E_{\alpha_i}^{(r)}$, $i \in I$, $r \geq 0$.

For $i \in I$, let T_i ($i \in I$) be the $\mathbf{Q}(q)$ -algebra automorphisms of \mathbf{U} defined as follows (see [L2]):

$$\begin{aligned} T_i(E_{\alpha_i}^{(m)}) &= (-1)^m q^{-m(m-1)} K_i^{-m} F_{\alpha_i}^{(m)}, \\ T_i(F_{\alpha_i}^{(m)}) &= (-1)^m q^{m(m-1)} E_{\alpha_i}^{(m)} K_i^m, \\ T_i(E_{\alpha_j}^{(m)}) &= \sum_{r=0}^{-ma_{ij}} (-1)^r q^{-r} E_{\alpha_i}^{(r)} E_{\alpha_j}^{(m)} E_{\alpha_i}^{(-ma_{ij}-r)} \quad \text{if } i \neq j, \\ T_i(F_{\alpha_j}^{(m)}) &= \sum_{r=0}^{-ma_{ij}} (-1)^r q^r F_{\alpha_i}^{(-ma_{ij}-r)} F_{\alpha_j}^{(m)} F_{\alpha_i}^{(r)} \quad \text{if } i \neq j. \end{aligned}$$

Recall that $\mathbf{i} = s_{i_1} s_{i_2} \cdots s_{i_r}$. For $j = 1, 2, \dots, r$, $m \in \mathbf{Z}$, $m \geq 0$, define a set of root vectors by

$$E_{\beta_j}^{(m)} = T_{i_1} T_{i_2} \cdots T_{i_{j-1}} (E_{\alpha_{i_j}}^{(m)}).$$

For $\mathbf{c} = (c_1, \dots, c_r) \in \mathbf{N}^r$, we set $E_{\mathbf{c}} = E_{\beta_1}^{(c_1)} E_{\beta_2}^{(c_2)} \cdots E_{\beta_r}^{(c_r)}$. The elements $E_{\mathbf{c}}$ are clearly homogenous and $|E_{\mathbf{c}}| = \sum_{j=1}^r c_j \beta_j$. The following result is proved in [L3, Corollary 40.2.2].

Proposition 1.2. *The set $\{E_{\mathbf{c}} : \mathbf{c} \in \mathbf{N}^r\}$ is a $\mathbf{Q}(q)$ -basis of \mathbf{U}^+ and an \mathcal{A} -basis of ${}_{\mathcal{A}}\mathbf{U}^+$. \square*

2. MONOMIAL BASIS

In this section we construct a monomial basis of \mathbf{U}^+ , i.e, a basis consisting of products of $E_{\alpha_i}^{(s)}$, $i \in I$, $s \geq 0$. We also give a simple proof for the existence and uniqueness of the canonical basis of \mathbf{U}^+ . Recall that we have fixed a reduced expression $\mathbf{j} = s_{j_1} s_{j_2} \cdots s_{j_r}$ of w_0 . For $\mathbf{c} = (c_1, \dots, c_r) \in \mathbf{N}^r$ we set

$$M_{\mathbf{c}} = E_{j_1}^{(c_1)} E_{j_2}^{(c_2)} \cdots E_{j_r}^{(c_r)}.$$

Let $>$ be the lexicographic ordering on \mathbf{N}^r such that

$$(1, 0, \dots, 0) > (0, 1, 0, \dots, 0) > \cdots > (0, 0, \dots, 1).$$

Theorem 1. (i) For any $\mathbf{c} \in \mathbf{N}^r$ there exists $f(\mathbf{c}) \in \mathbf{N}^r$ such that

$$M_{f(\mathbf{c})} = E_{\mathbf{c}} + \sum_{\substack{\mathbf{d} \in \mathbf{N}^r \\ \mathbf{d} > \mathbf{c}}} \xi_{\mathbf{d}} E_{\mathbf{d}},$$

where $\xi_{\mathbf{d}} \in \mathcal{A}$.

(ii) The set $\{M_{f(\mathbf{c})} : \mathbf{c} \in \mathbf{N}^r\}$ is an \mathcal{A} -basis of ${}_{\mathcal{A}}\mathbf{U}^+$.

We postpone the proof of the theorem and deduce first the result on canonical bases. Let \mathcal{L} be the $\mathbf{Z}[q^{-1}]$ -lattice of \mathbf{U}^+ spanned by the set $\{E_{\mathbf{c}} : \mathbf{c} \in \mathbf{N}^r\}$.

Theorem 2 (L3,K2). For each $\mathbf{c} \in \mathbf{N}^r$ there exist a unique $b_{\mathbf{c}}$ in the lattice \mathcal{L} such that $b_{\mathbf{c}}$ is bar-invariant and

$$b_{\mathbf{c}} = E_{\mathbf{c}} + \sum_{\substack{\mathbf{d} \in \mathbf{N}^r \\ \mathbf{d} > \mathbf{c}}} \zeta_{\mathbf{d}} E_{\mathbf{d}},$$

where $\zeta_{\mathbf{d}} \in q^{-1}\mathbf{Z}[q^{-1}]$. The set

$$\mathbf{B} = \{b_{\mathbf{c}} : \mathbf{c} \in \mathbf{N}^r\}$$

is an \mathcal{A} -basis of ${}_{\mathcal{A}}\mathbf{U}^+$ and is called the canonical basis or the global crystal basis of \mathbf{U}^+ .

Proof. For each $\mathbf{c} \in \mathbf{N}^r$, observe that the set

$$\mathbf{S}_{\mathbf{c}} = \{\mathbf{d} \in \mathbf{N}^r : \mathbf{d} \geq \mathbf{c}, |E_{\mathbf{c}}| = |E_{\mathbf{d}}|\}$$

is finite and not empty. Let $\mathbf{c} = \mathbf{c}_0 < \mathbf{c}_1 < \cdots < \mathbf{c}_m$ be the elements of $\mathbf{S}_{\mathbf{c}}$. If $\mathbf{c}_0 = \mathbf{c}_m$ then by Theorem 1

$$E_{\mathbf{c}} = M_{f(\mathbf{c})}$$

and there is nothing to prove. Otherwise, by Theorem 1 we can write

$$M_{f(\mathbf{c})} = E_{\mathbf{c}} + \sum_{1 \leq k \leq m} \xi_k E_{\mathbf{c}_k},$$

for some $\xi_k \in \mathbf{Z}[q, q^{-1}]$. Let ξ'_1 be the unique bar-invariant element of \mathcal{A} such that

$$\xi_1 - \xi'_1 = \eta_1 \in q^{-1}\mathbf{Z}[q^{-1}].$$

Applying Theorem 1, we get

$$M_{f(\mathbf{c})} - \xi'_1 M_{f(\mathbf{c}_1)} = E_{\mathbf{c}} + \eta_1 E_{\mathbf{c}_1} + \sum_{2 \leq k \leq m} \xi_{k,2} E_{\mathbf{c}_k},$$

where $\xi_{k,2} \in \mathbf{Z}[q, q^{-1}]$ for all $2 \leq k \leq m$. Next, let ξ'_2 be the unique bar-invariant element of $\mathbf{Z}[q, q^{-1}]$ such that

$$\xi_{2,2} - \xi'_2 = \eta_2 \in q^{-1}\mathbf{Z}[q^{-1}].$$

As before, we can find elements $\xi_{k,3} \in \mathbf{Z}[q, q^{-1}]$ for $3 \leq k \leq m$ such that,

$$M_{f(\mathbf{c})} - \xi'_1 M_{f(\mathbf{c}_1)} - \xi'_2 M_{f(\mathbf{c}_2)} = E_{\mathbf{c}} + \eta_1 E_{\mathbf{c}_1} + \eta_2 E_{\mathbf{c}_2} + \sum_{3 \leq k \leq m} \xi_{k,3} E_{\mathbf{c}_k}.$$

Repeating this process we find finally that

$$M_{f(\mathbf{c})} - \sum_{1 \leq k \leq m} \xi'_k M_{f(\mathbf{c}_k)} = E_{\mathbf{c}} + \sum_{1 \leq k \leq m} \eta_k E_{\mathbf{c}_k},$$

with $\eta_k \in q^{-1}\mathbf{Z}[q^{-1}]$. Since the left-hand side in the previous equation is obviously bar-invariant, the result follows by taking

$$b_{\mathbf{c}} = M_{f(\mathbf{c})} - \sum_{1 \leq k \leq m} \xi'_k M_{f(\mathbf{c}_k)}.$$

□

We now turn to the proof of Theorem 1. We assume (i) and prove (ii). Suppose that

$$\sum_{\mathbf{c}} \xi_{\mathbf{c}} M_{f(\mathbf{c})} = 0,$$

for some $\xi_{\mathbf{c}} \in \mathcal{A}$. Choose, if possible, $\mathbf{c}_0 \in \mathbf{N}^r$ minimal such that $\xi_{\mathbf{c}_0} \neq 0$. Using Theorem 1(i) we get that

$$\xi_{\mathbf{c}_0} E_{\mathbf{c}_0} + \sum_{\mathbf{c} > \mathbf{c}_0} \xi_{\mathbf{c}} E_{\mathbf{c}} = 0.$$

But this contradicts Proposition 1.2 and hence $\xi_{\mathbf{c}_0} = 0$ proving that the elements $M_{f(\mathbf{c})} \in {}_{\mathcal{A}}\mathbf{U}^+$ are linearly independent.

For each $\eta \in \mathbf{N}[I]$, set

$${}_{\mathcal{A}}\mathbf{U}_{\eta}^+ = {}_{\mathcal{A}}\mathbf{U}^+ \cap \mathbf{U}_{\eta}^+.$$

Let

$$S_{\eta} = \{\mathbf{c} \in \mathbf{N}^r : E_{\mathbf{c}} \in {}_{\mathcal{A}}\mathbf{U}_{\eta}^+\},$$

and let $\mathbf{c}_0 < \mathbf{c}_1 < \dots < \mathbf{c}_m$ be the elements of S_{η} . From Theorem 1(i) it is clear that

$$M_{f(\mathbf{c}_m)} = E_{\mathbf{c}_m}.$$

An obvious downward induction on $|S_{\eta}|$ proves that $E_{\mathbf{c}_k}$ is in the span of $M_{f(\mathbf{d})}$ for $\mathbf{d} \in \mathbf{N}^r$ and the result follows by Proposition 1.2.

The rest of the paper is devoted to proving Theorem 1(i). We need the following result proved in [L1, Section 6]. Let $\alpha \in R^+$ be such that

$$\alpha - \alpha_n = \sum r_i \alpha_i, \quad r_i \in \mathbf{Z}, \quad r_i \geq 0.$$

Fix $k < n$ and let

$$R_{\alpha,k}^+ = \{\beta \in R^+ : \beta = r\alpha + s\alpha_k, r, s \in \mathbf{Z}\}.$$

Lemma 2.1. *Let $a, b \in \mathbf{N}$.*

(i) *If $R_{\alpha,k}^+ = \{\alpha, \alpha_k\}$, then*

$$E_{\alpha_k}^{(a)} E_{\alpha}^{(b)} = E_{\alpha}^{(b)} E_{\alpha_k}^{(a)}.$$

(ii) *If $R_{\alpha,k}^+ = \{\alpha, \alpha + \alpha_k, \alpha_k\}$, then $d_{\alpha} = d_{\alpha_k} = d$ and*

$$\begin{aligned} E_{\alpha+\alpha_k}^{(a)} E_{\alpha}^{(b)} &= q^{dab} E_{\alpha}^{(b)} E_{\alpha+\alpha_k}^{(a)}, \\ E_{\alpha_k}^{(a)} E_{\alpha+\alpha_k}^{(b)} &= q^{dab} E_{\alpha+\alpha_k}^{(b)} E_{\alpha_k}^{(a)}, \\ E_{\alpha_k}^{(a)} E_{\alpha}^{(b)} &= \sum_{r \in \mathbf{N}} q^{-d(a-r)(b-r)} E_{\alpha}^{(b-r)} E_{\alpha+\alpha_k}^{(r)} E_{\alpha_k}^{(a-r)}. \end{aligned}$$

(iii) *If $R_{\alpha,k}^+ = \{\alpha, \alpha + \alpha_k, \alpha + 2\alpha_k, \alpha_k\}$, then $d_{\alpha} = 2$, $d_{\alpha_k} = 1$ and*

$$\begin{aligned} E_{\alpha+\alpha_k}^{(a)} E_{\alpha}^{(b)} &= q^{2ab} E_{\alpha}^{(b)} E_{\alpha+\alpha_k}^{(a)}, \\ E_{\alpha+2\alpha_k}^{(a)} E_{\alpha+\alpha_k}^{(b)} &= q^{2ab} E_{\alpha+\alpha_k}^{(b)} E_{\alpha+2\alpha_k}^{(a)}, \\ E_{\alpha_k}^{(a)} E_{\alpha+2\alpha_k}^{(b)} &= q^{2ab} E_{\alpha+2\alpha_k}^{(b)} E_{\alpha_k}^{(a)}, \end{aligned}$$

$$\begin{aligned}
E_{\alpha+2\alpha_k}^{(a)} E_{\alpha}^{(b)} &= \sum_{r \in \mathbf{N}} q^{2r(b-r)+2r(a-r)} \prod_{h=1}^r (q^{4h-2} - 1) E_{\alpha}^{(b-r)} E_{\alpha+\alpha_k}^{(2r)} E_{\alpha+2\alpha_k}^{(a-r)}, \\
E_{\alpha_k}^{(a)} E_{\alpha+\alpha_k}^{(b)} &= \sum_{r \in \mathbf{N}} q^{r(b-r)+r(a-r)-r} \prod_{h=1}^r (q^{2h} + 1) E_{\alpha+\alpha_k}^{(b-r)} E_{\alpha+2\alpha_k}^{(r)} E_{\alpha_k}^{(a-r)}, \\
E_{\alpha_k}^{(a)} E_{\alpha}^{(b)} &= \sum_{r,t \in \mathbf{N}} q^{-2(b-r-t)(a-r-t)-(a-r-2t)r} E_{\alpha}^{(b-r-t)} E_{\alpha+\alpha_k}^{(r)} E_{\alpha+2\alpha_k}^{(t)} E_{\alpha_k}^{(a-r-2t)}.
\end{aligned}$$

□

The proof of Theorem 1(i) proceeds by induction on the rank of \mathfrak{g} . For rank one there is nothing to prove, and the rank two case is contained in the next Lemma.

Lemma 2.2. (i) *If $\mathfrak{g} = A_2$, and $\mathbf{c} = (c_1, c_2, c_3)$, then*

$$M_{f(\mathbf{c})} = E_1^{(c_2)} E_2^{(c_1+c_2)} E_1^{(c_3)}.$$

(ii) *If $\mathfrak{g} = B_2$, and $\mathbf{c} = (c_1, c_2, c_3, c_4)$, then,*

$$M_{f(\mathbf{c})} = E_2^{(c_1)} E_1^{(c_2+2c_3)} E_2^{(c_2+c_3)} E_1^{(c_4)}.$$

(iii) *If \mathfrak{g} is of type G_2 , and $\mathbf{c} = (c_1, c_2, \dots, c_6)$, then*

$$M_{f(\mathbf{c})} = E_2^{(c_1)} E_1^{(c_2+3c_3)} E_2^{(c_2+2c_3)} E_1^{(2c_4+3c_5)} E_2^{(c_4+c_5)} E_1^{(c_6)}.$$

Proof. If \mathfrak{g} is of type A_2 or B_2 , the result follows by using the previous lemma. For G_2 , see [X1]. □

Turning to the general case, let $l = l_n$ be the number of positive roots β such that $\beta - \alpha_n = \sum_{i=1}^n n_i \alpha_i$, where $n_i \geq 0$ for all $i = 1, \dots, n$. For any $\mathbf{c} \in \mathbf{N}^r$, we write

$$\mathbf{c} = \mathbf{c}' + \mathbf{c}'',$$

where $c'_k = 0$ if $k > l$ and $c''_k = 0$ if $k \leq l$.

Lemma 2.3. *Let $\mathbf{c} \in \mathbf{N}^r$ be such that $\mathbf{c}'' = 0$. Then there exists $f(\mathbf{c})$ such that*

$$M_{f(\mathbf{c})} = E_{\mathbf{c}} + \sum_{\substack{\mathbf{d} \in \mathbf{N}^r \\ \mathbf{d}' > \mathbf{c}'}} \xi_{\mathbf{d}} E_{\mathbf{d}} \quad \xi_{\mathbf{d}} \in \mathbf{Z}[q, q^{-1}].$$

Assuming the lemma we complete the proof as follows.

The element $E_{\mathbf{c}'}$ can be regarded as an element of $\mathbf{U}_q(\mathfrak{g}_{n-1})$ where \mathfrak{g}_{n-1} is the simple Lie algebra associated to the $(n-1) \times (n-1)$ Cartan matrix obtained by dropping the n^{th} row and the n^{th} column. Hence by induction, we have a monomial $M_{f(\mathbf{c}'')}$ such that

$$M_{f(\mathbf{c}'')} = E_{\mathbf{c}''} + \sum_{\mathbf{d} > \mathbf{c}''} \eta_{\mathbf{d}} E_{\mathbf{d}},$$

where $\eta_{\mathbf{d}} \in \mathcal{A}$ and $d_k = 0$ if $k \leq l$. Let $M_{f(\mathbf{c}'')}$ be the monomial defined in the preceding lemma and set

$$M_{f(\mathbf{c})} = M_{f(\mathbf{c}')} M_{f(\mathbf{c}'')}.$$

We get

$$\begin{aligned}
M_{f(\mathbf{c})} &= M_{f(\mathbf{c}')}M_{f(\mathbf{c}'')} \\
&= E_{\mathbf{c}'}E_{\mathbf{c}''} + E_{\mathbf{c}'} \sum_{\mathbf{d} > \mathbf{c}''} \eta_{\mathbf{d}} E_{\mathbf{d}} + \sum_{\substack{\mathbf{d} \in \mathbf{N}^r \\ \mathbf{d}' > \mathbf{c}'}} \xi_{\mathbf{d}} E_{\mathbf{d}} E_{\mathbf{c}''} + \sum_{\substack{\mathbf{d} \in \mathbf{N}^r \\ \mathbf{d}' > \mathbf{c}'}} \xi_{\mathbf{d}} E_{\mathbf{d}} \sum_{\mathbf{d}_1 > \mathbf{c}''} \eta_{\mathbf{d}_1} E_{\mathbf{d}_1} \\
&= E_{\mathbf{c}} + \sum_{\substack{\mathbf{d} \in \mathbf{N}^r \\ \mathbf{d} > \mathbf{c}}} \zeta_{\mathbf{d}} E_{\mathbf{d}}, \quad \zeta_{\mathbf{d}} \in q^{-1} \mathbf{Z}[q^{-1}]. \quad \square
\end{aligned}$$

It remains to prove Lemma 2.3. Note that if $\mathbf{c}'' = 0$, it is enough to prove that $\mathbf{d} > \mathbf{c}$ since this together with the fact that $|E_{\mathbf{d}}| = |E_{\mathbf{c}}|$ implies that $\mathbf{d}' > \mathbf{c}'$. Let j_1, j_2, \dots, j_l be the first l -indices of the reduced expression \mathbf{j} of w_0 . Then using Lemma 2.1 repeatedly, it is not hard to see that,

$$M_{f(\mathbf{c})} = E_{j_1}^{(k_1)} E_{j_2}^{(k_2)} \dots E_{j_l}^{(k_l)},$$

where $\mathbf{k} = (k_1, k_2, \dots, k_l)$ is related to \mathbf{c} as follows:

$$k_i = c_n + c_{n-1} + \dots + c_{n-i+1}$$

if \mathfrak{g} is of type A_n ;

$$k_i = \begin{cases} c_i + c_{2n-i+1} + c_{2n-i+2} + \dots + c_{2n-1} & \text{if } 2 \leq i \leq n-1, \\ c_n + 2c_{n+1} + \dots + 2c_{2n-1} & \text{if } i = n, \\ c_{n-j+1} + \dots + c_{2n-1} & \text{if } 1 \leq i-n = j \leq n-1, \end{cases}$$

if \mathfrak{g} is of type B_n ;

$$k_i = \begin{cases} c_i + c_{2n-i+1} + c_{2n-i+2} + \dots + c_{2n-1} & \text{if } 2 \leq i \leq n-1, \\ 2c_n + c_{n+1} + \dots + c_{2n-1} & \text{if } i = n+1, \\ c_{n-j+1} + \dots + c_{n-1} + 2c_n + c_{n+1} + \dots + c_{2n-1} & \text{if } 2 \leq i-n = j \leq n-1, \end{cases}$$

if \mathfrak{g} is of type C_n ;

$$k_i = \begin{cases} c_1 & \text{if } i = 1, \\ c_i + c_{2n-i+1} + c_{2n-i+2} + \dots + c_{2n-2} & \text{if } 2 \leq i \leq n, \\ c_{n-j} + \dots + c_{2n-2} & \text{if } 1 \leq i-n = j \leq n-2. \end{cases}$$

if \mathfrak{g} is of type D_n .

The formulae for the exceptional algebras are naturally much more complicated to write down. We give as an example the case of E_6 , and omit the other cases. If \mathfrak{g} is of type E_6 , then

$$k_1 = c_9, \quad k_2 = c_5 + c_9, \quad k_3 = c_3 + c_5 + c_9,$$

$$k_4 = c_4 + c_6 + c_8 + c_{10} + c_{16}, \quad k_5 = c_2 + c_3 + c_5 + c_9, \quad k_6 = c_4 + c_7 + c_{11}$$

$$k_7 = c_6 + c_7 + c_{12} + c_{14} + c_{15} + c_{16}, \quad k_8 = c_{10} + c_{11} + \dots + c_{15} + c_{16},$$

$$k_9 = c_1 + c_2 + c_3 + c_5 + c_9, \quad k_{10} = c_4 + c_8 + c_{13} + c_{14} + c_{15} + c_{16},$$

$$k_{11} = c_6 + c_8 + c_{12} + c_{13} + c_{14} + c_{15} + c_{16},$$

$$k_{12} = c_7 + c_8 + c_{11} + c_{12} + c_{13} + c_{14} + c_{15} + c_{16},$$

$$k_{13} = c_8 + c_{10} + c_{11} + c_{12} + c_{13} + c_{14} + c_{15} + c_{16},$$

$$k_{14} = c_7 + c_8 + c_{10} + c_{11} + c_{12} + c_{13} + c_{14} + c_{15} + c_{16},$$

$$k_{15} = c_6 + c_7 + c_8 + c_{10} + c_{11} + c_{12} + c_{13} + c_{14} + c_{15} + c_{16},$$

$$k_{16} = c_4 + c_7 + c_8 + c_{10} + c_{11} + c_{12} + c_{13} + c_{14} + c_{15} + c_{16}.$$

As two examples we write down the details of proof of Lemma 2.3 for type A_n, B_n . We shall write $x \equiv y \pmod{(> \mathbf{c})}$ if $x - y$ is a $\mathbf{Z}[q, q^{-1}]$ -linear combination of $E_{\mathbf{d}}$ ($\mathbf{d} > \mathbf{c}$).

For type A_n we denote by $E_{i,j}$ ($i > j$) the root vector corresponding to $\alpha_i + \alpha_{i-1} + \cdots + \alpha_j$. We have

$$\begin{aligned}
M_{f(\mathbf{c})} &= E_1^{(c_n)} E_2^{(c_n+c_{n-1})} \cdots E_n^{(c_n+\cdots+c_1)} \\
&= E_1^{(c_n)} E_2^{(c_n+c_{n-1})} \cdots E_{n-2}^{(c_n+\cdots+c_3)} \sum_{\substack{r_n, s_n, t_n \in \mathbf{N} \\ r_n + s_n = c_n + \cdots + c_1 \\ s_n + t_n = c_n + \cdots + c_2}} v^{-r_n t_n} E_n^{(r_n)} E_{n,n-1}^{(s_n)} E_{n-1}^{(t_n)} \\
&= \sum_{\substack{r_n, s_n, t_n \in \mathbf{N} \\ r_n + s_n = c_n + \cdots + c_1 \\ s_n + t_n = c_n + \cdots + c_2}} v^{-r_n t_n} E_1^{(c_n)} E_2^{(c_n+c_{n-1})} \cdots E_{n-2}^{(c_n+\cdots+c_3)} E_n^{(r_n)} E_{n,n-1}^{(s_n)} E_{n-1}^{(t_n)} \\
&= \sum_{\substack{r_n, s_n, t_n \in \mathbf{N} \\ r_n + s_n = c_n + \cdots + c_1 \\ s_n + t_n = c_n + \cdots + c_2}} v^{-r_n t_n} E_n^{(r_n)} E_1^{(c_n)} E_2^{(c_n+c_{n-1})} \cdots E_{n-2}^{(c_n+\cdots+c_3)} E_{n,n-1}^{(s_n)} E_{n-1}^{(t_n)} \\
&\equiv E_n^{(c_1)} E_1^{(c_n)} E_2^{(c_n+c_{n-1})} \cdots E_{n-2}^{(c_n+\cdots+c_3)} E_{n,n-1}^{(c_2)} \pmod{(> \mathbf{c})} \\
&= E_1^{(c_n)} E_2^{(c_n+c_{n-1})} \cdots E_{n-3}^{(c_n+\cdots+c_4)} \\
&\quad \times \sum_{\substack{r_{n-1}, s_{n-1}, t_{n-1} \in \mathbf{N} \\ r_{n-1} + s_{n-1} = c_n + \cdots + c_2 \\ s_{n-1} + t_{n-1} = c_n + \cdots + c_3}} v^{-r_{n-1} t_{n-1}} E_{n,n-1}^{(r_{n-1})} E_{n,n-2}^{(s_{n-1})} E_{n-2}^{(t_{n-1})} \\
&\equiv E_n^{(c_1)} E_{n,n-1}^{(c_2)} E_1^{(c_n)} E_2^{(c_n+c_{n-1})} \cdots E_{n-3}^{(c_n+\cdots+c_4)} E_{n,n-2}^{(c_3)} \pmod{(> \mathbf{c})} \\
&\dots \\
&\equiv E_n^{(c_1)} E_{n,n-1}^{(c_2)} \cdots E_{n,3}^{(c_{n-2})} E_1^{(c_n)} E_{n,2}^{(c_n+c_{n-1})} \pmod{(> \mathbf{c})} \\
&= E_n^{(c_1)} E_{n,n-1}^{(c_2)} \cdots E_{n,3}^{(c_{n-2})} \sum_{\substack{r_2, s_2, t_2 \in \mathbf{N} \\ r_2 + s_2 = c_n + c_{n-1} \\ s_2 + t_2 = c_n}} v^{-r_2 t_2} E_{n,2}^{(r_2)} E_{n,1}^{(s_2)} E_1^{(t_2)} \\
&\equiv E_n^{(c_1)} E_{n,n-1}^{(c_2)} \cdots E_{n,3}^{(c_{n-2})} E_{n,2}^{(c_{n-1})} E_{n,1}^{(c_n)} \pmod{(> \mathbf{c})}.
\end{aligned}$$

This proves Lemma 2.3 for type A_n .

For type B_n , we shall write $E_{i,j}$ ($i \geq j$) for the root vector corresponding to $\alpha_i + \alpha_{i-1} + \cdots + \alpha_j$ and $E'_{n,j}$ ($n > j$) for the root vector corresponding to $\alpha_n + \alpha_{n-1} + \cdots + \alpha_{j+1} + 2\alpha_j + \cdots + 2\alpha_1$. Note that $\alpha_2, \dots, \alpha_n$ generate a root system of type A_{n-1} .

$$\begin{aligned}
M_{f(\mathbf{c})} &= E_n^{(k_1)} E_{n-1}^{(k_2)} \cdots E_1^{(k_n)} \cdots E_n^{(k_{2n-1})} \\
&\equiv E_n^{(k_1)} E_{n-1}^{(k_2)} \cdots E_1^{(k_n)} E_2^{(k_{n+1})} \cdots E_{n-2}^{(k_{2n-3})} E_n^{(c_2)} E_{n,n-1}^{(k_{2n-2})} \pmod{(> \mathbf{c})} \\
&\equiv E_n^{(k_1)} E_{n-1}^{(k_2)} E_n^{(c_2)} E_{n-2}^{(k_3)} \cdots E_1^{(k_n)} E_2^{(k_{n+1})} \cdots E_{n-2}^{(k_{2n-3})} E_{n,n-1}^{(k_{2n-2})} \pmod{(> \mathbf{c})} \\
&\equiv E_n^{(c_1)} E_{n,n-1}^{(c_2)} E_{n-1}^{(c_{2n-1})} E_{n-2}^{(k_3)} \cdots E_1^{(k_n)} E_2^{(k_{n+1})} \cdots E_{n-2}^{(k_{2n-3})} \\
&\quad \times E_{n,n-1}^{(k_{2n-2})} \pmod{(> \mathbf{c})} \\
&\equiv E_n^{(c_1)} E_{n,n-1}^{(c_2)} E_{n-1}^{(c_{2n-1})} E_{n-2}^{(k_3)} \cdots E_1^{(k_n)} E_2^{(k_{n+1})} \cdots E_{n-3}^{(k_{2n-4})} \\
&\quad \times E_{n,n-1}^{(c_3)} E_{n,n-2}^{(k_{2n-3})} \pmod{(> \mathbf{c})} \\
&\equiv E_n^{(c_1)} E_{n,n-1}^{(c_2)} E_{n,n-2}^{(c_3)} E_{n-1}^{(c_{2n-1})} E_{n-2}^{(c_{2n-2}+c_{2n-1})} E_{n-3}^{(k_4)} \cdots E_1^{(k_n)} \\
&\quad \times E_2^{(k_{n+1})} \cdots E_{n-3}^{(k_{2n-4})} E_{n,n-2}^{(k_{2n-3})} \pmod{(> \mathbf{c})} \\
&\dots \\
&\equiv E_n^{(c_1)} E_{n,n-1}^{(c_2)} \cdots E_{n,2}^{(c_{n-1})} E_{n-1}^{(c_{2n-1})} E_{n-2}^{(c_{2n-2}+c_{2n-1})} \cdots E_2^{(c_{n+2}+\cdots+c_{2n-1})} E_1^{(k_n)} \\
&\quad \times E_{n,2}^{(c_n+\cdots+c_{2n-1})} \pmod{(> \mathbf{c})} \\
&\equiv E_n^{(c_1)} E_{n,n-1}^{(c_2)} \cdots E_{n,2}^{(c_{n-1})} E_{n-1}^{(c_{2n-1})} E_{n-2}^{(c_{2n-2}+c_{2n-1})} \cdots E_2^{(c_{n+2}+\cdots+c_{2n-1})} E_{n,1}^{(c_n)} \\
&\quad \times E'_{n,1}^{(c_{n+1}+\cdots+c_{2n-1})} \pmod{(> \mathbf{c})} \\
&\equiv E_n^{(c_1)} E_{n,n-1}^{(c_2)} \cdots E_{n,2}^{(c_{n-1})} E_{n,1}^{(c_n)} E_{n-1}^{(c_{2n-1})} E_{n-2}^{(c_{2n-2}+c_{2n-1})} \cdots E_2^{(c_{n+2}+\cdots+c_{2n-1})} \\
&\quad \times E'_{n,1}^{(c_{n+1}+\cdots+c_{2n-1})} \pmod{(> \mathbf{c})} \\
&\equiv E_n^{(c_1)} E_{n,n-1}^{(c_2)} \cdots E_{n,2}^{(c_{n-1})} E_{n,1}^{(c_n)} E_{n-1}^{(c_{2n-1})} E_{n-2}^{(c_{2n-2}+c_{2n-1})} \cdots E_3^{(c_{n+3}+\cdots+c_{2n-1})} \\
&\quad \times E'_{n,1}^{(c_{n+1})} E'_{n,2}^{(c_{n+2}+\cdots+c_{2n-1})} \pmod{(> \mathbf{c})} \\
&\equiv E_n^{(c_1)} E_{n,n-1}^{(c_2)} \cdots E_{n,2}^{(c_{n-1})} E_{n,1}^{(c_n)} E'_{n,1}^{(c_{n+1})} E_{n-1}^{(c_{2n-1})} E_{n-2}^{(c_{2n-2}+c_{2n-1})} \cdots \\
&\quad \times E_3^{(c_{n+3}+\cdots+c_{2n-1})} E'_{n,2}^{(c_{n+2}+\cdots+c_{2n-1})} \pmod{(> \mathbf{c})} \\
&\dots \\
&\equiv E_n^{(c_1)} E_{n,n-1}^{(c_2)} \cdots E_{n,2}^{(c_{n-1})} E_{n,1}^{(c_n)} E'_{n,1}^{(c_{n+1})} \cdots E'_{n,n-1}^{(c_{2n-1})} \pmod{(> \mathbf{c})}.
\end{aligned}$$

For other types the proof is completely similar.

Finally we conjecture for each reduced expression of w_0 there is a corresponding monomial basis of \mathbf{U}^+ .

Acknowledgement: N.X. was supported by the National Natural Science Foundation of China. N.X. also would like to thank Newton Institute at Cambridge University for financial support and hospitality during his visit to the institute.

REFERENCES

- [BCP] Beck, J., Chari, V., Pressley, A.N., *An algebraic characterization of the affine canonical basis*, to appear, Duke. Math. J.
- [GL] Grojnowski, I., Lusztig, G., *A comparison of bases of quantized enveloping algebras*, Linear algebraic groups and their representations, Contemp. Math. **153** (1992), 11-19.
- [K1] Kashiwara, M., *Crystallizing the q -analogue of universal enveloping algebras*, Comm. Math. Phys. **133** (1990), 249–260.
- [K2] Kashiwara, M., *On crystal bases of the q -analogue of universal enveloping algebras*, Duke Math. J. **63** (1991), 465-516
- [L1] Lusztig, G., *Quantum groups at roots of 1*, Geom. Ded. **35** (1990), 89-114.
- [L2] Lusztig, G., *Canonical bases arising from quantized enveloping algebras*, JAMS **3** (1990), 447-498.
- [L3] Lusztig, G., *Introduction to quantum groups*, Progress in Mathematics 110, Birkhäuser, Boston · Basel · Berlin, 1993.
- [X1] Xi, N., *Bases of quantized enveloping algebras*, preprint, 1995.
- [X2] Xi, N., *A commutation formula for root vectors in quantized enveloping algebras*, Pacific J. Math., to appear.

V.C. AND N.X. DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, RIVERSIDE, CA 92521, USA

N.X. INSTITUTE OF MATHEMATICS, CHINESE ACADEMY OF SCIENCES, BEIJING 100080, CHINA
E-mail address: chari@math.ucr.edu, nanhua@math.ucr.edu, nanhua@math08.math.ac.cn