The twisted quantum affine algebra $U_q(A_2^{(2)})$ and correlation functions of the Izergin-Korepin model

Bo-Yu Hou ¹, Wen-Li Yang ^{1, 2} and Yao-Zhong Zhang ²
¹ Institute of Modern Physics, Northwest University, Xian 710069, China
² Department of Mathematics, University of Queensland, Brisbane, Qld 4072, Australia

We derive the exchange relations of the vertex operators of $U_q(A_2^{(2)})$ and show that these vertex operators give the bosonization of the Izergin-Korepin model. We give an integral expression of the correlation functions of the Izergin-Korepin model and derive the difference equations which they satisfy.

I. INTRODUCTION

It is well-known that quantum affine algberas play an essential role in the studies of low-dimensional massive integrable models such as quantum spin chains, since they provide the symmetry algberas of these models. Based on the works of q-vertex operators [1] [2] and level-one highest weight representations of the quantum affine algebra $U_q(\widehat{sl}_2)$, the Kyoto group [3] [4] developed a new method which enabled the group to diagonalize the XXZ spin- $\frac{1}{2}$ chain directly in the thermodynamic limit and moreover compute the correlation functions of the spin operators. This approach was later generalized to higher spin XXZ chains [5–7], vertex models with $U_q(\widehat{sl}_n)$ symmetries [8], the face type statistical mechanics models [9,10], and more recently to integrable models with quantum affine superalgbera symmetry [11].

In this paper, we will extend the above programme further to the Izergin-Korepin 19-vertex model [12]—[16] which has twisted quantum affine algebra $U_q(A_2^{(2)})$ as its non-abelian symmetry. The bosonic realization of $U_q(A_2^{(2)})$ and its level-one vertex operators were constructed recently [17]. In section 3, we use the results of [17] to calculate the exchange relations of the q-vertex operators. We show that these vertex operators satisfy the Faddeev-Zamolodchikov algebra with the R-matrix of $U_q(A_2^{(2)})$ as its constructure constant. A Miki's construction of $U_q(A_2^{(2)})$ is also given. In section 4, generalizing the Kyoto group's work [3] to the case of twisted quantum affine algebra $U_q(A_2^{(2)})$, we give the bosonization of the Izergin-Korepin model . In section 5, we compute the correlation functions of the local operators (such as the spin operator S_z) and give an integral expression of the correlation functions. A set of difference equations satisfied by the correlation functions have also been derived.

II. PRELIMINARIES

In this section, we briefly review the bosonization of the twisted quantum affine algebra $U_q(A_2^{(2)})$ at level one [17].

A. Quantum affine algebra $U_q(A_2^{(2)})$

The symmetric Cartan matrix of the twisted affine Lie algebra $A_2^{(2)}$ is

$$(a_{ij}) = \left(\begin{array}{cc} 8 & -4 \\ -4 & 2 \end{array}\right)$$

where i, j = 0, 1. Quantum affine algebra $U_q(A_2^{(2)})$ is a q-analogue of the universal enveloping algebra of $A_2^{(2)}$ generated by the Chevalley generators $\{e_i, f_i, t_i^{\pm 1}, d | i = 0, 1\}$, where d is the usual derivation operator. The defining relations are [18] [19]

$$\begin{split} t_i t_j &= t_j t_i, \quad t_i d = d t_i, \quad [d, e_i] = \delta_{i,0} e_i, \quad [d, f_i] = -\delta_{i,0} f_i, \\ t_i e_j t_i^{-1} &= (q^{\frac{1}{2}})^{a_{ij}} e_j, \quad t_i f_j t_i^{-1} = (q^{\frac{1}{2}})^{-a_{ij}} f_j, \\ [e_i, f_j] &= \delta_{ij} \frac{t_i - t_i^{-1}}{q_i - q_i^{-1}}, \end{split}$$

$$\sum_{r=0}^{1-a_{ij}} (-1)^r \begin{bmatrix} 1-a_{ij} \\ r \end{bmatrix}_{q_i} (e_i)^r e_j(e_i)^{1-a_{ij}-r} = 0 \quad , \quad \text{if } i \neq j,$$

$$\sum_{r=0}^{1-a_{ij}} (-1)^r \begin{bmatrix} 1-a_{ij} \\ r \end{bmatrix}_{q_i} (f_i)^r f_j(f_i)^{1-a_{ij}-r} = 0 \quad , \quad \text{if } i \neq j,$$

where $q_1 = q^{\frac{1}{2}}$, $q_0 = q^2$, $t_i = q_i^{h_i}$, and

$$[n]_{q_i} = \frac{q_i^n - q_i^{-n}}{q_i - q_i^{-1}} , \quad [n]_{q_i}! = [n]_{q_i}[n-1]_{q_i} \cdots [1]_{q_i} ,$$

$$\begin{bmatrix} n \\ r \end{bmatrix}_{q_i} = \frac{[n]_{q_i}!}{[n-r]_{q_i}![r]_{q_i}!}$$

 $U_q(A_2^{(2)})$ is a quasi-triangular Hopf algebra endowed with Hopf algebra structure:

$$\Delta(t_i) = t_i \otimes t_i, \quad \Delta(e_i) = e_i \otimes 1 + t_i \otimes e_i, \quad \Delta(f_i) = f_i \otimes t_i^{-1} + 1 \otimes f_i, \tag{II.1}$$

$$\epsilon(t_i) = 1, \quad \epsilon(e_i) = \epsilon(f_i) = 0,$$
(II.2)

$$S(e_i) = -t_i^{-1}e_i, \quad S(f_i) = -f_it_i, \quad S(t_i^{\pm 1}) = t_i^{\mp 1}, \quad S(d) = -d.$$
 (II.3)

 $U_q(A_2^{(2)})$ can also be realized by the Drinfeld generators [17] [20] $\{d, X_m^{\pm}, a_n, K^{\pm 1}, \gamma^{\pm 1/2} | m \in \mathbf{Z}, n \in \mathbf{Z}_{\neq 0}\}$. The relations read

$$\gamma$$
 is central, $[K, a_n] = 0$, $[d, K] = 0$, $[d, a_n] = na_n$, (II.4)

$$[a_m, a_n] = \delta_{m+n,0} \frac{\{[4n]_{q_1} - (-1)^n [2n]_{q_1}\} (\gamma^m - \gamma^{-m})}{m(q_1 - q_1^{-1})},$$
(II.5)

$$KX_m^{\pm} = q^{\pm 1}X_m^{\pm}K, \quad [d, X_m^{\pm}] = mX_m^{\pm},$$
 (II.6)

$$[a_m, X_n^{\pm}] = \pm \frac{\{[4m]_{q_1} - (-1)^m [2m]_{q_1}\}}{m} \gamma^{\mp |m|/2} X_{n+m}^{\pm}, \tag{II.7}$$

$$[X_m^+, X_n^-] = \frac{1}{q_1 - q_1^{-1}} (\gamma^{(m-n)/2} \psi_{m+n}^+ - \gamma^{-(m-n)/2} \psi_{m+n}^-), \tag{II.8}$$

$$(z - wq^{\pm 2})(z + wq^{\mp 1})X^{\pm}(z)X^{\pm}(w) = (zq^{\pm 2} - w)(zq^{\mp 1} + w)X^{\pm}(w)X^{\pm}(z) \quad , \tag{II.9}$$

where the corresponding Drinfeld currents $\psi^{\pm}(z)$ and $X^{\pm}(z)$ are defined by

$$\psi^{+}(z) = \sum_{m=0}^{\infty} \psi_{m}^{+} z^{-m} = Kexp\{(q_{1} - q_{1}^{-1}) \sum_{k=1}^{\infty} a_{k} z^{-k}\} ,$$

$$\psi^{-}(z) = \sum_{m=0}^{\infty} \psi_{-m}^{-} z^{m} = K^{-1} exp\{-(q_{1} - q_{1}^{-1}) \sum_{k=1}^{\infty} a_{-k} z^{k}\} ,$$

$$X^{\pm}(z) = \sum_{n \in \mathbb{Z}} X_{m}^{\pm} z^{-m} .$$

The Chevalley generators are related to the Drinfeld generators by the formulae:

$$t_1 = K, \quad e_1 = X_0^+, \quad t_0 = \gamma K^{-2}, \quad f_1 = X_0^-,$$
 (II.10)

$$e_0 = K^{-2}[X_0^-, X_1^-]_q, \quad f_0 = \frac{1}{[4]_{q_1}^2} [X_{-1}^+, X_0^+]_{q^{-1}} K^2$$
 (II.11)

B. Bosonization of $U_q(A_2^{(2)})$ at level one

Let us introduce the bosonic q-oscillators [17] $\{a_n, Q, P | n \in \mathbb{Z} - \{0\}\}$ which satisfy the commutation relations

$$[a_m, a_n] = \delta_{m+n,0} \frac{1}{m} \{ [4m]_{q_1} - (-1)^m [2m]_{q_1} \} [2m]_{q_1},$$

$$[P, a_m] = [Q, a_m] = 0 , [P, Q] = 1,$$

Set $F_1 = \bigoplus_{n \in \mathbb{Z}} C[a_{-1}, a_{-2}, \ldots] e^{\frac{Q}{2} + nQ} |0\rangle$, where the Fock vacuum vector $|0\rangle$ is defined by

$$a_n|0>=0$$
 for $n>0$, $P|0>=0$.

Then, on F_1 the Drinfeld currents of $U_q(A_2^{(2)})$ at level-one are realized by the free boson fields as [17]

$$\begin{split} &\gamma = q \quad , \quad K = q^P \quad , \\ &\psi^+(z) = q^P exp\{(q_1 - q_1^{-1}) \sum_{n=1}^\infty a_n z^{-n}\} \quad , \\ &\psi^-(z) = q^{-P} exp\{-(q_1 - q_1^{-1}) \sum_{n=1}^\infty a_{-n} z^n\} \quad , \\ &X^\pm(z) = exp\{\pm \sum_{n=1}^\infty \frac{a_{-n}}{[2n]_{q_1}} q^{\mp \frac{n}{2}} z^n\} exp\{\mp \sum_{n=1}^\infty \frac{a_n}{[2n]_{q_1}} q^{\mp \frac{n}{2}} z^{-n}\} e^{\pm Q} z^{\pm P + \frac{1}{2}} \quad . \end{split} \tag{II.12}$$

By the relations between Chevalley basis and Drinfeld basis (II.10—II.11), one can check that

$$t_1 e^{\frac{Q}{2}} |0> = q^{\frac{1}{2}} e^{\frac{Q}{2}} |0> = (q^{\frac{1}{2}})^{<\alpha_1, \Lambda_1>} e^{\frac{Q}{2}} |0>,$$

$$t_0 e^{\frac{Q}{2}} |0> = e^{\frac{Q}{2}} |0> = (q^{\frac{1}{2}})^{<\alpha_0, \Lambda_1>} e^{\frac{Q}{2}} |0>,$$

$$e_i e^{\frac{Q}{2}} |0> = 0 , i = 0, 1,$$

where α_1 , $\alpha_0 = -2\alpha_1 + \delta$ are the simple roots of $A_2^{(2)}$, Λ_1 is one basical fundamental weight, and δ is the imaginary root. The Fock space F_1 coincides with the level-one irreducible highest weight module $V(\Lambda_1)$ with the highest weight vector given by $|\Lambda_1\rangle = e^{\frac{Q}{2}}|0\rangle$.

C. Level-one vertex operators

Let V be the 3-dimensional evaluation representation of $U_q(A_2^{(2)})$, $\{v_1, v_0, v_{-1}\}$ be the basis vectors of V, and $E_{i,j}$ be the 3 × 3 matrix whose (i,j)-element is unity and zero otherwise. Then the 3-dimensional level-0 representation V_z of $U_q(A_2^{(2)})$ is given by

$$\begin{split} e_1 &= \alpha^{-1}[2]_{q_1} E_{1,0} + \alpha E_{0,-1}, \quad e_0 = z E_{-1,1}, \\ f_1 &= \alpha^{-1}[2]_{q_1} E_{-1,0} + \alpha E_{0,1} \quad , \quad f_0 = z^{-1} E_{1,-1}, \\ t_1 &= q_1^2 E_{1,1} + q_1^{-2} E_{-1,-1} + E_{0,0}, \quad t_0 = q_0 E_{-1,-1} + q_0^{-1} E_{1,1} + E_{0,0} \quad , \end{split}$$

where $\alpha = \{[2]_{q_1} q^{-\frac{1}{2}}\}^{\frac{1}{2}}$.

We define the dual modules V_z^{*b} of V_z by $\pi_{V^{*b}}(a) = \pi_V(b(a))^t$, $\forall a \in U_q(A_2^{(2)})$, where t is the transposition operation. b is the anti-automorphism defined by

$$b(x) = (-q)^{\hat{\rho}} S(x) (-q)^{-\hat{\rho}} \stackrel{def}{=} (-q)^{3d} (-q^{\frac{1}{2}})^{\frac{1}{2}h_1} S(x) (-q)^{-3d} (-q^{\frac{1}{2}})^{-\frac{1}{2}h_1}, \quad \forall x \in U_q(A_2^{(2)}) ,$$
 (II.13)

where $2\hat{\rho} = 6d + \frac{1}{2}h_1$. A convenient feature of b is that $b^2 = id$ (since $S^2(x) = q^{-2\hat{\rho}}xq^{2\hat{\rho}}$) and that the following isomorphism holds:

$$C: V_z \longrightarrow V_z^{*b} \quad , \quad v_i \otimes z^n \longrightarrow v_{-i}^* \otimes z^n.$$
 (II.14)

Throughout, we denote by $V(\lambda)$ a level-one irreducible highest weight $U_q(A_2^{(2)})$ -module with highest weight λ . Consider the following intertwiners of $U_q(A_2^{(2)})$ -modules:

$$\Phi_{\lambda}^{\mu V}(z): V(\lambda) \longrightarrow V(\mu) \otimes V_z, \qquad \Phi_{\lambda}^{\mu V^*}(z): V(\lambda) \longrightarrow V(\mu) \otimes V_z^{*b},
\Psi_{\lambda}^{V\mu}(z): V(\lambda) \longrightarrow V_z \otimes V(\mu), \qquad \Psi_{\lambda}^{V^*\mu}(z): V(\lambda) \longrightarrow V_z^{*b} \otimes V(\mu).$$

They are intertwiners in the sense that for any $x \in U_q(A_2^{(2)})$,

$$\Theta(z) \cdot x = \Delta(x) \cdot \Theta(z), \quad \Theta(z) = \Phi(z), \Phi^*(z), \Psi(z), \Psi^*(z). \tag{II.15}$$

 $\Phi(z)$ ($\Phi^*(z)$) is called type I (dual) vertex operator and $\Psi(z)$ ($\Psi^*(z)$) type II (dual) vertex operator. We expand the vertex operators as

$$\Phi(z) = \sum_{j=1,0,-1} \Phi_j(z) \otimes v_j , \quad \Phi^*(z) = \sum_{j=1,0,-1} \Phi_j^*(z) \otimes v_j^*,$$
 (II.16)

$$\Psi(z) = \sum_{j=1,0,-1} v_j \otimes \Psi_j(z) , \quad \Psi^*(z) = \sum_{j=1,0,-1} v_j^* \otimes \Psi_j^*(z).$$
 (II.17)

Define the operators $\Phi_j(z), \Phi_j^*(z), \Psi_j(z)$ and $\Psi_j^*(z)$ (j = 1, 0, -1) acting on the Fock space F_1 by

$$\Phi_{-1}(z) = exp\{\sum_{n=1}^{\infty} \frac{[2n]_{q_1}}{n} q^{\frac{9}{2}n} \omega_{-n}(-z)^n\} exp\{\sum_{n=1}^{\infty} \frac{[2n]_{q_1}}{n} q^{-\frac{7}{2}n} \omega_n(-z)^{-n}\} e^Q(zq^4)^{P+\frac{1}{2}}, \tag{II.18}$$

$$\Phi_0(z) = \left\{ \frac{q^{-\frac{1}{2}}}{[2]_{q_1}} \right\}^{\frac{1}{2}} [\Phi_{-1}(z), f_1]_q , \quad \Phi_1(z) = \left\{ \frac{q^{\frac{1}{2}}}{[2]_{q_1}} \right\}^{\frac{1}{2}} [\Phi_0(z), f_1] , \quad (II.19)$$

$$\Psi_1(z) = exp\{-\sum_{n=1}^{\infty} \frac{[2n]_{q_1}}{n} q^{\frac{1}{2}n} \omega_{-n} z^n\} exp\{-\sum_{n=1}^{\infty} \frac{[2n]_{q_1}}{n} q^{-\frac{3}{2}n} \omega_n z^{-n}\} e^{-Q} (-zq)^{-P+\frac{1}{2}}, \tag{II.20}$$

$$\Psi_0(z) = \left\{ \frac{q^{\frac{-1}{2}}}{[2]_{q_1}} \right\}^{\frac{1}{2}} [\Psi_1(z), e_1]_q , \quad \Psi_1(z) = \left\{ \frac{q^{\frac{1}{2}}}{[2]_{q_1}} \right\}^{\frac{1}{2}} [\Psi_0(z), e_1] , \qquad (II.21)$$

$$\Phi_i^*(z) = \Phi_{-i}(-zq^{-3}) , \Psi_i^*(z) = \Psi_{-i}(-zq^3),$$
(II.22)

where $[a,b]_x = ab - xba$ and the bosonic q-oscillators ω_m are defined by

$$\omega_m = -\frac{m}{\{[4m]_{q_1} - (-1)^m [2m]_{q_1}\}[2m]_{q_1}} a_m ,$$

such that $[a_n, \omega_m] = \delta_{m+n,0}$.

According to [17], the operators $\Phi(z)$, $\Phi^*(z)$, $\Psi(z)$ and $\Psi^*(z)$ defined in (II.18—II.22) are the only vertex operators of $U_q(A_2^{(2)})$ which intertwine the level-one irreducible highest weight $U_q(A_2^{(2)})$ -modules:

$$\Phi(z): V(\Lambda_1) \longrightarrow V(\Lambda_1) \otimes V_z, \quad \Phi^*(z): V(\Lambda_1) \longrightarrow V(\Lambda_1) \otimes V_z^{*b},
\Psi(z): V(\Lambda_1) \longrightarrow V_z \otimes V(\Lambda_1), \quad \Psi^*(z): V(\Lambda_1) \longrightarrow V_z^{*b} \otimes V(\Lambda_1).$$
(II.23)

III. EXCHANGE RELATIONS OF VERTEX OPERATORS

In this section, we derive the exchange relations of the type I and type II vertex operators of $U_q(A_2^{(2)})$.

A. The R-matrix

We introduce some abbreviations:

$$(z; p_1, p_2, ..., p_m) = \prod_{\{l_1, l_2, ..., l_m\} = 0}^{\infty} (1 - z p_1^{l_1} p_2^{l_2} \cdots p_m^{l_m}) ,$$

$$\Theta_n(z) = (z; p)(pz^{-1}; p)(p; p) , \{z\} = (z; q^6, q^6) .$$

Let $\overline{R}(z) \in End(V \otimes V)$ be the R-matrix of $U_q(A_2^{(2)})$:

$$\overline{R}(z)(v_i \otimes v_j) = \sum_{k,l} \overline{R}_{kl}^{i,j}(z)v_k \otimes v_l, \quad \forall v_i, v_j, v_k, v_l \in V,$$
(III.24)

where the matrix elements are given by

$$\begin{split} \overline{R}_{1,1}^{1,1}(\frac{z_1}{z_2}) &= \overline{R}_{-1,-1}^{-1,-1}(\frac{z_1}{z_2}) = 1, \quad \overline{R}_{0,-1}^{-1,0}(\frac{z_1}{z_2}) = \overline{R}_{1,0}^{0,1}(\frac{z_1}{z_2}) = \frac{(q-q^{-1})z_2}{z_1q-z_2q^{-1}}, \\ \overline{R}_{-1,0}^{-1,0}(\frac{z_1}{z_2}) &= \overline{R}_{0,-1}^{0,-1}(\frac{z_1}{z_2}) = \overline{R}_{1,0}^{1,0}(\frac{z_1}{z_2}) = \overline{R}_{0,1}^{0,1}(\frac{z_1}{z_2}) = \frac{z_1-z_2}{z_1q-z_2q^{-1}}, \\ \overline{R}_{-1,0}^{0,-1}(\frac{z_1}{z_2}) &= \overline{R}_{0,1}^{1,0}(\frac{z_1}{z_2}) = \frac{(q-q^{-1})z_1}{z_1q-z_2q^{-1}}, \quad \overline{R}_{-1,1}^{-1,1}(\frac{z_1}{z_2}) = \overline{R}_{1,-1}^{1,-1}(\frac{z_1}{z_2}) = \frac{(z_1-z_2)q^2(z_1q+z_2)}{(z_1q-z_2q^{-1})(z_1q^4+z_2q)}, \\ \overline{R}_{-1,1}^{1,-1}(\frac{z_1}{z_2}) &= \frac{(q^2-1)z_1\{z_2+z_2q^2+z_1q^3-z_1q^2\}}{(z_1q-z_2q^{-1})(z_1q^4+z_2q)}, \quad \overline{R}_{1,-1}^{-1,1}(\frac{z_1}{z_2}) = \frac{(q^2-1)z_2\{z_2-z_2q+z_1q^3+z_1q\}}{(z_1q-z_2q^{-1})(z_1q^4+z_2q)}, \\ \overline{R}_{0,0}^{1,-1}(\frac{z_1}{z_2}) &= \overline{R}_{-1,1}^{0,0}(\frac{z_1}{z_2}) = \frac{(q-q^{-1})q^{\frac{7}{2}}z_1(z_1-z_2)}{(z_1q-z_2q^{-1})(z_1q^4+z_2q)}, \\ \overline{R}_{0,0}^{0,1}(\frac{z_1}{z_2}) &= \overline{R}_{1,-1}^{0,0}(\frac{z_1}{z_2}) = \frac{(q^{-1}-q)q^{\frac{3}{2}}z_2(z_1-z_2)}{(z_1q-z_2q^{-1})(z_1q^4+z_2q)}, \\ \overline{R}_{0,0}^{0,0}(\frac{z_1}{z_2}) &= \frac{z_1-z_2}{z_1q-z_2q^{-1}} + \frac{(q-q^{-1})(q+q^4)z_2z_1}{(z_1q-z_2q^{-1})(z_1q^4+z_2q)}, \\ R_{bl}^{ij} &= 0 \quad , \quad \text{otherwise}. \end{split}$$

Define the R-matrices $R^I(z),\,R^{II}(z)$ and $R^U(z)$:

$$R^{I}(z) = r(z)\overline{R}(z)$$
 , $R^{II}(z) = \overline{r}(z)\overline{R}(z)$, $R^{U}(z) = \rho(z)\overline{R}(z)$, (III.25)

where

$$\begin{split} r(z) &= z^{-1} \frac{(-q^3 z; q^6)(q^2 z; q^6)(q^6 z^{-1}; q^6)(-q^5 z^{-1}; q^6)}{(-q^3 z^{-1}; q^6)(q^2 z^{-1}; q^6)(q^6 z; q^6)(-q^5 z; q^6)} \ , \\ \overline{r}(z) &= \frac{(q^4 z; q^6)(-q^3 z; q^6)(-qz^{-1}; q^6)(q^6 z^{-1}; q^6)}{(-q^3 z^{-1}; q^6)(q^4 z^{-1}; q^6)(q^6 z; q^6)(-qz; q^6)} \ , \\ \rho(z) &= \frac{r(z)}{\tau(zq^{-1})} \ , \ \text{and} \ \tau(z) = z^{-1} \frac{\Theta_{q^6}(qz)\Theta_{q^6}(-q^2z)}{\Theta_{q^6}(q^5 z)\Theta_{q^6}(-q^4z)} \end{split}$$

Then the R-matrices satisfy Yang-Baxter equation(YBE) on $V \otimes V \otimes V$

$$R_{12}^{i}(z)R_{13}^{i}(zw)R_{23}^{i}(w) = R_{23}^{i}(w)R_{13}^{i}(zw)R_{12}^{i}(z), \text{ for } i = I, II, U$$

and moreover enjoys: (i) the initial condition, $R^{i}(1) = P$ for i = I, II, U, with P being the permutation operator; (ii) the unitarity condition,

$$R_{12}^{i}(\frac{z}{w})R_{21}^{i}(\frac{w}{z}) = 1$$
, for $i = I, II, U$

where $R_{21}^{i}(z) = PR_{12}^{i}(z)P$; and (iii) the crossing relations

$$(R^i)_{m,n}^{k,l}(z) = (-q^{\frac{1}{2}})^{-\rho_l}(R^{(i)})_{-l,m}^{-n,k}(-z^{-1}q^{-3})(-q^{\frac{1}{2}})^{\rho_n}, \ \ \text{for} \ \ i=I,II.$$

Here and throughout,

$$\rho_1 = -1$$
 , $\rho_0 = 0$, $\rho_{-1} = 1$. (III.26)

B. The Faddeev-Zamolodchikov algebra

Now, we are in the position to calculate the exchange relations of the type I and type II vertex operators of $U_q(A_2^{(2)})$ given in (II.18—II.22).

Firstly we bosonize the derivation operator d:

$$d = \sum_{n=1}^{\infty} n a_{-n} \omega_n - \frac{P^2}{2} . {(III.27)}$$

One can easily check that this d operator obeys the following commutation relations:

$$q^{d}X^{\pm}(z)q^{-d} = X^{\pm}(zq^{-1}) , \quad q^{d}\Phi_{i}(z)q^{-d} = \Phi_{i}(zq^{-1}) , \quad q^{d}\Psi_{i}(z)q^{-d} = \Psi_{i}(zq^{-1}) , \quad (\text{III}.28)$$

as required.

Define

$$\oint dz f(z) = Res(f) = f_{-1} \ , \ \text{ for formal series function } f(z) = \sum_{n \in \mathbb{Z}} f_n z^n.$$

Then the Chevalley generators of $U_q(A_2^{(2)})$ can be expressed by the integrals,

$$e_{1} = \oint z^{-1}dz X^{+}(z), \quad f_{1} = \oint z^{-1}dz X^{-}(z),$$

$$e_{0} = K^{-2} \oint \oint dz w^{-1} dw \ [X^{-}(w), X^{-}(z)]_{q},$$

$$f_{0} = \frac{1}{[4]_{q_{1}}} \oint \oint z^{-2} dz w^{-1} dw \ [X^{+}(z), X^{+}(w)]_{q^{-1}}.$$

From the normal order relations in appendix A, one can also obtain the integral expression of the vertex operators defined in (II.18—II.22)

$$\Phi_0(z) = \{ \frac{q^{-\frac{1}{2}}}{[2]_{q_1}} \}^{\frac{1}{2}} \oint \frac{dw}{w} \frac{(q^{-1} - q)}{zq^3(1 + \frac{zq^5}{w})(1 + \frac{w}{zq^3})} : \Phi_{-1}(z)X^-(w) :, \tag{III.29}$$

$$\Phi_{1}(z) = \frac{1 - q^{2}}{[2]_{q_{1}}} \oint \frac{dw_{1}}{w_{1}} \oint \frac{dw}{w} \frac{(1 - \frac{w}{w_{1}})\{w_{1}^{2}(1 - \frac{w}{w_{1}q^{2}})(1 + \frac{wq}{w_{1}})(1 + \frac{zq^{5}}{w_{1}}) - zwq^{2}(1 + \frac{w_{1}}{zq^{3}})(1 - \frac{wq^{2}}{w_{1}})(1 + \frac{w_{1}q}{w})\}}{z^{3}q^{12}(1 + \frac{zq^{5}}{w_{1}})(1 + \frac{wq}{w_{1}})(1 + \frac{w_{1}q^{5}}{w})(1 + \frac{w}{zq^{5}})(1 + \frac{w}{zq^{5}})(1 + \frac{w}{zq^{5}})} \times : \Phi_{-1}(z)X^{-}(w)X^{-}(w_{1}):, \tag{III.30}$$

$$\Psi_0(z) = \left\{ \frac{q^{-\frac{1}{2}}}{[2]_{q_1}} \right\}^{\frac{1}{2}} \oint \frac{dw}{w} \frac{(1 - q^2)}{wq(1 - \frac{w}{zq^2})(1 - \frac{z}{w})} : \Psi_1(z)X^+(w) :, \tag{III.31}$$

$$\Psi_{-1}(z) = \frac{1 - q^2}{[2]_{q_1}} \oint \frac{dw_1}{w_1} \oint \frac{dw}{w} \frac{(1 - \frac{w}{w_1})\{(1 - \frac{w_1}{wq^2})(1 - \frac{z}{w_1})(1 + \frac{w}{w_1q})w_1 - zq(1 - \frac{w}{w_1q^2})(1 - \frac{w_1}{zq^2})(1 + \frac{w_1}{wq})\}}{zwq(1 - \frac{w}{zq^2})(1 - \frac{z}{w})(1 - \frac{w_1}{zq^2})(1 + \frac{w_1}{wq})(1 - \frac{z}{w_1})(1 + \frac{w}{w_1q})} \times : \Psi_1(z)X^+(w)X^+(w_1) :.$$
(III.32)

By the technique proposed in [10], using the above integral expressions and the relations given in appendix A and appendix B, and after tedious calcualtions, we can show that the bosonic vertex operators defined in (II.18—II.22) satisfy the Faddeev-Zamolodchikov (ZF) algebra

$$\Phi_j(z_2)\Phi_i(z_1) = \sum_{kl} R^I(\frac{z_1}{z_2})_{i,j}^{k,l} \Phi_k(z_1) \Phi_l(z_2), \tag{III.33}$$

$$\Psi_i^*(z_1)\Psi_j^*(z_2) = -\sum_{kl} R^{II}(\frac{z_1}{z_2})_{kl}^{ij}\Psi_l^*(z_2)\Psi_k^*(z_1), \tag{III.34}$$

$$\Psi_i^*(z_1)\Phi_j(z_2) = \tau(\frac{z_1}{z_2})\Phi_j(z_2)\Psi_i^*(z_1) \quad . \tag{III.35}$$

And moreover, the bosonic vertex operators define in (II.18—II.22) have the following invertibility relations,

$$\Phi_i(z)\Phi_j^*(z) = g\delta_{ij}(-q^{\frac{1}{2}})^{\rho_i}, \quad g = \frac{(q^6; q^6)(-q^5; q^6)}{(-q^3; q^6)(q^2; q^6)}, \tag{III.36}$$

$$\sum_{k} (-q^{\frac{1}{2}})^{-\rho_k} \Phi_k^*(z) \Phi_k(z) = g, \tag{III.37}$$

and

$$\Psi_{i}(z_{1})\Psi_{j}^{*}(z_{2}) = \frac{\overline{g}\delta_{ij}(-q^{\frac{1}{2}})^{-\rho_{i}}}{1 - \frac{z_{2}}{z_{1}}} + \text{regular term} \quad \text{when } z_{1} \longrightarrow z_{2},
\overline{g} = \frac{q(q^{4}; q^{6})(-q^{3}; q^{6})}{(-q; q^{6})(q^{6}; q^{6})} .$$
(III.38)

In the derivation of the above relations the following fact is helpful,

$$: \Phi_{-1}(z)\Phi_{-1}(-zq^{-3})X^{-}(zq^{2})X^{-}(-zq^{3})(zq^{2})^{-1}(-zq^{3})^{-1} := id.$$

$$: \Psi_{1}(z)\Psi_{1}(-zq^{3})X^{+}(zq^{2})X^{+}(-zq^{3})(zq^{2})^{-1}(-zq^{3})^{-1} := id.$$

C. Miki's construction of $U_q(A_2^{(2)})$

We generalize the Miki's construction to the twisted quantum affine algebra $U_q(A_2^{(2)})$ case. Define

$$(L^{+}(z))_{i}^{j} = \Phi_{i}(zq^{\frac{1}{2}})\Psi_{j}^{*}(zq^{-\frac{1}{2}}),$$

$$(L^{-}(z))_{i}^{j} = \Phi_{i}(zq^{-\frac{1}{2}})\Psi_{i}^{*}(zq^{\frac{1}{2}}).$$

Then from the Faddeev-Zamolodchikov algebra in (III.33—III.35) and the identity

$$-\frac{r(z)}{\overline{r}(z)} = \frac{\tau(zq^{-1})}{\tau(z^{-1}q^{-1})} = \frac{\tau(zq)}{\tau(z^{-1}q)} ,$$

we can verify by straightforward computations that the L-operators $L^{\pm}(z)$ give a realization of the Reshetikhin-Semenov-Tian-Shansky (RS) algebra [21] at level one in the twisted quantum affine algebra $U_q(A_2^{(2)})$

$$\begin{split} R^U(\frac{z}{w})L_1^{\pm}(z)L_2^{\pm}(w) &= L_2^{\pm}(w)L_1^{\pm}(z)R^U(\frac{z}{w}),\\ R^U(\frac{z^+}{w^-})L_1^{+}(z)L_2^{-}(w) &= L_2^{-}(w)L_1^{+}(z)R^U(\frac{z^-}{w^+}), \end{split}$$

where $L_1^{\pm}(z) = L^{\pm}(z) \otimes 1$, $L_2^{\pm}(z) = 1 \otimes L^{\pm}(z)$ and $z^{\pm} = zq^{\pm\frac{1}{2}}$. We remark that the Drinfeld bases (II.4—II.9) of the twisted quantum affine algebra $U_q(A_2^{(2)})$ can also be derived by using the twisted RS algebra and the corresponding Gauss decomposition [22].

IV. THE IZERGIN-KOREPIN MODEL

In this section, we give a mathematical definition of the Izergin-Korepin model on an infinite lattice.

A. Space of states

By means of the R-matrix (III.24) of $U_q(A_2^{(2)})$, one can define the Izergin-Korepin model on the infinte lattice $\cdots \otimes V \otimes V \otimes V \cdots$. Let h be the operator on $V \otimes V$ such that

$$PR(\frac{z_1}{z_2}) = 1 + uh + \cdots, \qquad u \longrightarrow 0,$$

P: the permutation operator, $e^u \equiv \frac{z_1}{z_2}$.

The Hamiltonian H of the Izergin-Korepin model is defined by [14]

$$H = \sum_{l \in \mathbb{Z}} h_{l+1,l}. \tag{IV.39}$$

H acts formally on the infinite tensor product,

$$\cdots V \otimes V \otimes V \cdots$$
 (IV.40)

It can be easily checked that

$$[U_q'(A_2^{(2)}), H] = 0,$$

where $U_q'(A_2^{(2)})$ is the subalgebra of $U_q(A_2^{(2)})$ with the derivation operator d being dropped. So $U_q'(A_2^{(2)})$ plays the role of infinite dimensional non-abelian symmetries of the Izergin-Korepin model on the infinite lattice. Since the level-one vertex operators only exist between $V(\Lambda_1)$ and itself, following [3], we can replace the infinite tensor product (IV.40) by the level-0 $U_q(A_2^{(2)})$ -module,

$$\mathbf{H} = \operatorname{Hom}(V(\Lambda_1), V(\Lambda_1)) \cong V(\Lambda_1) \otimes V(\Lambda_1)^{*b}$$

where $V(\Lambda_1)$ is level-one irreducible highest weight $U_q(A_2^{(2)})$ -module and $V(\Lambda_1)^{*b}$ is the dual module of $V(\Lambda_1)$. By theorem 2, this homomorphism can be realized by applying the type I vertex operators repeatedly. So we shall make the (hypothetical) identification:

"the space of physical states" = $V(\Lambda_1) \otimes V(\Lambda_1)^{*b}$.

Namely, we take

$$\mathbf{H} \equiv End(V(\Lambda_1))$$

as the space of states of the Izergin-Korepin model on the infinite lattice. The left action of $U_q(A_2^{(2)})$ on **H** is defined by

$$x.f = \sum x_{(1)} \circ f \circ b(x_{(2)}), \quad \forall x \in U_q(A_2^{(2)}), \ f \in \mathbf{H},$$

where we have used notation $\Delta(x) = \sum x_{(1)} \otimes x_{(2)}$. A linear operator of the form $O = A \otimes B$ $(A \in End(V(\Lambda_1)))$ and $B \in End(V(\Lambda_1)^{*b})$ operate on a state f as $f \to A \circ f \circ B^t$.

Note that **H** has the unique canonical element which is referred as to the physical vacuum [4] and denoted by $|vac\rangle$

$$|vac> = \chi^{-\frac{1}{2}}(-q)^{-\hat{\rho}},$$
 (IV.41)

where χ coincide with the character of $U_q(A_2^{(2)})$ -module $V(\Lambda_1)$

$$\chi = tr_{V(\Lambda_1)}(q^{-2\hat{\rho}}) = q^{\frac{1}{4}}(1+q) \prod_{n=1}^{\infty} (1+q^{6n-1})(1+q^{6n+1}).$$

The proof of the above character formula is given in appendix C.

B. Local structure and local operators

Following Jimbo and Miwa [4], we use the type I vertex operators and their variants to incorporate the local structure into the space of physical states **H**, that is to formulate the action of local operators of the Izergin-Korepin model on the infinite tensor product (IV.40) in terms of their actions on **H**.

Using the isomorphisms (c.f. theorem 2)

$$\Phi(z): V(\Lambda_1) \longrightarrow V(\Lambda_1) \otimes V_z,
\Phi^{*,t}(-zq^3): V_z \otimes V(\Lambda_1)^{*b} \longrightarrow V(\Lambda_1)^{*b},$$
(IV.42)

were t is the transposition on the quantum space, we have the following identification:

$$V(\Lambda_1) \otimes V(\Lambda_1)^{*b} \stackrel{\sim}{\to} V(\Lambda_1) \otimes V_z \otimes V(\Lambda_1)^{*b} \stackrel{\sim}{\to} V(\Lambda_1) \otimes V(\Lambda_1)^{*b}$$

The resulting isomorphism can be identified with the translation (or shift) operator defined by

$$T = g^{-1} \sum_{i} \Phi_{i}(1) \otimes \Phi_{i}^{*,t}(-q^{3}).$$

Its inverse is given by

$$T^{-1} = g^{-1} \sum_{i} \Phi_{i}^{*}(1) \otimes \Phi_{i}^{t}(-q^{-3}).$$

Thus we can define the local operators on V as operators on H [4]. Let us label the tensor components from the middle as $1, 2, \cdots$ for the left half and as $0, -1, -2, \cdots$ for the right half. The operators acting on the site 1 are defined by

$$E_{i,j} \stackrel{\text{def}}{=} E_{i,j}^{(1)} = g^{-1} (-q^{-\frac{1}{2}})^{-\rho_j} \Phi_i^*(1) \Phi_j(1) \otimes id.$$
 (IV.43)

In particular, we have spin operator S_z

$$S_z = g^{-1} \{ \Phi_1^*(1) \Phi_1(1) (-q^{-\frac{1}{2}}) + \Phi_0^*(1) \Phi_0(1) - \Phi_{-1}^*(1) \Phi_{-1}(1) (-q^{-\frac{1}{2}})^{-1} \} \otimes id.$$

More generally we set

$$E_{i,i}^{(n)} = T^{-(n-1)}E_{i,j}T^{n-1} \quad (n \in \mathbb{Z}).$$
(IV.44)

Then, from the invertibility relations of the type I vertex operators of $U_q(A_2^{(2)})$, we can show that the local operators $E_{ij}^{(n)}$ acting on **H** satisfy the following relations:

$$E_{i,j}^{(m)} E_{k,l}^{(n)} = \begin{cases} \delta_{jk} E_{i,l}^{(n)} & \text{if} \quad m = n \\ E_{k,l}^{(n)} E_{i,j}^{(m)} & \text{if} \quad m \neq n. \end{cases}$$
(IV.45)

Moreover, we can define transfer matrix of the Izergin-Korepin model on the infinite lattice as

$$T(z) = g^{-1} \sum_{i} \Phi_{i}(z) \otimes \Phi_{i}^{*,t}(-zq^{3})$$
 (IV.46)

The commutativity of the transfer matrix ,[T(z),T(w)]=0, follows from the ZF algebraic relations (III.33). The translation operator is T=T(1), as expected. Comparing with the definition of Hamiltonian of the Izergin-Korepin model (IV.39), the action of such a Hamiltonian on **H** can be given by

$$H = z \frac{d}{dz} \{ lnT(z) \} |_{z=1}$$
 (IV.47)

As is expected from the physical point of view, the vacuum vector $|vac\rangle$ is translationally invariant and singlet (i.e. belong to the trivial representation of $U_q(A_2^{(2)})$)

$$T|vac> = |vac>,$$

 $x.|vac> = \epsilon(x)|vac>.$ (IV.48)

This is proved as follows.

$$\begin{split} T(z)|vac> &= \chi^{-\frac{1}{2}}g^{-1}\sum_{i}\Phi_{i}(z)(-q)^{-\hat{\rho}}\Phi_{i}^{*}(-zq^{3})\\ &\stackrel{def}{=} \chi^{-\frac{1}{2}}g^{-1}\sum_{i}\Phi_{i}(z)(-q)^{-3d}(-q^{\frac{1}{2}})^{-\frac{1}{2}h_{1}}\Phi_{i}^{*}(-zq^{3})\\ &= \chi^{-\frac{1}{2}}g^{-1}\sum_{i}\Phi_{i}(z)\Phi_{i}^{*}(zq^{6})(-q^{\frac{1}{2}})^{\rho_{i}}(-q)^{-\hat{\rho}}\\ &\stackrel{def}{=} \chi^{-\frac{1}{2}}g^{-1}\sum_{i}\Phi_{-i}^{*}(-zq^{3})\Phi_{-i}(-zq^{3})(-q^{\frac{1}{2}})^{-\rho_{-i}}(-q)^{-\hat{\rho}}\\ &= \chi^{-\frac{1}{2}}(-q)^{-\hat{\rho}} = |vac>, \end{split}$$

where we have used the fact $\rho_i = -\rho_{-i}$, (II.13) and (III.36). Similary,

$$x.|vac\rangle = \chi^{-\frac{1}{2}} \sum_{x_{(1)}} (-q)^{\hat{\rho}} b(x_{(2)})$$
$$= \chi^{-\frac{1}{2}} \sum_{x_{(1)}} S(x_{(2)}) (-q)^{\hat{\rho}}$$
$$= \epsilon(x)|vac\rangle$$

In the third line we have used the axioms of Hopf algebra [4]

$$m \circ (id \otimes S) \circ \Delta = \epsilon.$$

This completes the proof.

For any local operator O on \mathbf{H} defined in (IV.44), its vacuum expectation value is given by

$$< vac|O|vac> \stackrel{def}{=} < O> = \frac{tr_{V(\Lambda_1)}(q^{-2\hat{\rho}}O)}{tr_{V(\Lambda_1)}(q^{-2\hat{\rho}})} = \frac{tr_{V(\Lambda_1)}(q^{-6d-\frac{1}{2}h_1}O)}{tr_{V(\Lambda_1)}(q^{-6d-\frac{1}{2}h_1})},$$

where the normalization < vac|vac> = 1 has been chosen. We shall denote the correlator < vac|O|vac> by < O>. By the proposition 4 and the definition of the local operators $E_{i,j}^{(n)}$ (IV.44), we have

$$< E_{i,j}^{(n)} > = < vac|T^{-(n-1)}E_{i,j}T^{(n-1)}|vac> = < vac|E_{i,j}|vac> = < E_{i,j} >$$

C. The n-particle states and form factors

In order to construct the general eigenstates of the transfer matrix of the Izergin-Korepin model (IV.46), we employ the type II vertex operators. Define the n-particle states

$$|\xi_{n},\xi_{n-1},\cdots,\xi_{1}\rangle_{i_{n},i_{n-1},\cdots,i_{1}} = \overline{g}^{-\frac{n}{2}}\Psi_{i_{n}}^{*}(\xi_{n})\Psi_{i_{n-1}}^{*}(\xi_{n-1})\cdots\Psi_{i_{1}}^{*}(\xi_{1})|vac\rangle$$

$$= \overline{g}^{-\frac{n}{2}}\chi^{-\frac{1}{2}}\Psi_{i_{n}}^{*}(\xi_{n})\Psi_{i_{n-1}}^{*}(\xi_{n-1})\cdots\Psi_{i_{1}}^{*}(\xi_{1})(-q)^{-\hat{\rho}}, \qquad (IV.49)$$

and its dual states

$$|i_n, i_{n-1}, \dots, i_1| < \xi_n, \xi_{n-1}, \dots, \xi_1| = \overline{g}^{-\frac{n}{2}} \chi^{-\frac{1}{2}} (-q)^{-\hat{\rho}} \Psi_{i_1}(\xi_1) \Psi_{i_2}(\xi_2) \dots \Psi_{i_n}(\xi_n).$$
 (IV.50)

Using the commutation relation of the type I vertex operators, (III.35), one can verify that

$$T(z)|\xi_{n},\dots,\xi_{1}\rangle_{i_{n},i_{n-1},\dots,i_{1}} = g^{-1}\overline{g}^{-\frac{n}{2}}\chi^{-\frac{1}{2}}\sum_{i}\Phi_{i}(z)\Psi_{i_{n}}^{*}(\xi_{n})\Psi_{i_{n-1}}^{*}(\xi_{n-1})\dots\Psi_{i_{1}}^{*}(\xi_{1})(-q)^{-\hat{\rho}}\Phi_{i}^{*}(-zq^{3})$$

$$= g^{-1}\overline{g}^{-\frac{n}{2}}\chi^{-\frac{1}{2}}\sum_{i}\Phi_{i}(z)\Psi_{i_{n}}^{*}(\xi_{n})\Psi_{i_{n-1}}^{*}(\xi_{n-1})\dots\Psi_{i_{1}}^{*}(\xi_{1})\Phi_{i}^{*}(zq^{6})(-q^{\frac{1}{2}})^{\rho_{i}}(-q)^{-\hat{\rho}}$$

$$= g^{-1}\overline{g}^{-\frac{n}{2}}\chi^{-\frac{1}{2}}\sum_{i}\prod_{j=1}^{n}\tau(\xi_{j}/z)\Phi_{i}(z)\Phi_{i}^{*}(zq^{6})(-q^{\frac{1}{2}})^{\rho_{i}}\Psi_{i_{n}}^{*}(\xi_{n})\Psi_{i_{n-1}}^{*}(\xi_{n-1})$$

$$\times\dots\Psi_{i_{1}}^{*}(\xi_{1})(-q)^{-\hat{\rho}}$$

$$= \prod_{i=1}^{n}\tau(\xi_{j}/z)|\xi_{n},\dots,\xi_{1}\rangle_{i_{n},i_{n-1},\dots,i_{1}}.$$

Here we have used $\tau(zq^6) = \tau(z)$. Therefore the n-particle states (IV.49) are the eigenstates of transfer matrix T(z). Likewise for its dual states (IV.50). Hence, the Hamiltonian H (IV.47) on the n-particle states are given by

$$H|\xi_n,\dots,\xi_1>_{i_n,i_{n-1},\dots,i_1}=\sum_{i=1}^n\epsilon(\xi_j)|\xi_n,\dots,\xi_1>_{i_n,i_{n-1},\dots,i_1},$$

where

$$\epsilon(z) = -z \frac{d}{z} ln\tau(z).$$

These results coincide with the energy of the elementary excitations derived from the Bethe ansatz method [23]. In much the same way as the correlation functions, the form factors of a local operator O of the form $O = A \otimes B$ can be given by

$$< vac|O(\xi_n, \dots, \xi_1) >_{i_n, i_{n-1}, \dots, i_1} = \frac{\overline{g}^{-\frac{n}{2}} tr_{V(\Lambda_1)}((-q)^{-\hat{\rho}} A \Psi_{i_n}^*(\xi_n) \Psi_{i_{n-1}}^*(\xi_{n-1}) \dots \Psi_{i_1}^*(\xi_1)(-q)^{-\hat{\rho}} B^t)}{tr_{V(\Lambda_1)}(q^{-2\hat{\rho}})}$$

In particular, for the spin operator S_z , we have

$$< vac|S_z|\xi_n, \dots, \xi_1>_{i_n, i_{n-1}, \dots, i_1}$$

$$= g^{-1} \frac{\overline{g}^{-\frac{n}{2}} tr_{V(\Lambda_1)} (q^{-2\hat{\rho}} \{\Phi_1^*(1)\Phi_1(1)(-q^{\frac{1}{2}}) + \Phi_0^*(1)\Phi_0(1) - \Phi_{-1}^*(1)\Phi_{-1}(1)(-q^{\frac{1}{2}})^{-1} \} \Psi_{i_n}^*(\xi_n) \Psi_{i_{n-1}}^*(\xi_{n-1}) \cdots \Psi_{i_1}^*(\xi_1))}{tr_{V(\Lambda_1)} (q^{-2\hat{\rho}})}$$

V. CORRELATION FUNCTIONS

The aim of this section is to calculate $\langle E_{mn} \rangle$. The generalization to the calculation of the multi-point functions and the form factors is straightforward.

Set

$$P_n^m(z_1, z_2|q) = \frac{tr_{V(\Lambda_1)}(q^{-2\hat{\rho}}\Phi_m^*(z_1)\Phi_n(z_2))}{tr_{V(\Lambda_1)}(q^{-2\hat{\rho}})},$$

then $\langle E_{mn} \rangle = g^{-1} P_n^m(z, z|q) (-q^{\frac{1}{2}})^{-\rho_n}$. Using the Clavelli-Shapiro technique [24] which we will present it in appendix C, we get

$$P_n^m(z_1, z_2|q) \stackrel{def}{=} \delta_{mn} P_m(z_1, z_2|q)$$

$$= \frac{\delta_{mn} (C_{-1})^2 (C^{(-)})^2}{\gamma} G(-\frac{z_2 q^3}{z_1}) \oint \frac{dw}{w} \oint \frac{dw_1}{w_1} \theta(\frac{-z_1 z_2 q^5}{w w_1}) I_m(z_1, z_2; w, w_1), \tag{V.51}$$

where $\theta(z)$ is the elliptic function,

$$\theta(z) \stackrel{def}{=} \sum_{n \in \mathbb{Z}} q^{3(n+\frac{1}{2})^2} z^n \quad ,$$

$$\begin{split} I_1(z_1,z_2;w,w_1) &= (\frac{-z_1z_2q^5}{ww_1})^{\frac{1}{2}}W_1(-\frac{wq^3}{z_1})W_1(-\frac{w_1q^3}{z_1})\{\frac{w}{z_2q^4}W_1(\frac{w}{z_2})W_1(\frac{w_1}{z_2})H(\frac{w_1}{w}) + \frac{z_2q^5}{w}W_2(\frac{z_2}{w_1})W_2(\frac{z_2}{w})H(\frac{w}{w_1})\\ &- qW_2(\frac{z_2}{w})W_1(\frac{w_1}{z_2})H(\frac{w_1}{w}) - W_2(\frac{z_2}{w_1})W_1(\frac{w}{z_2})H(\frac{w}{w_1})\} \quad, \\ I_0(z_1,z_2;w,w_1) &= (\frac{-z_1z_2q^5}{ww_1})^{\frac{1}{2}}W_1(-\frac{wq^3}{z_1})W_2(\frac{z_2}{w})H(\frac{w_1}{w})\{\frac{z_1q^2}{w}W_2(-\frac{z_1}{wq^3})W_1(\frac{w_1}{z_2}) - \frac{z_2q^5}{w_1}W_1(-\frac{w_1q^3}{z_1})W_2(\frac{z_2}{w_1})\\ &+ W_1(-\frac{wq^3}{z_1})W_1(\frac{w_1}{z_2}) + q^2(\frac{-z_1z_2q^5}{ww_1})^{\frac{1}{2}}W_2(-\frac{z_1}{wq^3})W_2(\frac{z_2}{w_1})\} \quad, \\ I_{-1}(z_1,z_2;w,w_1) &= -(\frac{-z_1z_2q^5}{ww_1})^{\frac{3}{2}}W_2(\frac{z_2}{w})W_2(\frac{z_2}{w_1})\{qW_2(-\frac{z_1}{wq^3})W_1(-\frac{w_1q^3}{z_1})H(\frac{w_1}{w}) + W_2(-\frac{z_1}{w_1q^3})H(\frac{w}{w_1})\\ &+ \frac{w}{z_1q}W_1(-\frac{wq^3}{z_1})W_1(-\frac{w_1q^3}{z_1})H(\frac{w_1}{w}) + \frac{z_1q^2}{w}W_2(-\frac{z_1}{w_1q^3})W_2(-\frac{z_1}{wq^3})H(\frac{w}{w_1})\} \end{split}$$

and the functions G(z), $W_1(z)$, $W_2(z)$, H(z) and the constant C_{-1} , $C^{(-)}$ are given in appendix D (V.53— V.56). In particular, the correlation function of spin operator S_z is given by

$$< vac|S_z|vac > = P_1(z, z|q)(-q^{-\frac{1}{2}}) + P_0(z, z|q) - P_{-1}(z, z|q)(-q^{-\frac{1}{2}})^{-1}$$
 (V.52)

where $P_m(z_1, z_2|q)$ are defined in (V.51).

We now derive the difference equations satisfied by these one-point functions. Noting (III.28) and using the cyclicity of trace, we get the difference equations of $P_m(z_1, z_2|q)$

$$\begin{split} P_m(z_1,z_2|q) &= \chi^{-1} tr_{V(\Lambda_1)}(q^{-6d-\frac{1}{2}h_1} \Phi_m^*(z_1) \Phi_m(z_2)) \\ &= \chi^{-1} tr_{V(\Lambda_1)}(q^{-6d-\frac{1}{2}h_1} \Phi_{-m}(-z_1q^{-3}) \Phi_m(z_2)) \\ &= \chi^{-1} \sum_n R^I(z_2,-z_1q^{-3})_{m,-m}^{n,-n} tr_{V(\Lambda_1)}(q^{-6d-\frac{1}{2}h_1} \Phi_n(z_2) \Phi_n^*(z_1)) \\ &= \chi^{-1} \sum_n q^{-\rho_n} R^I(z_2,-z_1q^{-3})_{m,-m}^{n,-n} tr_{V(\Lambda_1)}(\Phi_n(z_2q^6)q^{-6d-\frac{1}{2}h_1} \Phi_n^*(z_1)) \\ &= \sum_n q^{-\rho_n} R^I(z_2,-z_1q^{-3})_{m,-m}^{n,-n} P_n(z_1,z_2q^6|q) \quad . \end{split}$$

ACKNOWLEDGEMENTS

W.L.Yang would like to thank Prof.Fan for fruitful discussions.Y.-Z.Zhang thanks Australian Research Council IREX programme for an Asia-Pacific Link Award and Institute of Modern Physics of Northwest University for hospitality. The financial support from the National Natural Science Foundation of China and Australian Research Council large, small, QEII fellowship grants is also gratefully acknowledge.

APPENDIX A.

In this apppendix, we give the normal order relations of fundmental bosonic fields:

$$\begin{split} &\Phi_{-1}(z)\Phi_{-1}(w) = zq^4g(w/z):\Phi_{-1}(z)\Phi_{-1}(w): \quad, \\ &\Phi_{-1}(z)X^-(w) = \frac{1}{zq^4(1+\frac{w}{zq^3})}:\Phi_{-1}(z)X^-(w): \quad, \\ &X^-(w)\Phi_{-1}(z) = \frac{1}{w(1+\frac{zq^5}{w})}:\Phi_{-1}(z)X^-(w): \quad, \\ &\Phi_{-1}(z)X^+(w) = (zq^4+w):\Phi_{-1}(z)X^+(w):X^+(w)\Phi_{-1}(z) \quad, \\ &\Psi_1(z)\Psi_1(w) = -zq\overline{g}(w/z):\Psi_1(z)\Psi_1(w): \quad, \\ &\Psi_1(z)\Phi_{-1}(w) = -(zq)^{-1}h_1(w/z):\Psi_1(z)\Phi_{-1}(w): \quad, \end{split}$$

$$\begin{split} &\Phi_{-1}(z)\Psi_{1}(w)=(zq^{4})^{-1}h_{2}(w/z):\Phi_{-1}(z)\Psi_{1}(w):\ ,\\ &\Psi_{1}(z)X^{-}(w)=(w-zq):\Phi_{-1}(z)X^{-}(w):\ ,\ X^{-}(w)\Psi_{1}(z),\\ &X^{+}(w)\Psi_{1}(z)=\frac{1}{w-z}:\Psi_{1}(z)X^{+}(w):\ ,\\ &\Psi_{1}(z)X^{+}(w)=-\frac{1}{zq(1-\frac{w}{zq^{2}})}:\Psi_{1}(z)X^{+}(w):\ , \end{split}$$

where

$$g(z) = \frac{(-q^3z; q^6)(q^2z; q^6)}{(q^6z; q^6)(-q^5z; q^6)} , \quad \overline{g}(z) = \frac{(-qz; q^6)(z; q^6)}{(q^4z; q^6)(-q^3z; q^6)},$$

$$h_1(z) = \frac{(-q^8z; q^6)(q^7z; q^6)}{(q^5z; q^6)(-q^4z; q^6)} , \quad h_2(z) = \frac{(-q^2z; q^6)(qz; q^6)}{(q^{-1}z; q^6)(-q^{-2}z; q^6)}.$$

and

$$\begin{split} X^+(z)X^+(w) &= \frac{(z-w)(z-wq^{-2})}{z+wq^{-1}} : X^+(z)X^+(w) : \quad , \\ X^-(z)X^-(w) &= \frac{(z-w)(z-wq^2)}{z+wq} : X^-(z)X^-(w) : \quad , \\ X^+(z)X^-(w) &= \frac{z+w}{(z-wq)(z-wq^{-1})} : X^+(z)X^+(w) : \quad , \\ X^-(w)X^+(z) &= \frac{z+w}{(w-zq)(w-zq^{-1})} : X^+(z)X^+(w) : \quad . \end{split}$$

APPENDIX B.

By means of the bosonic realization of $U_q(A_2^{(2)})$, the integral expressions of the vertex operators and the technique given in Ref. [10], one can check the following relations

• For the type I vertex operators

$$\begin{split} [\Phi_1(z),f_1]_{q^{-1}} &= 0, \quad \Phi_1(z) = \frac{1}{\alpha} [\Phi_0(z),f_1], \quad \Phi_0(z) = \frac{\alpha}{[2]_{q_1}} [\Phi_{-1}(z),f_1]_q, \\ [\Phi_1(z),e_1] &= \frac{[2]_{q_1}}{\alpha} t_1 \Phi_0(z), \quad [\Phi_0(z),e_1] = \alpha t_1 \Phi_{-1}(z), \quad [\Phi_{-1}(z),e_1] = 0, \\ \Phi_1(z)t_1 &= q t_1 \Phi_1(z), \quad \Phi_0(z)t_1 = t_1 \Phi_0(z), \quad \Phi_{-1}(z)t_1 = q^{-1} t_1 \Phi_{-1}(z), \end{split}$$

where we take $\alpha = \{[2]_{q_1}q^{-\frac{1}{2}}\}^{\frac{1}{2}}$.

• For the type II vertex operators

$$\begin{split} \Psi_{-1}(z) &= \frac{1}{\alpha} [\Psi_0(z), e_1], \quad \Psi_0(z) = \frac{\alpha}{[2]_{q_1}} [\Psi_1(z), e_1]_q, \quad [\Psi_{-1}(z), e_1]_{q^{-1}} = 0, \\ [\Psi_1(z), f_1] &= 0, \quad [\Psi_0(z), f_1] = \alpha t_1^{-1} \Psi_1(z), \quad [\Psi_{-1}(z), f_1] = \frac{[2]_{q_1}}{\alpha} t_1^{-1} \Psi_0(z), \\ \Psi_1(z) t_1 &= q t_1 \Psi_1(z), \quad \Psi_0(z) t_1 = t_1 \Psi_0(z), \quad \Psi_{-1}(z) t_1 = q^{-1} t_1 \Psi_{-1}(z). \end{split}$$

APPENDIX C.

In computating the correlation functions, one encounters the trace of the form

$$tr(x^{-d}e^{\sum_{m=1}^{\infty}A_{m}a_{-m}}e^{\sum_{m=1}^{\infty}B_{m}a_{m}})f^{P})$$

where A_m , B_m and f are all some coefficients. We can calculate the contributions from the oscillators modes and the zero modes separately. The trace over the oscillator modes can be carried out as follows by using the Clavelli-Shapiro technique [24]. Let us introduce the extra oscillators a'_m which commutate with a_m . a'_m satisfy the same commutation relations as those satisfied by a_m . Introduce the operators

$$H_m = \frac{a_m \otimes 1}{1 - x^m} + 1 \otimes a'_{-m} \quad , \quad m > 0 \quad ,$$
 $H_m = a_m \otimes 1 + \frac{1 \otimes a'_{-m}}{x^m - 1} \quad , \quad m < 0 \quad ,$

which act on the space of the tensor Fock spaces of $\{a_m\}$ and $\{a'_m\}$. Then for any bosonic operator $O(a_m)$, one can show

$$tr(x^{-d}O(a_m)) = \frac{\langle 0|O(H_m)|0\rangle}{\prod_{n=1}(1-x^n)}$$

providing that d satisfies the derivation properties (II.4). We write $< 0|O(H_m)|0> \equiv << O(a_m) >>$. Then by the Wick theorem, one obtains

$$\begin{split} &<<\Phi_{-1}(z)\Phi_{-1}(w)>>=C_{-1}C_{-1}G(w/z)\ ,\\ &<<\Phi_{-1}(z)X^-(w)>>=C_{-1}C^{(-)}W_1(w/z)\ ,\\ &<< X^-(z)\Phi_{-1}(w)>>=C_{-1}C^{(-)}W_2(w/z)\ ,\\ &<< X^-(z)X^-(w)>>=C^{(-)}C^{(-)}H(w/z)\ , \end{split}$$

where

$$C_{-1} = \frac{\{-q^9\}\{q^8\}}{\{q^{12}\}\{-q^{11}\}} \quad , \quad C^{(-)} = \frac{(q^6; q^6)(q^8; q^6)}{(-q^7; q^6)} \quad , \tag{V.53}$$

$$G(z) = \frac{\{-q^3z\}\{q^2z\}\{-q^9z^{-1}\}\{q^8z^{-1}\}}{\{q^6z\}\{-q^5z\}\{q^{12}z^{-1}\}\{-q^{11}z^{-1}\}} , \qquad (V.54)$$

$$W_1(z) = \frac{(-q^{11}z^{-1}; q^6)}{(-q^{-3}z; q^6)} \quad , \quad W_2(z) = \frac{(-q^3z^{-1}; q^6)}{(-q^5z; q^6)} \quad , \tag{V.55}$$

$$H(z) = \frac{(q^2 z; q^6)(z; q^6)(q^8 z^{-1}; q^6)(q^6 z^{-1}; q^6)}{(-qz; q^6)(-q^7 z^{-1}; q^6)}$$
 (V.56)

Now , we use the above technique to calculate the character of $U_q(A_2^{(2)})$ -module $V(\Lambda_1)$

$$\chi = tr_{V(\Lambda_1)}(q^{-2\hat{\rho}}) = tr_{F_1}(q^{-6d}q^{-\frac{1}{2}h_1}) = tr_{F_1}(q^{-6d}q^{-P})$$

$$= \frac{1}{\prod_{n=1}^{\infty} (1 - q^{6n})} \sum_{n \in \mathbb{Z}} q^{3(n + \frac{1}{2})^2} q^{-(n + \frac{1}{2})}$$

$$= q^{\frac{1}{4}}(1 + q) \prod_{n=1}^{\infty} (1 + q^{6n-1})(1 + q^{6n+1}) .$$

We have used the Jacobi tripe product identity [25]

$$\sum_{n \in \mathbb{Z}} q^{(n+v-\frac{1}{2})^2} t^n = q^{\frac{1}{12}} q^{v^2-v+\frac{1}{6}} \prod_{n=1}^{\infty} (1-q^{2n}) (1+q^{2(n+v-1)}t) (1+q^{2(n-v)}t^{-1})$$

- [1] I.B.Frenkel and N.Yu.Reshetikhin, Commun. Math. Phys. 146 (1992), 1.
- [2] I.B.Frenkel and N.Jing, Proc. Nat'l. Acad. Sci. USA 85 (1988), 9373.
- [3] B.Davies, O.Foda, M.Jimbo, T.Miwa and A.Nakayashiki, Commun. Math. Phys. 151 (1993), 89.
- [4] M.Jimbo and T.Miwa, Algebraic Analysis of Solvable Lattice Models, CBMS Regional Conference Series in Mathematics, Vol.85, AMS, 1994.
- [5] M.Idzumi, Int. J. Mod. Phys. A9 (1994), 449.
- [6] H.Bougurzi and R.Weston, Nucl. Phys. B417 (1994), 439.
- [7] J.Hong, S.J.Kang, T.Miwa and R.Weston, J. Phys. A31 (1998), L515.
- [8] Y.Koyama, Commun. Math. Phys. 164(1994), 277.
- [9] S.Lukyanov and Y.Pugai, Nucl. Phys. **B473**(1996), 631.
- [10] Y.Asai, M.Jimbo, T.Miwa and Y.Pugai, J. Phys. A29 (1996), 6595.
- [11] W.L.Yang and Y.-Z.Zhang, Highest weight representations of $U_q(\widehat{sl}(2|1))$ and correlation functions of q-deformed super-symmetric t-J model, e-print math.QA/9812027 (1998), Nucl. Phys. **B** (1999), in press.
- [12] A.G.Izergin and V.E.Korepin, Commun. Math. Phys. **79** (1981), 303.
- [13] N.Yu.Reshetikhin, Lett. Math. Phys. 7 (1993), 205.
- [14] L.Mezincescu and R.I.Nepomechie, Nucl. Phys. 372 (1992), 597; Int. J. Mod. Phys. A6 (1991), 5231.
- [15] A.Kuniba, Nucl. Phys. B335 (1991), 801.
- [16] M.J.Martins, Nucl. Phys. **B450** (1995), 768.
- [17] N.H.Jing and K.C.Misra, Vertex operators for twisted quantum affine algebras, e-print q-alg/9701034.
- [18] V.Chari and A.Pressley, Commun. Math. Phys. 196 (1998),461.
- [19] G.W.Delius, M.D.Gould and Y.-Z.Zhang, Int. J. Mod. Phys. A11 (1996),3415.
- [20] V.G.Drinfeld, Sov. Math. Dokl. 36 (1988) 212.
- [21] N.Yu.Reshetikhin, M.A.Semenov-Tian-Shansky, Lett. Math. Phys. 19 (1990), 133.
- [22] W.L.Yang and Y.-Z.Zhang, The twisted quantum affine algebra $U_q(A_2^{(2)})$ and its bosonization at any level., in preparation.
- [23] V.I.Vichirko and N.Yu. Reshetikhin, Thero. Math. Phys. Vol.56, No.2 (1983), 260.
- [24] L.Clavelli and J.A.Shapiro, Nucl. Phys. **B57** (1973), 490.
- [25] D. Mumford, Tata lectures on the theta function (Birhauser, Basel, 1983).