

A geometric estimate for a periodic Schrödinger operator whose potential is the curvature of a spherical curve. *

THOMAS FRIEDRICH (Berlin)

September 4, 2018

Abstract

We estimate from below by geometric data the eigenvalues of the periodic Sturm-Liouville operator $-4\frac{d^2}{ds^2} + \kappa^2(s)$ with potential given by the curvature of a closed curve.

Subj. Class.: Differential geometry.

1991 MSC: 58G25, 53A05.

Keywords: Dirac operator, spectrum, surfaces, Schrödinger operators, Fenchel inequality.

1 Introduction

Let $X^3(c)$ be a 3-dimensional space form of constant curvature $c = 0$ or 1 and admitting a real Killing spinor with respect to some spin structure. Consider a compact, oriented and immersed surface $M^2 \subset X^3(c)$ with mean curvature H . The spin structure of $X^3(c)$ induces a spin structure on M^2 . Denote by D the corresponding Dirac operator acting on spinor fields defined over the surface M^2 . The first eigenvalue $\lambda_1^2(D)$ of the operator D^2 and the first eigenvalue μ_1 of the Schrödinger operator $\Delta + H^2 + c$ are related by the inequality

$$\lambda_1^2(D) \leq \mu_1(\Delta + H^2 + c).$$

Equality holds if and only if the mean curvature H is constant (see [1], [5]). Moreover, the Killing spinor defines a map $f \mapsto \Phi(f)$ of the space $L^2(M^2)$ of functions into the space $L^2(M^2; S)$ of spinors such that

$$\|D(\Phi(f))\|_{L^2}^2 = \langle \Delta f + H^2 f + cf, f \rangle_{L^2}.$$

In particular, the mentioned inequality holds for all eigenvalues, i.e.,

*Supported by the SFB 288 of the DFG.

$$\lambda_k^2(D) \leq \mu_k(\Delta + H + c).$$

This inequality was used in order to estimate the first eigenvalue of the Dirac operator defined on special surfaces of Euclidean space (see [1]). On the other hand, in case we know $\lambda_1^2(D)$, the inequality yields a lower bound for the spectrum of the Schrödinger operator $\Delta + H^2 + c$. For example, for any Riemannian metric g on the 2-dimensional sphere S^2 the inequality

$$\lambda_1^2(D) \geq \frac{4\pi}{\text{vol}(S^2, g)}$$

holds (see [2], [6]). Consequently, we obtain

$$\frac{4\pi}{\text{vol}(M^2, g)} \leq \mu_1(\Delta + H^2)$$

for any surface $M^2 \hookrightarrow \mathbb{R}^3$ of genus zero in Euclidean space \mathbb{R}^3 . In this note we expose the described idea and, in particular, we estimate the spectrum of special periodic Schrödinger operators where the potential is given by the curvature κ of a spherical curve.

2 The one-dimensional case

First of all, let us consider the 1-dimensional case, i.e., a curve γ of length L in a two-dimensional space form $X^2(c)$. Let Φ be a Killing spinor of length one on $X^2(c)$:

$$\nabla_T \Phi = \frac{1}{2}c \cdot T \cdot \Phi.$$

The restriction $\varphi = \Phi|_\gamma$ defines a pair of spinors on γ and the spinor field $\psi = f \cdot \varphi$ satisfies the equation:

$$|D\psi|^2 = |\dot{f}|^2 + f^2 \left(\frac{c}{4} + \frac{1}{4}\kappa_g^2 \right),$$

where κ_g is the curvature of the curve γ in $X^2(c)$. Therefore, we obtain

$$\lambda_k^2(D) \leq \mu_k \left(-\frac{d^2}{ds^2} + \frac{c}{4} + \frac{1}{4}\kappa_g^2 \right).$$

Suppose now that the spin structure on γ induced by the spin structure of $X^2(c)$ is non-trivial. Then we have $\lambda_{k+1}^2(D) = \frac{4\pi^2}{L^2}(k + 1/2)^2$ (see [4]) and, in particular, we obtain

$$\frac{4\pi^2}{L^2} \left(k + \frac{1}{2} \right)^2 \leq \mu_{k+1} \left(-\frac{d^2}{ds^2} + \frac{c}{4} + \frac{1}{4}\kappa_g^2 \right).$$

Theorem 1: *Let $\gamma \subset \mathbb{R}^3$ be a plane or spherical curve and denote by $\kappa^2 = c + \kappa_g^2$ the square of its curvature. Suppose that the induced spin structure on γ is non-trivial, i.e., the tangent vector field has an odd rotation number. Then the inequality*

$$\frac{4\pi^2}{L^2} \leq \mu_1 \left(-4 \frac{d^2}{ds^2} + \kappa^2 \right)$$

holds, where μ_1 is the first eigenvalue of the periodic Sturm-Liouville operator on the interval $[0, L]$. Moreover, equality occurs if and only if the curvature is constant.

Remark: No geometric lower bound for the Sturm-Liouville operator $-4 \frac{d^2}{ds^2} + \kappa^2$ with potential defined by the square of the curvature $\kappa(s)$ of a closed curve γ in Euclidean space seems to be known. We conjecture that the estimate given in Theorem 1 holds for any closed curve in \mathbb{R}^3 . Let us compare this inequality with the well-known Fenchel-Milnor inequality

$$2\pi \leq \oint_{\gamma} \kappa.$$

Thus, by the Cauchy-Schwarz inequality we obtain

$$\frac{4\pi^2}{L^2} \leq \frac{1}{L} \oint_{\gamma} \kappa^2.$$

Moreover, using the test function $f \equiv 1$, we have

$$\mu_1 \left(-4 \frac{d^2}{ds^2} + \kappa^2 \right) \leq \frac{1}{L} \oint_{\gamma} \kappa^2.$$

Suppose that γ is a simple curve in \mathbb{R}^3 and denote by ρ the minimal number of generators of the fundamental group $\pi_1(\mathbb{R}^3 \setminus \gamma)$. Then we have

$$2\pi\rho \leq \oint_{\gamma} \kappa.$$

In the spirit of this remark one should be able to prove the stronger inequality

$$\frac{4\pi^2}{L^2} \rho^2 \leq \mu_1 \left(-4 \frac{d^2}{ds^2} + \kappa^2 \right)$$

in case of a simple curve in \mathbb{R}^3 .

Examples: We calculated the eigenvalue μ_1 for some classical curves in \mathbb{R}^3 :

a.) *The lemniscate:* $x = \sin(t)$, $y = \cos(t) \sin(t)$.

$$4\pi^2/L^2 = 1.06193, \quad \mu_1 = 3.7315, \quad \frac{1}{L} \oint_{\gamma} \kappa^2 = 4.36004.$$

b.) *The trefoil:* $x = \sin(3t) \cos(t)$, $y = \sin(3t) \sin(t)$.

$$4\pi^2/L^2 = 0.221, \quad \mu_1 = 5.21, \quad \frac{1}{L} \oint_{\gamma} \kappa^2 = 8.16.$$

c.) *Viviani's curve*: $x = 1 + \cos(t)$, $y = \sin(2t)$, $z = 2 \sin(t)$.

$$4\pi^2/L^2 = 0.169071, \quad \mu_1 = 0.5335, \quad \frac{1}{L} \oint_{\gamma} \kappa^2 = 0.567803.$$

d.) *Torus knot*: $x = (8 + 3 \cos(5t)) \cos(2t)$, $y = (8 + 3 \cos(5t)) \sin(2t)$,
 $z = 5 \sin(5t)$.

$$4\pi^2/L^2 = 0.00146034, \quad \mu_1 = 0.03232, \quad \frac{1}{L} \oint_{\gamma} \kappa^2 = 0.0333803.$$

e.) *The spherical spiral*: $x = \cos(t) \cos(4t)$, $y = \cos(t) \sin(4t)$, $z = \sin(t)$.

$$4\pi^2/L^2 = 0.127036, \quad \mu_1 = 1.744, \quad \frac{1}{L} \oint_{\gamma} \kappa^2 = 4.93147.$$

3 The two-dimensional Schrödinger operator

We generalize this inequality to the case of the two-dimensional periodic Schrödinger operator

$$P_{A,L} = - \left(1 + \frac{A^2}{L^2} \right) \frac{\partial}{\partial t^2} - 4 \frac{\partial^2}{\partial s^2} - \frac{4A}{L} \frac{\partial}{\partial t} \frac{\partial}{\partial s} + \kappa^2(s)$$

defined on $[0, 2\pi] \times [0, L]$:

Theorem 2: *Let $\gamma \subset S^2 \subset \mathbb{R}^3$ be a closed, simple curve of length L bounding a region of area A , and denote by κ its curvature. Then the spectrum of the two-dimensional periodic Schrödinger operator $P_{A,L}$ is bounded by*

$$\frac{4\pi^2}{L^2} \leq \mu_1(P_{A,L}).$$

Equality holds if and only if the curvature of γ is constant.

In general, let us consider a Riemannian manifold (Y^n, g) of dimension n as well as an S^1 -principal fibre bundle $\pi : P \rightarrow Y^n$ over Y^n . Denote by \vec{V} the vertical vector field on P induced by the action of the group S^1 on the total space P , i.e.,

$$\vec{V}(p) = \frac{d}{dt} (p \cdot e^{it})_{t=0}, \quad p \in P.$$

A connection Z in the bundle P defines a decomposition of the tangent bundle $T(P) = T^v(P) \oplus T^h(P)$ into its vertical and horizontal subspace. We introduce a Riemannian metric g^* on the total space P , requiring that

a) $g^*(\vec{V}, \vec{V}) = 1,$

b) $g^*(T^v, T^h) = 0,$

c) the differential $d\pi$ maps $T^h(P)$ isometrically onto $T(Y^n)$.

A closed curve $\gamma : [0, L] \rightarrow Y^n$ of length L defines a torus $H(\gamma) := \pi^{-1}(\gamma) \subset P$ and we want to study the isometry class of this flat torus in P . Let $\alpha = e^{i\Theta} \in S^1$ be the holonomy of the connection Z along the closed curve γ . Consider a horizontal lift $\hat{\gamma} : [0, L] \rightarrow P$ of the curve γ . Then

$$\hat{\gamma}(L) = \hat{\gamma}(0)e^{i\Theta}$$

holds. Consequently, the formula

$$\Phi(t, s) = \hat{\gamma}(s)e^{-i\Theta s/L}e^{it}$$

defines a parametrization $\Phi : [0, 2\pi] \times [0, L] \rightarrow H(\gamma)$ of the torus $H(\gamma)$. Since

$$\frac{\partial \Phi}{\partial t} = \vec{V} \quad , \quad \frac{\partial \Phi}{\partial s} = dR_{e^{it}e^{-i\Theta s/L}}(\dot{\hat{\gamma}}(s)) - \frac{\Theta}{L}\vec{V},$$

we obtain

$$g^* \left(\frac{\partial \Phi}{\partial t}, \frac{\partial \Phi}{\partial t} \right) = 1, \quad g^* \left(\frac{\partial \Phi}{\partial t}, \frac{\partial \Phi}{\partial s} \right) = -\frac{\Theta}{L}, \quad g^* \left(\frac{\partial \Phi}{\partial s}, \frac{\partial \Phi}{\partial s} \right) = 1 + \frac{\Theta^2}{L^2},$$

i.e., the torus $H(\gamma)$ is isometric to the flat torus $(\mathbb{R}^2/\Gamma_o, g^*)$, where Γ_o is the orthogonal lattice $\Gamma_o = 2\pi \cdot \mathbb{Z} \oplus L \cdot \mathbb{Z}$ and the metric g^* has the non-diagonal form

$$g^* = \begin{pmatrix} 1 & -\frac{\Theta}{L} \\ -\frac{\Theta}{L} & 1 + \frac{\Theta^2}{L^2} \end{pmatrix}.$$

Using the transformation

$$x = -\frac{\Theta}{L}s + t \quad , \quad y = s,$$

we see that $H(\gamma)$ is isometric to the flat torus $(\mathbb{R}^2/\Gamma, dx^2 + dy^2)$, where the lattice Γ is generated by the two vectors

$$v_1 = \begin{pmatrix} 2\pi \\ 0 \end{pmatrix} \quad , \quad v_2 = \begin{pmatrix} \Theta \\ L \end{pmatrix}.$$

In case the closed curve $\gamma : [0, L] \rightarrow Y^n$ is the oriented boundary of an oriented compact surface $M^2 \subset Y^n$, we can calculate the holonomy $\alpha = e^{i\Theta}$ along the curve γ . Indeed, let Ω^Z be the curvature form of the connection Z . Ω^Z is a 2-form defined on the manifold Y^n with values in the Lie algebra of the group S^1 , i.e., with values in $i \cdot \mathbb{R}^1$. The parameter Θ is given by the integral

$$\Theta = i \int_{M^2} \Omega^Z.$$

Let us consider the Hopf fibration $\pi : S^3 \rightarrow S^2$, where

$$S^3 = \{(z_1, z_2) \in \mathbb{C}^2 : |z_1|^2 + |z_2|^2 = 1\}$$

is the 3-dimensional sphere of radius 1. The connection Z is given by the formula

$$Z = \frac{1}{2} \{ \bar{z}_1 dz_1 - z_1 d\bar{z}_1 + \bar{z}_2 dz_2 - z_2 d\bar{z}_2 \}$$

and its curvature form ($\omega = z_1/z_2$)

$$\Omega^Z = -\frac{d\omega \wedge d\bar{\omega}}{(1 + |\omega|^2)^2} = -\frac{i}{2} dS^2$$

essentially coincides with one half of the volume form of the unit sphere S^2 of radius 1. However, the differential $d\pi : T^h(S^3) \rightarrow T(S^2)$ multiplies the length of a vector by two, i.e., the Hopf fibration is a Riemannian submersion in the sense described before if we fix the metric of the sphere $S^2(\frac{1}{2}) = \{x \in \mathbb{R}^3 : |x| = \frac{1}{2}\}$ on S^2 . Consequently, in case of a closed simple curve $\gamma \subset S^2$ bounding a region of area A , the Hopf torus $H(\gamma) \subset S^3$ is isometric to the flat torus \mathbb{R}^2/Γ and the lattice Γ is generated by the two vectors

$$v_1 = \begin{pmatrix} 2\pi \\ 0 \end{pmatrix}, \quad v_2 = \begin{pmatrix} A/2 \\ L/2 \end{pmatrix}.$$

The mean curvature H of the torus $H(\gamma) \subset S^3$ coincides with the geodesic curvature κ_g of the curve $\gamma \subset S^2 \subset \mathbb{R}^3$ (see [7], [8]). We apply now the inequality

$$\lambda_1^2(D) \leq \mu_1(\Delta + H^2 + 1)$$

to the Hopf torus $H(\gamma) \subset S^3$. Then we obtain the estimate

$$\lambda_1^2(D) \leq \mu_1(P_{A,L}),$$

where D is the Dirac operator on the flat torus \mathbb{R}^2/Γ with respect to the induced spin structure. All spin structures of a 2-dimensional torus are classified by pairs $(\varepsilon_1, \varepsilon_2)$ of numbers $\varepsilon_i = 0, 1$. If γ is a simple curve in S^2 , the induced spin structure on the Hopf torus $H(\gamma)$ is non-trivial and given by the pairs $(\varepsilon_1, \varepsilon_2) = (0, 1)$. The spectrum of the Dirac operator for all flat tori is well-known (see [4]): The dual lattice Γ^* is generated by

$$v_1^* = \begin{pmatrix} \frac{1}{2\pi} \\ -\frac{A}{2\pi L} \end{pmatrix}, \quad v_2^* = \begin{pmatrix} 0 \\ \frac{2}{L} \end{pmatrix}$$

and the eigenvalues of D^2 are given by

$$\begin{aligned} \lambda^2(k, l) &= 4\pi^2 \left\| kv_1^* + \left(l + \frac{1}{2}\right)v_2^* \right\|^2 = \\ &= k^2 + \frac{4\pi^2}{L^2} \left((2l + 1) - k\frac{A}{2\pi} \right)^2. \end{aligned}$$

We minimize $\lambda^2(k, l)$ on the integral lattice \mathbb{Z}^2 . The isoperimetric inequality $4\pi A - A^2 \leq L^2$ and $A \leq \text{vol}(S^2) = 4\pi$ yield the result that $\lambda^2(k, l)$ attains its minimum at $(k, l) = (0, 1)$, i.e.,

$$\frac{4\pi^2}{L^2} \leq \lambda^2(k, l).$$

Remark 1: We replace the Hopf fibration by the S^1 -principal fibre bundle of Chern class $m \geq 0$. The corresponding total space is the Lens space $L(m, 1)$ and we have the commutative diagram

$$\begin{array}{ccc} S^3 & \longrightarrow & L(m, 1) \\ & \searrow \pi & \swarrow \pi_m \\ & & S^2 \end{array}$$

Let $H_m(\gamma) \subset L(m, 1)$ be the Hopf torus. $H_m(\gamma)$ is isometric to \mathbb{R}^2/Γ_m , where the lattice Γ_m is generated by the vectors

$$v_1 = \begin{pmatrix} 2\pi/m \\ 0 \end{pmatrix}, \quad v_2 = \begin{pmatrix} A/2 \\ L/2 \end{pmatrix}.$$

Moreover, the Lens space $L(m, 1)$ admits a unique spin structure with a Killing spinor (see [3]). Even in case of $m \neq 1$, the induced spin structure on $H_m(\gamma)$ is described by the parameters $(\varepsilon_1, \varepsilon_2) = (0, 1)$. Since the local geometry of $H_m(\gamma)$ in $L(m, 1)$ essentially coincides with the geometry of $H(\gamma)$ in S^3 , we obtain the inequality

$$\frac{4\pi^2}{L^2} = \min_{(k,l)} \left\{ k^2 m^2 + \frac{4\pi^2}{L^2} \left((2l+1) - k \frac{mA}{2\pi} \right)^2 \right\} \leq \mu_1 \left(-4 \frac{d^2}{ds^2} + \kappa^2 \right).$$

Consequently, the investigation of the two-dimensional Schrödinger operator in case of $m \neq 1$ yields the same result for the Sturm-Liouville operator as above.

Remark 2: Suppose now that equality holds for some curve $\gamma \subset S^2$. We consider the corresponding Hopf torus $H(\gamma) \subset S^3$ and then we obtain

$$\lambda_1^2(D) = \mu_1(\Delta + H^2 + 1).$$

Therefore, the mean curvature $H = \kappa$ is constant, i.e., γ is a curve on S^2 of constant curvature κ . Consequently, γ is a circle in a 2-dimensional plane. Denote by r its radius. Then

$$\kappa^2 = \frac{1}{r^2}, \quad L = 2\pi r, \quad A = 2\pi(1 - \sqrt{1 - r^2}),$$

and the inequality

$$\frac{4\pi^2}{L^2} \leq \kappa^2$$

is an equality for all $r \neq 0$.

References

- [1] I. Agricola, Th. Friedrich. Upper bounds for the first eigenvalue of the Dirac operator on surfaces, *Journ. Geom. Phys.* 30 (1999), 1-22.
- [2] Chr. Bär. Lower eigenvalues estimates for Dirac operators, *Math. Ann.* 293 (1992), 39-46.
- [3] Th. Friedrich. Der erste Eigenwert des Dirac-Operators einer kompakten Riemannschen Mannigfaltigkeit nichtnegativer Skalar­krümmung, *Math. Nachr.* 97 (1980), 117-146.
- [4] Th. Friedrich. Zur Abhängigkeit des Dirac-Operators von der Spin-Struktur, *Coll. Math.* vol. XLVIII (1984), 57-62.
- [5] Th. Friedrich. On the spinor representation of surfaces in Euclidean 3-spaces, *Journ. Geom. Phys.* 28 (1998), 143-157.
- [6] J. Lott. Eigenvalue bounds for the Dirac operator, *Pac. Journ. Math.* 125 (1986), 117-128.
- [7] U. Pinkall. Hopf tori in S^3 , *Invent. Math.* 81 (1985), 379-386.
- [8] T.J. Willmore. *Riemannian Geometry*, Clarendon Press Oxford 1996.

THOMAS FRIEDRICH

Humboldt-Universität zu Berlin, Institut für Mathematik, Sitz: Ziegelstraße 13a,
Unter den Linden 6, D-10099 Berlin

e-mail: friedric@mathematik.hu-berlin.de