

# Decomposition of the vertex operator algebra $V_{\sqrt{2}D_l}$

Chongying Dong<sup>1</sup>

Department of Mathematics, University of California  
Santa Cruz, CA 95064

Ching Hung Lam

Institute of Mathematics, University of Tsukuba  
Tsukuba 305-8571, Japan

Hiromichi Yamada

Department of Mathematics, Hitotsubashi University  
Kunitachi, Tokyo 186-8601, Japan

## 1 Introduction

A weight two vector  $v$  of a vertex operator algebra is called a conformal vector with central charge  $c$  if the component operators  $L_v(n)$  for  $n \in \mathbb{Z}$  of  $Y(v, z) = \sum_{n \in \mathbb{Z}} L_v(n)z^{-n-2}$  satisfy the Virasoro algebra relation with central charge  $c$ . In this case, the vertex operator subalgebra  $\text{Vir}(v)$  generated by  $v$  is isomorphic to a Virasoro vertex operator algebra with central charge  $c$  ([FZ], [M]).

Let  $V_{\sqrt{2}R}$  be the vertex operator algebra associated with  $\sqrt{2}$  times an ordinary root lattice  $R$  of type  $A_l$ ,  $D_l$ , or  $E_l$ . In [DLMN], several sets of mutually orthogonal conformal vectors in  $V_{\sqrt{2}R}$  were constructed and studied. It was shown that the Virasoro element of  $V_{\sqrt{2}R}$  can be written as a sum of  $l + 1$  mutually orthogonal conformal vectors  $\omega^i$ ,  $1 \leq i \leq l + 1$ . Since these conformal vectors are mutually orthogonal, the subalgebra  $T$  generated by them is a tensor product  $T = \otimes_{i=1}^{l+1} \text{Vir}(\omega^i)$  and  $\text{Vir}(\omega^i)$  is isomorphic to  $L(c_i, 0)$  with  $c_i$  the central charge of  $\omega^i$ . Here, we denote by  $L(c, h)$  the irreducible highest weight module for the Virasoro algebra with highest weight  $h \in \mathbb{C}$  and central charge  $c$ . Moreover,  $V_{\sqrt{2}R}$  is completely reducible as a  $T$ -module and each irreducible direct summand is in the form  $\otimes_{i=1}^{l+1} L(c_i, h_i)$ . Hence one can study the structure of the vertex operator algebra  $V_{\sqrt{2}R}$  from a point of view that  $V_{\sqrt{2}R}$  is a  $T$ -module. Along this line, the decomposition of  $V_{\sqrt{2}A_2}$  and  $V_{\sqrt{2}A_3}$  into a direct sum of irreducible  $T$ -modules has been determined in [KMY] and [DLY]. See also [DMZ], in which the moonshine module  $V^\natural$  is studied as a module for  $L(\frac{1}{2}, 0)^{\otimes 48}$ .

In this paper we determine the decomposition of  $V_{\sqrt{2}D_l}$  into a sum of irreducible  $T$ -modules for general  $l$ . There are many ways to choose a set of the mutually orthogonal

---

<sup>1</sup>Supported by NSF grant DMS-9700923 and a research grant from the Committee on Research, UC Santa Cruz.

conformal vectors in [DLMN] (see Remark 3.1). In this paper, we shall choose a set of conformal vectors so that  $c_1 = 1/2$ ,  $c_2 = 7/10$ ,  $c_3 = 4/5$ , and  $c_i = 1$  for  $4 \leq i \leq l + 1$ . Such a choice enables us to use the ideas and methods for the decomposition of  $V_{\sqrt{2}A_3}$  in [DLY] with suitable modifications.

We briefly explain how the decomposition of  $V_{\sqrt{2}D_l}$  is obtained. We consider a sublattice  $N$  of  $L = A_1^{\oplus l}$  isomorphic to  $\sqrt{2}D_l$  in a standard way. We show that there exists an automorphism  $\rho$  of the vertex operator algebra  $V_L$  which maps  $V_N$  onto  $V_L^+$  (the fixed point subspace of the involution of  $V_L$  induced from the involution  $\beta \mapsto -\beta$  of  $L$ ). Thus, the decomposition of  $V_N$  as a  $T$ -module is equivalent to the decomposition of  $V_L^+$  as a  $\tilde{T}$ -module, where  $\tilde{T} = \rho(T)$ . The decomposition of  $V_L^+$  as a  $\tilde{T}$ -module is relatively easy and is completely determined in Section 4. For the purpose of completeness we also review the necessary results on the decomposition of a vertex operator algebra associated with a positive definite rank one lattice as a module for  $L(1, 0)$  in Section 5, most of which are taken from [DG] and [KR].

The authors would like to thank Masaaki Kitazume and Masahiko Miyamoto for helpful advice.

## 2 Some automorphisms of $V_L$

We shall use standard notation for the vertex operator algebra  $V_L = M(1) \otimes \mathbb{C}\{L\}$  associated with a positive definite even lattice  $L$  [FLM]. Note that if the square length of any element in  $L$  is a multiple of 4 or if  $L$  is a rank one lattice then the central extension  $\hat{L}$  of  $L$  by the cyclic group of order two is split and  $\mathbb{C}\{L\}$  in the notation of [FLM] is isomorphic to the group algebra  $\mathbb{C}[L]$ .

In [DLY] three automorphisms  $\theta_1$ ,  $\theta_2$ ,  $\sigma$  of order two of the vertex operator algebra  $V_{\mathbb{Z}\alpha}$  associated with a rank one lattice  $\mathbb{Z}\alpha$ , where  $\langle \alpha, \alpha \rangle = 2$ , are considered. They are determined by

$$\begin{aligned} \theta_1 : \alpha(-1) &\mapsto \alpha(-1), & e^\alpha &\mapsto -e^\alpha, & e^{-\alpha} &\mapsto -e^{-\alpha}, \\ \theta_2 : \alpha(-1) &\mapsto -\alpha(-1), & e^\alpha &\mapsto e^{-\alpha}, & e^{-\alpha} &\mapsto e^\alpha, \\ \sigma : \alpha(-1) &\mapsto e^\alpha + e^{-\alpha}, & e^\alpha + e^{-\alpha} &\mapsto \alpha(-1), & e^\alpha - e^{-\alpha} &\mapsto -(e^\alpha - e^{-\alpha}). \end{aligned}$$

The automorphism  $\theta_1$  maps  $u \otimes e^\beta$  to  $(-1)^{\langle \alpha, \beta \rangle / 2} u \otimes e^\beta$  for  $u \in M(1)$  and  $\beta \in \mathbb{Z}\alpha$  and  $\theta_2$  is the automorphism induced from the isometry  $\beta \mapsto -\beta$  of  $\mathbb{Z}\alpha$ . Note also that

$$\sigma\theta_1\sigma = \theta_2, \quad \sigma(\alpha(-1)^2) = \alpha(-1)^2 \quad \text{and} \quad \sigma(e^{\pm\alpha}) = (\alpha(-1) \mp (e^\alpha - e^{-\alpha}))/2.$$

Let  $L$  be a lattice with basis  $\{\alpha_1, \alpha_2, \dots, \alpha_l\}$  such that  $\langle \alpha_i, \alpha_j \rangle = 2\delta_{ij}$ . Then,  $L = \mathbb{Z}\alpha_1 \oplus \dots \oplus \mathbb{Z}\alpha_l$  and each  $\mathbb{Z}\alpha_i$  is isomorphic to a root lattice of type  $A_1$ . Since  $\langle \alpha_i, \alpha_j \rangle = 0$  for  $i \neq j$ , the vertex operator algebra  $V_L$  is a tensor product  $V_L = V_{\mathbb{Z}\alpha_1} \otimes \dots \otimes V_{\mathbb{Z}\alpha_l}$  of  $V_{\mathbb{Z}\alpha_i}$ 's. Using the automorphisms  $\theta_1$ ,  $\theta_2$ , and  $\sigma$  of  $V_{\mathbb{Z}\alpha_i}$  described above, we can define three automorphisms  $\psi_1$ ,  $\psi_2$ , and  $\tau$  of  $V_L$  of order two by

$$\psi_1 = \theta_1 \otimes \dots \otimes \theta_1, \quad \psi_2 = \theta_2 \otimes \dots \otimes \theta_2, \quad \tau = \sigma \otimes \dots \otimes \sigma.$$

Then

$$\psi_1(u \otimes e^\beta) = (-1)^{\langle \alpha_1 + \alpha_2 + \dots + \alpha_i, \beta \rangle / 2} u \otimes e^\beta \quad (2.1)$$

for  $u \in M(1)$  and  $\beta \in L$ ,  $\psi_2$  is the automorphism induced from the isometry  $\beta \mapsto -\beta$  of  $L$ , and  $\tau\psi_1\tau = \psi_2$ .

We shall calculate the images of some elements in  $V_L$  under the automorphism  $\tau$ , which will be used in the next section. If  $i \neq j$ , then in  $V_{\mathbb{Z}\alpha_i} \otimes V_{\mathbb{Z}\alpha_j}$  we have

$$\begin{aligned} \tau(\alpha_i(-1)\alpha_j(-1)) &= \sigma(\alpha_i(-1))\sigma(\alpha_j(-1)) \\ &= (e^{\alpha_i} + e^{-\alpha_i})(e^{\alpha_j} + e^{-\alpha_j}) \\ &= e^{\alpha_i + \alpha_j} + e^{\alpha_i - \alpha_j} + e^{-\alpha_i + \alpha_j} + e^{-\alpha_i - \alpha_j}. \end{aligned}$$

Since  $(\alpha_i \pm \alpha_j)(-1)^2 = \alpha_i(-1)^2 + \alpha_j(-1)^2 \pm 2\alpha_i(-1)\alpha_j(-1)$  in  $V_{\mathbb{Z}\alpha_i} \otimes V_{\mathbb{Z}\alpha_j}$  and  $\sigma(\alpha_i(-1)^2) = \alpha_i(-1)^2$ , it follows that

$$\tau((\alpha_i \pm \alpha_j)(-1)^2) = \alpha_i(-1)^2 + \alpha_j(-1)^2 \pm 2\tau(\alpha_i(-1)\alpha_j(-1)).$$

We also have

$$\begin{aligned} \tau(e^{\alpha_i \pm \alpha_j}) &= \sigma(e^{\alpha_i})\sigma(e^{\pm \alpha_j}) \\ &= \frac{1}{4}(\alpha_i(-1) - (e^{\alpha_i} - e^{-\alpha_i}))(\alpha_j(-1) \mp (e^{\alpha_j} - e^{-\alpha_j})) \end{aligned}$$

in  $V_{\mathbb{Z}\alpha_i} \otimes V_{\mathbb{Z}\alpha_j}$ . Likewise

$$\tau(e^{-(\alpha_i \pm \alpha_j)}) = \frac{1}{4}(\alpha_i(-1) + (e^{\alpha_i} - e^{-\alpha_i}))(\alpha_j(-1) \pm (e^{\alpha_j} - e^{-\alpha_j})),$$

and thus

$$\tau(e^{\alpha_i \pm \alpha_j} + e^{-(\alpha_i \pm \alpha_j)}) = \frac{1}{2}(\alpha_i(-1)\alpha_j(-1) \pm e^{\alpha_i + \alpha_j} \mp e^{\alpha_i - \alpha_j} \mp e^{-\alpha_i + \alpha_j} \pm e^{-\alpha_i - \alpha_j}).$$

For  $\beta = \pm(\alpha_i \pm \alpha_j)/\sqrt{2}$ , set

$$w^\pm(\beta) = \frac{1}{2}\beta(-1)^2 \pm (e^{\sqrt{2}\beta} + e^{-\sqrt{2}\beta}). \quad (2.2)$$

Then,  $w^\pm(\beta) = w^\pm(-\beta)$ . From the above calculation, we have

$$\begin{aligned} \tau(w^-(\alpha_i \pm \alpha_j)/\sqrt{2}) &= w^\pm((\alpha_i - \alpha_j)/\sqrt{2}), \\ \tau(w^+(\alpha_i \pm \alpha_j)/\sqrt{2}) &= w^\pm((\alpha_i + \alpha_j)/\sqrt{2}). \end{aligned} \quad (2.3)$$

### 3 Conformal vectors

Let us consider a sublattice

$$N = \sum_{i,j=1}^l \mathbb{Z}(\alpha_i \pm \alpha_j)$$

of  $L$ , which is isomorphic to the root lattice of type  $\sqrt{2}D_l$ . We choose the following elements as the simple roots of type  $D_l$ :

$$\begin{aligned} \beta_1 &= (\alpha_1 + \alpha_2)/\sqrt{2}, & \beta_2 &= (-\alpha_2 + \alpha_3)/\sqrt{2}, & \beta_3 &= (-\alpha_1 + \alpha_2)/\sqrt{2}, \\ \beta_i &= (-\alpha_i + \alpha_{i+1})/\sqrt{2} & \text{for } & 3 \leq i \leq l-1. \end{aligned}$$

Then

$$\Phi_l^+ = \{(\alpha_i + \alpha_j)/\sqrt{2}, (-\alpha_i + \alpha_j)/\sqrt{2} \mid 1 \leq i < j \leq l\}$$

is the set of positive roots. Using the notation  $w^\pm(\beta)$  defined in (2.2), we set

$$\begin{aligned} s^1 &= \frac{1}{4}w^-(\beta_1), \\ s^2 &= \frac{1}{5}(w^-(\beta_1) + w^-(\beta_2) + w^-(\beta_1 + \beta_2)), \\ s^r &= \frac{1}{2r} \sum_{1 \leq i < j \leq r} \left( w^- \left( (\alpha_i + \alpha_j)/\sqrt{2} \right) + w^- \left( (-\alpha_i + \alpha_j)/\sqrt{2} \right) \right), \quad 3 \leq r \leq l, \\ \omega &= \frac{1}{4(l-1)} \sum_{\beta \in \Phi_l^+} \beta(-1)^2. \end{aligned} \tag{3.1}$$

It was shown by [DLMN] that the elements

$$\omega^1 = s^1, \quad \omega^i = s^i - s^{i-1}, \quad 2 \leq i \leq l, \quad \omega^{l+1} = \omega - s^l \tag{3.2}$$

are mutually orthogonal conformal vectors. Their central charges  $c(\omega^i)$  are as follows:

$$c(\omega^1) = 1/2, \quad c(\omega^2) = 7/10, \quad c(\omega^3) = 4/5, \quad \text{and} \quad c(\omega^i) = 1 \text{ for } 4 \leq i \leq l+1.$$

The subalgebra  $\text{Vir}(\omega^i)$  of the vertex operator algebra  $V_N$  generated by  $\omega^i$  is isomorphic to the Virasoro vertex operator algebra  $L(c(\omega^i), 0)$  which is the irreducible highest weight module for the Virasoro algebra with central charge  $c(\omega^i)$  and highest weight 0. Since  $\omega^1, \omega^2, \dots, \omega^{l+1}$  are mutually orthogonal, the subalgebra  $T$  of  $V_N$  generated by these conformal vectors is a tensor product of  $\text{Vir}(\omega^i)$ 's, namely,

$$\begin{aligned} T &= \text{Vir}(\omega^1) \otimes \cdots \otimes \text{Vir}(\omega^{l+1}) \\ &\cong L(c(\omega^1), 0) \otimes \cdots \otimes L(c(\omega^{l+1}), 0). \end{aligned}$$

As a  $T$ -module  $V_N$  is completely reducible. Our purpose in this paper is to determine all the irreducible direct summands of  $V_N$ .

**Remark 3.1.** In [DLMN] it is shown that corresponding to a chain  $\cdots \subset \Phi'' \subset \Phi' \subset \Phi$  of root systems of type  $A$ ,  $D$ , or  $E$ , one has a decomposition of the Virasoro element of the vertex operator algebra  $V_{\sqrt{2}R}$ , where  $R$  denotes a root lattice of type  $\Phi$ , into a sum of mutually orthogonal conformal vectors (see also [M, Lemma 5.1]). In our notation  $\Phi_l = \Phi_l^+ \cup -\Phi_l^+$  is a root system of type  $D_l$  and  $\omega$  is the Virasoro element of  $V_N$ . The conformal vectors  $\omega^1, \dots, \omega^{l+1}$  in (3.2) correspond to the chain  $\Phi_1 \subset \Phi_2 \subset \cdots \subset \Phi_l$ , where  $\Phi_1 = \{\pm\beta_1\}$ ,  $\Phi_2 = \{\pm\beta_1, \pm\beta_2, \pm(\beta_1 + \beta_2)\}$ , and

$$\Phi_r = \{\pm(\alpha_i + \alpha_j)/\sqrt{2}, \pm(\alpha_i - \alpha_j)/\sqrt{2} \mid 1 \leq i < j \leq r\}, \quad 3 \leq r \leq l.$$

Note that  $\Phi_r$  is a root system of type  $A_r$  for  $1 \leq r \leq 3$  and of type  $D_r$  for  $4 \leq r \leq l$ .

There is another choice of conformal vectors. Namely, if one considers a chain of root systems of type  $A_1 \subset A_2 \subset \cdots \subset A_{l-1} \subset D_l$ , then the central charges of the corresponding mutually orthogonal conformal vectors  $\omega'^1, \dots, \omega'^{l+1}$  are given by

$$c(\omega'^r) = 1 - \frac{6}{(r+2)(r+3)}, \quad 1 \leq r \leq l-1, \quad c(\omega'^l) = \frac{2(l-1)}{l+2}, \quad \text{and} \quad c(\omega'^{l+1}) = 1.$$

The images of the conformal vectors  $\omega^1, \dots, \omega^{l+1}$  under the automorphism  $\tau$  of  $V_L$  can be calculated by (2.3). In fact, we have

**Lemma 3.2.** *Let  $s^1, s^2$ , and  $s^r$ ,  $3 \leq r \leq l$  be defined as in (3.1). Then,*

$$\begin{aligned} \tau(s^1) &= \frac{1}{4}w^+(\beta_3), \\ \tau(s^2) &= \frac{1}{5}(w^+(\beta_3) + w^-(\beta_2) + w^+(\beta_2 + \beta_3)), \\ \tau(s^r) &= \frac{1}{2r} \sum_{1 \leq i < j \leq r} (w^+((\alpha_i - \alpha_j)/\sqrt{2}) + w^-((\alpha_i - \alpha_j)/\sqrt{2})) \\ &= \frac{1}{4r} \sum_{1 \leq i < j \leq r} (\alpha_i - \alpha_j)(-1)^2, \quad 3 \leq r \leq l, \\ \tau(\omega) &= \omega. \end{aligned}$$

Next we shall consider the images of these elements under the automorphism  $\varphi$  of  $V_L$ , where  $\varphi$  is defined by

$$\varphi : u \otimes e^\beta \longmapsto (-1)^{\langle \alpha_2 + \alpha_3, \beta \rangle / 2} u \otimes e^\beta$$

for  $u \in M(1)$  and  $\beta \in L$ . The automorphism  $\varphi$  acts as  $\theta_2$  on  $V_{\mathbb{Z}\alpha_2}$  and  $V_{\mathbb{Z}\alpha_3}$  and acts as the identity on  $V_{\mathbb{Z}\alpha_i}$  for  $i \neq 2, 3$ . Set  $\rho = \varphi\tau$ .

**Remark 3.3.** The automorphism  $\rho$  here is slightly different from the automorphism  $\rho$  of [DLY]. In [DLY]  $\rho$  is defined to be a composite of  $\theta_2 \otimes 1 \otimes 1$  and  $\varphi\tau$ . The symmetry  $\alpha_1 \longleftrightarrow -\alpha_1$  and  $\alpha_i \longleftrightarrow \alpha_i$  for  $2 \leq i \leq l$  induces a  $\mathbb{Z}_2$ -symmetry of the Dynkin diagram of type  $D_l$ , and the restriction of the automorphism  $\theta_2 \otimes 1 \otimes \cdots \otimes 1$  of  $V_L$  to  $V_N$  corresponds to this symmetry. In this paper we do not consider it.

The following lemma is easily obtained from Lemma 3.2 and the definition of  $\varphi$ .

**Lemma 3.4.** *We have*

$$\begin{aligned}\rho(s^1) &= \frac{1}{4}w^-(\beta_3), & \rho(s^2) &= \frac{1}{5}(w^-(\beta_3) + w^-(\beta_2) + w^-(\beta_2 + \beta_3)), \\ \rho(s^r) &= \tau(s^r), \quad 3 \leq r \leq l, & \rho(\omega) &= \omega.\end{aligned}$$

Let  $\tilde{\omega}^i = \rho(\omega^i)$  and set

$$\begin{aligned}\gamma_r &= \alpha_1 + \alpha_2 + \cdots + \alpha_r - r\alpha_{r+1}, \quad 1 \leq r \leq l-1, \\ \gamma_l &= \alpha_1 + \alpha_2 + \cdots + \alpha_l.\end{aligned}\tag{3.3}$$

**Lemma 3.5.** (1) *The vectors  $\tilde{\omega}^1$ ,  $\tilde{\omega}^2$ , and  $\tilde{\omega}^3$  are the mutually orthogonal conformal vectors of  $V_{\mathbb{Z}(\alpha_1 - \alpha_2) + \mathbb{Z}(\alpha_2 - \alpha_3)} \cong V_{\sqrt{2}A_2}$  defined in [DLMN].*

$$(2) \quad \tilde{\omega}^{r+1} = \frac{1}{4r(r+1)}\gamma_r(-1)^2 \quad \text{for } 3 \leq r \leq l-1.$$

$$(3) \quad \tilde{\omega}^{l+1} = \frac{1}{4l}\gamma_l(-1)^2.$$

**Proof** (1) is shown in [DLY, Lemma 3.6]. For  $3 \leq r \leq l-1$  we have

$$\begin{aligned}\rho(s^{r+1}) - \rho(s^r) &= \frac{1}{4(r+1)} \sum_{1 \leq i < j \leq r+1} (\alpha_i - \alpha_j)(-1)^2 - \frac{1}{4r} \sum_{1 \leq i < j \leq r} (\alpha_i - \alpha_j)(-1)^2 \\ &= \frac{1}{4r(r+1)} \left( r \sum_{1 \leq i \leq r} (\alpha_i - \alpha_{r+1})(-1)^2 - \sum_{1 \leq i < j \leq r} (\alpha_i - \alpha_j)(-1)^2 \right) \\ &= \frac{1}{4r(r+1)} (\alpha_1 + \alpha_2 + \cdots + \alpha_r - r\alpha_{r+1})(-1)^2.\end{aligned}$$

We also have

$$\begin{aligned}\omega - \rho(s^l) &= \frac{1}{8(l-1)} \sum_{1 \leq i < j \leq l} ((\alpha_i + \alpha_j)(-1)^2 + (\alpha_i - \alpha_j)(-1)^2) \\ &\quad - \frac{1}{4l} \sum_{1 \leq i < j \leq l} (\alpha_i - \alpha_j)(-1)^2 \\ &= \frac{1}{4(l-1)} \sum_{1 \leq i < j \leq l} (\alpha_i(-1)^2 + \alpha_j(-1)^2) \\ &\quad - \frac{1}{4l} \sum_{1 \leq i < j \leq l} (\alpha_i(-1)^2 + \alpha_j(-1)^2 - 2\alpha_i(-1)\alpha_j(-1)) \\ &= \frac{1}{4l} (\alpha_1 + \alpha_2 + \cdots + \alpha_l)(-1)^2. \quad \square\end{aligned}$$

Note that for  $3 \leq r \leq l$ , the element  $\tilde{\omega}^{r+1}$  is the Virasoro element of the vertex operator algebra  $V_{\mathbb{Z}\gamma_r}$  associated with a rank one lattice  $\mathbb{Z}\gamma_r$ .

Set  $U^\pm = \{v \in U \mid \psi_2(v) = \pm v\}$  for any  $\psi_2$ -invariant subspace  $U$  of  $V_L$ .

**Lemma 3.6.** (1)  $N = \{\beta \in L \mid \langle \alpha_1 + \cdots + \alpha_l, \beta \rangle \equiv 0 \pmod{4}\}$ .

(2)  $V_N = \{v \in V_L \mid \psi_1(v) = v\}$ .

(3)  $\rho(V_N) = V_L^+$ .

**Proof** (1) is clear from the definition of  $N$ . (1) and (2.1) imply (2). Since  $\tau\psi_1 = \psi_2\tau$  and the automorphisms  $\psi_2$  and  $\varphi$  commute, we have  $\rho\psi_1 = \psi_2\rho$ . Hence (3) holds.  $\square$

The last assertion of the above lemma implies that the decomposition of  $V_N$  into a direct sum of irreducible  $T$ -modules is equivalent to that of  $V_L^+$  as a  $\tilde{T}$ -module, where  $\tilde{T} = \rho(T)$  is of the form

$$\begin{aligned} \tilde{T} &= \text{Vir}(\tilde{\omega}^1) \otimes \cdots \otimes \text{Vir}(\tilde{\omega}^{l+1}) \\ &\cong L\left(\frac{1}{2}, 0\right) \otimes L\left(\frac{7}{10}, 0\right) \otimes L\left(\frac{4}{5}, 0\right) \otimes L(1, 0) \otimes \cdots \otimes L(1, 0). \end{aligned}$$

The decomposition of  $V_L^+$  as a  $\tilde{T}$ -module will be accomplished in the next section.

## 4 Decomposition of $V_L^+$ as a $\tilde{T}$ -module

In this section, we shall study the decomposition of  $V_L^+$  into a direct sum of irreducible  $\tilde{T}$ -modules. As in [DLY] we set

$$E = \mathbb{Z}(\alpha_1 - \alpha_2) + \mathbb{Z}(\alpha_2 - \alpha_3) \quad \text{and} \quad D = E + \mathbb{Z}\gamma_3 + \cdots + \mathbb{Z}\gamma_l.$$

The elements  $\alpha_1 - \alpha_2, \alpha_2 - \alpha_3, \gamma_3, \dots, \gamma_l$  form a basis of the lattice  $D$ . Since  $\gamma_r = \sum_{i=1}^r i(\alpha_i - \alpha_{i+1})$ , we can take

$$\{\alpha_1 - \alpha_2, \alpha_2 - \alpha_3, 3(\alpha_3 - \alpha_4), \dots, (l-2)(\alpha_{l-2} - \alpha_{l-1}), \gamma_{l-1}, \gamma_l\}$$

as another basis. The lattices  $E, \mathbb{Z}\gamma_3, \dots, \mathbb{Z}\gamma_l$  are mutually orthogonal, so the vertex operator algebra  $V_D$  associated with the lattice  $D$  is a tensor product  $V_D = V_E \otimes V_{\mathbb{Z}\gamma_3} \otimes \cdots \otimes V_{\mathbb{Z}\gamma_l}$ .

Next, we want to describe the cosets of  $D$  in  $L$ . Set

$$\xi_r = \frac{1}{r(r+1)}\gamma_r, \quad 1 \leq r \leq l-1, \quad \text{and} \quad \xi_l = \frac{1}{l}\gamma_l.$$

Clearly  $\langle \gamma_r, \xi_r \rangle = 2$  and we have

$$-\xi_1 + \xi_2 = \frac{1}{3}(-(\alpha_1 - \alpha_2) + (\alpha_2 - \alpha_3)). \quad (4.1)$$

Moreover,

$$\alpha_r - \alpha_{r+1} = \frac{1}{r}(-\gamma_{r-1} + \gamma_r) = -(r-1)\xi_{r-1} + (r+1)\xi_r,$$

and so

$$-\xi_1 + \xi_2 + \cdots + \xi_l = \alpha_2. \quad (4.2)$$

To simplify the notation, we set  $\eta = -\xi_1 + \xi_2$ .

**Lemma 4.1.**  $|D + \mathbb{Z}\alpha_2 : D|$  is equal to the least common multiple of  $3, 4, \dots, l$ .

**Proof** Since  $\alpha_1 - \alpha_2, \alpha_2 - \alpha_3, \gamma_3, \dots, \gamma_l$  are linearly independent over  $\mathbb{Q}$ , the assertion follows from (4.1) and (4.2).  $\square$

Lemma 4.1 is valid even for  $l = 3$  if we consider a lattice of type  $A_3$  as of type  $D_3$ .

Note that  $D + \mathbb{Z}\alpha_2 = L$  for  $3 \leq l \leq 5$ . Indeed, the coset  $D + \alpha_2$  contains  $\alpha_1, \alpha_2$ , and  $\alpha_3$ . Moreover,  $\alpha_4 \in D + 9\alpha_2$  if  $l = 4$ , and  $\alpha_4 \in D + 21\alpha_2$  and  $\alpha_5 \in D + 36\alpha_2$  if  $l = 5$ . Hence  $n\alpha_2, 0 \leq n \leq d - 1$ , where  $d$  denotes the least common multiple of  $3, 4, \dots, l$ , form a complete system of representatives of the cosets of  $D$  in  $L$  in these three cases. However,  $D + \mathbb{Z}\alpha_2 \neq L$  for  $l \geq 6$ . We shall use the following elements to describe all the cosets of  $D$  in  $L$ . For  $m_3, \dots, m_{l-2}, n \in \mathbb{Z}$  we let

$$\begin{aligned} \lambda &= \lambda(m_3, m_4, \dots, m_{l-2}, n) \\ &= m_3(\alpha_3 - \alpha_4) + m_4(\alpha_4 - \alpha_5) + \dots + m_{l-2}(\alpha_{l-2} - \alpha_{l-1}) + n\alpha_2 \\ &\equiv (m_3 + n)\eta + \sum_{r=3}^{l-3} ((r+1)m_r - rm_{r+1} + n)\xi_r \\ &\quad + ((l-1)m_{l-2} + n)\xi_{l-2} + n\xi_{l-1} + n\xi_l \pmod{D}. \end{aligned} \tag{4.3}$$

The last congruence modulo  $D$  comes from (4.1), (4.2), and the fact that  $\alpha_r - \alpha_{r+1} = -(r-1)\xi_{r-1} + (r+1)\xi_r$ .

**Lemma 4.2.** (1)  $\{\lambda = \lambda(m_3, \dots, m_{l-2}, n) \mid 0 \leq m_r \leq r-1, 0 \leq n \leq l(l-1) - 1\}$  forms a complete system of representatives of the cosets of  $D$  in  $L$ .

(2) Every element in the coset  $D + \lambda$  can be uniquely written in the form

$$\begin{aligned} (\nu + (m_3 + n)\eta) &+ \sum_{i=3}^{l-3} (\mu_i + ((i+1)m_i - im_{i+1} + n)\xi_i) \\ &+ (\mu_{l-2} + ((l-1)m_{l-2} + n)\xi_{l-2}) + (\mu_{l-1} + n\xi_{l-1}) + (\mu_l + n\xi_l) \end{aligned}$$

for  $\nu \in E$  and  $\mu_i \in \mathbb{Z}\gamma_i$ .

**Proof** Let  $M_r = \mathbb{Z}(\alpha_1 - \alpha_2) + \dots + \mathbb{Z}(\alpha_r - \alpha_{r+1}) + \mathbb{Z}\gamma_{r+1} + \dots + \mathbb{Z}\gamma_l$ . Then  $D = M_2 \subset M_3 \subset \dots \subset M_{l-2} \subset L$ . Note that

$$-l(\alpha_1 - \alpha_2) + \sum_{i=2}^{l-2} l(l-i-1)(\alpha_i - \alpha_{i+1}) + \gamma_{l-1} + (l-1)\gamma_l = l(l-1)\alpha_2. \tag{4.4}$$

Since  $\alpha_i \in M_{l-2} + \alpha_2$  for  $1 \leq i \leq l-1$  and  $\alpha_l \in M_{l-2} + (l-1)^2\alpha_2$ ,  $L = M_{l-2} + \mathbb{Z}\alpha_2$  and  $\{n\alpha_2 \mid 0 \leq n \leq l(l-1) - 1\}$  is a complete system of representatives of the cosets of  $M_{l-2}$  in  $L$ . Furthermore,  $\gamma_r = \sum_{i=1}^r i(\alpha_i - \alpha_{i+1})$  implies that  $\{m_r(\alpha_r - \alpha_{r+1}) \mid 0 \leq m_r \leq r-1\}$  is a complete system of representatives of the cosets of  $M_{r-1}$  in  $M_r$ . Hence (1) holds. For



$i = 1, 2$ , the elements  $\alpha_i - \alpha_{i+1}, \gamma_3, \dots, \gamma_l$  are mutually orthogonal, and so each element of  $D + \lambda$  is uniquely decomposed into an orthogonal sum of the form stated in (2).  $\square$

By Lemma 4.2, we have  $V_L = \bigoplus_{\lambda} V_{D+\lambda}$  where the sum runs through the coset representatives of  $D$  in  $L$ , and

$$V_{D+\lambda} = V_{E+(m_3+n)\eta} \otimes \left( \bigotimes_{r=3}^{l-3} V_{\mathbb{Z}\gamma_r + ((r+1)m_r - rm_{r+1} + n)\xi_r} \right) \otimes V_{\mathbb{Z}\gamma_{l-2} + ((l-1)m_{l-2} + n)\xi_{l-2}} \otimes V_{\mathbb{Z}\gamma_{l-1} + n\xi_{l-1}} \otimes V_{\mathbb{Z}\gamma_l + n\xi_l} \quad (4.5)$$

for  $\lambda = \lambda(m_3, \dots, m_{l-2}, n)$ . Since  $\langle E, \eta \rangle = 2\mathbb{Z}$ , the tensor factor of the form  $V_{E+k\eta}$  is an irreducible module for  $V_E$  [D]. Likewise,  $V_{\mathbb{Z}\gamma_i + k\xi_i}$  is an irreducible module for  $V_{\mathbb{Z}\gamma_i}$ . Thus (4.5) holds as a module for  $V_D = V_E \otimes V_{\mathbb{Z}\gamma_3} \otimes \dots \otimes V_{\mathbb{Z}\gamma_l}$ .

**Lemma 4.3.** *For  $\lambda = \lambda(m_3, \dots, m_{l-2}, n)$ ,  $0 \leq m_r \leq r-1$ ,  $0 \leq n \leq l(l-1)-1$ , we have  $D + \lambda = D - \lambda$  if and only if  $m_3, \dots, m_{l-2}$ , and  $n$  satisfy one of the following conditions.*

- (1)  $n = 0$ , and  $m_r = 0$  if  $r$  is odd and  $m_r = 0$  or  $r/2$  if  $r$  is even.
- (2)  $n = l(l-1)/2$  and  $2m_r + l(l-1) \equiv 0 \pmod{r}$ . Such an  $m_r$  is unique if  $r$  is odd and there are exactly two such  $m_r$  if  $r$  is even.

**Proof** Suppose  $D + \lambda = D - \lambda$ , or equivalently  $2\lambda \in D$ . Then  $2n\alpha_2 \in M_{l-2}$  since  $\lambda$  and  $n\alpha_2$  are congruent modulo  $M_{l-2}$ . Now the quotient group  $L/M_{l-2}$  is a cyclic group of order  $l(l-1)$  generated by the coset  $M_{l-2} + \alpha_2$ . Thus  $n = 0$  or  $l(l-1)/2$ .

First, we assume that  $n = 0$ . In this case, since  $\alpha_1 - \alpha_2, \alpha_2 - \alpha_3, 3(\alpha_3 - \alpha_4), \dots, (l-2)(\alpha_{l-2} - \alpha_{l-1}), \gamma_{l-1}, \gamma_l$  form a basis of the lattice  $D$ , we have  $2\lambda \in D$  if and only if  $2m_r$  is divisible by  $r$  for  $3 \leq r \leq l-2$ .

Next, assume that  $n = l(l-1)/2$ . In this case

$$\begin{aligned} 2\lambda &= 2m_3(\alpha_3 - \alpha_4) + \dots + 2m_{l-2}(\alpha_{l-2} - \alpha_{l-1}) + l(l-1)\alpha_2 \\ &= -l(\alpha_1 - \alpha_2) + l(l-3)(\alpha_2 - \alpha_3) \\ &\quad + \sum_{r=3}^{l-2} (2m_r + l(l-r-1))(\alpha_r - \alpha_{r+1}) + \gamma_{l-1} + (l-1)\gamma_l \end{aligned}$$

by (4.4). Hence  $2\lambda \in D$  if and only if  $2m_r + l(l-r-1)$  is divisible by  $r$  for  $3 \leq r \leq l-2$ .  $\square$

The automorphism  $\psi_2$  fixes the conformal vectors  $\tilde{\omega}^1, \dots, \tilde{\omega}^{l+1}$ , and so  $\tilde{T} \subset V_D^+$ . In particular,  $\psi_2$  is a  $\tilde{T}$ -module isomorphism. We have  $\psi_2(V_{D+\lambda}) = V_{D-\lambda}$ , and thus  $V_{D-\lambda}$  is isomorphic to  $V_{D+\lambda}$  as a  $\tilde{T}$ -module. If  $D + \lambda \neq D - \lambda$ , the fixed point subspace  $(V_{D+\lambda} \oplus V_{D-\lambda})^+$  in  $V_{D+\lambda} \oplus V_{D-\lambda}$  is equal to  $\{v + \psi_2(v) \mid v \in V_{D+\lambda}\}$  and it is isomorphic to  $V_{D+\lambda}$ .

If  $D + \lambda = D - \lambda$  for  $\lambda = \lambda(m_3, \dots, m_{l-2}, n)$ , then  $m_3, \dots, m_{l-2}$ , and  $n$  satisfy the conditions in Lemma 4.3. In this case (4.5) is in the following form:

$$V_{D+\lambda} = V_E \otimes V_{\mathbb{Z}\gamma_3 + b_3\gamma_3} \otimes \dots \otimes V_{\mathbb{Z}\gamma_l + b_l\gamma_l},$$

with  $b_i \in \{0, \frac{1}{2}\}$  for  $3 \leq i \leq l-1$  and  $b_l = 0$  or  $b_l \in \{0, \frac{1}{2}\}$  depending on whether  $l$  is odd or even.

For  $\varepsilon = (\varepsilon_0, \varepsilon_3, \dots, \varepsilon_l)$  with  $\varepsilon_i = +$  or  $-$ , set

$$V_{D+\lambda}^\varepsilon = V_E^{\varepsilon_0} \otimes V_{\mathbb{Z}\gamma_3+b_3\gamma_3}^{\varepsilon_3} \otimes \cdots \otimes V_{\mathbb{Z}\gamma_l+b_l\gamma_l}^{\varepsilon_l}. \quad (4.6)$$

Then

$$V_{D+\lambda}^+ = \bigoplus_\varepsilon V_{D+\lambda}^\varepsilon, \quad (4.7)$$

where  $\varepsilon$  runs over all  $\varepsilon = (\varepsilon_0, \varepsilon_3, \dots, \varepsilon_l)$  such that even number of  $\varepsilon_i$ 's are  $-$ .

The decomposition of  $V_E^\pm$  and  $V_{E+\eta}$  as  $\tilde{T}'$ -modules, where  $\tilde{T}' = \text{Vir}(\tilde{\omega}^1) \otimes \text{Vir}(\tilde{\omega}^2) \otimes \text{Vir}(\tilde{\omega}^3)$ , can be found in [KMY, Lemma 4.1] and [DLY, (3.2), (3.3)]. In fact,  $V_E^+$  is a direct sum of four irreducible  $\tilde{T}'$ -modules, which are isomorphic to

$$\begin{aligned} &L\left(\frac{1}{2}, 0\right) \otimes L\left(\frac{7}{10}, 0\right) \otimes L\left(\frac{4}{5}, 0\right), & L\left(\frac{1}{2}, 0\right) \otimes L\left(\frac{7}{10}, \frac{3}{5}\right) \otimes L\left(\frac{4}{5}, \frac{7}{5}\right), \\ &L\left(\frac{1}{2}, \frac{1}{2}\right) \otimes L\left(\frac{7}{10}, \frac{1}{10}\right) \otimes L\left(\frac{4}{5}, \frac{7}{5}\right), & L\left(\frac{1}{2}, \frac{1}{2}\right) \otimes L\left(\frac{7}{10}, \frac{3}{2}\right) \otimes L\left(\frac{4}{5}, 0\right), \end{aligned} \quad (4.8)$$

and  $V_E^-$  is a direct sum of four irreducible  $\tilde{T}'$ -modules, which are isomorphic to

$$\begin{aligned} &L\left(\frac{1}{2}, 0\right) \otimes L\left(\frac{7}{10}, \frac{3}{5}\right) \otimes L\left(\frac{4}{5}, \frac{2}{5}\right), & L\left(\frac{1}{2}, \frac{1}{2}\right) \otimes L\left(\frac{7}{10}, \frac{1}{10}\right) \otimes L\left(\frac{4}{5}, \frac{2}{5}\right), \\ &L\left(\frac{1}{2}, 0\right) \otimes L\left(\frac{7}{10}, 0\right) \otimes L\left(\frac{4}{5}, 3\right), & L\left(\frac{1}{2}, \frac{1}{2}\right) \otimes L\left(\frac{7}{10}, \frac{3}{2}\right) \otimes L\left(\frac{4}{5}, 3\right). \end{aligned} \quad (4.9)$$

Similarly,  $V_{E+\eta}$  is a direct sum of four irreducible  $\tilde{T}'$ -modules, which are isomorphic to

$$\begin{aligned} &L\left(\frac{1}{2}, 0\right) \otimes L\left(\frac{7}{10}, 0\right) \otimes L\left(\frac{4}{5}, \frac{2}{3}\right), & L\left(\frac{1}{2}, 0\right) \otimes L\left(\frac{7}{10}, \frac{3}{5}\right) \otimes L\left(\frac{4}{5}, \frac{1}{15}\right), \\ &L\left(\frac{1}{2}, \frac{1}{2}\right) \otimes L\left(\frac{7}{10}, \frac{1}{10}\right) \otimes L\left(\frac{4}{5}, \frac{1}{15}\right), & L\left(\frac{1}{2}, \frac{1}{2}\right) \otimes L\left(\frac{7}{10}, \frac{3}{2}\right) \otimes L\left(\frac{4}{5}, \frac{2}{3}\right). \end{aligned} \quad (4.10)$$

For  $3 \leq i \leq l$ , the decomposition of  $V_{\mathbb{Z}\gamma_i}^\pm$  and  $V_{\mathbb{Z}\gamma_i+\gamma_i/2}$  as  $\text{Vir}(\tilde{\omega}^{i+1})$ -modules is given in Appendix (see also [DG]).

We divide a complete system of representatives of the cosets of  $D$  in  $L$  into three subsets  $\Lambda_1$ ,  $\Lambda_2$ , and  $-\Lambda_2$  so that  $D + \lambda = D - \lambda$  if and only if  $\lambda \in \Lambda_1$ . Then

$$V_L = \left( \bigoplus_{\lambda \in \Lambda_1} V_{D+\lambda} \right) \oplus \left( \bigoplus_{\lambda \in \Lambda_2} V_{D+\lambda} \right) \oplus \left( \bigoplus_{\lambda \in \Lambda_2} V_{D-\lambda} \right).$$

By the above argument, we conclude that

**Theorem 4.4.** *As  $\tilde{T}$ -modules,*

$$V_L^+ \cong \left( \bigoplus_{\lambda \in \Lambda_1} V_{D+\lambda}^+ \right) \oplus \left( \bigoplus_{\lambda \in \Lambda_2} V_{D+\lambda} \right).$$

Furthermore, the decomposition of  $V_{D+\lambda}^+$ ,  $\lambda \in \Lambda_1$ , and  $V_{D+\lambda}$ ,  $\lambda \in \Lambda_2$ , into a direct sum of irreducible  $\tilde{T}$ -modules is given by (4.5) through (4.10) and (5.1) through (5.7).

By the same method, we also have the decomposition of  $V_L^- \cong (\oplus_{\lambda \in \Lambda_1} V_{D+\lambda}^-) \oplus (\oplus_{\lambda \in \Lambda_2} V_{D+\lambda})$  into a direct sum of irreducible  $\tilde{T}$ -modules.

We illustrate the first two cases, namely  $l = 3$  and  $4$ , and leave the details for the other cases to the interested reader. In case of  $l = 3$ ,  $N = \sqrt{2}A_3$  and the decomposition of  $V_L^+$  as a  $\tilde{T}$ -module given by Theorem 4.4 is identical with the decomposition of  $V_{\sqrt{2}A_3}$  as a  $T$ -module obtained in [DLY, Theorem 3.7].

If  $l = 4$ , then  $D = E + \mathbb{Z}\gamma_3 + \mathbb{Z}\gamma_4$ , where  $\gamma_3 = \alpha_1 + \alpha_2 + \alpha_3 - 3\alpha_4$  and  $\gamma_4 = \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4$ . By Lemma 4.2,  $\{\lambda = \lambda(n) = n\alpha_2 \mid 0 \leq n \leq 11\}$  is a complete system of coset representatives of  $D$  in  $L$  with  $\Lambda_1 = \{0, 6\alpha_2\}$  and  $\Lambda_2 = \{j\alpha_2 \mid 1 \leq j \leq 5\}$ . Thus

$$V_L^+ \cong V_D^+ \oplus V_{D+6\alpha_2}^+ \oplus (\oplus_{j=1}^5 V_{D+j\alpha_2}).$$

Now,  $\alpha_2 = \eta + \xi_3 + \xi_4$  and  $V_{D+j\alpha_2} = V_{E+j\eta} \otimes V_{\mathbb{Z}\gamma_3+j\xi_3} \otimes V_{\mathbb{Z}\gamma_4+j\xi_4}$ . Furthermore,

$$\begin{aligned} V_D^+ &= (V_E^+ \otimes V_{\mathbb{Z}\gamma_3}^+ \otimes V_{\mathbb{Z}\gamma_4}^+) \oplus (V_E^+ \otimes V_{\mathbb{Z}\gamma_3}^- \otimes V_{\mathbb{Z}\gamma_4}^-) \\ &\quad \oplus (V_E^- \otimes V_{\mathbb{Z}\gamma_3}^+ \otimes V_{\mathbb{Z}\gamma_4}^-) \oplus (V_E^- \otimes V_{\mathbb{Z}\gamma_3}^- \otimes V_{\mathbb{Z}\gamma_4}^+), \end{aligned}$$

$$\begin{aligned} V_{D+6\alpha_2}^+ &= (V_E^+ \otimes V_{\mathbb{Z}\gamma_3+\gamma_3/2}^+ \otimes V_{\mathbb{Z}\gamma_4+\gamma_4/2}^+) \oplus (V_E^+ \otimes V_{\mathbb{Z}\gamma_3+\gamma_3/2}^- \otimes V_{\mathbb{Z}\gamma_4+\gamma_4/2}^-) \\ &\quad \oplus (V_E^- \otimes V_{\mathbb{Z}\gamma_3+\gamma_3/2}^+ \otimes V_{\mathbb{Z}\gamma_4+\gamma_4/2}^-) \oplus (V_E^- \otimes V_{\mathbb{Z}\gamma_3+\gamma_3/2}^- \otimes V_{\mathbb{Z}\gamma_4+\gamma_4/2}^+). \end{aligned}$$

Hence the decomposition of  $V_L^+$  into a direct sum of irreducible  $\tilde{T}$ -modules follows from (4.8) through (4.10) and (5.1) through (5.7).

## 5 Appendix

We review the decomposition of a vertex operator algebra associated with a rank one lattice and its modules into a direct sum of irreducible  $L(1, 0)$ -modules. Most materials discussed here can be found in [DG], [KR].

Let  $\mathbb{Z}\gamma$  be a rank one positive definite even lattice with  $\langle \gamma, \gamma \rangle = 2n$  and  $n$  a positive integer. Set  $\xi = \gamma/2n$ . Then  $V_{\mathbb{Z}\gamma+a\xi}$ ,  $0 \leq a \leq 2n - 1$ , are the inequivalent irreducible modules for the vertex operator algebra  $V_{\mathbb{Z}\gamma}$  [D]. We remark that  $\mathbb{Z}\gamma$  is denoted by  $L_{2n}$  in [DG]. The element  $\omega = \gamma(-1)^2/4n$  is the Virasoro element of  $V_{\mathbb{Z}\gamma}$ . The subalgebra  $\text{Vir}(\omega)$  generated by  $\omega$  is isomorphic to the Virasoro vertex operator algebra  $L(1, 0)$  of central charge 1. Note that

$$V_{\mathbb{Z}\gamma+a\xi} = \oplus_{m \in \mathbb{Z}} M(1) \otimes e^{(m+a/2n)\gamma},$$

and each direct summand  $M(1) \otimes e^{(m+a/2n)\gamma}$  is completely reducible as a  $\text{Vir}(\omega)$ -module ([KR, Proposition 3.1]).

If  $W$  is a  $\text{Vir}(\omega)$ -module, the  $q$ -dimension of  $W$  is defined by  $\dim_q W = \sum_n (\dim W_n) q^n$ . Let  $L(1, h)$  be the irreducible highest weight module for  $L(1, 0)$  with highest weight  $h \in \mathbb{C}$ . Its  $q$ -dimension is  $\dim_q L(1, h) = (q^{m^2/4} - q^{(m+2)^2/4})/\phi(q)$  if  $h = m^2/4$  for some integer

$m$  and otherwise  $\dim_q L(1, h) = q^h / \phi(q)$ , where  $\phi(q) = \prod_{j=1}^{\infty} (1 - q^j)$  (see [DG]). On the other hand, the weight of  $e^{(m+a/2n)\gamma}$  is

$$\frac{1}{2} \langle (m + \frac{a}{2n})\gamma, (m + \frac{a}{2n})\gamma \rangle = \frac{1}{4n} (2mn + a)^2.$$

Then  $\dim_q M(1) \otimes e^{(m+a/2n)\gamma} = q^{(2mn+a)^2/4n} / \phi(q)$ . Comparing this with  $\dim_q L(1, h)$ , we know how  $M(1) \otimes e^{(m+a/2n)\gamma}$  decomposes into a direct sum of irreducible modules  $L(1, h)$ 's.

As usual we consider the automorphism  $\theta$  of order two induced from the automorphism  $\beta \mapsto -\beta$  of the underlying lattice. Since  $\theta$  fixes  $\omega$ , it is in fact an automorphism for  $\text{Vir}(\omega)$ -modules. In particular,  $\theta(M(1) \otimes e^{k\gamma}) = M(1) \otimes e^{-k\gamma}$  is isomorphic to  $M(1) \otimes e^{k\gamma}$  as  $\text{Vir}(\omega)$ -modules. If  $U$  is a  $\theta$ -invariant subspace, the eigenspace with eigenvalue  $\pm 1$  is denoted by  $U^{\pm}$ . Then  $U = U^+ \oplus U^-$ .

Note that  $(2mn + a)^2/4n = j^2/4$  for some integer  $j$  if and only if  $n = k^2$  for some nonnegative integer  $k$  and  $a = 0, n$ . We divide the calculation into three cases.

Case 1:  $n = k^2$  for some  $0 < k \in \mathbb{Z}$  and  $a = 0$ . In this case the weight of  $e^{(m+a/2n)\gamma} = e^{m\gamma}$  is  $m^2 k^2$  and

$$M(1) \otimes e^{m\gamma} = \bigoplus_{p=0}^{\infty} L(1, (mk + p)^2).$$

As a special case, we have  $M(1) = \bigoplus_{p=0}^{\infty} L(1, p^2)$  when  $m = 0$ . Moreover,  $\theta(M(1) \otimes e^{m\gamma}) = M(1) \otimes e^{-m\gamma}$  is isomorphic to  $M(1) \otimes e^{m\gamma}$  as a  $\text{Vir}(\omega)$ -module and

$$V_{\mathbb{Z}\gamma} = \left( \bigoplus_{m=1}^{\infty} M(1) \otimes e^{-m\gamma} \right) \oplus M(1) \oplus \left( \bigoplus_{m=1}^{\infty} M(1) \otimes e^{m\gamma} \right),$$

and we have (see [DG])

$$V_{\mathbb{Z}\gamma}^+ \cong \left( \bigoplus_{p=0}^{\infty} L(1, 4p^2) \right) \oplus \left( \bigoplus_{m=1}^{\infty} \bigoplus_{p=0}^{k-1} mL(1, (mk + p)^2) \right), \quad (5.1)$$

$$V_{\mathbb{Z}\gamma}^- \cong \left( \bigoplus_{p=0}^{\infty} L(1, (2p + 1)^2) \right) \oplus \left( \bigoplus_{m=1}^{\infty} \bigoplus_{p=0}^{k-1} mL(1, (mk + p)^2) \right). \quad (5.2)$$

Case 2:  $n = k^2$  for some  $0 < k \in \mathbb{Z}$  and  $a = n$ . The weight of  $e^{(m+a/2n)\gamma} = e^{(m+1/2)\gamma}$  is  $(2m + 1)^2 k^2/4$  and

$$q^{(2m+1)^2 k^2/4} = \sum_{p=0}^{\infty} (q^{((2m+1)k+2p)^2/4} - q^{(2m+1)k+2p+2)^2/4})$$

in the algebra  $\mathbb{C}[[q^{1/4}]]$  of formal power series. Comparing the  $q$ -dimension, we have

$$M(1) \otimes e^{(m+1/2)\gamma} \cong \bigoplus_{p=0}^{\infty} L(1, \frac{1}{4}((2m + 1)k + 2p)^2).$$

Since  $\theta(M(1) \otimes e^{(m+1/2)\gamma}) = M(1) \otimes e^{(-m-1+1/2)\gamma}$  and

$$V_{\mathbb{Z}\gamma+\gamma/2} = \left( \bigoplus_{m=0}^{\infty} M(1) \otimes e^{(m+1/2)\gamma} \right) \oplus \left( \bigoplus_{m=0}^{\infty} M(1) \otimes e^{(-m-1+1/2)\gamma} \right),$$

it follows that

$$\begin{aligned} V_{\mathbb{Z}\gamma+\gamma/2}^+ &\cong V_{\mathbb{Z}\gamma+\gamma/2}^- \cong \bigoplus_{m=0}^{\infty} M(1) \otimes e^{(m+1/2)\gamma} \\ &\cong \bigoplus_{m=0}^{\infty} \bigoplus_{p=0}^{\infty} L(1, \frac{1}{4}((2m+1)k+2p)^2). \end{aligned} \quad (5.3)$$

Case 3:  $(2mn+a)^2/n \neq j^2$  for any integer  $j$ . In this case

$$M(1) \otimes e^{(m+a/2n)\gamma} \cong L(1, \frac{1}{4n}(2mn+a)^2)$$

is an irreducible  $\text{Vir}(\omega)$ -module. Now  $V_{\mathbb{Z}\gamma+a\xi}$  is invariant under  $\theta$  if and only if  $a=0$  or  $n$ . If  $a=0$ , then  $M(1) \otimes e^{m\gamma} \cong L(1, m^2n)$  for  $m \neq 0$  and

$$V_{\mathbb{Z}\gamma}^+ \cong \left( \bigoplus_{p=0}^{\infty} L(1, 4p^2) \right) \oplus \left( \bigoplus_{m=1}^{\infty} L(1, m^2n) \right), \quad (5.4)$$

$$V_{\mathbb{Z}\gamma}^- \cong \left( \bigoplus_{p=0}^{\infty} L(1, (2p+1)^2) \right) \oplus \left( \bigoplus_{m=1}^{\infty} L(1, m^2n) \right) \quad (5.5)$$

(see [DG, Lemma 2.10]). If  $a=n$ , then we have

$$V_{\mathbb{Z}\gamma+\gamma/2}^+ \cong V_{\mathbb{Z}\gamma+\gamma/2}^- \cong \bigoplus_{m=0}^{\infty} L(1, \frac{1}{16}n(4m+1)^2) \quad (5.6)$$

as  $\text{Vir}(\omega)$ -modules. Finally, if  $a \neq 0$ ,  $n$ , then

$$V_{\mathbb{Z}\gamma+a\xi} \cong \bigoplus_{m \in \mathbb{Z}} L(1, \frac{1}{4n}(2mn+a)^2). \quad (5.7)$$

## References

- [D] C. Dong, Vertex algebras associated with even lattices, *J. Algebra* **161** (1993), 245-265.
- [DG] C. Dong and R. L. Griess, Jr., Rank one lattice type vertex operator algebras and their automorphism groups, *J. Algebra* **208** (1998), 262-275.
- [DLMN] C. Dong, H. Li, G. Mason and S. P. Norton, Associative subalgebras of the Griess algebra and related topics, in: *Proc. of the Conference on the Monster and Lie algebras at The Ohio State University, May 1996*, ed. by J. Ferrar and K. Harada, Walter de Gruyter, Berlin-New York, 1998, 27-42.
- [DLY] C. Dong, C. Lam and H. Yamada, Decomposition of the vertex operator algebra  $V_{\sqrt{2}A_3}$ , to appear in *J. Algebra*.
- [DMZ] C. Dong, G. Mason and Y. Zhu, Discrete series of the Virasoro algebra and the Moonshine module, *Proc. Symp. Pure Math., Amer. Math. Soc.* **56** II (1994), 295-316.

- [FLM] I. B. Frenkel, J. Lepowsky and A. Meurman, *Vertex Operator Algebras and the Monster*, Pure and Applied Math., Vol. **134**, Academic Press, 1988.
- [FZ] I. Frenkel and Y. Zhu, Vertex operator algebras associated to representations of affine and Virasoro algebras, *Duke Math. J.* **66** (1992), 123-168.
- [KMY] M. Kitazume, M. Miyamoto and H. Yamada, Ternary codes and vertex operator algebras, to appear in *J. Algebra*.
- [KR] V. Kac and A. Raina, *Bombay Lectures on Highest Weight Representations of Infinite Dimensional Lie Algebras*, World Scientific, 1987.
- [M] M. Miyamoto, Griess algebras and conformal vectors in vertex operator algebras, *J. Algebra* **179** (1996), 523-548.