

# Isomorphism classes of $A$ -hypergeometric systems

Mutsumi Saito

December 27, 1999

## Abstract

For a finite set  $A$  of integral vectors, Gel'fand, Kapranov and Zelevinskii defined a system of differential equations with a parameter vector as a  $D$ -module, which system is called an  $A$ -hypergeometric (or a  $GKZ$  hypergeometric) system. Classifying the parameters according to the  $D$ -isomorphism classes of their corresponding  $A$ -hypergeometric systems is one of the most fundamental problems in the theory. In this paper we give a combinatorial answer for the problem, and illustrate it in two particularly simple cases: the normal case and the monomial curve case.

## 1 Introduction

For a finite set  $A$  of integral vectors, Gel'fand, Kapranov and Zelevinskii defined a system of differential equations with a parameter vector as a  $D$ -module, which system is called an  $A$ -hypergeometric (or a  $GKZ$  hypergeometric) system ([5]). Many authors studied  $D$ -invariants of the  $A$ -hypergeometric systems: In Cohen-Macaulay case, Gel'fand, Kapranov and Zelevinskii determined the characteristic cycles ([6]) and proved the irreducibility of the monodromy representations for nonresonant parameters ([4]); Adolphson proved the rank of an  $A$ -hypergeometric system equals the volume of the convex hull of  $A$  in the semi-nonresonant case ([1]); The author, Sturmfels and Takayama scrutinized the ranks in [13]; Cattani, D'Andrea, and Dickenstein determined rational solutions and algebraic solutions in monomial curve case ([2]), and recently Cattani, Dickenstein, and Sturmfels in [3] considered when an  $A$ -hypergeometric system has a rational solution other than Laurent polynomial solutions.

The purpose of this paper is to classify  $A$ -hypergeometric systems with respect to  $D$ -isomorphisms. This is one of the most fundamental problems in the theory. Under the assumption that the finite set  $A$  lies in a hyperplane off the origin, we shall give a combinatorial answer for this problem, and illustrate it in two particularly simple cases: the normal case and the monomial curve case.

Throughout the paper, we consider the finite set  $A$  fixed. In Section 2, we define a finite set  $E_\tau(\beta)$  for a parameter  $\beta$  and a face  $\tau$  of the cone generated by  $A$ . Then our main theorem (Theorem 2.1) states that two  $A$ -hypergeometric

systems corresponding to parameters  $\beta$  and  $\beta'$  are  $D$ -isomorphic if and only if  $E_\tau(\beta)$  equals  $E_\tau(\beta')$  for all faces  $\tau$ . In Section 2, we prove the only-if-part of the theorem and state some basic properties of the set  $E_\tau(\beta)$ .

Sections 3 and 4 are devoted to the study of the algebra of contiguity operators, which algebra is called the *symmetry algebra*. In Section 3, we summarize some known facts on the symmetry algebra. We introduce the *b-ideals* in Section 4 and prove their elements correspond to contiguity operators. Furthermore we describe each *b-ideal* in terms of the standard pairs of a certain monomial ideal. Using this description, we give the proof of the if-part of our main theorem in the end of Section 4.

In Sections 5 and 6, we illustrate our main theorem in the normal case and the monomial curve case respectively, since the theorem reduces to relatively simple forms in both cases.

The author is very grateful to Professor Nobuki Takayama for his fruitful comments.

## 2 Main theorem

We work over a field  $\mathbf{k}$  of characteristic zero. Let  $A = (a_1, \dots, a_n) = (a_{ij})$  be an integer  $d \times n$ -matrix of rank  $d$ . We assume that all  $a_j$  belong to one hyperplane off the origin in  $\mathbf{Q}^d$ . We denote by  $I_A$  the toric ideal in  $\mathbf{k}[\partial] = \mathbf{k}[\partial_1, \dots, \partial_n]$ , that is

$$I_A = \langle \partial^u - \partial^v \mid Au = Av, u, v \in \mathbf{N}^n \rangle \subset \mathbf{k}[\partial].$$

For a column vector  $\beta = {}^t(\beta_1, \dots, \beta_d) \in \mathbf{k}^d$ , let  $H_A(\beta)$  denote the left ideal of the Weyl algebra

$$D = \mathbf{k}\langle x_1, \dots, x_n, \partial_1, \dots, \partial_n \rangle$$

generated by  $I_A$  and  $\sum_{j=1}^n a_{ij}\theta_j - \beta_i$  ( $i = 1, \dots, d$ ) where  $\theta_j = x_j\partial_j$ . The quotient  $M_A(\beta) = D/H_A(\beta)$  is called the *A-hypergeometric system with parameter  $\beta$* .

We denote the set  $\{a_1, \dots, a_n\}$  by  $A$  as well. Let  $\tau$  be a face of the cone

$$\mathbf{Q}_{\geq 0}A = \left\{ \sum_{j=1}^n c_j a_j \mid c_j \in \mathbf{Q}_{\geq 0} \right\}. \quad (2.1)$$

For a parameter  $\beta \in \mathbf{k}^d$ , we consider the following set:

$$E_\tau(\beta) := \{ \lambda \in \mathbf{k}(A \cap \tau) / \mathbf{Z}(A \cap \tau) \mid \beta - \lambda \in \mathbf{N}A + \mathbf{Z}(A \cap \tau) \}. \quad (2.2)$$

Here  $\mathbf{N} = \{0, 1, 2, \dots\}$  and we agree that  $\mathbf{k}(A \cap \tau) = \mathbf{Z}(A \cap \tau) = \{0\}$  when  $\tau = \{0\}$ .

The following is the main theorem in this paper.

**Theorem 2.1** *The A-hypergeometric systems  $M_A(\beta)$  and  $M_A(\beta')$  are isomorphic as D-modules if and only if  $E_\tau(\beta) = E_\tau(\beta')$  for all faces  $\tau$  of the cone  $\mathbf{Q}_{\geq 0}A$ .*

Before the proof, we recall the formal series solutions  $\phi_v$  defined in [13]. For  $v \in \mathbf{k}^n$ , its *negative support*  $\text{nsupp}(v)$  is the set of indices  $i$  with  $v_i \in \mathbf{Z}_{<0}$ . When  $\text{nsupp}(v)$  is minimal with respect to inclusions among  $\text{nsupp}(v+u)$  with  $u \in \mathbf{Z}^n$  and  $Au = 0$ ,  $v$  is said to have *minimal negative support*. For  $v$  with minimal negative support, we define a formal series

$$\phi_v = \sum_{u \in N_v} \frac{[v]_{u_-}}{[v+u]_{u_+}} x^{v+u}. \quad (2.3)$$

Here

$$N_v = \{ u \in \mathbf{Z}^n \mid Au = 0, \text{nsupp}(v) = \text{nsupp}(v+u) \},$$

and  $u_+, u_- \in \mathbf{N}^n$  satisfy  $u = u_+ - u_-$  with disjoint supports, and  $[v]_w = \prod_{j=1}^n v_j(v_j-1)\cdots(v_j-w_j+1)$  for  $w \in \mathbf{N}^n$ . Proposition 3.4.13 of [13] states that the series  $\phi_v$  is a formal solution of  $M_A(Av)$ .

**Proof.** Here we prove the only-if-part of the theorem. The proof of the if-part will be given in the end of Section 4.

We suppose that  $\lambda \in E_\tau(\beta) \setminus E_\tau(\beta')$ , and we shall prove  $M_A(\beta)$  and  $M_A(\beta')$  are not isomorphic.

Represent  $\lambda$  as  $\sum_{a_j \in \tau} l_j a_j$ . Consider the direct product

$$R_{\tau,\lambda} := \prod_{u \in \mathbf{Z}^n, u_j \in \mathbf{N} (a_j \notin \tau)} \mathbf{k}x^{l+u}.$$

Here we put  $l_j = 0$  for  $a_j \notin \tau$ . Note that  $R_{\tau,\lambda}$  has the natural  $D$ -module structure. There exists  $u \in \mathbf{Z}^n$  with  $u_j \in \mathbf{N}$  ( $a_j \notin \tau$ ) such that  $\beta = A(l+u)$  and  $l+u$  has minimal negative support. Then the series  $\phi_{l+u} \in R_{\tau,\lambda}$  is a formal solution of  $M_A(\beta)$ , and hence  $\text{Hom}_D(M_A(\beta), R_{\tau,\lambda}) \neq 0$ . On the other hand,  $\text{Hom}_D(M_A(\beta'), R_{\tau,\lambda}) = 0$  since  $A(l+u) \neq \beta'$  for any  $u \in \mathbf{Z}^n$  with  $u_j \in \mathbf{N}$  ( $a_j \notin \tau$ ). Therefore  $M_A(\beta)$  and  $M_A(\beta')$  are not isomorphic.  $\square$

In the remainder of this section, we collect some properties of the set  $E_\tau(\beta)$ . We call a face of  $\mathbf{Q}_{\geq 0}A$  of dimension  $d-1$ , a facet. Recall that for a facet  $\sigma$  the linear form  $F_\sigma$  satisfying the following conditions is unique and called the *primitive integral support function*:

1.  $F_\sigma(\mathbf{Z}A) = \mathbf{Z}$ ,
2.  $F_\sigma(a_j) \geq 0$  for all  $j = 1, \dots, n$ ,
3.  $F_\sigma(a_j) = 0$  for all  $a_j \in \sigma$ .

**Proposition 2.2** 1. Each  $E_{\mathbf{Q}_{\geq 0}A}(\beta)$  consists of one element. The equality  $E_{\mathbf{Q}_{\geq 0}A}(\beta) = E_{\mathbf{Q}_{\geq 0}A}(\beta')$  means  $\beta - \beta' \in \mathbf{Z}A$ .

2.  $E_{\{0\}}(\beta) = \{0\}$  or  $\emptyset$ .  $E_{\{0\}}(\beta) = \{0\}$  if and only if  $\beta \in \mathbf{N}A$ .
3. For a facet  $\sigma$ ,  $E_\sigma(\beta) \neq \emptyset$  if and only if  $F_\sigma(\beta) \in F_\sigma(\mathbf{N}A)$ .

4. For faces  $\tau \subset \sigma$ , there exists a natural map from  $E_\tau(\beta)$  to  $E_\sigma(\beta)$ . In particular, if  $E_\tau(\beta) \neq \emptyset$ , then  $E_\sigma(\beta) \neq \emptyset$ .
5. For any  $\chi \in \mathbf{NA}$ , there exists a natural inclusion from  $E_\tau(\beta)$  to  $E_\tau(\beta + \chi)$ .

**Proof.** All statements follow directly from the definition of  $E_\tau(\beta)$ .  $\square$

**Proposition 2.3** 1.

$$|E_\tau(\beta)| \leq [(\mathbf{Q}(A \cap \tau)) \cap \mathbf{ZA} : \mathbf{Z}(A \cap \tau)]. \quad (2.4)$$

2. Assume  $(\mathbf{Q}(A \cap \tau)) \cap \mathbf{ZA} = \mathbf{Z}(A \cap \tau)$ . If  $\beta - \beta' \in \mathbf{ZA}$ , and if neither  $E_\tau(\beta)$  nor  $E_\tau(\beta')$  is empty, then  $E_\tau(\beta) = E_\tau(\beta')$ .

**Proof.**

1. Let  $\lambda, \lambda' \in E_\tau(\beta)$ . Then  $\lambda - \lambda' \in (\mathbf{k}(A \cap \tau)) \cap \mathbf{ZA}$ . By Cramér's formula,  $(\mathbf{k}(A \cap \tau)) \cap \mathbf{ZA} = (\mathbf{Q}(A \cap \tau)) \cap \mathbf{ZA}$ .
2. Let  $E_\tau(\beta) = \{\lambda\}$ ,  $E_\tau(\beta') = \{\lambda'\}$ . Since  $\beta - \beta' \in \mathbf{ZA}$ , there exist  $\chi, \chi' \in \mathbf{NA}$  such that  $\beta + \chi = \beta' + \chi'$ . Then  $\{\lambda\} = E_\tau(\beta + \chi) = E_\tau(\beta' + \chi') = \{\lambda'\}$  by Proposition 2.2 (5).

$\square$

**Example 2.4** Let

$$A = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 2 \\ 0 & 1 & 1 & 0 \end{pmatrix}.$$

There are four facets:

$$\sigma_{12} : = \mathbf{Q}_{\geq 0}a_1 + \mathbf{Q}_{\geq 0}a_2, \quad (2.5)$$

$$\sigma_{23} : = \mathbf{Q}_{\geq 0}a_2 + \mathbf{Q}_{\geq 0}a_3, \quad (2.6)$$

$$\sigma_{34} : = \mathbf{Q}_{\geq 0}a_3 + \mathbf{Q}_{\geq 0}a_4, \quad (2.7)$$

$$\sigma_{14} : = \mathbf{Q}_{\geq 0}a_1 + \mathbf{Q}_{\geq 0}a_4, \quad (2.8)$$

and four one-dimensional faces:  $\mathbf{Q}_{\geq 0}a_1, \dots, \mathbf{Q}_{\geq 0}a_4$ . For all faces  $\tau$  but  $\sigma_{14}$ , the indices  $[(\mathbf{Q}(A \cap \tau)) \cap \mathbf{ZA} : \mathbf{Z}(A \cap \tau)]$  are one. Hence for  $\beta \in \mathbf{NA}$ ,  $E_\tau(\beta) = \{0\}$  for all faces  $\tau \neq \sigma_{14}$ . The quotient  $(\mathbf{Q}(A \cap \sigma_{14})) \cap \mathbf{ZA} / \mathbf{Z}(A \cap \sigma_{14})$  has two elements and can be represented by 0 and  ${}^t(1, 1, 0)$ . Since  $a_2 - {}^t(1, 1, 0) = a_3 - a_4$ , and  $a_3 - {}^t(1, 1, 0) = a_2 - a_1$ , we obtain  $E_{\sigma_{14}}(a_2) = E_{\sigma_{14}}(a_3) = \{0, {}^t(1, 1, 0)\}$ . Proposition 2.2 (5) implies that for  $\beta \in \mathbf{NA}$ ,  $E_{\sigma_{14}}(\beta) = \{0\}$  if and only if  $\beta \in \mathbf{Na}_1 + \mathbf{Na}_4$ , otherwise  $E_{\sigma_{14}}(\beta) = \{0, {}^t(1, 1, 0)\}$ . Therefore  $\mathbf{NA}$  splits into two isomorphism classes in this case.

Recall that a parameter  $\beta$  is said to be *nonresonant* (respectively *semi-nonresonant*) if  $\beta \notin \mathbf{Z}A + \mathbf{k}(A \cap \sigma)$  (respectively  $\beta \notin (\mathbf{Z}A \cap \mathbf{Q}_{\geq 0}A) + \mathbf{k}(A \cap \sigma)$ ) for any facet  $\sigma$ , or equivalently, if  $F_\sigma(\beta) \notin \mathbf{Z}$  (respectively  $F_\sigma(\beta) \notin \mathbf{N}$ ) for any facet  $\sigma$ . Hence the nonresonance implies the semi-nonresonance.

**Proposition 2.5** *If  $\beta$  is semi-nonresonant, then  $E_\tau(\beta) = \emptyset$  for all proper faces  $\tau$  of  $\mathbf{Q}_{\geq 0}A$ .*

**Proof.** The semi-nonresonance clearly implies  $E_\sigma(\beta) = \emptyset$  for all facets  $\sigma$ . Proposition 2.2 (4) finishes the proof.  $\square$

**Corollary 2.6** *Let  $\beta$  and  $\beta'$  be semi-nonresonant. Then  $M_A(\beta)$  and  $M_A(\beta')$  are isomorphic if and only if  $\beta - \beta' \in \mathbf{Z}A$ .*

**Proposition 2.7** *If a parameter  $\beta$  satisfies*

$$E_\tau(\beta) = \emptyset \quad \text{for all proper faces } \tau, \quad (2.9)$$

then

1. for any  $\chi \in \mathbf{N}A$ ,  $M_A(\beta - \chi)$  is isomorphic to  $M_A(\beta)$ .
2. Recall that all elements of  $A$  lie on one hyperplane  $H$  off the origin. We normalize the volume of a polytope on  $H$  so that a simplex whose vertices affinely span the lattice  $H \cap \mathbf{Z}A$  has volume one. Then the rank of  $M_A(\beta)$ , i.e. the rank of the solution sheaf of  $M_A(\beta)$ , equals the volume of the convex hull of  $A$ .

**Proof.** (1) By Proposition 2.2 (5),  $E_\tau(\beta - \chi) = \emptyset$  for all proper faces  $\tau$ . Hence by Proposition 2.2 (1), we deduce the statement from Theorem 2.1.

The proof of (2) is the same as that of Theorem 4.5.2 of [13] (p. 185).  $\square$

### 3 Symmetry algebra

We consider the algebra of contiguity operators, which algebra is called the symmetry algebra. It controls isomorphisms among  $A$ -hypergeometric systems with different parameters. We have investigated the symmetry algebra of normal  $A$ -hypergeometric systems in [11]. The proofs of some results in [11] remain valid without the normality condition. In this section, we summarize such results.

Let

$$\tilde{S} := \{ P \in D \mid I_A P \subset D I_A \}.$$

Then  $\tilde{S}$  is an associative algebra and  $\tilde{S} \cap D I_A$  is its two-sided ideal. We call  $S := \tilde{S} / \tilde{S} \cap D I_A$  the *symmetry algebra* of  $A$ -hypergeometric systems. The symmetry algebra  $S$  is nothing but the algebra  $\text{End}_D(D/D I_A)$ . We remark that  $D/D I_A$  can be considered as the system of differential equations for the generating functions of  $A$ -hypergeometric functions.

In what follows, we denote simply by  $P$ , the element of  $D/DI_A$  represented by  $P \in D$ . For  $\chi \in \mathbf{NA}$ , all  $\partial^u$  with  $Au = \chi$  represent the same element in  $D/DI_A$ . Hence we sometimes denote it by  $\partial^\chi$ .

**Proposition 3.1** 1.  $\partial_1, \dots, \partial_n \in S$ .

2.  $\sum_{j=1}^n a_{ij}\theta_j \in S$  for all  $i = 1, \dots, d$ .

3. The morphism from the polynomial ring  $\mathbf{k}[s] = \mathbf{k}[s_1, \dots, s_d]$  to  $S$  mapping  $s_i$  to  $\sum_{j=1}^n a_{ij}\theta_j$  ( $i = 1, \dots, d$ ) is injective.

**Proof.** See Lemma 1.1 in [11] for (1) and (2), and Corollary 1.3 in [11] for (3).  $\square$

We consider that  $\mathbf{ZA}$  is the character group of the algebraic torus  $T = \{(t_1, \dots, t_d) \mid t_1, \dots, t_d \in \mathbf{k}^\times\}$ . Let  $N$  be the dual group of  $\mathbf{ZA}$ , and  $s_1, \dots, s_d$  the basis of  $\mathbf{k} \otimes_{\mathbf{Z}} N$  dual to the standard basis of  $\mathbf{k}^d = \mathbf{k} \otimes_{\mathbf{Z}} \mathbf{ZA}$ . Under the identification of  $\mathbf{k} \otimes_{\mathbf{Z}} N$  with the Lie algebra of  $T$  ([8]), each  $s_i$  equals  $t_i \frac{\partial}{\partial t_i}$ . The morphism in Proposition 3.1 (3) is induced from the differential of the injective morphism:

$$T \ni t \mapsto (t^{a_1}, \dots, t^{a_n}) \in (\mathbf{k}^\times)^n. \quad (3.10)$$

We thus consider  $\mathbf{k}[s]$  as a subspace of  $S$  and, accordingly, as a subspace of  $D/DI_A$ . For each  $\chi \in \mathbf{ZA}$ , we define the weight space  $S_\chi$  with weight  $\chi$  by

$$S_\chi := \{P \in S \mid [s, P] = \chi(s)P \quad (\forall s \in N)\}.$$

Here the bracket  $[P, Q]$  means  $PQ - QP$ .

**Remark 3.2** Note that the multiplication by  $P \in S_\chi$  from the right defines a  $D$ -homomorphism from  $M_A(\beta + \chi)$  to  $M_A(\beta)$ . Hence  $P(\psi_\beta)$  is a solution of  $M_A(\beta + \chi)$  for a solution  $\psi_\beta$  of  $M_A(\beta)$ . In this sense, the operator  $P$  is a contiguity operator shifting parameters by  $\chi$ .

**Theorem 3.3** 1. The symmetry algebra  $S$  has no zero-divisors.

2. The symmetry algebra  $S$  has the following weight space decomposition:

$$S = \bigoplus_{\chi \in \mathbf{ZA}} S_\chi. \quad (3.11)$$

3. The weight space  $S_0$  equals the polynomial ring  $\mathbf{k}[s]$ .

4. For each  $\chi \in \mathbf{NA}$ , the weight space  $S_{-\chi}$  equals  $\mathbf{k}[s]\partial^\chi$ .

**Proof.** See Lemma 1.4, and Propositions 2.3, 2.4, 2.9 in [11].  $\square$

The following proposition will be used in the next section.

**Proposition 3.4** (Proposition 2.6 in [11]) *The natural morphism*

$$D/DI_A \longrightarrow \mathbf{k}\langle x, \partial^\pm \rangle / \mathbf{k}\langle x, \partial^\pm \rangle I_A$$

is injective where  $\mathbf{k}\langle x, \partial^\pm \rangle$  is the algebra generated by  $D$  and  $\partial_1^{-1}, \dots, \partial_n^{-1}$  with relations  $[x_i, \partial_j^{-1}] = \delta_{ij} \partial_j^{-2}$  ( $i, j = 1, \dots, n$ ).

## 4 $b$ -Ideals

We have seen in Theorem 3.3 that the symmetry algebra  $S$  has a weight decomposition with respect to  $\mathbf{Z}A$ , and that each  $S_\chi$  for  $-\chi \in \mathbf{N}A$  is the free  $\mathbf{k}[s]$ -module of rank one with basis  $\partial^{-\chi}$ . Next we wish to compute the weight space  $S_\chi$  for arbitrary  $\chi$ . Suppose that  $E \in S_\chi$  and  $\chi = \chi_+ - \chi_-$  with  $\chi_+, \chi_- \in \mathbf{N}A$ . Then the operator  $E\partial^{\chi_+}$  belongs to  $S_{-\chi_-}$ . Hence by Theorem 3.3 (4), there exists a polynomial  $b \in \mathbf{k}[s]$  such that  $E\partial^{\chi_+} = b\partial^{\chi_-}$ . Such polynomials  $b$  varying  $E \in S_\chi$  form an ideal of  $\mathbf{k}[s]$ . We shall define the  $b$ -ideal  $B_\chi$  below to be such an ideal.

Fix any  $\chi \in \mathbf{Z}A$ , and define an ideal  $I_\chi$  of  $\mathbf{k}[\partial]$  by

$$I_\chi := I_A + M_\chi \tag{4.12}$$

where

$$M_\chi := \langle \partial^u \mid Au \in \chi + \mathbf{N}A \rangle. \tag{4.13}$$

Define the ideal  $B_\chi$  of  $b$ -polynomials by

$$B_\chi := \mathbf{k}[s] \cap DI_\chi. \tag{4.14}$$

**Proposition 4.1** *Let  $\chi = \chi_+ - \chi_-$  with  $\chi_+, \chi_- \in \mathbf{N}A$ . For  $b \in B_\chi$ , there exists a unique operator  $E \in S_\chi$  such that  $b\partial^{\chi_-} = E\partial^{\chi_+}$ . The operator  $E$  is independent of the expression  $\chi = \chi_+ - \chi_-$ .*

*Moreover any operator in  $S_\chi$  can be obtained in this way.*

**Proof.** Since  $b\partial^{\chi_-} \in DI_\chi \partial^{\chi_-} \subset DI_A + D\partial^{\chi_+}$ , there exists an operator  $E \in D$  such that  $b\partial^{\chi_-} = E\partial^{\chi_+}$ . The uniqueness, the independence, and  $E \in S_\chi$  follow from the equality  $E = b\partial^\chi$  in  $\mathbf{k}\langle x, \partial^\pm \rangle$  and Proposition 3.4.

Let  $E \in S_\chi$  and  $\chi = \chi_+ - \chi_-$  with  $\chi_+, \chi_- \in \mathbf{N}A$ . Then  $E\partial^{\chi_+} \in S_{-\chi_-}$ . By Theorem 3.3 (4), there exists a polynomial  $b \in \mathbf{k}[s]$  such that  $E\partial^{\chi_+} = b\partial^{\chi_-}$ . Then  $b \in I_\chi$  and thus  $b \in B_\chi$ .  $\square$

We have the following algorithm of obtaining the operator  $E \in S_\chi$  corresponding to  $b \in B_\chi$ , which generalizes Algorithm 3.4 in [12].

**Algorithm 4.2** *Let  $\chi = Au - Av$  and  $u, v \in \mathbf{N}^n$ .*

*Input: a polynomial  $b \in B_\chi$ .*

*Output: an operator  $E \in S_\chi$  with  $E\partial^u = b\partial^v$ .*

1. For  $i = 1, \dots, n$ , compute a Gröbner basis  $\mathcal{G}_i$  of  $I_A$  with respect to any reverse lexicographic term order with lowest variable  $\partial_i$ .
2. Expand  $b(\sum_j a_{1j}\theta_j, \dots, \sum_j a_{nj}\theta_j)\partial^v$  in  $\mathbf{Q}\langle x, \partial \rangle$  into a  $\mathbf{Q}$ -linear combination of monomials  $x^l\partial^m$ .
3.  $i := 1$ ,  $E :=$  the output of Step 2.  
While  $i \leq n$ , do
  - (a) Reduce  $E$  modulo  $\mathcal{G}_i$  in  $\mathbf{Q}\langle x, \partial \rangle$ .
  - (b) The output of Step 3-(a) has  $\partial_i^{u_i}$  as a right factor. Divide it by  $\partial_i^{u_i}$ .
  - (c)  $i := i + 1$ ,  $E :=$  the output of Step 3-(b).

The proof of the correctness is completely analogous to that of Algorithm 3.4 in [12].

We thus reduce the study of  $S_\chi$  to that of  $B_\chi$ , and for the study of  $B_\chi = \mathbf{k}[s] \cap DI_\chi$ , we study  $\mathbf{k}[\theta] \cap DI_\chi$  first. Since  $M_\chi$  is the largest monomial ideal in  $I_\chi$ , we have by Lemma 4.4.4 in [13],

**Proposition 4.3**

$$\mathbf{k}[\theta] \cap DI_\chi = \widetilde{M}_\chi \quad (4.15)$$

where  $\widetilde{M}_\chi$  is the distraction of  $M_\chi$ , i.e.,  $\widetilde{M}_\chi = \mathbf{k}[\theta] \cap DM_\chi$ .

For the study of  $\widetilde{M}_\chi$ , we recall the standard pairs of a monomial ideal. Let  $M$  be a monomial ideal of  $\mathbf{k}[\partial]$ . Then a pair  $(u, \tau)$  with  $u \in \mathbf{N}^n$  and  $\tau \subset \{1, \dots, n\}$  is called a *standard pair* of  $M$  if it satisfies the following conditions:

1.  $u_j = 0$  for all  $j \in \tau$ . (We abbreviate this to  $u \in \mathbf{N}^{\tau^c}$ , where  $c$  stands for taking the complement.)
2. There exists no  $v \in \mathbf{N}^\tau$  such that  $\partial^{u+v} \in M$ .
3. For each  $j \notin \tau$ , there exists  $v \in \mathbf{N}^{\tau \cup \{j\}}$  such that  $\partial^{u+v} \in M$ .

For an algorithm of obtaining the set of standard pairs, see [7]. Let  $\mathcal{S}(M_\chi)$  denote the set of standard pairs of  $M_\chi$ . By Corollary 3.2.3 in [13], the distraction  $\widetilde{M}_\chi$  is described as follows:

$$\widetilde{M}_\chi = \bigcap_{(u, \tau) \in \mathcal{S}(M_\chi)} \langle \theta_i - u_i \mid i \notin \tau \rangle. \quad (4.16)$$



**Lemma 4.4** *Let  $(u, \tau)$  be a standard pair of  $M_\chi$ . Then  $A\mathbf{Q}_{\geq 0}^\tau := \sum_{j \in \tau} \mathbf{Q}_{\geq 0} a_j$  is a proper face of  $\mathbf{Q}_{\geq 0}A$ , and moreover  $\tau = \{i \mid a_i \in A\mathbf{Q}_{\geq 0}^\tau\}$ .*

**Proof.** Suppose that  $A\mathbf{Q}_{\geq 0}^\tau$  is not contained in any facet of  $\mathbf{Q}_{\geq 0}A$ . Then there exists  $\gamma \in A\mathbf{N}^\tau := \sum_{j \in \tau} \mathbf{N}a_j$  such that  $F_\sigma(\gamma) > 0$  for all facets  $\sigma$ . Then  $F_\sigma(Au + m\gamma) \gg 0$  for  $m \gg 0$  and all facets  $\sigma$ . By Lemma 1 in the appendix of [14],  $Au + m\gamma \in \chi + \mathbf{N}A$  for  $m \gg 0$ . This contradicts  $(u, \tau)$  is a standard pair of  $M_\chi$ .

Next we claim  $(A\mathbf{Q}_{\geq 0}^{\tau^c}) \cap (A\mathbf{Q}^\tau) = \{0\}$ , which implies the lemma. Suppose  $(A\mathbf{Q}_{\geq 0}^{\tau^c}) \cap (A\mathbf{Q}^\tau) \neq \{0\}$ . Let  $v \in \mathbf{N}^{\tau^c}$  be a nonzero element satisfying  $Av \in A\mathbf{Z}^\tau$ . Then there exists  $w \in \mathbf{N}^\tau$  such that  $Aw \in Av + A\mathbf{N}^\tau$ . Since  $A(u + mw) \notin M_\chi$  for any  $m \in \mathbf{N}$ ,  $(Au + A\mathbf{N}^{\tau \cup \tau'}) \cap M_\chi = \emptyset$  for  $\tau' = \{i \mid v_i \neq 0\}$ . This contradicts  $(u, \tau)$  is a standard pair of  $M_\chi$  again.  $\square$

Thanks to Lemma 4.4, we regard the set  $\tau$  of a standard pair  $(u, \tau)$  as the proper face  $A\mathbf{Q}^\tau$  of  $\mathbf{Q}_{\geq 0}A$ .

For an ideal  $I$  of  $\mathbf{k}[s]$ , we denote by  $V(I)$  the zero set of  $I$ . Proposition 4.3 and the equation (4.16) give the following prime decomposition of  $B_\chi$  and irreducible decomposition of the zero set  $V(B_\chi)$ .

**Theorem 4.5** 1.

$$B_\chi = \bigcap_{(u, \tau) \in \mathcal{S}(M_\chi)} \langle F_\sigma - F_\sigma(Au) \mid \sigma : \text{facet} \supset \tau \rangle. \quad (4.17)$$

2.

$$V(B_\chi) = \bigcup_{(u, \tau) \in \mathcal{S}(M_\chi)} (Au + \mathbf{k}(A \cap \tau)). \quad (4.18)$$

**Proof.** From (4.16), we only need to show

$$\mathbf{k}[s] \cap \langle \theta_i - u_i \mid i \notin \tau \rangle = \langle F_\sigma - F_\sigma(Au) \mid \sigma \supset \tau \rangle. \quad (4.19)$$

First we have

$$V(\mathbf{k}[s] \cap \langle \theta_i - u_i \mid i \notin \tau \rangle) = Au + \mathbf{k}(A \cap \tau) = V(\langle F_\sigma - F_\sigma(Au) \mid \sigma \supset \tau \rangle). \quad (4.20)$$

Hence

$$\begin{aligned} \mathbf{k}[s] \cap \langle \theta_i - u_i \mid i \notin \tau \rangle &\supset \langle F_\sigma - F_\sigma(Au) \mid \sigma \supset \tau \rangle \\ &= I(V(\langle F_\sigma - F_\sigma(Au) \mid \sigma \supset \tau \rangle)) \\ &= I(V(\mathbf{k}[s] \cap \langle \theta_i - u_i \mid i \notin \tau \rangle)) \end{aligned} \quad (4.21)$$

where  $I$  stands for taking the defining ideal. On the other hand,  $J \subset I(V(J))$  is automatic for any ideal  $J$ . We therefore obtain (4.19).  $\square$

**Proposition 4.6** 1.

$$V(B_{\chi+\chi'}) \subset V(B_\chi) \cup (V(B_{\chi'}) + \chi) \quad \text{for } \chi, \chi' \in \mathbf{ZA}. \quad (4.22)$$

2.

$$V(B_{\chi+\chi'}) = V(B_\chi) \cup (V(B_{\chi'}) + \chi) \quad \text{for } \chi, \chi' \in \mathbf{NA}. \quad (4.23)$$

**Proof.**

1. Let  $p_\chi \in B_\chi$ ,  $p_{\chi'} \in B_{\chi'}$ , and  $P_\chi \in S_\chi$ ,  $P_{\chi'} \in S_{\chi'}$  be in the correspondence in Proposition 4.1. Then

$$\begin{aligned} P_\chi P_{\chi'} \partial^{\chi'} \partial^{\chi+} &= P_\chi p_{\chi'}(s) \partial^{\chi'} \partial^{\chi+} \\ &= p_{\chi'}(s - \chi) P_\chi \partial^{\chi+} \partial^{\chi'} \\ &= p_{\chi'}(s - \chi) p_\chi(s) \partial^{\chi-} \partial^{\chi'}. \end{aligned} \quad (4.24)$$

Hence  $p_{\chi'}(s - \chi) p_\chi(s) \in B_{\chi+\chi'}$ .

2. Let  $p_{\chi+\chi'} \in B_{\chi+\chi'}$  and  $P_{\chi+\chi'} \in S_{\chi+\chi'}$  be in the correspondence in Proposition 4.1. Then

$$p_{\chi+\chi'} = P_{\chi+\chi'} \partial^{\chi'} \cdot \partial^\chi.$$

Hence  $p_{\chi+\chi'}(s) \in B_\chi$ .

Furthermore

$$p_{\chi+\chi'}(s + \chi) \partial^\chi = \partial^\chi p_{\chi+\chi'}(s) = \partial^\chi P_{\chi+\chi'} \partial^{\chi'} \partial^\chi.$$

Hence  $p_{\chi+\chi'}(s + \chi) = \partial^\chi P_{\chi+\chi'} \partial^{\chi'}$ , which implies  $p_{\chi+\chi'}(s + \chi) \in B_{\chi'}$ .

□

**Proposition 4.7** Let  $\chi \in \mathbf{ZA}$ . Let  $p_\chi \in B_\chi$ ,  $p_{-\chi} \in B_{-\chi}$ , and  $P_\chi \in S_\chi$ ,  $P_{-\chi} \in S_{-\chi}$  be in the correspondence in Proposition 4.1. Then

$$P_{-\chi} P_\chi = p_\chi(s + \chi) p_{-\chi}(s). \quad (4.25)$$

**Proof.**

$$\begin{aligned} P_{-\chi} P_\chi \partial^{\chi+} &= P_{-\chi} p_\chi(s) \partial^{\chi-} \\ &= p_\chi(s + \chi) P_{-\chi} \partial^{\chi-} \\ &= p_\chi(s + \chi) p_{-\chi}(s) \partial^{\chi+}. \end{aligned} \quad (4.26)$$

Divide it by  $\partial^{\chi+}$  to obtain the conclusion. □

For  $\chi \in \mathbf{Z}A$ , define an ideal  $B_{-\chi, \chi}$  by

$$B_{-\chi, \chi} := \langle p_\chi(s + \chi)p_{-\chi}(s) \mid p_\chi \in B_\chi, p_{-\chi} \in B_{-\chi} \rangle. \quad (4.27)$$

Then the following proposition is immediate from the definition of  $B_{-\chi, \chi}$ .

**Proposition 4.8** 1.

$$V(B_{-\chi, \chi}) = (V(B_\chi) - \chi) \cup V(B_{-\chi}). \quad (4.28)$$

2.

$$V(B_{-\chi, \chi}) = V(B_{\chi, -\chi}) - \chi. \quad (4.29)$$

**Theorem 4.9** Let  $\chi \in \mathbf{Z}A$ . If  $\beta \notin V(B_{-\chi, \chi})$ , then two  $A$ -hypergeometric systems  $M_A(\beta)$  and  $M_A(\beta + \chi)$  are isomorphic.

**Proof.** First note that  $\beta \notin V(B_{-\chi, \chi})$  is equivalent to  $\beta + \chi \notin V(B_{\chi, -\chi})$  by Proposition 4.8. Take polynomials  $p_\chi \in B_\chi$  and  $p_{-\chi} \in B_{-\chi}$  such that  $p_\chi(\beta + \chi)p_{-\chi}(\beta) \neq 0$ . Let  $P_\chi \in S_\chi$ ,  $P_{-\chi} \in S_{-\chi}$  be in the correspondence in Proposition 4.1. Then by Proposition 4.7, we have the following equalities:

$$P_{-\chi}P_\chi = p_\chi(s + \chi)p_{-\chi}(s), \quad (4.30)$$

$$P_\chi P_{-\chi} = p_{-\chi}(s - \chi)p_\chi(s). \quad (4.31)$$

The multiplications by  $P_{-\chi}$ ,  $P_\chi$  respectively induce homomorphisms:

$$f : M_A(\beta) \longrightarrow M_A(\beta + \chi), \quad (4.32)$$

$$g : M_A(\beta + \chi) \longrightarrow M_A(\beta). \quad (4.33)$$

Then

$$g \circ f = p_\chi(\beta + \chi)p_{-\chi}(\beta)id_{M_A(\beta)} \quad (4.34)$$

and

$$f \circ g = p_{-\chi}((\beta + \chi) - \chi)p_\chi(\beta + \chi)id_{M_A(\beta + \chi)} \quad (4.35)$$

$$= p_{-\chi}(\beta)p_\chi(\beta + \chi)id_{M_A(\beta + \chi)}. \quad (4.36)$$

Hence  $f$  and  $g$  are isomorphisms.  $\square$

Now we are ready to prove the if-part of our main theorem.

**Proof of the if-part of Theorem 2.1.**

We suppose that  $E_\tau(\beta) = E_\tau(\beta')$  for all faces. Let  $\chi := \beta' - \beta$ . We claim  $\beta \notin V(B_{-\chi})$ . Assume the contrary. Then by Theorem 4.5, there exists a standard pair  $(u, \tau) \in \mathcal{S}(M_{-\chi})$  such that  $\beta - Au \in \mathbf{k}(A \cap \tau)$ . The equality  $E_\tau(\beta) = E_\tau(\beta')$  implies that there exists  $v \in \mathbf{N}^n$  such that  $\beta - \beta' = A(u - v)$ . Hence the intersection of  $Au + \mathbf{N}(A \cap \tau)$  with  $(\beta - \beta') + \mathbf{N}A$  is not empty. This contradicts the standardness of  $(u, \tau)$ . We have thus proved  $\beta \notin V(B_{-\chi})$ . By symmetry we have  $\beta' \notin V(B_\chi)$ , which is equivalent to  $\beta \notin V(B_\chi) - \chi$ . Hence  $\beta \notin V(B_{-\chi, \chi})$  by Proposition 4.8. From Theorem 4.9 we conclude  $M_A(\beta)$  is isomorphic to  $M_A(\beta')$ .  $\square$

As a corollary of the proof of the if-part of Theorem 2.1, we obtain the following.

**Corollary 4.10** *If two  $A$ -hypergeometric systems  $M_A(\beta)$  and  $M_A(\beta')$  are isomorphic, then there exists an operator  $P \in S_{\beta'-\beta}$  such that the multiplication by  $P$  from the right induces an isomorphism from  $M_A(\beta)$  to  $M_A(\beta')$ .*

## 5 Normal case

In this section, we consider the normal case:

$$\mathbf{N}A = \mathbf{Z}A \cap \mathbf{Q}_{\geq 0}A. \quad (5.37)$$

Many important examples are known to be normal, such as Aomoto-Gel'fand systems, the  $A$ -hypergeometric systems corresponding to  ${}_{p+1}F_p$ , Lauricella functions, etc. (see [9], [10]). It will turn out below that the parameter space can be classified in terms of the primitive integral support functions  $F_\sigma$  in the normal case.

**Lemma 5.1** *Assume  $A$  to be normal. Then we have the following.*

1.  $(\mathbf{Q}(A \cap \tau)) \cap \mathbf{Z}A$  equals  $\mathbf{Z}(A \cap \tau)$  for all faces  $\tau$ .
2.  $F_\sigma(\mathbf{N}A) = \mathbf{N}$  for all facets  $\sigma$ .
3. For a face  $\tau$ ,

$$\mathbf{N}A + \mathbf{Z}(A \cap \tau) = \mathbf{Z}A \cap \bigcap_{\sigma: \text{facet} \supset \tau} (\mathbf{N}A + \mathbf{k}(A \cap \sigma)). \quad (5.38)$$

**Proof.** (1) Let  $\chi \in (\mathbf{Q}(A \cap \tau)) \cap \mathbf{Z}A$ . Add a vector  $\chi' \in \mathbf{N}(A \cap \tau)$  to  $\chi$  so that  $\chi + \chi' \in \mathbf{Q}_{\geq 0}(A \cap \tau)$ . By the normality, we see  $\chi + \chi' \in \mathbf{N}(A \cap \tau)$ . Hence  $\chi$  belongs to  $\mathbf{Z}(A \cap \tau)$ .

(2) Let  $\chi \in \mathbf{Z}A$  satisfy  $F_\sigma(\chi) = 1$ . For  $\sigma' \neq \sigma$ , there exists  $a_j \in \sigma \setminus \sigma'$ . Hence there exists  $\chi' \in \mathbf{N}(A \cap \sigma)$  such that  $F_{\sigma'}(\chi + \chi') \geq 0$  for all facets  $\sigma'$ . By the normality,  $\chi + \chi' \in \mathbf{N}A$ . Since  $F_\sigma(\chi + \chi') = 1$ , we obtain  $F_\sigma(\mathbf{N}A) = \mathbf{N}$ .

(3) Let  $\chi \in \mathbf{Z}A$  satisfy  $F_\sigma(\chi) \geq 0$  for all facets containing  $\tau$ . For a facet  $\sigma$  not containing the face  $\tau$ , there exists  $a_j \in \tau \setminus \sigma$ . Hence there exists a vector  $\chi' \in \mathbf{N}(A \cap \tau)$  such that  $F_\sigma(\chi + \chi') \geq 0$  for all facets  $\sigma$  of the cone  $\mathbf{Q}_{\geq 0}A$ . By the normality,  $\chi + \chi' \in \mathbf{N}A$ , and thus  $\chi \in \mathbf{N}(A \setminus A \cap \tau) + \mathbf{Z}(A \cap \tau)$ .  $\square$

**Theorem 5.2** *Let  $\beta, \beta' \in \mathbf{k}^d$ . Then  $M_A(\beta) \simeq M_A(\beta')$  if and only if  $\beta - \beta' \in \mathbf{Z}A$  and  $\{\sigma : \text{facet}, F_\sigma(\beta) \in \mathbf{N}\} = \{\sigma : \text{facet}, F_\sigma(\beta') \in \mathbf{N}\}$ .*

**Proof.** By Proposition 2.2 (3), the only-if-part follows from Theorem 2.1.

Next we prove the if-part. Suppose  $\beta - \beta' \in \mathbf{Z}A$  and  $\{\sigma : \text{facet}, F_\sigma(\beta) \in \mathbf{N}\} = \{\sigma : \text{facet}, F_\sigma(\beta') \in \mathbf{N}\}$ . By Lemma 5.1 (1), (2), and Propositions 2.2, 2.3, we obtain  $E_\sigma(\beta) = E_\sigma(\beta')$  for all facets. By Lemma 5.1 (3), the if-part follows from Theorem 2.1.  $\square$

**Example 5.3** *Let*

$$A = \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & -1 \end{pmatrix}.$$

Let  $\beta \in \mathbf{ZA} = \mathbf{Z}^d$ . Then by Theorem 5.2, the  $A$ -hypergeometric system  $M_A(\beta)$  is isomorphic to

$$\begin{aligned} M_A({}^t(0, 0, 0)) & \text{ if } \beta_1 \geq 0, \beta_2 \geq 0, \beta_1 + \beta_3 \geq 0, \beta_2 + \beta_3 \geq 0, \\ M_A({}^t(-1, 0, 1)) & \text{ if } \beta_1 < 0, \beta_2 \geq 0, \beta_1 + \beta_3 \geq 0, \beta_2 + \beta_3 \geq 0, \\ M_A({}^t(0, -1, 1)) & \text{ if } \beta_1 \geq 0, \beta_2 < 0, \beta_1 + \beta_3 \geq 0, \beta_2 + \beta_3 \geq 0, \\ M_A({}^t(0, 1, -1)) & \text{ if } \beta_1 \geq 0, \beta_2 \geq 0, \beta_1 + \beta_3 < 0, \beta_2 + \beta_3 \geq 0, \\ M_A({}^t(1, 0, -1)) & \text{ if } \beta_1 \geq 0, \beta_2 \geq 0, \beta_1 + \beta_3 \geq 0, \beta_2 + \beta_3 < 0, \\ M_A({}^t(-1, -1, 1)) & \text{ if } \beta_1 < 0, \beta_2 < 0, \beta_1 + \beta_3 \geq 0, \beta_2 + \beta_3 \geq 0, \\ M_A({}^t(-1, 0, 0)) & \text{ if } \beta_1 < 0, \beta_2 \geq 0, \beta_1 + \beta_3 < 0, \beta_2 + \beta_3 \geq 0, \\ M_A({}^t(0, -1, 0)) & \text{ if } \beta_1 \geq 0, \beta_2 < 0, \beta_1 + \beta_3 \geq 0, \beta_2 + \beta_3 < 0, \\ M_A({}^t(0, 0, -1)) & \text{ if } \beta_1 \geq 0, \beta_2 \geq 0, \beta_1 + \beta_3 < 0, \beta_2 + \beta_3 < 0, \\ M_A({}^t(-2, -1, 1)) & \text{ if } \beta_1 < 0, \beta_2 < 0, \beta_1 + \beta_3 < 0, \beta_2 + \beta_3 \geq 0, \\ M_A({}^t(-1, -2, 1)) & \text{ if } \beta_1 < 0, \beta_2 < 0, \beta_1 + \beta_3 \geq 0, \beta_2 + \beta_3 < 0, \\ M_A({}^t(-1, 0, -1)) & \text{ if } \beta_1 < 0, \beta_2 \geq 0, \beta_1 + \beta_3 < 0, \beta_2 + \beta_3 < 0, \\ M_A({}^t(0, -1, -1)) & \text{ if } \beta_1 \geq 0, \beta_2 < 0, \beta_1 + \beta_3 < 0, \beta_2 + \beta_3 < 0, \\ M_A({}^t(-1, -1, 0)) & \text{ if } \beta_1 < 0, \beta_2 < 0, \beta_1 + \beta_3 < 0, \beta_2 + \beta_3 < 0. \end{aligned}$$

## 6 Monomial curve case

In this section, we consider  $d = 2$  case. Let

$$A = \begin{pmatrix} 1 & 1 & 1 & \cdots & 1 & 1 \\ 0 & i_2 & i_3 & \cdots & i_{n-1} & i_n \end{pmatrix}$$

with  $0 < i_2 < i_3 < \cdots < i_n$  relative prime integers. Put  $F_{\sigma_1}(s) = s_2$  and  $F_{\sigma_2}(s) = i_n s_1 - s_2$ .

We denote by  $\mathcal{E}(A)$  the set of holes, i.e.,

$$\mathcal{E}(A) : = ((\mathbf{NA} + \mathbf{Z}a_1) \cap (\mathbf{NA} + \mathbf{Z}a_n)) \setminus \mathbf{NA} \quad (6.39)$$

$$\begin{aligned} & = \{ \beta \mid E_{\mathbf{Q}_{\geq 0}A}(\beta) = \{0\}, E_{\sigma_1}(\beta) = \{0\}, \\ & \quad E_{\sigma_2}(\beta) = \{0\}, E_{\{0\}}(\beta) = \emptyset \}. \end{aligned} \quad (6.40)$$

The rank of  $M_A(\beta)$  is  $d$  or  $d + 1$ , and it equals  $d + 1$  if and only if  $\beta \in \mathcal{E}(A)$  (see [2], [13]).

**Lemma 6.1** *For any face  $\tau$ ,*

$$\mathbf{ZA} \cap (\mathbf{k}(A \cap \tau)) = \mathbf{Z}(A \cap \tau). \quad (6.41)$$

**Proof.** When  $\tau$  is the whole cone  $\mathbf{Q}_{\geq 0}A$  or the origin  $\{0\}$ , the statement is trivial.

Note that  $\beta$  belongs to  $\mathbf{ZA}$  if and only if  $F_{\sigma_1}(\beta) \in \mathbf{Z}$ ,  $F_{\sigma_2}(\beta) \in \mathbf{Z}$ , and  $F_{\sigma_1}(\beta) + F_{\sigma_2}(\beta) \in i_n \mathbf{Z}$ . Suppose  $\beta \in \mathbf{ZA} \cap (\mathbf{k}(A \cap \sigma_1))$ . Then  $F_{\sigma_2}(\beta) \in i_n \mathbf{Z}$ . When  $F_{\sigma_2}(\beta) = di_n$ , we have  $\beta = da_n$ .  $\square$

**Corollary 6.2**

$$\mathcal{E}(A) = \{ \beta \in \mathbf{ZA} \mid F_{\sigma_1}(\beta) \in F_{\sigma_1}(\mathbf{NA}), F_{\sigma_2}(\beta) \in F_{\sigma_2}(\mathbf{NA}) \} \setminus \mathbf{NA}. \quad (6.42)$$

**Proof.** This is immediate from Lemma 6.1.  $\square$

Theorem 2.1 in the monomial curve case is as follows.

**Theorem 6.3** *Let  $\beta, \beta' \in \mathbf{k}^d$ .*

1. *Suppose  $\beta \notin \mathcal{E}(A)$ . Then  $M_A(\beta')$  is isomorphic to  $M_A(\beta)$  if and only if  $\beta - \beta' \in \mathbf{ZA}$ ,  $\beta' \notin \mathcal{E}(A)$ , and  $\{\sigma_i : F_{\sigma_i}(\beta) \in F_{\sigma_i}(\mathbf{NA})\} = \{\sigma_i : F_{\sigma_i}(\beta') \in F_{\sigma_i}(\mathbf{NA})\}$ .*
2. *Suppose  $\beta \in \mathcal{E}(A)$ . Then  $M_A(\beta')$  is isomorphic to  $M_A(\beta)$  if and only if  $\beta \in \mathcal{E}(A)$ .*

**Proof.** (2) directly follows from Theorem 2.1.

The only-if-part of (1) follows from Theorem 2.1 by Proposition 2.2 (3). Next suppose that  $\beta - \beta' \in \mathbf{ZA}$ ,  $\beta, \beta' \notin \mathcal{E}(A)$ , and that  $\{\sigma_i : F_{\sigma_i}(\beta) \in F_{\sigma_i}(\mathbf{NA})\} = \{\sigma_i : F_{\sigma_i}(\beta') \in F_{\sigma_i}(\mathbf{NA})\}$ . Then by Lemma 6.1, Proposition 2.2 (3), and Proposition 2.3 (2), we have  $E_{\sigma_i}(\beta) = E_{\sigma_i}(\beta')$  for  $i = 1, 2$ . Moreover we know  $E_{\{0\}}(\beta), E_{\{0\}}(\beta') = \emptyset$  from Proposition 2.2 (2). Hence  $M_A(\beta)$  and  $M_A(\beta')$  are isomorphic by Theorem 2.1.  $\square$

**Example 6.4** ([13, Chapter 4]) *Let*

$$A = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 \\ 0 & 2 & 4 & 7 & 9 \end{pmatrix}.$$

*Then*

$$F_{\sigma_1}(\mathbf{NA}) = \{0, 2, 4, 6, 7, 8, 9, \dots\}, \quad (6.43)$$

*and*

$$F_{\sigma_2}(\mathbf{NA}) = \{0, 2, 4, 5, 6, 7, 8, 9, \dots\}, \quad (6.44)$$

*Parameters in  $\mathbf{ZA} = \mathbf{Z}^2$  are decomposed into five parts according to the isomorphism classes of their corresponding A-hypergeometric systems:*

1.  $\mathbf{NA}$ ,
2.  $\{ {}^t(\beta_1, \beta_2) \mid \beta_2 \in F_{\sigma_1}(\mathbf{NA}), 9\beta_1 - \beta_2 \notin F_{\sigma_2}(\mathbf{NA}) \}$ ,
3.  $\{ {}^t(\beta_1, \beta_2) \mid \beta_2 \notin F_{\sigma_1}(\mathbf{NA}), 9\beta_1 - \beta_2 \in F_{\sigma_2}(\mathbf{NA}) \}$ ,
4.  $\{ {}^t(\beta_1, \beta_2) \mid \beta_2 \notin F_{\sigma_1}(\mathbf{NA}), 9\beta_1 - \beta_2 \notin F_{\sigma_2}(\mathbf{NA}) \}$ ,
5.  $\mathcal{E}(A) = \{ {}^t(2, 10), {}^t(2, 12), {}^t(3, 19) \}$  : *the set of holes.*

## 7 Final remark

Thanks to Theorem 2.1, all  $D$ -invariants of  $A$ -hypergeometric systems can be described in terms of  $E_\tau(\beta)$ ; the characteristic cycles (in particular, the rank), the monodromy representations, etc. One of most recent results is given by Tsushima ([15]) on Laurent polynomial solutions. He has proved that the vector space of Laurent polynomial solutions of  $M_A(\beta)$  has a basis consisting of canonical series whose negative support corresponds to a face  $\tau$  of  $\mathbf{Q}_{\geq 0}A$  such that  $\dim \tau = |\{a_j \mid a_j \in \tau\}|$ , and that  $0 \in E_\tau(\beta)$  but  $0 \notin E_{\tau'}(\beta)$  for any  $\tau' \subset \tau$ . In particular, the dimension of the vector space of Laurent polynomial solutions equals the cardinality of the set of such faces. This is a generalization of the corresponding result by Cattani, D'Andrea and Dickenstein ([2]) in the monomial curve case.

## References

- [1] Adolphson, A.(1994): Hypergeometric functions and rings generated by monomials. *Duke Mathematical Journal* **73**, 269–290.
- [2] Cattani, E., D'Andrea, C., Dickenstein, A. (1998): The  $\mathcal{A}$ -hypergeometric system associated with a monomial curve. *Duke Mathematical Journal* **99**, 179–207.
- [3] Cattani, E., Dickenstein, A., Sturmfels, B. (1999): Rational hypergeometric functions. *math.AG/9911030*.
- [4] Gel'fand, I.M., Kapranov, M.M., Zelevinskii, A.V. (1990): Generalized Euler integrals and  $A$ -hypergeometric functions. *Advances in Mathematics* **84**, 255–271.
- [5] Gel'fand, I.M., Zelevinskii, A.V., Kapranov, M.M. (1988): Equations of hypergeometric type and Newton polyhedra. *Soviet Mathematics Doklady* **37**, 678–683.
- [6] Gel'fand, I.M., Zelevinskii, A.V., Kapranov, M.M. (1989): Hypergeometric functions and toral manifolds. *Functional Analysis and its Applications* **23**, 94–106.
- [7] Hosten, S., Thomas, R.R. (1998): Standard pairs and group relaxations in integer programming, to appear in *Journal of Pure and Applied Algebra*.
- [8] Oda, T. (1988): *Convex Bodies and Algebraic Geometry: An Introduction to the Theory of Toric Varieties*. *Ergebnisse der Mathematik und ihrer Grenzgebiete, 3. Folge, Band 15*, Springer, Heidelberg.
- [9] Saito, M. (1994): Normality of affine toric varieties associated with Hermitian symmetric spaces. *Journal of Mathematical Society of Japan* **46**, 699–724.

- [10] Saito, M. (1995): Contiguity Relations for the Lauricella Functions. *Funkcialaj Ekvacioj* **38**, 37–58.
- [11] Saito, M. (1996): Symmetry algebras of normal  $\mathcal{A}$ -hypergeometric systems. *Hokkaido Mathematical Journal* **25**, 591–619.
- [12] Saito, M., Sturmfels, B., Takayama, N. (1999): Hypergeometric polynomials and integer programming. *Compositio Mathematica*, **155**, 185–204.
- [13] Saito, M., Sturmfels, B., Takayama, N. (1999): *Gröbner deformations of hypergeometric differential equations*. Algorithms and Computation in Mathematics **6**, Springer, Berlin, Heidelberg, New York.
- [14] Saito, M., Takayama, N. (1994): Restrictions of  $\mathcal{A}$ -hypergeometric systems and connection formulas of the  $\Delta_1 \times \Delta_{n-1}$ -hypergeometric function. *International Journal of Mathematics* **5**, 537–560.
- [15] Tsushima, T. : Laurent polynomial solutions of  $\mathcal{A}$ -hypergeometric systems. in preparation.

Department of Mathematics  
Hokkaido University  
Sapporo, 060-0810  
Japan  
e-mail: saito@math.sci.hokudai.ac.jp