Double scaling and intermittency in shear dominated flows

C.M. Casciola, *R. Benzi, [†]P. Gualtieri^{*}, B. Jacob^{*}, & R. Piva^{*}.

November 13, 2018

1 Abstract

The Refined Kolmogorov Similarity Hypothesis is a valuable tool for the description of intermittency in isotropic conditions. For flows in presence of a substantial mean shear, the nature of intermittency changes since the process of energy transfer is affected by the turbulent kinetic energy production associated with the Reynolds stresses. In these conditions a new form of refined similarity law has been found able to describe the increased level of intermittency which characterizes shear dominated flows. Ideally a length scale associated with the mean shear separates the two ranges, i.e. the classical Kolmogorov-like inertial range, below, and the shear dominated range, above. However, the data analyzed in previous papers correspond to conditions where the two scaling regimes can only be observed individually.

In the present letter we give evidence of the coexistence of the two regimes and support the conjecture that the statistical properties of the dissipation field are practically insensible to the mean shear. This allows for a theoretical prediction of the scaling exponents of structure functions in the shear dominated range based on the known intermittency corrections for isotropic flows. The prediction is found to closely match the available numerical and experimental data.

2 Introduction

At large Reynolds number (*Re*) turbulent flows are characterized by strong non Gaussian intermittent fluctuations. For homogeneous isotropic turbulence, a quantitative measure of intermittency can be given by using the structure functions $\langle \delta V^p(r) \rangle$ where

$$\delta V(\vec{r}) = \left[\vec{u}(\vec{x} + \vec{r}) - \vec{u}(\vec{x})\right] \cdot \frac{\vec{r}}{r}.$$
(1)

Then, the generalized dimensionless flatness

$$F_p(r) = \frac{\langle \delta V^p(r) \rangle}{\langle \delta V^2(r) \rangle^{p/2}}$$

$$\tag{2}$$

exhibits intermittency, in the sense that $F_p(r) \to \infty$ for $r \to 0$ and $Re \to \infty$. For $\eta \ll r \ll L_0$, where η is the Kolmogorov dissipation length and L_0 is the integral scale of turbulence, structure functions show scaling behavior, i.e. $\langle \delta V^p(r) \rangle \propto r^{\zeta(p)}$, where $\zeta(p)$ are anomalous scaling exponents $(\zeta(p) \neq p/q\zeta(q))$, and $\zeta(3) = 1$ due to the Kolmogorov four-fifth equation [1]. It is a remarkable result, obtained in the last ten years, that $\zeta(p)$ are observed to be universal, i.e. independent of the Reynolds number and of the forcing mechanism, for homogeneous and isotropic turbulence [2].

Much less information is available for non isotropic turbulence. Recently, a number of experimental [3] and numerical investigations [4] in shear flow turbulence have shown that intermittency increases when the shear strongly affects the energy cascade. In the language of scaling exponents, an increase of intermittency means that $\zeta(p)$ are different from those observed in homogeneous and isotropic turbulence. Based on DNS of turbulent channel flow, it

^{*}Dip. Mecc. Aeron., Università di Roma "La Sapienza", via Eudossiana 18, 00184, Roma, Italy.

[†]Dip. di Fisica and INFN, Università di Roma "Tor Vergata", Via della Ricerca scientifica 1, 00133, Roma, Italy.

was recently proposed [5] that the increase of intermittency is due to the breaking of the Kolmogorov Refined Similarity Hypothesis (RKSH), which for homogeneous and isotropic turbulence reads [6]

$$<\delta V^p(r) > \propto <\epsilon_r^{p/3} > r^{p/3} \tag{3}$$

where

$$\epsilon_r = \frac{1}{\mathcal{B}(r)} \int_{\mathcal{B}(r)} \epsilon_{loc}(\vec{x}) d^3x \tag{4}$$

and $\mathcal{B}(r)$ is a volume of characteristic size r while $\epsilon_{loc}(\vec{x})$ is the instantaneous local rate of energy dissipation. Equation (3), in its Extended Self Similarity (ESS) formulation

$$<\delta V^p(r)>\propto \frac{<\epsilon_r^{p/3}>}{<\epsilon_r>^{p/3}}<\delta V^3>^{p/3}$$
(5)

has been successfully checked for a wide range of Reynolds numbers in different homogeneous and isotropic turbulent flows [7].

For shear flow turbulence, it has been suggested that equation (5) should be replaced [5] by

$$<\delta V^{p}(r) > \propto \frac{<\epsilon_{r}^{p/2}>}{<\epsilon_{r}>^{p/2}} < \delta V^{2}>^{p/2}$$
(6)

for $r \gg L_s$ where $L_s = \sqrt{\overline{\epsilon}/S^3}$, S being the mean shear in the system. Following these considerations, the shear scale ideally separates, with regard to the scaling properties of structure functions, the range of scales where turbulent kinetic energy production prevails $(r \gg L_s)$ from the range of scales characterized by purely inertial energy transfer $(r \ll L_s)$. Equation (6) is able to explain the observed increase in intermittency, by assuming that the scaling properties of $\langle \epsilon_r^{r} \rangle$ are not changed by the mean shear S.

It is the aim of this letter to present and discuss numerical and experimental evidence that the two scaling regimes, predicted by eq. (5) and (6), are indeed observed simultaneously in the range below and above L_s respectively. Moreover, the results confirm the conjecture that the scaling properties of the energy dissipation are not modified appreciably with respect to homogeneous and isotropic turbulence.

3 Data sets and analysis

We discuss two sets of data, one obtained by a long and highly resolved DNS of homogeneous shear flow turbulence [8], the other by hot wire measurements in the logarithmic region of a turbulent boundary layer [9].

Concerning the homogeneous shear flow, we have considered a turbulent flow with an imposed mean velocity gradient S free from boundaries. The Navier-Stokes equations, written in terms of velocity fluctuations, are solved by using an efficient pseudo-spectral method with a third order Runge-Kutta scheme for time advancement following the transformation of variable proposed by Rogallo [10]. As shown by Pumir [11] and recently confirmed by the present authors [8], the flow reaches a statistical steady state characterized by large fluctuations of the turbulent kinetic energy. The growth of turbulent kinetic energy is associated to large values of the Reynolds stresses, produced by a well defined system of streamwise vortices via a lift-up mechanism [12]. In this flow, because of shear scale fluctuations due to the mentioned behavior of both turbulent kinetic energy and Reynolds stresses, the crossover between the two scaling ranges is not sharply defined. In fact, we observe an overlapping of the two scaling regimes, and the resulting scaling shows an effective slope. In order to reduce as much as possible the fluctuations of the shear scale, a conditional sampling is introduced by considering only flow configurations where the production term exceeds a given threshold. Among these configurations, only those corresponding to a large value of turbulent kinetic energy ($\mathcal{E} > \alpha \mathcal{E}_{rms}$) are retained to reduce the mean value of the ratio $L_s/\eta \propto \Omega^{3/4}$ (Ω being the mean enstrophy).

Concerning experiments, we analyze the velocity data on a flat plate boundary layer measured in a wind tunnel (test section length of 150cm) operated at 11.9m/s. The boundary layer thickness is $\simeq 25mm$ and the Reynolds number based on the momentum thickness is about 2200. The boundary layer has the expected logarithmic region with the usual log-law constants [13]. Hot wire measurements were performed at several distances from the plate, using a constant temperature anemometer. The data acquisition was long enough to achieve convergence of the sixth order structure function.

4 Double scaling regime

We begin by analyzing the DNS of the homogeneous shear flow. We have strong evidence that for $r > L_s$, eq. (5) fails and the new form of RKSH is established, as reported in [14], [8] and further discussed in the following. Furthermore, the statistical properties of energy dissipation $< \epsilon_r^q >$ are not distinguishable from those observed in homogeneous and isotropic turbulence. The last statement can be directly checked by looking at figure 1 where we plot $< \epsilon_r^3 >$ versus $< \epsilon_r^2 >$ both for homogeneous shear flow and isotropic turbulence while, in the inset of the same figure, we plot $< \epsilon_r^2 >$ versus r/η for both cases. At variance with DNS, a direct measurement of $< \epsilon_r^q >$ is not available for the experimental data and we are not fully confident in the one dimensional surrogate of ϵ_{loc} as a direct measure of the local rate of energy dissipation, being the flow strongly anisotropic. Nevertheless by using the one dimensional surrogate, we can practically reproduce the results shown in figure 1.

At any rate, to be cautious, we may avoid the explicit use of the energy dissipation by plotting structure functions in the form suggested by Ruiz-Chavarria et. al. [15]. Specifically, here, we introduce indicators based on $\langle \delta V^p(r) \rangle$ to detect the two scaling regions and to compare our findings with the predictions made in eq. (5) and (6). We remark that, both for numerical and experimental data, the Reynolds number is not large enough to observe the scaling of $\langle \delta V^p(r) \rangle$ and $\langle \epsilon_r^q \rangle$ with respect to separation. Thus we employ the ESS as a valuable tool to estimate the scaling exponents. In particular this implies that the exponents $\tau(q)$ are defined by the relation $\langle \epsilon_r^q \rangle \propto \langle \delta V^3 \rangle^{\tau(q)}$.

Following eq. (5) and (6) and the above discussion, we compute both for the DNS and the experimental data the quantity $\sigma_p \equiv \langle \delta V^p \rangle / \langle \delta V^2 \rangle^{p/2}$ and $\rho_p \equiv \langle \delta V^p \rangle / \langle \delta V^3 \rangle^{p/3}$ which are expected to satisfy the relations:

$$\sigma_p \propto \begin{cases} <\delta V^3 >^{\tau(p/2)} & r \gg L_s \\ <\delta V^3 >^{\tau(p/3)-\tau(2/3)p/2} & r \ll L_s \end{cases}$$
(7)

and

$$\rho_p \propto \begin{cases} <\delta V^3 >^{\tau(p/2)-\tau(3/2)p/3} & r \gg L_s \\ <\delta V^3 >^{\tau(p/3)} & r \ll L_s. \end{cases}$$
(8)

Equations (7) and (8) allows us to compare the ESS exponents of σ_p and ρ_p against the exponents predicted by equations (5) and (6). Let us remark that eq. (7) and (8) are also based on the assumption that $\tau(q)$ are the same both for shear dominated flows and homogeneous and isotropic turbulence, as supported by the DNS data for $\langle \epsilon_r^q \rangle$ shown in figure 1.

In figure 2 we plot $\log \sigma_6$ against $\log \langle \delta V^3 \rangle$ for the data of the homogeneous shear flow and the turbulent boundary layer. The fits for $r \gg L_s$ are in close agreement with the value of $\tau(3) = -0.59$ expected from homogeneous and isotropic turbulence. In the inset of figure 2 we show the local slope $d[\log \sigma_6]/d[\log \langle \delta V^3 \rangle]$ computed from the homogeneous shear flow dataset. The two dashed lines indicate the numbers $\tau(2) - 3\tau(2/3)$ and $\tau(3)$, i.e. the expected scaling exponents for $r \ll L_s$ and $r \gg L_s$ respectively.

Figure 2 shows the main result of this letter, i.e. the clear evidence of the two scaling regions below and above L_s . Specifically, the experimental and numerical evidence of the coexistence of two different intermittent regions in shear flow turbulence. As a further check of the theory, in figure 3, we show $\log \rho_6$ versus $\log \langle \delta V^3 \rangle$ both for the homogeneous shear flow and the turbulent boundary layer while in the inset we show the local slope $d[\log \rho_6]/d[\log \langle \delta V^3 \rangle]$. Also for the variable ρ_6 we can claim a very good agreement between the observed experimental and numerical results against theoretical predictions. Our results concerning the double scaling of structure functions are consistent and complementary with the generalized structure function $\langle \delta V^3 + \alpha r \delta V^2 \rangle^{p/3} \rangle$ proposed by Toschi et. al. [16].

Finally we remark that, by denoting $\zeta(p)$ the anomalous exponents in shear flow turbulence and $\zeta(p)$ the anomalous exponent in homogeneous isotropic turbulence, the ESS estimate becomes

$$\hat{\zeta}(p) = \frac{p}{3} \left[1 - \tau \left(\frac{3}{2} \right) \right] - \frac{p}{2} + \zeta \left(\frac{3p}{2} \right).$$
(9)

The first term comes from the fact that in shear turbulence, we may express

$$<\delta V^3 > \propto <\epsilon_r^{3/2} > <\delta V^3 >^{3\zeta(2)/2} \tag{10}$$

which implies $\zeta(2) = 2/3[1 - \tau(3/2)]$. Equation (9) provides a theoretical estimation for the scaling exponents of structure functions in shear dominated flows by using the intermittency corrections of isotropic turbulence. The values given by eq. (9) are compared against their direct measure in the DNS of the homogeneous shear flow in table 1.

5 Final remarks

We have analyzed scaling properties of velocity fluctuations in shear dominated flows considering a DNS of a homogeneous shear flow and hot wire measurements in the logarithmic region of a turbulent boundary layer. When dealing with DNS data a conditional statistical analysis has allowed us to consider only shear dominated samples and to reduce the fluctuations of the shear scale resulting in a well defined scaling behavior below and above L_s respectively. To by pass the use of the one dimensional surrogate of energy dissipation, in the experiments, we have evaluated the refined similarity laws only in terms of structure functions.

The analyzed data show clearly the coexistence of a double scaling behavior of structure functions across the shear scale. Actually the scaling exponents computed below and above the shear scale are in a very good agreement with the theoretical values provided by the classical RKSH [6], [7], and the new refined similarity law [5] respectively. Since the statistical properties of the energy dissipation are weakly affected by the shear, intermittency corrections of isotropic flows can be successfully employed in the new form of similarity law, see table 1, to predict the scaling exponents of structure functions in the shear dominated range.

Table 1: Scaling exponents of structure functions (DNS) above and below the shear scale L_s . Data are compared with those of homogeneous and isotropic turbulence and with the prediction of eq. (9).

p	1	2	3	4	5	6
$r < L_s$	0.36	0.69	1.00	1.28	1.54	1.78
$r > L_s$	0.38	0.72	1.00	1.23	1.42	1.56
hom. iso	0.36	0.69	1.00	1.28	1.54	1.78
eq. (9)	0.39	0.73	1.00	1.23	1.42	1.58

References

- Kolmogorov, The local structure of turbulence in incompressible viscous fluid for very large Reynolds number, Dolk. Akad. SSSR 30, 301, 1941, reprinted in Proc. R. Soc. Lond., A 434, 15-19, 1991.
- [2] Benzi, Ciliberto, Baudet, Ruiz Chavarria, On the scaling of three dimensional homogeneous and isotropic turbulence, Physica D, 80, 385-398, 1995.
- [3] Anselmet, Gagne, Hopfinger, Antonia, High order velocity structure functions in turbulent shear flow, J. Fluid Mech., 140, 63, 1984.
- [4] Toschi, Amati, Succi, Benzi, Piva, Intermittency and structure functions in channel flow turbulence, Phys. Rev. Lett, 82(25), 5044-5049, 1999.
- [5] Benzi, Amati, Casciola, Toschi, Piva, Intermittency and scaling laws for wall bounded turbulence, Phys. Fluids, 11(6), 1-3, 1999.
- [6] Kolmogorov, A refinement of previous hypothesis concerning the local structure of turbulence in incompressible viscous fluid for very large Reynolds number, J. Fluid. Mech. 13, 82-85, 1962.
- [7] Benzi, Biferale, Ciliberto, Struglia, Tripicccione, Generalized scaling in fully developed turbulence, Physica D, 96, 162-181, 1996.
- [8] Gualtieri, Casciola, Benzi, Amati, Piva, Scaling laws and intermittency in homogeneous shear flow, Submitted to Phys. Fluids., Nov. 2000.
- [9] Jacob, Olivieri, Casciola, Experimental assessment of a new form of scaling law for near wall turbulence, Submitted to Phys. Fluids, Gen. 2001.
- [10] Rogallo, Numerical experiments in homogeneous turbulence, NASA T.M., 81315, 1981.
- [11] Pumir, Turbulence in homogeneous shear flow, Phys. Fluids, 8(11), 3112-3127, 1996.
- [12] Kida and Tanaka, Dynamics of vortical structures in homogeneous shear flow, J. Fluid Mech., 274, 43-68, 1994.
- [13] Monin, Yaglom, Statistical fluid mechanics. Vol. 1,2, The MIT press, 1975.
- [14] Benzi, Casciola, Gualtieri, Piva, Numerical evidence of a new similarity law in shear dominated flows, to appear in Comp. and Math.
- [15] Ruiz-Chavarria, Ciliberto, Baudet, Leveque, Scaling properties of the streamwise component of velocity in a turbulent boundary layer, Physica D, 141, 183-198, 2000.
- [16] Toschi, Leveque, Ruiz-Chavarria, Shear effects in non-homogeneous turbulence, Phys. Rev. Lett, 85(7), 1436-1439, 2000.

FIGURES



Figure 1: $\langle \epsilon_r^3 \rangle = vs. \quad \langle \epsilon_r^2 \rangle$ in the homogeneous shear flow (circles) and in homogeneous isotropic turbulence (diamonds). In the inset $\langle \epsilon_r^2 \rangle = vs. \quad r/\eta$ for the two cases.



Figure 2: $\log \sigma_6 \quad vs. \quad \log < \delta V^3 > \text{in the homogeneous shear flow (circles) and in the turbulent boundary layer at } y^+ = 115 \text{ (triangles). DNS and experimental data are fitted at scales } r > L_s \text{ by power laws with a slope } s = -0.58 \text{ and } s = -0.59 \text{ respectively. In the inset, the local slope } d[\log \sigma_6]/d[\log < \delta V^3 >] \quad vs. \quad r/\eta \text{ in the homogeneous shear flow obtained by considering the conditional sampling with } \alpha = 1.3 \text{ (solid line). The dotted lines corresponds to the two scalings given by eq. (7) at scales <math>r < L_s (-0.3)$ and $r > L_s (-0.59)$ using the values of $\tau(q)$ for isotropic turbulence.



Figure 3: $\log \rho_6 \quad vs. \quad \log < \delta V^3 > \text{in the homogeneous shear flow (circles) and in the turbulent boundary layer at <math>y^+ = 115$ (triangles). Both DNS and experimental data are fitted at scales $r < L_s$ by a power law with a slope s = -0.22. In the inset, the local slope $d[\log \rho_6]/d[\log < \delta V^3 >] \quad vs. \quad r/\eta$ for the homogeneous shear flow obtained by considering the conditional sampling with $\alpha = 1.3$ (solid line). The dotted lines corresponds to the two scaling given by eq. (8) at scales $r < L_s$ (-0.22) and $r > L_s$ (-0.41) using the values of $\tau(q)$ for isotropic turbulence.