Complex version KdV equation and the periods solution

Yang Lei, Yang Kongqing, Luo Honggang

(Department of physics, Lanzhou University, Gansu 730000, China)

 $(29/6\ 1999)$

In this paper, the complex version KdV equation is discussed. The corresponding coupled equations is a integrable system in the sense of the bi-Hamiltonian structure, so the complex version KdV equation is integrable. A new spectral form is given, the periodic solution of the complex version KdV equation is obtained. It is showed that the periodic solution is the classical solution.

I. INTRODUCTION

Since 80's, the coupled KdV equations as an important mathematical model has been studied widely. In 1981, Fuchssteiner[1] made a detailed study of four coupled KdV equations and gave the bi-Hamiltonian structure of them. One coupled KdV equations is the complexly-coupled KdV equations

$$u_t = u_{xxx} + 6uu_x + 6\varphi\varphi_x, \varphi_t = \varphi_{xxx} + 6u\varphi_x + 6u_x\varphi.$$
(1)

Later, Oevel[2] pointed out that ".....inserting a complex ansatz $u + i\varphi$ into the KdV.....it is a complex version of the KdV poses two conservation laws in every order. On the following day, Hirota and Satsuma introduced a coupled KdV equations[3], the integrability of the equations was discussed by the bi-Hamiltonian structure[1,2] and Lax pair[4]. Recently, more coupled KdV equations and their bi-Hamiltonian structure has been presented [5,6].

In this paper, the complex version KdV equation is considered,

$$u_t + \alpha u u_x + \beta u_{xxx} = 0, \tag{2}$$

where x is a one-dimension space, t is a one-dimension time, α and β are coefficients, which play important role in our numerical results, u(x,t) = p(x,t) + iq(x,t) is an unknown variable, p(x,t) is the real part of u(x,t), q(x,t)is the imaginary part of u(x,t). Extending the lax pair of the KdV equation to complex field, $L = -D^2 + p + iq$, $M = 6(p + iq)D - 4D^3 + 3p_x + 3iq_x$, the complex version KdV equation can be derived from the lax equation $L_t = [M, L]$. And the complex lax pair binds with the complex potential scattering and inverse scatting problem of the schrödinger equation, which had been studied in a optical problem[7]. Here, let $\alpha = \beta = 1$, substituting the u(x,t) = p(x,t) + iq(x,t) into the equation(2), separating the real part and the imaginary part from the equation(2), a coupled KdV equations is obtained

$$p_t + p_{xxx} + pp_x - qq_x = 0, q_t + q_{xxx} + pq_x + p_xq = 0.$$
(3)

The equation(3) are equal to the equation(2) completely and has a little of differences with the Fuchssteiner's equation(1). Under the transformation: $p \to 6u$, $q \to 6i\phi$, $x \to -x$, the equation(3) are exactly equivalent to equation(1).

II. METHOD DESCRIBING AND THE PERIODIC SOLUTION

In general, the nonlinear evolution equation is written as $u_t(x,t) = F(u(x,t), u_x(x,t), \ldots)$, here F(u) is a non-linear function of $u(x,t), u_x(x,t), \ldots$, which is a nonlinear operator involving only the spatial derivatives. Take the Fourier expansion,

$$u(x,t) = \sum_{n=-\infty}^{\infty} u_n(t)e^{ik_nx}$$
(4)

where $k_n = 2n\pi/L$ and L is a period interval in space. Let expression (4) come in with $u_t(x,t) = F(u(x,t))$, receive the infinite reference ordinary differential equations(ODE), which is highly nonlinearity and complexity in nature, the exact solutions of the ODEs can't be obtained. In this paper, a new spectral form is given, expanding u(x,t) on the sub-set of Fourier basis $\{e^{ik_nx}, n = 0, 1, 2, ...\}$, namely,

$$u(x,t) = \sum_{n=0}^{\infty} u_n(t)e^{ik_nx}$$
(5)

where $k_n = 2n\pi/L$. Substitute the formula (5) into $u_t(x,t) = F(u(x,t))$, thus

$$\sum_{n=0}^{\infty} \frac{du_n}{dt} e^{ik_n x} = F(u). \tag{6}$$

Multiplication of equations (6) by $e^{-ik_m x}$, and integration over a period $2\pi/L$ gives

$$\sum_{n=0}^{\infty} a_{mn} \frac{du_n}{dt} = p_m(u),\tag{7}$$

where $a_{mn} = \delta_{mn}$, $p_m(u) = \frac{L}{2\pi} \int_0^{2\pi/L} F(u) e^{-ik_m x} dx$. Apparently, the equation(5) with (6) and (7) reduces to a ordinary differential equations,

$$\frac{du_m}{dt} = p_m(u) \qquad for \quad m = 0, 1, 2, \cdots,$$
(8)

where p_m is a function of the u_m , for the KdV equation $F(u) = -[\alpha u u_x + \beta u_{xxx}]$, the (8) can be worked explicitly out,

$$\frac{du_m(t)}{dt} = i\alpha \sum_{\substack{k,l=0\\k+l=m}}^m u_k(t) lu_l(t) - i\beta m^3 u_m(t) \quad for \quad m = 0, 1, 2, \cdots.$$
(9)

Therefore, underlying the basis (4) the KdV equation becomes to a hierarchy of infinite first-order ordinary differential equations. It is easy to solve these ordinary differential equations by the standard integral method,

$$u_m(t) = e^{-i(-m\alpha u_0 + \beta m^3)t} \left[\int i\alpha \sum_{\substack{k,l=1\\k+l=m}}^{m-1} u_k(t') lu_l(t') e^{-i(-m\alpha u_0 + \beta m^3)t'} dt' + C_m \right],$$
(10)

where C_m is integration constant, and u_0 also any constant from $\frac{du_0(t)}{dt} = 0$. When taking $C_1 = 1$, $C_m(m \ge 2)$ for zero,

$$u(x,t) = \sum_{n=1}^{\infty} c_n e^{int} e^{-inx},$$

$$c_n = \frac{6}{n(n^2-1)} \sum_{\substack{k,l=1\\k+l=n}}^{n-1} c_k l c_l \qquad (n \ge 2),$$
(11)

here $c_1 = 1$, c_n is the coefficient. Although the base of the sub-space $\{e^{ik_nx}, n = 0, 1, 2, ...\}$ is incomplete, it is closed under the operation of the KdV equation. Under the closed condition and Sobolev theorem of embedding, one can show that the solution is the classical solution of the corresponding equation (Appendix A and B give the proof).

III. CONCLUSION

In conclusion, based on the incomplete basis $\{e^{-inx}, n = 0, 1, 2, \dots\}$, the complex version KdV equation(2) is discussed. Under the operation of the KdV equation, the basis is closed and forms an invariant set. The classical periodic solution was obtained in this invariant set. In Appendixes, it is showed that this solution is the classical solution of the KdV equation.

IV. APPENDIX A: THE PROOF OF THE CONVERGENCE OF THE PERIODIC SOLUTION

The convergence of the periodic solution

$$u(x,t) = \sum_{n=1}^{\infty} c_n e^{int} e^{-inx},$$

$$c_n = \frac{6}{n(n^2-1)} \sum_{\substack{k,l=1\\k+l=n}}^{n-1} c_k lc_l \qquad (n \ge 2),$$

here $c_1 = 1$, is discussed as follows: Considering the convergence of module of coefficients. It is obvious that

$$0 \le |c_n| \le \frac{6}{n(n^2 - 1)} \sum_{\substack{k, l = 1 \\ k+l = n}}^{n-1} |c_k| |c_l| \le 1 \text{ for } n = 1, 2, \cdots.$$

Then for arbitrary given $0 < \epsilon < 1/2$, one can show inductively that $|c_n| \leq \frac{A}{n^{3/2+\epsilon}}$, where $A \geq 1$ is a constant depending on ϵ . Note that

$$S_{\epsilon} = \sum_{n=1}^{\infty} \frac{1}{n^{3/2+\epsilon}} < \infty$$

and inequality

$$\sum_{k=1}^{n} \frac{1}{k^{\alpha}} \le \sum_{k=1}^{n} \int_{k}^{k+1} \frac{2}{x^{\alpha}} \, dx = 2 \int_{1}^{n+1} \frac{1}{x^{\alpha}} \, dx \le \frac{2(n+1)^{1-\alpha}}{1-\alpha}.$$

Without loss of generality, take $A \ge 64$. it is obvious that

$$B_{n} = \frac{6n^{2(3/2+\epsilon)}}{n(n^{2}-1)} \left\{ \sum_{l=1}^{\left[\frac{n-1}{2}\right]} \frac{1}{l^{1/2+\epsilon}(n-l)^{3/2+\epsilon}} + \sum_{l=\left[\frac{n-1}{2}+1\right]}^{n-1} \frac{1}{l^{1/2+\epsilon}(n-l)^{3/2+\epsilon}} \right\}$$
$$\leq \frac{6n^{2(3/2+\epsilon)}}{n(n^{2}-1)} \left\{ \frac{2^{3/2+\epsilon}}{n^{3/2+\epsilon}} \sum_{l=1}^{n-1} \frac{1}{l^{1/2+\epsilon}} + \frac{2^{1/2+\epsilon}}{n^{1/2+\epsilon}} \sum_{l=1}^{n-1} \frac{1}{(n-l)^{3/2+\epsilon}} \right\}$$
$$\leq \frac{6n^{2(3/2+\epsilon)}}{n(n^{2}-1)} \left\{ \frac{2^{5/2+\epsilon}(n+1)^{1/2-\epsilon}}{(1/2-\epsilon)n^{3/2+\epsilon}} + \frac{2^{1/2+\epsilon}S_{n}}{n^{1/2+\epsilon}} \right\}.$$

where $B_n = \frac{6n^{2(3/2+\epsilon)}}{n(n^2-1)} \sum_{l=1}^{n-1} \frac{1}{l^{1/2+\epsilon}(n-l)^{3/2+\epsilon}}$. It is obvious that $B_n \to 0$ at $n \to +\infty$, so there exists n_0 to make $B_n \leq 1$ at $n \geq n_0$. Thus take $A = max\{64, n_0^{3/2+\epsilon}\}$, for given A. It is natural to draw a conclusion that the periodic solution (11) is convergent.

V. APPENDIX B: THE PROOF OF THE CLASSICAL SOLUTION

The first proof, the solution(11) is weak solution.

For $0 \le t_1 < t_2 \le T$, $Q_{t_1,t_2} = \mathbf{R} \times [t_1,t_2]$. Definition: $u \in C(Q_T)$ is periodic weak solution with periodicity P of the KdV equation, if u satisfy

$$\int_{t_1}^{t_2} \int_0^P \{ u(-\varphi_t + \alpha \varphi_{xx} - \beta \varphi_{xxx}) - \frac{1}{2} u^2 \varphi_x \} dx dt = 0,$$

where $\varphi(x,t) \in C^{\infty}(Q_{t_1,t_2}), \varphi(\cdot,t_1) = \varphi(\cdot,t_2) = 0$, and φ is a periodic function with periodicity P for variable x. The following we take $P = 2\pi$. Assuming $R_N(x,t) = u(x,t) - \sum_{n=0}^N u_n(t)e^{-inx}$, from Appendix A, it reads $R_N(x,t) \to 0$, as $N \to \infty$. Definition I_N is

$$I_N = \int_{t_1}^{t_2} \int_0^{2\pi} \{\varphi(D_t u_N + \alpha D_{xx} u_N + \beta D_{xxx} u_N + \frac{1}{2} D_x(u_N^2))\} dxdt,$$

where subscript x, t denote differential, $u_N(x, t) = \sum_{n=0}^{N} u_n(t)e^{-inx}$. Apparently, I_N tends zero as $N \to \infty$. Thus it's show that the solution of the KdV equation is a weak solution by definition.

The second proof, the weak solution is classical solution.

Note the solution (11), following two conclusions have been showed: (i), $\forall A > 0, 0 < \epsilon < \frac{1}{2}, |c_n| \le \frac{A}{n^{\frac{3}{2}+\epsilon}} (n = 1, 2, \cdots).$ (ii), For arbitrary $\eta \in C_0^{\infty}(-\infty, \infty), \phi(x, t) \in C^{\infty}(\mathbf{R}^2)$, where $\phi(x, t)$ is a periodic function with respect to the variable x, u(x, t) satisfies

$$\int_{-\infty}^{\infty} dt \int_{0}^{2\pi} \{ u(x,t) [-(\eta^2 \phi)_t + \eta^2 (\alpha \phi_{xx} - \beta \phi_{xxx})] - \frac{1}{2} \eta^2 u^2 \phi_x \} dx = 0$$

Based on the conclusions (i) and (ii), a standard step can be applied, it is easy to show that the formal solution (11) is infinitely differentiable and must be the classical solutions of the KdV equation at t > 0. (The similar conclusion can be draw at t < 0.)

VI. ACKNOWLEDGMENTS

The author is grateful to Professors Ya-Zhe Chen and Ming-Liang Wang for valuable discussions. The corresponding author email address: yangkq@lzu.edu.cn (Yang Kongqing).

References

- [1] Benno.Fuchssteiner, Progress. Theo. Phys, 68 (1982) 1082.
- [2] W.Oevel, *Phys.Lett.A*, **94** (1983) 404.
- [3] R.Hirota, J.Satsuma, *Phys.Lett.A*, **85** (1981) 407.
- [4] R.Dodd, A.Fordy, *Phys.Lett.A*, **89** (1982) 168.
- [5] WenXiu.Ma, J.Phys.A, 31 (1998) 7585.
- [6] Prodyot.Kumar.Roy, Phys.Lett.A, 249 (1998) 55.
- [7] Keith A.Brueckner, Advances in Theoretical Physics (volume 1), Academic Press Inc.