

Riccati Solutions of Discrete Painlevé Equations with Weyl Group Symmetry of Type $E_8^{(1)}$

Mikio Murata, Hidetaka Sakai and Jin Yoneda
Graduate School of Mathematical Sciences,
The University of Tokyo,
Komaba, Tokyo 153-8914, Japan.

Abstract

We present a special solutions of the discrete Painlevé equations associated with $A_0^{(1)}$, $A_0^{(1)*}$ and $A_0^{(1)**}$ -surface. These solutions can be expressed by solutions of linear difference equations. Here the $A_0^{(1)}$ -surface discrete Painlevé equation is the most generic difference equation, as all discrete Painlevé equations can be obtained by its degeneration limit. These special solutions exist when the parameters of the discrete Painlevé equation satisfy a particular constraint. We consider that these special functions belong to the hypergeometric family although they seems to go beyond the known discrete and q -discrete hypergeometric functions. We also discuss the degeneration scheme of these solutions.

1 Introduction

One of the authors presented the list of all difference Painlevé equations from the view point of algebraic geometry. For a rational surface, we can obtain a birational representation of certain affine Weyl group as its symmetry. We regard a translation part of the symmetry as a difference system. Discrete Painlevé equations were classified on the basis of the types of rational surfaces, and some new equations were discovered by this classification ([12]).

Let X be a smooth projective surface. We denote by \mathcal{K}_X the canonical divisor class on X , and by $|- \mathcal{K}_X|$ the set of all positive divisors on X such that is linearly equivalent to $- \mathcal{K}_X$. We call an element of $|- \mathcal{K}_X|$ an anti-canonical divisor. Generalized Halphen surfaces are smooth projective rational surfaces with an anti-canonical divisor of canonical type. If $|- \mathcal{K}_X|$ has a unique divisor D , then X is classified according to the type R of D , where R is in the Table 2. The list of R can be obtained from the list of sublattice of $Q(E_8^{(1)})$ which are indecomposable and of affine type. We call a surface X an R -surface.

Table 3: The Painlevé differential equations

the type of surface	$D_4^{(1)}$	$D_5^{(1)}$	$D_6^{(1)}$	$D_7^{(1)}$	$D_8^{(1)}$	$E_6^{(1)}$	$E_7^{(1)}$	$E_8^{(1)}$
Painlevé equation	P_{VI}	P_V	$P_{III}^{D_6^{(1)}}$	$P_{III}^{D_7^{(1)}}$	$P_{III}^{D_8^{(1)}}$	P_{IV}	P_{II}	P_I

exists or not. For example, $A_3^{(1)}$ -surface has Weyl group symmetry of type $D_5^{(1)}$ but there is no surface with Weyl group symmetry of type $D_6^{(1)}$. So in this paper, we distinguish each discrete Painlevé equations by the rational surface on which the equation is defined, and call, for example, the $A_0^{(1)}$ -surface discrete Painlevé equation ($dP(A_0^{(1)})$).

We know many results about special solutions with respect to Painlevé differential equations whose parameters satisfy particular conditions ([7], etc.). These solutions are divided into algebraic solutions and what we call Riccati solutions. Riccati solutions are expressed by solutions of linear equations of the second order. For example, the sixth Painlevé equation P_{VI} has special solutions expressed in terms of the Gauss hypergeometric functions. The discrete Painlevé equations also possess Riccati solutions for particular values of the parameter. For example, the $A_3^{(1)}$ -surface discrete Painlevé equation (q - P_{VI}) has solutions expressed by the q -hypergeometric functions.

In [11], A. Ramani, B. Grammaticos, T. Tamizhmani and K. M. Tamizhmani presented special solutions of the $A_1^{(1)}$ -surface discrete Painlevé equation and its degenerations. But they did not mention about the discrete Painlevé equations corresponding to the surfaces of types $A_0^{(1)}$, $A_0^{(1)*}$ and $A_0^{(1)**}$.

In this paper, we obtain special solutions for these discrete Painlevé equations. We also discuss the degeneration scheme of these equations.

The paper is organized as follows:

In Sect. 2, we show a geometrical construction of the $A_0^{(1)}$ -surface discrete Painlevé equation and a Riccati solution of this equation in Theorem 2.

In Sect. 3, we obtain the $A_0^{(1)*}$ -surface discrete Painlevé equation in geometric approach, and a special solution of this equation. And we show a degeneration scheme of these equations in Theorem 3, 5.

In Sect. 4, we obtain the $A_0^{(1)**}$ -surface discrete Painlevé equation in geometric approach, and a special solution of this equation in a similar way. And we show a degeneration scheme of these equations in Theorem 8.

In Sect. 5, we show as Theorem 9 a degeneration scheme between a special solution of the $A_0^{(1)*}$ -surface discrete Painlevé equation in Sect. 3 and a special solution of the $A_1^{(1)}$ -surface discrete Painlevé equation.

2 Elliptic type

2.1 Discrete Painlevé equation

We present the $A_0^{(1)}$ -surface discrete Painlevé equation ($dP(A_0^{(1)})$). This system is equivalent to ell. P derived in [12]. In this paper, all 2×2 matrices represent $\text{PGL}(2)$ -action on \mathbb{P}^1 , i.e., $w = \begin{pmatrix} a & b \\ c & d \end{pmatrix} z$ means $w = \frac{az+b}{cz+d}$.

The $A_0^{(1)}$ -surface discrete Painlevé equation are the following difference system for unknown functions $f(t)$, $g(t)$:

$$(2.1) \quad \begin{aligned} \bar{g} &= M \left(f, c_7, c_8, t - \frac{1}{4} \sum_{i=1}^6 c_i \right) M \left(f, c_5, c_6, t - \frac{1}{4} \sum_{i=1}^4 c_i \right) \\ &\quad \cdot M \left(f, c_3, c_4, t - \frac{1}{4}(c_1 + c_2) \right) M(f, c_1, c_2, t) g, \end{aligned}$$

$$(2.2) \quad \begin{aligned} \underline{f} &= M \left(g, d_7, d_8, t - \frac{1}{4} \sum_{i=1}^6 d_i \right) M \left(g, d_5, d_6, t - \frac{1}{4} \sum_{i=1}^4 d_i \right) \\ &\quad \cdot M \left(g, d_3, d_4, t - \frac{1}{4}(d_1 + d_2) \right) M(g, d_1, d_2, t) f, \end{aligned}$$

where $\bar{g} = g(t + \lambda)$, $\underline{f} = f(t - \lambda)$ and

$$(2.3) \quad \begin{aligned} &M(h, \kappa_1, \kappa_2, s) \\ &= \begin{pmatrix} -\wp(2s - \frac{-\kappa_1 + \kappa_2}{2}) & \wp(2s - \frac{\kappa_1 - \kappa_2}{2}) \\ -1 & 1 \end{pmatrix} \\ &\cdot \begin{pmatrix} (h - \wp(\kappa_2))(\wp(2s) - \wp(2s - \kappa_2))(\wp(2s - \frac{\kappa_1 + \kappa_2}{2}) - \wp(2s - \frac{\kappa_1 - \kappa_2}{2})) & 0 \\ 0 & (h - \wp(\kappa_1))(\wp(2s) - \wp(2s - \kappa_1))(\wp(2s - \frac{\kappa_1 + \kappa_2}{2}) - \wp(2s - \frac{-\kappa_1 + \kappa_2}{2})) \end{pmatrix} \\ &\cdot \begin{pmatrix} 1 & -\wp(2s - \kappa_1) \\ 1 & -\wp(2s - \kappa_2) \end{pmatrix}. \end{aligned}$$

Here b_i ($i = 1, \dots, 8$) are constant parameters and set $\lambda = \frac{1}{2} \sum_{i=1}^8 b_i$, $c_i = b_i + t$, $d_i = t - b_i$. Note that we will regard $(f(t), g(t))$ as inhomogeneous coordinates of $\mathbb{P}^1 \times \mathbb{P}^1$.

We derive this discrete equation as a translation of $W(E_8^{(1)})$ again.

We construct $A_0^{(1)}$ -surface by blowing up $\mathbb{P}^1 \times \mathbb{P}^1$ at eight points p_i ($i = 1, \dots, 8$). For generic eight points there is an elliptic curve which passes through these eight points. We parameterize these eight points and the curve as follows:

$$(2.4) \quad (f + g + \wp(2t))(4\wp(2t)fg - g_3) = \left(fg + \wp(2t)(f + g) + \frac{g_2}{4} \right)^2,$$

$$(2.5) \quad p_i: (\wp(b_i + t), \wp(t - b_i)) \quad (i = 1, \dots, 8),$$

Remark 2.1. We can parameterize an isomorphism class of surfaces by using the period mapping. The period mapping maps the elements of the second homology to \mathbb{C} .

Let ω be a meromorphic 2-form on X with $\text{div}(\omega) = -D$. Then ω determines the period mapping $\hat{\chi}: H_2(X - D, \mathbb{Z}) \rightarrow \mathbb{C}$ which sends $\Gamma \in H_2(X - D, \mathbb{Z})$ to $\int_{\Gamma} \omega$.

Now, there exists the short exact sequence:

$$0 \rightarrow H_1(D, \mathbb{Z}) \rightarrow H_2(X - D, \mathbb{Z}) \rightarrow Q(E_8^{(1)}) \rightarrow 0,$$

where $Q(E_8^{(1)}) = \sum_{i=0}^8 \mathbb{Z}\alpha_i$ is the root lattice of type $E_8^{(1)}$. So we obtain the mapping

$$\chi: Q(E_8^{(1)}) \rightarrow \mathbb{C} \quad \text{mod } \hat{\chi}(H_1(D, \mathbb{Z}))$$

through the period mapping $\hat{\chi}$. In the case, the parameterization is the following.

$$(2.6) \quad \begin{aligned} \chi(\alpha_1) &= -4t, & \chi(\alpha_2) &= b_1 + b_2 + 2t, & \chi(\alpha_i) &= b_i - b_{i-1} \quad (i = 3, \dots, 7), \\ \chi(\alpha_8) &= b_2 - b_1, & \chi(\alpha_0) &= b_8 - b_7. \end{aligned}$$

Here $Q(E_8^{(1)})$ is realized in $\text{Pic}(X) = H_2(X, \mathbb{Z})$. And α_i 's are represented by the elements of the Picard group as follows:

$$(2.7) \quad \begin{aligned} \alpha_1 &= H_1 - H_0, & \alpha_2 &= H_0 - E_1 - E_2, & \alpha_i &= E_{i-1} - E_i \quad (i = 3, \dots, 7), \\ \alpha_8 &= E_1 - E_2, & \alpha_0 &= E_7 - E_8. \end{aligned}$$

We denote the total transform of $f = \text{constant}$, (or $g = \text{constant}$) on X by H_0 (or H_1 respectively) and the total transform of the point p_i by E_i . The Picard group $\text{Pic}(X)$ and the canonical divisor \mathcal{K}_X are

$$\text{Pic}(X) = \mathbb{Z}H_0 + \mathbb{Z}H_1 + \sum_{i=1}^8 \mathbb{Z}E_i, \quad \mathcal{K}_X = -2H_0 - 2H_1 + \sum_{i=1}^8 E_i,$$

where the intersection numbers of pairs of base elements are

$$H_i \cdot H_j = 1 - \delta_{i,j}, \quad E_i \cdot E_j = -\delta_{i,j}, \quad H_i \cdot E_j = 0, \quad \text{where } \delta_{i,j} = \begin{cases} 1 & i = j, \\ 0 & i \neq j. \end{cases}$$

□

Generators of affine Weyl group $W(E_8^{(1)}) = \langle w_i \ (i = 0, 1, \dots, 8) \rangle$ act on these coordinates

and parameters. We give a representation of these actions in order to construct $dP(A_0^{(1)})$.

$$\begin{aligned}
w_2: & \left(\begin{array}{cccc} b_1 & b_2 & b_3 & b_4 \\ b_5 & b_6 & b_7 & b_8 \end{array}, t, g \right) \\
& \mapsto \left(\begin{array}{cccc} b_1 - 3\frac{2t+b_1+b_2}{4} & b_2 - 3\frac{2t+b_1+b_2}{4} & b_3 + \frac{2t+b_1+b_2}{4} & b_4 + \frac{2t+b_1+b_2}{4} \\ b_5 + \frac{2t+b_1+b_2}{4} & b_6 + \frac{2t+b_1+b_2}{4} & b_7 + \frac{2t+b_1+b_2}{4} & b_8 + \frac{2t+b_1+b_2}{4} \end{array}, t - \frac{2t+b_1+b_2}{4}, \tilde{g} \right), \\
w_1: & (t, f, g) \mapsto (-t, g, f), \quad w_i: (b_{i-1}, b_i) \mapsto (b_i, b_{i-1}) \quad (i = 3, \dots, 7), \\
w_8: & (b_1, b_2) \mapsto (b_2, b_1), \quad w_0: (b_7, b_8) \mapsto (b_8, b_7),
\end{aligned}$$

where \tilde{g} is given by

$$\begin{aligned}
& \frac{\tilde{g} - \wp\left(2t - \frac{b_1-b_2}{2}\right)}{\tilde{g} - \wp\left(2t - \frac{-b_1+b_2}{2}\right)} \\
& = \frac{f - \wp(b_2 + t) \wp\left(t - \frac{b_1+b_2}{2}\right) - \wp\left(2t - \frac{b_1-b_2}{2}\right) \wp(2t) - \wp(t - b_2) g - \wp(t - b_1)}{f - \wp(b_1 + t) \wp\left(t - \frac{b_1+b_2}{2}\right) - \wp\left(2t - \frac{-b_1+b_2}{2}\right) \wp(2t) - \wp(t - b_1) g - \wp(t - b_2)}.
\end{aligned}$$

Notice that we can rewrite the action of w_2 into the following form:

$$w_2: (c_1, c_2, t, g) \mapsto (-c_2, -c_1, t - \frac{c_1+c_2}{4}, M(f, c_1, c_2, t)g),$$

where we use the notation $c_i = b_i + t$ and $M(f, c_1, c_2, t)$ is defined by (2.3).

By taking a translation of $W(E_8^{(1)})$, we obtain a nonlinear difference equation. The translation can be described by a product of simple reflections w_i 's.

$$\begin{aligned}
dP(A_0^{(1)}) & = w_1 \circ w_2 \circ w_3 \circ w_8 \circ w_4 \circ w_3 \circ w_2 \circ w_5 \circ w_4 \circ w_3 \circ w_8 \circ w_6 \circ w_5 \circ w_4 \circ w_3 \circ \\
(2.8) \quad & \circ w_2 \circ w_7 \circ w_6 \circ w_5 \circ w_4 \circ w_3 \circ w_8 \circ w_0 \circ w_7 \circ w_6 \circ w_5 \circ w_4 \circ w_3 \circ w_2 \circ w_1 \circ \\
& \circ w_2 \circ w_3 \circ w_4 \circ w_5 \circ w_6 \circ w_7 \circ w_0 \circ w_8 \circ w_3 \circ w_4 \circ w_5 \circ w_6 \circ w_7 \circ w_2 \circ w_3 \circ \\
& \circ w_4 \circ w_5 \circ w_6 \circ w_8 \circ w_3 \circ w_4 \circ w_5 \circ w_2 \circ w_3 \circ w_4 \circ w_8 \circ w_3 \circ w_2 :
\end{aligned}$$

$$(2.9) \quad (b_i, t, f, g) \mapsto (b_i, t + \lambda, \bar{f}, \bar{g}) \quad (i = 1, \dots, 8), \quad \lambda = \frac{1}{2} \sum_{i=1}^8 b_i,$$

where mappings of f, g are defined by (2.1), (2.2).

Remark 2.2. In [12], we obtain $A_0^{(1)}$ -surface by blowing up \mathbb{P}^2 with the centers at nine points.

$$(2.10) \quad y^2 z = 4x^3 - g_2 x^2 z - g_3 z^3,$$

$$\begin{aligned}
(2.11) \quad & p_i: (\wp(\theta_i) : \wp'(\theta_i) : 1) \quad (i = 1, \dots, 9), \\
& \chi(\alpha_i) = \theta_{i+1} - \theta_i \quad (i = 1, \dots, 7), \quad \chi(\alpha_8) = \theta_1 + \theta_2 + \theta_3, \quad \chi(\alpha_0) = \theta_9 - \theta_8.
\end{aligned}$$

Both parameters and coordinates correspond as follows:

$$(2.12) \quad b_1 = -\frac{3}{4}(\theta_1 + \theta_2), \quad b_i = \theta_{i+1} + \frac{1}{4}(\theta_1 + \theta_2) \quad (i = 2, \dots, 8), \quad t = \frac{1}{4}(\theta_1 - \theta_2),$$

$$(2.13) \quad f = \frac{(4\wp(\frac{\theta_1}{2})^3 - 3g_2\wp(\frac{\theta_1}{2}) - 4g_3)x - 2\wp(\frac{\theta_1}{2})\wp'(\frac{\theta_1}{2})y - \left(g_2\wp(\frac{\theta_1}{2})^2 + 6g_3\wp(\frac{\theta_1}{2}) + \frac{g_2^2}{4}\right)z}{- (12\wp(\frac{\theta_1}{2})^2 - g_2)x - 2\wp'(\frac{\theta_1}{2})y + (4\wp(\frac{\theta_1}{2})^3 + g_2\wp(\frac{\theta_1}{2}) + 2g_3)z},$$

$$(2.14) \quad g = \frac{(4\wp(\frac{\theta_2}{2})^3 - 3g_2\wp(\frac{\theta_2}{2}) - 4g_3)x - 2\wp(\frac{\theta_2}{2})\wp'(\frac{\theta_2}{2})y - \left(g_2\wp(\frac{\theta_2}{2})^2 + 6g_3\wp(\frac{\theta_2}{2}) + \frac{g_2^2}{4}\right)z}{- (12\wp(\frac{\theta_2}{2})^2 - g_2)x - 2\wp'(\frac{\theta_2}{2})y + (4\wp(\frac{\theta_2}{2})^3 + g_2\wp(\frac{\theta_2}{2}) + 2g_3)z}.$$

We note that the next formula holds for any θ, φ .

$$(2.15) \quad \wp(\theta + \varphi) = \frac{(4\wp(\varphi)^3 - 3g_2\wp(\varphi) - 4g_3)\wp(\theta) - 2\wp(\varphi)\wp'(\varphi)\wp'(\theta) - \left(g_2\wp(\varphi)^2 + 6g_3\wp(\varphi) + \frac{g_2^2}{4}\right)}{- (12\wp(\varphi)^2 - g_2)\wp(\theta) - 2\wp'(\varphi)\wp'(\theta) + (4\wp(\varphi)^3 + g_2\wp(\varphi) + 2g_3)}.$$

This formula is a kind of additive formula of \wp -function. \square

Note that points on the elliptic curve (2.4) move to points on this elliptic curve.

Proposition 1. $dP(A_0^{(1)})$ has the following trivial solution:

$$(2.16) \quad f = \wp(q + 2t^2/\lambda + t), \quad g = \wp(-q - 2t^2/\lambda + t),$$

where q is a constant determined by initial condition.

Proof. We suppose that the solution's form is $f = \wp(p + t)$, $g = \wp(t - p)$, where $p = p(t)$. We input them into (2.1), (2.2). We note that the following identity holds for arbitrary c ,

$$(2.17) \quad \frac{\wp\left(2t - \frac{c_1+c_2}{2} - c\right) - \wp\left(2t - \frac{c_1+c_2}{2} + c_2\right)}{\wp\left(2t - \frac{c_1+c_2}{2} - c\right) - \wp\left(2t - \frac{c_1+c_2}{2} + c_1\right)} \frac{\wp(c) - \wp(c_1)}{\wp(c) - \wp(c_2)} \frac{\wp(2t - c) - \wp(2t - c_2)}{\wp(2t - c) - \wp(2t - c_1)} \\ = \frac{\wp\left(2t - \frac{c_1+c_2}{2}\right) - \wp\left(2t - \frac{c_1+c_2}{2} + c_2\right)}{\wp\left(2t - \frac{c_1+c_2}{2}\right) - \wp\left(2t - \frac{c_1+c_2}{2} + c_1\right)} \frac{\wp(2t) - \wp(2t - c_2)}{\wp(2t) - \wp(2t - c_1)},$$

because both sides equal to

$$\frac{\sigma\left(2t - \frac{c_1+c_2}{2} + c_1\right)^2 \sigma(c_2)^2 \sigma(2t - c_1)^2}{\sigma\left(2t - \frac{c_1+c_2}{2} + c_2\right)^2 \sigma(c_1)^2 \sigma(2t - c_2)^2}.$$

Since (2.17) is equivalent to

$$(2.18) \quad \wp\left(2t - \frac{c_1+c_2}{2} - c\right) = M(\wp(c), c_1, c_2, t)\wp(2t - c),$$

we obtain

$$(2.19) \quad \wp(t + \lambda - \bar{p}) = \wp(-3t - \lambda - p),$$

$$(2.20) \quad \wp(t - \lambda + \underline{p}) = \wp(-3t + \lambda + p).$$

The compatibility condition of them leads one difference equation,

$$(2.21) \quad \bar{p} = p + 4t + 2\lambda.$$

So that,

$$(2.22) \quad p = q + 2t^2/\lambda.$$

where q is a constant determined by initial condition. □

2.2 Linear equation

We derive the following theorem in this section.

Theorem 2. *A solution of the following system of equations is written by a solution of linear equation;*

$$(2.23) \quad \begin{vmatrix} fg & g & f & 1 \\ \wp(b_1 + t)\wp(t - b_1) & \wp(t - b_1) & \wp(b_1 + t) & 1 \\ \wp(b_3 + t)\wp(t - b_3) & \wp(t - b_3) & \wp(b_3 + t) & 1 \\ \wp(b_5 + t)\wp(t - b_5) & \wp(t - b_5) & \wp(b_5 + t) & 1 \end{vmatrix} = 0,$$

$$(2.24) \quad \begin{vmatrix} f\bar{g} & \bar{g} & f & 1 \\ \wp(b_8 + \bar{t})\wp(\bar{t} - b_8) & \wp(\bar{t} - b_8) & \wp(b_8 + \bar{t}) & 1 \\ \wp(b_6 + \bar{t})\wp(\bar{t} - b_6) & \wp(\bar{t} - b_6) & \wp(b_6 + \bar{t}) & 1 \\ \wp(b_4 + \bar{t})\wp(\bar{t} - b_4) & \wp(\bar{t} - b_4) & \wp(b_4 + \bar{t}) & 1 \end{vmatrix} = 0,$$

where $\bar{t} = t + \lambda$. When $b_1 + b_3 + b_5 + b_7 = 0$, a solution of this system is a special solution of $dP(A_0^{(1)})$ (2.1), (2.2). □

Transforming these equations,

$$(2.25) \quad \bar{g} = \frac{\begin{vmatrix} \wp(b_8 + \bar{t})\wp(\bar{t} - b_8) & \wp(\bar{t} - b_8) & 1 \\ \wp(b_6 + \bar{t})\wp(\bar{t} - b_6) & \wp(\bar{t} - b_6) & 1 \\ \wp(b_4 + \bar{t})\wp(\bar{t} - b_4) & \wp(\bar{t} - b_4) & 1 \end{vmatrix} f + \begin{vmatrix} \wp(b_8 + \bar{t})\wp(\bar{t} - b_8) & \wp(b_8 + \bar{t}) & \wp(\bar{t} - b_8) \\ \wp(b_6 + \bar{t})\wp(\bar{t} - b_6) & \wp(b_6 + \bar{t}) & \wp(\bar{t} - b_6) \\ \wp(b_4 + \bar{t})\wp(\bar{t} - b_4) & \wp(b_4 + \bar{t}) & \wp(\bar{t} - b_4) \end{vmatrix}}{\begin{vmatrix} \wp(b_8 + \bar{t}) & \wp(\bar{t} - b_8) & 1 \\ \wp(b_6 + \bar{t}) & \wp(\bar{t} - b_6) & 1 \\ \wp(b_4 + \bar{t}) & \wp(\bar{t} - b_4) & 1 \end{vmatrix} f + \begin{vmatrix} \wp(b_8 + \bar{t})\wp(\bar{t} - b_8) & \wp(b_8 + \bar{t}) & 1 \\ \wp(b_6 + \bar{t})\wp(\bar{t} - b_6) & \wp(b_6 + \bar{t}) & 1 \\ \wp(b_4 + \bar{t})\wp(\bar{t} - b_4) & \wp(b_4 + \bar{t}) & 1 \end{vmatrix}},$$

(2.26)

$$f = \frac{\begin{vmatrix} \wp(b_1+t)\wp(t-b_1) & \wp(b_1+t) & 1 \\ \wp(b_3+t)\wp(t-b_3) & \wp(b_3+t) & 1 \\ \wp(b_5+t)\wp(t-b_5) & \wp(b_5+t) & 1 \end{vmatrix} g + \begin{vmatrix} \wp(b_1+t)\wp(t-b_1) & \wp(t-b_1) & \wp(b_1+t) \\ \wp(b_3+t)\wp(t-b_3) & \wp(t-b_3) & \wp(b_3+t) \\ \wp(b_5+t)\wp(t-b_5) & \wp(t-b_5) & \wp(b_5+t) \end{vmatrix}}{\begin{vmatrix} \wp(t-b_1) & \wp(b_1+t) & 1 \\ \wp(t-b_3) & \wp(b_3+t) & 1 \\ \wp(t-b_5) & \wp(b_5+t) & 1 \end{vmatrix} g + \begin{vmatrix} \wp(b_1+t)\wp(t-b_1) & \wp(t-b_1) & 1 \\ \wp(b_3+t)\wp(t-b_3) & \wp(t-b_3) & 1 \\ \wp(b_5+t)\wp(t-b_5) & \wp(t-b_5) & 1 \end{vmatrix}}.$$

Eliminating f , we can obtain a difference equation of the first order with respect to the variable g . When g is expressed by homogeneous coordinate ($g = \frac{g_1}{g_2}$), $\bar{g} = \frac{Ag+B}{Cg+D}$ leads to $\frac{\bar{g}_1}{\bar{g}_2} = \frac{Ag_1+Bg_2}{Cg_1+Dg_2}$. The solution g is represented by solution of linear equation:

$$\begin{pmatrix} \bar{g}_1 \\ \bar{g}_2 \end{pmatrix} = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \begin{pmatrix} g_1 \\ g_2 \end{pmatrix}.$$

Now we demonstrate Theorem 2. On the condition $b_1 + b_3 + b_5 + b_7 = 0$, we consider the curve $I = 0$, where

(2.27)

$$I = \begin{vmatrix} fg & g & f & 1 \\ \wp(b_1+t)\wp(t-b_1) & \wp(t-b_1) & \wp(b_1+t) & 1 \\ \wp(b_3+t)\wp(t-b_3) & \wp(t-b_3) & \wp(b_3+t) & 1 \\ \wp(b_5+t)\wp(t-b_5) & \wp(t-b_5) & \wp(b_5+t) & 1 \end{vmatrix} = \begin{vmatrix} fg & g & f & 1 \\ \wp(c_1)\wp(2t-c_1) & \wp(2t-c_1) & \wp(c_1) & 1 \\ \wp(c_3)\wp(2t-c_3) & \wp(2t-c_3) & \wp(c_3) & 1 \\ \wp(c_5)\wp(2t-c_5) & \wp(2t-c_5) & \wp(c_5) & 1 \end{vmatrix}.$$

Remark 2.3. On this condition, p_i ($i = 1, 3, 5, 7$) lie on $I = 0$, because the following formula holds,

$$(2.28) \quad \begin{vmatrix} \wp(b_1+t)\wp(t-b_1) & \wp(t-b_1) & \wp(b_1+t) & 1 \\ \wp(b_3+t)\wp(t-b_3) & \wp(t-b_3) & \wp(b_3+t) & 1 \\ \wp(b_5+t)\wp(t-b_5) & \wp(t-b_5) & \wp(b_5+t) & 1 \\ \wp(b_7+t)\wp(t-b_7) & \wp(t-b_7) & \wp(b_7+t) & 1 \end{vmatrix} = 0.$$

The curve $I = 0$ has the divisor class $C = H_0 + H_1 - E_1 - E_3 - E_5 - E_7$, and the self-intersection number of C is -2 . In this case, we can restrict affine Weyl group action on the curve $I = 0$, which is isomorphic to \mathbb{P}^1 , and the translation is automorphism on \mathbb{P}^1 namely homographic transformation. \square

Calculating w_2 acting I , we set \tilde{I} as follows:

$$(2.29) \quad \tilde{I} = \begin{vmatrix} f\tilde{g} & \tilde{g} & f & 1 \\ \wp(c_2)\wp(2\tilde{t}+c_2) & \wp(2\tilde{t}+c_2) & \wp(c_2) & 1 \\ \wp(c_3)\wp(2\tilde{t}-c_3) & \wp(2\tilde{t}-c_3) & \wp(c_3) & 1 \\ \wp(c_5)\wp(2\tilde{t}-c_5) & \wp(2\tilde{t}-c_5) & \wp(c_5) & 1 \end{vmatrix},$$

where $\tilde{t} = t - \frac{c_1+c_2}{4}$. Then

$$(2.30) \quad \tilde{I} = I \left(\frac{\wp(2\tilde{t}) - \wp(2\tilde{t} + c_2)}{\wp(2\tilde{t}) - \wp(2\tilde{t} + c_1)} \frac{\wp(2t) - \wp(2t - c_2)}{\wp(2t) - \wp(2t - c_1)} \right)^2 (f - \wp(c_2))(\wp(c_2) - \wp(c_3))(\wp(c_2) - \wp(c_5)) \cdot (\wp(2t - c_1) - \wp(2t - c_2))(\wp(2\tilde{t} + c_2) - \wp(2\tilde{t} + c_1))^2 \cdot \left(\begin{array}{c|c|c|c} f & g & 1 & \\ \wp(c_1) & \wp(2t - c_1) & 1 & \\ \wp(c_2) & \wp(2t - c_2) & 1 & \end{array} \middle| \begin{array}{c|c|c|c} \wp(c_1) & \wp(2t - c_1) & 1 & \\ \wp(c_2) & \wp(2t - c_2) & 1 & \\ \wp(c_3) & \wp(2t - c_3) & 1 & \end{array} \middle| \begin{array}{c|c|c|c} \wp(c_1) & \wp(2t - c_1) & 1 & \\ \wp(c_2) & \wp(2t - c_2) & 1 & \\ \wp(c_5) & \wp(2t - c_5) & 1 & \end{array} \right)^{-1}.$$

Therefore $I = 0 \implies \tilde{I} = 0$. Similarly we consider the curve $\mathcal{I} = 0$,

$$(2.31) \quad \mathcal{I} = \begin{vmatrix} f\bar{g} & \bar{g} & f & 1 \\ \wp(b_8 + t)\wp(\bar{t} - b_8) & \wp(\bar{t} - b_8) & \wp(b_8 + t) & 1 \\ \wp(b_6 + t)\wp(\bar{t} - b_6) & \wp(\bar{t} - b_6) & \wp(b_6 + t) & 1 \\ \wp(b_4 + t)\wp(\bar{t} - b_4) & \wp(\bar{t} - b_4) & \wp(b_4 + t) & 1 \end{vmatrix}.$$

Then $I = 0 \implies \mathcal{I} = 0$ holds, and we consider the curve $\bar{I} = 0$,

$$(2.32) \quad \bar{I} = \begin{vmatrix} \bar{f}\bar{g} & \bar{g} & \bar{f} & 1 \\ \wp(b_1 + \bar{t})\wp(\bar{t} - b_1) & \wp(\bar{t} - b_1) & \wp(b_1 + \bar{t}) & 1 \\ \wp(b_3 + \bar{t})\wp(\bar{t} - b_3) & \wp(\bar{t} - b_3) & \wp(b_3 + \bar{t}) & 1 \\ \wp(b_5 + \bar{t})\wp(\bar{t} - b_5) & \wp(\bar{t} - b_5) & \wp(b_5 + \bar{t}) & 1 \end{vmatrix}.$$

Then $I = 0 \implies \bar{I} = 0$. This means that f, g which satisfy $I = 0$ are a special solution on the condition $b_1 + b_3 + b_5 + b_7 = 0$.

3 Multiplicative type

We can obtain $A_0^{(1)*}$ -surface from $A_0^{(1)}$ -surface by degeneration. By the same process the $A_0^{(1)*}$ -surface discrete Painlevé equation and the linear equation also can be obtained.

3.1 Discrete Painlevé equation

We discuss about the following theorem in this section. The equations obtained by the degeneration in the theorem coincide with the equations in [8].

Put $g_2 = \frac{4}{3}(1 + 3\varepsilon^2)$, $g_3 = -\frac{8}{27}(1 - 9\varepsilon^2)$ in the $A_0^{(1)}$ -surface discrete Painlevé equation (2.1), (2.2) and let ε tend to 0. Then we obtain the $A_0^{(1)*}$ -surface discrete Painlevé equation. Moreover the change of the variables and parameters $e^{2t} = t_1$, $e^{2\lambda} = \lambda_1$, $f = \frac{1}{3} \frac{f_1+10}{f_1-2}$, $g = \frac{1}{3} \frac{g_1+10}{g_1-2}$, $e^{2b_i} = \beta_i$ yields the expression in the following theorem. In the expression we replace

again t_1 by t and $\lambda_1, f_1, g_1, \beta_i$ by λ, f, g, b_i , respectively. For the sake of simplification of notation, the replacement process will be written as follows:

$$e^{2t} \rightarrow t, \quad e^{2\lambda} \rightarrow \lambda, \quad f \rightarrow \frac{1}{3} \frac{f+10}{f-2}, \quad g \rightarrow \frac{1}{3} \frac{g+10}{g-2}, \quad e^{2b_i} \rightarrow b_i.$$

These are summarized as follows:

Theorem 3. *Make the substitution: $g_2 = \frac{4}{3}(1 + 3\varepsilon^2)$, $g_3 = -\frac{8}{27}(1 - 9\varepsilon^2)$ in $dP(A_0^{(1)})$ (2.1), (2.2). Take the limit $\varepsilon \rightarrow 0$. Moreover by the change of variables and parameters: $e^{2t} \rightarrow t, e^{2\lambda} \rightarrow \lambda, f \rightarrow \frac{1}{3} \frac{f+10}{f-2}, g \rightarrow \frac{1}{3} \frac{g+10}{g-2}, e^{2b_i} \rightarrow b_i$, we obtain $dP(A_0^{(1)*})$:*

$$(3.1) \quad \frac{(\bar{g}t^2\lambda - f)(gt^2 - f) - (t^4\lambda^2 - 1)(t^4 - 1)}{(\bar{g}/(t^2\lambda) - f)(g/t^2 - f) - (1 - 1/(t^4\lambda^2))(1 - 1/t^4)}$$

$$= \lambda^2(f^4 - m_1tf^3 + (m_2t^2 - 3 - m_8t^8)f^2 + (m_7t^7 - m_3t^3 + 2m_1t)f$$

$$+ m_8t^8 - m_6t^6 + m_4t^4 - m_2t^2 + 1)$$

$$\cdot (m_8f^4 - m_7f^3/t + (m_6/t^2 - 3m_8 - 1/t^8)f^2 + (m_1/t^7 - m_5/t^3 + 2m_7/t)f$$

$$+ 1/t^8 - m_2/t^6 + m_4/t^4 - m_6/t^2 + m_8)^{-1},$$

$$(3.2) \quad \frac{(ft^2/\lambda - g)(ft^2 - g) - (t^4/\lambda^2 - 1)(t^4 - 1)}{(f\lambda/t^2 - g)(f/t^2 - g) - (1 - \lambda^2/t^4)(1 - 1/t^4)}$$

$$= \frac{1}{\lambda^2}(m_8g^4 - m_7tg^3 + (m_6t^2 - 3m_8 - t^8)g^2 + (m_1t^7 - m_5t^3 + 2m_7t)g$$

$$+ t^8 - m_2t^6 + m_4t^4 - m_6t^2 + m_8)$$

$$\cdot (g^4 - m_1g^3/t + (m_2/t^2 - 3 - m_8/t^8)g^2 + (m_7/t^7 - m_3/t^3 + 2m_1/t)g$$

$$+ m_8/t^8 - m_6/t^6 + m_4/t^4 - m_2/t^2 + 1)^{-1},$$

where $\bar{g} = g(t\lambda)$, $\underline{f} = f(t/\lambda)$, $\lambda = \sqrt{\prod_{i=1}^8 b_i}$, m_i is the i -th elementary symmetric function of b_j ($j = 1, \dots, 8$). \square

The above theorem shows that we can obtain $dP(A_0^{(1)*})$ from $dP(A_0^{(1)})$. But, for readers' convenience, we describe geometrical construction of this discrete equation similar to the previous section.

We construct $A_0^{(1)*}$ -surface by blowing up $\mathbb{P}^1 \times \mathbb{P}^1$ at eight points. These eight points and a curve which these points lie on are as follows:

$$(3.3) \quad f^2 + g^2 - \left(t^2 + \frac{1}{t^2}\right)fg + \left(t^2 - \frac{1}{t^2}\right)^2 = 0,$$

$$(3.4) \quad p_i: \left(b_it + \frac{1}{b_it}, \frac{t}{b_i} + \frac{b_i}{t}\right) \quad (i = 1, \dots, 8),$$

$$(3.5) \quad e^{2\chi(\alpha_1)} = t^{-4}, \quad e^{2\chi(\alpha_2)} = b_1b_2t^2, \quad e^{2\chi(\alpha_i)} = b_i/b_{i-1} \quad (i = 3, \dots, 7),$$

$$e^{2\chi(\alpha_8)} = b_2/b_1, \quad e^{2\chi(\alpha_0)} = b_8/b_7.$$

Generators of affine Weyl group $W(E_8^{(1)}) = \langle w_i \ (i = 0, 1, \dots, 8) \rangle$ act on these coordinates and parameters. We give a representation of these actions in order to construct $dP(A_0^{(1)*})$.

$$\begin{aligned}
w_2: & \left(\begin{array}{cccc} b_1 & b_2 & b_3 & b_4 \\ b_5 & b_6 & b_7 & b_8 \end{array}, t, g \right) \\
& \mapsto \left(\begin{array}{cccc} b_1/(\sqrt[4]{b_1 b_2 t^2})^3 & b_2/(\sqrt[4]{b_1 b_2 t^2})^3 & b_3 \sqrt[4]{b_1 b_2 t^2} & b_4 \sqrt[4]{b_1 b_2 t^2} \\ b_5 \sqrt[4]{b_1 b_2 t^2} & b_6 \sqrt[4]{b_1 b_2 t^2} & b_7 \sqrt[4]{b_1 b_2 t^2} & b_8 \sqrt[4]{b_1 b_2 t^2} \end{array}, t/\sqrt[4]{b_1 b_2 t^2}, \tilde{g} \right), \\
w_1: & (t, f, g) \mapsto (1/t, g, f), \quad w_i: (b_{i-1}, b_i) \mapsto (b_i, b_{i-1}) \quad (i = 3, \dots, 7), \\
w_8: & (b_1, b_2) \mapsto (b_2, b_1), \quad w_0: (b_7, b_8) \mapsto (b_8, b_7),
\end{aligned}$$

where \tilde{g} is given by

$$\frac{\tilde{g} - \left(\sqrt{\frac{b_2}{b_1} t^2} + \sqrt{\frac{b_1}{b_2} \frac{1}{t^2}} \right)}{\tilde{g} - \left(\sqrt{\frac{b_1}{b_2} t^2} + \sqrt{\frac{b_2}{b_1} \frac{1}{t^2}} \right)} = \frac{f - \left(b_2 t + \frac{1}{b_2 t} \right) g - \left(\frac{t}{b_1} + \frac{b_1}{t} \right)}{f - \left(b_1 t + \frac{1}{b_1 t} \right) g - \left(\frac{t}{b_2} + \frac{b_2}{t} \right)}.$$

Notice that we can rewrite the action of w_2 for $c_i = b_i t$ into the following form:

$$w_2: (c_1, c_2, t, g) \mapsto (1/c_2, 1/c_1, t/\sqrt[4]{c_1 c_2}, \tilde{g}).$$

Here we put Γ and $\tilde{\Gamma}$ as

$$\Gamma = \frac{gt^2 - f}{t^4 - 1}, \quad \tilde{\Gamma} = \frac{\tilde{g} \frac{t^2}{\sqrt{c_1 c_2}} - f}{\frac{t^4}{c_1 c_2} - 1},$$

then the relation between Γ and $\tilde{\Gamma}$ can be simply represented as

$$(3.6) \quad \tilde{\Gamma} = \frac{(c_1 c_2 f - c_1 - c_2) \Gamma - (c_1 c_2 - 1)}{(c_1 c_2 - 1) \Gamma + (f - c_1 - c_2)}.$$

So this transformation can be represented by $\text{PGL}(2)$ -action:

$$(3.7) \quad \tilde{\Gamma} = \begin{pmatrix} c_1 c_2 f - c_1 - c_2 & -(c_1 c_2 - 1) \\ c_1 c_2 - 1 & f - c_1 - c_2 \end{pmatrix} \Gamma.$$

By taking a translation of $W(E_8^{(1)})$, we obtain a nonlinear difference equation. The translation can be described by a product of simple reflections w_i 's. This representation is same as the case of $dP(A_0^{(1)})$, that is (2.8).

Now we calculate \bar{g} from g and f . Putting $\Gamma, \hat{\Gamma}$,

$$\Gamma = \frac{gt^2 - f}{t^4 - 1}, \quad \hat{\Gamma} = \frac{\bar{g} \frac{t^2}{\sqrt{\prod_{i=1}^8 c_i}} - f}{\frac{t^4}{\prod_{i=1}^8 c_i} - 1} = \frac{\bar{g} \frac{1}{t^2 \lambda} - f}{\frac{1}{t^4 \lambda^2} - 1},$$

\bar{g} can be described as follows:

$$\begin{aligned}
(3.8) \quad \dot{\Gamma} &= \begin{pmatrix} c_7 c_8 f - c_7 - c_8 & -(c_7 c_8 - 1) \\ c_7 c_8 - 1 & f - c_7 - c_8 \end{pmatrix} \begin{pmatrix} c_5 c_6 f - c_5 - c_6 & -(c_5 c_6 - 1) \\ c_5 c_6 - 1 & f - c_5 - c_6 \end{pmatrix} \\
&\cdot \begin{pmatrix} c_3 c_4 f - c_3 - c_4 & -(c_3 c_4 - 1) \\ c_3 c_4 - 1 & f - c_3 - c_4 \end{pmatrix} \begin{pmatrix} c_1 c_2 f - c_1 - c_2 & -(c_1 c_2 - 1) \\ c_1 c_2 - 1 & f - c_1 - c_2 \end{pmatrix} \Gamma \\
&= \begin{pmatrix} t^8 D & -(t^8 D - N)/f \\ (t^8 D - N)/f & N \end{pmatrix} \Gamma.
\end{aligned}$$

Here N and D are as follows:

$$\begin{aligned}
(3.9) \quad N &= f^4 - m_1 t f^3 + (m_2 t^2 - 3 - m_8 t^8) f^2 + (m_7 t^7 - m_3 t^3 + 2m_1 t) f \\
&\quad + m_8 t^8 - m_6 t^6 + m_4 t^4 - m_2 t^2 + 1,
\end{aligned}$$

$$\begin{aligned}
(3.10) \quad D &= m_8 f^4 - m_7 f^3/t + (m_6/t^2 - 3m_8 - 1/t^8) f^2 + (m_1/t^7 - m_5/t^3 + 2m_7/t) f \\
&\quad + 1/t^8 - m_2/t^6 + m_4/t^4 - m_6/t^2 + m_8,
\end{aligned}$$

where m_i is the i -th elementary symmetric function of b_j 's. This equation can be modified

$$t^8 \frac{\Gamma(\dot{\Gamma} - f) + 1}{\dot{\Gamma}(\Gamma - f) + 1} = \frac{N}{D}.$$

This equation is (3.1).

Similarly putting $\Phi, \dot{\Phi}$,

$$\Phi = \frac{f \frac{1}{t^2} - g}{\frac{1}{t^4} - 1}, \quad \dot{\Phi} = \frac{f \frac{1}{-t^2 \sqrt{\prod_{i=1}^8 d_i}} - g}{\frac{1}{t^4 \prod_{i=1}^8 d_i} - 1} = \frac{f \frac{t^2}{\lambda} - g}{\frac{t^4}{\lambda^2} - 1},$$

\underline{f} can be described as follows:

$$\begin{aligned}
(3.11) \quad \dot{\Phi} &= \begin{pmatrix} d_7 d_8 g - d_7 - d_8 & -(d_7 d_8 - 1) \\ d_7 d_8 - 1 & g - d_7 - d_8 \end{pmatrix} \begin{pmatrix} d_5 d_6 g - d_5 - d_6 & -(d_5 d_6 - 1) \\ d_5 d_6 - 1 & g - d_5 - d_6 \end{pmatrix} \\
&\cdot \begin{pmatrix} d_3 d_4 g - d_3 - d_4 & -(d_3 d_4 - 1) \\ d_3 d_4 - 1 & g - d_3 - d_4 \end{pmatrix} \begin{pmatrix} d_1 d_2 g - d_1 - d_2 & -(d_1 d_2 - 1) \\ d_1 d_2 - 1 & g - d_1 - d_2 \end{pmatrix} \Phi \\
&= \begin{pmatrix} \nabla/t^8 & -(\nabla/t^8 - \Delta)/g \\ (\nabla/t^8 - \Delta)/g & \Delta \end{pmatrix} \Phi.
\end{aligned}$$

Here ∇ and Δ are as follows:

$$\begin{aligned}
(3.12) \quad \nabla &= m_8 g^4 - m_7 t g^3 + (m_6 t^2 - 3m_8 - t^8) g^2 + (m_1 t^7 - m_5 t^3 + 2m_7 t) g \\
&\quad + t^8 - m_2 t^6 + m_4 t^4 - m_6 t^2 + m_8,
\end{aligned}$$

$$\begin{aligned}
(3.13) \quad \Delta &= g^4 - m_1 g^3/t + (m_2/t^2 - 3 - m_8/t^8) g^2 + (m_7/t^7 - m_3/t^3 + 2m_1/t) g \\
&\quad + m_8/t^8 - m_6/t^6 + m_4/t^4 - m_2/t^2 + 1.
\end{aligned}$$

This equation can be modified

$$t^8 \frac{\dot{\Phi}(\Phi - g) + 1}{\Phi(\dot{\Phi} - g) + 1} = \frac{\nabla}{\Delta}.$$

This is (3.2).

Remark 3.1. In [12], we obtain $A_0^{(1)*}$ -surface by blowing up \mathbb{P}^2 with the centers at nine points.

$$(3.14) \quad y^2 z = 4x^2(x + z),$$

$$p_i: \left(\frac{1}{\sinh^2 \theta_i} : \frac{-2 \cosh \theta_i}{\sinh^3 \theta_i} : 1 \right) \quad (i = 1, \dots, 9),$$

$$(3.15) \quad \chi(\alpha_i) = \theta_{i+1} - \theta_i \quad (i = 1, \dots, 7), \quad \chi(\alpha_8) = \theta_1 + \theta_2 + \theta_3, \quad \chi(\alpha_0) = \theta_9 - \theta_8.$$

Both parameters and coordinates correspond as follows:

$$(3.16) \quad b_1 = \exp\left(-\frac{3}{2}(\theta_1 + \theta_2)\right), \quad b_i = \exp\left(2\theta_{i+1} + \frac{1}{2}(\theta_1 + \theta_2)\right) \quad (i = 2, \dots, 8),$$

$$t = \exp\left(\frac{1}{2}(\theta_1 - \theta_2)\right),$$

$$(3.17) \quad f = \frac{-2(1 - 6e^{2\theta_1} + e^{4\theta_1})x - (1 - e^{4\theta_1})y + 16e^{2\theta_1}z}{e^{\theta_1}(2(1 + e^{2\theta_1})x - (1 - e^{2\theta_1})y)},$$

$$(3.18) \quad g = \frac{-2(1 - 6e^{2\theta_2} + e^{4\theta_2})x - (1 - e^{4\theta_2})y + 16e^{2\theta_2}z}{e^{\theta_2}(2(1 + e^{2\theta_2})x - (1 - e^{2\theta_2})y)}.$$

□

Similar to the case of $A_0^{(1)}$, $dP(A_0^{(1)*})$ has a trivial solution.

Proposition 4. $dP(A_0^{(1)*})$ has the following trivial solution:

$$(3.19) \quad f = tq \exp\left(\frac{2(\log t)^2}{\log \lambda}\right) + \frac{1}{tq \exp\left(\frac{2(\log t)^2}{\log \lambda}\right)}, \quad g = \frac{t}{q \exp\left(\frac{2(\log t)^2}{\log \lambda}\right)} + \frac{q \exp\left(\frac{2(\log t)^2}{\log \lambda}\right)}{t},$$

where q is determined by initial condition.

3.2 Linear equation

We present a special solution of $dP(A_0^{(1)*})$ in this section.

Theorem 5. *By the limiting process: $g_2 = \frac{4}{3}(1 + 3\varepsilon^2)$, $g_3 = -\frac{8}{27}(1 - 9\varepsilon^2)$ ($\varepsilon \rightarrow 0$) in (2.23), (2.24), and the change of variables and parameters: $e^{2t} \rightarrow t$, $e^{2\lambda} \rightarrow \lambda$, $f \rightarrow \frac{1}{3} \frac{f+10}{f-2}$, $g \rightarrow$*

$\frac{1}{3} \frac{g+10}{g-2}, e^{2b_i} \rightarrow b_i$, we obtain the system of equations:

$$(3.20) \quad \begin{vmatrix} fg & g & f & 1 \\ \left(b_1 t + \frac{1}{b_1 t}\right) \left(\frac{t}{b_1} + \frac{b_1}{t}\right) & \frac{t}{b_1} + \frac{b_1}{t} & b_1 t + \frac{1}{b_1 t} & 1 \\ \left(b_3 t + \frac{1}{b_3 t}\right) \left(\frac{t}{b_3} + \frac{b_3}{t}\right) & \frac{t}{b_3} + \frac{b_3}{t} & b_3 t + \frac{1}{b_3 t} & 1 \\ \left(b_5 t + \frac{1}{b_5 t}\right) \left(\frac{t}{b_5} + \frac{b_5}{t}\right) & \frac{t}{b_5} + \frac{b_5}{t} & b_5 t + \frac{1}{b_5 t} & 1 \end{vmatrix} = 0,$$

$$(3.21) \quad \begin{vmatrix} f\bar{g} & \bar{g} & f & 1 \\ \left(b_8 t + \frac{1}{b_8 t}\right) \left(\frac{\bar{t}}{b_8} + \frac{b_8}{\bar{t}}\right) & \frac{\bar{t}}{b_8} + \frac{b_8}{\bar{t}} & b_8 t + \frac{1}{b_8 t} & 1 \\ \left(b_6 t + \frac{1}{b_6 t}\right) \left(\frac{\bar{t}}{b_6} + \frac{b_6}{\bar{t}}\right) & \frac{\bar{t}}{b_6} + \frac{b_6}{\bar{t}} & b_6 t + \frac{1}{b_6 t} & 1 \\ \left(b_4 t + \frac{1}{b_4 t}\right) \left(\frac{\bar{t}}{b_4} + \frac{b_4}{\bar{t}}\right) & \frac{\bar{t}}{b_4} + \frac{b_4}{\bar{t}} & b_4 t + \frac{1}{b_4 t} & 1 \end{vmatrix} = 0,$$

where $\bar{t} = t\lambda$.

A solution of this system is a special solution of $dP(A_0^{(1)*})$ with $b_1 b_3 b_5 b_7 = 1$. \square

We can easily check that the equations (3.20), (3.21) define a special solution of $dP(A_0^{(1)*})$ similar to the case of $A_0^{(1)}$.

Transforming these equations,

$$(3.22) \quad \bar{g} = \left(f \left((b_2 + b_4 + b_6 + b_8) t - \left(\frac{1}{b_2} + \frac{1}{b_4} + \frac{1}{b_6} + \frac{1}{b_8} \right) \frac{1}{t} \right) \right. \\ \left. - (b_2 b_4 + b_2 b_6 + b_2 b_8 + b_4 b_6 + b_4 b_8 + b_6 b_8) \left(t^2 - \frac{1}{t^2} \right) + \left(t^2 \bar{t}^2 - \frac{1}{t^2 \bar{t}^2} \right) \right) \\ \left/ \left(f \left(t\bar{t} - \frac{1}{t\bar{t}} \right) - \left(\left(\frac{1}{b_2} + \frac{1}{b_4} + \frac{1}{b_6} + \frac{1}{b_8} \right) \bar{t} - (b_2 + b_4 + b_6 + b_8) \frac{1}{\bar{t}} \right) \right), \right.$$

$$(3.23) \quad f = \left(g \left(\left(\frac{1}{b_1} + \frac{1}{b_3} + \frac{1}{b_5} + \frac{1}{b_7} \right) t - (b_1 + b_3 + b_5 + b_7) \frac{1}{t} \right) \right. \\ \left. - (b_1 b_3 + b_1 b_5 + b_1 b_7 + b_3 b_5 + b_3 b_7 + b_5 b_7) \left(t^2 - \frac{1}{t^2} \right) + \left(t^4 - \frac{1}{t^4} \right) \right) \\ \left/ \left(g \left(t^2 - \frac{1}{t^2} \right) - \left((b_1 + b_3 + b_5 + b_7) t - \left(\frac{1}{b_1} + \frac{1}{b_3} + \frac{1}{b_5} + \frac{1}{b_7} \right) \frac{1}{t} \right) \right)$$

Eliminating f , we obtain a difference equation of the first order with respect to the variable g .

4 Additive type

We can obtain $A_0^{(1)**}$ -surface from $A_0^{(1)*}$ -surface by a degeneration process. By the same process the $A_0^{(1)**}$ -surface discrete Painlevé equation and linear equation also can be obtained.

4.1 Discrete Painlevé equation

We discuss about the following theorem in this section.

Theorem 6 (ORG[8]). *By the limiting process: $t \rightarrow e^{\varepsilon t}$, $\lambda \rightarrow 1 + \varepsilon\lambda$, $f \rightarrow 2 + \varepsilon^2 f$, $g \rightarrow 2 + \varepsilon^2 g$, $b_i \rightarrow e^{\varepsilon b_i}$, ($\varepsilon \rightarrow 0$), we obtain the $dP(A_0^{(1)**})$ from $dP(A_0^{(1)*})$ (3.1), (3.2):*

$$(4.1) \quad \frac{(f - \bar{g} + (2t + \lambda)^2)(f - g + 4t^2) + 4f(2t + \lambda)2t}{2t(f - \bar{g} + (2t + \lambda)^2) + (2t + \lambda)(f - g + 4t^2)} = 2 \frac{f^4 + S_2 f^3 + S_4 f^2 + S_6 f + S_8}{S_1 f^3 + S_3 f^2 + S_5 f + S_7},$$

$$(4.2) \quad \frac{(g - \underline{f} + (2t - \lambda)^2)(g - f + 4t^2) + 4g(2t - \lambda)2t}{2t(g - \underline{f} + (2t - \lambda)^2) + (2t - \lambda)(g - f + 4t^2)} = 2 \frac{g^4 + \Sigma_2 g^3 + \Sigma_4 g^2 + \Sigma_6 g + \Sigma_8}{\Sigma_1 g^3 + \Sigma_3 g^2 + \Sigma_5 g + \Sigma_7},$$

where $\bar{g} = g(t + \lambda)$, $\underline{f} = f(t - \lambda)$, $\lambda = \frac{1}{2} \sum_{i=1}^8 b_i$, S_i is the i -th elementary symmetric function of the quantities $c_j = b_j + t$ ($j = 1, \dots, 8$), Σ_i is the i -th elementary symmetric function of the quantities $d_j = t - b_j$. \square

The above theorem shows that we can obtain $dP(A_0^{(1)**})$ from $dP(A_0^{(1)*})$. But we describe geometrical construction of this discrete equation similar to the previous section.

We construct $A_0^{(1)**}$ -surface by blowing up $\mathbb{P}^1 \times \mathbb{P}^1$ at eight points. These eight points and a curve which these points lie on are as follows:

$$(4.3) \quad (f - g)^2 - 8t^2(f + g) + 16t^4 = 0,$$

$$(4.4) \quad p_i: ((b_i + t)^2, (t - b_i)^2) \quad (i = 1, \dots, 8),$$

$$(4.5) \quad \begin{aligned} \chi(\alpha_1) = -2t, \quad \chi(\alpha_2) = \frac{1}{2}(b_1 + b_2) + t, \quad \chi(\alpha_i) = \frac{1}{2}(b_i - b_{i-1}) \quad (i = 3, \dots, 7) \\ \chi(\alpha_8) = \frac{1}{2}(b_2 - b_1), \quad \chi(\alpha_0) = \frac{1}{2}(b_8 - b_7). \end{aligned}$$

Generators of affine Weyl group $W(E_8^{(1)}) = \langle w_i \ (i = 0, 1, \dots, 8) \rangle$ act on these coordinates and parameters. We give a representation of these actions in order to construct $dP(A_0^{(1)**})$.

$$\begin{aligned} w_2: & \left(\begin{array}{cccc} b_1 & b_2 & b_3 & b_4 \\ b_5 & b_6 & b_7 & b_8 \end{array}, t, g \right) \\ & \mapsto \left(\begin{array}{cccc} b_1 - 3\frac{2t+b_1+b_2}{4} & b_2 - 3\frac{2t+b_1+b_2}{4} & b_3 + \frac{2t+b_1+b_2}{4} & b_4 + \frac{2t+b_1+b_2}{4} \\ b_5 + \frac{2t+b_1+b_2}{4} & b_6 + \frac{2t+b_1+b_2}{4} & b_7 + \frac{2t+b_1+b_2}{4} & b_8 + \frac{2t+b_1+b_2}{4} \end{array}, t - \frac{2t+b_1+b_2}{4}, \tilde{g} \right), \end{aligned}$$

$$w_1: (t, f, g) \mapsto (-t, g, f), \quad w_i: (b_{i-1}, b_i) \mapsto (b_i, b_{i-1}) \quad (i = 3, \dots, 7),$$

$$w_8: (b_1, b_2) \mapsto (b_2, b_1), \quad w_0: (b_7, b_8) \mapsto (b_8, b_7),$$

where \tilde{g} is given by

$$\frac{\tilde{g} - (2t - \frac{b_1 - b_2}{2})^2}{\tilde{g} - (2t - \frac{-b_1 + b_2}{2})^2} = \frac{f - (b_2 + t)^2}{f - (b_1 + t)^2} \frac{g - (t - b_1)^2}{g - (t - b_2)^2}.$$

Notice that we can rewrite the action of w_2 for $c_i = b_i + t$ into the following form:

$$w_2: (c_1, c_2, t, g) \mapsto (-c_2, -c_1, t - \frac{c_1+c_2}{4}, \tilde{g}).$$

Here we put $\Gamma, \tilde{\Gamma}$,

$$\Gamma = \frac{f - g + 4t^2}{4t}, \quad \tilde{\Gamma} = \frac{f - \tilde{g} + (2t - \frac{c_1+c_2}{2})^2}{4t - c_1 - c_2},$$

then the relation between Γ and $\tilde{\Gamma}$ can be simply represented by PGL(2)-action:

$$(4.6) \quad \tilde{\Gamma} = \begin{pmatrix} f + c_1c_2 & -(c_1 + c_2)f \\ -(c_1 + c_2) & f + c_1c_2 \end{pmatrix} \Gamma.$$

By taking a translation of $W(E_8^{(1)})$, we obtain a nonlinear difference equation. The translation can be described by a product of simple reflections w_i 's. This representation is same as the case of $dP(A_0^{(1)})$, that is (2.8).

Now we calculate \bar{g} from g and f . Putting $\Gamma, \acute{\Gamma}$,

$$\Gamma = \frac{f - g + 4t^2}{4t}, \quad \acute{\Gamma} = \frac{f - \bar{g} + (2t - \frac{1}{2} \sum_{i=1}^8 c_i)^2}{4t - \sum_{i=1}^8 c_i} = \frac{f - \bar{g} + (2t + \lambda)^2}{-4t - 2\lambda},$$

\bar{g} can be described as follows.

$$(4.7) \quad \begin{aligned} \acute{\Gamma} &= \begin{pmatrix} f + c_7c_8 & -(c_7 + c_8)f \\ -(c_7 + c_8) & f + c_7c_8 \end{pmatrix} \begin{pmatrix} f + c_5c_6 & -(c_5 + c_6)f \\ -(c_5 + c_6) & f + c_5c_6 \end{pmatrix} \\ &\cdot \begin{pmatrix} f + c_3c_4 & -(c_3 + c_4)f \\ -(c_3 + c_4) & f + c_3c_4 \end{pmatrix} \begin{pmatrix} f + c_1c_2 & -(c_1 + c_2)f \\ -(c_1 + c_2) & f + c_1c_2 \end{pmatrix} \Gamma \\ &= \begin{pmatrix} f^4 + S_2f^3 + S_4f^2 + S_6f + S_8 & -(S_1f^3 + S_3f^2 + S_5f + S_7)f \\ -(S_1f^3 + S_3f^2 + S_5f + S_7) & f^4 + S_2f^3 + S_4f^2 + S_6f + S_8 \end{pmatrix} \Gamma, \end{aligned}$$

where S_i is the i -th elementary symmetric function of c_j 's. This equation can be modified

$$\frac{\acute{\Gamma}\Gamma - f}{\acute{\Gamma} - \Gamma} = \frac{f^4 + S_2f^3 + S_4f^2 + S_6f + S_8}{S_1f^3 + S_3f^2 + S_5f + S_7}.$$

This equation is (4.1).

Similarly putting $\Phi, \acute{\Phi}$,

$$\Phi = \frac{g - f + 4t^2}{-4t}, \quad \acute{\Phi} = \frac{g - \underline{f} + (-2t + \frac{1}{2} \sum_{i=1}^8 d_i)^2}{-4t + \sum_{i=1}^8 d_i} = \frac{g - \underline{f} + (2t - \lambda)^2}{4t - 2\lambda},$$

\underline{f} can be described as follows.

$$\begin{aligned}
(4.8) \quad \underline{\Phi} &= \begin{pmatrix} g + d_7d_8 & (d_7 + d_8)g \\ d_7 + d_8 & g + d_7d_8 \end{pmatrix} \begin{pmatrix} g + d_5d_6 & (d_5 + d_6)g \\ d_5 + d_6 & g + d_5d_6 \end{pmatrix} \\
&\cdot \begin{pmatrix} g + d_3d_4 & (d_3 + d_4)g \\ d_3 + d_4 & g + d_3d_4 \end{pmatrix} \begin{pmatrix} g + d_1d_2 & (d_1 + d_2)g \\ d_1 + d_2 & g + d_1d_2 \end{pmatrix} \Phi \\
&= \begin{pmatrix} g^4 + S_2g^3 + \Sigma_4g^2 + \Sigma_6g + \Sigma_8 & (\Sigma_1g^3 + \Sigma_3g^2 + \Sigma_5g + \Sigma_7)g \\ \Sigma_1g^3 + \Sigma_3g^2 + \Sigma_5g + \Sigma_7 & g^4 + \Sigma_2g^3 + \Sigma_4g^2 + \Sigma_6g + \Sigma_8 \end{pmatrix} \Phi,
\end{aligned}$$

where $d_i = t - b_i$ ($i = 1, \dots, 8$), and Σ_i is the i -th elementary symmetric function of d_j 's. This equation can be modified

$$\frac{\underline{\Phi}\Phi - g}{\Phi - \underline{\Phi}} = \frac{g^4 + \Sigma_2g^3 + \Sigma_4g^2 + \Sigma_6g + \Sigma_8}{\Sigma_1g^3 + \Sigma_3g^2 + \Sigma_5g + \Sigma_7}.$$

This is (4.2).

Remark 4.1. In [12], we obtain $A_0^{(1)**}$ -surface by blowing up \mathbb{P}^2 with the centers at nine points.

$$\begin{aligned}
(4.9) \quad &y^2z = 4x^3, \\
&p_i: (a_i : -2 : a_i^3) \quad (i = 1, \dots, 9), \\
&\chi(\alpha_i) = (a_{i+1} - a_i)/\lambda \quad (i = 1, \dots, 7), \quad \chi(\alpha_8) = (a_1 + a_2 + a_3)/\lambda, \\
(4.10) \quad &\chi(\alpha_0) = (a_9 - a_8)/\lambda, \quad \lambda = \sum_{i=1}^9 a_i.
\end{aligned}$$

Both parameters and coordinates correspond as follows:

$$(4.11) \quad b_1 = -\frac{3}{2}(a_1 + a_2), \quad b_i = 2a_{i+1} + \frac{1}{2}(a_1 + a_2) \quad (i = 2, \dots, 8), \quad t = \frac{1}{2}(a_1 - a_2),$$

$$(4.12) \quad f = \frac{-6a_1^2x + a_1^3y + 8z}{2x + a_1y}, \quad g = \frac{-6a_2^2x + a_2^3y + 8z}{2x + a_2y}.$$

□

Proposition 7. $dP(A_0^{(1)**})$ has the following trivial solution:

$$(4.13) \quad f = (q + 2t^2/\lambda + t)^2, \quad g = (-q - 2t^2/\lambda + t)^2,$$

where q is determined by initial condition.

4.2 Linear equation

We present a special solution of $dP(A_0^{(1)**})$ in this section.

Theorem 8. *By the limiting process: $t \rightarrow e^{\varepsilon t}$, $\lambda \rightarrow 1 + \varepsilon\lambda$, $f \rightarrow 2 + \varepsilon^2 f$, $g \rightarrow 2 + \varepsilon^2 g$, $b_i \rightarrow e^{\varepsilon b_i}$ ($\varepsilon \rightarrow 0$) in (3.20), (3.21), we obtain the system of equations:*

$$(4.14) \quad \begin{vmatrix} fg & g & f & 1 \\ (b_1 + t)^2(t - b_1)^2 & (t - b_1)^2 & (b_1 + t)^2 & 1 \\ (b_3 + t)^2(t - b_3)^2 & (t - b_3)^2 & (b_3 + t)^2 & 1 \\ (b_5 + t)^2(t - b_5)^2 & (t - b_5)^2 & (b_5 + t)^2 & 1 \end{vmatrix} = 0,$$

$$(4.15) \quad \begin{vmatrix} f\bar{g} & \bar{g} & f & 1 \\ (b_8 + t)^2(\bar{t} - b_8)^2 & (\bar{t} - b_8)^2 & (b_8 + t)^2 & 1 \\ (b_6 + t)^2(\bar{t} - b_6)^2 & (\bar{t} - b_6)^2 & (b_6 + t)^2 & 1 \\ (b_4 + t)^2(\bar{t} - b_4)^2 & (\bar{t} - b_4)^2 & (b_4 + t)^2 & 1 \end{vmatrix} = 0,$$

where $\bar{t} = t + \lambda$.

A solution of this system is a special solution of $dP(A_0^{(1)**})$ with $b_1 + b_3 + b_5 + b_7 = 0$. \square

We can easily check that the equations (4.14),(4.15) define a special solution of $dP(A_0^{(1)**})$.

Transforming these equations,

$$(4.16) \quad \bar{g} = \frac{-A_{2468}(-\frac{t+\bar{t}}{2})f + B_{2468}(\frac{t+\bar{t}}{2})}{128(t+\bar{t})f - A_{2468}(\frac{t+\bar{t}}{2})}, \quad f = \frac{A_{1357}(t)g + B_{1357}(t)}{256tg + A_{1357}(-t)},$$

where

$$\begin{aligned} A_{ijkl}(t) &= -8(b_i + b_j - b_k - b_l)(b_i - b_j + b_k - b_l)(b_i - b_j - b_k + b_l) \\ &\quad + 16((b_i + b_j - b_k - b_l)^2 + (b_i - b_j + b_k - b_l)^2 + (b_i - b_j - b_k + b_l)^2)t - 256t^3, \\ B_{ijkl}(t) &= -(3b_i - b_j - b_k - b_l)(3b_j - b_i - b_k - b_l)(3b_k - b_i - b_j - b_l)(3b_l - b_i - b_j - b_k)t \\ &\quad - 32((b_i + b_j - b_k - b_l)^2 + (b_i - b_j + b_k - b_l)^2 + (b_i - b_j - b_k + b_l)^2)t^3 + 768t^5. \end{aligned}$$

Eliminating f , we obtain a difference equation of the first order with respect to the variable g .

5 Riccati solution of $dP(A_1^{(1)})$

We can obtain $A_1^{(1)}$ -surface from $A_0^{(1)*}$ -surface by a degeneration process. By the same process the $A_1^{(1)}$ -surface discrete Painlevé equation and linear equation also can be obtained.

Theorem 9. *By the limiting process: $t \rightarrow \sqrt{t}$, $\lambda \rightarrow \sqrt{\lambda}$, $f \rightarrow \frac{fp^{1/4}}{\varepsilon\sqrt{t}}$, $g \rightarrow \frac{\sqrt{t}p^{1/4}}{\varepsilon g}$, $b_i \rightarrow b_i p^{1/4}/\varepsilon$ ($i = 1, \dots, 4$), $b_i \rightarrow \varepsilon p^{1/4}/b_i$ ($i = 5, \dots, 8$) ($\varepsilon \rightarrow 0$) in (3.20), (3.21), where $p = \sqrt{b_1 b_2 b_3 b_4 / (b_5 b_6 b_7 b_8)}$, we obtain the system of equations:*

$$(5.1) \quad \begin{vmatrix} fg & g & f & 1 \\ t^2 & t/b_1 & b_1 t & 1 \\ t^2 & t/b_3 & b_3 t & 1 \\ 1 & 1/b_5 & b_5 & 1 \end{vmatrix} = 0, \quad \begin{vmatrix} f\bar{g} & \bar{g} & f & 1 \\ 1 & 1/b_8 & b_8 & 1 \\ 1 & 1/b_6 & b_6 & 1 \\ t\bar{t} & \bar{t}/b_4 & b_4 t & 1 \end{vmatrix} = 0,$$

where $\bar{t} = t\lambda$.

A solution of this system is a special solution of $dP(A_1^{(1)})$ with $b_5 b_7 = b_1 b_3 p$. \square

We introduce this theorem in this section.

By the limiting procedure on the $A_0^{(1)*}$ -surface discrete Painlevé equation (3.1), (3.2), we obtain the $A_1^{(1)}$ -surface discrete Painlevé equation ([8]).

$$(5.2) \quad \frac{(f\bar{g} - t\bar{t})(fg - t^2)}{(f\bar{g} - 1)(fg - 1)} = \frac{(f - b_1 t)(f - b_2 t)(f - b_3 t)(f - b_4 t)}{(f - b_5)(f - b_6)(f - b_7)(f - b_8)},$$

$$(5.3) \quad \frac{(fg - t^2)(fg - t\bar{t})}{(fg - 1)(\underline{fg} - 1)} = \frac{(g - t/b_1)(g - t/b_2)(g - t/b_3)(g - t/b_4)}{(g - 1/b_5)(g - 1/b_6)(g - 1/b_7)(g - 1/b_8)}.$$

We can construct $dP(A_1^{(1)})$ by geometrical approach similar to the above discussions, but we only show the construction of surface. Here $A_1^{(1)}$ -surface is obtained by blowing up $\mathbb{P}^1 \times \mathbb{P}^1$ at eight points. These eight points and a curve which these points lie on are as follows:

$$(5.4) \quad (fg - t^2)(fg - 1) = 0,$$

$$p_i: \left(b_i t, \frac{t}{b_i} \right) \quad (i = 1, \dots, 4), \quad p_i: \left(b_i, \frac{1}{b_i} \right) \quad (i = 5, \dots, 8).$$

By the same limiting procedure on the system of equations (3.20), (3.21), we obtain the system of equations (5.1), or

$$(5.5) \quad \bar{g} = \frac{f(1 - t\bar{t}) + t(\bar{t}(b_6 + b_8) - b_2 - b_4)}{g(b_6 + b_8 - t(b_2 + b_4)) + b_6 b_8 (t\bar{t} - 1)},$$

$$(5.6) \quad f = \frac{g b_5 b_7 (t^2 - 1) + t(b_1 + b_3 - t(b_5 + b_7))}{g(t(b_1 + b_3) - b_5 - b_7) + 1 - t^2}.$$

These equations coincide with the equations in [11].

6 Discussion

In the paper, we derive the linear equations as the special solutions of discrete Painlevé equations. So that we show that these equations belong to the hypergeometric family, we

will seek the series solutions of them like the q -hypergeometric series. If there exists the series solutions of $dP(A_0^{(1)})$, it should be called elliptic-hypergeometric series.

Acknowledgement. The authors would like to thank K. Okamoto and M. Jimbo for discussions and advice. The authors are also grateful to Y. Ohta, T. Takebe, T. Takenawa and T. Tsuda for useful comments.

References

- [1] B. Grammaticos and A. Ramani, On a novel q -discrete analogue of the Painlevé VI equation, *Phys. Lett. A* **257** (1999), no.5-6, 288–292.
- [2] B. Grammaticos, A. Ramani and V. G. Papageorgiou, Do integrable mappings have the Painlevé property?, *Phys. Rev. Lett.* **67** (1991), 1825–1828.
- [3] B. Grammaticos, Y. Ohta, A. Ramani and H. Sakai, Degeneration through coalescence of the q -Painlevé VI equations, *J. Phys. A : Math. Gen.* **31** (1998), 3545–3558.
- [4] M. Jimbo and H. Sakai, A q -analog of the sixth Painlevé equation, *Lett. Math. Phys.* **38** (1996), 145–154.
- [5] V. Kac, Infinite dimensional Lie algebras, 3rd ed., *Cambridge University Press* (1990).
- [6] M. Noumi and Y. Yamada, Affine Weyl groups, discrete dynamical systems and Painlevé equations, *Comm. Math. Phys.* **199** (1998), no.2, 281–295.
- [7] K. Okamoto, Studies on the Painlevé equations I. *Annali di Matematica pura ed applicata* **CXLVI** (1987), 337–381; II. *Jap. J. Math.* **13** (1987), 47–76; III. *Math. Ann.* **275** (1986), 221–255; IV. *Funkcial. Ekvac. Ser. Int.* **30** (1987), 305–332.
- [8] Y. Ohta, A. Ramani and B. Grammaticos, An affine Weyl group approach to the 8-parameter discrete Painlevé equation, *J. Phys. A : Math. Gen.* **34** (2001), 10523–10532.
- [9] P. Painlevé, Sur les équations différentielles du second ordre dont l’intégrale générale est uniforme, *Oeuvre t.* **III**, 187–271.
- [10] A. Ramani, B. Grammaticos and J. Hietarinta, Discrete versions of the Painlevé equations, *Phys. Rev. Lett.* **67** (1991), 1829–1832.
- [11] A. Ramani, B. Grammaticos, T. Tamizhmani and K. M. Tamizhmani, Special function solutions of the discrete Painlevé equations, *Comput. Math. Appl.* **42** (2001), no.3-5, 603–614.

- [12] H. Sakai, Rational Surfaces Associated with Affine Root Systems and Geometry of the Painlevé Equations, *Comm. Math. Phys.* **220** (2001), 165–229.